
ACCELERATED ZERO-ORDER SGD UNDER HIGH-ORDER SMOOTHNESS AND OVERPARAMETERIZED REGIME

Georgii Bychkov

Lomonosov MSU, ISP RAS
Moscow, Russia

georgy.bychkov@graphics.cs.msu.ru

Darina Dvinskikh

HSE University, MIPT, ISP RAS
Moscow, Russia

dmdvinskikh@hse.ru

Anastasia Antsiferova

MSU Institute for Artificial Intelligence, ISP RAS
Moscow, Russia

aantsiferova@graphics.cs.msu.ru

Alexander Gasnikov

Innopolis University, MIPT, IITP RAS
Innopolis, Republic of Tatarstan, Russia

gasnikov@yandex.ru

Aleksandr Lobanov

MIPT, Skoltech, ISP RAS
Dolgoprudny, Moscow Oblast, Russia
lobbsasha@mail.ru

November 22, 2024

ABSTRACT

We present a novel gradient-free algorithm to solve a convex stochastic optimization problem, such as those encountered in medicine, physics, and machine learning (e.g., adversarial multi-armed bandit problem), where the objective function can only be computed through numerical simulation, either as the result of a real experiment or as feedback given by the function evaluations from an adversary. Thus we suppose that only a black-box access to the function values of the objective is available, possibly corrupted by adversarial noise: deterministic or stochastic. The noisy setup can arise naturally from modeling randomness within a simulation or by computer discretization, or when exact values of function are forbidden due to privacy issues, or when solving non-convex problems as convex ones with an inexact function oracle. By exploiting higher-order smoothness, fulfilled, e.g., in logistic regression, we improve the performance of zero-order methods developed under the assumption of classical smoothness (or having a Lipschitz gradient). The proposed algorithm enjoys optimal oracle complexity and is designed under an overparameterization setup, i.e., when the number of model parameters is much larger than the size of the training dataset. Overparameterized models fit to the training data perfectly while also having good generalization and outperforming underparameterized models on unseen data. We provide convergence guarantees for the proposed algorithm under both types of noise. Moreover, we estimate the maximum permissible adversarial noise level that maintains the desired accuracy in the Euclidean setup, and then we extend our results to a non-Euclidean setup. Our theoretical results are verified on the logistic regression problem.

1 Introduction

In this paper we focus on the following stochastic convex optimization problem

$$\min_{x \in \mathbb{R}^d} \{f(x) := \mathbb{E}_\xi f(x, \xi)\}, \tag{1}$$

where f is convex in x , and \mathbb{E}_ξ is the expectation, with respect to ξ with unknown distribution.

Motivated by different applications where gradient information is unavailable or calculating gradients is prohibitively expensive, we seek to solve (1) only by accessing noisy evaluations of $f(x, \xi)$ [1, 2, 3, 4], possibly adversarial noisy. Methods that query only function values for given inputs, rather than gradients, are known as zero-order, or gradient-free methods, [1, 5, 6, 7, 8, 9, 10, 11, 12] and are actively studied due to the vast number of applications. Indeed, in medicine, biology, and physics, the objective function can only be computed through numerical simulation or as the result of a real experiment, i.e., automatic differentiation cannot be used to compute function derivatives. We also mention classical problem of adversarial multi-armed bandit [13, 14, 15], where a learner receives a feedback given by the function evaluations from an adversary. Usually, a black-box function we are optimizing, is affected by stochastic or computational noise. This noise can arise naturally from modeling randomness within a simulation or by computer discretization. The Noisy setup is also useful in cases where exact values of function are forbidden due to privacy issues. Nowadays, there are a lot of adversarial attacks that influence oracle output. An attacker can implement a target function with backdoor attacks [16]. Thus, we address two concepts of adversarial noises: *adversarial deterministic noise* [17] and *adversarial stochastic noise* [18]. Adversarial stochastic noise makes the gradient approximation unbiased, so it will only accumulate in the variance. Adversarial deterministic noise, on the other hand, accumulates in both bias and variance. Adversarial deterministic noise has a stronger effect on the optimization process than adversarial stochastic noise.

There is a large amount of literature on zero-order optimization, to name a few we mention [7] for the smooth case (differential and with Lipschitz-continuous gradients), for the non-smooth case we refer to [8]. These papers use the exact function evaluations that can be infeasible in some applications. Moreover, non-convex problems are often considered as convex ones with an inexact function oracle. Zero-order methods with noisy function evaluations are studied in [1, 2, 19, 11, 3, 4]. In contrast to these works we consider overparametrized regime [20, 21, 22], i.e., when the number of model parameters is much larger than the size of the training dataset. Overparametrized models fit to the training data perfectly while also having good generalization and outperforming underparameterized models on unseen data. For instance, as stated in State of AI report, DeepMind researchers found that language models are significantly undertrained and there is a lack of labeled training data [23, 24]. Overparametrized models were studied in [25] for smooth and convex functions. In contrast to [25], we consider higher-order smoothness [26, 27, 28], fulfilled e.g., in logistic regression, to improve the performance of zero-order methods. To tackle higher-order smoothness, we use kernel-based gradient approximation [27]. We summarize our contributions as follows:

- We propose a novel gradient-free algorithm enjoying optimal oracle complexity under certain batch sizes.
- We estimate the maximum permissible adversarial noise level that maintains the desired accuracy in the Euclidean setup, and then we extend our results to a non-Euclidean setup.
- We confirm our theoretical results on machine learning problem. The numerical experiments demonstrate the superiority of the proposed method over AZO-SGD [25], designed for smooth objective functions.

Paper organization. The structure of the paper is the following. In Section 2, we provide background material, notation, and assumptions. In Section 3, we present the main algorithm to solve (1) along with the convergence guarantees. Section 4 provides the numerical experiments demonstrating the superiority of our novel algorithm on the logistic regression. In Section 5, we summarize the paper and formulate conclusions.

2 Preliminaries

In this section, we provide all the notation and assumptions necessary to prove the convergence of our novel algorithm.

Notation. We use $\|x\|_p$ to denote the ℓ_p -norm, for the Euclidean norm we use the shorthand notation $\|x\|_2 = \|x\|$. We use $\langle x, y \rangle := \sum_{k=1}^d x_k y_k$ as the standard inner product of $x, y \in \mathbb{R}^d$, where x_i and y_i are the i -th component of x and y respectively. We denote by $S_p^d(r) := \{x \in \mathbb{R}^d : \|x\|_p = r\}$ and $B_p^d(r) := \{x \in \mathbb{R}^d : \|x\|_p \leq r\}$ the sphere and the ball in the ℓ_p -norm respectively. Symbol \lesssim stands for the asymptotic inequality.

Assumption 1 (L -smoothness). $f(x, \xi)$ is L -Lipschitz smooth, or has L -Lipschitz continuous gradient, if $f(x, \xi)$ is continuously differentiable, with respect to x , and its gradient satisfies Lipschitz condition for any ξ and $x, y \in \mathbb{R}^d$

$$\|\nabla f(x, \xi) - \nabla f(y, \xi)\| \leq L\|x - y\|.$$

The next assumption is an extension of Assumption 1.

Assumption 2 (Higher-order smoothness). Let l denote the maximal integer number strictly less than β . Let $\mathcal{F}_\beta(L)$ denote the set of all functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ which are differentiable l times and for any ξ and $x, y \in \mathbb{R}^d$ the Hölder-type

condition:

$$\left| f(x, \xi) - \sum_{0 \leq |n| \leq l} \frac{1}{n!} D^n f(y, \xi) (x - y)^n \right| \leq L_\beta \|x - y\|,$$

where $L_\beta > 0$, the sum is over multi-index $n = (n_1, \dots, n_d) \in \mathbb{N}^d$, the notation $n! = n_1! \dots n_d!$, and

$$D^n f(y, \xi) v^n = \frac{\partial^{|n|} f(y, \xi)}{\partial^{n_1} x_1 \dots \partial^{n_d} x_d} v_1^{n_1} \dots v_d^{n_d},$$

where $|n| = n_1 + \dots + n_d$, for all $v = (v_1, \dots, v_d) \in \mathbb{R}^d$.

When $\beta = 2$, Assumption 2 turns into Assumption 1.

The next assumption means that in overparametrized regime the variance of the stochastic gradient estimates at optimal point $x^* = \arg \min_{x \in \mathbb{R}^d} f(x)$ can be upper bounded by using the minimum value $f(x^*)$.

Assumption 3 ([29], Lemma 3). There exists $\sigma_*^2 \geq 0$ such that

$$\mathbb{E}[\|\nabla f(x^*, \xi) - \nabla f(x^*)\|^2] \leq \sigma_*^2.$$

We will use this assumption for the gradient estimates.

3 Main results: the AZO-SGD-HS algorithm

In this section, we present a novel algorithm, named AZO-SGD-HS (see Algorithm 1) and provide its theoretical guarantees of convergence when the adversarial noise is deterministic and when it is stochastic. We also report on the maximum permissible adversarial noise level Δ .

3.1 Deterministic adversarial noise

Let us suppose that a zero-order oracle is corrupted by a deterministic noise $\delta(x)$, bounded by a some constant $\Delta : |\delta(x)| \leq \Delta$ for all $x \in \mathbb{R}^n$

$$f_\delta(x, \xi) = f(x, \xi) + \delta(x). \quad (2)$$

Then by using two-point noisy zero-order oracle (2) the gradient can be estimated as follows

$$\mathbf{g}(x, \mathbf{e}, r, \xi) = \frac{d}{2h} (f_\delta(x + h r \mathbf{e}, \xi) - f_\delta(x - h r \mathbf{e}, \xi)) K(r) \mathbf{e}, \quad (3)$$

where $h > 0$ is a smoothing parameter, $\mathbf{e} \in S_2^d(1)$ is a vector uniformly distributed on the Euclidean unit sphere, $K : [-1, 1] \rightarrow \mathbb{R}$ is a kernel function that satisfies

$$E[K(u)] = 0, \quad E[uK(u)] = 1, \quad E[|u|^\beta |K(u)|] < \infty \text{ and for all } j = 2, \dots, l \quad E[u_j K(u)] = 0.$$

The next theorem states the convergence guarantee for AZO-SGD-HS with two-point zero-order oracle (2). We will use the shorthand notations $\kappa_\beta = \int |u|^\beta |K(u)| du$ and $\kappa = \int |K(u)|^2 du$.

Theorem 1. *Let $f(\cdot, \cdot)$ satisfy Assumption 2 with parameter β . Let smoothing parameter be $h \leq (\varepsilon / (\kappa_\beta LR))^{1/(\beta-1)}$ then Assumption 3 is satisfied with gradient approximation (4). Let the maximum admissible level of adversarial deterministic noise be*

$$\Delta \leq \frac{\varepsilon^{1+1/(\beta-1)}}{d \kappa_\beta^{1/(\beta-1)} R^{1+1/(\beta-1)} L^{1/(\beta-1)}}.$$

Let x_N^{ag} be the output of the AZO-SGD-HS then

$$\mathbb{E}[f(x_N^{ag}) - f(x^*)] \leq \varepsilon$$

in at most the following number of iterations N and oracle calls T , where R will denote a radius of a solution x^* : $\|x^* - x_0\| \leq R$, and x_0 is a starting point:

- $B = 1$

$$N = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d \kappa \sigma_*^2 R^2}{\varepsilon^2} \right) \right), \quad T = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d \kappa \sigma_*^2 R^2}{\varepsilon^2} \right) \right)$$

Algorithm 1 Accelerated zero-order SGD for high-order smoothness (AZO-SGD-HS)**1: Inputs:**

Starting points $x_0^{ag} = x_0 = \vec{0} \in \mathbb{R}^d$, number of iterations N , batch size B , smoothing parameter $h > 0$, kernel $K(\cdot)$.

2: for $k = 0$ to $N - 1$ **do**

3: $\beta_k = 1 + \frac{k}{6}$; $\gamma_k = \gamma(k + 1)$ where $\gamma = \min \left\{ \frac{1}{12L}, \frac{B}{24L(N+1)}, \sqrt{\frac{BR^2}{L\beta^*N^3}} \right\}$.

4: $x_k^{md} = \beta_k^{-1}x_k + (1 - \beta_k^{-1})x_k^{ag}$.

5: Sample $\mathbf{e}_i \in S_2^d(1)$, $r_i = \text{Uniform}[-1, 1]$ and ξ_i for $i = 1, \dots, B$.

6: Compute batched gradient approximation

$$g_k = \frac{1}{B} \sum_{i=1}^B \mathbf{g}(x_k^{md}, \mathbf{e}_i, r_i, \xi), \quad (4)$$

where \mathbf{g} is defined in

- deterministic noise: (3)
- stochastic noise: (6)

7: $\bar{x}_{k+1} = x_k - \gamma g_k$.

8: $x_{k+1} = \min \left\{ 1, \frac{R}{\|\bar{x}_{k+1}\|} \right\} \bar{x}_{k+1}$.

9: $x_{k+1}^{ag} = \beta_k^{-1}x_{k+1} + (1 - \beta_k^{-1})x_k^{ag}$.

10: end for

11: return x_N^{ag} .

- $N > B > 1$

$$N = \mathcal{O} \left(\max \left(\frac{LR^2}{B\varepsilon}, \sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2R^2}{B\varepsilon^2} \right) \right)$$

$$T = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, B\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2R^2}{\varepsilon^2} \right) \right)$$

- $B = N$

$$N = \mathcal{O} \left(\max \left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{\sqrt{d}\kappa\sigma_*R}{\varepsilon} \right) \right), T = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2R^2}{\varepsilon^2} \right) \right)$$

- $B > N$

$$N = \mathcal{O} \left(\sqrt{\frac{LR^2}{\varepsilon}} \right), T = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2R^2}{\varepsilon^2}, \frac{\kappa_\beta^{2/(\beta-1)} \kappa d^2 L^{2/(\beta-1)} \Delta^2 R^{2+2/(\beta-1)}}{\varepsilon^{2+2/(\beta-1)}} \right) \right).$$

Sketch of the proof. To find the upper bounds for the maximum admissible level of adversarial noise Δ , smoothing parameter h , the asymptotic for the number of iteration N , and the total number of oracle calls T . The proof consists in upper bounding the bias and second moment of the gradient approximation.

Step 1. (bias of gradient approximation).

We calculate the bias of the gradient approximation $\|\mathbb{E}[\mathbf{g}(x_k, \mathbf{e}, r, \xi)] - \nabla f(x_k)\|$.

$$\begin{aligned}
b &= \|\mathbb{E}[\mathbf{g}(x_k, \mathbf{e}, r, \xi)] - \nabla f(x_k)\| = \left\| \mathbb{E} \left[\frac{d}{2h} (f_\delta(x_k + h\mathbf{r}\mathbf{e}, \xi) - f_\delta(x_k - h\mathbf{r}\mathbf{e}, \xi)) K(r) \mathbf{e} \right] - \nabla f(x_k) \right\| \\
&\stackrel{\textcircled{1}}{=} \left\| \mathbb{E} \left[\frac{d}{2h} (f(x_k + h\mathbf{r}\mathbf{e}, \xi)) K(r) \mathbf{e} + \delta(x_k + h\mathbf{e}) \right] - \nabla f(x_k) \right\| \\
&\stackrel{\textcircled{2}}{\leq} \left\| \mathbb{E} \left[\frac{d}{2h} (f(x_k + h\mathbf{r}\mathbf{e}, \xi)) K(r) \mathbf{e} \right] - \nabla f(x_k) \right\| + \frac{d\Delta}{h} \\
&\stackrel{\textcircled{3}}{=} \|\mathbb{E}[(\nabla f(x_k + h\mathbf{r}\mathbf{u}, \xi) r K(r)) - \nabla f(x_k)]\| + \frac{d\Delta}{h} \\
&\stackrel{\textcircled{4}}{=} \sup_{z \in S_2^d(1)} \mathbb{E}[(\nabla_z f(x_k + h\mathbf{r}\mathbf{u}, \xi) - \nabla_z f(x_k)) r K(r)] + \frac{d\Delta}{h} \\
&\stackrel{\textcircled{5}}{\leq} \kappa_\beta h^{\beta-1} \frac{L}{(l-1)!} \mathbb{E}[\|\mathbf{u}\|^{\beta-1}] + \frac{d\Delta}{h} \leq \kappa_\beta h^{\beta-1} \frac{L}{(l-1)!} \frac{d}{d+\beta-1} + \frac{d\Delta}{h} \lesssim \kappa_\beta L h^{\beta-1} + \frac{d\Delta}{h},
\end{aligned}$$

where $\mathbf{u} \in B_2^d(1)$, $\textcircled{1}$: distribution of \mathbf{e} is symmetric; $\textcircled{2}$: triangle inequality and bounded noise $|\delta(x)| \leq \Delta$; $\textcircled{3}$: a version of the Stokes' theorem [30] (see Section 13.3.5, Exercise 14a); $\textcircled{4}$: norm of the gradient is the supremum of directional derivatives $\nabla_z f(x) = \lim_{\varepsilon \rightarrow 0} \frac{f(x+\varepsilon z) - f(x)}{\varepsilon}$; $\textcircled{5}$: Taylor expansion.

Step 2. (bounding the second moment of gradient approximation).

In this step, we calculate the second moment of the gradient approximation $\mathbb{E}\|\mathbf{g}(x^*, \mathbf{e}, r, \xi)\|^2$.

$$\begin{aligned}
\zeta^2 &= \mathbb{E}[\|\mathbf{g}(x^*, \mathbf{e}, r, \xi)\|^2] = \mathbb{E} \left[\left\| \frac{d}{2h} (f_\delta(x^* + h\mathbf{r}\mathbf{e}, \xi) - f_\delta(x^* - h\mathbf{r}\mathbf{e}, \xi)) K(r) \mathbf{e} \right\|^2 \right] \\
&= \frac{d^2}{4h^2} \mathbb{E}[(f_\delta(x^* + h\mathbf{r}\mathbf{e}, \xi) - f_\delta(x^* - h\mathbf{r}\mathbf{e}, \xi)) K(r)]^2 \\
&= \frac{d^2}{4h^2} \mathbb{E}[(f(x^* + h\mathbf{r}\mathbf{e}, \xi) - f(x^* - h\mathbf{r}\mathbf{e}, \xi) + \delta(x^* + h\mathbf{r}\mathbf{e}) - \delta(x^* - h\mathbf{r}\mathbf{e}))^2 (K(r))^2] \\
&\stackrel{\textcircled{1}}{\leq} \frac{d^2 \kappa}{2h^2} (\mathbb{E}(f(x^* + h\mathbf{r}\mathbf{e}, \xi) - f(x^* - h\mathbf{r}\mathbf{e}, \xi))^2 + 2\Delta^2) \\
&\stackrel{\textcircled{2}}{\leq} \frac{d^2 \kappa}{2h^2} \left(\frac{h^2}{d} \mathbb{E}(\|\nabla f(x^* + h\mathbf{r}\mathbf{e}, \xi) + \nabla f(x^* - h\mathbf{r}\mathbf{e}, \xi)\|^2) + 2\Delta^2 \right) \\
&= \frac{d^2 \kappa}{2h^2} \left(\frac{h^2}{d} \mathbb{E}(\|\nabla f(x^* + h\mathbf{r}\mathbf{e}, \xi) + \nabla f(x^* - h\mathbf{r}\mathbf{e}, \xi) \pm 2\nabla f(x^*, \xi)\|^2) + 2\Delta^2 \right) \\
&\stackrel{\textcircled{3}}{\leq} 4d\kappa \|\nabla f(x^*, \xi)\|^2 + 4d\kappa L^2 h^2 \mathbb{E}[\|\mathbf{e}\|^2] + \frac{d^2 \kappa \Delta^2}{h^2} \\
&\stackrel{\textcircled{4}}{\leq} 4d\kappa \sigma_*^2 + 4d\kappa L^2 h^2 \mathbb{E}[\|\mathbf{e}\|^2] + \frac{d^2 \kappa \Delta^2}{h^2} = 4d\kappa \sigma_*^2 + 4d\kappa L^2 h^2 + \frac{d^2 \kappa \Delta^2}{h^2},
\end{aligned}$$

$\textcircled{1}$: inequality of squared norm of the sum, inequality between positive random variables and independence of the noise; $\textcircled{2}$: Wirtinger-Poincaré inequality; $\textcircled{3}$: L-smoothness function; $\textcircled{4}$: overparametrization assumption. □

For the complete proof of this theorem, we refer to Appendix A.1.

The Theorem 1 shows that the iteration complexity is seriously affected by chosen B , and a large enough value can make the algorithm have an effective iterative complexity of $\mathcal{O}\left(\sqrt{LR^2/\varepsilon}\right)$. Total oracle calls are generally worse than AZO-SGD due to the large B needed for optimal iteration complexity. Smoothing parameter value increases with the growth of β that is caused by use of kernel approximation that efficiently solves the task. The theorem gives an estimation of the maximum allowable level of adversarial noise. The bound matches the corresponding value from AZO-SGD in the case of $\beta = 2$. Moreover, with the growth of smoothness, the restriction on the noise level becomes weaker, allowing for better-quality solutions.

Remark 1. Theorem 1 can be generalized to a more general class of problems by using the ℓ_p -norm. The number of iterations N , the maximum admissible level of adversarial noise Δ and the smoothing parameter h remain the same. The total number of oracle calls for $B > N$ changes to

$$\begin{aligned} T &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{\kappa \kappa'(p, d) d \sigma_*^2 R^2}{\varepsilon^2}, \frac{\kappa_\beta^{2/(\beta-1)} \kappa'(p, d) \kappa d^2 \Delta^2 L^{2/(\beta-1)} R^{2+2/(\beta-1)}}{\varepsilon^{2+2/(\beta-1)}} \right) \right) \\ &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{\kappa \min\{q, \ln d\} d^{2-2/p} \sigma_*^2 R^2}{\varepsilon^2}, \frac{\min\{q, \ln d\} \kappa_\beta^{2/(\beta-1)} \kappa d^{3-2/p} \Delta^2 L^{2/(\beta-1)} R^{2+2/(\beta-1)}}{\varepsilon^{2+2/(\beta-1)}} \right) \right). \end{aligned}$$

This remark means that oracle complexity can be improved. For instance, optimizing in the ℓ_1 -norm can reduce the number of iterations by a factor of $d/\ln d$, which can be a significant speedup for large d . The proof of this remark is given in Appendix A.2.

3.2 Stochastic adversarial noise

Let us suppose that a zero-order oracle is corrupted by a stochastic noise δ . At each iteration we request two function evaluations on the inputs x_t and x'_t

$$f_\delta(x_t, \xi) = f(x_t, \xi) + \delta_t \quad \text{and} \quad f_\delta(x'_t, \xi) = f(x'_t, \xi) + \delta'_t. \quad (5)$$

Then the gradient can be approximated as follows

$$\mathbf{g}(x, \mathbf{e}, r, \xi) = \frac{d}{2h} (f(x + h r \mathbf{e}, \xi) - f(x - h r \mathbf{e}, \xi) + (\delta - \delta')) K(r) \mathbf{e}. \quad (6)$$

Assumption 4 (Stochastic adversarial noise [31]). For all $t = 1, 2, \dots$, it holds

$$\mathbb{E}[\delta_t^2] \leq \Delta^2, \quad \text{and} \quad \mathbb{E}[\delta'_t{}^2] \leq \Delta^2,$$

δ_t, δ'_t are independent.

The next theorem states the convergence guarantee for AZO-SGD-HS with zero-order oracle (5). We will use the shorthand notations $\kappa_\beta = \int |u|^\beta |K(u)| du$ and $\kappa = \int |K(u)| du$.

Theorem 2. *Let f satisfy Assumption 2 with parameter β , the adversarial noise is stochastic and follows Assumption 4. Let smoothing parameter be $h \leq (\varepsilon/(\kappa_\beta LR))^{1/(\beta-1)}$ then Assumption 3 is satisfied with gradient approximation (4). Let the maximum admissible level of stochastic adversarial noise be*

- $B \leq N$

$$\Delta \leq \frac{\sigma_* \varepsilon^{1/(\beta-1)}}{\kappa_\beta^{1/(\beta-1)} \sqrt{d} R^{1/(\beta-1)} L^{1/(\beta-1)}}$$

- $B > N$

$$\Delta \leq \frac{\varepsilon^{3/4+1/(\beta-1)} \sqrt{B} L^{1/4-1/(\beta-1)}}{\sqrt{\kappa \kappa'(p, d) d} \kappa_\beta^{1/(\beta-1)} R^{1/2+1/(\beta-1)}}.$$

Let x_N^{ag} be the output of the AZO-SGD-HS then

$$\mathbb{E}[f(x_N^{ag}) - f^*] \leq \varepsilon$$

in at most the following number of iterations N and oracle calls T , where R will denote a radius of a solution x^* : $\|x^* - x_0\| \leq R$, and x_0 is a starting point:

- $B = 1$

$$N = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d \kappa \sigma_*^2 R^2}{\varepsilon^2} \right) \right), \quad T = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d \kappa \sigma_*^2 R^2}{\varepsilon^2} \right) \right)$$

- $N > B > 1$

$$N = \mathcal{O} \left(\max \left(\frac{LR^2}{B\varepsilon}, \sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2 R^2}{B\varepsilon^2} \right) \right)$$

$$T = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, B\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2} \right) \right)$$

- $B = N$

$$N = \mathcal{O} \left(\max \left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{\sqrt{d\kappa\sigma_*} R}{\varepsilon} \right) \right), T = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2} \right) \right)$$

- $B > N$

$$N = \mathcal{O} \left(\sqrt{\frac{LR^2}{\varepsilon}} \right), T = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}, \frac{\kappa_\beta^{2/(\beta-1)} \kappa d^2 \Delta^2 L^{2/(\beta-1)} R^{2+2/(\beta-1)}}{\varepsilon^{2+2/(\beta-1)}} \right) \right).$$

Sketch of the proof.

Step 1 (bias of gradient approximation)

In this step, we calculate the bias of the gradient approximation $\|\mathbb{E}[g(x_k, \xi, \mathbf{e})] - \nabla f(x_k)\|$. The main difference between them is

$$\begin{aligned} b &= \|\mathbb{E}[g(x_k, \mathbf{e}, r, \xi)] - \nabla f(x_k)\| = \left\| \mathbb{E} \left[d \frac{f_\delta(x_k + hr\mathbf{e}, \xi) - f_\delta(x_k - hr\mathbf{e}, \xi)}{2h} K(r)\mathbf{e} \right] - \nabla f(x_k) \right\| \\ &\stackrel{\textcircled{1}}{=} \left\| \mathbb{E} \left[d \frac{f(x_k + hr\mathbf{e}, \xi)}{h} K(r)\mathbf{e} \right] + \mathbb{E} \left[\frac{\delta - \delta'}{2h} K(r)\mathbf{e} \right] - \nabla f(x_k) \right\| \\ &\stackrel{\textcircled{2}}{=} \left\| \mathbb{E} \left[d \frac{f(x_k + hr\mathbf{e}, \xi)}{h} K(r)\mathbf{e} \right] - \nabla f(x_k) \right\| \stackrel{\textcircled{3}}{=} \|\mathbb{E}[(\nabla f(x_k + hr\mathbf{u}, \xi)rK(r)) - \nabla f(x_k)]\| \\ &\stackrel{\textcircled{4}}{\leq} \sup_{z \in S_2^d(1)} \mathbb{E}[(\nabla_z f(x_k + hr\mathbf{u}, \xi) - \nabla_z f(x_k))rK(r)] \\ &\stackrel{\textcircled{5}}{\leq} \kappa_\beta h^{\beta-1} \frac{L}{(l-1)!} \mathbb{E}[\|\mathbf{u}\|^{\beta-1}] \leq \kappa_\beta h^{\beta-1} \frac{L}{(l-1)!} \frac{d}{d+\beta-1} \lesssim \kappa_\beta L h^{\beta-1}, \end{aligned}$$

where $u \in B_2^d(1)$, $\textcircled{1}$: distribution of \mathbf{e} is symmetric; $\textcircled{2}$: independence of the stochastic noise; $\textcircled{3}$: a version of the Stokes' theorem [30] (see Section 13.3.5, Exercise 14a); $\textcircled{4}$: norm of the gradient is the supremum of directional derivatives $\nabla_z f(x) = \lim_{\varepsilon \rightarrow 0} \frac{f(x+\varepsilon z) - f(x)}{\varepsilon}$; $\textcircled{5}$: Taylor expansion.

Step 2 (bounding the second moment of gradient approximation)

In this step, we calculate the second moment of the gradient approximation $\mathbb{E}[\|g(x^*, \xi, \mathbf{e})\|^2]$. The calculations are done similarly to the deterministic case.

$$\begin{aligned}
\zeta^2 &= \mathbb{E}[\|\mathbf{g}(x^*, \mathbf{e}, r, \xi)\|^2] = \mathbb{E}\left[\left\|d\frac{f_\delta(x^* + hr\mathbf{e}, \xi) - f_\delta(x^* - hr\mathbf{e}, \xi)}{2h}K(r)\mathbf{e}\right\|^2\right] \\
&= \frac{d^2}{4h^2}\mathbb{E}[(f_\delta(x^* + hr\mathbf{e}, \xi) - f_\delta(x^* - hr\mathbf{e}, \xi))K(r)]^2 \\
&= \frac{d^2}{4h^2}\mathbb{E}[(f(x^* + hr\mathbf{e}, \xi) - f(x^* - hr\mathbf{e}, \xi) + (\delta - \delta'))^2(K(r))^2] \\
&\stackrel{\textcircled{1}}{\leq} \frac{d^2\kappa}{2h^2}(\mathbb{E}(f(x^* + hr\mathbf{e}, \xi) - f(x^* - hr\mathbf{e}, \xi))^2 + 2\Delta^2) \\
&\stackrel{\textcircled{2}}{\leq} \frac{d^2\kappa}{2h^2}\left(\frac{h^2}{d}\mathbb{E}(\|\nabla f(x^* + hr\mathbf{e}, \xi) + \nabla f(x^* - hr\mathbf{e}, \xi)\|^2) + 2\Delta^2\right) \\
&= \frac{d^2\kappa}{2h^2}\left(\frac{h^2}{d}\mathbb{E}(\|\nabla f(x^* + hr\mathbf{e}, \xi) + \nabla f(x^* - hr\mathbf{e}, \xi) \pm 2\nabla f(x^*, \xi)\|^2) + 2\Delta^2\right) \\
&\stackrel{\textcircled{3}}{\leq} 4d\kappa\|\nabla f(x^*, \xi)\|^2 + 4d\kappa L^2 h^2 \mathbb{E}[\|\mathbf{e}\|^2] + \frac{d^2\kappa\Delta^2}{h^2} \\
&\stackrel{\textcircled{4}}{\leq} 4d\kappa\sigma_*^2 + 4d\kappa L^2 h^2 \mathbb{E}[\|\mathbf{e}\|^2] + \frac{d^2\kappa\Delta^2}{h^2} = 4d\kappa\sigma_*^2 + 4d\kappa L^2 h^2 + \frac{d^2\kappa\Delta^2}{h^2},
\end{aligned}$$

①: inequality of squared norm of the sum, inequality between positive random variables and independence of the noise; ②: Wirtinger-Poincare inequality; ③: L-smoothness function; ④: overparametrization assumption. \square

For the complete proof we refer to Appendix A.3.

Compared to Theorem 1, the Theorem 2 shows similar results on iterative complexity N , total oracle calls T , and smoothing parameter h . The estimation on the maximum allowable level of adversarial noise in this case explicitly depends on batch size B , which demonstrates the effect of overbatching that already occurred for similar problems as in [32]. The bound allows to significantly increase the quality of the solution with the increase in batch size.

Remark 2. Theorem 2 was proven for the Euclidean case but it can be generalized to the ℓ_p -norm. In this case, the number of iterations N and smoothing parameter h remain the same. The total number of oracle calls changes to

$$\begin{aligned}
T &= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\kappa'(p, d)\sigma_*^2 R^2}{\varepsilon^2}, \frac{\kappa_\beta^{2/(\beta-1)}\kappa\kappa'(p, d)d^2\Delta^2 L^{2/(\beta-1)}R^{2+2/(\beta-1)}}{\varepsilon^{2+2/(\beta-1)}}\right)\right) \\
&= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{\kappa\min\{q, \ln d\}d^{2-2/p}\sigma_*^2 R^2}{\varepsilon^2}, \frac{\kappa_\beta^{2/(\beta-1)}\kappa\min\{q, \ln d\}d^{3-2/p}\Delta^2 L^{2/(\beta-1)}R^{2+2/(\beta-1)}}{\varepsilon^{2+2/(\beta-1)}}\right)\right)
\end{aligned}$$

and for $B > N$, the maximum admissible level of adversarial noise Δ changes to

$$\Delta \leq \frac{\varepsilon^{3/4+1/(\beta-1)}\sqrt{BL}^{1/4-1/(\beta-1)}}{\sqrt{\kappa\min\{q, \ln d\}d^{3/2-1/p}\kappa_\beta^{1/(\beta-1)}R^{1/2+1/(\beta-1)}}}.$$

The proof of this remark is given in Appendix A.4. The remark also makes it possible to reduce the total number of oracle calls in the ℓ_1 -norm. However, this reduces the maximum adversarial noise level that can potentially worsen the results. In contrast, using the ℓ_∞ norm increases the total number of oracle calls and the maximum admissible level of adversarial noise, resulting in a trade-off between these parameters.

4 Experiments

We evaluate the effectiveness of AZO-SGD-HS by comparing it with AZO-SGD [25], which does not utilize the property of high-order smoothness, on the logistic regression

$$\min_{w \in \mathbb{R}^{d+1}} \sum_{i=1}^n -(y_i \log(p_i) + (1 - y_i) \log(1 - p_i)), \quad (7)$$

where $p_i = 1/(1 + e^{(w, X_i)})$, X_i is the i -th example from the training dataset, and y_i is the correct label of X_i , n is the number of examples from training dataset, w is the vector of parameters. The problem is convex w.r.t. w . Problem (7) is smooth with parameter $L = \lambda_{\max}(X^\top X)/(4n)$, where $\lambda_{\max}(X^\top X)$ is the maximum eigenvalue of $X^\top X$. The high-order smoothness condition is also satisfied. The results are presented in Figure 1.




Figure 1: Comparison of the proposed algorithm AZO-SGD-HS with AZO-SGD from [25]

Synthetic data. For binary classification, we sampled $n = 100$ examples with $d = 1000$ features. The setup $d \gg n$ enjoys overparametrization. The dataset was sampled with the `sklearn.datasets.make_classification` procedure.

Setup. The parameters of the algorithms were chosen as follows: number of iterations $N = 1000$, adversarial noise level $\Delta = 0.0001$, $\beta = 4$ for AZO-SGD-HS and theoretically calculated $L = 4.3$ for AZO-SGD [25]. We compare the methods under batch size $B = 2000$. For demonstration of overbatching effect we considered $B = \{100, 200, 500, 1000, 2000, 5000\}$. To model deterministic adversarial noise we decreased the weights by a noisy normed gradient of the loss function w.r.t weights. To model stochastic adversarial noise, we used normal distribution clipped to $[-\Delta, \Delta]$ with variance equal to Δ^2 . The kernel approximation was based on the Legendre polynomials [27]

$$K(r) = \sum_{m=0}^{\beta} p'_m(0)p_m(r), \quad p_m(u) = \sqrt{2m+1}L_m(u),$$

where L_m is the m -th Legendre polynomial. For $\beta = 4$

$$K(r) = \frac{15r}{4}(5 - 7r^2).$$

5 Conclusion

In this paper, we proposed a new gradient-free algorithm (AZO-SGD-HS) utilizing a higher-order smoothness condition and designed for the overparametrized regime. The convergence of the proposed method was studied in the Euclidean case and then was generalized to the non-Euclidean setup. The algorithm achieves optimal oracle call complexity, when the batch size B is large enough. The convergence guarantee as well as the superiority of the proposed method over AZO-SGD [25] were confirmed by numerical experiments.

References

- [1] Boris Polyak. *Introduction to Optimization*. New York, Optimization Software, 1987.
- [2] ON Granichin and BT Polyak. Randomized algorithms of an estimation and optimization under almost arbitrary noises. *M.: Nauka*, 2003.
- [3] Andrej Risteski and Yuanzhi Li. Algorithms and matching lower bounds for approximately-convex optimization. *Advances in Neural Information Processing Systems*, 29:4745–4753, 2016.
- [4] Artem Vasin, Alexander Gasnikov, and Vladimir Spokoiny. Stopping rules for accelerated gradient methods with additive noise in gradient, 2021.
- [5] James C. Spall. *Introduction to Stochastic Search and Optimization*. John Wiley & Sons, Inc., New York, NY, USA, 1 edition, 2003.
- [6] A. Conn, K. Scheinberg, and L. Vicente. *Introduction to Derivative-Free Optimization*. Society for Industrial and Applied Mathematics, 2009.
- [7] John C. Duchi, Michael I. Jordan, Martin J. Wainwright, and Andre Wibisono. Optimal rates for zero-order convex optimization: The power of two function evaluations. *IEEE Trans. Information Theory*, 61(5):2788–2806, 2015.
- [8] Ohad Shamir. An optimal algorithm for bandit and zero-order convex optimization with two-point feedback. *Journal of Machine Learning Research*, 18:52:1–52:11, 2017. First appeared in arXiv:1507.08752.
- [9] Yurii Nesterov and Vladimir Spokoiny. Random gradient-free minimization of convex functions. *Found. Comput. Math.*, 17(2):527–566, April 2017. First appeared in 2011 as CORE discussion paper 2011/16.
- [10] A. V. Gasnikov, E. A. Krymova, A. A. Lagunovskaya, I. N. Usmanova, and F. A. Fedorenko. Stochastic online optimization. single-point and multi-point non-linear multi-armed bandits. convex and strongly-convex case. *Automation and Remote Control*, 78(2):224–234, Feb 2017. arXiv:1509.01679.
- [11] Aleksandr Beznosikov, Abdurakhmon Sadiev, and Alexander Gasnikov. Gradient-free methods with inexact oracle for convex-concave stochastic saddle-point problem. In *International Conference on Mathematical Optimization Theory and Operations Research*, pages 105–119. Springer, 2020.
- [12] Alexander Gasnikov, Anton Novitskii, Vasilii Novitskii, Farshed Abdukhakimov, Dmitry Kamzolov, Aleksandr Beznosikov, Martin Takac, Pavel Dvurechensky, and Bin Gu. The power of first-order smooth optimization for black-box non-smooth problems. *arXiv preprint arXiv:2201.12289*, 2022.
- [13] Abraham D Flaxman, Adam Tauman Kalai, and H Brendan McMahan. Online convex optimization in the bandit setting: gradient descent without a gradient. *arXiv preprint cs/0408007*, 2004.
- [14] Peter Bartlett, Varsha Dani, Thomas Hayes, Sham Kakade, Alexander Rakhlin, and Ambuj Tewari. High-probability regret bounds for bandit online linear optimization. In *Proceedings of the 21st Annual Conference on Learning Theory-COLT 2008*, pages 335–342. Omnipress, 2008.
- [15] Sébastien Bubeck and Nicoló Cesa-Bianchi. Regret analysis of stochastic and nonstochastic multi-armed bandit problems. *Foundations and Trends® in Machine Learning*, 5(1):1–122, 2012.
- [16] Yunjie Ge, Qian Wang, Baolin Zheng, Xinlu Zhuang, Qi Li, Chao Shen, and Cong Wang. Anti-distillation backdoor attacks: Backdoors can really survive in knowledge distillation. In *Proceedings of the 29th ACM International Conference on Multimedia*, pages 826–834, 2021.
- [17] Darina Dvinskikh, Vladislav Tominin, Yaroslav Tominin, and Alexander Gasnikov. Gradient-free optimization for non-smooth saddle point problems under adversarial noise. *arXiv preprint arXiv:2202.06114*, 2022.
- [18] Aleksandr Lobanov. Stochastic adversarial noise in the “black box” optimization problem. In *International Conference on Optimization and Applications*, pages 60–71. Springer, 2023.
- [19] Anastasia Sergeevna Bayandina, Alexander V Gasnikov, and Anastasia A Lagunovskaya. Gradient-free two-point methods for solving stochastic nonsmooth convex optimization problems with small non-random noises. *Automation and Remote Control*, 79(8):1399–1408, 2018.

- [20] Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. *Advances in neural information processing systems*, 31, 2018.
- [21] Zeyuan Allen-Zhu, Yuanzhi Li, and Yingyu Liang. Learning and generalization in overparameterized neural networks, going beyond two layers. *Advances in neural information processing systems*, 32, 2019.
- [22] Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine-learning practice and the classical bias–variance trade-off. *Proceedings of the National Academy of Sciences*, 116(32):15849–15854, 2019.
- [23] Jordan Hoffmann, Sebastian Borgeaud, Arthur Mensch, Elena Buchatskaya, Trevor Cai, Eliza Rutherford, Diego de Las Casas, Lisa Anne Hendricks, Johannes Welbl, Aidan Clark, et al. Training compute-optimal large language models. *arXiv preprint arXiv:2203.15556*, 2022.
- [24] Ronen Eldan and Yuanzhi Li. Tinstories: How small can language models be and still speak coherent english? *arXiv preprint arXiv:2305.07759*, 2023.
- [25] Aleksandr Lobanov and Alexander Gasnikov. Accelerated zero-order sgd method for solving the black box optimization problem under “overparametrization” condition. In *International Conference on Optimization and Applications*, pages 72–83. Springer, 2023.
- [26] Boris Teodorovich Polyak and Aleksandr Borisovich Tsybakov. Optimal order of accuracy of search algorithms in stochastic optimization. *Problemy Peredachi Informatsii*, 26(2):45–53, 1990.
- [27] Francis Bach and Vianney Perchet. Highly-smooth zero-th order online optimization. In *Conference on Learning Theory*, pages 257–283. PMLR, 2016.
- [28] Arya Akhavan, Evgenii Chzhen, Massimiliano Pontil, and Alexandre B Tsybakov. Gradient-free optimization of highly smooth functions: improved analysis and a new algorithm. *arXiv preprint arXiv:2306.02159*, 2023.
- [29] Blake E Woodworth and Nathan Srebro. An even more optimal stochastic optimization algorithm: minibatching and interpolation learning. *Advances in Neural Information Processing Systems*, 34:7333–7345, 2021.
- [30] Vladimir Antonovich Zorich and Octavio Paniagua. *Mathematical analysis II*, volume 220. Springer, 2016.
- [31] Arya Akhavan, Evgenii Chzhen, Massimiliano Pontil, and Alexandre Tsybakov. A gradient estimator via 11-randomization for online zero-order optimization with two point feedback. *Advances in Neural Information Processing Systems*, 35:7685–7696, 2022.
- [32] Aleksandr Lobanov, Nail Bashirov, and Alexander Gasnikov. The “black-box” optimization problem: Zero-order accelerated stochastic method via kernel approximation. *Journal of Optimization Theory and Applications*, pages 1–36, 2024.

A Appendix

A.1 Proof of Theorem 1

We reuse the bias and the second moment of gradient approximation from the main part of the paper. They are:

$$\begin{aligned}
 b &\lesssim \kappa_\beta L h^{\beta-1} + \frac{d\Delta}{h} \\
 \zeta^2 &\leq 4d\kappa\sigma_*^2 + 4d\kappa L^2 h^2 + \frac{d^2\kappa\Delta^2}{h^2}.
 \end{aligned}$$

Convergence

We use the bound for biased oracle from [25] and substitute the bias and the second moment found earlier.

$$\begin{aligned}
\mathbb{E}[f(x_N^{ag}) - f^*] &= c \left(\frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\zeta R}{\sqrt{BN}} + bR + \frac{b^2}{2L}N \right) \\
&\lesssim \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\left(\sqrt{d\kappa\sigma_*^2} + \sqrt{d\kappa L^2 h^2} + \sqrt{\frac{d^2 \kappa \Delta^2}{h^2}} \right) R}{\sqrt{BN}} + \left(\kappa_\beta L h^{\beta-1} + \frac{d\Delta}{h} \right) R \\
&+ \frac{\left(\kappa_\beta L h^{\beta-1} + \frac{d\Delta}{h} \right)^2}{L} N \leq \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\left(\sqrt{d\kappa\sigma_*} + \sqrt{d\kappa} L h + \sqrt{\kappa} \frac{d\Delta}{h} \right) R}{\sqrt{BN}} \\
&+ \left(\kappa_\beta L h^{\beta-1} + \frac{d\Delta}{h} \right) R + \frac{\kappa_\beta^2 L^2 h^{2(\beta-1)} + \frac{d^2 \Delta^2}{h^2}}{L} N \\
&= \underbrace{\frac{LR^2}{N^2}}_{\textcircled{1}} + \underbrace{\frac{LR^2}{BN}}_{\textcircled{2}} + \underbrace{\frac{\sqrt{d\kappa\sigma_*} R}{\sqrt{BN}}}_{\textcircled{3}} + \underbrace{\frac{\sqrt{d\kappa} L h R}{\sqrt{BN}}}_{\textcircled{4}} + \underbrace{\frac{\sqrt{\kappa} d \Delta R}{h \sqrt{BN}}}_{\textcircled{5}} + \underbrace{\kappa_\beta L h^{\beta-1} R}_{\textcircled{6}} + \underbrace{\frac{d\Delta}{h} R}_{\textcircled{7}} + \underbrace{\kappa_\beta^2 L h^{2(\beta-1)}}_{\textcircled{8}} + \underbrace{\frac{d^2 \Delta^2 N}{h^2 L}}_{\textcircled{9}}.
\end{aligned}$$

We consider several cases for the B values and find all parameter values for all cases

Case 1: $B = 1$

$$\begin{aligned}
\mathbb{E}[f(x_N^{ag}) - f^*] &= \frac{LR^2}{N^2} + \frac{LR^2}{N} + \frac{\sqrt{d\kappa\sigma_*} R}{\sqrt{N}} + \frac{\sqrt{d\kappa} L h R}{\sqrt{N}} + \frac{\sqrt{\kappa} d \Delta R}{h \sqrt{N}} + \kappa_\beta L h^{\beta-1} R \\
&+ \frac{d\Delta}{h} R + \kappa_\beta^2 L h^{2(\beta-1)} + \frac{d^2 \Delta^2 N}{h^2 L} \leq 9\varepsilon.
\end{aligned}$$

We bound all terms by ε and find bounds for needed parameters.

From $\textcircled{1}$, $\textcircled{2}$ and $\textcircled{3}$, we find N

$$\begin{aligned}
\frac{LR^2}{N^2} \leq \varepsilon &\implies N \geq \sqrt{\frac{LR^2}{\varepsilon}} = \mathcal{O}\left(\sqrt{\frac{LR^2}{\varepsilon}}\right) \\
\frac{LR^2}{N} \leq \varepsilon &\implies N \geq \frac{LR^2}{\varepsilon} \\
\frac{\sqrt{d\kappa\sigma_*} R}{\sqrt{N}} \leq \varepsilon &\implies N \geq \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}.
\end{aligned}$$

Note that $\frac{LR^2}{\varepsilon} > \sqrt{\frac{LR^2}{\varepsilon}}$. That leads to the following expression for N

$$N = \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}\right)\right).$$

From $\textcircled{4}$, $\textcircled{6}$ and $\textcircled{8}$, we find h

$$\begin{aligned}
\frac{\sqrt{d\kappa}LhR}{\sqrt{N}} \leq \varepsilon &\implies h \leq \frac{\sqrt{N}\varepsilon}{\sqrt{d\kappa}LR} = \max\left(\frac{\sqrt{\frac{LR^2}{\varepsilon}}\varepsilon}{\sqrt{d\kappa}LR}, \frac{\sqrt{\frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}}\varepsilon}{\sqrt{d\kappa}LR}\right) = \max\left(\frac{\varepsilon^{1/2}}{\sqrt{d\kappa}L}, \frac{\sigma_*}{L}\right) \\
\kappa_\beta^2 Lh^{2(\beta-1)} \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}} \\
\kappa_\beta Lh^{\beta-1}R \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\
h &= \min\left(\max\left(\frac{\varepsilon^{1/2}}{\sqrt{d\kappa}L}, \frac{\sigma_*}{L}\right), \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}}, \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\right) = \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}.
\end{aligned}$$

From ⑤, ⑦ and ⑨, we find Δ :

$$\begin{aligned}
\frac{\sqrt{\kappa}d\Delta R}{h\sqrt{N}} \leq \varepsilon &\implies \Delta \leq \frac{\varepsilon h\sqrt{N}}{\sqrt{\kappa}dR} \leq \max\left(\frac{\varepsilon\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\sqrt{\frac{LR^2}{\varepsilon}}}{\sqrt{\kappa}dR}, \frac{\varepsilon\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\sqrt{\frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}}}{\sqrt{\kappa}dR}\right) \\
&= \max\left(\frac{\varepsilon^{1/2+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{\sqrt{\kappa}d(\kappa_\beta R)^{\frac{1}{\beta-1}}}, \frac{\varepsilon^{\frac{1}{\beta-1}}\sigma_*}{\sqrt{d\kappa_\beta^{\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}}}\right) = \frac{\varepsilon^{\frac{1}{\beta-1}}\sigma_*}{\sqrt{d\kappa_\beta^{\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}}} \\
\frac{d\Delta}{h}R \leq \varepsilon &\implies \Delta \leq \frac{h\varepsilon}{dR} \leq \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\varepsilon}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} = \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} \\
\frac{d^2\Delta^2 N}{h^2 L} \leq \varepsilon &\implies \Delta \leq \frac{\sqrt{\varepsilon}Lh}{\sqrt{N}d} \leq \frac{\sqrt{\varepsilon}L\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}}{\sqrt{\frac{LR^2}{\varepsilon}}d} = \frac{\varepsilon^{1/2+1/4+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}-1/4}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}} \\
&= \frac{\varepsilon^{3/4+\frac{1}{\beta-1}}L^{1/4-\frac{1}{\beta-1}}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}} \\
\Delta &= \min\left\{\frac{\varepsilon^{\frac{1}{\beta-1}}\sigma_*}{\sqrt{d\kappa_\beta^{\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}}}, \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}}, \frac{\varepsilon^{3/4+\frac{1}{\beta-1}}L^{1/4-\frac{1}{\beta-1}}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}}\right\} \\
&= \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}}.
\end{aligned}$$

Then, we can find total number of oracle calls

$$T = N \cdot B = \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}\right)\right).$$

Case 2: $N > B > 1$

$$\begin{aligned}
\mathbb{E}[f(x_N^{ag}) - f^*] &= \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\sqrt{d\kappa}\sigma_*R}{\sqrt{BN}} + \frac{\sqrt{d\kappa}LhR}{\sqrt{BN}} + \frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} + \kappa_\beta Lh^{\beta-1}R \\
&+ \frac{d\Delta}{h}R + \kappa_\beta^2 Lh^{2(\beta-1)} + \frac{d^2\Delta^2 N}{h^2 L} \leq 9\varepsilon.
\end{aligned}$$

From ①, ②, ③, we find N

$$\begin{aligned}
\frac{LR^2}{N^2} \leq \varepsilon &\implies N \geq \sqrt{\frac{LR^2}{\varepsilon}} \\
\frac{LR^2}{BN} \leq \varepsilon &\implies N \geq \frac{LR^2}{B\varepsilon} \\
\frac{\sqrt{d}\kappa\sigma_*R}{\sqrt{BN}} \leq \varepsilon &\implies N \geq \frac{d\kappa\sigma_*^2R^2}{B\varepsilon^2} \\
N &= \mathcal{O}\left(\max\left(\frac{LR^2}{B\varepsilon}, \sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2R^2}{B\varepsilon^2}\right)\right).
\end{aligned}$$

From ④, ⑥, ⑧, we find h

$$\begin{aligned}
\frac{\sqrt{d}\kappa LhR}{\sqrt{BN}} \leq \varepsilon &\implies h \leq \frac{\varepsilon\sqrt{BN}}{\sqrt{d}\kappa LR} \leq \frac{\varepsilon\sqrt{BN}}{\sqrt{d}\kappa LR} \leq \max\left(\frac{\varepsilon\sqrt{\frac{LR^2}{\varepsilon}}}{\sqrt{d}\kappa LR}, \frac{\varepsilon\sqrt{B\sqrt{\frac{LR^2}{\varepsilon}}}}{\sqrt{d}\kappa LR}, \frac{\varepsilon\sqrt{B\frac{d\kappa\sigma_*^2R^2}{B\varepsilon^2}}}{\sqrt{d}\kappa LR}\right) \\
&= \max\left(\sqrt{\frac{\varepsilon}{d}\kappa L}, \frac{\varepsilon^{3/4}\sqrt{B}}{\sqrt{d}\kappa RL^{3/4}}, \frac{\sigma_*}{L}\right) \\
\kappa_\beta^2 Lh^{2(\beta-1)} \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}} \\
\kappa_\beta Lh^{\beta-1}R \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\
h &= \min\left(\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}, \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}}, \max\left(\sqrt{\frac{\varepsilon}{d}\kappa L}, \frac{\varepsilon^{3/4}\sqrt{B}}{\sqrt{d}\kappa RL^{3/4}}, \frac{\sigma_*}{L}\right)\right) = \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}.
\end{aligned}$$

From ⑤, ⑦, ⑨, we find Δ

$$\begin{aligned}
\frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} \leq \varepsilon &\implies \Delta \leq \frac{h\sqrt{BN}\varepsilon}{\sqrt{\kappa}dR} \leq \max\left(\frac{h\left(\sqrt{\frac{LR^2}{\varepsilon}}\right)\varepsilon}{\sqrt{\kappa}dR}, \frac{h\left(\sqrt{B\sqrt{\frac{LR^2}{\varepsilon}}}\right)\varepsilon}{\sqrt{\kappa}dR}\right) \\
&= \max\left(\frac{h(\sqrt{L}\varepsilon)}{\sqrt{\kappa}d}, \frac{h(\sqrt{B\sqrt{L}})\varepsilon^{3/4}}{\sqrt{\kappa}dR^{1/2}}\right) \\
&= \max\left(\frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}(\sqrt{L}\varepsilon)}{\sqrt{\kappa}d}, \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}(\sqrt{B\sqrt{L}})\varepsilon^{3/4}}{\sqrt{\kappa}dR^{1/2}}\right) \\
&= \max\left(\frac{\varepsilon^{1/2+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}}, \frac{L^{1/4-\frac{1}{\beta-1}}\sqrt{B}\varepsilon^{3/4+\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{1/2+\frac{1}{\beta-1}}}\right) = \frac{\varepsilon^{1/2+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}} \\
\frac{d\Delta}{h}R \leq \varepsilon &\implies \Delta \leq \frac{h\varepsilon}{dR} \leq \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\varepsilon}{dR} = \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} \\
\frac{d^2\Delta^2N}{h^2L} \leq \varepsilon &\implies \Delta \leq \frac{\sqrt{\varepsilon}Lh}{\sqrt{Nd}} \leq \frac{\sqrt{\varepsilon}L\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}}{\sqrt{\frac{LR^2}{\varepsilon}}d} = \frac{\varepsilon^{1/2+1/4+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}-1/4}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}} \\
&= \frac{\varepsilon^{3/4+\frac{1}{\beta-1}}L^{1/4-\frac{1}{\beta-1}}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}} \\
\Delta &= \min\left(\frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}}, \frac{\varepsilon^{1/2+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}}, \frac{\varepsilon^{3/4+\frac{1}{\beta-1}}L^{1/4-\frac{1}{\beta-1}}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}}\right) \\
&= \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} \\
T = N \cdot B &= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, B\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa_\beta^2 R^2}{\varepsilon^2}\right)\right).
\end{aligned}$$

Case 3: $B = N$

$$\begin{aligned}
\mathbb{E}[f(x_N^{ag}) - f^*] &= \frac{LR^2}{N^2} + \frac{LR^2}{N^2} + \frac{\sqrt{d\kappa}\sigma_*R}{N} + \frac{\sqrt{d\kappa}LhR}{N} + \frac{\sqrt{\kappa}d\Delta R}{hN} + \kappa_\beta Lh^{\beta-1}R \\
&+ \frac{d\Delta}{h}R + \kappa_\beta^2 Lh^{2(\beta-1)} + \frac{d^2\Delta^2N}{h^2L} \leq 9\varepsilon.
\end{aligned}$$

From ①, ③, we find N

$$\begin{aligned}
\frac{LR^2}{N^2} \leq \varepsilon &\implies N \geq \sqrt{\frac{LR^2}{\varepsilon}} \\
\frac{\sqrt{d\kappa}\sigma_*R}{N} \leq \varepsilon &\implies N \geq \frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon} \\
N &= \max\left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon}\right).
\end{aligned}$$

From ④, ⑥, ⑧, we find h

$$\begin{aligned}
\frac{\sqrt{d\kappa}LhR}{N} \leq \varepsilon &\implies h \leq \frac{\varepsilon N}{\sqrt{d\kappa}LR} = \max\left(\frac{\varepsilon}{\sqrt{d\kappa}LR}\sqrt{\frac{LR^2}{\varepsilon}}, \frac{\varepsilon}{\sqrt{d\kappa}LR}\frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon}\right) \\
&= \max\left(\sqrt{\frac{\varepsilon}{d\kappa L}}, \frac{\sigma_*}{L}\right) \\
\kappa_\beta^2 Lh^{2(\beta-1)} \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}} \\
\kappa_\beta Lh^{\beta-1}R \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\
h &= \min\left(\max\left(\sqrt{\frac{\varepsilon}{d\kappa L}}, \frac{\sigma_*}{L}\right), \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}}, \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\right) = \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}.
\end{aligned}$$

From ⑤, ⑦, ⑨, we find Δ

$$\begin{aligned}
\frac{\sqrt{\kappa}d\Delta R}{hN} \leq \varepsilon &\implies \Delta \leq \frac{hN\varepsilon}{\sqrt{\kappa}dR} = \max\left(\frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\varepsilon}{\sqrt{\kappa}dR}\sqrt{\frac{LR^2}{\varepsilon}}, \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\varepsilon}{\sqrt{\kappa}dR}\frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon}\right) \\
&= \max\left(\frac{\varepsilon^{1/2+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}}, \frac{\sigma_*\varepsilon^{\frac{1}{\beta-1}}}{\kappa_\beta^{\frac{1}{\beta-1}}\sqrt{d}R^{\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}}\right) = \frac{\sigma_*\varepsilon^{\frac{1}{\beta-1}}}{\kappa_\beta^{\frac{1}{\beta-1}}\sqrt{d}R^{\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} \\
\frac{d\Delta}{h}R \leq \varepsilon &\implies \Delta \leq \frac{h\varepsilon}{dR} \leq \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\varepsilon}{dR} = \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} \\
\frac{d^2\Delta^2N}{h^2L} \leq \varepsilon &\implies \Delta \leq \frac{\sqrt{\varepsilon}Lh}{\sqrt{Nd}} \leq \min\left(\frac{\sqrt{\varepsilon}L\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}}{\sqrt{\frac{LR^2}{\varepsilon}}d}, \frac{\sqrt{\varepsilon}L\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}}{\sqrt{\frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon}}d}\right) \\
&= \min\left(\frac{\varepsilon^{1/2+1/4+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}-1/4}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}}, \frac{\varepsilon^{1+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{d^{5/4}\kappa_\beta^{\frac{1}{\beta-1}}\sqrt{\sqrt{\kappa}\sigma_*R}^{1/2+\frac{1}{\beta-1}}}\right) \\
&= \min\left(\frac{\varepsilon^{3/4+\frac{1}{\beta-1}}L^{1/4-\frac{1}{\beta-1}}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}}, \frac{\varepsilon^{1+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{d^{5/4}\kappa_\beta^{\frac{1}{\beta-1}}\sqrt{\sqrt{\kappa}\sigma_*R}^{1/2+\frac{1}{\beta-1}}}\right) = \frac{\varepsilon^{1+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{d^{5/4}\kappa_\beta^{\frac{1}{\beta-1}}\sqrt{\sqrt{\kappa}\sigma_*R}^{1/2+\frac{1}{\beta-1}}} \\
\Delta &= \min\left(\frac{\varepsilon^{1+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{d^{5/4}\kappa_\beta^{\frac{1}{\beta-1}}\sqrt{\sqrt{\kappa}\sigma_*R}^{1/2+\frac{1}{\beta-1}}}, \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}}, \frac{\sigma_*\varepsilon^{\frac{1}{\beta-1}}}{\kappa_\beta^{\frac{1}{\beta-1}}\sqrt{d}R^{\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}}\right) \\
&= \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} \\
T = N \cdot B &= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2R^2}{\varepsilon^2}\right)\right).
\end{aligned}$$

Case 4: $B > N$

$$\begin{aligned} \mathbb{E}[f(x_N^{ag}) - f^*] &= \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\sqrt{d\kappa}\sigma_*R}{\sqrt{BN}} + \frac{\sqrt{d\kappa}LhR}{\sqrt{BN}} + \frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} + \kappa_\beta Lh^{\beta-1}R \\ &+ \frac{d\Delta}{h}R + \kappa_\beta^2 Lh^{2(\beta-1)} + \frac{d^2\Delta^2 N}{h^2 L} \leq 9\varepsilon. \end{aligned}$$

From ①, ②, ③, we find N and B . Note that if $B > N$ then $\frac{LR^2}{N^2} > \frac{LR^2}{BN}$

$$\begin{aligned} \frac{LR^2}{N^2} \leq \varepsilon &\implies B > N \geq \sqrt{\frac{LR^2}{\varepsilon}} \\ \frac{\sqrt{d\kappa}\sigma_*R}{\sqrt{BN}} \leq \varepsilon &\implies B \geq \frac{d\kappa\sigma_*^2 R^2}{N\varepsilon^2} = \mathcal{O}\left(\frac{d\kappa\sigma_*^2 R}{L^{1/2}\varepsilon^{3/2}}\right). \end{aligned}$$

From ④, ⑥, ⑧, we find h

$$\begin{aligned} \kappa_\beta^2 Lh^{2(\beta-1)} \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}} \\ \kappa_\beta Lh^{\beta-1}R \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\ \frac{\sqrt{d\kappa}LhR}{\sqrt{BN}} \leq \varepsilon &\implies h \leq \frac{\varepsilon\sqrt{BN}}{\sqrt{d\kappa}LR} \leq \frac{\varepsilon\sqrt{BN}}{\sqrt{d\kappa}LR} \leq \frac{\varepsilon\sqrt{B}\sqrt[4]{\frac{LR^2}{\varepsilon}}}{\sqrt{d\kappa}LR} = \frac{\varepsilon^{3/4}\sqrt{B}}{\sqrt{d\kappa}L^{3/4}R^{1/2}} \\ h &= \min\left(\left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}}, \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}, \frac{\varepsilon^{3/4}\sqrt{B}}{\sqrt{d\kappa}L^{3/4}R^{1/2}}\right) = \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}. \end{aligned}$$

From ⑤, ⑦, ⑨, we find Δ

$$\begin{aligned} \frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} \leq \varepsilon &\implies \Delta \leq \frac{h\sqrt{BN}\varepsilon}{\sqrt{\kappa}dR} \leq \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \left(\sqrt{B}\sqrt[4]{\frac{LR^2}{\varepsilon}}\right)\varepsilon}{\sqrt{\kappa}dR} = \frac{\varepsilon^{3/4+\frac{1}{\beta-1}}\sqrt{B}L^{1/4-\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{1/2+\frac{1}{\beta-1}}} \\ \frac{d\Delta}{h}R \leq \varepsilon &\implies \Delta \leq \frac{h\varepsilon}{dR} \leq \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\varepsilon}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} = \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} \\ \frac{d^2\Delta^2 N}{h^2 L} \leq \varepsilon &\implies \Delta \leq \frac{\sqrt{\varepsilon}Lh}{\sqrt{N}d} \leq \frac{\sqrt{\varepsilon}L\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}}{\sqrt[4]{\frac{LR^2}{\varepsilon}}d} = \frac{\varepsilon^{1/2+1/4+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}-1/4}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}} = \\ &= \frac{\varepsilon^{3/4+\frac{1}{\beta-1}}L^{1/4-\frac{1}{\beta-1}}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}} \\ \Delta &= \min\left\{\frac{\varepsilon^{3/4+\frac{1}{\beta-1}}\sqrt{B}L^{1/4-\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{1/2+\frac{1}{\beta-1}}}, \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}}, \frac{\varepsilon^{3/4+\frac{1}{\beta-1}}L^{1/4-\frac{1}{\beta-1}}}{dR^{1/2+\frac{1}{\beta-1}}\kappa_\beta^{\frac{1}{\beta-1}}}\right\} \\ &= \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}}. \end{aligned}$$

We can get the batch size B via the maximum noise level Δ from ⑤:

$$\frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} \leq \varepsilon \implies B \geq \frac{\kappa d^2 \Delta^2 R^2}{\varepsilon^2 h^2 N} \geq \frac{\kappa d^2 \Delta^2 R^2}{\varepsilon^2 \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{2}{\beta-1}} \sqrt{\frac{LR^2}{\varepsilon}}} \geq \frac{\kappa_\beta^{\frac{2}{\beta-1}} \kappa d^2 \Delta^2 R^{1+\frac{2}{\beta-1}}}{\varepsilon^{3/2+\frac{2}{\beta-1}} L^{1/2-\frac{2}{\beta-1}}}.$$

That leads to the following B

$$B = \mathcal{O} \left(\max \left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2 R}{L^{1/2}\varepsilon^{3/2}}, \frac{\kappa_\beta^{\frac{2}{\beta-1}} \kappa d^2 \Delta^2 R^{1+\frac{2}{\beta-1}}}{\varepsilon^{3/2+\frac{2}{\beta-1}} L^{1/2-\frac{2}{\beta-1}}} \right) \right)$$

$$T = B \cdot N = \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}, \frac{\kappa_\beta^{\frac{2}{\beta-1}} \kappa d^2 L^{\frac{2}{\beta-1}} \Delta^2 R^{2+\frac{2}{\beta-1}}}{\varepsilon^{2+\frac{2}{\beta-1}}} \right) \right).$$

That ends the proof. \square

A.2 Proof of Remark 1

Bias of gradient approximation

$$b = \|\mathbb{E}[\mathbf{g}(x_k, \mathbf{e}, r, \xi)] - \nabla f(x_k)\|_p = \left\| \mathbb{E} \left[d \frac{f_\delta(x_k + hr\mathbf{e}, \xi) - f_\delta(x_k - hr\mathbf{e}, \xi)}{2h} K(r)\mathbf{e} \right] - \nabla f(x_k) \right\|_p$$

$$\stackrel{\textcircled{1}}{=} \left\| \mathbb{E} \left[d \frac{f(x_k + hr\mathbf{e}, \xi)}{h} K(r)\mathbf{e} + \delta(x_k + h\mathbf{e}) \right] - \nabla f(x_k) \right\|_p$$

$$\stackrel{\textcircled{2}}{\leq} \left\| \mathbb{E} \left[d \frac{f(x_k + hr\mathbf{e}, \xi)}{h} K(r)\mathbf{e} \right] - \nabla f(x_k) \right\|_p + \frac{d\Delta}{h}$$

$$\stackrel{\textcircled{3}}{=} \|\mathbb{E}[(\nabla f(x_k + hr\mathbf{u}, \xi)rK(r)) - \nabla f(x_k)]\|_p + \frac{d\Delta}{h}$$

$$\stackrel{\textcircled{4}}{=} \sup_{z \in S_p^d(1)} \mathbb{E}[(\nabla_z f(x_k + hr\mathbf{u}, \xi) - \nabla_z f(x_k))rK(r)] + \frac{d\Delta}{h}$$

$$\stackrel{\textcircled{5}}{\leq} \kappa_\beta h^{\beta-1} \frac{L}{(l-1)!} \mathbb{E}[\|\mathbf{u}\|_p^{\beta-1}] + \frac{d\Delta}{h} \lesssim \kappa_\beta L h^{\beta-1} + \frac{d\Delta}{h},$$

where $\mathbf{u} \in B_2^d(1)$, $\textcircled{1}$: distribution of e is symmetric; $\textcircled{2}$: independence of the stochastic noise; $\textcircled{3}$: a version of the Stokes' theorem [30] (see Section 13.3.5, Exercise 14a); $\textcircled{4}$: norm of the gradient is the supremum of directional derivatives $\nabla_z f(x) = \lim_{\varepsilon \rightarrow 0} \frac{f(x+\varepsilon z) - f(x)}{\varepsilon}$; $\textcircled{5}$: Taylor expansion.

Bounding second moment of gradient approximation

Note that $\mathbb{E}[\|\mathbf{e}\|_p^2] \leq \kappa'(p, d) = \min\{q, \ln d\}d^{2/q-1} = \min\{q, \ln d\}d^{1-2/p}$ and

$$\begin{aligned}
\zeta^2 &= \mathbb{E}[\|\mathbf{g}(x^*, \mathbf{e}, r, \xi)\|_p^2] = \mathbb{E}\left[\left\|d\frac{f_\delta(x^* + h\mathbf{r}\mathbf{e}, \xi) - f_\delta(x^* - h\mathbf{r}\mathbf{e}, \xi)}{2h}K(r)\mathbf{e}\right\|_p^2\right] \\
&= \frac{d^2}{4h^2}\mathbb{E}[\|(f_\delta(x^* + h\mathbf{r}\mathbf{e}, \xi) - f_\delta(x^* - h\mathbf{r}\mathbf{e}, \xi))K(r)\|_p^2] \\
&= \frac{d^2}{4h^2}\mathbb{E}[(f(x^* + h\mathbf{r}\mathbf{e}, \xi) - f(x^* - h\mathbf{r}\mathbf{e}, \xi) + \delta(x^* + h\mathbf{r}\mathbf{e}) - \delta(x^* - h\mathbf{r}\mathbf{e}))^2K^2(r)\|\mathbf{e}\|_p^2] \\
&\stackrel{\textcircled{1}}{\leq} \frac{d^2\kappa}{2h^2}\mathbb{E}[(f(x^* + h\mathbf{r}\mathbf{e}, \xi) - f(x^* - h\mathbf{r}\mathbf{e}, \xi))^2\|\mathbf{e}\|_p^2] + \mathbb{E}[(\delta(x^* + h\mathbf{r}\mathbf{e}) - \delta(x^* - h\mathbf{r}\mathbf{e}))^2\|\mathbf{e}\|_p^2] \\
&\stackrel{\textcircled{2}}{\leq} \frac{d^2\kappa}{2h^2}(\mathbb{E}[(f(x^* + h\mathbf{r}\mathbf{e}, \xi) - f(x^* - h\mathbf{r}\mathbf{e}, \xi))^2\|\mathbf{e}\|_p^2] + 2\Delta^2\mathbb{E}[\|\mathbf{e}\|_p^2]) \\
&\leq \frac{d^2\kappa\kappa'(p, d)}{2h^2}(\mathbb{E}[(f(x^* + h\mathbf{r}\mathbf{e}, \xi) - f(x^* - h\mathbf{r}\mathbf{e}, \xi))^2] + 2\Delta^2) \\
&\stackrel{\textcircled{3}}{\leq} \frac{d^2\kappa\kappa'(p, d)}{2h^2}\left(\frac{h^2}{d}\mathbb{E}[\|\nabla f(x^* + h\mathbf{r}\mathbf{e}, \xi) + \nabla f(x^* - h\mathbf{r}\mathbf{e}, \xi)\|_2^2] + 2\Delta^2\right) \\
&= \frac{d^2\kappa\kappa'(p, d)}{2h^2}\left(\frac{h^2}{d}\mathbb{E}[\|\nabla f(x^* + h\mathbf{r}\mathbf{e}, \xi) + \nabla f(x^* - h\mathbf{r}\mathbf{e}, \xi) \pm 2\nabla f(x^*, \xi)\|_2^2] + 2\Delta^2\right) \\
&\stackrel{\textcircled{4}}{\leq} 4d\kappa\kappa'(p, d)\|\nabla f(x^*, \xi)\|_2^2 + 4d\kappa\kappa'(p, d)L^2h^2\mathbb{E}[\|\mathbf{e}\|_2^2] + \frac{d^2\kappa\kappa'(p, d)\Delta^2}{h^2} \\
&\stackrel{\textcircled{5}}{\leq} 4d\kappa\kappa'(p, d)\sigma_*^2 + 4d\kappa\kappa'(p, d)L^2h^2\mathbb{E}[\|\mathbf{e}\|_2^2] + \frac{d^2\kappa\Delta^2}{h^2} \\
&= 4d\kappa\kappa'(p, d)\sigma_*^2 + 4d\kappa\kappa'(p, d)L^2h^2 + \frac{d^2\kappa\kappa'(p, d)\Delta^2}{h^2},
\end{aligned}$$

①: inequality of squared norm of the sum and inequality between positive random variables; ②: bounded noise $|\delta| < \Delta$; ③: Wirtinger-Poincare inequality; ④: L-smoothness function; ⑤: overparametrisation assumption.

Convergence

$$\begin{aligned}
\mathbb{E}[f(x_N^{ag}) - f^*] &= c\left(\frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\zeta R}{\sqrt{BN}} + bR + \frac{b^2}{2L}N\right) \\
&\lesssim \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\left(\sqrt{d\kappa\kappa'(p, d)\sigma_*^2} + \sqrt{d\kappa\kappa'(p, d)L^2h^2} + \sqrt{\frac{d^2\kappa\kappa'(p, d)\Delta^2}{h^2}}\right)R}{\sqrt{BN}} \\
&+ \left(\kappa_\beta Lh^{\beta-1} + \frac{d\Delta}{h}\right)R + \frac{(\kappa_\beta Lh^{\beta-1} + \frac{d\Delta}{h})^2}{L}N \leq \frac{LR^2}{N^2} + \frac{LR^2}{BN} \\
&+ \frac{\left(\sqrt{d\kappa\kappa'(p, d)\sigma_*} + \sqrt{d\kappa\kappa'(p, d)Lh} + \sqrt{\kappa\kappa'(p, d)\frac{d\Delta}{h}}\right)R}{\sqrt{BN}} + \left(\kappa_\beta Lh^{\beta-1} + \frac{d\Delta}{h}\right)R \\
&+ \frac{\kappa_\beta^2 L^2 h^{2(\beta-1)} + \frac{d^2\Delta^2}{h^2}}{L}N = \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\sqrt{d\kappa\kappa'(p, d)\sigma_*}R}{\sqrt{BN}} \\
&+ \frac{\sqrt{d\kappa\kappa'(p, d)Lh}R}{\sqrt{BN}} + \frac{\sqrt{\kappa\kappa'(p, d)d\Delta}R}{h\sqrt{BN}} + \kappa_\beta Lh^{\beta-1}R + \frac{d\Delta}{h}R + \kappa_\beta^2 Lh^{2(\beta-1)} + \frac{d^2\Delta^2 N}{h^2 L}.
\end{aligned}$$

By performing the same transformations as in Theorem 1, we get the following parameters:

Case 1: $B = 1$

$$\begin{aligned}
 N &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2} \right) \right) \\
 h &= \left(\frac{\varepsilon}{\kappa_\beta LR} \right)^{\frac{1}{\beta-1}} \\
 \Delta &= \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}} R^{1+\frac{1}{\beta-1}} L^{\frac{1}{\beta-1}}} \\
 T &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2} \right) \right).
 \end{aligned}$$

Case 2: $N > B > 1$

$$\begin{aligned}
 N &= \mathcal{O} \left(\max \left(\frac{LR^2}{B\varepsilon}, \sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{B\varepsilon^2} \right) \right) \\
 h &= \left(\frac{\varepsilon}{\kappa_\beta LR} \right)^{\frac{1}{\beta-1}} \\
 \Delta &= \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}} R^{1+\frac{1}{\beta-1}} L^{\frac{1}{\beta-1}}} \\
 T &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, B\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2} \right) \right).
 \end{aligned}$$

Case 3: $B = N$

$$\begin{aligned}
 N &= \mathcal{O} \left(\max \left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{\sqrt{d\kappa\kappa'(p,d)\sigma_*} R}{\varepsilon} \right) \right) \\
 h &= \left(\frac{\varepsilon}{\kappa_\beta LR} \right)^{\frac{1}{\beta-1}} \\
 \Delta &= \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}} R^{1+\frac{1}{\beta-1}} L^{\frac{1}{\beta-1}}} \\
 T &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2} \right) \right).
 \end{aligned}$$

Case 4: $B > N$

$$\begin{aligned}
N &= \mathcal{O}\left(\sqrt{\frac{LR^2}{\varepsilon}}\right) \\
B &= \mathcal{O}\left(\max\left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R}{\sqrt{L}\varepsilon^{3/2}}, \frac{\kappa_\beta^{\frac{2}{\beta-1}}\kappa\kappa'(p,d)d^2\Delta^2 R^{1+\frac{2}{\beta-1}}}{\varepsilon^{3/2+\frac{2}{\beta-1}}L^{1/2-\frac{2}{\beta-1}}}\right)\right) \\
h &= \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\
\Delta &= \frac{\varepsilon^{1+\frac{1}{\beta-1}}}{d\kappa_\beta^{\frac{1}{\beta-1}}R^{1+\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}} \\
T &= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{\kappa\kappa'(p,d)d\sigma_*^2 R^2}{\varepsilon^2}, \frac{\kappa_\beta^{\frac{2}{\beta-1}}\kappa'(p,d)\kappa d^2\Delta^2 R^{1+\frac{2}{\beta-1}}}{\varepsilon^{3/2+\frac{2}{\beta-1}}L^{1/2-\frac{2}{\beta-1}}}\right)\right) = \\
&= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{\kappa \min\{q, \ln d\}d^{2-2/p}\sigma_*^2 R^2}{\varepsilon^2}, \frac{\min\{q, \ln d\}\kappa_\beta^{\frac{2}{\beta-1}}\kappa d^{3-2/p}\Delta^2 R^{1+\frac{2}{\beta-1}}}{\varepsilon^{3/2+\frac{2}{\beta-1}}L^{1/2-\frac{2}{\beta-1}}}\right)\right).
\end{aligned}$$

By substituting $\kappa'(p, d)$ into formulas, we prove the remark. \square

A.3 Proof of Theorem 2

We reuse the bias and the second moment of gradient approximation from the main part of the paper. They are:

$$\begin{aligned}
b &\lesssim \kappa_\beta Lh^{\beta-1} \\
\zeta^2 &\leq 4d\kappa\sigma_*^2 + 4d\kappa L^2 h^2 + \frac{d^2\kappa\Delta^2}{h^2}.
\end{aligned}$$

Convergence

We reemploy the scheme for bounding $\mathbb{E}[f(x_N^{ag}) - f^*]$ from Theorem 1.

$$\begin{aligned}
\mathbb{E}[f(x_N^{ag}) - f^*] &= c\left(\frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\zeta R}{\sqrt{BN}} + bR + \frac{b^2}{2L}N\right) \\
&\lesssim \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\left(\sqrt{d\kappa\sigma_*^2} + \sqrt{d\kappa L^2 h^2} + \sqrt{\frac{d^2\kappa\Delta^2}{h^2}}\right)R}{\sqrt{BN}} + (\kappa_\beta Lh^{\beta-1})R + \frac{(\kappa_\beta Lh^{\beta-1})^2}{L}N \\
&\leq \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\left(\sqrt{d\kappa\sigma_*} + \sqrt{d\kappa Lh} + \sqrt{\kappa\frac{d\Delta}{h}}\right)R}{\sqrt{BN}} + (\kappa_\beta Lh^{\beta-1})R + \frac{\kappa_\beta^2 L^2 h^{2(\beta-1)}}{L}N \\
&= \underbrace{\frac{LR^2}{N^2}}_{\textcircled{1}} + \underbrace{\frac{LR^2}{BN}}_{\textcircled{2}} + \underbrace{\frac{\sqrt{d\kappa\sigma_*}R}{\sqrt{BN}}}_{\textcircled{3}} + \underbrace{\frac{\sqrt{d\kappa Lh}R}{\sqrt{BN}}}_{\textcircled{4}} + \underbrace{\frac{\sqrt{\kappa d\Delta}R}{h\sqrt{BN}}}_{\textcircled{5}} + \underbrace{\kappa_\beta Lh^{\beta-1}R}_{\textcircled{6}} + \underbrace{\kappa_\beta^2 Lh^{2(\beta-1)}}_{\textcircled{7}}.
\end{aligned}$$

Case 1: $B = 1$

$$\mathbb{E}[f(x_N^{ag}) - f^*] = \frac{LR^2}{N^2} + \frac{LR^2}{N} + \frac{\sqrt{d\kappa\sigma_*}R}{\sqrt{N}} + \frac{\sqrt{d\kappa Lh}R}{\sqrt{N}} + \frac{\sqrt{\kappa d\Delta}R}{h\sqrt{N}} + \kappa_\beta Lh^{\beta-1}R + \kappa_\beta^2 Lh^{2(\beta-1)} \leq 7\varepsilon.$$

From ①, ②, ③, we find N

$$\begin{aligned}\frac{LR^2}{N^2} \leq \varepsilon &\implies N \geq \sqrt{\frac{LR^2}{\varepsilon}} \\ \frac{LR^2}{N} \leq \varepsilon &\implies N \geq \frac{LR^2}{\varepsilon} \\ \frac{\sqrt{d\kappa}\sigma_*R}{\sqrt{N}} \leq \varepsilon &\implies N \geq \frac{d\kappa\sigma_*^2R^2}{\varepsilon^2} \\ N &= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2R^2}{\varepsilon^2}\right)\right).\end{aligned}$$

From ④, ⑥, ⑦, we find h

$$\begin{aligned}\frac{\sqrt{d\kappa}LhR}{\sqrt{N}} \leq \varepsilon &\implies h \leq \frac{\sqrt{N}\varepsilon}{\sqrt{d\kappa}LR} \leq \max\left(\frac{\sqrt{\frac{LR^2}{\varepsilon}}\varepsilon}{\sqrt{d\kappa}LR}, \frac{\sqrt{\frac{d\kappa\sigma_*^2R^2}{\varepsilon^2}}\varepsilon}{\sqrt{d\kappa}LR}\right) = \max\left(\frac{\varepsilon^{1/2}}{\sqrt{d\kappa}L}, \frac{\sigma_*}{L}\right) \\ \kappa_\beta^2Lh^{2(\beta-1)} \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta^2L}\right)^{\frac{1}{2(\beta-1)}} \\ \kappa_\beta Lh^{\beta-1}R \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\ h &= \min\left(\max\left(\frac{\varepsilon^{1/2}}{\sqrt{d\kappa}L}, \frac{\sigma_*}{L}\right), \left(\frac{\varepsilon}{\kappa_\beta^2L}\right)^{\frac{1}{2(\beta-1)}}, \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\right) = \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}.\end{aligned}$$

From ⑤, we find Δ

$$\begin{aligned}\frac{\sqrt{\kappa}d\Delta R}{h\sqrt{N}} \leq \varepsilon &\implies \Delta \leq \frac{\varepsilon h\sqrt{N}}{\sqrt{\kappa}dR} \leq \max\left(\frac{\varepsilon\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\sqrt{\frac{LR^2}{\varepsilon}}}{\sqrt{\kappa}dR}, \frac{\varepsilon\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\sqrt{\frac{d\kappa\sigma_*^2R^2}{\varepsilon^2}}}{\sqrt{\kappa}dR}\right) \\ &= \max\left(\frac{\varepsilon^{1/2+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{\sqrt{\kappa}d(\kappa_\beta R)^{\frac{1}{\beta-1}}}, \frac{\varepsilon^{\frac{1}{\beta-1}}\sigma_*}{\sqrt{d\kappa}^{\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}}\right) = \frac{\varepsilon^{\frac{1}{\beta-1}}\sigma_*}{\sqrt{d\kappa}^{\frac{1}{\beta-1}}L^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}} \\ T = B \cdot N &= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2R^2}{\varepsilon^2}\right)\right).\end{aligned}$$

Case 2: $N > B > 1$

$$\mathbb{E}[f(x_N^{ag}) - f^*] = \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\sqrt{d\kappa}\sigma_*R}{\sqrt{BN}} + \frac{\sqrt{d\kappa}LhR}{\sqrt{BN}} + \frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} + \kappa_\beta Lh^{\beta-1}R + \kappa_\beta^2Lh^{2(\beta-1)} \leq 7\varepsilon.$$

From ①, ②, ③, we find N

$$\begin{aligned}\frac{LR^2}{N^2} \leq \varepsilon &\implies N \geq \sqrt{\frac{LR^2}{\varepsilon}} \\ \frac{LR^2}{BN} \leq \varepsilon &\implies N \geq \frac{LR^2}{B\varepsilon} \\ \frac{\sqrt{d\kappa}\sigma_*R}{\sqrt{BN}} \leq \varepsilon &\implies N \geq \frac{d\kappa\sigma_*^2R^2}{B\varepsilon^2} \\ N &= \mathcal{O}\left(\max\left(\frac{LR^2}{B\varepsilon}, \sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2R^2}{B\varepsilon^2}\right)\right).\end{aligned}$$

From ④, ⑥, ⑦, we find h

$$\begin{aligned}
\frac{\sqrt{d\kappa}LhR}{\sqrt{BN}} \leq \varepsilon &\implies h \leq \frac{\sqrt{BN}\varepsilon}{\sqrt{d\kappa}LR} = \max\left(\frac{\sqrt{B\frac{LR^2}{B\varepsilon}}\varepsilon}{\sqrt{d\kappa}LR}, \frac{\sqrt{B\sqrt{\frac{LR^2}{\varepsilon}}}\varepsilon}{\sqrt{d\kappa}LR}, \frac{\sqrt{B\frac{d\kappa\sigma_*^2R^2}{B\varepsilon^2}}\varepsilon}{\sqrt{d\kappa}LR}\right) \\
&= \max\left(\sqrt{\frac{\varepsilon}{d\kappa}L}, \frac{\varepsilon^{3/4}B^{1/2}}{\sqrt{d\kappa}L^{3/4}R^{1/2}}, \frac{\sigma_*}{L}\right) \\
\kappa_\beta^2Lh^{2(\beta-1)} \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta^2L}\right)^{\frac{1}{2(\beta-1)}} \\
\kappa_\beta Lh^{\beta-1}R \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\
h &= \min\left(\max\left(\sqrt{\frac{\varepsilon}{d\kappa}L}, \frac{\varepsilon^{3/4}B^{1/2}}{\sqrt{d\kappa}L^{3/4}R^{1/2}}, \frac{\sigma_*}{L}\right), \left(\frac{\varepsilon}{\kappa_\beta^2L}\right)^{\frac{1}{2(\beta-1)}}, \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\right) = \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}.
\end{aligned}$$

From ⑤, we find Δ

$$\begin{aligned}
\frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} \leq \varepsilon &\implies \Delta \leq \frac{h\sqrt{BN}\varepsilon}{\sqrt{\kappa}dR} \\
&\leq \max\left(\frac{h\left(\sqrt{\frac{LR^2}{\varepsilon}}\right)\varepsilon}{\sqrt{\kappa}dR}, \frac{h\left(\sqrt{B\sqrt{\frac{LR^2}{\varepsilon}}}\right)\varepsilon}{\sqrt{\kappa}dR}, \frac{h\left(\sqrt{B\frac{d\kappa\sigma_*^2R^2}{B\varepsilon^2}}\right)\varepsilon}{\sqrt{\kappa}dR}\right) \\
&= \max\left(\frac{h\left(\sqrt{L\varepsilon}\right)}{\sqrt{\kappa}d}, \frac{h\left(\sqrt{B\sqrt{L}}\right)\varepsilon^{3/4}}{\sqrt{\kappa}dR^{1/2}}, \frac{\sigma_*h}{\sqrt{d}}\right) \\
&= \max\left(\frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\left(\sqrt{L\varepsilon}\right)}{\sqrt{\kappa}d}, \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\left(\sqrt{B\sqrt{L}}\right)\varepsilon^{3/4}}{\sqrt{\kappa}dR^{1/2}}, \frac{\sigma_*}{\sqrt{d}}\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\right) \\
&= \max\left(\frac{\varepsilon^{1/2+\frac{1}{\beta-1}}L^{1/2-\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{\frac{1}{\beta-1}}}, \frac{L^{1/4-\frac{1}{\beta-1}}\sqrt{B}\varepsilon^{3/4+\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}}R^{1/2+\frac{1}{\beta-1}}}, \frac{\sigma_*}{\sqrt{d}}\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\right) = \frac{\sigma_*}{\sqrt{d}}\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\
T = N \cdot B &= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, B\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2R^2}{\varepsilon^2}\right)\right).
\end{aligned}$$

Case 3: $B = N$

$$\mathbb{E}[f(x_N^{ag}) - f^*] = \frac{LR^2}{N^2} + \frac{LR^2}{N^2} + \frac{\sqrt{d\kappa}\sigma_*R}{N} + \frac{\sqrt{d\kappa}LhR}{N} + \frac{\sqrt{\kappa}d\Delta R}{hN} + \kappa_\beta Lh^{\beta-1}R + \kappa_\beta^2Lh^{2(\beta-1)} \leq 7\varepsilon$$

From ①, ②, ③, we find N

$$\begin{aligned}
\frac{LR^2}{N^2} \leq \varepsilon &\implies N \geq \sqrt{\frac{LR^2}{\varepsilon}} \\
\frac{\sqrt{d\kappa}\sigma_*R}{N} \leq \varepsilon &\implies N \geq \frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon} \\
N &= \mathcal{O}\left(\max\left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon}\right)\right).
\end{aligned}$$

From ④, ⑥, ⑦, we find h

$$\begin{aligned}
\frac{\sqrt{d\kappa}LhR}{N} \leq \varepsilon &\implies h \leq \frac{\varepsilon N}{\sqrt{d\kappa}LR} = \max\left(\frac{\varepsilon}{\sqrt{d\kappa}LR} \sqrt{\frac{LR^2}{\varepsilon}}, \frac{\varepsilon}{\sqrt{d\kappa}LR} \frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon}\right) \\
&= \max\left(\sqrt{\frac{\varepsilon}{d\kappa L}}, \frac{\sigma_*}{L}\right) \\
\kappa_\beta^2 Lh^{2(\beta-1)} \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}} \\
\kappa_\beta Lh^{\beta-1}R \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\
h &= \min\left(\max\left(\sqrt{\frac{\varepsilon}{d\kappa L}}, \frac{\sigma_*}{L}\right), \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}}, \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}\right) = \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}.
\end{aligned}$$

From ⑤, we find Δ

$$\begin{aligned}
\frac{\sqrt{\kappa}d\Delta R}{hN} \leq \varepsilon &\implies \Delta \leq \frac{hN\varepsilon}{\sqrt{\kappa}dR} = \max\left(\frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \varepsilon}{\sqrt{\kappa}dR} \sqrt{\frac{LR^2}{\varepsilon}}, \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \varepsilon}{\sqrt{\kappa}dR} \frac{\sqrt{d\kappa}\sigma_*R}{\varepsilon}\right) \\
&= \max\left(\frac{\varepsilon^{1/2+\frac{1}{\beta-1}} L^{1/2-\frac{1}{\beta-1}}}{\sqrt{\kappa}d\kappa_\beta^{\frac{1}{\beta-1}} R^{\frac{1}{\beta-1}}}, \frac{\sigma_*\varepsilon^{\frac{1}{\beta-1}}}{\kappa_\beta^{\frac{1}{\beta-1}} \sqrt{dR}^{\frac{1}{\beta-1}} L^{\frac{1}{\beta-1}}}\right) = \frac{\sigma_*\varepsilon^{\frac{1}{\beta-1}}}{\kappa_\beta^{\frac{1}{\beta-1}} \sqrt{dR}^{\frac{1}{\beta-1}} L^{\frac{1}{\beta-1}}} \\
T = N \cdot B &= \mathcal{O}\left(\max\left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}\right)\right).
\end{aligned}$$

Case 4: $B > N$

$$\mathbb{E}[f(x_N^{ag}) - f^*] = \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\sqrt{d\kappa}\sigma_*R}{\sqrt{BN}} + \frac{\sqrt{d\kappa}LhR}{\sqrt{BN}} + \frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} + \kappa_\beta Lh^{\beta-1}R + \kappa_\beta^2 Lh^{2(\beta-1)} \leq 7\varepsilon.$$

From ①, ②, ③, we find N and B . Note that if $B > N$ then $\frac{LR^2}{N^2} > \frac{LR^2}{BN}$

$$\begin{aligned}
\frac{LR^2}{N^2} \leq \varepsilon &\implies B > N \geq \sqrt{\frac{LR^2}{\varepsilon}} \\
\frac{\sqrt{d\kappa}\sigma_*R}{\sqrt{BN}} \leq \varepsilon &\implies B \geq \frac{d\kappa\sigma_*^2 R^2}{N\varepsilon^2} = \mathcal{O}\left(\frac{d\kappa\sigma_*^2 R}{L^{1/2}\varepsilon^{3/2}}\right).
\end{aligned}$$

From ④, ⑥, ⑦, we find h

$$\begin{aligned}
\frac{\sqrt{d\kappa}LhR}{\sqrt{BN}} \leq \varepsilon &\implies h \leq \frac{\varepsilon\sqrt{BN}}{\sqrt{d\kappa}LR} \leq \frac{\varepsilon\sqrt{B}^4 \sqrt{\frac{LR^2}{\varepsilon}}}{\sqrt{d\kappa}LR} = \frac{\varepsilon^{3/4}\sqrt{B}}{\sqrt{d\kappa}L^{3/4}R^{1/2}} \\
\kappa_\beta^2 Lh^{2(\beta-1)} \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}} \\
\kappa_\beta Lh^{\beta-1}R \leq \varepsilon &\implies h \leq \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \\
h &= \min\left(\left(\frac{\varepsilon}{\kappa_\beta^2 L}\right)^{\frac{1}{2(\beta-1)}}, \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}, \frac{\varepsilon^{3/4}\sqrt{B}}{\sqrt{d\kappa}L^{3/4}R^{1/2}}\right) = \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}}.
\end{aligned}$$

From ⑤, we find Δ

$$\begin{aligned} \frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} \leq \varepsilon &\implies \Delta \leq \frac{h\sqrt{BN}\varepsilon}{\sqrt{\kappa}dR} \leq \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \left(\sqrt{B}^4 \sqrt{\frac{LR^2}{\varepsilon}}\right) \varepsilon}{\sqrt{\kappa}dR} \\ &= \frac{\left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{1}{\beta-1}} \left(\sqrt{B}^4 \sqrt{\frac{LR^2}{\varepsilon}}\right) \varepsilon}{\sqrt{\kappa}dR} = \frac{\varepsilon^{3/4 + \frac{1}{\beta-1}} \sqrt{B} L^{1/4 - \frac{1}{\beta-1}}}{\sqrt{\kappa} d \kappa_\beta^{\frac{1}{\beta-1}} R^{1/2 + \frac{1}{\beta-1}}} \end{aligned}$$

or we can get the batch size B via the maximum noise level Δ :

$$\begin{aligned} \frac{\sqrt{\kappa}d\Delta R}{h\sqrt{BN}} \leq \varepsilon &\implies B \geq \frac{\kappa d^2 \Delta^2 R^2}{\varepsilon^2 h^2 N} \geq \frac{\kappa d^2 \Delta^2 R^2}{\varepsilon^2 \left(\frac{\varepsilon}{\kappa_\beta LR}\right)^{\frac{2}{\beta-1}} \sqrt{\frac{LR^2}{\varepsilon}}} \\ &= \frac{\kappa_\beta^{\frac{2}{\beta-1}} \kappa d^2 \Delta^2 R^{1 + \frac{2}{\beta-1}}}{\varepsilon^{3/2 + \frac{2}{\beta-1}} L^{1/2 - \frac{2}{\beta-1}}} \\ B &= \mathcal{O} \left(\max \left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\sigma_*^2 R}{L^{1/2}\varepsilon^{3/2}}, \frac{\kappa_\beta^{\frac{2}{\beta-1}} \kappa d^2 \Delta^2 R^{1 + \frac{2}{\beta-1}}}{\varepsilon^{3/2 + \frac{2}{\beta-1}} L^{1/2 - \frac{2}{\beta-1}}} \right) \right) \\ T = N \cdot B &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\sigma_*^2 R^2}{\varepsilon^2}, \frac{\kappa_\beta^{\frac{2}{\beta-1}} \kappa d^2 \Delta^2 L^{\frac{2}{\beta-1}} R^{2 + \frac{2}{\beta-1}}}{\varepsilon^{2 + \frac{2}{\beta-1}}} \right) \right). \end{aligned}$$

That ends the proof. □

A.4 Proof of Remark 2

Bias of gradient approximation

$$\begin{aligned} b &= \|\mathbb{E}[\mathbf{g}(x_k, \mathbf{e}, r, \xi)] - \nabla f(x_k)\|_p = \left\| \mathbb{E} \left[d \frac{f_\delta(x_k + hr\mathbf{e}, \xi) - f_\delta(x_k - hr\mathbf{e}, \xi)}{2h} K(r)\mathbf{e} \right] - \nabla f(x_k) \right\|_p \\ &\stackrel{\textcircled{1}}{=} \left\| \mathbb{E} \left[d \frac{f(x_k + hr\mathbf{e}, \xi)}{h} K(r)\mathbf{e} + \delta(x_k + h\mathbf{e}) \right] - \nabla f(x_k) \right\|_p \\ &\stackrel{\textcircled{2}}{\leq} \left\| \mathbb{E} \left[d \frac{f(x_k + hr\mathbf{e}, \xi)}{h} K(r)\mathbf{e} \right] - \nabla f(x_k) \right\|_p \\ &\stackrel{\textcircled{3}}{=} \|\mathbb{E}[(\nabla f(x_k + hr\mathbf{u}, \xi)rK(r)) - \nabla f(x_k)]\|_p \\ &\stackrel{\textcircled{4}}{=} \sup_{z \in S_p^d(1)} \mathbb{E}[(\nabla_z f(x_k + hr\mathbf{u}, \xi) - \nabla_z f(x_k))rK(r)] \\ &\stackrel{\textcircled{5}}{\leq} \kappa_\beta h^{\beta-1} \frac{L}{(l-1)!} \mathbb{E}[\|\mathbf{u}\|_p^{\beta-1}] \lesssim \kappa_\beta L h^{\beta-1}, \end{aligned}$$

where $\mathbf{u} \in B_2^d(1)$, ①: distribution of e is symmetric; ②: independence of the stochastic noise; ③: a version of the Stokes' theorem [30] (see Section 13.3.5, Exercise 14a); ④: norm of the gradient is the supremum of directional derivatives $\nabla_z f(x) = \lim_{\varepsilon \rightarrow 0} \frac{f(x+\varepsilon z) - f(x)}{\varepsilon}$; ⑤: Taylor expansion.

Bounding second moment of gradient approximation

The calculations can be taken from Remark 1 without changes.

Convergence

$$\begin{aligned}
\mathbb{E}[f(x_N^{ag}) - f^*] &= c \left(\frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\zeta R}{\sqrt{BN}} + bR + \frac{b^2}{2L}N \right) \\
&\lesssim \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\left(\sqrt{d\kappa\kappa'(p,d)\sigma_*^2} + \sqrt{d\kappa\kappa'(p,d)L^2h^2} + \sqrt{\frac{d^2\kappa\kappa'(p,d)\Delta^2}{h^2}} \right) R}{\sqrt{BN}} + (\kappa_\beta Lh^{\beta-1})R \\
&+ \frac{(\kappa_\beta Lh^{\beta-1})^2}{L}N \leq \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\left(\sqrt{d\kappa\kappa'(p,d)\sigma_*} + \sqrt{d\kappa\kappa'(p,d)Lh} + \sqrt{\kappa\kappa'(p,d)\frac{d\Delta}{h}} \right) R}{\sqrt{BN}} \\
&+ (\kappa_\beta Lh^{\beta-1})R + \frac{\kappa_\beta^2 L^2 h^{2(\beta-1)}}{L}N = \frac{LR^2}{N^2} + \frac{LR^2}{BN} + \frac{\sqrt{d\kappa\kappa'(p,d)\sigma_*}R}{\sqrt{BN}} \\
&+ \frac{\sqrt{d\kappa\kappa'(p,d)Lh}R}{\sqrt{BN}} + \frac{\sqrt{\kappa\kappa'(p,d)d\Delta}R}{h\sqrt{BN}} + \kappa_\beta Lh^{\beta-1}R + \kappa_\beta^2 Lh^{2(\beta-1)}.
\end{aligned}$$

By performing the same transformations as in Theorem 2, we get the following parameters:

Case 1: $B = 1$

$$\begin{aligned}
N &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2} \right) \right) \\
h &\leq \left(\frac{\varepsilon}{\kappa_\beta LR} \right)^{\frac{1}{\beta-1}} \\
\Delta &\leq \frac{\varepsilon^{\frac{1}{\beta-1}} \sigma_*}{\sqrt{d\kappa_\beta^{\frac{1}{\beta-1}} L^{\frac{1}{\beta-1}} R^{\frac{1}{\beta-1}}}} \\
T &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2} \right) \right).
\end{aligned}$$

Case 2: $N > B > 1$

$$\begin{aligned}
N &= \mathcal{O} \left(\max \left(\frac{LR^2}{B\varepsilon}, \sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{B\varepsilon^2} \right) \right) \\
h &\leq \left(\frac{\varepsilon}{\kappa_\beta LR} \right)^{\frac{1}{\beta-1}} \\
\Delta &\leq \frac{\varepsilon^{\frac{1}{\beta-1}} \sigma_*}{\sqrt{d\kappa_\beta^{\frac{1}{\beta-1}} L^{\frac{1}{\beta-1}} R^{\frac{1}{\beta-1}}}} \\
T &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, B\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2} \right) \right).
\end{aligned}$$

Case 3: $B = N$

$$\begin{aligned}
 N &= \mathcal{O} \left(\max \left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{\sqrt{d\kappa\kappa'(p,d)\sigma_*R}}{\varepsilon} \right) \right) \\
 h &\leq \left(\frac{\varepsilon}{\kappa_\beta LR} \right)^{\frac{1}{\beta-1}} \\
 \Delta &\leq \frac{\sigma_*\varepsilon^{\frac{1}{\beta-1}}}{\kappa_\beta^{\frac{1}{\beta-1}} \sqrt{d} R^{\frac{1}{\beta-1}} L^{\frac{1}{\beta-1}}} \\
 T &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2} \right) \right).
 \end{aligned}$$

Case 4: $B > N$

$$\begin{aligned}
 N &= \mathcal{O} \left(\sqrt{\frac{LR^2}{\varepsilon}} \right) \\
 B &= \mathcal{O} \left(\max \left(\sqrt{\frac{LR^2}{\varepsilon}}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R}{L^{1/2}\varepsilon^{3/2}}, \frac{\kappa_\beta^{\frac{2}{\beta-1}} \kappa\kappa'(p,d)d^2\Delta^2 R^{1+\frac{2}{\beta-1}}}{\varepsilon^{3/2+\frac{2}{\beta-1}} L^{1/2-\frac{2}{\beta-1}}} \right) \right) \\
 h &\leq \left(\frac{\varepsilon}{\kappa_\beta LR} \right)^{\frac{1}{\beta-1}} \\
 \Delta &\leq \frac{\varepsilon^{3/4+\frac{1}{\beta-1}} \sqrt{B} L^{1/4-\frac{1}{\beta-1}}}{\sqrt{\kappa\kappa'(p,d)} d \kappa_\beta^{\frac{1}{\beta-1}} R^{1/2+\frac{1}{\beta-1}}} \\
 T &= \mathcal{O} \left(\max \left(\frac{LR^2}{\varepsilon}, \frac{d\kappa\kappa'(p,d)\sigma_*^2 R^2}{\varepsilon^2}, \frac{\kappa_\beta^{\frac{2}{\beta-1}} \kappa\kappa'(p,d)d^2\Delta^2 L^{\frac{2}{\beta-1}} R^{2+\frac{2}{\beta-1}}}{\varepsilon^{2+\frac{2}{\beta-1}}} \right) \right).
 \end{aligned}$$

By substituting $\kappa'(p, d)$ into formulas, we prove the remark. □