Volume 3 (2024), Pages 35–43

# Computing the permanental polynomial of 4k-intercyclic bipartite graphs

Ravindra B. Bapat, Ranveer Singh, and Hitesh Wankhede\*

(Communicated by Sudipta Mallik)

#### Abstract

Let G be a bipartite graph with adjacency matrix A(G). The characteristic polynomial  $\phi(G,x) = \det(xI - A(G))$  and the permanental polynomial  $\pi(G,x) = \operatorname{per}(xI - A(G))$  are both graph invariants used to distinguish graphs. For bipartite graphs, we define the modified characteristic polynomial, which is obtained by changing the signs of some of the coefficients of  $\phi(G,x)$ . For 4k-intercyclic bipartite graphs, i.e., those for which the removal of any 4k-cycle results in a  $C_{4k}$ -free graph, we provide an expression for  $\pi(G,x)$  in terms of the modified characteristic polynomial of the graph and its subgraphs. Our approach is purely combinatorial in contrast to the Pfaffian orientation method found in the literature to compute the permanental polynomial.

## 1 Introduction and preliminaries

We consider simple and undirected graphs. Let G be a graph with the vertex set  $V(G) = \{v_1, v_2, \ldots, v_n\}$ . The adjacency matrix  $A(G) = (a_{i,j})$  of a graph G is defined such that  $a_{i,j} = 1$  if  $v_i$  and  $v_j$  are adjacent and 0 otherwise where  $i, j \in \{1, 2, \ldots, n\}$ . The determinant and the permanent of A(G), are defined as

$$\det(A(G)) = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \prod_{i=1}^n a_{i,\sigma(i)} \text{ and } \operatorname{per}(A(G)) = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i,\sigma(i)},$$

respectively, where  $S_n$  is the set of all permutations of the set  $\{1, 2, ..., n\}$  and  $\operatorname{sgn}(\sigma)$  is the signature of the permutation  $\sigma$ . While the determinant can be computed in polynomial time using the Gaussian elimination method, and the fastest known algorithm runs in  $\mathcal{O}(n^{2.371552})$  time [3, 19], computing the permanent is notoriously difficult, as it is #P-complete [18]. The

MSC2020: 05C31, 05C50, 05C85; Keywords: Permanental polynomial, 4k-intercyclic bipartite graph Received Mar 27, 2024; Revised Nov 16, 2024; Accepted Nov 18, 2024; Published Nov 20, 2024  $\bigcirc$  The author(s). Released under the CC BY 4.0 International License

<sup>\*</sup> Corresponding author

"Permanent vs. Determinant Problem" about symbolic matrices in computational complexity theory is as follows: "Can we express the permanent of a matrix as the determinant of a (possibly polynomially larger) matrix?" For an upper bound on the size of the larger matrix, see [10], and for a survey on lower bounds, see [1]. The problem "Given a (0,1)-matrix A, under what conditions, changing the sign of some the nonzero entries yields a matrix B such that per(A) = det(B)?" is famously known as "Polya Permanent Problem," [15] and it is equivalent to twenty-three other problems listed in [13]. Immanants are matrix functions that generalize determinant and permanent, and their complexity dichotomy was also recently studied [7].

The characteristic polynomial and the permanental polynomial of graph G on n vertices are univariate polynomials defined as

$$\phi(G, x) = \det(xI - A(G))$$
 and  $\pi(G, x) = \operatorname{per}(xI - A(G))$ ,

respectively, where I is the identity matrix of order n. The characteristic and the permanental polynomials are graph invariants, and they can be used to distinguish graphs towards Graph Isomorphism Problem [17]. But the permanental polynomial is not studied in great detail as compared to the characteristic polynomial, probably due to its computational difficulty. Some computational evidence suggests that the permanental polynomial is better than the characteristic polynomial while distinguishing graphs [8, 12]. We are interested in finding ways to compute the permanental polynomial efficiently; one way to do that is by expressing the permanental polynomial in terms of the characteristic polynomial. For an excellent survey on the permanental polynomial, we refer to [11].

Let

$$\phi(G, x) = \sum_{i=0}^{n} a_i x^{n-i} \text{ and } \pi(G, x) = \sum_{i=0}^{n} b_i x^{n-i}.$$

The interpretation of these coefficients is given as

$$a_i = \sum_{U_i} (-1)^{p(U_i)} 2^{c(U_i)} [16] , b_i = (-1)^i \sum_{U_i} 2^{c(U_i)} [14] ,$$
 (1.1)

where the summation is taken over all the Sachs subgraphs  $U_i$  (subgraphs whose components are either cycles or edges) of G on i vertices,  $p(U_i)$  denotes the number of components in  $U_i$ , and  $c(U_i)$  denotes the number of components in  $U_i$  that are cycles. It is easy to see the following.

**Proposition 1.1.** [5, 6] A graph G is bipartite if and only if  $a_i = b_i = 0$  for each odd i.

Hence, for a bipartite graph G, we have

$$\phi(G, x) = \sum_{i=0,2,4,\dots} a_i x^{n-i} \text{ and } \pi(G, x) = \sum_{i=0,2,4,\dots} b_i x^{n-i}.$$
 (1.2)

Defining  $f_i = b_i - (-1)^{i/2}a_i$  for each nonnegative even integer i, we introduce a modified characteristic polynomial and also a new graph polynomial for bipartite graphs as

$$\phi_p(G, x) = \sum_{i=0,2,4,\dots} (-1)^{i/2} a_i x^{n-i} \text{ and } f(G, x) = \sum_{i=0,2,4,\dots} f_i x^{n-i},$$

respectively, such that we have

$$\pi(G, x) = \phi_p(G, x) + f(G, x).$$
 (1.3)

We denote a cycle of length k by  $C_k$ . For each positive integer k, we refer to a  $C_{4k}$  by a 4k-cycle. A graph is called  $C_{4k}$ -free if it does not contain a 4k-cycle for all positive integers k. A graph is called 4k-intercyclic if it does not contain two vertex-disjoint 4k-cycles (these 4k-cycles could be of different length). Equivalently, after removal of the vertices of any 4k-cycle from a 4k-intercyclic graph, the resultant graph is  $C_{4k}$ -free (see Figure 1 for an example).




Figure 1: Example of a 4k-intercyclic bipartite graph.

In 1985, Borowiecki proved the following.

**Theorem 1.2.** [5] Let G be a bipartite graph with the spectrum  $\{\lambda_1, \lambda_2, ..., \lambda_n\}$ . Then, G is  $C_{4k}$ -free if and only if its per-spectrum is  $\{i\lambda_1, i\lambda_2, ..., i\lambda_n\}$ .

By inspecting the proof of this theorem, we notice that a bipartite graph G is  $C_{4k}$ -free if and only if  $\pi(G, x) = \phi_p(G, x)$  (see Corollary 2.4). As a result, the permanental polynomial of  $C_{4k}$ -free bipartite graphs can be computed directly through the modified characteristic polynomial. Yan and Zhang, in 2004, found that the permanental polynomial of a larger class of bipartite graphs can be computed using the characteristic polynomial of some oriented graph. They proved the following.

**Theorem 1.3.** [20] Let G be a bipartite graph with n vertices containing no subgraph that is an even subdivision of  $K_{2,3}$ . Then there exists an orientation  $G^e$  of G such that  $\pi(G,x) = \det(xI - A(G^e))$ , where  $A(G^e)$  denotes the skew adjacency matrix of  $G^e$ .

Later Zhang and Li, in 2012, proved the converse of this statement.

**Theorem 1.4.** [21] There exists an orientation  $G^e$  of a bipartite graph G such that  $\pi(G, x) = \det(xI - A(G^e))$  if and only if G contains no even subdivision of  $K_{2,3}$ .

For the definition of an even subdivision of a graph, see [20, 21]. Zhang and Li also show that bipartite graphs that do not contain an even subdivision of  $K_{2,3}$  are planar and admit Pfaffian orientation. They also give characterization and recognition of such graphs, which leads to a polynomial time algorithm to compute the permanental polynomial of such bipartite graphs. Next, we reformulate Theorem 1.4.

<sup>&</sup>lt;sup>1</sup>The spectrum and the per-spectrum of a graph G are the multiset of all roots of its characteristic polynomial and the permanental polynomial, respectively, and i is an imaginary unit.

**Theorem 1.5.** There exists an orientation  $G^e$  of a graph G such that  $\pi(G, x) = \det(xI - A(G^e))$  if and only if G is a bipartite graph containing no even subdivision of  $K_{2,3}$ .

Proof. Suppose that there is an orientation  $G^e$  such that  $\pi(G,x) = \det(xI - A(G^e))$  holds. It is enough to show that G is bipartite. The conclusion of the theorem then follows by Theorem 1.4. Since the skew-adjacency matrix  $A(G^e)$  is skew-symmetric, it has purely imaginary eigenvalues. Hence, the permanental polynomial can be expressed as  $\pi(G,x) = (x - i\lambda_1)(x - i\lambda_2) \dots (x - i\lambda_n)$  for some real numbers  $\lambda_1, \lambda_2, \dots, \lambda_n$ . When k is odd, the imaginary unit i is a factor of the coefficient  $b_k$  of  $x^{n-k}$ . Since the coefficients of  $\pi(G,x)$  must be real, it follows that  $b_k = 0$  for all odd k, and by Proposition 1.1, G is bipartite.  $\square$ 

Theorem 1.5 suggests that the orientation approach in computing the permanental polynomial only works for the class of bipartite graphs that do not contain an even subdivision of  $K_{2,3}$ . In this article, we give a formula to compute  $\pi(G,x)$  for the class of 4k-intercyclic bipartite graphs (a superset of the class of  $C_{4k}$ -free bipartite graphs). This is done by expressing f(G,x) in terms of the modified characteristic polynomial of the subgraphs of G. Our approach is combinatorial rather than based on Pfaffian orientation. Note that the class of 4k-intercyclic bipartite graphs is different from and not a subset of the class of bipartite graphs that do not contain an even subdivision of  $K_{2,3}$ .

We would also like to mention that our result seem to be in the same spirit as Polya's scheme completed by Galluccio and Loebl in 1999. They proved that the generating function of the perfect matchings of a graph of genus g may be written as a linear combination of  $4^g$  Pfaffians, and as a consequence obtained the following result.

**Theorem 1.6.** [9] Let A be a square matrix. Then per(A) may be expressed as a linear combination of terms of the form  $det(A^i)$ ,  $i = 1, ..., 4^g$ , where each  $A^i$  is obtained from A by changing the sign of some entries and g is the genus of the bipartite graph corresponding to the biadjacency matrix A.

## 2 Main result and application

**Theorem 2.1.** Let G be a 4k-intercyclic bipartite graph. Then,

$$\pi(G, x) = \phi_p(G, x) + 4 \sum_{R \in \mathcal{C}_{4k}(G)} \phi_p(G \backslash R, x),$$

where  $C_{4k}(G)$  denotes the set of all 4k-cycles in G.

To prove this theorem, we need the following lemma.

**Lemma 2.2.** Let G be a bipartite graph. Then, for each nonnegative even integer i, we have

$$f_i = \sum_{j=1,3,5,\dots} 2^{j+1} \sum_{\substack{U_i \text{ containing} \\ \text{exactly } j \text{ } 4k \text{ -cycles}}} 2^{t(U_i)},$$

where  $U_i$  denotes a Sachs subgraph on i vertices, and  $t(U_i)$  is the number of (4k+2)-cycles in  $U_i$ .

Proof of Lemma 2.2. In a bipartite graph, there can be two types of cycles: 4k-cycles or (4k+2)-cycles. Let  $U_i$  be any Sachs subgraph on i vertices, and  $s(U_i)$  and  $t(U_i)$  be the number of 4k-cycles and (4k+2)-cycles in it respectively. Then,  $U_i$  can be expressed as

$$U_i = \{C_{4k_1} \cup \dots \cup C_{4k_{s(U_i)}}\} \cup \{C_{4l_1+2} \cup \dots \cup C_{4l_{t(U_i)}+2}\} \cup \underbrace{\{K_2 \cup \dots \cup K_2\}}_{r(U_i)-\text{times}},$$

such that  $p(U_i) = s(U_i) + t(U_i) + r(U_i), c(U_i) = s(U_i) + t(U_i)$  and

$$i = 4(k_1 + \dots + k_{s(U_i)}) + 4(l_1 + \dots + l_{t(U_i)}) + 2(t(U_i) + r(U_i)).$$

Check that  $s(U_i) + t(U_i) + r(U_i) \equiv i/2 + s(U_i) \pmod{2}$ . Using this fact, the coefficients of the characteristic polynomial and the permanental polynomial given in Equation 1.1 can be written as

$$(-1)^{i/2}a_i = \sum_{U_i} (-1)^{s(U_i)} 2^{s(U_i) + t(U_i)}$$
 and  $b_i = \sum_{U_i} 2^{s(U_i) + t(U_i)}$ ,

respectively  $((-1)^i = 1 \text{ since } i \text{ is even})$ . Since  $f_i = b_i - (-1)^{i/2}a_i$ , we get

$$f_{i} = \sum_{U_{i}} (1 - (-1)^{s(U_{i})}) 2^{s(U_{i}) + t(U_{i})} = \sum_{\substack{U_{i} \text{ containing an odd} \\ \text{number of } 4k \text{-cycles}}} 2^{s(U_{i}) + 1} 2^{t(U_{i})}$$

$$= \sum_{U_{i}} 2^{j+1} \sum_{U_{i}} 2^{t(U_{i})}.$$
(2.1)

$$= \sum_{j=1,3,5,\dots} 2^{j+1} \sum_{\substack{U_i \text{ containing} \\ \text{exactly } j \text{ $4k$-cycles}}} 2^{t(U_i)}.$$

Note that the contribution in Equation 2.1 of the Sachs subgraphs in which we have exactly an even number of 4k-cycles vanishes.

Proof of Theorem 2.1. Since G is 4k-intercyclic, the subgraph  $G \setminus R$  is  $C_{4k}$ -free for any  $R \in C_{4k}(G)$ . Similarly, any Sachs subgraph  $U_i$  of G can contain at most one 4k-cycle, that is,  $s(U_i) \leq 1$ . Using Lemma 2.2, we have

$$f_i = 4 \sum_{\substack{U_i \text{ containing} \\ \text{exactly one } 4k\text{-cycle}}} 2^{t(U_i)} = 4 \sum_{\substack{R \in \mathcal{C}_{4k}(G) \\ \text{containing } R}} 2^{t(U_i)}$$

for each nonnegative even integer i. For any fixed  $R \in \mathcal{C}_{4k}(G)$ , there is a one-to-one correspondence between the Sachs subgraphs in G containing R and the Sachs subgraphs in  $G \setminus R$ . To see this, consider a Sachs subgraph  $U_i$  in G containing R. Let  $l_R = i - |V(R)|$  and  $W_{l_R} = U_i \setminus R$  be the unique subgraph of  $G \setminus R$  corresponding to  $U_i$ . Since G is 4k-intercyclic, removing R ensures that  $W_{l_R}$  is a Sachs subgraph of  $G \setminus R$  since both  $G \setminus R$  and  $U_i \setminus R$  do not contain 4k-cycles. Conversely, adding R back to a Sachs subgraph  $W_{l_R}$  in  $G \setminus R$  uniquely reconstructs a corresponding subgraph  $U_i$  in G. This establishes a one-to-one correspondence. Moreover, the number of cycles in  $W_{l_R}$  is the same as the number (4k + 2)-cycles in  $U_i$ , i.e.,  $c(W_{l_R}) = t(U_i)$ . As a result, we have

$$f_i = 4 \sum_{R \in \mathcal{C}_{4k}(G)} \sum_{W_{l_R}} 2^{c(W_{l_R})}.$$

Now consider the polynomial

$$\begin{split} f(G,x) &= \sum_{i=0,2,4,\dots} f_i x^{n-i} = 4 \sum_{i=0,2,4,\dots} \sum_{R \in \mathcal{C}_{4k}(G)} \sum_{Wl_R} 2^{c(W_{l_R})} x^{n-i} \\ &= 4 \sum_{R \in \mathcal{C}_{4k}(G)} \sum_{l_R = 0,2,4,\dots} \sum_{Wl_R} 2^{c(W_{l_R})} x^{(n-|V(R)|) - l_R} \\ &= 4 \sum_{R \in \mathcal{C}_{4k}(G)} \pi(G \backslash R, x), \end{split}$$

where the second last step follows from the rearrangement of sums, and the last step by Equation 1.1, 1.2 and the fact that  $G\backslash R$  is a  $C_{4k}$ -free bipartite graph on n-|V(R)| vertices. By Equation 1.3, we conclude

$$\pi(G, x) = \phi_p(G, x) + 4 \sum_{R \in \mathcal{C}_{ak}(G)} \pi(G \backslash R, x).$$

Since  $G \setminus R$  is  $C_{4k}$ -free, the application of this expression to it leads to  $\pi(G \setminus R, x) = \phi_p(G \setminus R, x)$  proving the theorem.

**Example 2.3.** Consider the 4k-intercyclic graph G shown in Figure 1. It contains three 4-cycles and two 8-cycles, and removal of each of them from the graph yields the following subgraphs:  $P_5 \cup K_1$ ,  $P_4 \cup K_2$ ,  $P_4 \cup K_1 \cup K_1$ ,  $K_2$  and  $K_1 \cup K_1$ , respectively. Then, using Theorem 2.1,

$$\pi(G,x) = \phi_p(G,x) + 4(\phi_p(P_5 \cup K_1, x) + \phi_p(P_4 \cup K_2, x) + \phi_p(P_4 \cup K_1 \cup K_1, x) + \phi_p(K_2, x) + \phi_p(K_1 \cup K_1, x)).$$

We need to do the following computations:  $\phi_p(G, x) = x^{10} + 12x^8 + 40x^6 + 47x^4 + 18x^2 + 1$ ,  $\phi_p(P_5 \cup K_1, x) = x^6 + 4x^4 + 3x^2$ ,  $\phi_p(P_4 \cup K_2, x) = x^6 + 4x^4 + 4x^2 + 1$ ,  $\phi_p(P_4 \cup K_1, x) = x^6 + 3x^4 + x^2$ ,  $\phi_p(K_2) = x^2 + 1$ ,  $\phi_p(K_1 \cup K_1, x) = x^2$ . Hence, we get  $\pi(G, x) = x^{10} + 12x^8 + 52x^6 + 91x^4 + 58x^2 + 9$ . Note that Theorem 1.3 and 1.4 are not applicable for this graph G as it contains  $K_{2,3}$ .

The following corollary shows that Theorem 2.1 is a generalization of Borowiecki's proof idea for computational purposes at least.

Corollary 2.4. [5] A bipartite graph G is  $C_{4k}$ -free if and only if  $\pi(G, x) = \phi_p(G, x)$ .

*Proof.* The forward implication easily follows from Theorem 2.1. Suppose  $\pi(G, x) = \phi_p(G, x)$  holds, then from Equation 2.1, we have

$$f_i = \sum_{\substack{U_i \text{ with an odd} \\ \text{number of } 4k\text{-cycles}}} 2^{s(U_i)+1} 2^{t(U_i)} = 0,$$

for each *i*. Suppose, on the contrary, that G contains a 4k-cycle for some k. Then, there exists a Sachs subgraph  $U_{4k} = C_{4k}$ , and it contains an odd number of 4k-cycles. Hence,  $f_{4k} \neq 0$ , and we get a contradiction which concludes that G is  $C_{4k}$ -free.

By Theorem 2.1, the computation of permanental polynomial of a 4k-intercyclic bipartite graph requires listing all the 4k-cycles in it. For any graph on n vertices, all cycles of length up to  $\log n$  can be found in polynomial time using the color coding method of Alon, Yuster and Zwick [2]. By an algorithm of Birmelé et al. [4], listing all cycles in a graph requires  $\mathcal{O}(m + \sum_{c \in \mathcal{C}(G)} |c|)$  time, where m is the number of edges and  $\mathcal{C}(G)$  is the set of all cycles. Hence, the permanental polynomial of a 4k-intercyclic bipartite graph can be computed in polynomial time if the length of the largest 4k cycle is bounded by  $\log n$  or if the number of cycles is bounded by a polynomial in n.

We now discuss an application of Theorem 2.1 in constructing cospectral graphs. Recall that two graphs  $G_1$  and  $G_2$  are said to be cospectral if they have the same characteristic polynomial, that is,  $\phi(G_1, x) = \phi(G_2, x)$ . Similarly, we say that they are per-cospectral if they have the same permanental polynomial, that is,  $\pi(G_1, x) = \pi(G_2, x)$ . Since  $\phi_p(G, x)$  can be recovered from  $\phi(G, x)$ , it follows from Corollary 2.4 that two  $C_{4k}$ -free bipartite graphs  $G_1$  and  $G_2$  are cospectral if and only if they are per-cospectral [5]. Next, we give a general procedure to construct 4k-intercyclic bipartite graphs that are simultaneously cospectral as well as per-cospectral. Let p be some polynomial. Then, corresponding to p, we define a class of 4k-intercyclic bipartite graphs  $\mathcal{G}_p = \{G \mid f(G, x) = p\}$ .

**Theorem 2.5.** Let  $G_1, G_2 \in \mathcal{G}_p$  for some polynomial p. Then,  $G_1$  and  $G_2$  are cospectral if and only if they are per-cospectral.

```
Proof. Write \pi(G_1, x) = \phi_p(G_1, x) + f(G_1, x), and \pi(G_2, x) = \phi_p(G_2, x) + f(G_2, x). But since G_1, G_2 \in \mathcal{G}_p, we have f(G_1, x) = f(G_2, x). Hence, \pi(G_1, x) - \pi(G_2, x) = \phi_p(G_1, x) - \phi_p(G_2, x). By definition, \phi_p(G_1, x) = \phi_p(G_2, x) if and only if \phi(G_1, x) = \phi(G_2, x). Then, it follows that \phi(G_1, x) = \phi(G_2, x) if and only if \pi(G_1, x) = \pi(G_2, x).
```

Three such classes of 4k-intercyclic bipartite graphs are given below.

- 1. The class of all  $C_{4k}$ -free bipartite graphs. Let p = 0, then  $\mathcal{G}_0 = \{G \mid f(G, x) = 0\}$ . By Corollary 2.4, every graph in this class is  $C_{4k}$ -free.
- 2. The class of all bipartite graphs with exactly l  $C_4$ 's and no other cycle such that  $G \setminus C$  is an edgeless graph for any cycle C, where l is some positive integer. Let  $p = 4lx^{n-4}$  for a given n and l, then  $\mathcal{G}_{4lx^{n-4}} = \{G \mid f(G,x) = 4lx^{n-4}\}$ . By Theorem 2.1, we have  $\sum_{R \in C_{4k}(G)} \phi_p(G \setminus R, x) = lx^{n-4}$ . Now the degree  $x^{n-4}$  is achieved in  $\phi_p(G \setminus R, x)$  only when R is  $C_4$ . Hence, there are l  $C_4$ 's in G such that  $G \setminus C$  is an edgeless graph for any cycle C.
- 3. The class of all unicyclic bipartite graphs with a  $C_4$  such that any two graphs in this class are cospectral after the removal of  $C_4$ . This follows easily from Theorem 2.1.

For any such class  $\mathcal{G}_p$  of 4k-intercyclic bipartite graphs, the permanental polynomial is not any more useful than the characteristic polynomial in distinguishing them.

## Acknowledgments

This work is supported by the Department of Science and Technology (Govt. of India)

through project DST/04/2019/002676. The second author acknowledges the support from ANRF, SRG/2022/002219. A preliminary version of this article was presented as a poster by the second author at the European Conference on Combinatorics, Graph Theory, and Applications (EuroComb 2023) held in Prague. Part of this work was done while the last author was at IIT Indore. We would like to thank the anonymous reviewers for their constructive feedback, which greatly improved the presentation of this article. One of these comments brought to our attention a result by Galluccio and Loebl (Theorem 1.6). We also thank Noga Alon and George Manoussakis for helpful email conversation on listing cycles.

#### References

- [1] Manindra Agrawal, *Determinant versus permanent*, in Proceedings of the 25th International Congress of Mathematicians, ICM 2006, volume 3, 985–997 (2006).
- [2] Noga Alon, Raphael Yuster, and Uri Zwick, *Color-coding*, Journal of the ACM (JACM) 42 (1995) 844–856.
- [3] Alfred V. Aho and John E. Hopcroft, *The design and analysis of computer algorithms*, Pearson Education India (1974).
- [4] Etienne Birmelé, Rui Ferreira, Roberto Grossi, Andrea Marino, Nadia Pisanti, Romeo Rizzi, and Gustavo Sacomoto, *Optimal listing of cycles and st-paths in undirected graphs*, in Proceedings of the twenty-fourth annual ACM-SIAM symposium on Discrete algorithms, 1884–1896 (2013).
- [5] Mieczyslaw Borowiecki, On spectrum and per-spectrum of graphs, Publ. Inst. Math.(Beograd) 38 (1985) 31–33.
- [6] Dragoš M. Cvetković, Michael Doob, and Horst Sachs, Spectra of graphs: theory and application, Academic press (1979).
- [7] Radu Curticapean, A full complexity dichotomy for immanant families, in Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing, 1770–1783 (2021).
- [8] Matthias Dehmer, Frank Emmert-Streib, Bo Hu, Yongtang Shi, Monica Stefu, and Shailesh Tripathi, *Highly unique network descriptors based on the roots of the permanental polynomial*, Information Sciences 408 (2017) 176–181.
- [9] Anna Galluccio and Martin Loebl, On the theory of Pfaffian orientations. I. Perfect matchings and permanents, the Electronic Journal of Combinatorics (1999) R6–R6.
- [10] Bruno Grenet, An upper bound for the permanent versus determinant problem, Theory of Computing (2011).
- [11] Wei Li, Shunyi Liu, Tingzeng Wu, and Heping Zhang, On the permanental polynomials of graphs, Graph Polynomials (2016) 101–121, Chapman and Hall/CRC.

- [12] Shunyi Liu and Jinjun Ren, Enumeration of copermanental graphs, arXiv preprint arXiv:1411.0184 (2014).
- [13] William McCuaig, *Pólya's permanent problem*, the Electronic Journal of Combinatorics (2004) R79–R79.
- [14] Russell Merris, Kenneth R. Rebman, and William Watkins, *Permanental polynomials of graphs*, Linear Algebra and Its Applications 38 (1981) 273–288.
- [15] György Pólya, Aufgabe 424, Archiv der Mathematik und Physik 20 (1913) 271.
- [16] Horst Sachs, Beziehungen zwischen den in einem Graphen enthaltenen Kreisen und seinem charakteristischen Polynom, Publ. Math. Debrecen 11 (1964) 119–134.
- [17] Edwin R. Van Dam and Willem H. Haemers, Which graphs are determined by their spectrum?, Linear Algebra and its applications 373 (2003) 241–272.
- [18] Leslie G. Valiant, *The complexity of computing the permanent*, Theoretical computer science 8 (1979) 189–201.
- [19] Virginia Vassilevska Williams, Yinzhan Xu, Zixuan Xu, and Renfei Zhou, New bounds for matrix multiplication: from alpha to omega, in Proceedings of the 2024 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA), 3792–3835 (2024).
- [20] Weigen Yan and Fuji Zhang, On the permanental polynomials of some graphs, Journal of Mathematical Chemistry 35 (2004) 175–188.
- [21] Heping Zhang and Wei Li, Computing the permanental polynomials of bipartite graphs by Pfaffian orientation, Discrete Applied Mathematics 160 (2012) 2069–2074.

#### **Contact Information**

Ravindra B. Bapat Indian Statistical Institute, rbb@isid.ac.in New Delhi 110016, India.

https://orcid.org/0000-0002-1127-2912

Ranveer Singh Indian Institute of Technology Indore, ranveer@iiti.ac.in Indore 453552, India.

https://orcid.org/0000-0002-4566-0331

Hitesh Wankhede The Institute of Mathematical Sciences (HBNI), hiteshwankhede9@gmail.com Chennai 600113, India.

https://orcid.org/0000-0003-3431-6305