

ON MEAN FIELD GAMES IN INFINITE DIMENSION

SALVATORE FEDERICO, FAUSTO GOZZI, AND ANDRZEJ ŚWIĘCH

ABSTRACT. We study a Mean Field Games (MFG) system in a real, separable infinite dimensional Hilbert space. The system consists of a second order parabolic type equation, called Hamilton-Jacobi-Bellman (HJB) equation in the paper, coupled with a nonlinear Fokker-Planck (FP) equation. Both equations contain a Kolmogorov operator. Solutions to the HJB equation are interpreted in the mild solution sense and solutions to the FP equation are interpreted in an appropriate weak sense. We prove well-posedness of the considered MFG system under certain conditions. The existence of a solution to the MFG system is proved using Tikhonov's fixed point theorem in a proper space. Uniqueness of solutions is obtained under typical separability and Lasry-Lions type monotonicity conditions.

Keywords: Mean field games, PDE in infinite dimensional spaces, Hamilton-Jacobi-Bellman equation, Fokker-Planck equation, Wasserstein space, mild solution, stochastic differential equation in infinite dimension.

MSC (2010): 35Q89; 35R15; 49N80; 35Q84; 49L12; 60H15; 91A16; 93E20.

1. INTRODUCTION

In this paper, we are concerned with a class of Mean Field Games (MFG) problems in a real separable Hilbert space $(H, \langle \cdot, \cdot \rangle)$. The analytical framework is the following. Let $\mathcal{P}_1(H)$ denote the space of Borel probability measures on H having finite first moment and let $T > 0$. We consider the following system of coupled Hamilton-Jacobi-Bellman (HJB) and nonlinear Fokker-Planck (FP) equations for functions $v : [0, T] \times H \rightarrow \mathbb{R}$ and $m : [0, T] \rightarrow \mathcal{P}_1(H)$,

$$\text{(HJB)} \quad -\partial_t v(t, x) - Lv(t, x) + \mathcal{H}(x, Dv(t, x), m(t)) = 0, \quad v(T, \cdot) = G(\cdot, m(T)),$$

$$\text{(FP)} \quad \partial_t m(t) - L^* m(t) - \operatorname{div}(\mathcal{H}_p(x, Dv(t, x), m(t)) m(t)) = 0, \quad m(0) = m_0,$$

where

- (i) $m_0 \in \mathcal{P}_1(H)$, $G : H \times \mathcal{P}_1(H) \rightarrow \mathbb{R}$;
- (ii) L is the operator formally defined on functions $\phi : H \rightarrow \mathbb{R}$ by

$$L\phi(x) = \langle Ax, D\phi(x) \rangle + \frac{1}{2} \operatorname{Tr}[D^2\phi(x)],$$

where $A : D(A) \subseteq H \rightarrow H$ is a closed and densely defined linear operator;

- (iii) L^* is, formally, the adjoint of L and the operator \mathbf{div} is, formally, the opposite of the adjoint of the gradient: both are used here only for notational convenience as the rigorous definition of a solution to **(FP)** will not involve them, see Section 3 below;
- (iv) $\mathcal{H} : H \times H \times \mathcal{P}_1(H) \rightarrow \mathbb{R}$ and \mathcal{H}_p denotes the Fréchet derivative of \mathcal{H} with respect to the second argument.

We point out that we slightly abuse terminology here, since the first equation is not necessarily a Hamilton-Jacobi-Bellman equation as the function \mathcal{H} can be very general. However, since our primary motivation comes from this case, we will call it an HJB equation.

The theory of MFG originated from the pioneering works of Lasry and Lions, and of Huang, Caines, and Malhamé (see [29, 30, 31, 32]). This theory was primarily developed to provide an analytical framework for (stochastic) differential games involving a large number of symmetric players. In the limit, as the number of players approaches infinity and assuming that the actions of individual players do not directly influence the overall population dynamics, the concept of Nash equilibrium leads to the study of a coupled system of PDEs of the type given by equations **(HJB)** and **(FP)**. Since the inception of this theory, substantial progress has been made in finite-dimensional settings, both theoretically and practically. Notable references for this research area include the book [10] and the survey paper [8]. However, contributions to the theory of Mean Field Games (MFG) in infinite-dimensional spaces remain surprisingly limited. To our knowledge, the literature consists of only a few essential papers, such as [23], which addresses systemic risk by taking the limit of the N -player model in [11], and recent works [21, 33], both of which focus on the linear-quadratic case. This scarcity is somewhat in contrast with the established theory of stochastic optimal control in infinite-dimensional spaces and its connection with Hamilton-Jacobi-Bellman (HJB) equations, which has been well developed over several decades, beginning with foundational works by Barbu and Da Prato (see, e.g., [2]). For a general overview of this subject and its applications, we refer to the recent monograph [19], and see also [17, 35].

In this paper, we aim to bridge this gap by providing a preliminary contribution to the study of MFG in infinite dimensions beyond the linear-quadratic case. Building upon existing results for both HJB and Fokker-Planck equations in infinite-dimensional settings, we investigate the nonlinear coupled system of MFG equations **(HJB)**–**(FP)**. Our main contribution is the proof of well-posedness of this system under certain conditions. The HJB equation **(HJB)** is interpreted in the mild form, allowing us to use well-known theories of mild solutions to partial differential equations in Hilbert spaces. However, this approach requires stronger assumptions on the operator A to ensure certain smoothing properties of the associated Ornstein-Uhlenbeck transition semigroup.

The Fokker-Planck equation **(FP)** is interpreted in a weak form using a special class of test functions. Existence of a solution to the system **(HJB)**–**(FP)** is established with a typical MFG fixed-point argument via Tikhonov's fixed-point theorem, which requires the existence of a unique weak solution to the linear **(FP)** when the term $\mathcal{H}_p(x, Dv(t, x), m(t))$ is fixed. Fokker-Planck-type equations in Hilbert spaces have been widely studied, with various results available on existence and uniqueness (see [3, 4, 5, 6, 15, 37, 39]). To show that our linear **(FP)** has a weak solution, we follow a common approach of linking such an equation to a stochastic differential equation (SDE) in H and noting that the laws of solutions to this SDE are weak solutions of the linear **(FP)**. We apply results from [14] on SDEs in Hilbert spaces with bounded measurable drift, adapting them to our setting. Uniqueness of weak solutions then follows easily from a result in [5], chosen here for its clarity and accessibility. However, this approach imposes additional restrictions on the operator A , ultimately leading to fairly strict assumptions on A .

Uniqueness of solutions to the MFG system **(HJB)**–**(FP)** is obtained under typical monotonicity and separability conditions on \mathcal{H} (see [8] and [32]). To carry out the uniqueness argument, which is now standard in the finite-dimensional case, we approximate the mild solution of the Kolmogorov equation obtained by subtracting two different solutions to the **(HJB)** equations in the **(HJB)**–**(FP)** system by constructing smooth solutions of approximating Kolmogorov equations that belong to the class of test functions used in the definition of a weak solution for the Fokker-Planck equation. Since this is the first study on the subject, we chose not to address the most general case to avoid obscuring the main ideas with excessive technical detail. A possible

generalization could involve a more flexible operator L , in which the term $\text{Tr}[D^2\phi(x)]$, associated with the cylindrical Wiener process W in the underlying SDE, is replaced by $\text{Tr}[QD^2\phi(x)]$ for some self-adjoint operator Q (corresponding to a more general additive noise in the SDE). This could potentially relax the assumptions on A . Another interesting direction would be to interpret equation (HJB) in the viscosity sense, which could also include the case of first-order MFG. We plan to investigate this in future work.

Regarding possible applications, it would be interesting to try to relax the assumptions on the operator A to cover potentially interesting cases. In particular, it would be interesting to treat the case of first order operators A , some of which appear in financial and economic applications, such as models of systemic risk with delays (as in [23]) or models of optimal investment with vintage capital (see [20, 22] in the case of a single firm with vintage structure; [7] in the case of a continuum of homogenous firms but without vintage structure).

The plan of the paper is the following. In Section 2, we introduce the basic notation and assumptions, presenting preliminary technical results on the well-posedness of the SDE related to the (linear) Fokker-Planck equation, along with some estimates for its solutions. Section 3 is dedicated to the existence and uniqueness of weak solutions for the Fokker-Planck equation (3.1). In Section 4, we define the notion of a mild solution to (HJB) and collect some foundational results on mild solutions. In Section 5, we show the existence of solutions to the MFG system (HJB)–(FP), while the uniqueness of solutions is established in Section 6. An example that satisfies the manuscript’s assumptions is discussed in Section 7. Finally, in Appendix A, we present a result on the compactness of sets in spaces of probability measures on a Hilbert space, and in Appendix B, we discuss essential concepts and results related to Kolmogorov equations in Hilbert spaces that are needed in this paper.

2. PRELIMINARIES AND ASSUMPTIONS

We recall that H is a real separable Hilbert space with inner product $\langle \cdot, \cdot \rangle$. We will write $|\cdot|$ for the norm in H ; if needed we stress H in the notation writing $|\cdot|_H$. We always identify H with its dual space. For $R > 0$, we denote $B_R := \{x \in H : |x| \leq R\}$. Throughout the paper, \mathbb{N} will denote the set of natural numbers without 0.

2.1. Basic spaces and notations. We introduce the notation for various functional and operator spaces used in the paper. Below, by I we denote an interval in \mathbb{R} . Given a function $u : I \times H \rightarrow \mathbb{R}$, $(t, x) \mapsto u(t, x)$, we will write $\partial_t u$ for the partial derivative of u with respect to t and Du, D^2u , respectively, for the first and second order Fréchet derivatives of u with respect to the x -variable. For a bounded function between normed spaces, we denote its sup-norm in the usual way by $\|\cdot\|_\infty$.

Below is a list of basic spaces used in the paper.

- (i) $\mathcal{L}(H)$ is the space of bounded linear operators from H to H .
- (ii) $\mathcal{L}^+(H)$ is the space of bounded linear self-adjoint nonnegative operators from H to H .
- (iii) $\mathcal{L}_1(H)$ is the space of bounded linear trace-class operators from H to H .
- (iv) $\mathcal{L}_1^+(H) = \mathcal{L}^+(H) \cap \mathcal{L}_1(H)$.
- (v) $C_b(H)$ is the space of continuous and bounded functions $u : H \rightarrow \mathbb{R}$.
- (vi) $C_b(I \times H)$ is the space of continuous and bounded functions $u : I \times H \rightarrow \mathbb{R}$.
- (vii) $C_b(I \times H; H)$ is the space of continuous and bounded functions $u : I \times H \rightarrow H$.
- (viii) $C_b(I \times H; \mathcal{L}(H))$ (respectively, $C_b(I \times H; \mathcal{L}_1(H))$) is the space of continuous and bounded functions $u : I \times H \rightarrow \mathcal{L}(H)$ (respectively, $u : I \times H \rightarrow \mathcal{L}_1(H)$).

- (ix) $C_b^{0,1}(I \times H)$ is the space of continuous and bounded functions $u : I \times H \rightarrow \mathbb{R}$, which are Fréchet differentiable with respect to the variable x and such that $Du \in C_b(I \times H; H)$.
- (x) $UC_b(H)$ (respectively, $UC_b(H; H)$, $UC_b(H; \mathcal{L}_1(H))$) is the space of bounded and uniformly continuous functions $u : H \rightarrow \mathbb{R}$ (respectively, $u : H \rightarrow H$, $u : H \rightarrow \mathcal{L}_1(H)$).
- (xi) $UC_b^2(H) = \{u : H \rightarrow \mathbb{R} : u \in UC_b(H), Du \in UC_b(H; H), D^2u \in UC_b(H; \mathcal{L}(H))\}$.
- (xii) Finally,

$$C_b^{1,2}(I \times H) = \{u \in C_b(I \times H) : \partial_t u \in C_b(I \times H), Du \in C_b(I \times H; H), D^2u \in C_b(I \times H; \mathcal{L}(H))\}.$$

We denote by $\mathcal{M}(H)$ the topological vector space of finite signed measures on H endowed with the weak topology, i.e., the locally convex topology on $\mathcal{M}(H)$ induced by the family of seminorms

$$(2.1) \quad |\mu|_f := \left| \int_H f(x) \mu(dx) \right|, \quad f \in C_b(H).$$

By $\mathcal{P}(H)$ we denote the space of Borel probability measures on H . The weak topology defined on it is the one inherited by the weak topology of $\mathcal{M}(H)$ defined above. We denote by $\mathcal{P}_1(H)$ the subset of $\mathcal{P}(H)$ of the probability measures having finite first moment. In $\mathcal{P}_1(H)$ we have the Monge-Kantorovich distance (see, e.g., [1, 38])

$$(2.2) \quad \mathbf{d}_1(\mu, \nu) = \inf_{\gamma \in \Gamma(\mu, \nu)} \int_{H \times H} |x - y| \gamma(dx, dy),$$

where $\Gamma(\mu, \nu)$ is the set of all Borel probability measures on $H \times H$ with first and second marginals μ and ν , respectively. Unless differently specified, the space $\mathcal{P}_1(H)$ will be considered endowed with the topology induced by \mathbf{d}_1 which makes it a complete metric space. We notice that, if $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space and $X, Y \in L^1(\Omega; H)$ are such that $\mathcal{L}(X) = \mu$, $\mathcal{L}(Y) = \nu$, then $\gamma = (X \times Y)_\# \mathbb{P} \in \Gamma(\mu, \nu)$, so that

$$\mathbf{d}_1(\mu, \nu) \leq \int_{H \times H} |x - y| \gamma(dx, dy) = \int_{\Omega} |X(\omega) - Y(\omega)| d\mathbb{P}(\omega) = \mathbb{E}[|X - Y|].$$

We will deal with the space

$$(2.3) \quad \mathcal{S} = C([0, T]; (\mathcal{P}_1(H), \mathbf{d}_1)).$$

It is a complete metric space endowed with the metric

$$\rho_\infty(\mu, \nu) = \sup_{t \in [0, T]} \mathbf{d}_1(\mu(t), \nu(t)).$$

We will also use the symbol ρ_∞ to denote the induced metric topology.

2.2. Weighted spaces. Given any $\gamma > 0$, we define

$$(2.4) \quad C_{b,\gamma}([0, T] \times H) := \{f : [0, T] \times H \rightarrow \mathbb{R} \text{ s.t. } f \in C_b([0, \tau] \times H) \forall \tau \in (0, T) \\ \text{and } (t, x) \mapsto (T - t)^\gamma f(t, x) \text{ belongs to } C_b([0, T] \times H)\}.$$

Similarly, we define

$$(2.5) \quad C_{b,\gamma}([0, T] \times H; H) := \{w : [0, T] \times H \rightarrow H \text{ s.t. } w \in C_b([0, \tau] \times H; H) \forall \tau \in (0, T) \\ \text{and } (t, x) \mapsto (T - t)^\gamma w(t, x) \text{ belongs to } C_b([0, T] \times H; H)\}.$$

The spaces above are Banach spaces when endowed with the norm (see e.g. [24, 27] and [12])

$$\|f\|_{C_{b,\gamma}([0, T] \times H)} := \sup_{(t, x) \in [0, T] \times H} (T - t)^\gamma |f(t, x)|,$$

$$\|f\|_{C_{b,\gamma}([0,T]\times H)} := \sup_{(t,x)\in[0,T]\times H} (T-t)^\gamma |f(t,x)|_H.$$

We will often simply write $\|f\|_{C_{b,\gamma}}$. Moreover, we define

$$(2.6) \quad C_{b,\gamma}^{0,1}([0,T]\times H) := \left\{ f \in C_b([0,T]\times H) : f \in C_b^{0,1}([0,\tau]\times H) \forall \tau \in (0,T) \right. \\ \left. \text{and } (t,x) \mapsto (T-t)^\gamma Df(t,x) \text{ belongs to } C_b([0,T]\times H; H) \right\}.$$

The space above is a Banach space when endowed with the norm (see e.g. [24, 27] and [12])

$$\|f\|_{C_{b,\gamma}^{0,1}([0,T]\times H)} := \|f\|_\infty + \sup_{(t,x)\in[0,T]\times H} (T-t)^\gamma |Df(t,x)|_H.$$

We will sometimes simply write $\|f\|_{C_{b,\gamma}^{0,1}}$.

2.3. The operator A and the generated semigroup. Concerning the operator A , we assume the following.

Assumption 2.1. $A : D(A) \subseteq H \rightarrow H$ is closed, densely defined, negative and self-adjoint. Moreover there exists $\delta > 0$ such that $(-A)^{-1+\delta} \in \mathcal{L}_1^+(H)$.

Given Assumption 2.1, A^{-1} is compact and self-adjoint. Therefore, there exist an orthonormal basis $\{\mathbf{e}_k\}_{k \in \mathbb{N}}$ and a decreasing sequence $(\lambda_k)_{k \in \mathbb{N}} \subset (-\infty, 0)$ such that $\lambda_n \rightarrow -\infty$, with respect to which A is diagonal:

$$A\mathbf{e}_k = \lambda_k \mathbf{e}_k, \quad k \in \mathbb{N},$$

and

$$(2.7) \quad \sum_{k \in \mathbb{N}} |\lambda_k|^{-1+\delta} < \infty.$$

Moreover, A generates an analytic C_0 -semigroup of contractions $\{e^{tA}\}_{t \geq 0} \subseteq \mathcal{L}(H)$ and

$$(2.8) \quad e^{tA} \mathbf{e}_k = e^{\lambda_k t} \mathbf{e}_k, \quad k \in \mathbb{N}.$$

Remark 2.2.

(i) An example of A that satisfies the above Assumption 2.1 is the Laplace operator ∂_{xx} in $H = L^2([0, 1])$ with zero Dirichlet boundary condition.

(ii) It can be verified from (2.7) that Assumption 2.1 implies typical conditions on the operator A ensuring existence and uniqueness of mild solutions of equations in infinite dimensional spaces (see [19, Ch. 4]), that is:

(a) One has

$$Q_t := \int_0^t e^{sA} e^{sA^*} ds \in \mathcal{L}_1^+(H) \quad \forall t > 0$$

and there exists $\alpha > 0$ such that

$$Q_t^{(\alpha)} := \int_0^t s^{-\alpha} e^{sA} e^{sA^*} ds \in \mathcal{L}_1^+(H) \quad \forall t > 0.$$

(b) $e^{tA}(H) \subset Q_t^{1/2}(H)$ for every $t > 0$ and there exist $C_0 > 0$ and $\gamma \in (0, 1)$ such that

$$|Q_t^{-1/2} e^{tA}| \leq C_0 t^{-\gamma}, \quad \forall t \geq 0,$$

where $Q_t^{-1/2}$ is the pseudoinverse of $Q_t^{1/2}$ (see [19, Def. B.1]).

2.4. The probabilistic framework and the reference SDE. We now define a probabilistic framework. We consider a complete filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$ supporting a sequence $\{\beta_k(t)\}_{k \in \mathbb{N}}$ of independent one dimensional Brownian motions. Given the orthonormal basis $\{e_k\}_{k \in \mathbb{N}}$ of Assumption 2.1, we consider the cylindrical Wiener process $\{W_t\}_{t \geq 0}$, formally written as

$$W_t = \sum_{k \in \mathbb{N}} \beta_k(t) e_k.$$

Under Assumption 2.1, the stochastic convolution (see Remark 2.2(ii) and [18, Ch. 5])

$$(2.9) \quad W_t^A := \int_0^t e^{(t-s)A} dW_s, \quad t \geq 0,$$

is a well-defined centered Gaussian process in the Hilbert space H with covariance operator

$$Q_t := \int_0^t e^{2sA} ds \in \mathcal{L}_1^+(H)$$

admitting a continuous version. The process

$$X_t^{0,x} := e^{tA}x + W_t^A, \quad t \geq 0,$$

can be seen as a generalized (called *mild*) solution to the SDE

$$dX_t = AX_t dt + dW_t, \quad X_0 = x \in H.$$

$X_t^{0,x}$ is called the Ornstein-Uhlenbeck process and is associated to the Markov transition semigroup

$$(2.10) \quad [R_t \varphi](x) = \mathbb{E} \left[\varphi(X_t^{0,x}) \right] = \int_H \varphi(y) \mathcal{N}(e^{tA}x, Q_t)(dy), \quad \varphi \in C_b(H),$$

called the Ornstein-Uhlenbeck semigroup. We will use it later to define the concept of a mild solution to (HJB).

To construct a weak solution to a linear Fokker-Planck equation we will consider the SDE

$$(2.11) \quad dX_t = AX_t dt + w(t, X_t) dt + dW_t, \quad \mathcal{L}(X_0) = m_0.$$

We will need the following assumption.

Assumption 2.3. (i) $\int_H |x|^4 m_0(dx) < \infty$.

(ii) $w \in C_b((0, T) \times H; H)$.

Theorem 2.4. *Let $m_0 \in \mathcal{P}_1(H)$ and let Assumptions 2.1 and 2.3(ii) hold. There exists a unique weak-mild solution to (2.11) in the following sense.*

(i) *There exist a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbf{P})$, an \mathcal{F}_0 -measurable random variable X_0 with $\mathcal{L}(X_0) = m_0$, a cylindrical Wiener process $(W_t)_{t \in [0, T]}$ with respect to $(\mathcal{F}_t)_{t \in [0, T]}$, and an adapted process $(X_t^w)_{t \in [0, T]}$ such that*

$$X_t^w = e^{tA}X_0 + \int_0^t e^{(t-s)A} w(s, X_s^w) ds + W_t^A, \quad \forall t \geq 0;$$

(ii) *If another weak-mild solution \widehat{X}^w exists, then $\mathcal{L}(X_t^w) = \mathcal{L}(\widehat{X}_t^w)$ for every $t \in [0, T]$.*

Proof. When w does not depend on t , the result is proved [14, Th. 13] and the proof works in the same way if w depends on t . We just need to explain how our setup fits into the framework of [14].

Given our filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$ with the Wiener process W , let $(\Omega_1, \mathcal{F}_1, \mathbf{P}_1)$ be another complete atomless probability space. For any $m_0 \in \mathcal{P}_1(H)$, since $(\Omega_1, \mathcal{F}_1, \mathbf{P}_1)$ is

nonatomic, there exists an \mathcal{F}_1 -measurable random variable $\xi_0 : \Omega_1 \rightarrow X$ such that $m_0 = \mathcal{L}(\xi_0)$ (see, e.g., [9, Lemma 5.29]). We define a new probability space $(\Omega_2, \mathcal{F}_2, \mathbb{P}_2)$ as follows:

$$\Omega_2 = \Omega \times \Omega_1, \quad \mathbb{P}_2 = \mathbb{P} \otimes \mathbb{P}_1 \text{ on } \mathcal{F} \otimes \mathcal{F}_1,$$

$$\mathcal{F}_2 = \text{completion of } \mathcal{F} \otimes \mathcal{F}_1 \text{ with respect to } \mathbb{P}_2 = \mathbb{P} \otimes \mathbb{P}_1.$$

We then define (\mathcal{F}_t^2) to be the augmentation of $\mathcal{F}_t \otimes \mathcal{F}_1$. The cylindrical Wiener process $\{W_t\}_{t \in [0, T]}$ and the random variable ξ_0 extend naturally to the space $(\Omega_2, \mathcal{F}_2, (\mathcal{F}_t^2)_{t \in [0, T]}, \mathbb{P}_2)$. We denote them by W^2 and X_0 respectively. The random variable $X_0 : \Omega \rightarrow H$ is \mathcal{F}_0^2 -measurable, $m_0 = \mathcal{L}(X_0)$, and $\{W_t^2\}_{t \in [0, T]}$ is a cylindrical Wiener process in the filtered probability space $(\Omega_2, \mathcal{F}_2, (\mathcal{F}_t^2)_{t \in [0, T]}, \mathbb{P}_2)$.

Now, following the arguments of [14], the weak-mild solution is constructed starting from the space $(\Omega_2, \mathcal{F}_2, (\mathcal{F}_t^2)_{t \in [0, T]}, \mathbb{P}_2)$ using the Girsanov Theorem by introducing a new cylindrical Wiener process $\{\tilde{W}_t^2\}_{t \in [0, T]}$ and replacing \mathbb{P}_2 by $\tilde{\mathbb{P}}_2$, where

$$d\tilde{\mathbb{P}}_2 = M_T d\mathbb{P}_2,$$

and $(M_s)_{s \in [0, T]}$ is an appropriate martingale such that $M_0 \equiv 1$. If $f \in C_b(H)$, we then have

$$\tilde{\mathbb{E}}[f(X_0)] = \mathbb{E}[M_T f(X_0)] = \mathbb{E}[\mathbb{E}[M_T f(X_0) | \mathcal{F}_0^2]] = \mathbb{E}[f(X_0) \mathbb{E}[M_T | \mathcal{F}_0^2]] = \mathbb{E}[f(X_0)].$$

Thus, the law of X_0 in the new probability space is the same. This puts us in the framework of [14] and we can now proceed as in the proof of [14, Th. 13]. \square

We provide some estimates for the laws $\mathcal{L}(X_t^w)$ of the solution to (2.11) that will be used afterwards.

Proposition 2.5. *Let Assumptions 2.1 and 2.3 hold and let X_t^w be the unique weak-mild solution to (2.11).*

(i) *There exists c_0 independent of $\|w\|_\infty$ such that*

$$\int_H |x|^4 \mathcal{L}(X_t^w)(dx) \leq c_0 \left(1 + \int_H |x|^4 m_0(dx) + \|w\|_\infty^4 \right), \quad \forall t \in [0, T].$$

(ii) *For each $R > 0$, there exists a modulus of continuity $\omega_{m_0, R}$ such that, if $|w|_\infty \leq R$, it holds*

$$\mathbf{d}_1(\mathcal{L}(X_t^w), \mathcal{L}(X_s^w)) \leq \omega_{m_0, R}(|t - s|), \quad \forall s, t \in [0, T].$$

Proof. (i) Using the definition of mild solution, the fact that $|e^{tA}| \leq 1$ and the estimate on the stochastic convolution W_t^A [19, Prop. 1.144], we have, for some $c_1 > 0$,

$$\begin{aligned} \mathbb{E}[|X_t^w|^4] &\leq \mathbb{E} \left[\left| e^{tA} X_0 + \int_0^t e^{(t-s)A} w(s, X_s^w) ds + W_t^A \right|^4 \right] \\ &\leq c_1 (\mathbb{E}[|X_0|^4] + T^4 \|w\|_\infty^4 + 1), \quad \forall 0 \leq t \leq T, \end{aligned}$$

which implies (i).

(ii) Let $0 \leq s \leq t \leq T$. Using the definition of mild solution, we have the existence of some $c_1 > 0$, independent of s, t such that

$$\begin{aligned} \mathbf{d}_1(\mathcal{L}(X_t^w), \mathcal{L}(X_s^w)) &\leq \mathbb{E}[|X_t^w - X_s^w|] \leq c_1 \left(\mathbb{E} \left[\left| (e^{(t-s)A} - I) X_0 \right| \right] \right. \\ &\quad + \mathbb{E} \left[\left| (e^{(t-s)A} - I) \int_0^s e^{(s-r)A} w(r, X_r^w) dr \right| + \int_s^t \left| e^{(t-r)A} w(r, X_r^w) \right| dr \right] \\ &\quad \left. + \mathbb{E}[|W_t^A - W_s^A|] \right). \end{aligned}$$

We proceed to estimate the terms in the right hand side.

- (a) Recalling that $|e^{\alpha A}| \leq 1$ for $\alpha \geq 0$, we have, for every $0 \leq s \leq t \leq T$,

$$\mathbb{E} \left[\int_s^t \left| e^{(t-r)A} w(r, X_r^w) \right| dr \right] \leq (t-s) \|w\|_\infty.$$

- (b) Considering that $(e^{tA})_{t \in [0, T]}$ is strongly continuous and using dominated convergence, we have

$$\mathbb{E} \left[\left| (e^{(t-s)A} - I) X_0 \right| \right] \rightarrow 0, \quad \text{as } s \rightarrow t.$$

Hence, there exists a modulus of continuity $\omega_{m_0, A}$, independent of w such that, for every $0 \leq s \leq t \leq T$,

$$\mathbb{E} \left[\left| (e^{(t-s)A} - I) X_0 \right| \right] \leq \omega_{m_0, A}(|t-s|).$$

- (c) Let $R > 0$. Since A generates a semigroup of contractions, for every $0 < \eta < s \leq T$ and w such that $\|w\|_\infty \leq R$, we have (P-a.s.)

$$\int_0^{s-\eta} e^{(s-r)A} w(r, X_r^w) dr \in e^{\eta A}(B_{TR}) =: K_{\eta, R}.$$

Assumption 2.1 guarantees that the semigroup is compact so $K_{\eta, R}$ is compact. Considering also that the semigroup is strongly continuous, we have for $0 \leq s \leq t \leq T$

$$(2.12) \quad \sup_{x \in K_{\eta, R}} |(e^{(t-s)A} - I)x| \leq \omega_{\eta, A, R}(t-s),$$

for some modulus of continuity $\omega_{\eta, A, R}$. Using (2.12), it follows that, for every $0 \leq \eta \leq s \leq t \leq T$

$$\begin{aligned} & \mathbb{E} \left[\left| (e^{(t-s)A} - I) \int_0^s e^{(s-r)A} w(r, X_r^w) dr \right| \right] \\ & \leq \mathbb{E} \left[\left| (e^{(t-s)A} - I) \int_0^{s-\eta} e^{(s-r)A} w(r, X_r^w) dr \right| + \left| (e^{(t-s)A} - I) \int_{s-\eta}^s e^{(s-r)A} w(r, X_r^w) dr \right| \right] \\ & \leq \omega_{\eta, A, R}(t-s) + 2\eta R. \end{aligned}$$

Now, define

$$\tilde{\omega}_{A, R}(r) := \inf_{0 < \eta \leq T} \left\{ \omega_{\eta, A, R}(r) + 2\eta R \right\}, \quad r > 0.$$

Then, $\tilde{\omega}_{A, R}(0^+) = 0$ and $\tilde{\omega}_{A, R}$ is concave; so, it is a modulus of continuity. Taking now the infimum over $0 < \eta \leq T$ in the inequality above, we get, for every $0 \leq s \leq t \leq T$,

$$\mathbb{E} \left[\left| (e^{(t-s)A} - I) \int_0^s e^{(s-r)A} w(r, X_r^w) dr \right| \right] \leq \tilde{\omega}_{A, R}(|t-s|).$$

- (d) The map $[0, T] \rightarrow L^2(\Omega, \mathbb{P})$, $t \mapsto W_t^A$ is continuous (see [18, Th. 5.2]); hence, also the map $[0, T] \rightarrow L^1(\Omega, \mathbb{P})$, $t \mapsto W_t^A$ is continuous. Hence, since $[0, T]$ is compact, there exists a modulus of continuity $\hat{\omega}_A$ such that, for every $0 \leq s \leq t \leq T$,

$$\mathbb{E} [|W_t^A - W_s^A|] \leq \hat{\omega}_A(|t-s|).$$

We now combine all the estimates of points (a)–(d) above to obtain Claim (ii). \square

2.5. **Assumptions on \mathcal{H} and G .** We make the following assumptions about \mathcal{H} and G .

Assumption 2.6.

(i) The function \mathcal{H} is continuous, the function $\mathcal{H}(\cdot, 0, \mu_0)$ is bounded for some $\mu_0 \in \mathcal{P}_1(H)$, and there exists $C > 0$ such that

$$|\mathcal{H}(x, p, \mu) - \mathcal{H}(x, p', \mu')| \leq C(|p - p'| + \mathbf{d}_1(\mu, \mu')), \quad \forall x \in H, p, p' \in H, \mu, \mu' \in \mathcal{P}_1(H).$$

(ii) \mathcal{H} is differentiable with respect to the second variable, \mathcal{H}_p is continuous and bounded, and there exists $C > 0$ such that

$$|\mathcal{H}_p(x, p, \mu) - \mathcal{H}_p(x, p', \mu')| \leq C(|p - p'| + \mathbf{d}_1(\mu, \mu')), \quad \forall x \in H, p, p' \in H, \mu, \mu' \in \mathcal{P}_1(H).$$

(iii) G is continuous and such that the function $\mu \mapsto G(\cdot, \mu)$ is continuous as a map from $\mathcal{P}_1(H)$ to $C_b(H)$.

Remark 2.7. Requiring Lipschitz continuity of \mathcal{H} and \mathcal{H}_p with respect to p and μ variables is a little restrictive. However, Assumption 2.6 still covers many basic cases, especially those having separated structure. An example of \mathcal{H} coming from a stochastic optimal control problem is provided in Section 7.

3. THE FOKKER-PLANCK EQUATION: WEAK SOLUTIONS

In this section, we study (FP) when the term $\mathcal{H}_p(x, Dv(t, x), m(t))$ is frozen. For that, we fix $w \in C_b([0, T] \times H; H)$ and consider the linear Fokker-Planck equation in the space $\mathcal{P}_1(H)$

$$(3.1) \quad \begin{cases} \partial_t m(t) - L^* m(t) - \operatorname{div}(w(t, x)m(t)) = 0, \\ m(0) = m_0 \in \mathcal{P}_1(H). \end{cases}$$

We will interpret (3.1) in a weak sense. For that, we define the set of test functions

$$(3.2) \quad \mathcal{D}_T = \left\{ \varphi \in C_b^{1,2}([0, T] \times H) : A^* D\varphi(t, x) \in C_b([0, T] \times H; H), D^2\varphi \in C_b([0, T] \times H; \mathcal{L}_1(H)) \right\}.$$

To provide a rigorous definition of solution to (3.1), we introduce the following operator¹

$$(3.3) \quad (L_0\phi)(x) = \langle x, A^* D\phi(x) \rangle + \frac{1}{2} \operatorname{Tr} [D^2\phi(x)],$$

defined on

$$(3.4) \quad D(L_0) = \left\{ \phi \in UC_b^2(H) : D^2\phi \in UC_b(H; \mathcal{L}_1(H)), A^* D\phi \in UC_b(H; H) \right\}.$$

The formal operators L^* and div in (3.1) are interpreted similarly as for a Fokker-Planck equation in the case $H = \mathbb{R}^n$ when using the class of test functions \mathcal{D}_T . The definition of *weak solution* to (3.1) given below is the same as the one of [5]; however, we use a larger class of test functions.

Definition 3.1. A map $m : [0, T] \rightarrow \mathcal{P}_1(H)$ such that $t \mapsto m(t)(I)$ is measurable for each $I \in \mathcal{B}(H)$ ² is called a *weak solution* to (3.1) if $m(0) = m_0$ and

$$(3.5) \quad \begin{aligned} & \int_H \varphi(t, x)m(t, dx) - \int_H \varphi(0, x)m(0, dx) \\ &= \int_0^t \left(\int_H [\partial_t \varphi(s, x) + L_0\varphi(s, x) - \langle w(s, x), D\varphi(s, x) \rangle] m(s, dx) \right) ds \end{aligned}$$

for every $t \in [0, T]$ and $\varphi \in \mathcal{D}_T$.

¹Note that, as explained in [19, Section B.5.2], the operator L_0 is closable in a suitable weak sense (so called \mathcal{K} -closable) and its closure (called \mathcal{K} -closure there) is the generator (again in a suitable weak sense) of the semigroup R_t in (2.10).

²That is, in the terminology of [5], m is a probability kernel.

The connection between (3.1) and (2.11) is provided by the following.

Proposition 3.2. *Let Assumptions 2.1 and 2.3(ii) hold and let X^w be the unique weak-mild solution to (2.11) provided by Theorem 2.4. Then, $\mathcal{L}(X^w)$ is a weak solution to (2.11).*

Proof. This is just an application of Dynkin's formula for regular functions: see, e.g, in [19, Prop. 1.164], by interpreting $(w(s, X_s^w))_{s \in [0, T]}$ as the process $(f_s)_{s \in [0, T]}$ there. \square

Concerning uniqueness of solutions to (3.1), we have the following.

Proposition 3.3. *Let Assumptions 2.1 and 2.3 hold. Then (3.1) admits at most one weak solution in the class*

$$\mathcal{S}_4 := \left\{ m \in \mathcal{S} : \int_0^T \int_H |x|^4 m(t, dx) dt < \infty \right\}.$$

Proof. The claim follows from [5, Th. 4.1] once we prove that a solution in our sense is also a solution in the sense of [5, Th. 4.1]. To do that we first notice that, in [5, Formula (1.5)], the definition of solution is given using the set test functions (called $\mathcal{E}_A(H)$ there) which is the linear span of the real parts of the functions of the form

$$(t, x) \mapsto \phi(t) e^{i\langle x, h(t) \rangle}, \quad \phi \in C^1([0, T]), \quad h \in C^1([0, T], D(A^*)).$$

Given the above, we observe that such a set is clearly contained in the set \mathcal{D}_T defined in (3.2). Finally, we point out that in [5] the Kolmogorov operator L is defined differently (corresponding to our $\partial_t + L_0 - \langle w, D \rangle$); still, equality (1.5) in [5] corresponds to our (3.5). \square

We combine and summarize the above two results in the following theorem.

Theorem 3.4. *Let Assumptions 2.1 and 2.3 hold. Then (3.1) admits a unique weak solution in the class \mathcal{S}_4 , which coincides with the law of the unique weak-mild solution X^w to (2.11).*

Proof. By Proposition 2.5(ii), our solution provided by Proposition 3.2 belongs to the class \mathcal{S}_4 and the uniqueness follows from Proposition 3.3. \square

4. THE HJB EQUATION: MILD SOLUTIONS

In this section we deal with equation (HJB) for a given fixed $m \in \mathcal{S}$. We introduce the concept of a *mild solution* to (HJB) (see [19, Section 4.4.1.2]).

Definition 4.1 (Mild solution to (HJB)). *Recall (2.4). A mild solution to (HJB) is a function $v \in C_{b, \gamma}^{0,1}([0, T] \times H)$ such that, for all $(t, x) \in [0, T] \times H$,*

$$v(t, x) = [R_{T-t} G(\cdot, m(T))](x) - \int_t^T [R_{s-t} \mathcal{H}(\cdot, Dv(s, \cdot), m(s))](x) ds.$$

Theorem 4.2. *Let $m \in \mathcal{S}$ and let Assumptions 2.1 and 2.6 hold.*

- (i) *There exists a unique mild solution u to (HJB).*
- (ii) *If $m_n \rightarrow m$ in \mathcal{S} , denoting by v_n and v the corresponding mild solutions to (HJB), we have*

$$\|Dv_n - Dv\|_{C_{b, \gamma}} \rightarrow 0.$$

Proof. (i) This part is contained in [19, Theorem 4.90].³

³More precisely, it is enough to take in that theorem, $m = 0$, $\gamma_G(t) = t^{-\gamma}$, $G = \mathbf{Id}$. We warn that G in this footnote refers to the operator used in [19, Theorem 4.90], which has nothing to do with our function G .

- (ii) Using the smoothing properties of the Ornstein-Uhlenbeck semigroup (see e.g. [19, Theorem 4.37]), we have the representation of the gradients

$$Dv(t, x) = D[R_{T-t}G(\cdot, m(T))](x) - \int_t^T D[R_{s-t}\mathcal{H}(\cdot, Dv(s, \cdot), m(s))](x)ds.$$

$$Dv_n(t, x) = D[R_{T-t}G(\cdot, m_n(T))](x) - \int_t^T D[R_{s-t}\mathcal{H}(\cdot, Dv_n(s, \cdot), m_n(s))](x)ds.$$

Hence, fixing $t \in [0, T)$ and considering [19, Th. 4.37] and Assumption 2.6, we have, for some $C > 0$,

$$\begin{aligned} \|Dv(t, \cdot) - Dv_n(t, \cdot)\|_\infty &\leq \|D[R_{T-t}(G(\cdot, m(T)) - G(\cdot, m_n(T)))]\|_\infty \\ &\quad + \int_t^T \left\| D \left[R_{s-t}(\mathcal{H}(\cdot, Dv(s, \cdot), m(s)) - \mathcal{H}(\cdot, Dv_n(s, \cdot), m_n(s))) \right] \right\|_\infty ds \\ &\leq C \left((T-t)^{-\gamma} \|G(\cdot, m(T)) - G(\cdot, m_n(T))\|_\infty \right. \\ &\quad \left. + \int_t^T (s-t)^{-\gamma} \left(\|Dv(s, \cdot) - Dv_n(s, \cdot)\|_\infty + \mathbf{d}_1(m(s), m_n(s)) \right) ds \right) \\ &\leq C \left(\eta_n (T-t)^{-\gamma} + \rho_\infty(m, m_n) T^{1-\gamma} + \int_t^T (s-t)^{-\gamma} \|Dv(s, \cdot) - Dv_n(s, \cdot)\|_\infty ds \right), \end{aligned}$$

where, by Assumption 2.6(iii),

$$\eta_n := \|G(\cdot, m(T)) - G(\cdot, m_n(T))\|_\infty \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

By a suitable generalization of Gronwall's Lemma (see [19, Prop. D.30]), we get

$$\|Dv(t, \cdot) - Dv_n(t, \cdot)\|_\infty \leq C(T-t)^{-\gamma} (\eta_n + T\rho_\infty(m, m_n)).$$

Multiplying both sides by $(T-t)^\gamma$ and taking the supremum over $t \in [0, T)$, we get

$$\|Dv - Dv_n\|_{C_{b,\gamma}} \leq C (\eta_n + T\rho_\infty(m, m_n))$$

and the claim follows. \square

5. THE MFG SYSTEM: EXISTENCE OF SOLUTIONS

We turn to the analysis of the coupled MFG system (HJB)-(FP). Let Assumptions 2.3(i) and 2.6 hold. Given $m \in \mathcal{S}$, we denote by $v^{(m)}$ the unique mild solution to (HJB). Moreover, recalling the notation X^w used in Section 3, we set

$$X_t^{(m)} := X_t^{\mathcal{H}_p(\cdot, Dv^{(m)}(\cdot, \cdot), m(\cdot))}.$$

In the remainder of this section, $\{\mathbf{e}_k\}_{k \in \mathbb{N}}$ is the orthonormal basis provided by Assumption 2.1.

Lemma 5.1. *Let Assumptions 2.1 and 2.3 holds. For $n \in \mathbb{N}$, set*

$$\alpha_n := (2|\lambda_n|)^{-1}, \quad \beta_n := \int_H \langle x, \mathbf{e}_n \rangle^2 m_0(dx).$$

(i) *We have*

$$\int_H \sup_{t \in [0, T]} \langle e^{tA} x, \mathbf{e}_n \rangle^2 m_0(dx) \leq \beta_n, \quad \forall n \in \mathbb{N}.$$

(ii) We have

$$\sup_{t \in [0, T], x \in H} \int_0^t \langle e^{(t-s)A} w(s, x), \mathbf{e}_n \rangle^2 ds \leq \alpha_n \cdot \sup_{s \in (0, T), x \in H} \langle w(s, x), \mathbf{e}_n \rangle^2 \leq \alpha_n \|w\|_\infty^2, \quad \forall n \in \mathbb{N}.$$

(iii) We have

$$\sup_{t \in [0, T]} \mathbf{E} \langle W_t^A, \mathbf{e}_n \rangle^2 \leq \alpha_n, \quad \forall n \in \mathbb{N}.$$

Proof. (i) In fact,

$$\begin{aligned} \int_H \sup_{t \in [0, T]} \langle e^{tA} x, \mathbf{e}_i \rangle^2 m_0(dx) &= \int_H \sup_{t \in [0, T]} \langle x, e^{tA} \mathbf{e}_n \rangle^2 m_0(dx) \\ &= \int_H \left(\sup_{t \in [0, T]} e^{2\lambda_n t} \right) \langle x, \mathbf{e}_n \rangle^2 m_0(dx) \leq \int_H \langle x, \mathbf{e}_n \rangle^2 m_0(dx) = \beta_n. \end{aligned}$$

(ii) We compute,

$$\begin{aligned} \sup_{t \in [0, T], x \in H} \int_0^t \langle e^{(t-s)A} w(s, x), \mathbf{e}_n \rangle^2 ds &= \sup_{t \in [0, T], x \in H} \int_0^t \langle w(s, x), e^{(t-s)A} \mathbf{e}_n \rangle^2 ds \\ &= \sup_{t \in [0, T], x \in H} \int_0^t e^{2(t-s)\lambda_n} \langle w(s, x), \mathbf{e}_n \rangle^2 ds \leq \sup_{s \in (0, T), x \in H} \langle w(s, x), \mathbf{e}_n \rangle^2 \cdot \sup_{t \in [0, T]} \int_0^t e^{2(t-s)\lambda_n} ds \\ &\leq \alpha_n \cdot \sup_{s \in (0, T), x \in H} \langle w(s, x), \mathbf{e}_n \rangle^2 \leq \alpha_n \|w\|_\infty^2. \end{aligned}$$

(iii) Recall that, by Assumption 2.1, we have

$$\text{Cov}(W_t^A) = Q_t = \int_0^t e^{2sA} ds.$$

Then,

$$\sup_{t \in [0, T]} \mathbf{E} \langle W_t^A, \mathbf{e}_n \rangle^2 = \sup_{t \in [0, T]} \langle Q_t \mathbf{e}_n, \mathbf{e}_n \rangle \leq \langle Q_T \mathbf{e}_n, \mathbf{e}_n \rangle = (2|\lambda_n|)^{-1}.$$

□

Set⁴

$$a_n := 3 \left(\beta_n + \alpha_n + \alpha_n \|\mathcal{H}_p\|_\infty^2 \right).$$

Then, by Assumptions 2.1 and 2.3(i), we have

$$(5.1) \quad \mathbf{a} = (a_n)_{n \in \mathbb{N}} \in \ell_1.$$

Let us consider the map

$$(5.2) \quad \Psi : \mathcal{S} \rightarrow \mathcal{S}, \quad \Psi(m)(t) := \mathcal{L}(X_t^{(m)}).$$

This map is well defined due to Proposition 2.5.

Proposition 5.2. *Let Assumptions 2.1, 2.3(i) and 2.6 hold. Then, for each $m \in \mathcal{S}$, $n \in \mathbb{N}$, and $t \in [0, T]$, we have*

$$\int_H |\langle x, \mathbf{e}_n \rangle|^2 \Psi(m)(t, dx) \leq a_n.$$

⁴Note that a_n only depends on the data A , \mathcal{H} , and on the initial measure m_0 .

Proof. We have, using the equation for $X_t^{(m)}$ and denoting $w(s, x) = \mathcal{H}_p(x, Dv^{(m)}(s, x), m(s))$,

$$\begin{aligned} \int_H |\langle x, \mathbf{e}_n \rangle|^2 \Psi(m)(t, dx) &= \mathbb{E} \langle X_t^{(m)}, \mathbf{e}_n \rangle^2 \\ &\leq 3\mathbb{E} \left[\langle e^{tA} X_0, \mathbf{e}_n \rangle^2 + \int_0^t \langle e^{(t-s)A} w(s, X_s^{(m)}), \mathbf{e}_n \rangle^2 ds + \langle W_t^A, \mathbf{e}_n \rangle^2 \right] \end{aligned}$$

Applying Lemma 5.1(i), we get the following estimate for the first term

$$\mathbb{E} [\langle e^{tA} X_0, \mathbf{e}_n \rangle^2] = \int_H \langle e^{tA} x, \mathbf{e}_n \rangle^2 m_0(dx) \leq \beta_n.$$

As for the second term, we have by Lemma 5.1(ii)

$$\mathbb{E} \left[\int_0^t \langle e^{(t-s)A} w(s, X_s^{(m)}), \mathbf{e}_n \rangle^2 ds \right] \leq \sup_{x \in H} \int_0^t \langle e^{(t-s)A} w(s, x), \mathbf{e}_n \rangle^2 ds \leq \alpha_n \|\mathcal{H}_p\|_\infty^2.$$

The last term is estimated by Lemma 5.1(iii), which gives

$$\mathbb{E} [\langle W_t^A, \mathbf{e}_n \rangle^2] \leq \alpha_n.$$

The claim now follows from the definition of a_n . \square

Let c_0 be the constant of Proposition 2.5. We set

$$\hat{c}_{m_0} := 1 + c_0 \left(1 + \int_H |x|^4 m_0(dx) + \|\mathcal{H}_p\|_\infty^4 \right)$$

and

$$(5.3) \quad \mathcal{R}_{m_0} := \left\{ \mu \in \mathcal{P}_1(H) : \int_H |x|^4 \mu(dx) \leq \hat{c}_{m_0} \right\}.$$

Observe that, by Lemma A.1 and by [38, Remark 7.13 (ii)], the metric \mathbf{d}_1 metrizes on \mathcal{R}_{m_0} the topology of weak convergence in $\mathcal{P}(H)$. In view of Proposition 5.2, we also define

$$\mathcal{Q}_{m_0} := \left\{ \mu \in \mathcal{R}_{m_0} : \int_H \langle x, \mathbf{e}_k \rangle^2 \mu(dx) \leq a_k \right\}.$$

By Proposition A.2, \mathcal{Q}_{m_0} is \mathbf{d}_1 -compact. We now consider a subset of \mathcal{S} of functions which take values in the set \mathcal{Q}_{m_0} and are \mathbf{d}_1 -equi-uniformly continuous. More precisely, recalling Proposition 2.5(ii) we define

$$(5.4) \quad \omega_{m_0} := \omega_{m_0, \|\mathcal{H}_p\|_\infty}$$

and consider the subset $\mathcal{C}_{m_0} \subset \mathcal{S}$ defined as

$$(5.5) \quad \mathcal{C}_{m_0} := \left\{ m : [0, T] \rightarrow \mathcal{Q}_{m_0} : m(0) = m_0, \mathbf{d}_1(m(t), m(s)) \leq \omega_{m_0}(|t - s|) \right\}.$$

Lemma 5.3. *Let Assumptions 2.1, 2.3(i) and 2.6 hold. The set \mathcal{C}_{m_0} is convex and compact in $(\mathcal{S}, \rho_\infty)$.*

Proof. Relative compactness. Relative compactness follows from Proposition A.2 and the Arzelà-Ascoli's theorem for functions with values in complete metric spaces.

Closedness. Let $\{m_n\}_{n \in \mathbb{N}} \subset \mathcal{C}_{m_0}$ be such that $m_n \rightarrow m \in \mathcal{S}$. This means that

$$\sup_{[0, T]} \mathbf{d}_1(m_n(t), m(t)) \rightarrow 0.$$

Clearly $m(0) = m_0$. Moreover, $m_n(t)$ converges weakly to $m(t)$ for all $t \in [0, T]$, so

$$\int_H (|x|^4 \wedge K) m(t, dx) = \lim_{n \rightarrow \infty} \int_H (|x|^4 \wedge K) m_n(t, dx) \leq \hat{c}_{m_0}, \quad \forall t \in [0, T], \quad \forall K > 0.$$

Hence, by dominated convergence

$$\int_H |x|^4 m(t, dx) = \lim_{K \rightarrow \infty} \int_H (|x|^4 \wedge K) m(t, dx) \leq \hat{c}_{m_0}.$$

Similarly, one shows that

$$\sup_{t \in [0, T]} \int_H \langle x, \mathbf{e}_k \rangle^2 m(t, dx) \leq a_k \quad \forall k \in \mathbb{N}.$$

Next, for $t, s \in [0, T]$, we have

$$\begin{aligned} \mathbf{d}_1(m(t), m(s)) &\leq \mathbf{d}_1(m_n(t), \mu(t)) + \mathbf{d}_1(m_n(s), m(s)) + \mathbf{d}_1(m_n(t), m_n(s)) \\ &\leq \mathbf{d}_1(m_n(t), m(t)) + \mathbf{d}_1(m_n(s), m(s)) + \omega_{m_0}(|t - s|) \end{aligned}$$

Taking the limit as $n \rightarrow \infty$ above, we get

$$\mathbf{d}_1(m(t), m(s)) \leq \omega_{m_0}(|t - s|).$$

Convexity. Let $m_1, m_2 \in \mathcal{C}_{m_0}$, $\lambda \in [0, 1]$, and set $m_\lambda := \lambda m_1 + (1 - \lambda)m_2$. The only nontrivial claim to verify is that, for $t, s \in [0, T]$,

$$\mathbf{d}_1(m_\lambda(t), m_\lambda(s)) \leq \omega_{m_0}(|t - s|).$$

We will use the fact (see, e.g., [38, Rem. 7.5(i)] or [9, Cor. 5.4]) that for $\mu, \nu \in \mathcal{P}_1(H)$

$$\mathbf{d}_1(\mu, \nu) = \sup_{f \in \text{Lip}_1} \int_H f(x)(\mu(dx) - \nu(dx)),$$

where Lip_1 is the set of Lipschitz functions on H with Lipschitz constant 1. Then

$$\begin{aligned} \mathbf{d}_1(m_\lambda(t), m_\lambda(s)) &= \sup_{f \in \text{Lip}_1} \int_H f(x)(m_\lambda(t, dx) - m_\lambda(s, dx)) \\ &\leq \lambda \sup_{f \in \text{Lip}_1} \int_H f(x)(m_1(t, dx) - m_1(s, dx)) + (1 - \lambda) \sup_{f \in \text{Lip}_1} \int_H f(x)(m_2(t, dx) - m_2(s, dx)) \\ &= \lambda \mathbf{d}_1(m_1(t), m_1(s)) + (1 - \lambda) \mathbf{d}_1(m_2(t), m_2(s)) \leq \omega_{m_0}(|t - s|), \end{aligned}$$

the claim. \square

In order to use a fixed point theorem to prove existence of solutions to the MFG system, we need to embed properly \mathcal{C}_{m_0} into a topological vector space. We consider the vector space $C([0, T]; \mathcal{M}(H))$, where $\mathcal{M}(H)$ is endowed with the weak topology induced by the family of seminorms (2.1). On the space $C([0, T]; \mathcal{M}(H))$, we define a suitable topology as follows. Define the family of seminorms

$$|m(\cdot)|_f := \sup_{t \in [0, T]} \left| \int_H f(x) m(t, dx) \right|, \quad f \in C_b(H).$$

This family of seminorms induces on $C([0, T]; \mathcal{M}(H))$ a topology, that we denote by τ_{uw} and call the *uniform-weak topology*, that makes $C([0, T]; \mathcal{M}(H))$ a locally convex topological vector space.

Lemma 5.4. *Let Assumptions 2.1, 2.3(i) and 2.6 hold. We have the following inclusion:*

$$\mathcal{C}_{m_0} \subset C([0, T]; \mathcal{M}(H)).$$

Moreover, the topology induced by ρ_∞ on \mathcal{C}_{m_0} is the same as the one induced by τ_{uw} .

Proof. In the following we use the notation $m(\cdot)$ to denote elements of \mathcal{C}_{m_0} in order to avoid confusion.

We first prove the inclusion. Obviously, every function in \mathcal{C}_{m_0} can also be seen as a function with values in $\mathcal{M}(H)$. The inclusion then holds if we prove that every function in \mathcal{C}_{m_0} is continuous as a function from $[0, T]$ to $\mathcal{M}(H)$. Indeed, let $m(\cdot) \in \mathcal{C}_{m_0}$; then, for any $t_0 \in [0, T]$ and any sequence $t_n \rightarrow t_0$ one has $\mathbf{d}_1(m(t_n), m(t_0)) \rightarrow 0$ as $n \rightarrow \infty$. This implies that, in particular, $m(t_n)$ converges weakly to $m(t_0)$ when $n \rightarrow \infty$, which completes the proof.

(i) $\tau_{\mathbf{uw}}$ stronger than ρ_∞ . Let $\bar{m}(\cdot) \in \mathcal{C}_{m_0}$, $\varepsilon > 0$ and consider the ρ_∞ -neighborhood

$$\mathcal{U}(\bar{m}(\cdot)) := \left\{ m(\cdot) \in \mathcal{C}_{m_0} : \sup_{t \in [0, T]} \mathbf{d}_1(m(t), \bar{m}(t)) < \varepsilon \right\}.$$

By the equiuniform \mathbf{d}_1 -continuity of the elements of \mathcal{C}_{m_0} , there exists $\{t_1, \dots, t_N\} \subset [0, T]$ such that

$$\mathcal{U}_N(\bar{m}(\cdot)) := \left\{ m(\cdot) \in \mathcal{C}_{m_0} : \mathbf{d}_1(m(t_i), \bar{m}(t_i)) < \varepsilon/2, \quad \forall i = 1, \dots, N \right\} \subseteq \mathcal{U}(\bar{m}(\cdot)).$$

Since \mathbf{d}_1 metrizes the weak topology in \mathcal{Q}_{m_0} , for each $i = 1, \dots, N$ we have the existence of $\delta > 0$ and a family $\{f_{i,1}, \dots, f_{i,M}\} \subset C_b(H)$ such that

$$\begin{aligned} \mathcal{V}_{t_i}(\bar{m}(t_i)) &:= \{ \mu \in \mathcal{Q}_{m_0} : |\mu - \bar{m}(t_i)|_{f_{i,j}} < \delta, \quad \forall j = 1, \dots, M \} \\ &\subseteq \{ \mu \in \mathcal{Q}_{m_0} : \mathbf{d}_1(\mu, \bar{m}(t_i)) < \varepsilon/2 \} =: \mathcal{U}_{t_i}(\bar{m}(t_i)) \end{aligned}$$

Define now the $\tau_{\mathbf{uw}}$ -neighborhood of $\bar{m}(\cdot)$

$$\mathcal{V}(\bar{m}(\cdot)) := \left\{ \mu \in \mathcal{C}_{m_0} : |m(\cdot) - \bar{m}(\cdot)|_{f_{i,j}} < \delta, \quad \forall j = 1, \dots, M, \quad \forall i = 1, \dots, N \right\}.$$

Then,

$$\begin{aligned} \mathcal{V}(\bar{m}(\cdot)) &\subseteq \{ m(\cdot) \in \mathcal{C}_{m_0} : m(t_i) \in \mathcal{V}_{t_i}(\bar{m}(t_i)) \quad \forall i = 1, \dots, N \} \\ &\subseteq \{ m(\cdot) \in \mathcal{C}_{m_0} : m(t_i) \in \mathcal{U}_{t_i}(\bar{m}(t_i)), \quad \forall i = 1, \dots, N \} \\ &= \mathcal{U}_N(\bar{m}(\cdot)) \subseteq \mathcal{U}(\bar{m}(\cdot)), \end{aligned}$$

concluding the proof of this part.

(ii) ρ_∞ stronger than $\tau_{\mathbf{uw}}$. First, let us show that, for each $f \in C_b(H)$, the family

$$(5.6) \quad \left\{ [0, T] \rightarrow \mathbb{R}, \quad t \mapsto \int_H f(x) m(t, dx) \right\}_{m(\cdot) \in \mathcal{C}_{m_0}}$$

is equiuniformly continuous; that is, there exists a modulus of continuity ω_f such that

$$(5.7) \quad \left| \int_H f(x) (m(t, dx) - m(s, dx)) \right| \leq \omega_f(|t - s|), \quad \forall t, s \in [0, T], \quad \forall m \in \mathcal{C}_{m_0}.$$

Assume, by contradiction, that this is not true. Then, we may find $\eta > 0$ and suitable sequences $\{s_n\}, \{t_n\} \subset [0, T]$, $\{m_n(\cdot)\} \subset \mathcal{C}_{m_0}$, such that

$$(5.8) \quad \left| \int_H f(x) (m_n(t_n, dx) - m_n(s_n, dx)) \right| \geq \eta, \quad |s_n - t_n| \rightarrow 0.$$

Now, on the one hand, by \mathbf{d}_1 -equiuniform continuity of the functions belonging to \mathcal{C}_{m_0} , we have

$$(5.9) \quad \mathbf{d}_1(m_n(s_n), m_n(t_n)) \rightarrow 0;$$

on the other hand, by weak compactness of \mathcal{Q}_{m_0} , up to passing to subsequences, we have

$$(5.10) \quad m_n(s_n) \xrightarrow{w} \mu_1, \quad m_n(t_n) \xrightarrow{w} \mu_2$$

for some $\mu_1, \mu_2 \in \mathcal{Q}_{m_0}$. By (5.9), we must have

$$(5.11) \quad \mu_1 = \mu_2.$$

Then, letting $n \rightarrow \infty$ in (5.8) and using (5.10) and (5.11), we get

$$0 < \eta \leq \lim_{n \rightarrow \infty} \left| \int_H f(x)(m_n(t_n, dx) - m_n(s_n, dx)) \right| = 0,$$

a contradiction. This proves (5.6).

Now, let $\bar{m}(\cdot) \in \mathcal{C}_{m_0}$, $\varepsilon > 0$, $\{f_1, \dots, f_N\} \subset C_b(H)$ and consider the $\tau_{\mathbf{u}\mathbf{w}}$ -neighborhood

$$\mathcal{V}(\bar{m}(\cdot)) := \left\{ m(\cdot) \in \mathcal{C}_{m_0} : |m(t) - \bar{m}(t)|_{f_i} < \varepsilon \quad \forall t \in [0, T], \forall i = 1, \dots, N \right\}.$$

By (5.6), we have the existence of $\{t_1, \dots, t_M\} \subset [0, T]$ such that

$$\begin{aligned} \mathcal{V}_N(\bar{m}(\cdot)) &:= \left\{ m(\cdot) \in \mathcal{C}_{m_0} : |m(t_j) - \bar{m}(t_j)|_{f_i} < \varepsilon/2, \quad \forall i = 1, \dots, N, \forall j = 1, \dots, M \right\} \\ &\subseteq \mathcal{V}(\bar{m}(\cdot)). \end{aligned}$$

Since \mathbf{d}_1 metrizes the weak topology on \mathcal{Q}_{m_0} , for each $j = 1, \dots, M$ we have the existence of $\delta_j > 0$ such that

$$\mathbf{d}_1(m(t_j), \bar{m}(t_j)) < \delta_j \Rightarrow |m(t_j) - \bar{m}(t_j)|_{f_i} < \varepsilon/2, \quad \forall i = 1, \dots, N.$$

Let now $\delta := \min\{\delta_1, \dots, \delta_M\} > 0$. Then,

$$\begin{aligned} \mathcal{U}(\bar{m}(\cdot)) &:= \left\{ m(\cdot) \in \mathcal{C}_{m_0} : \sup_{t \in [0, T]} \mathbf{d}_1(m(t), \bar{m}(t)) < \delta \right\} \\ &\subseteq \left\{ m(\cdot) \in \mathcal{C}_{m_0} : \mathbf{d}_1(m(t_j), \bar{m}(t_j)) < \delta \quad \forall j = 1, \dots, M \right\} \\ &\subseteq \mathcal{V}_N(\bar{m}(\cdot)) \subseteq \mathcal{V}(\bar{m}(\cdot)), \end{aligned}$$

concluding the proof. □

We now define a solution to the MFG system **(HJB)**–**(FP)**.

Definition 5.5. *A pair (v, m) is a solution to the MFG system **(HJB)**–**(FP)** if v is a mild solution to **(HJB)** (Definition 4.1) and m is a weak solution of (3.1) with $w(t, x) = \mathcal{H}_p(x, Dv(t, x), m(t))$ (Definition 3.1).*

Theorem 5.6. *Let Assumptions 2.1, 2.3(i), and 2.6 hold. Then, the MFG system **(HJB)**–**(FP)** admits a solution.*

Proof. By construction, if m^* is a fixed point of the map Ψ defined in (5.2), then the couple $(v^{(m^*)}, m^*)$ is a solution to the MFG system **(HJB)**–**(FP)**. We are therefore going to show that Ψ admits a fixed point.

We consider \mathcal{C}_{m_0} embedded in the topological vector space $(C([0, T]; \mathcal{M}(H)), \tau_{\mathbf{u}\mathbf{w}})$. By Lemma 5.4 we can indifferently use on \mathcal{C}_{m_0} the $\tau_{\mathbf{u}\mathbf{w}}$ or the ρ_∞ topology. In what follows, the topological notions are referred, indifferently, to one of those equivalent topologies. We want to apply Tikhonov's fixed point theorem. First of all, we notice that, by the definitions of \hat{c}_{m_0} and \mathbf{a} , using Proposition 2.5 and Proposition 5.2, we have

$$(5.12) \quad \Psi(\mathcal{C}_{m_0}) \subseteq \mathcal{C}_{m_0}.$$

Since \mathcal{C}_{m_0} is compact and convex, and since (5.12) holds, in order to conclude the existence of a fixed point of Ψ in \mathcal{C}_{m_0} , we need to show that $\Psi|_{\mathcal{C}_{m_0}}$ is continuous. The remaining part of the proof is devoted to this goal.

Let $\{m_n\} \subset \mathcal{C}_{m_0}$ be such that $m_n \rightarrow m \in \mathcal{C}_{m_0}$. Since \mathcal{C}_{m_0} is compact, from each subsequence (m_{n_k}) , one can extract a sub-subsequence such that $\Psi(m_{n_{k_h}}) \rightarrow \hat{m}$ for some $\hat{m} \in \mathcal{C}_{m_0}$. Set

$$w^{(m)} := \mathcal{H}_p(\cdot, Dv^{(m)}(\cdot, \cdot), m(\cdot)), \quad w^{(m_{n_{k_h}})} := \mathcal{H}_p(\cdot, Dv^{(m_{n_{k_h}})}(\cdot, \cdot), m_{n_{k_h}}(\cdot)).$$

For each $h \in \mathbb{N}$,

$$\begin{aligned} & \int_H \varphi(t, x) \Psi(m_{n_{k_h}})(t, dx) - \int_H \varphi(0, x) m_0(dx) \\ &= \int_0^t \left(\int_H \left[\partial_t \varphi(s, x) + L_0 \varphi(s, x) - \langle w^{(m_{n_{k_h}})}(s, x), D\varphi(s, x) \rangle \right] \Psi(m_{n_{k_h}})(s, dx) \right) ds \end{aligned}$$

for every $t \in (0, T]$ and $\varphi \in D_T$. Considering that, by Theorem 4.2(ii), we have $w^{(m_{n_{k_h}})} \rightarrow w^{(m)}$ pointwise as $h \rightarrow \infty$, we take the limit as $h \rightarrow \infty$ using dominated convergence theorem in the equation above to obtain

$$\begin{aligned} & \int_H \varphi(t, x) \hat{m}(t, dx) - \int_H \varphi(0, x) m_0(dx) \\ &= \int_0^t \left(\int_H \left[\partial_t \varphi(s, x) + L_0 \varphi(s, x) - \langle w^{(m)}(s, x), D\varphi(s, x) \rangle \right] \hat{m}(s, dx) \right) ds. \end{aligned}$$

On the other hand, we also have

$$\begin{aligned} & \int_H \varphi(t, x) \mathcal{L}(X^{(m)})(t, dx) - \int_H \varphi(0, x) m_0(dx) \\ &= \int_0^t \left(\int_H \left[\partial_t \varphi(s, x) + L_0 \varphi(s, x) - \langle w^{(m)}(s, x), D\varphi(s, x) \rangle \right] \mathcal{L}(X^{(m)})(s, dx) \right) ds. \end{aligned}$$

Hence, by Proposition 3.3, we must have $\hat{m} = \mathcal{L}(X^{(m)})$. By the arbitrariness of $k \mapsto n_k$, it follows that

$$\Psi(m_n) \rightarrow \hat{m} = \mathcal{L}(X^{(m)}) = \Psi(m) \quad \text{in } (\mathcal{S}, \rho_\infty),$$

which gives the continuity of Ψ . \square

6. THE MFG SYSTEM: EXISTENCE AND UNIQUENESS OF SOLUTIONS

In this section, we establish the uniqueness of solutions for our MFG system **(HJB)**–**(FP)**. It is well known, even in finite-dimensional state spaces (see, e.g., [8, Rem. 1.4, p.32]), that uniqueness of a solution for this system cannot generally be guaranteed. A typical approach to ensure uniqueness is to impose certain monotonicity conditions on the data. One of the most common monotonicity conditions is the *Lasry-Lions monotonicity condition* (see, for example, [30, 31, 32] and [8, Section 1.3.2]). For other monotonicity conditions that ensure uniqueness in finite-dimensional MFG systems, readers can refer to [28, 34] and the references therein. Another typical feature of uniqueness results is the separability of the Hamiltonian function \mathcal{H} (as assumed in Assumption 6.1(i) here). Well-posedness of MFG systems without separability has been studied in [34]. Here, we assume separability and use a variant of the Lasry-Lions monotonicity condition from [8, Theorem 1.4]. Our assumptions are as follows.

Assumption 6.1.

(i) The Hamiltonian \mathcal{H} has a separated form, i.e.

$$\mathcal{H}(x, p, \mu) = \mathcal{H}^0(x, p) - F(x, \mu),$$

for some continuous functions

$$\mathcal{H}^0 : H \times H \rightarrow \mathbb{R}, \quad F : H \times \mathcal{P}_1(H) \rightarrow \mathbb{R}.$$

(ii) The map \mathcal{H}^0 is convex in p .

(iii) The functions F and G are monotone in $\mu \in \mathcal{P}_1(H)$ in the following sense

$$\int_H [F(x, \mu_1) - F(x, \mu_2)](\mu_1 - \mu_2)(dx) \geq 0 \quad \forall \mu_1, \mu_2 \in \mathcal{P}_1(H),$$

$$\int_H [G(x, \mu_1) - G(x, \mu_2)](\mu_1 - \mu_2)(dx) \geq 0 \quad \forall \mu_1, \mu_2 \in \mathcal{P}_1(H).$$

(iv) One of the following is satisfied

(a)

$$\int_H [F(x, \mu_1) - F(x, \mu_2)](\mu_1 - \mu_2)(dx) > 0 \quad \forall \mu_1, \mu_2 \in \mathcal{P}_1(H), \mu_1 \neq \mu_2;$$

(b)

$$\int_H [F(x, \mu_1) - F(x, \mu_2)](\mu_1 - \mu_2)(dx) = 0 \quad \implies \quad F(\cdot, \mu_1) = F(\cdot, \mu_2),$$

$$\int_H [G(x, \mu_1) - G(x, \mu_2)](\mu_1 - \mu_2)(dx) = 0 \quad \implies \quad G(\cdot, \mu_1) = G(\cdot, \mu_2);$$

(c)

$$\mathcal{H}^0(x, p_1) - \mathcal{H}^0(x, p_2) - \langle \mathcal{H}_p^0(x, p_2), p_1 - p_2 \rangle = 0$$

$$\implies \quad \mathcal{H}_p^0(x, p_1) = \mathcal{H}_p^0(x, p_2), \quad \forall x \in H, p_1, p_2 \in H$$

Theorem 6.2. *Let Assumptions 2.3(i), 2.6, and 6.1 hold. Then, there exists a unique solution to the MFG system (HJB)–(FP).*

Proof. Existence. Existence of a solution was proved in Theorem 5.6.

Uniqueness. The proof is an adaptation to our Hilbert space setting of a typical proof of uniqueness in finite dimensional spaces (see, e.g., [8, Theorem 1.4]).

Let (v_1, m_1) and (v_2, m_2) be two solutions of our MFG system. We set $\bar{v} = v_1 - v_2$ and $\bar{m} = m_1 - m_2$. Then, \bar{v} solves the integral equation

$$\bar{v}(t, x) = R_{T-t}[G(\cdot, m_1(T)) - G(\cdot, m_2(T))](x) - \int_t^T R_{s-t}[\mathcal{H}^0(\cdot, Du_1(s, \cdot)) - \mathcal{H}^0(\cdot, Du_2(s, \cdot))](x) ds$$

$$- \int_t^T R_{s-t}[F(\cdot, m_1(s)) - F(\cdot, m_2(s))](x) ds,$$

which means that \bar{v} is a mild solution to the equation

$$\begin{cases} -\partial_t \bar{v} - L\bar{v} + \mathcal{H}^0(x, Dv_1(t, x)) - \mathcal{H}^0(x, Dv_2(t, x)) - (F(x, m_1(t)) - F(x, m_2(t))) = 0, \\ \bar{v}(T, x) = G(x, m_1(T)) - G(x, m_2(T)). \end{cases}$$

On the other hand, observing that that $\bar{m}_0 \equiv 0$, we get that \bar{m} satisfies, for every $\varphi \in \mathcal{D}_T$,

(6.1)

$$\int_H \varphi(t, x) \bar{m}(t, dx) - \int_H \varphi(0, x) \bar{m}(0, dx) = \int_0^T \int_H [\partial_t \varphi(s, x) + L_0 \varphi(s, x)] \bar{m}(s, dx)$$

$$- \int_0^T \left(\int_H [\langle \mathcal{H}_p^0(x, Dv_1(s, x)), D\varphi(s, x) \rangle] m_1(s, dx) - \int_H [\langle \mathcal{H}_p^0(x, Dv_2(s, x)), D\varphi(s, x) \rangle] m_2(s, dx) \right) ds.$$

We now set

$$f(s, x) := \mathcal{H}^0(x, Dv_1(s, x)) - \mathcal{H}^0(x, Dv_2(s, x)) - (F(x, m_1(s)) - F(x, m_2(s))).$$

By Assumption 2.6, by Proposition 2.5, and by the definition of solution to (HJB)-(FP), we have $f \in C_{b,\gamma}([0, T] \times H)$, where the latter space is defined in (2.4). Then, we consider the Kolmogorov equation

$$\begin{cases} \partial_t v(t, x) + Lv(t, x) = f(t, x), \\ v(T, x) = \phi(x) := G(x, m_1(T)) - G(x, m_2(T)), \end{cases}$$

By Theorem B.4, \bar{v} is also a \mathcal{K} -strong solution of the same equation and moreover, we can choose the approximating data ϕ_n, f_n such that the solutions \bar{v}_n of the approximating Kolmogorov equations

$$\begin{cases} \partial_t \bar{v}_n(t, x) + L_0 \bar{v}_n(t, x) = f_n(t, x), \\ \bar{v}_n(T, x) = \phi_n(x) \end{cases}$$

belong to \mathcal{D}_T for all $n \in \mathbb{N}$. Now, using $\varphi = \bar{v}_n$ in (6.1), we get

$$(6.2) \quad \begin{aligned} \int_H \bar{v}_n(T, x) \bar{m}(t, dx) &= \int_0^T \left(\int_H [\partial_t \bar{v}_n(s, x) + L_0 \bar{v}_n(s, x)] \bar{m}(s, dx) \right) ds \\ &- \int_0^T \left(\int_H [\langle \mathcal{H}_p^0(x, Dv_1(s, x)), D\bar{v}_n(s, x) \rangle] m_1(s, dx) - \int_H [\langle \mathcal{H}_p^0(x, Dv_2(s, x)), D\bar{v}_n(s, x) \rangle] m_2(s, dx) \right) ds. \end{aligned}$$

On the other hand, for every $s \in [0, T]$, we can integrate with respect to the measure $\bar{m}(s)$ the Kolmogorov equations solved by the \bar{v}_n 's, to get

$$\int_H (\partial_t \bar{v}_n(s, x) + L_0 \bar{v}_n(s, x)) \bar{m}(s, dx) = \int_H f_n(s, x) \bar{m}(s, dx).$$

Plugging this equality into (6.2), we obtain

$$(6.3) \quad \begin{aligned} \int_H \bar{v}_n(T, x) \bar{m}(t, dx) &- \int_0^T \left(\int_H f_n(s, x) \bar{m}(s, dx) \right) ds = \\ &- \int_0^T \left(\int_H [\langle \mathcal{H}_p^0(x, Dv_1(s, x)), D\bar{v}_n(s, x) \rangle] m_1(s, dx) - \int_H [\langle \mathcal{H}_p^0(x, Dv_2(s, x)), D\bar{v}_n(s, x) \rangle] m_2(s, dx) \right) ds. \end{aligned}$$

We now need to pass to the limit in (6.3) as $n \rightarrow \infty$. By the definition of \mathcal{K} -convergence in $C_{b,\gamma}([0, T] \times H)$ (Definition B.2), there is $M \in \mathbb{R}$ such that for every $(s, x) \in [0, T] \times H$,

$$\sup_{n \geq 1} |f_n(s, x) - f(s, x)| = (T - t)^{-\gamma} \sup_{n \geq 1} \left[(T - s)^\gamma |f_n(s, x) - f(s, x)| \right] \leq M(T - s)^{-\gamma}.$$

Hence, since $(T - s)^{-\gamma} \in L^1(0, T)$ and $\lim_{n \rightarrow \infty} |f_n(s, x) - f(s, x)| = 0$ for every $(s, x) \in [0, T] \times H$, we obtain

$$\lim_{n \rightarrow \infty} \int_0^T \left(\int_H |f_n(s, x) - f(s, x)| \bar{m}(s, dx) \right) ds = 0$$

by the Dominated Convergence Theorem. The convergence of the other terms follows directly from the Dominated Convergence Theorem, since the functions involved are uniformly bounded.

Therefore, passing to the limit in (6.3), we get

$$(6.4) \quad \begin{aligned} 0 = & - \int_H \bar{v}(T, x) \bar{m}(t, dx) + \int_0^T \left(\int_H f(s, x) \bar{m}(s, dx) \right) ds \\ & - \int_0^T \left(\int_H [\langle \mathcal{H}_p^0(x, Dv_1(s, x)), D\bar{v}(s, x) \rangle] m_1(s, dx) - \int_H [\langle \mathcal{H}_p^0(x, Dv_2(s, x)), D\bar{v}(s, x) \rangle] m_2(s, dx) \right) ds. \end{aligned}$$

Hence, using the definition of f , we get

$$\begin{aligned} 0 = & \int_H \bar{u}(T, x) \bar{m}(t, dx) + \int_0^T \int_H [F(x, m_1(s)) - F(x, m_2(s))] \bar{m}(s, dx) \\ & + \int_0^T \int_H [\mathcal{H}^0(x, Dv_2(s, x)) - \mathcal{H}^0(x, Dv_1(s, x)) - \langle \mathcal{H}_p^0(x, Dv_1(s, x)), -D\bar{v}(s, x) \rangle] m_1(s, dx) \\ & + \int_0^T \int_H [\mathcal{H}^0(x, Dv_1(s, x)) - \mathcal{H}^0(x, Dv_2(s, x)) - \langle \mathcal{H}_p^0(x, Dv_2(s, x)), D\bar{v}(s, x) \rangle] m_2(s, dx). \end{aligned}$$

By Assumption 6.1(ii)-(iii) all terms above are nonnegative, so they all must be equal to 0. We now conclude as follows.

- (a) If case (a) of Assumption 6.1(iv) holds, we have $m_1 = m_2$ and then uniqueness of mild solutions of equation (HJB) gives $v_1 = v_2$.
- (b) If case (b) of Assumption 6.1(iv) holds, we first use uniqueness of mild solutions of equation (HJB) which holds since $F(x, m_1(s)) = F(x, m_2(s))$ and $G(x, m_1(T)) = G(x, m_2(T))$ to obtain $v_1 = v_2$ and then conclude that $m_1 = m_2$ by uniqueness of weak solutions of (3.5) with $w(s, x) := \mathcal{H}_p^0(x, Dv_1(s, x)) = \mathcal{H}_p^0(x, Dv_2(s, x))$.
- (c) If case (c) of Assumption 6.1(iv) holds, we first obtain $\mathcal{H}_p^0(x, Dv_1(s, x)) = \mathcal{H}_p^0(x, Dv_2(s, x))$, $m_1(s)$ and $m_2(s)$ a.e. for $s \in [0, T]$. This implies $m_1 = m_2$ by uniqueness of weak solutions of (3.5) with $w(s, x) = \mathcal{H}_p^0(x, Dv_1(s, x))$ (or equivalently with $w(s, x) = \mathcal{H}_p^0(x, Dv_2(s, x))$) and then conclude that $v_1 = v_2$ using uniqueness of mild solutions of equation (HJB). □

7. EXAMPLES FROM STOCHASTIC OPTIMAL CONTROL

Typically, the Hamiltonian \mathcal{H} comes from stochastic optimal control problems. Consider the following H -valued stochastic optimal control problem for which the value function

$$v(t, x) = \inf_{\alpha \in \mathcal{A}_t} \mathbf{E} \left[\int_t^T f(X_s^{t,x,\alpha(\cdot)}, \alpha_s, m(s)) ds + g(X_T^{t,x,\alpha(\cdot)}, m(T)) \right],$$

where

$$\mathcal{A}_t = \{ \alpha : [t, T] \times \Omega \rightarrow \Lambda \text{ (}\mathcal{F}_s^t\text{) - progressively measurable} \}$$

for some reference probability space $(\Omega, \mathcal{F}, \mathcal{F}_s^t, \mathbf{P}, W)$, Λ is a complete separable metric space, $m(\cdot) \in \mathcal{S}$, and $X^{t,x,\alpha(\cdot)}$ solves, in mild sense, on $[t, T]$ the infinite dimensional SDE

$$dX_s = (AX_s + b(X_s, \alpha_s, m(s))) ds + dW_s, \quad X_t = x.$$

We then have

$$\mathcal{H}(x, p, \mu) = \sup_{\alpha \in \Lambda} \{ -b(x, \alpha, \mu)p - f(x, \alpha, \mu) \}.$$

We make the following assumptions:

- (i) $\Lambda = B_R \subset H$;
- (ii) $b(x, \alpha, \mu) = -\alpha + b_0(x)$, where $b_0 : H \rightarrow \mathbb{R}$ is bounded and Lipschitz continuous;

- (iii) $f(x, \alpha, \mu) = f_0(x, \mu) + f_1(|\alpha|)$, where $f_0 : H \times \mathcal{P}_1(H)$ is continuous in all variables, $f_0(\cdot, \delta_0)$ is bounded, f_0 is Lipschitz continuous in m with respect to \mathbf{d}_1 , uniformly in x , and $f_1 \in C_{\text{loc}}^{1,1}(\mathbb{R})$ is symmetric and uniformly convex⁵;
- (iv) $g : H \times \mathcal{P}_1(H) \rightarrow \mathbb{R}$ is continuous and $g(\cdot, \delta_0)$ is bounded.

Then,

$$\mathcal{H}(x, p, \mu) = \sup_{|\alpha| \leq R} \{ \langle \alpha, p \rangle - f_1(|\alpha|) \} - \langle b_0(x), p \rangle - f_0(x, \mu) =: \mathcal{H}^1(p) - \langle b_0(x), p \rangle - f_0(x, \mu)$$

and in this case

$$\mathcal{H}^0(x, p) = \mathcal{H}^1(p) - \langle b_0(x), p \rangle, \quad F(x, \mu) = f_0(x, \mu), \quad G(x, \mu) = g(x, \mu).$$

It is easy to see that Assumption 2.6(i) is satisfied. Regarding Assumption 2.6(ii), we compute explicitly that

$$\mathcal{H}^1(p) = \begin{cases} (f_1')^{-1}(|p|)|p| - f_1((f_1')^{-1}(|p|)), & \text{if } |p| < f_1'(R), \\ R|p| - f_1(R), & \text{if } |p| \geq f_1'(R). \end{cases}$$

Since $f_1 \in C^{1,1}(\mathbb{R})$ and is uniformly convex, we deduce that $(f_1')^{-1}$ is Lipschitz continuous, so H is Lipschitz continuous in p too. Moreover, we have

$$D\mathcal{H}^1(p) = \begin{cases} (f_1')^{-1}(|p|)\frac{p}{|p|}, & \text{if } |p| < f_1'(R), \\ R\frac{p}{|p|}, & \text{if } |p| \geq f_1'(R), \end{cases}$$

and it is clear that this function is Lipschitz continuous. Hence, $\mathcal{H}_p^0(x, p) = D\mathcal{H}^1(p) - b_0(x)$ is bounded and Lipschitz continuous in p , uniformly in $x \in H$.

Examples of functions satisfying Assumption 6.1(iii) when $H = \mathbb{R}^d$ are for instance in Section 3.4.2 of [10], volume I, and similar examples also work in a real separable Hilbert space H and satisfy the conditions imposed in this paper. We present two examples.

Example 7.1. Let $h_1 : H \times \mathcal{P}_1(H) \rightarrow \mathbb{R}$ and $h_2 : H \times \mathcal{P}_1(H) \rightarrow H$ be Lipschitz continuous and bounded. We define

$$F_1(x, \mu) := h_1(x) \int_H h_1(y) \mu(dy), \quad F_2(x, \mu) := \left\langle h_2(x), \int_H h_2(y) \mu(dy) \right\rangle.$$

We then have

$$\int_H [F_i(x, \mu_1) - F_i(x, \mu_2)] (\mu_1 - \mu_2)(dx) = \left| \int_H h_i(y) m(dy) \right|^2, \quad i = 1, 2.$$

Hence, Assumption 6.1(iii) and Assumption 6.1(iv)(b) are satisfied for F_1 and F_2 . It is also easy to see that, for every $x \in H$,

$$|F_i(x, \mu_1) - F_i(x, \mu_2)| \leq C_i \mathbf{d}_1(\mu_1, \mu_2), \quad i = 1, 2,$$

where C_i is the Lipschitz constant of h_i . Thus both functions satisfy Assumption 2.6(i).

Example 7.2. Let $\ell : H \times [0, \infty) \rightarrow \mathbb{R}$ be bounded, continuous, and such that $\ell(x, \cdot)$ is strictly increasing for every $x \in H$ and is Lipschitz continuous with Lipschitz constant independent of x .

⁵This means that there exists $\eta > 0$ such that $s \mapsto f_1(s) - \eta s^2$ is convex.

Let $\rho : H \rightarrow [0, \infty)$ be bounded and Lipschitz continuous. Let ν be a positive finite measure on H which is full⁶. We define

$$F(x, \mu) := \int_H \ell(z, \rho * \mu(z)) \rho(z - x) \nu(dz),$$

where $\rho * m$ denotes the convolution of ρ with μ , that is

$$\rho * \mu(z) = \int_H \rho(z - u) \nu(du).$$

We notice first that this convolution operation is Lipschitz continuous in μ and z . Indeed, let $z, y \in H$ and $\mu_1, \mu_2 \in \mathcal{P}_1(H)$ and let $\gamma \in \Gamma(\mu_1, \mu_2)$. Then,

$$\begin{aligned} |\rho * \mu_1(z) - \rho * \mu_2(y)| &= \left| \int_H \rho(z - u) \mu_1(du) - \int_H \rho(y - w) \mu_2(dw) \right| \\ &= \left| \int_H (\rho(z - u) - \rho(y - w)) \gamma(du, dw) \right| \leq C \int_H |(z - u) - (y - w)| \gamma(du, dw) \\ &\leq C|z - y| + C \int_H |u - w| \gamma(du, dw). \end{aligned}$$

Taking the infimum over all $\gamma \in \Gamma(\mu_1, \mu_2)$ we thus obtain

$$|\rho * \mu_1(z) - \rho * \mu_2(y)| \leq C(|z - y| + \mathbf{d}_1(\mu_1, \mu_2)).$$

We now have

$$\begin{aligned} &\int_H [F(x, \mu_1) - F(x, \mu_2)] (\mu_1 - \mu_2)(dx) \\ &= \int_H [\ell(z, \rho * \mu_1(z)) - \ell(z, \rho * \mu_2(z))] \int_H \rho(z - x) (\mu_1(dx) - \mu_2(dx)) \nu(dz) \\ &= \int_H [\ell(z, \rho * \mu_1(z)) - \ell(z, \rho * \mu_2(z))] [\rho * \mu_1(z) - \rho * \mu_2(z)] \nu(dz) \geq 0, \end{aligned}$$

because $\ell(z, \cdot)$ is increasing. Moreover, if the last expression is equal to 0, using that $\ell(z, \cdot)$ is strictly increasing, we conclude that $\rho * \mu_1(z) - \rho * \mu_2(z) = 0, \mu$ a.e.. However, since this function is continuous and μ is full, this implies that $\rho * \mu_1(z) = \rho * \mu_2(z)$ for every $z \in H$. Hence, $F(\cdot, \mu_1) = F(\cdot, \mu_2)$ on H . Therefore, F satisfies Assumption 6.1(iii) and Assumption 6.1(iv)(b). Finally, we compute

$$\begin{aligned} |F(x, \mu_1) - F(x, \mu_2)| &\leq C \int_H |\ell(z, \rho * \mu_1(z)) - \ell(z, \rho * \mu_2(z))| \mu(dz) \\ &\leq C \int_H |\rho * \mu_1(z) - \rho * \mu_2(z)| \mu(dz) \leq C \mathbf{d}_1(\mu_1, \mu_2) \end{aligned}$$

which means that F satisfies Assumption 2.6(i).

APPENDIX A. COMPACTNESS IN THE SPACE OF PROBABILITY MEASURES

We provide a result about compactness of sets in spaces of probability measures on H which is used in the paper.

⁶That is $\nu(D) > 0$ for every open set D in H . An example of a measure which is full is the nondegenerate Gaussian measure with mean $a \in H$ and covariance operator $Q \in \mathcal{L}_1^+(H)$, see [13, Proposition 1.25] (nondegeneracy means that $\ker Q = \{0\}$).

Lemma A.1. *Let $L > 0$ and*

$$\mathcal{Q}_L := \left\{ \mu \in \mathcal{P}_1(H) : \int_H |x|^2 \mu(dx) \leq L \right\}.$$

Then

$$\lim_{R \rightarrow \infty} \sup_{\mu \in \mathcal{Q}_L} \int_{B_R^c} |x| \mu(dx) \rightarrow 0.$$

Proof. Since

$$\sup_{\mu \in \mathcal{Q}_L} \int_{B_R^c} |x| \mu(dx) \leq \sup_{\mu \in \mathcal{Q}_L} \frac{1}{R} \int_{B_R^c} |x|^2 \mu(dx) \leq \frac{1}{R} \sup_{\mu \in \mathcal{Q}_L} \int_H |x|^2 \mu(dx) = \frac{L}{R},$$

and the claim follows. \square

Proposition A.2. *Let $S \subset \mathcal{P}_1(H)$. If, for some constant $c_1 > 0$, we have*

$$\sup_{\mu \in S} \int_H |x|^2 \mu(dx) \leq c_1 \quad \text{and} \quad \lim_{N \rightarrow \infty} \sup_{\mu \in S} \sum_{i=N}^{\infty} \int_H \langle x, \mathbf{e}_i \rangle^2 \mu(dx) = 0,$$

then S is relatively compact with respect to \mathbf{d}_1 .

Proof. Let $\{\mu_n\} \subseteq S$. By [36, Ch. VI, Th. 2.2]⁷, the set S is tight. Hence, by Prokhorov's theorem, it is weakly relatively compact in $\mathcal{P}_1(H)$; that is, there exists $\mu \in \mathcal{P}_1(H)$ and a subsequence, still labeled by $\{\mu_n\}$, such that

$$\int_H \phi(x) \mu_n(dx) \rightarrow \int_H \phi(x) \mu(dx), \quad \forall \phi \in C_b(H).$$

The claim follows using Lemma A.1 and [38, Th. 7.12(ii)]. \square

APPENDIX B. CLASSICAL AND STRONG SOLUTIONS TO KOLMOGOROV EQUATIONS

In this section, we define classical, mild, and \mathcal{K} -strong solutions of backward Kolmogorov equations and recall an approximation result used in Section 6. The material is mainly taken from [19, Section B.7] and is presented, for the reader's convenience, in a simplified form needed here. We first consider the following terminal value problem (Kolmogorov equation) in the Hilbert space H :

$$(B.1) \quad \begin{cases} \partial_t u(t, x) + L_0 u(t, x) = f(t, x), & (t, x) \in [0, T) \times H, \\ u(T, x) = \varphi(x), & x \in H, \end{cases}$$

where L_0 is given in (3.3), $\varphi \in C_b(H)$ and $f \in C_{b,\gamma}([0, T) \times H)$ (recall that $C_{b,\gamma}([0, T) \times H)$ was defined in (2.4)). Since, as recalled in a footnote in Section 3, the closure (in a suitable weak sense) of the operator L_0 is the generator (again in a suitable weak sense) of the semigroup R_t in (2.10), we can formally rewrite (B.1) in the following mild form:

$$(B.2) \quad u(t, x) = R_{T-t}[\varphi](x) + \int_t^T R_{s-t}[f(s, \cdot)](x) ds.$$

We call the function u above, defined by the right hand side, the *mild solution* of (B.1). Recalling the definition of L_0 in (3.3), we define on $D(L_0)$ the norm

$$(B.3) \quad \|\phi\|_{D(L_0)} := \|\phi\|_0 + \|D\phi\|_0 + \|A^* D\phi\|_0 + \sup_{x \in H} \|D^2 \phi(x)\|_{\mathcal{L}_1(H)}.$$

⁷In [36, Ch. VI, Th. 2.2] it is stated that the second condition alone guarantees relative compactness of S . This is clearly not true, as can be seen by simply taking as S a non compact sequence concentrated on the line generated by \mathbf{e}_1 . However the proof of this theorem there remains valid if we add the assumption of uniform boundedness of second moments.

Arguing as in Theorem 2.7 of [16], it can be proved that $D(L_0)$, endowed with the above norm, is a Banach space⁸. We now recall a notion of classical solution to (B.1) (see [19, Definition B.82])⁹.

Definition B.1. *We say that $u \in C_b([0, T] \times H)$ is a classical solution to (B.1) in $D(L_0)$ if*

$$(B.4) \quad \begin{cases} u(\cdot, x) \in C^1([0, T]), \quad \forall x \in H, \\ u(t, \cdot) \in D(L_0) \text{ for any } t \in [0, T] \text{ and } \sup_{t \in [0, T]} \|u(t, \cdot)\|_{D(L_0)} < \infty, \\ Du, A^*Du \in C_b([0, T] \times H, H), \quad D^2u \in C_b([0, T] \times H, \mathcal{L}_1(H)) \end{cases}$$

and u satisfies (B.1) for every $(t, x) \in [0, T] \times H$.

We stress that Definition B.1 implicitly implies that $\varphi \in D(L_0)$ and $f \in C([0, T] \times H)$.

To introduce a notion of a strong solution, called \mathcal{K} -strong solution, we first recall a special case of the definition of \mathcal{K} -convergence which is needed in this paper (see [19, Definition B.84]).

Definition B.2. *Let Z be a real Hilbert space. Given $(f_n)_{n \in \mathbb{N}} \subset C_{b,\gamma}([0, T] \times H, Z)$, we say that f_n is \mathcal{K} -convergent to $f \in C_{b,\gamma}([0, T] \times H, Z)$ if*

$$(B.5) \quad \begin{cases} \sup_{n \in \mathbb{N}} \|f_n\|_{C_{b,\gamma}([0, T] \times H, Z)} < \infty, \\ \lim_{n \rightarrow \infty} \sup_{(t,x) \in (0, T] \times K} (T-t)^\gamma |f_n(t, x) - f(t, x)| = 0, \end{cases}$$

for every compact set $K \subset H$. In this case, we write $\mathcal{K} - \lim_{n \rightarrow \infty} f_n = f$ in $C_{b,\gamma}([0, T] \times H, Z)$.

The definition of a \mathcal{K} -strong solution to (B.1) provided below is a special case of a more general definition that can be found in [19, Definition B.85].

Definition B.3. *Let $\gamma \in (0, 1)$. Let $\varphi \in C_b(H)$ and $f \in C_{b,\gamma}([0, T] \times H)$. We say that a function $u \in C_b([0, T] \times H)$ is a \mathcal{K} -strong solution in $D(L_0)$ of (B.1) if $u(t, \cdot)$ is Fréchet differentiable for each $t \in [0, T)$ and there exist sequences $(u_n)_{n \in \mathbb{N}} \subset C_b([0, T] \times H)$, $(\varphi_n)_{n \in \mathbb{N}} \subset D(L_0)$, $(f_n)_{n \in \mathbb{N}} \subset C_{b,\gamma}([0, T] \times H)$ such that:*

(i) *For every $n \in \mathbb{N}$, u_n is a classical solution in $D(L_0)$ (cf. Definition B.1) to*

$$(B.6) \quad \begin{cases} w_t = L_0 w + f_n, \\ w(0) = \varphi_n. \end{cases}$$

(ii) *The following limits hold*

$$\begin{cases} \mathcal{K} - \lim_{n \rightarrow \infty} \varphi_n = \varphi, & \text{in } C_b(H), \\ \mathcal{K} - \lim_{n \rightarrow \infty} u_n = u, & \text{in } C_b([0, T] \times H), \end{cases}$$

and

$$\begin{cases} \mathcal{K} - \lim_{n \rightarrow \infty} f_n = f, & \text{in } C_{b,\gamma}([0, T] \times H) \\ \mathcal{K} - \lim_{n \rightarrow \infty} Du_n = Du & \text{in } C_{b,\gamma}([0, T] \times H, H). \end{cases}$$

We end this section with an approximation result which is a stronger version of [19, Theorem B.95(i)].

⁸The definition of the domain of the operator studied in this paper is slightly different, but the arguments of the mentioned reference can be adapted easily. We also notice that the Banach space structure is not essential for our purposes, even if it simplifies the notation.

⁹We also observe that there are other definitions of classical solutions in the literature, see e.g. Section 6.2 of [17, Section 6.2], [25, Definition 4.6], [26, Definition 4.1]. We refer interested readers to [19, Section B.7.1] for a detailed overview on that.

Theorem B.4. $\phi \in C_b(H)$ and $f \in C_{b,\gamma}([0, T] \times H)$. Then the mild solution u (defined in (B.2)) of equation (B.1), is also a \mathcal{K} -strong solution of to (B.1). Moreover the sequence (u_n) can be chosen so that $u_n \in \mathcal{D}_T$ for all $n \in \mathbb{N}$.

Proof. We can apply [19, Theorem B.95 (i)] (in the case when, in the notation there, $m = 0$, $\eta(t) = t^{-\gamma}$), to conclude that u is also a \mathcal{K} -strong solution of the same equation. Moreover, from the proof of [19, Theorem B.95 (i)] one can easily see that the approximating data φ_n , f_n , and consequently the approximating solutions u_n , can be chosen so that $\varphi_n \in D(L_0)$ and $f_n, \partial_t \bar{v}_n \in C_b([0, T] \times H)$ for all $n \in \mathbb{N}$. As a consequence, we get that $u_n \in \mathcal{D}_T$ for all $n \in \mathbb{N}$. \square

Acknowledgements. Salvatore Federico is a member of the GNAMPA-INdAM group and of the PRIN project 2022BEMMLZ “Stochastic control and games and the role of information”, financed by the Italian Ministry of University and Research. Fausto Gozzi has been supported by the Italian Ministry of University and Research (MIUR), in the framework of PRIN projects 2017FKHBA8 001 (The Time-Space Evolution of Economic Activities: Mathematical Models and Empirical Applications) and 20223PNJ8K (Impact of the Human Activities on the Environment and Economic Decision Making in a Heterogeneous Setting: Mathematical Models and Policy Implications).

REFERENCES

- [1] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient Flows in Metric Spaces and in the Space of Probability Measures*. Springer Science & Business Media, 2008.
- [2] V. Barbu and G. Da Prato. *Hamilton-Jacobi Equations in Hilbert Spaces*, Pitman Research Notes in Mathematics Series, vol. 86. Longman, Boston, 1983.
- [3] V. Bogachev, G. Da Prato, and M. Röckner. Existence and uniqueness of solutions for Fokker–Planck equations on Hilbert spaces. *J. Evol. Equ.*, Vol. 10, pp. 487–509, 2010.
- [4] V. Bogachev, G. Da Prato, M. Röckner, and W. Stannat. Uniqueness of solutions to weak parabolic equations for measures. *Bull. Lond. Math. Soc.*, Vol. 39(4), pp. 631–640, 2007.
- [5] V. Bogachev, G. Da Prato, and M. Röckner. Uniqueness for Solutions of Fokker–Planck Equations on Infinite-Dimensional Spaces. *Commun. Partial Differ. Equ.*, Vol. 36(6), pp. 925–939, 2011.
- [6] V. Bogachev, G. Da Prato, and M. Röckner. Fokker–Planck Equations and Maximal Dissipativity for Kolmogorov Operators with Time-Dependent Singular Drifts in Hilbert Spaces. *J. Funct. Anal.*, Vol. 256(4), pp. 1269–1298, 2009.
- [7] A. Calvia, S. Federico, G. Ferrari, and F. Gozzi. A mean-field model of optimal investment. *arXiv: 2404.02871*, 2024.
- [8] P. Cardaliaguet and A. Porretta. An introduction to mean field game theory. In *Mean field games, Lecture Notes in Math.*, Vol. 2281, pp. 1–158. Springer, Cham, 2020.
- [9] R. Carmona and F. Delarue. *Mean Field Games*. Progress in Nonlinear Differential Equations and Their Applications. Springer, Cham, 2017.
- [10] R. Carmona and F. Delarue. *Probabilistic Theory of Mean Field Games with Applications I-II*. Springer, 2018.
- [11] R. Carmona, J. P. Fouque, S. M. Mousavi, and L.-H. Sun. Systemic risk and stochastic games with delay. *J. Optimiz. Theory App.*, Vol. 179, pp. 366–399, 2018.
- [12] S. Cerrai. Weakly continuous semigroups in the space of functions with polynomial growth. *Dynam. Syst. Appl.*, Vol. 4, pp. 351–372, 1995.
- [13] G. Da Prato. *An introduction to infinite-dimensional analysis*. Springer Science & Business Media, 2006.
- [14] G. Da Prato, F. Flandoli, E. Priola, and M. Röckner. Strong Uniqueness for Stochastic Evolution Equations in Hilbert Spaces Perturbed by a Bounded Measurable Drift. *Ann. Prob.*, Vol. 41(5), pp. 3306–3344, 2013.
- [15] G. Da Prato, F. Flandoli, and M. Röckner. Fokker–Planck equations for SPDE with non-trace-class noise. *Comm. Math. Stat.*, Vol. 1, pp. 281–304, 2013.
- [16] G. Da Prato and J. Zabczyk. Regular Densities of Invariant Measures in Hilbert Spaces. *J. Funct. Anal.*, Vol. 130(2), pp. 427–449, 1995.
- [17] G. Da Prato and J. Zabczyk. *Second Order Partial Differential Equations in Hilbert Spaces*, London Mathematical Society Lecture Note Series, Vol. 293. Cambridge University Press, Cambridge, 2002.
- [18] G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions*, Encyclopedia of Mathematics and its Applications, Vol. 152. Cambridge University Press, Cambridge, 2014.

- [19] G. Fabbri, F. Gozzi, and A. Święch. *Stochastic Optimal Control in Infinite Dimension*, Probability Theory and Stochastic Modeling, Vol. 82. Springer, Switzerland, 2017.
- [20] S. Faggian, F. Gozzi, and P. M. Kort. Optimal investment with vintage capital: Equilibrium distributions. *J. Math. Econ.*, Vol. 96, 2021.
- [21] S. Federico, F. Gozzi, and D. Ghilli. Linear-Quadratic Mean Field Games in Hilbert Spaces. *arXiv: 2402.14935*, 2024.
- [22] G. Feichtinger, R. F. Hartl, P. M. Kort, and V. M. Veliov. Anticipation effects of technological progress on capital accumulation: a vintage capital approach. *J. Econ. Theory*, Vol. 126(1), pp. 143–164, 2006.
- [23] J.-P. Fouque and Z. Zhang. Mean field game with delay: a toy model. *Risks*, Vol. 6(3), 2018.
- [24] F. Gozzi. Regularity of Solutions of a Second Order Hamilton-Jacobi Equation and Application to a Control Problem. *Comm. Part. Diff. Equ.*, Vol. 20(5-6), pp. 775–826, 1995.
- [25] F. Gozzi. Strong Solutions for Kolmogorov Equation in Hilbert Spaces. In G. Da Prato and J.-P. Zolesio, editors, *Partial Differential Equation Methods in Control and Shape Analysis*, Lecture Notes in Pure and Applied Mathematics, Vol. 188, pp. 163–187. Dekker, New York, 1997.
- [26] F. Gozzi. *Second Order Hamilton-Jacobi Equations in Hilbert Spaces and Stochastic Optimal Control*. PhD thesis, Scuola Normale Superiore, Pisa, 1998.
- [27] F. Gozzi. Second Order Hamilton-Jacobi Equations in Hilbert Spaces and Stochastic Optimal Control. In G. Da Prato and L. Tubaro, editors, *Stochastic Partial Differential Equations and Applications*, Lecture Notes in Pure and Applied Mathematics, Vol. 227, pp. 255–285. Dekker, New York, 2002.
- [28] P. J. Graber and A. R. Mészáros. On monotonicity conditions for mean field games. *J. Funct. Anal.*, Vol. 285(9), 2023.
- [29] M. Huang, P. E. Caines, and R. P. Malhamé. Large-Population Cost-Coupled LQG Problems with Nonuniform Agents: Individual-Mass Behavior and Decentralized ε -Nash Equilibria. *IEEE Trans. Autom. Control.*, Vol. 52(9), pp. 1560–1571, 2007.
- [30] J. M. Lasry and P.-L. Lions. Jeux à champ moyen. I – Le Cas Stationnaire. *C. R. Math.*, Vol. 343(9), pp. 619–625, 2006.
- [31] J. M. Lasry and P.-L. Lions. Jeux à champ moyen. II – Horizon Fini et Contrôle Optimal. *C. R. Math.*, Vol. 343(10), pp. 679–684, 2006.
- [32] J. M. Lasry and P.-L. Lions. Mean field games. *Jpn. J. Math.*, Vol. 2(1), pp. 229–260, 2007.
- [33] H. Liu and D. Firoozi. Hilbert Space-Valued LQ Mean Field Games: An Infinite-Dimensional Analysis. *arXiv: 2403.01012*, 2024.
- [34] A. R. Mészáros and C. Mou. Mean field games systems under displacement monotonicity. *SIAM J. Math. Anal.*, Vol. 56(1), pp. 529–553, 2024.
- [35] M. Nisio. *Stochastic control theory*. ISI Lecture Notes, Vol. 9, 2015.
- [36] K. R. Parthasarathy. *Probability Measures on Metric Spaces, Probability and Mathematical Statistics, Vol. 3*. Academic Press, New York, 1967.
- [37] M. Röckner, R. Zhu, and X. Zhu. A note on stochastic semilinear equations and their associated Fokker–Planck equations. *J. Math. Anal. App.*, Vol. 415(1), pp. 83–109, 2014.
- [38] C. Villani. *Topics in Optimal Transportation*, Graduate Studies in Mathematics, Vol. 58. American Mathematical Society, 2003.
- [39] S. Wiesinger. Uniqueness for solutions of Fokker–Planck equations related to singular SPDE driven by Lévy and cylindrical Wiener noise. *J. Evol. Equ.*, Vol. 13(2), pp. 369–394, 2013.

S. FEDERICO: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ “ALMA MATER” DI BOLOGNA, PIAZZA DI PORTA SAN DONATO, BOLOGNA, ITALY

Email address: s.federico@unibo.it

F. GOZZI: DIPARTIMENTO DI ECONOMIA E FINANZA, LUISS UNIVERSITY, VIALE ROMANIA 32, ROME, ITALY

Email address: fgozzi@luiss.it

A. ŚWIĘCH: SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332, USA

Email address: swiech@math.gatech.edu