

# Sparsifying Suprema of Gaussian Processes

Anindya De\*   Shivam Nadimpalli†   Ryan O’Donnell‡   Rocco A. Servedio§

November 25, 2024

## Abstract

We give a dimension-independent sparsification result for suprema of centered Gaussian processes: Let  $T$  be any (possibly infinite) bounded set of vectors in  $\mathbb{R}^n$ , and let  $\{\mathbf{X}_t\}_{t \in T}$  be the canonical Gaussian process on  $T$ . We show that there is an  $O_\varepsilon(1)$ -size subset  $S \subseteq T$  and a set of real values  $\{c_s\}_{s \in S}$  such that  $\sup_{s \in S} \{\mathbf{X}_s + c_s\}$  is an  $\varepsilon$ -approximator of  $\sup_{t \in T} \mathbf{X}_t$ . Notably, the size of  $S$  is completely independent of both the size of  $T$  and of the ambient dimension  $n$ .

We use this to show that every norm is essentially a junta when viewed as a function over Gaussian space: Given any norm  $\nu(x)$  on  $\mathbb{R}^n$ , there is another norm  $\psi(x)$  which depends only on the projection of  $x$  along  $O_\varepsilon(1)$  directions, for which  $\psi(\mathbf{g})$  is a multiplicative  $(1 \pm \varepsilon)$ -approximation of  $\nu(\mathbf{g})$  with probability  $1 - \varepsilon$  for  $\mathbf{g} \sim N(0, I_n)$ .

We also use our sparsification result for suprema of centered Gaussian processes to give a sparsification lemma for convex sets of bounded geometric width: Any intersection of (possibly infinitely many) halfspaces in  $\mathbb{R}^n$  that are at distance  $O(1)$  from the origin is  $\varepsilon$ -close, under  $N(0, I_n)$ , to an intersection of only  $O_\varepsilon(1)$  many halfspaces.

We describe applications to agnostic learning and tolerant property testing.

---

\*University of Pennsylvania. Email: [anindya@seas.upenn.edu](mailto:anindya@seas.upenn.edu).

†MIT. Email: [shivamn@mit.edu](mailto:shivamn@mit.edu).

‡Carnegie Mellon University. Email: [odonnell@cs.cmu.edu](mailto:odonnell@cs.cmu.edu).

§Columbia University. Email: [ras2105@columbia.edu](mailto:ras2105@columbia.edu).

# 1 Introduction

A *Gaussian process* is a joint distribution  $\{\mathbf{X}_t\}_{t \in T}$  over (possibly infinitely many) random variables such that any finite subset of them is distributed as a multivariate Gaussian distribution. Gaussian processes are central objects of study in probability theory [Mil71, Dud67, Sud69, Fer75, Tal87, Gor88, Gor92, Tal95, vH18], and their relevance to theoretical computer science and data science has become increasingly clear over the past decade or so; see e.g. [Sto09, DLP12, OH10, PV12, Sto13, Mek15, Nel16, ORS18, BST19, BDOS21]. We refer the reader to [vH16, Ver18, Tal22] for further background on Gaussian processes and their applications.

Much of the interest in Gaussian processes centers around understanding the behavior of their *suprema*. The supremum of a Gaussian process  $\{\mathbf{X}_t\}_{t \in T}$  is the random variable  $\sup_{t \in T} \mathbf{X}_t$ . Since the index set  $T$  of a Gaussian process  $\{\mathbf{X}_t\}_{t \in T}$  may be arbitrarily large or even infinite, the supremum of  $\{\mathbf{X}_t\}_{t \in T}$  can be quite difficult to understand. A crowning achievement in the study of Gaussian processes is the theory of *majorizing measures*, developed by Fernique, Talagrand, and others, which gives a characterization of the supremum of any centered Gaussian process up to constant factors in terms of the *geometry* of the index set  $T$  (see [Fer75, Tal87, LT91, Tal22] and the references therein).

## 1.1 Main Results

Given a bounded set of vectors  $T \subset \mathbb{R}^n$ , we will write  $\{\mathbf{X}_t = t \cdot \mathbf{g}\}_{t \in T}$  for the *canonical Gaussian process on  $T$* , where  $\mathbf{g}$  is distributed according to the  $n$ -dimensional standard Gaussian distribution  $N(0, I_n)$ . It is a standard fact that every finite centered Gaussian process (i.e. a Gaussian process where each component random variable has mean 0) can be realized as a canonical Gaussian process; as such, we will focus exclusively on canonical Gaussian processes.<sup>1</sup>

The main contribution of this work is a *dimension-independent* sparsification result for the suprema of centered Gaussian processes:

**Theorem 1.** Let  $T \subset \mathbb{R}^n$  be a bounded set of vectors and let  $\{\mathbf{X}_t\}_{t \in T}$  be the canonical Gaussian process on  $T$ . For any  $\varepsilon > 0$ , there is a subset  $S \subseteq T$  with

$$|S| = 2^{2^{O(\frac{1}{\varepsilon})}}$$

and a collection of real values  $\{c_s\}_{s \in S}$  such that

$$\mathbf{E} \left[ \left| \sup_{t \in T} \mathbf{X}_t - \sup_{s \in S} \{\mathbf{X}_s + c_s\} \right| \right] \leq \varepsilon \cdot \mathbf{E} \left[ \sup_{t \in T} \mathbf{X}_t \right]. \quad (1)$$

In words, we show that there is a constant-dimensional (non-centered) Gaussian process  $\{\mathbf{X}_s + c_s\}_{s \in S}$ , where  $S \subseteq T$ , that is a high-accuracy approximator (in an  $L_1$ -sense) of the original Gaussian process. With some additional work, it is possible to obtain a *centered* approximator to the original Gaussian process, although the vectors defining this approximator no longer lie in the original set  $T$ ; we record this as [Corollary 23](#). The proof of [Theorem 1](#) relies on Talagrand’s *majorizing measures* theorem [Tal87, Tal22]; we give a technical overview in [Section 1.2](#) and prove [Theorem 1](#) in [Section 3](#).

Next, we obtain a “junta theorem” for norms over Gaussian space as a consequence of [Theorem 1](#). As a motivating example, if  $\nu(x) = \|x\|$  is the  $\ell_2$  norm on  $\mathbb{R}^n$ , then it is easy to see via standard

---

<sup>1</sup>See [Section 2.2](#) for further background on Gaussian processes.

concentration bounds (see, for example, [Proposition 6](#)) that the norm  $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$  given by

$$\psi(x) = \varepsilon\sqrt{n} \left( \sup_{1 \leq i \leq 2^{\Theta(1/\varepsilon^2)}} |x_i| \right)$$

multiplicatively approximates  $\nu(x)$  within a  $(1 \pm \varepsilon)$ -factor on  $(1 - \varepsilon)$ -fraction of  $N(0, I_n)$ . Note that  $\psi(\cdot)$  only depends on  $2^{\Theta(1/\varepsilon^2)}$  directions, independent of the ambient dimension  $n$ . The following theorem says a similar phenomenon holds for *every* norm over Gaussian space:

**Theorem 2.** Fix any norm  $\nu : \mathbb{R}^n \rightarrow \mathbb{R}$ . For  $\varepsilon \in (0, 0.5)$ , there exists another norm  $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$  for which

$$\Pr_{g \sim N(0, I_n)} \left[ \nu(g) \in [(1 - \varepsilon)\psi(g), (1 + \varepsilon)\psi(g)] \right] \geq 1 - \varepsilon \quad (2)$$

where the norm  $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$  is a “ $2^{2^{O(1/\varepsilon^3)}}$ -subspace junta,” i.e.  $\psi(x)$  depends only on the projection of the point  $x$  onto a  $2^{2^{O(1/\varepsilon^3)}}$ -dimensional subspace.

[Theorem 2](#) is in fact an immediate consequence of a “junta theorem” for Gaussian processes whose index set  $T$  is symmetric (i.e.  $t \in T$  whenever  $-t \in T$ ), namely [Corollary 24](#). (See [Remark 25](#) for more on this.)

We also use our sparsification result, [Theorem 1](#), to obtain a general sparsification lemma for convex sets of bounded “geometric width” (which was the original impetus for this work). Formally, given two measurable sets  $K, L \subseteq \mathbb{R}^n$ , let

$$\text{dist}_G(K, L) := \Pr_{g \sim N(0, I_n)} [g \in K \triangle L]$$

denote the *Gaussian distance* between  $K$  and  $L$ , where  $K \triangle L = (K \setminus L) \cup (L \setminus K)$  is the symmetric difference of the sets  $K$  and  $L$ . We prove the following:

**Theorem 3.** Suppose  $K \subseteq \mathbb{R}^n$  is an intersection of arbitrarily many halfspaces that are at distance at most  $r \geq 1$  from the origin, i.e. there exists  $T \subseteq \mathbb{S}^{n-1}$  and a collection of non-negative numbers  $\{r_t\}_{t \in T}$  such that

$$K = \bigcap_{t \in T} \{x \in \mathbb{R}^n : t \cdot x \leq r_t\} \quad \text{where } r_t \leq r \text{ for all } t \in T.$$

For  $0 < \varepsilon < 0.5$ , there exists  $L \subseteq \mathbb{R}^n$  which is an intersection of

$$2^{\exp\left(\tilde{O}\left(\frac{1}{\varepsilon^2}\right) \cdot r^4\right)} \text{ halfspaces}$$

such that  $\text{dist}_G(K, L) \leq \varepsilon$ .

[Theorem 3](#) can be viewed as a convex-set analogue of recently-established sparsification lemmas for “narrow” CNFs in the setting of Boolean functions [[LSZ19](#), [LWZ21](#)]. [Theorem 3](#) immediately yields tolerant testing and agnostic learning results for intersections of narrow halfspaces with respect to the Gaussian distribution; we describe these applications in [Section 5](#).

Finally, we also establish  $2^{(1/\varepsilon)^{\Omega(1)}}$  lower bounds for these sparsification problems (see [Section 1.3](#) for a brief discussion and [Section 6](#) for the details of these lower bounds).

## 1.2 Techniques

We begin by sketching the key ideas behind our main theorem, [Theorem 1](#). To simplify the ensuing discussion, we first rescale the vectors so that  $\mathbf{E}[\sup_{t \in T} \mathbf{X}_t] = 1$ . Now the goal is to show the existence of a subset  $S \subseteq T$  and suitable constants  $\{c_s\}_{s \in S}$ , such that

$$\mathbf{E}\left[\left|\sup_{t \in T} \mathbf{X}_t - \sup_{s \in S} \{\mathbf{X}_s + c_s\}\right|\right] \leq \varepsilon.$$

A natural idea then is to use a clustering of the vectors in  $T$ . In particular, as a first attempt, consider a  $\delta$ -cover of  $T$  for suitably small  $\delta$ , i.e. let  $\mathcal{P}$  be a partition of  $T$  such that each  $P \in \mathcal{P}$  has (Euclidean) diameter  $\delta$ . For each part  $P$  let  $s_P$  be an arbitrarily chosen representative vector. Then, using the fact that the supremum of a Gaussian process has subgaussian tails ([Proposition 10](#)), it follows that

$$\mathbf{E}\left[\left|\sup_{t \in P} \mathbf{X}_t - \{\mathbf{X}_{s_P} + c_P\}\right|\right] = O(\delta) \quad \text{where } c_P := \mathbf{E}\left[\sup_{t \in P} \mathbf{X}_t - \mathbf{X}_{s_P}\right]. \quad (3)$$

In words, [Equation \(3\)](#) means that within each part  $P$ , the supremum of the Gaussian process  $\sup_{t \in P} \mathbf{X}_t$  can be approximated by a single (non-centered) Gaussian random variable  $\mathbf{X}_{s_P} + c_P$ . Crucially, the error in [Equation \(3\)](#) is just dependent on the diameter of  $P$  and is independent of the number of vectors in  $P$  and the ambient dimension of the vectors in  $T$ .

A natural way to approximate  $\sup_{t \in T} \mathbf{X}_t$  is to then take the supremum of the random variables  $\mathbf{X}_{s_P} + c_P$  as  $P$  ranges over  $\mathcal{P}$ . Furthermore, another application of [Proposition 10](#) shows that

$$\mathbf{E}\left[\left|\sup_{t \in T} \mathbf{X}_t - \sup_{P \in \mathcal{P}} \{\mathbf{X}_{s_P} + c_P\}\right|\right] = O\left(\delta \sqrt{\log M}\right), \quad (4)$$

where  $M$  is the number of parts in the partition  $\mathcal{P}$ . Despite the fact that the error term only grows as  $O(\sqrt{\log M})$ , this dependence turns out to be inadequate for the above approach to work. In particular, there are Gaussian processes  $\{\mathbf{X}_t\}_{t \in T}$  such that  $\mathbf{E}[\sup_{t \in T} \mathbf{X}_t] = 1$  and yet for every  $\delta > 0$ , a  $\delta$ -covering of the space has size  $M = 2^{\Theta(1/\delta^2)}$ , which means that the R.H.S. of [Equation \(4\)](#) is an absolute constant.<sup>2</sup>

To circumvent this barrier, we consider a *multiscale* clustering instead of using a clustering of the entire space with balls of a single radius  $\delta$ ; this naturally leads us to Talagrand’s celebrated *majorizing measures* theorem [[Tal87](#)]. (The modern treatment of this theorem is in terms of the so-called *generic chaining* [[Tal22](#)].) For a Gaussian process  $\{\mathbf{X}_t\}_{t \in T}$  with  $\mathbf{E}[\sup_{t \in T} \mathbf{X}_t] = 1$ , the generic chaining formulation of the majorizing measures theorem can be seen as giving a hierarchical clustering of the underlying set of vectors with two key properties:

1. The  $h^{\text{th}}$  level of the clustering is a partition of the space with at most  $2^{2^h}$  parts.
2. For any  $t \in T$  and any  $h$ , let  $\mathcal{A}_h(t)$  be the piece of the partition in which  $t$  lies. Then, for every  $t \in T$ , we have that  $\sum_{h \geq 0} 2^{h/2} \cdot \text{diam}(\mathcal{A}_h(t)) = O(1)$ .

Note that the above implies that for every  $t$ , as  $h \rightarrow \infty$ , we have  $\text{diam}(\mathcal{A}_h(t)) = 2^{-h/2} \cdot o_h(1)$ . However, the rate of convergence of the sequence  $\text{diam}(\mathcal{A}_h(t)) \cdot 2^{h/2}$  can depend on  $t$ . If the convergence were uniform (i.e. independent of  $t$ ), then it is easy to show the following: Consider

---

<sup>2</sup>We note that by the Sudakov “minoration principle” [[Sud69](#)] (alternatively, see Section 7.4 of [[Ver18](#)]) and the fact that  $\mathbf{E}[\sup_{t \in T} \mathbf{X}_t] = 1$ , it follows that there is a  $\delta$ -cover of size  $2^{O(1/\delta^2)}$ .

the partitioning defined by the  $h^{\text{th}}$  level where  $h = 1/\varepsilon$ ; such a partitioning would have  $M = 2^{2^h}$  parts. Furthermore, by the uniform convergence assumption, the diameter of each part would be bounded by  $\delta \leq (1/h) \cdot 2^{-h/2}$ . Thus, the total error would be  $O(\delta\sqrt{\log M}) = O(\varepsilon)$ . So in this idealized setting of uniform convergence we would be able to achieve a total error of  $\varepsilon$  using  $2^{2^{O(1/\varepsilon)}}$  clusters.

While this idealized setting is not always achievable, we can still use the majorizing measures theorem to achieve the same asymptotic guarantee. In particular, we use the hierarchical clustering given by the majorizing measures theorem to obtain a partition of  $T$  with  $2^{2^{O(1/\varepsilon)}}$  parts such that the total error of our approximator is  $O(\varepsilon)$ ; see [Claim 22](#) and the paragraph preceding it for an explicit description of the clustering procedure. This completes an overview of the technical ideas behind our main result, [Theorem 1](#).

The Gaussian process guaranteed to exist by [Theorem 1](#) is non-centered; by simulating each of the constants  $c_s$  in [Theorem 1](#) using auxiliary Gaussian random variables, we show in [Corollary 23](#) how to obtain a centered Gaussian process that approximates the supremum of the original Gaussian process.

We now briefly describe the technical ingredients in two other sparsification results in this paper, namely sparsification of norms ([Theorem 2](#)) and of bounded width polytopes ([Theorem 3](#)).

**A Junta Theorem for Norms over Gaussian Space.** Recall that [Theorem 2](#) informally says that every norm over Gaussian space is essentially a junta. We obtain [Theorem 2](#) as a consequence of a multiplicative sparsification lemma for the supremum of a Gaussian process on a *symmetric* set ([Corollary 24](#)).<sup>3</sup>

Let  $\nu(\cdot)$  be any norm over  $\mathbb{R}^n$ ; thanks to homogeneity of norms, we may assume that  $\mathbf{E}[\nu(\mathbf{g})] = 1$  for  $\mathbf{g} \sim N(0, I_n)$ . Since  $\nu$  is a norm, there exists a symmetric set  $T \subseteq \mathbb{R}^n$  such that

$$\nu(x) = \sup_{t \in T} x \cdot t.$$

In particular, the set  $T$  corresponds to the unit ball of the *dual norm*  $\nu^\circ$  [[Tko18](#)]. Consequently, our earlier assumption can be restated as

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [\nu(\mathbf{g})] = \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \sup_{t \in T} t \cdot \mathbf{g} \right] = 1.$$

Building on [Theorem 1](#) and [Corollary 23](#), we show the existence of a symmetric set  $S \subseteq \mathbb{R}^n$  with

$$|S| \leq \left( \frac{1}{\varepsilon} \right)^{\exp\left(O\left(\frac{1}{\varepsilon^3}\right)\right)}$$

such that

$$\Pr \left[ \left| \sup_{t \in T} t \cdot \mathbf{g} - \sup_{s \in S} s \cdot \mathbf{g} \right| \leq \varepsilon \right] \geq 1 - \varepsilon.$$

An anti-concentration lemma for the supremum of a symmetric Gaussian process ([Lemma 26](#))—which we prove using the celebrated  $S$ -inequality of Latała and Oleszkiewicz [[LO99](#)]<sup>4</sup>—allows us to convert this additive bound into a multiplicative guarantee, yielding [Corollary 24](#). Finally, as the set  $S$  is symmetric, the function  $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$  defined as  $\psi(x) = \sup_{s \in S} s \cdot x$  is a norm that depends on at most  $|S|$  directions, completing the proof.

<sup>3</sup>A set  $T \subseteq \mathbb{R}^n$  is *symmetric* if  $-t \in T$  whenever  $t \in T$ .

**Sparsifying Intersections of Narrow Halfspaces.** For the purposes of this discussion, assume that we have a convex set  $K$  which uniformly has “unit width,” i.e.

$$K = \bigcap_{t \in T} \{x \in \mathbb{R}^n : t \cdot x \leq 1\}$$

where each  $t$  is a unit vector in  $\mathbb{S}^{n-1}$ . Consider the Gaussian process  $\{\mathbf{X}_t\}_{t \in T}$  where  $\mathbf{X}_t := \mathbf{g} \cdot t$ , and note that  $K = \{x : f_T(x) \leq 1\}$  where  $f_T(x) := \sup_{t \in T} t \cdot x$ . Thanks to [Theorem 1](#), we know that there is a subset  $S \subset T$  (with  $|S| = 2^{2^{O(1/\varepsilon)}}$ ) and suitable constants  $\{c_s\}_{s \in S}$  such that

$$\mathbf{E} \left[ \left| \sup_{t \in T} t \cdot \mathbf{g} - \sup_{s \in S} \{s \cdot \mathbf{g} + c_s\} \right| \right] = \varepsilon \cdot \mathbf{E} \left[ \sup_{t \in T} t \cdot \mathbf{g} \right].$$

Thus, a natural way to define an approximator to the convex set  $K$  is to consider the polytope  $J$ , defined as

$$J = \{x \in \mathbb{R}^n : \text{for all } s \in S, s \cdot x + c_s \leq 1\}.$$

The only remaining step to show that  $J$  is close to  $K$  (in the sense that the Gaussian volume of their symmetric difference is small) is to argue that the function  $f_T(x)$  is anti-concentrated, which follows from a result of Chernozhukov et al. [[CCK14](#)].

While in general the set  $T$  need not be a subset of  $\mathbb{S}^{n-1}$  (i.e. all the vectors need not have unit norm), it is not too difficult to adapt the above argument to the more general setting; we defer details of this to [Section 4](#).

### 1.3 Discussion and Applications

We first provide some broader context by recalling similar-in-spirit results in the discrete setting of  $\{0, 1\}^n$ , and then discuss some of the qualitative aspects and applications of our Gaussian-space sparsifiers.

**Analogy with Sparsification of CNFs and Submodular Functions over  $\{0, 1\}^n$ .** Our sparsification result for convex sets of bounded geometric width ([Definition 19](#)), [Theorem 3](#), can be viewed as a Gaussian space analogue of recent results from Boolean function analysis on sparsifying *CNF formulas* of bounded width. Recall that a width- $w$  CNF formula is a conjunction of an arbitrary number of clauses (Boolean disjunctions), each of which contains at most  $w$  literals and hence “lops off” at least a  $2^{-w}$  fraction of all points in  $\{0, 1\}^n$ . This is analogous to how a convex set of geometric width  $k$  is an intersection of an arbitrary number of halfspaces in  $\mathbb{R}^n$ , each at distance at most  $k$  from the origin and hence “lopping off” at least an  $\approx \exp(-k^2/2)$  fraction of Gaussian space under  $N(0, I_n)$ .

In the Boolean context, the state-of-the art result of Lovett et al. [[LWZ21](#)] gives, for any initial width- $w$  CNF over  $\{0, 1\}^n$ , an  $\varepsilon$ -approximating width- $w$  CNF consisting of  $s = (2+w^{-1} \log(1/\varepsilon))^{O(w)}$  clauses—note that this size bound is completely independent of the dimension  $n$  and the number of clauses in the initial CNF formula. Our [Theorem 3](#) similarly gives a sparsifying polytope using a number of halfspaces that depends only on the error parameter  $\varepsilon$  and the geometric width of the original convex set, with no dependence on the ambient dimension  $n$  or the number of halfspaces in the original convex set. (Indeed, obtaining such a Gaussian-space analogue of the Boolean CNF sparsification result was the initial impetus for this work.)

We remark that our norm sparsification result, [Theorem 2](#), can also be viewed as analogous to known results [[FKV13](#), [FV16](#)] giving junta approximators for *submodular functions* over  $\{0, 1\}^n$ . (Recall that submodular functions are often viewed as being a discrete analogue of convex functions [[FKV13](#), [Lov82](#)] and that every norm over  $\mathbb{R}^n$  is a convex function.)

**Sparsification Lower Bounds.** In light of our positive results for sparsification, it is natural to ask about lower bounds: How close to optimal are the quantitative upper bounds that we establish? As mentioned above, there is an upper bound of  $(2 + w^{-1} \log(1/\varepsilon))^{O(w)}$  clauses for the analogous Boolean question of sparsifying width- $w$  CNFs over  $\{0, 1\}^n$ , and this is known to be best possible up to the hidden constant in the  $O(w)$  exponent [LWZ21]. So for constant width, in the Boolean setting the correct dependence on the error parameter  $\varepsilon$  is *poly-logarithmic*, while our upper bound given in [Theorem 3](#) is *doubly exponential*—a difference of three exponentials! This contrast naturally motivates the question of whether stronger quantitative results can be obtained in our setting.

In [Section 6.2](#) we show that the quantitative parameters of our sparsifiers are not too far from the best possible: roughly speaking, we give a  $2^{(1/\varepsilon)^{\Omega(1)}}$  lower bound, which is only one exponential away from our doubly-exponential upper bound. In particular, our lower bound shows that sparsifying constant-geometric-width polytopes over  $\mathbb{R}^n$  inherently requires a much worse  $\varepsilon$ -dependence than sparsification of constant-width CNF formulas over  $\{0, 1\}^n$ : the former requires  $2^{\Omega(1/\varepsilon)}$  halfspaces, while the latter can be achieved using only  $(\log(1/\varepsilon))^{O(1)}$  clauses.

**On Centered and Non-Centered Sparsifiers.** Recalling [Equation \(1\)](#), since  $S \subseteq T$  the sparsified Gaussian process  $\{\mathbf{X}_s + c_s\}_{s \in S}$  can be viewed as a non-centered (because of the  $c_s$ 's) sub-process of the original centered Gaussian process  $\{\mathbf{X}_t\}_{t \in T}$ . In [Corollary 23](#) we show that our main sparsification result, [Theorem 1](#), can be easily modified to give a sparsifier  $S$  of essentially the same size which is a centered process; however, this sparsifier  $S$  is not a sub-process of the original Gaussian process (i.e.  $S$  is no longer a subset of  $T$ ). This tradeoff is unavoidable: In [Section 6.1](#) we give a simple example showing that any sparsifier whose size is independent of the ambient dimension cannot both be a sub-process and be centered (and as we explain in [Section 6.1](#), analogous tradeoffs are similarly unavoidable for [Theorem 2](#) and [Theorem 3](#)).

**Algorithmic Applications.** The class of convex sets which can be expressed as intersections of  $k$  halfspaces has been extensively studied in Boolean function analysis, in particular from the vantage point of learning theory and property testing. It is known that over the Gaussian space, this class is both (a) tolerantly testable with query complexity independent of the ambient dimension  $n$  [DMN21], and (b) agnostically learnable with a quasipolynomial dependence on  $k$  and a polynomial dependence on  $n$  [KOS08, DKK<sup>+</sup>24]. [Theorem 3](#), which shows that bounded-width polytopes can be approximated by an intersection of  $k$  halfspaces where  $k$  depends only on the width and target error, lets us obtain qualitatively similar tolerant testing and agnostic learning results for bounded width polytopes over the Gaussian space. This is described in more detail in [Section 5](#).

## 2 Preliminaries

We use boldfaced letters such as  $\mathbf{x}, \mathbf{X}$ , etc. to denote random variables (which may be real- or vector-valued; the intended type will be clear from the context). We write  $\mathbf{x} \sim \mathcal{D}$  to indicate that the random variable  $\mathbf{x}$  is distributed according to probability distribution  $\mathcal{D}$ .

We will frequently identify a set  $K \subseteq \mathbb{R}^n$  with its 0/1-valued indicator function. Given  $t \in \mathbb{R}$  and a set  $K \subseteq \mathbb{R}^n$ , we write  $tK$  for the  $t$ -dilation of  $K$ , i.e.  $tK := \{tx : x \in K\}$ . Unless explicitly stated otherwise,  $\|\cdot\|$  denotes the  $\ell_2$ -norm. We write  $\mathbb{S}^{n-1}$  to denote the unit  $\ell_2$ -sphere in  $\mathbb{R}^n$ , i.e.

$$\mathbb{S}^{n-1} := \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

Throughout,  $\{e_i\}_{i \in [n]}$  will denote the collection of standard basis vectors in  $\mathbb{R}^n$ . We will write  $\log$  to denote logarithm base two and write  $\ln$  to denote the natural logarithm  $\log_e$ . Finally,  $[n] := \{1, \dots, n\}$ .

## 2.1 Gaussian Random Variables

Identifying  $0 \equiv 0^n$  and writing  $I_n$  for the  $n \times n$  identity matrix,  $N(0, I_n)$  will denote the  $n$ -dimensional standard Gaussian distribution. We write  $\text{Vol}(K)$  to denote the Gaussian measure of a (Lebesgue measurable) set  $K \subseteq \mathbb{R}^n$ , i.e.

$$\text{Vol}(K) := \Pr_{\mathbf{g} \sim N(0, I_n)} [\mathbf{g} \in K].$$

We recall the following standard tail bound on Gaussian random variables:

**Proposition 4** (Proposition 2.1.2 of [Ver18]). Suppose  $\mathbf{g} \sim N(0, 1)$  is a one-dimensional Gaussian random variable. Then for all  $r > 0$ ,

$$\left(\frac{1}{r} - \frac{1}{r^3}\right) \cdot \varphi(r) \leq \Pr_{\mathbf{g} \sim N(0, 1)} [\mathbf{g} \geq r] \leq \frac{1}{r} \cdot \varphi(r),$$

where  $\varphi$  is the one-dimensional standard Gaussian density, which is given by

$$\varphi(x) := \frac{1}{\sqrt{2\pi}} e^{-x^2/2}.$$

We will use the following well-known bounds on the maximum of i.i.d. Gaussians:

**Proposition 5.** Suppose  $\mathbf{g} \sim N(0, I_n)$ . Then

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \max_{i \in [n]} \mathbf{g}_i \right] = \Theta(\sqrt{\ln n}) \quad \text{and} \quad \mathbf{Var}_{\mathbf{g} \sim N(0, I_n)} \left[ \max_{i \in [n]} \mathbf{g}_i \right] = \Theta\left(\frac{1}{\ln n}\right). \quad (5)$$

In particular for  $n$  large enough,

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \max_{i \in [n]} \mathbf{g}_i \right] \in \left[ \sqrt{2 \ln n} \left(1 - \frac{4}{\ln n}\right), \sqrt{2 \ln n} \left(1 + \frac{4}{\ln n}\right) \right]. \quad (6)$$

Furthermore, we have

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \max_{i \in [n]} |\mathbf{g}_i| \right] = \Theta(\sqrt{\ln n}) \quad \text{and} \quad \mathbf{Var}_{\mathbf{g} \sim N(0, I_n)} \left[ \max_{i \in [n]} |\mathbf{g}_i| \right] = O\left(\frac{1}{\ln n}\right). \quad (7)$$

See Appendix A.2 of [Cha14] for the bound on expectation in Equation (5) and Chapter 5.2 of [Cha14] for the bound on the variance. See Example 10.5.3 of [DN04] (or alternatively [Sas11]) for Equation (6). See Exercise 2.11 of [Wai19] for the bound on the expectation in Equation (7); the bound on the variance follows from Chapter 5.2 of [Cha14].

For  $L > 0$ , recall that a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is  $L$ -Lipschitz if  $|f(x) - f(y)| \leq L \cdot \|x - y\|$ . We will require the following concentration inequality for Lipschitz functions of Gaussian random variables:

**Proposition 6** (Theorem 5.2.2 of [Ver18]). Suppose  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is an  $L$ -Lipschitz function. Then

$$\Pr_{\mathbf{g} \sim N(0, I_n)} [ |f(\mathbf{g}) - \mathbf{E}[f]| \geq t ] \leq 2 \exp\left(-\frac{t^2}{2L^2}\right).$$



## 2.2 Gaussian Processes and Suprema

Recall that a *random process* is a collection of random variables  $\{\mathbf{X}_t\}_{t \in T}$  on the same probability space which are indexed by elements  $t$  of some set  $T$ .

**Definition 7** (Gaussian process). A random process  $\{\mathbf{X}_t\}_{t \in T}$  is called a *Gaussian process* if, for any finite subset  $T_0 \subseteq T$ , the random vector  $\{\mathbf{X}_t\}_{t \in T_0}$  has a multivariate Gaussian distribution. We say  $\{\mathbf{X}_t\}_{t \in T}$  is *centered* if  $\mathbf{E}[\mathbf{X}_t] = 0$  for all  $t \in T$ .

We assume throughout that all Gaussian processes are *separable*; we will not dwell on this technical point and refer the reader to Section 2.2 of [LT91] for a detailed discussion.

**Definition 8** (Canonical Gaussian process). Given  $T \subseteq \mathbb{R}^n$ , the centered random process  $\{\mathbf{X}_t\}_{t \in T}$  defined as

$$\mathbf{X}_t := \mathbf{g} \cdot t \quad \text{where } \mathbf{g} \sim N(0, I_n)$$

is a Gaussian process that we refer to as the *canonical Gaussian process on  $T$* .

It is a standard fact that any centered Gaussian process can be realized as a canonical process (see Section 7.1.2 of [Ver18]) and consequently we will focus on canonical Gaussian processes throughout this paper. The following notation will be convenient:

**Definition 9.** Given a set  $T \subseteq \mathbb{R}^n$ , we define

$$f_T(x) := \sup_{t \in T} x \cdot t.$$

We remark that if  $T$  is a closed convex set then  $f_T$  is known as the “support function” of  $T$  (though we will be interested in arbitrary bounded subsets  $T$  of  $\mathbb{R}^n$ , which need not be convex sets). Note that when  $\mathbf{g} \sim N(0, I_n)$ ,  $f_T(\mathbf{g}) = \sup_{t \in T} \mathbf{X}_t$  where  $\{\mathbf{X}_t\}_{t \in T}$  is the canonical Gaussian process on  $T$ . We will frequently rely on the fact that the supremum of a Gaussian process has subgaussian tails, which we recall below.

**Proposition 10** (Appendix A.5 of [Cha14]). Let  $\{\mathbf{X}_t\}_{t \in T}$  be a centered Gaussian process indexed by a set  $T \subseteq \mathbb{R}^n$  and let  $r \geq 0$ . Then

$$\Pr \left[ \left| \sup_{t \in T} \mathbf{X}_t - \mathbf{E} \left[ \sup_{t \in T} \mathbf{X}_t \right] \right| \geq r \right] \leq 2 \exp \left( \frac{-r^2}{2 \sup_{t \in T} \mathbf{Var} [\mathbf{X}_t]} \right).$$

For a canonical Gaussian process  $\{\mathbf{X}_t\}_{t \in T}$  as in **Proposition 10**, we have that  $\sup_{t \in T} \mathbf{Var} [\mathbf{X}_t] = \sup_{t \in T} \|t\|^2$ , and hence  $\sup_{t \in T} \mathbf{Var} [\mathbf{X}_t] \geq \text{diam}(T)^2/4$ . Finally, the magnitude of the canonical Gaussian process on  $T$  is captured by the following important geometric quantity:

**Definition 11** (Gaussian width). The *Gaussian width* of a set  $T \subseteq \mathbb{R}^n$  is defined as

$$w(T) := \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [f_T(\mathbf{g})] = \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \sup_{t \in T} \mathbf{g} \cdot t \right].$$

Gaussian width was originally introduced in geometric functional analysis and asymptotic convex geometry [AAGM15, AAGM21]. It is easy to see that  $w(T)$  is finite if and only if  $T$  is bounded; we refer the reader to Chapter 7 of [Ver18] for further information.

### 2.2.1 Anti-Concentration Bounds

We will require the following anti-concentration bound on  $f_T(\mathbf{g})$  obtained by [CCK14] (which is an easy consequence of an analogous statement for Gaussian random vectors obtained in [CCK15]):

**Theorem 12** (Theorem 2.1 of [CCK14]). Suppose  $T \subseteq \mathbb{S}^{n-1}$ . Let  $\{\mathbf{X}_t\}_{t \in T}$  be the canonical Gaussian process on  $T$ . For every  $\theta \in \mathbb{R}$  and  $\varepsilon > 0$ , we have

$$\Pr_{\mathbf{g} \sim N(0, I_n)} [ |f_T(\mathbf{g}) - \theta| \leq \varepsilon ] \leq 4\varepsilon(1 + w(T)) \quad (8)$$

where  $w(T)$  is as in Definition 11.

### 2.2.2 Talagrand's Majorizing Measures Theorem

We refer the reader to [vH16, Ver18, Tal22] for further background on the majorizing measures theorem as well as applications thereof.

**Definition 13** (Admissible sequence). Let  $T \subseteq \mathbb{R}^n$ . An *admissible sequence*  $\{\mathcal{A}_h\}_{h \in \mathbb{N}}$  of partitions of  $T$  is defined to be a collection of partitions with the following properties:

- (i)  $\mathcal{A}_0 = \{T\}$ ,
- (ii)  $\mathcal{A}_{h+1}$  is a refinement of  $\mathcal{A}_h$  for all  $h$ , and
- (iii)  $|\mathcal{A}_h| \leq 2^{2^h}$  for all  $h \in \mathbb{N}$ .

**Definition 14** (Talagrand's  $\gamma_2$  functional). Let  $T \subseteq \mathbb{R}^n$ . Given  $t \in T$  and a partition  $\mathcal{P}$  of  $T$ , we write  $\mathcal{P}(t)$  for the part of  $\mathcal{P}$  containing  $t$ . We define

$$\gamma_2(T) = \inf_{\text{admissible } \mathcal{A}} \sup_{t \in T} \sum_{h \geq 0} 2^{h/2} \cdot \text{diam}(\mathcal{A}_h(t))$$

where  $\text{diam}(S)$  denotes the diameter of the set  $S$ , i.e.  $\text{diam}(S) = \sup_{s_1, s_2 \in S} \|s_1 - s_2\|_2$ .

The *majorizing measures theorem*, with upper bound due to Fernique [Fer75] and lower bound due to Talagrand [Tal87], is the following:

**Theorem 15** (Theorem 2.10.1 of [Tal22]). Let  $T \subseteq \mathbb{R}^n$  and let  $\{\mathbf{X}_t\}_{t \in T}$  be a centered Gaussian process. Then there exists a universal constant  $L$  such that

$$\frac{1}{L} \cdot \gamma_2(T) \leq \mathbf{E} \left[ \sup_{t \in T} \mathbf{X}_t \right] \leq L \cdot \gamma_2(T).$$

### 2.3 Polyhedral Approximation Under the Gaussian Distance

We will be interested in approximating subsets of  $\mathbb{R}^n$  under the Gaussian distance:

**Definition 16.** Given two measurable sets  $K, L \subseteq \mathbb{R}^n$ , we define the *Gaussian distance between  $K$  and  $L$*  to be

$$\text{dist}_G(K, L) := \Pr_{\mathbf{x} \sim N(0, I_n)} [K(\mathbf{x}) \neq L(\mathbf{x})].$$

In other words,  $\text{dist}_G(K, L) = \text{Vol}(K \triangle L)$ , i.e. the Gaussian measure of the symmetric difference of the sets  $K$  and  $L$ .

Recall that every convex set can be written as an intersection of (possibly infinitely many) halfspaces. The following upper bound on the number of halfspaces needed to approximate a polytope under the Gaussian distance was obtained in [DNS24]:

**Theorem 17** (Theorem 19 of [DNS24]). Given a convex set  $K \subseteq \mathbb{R}^n$  and  $\varepsilon \in (0, 10^{-3})$ , there exists a set  $L \subseteq \mathbb{R}^n$  which is an intersection of  $(n/\varepsilon)^{O(n)}$  halfspaces such that  $\text{dist}_G(K, L) \leq \varepsilon$ .

[DNS24] also gave a mildly-exponential lower bound on the number of halfspaces needed to approximate the  $\ell_2$ -ball of radius  $\sqrt{n}$  to constant accuracy. The arguments of [DNS24] give the following lower bound:

**Theorem 18** (Theorem 63 of [DNS24]). Let  $B \subseteq \mathbb{R}^n$  be an origin-centered  $\ell_2$  ball with radius  $r \in [\sqrt{n} - 1, \sqrt{n} + 1]$ , so  $\text{Vol}(B) \in [-\tau, 1 - \tau]$  for an absolute constant  $\tau > 0$ . There exists an absolute constant  $\kappa \in (0, 0.5)$  such that any intersection of halfspaces  $L \subseteq \mathbb{R}^n$  with  $\text{dist}_G(B, L) \leq \kappa$  must have at least  $2^{\Omega(\sqrt{n})}$  halfspaces.

We note that the lower bound in **Theorem 18** is tight up to constant factors hidden by the  $\Omega(\cdot)$ -notation; we refer the reader to Section 5 of [DNS24] for a probabilistic construction of an intersection of  $2^{O(\sqrt{n})}$  halfspaces that approximates the  $\ell_2$ -ball of radius  $\sqrt{n}$  to constant accuracy.

We will be interested in convex sets that have bounded *geometric width*, which we define as follows:

**Definition 19** (Geometric width of a convex set). A convex set  $K \subseteq \mathbb{R}^n$  is said to have *geometric width at most  $r$*  if  $K$  can be expressed as an intersection of (possibly infinitely many) halfspaces, each of which is of the form  $\mathbf{1}\{u \cdot x \leq r'\}$  where  $\|u\| = 1$  and  $r' \leq r$ .

Thus the geometric width of a single halfspace  $H = \mathbf{1}\{u \cdot x \leq r\}$  where  $\|u\| = 1$  is  $r$ , and the geometric width of an origin-centered  $\ell_2$ -ball of radius  $r$  is  $r$ .

### 3 Sparsifying Gaussian Processes with Bounded Expected Suprema

In this section, which is at the heart of our technical contribution, we show that if a set  $T$  is bounded (equivalently, has bounded Gaussian width) then the supremum of the canonical Gaussian processes on  $T$  can be approximated by the supremum of a sparse or “low-dimensional” (non-centered) Gaussian process.

**Theorem 20** (Sparsifying suprema of Gaussian processes). Let  $\varepsilon \in (0, 0.5)$ . For any  $T \subseteq \mathbb{R}^n$  with  $0 < w(T) < \infty$ , there exists a set  $S \subseteq T$  with

$$|S| = 2^{2^{O\left(\frac{w(T)}{\varepsilon}\right)}}$$

and constants  $\{c_s\}_{s \in S}$  satisfying  $0 \leq c_s \leq w(T)$  such that

$$\mathbf{E}_{g \sim N(0, I_n)} \left[ \left| f_T(g) - \sup_{s \in S} \{g \cdot s + c_s\} \right| \right] \leq \varepsilon.$$

Note that **Theorem 1** is easily obtained from **Theorem 20** by rescaling the Gaussian process  $\{\mathbf{X}_t\}_{t \in T}$  by a factor of  $w(T)$  so that the resulting process has Gaussian width 1. Towards **Theorem 20**, we require the following technical lemma:

**Lemma 21.** Let  $\mathcal{P}$  be a partition of  $T$  into finitely many pieces. Define  $S \subseteq T$  as

$$S := \{s_P : P \in \mathcal{P}\}$$

where each  $s_P$  is an arbitrary (but fixed) representative element of the part  $P$ . Define

$$c_P := \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \sup_{t \in P} \mathbf{g} \cdot (t - s_P) \right].$$

Then for any  $\delta \geq 0.5 \cdot \max_{P \in \mathcal{P}} \text{diam}(P)$  we have

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \left| f_T(\mathbf{g}) - \sup_{P \in \mathcal{P}} \{ \mathbf{g} \cdot s_P + c_P \} \right| \right] \leq \delta \left( 1 + \sum_{P \in \mathcal{P}} \exp \left( \frac{-2\delta^2}{\text{diam}(P)^2} \right) \right).$$

*Proof.* For any  $g \in \mathbb{R}^n$ , we have

$$\begin{aligned} \left| f_T(g) - \sup_{P \in \mathcal{P}} \{g \cdot s_P + c_P\} \right| &= \left| \sup_{t \in T} g \cdot t - \sup_{P \in \mathcal{P}} \{g \cdot s_P + c_P\} \right| \\ &= \left| \sup_{P \in \mathcal{P}} \sup_{t \in P} g \cdot t - \sup_{P \in \mathcal{P}} \{g \cdot s_P + c_P\} \right| \\ &\leq \sup_{P \in \mathcal{P}} \left| \sup_{t \in P} g \cdot t - (g \cdot s_P + c_P) \right| \\ &= \sup_{P \in \mathcal{P}} \left| \sup_{t \in P} \{g \cdot (t - s_P) - c_P\} \right|, \end{aligned}$$

where the inequality is using convexity of the sup function and Jensen's inequality. Motivated by this, for each  $P \in \mathcal{P}$  let  $\{\mathbf{Y}_t^{(P)}\}_{t \in P}$  be the following centered Gaussian process:

$$\mathbf{Y}_t^{(P)} := \mathbf{g} \cdot (t - s_P) \quad \text{where } \mathbf{g} \sim N(0, I_n).$$

Note that

$$\mathbf{E} \left[ \sup_{t \in P} \mathbf{Y}_t^{(P)} \right] = c_P.$$

We thus have

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \left| f_T(\mathbf{g}) - \sup_{P \in \mathcal{P}} \{ \mathbf{g} \cdot s_P + c_P \} \right| \right] \leq \mathbf{E} \left[ \sup_{P \in \mathcal{P}} \left| \sup_{t \in P} \mathbf{Y}_t^{(P)} - \mathbf{E} \left[ \sup_{t \in P} \mathbf{Y}_t^{(P)} \right] \right| \right]. \quad (9)$$

In particular, we can write

$$\begin{aligned}
\text{R.H.S. of Equation (9)} &= \int_{r=0}^{\infty} \Pr \left[ \sup_{P \in \mathcal{P}} \left| \sup_{t \in P} \mathbf{Y}_t^{(P)} - \mathbf{E} \left[ \sup_{t \in P} \mathbf{Y}_t^{(P)} \right] \right| \geq r \right] dr \\
&\leq \delta + \int_{r=\delta}^{\infty} \Pr \left[ \sup_{P \in \mathcal{P}} \left| \sup_{t \in P} \mathbf{Y}_t^{(P)} - \mathbf{E} \left[ \sup_{t \in P} \mathbf{Y}_t^{(P)} \right] \right| \geq r \right] dr \\
&\leq \delta + \int_{r=\delta}^{\infty} \sum_{P \in \mathcal{P}} \Pr \left[ \left| \sup_{t \in P} \mathbf{Y}_t^{(P)} - \mathbf{E} \left[ \sup_{t \in P} \mathbf{Y}_t^{(P)} \right] \right| \geq r \right] dr \tag{10}
\end{aligned}$$

$$= \delta + \sum_{P \in \mathcal{P}} \int_{r=\delta}^{\infty} \Pr \left[ \left| \sup_{t \in P} \mathbf{Y}_t^{(P)} - \mathbf{E} \left[ \sup_{t \in P} \mathbf{Y}_t^{(P)} \right] \right| \geq r \right] dr \tag{11}$$

$$\leq \delta + \sum_{P \in \mathcal{P}} \int_{r=\delta}^{\infty} 2 \exp \left( \frac{-2r^2}{\text{diam}(P)^2} \right) dr \tag{12}$$

where Equation (10) follows via a union bound, Equation (11) uses Fubini's theorem, and Equation (12) follows from Proposition 10.

A change of variables together with the Gaussian tail bound from Proposition 4 gives

$$\begin{aligned}
\int_{r=\delta}^{\infty} \exp \left( \frac{-2r^2}{\text{diam}(P)^2} \right) dr &= \sqrt{\frac{\pi}{2}} \cdot \text{diam}(P) \cdot \Pr \left[ N(0, 1) \geq \frac{2\delta}{\text{diam}(P)} \right] \\
&\leq \frac{\text{diam}(P)^2}{4\delta} \cdot \exp \left( \frac{-2\delta^2}{\text{diam}(P)^2} \right).
\end{aligned}$$

By the assumption that  $2\delta \geq \text{diam}(P)$  for all  $P \in \mathcal{P}$ , combining this with Equation (12) gives

$$\text{R.H.S. of Equation (9)} \leq \delta \left( 1 + \sum_{P \in \mathcal{P}} \exp \left( \frac{-2\delta^2}{\text{diam}(P)^2} \right) \right)$$

which completes the proof.  $\square$

With Lemma 21 in hand, Theorem 20 follows via the lower bound from Talagrand's majorizing measures theorem (Theorem 15):

*Proof of Theorem 20.* Let  $\{\mathbf{X}_t\}_{t \in T}$  be the canonical Gaussian process on  $T$ . By Theorem 15, we have that

$$\frac{1}{L} \cdot \gamma_2(T) \leq w(T)$$

for some universal constant  $L$ . In particular, there exists an admissible sequence (cf. Definition 13)  $\mathcal{A}$  such that

$$\sum_{h \geq 0} 2^{h/2} \cdot \text{diam}(\mathcal{A}_h(t)) \leq L \cdot w(T) \quad \text{for all } t \in T. \tag{13}$$

We will construct an appropriate partition of  $T$  from the admissible sequence  $\mathcal{A}$  and then appeal to [Lemma 21](#) to complete the proof. Let  $\delta$  be a parameter we will set shortly. Construct  $\mathcal{P}$  as follows: In stage  $h = 1, 2, 3, \dots$ , we place into  $\mathcal{P}$  all parts  $P$  from  $\mathcal{A}_h$  satisfying

$$2 \cdot \text{diam}(P) \leq \delta \cdot 2^{-h/2}, \quad (14)$$

provided  $P \not\subseteq \bigcup \mathcal{P}$  already.

**Claim 22.** This process terminates with a partition  $\mathcal{P}$  of  $T$  by stage  $h = 1 + \lfloor 2L \cdot w(T)/\delta \rfloor$ .

*Proof.* Suppose otherwise. Then there exists some  $t \in T$  with

$$2^{h/2} \cdot \text{diam}(\mathcal{A}_h(t)) > \frac{\delta}{2} \quad \text{for all } 1 \leq h \leq 1 + \lfloor 2L \cdot w(T)/\delta \rfloor,$$

which contradicts [Equation \(13\)](#). □

As an immediate consequence of [Claim 22](#) and Item (iii) of [Definition 13](#) it follows that

$$|\mathcal{P}| \leq 2^{2^{1+\lfloor 2L \cdot w(T)/\delta \rfloor}}. \quad (15)$$

Next, note that at stage  $h$  of the above process, we add at most  $2^{2^h}$  parts to  $\mathcal{P}$  and each part  $P$  satisfies [Equation \(14\)](#). Consequently,

$$\begin{aligned} \sum_{P \in \mathcal{P}} \exp\left(\frac{-2\delta^2}{\text{diam}(P)^2}\right) &\leq \sum_{P \in \mathcal{P}} \exp\left(\frac{-\delta^2}{2 \cdot \text{diam}(P)^2}\right) \leq \sum_{h=1}^{\infty} 2^{2^h} \cdot \exp(-2^{h+1}) \\ &= \sum_{h=1}^{\infty} \exp(\ln 2 \cdot 2^h - 2 \cdot 2^h) \\ &\leq \sum_{h=1}^{\infty} \exp(-2^h) \leq 1. \end{aligned} \quad (16)$$

Set  $\delta := \varepsilon/2$ ; it thus follows from [Lemma 21](#) that for appropriate sequences of vectors  $\{s_P\}_{P \in \mathcal{P}}$  and constants  $\{c_P\}_{P \in \mathcal{P}}$ ,

$$\begin{aligned} \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \left| f_T(\mathbf{g}) - \sup_{P \in \mathcal{P}} \{\mathbf{g} \cdot s_P + c_P\} \right| \right] &\leq \delta \left( 1 + \sum_{P \in \mathcal{P}} \exp\left(\frac{-2\delta^2}{\text{diam}(P)^2}\right) \right) \\ &\leq \varepsilon \end{aligned}$$

thanks to [Equation \(16\)](#) and our choice of  $\delta$ . (Note that [Equation \(14\)](#) immediately implies that  $\delta \geq \sqrt{2} \cdot \text{diam}(P)$  for all  $P \in \mathcal{P}$ , as required by [Lemma 21](#).) Furthermore, taking  $S = \{s_P\}_{P \in \mathcal{P}}$ , we have by [Equation \(15\)](#) that

$$|S| = |\mathcal{P}| \leq 2^{2^{O\left(\frac{w(T)}{\varepsilon}\right)}}$$

as desired, concluding the proof of [Theorem 20](#). □

### 3.1 Sparsifying Using a Centered Gaussian Process

Observe that the sparsified process  $\{\mathbf{X}_s + c_s\}_{s \in S}$  guaranteed by [Theorem 20](#) is a non-centered Gaussian process. It is not too difficult to obtain a sparsifier of  $\{\mathbf{X}_t\}_{t \in T}$  corresponding to a *centered* Gaussian process, although the vectors used by this sparsifier no longer lie in the original set  $T$ :

**Corollary 23** (Sparsifying using a centered Gaussian process). Let  $\varepsilon \in (0, 0.5)$ . For any  $T \subseteq \mathbb{R}^n$  with  $w(T) < \infty$ , there exists a set  $S \subset \mathbb{R}^{n+A}$  with

$$|S| = 2^{2^{O\left(\frac{w(T)}{\varepsilon}\right)}} \quad \text{and} \quad A = \exp\left(\Theta\left(\frac{w(T)}{\varepsilon}\right)\right)$$

such that (concatenating each vector  $t \in T$  with the  $A$ -dimensional vector  $0^A$ , which lets us view  $f_T$  as  $f_T : \mathbb{R}^{n+A} \rightarrow \mathbb{R}$ )

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_{n+A})} [|f_T(\mathbf{g}) - f_S(\mathbf{g})|] \leq \varepsilon. \quad (17)$$

Moreover, each  $s \in S$  is of the form  $s = (s', \alpha e_i)$  for some  $i \in [A]$  and  $s' \in T$ . Furthermore  $(s', -\alpha e_i)$  belongs to  $S$  whenever  $(s', \alpha e_i)$  belongs to  $S$ .

Looking ahead, we will rely on the symmetry of coordinates in  $A$  in the following subsection ([Section 3.2](#)).

*Proof.* From [Theorem 20](#), for the function  $f_T(\cdot)$ , there is a set  $S'$  and a suitable set of constants  $\{c_{s'}\}_{s' \in S'}$  with  $c_{s'} \geq 0$  such that

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \left| f_T(\mathbf{g}) - \sup_{s' \in S'} \{\mathbf{g} \cdot s' + c_{s'}\} \right| \right] \leq \frac{\varepsilon}{2}, \quad (18)$$

where  $|S'| = 2^{2^{O\left(\frac{w(T)}{\varepsilon}\right)}}$ . The idea is to now replace each constant  $c_{s'}$  by a suitable maximum of independent Gaussian random variables. In particular, let  $A$  be a sufficiently large number that we will set shortly and let

$$\mu_A := \mathbf{E}_{\mathbf{g} \sim N(0, I_A)} \left[ \sup_{i \in [A]} |g_i| \right], \quad \sigma_A^2 := \mathbf{Var}_{\mathbf{g} \sim N(0, I_A)} \left[ \sup_{i \in [A]} |g_i| \right].$$

For every  $s' \in S'$ , define the set  $\text{Aux}(s') \subseteq \mathbb{R}^{n+A}$  as follows:

$$\text{Aux}(s') := \bigcup_{j \in [A]} \left\{ \left( s', \frac{c_{s'} e_j}{\mu_A} \right), \left( s', \frac{-c_{s'} e_j}{\mu_A} \right) \right\}.$$

Finally, we define the set  $S \subseteq \mathbb{R}^{n+A}$  as

$$S := \bigcup_{s' \in S'} \text{Aux}(s').$$

Note that it suffices to show that

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_{n+A})} \left[ \left| \sup_{s' \in S'} \{\mathbf{g} \cdot s' + c_{s'}\} - \sup_{s \in S} \mathbf{g} \cdot s \right| \right] \leq \frac{\varepsilon}{2}. \quad (19)$$

Indeed, [Equations \(18\)](#) and [\(19\)](#) immediately imply [Equation \(17\)](#) from the statement of the theorem.

We establish [Equation \(19\)](#) in the remainder of the proof. As in the proof of [Lemma 21](#), note that for any  $g \in \mathbb{R}^{n+A}$  we have

$$\begin{aligned}
\left| \sup_{s' \in S'} \{g \cdot s' + c_{s'}\} - \sup_{s \in S} g \cdot s \right| &= \left| \sup_{s' \in S'} \{g \cdot s' + c_{s'}\} - \sup_{s' \in S'} \sup_{s \in \text{Aux}(s')} g \cdot s \right| \\
&\leq \sup_{s' \in S'} \left| \{g \cdot s' + c_{s'}\} - \sup_{s \in \text{Aux}(s')} g \cdot s \right| \\
&= \sup_{s' \in S'} \left| c_{s'} - \sup_{j \in [A]} \frac{c_{s'} |g_j|}{\mu_A} \right| \\
&= \sup_{s' \in S'} \frac{c_{s'}}{\mu_A} \left| \mu_A - \sup_{j \in [A]} |g_j| \right| \\
&\leq \frac{w(T)}{\mu_A} \cdot \left| \mu_A - \sup_{j \in [A]} |g_j| \right|,
\end{aligned}$$

where we relied on the definition of  $\text{Aux}(s')$  in the second equality and on the fact that  $c_{s'} \leq w(T)$  in the final inequality. (The latter is readily verified from the proof of [Lemma 21](#).) Taking expectations with respect to  $\mathbf{g} \sim N(0, I_{n+A})$  then gives

$$\text{L.H.S. of (19)} \leq \frac{w(T)}{\mu_A} \mathbf{E}_{\mathbf{g} \sim N(0, I_A)} \left[ \left| \mu_A - \sup_{j \in [A]} |g_j| \right| \right] \leq \frac{w(T) \sigma_A}{\mu_A} = \Theta\left(\frac{w(T)}{\log A}\right)$$

where penultimate inequality relies on Jensen's inequality (more specifically, on the fact that  $\mathbf{E}[\mathbf{X}] \leq \mathbf{E}[\mathbf{X}^2]^{1/2}$  for a random variable  $\mathbf{X}$ ) and the final equality follows from [Proposition 5](#). In particular, taking

$$A = \exp\left(\Theta\left(\frac{w(T)}{\varepsilon}\right)\right)$$

immediately implies [Equation \(19\)](#), completing the proof.  $\square$

### 3.2 Multiplicative Approximation for Symmetric Processes

We will establish the following consequence of [Theorem 20](#):

**Corollary 24.** Suppose  $T \subseteq \mathbb{R}^n$  is symmetric (i.e.  $t \in T$  whenever  $-t \in T$ ) with  $w(T) < \infty$ . For  $\varepsilon \in (0, 0.5)$ , there exists another symmetric set  $S \subseteq \mathbb{R}^n$  with  $\dim(\text{span}(S)) \leq c_\varepsilon$ , where

$$c_\varepsilon := 2^{\exp\left(O\left(\frac{1}{\varepsilon^3}\right)\right)},$$

such that

$$\Pr_{\mathbf{g} \sim N(0, I_n)} \left[ f_T(\mathbf{g}) \in [(1 - \varepsilon)f_S(\mathbf{g}), (1 + \varepsilon)f_S(\mathbf{g})] \right] \geq 1 - \varepsilon.$$

In other words, we show that a centered Gaussian process on a symmetric index set can be multiplicatively approximated by a ‘‘low-dimensional’’ symmetric process. Furthermore, for  $n > c_\varepsilon$ , we will in fact have

$$|S| \leq 2^{\exp\left(O\left(\frac{1}{\varepsilon^3}\right)\right)}.$$



This will be immediate from the proof of [Corollary 24](#). (Note that when  $n \leq c_\varepsilon$ , [Corollary 24](#) holds trivially with  $T = S$ .)

**Remark 25.** Note that [Theorem 2](#) follows immediately from [Corollary 24](#). To see this, note that any norm  $\nu : \mathbb{R}^n \rightarrow \mathbb{R}$  can be written as  $\nu(x) = f_T(x)$  where  $T \subseteq \mathbb{R}^n$  is the symmetric convex set corresponding to the unit ball of the dual norm  $\nu^\circ$  (cf. [\[Tko18\]](#)). The function  $f_S : \mathbb{R}^n \rightarrow \mathbb{R}$  is then the approximating norm  $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$  from the statement of [Theorem 2](#).

### 3.2.1 Anti-Concentration of the Supremum for Symmetric Processes

Towards [Corollary 24](#), we will require the following anti-concentration lemma:

**Lemma 26.** Suppose  $T \subseteq \mathbb{R}^n$  is symmetric (i.e.  $t \in T$  whenever  $-t \in T$ ), and let  $\varepsilon \in (0, 0.5)$ . Then

$$\Pr_{\mathbf{g} \sim N(0, I_n)} [ |f_T(\mathbf{g})| \leq \varepsilon \cdot w(T) ] \leq 10\varepsilon.$$

Note that unlike the anti-concentration inequality due to Chernozhukov et al. [\[CCK14\]](#) (see [Theorem 12](#)), the above lemma does not require the index set  $T$  to be a subset of  $\mathbb{S}^{n-1}$ . On the other hand, [Theorem 12](#) guarantees anti-concentration of  $f_T(\cdot)$  everywhere and not just around 0. The proof of [Lemma 26](#) will rely on the well-known “ $S$ -inequality” due to Latała and Oleszkiewicz [\[LO99\]](#):

**Theorem 27** ( $S$ -inequality [\[LO99\]](#)). Let  $K \subseteq \mathbb{R}^n$  be a symmetric convex set, and suppose  $S \subseteq \mathbb{R}^n$  is a symmetric slab with  $\text{Vol}(S) = \text{Vol}(K)$ , i.e.

$$S = \{x \in \mathbb{R}^n : |x_1| \leq \theta\},$$

where  $\theta$  is chosen so as to ensure  $\text{Vol}(S) = \text{Vol}(K)$ . Then for any  $t \in [0, 1]$ , we have

$$\text{Vol}(tS) \leq \text{Vol}(tK).$$

*Proof of [Lemma 26](#).* Without loss of generality, we can rescale  $T$  so that  $\|t\| \leq 1$  for all  $t \in T$  and furthermore there exists  $t_0 \in T$  such that  $\|t_0\| = 1$ . Thanks to symmetry of  $T$ , we have

$$f_T(x) \geq |t_0 \cdot x| \quad \text{and so} \quad \text{med}(f_T(\mathbf{g})) \geq \text{med}(|t_0 \cdot \mathbf{g}|) \text{ for } \mathbf{g} \sim N(0, I_n),$$

where  $\text{med}(\mathbf{X})$  denotes the median of the random variable  $\mathbf{X}$ . As  $\|t_0\| = 1$ , we have  $|t_0 \cdot \mathbf{g}|$  is distributed according to a half-normal distribution and so

$$\text{med}(f_T) := \text{med}(f_T(\mathbf{g})) \geq 0.67 \tag{20}$$

where  $\mathbf{g} \sim N(0, I_n)$ .

Thanks to our rescaling of  $T$  and the fact that  $T$  is symmetric, it is easy to check that for  $x, y \in \mathbb{R}^n$ ,

$$|f_T(x) - f_T(y)| \leq f_T(x - y) = \sup_{t \in T} (x - y) \cdot t \leq \sup_{\|t\| \leq 1} (x - y) \cdot t = \|x - y\|,$$

and so the function  $f_T : \mathbb{R}^n \rightarrow \mathbb{R}$  is 1-Lipschitz. [Proposition 6](#) then readily implies that

$$w(T) = \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [f(\mathbf{g})] \leq \text{med}(f_T) + 4.$$

(See, for example, Problem 4.2(d) of [vH16].) Combining this with Equation (20), we get that  $w(T) \leq 10 \cdot \text{med}(f_T)$ . This lets us write

$$\begin{aligned} \Pr_{\mathbf{g} \sim N(0, I_n)} [ |f_T(\mathbf{g})| \leq \varepsilon \cdot w(T) ] &\leq \Pr_{\mathbf{g} \sim N(0, I_n)} [ |f_T(\mathbf{g})| \leq 10\varepsilon \cdot \text{med}(f_T) ] \\ &= \text{Vol} \left( 10\varepsilon \cdot \{x : f_T(x) \leq \text{med}(f_T)\} \right) \\ &\leq \text{Vol}(10\varepsilon \cdot \{x : |x_1| \leq c\}) \text{ for } c \text{ satisfying } \Phi(c) = 3/4 \\ &\leq 10\varepsilon, \end{aligned}$$

where the penultimate inequality relied on the  $S$ -inequality (Theorem 27). (We relied on the symmetry of the set  $T$  to conclude that the set  $\{x : f_T(x) \leq \text{med}(f_T)\}$  is a symmetric convex set with Gaussian volume  $1/2$ .)  $\square$

### 3.2.2 Proof of Corollary 24

If  $n \leq c_\varepsilon$ , then the result holds trivially with  $T = S$ . So we will assume for the remainder of the proof that  $n > c_\varepsilon$ .

**Step 0: Additive Approximation Suffices.** By scaling, we can assume that  $w(T) = 1$ . We will show that there exists a symmetric set  $S \subseteq \mathbb{R}^n$  such that

$$|S| \leq 2^{\exp\left(O\left(\frac{1}{\varepsilon^3}\right)\right)} \quad \text{and} \quad \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [ |f_T(\mathbf{g}) - f_S(\mathbf{g})| ] \leq \frac{\varepsilon^3}{40}. \quad (21)$$

To see why this suffices, note that by Markov's inequality,

$$\Pr_{\mathbf{g} \sim N(0, I_n)} \left[ |f_T(\mathbf{g}) - f_S(\mathbf{g})| \geq \frac{\varepsilon^2}{20} \right] \leq \frac{\varepsilon}{2}.$$

Also note that Lemma 26 implies

$$\Pr_{\mathbf{g} \sim N(0, I_n)} \left[ |f_T(\mathbf{g})| \geq \frac{\varepsilon}{20} \right] \geq 1 - \frac{\varepsilon}{2}.$$

Corollary 24 then immediately follows. The rest of the proof will establish Equation (21).

**Step 1: Non-Centered Symmetric Approximation.** Theorem 20 guarantees the existence of a set  $T' \subseteq T$  with

$$|T'| \leq 2^{\exp\left(O\left(\frac{1}{\varepsilon^3}\right)\right)}$$

and constants  $c_v$  for  $v \in T'$  satisfying  $0 < c_v \leq 1$  such that

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [ |h(\mathbf{g}) - f_T(\mathbf{g})| ] \leq \frac{\varepsilon^3}{160} \quad \text{where } h(x) := \sup_{v \in T'} v \cdot x + c_v.$$

Inspecting the proof of Theorem 20, it is easy to check that applying it to  $-T = T$  returns the collection of vectors  $-T'$  and associated constants  $c'_v$  for  $v \in -T'$  where  $c'_v = c_{-v}$  and satisfying the following:

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [ |h'(\mathbf{g}) - f_T(\mathbf{g})| ] \leq \frac{\varepsilon^3}{160} \quad \text{where } h'(x) := \sup_{v \in T'} -v \cdot x + c_v.$$

Note that if  $|x - a| \leq \delta_1$  and  $|x - b| \leq \delta_2$ , then  $|x - \max\{a, b\}| \leq \max\{\delta_1, \delta_2\} \leq \delta_1 + \delta_2$ . This motivates us to define

$$h''(x) := \sup_{v \in T'} \{v \cdot x + c_v, -v \cdot x + c_v\} = \sup_{v \in T' \cup -T'} \{v \cdot x + c_v\}.$$

It follows from the above observation that

$$\begin{aligned} \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [|f_T(\mathbf{g}) - h''(\mathbf{g})|] &\leq \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [|f_T(\mathbf{g}) - h(\mathbf{g})|] + \mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [|f_T(\mathbf{g}) - h'(\mathbf{g})|] \\ &\leq \frac{\varepsilon^3}{80}. \end{aligned} \tag{22}$$

Set  $S_0 := T' \cup -T'$  and note that

$$|S_0| \leq 2^{\exp\left(O\left(\frac{1}{\varepsilon^3}\right)\right)}.$$

Thus the function  $h''(\cdot)$ , which approximates  $f_T(\cdot)$ , is a the supremum of a non-centered Gaussian process on a symmetric set.

**Step 2: Obtaining a Centered Symmetric Approximator.** Recall from above that  $n > c_\varepsilon$ . Consequently, we can assume that  $|S_0| + A \leq n$  for  $A$  that we will define shortly. In particular, we can view  $\mathbb{R}^n$  as  $\mathbb{R}^{|S'|} \times \mathbb{R}^{A'}$  for some  $A' \geq A$ .

Repeating the argument used to prove [Corollary 23](#) on the set  $S_0$  with error parameter  $\frac{\varepsilon^2}{20}$  (and applying a suitable rotation, using the rotation-invariance of the Gaussian distribution), we get that there exists a set  $S \subseteq \mathbb{R}^{|S_0|} \times \mathbb{R}^{A'} = \mathbb{R}^n$  such that  $|S| = |S_0|$  and

$$A = \exp\left(O\left(\frac{1}{\varepsilon^3}\right)\right)$$

such that

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_{n+A})} [|h''(\mathbf{g}) - f_S(\mathbf{g})|] \leq \frac{\varepsilon^3}{80}, \tag{23}$$

and furthermore each  $s \in S$  is of the form  $(s', \alpha e_i)$  for some  $i \in [A]$  and  $s' \in S_0$ . [Corollary 23](#) also guarantees that  $(s', -\alpha e_i)$  belongs to  $S$  whenever  $(s', \alpha e_i)$  belongs to  $S$ . This, combined with the symmetry of  $S_0$ , implies that the set  $S$  is symmetric.

**Step 3: Putting Everything Together.** [Equation \(21\)](#) follows from [Equations \(22\)](#) and [\(23\)](#) and the triangle inequality. Furthermore, the set  $S$  is symmetric as discussed above and has the desired size as  $|S| = |S_0|$ . This completes the proof of [Corollary 24](#).  $\square$

## 4 Sparsifying Intersections of Narrow Halfspaces

As another consequence of [Theorem 20](#), we obtain dimension-free sparsification of intersections of halfspaces of bounded (geometric) width:

**Theorem 3.** Suppose  $K \subset \mathbb{R}^n$  is an intersection of arbitrarily many halfspaces that are at distance at most  $r \geq 1$  from the origin, i.e. there exists  $T \subseteq \mathbb{S}^{n-1}$  and a collection of non-negative numbers  $\{r_t\}_{t \in T}$  such that

$$K = \bigcap_{t \in T} \{x \in \mathbb{R}^n : t \cdot x \leq r_t\} \quad \text{where } r_t \leq r \text{ for all } t \in T.$$

For  $0 < \varepsilon < 0.5$ , there exists  $L \subseteq \mathbb{R}^n$  which is an intersection of

$$2^{\exp\left(\tilde{O}\left(\frac{1}{\varepsilon^2}\right) \cdot r^4\right)} \text{ halfspaces}$$

such that  $\text{dist}_G(K, L) \leq \varepsilon$ .

This can be viewed as a convex set analogue of similar sparsification results obtained for DNFs of bounded width over the Boolean hypercube [LV96, Tre04, GMR13, LWZ21], adding to the “emerging analogy between Boolean functions and convex sets” [DNS21, DNS22, DNS24, CDN<sup>+</sup>24]. We discuss applications to learning theory and property testing in Section 5, and consider the tightness of our bounds in Section 6.

Turning to the proof of Theorem 3, we first establish it for the special case of intersections of halfspaces of width *exactly*  $r$  for  $r \geq 1$ :

**Lemma 28** (Special case of Theorem 3 when  $r_t = r$  for all  $t$ ). Fix  $r \geq 1$  and suppose  $K \subseteq \mathbb{R}^n$  is a convex set of geometric width  $r$ , i.e. there exists  $T \subseteq (1/r)\mathbb{S}^{n-1}$  such that

$$K = \bigcap_{t \in T} \{x \in \mathbb{R}^n : t \cdot x \leq 1\}.$$

For  $0 < \varepsilon \leq 0.5$ , there exists a set  $L \subseteq \mathbb{R}^n$  which is an intersection of

$$2^{\exp\left(\tilde{O}\left(\frac{1}{\varepsilon^2}\right) \cdot r^4\right)} \text{ halfspaces}$$

such that  $\text{dist}_G(K, L) \leq \varepsilon$ .

*Proof of Lemma 28.* If  $\text{Vol}(K) \leq \varepsilon$  then we can take  $L = \emptyset$ , and if  $\text{Vol}(K) \geq 1 - \varepsilon$  then we can take  $L = \mathbb{R}^n$ . So we assume for the remainder of the argument that

$$\text{Vol}(K) \in (\varepsilon, 1 - \varepsilon). \tag{24}$$

Next, we will show that

$$w(T) \leq 1 + \sqrt{\frac{2 \ln\left(\frac{2}{\varepsilon}\right)}{r^2}}. \tag{25}$$

If  $w(T) < 1$ , then Equation (25) holds trivially; so suppose  $w(T) \geq 1$ . Note that  $K(x) = 1$  if and only if  $f_T(x) \leq 1$ . Let  $\{\mathbf{X}_t\}_{t \in T}$  be the canonical Gaussian process on  $T$ . By Equation (24),

$$\begin{aligned} \varepsilon \leq \text{Vol}(K) &= \Pr_{\mathbf{g} \sim N(0, I_n)} [f_T(\mathbf{g}) \leq 1] \\ &\leq \Pr_{\mathbf{g} \sim N(0, I_n)} [ |f_T(\mathbf{g}) - w(T)| \geq |w(T) - 1| ] \quad (\text{because } w(T) \geq 1) \\ &\leq 2 \exp\left(\frac{-(w(T) - 1)^2}{2 \sup_{t \in T} \mathbf{Var}[\mathbf{X}_t]}\right) \end{aligned} \tag{26}$$

$$\leq 2 \exp\left(\frac{-r^2(w(T) - 1)^2}{2}\right) \tag{27}$$

where Equation (26) follows from Proposition 10 and Equation (27) relies on the fact that because  $T \subseteq (1/r)\mathbb{S}^{n-1}$ , we have  $\mathbf{Var}[\mathbf{X}_t] = 1/r^2$  for all  $t \in T$ . Rearranging gives Equation (25) as claimed.

Let  $\eta_1, \eta_2 > 0$  be parameters that we will set later. Since  $T$  has bounded Gaussian width, we can now use [Theorem 20](#) to obtain a set  $S \subseteq T$  with

$$|S| \leq 2^{2^{O\left(\frac{w(T)}{\eta_1}\right)}} \quad (28)$$

and constants  $\{c_s\}_{s \in S}$  such that

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \left| f_T(\mathbf{g}) - \sup_{s \in S} \{\mathbf{g} \cdot s + c_s\} \right| \right] \leq \eta_1. \quad (29)$$

Motivated by this, we define the  $S$ -subspace junta  $J : \mathbb{R}^n \rightarrow \mathbb{R}$  to be  $J(\mathbf{g}) := \sup_{s \in S} \{\mathbf{g} \cdot s + c_s\}$ . (Recall that a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is said to be an “ $S$ -subspace junta” if there is a subspace  $V = \text{span}(S)$  of  $\mathbb{R}^n$  such that  $f(x)$  depends only on the projection of  $x$  on the subspace  $V$ ; see e.g. [\[VX11, DMN21\]](#).) By Markov’s inequality and [Equation \(29\)](#),

$$\mathbf{Pr}_{\mathbf{g} \sim N(0, I_n)} \left[ |f_T(\mathbf{g}) - J(\mathbf{g})| \geq \sqrt{\eta_1} \right] \leq \sqrt{\eta_1}. \quad (30)$$

We have by [Theorem 12](#) that

$$\begin{aligned} \mathbf{Pr}_{\mathbf{g} \sim N(0, I_n)} \left[ |f_T(\mathbf{g}) - 1| \leq \eta_2 \right] &= \mathbf{Pr}_{\mathbf{g} \sim N(0, I_n)} \left[ |f_{rT}(\mathbf{g}) - r| \leq r\eta_2 \right] \\ &\leq 4\eta_2 \cdot r(1 + w(rT)) \\ &= 4\eta_2 \cdot r(1 + r \cdot w(T)) \\ &\leq 4\eta_2 \cdot r \left( 1 + r + \sqrt{2 \ln \left( \frac{2}{\varepsilon} \right)} \right) \end{aligned} \quad (31)$$

$$\leq 4\eta_2 \cdot r \left( 2r + \sqrt{2 \ln \left( \frac{2}{\varepsilon} \right)} \right) \quad (32)$$

where the penultimate inequality relied on [Equation \(25\)](#) and the final inequality relied on the fact that  $r \geq 1$ . Set

$$\eta_1 = \frac{\eta_2^2}{4} \quad \text{and} \quad \eta_2 = \frac{\varepsilon}{8r \left( 2r + \sqrt{2 \ln \left( \frac{2}{\varepsilon} \right)} \right)}$$

so as to make [Equation \(32\)](#) equal to  $\varepsilon/2$ . It follows from [Equations \(30\)](#) and [\(32\)](#) that

$$\mathbf{Pr}_{\mathbf{g} \sim N(0, I_n)} \left[ \mathbf{1}\{f_T(\mathbf{g}) \leq 1\} \neq \mathbf{1}\{J(\mathbf{g}) \leq 1\} \right] \leq \varepsilon.$$

Note that  $\mathbf{1}\{J(\mathbf{g}) \leq 1\}$  is an intersection of  $|S|$  many halfspaces. In particular, note that

$$\frac{1}{\eta_1} = \frac{4}{\eta_2^2} \leq \tilde{O}\left(\frac{1}{\varepsilon^2}\right) \cdot r^4 \quad \text{and so} \quad |S| \leq 2^{\exp\left(\tilde{O}\left(\frac{1}{\varepsilon^2}\right) \cdot r^4\right)}$$

thanks to [Equations \(25\)](#) and [\(28\)](#), which completes the proof.  $\square$

We now prove [Theorem 3](#) using [Lemma 28](#).

*Proof of Theorem 3.* By a limiting argument, we can assume without loss of generality that  $K$  is an intersection of finitely many halfspaces. Let  $N = |T|$ . We further note that we can assume every  $r_t \geq -\sqrt{2 \ln(2/\varepsilon)}$ , since otherwise  $\text{Vol}(K) < \varepsilon$  and we can trivially  $\varepsilon$ -approximate  $K$  by taking  $L$  to be the empty set.

Let  $Q \in \mathbb{N}$  be a parameter that we will soon set; think of  $Q$  as a large number. For  $t \in T$ , define

$$\delta_t := 2r - r_t, \quad M_t := \exp\left((\delta_t Q)^2\right)$$

and set  $M := \sup_{t \in T} M_t$ . Because  $r_t \leq r$  and  $r > 0$ , it follows that  $\delta_t > 0$  for all  $t \in T$ .

For each halfspace  $H_t = \{x : x \cdot t \leq r_t\}$ , define the halfspaces

$$H_t^{(i)} := \left\{ (x, y) \in \mathbb{R}^n \times \mathbb{R}^M : x \cdot t + \frac{y_i}{\sqrt{2} \cdot Q} \leq 2r \right\} \quad \text{for } i \in [M_t].$$

Define the convex body  $K' \subseteq \mathbb{R}^{n+M}$  as

$$K' := \bigcap_{t \in T} \bigcap_{i \in [M_t]} H_t^{(i)}.$$

We will show that a suitable cross-section of  $K'$  is a good approximator to  $K$ . By [Proposition 5](#)—in particular, by [Equations \(5\) and \(6\)](#)—we have that for any  $t \in T$ ,

$$\mu_t := \mathbf{E}_{\mathbf{y}_i \sim N(0, I_M)} \left[ \sup_{i \in [M_t]} \frac{\mathbf{y}_i}{\sqrt{2} \cdot Q} \right] = \frac{\sqrt{2 \ln(M_t)}(1 \pm 4(\ln M_t)^{-1})}{\sqrt{2} \cdot Q} = \delta_t \left( 1 \pm \frac{4}{\ln M_t} \right), \quad (33)$$

and

$$\mathbf{Var}_{\mathbf{y}_i \sim N(0, I_M)} \left[ \sup_{i \in [M_t]} \frac{\mathbf{y}_i}{\sqrt{2} \cdot Q} \right] = \frac{\tau}{Q^2 \ln(M_t)} = \frac{\tau}{Q^4 \delta_t^2}$$

for  $\tau$  that is at most an absolute constant. By Chebyshev's inequality,

$$\mathbf{Pr}_{\mathbf{y}_i \sim N(0, I_M)} \left[ \left| \sup_{i \in [M_t]} \frac{\mathbf{y}_i}{\sqrt{2} \cdot Q} - \mu_t \right| \geq \frac{k\sqrt{\tau}}{Q^2 \delta_t} \right] \leq \frac{1}{k^2}. \quad (34)$$

Taking  $k = \sqrt{3N/\varepsilon}$  ensures that the above probability is at most  $\varepsilon/3N$ . We now record the first constraint on  $Q$ , which is that we must ensure that for all  $t \in T$  the following holds:

$$\frac{k\sqrt{\tau}}{Q^2 \delta_t} = \sqrt{\frac{3N\tau}{\varepsilon}} \cdot \frac{1}{Q^2 \delta_t} \leq \frac{\varepsilon}{100N}, \quad \text{i.e.} \quad Q \geq \left[ 3^{1/4} \sqrt{100\tau} \left( \frac{N}{\varepsilon} \right)^{3/4} \max_{t \in T} \sqrt{\frac{1}{\delta_t}} \right].$$

[Equation \(34\)](#) thus implies that

$$\mathbf{Pr}_{\mathbf{y}_i \sim N(0, I_M)} \left[ \left| \sup_{i \in [M_t]} \frac{\mathbf{y}_i}{\sqrt{2} \cdot Q} - \mu_t \right| \geq \frac{\varepsilon}{100N} \right] \leq \frac{\varepsilon}{3N}. \quad (35)$$

Next, viewing  $H_t$  as a halfspace in  $\mathbb{R}^{n+M}$ , note that for any  $t \in T$ :

$$\text{dist}_G \left( H_t, \bigcap_{i \in [M_t]} H_t^{(i)} \right) \leq \frac{\varepsilon}{3N} + \text{dist}_G \left( H_t, \left\{ x : x \cdot t + \mu_t \pm \frac{\varepsilon}{100N} \leq 2r \right\} \right) \quad (\text{Equation (35)})$$

$$= \frac{\varepsilon}{3N} + \text{dist}_G \left( H_t, \left\{ x : x \cdot t \leq r_t \pm \frac{\varepsilon}{100N} \pm \frac{4(2r - r_t)}{\ln M_t} \right\} \right) \quad (36)$$

where we used [Equation \(33\)](#) to obtain [Equation \(36\)](#) from the previous line. Now we record the second constraint on  $Q$ , which comes from requiring that

$$\frac{4(2r - r_t)}{\ln M_t} \leq \frac{\varepsilon}{100N};$$

recalling that  $M_t = \exp((\delta_t Q)^2)$ , this constraint is

$$Q \geq \max_{t \in T} \frac{20}{\delta_t} \cdot \sqrt{\frac{N(2r - r_t)}{\varepsilon}}.$$

Combining this with [Equation \(36\)](#), we get that

$$\text{dist}_G \left( H_t, \bigcap_{i \in [M_t]} H_t^{(i)} \right) \leq \frac{\varepsilon}{3N} + \frac{4\varepsilon}{100N} \leq \frac{112\varepsilon}{300N}. \quad (37)$$

In particular, a union bound over all  $N$  halfspaces immediately implies that

$$\text{dist}_G(K, K') \leq \frac{112\varepsilon}{300}. \quad (38)$$

Next, note that the geometric width of every halfspace in  $K'$  is exactly  $2r(1 + (1/(2Q^2)))^{-1/2}$ , as  $t \in \mathbb{S}^{n-1}$  for all  $t \in T$ . Note that

$$\frac{2r}{\sqrt{1 + (1/2Q^2)}} = \Theta(r)$$

as  $Q \geq 1$ . It thus follows from [Lemma 28](#) that there exists a set  $L' \subseteq \mathbb{R}^{n+M}$  that is an intersection of

$$2^{\exp\left(\tilde{O}\left(\frac{1}{\varepsilon^2}\right) \cdot r^4\right)} \text{ halfspaces}$$

such that  $\text{dist}_G(K', L') \leq 188\varepsilon/300$ . Together with [Equation \(38\)](#), this immediately implies that  $\text{dist}_G(K, L') \leq \varepsilon$  because of the triangle inequality.

Finally, note that since

$$\text{dist}_G(K, L') = \mathbf{E}_{\mathbf{x}, \mathbf{y}} \left[ \mathbf{1}\{K(\mathbf{x}) \neq L'(\mathbf{x}, \mathbf{y})\} \right] \leq \varepsilon,$$

there exists  $y \in \mathbb{R}^M$  such that  $\text{dist}_G(K, L'|_y) \leq \varepsilon$  where we write  $L'|_y$  for the cross-section of  $L'$  with the coordinates in  $\mathbb{R}^M$  set to  $y$ . Since taking a cross-section cannot increase the number of halfspaces, it follows that  $L := L'|_y$  has at most as many facets as  $L'$ , which completes the proof.  $\square$

## 5 Algorithmic Applications

In this section we briefly describe some algorithmic applications of our structural results. More precisely, we obtain polynomial-time agnostic learning results, and constant-query tolerant testing results, for new classes of convex sets under the Gaussian distribution; we do this by combining our new structural result, [Theorem 3](#) with known algorithms for learning and testing. Recall that [Theorem 3](#) tells us that if  $K$  is any convex set of geometric width at most  $r \geq 1$ , then for any  $\varepsilon > 0$  there is a set  $L = L(\varepsilon) \subseteq \mathbb{R}^n$ , which is an intersection of at most

$$2^{\exp\left(r^4 \cdot \tilde{O}\left(\frac{1}{\varepsilon^2}\right)\right)} \text{ halfspaces,}$$

such that  $\text{dist}_G(K, L) \leq \varepsilon$ .

The following notation will be helpful: For  $r > 0$ , let  $\text{Conv}_r$  denote the class of all convex sets  $K \subset \mathbb{R}^n$  that have geometric width at most  $r$ . Recall that such a set  $K$  is an intersection of (possibly infinitely many) halfspace of the form  $\mathbf{1}\{u \cdot x \leq r_u\}$ , where each  $r_u \leq r$ .

## 5.1 (Agnostically) Learning Convex Sets of Bounded Width

In [KOS08] Klivans et al. gave an algorithm which runs in time  $n^{O(\log(k)/\varepsilon^4)}$  and *agnostically learns* intersections of  $k$  halfspaces over  $\mathbb{R}^n$  to accuracy  $\varepsilon$  using only independent labeled examples drawn from  $N(0, I_n)$ . This means that for any target function  $f : \mathbb{R}^n \rightarrow \{0, 1\}$ , if there is some intersection of  $k$  halfspaces  $g : \mathbb{R}^n \rightarrow \{0, 1\}$  which satisfies  $\text{dist}_G(f, g) \leq \text{opt}$ , then with high probability the algorithm outputs a hypothesis  $h : \mathbb{R}^n \rightarrow \{0, 1\}$  which satisfies  $\text{dist}_G(f, h) \leq \text{opt} + \varepsilon$ .

For the rest of this section, set the parameter  $\alpha(r, \varepsilon)$  as

$$\alpha(r, \varepsilon) := 2^{\exp\left(r^4 \cdot \tilde{O}\left(\frac{1}{\varepsilon^2}\right)\right)}.$$

By [Theorem 3](#), for any function  $f' \in \text{Conv}_r$  there is an intersection of at most  $\alpha(r, \varepsilon/2)$  halfspaces, which we denote  $g$ , which is  $\varepsilon/2$ -close to  $f'$ . The [KOS08] result (applied with its “ $\varepsilon$ ” parameter set to  $\varepsilon/2$ ) thus immediately yields the following:

**Corollary 29.** Let  $r \geq 1$ . There is an algorithm which runs in time  $n^{O(\log(\alpha(r, \varepsilon/2))/\varepsilon^4)}$  and agnostically learns the class  $\text{Conv}_r$  to accuracy  $\varepsilon$  using only independent labeled examples drawn from  $N(0, I_n)$ .

Diakonikolas et al. [DKK<sup>+</sup>24] have recently shown that if the learning algorithm is allowed to make black-box queries to the unknown target function  $f$ , then intersections of  $k$  halfspaces can be agnostically learned in time  $\text{poly}(n) \cdot 2^{\text{poly}(\log(k)/\varepsilon)}$ . This immediately gives the following:

**Corollary 30.** Let  $r \geq 1$ . There is an algorithm which runs in time  $\text{poly}(n) \cdot 2^{\text{poly}(\log(\alpha(r, \varepsilon/2))/\varepsilon)}$  and agnostically learns the class  $\text{Conv}_r$  to accuracy  $\varepsilon$  under the standard  $N(0, I_n)$  distribution using black-box queries.

For constant  $\varepsilon$ , [Corollary 29](#) achieves  $\text{poly}(n)$ -time agnostic learning for convex sets of any constant geometric width, and [Corollary 30](#) achieves  $\text{poly}(n)$ -time agnostic learning (with queries) for convex sets of geometric width  $c \cdot (\log \log n)^{1/4}$  for an absolute constant  $c > 0$ .

## 5.2 (Tolerantly) Testing Convex Sets of Bounded Width

In [DMN21] De et al. gave a *tolerant testing* algorithm for the class of intersections of  $k$  halfspaces under the standard  $N(0, I_n)$  Gaussian distribution, which uses  $k^{\text{poly}(\log(k)/\varepsilon)}$  black-box queries. This means that for any target function  $f : \mathbb{R}^n \rightarrow \{0, 1\}$ , for any desired input parameters  $0 \leq \varepsilon_1 < \varepsilon_2$  with  $\varepsilon_2 - \varepsilon_1 = \varepsilon$ ,

- If there is some intersection of  $k$  halfspaces  $g : \mathbb{R}^n \rightarrow \{0, 1\}$  which satisfies  $\text{dist}_G(f, g) \leq \varepsilon_1$  then with high probability the algorithm outputs “accept,” and
- If every intersection of  $k$  halfspaces  $g$  has  $\text{dist}_G(f, g) \geq \varepsilon_2$  then with high probability the algorithm outputs “reject.”



We emphasize that the query complexity of the [DMN21] algorithm is completely independent of the ambient dimension  $n$ .

Combining the [DMN21] testing algorithm with [Theorem 3](#), we immediately get that the class  $\text{Conv}_r$  is tolerantly testable with query complexity just dependent on  $r$  and independent of the ambient dimension  $n$ . In particular, we have the following result:

**Theorem 31.** Let  $r \geq 1$ . For any input parameters  $0 \leq \varepsilon_1 < \varepsilon_2$ , there is an algorithm which makes  $k^{\text{poly}(\log(k)/\varepsilon)}$  queries to the target function  $f : \mathbb{R}^n \rightarrow \{0, 1\}$  and has the following guarantee:

- If there is a function  $g \in \text{Conv}_r$  such  $\text{dist}_G(f, g) \leq \varepsilon_1$  then with high probability the algorithm outputs “accept,” and
- If for every  $g \in \text{Conv}_r$ ,  $\text{dist}_G(f, g) \geq \varepsilon_2$  then with high probability the algorithm outputs “reject.”

Here  $\varepsilon = \varepsilon_2 - \varepsilon_1$  and  $k = 2^{\exp\left(r^4 \cdot \tilde{O}\left(\frac{1}{\varepsilon^2}\right)\right)}$ .

## 6 Lower Bounds

In this section we establish some lower bounds which complement the upper bounds given in [Theorem 1](#), [Theorem 2](#) (and consequently [Corollary 24](#)) and [Theorem 3](#). Before giving these results in [Section 6.2](#), we first briefly discuss the “non-proper” nature of our upper bounds in [Section 6.1](#).

### 6.1 The Impossibility of “Proper” Approximation

In this section we observe that dimension-independent “proper” variants of our approximation results [Theorem 1](#) (sparsifying the  $f_T$  function), [Theorem 2](#) (sparsifying a norm), and [Theorem 3](#) (sparsifying a polytope with bounded geometric width) cannot exist.

Let  $T$  be a bounded subset of  $\mathbb{R}^n$ , so  $w(T) < \infty$ , and consider the corresponding centered Gaussian process  $\{\mathbf{X}_t\}_{t \in T}$  and the associated function  $f_T(x) = \sup_{t \in T} t \cdot x$ . [Theorem 1](#) states that  $f_T$  is  $\varepsilon$ -approximated in  $L_1$ , under  $N(0, I_n)$ , by a function  $g(x) = \sup_{s \in S} \{s \cdot x + c_s\}$  where  $S \subseteq T$  satisfies  $|S| \leq (1/\varepsilon)^{\exp(O(w(T)/\varepsilon))}$ ; so the size of  $S$  depends only on  $w(T)$  and  $\varepsilon$ , and is completely independent of the ambient dimension  $n$ .

Even though  $S$  is a subset of  $T$ , the approximating function  $g$  is not a “proper” approximator because of the shifts  $c_s$ . It is natural to wonder whether those shifts are necessary: Can  $f_T$  be  $\varepsilon$ -approximated in  $L_1$  by a function of the form  $\sup_{s \in S} \{s \cdot x\}$ , where  $S$  is a subset of  $T$  of size  $O_{\varepsilon, w(T)}(1)$ , independent of  $n$ ? The answer to this question is no, as shown by the following simple example:

**Example 32.** Let  $a(n) := \mathbf{E}_{g \sim N(0, I_n)}[\sup\{g_1, \dots, g_n\}]$  be the expected supremum of  $n$  i.i.d. standard Gaussians. Sharpening the bound from [Proposition 5](#), it is well known (see e.g. Exercise 5.1 of [vH16]) that  $a(n) = (1 \pm o_n(1))\sqrt{2 \ln n}$ . Let  $T = \{\frac{1}{a(n)}e_1, \dots, \frac{1}{a(n)}e_n\}$ , so the Gaussian width of  $T$  is  $w(T) = 1$ . Now consider any candidate “proper” approximator  $f_S(x)$  for  $f_T(x)$  which is obtained by choosing a subset  $S \subset T$ ,  $|S| = m$  of  $m = n^{1-c}$  of the  $n$  vectors in  $T$ , where  $0 < c < 1$  is any fixed constant. Without loss of generality we may take  $S$  to be the first  $m$  vectors in  $T$ , so  $f_S(x) = \frac{1}{a(n)} \sup_{1 \leq i \leq m} \{x_i\}$  whereas  $f_T(x) = \frac{1}{a(n)} \sup_{1 \leq i \leq n} \{x_i\}$ . Since  $a(m) = (1 \pm o(1))\sqrt{2 \ln m} = (1 \pm o(1))\sqrt{2(1-c) \ln n}$ , we have  $\mathbf{E}[\sup_{t \in S} \mathbf{X}_t] = (1 \pm o(1))\sqrt{1-c}$ ,

and so applying [Proposition 10](#) to the canonical Gaussian processes  $\{\mathbf{X}_t\}_{t \in T}$  and  $\{\mathbf{X}_s\}_{s \in S}$ , we get that

$$\begin{aligned} \Pr_{\mathbf{g} \sim N(0, I_n)} \left[ f_T(\mathbf{g}) \in [1 - o(1), 1 + o(1)] \right] &\geq 1 - o(1), \\ \Pr_{\mathbf{g} \sim N(0, I_n)} \left[ f_S(\mathbf{g}) \in [\sqrt{1-c} - o(1), \sqrt{1-c} + o(1)] \right] &\geq 1 - o(1), \end{aligned}$$

so  $\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} [|f_T(\mathbf{g}) - f_S(\mathbf{g})|] \geq 1 - \sqrt{1-c} - o(1) = \Omega(1)$ . Thus, even if  $w(T) = 1$ , the size of  $S \subset T$  which is needed for  $f_S(x)$  to  $\varepsilon$ -approximate  $f_T(x)$  in  $L_1$  under  $N(0, I_n)$  is at least  $n^{1-c(\varepsilon)}$ , where  $c(\varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0$ .

Turning to [Theorem 3](#), recall that that result shows that any convex set  $K \subset \mathbb{R}^n$  with geometric width at most  $r$  can be  $\varepsilon$ -approximated by an intersection  $L$  of at most

$$2^{\exp\left(r^4 \tilde{O}\left(\frac{1}{\varepsilon^2}\right)\right)} \text{ halfspaces,}$$

and that moreover the halfspaces constituting  $L$  in our construction may not be supporting halfspaces of  $K$ . We remark that in the analogous context of width- $w$  CNF approximation over  $\{0, 1\}^n$ , the state-of-the art result of Lovett et al. [[LWZ21](#)] gives an  $\varepsilon$ -approximating width- $w$  CNF of size  $s = (2 + \frac{1}{w} \log(1/\varepsilon))^{O(w)}$  which is obtained by keeping  $s$  (carefully chosen) clauses in the original CNF and discarding the rest of them. Thus it is natural to wonder whether there is an analogue of [Theorem 3](#) in which (i) each halfspace constraint in the approximator is required to be one of the original supporting halfspaces of  $K$ , and yet (ii) the sparsity of the approximator is independent of  $n$ . It turns out that this is impossible, as witnessed by the following simple example which is reminiscent of [Example 32](#):

**Example 33.** Let  $K$  be a convex set in  $\mathbb{R}^{n+1}$  which is the intersection of  $n$  halfspaces, the  $i$ -th of which is

$$x_0 + \frac{1}{a(n)} x_i \leq \sqrt{1 + \frac{1}{a(n)^2}}, \quad (39)$$

so the geometric width of  $K$  is 1. Similar to [Example 32](#), for any constant  $0 < c < 1$  let  $L$  be the intersection of  $m = n^{1-c}$  of these  $n$  halfspaces; without loss of generality we can suppose the  $m$  halfspaces are given by [Equation \(39\)](#) for  $i = 1, \dots, m$ . Now, observe that  $x \in K$  iff  $x_0 + \frac{1}{a(n)} \sup\{x_1, \dots, x_n\} \leq \sqrt{1 + \frac{1}{a(n)^2}}$ . Since by [Proposition 10](#) we have  $\Pr[\sup\{\mathbf{g}_1, \dots, \mathbf{g}_n\} = (1 \pm o(1))a(n)] = 1 - o(1)$ , for a  $1 - o(1)$  fraction of outcomes of  $\mathbf{g}_1, \dots, \mathbf{g}_n$ , we have that  $\mathbf{g} = (\mathbf{g}_0, \dots, \mathbf{g}_n) \in K$  iff

$$\mathbf{g}_0 \leq \sqrt{1 + \frac{1}{a(n)^2}} - \frac{(1 \pm o(1))a(n)}{a(n)} = \pm o(1).$$

Turning to  $L$ , we have that  $x \in L$  iff  $x_0 + \frac{1}{a(n)} \sup\{x_1, \dots, x_m\} \leq \sqrt{1 + \frac{1}{a(n)^2}}$ . By [Proposition 10](#) we have  $\Pr[\sup\{\mathbf{g}_1, \dots, \mathbf{g}_m\} = (1 \pm o(1))a(m)] = 1 - o(1)$ , so for a  $1 - o(1)$  fraction of outcomes of  $\mathbf{g}_1, \dots, \mathbf{g}_n$ , we have that  $\mathbf{g} = (\mathbf{g}_0, \dots, \mathbf{g}_n) \in L$  iff

$$\mathbf{g}_0 \leq \sqrt{1 + \frac{1}{a(n)^2}} - \frac{(1 \pm o(1))a(m)}{a(n)} = 1 - (1 \pm o(1))\sqrt{1-c} \pm o(1).$$

It follows that for any constant  $0 < c < 1$ , we have  $\text{Vol}(K \Delta L)$  is at least an absolute constant, so  $L$  cannot be an  $\varepsilon$ -approximator of  $K$  for sufficiently small constant  $\varepsilon$ .

This example can be adapted to the context of [Theorem 2](#) (norm approximation); we leave the details to the interested reader.

## 6.2 Lower Bounds

In this section we give lower bounds on sparsification. To begin, a straightforward geometric argument shows that any intersection of halfspaces which  $\varepsilon$ -approximates the unit ball  $\{(x_1, x_2) : x_1^2 + x_2^2 \leq 1\}$  in  $\mathbb{R}^2$  must use at least  $(1/\varepsilon)^{\Omega(1)}$  halfspaces. This simple example already shows that approximating bounded-geometric-width convex sets with only  $\text{polylog}(1/\varepsilon)$  halfspaces is impossible (in contrast with known results for Boolean CNF sparsification, as discussed in [Section 1.3](#)).

In this context it is instructive to recall the upper bound from [\[DNS24\]](#) which we stated as [Theorem 17](#). This result tells us that for any fixed value of  $n$ , if  $\varepsilon$  is allowed to grow sufficiently small relative to  $n$ , then for any convex set  $K \subseteq \mathbb{R}^n$  (not necessarily of bounded width),  $(n/\varepsilon)^{O(n)}$  halfspaces suffice for  $\varepsilon$ -approximation; this is  $\text{poly}(1/\varepsilon)$  many halfspaces for any fixed value of  $n$ . But this is not the end of the story: if we take  $\varepsilon \rightarrow 0$  and allow the dimension  $n$  to vary with  $\varepsilon$ , then it is possible to give a lower bound which is *exponential* in  $1/\varepsilon$ :

**Theorem 34.** For all  $\varepsilon > 0$ , for all sufficiently large  $n \geq n(\varepsilon)$ , there is a convex body  $K \subset \mathbb{R}^n$  of geometric width 1, such that any  $\varepsilon$ -approximating convex set  $L \subset \mathbb{R}^n$  for  $K$  must be an intersection of at least  $2^{\Omega(1/\varepsilon)}$  many halfspaces.

*Proof.* The argument is a reduction to a recent lower bound of [\[DNS24\]](#) ([Theorem 18](#)) showing that any intersection of halfspaces that  $\Theta(1)$ -approximates the origin-centered ball of radius  $\approx \sqrt{n}$  must use  $2^{\Omega(\sqrt{n})}$  halfspaces.

Given  $\varepsilon > 0$ , let  $n = \lceil 1/\varepsilon^2 \rceil$ . For convenience we work in  $\mathbb{R}^{n+1}$ . Let  $K \subset \mathbb{R}^{n+1}$  be the body defined as

$$\bigcap_{u=(u_1, \dots, u_n) : \|u\|_2=1} \left\{ x \in \mathbb{R}^{n+1} : \frac{u_1 x_1 + \dots + u_n x_n}{\sqrt{n}} + x_{n+1} \leq 1 + \frac{1}{n} \right\}. \quad (40)$$

It is clear that the geometric width of  $K$  is exactly 1.

Suppose that  $L$  is an intersection of halfspaces such that  $\text{Vol}(L, K) \leq c/\sqrt{n}$  (here  $c > 0$  is a suitably small absolute constant that we will specify below). For  $\rho \in \mathbb{R}$ , write  $K_\rho$  to denote the  $n$ -dimensional cross-section of  $K$ ,

$$\bigcap_{u=(u_1, \dots, u_n) : \|u\|_2=1} \left\{ x \in \mathbb{R}^n : \frac{u_1 x_1 + \dots + u_n x_n}{\sqrt{n}} + \rho \leq 1 + \frac{1}{n} \right\},$$

obtained by fixing the last coordinate  $x_{n+1}$  to  $\rho$ , and likewise write  $L_\rho$  for the corresponding cross-section of  $L$ . Using the fact that the pdf of the standard  $N(0, 1)$  Gaussian is at least 0.3 everywhere on  $[-1/(2\sqrt{n}), 1/(2\sqrt{n})]$ , there must exist an outcome of  $\rho \in [-1/(2\sqrt{n}), 1/(2\sqrt{n})]$  such that  $\text{Vol}_n(K_\rho \triangle L_\rho) \leq 10c$  (in fact, most outcomes of  $\rho \in [-1/(2\sqrt{n}), 1/(2\sqrt{n})]$  must have this property).

Fix such an outcome of  $\rho \in [-1/(2\sqrt{n}), 1/(2\sqrt{n})]$ . The set  $K_\rho \subset \mathbb{R}^n$  is equivalent to

$$\bigcap_{u=(u_1, \dots, u_n) : \|u\|_2=1} \left\{ x \in \mathbb{R}^n : u_1 x_1 + \dots + u_n x_n \leq \sqrt{n} \cdot \left( 1 + \frac{1}{n} - \rho \right) \right\},$$

which is an origin-centered ball of radius between  $\sqrt{n} - 1$  and  $\sqrt{n} + 1$ . Choosing  $c = 10\kappa$  for the value  $\kappa$  from [Theorem 18](#) and applying that theorem, we get that  $L_\rho$  must be an intersection of  $2^{\Omega(\sqrt{n})} = 2^{\Omega(1/\varepsilon)}$  halfspaces, so  $L$  must also be an intersection of  $2^{\Omega(1/\varepsilon)}$  halfspaces, and [Theorem 34](#) is proved.  $\square$

We remark that [Theorem 18](#) directly gives that for a suitable absolute constant  $\kappa > 0$ , for any integer  $w \geq 1$ , there is a convex set  $K$  of geometric width  $w$  (namely the origin-centered ball of radius  $w$  in  $\mathbb{R}^n$ , where  $n \geq w^2$ ) such that any  $\kappa$ -approximating convex set  $L$  for  $K$  must be an intersection of at least  $2^{\Omega(w)}$  many halfspaces.

We also remark that the lower bound of [Theorem 34](#) easily yields a similar lower bound on sparsifying suprema of Gaussian processes:

**Corollary 35.** For all  $\varepsilon > 0$ , for all sufficiently large  $n \geq n(\varepsilon)$ , there is a set  $T \subseteq \mathbb{R}^n$  with  $w(T) = 1$  such that any set  $S$  and collection of associated real constants  $\{c_s\}_{s \in S}$  satisfying

$$\mathbf{E}_{\mathbf{g} \sim N(0, I_n)} \left[ \left| f_T(\mathbf{g}) - \sup_{s \in S} \{\mathbf{g} \cdot s + c_s\} \right| \right] \leq \varepsilon \quad (41)$$

must have  $|S| = 2^{\tilde{\Omega}(1/\sqrt{\varepsilon})}$ .

The idea behind [Corollary 35](#) is that if there existed a sparse set  $S$  of size  $2^{\tilde{\Omega}(1/\sqrt{\varepsilon})}$  with associated constants  $\{c_s\}_{s \in S}$  satisfying [Equation \(41\)](#), then following the proof of [Theorem 3](#) we would get an approximating convex set  $L$  for the set  $K$  in [Equation \(40\)](#) with  $2^{\Omega(1/\varepsilon)}$  halfspaces, contradicting [Theorem 34](#). We leave the details to the interested reader.

## Acknowledgements

A.D. is supported by NSF grants CCF-1910534 and CCF-2045128. S.N. is supported by NSF grants CCF-2106429, CCF-2211238, CCF-1763970, and CCF-2107187. R.A.S. is supported by NSF grants CCF-2106429 and CCF-2211238. This work was partially completed while some of the authors were visiting the Simons Institute for the Theory of Computing. The authors would also like to thank the anonymous FOCS 2024 reviewers for helpful comments and feedback.

## References

- [AAGM15] S. Artstein-Avidan, A. Giannopoulos, and V.D. Milman. *Asymptotic Geometric Analysis, Part I*, volume 202 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2015. [8](#)
- [AAGM21] Shiri Artstein-Avidan, Apostolos Giannopoulos, and Vitali D Milman. *Asymptotic geometric analysis, Part II*, volume 261. American Mathematical Society, 2021. [8](#)
- [BDOS21] Sander Borst, Daniel Dadush, Neil Olver, and Makrand Sinha. Majorizing Measures for the Optimizer. In *12th Innovations in Theoretical Computer Science Conference (ITCS 2021)*, pages 73:1–73:20, 2021. [1](#)
- [BST19] Nikhil Bansal, Ola Svensson, and Luca Trevisan. New notions and constructions of sparsification for graphs and hypergraphs. In *60th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2019*, pages 910–928, 2019. [1](#)
- [CCK14] Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Anti-concentration and honest, adaptive confidence bands. *The Annals of Statistics*, 42(5):1787–1818, 2014. [5](#), [9](#), [16](#)

- [CCK15] Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Comparison and anti-concentration bounds for maxima of Gaussian random vectors. *Probability Theory and Related Fields*, 162:47–70, 2015. 9
- [CDN<sup>+</sup>24] Xi Chen, Anindya De, Shivam Nadimpalli, Rocco A. Servedio, and Erik Waingarten. Lower Bounds for Convexity Testing. 2024. [arXiv:2410.17958](https://arxiv.org/abs/2410.17958). 19
- [Cha14] Sourav Chatterjee. *Superconcentration and related topics*, volume 15 of *Springer Monographs in Mathematics*. Springer, 2014. 7, 8
- [DKK<sup>+</sup>24] I. Diakonikolas, D. Kane, V. Kontonis, C. Tzamos, and N. Zarifis. Agnostically Learning Multi-index Models with Queries. In *Proceedings of the 65th IEEE Symposium on Foundations of Computer Science (FOCS)*, 2024. To appear. 6, 23
- [DLP12] J. Ding, J. Lee, and Y. Peres. Cover times, blanket times, and majorizing measures. *Annals of Mathematics*, 175:1409–1471, 2012. 1
- [DMN21] Anindya De, Elchanan Mossel, and Joe Neeman. Robust testing of low dimensional functions. In *STOC '21: 53rd Annual ACM SIGACT Symposium on Theory of Computing*, pages 584–597, 2021. 6, 20, 23, 24
- [DN04] Herbert A. David and Haikady N. Nagaraja. *Order statistics*. John Wiley & Sons, 2004. 7
- [DNS21] Anindya De, Shivam Nadimpalli, and Rocco A. Servedio. Quantitative correlation inequalities via semigroup interpolation. In *12th Innovations in Theoretical Computer Science Conference, ITCS 2021*, volume 185, pages 69:1–69:20, 2021. 19
- [DNS22] Anindya De, Shivam Nadimpalli, and Rocco A. Servedio. Convex Influences. In *13th Innovations in Theoretical Computer Science Conference, ITCS*, volume 215 of *LIPICs*, pages 53:1–53:21, 2022. 19
- [DNS24] Anindya De, Shivam Nadimpalli, and Rocco A. Servedio. Gaussian Approximation of Convex Sets by Intersections of Halfspaces. In *Proceedings of the 65th IEEE Symposium on Foundations of Computer Science (FOCS)*, 2024. To appear. 10, 19, 26
- [Dud67] R. M. Dudley. The sizes of compact subsets of Hilbert space and continuity of Gaussian processes. *Journal of Functional Analysis*, 1(3):290–330, 1967. 1
- [Fer75] Xavier M. Fernique. Regularite des trajectoires des fonctions aleatoires gaussiennes. In *Ecole d'Eté de Probabilités de Saint-Flour IV–1974*, pages 1–96, 1975. 1, 9
- [FKV13] Vitaly Feldman, Pravesh Kothari, and Jan Vondrák. Representation, approximation and learning of submodular functions using low-rank decision trees. In *COLT 2013 - The 26th Annual Conference on Learning Theory*, volume 30 of *JMLR Workshop and Conference Proceedings*, pages 711–740, 2013. 5
- [FV16] Vitaly Feldman and Jan Vondrák. Optimal bounds on approximation of submodular and XOS functions by juntas. *SIAM J. Comput.*, 45(3):1129–1170, 2016. 5
- [GMR13] P. Gopalan, R. Meka, and O. Reingold. DNF sparsification and a faster deterministic counting algorithm. *Computational Complexity*, 22(2):275–310, 2013. 19

- [Gor88] Yehoram Gordon. On Milman’s inequality and random subspaces which escape through a mesh in  $\mathbb{R}^n$ . In *Geometric Aspects of Functional Analysis: Israel Seminar (GAFA) 1986–87*, pages 84–106. Springer, 1988. 1
- [Gor92] Yehoram Gordon. Majorization of Gaussian processes and geometric applications. *Probability Theory and Related Fields*, 91(2):251–267, 1992. 1
- [KOS08] A. Klivans, R. O’Donnell, and R. Servedio. Learning geometric concepts via Gaussian surface area. In *Proc. 49th IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 541–550, 2008. 6, 23
- [LO99] Rafał Latała and Krzysztof Oleszkiewicz. Gaussian measures of dilatations of convex symmetric sets. *The Annals of Probability*, 27(4):1922–1938, 10 1999. 4, 16
- [Lov82] László Miklós Lovász. Submodular functions and convexity. In *International Symposium on Mathematical Programming*, 1982. 5
- [LSZ19] Shachar Lovett, Noam Solomon, and Jiapeng Zhang. From DNF compression to sunflower theorems via regularity. 2019. [arXiv:1903.00580](https://arxiv.org/abs/1903.00580). 2
- [LT91] M. Ledoux and M. Talagrand. *Probability in Banach Spaces*. Springer, 1991. 1, 8
- [LV96] M. Luby and B. Velickovic. On deterministic approximation of DNF. *Algorithmica*, 16(4/5):415–433, 1996. 19
- [LWZ21] Shachar Lovett, Kewen Wu, and Jiapeng Zhang. Decision list compression by mild random restrictions. *Journal of the ACM (JACM)*, 68(6):1–17, 2021. 2, 5, 6, 19, 25
- [Mek15] R. Meka. A Polynomial Time Approximation Scheme for Computing the Supremum of Gaussian Processes. *Annals of Applied Probability*, 25(2):465–476, 2015. 1
- [Mil71] Vitali D Milman. A new proof of A. Dvoretzky’s theorem on cross-sections of convex bodies. *Funkcional. Anal. i Prilozen*, 5:28–37, 1971. 1
- [Nel16] J. Nelson. Chaining introduction with some computer science applications. *Bulletin of the EATCS*, (120), 2016. 1
- [OH10] Samet Oymak and Babak Hassibi. New null space results and recovery thresholds for matrix rank minimization. 2010. [arXiv:1011.6326](https://arxiv.org/abs/1011.6326). 1
- [ORS18] Samet Oymak, Benjamin Recht, and Mahdi Soltanolkotabi. Isometric sketching of any set via the restricted isometry property. *Information and Inference: A Journal of the IMA*, 7(4):707–726, 2018. 1
- [PV12] Yaniv Plan and Roman Vershynin. Robust 1-bit compressed sensing and sparse logistic regression: A convex programming approach. *IEEE Transactions on Information Theory*, 59(1):482–494, 2012. 1
- [Sas11] Sasha. Answer to “Expectation of the maximum of gaussian random variables,” Mathematics StackExchange, December 2011. Available [here](#). 7
- [Sto09] Mihailo Stojnic. Various thresholds for  $\ell_1$ -optimization in compressed sensing. 2009. [arXiv:0907.3666](https://arxiv.org/abs/0907.3666). 1

- [Sto13] Mihailo Stojnic. Regularly random duality. 2013. 1
- [Sud69] V. Sudakov. Gauss and Cauchy measures and  $\varepsilon$ -entropy. In *Doklady Akademii Nauk*, volume 185, pages 51–53. Russian Academy of Sciences, 1969. 1, 3
- [Tal87] Michel Talagrand. Regularity of Gaussian processes. *Acta Mathematica*, 159(none):99–149, 1987. 1, 3, 9
- [Tal95] Michel Talagrand. Sections of smooth convex bodies via majorizing measures. *Acta Mathematica*, 175(2):273–300, 1995. 1
- [Tal22] Michel Talagrand. *Upper and lower bounds for stochastic processes: decomposition theorems*, volume 60. Springer Nature, 2022. 1, 3, 9
- [Tko18] Tomasz Tkocz. Asymptotic Convex Geometry Lecture Notes, 2018. URL: <https://www.math.cmu.edu/~ttkocz/teaching/1819/asympt-conv-geom-notes.pdf>. 4, 16
- [Tre04] L. Trevisan. A note on approximate counting for  $k$ -DNF. In *Proceedings of the Eighth International Workshop on Randomization and Computation*, pages 417–426, 2004. 19
- [Ver18] Roman Vershynin. *High-Dimensional Probability: An Introduction with Applications in Data Science*, volume 47. Cambridge University Press, 2018. 1, 3, 7, 8, 9
- [vH16] Ramon van Handel. Probability in high dimension. *Lecture Notes (Princeton University)*, 2016. URL: <https://web.math.princeton.edu/~rvan/APC550.pdf>. 1, 9, 17, 24
- [vH18] Ramon van Handel. Chaining, interpolation, and convexity. *Journal of the European Mathematical Society*, 20(10):2413–2435, 2018. 1
- [VX11] Santosh S. Vempala and Ying Xiao. Structure from local optima: Learning subspace juntas via higher order PCA. *CoRR*, abs/1108.3329, 2011. URL: <http://arxiv.org/abs/1108.3329>, arXiv:1108.3329. 20
- [Wai19] Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge university press, 2019. 7