

Totally Δ -modular IPs with two non-zeros in most rows

Stefan Kober¹ [0000-0003-2610-1494]

¹Université Libre de Bruxelles

Abstract. Integer programs (IPs) on constraint matrices with bounded sub-determinants are conjectured to be solvable in polynomial time. We give a strongly polynomial time algorithm to solve IPs where the constraint matrix has bounded subdeterminants and at most two non-zeros per row after removing a constant number of rows and columns. This result extends the work by Fiorini, Joret, Weltge & Yuditsky (J. ACM 72(1), 1-50 (2025)) by allowing for additional, unifying constraints and variables.

Keywords: Integer Programming · Parametrized Integer Programming · Sub-determinants · Structural Graph Theory · Dynamic Programming.

1 Introduction

In this work, we are concerned with integer programs (IPs) of the following form:

$$\min \{c^\top x : Mx \leq b, x \in \mathbb{Z}^n\}, \quad (\text{IP})$$

where $c \in \mathbb{Z}^n$, $M \in \mathbb{Z}^{m \times n}$, and $b \in \mathbb{Z}^m$ for $m, n \in \mathbb{N}$. In general, assuming $P \neq NP$, one cannot hope for a polynomial-time algorithm to solve (IP). Still, there are certain subclasses of IPs that can be solved efficiently. The most prominent example for this is IPs with *totally unimodular* (TU) constraint matrices, i.e., if M is TU. Recall that a matrix $M \in \mathbb{Z}^{m \times n}$ is TU, if all its *subdeterminants* are in $\{0, \pm 1\}$, where a subdeterminant is the determinant of a square submatrix of M .

Totally unimodular matrices possess the nice geometric property that the convex hull of the feasible solutions to (IP) is equal to the space of feasible solutions of its linear relaxation $\min \{c^\top x : Mx \leq b, x \in \mathbb{R}^n\}$. Therefore, we can solve such IPs by solving a single linear program. Such (combinatorial) linear programs can be solved in strongly polynomial time due to a result by Tardos [28]. This implies polynomial solvability for a collection of combinatorial optimization problems that can be stated as IPs on TU constraint matrices, such as the maximum-flow problem, the matching problem in bipartite graphs, or the stable set problem in bipartite graphs.

Almost 30 years ago, Shevchenko [27] asked for generalizations of this result to IPs on constraint matrices with bounded subdeterminants. We call a matrix M *totally Δ -modular*, if all its subdeterminants are in $[-\Delta, \Delta] \cap \mathbb{Z}$. A fundamental open question in combinatorial optimization is the following:

Can we solve (IP) in polynomial time, when M is totally Δ -modular?

In a breakthrough paper, Artmann, Weismantel & Zenklusen [3] answered the question in the affirmative for $\Delta = 2$. Generalizing their ideas to $\Delta \geq 3$ has proven to

be difficult, despite some progress, see e.g. Nägele, Santiago & Zenklusen [25]. Fiorini, Joret, Weltge & Yuditsky [16] fix Δ to be an arbitrary constant, and further require a structural condition on M , see Theorem 1.

Theorem 1 (Fiorini, Joret, Weltge & Yuditsky [16]). *Let $\Delta \in \mathbb{N}$. There is a polynomial-time algorithm to solve (IP) if M is totally Δ -modular and has at most two non-zeros per row (or column).*

Another example for restricting the structure of the constraint matrix are *nearly totally unimodular matrices* as studied by Aprile et al. [2]. The same idea has also proven helpful for a strongly polynomial-time algorithm for linear programming, see Dadush et al. [8], who also study the case of two non-zeros per row (or column).

We follow this second stream of research and investigate the question in the case where Δ is arbitrary but fixed, and there is a further structural condition on the constraint matrix. In the context of this work, we study the following parametrized problem, and provide a strongly polynomial-time algorithm. In particular, our result combines ideas from Fiorini et al. [16] and Aprile et al. [2].

Problem 2. Fix $\Delta, k \in \mathbb{N}$. Let $m_1, n_1 \in \mathbb{N}$ and $m_2, n_2 \leq k$, and define $m := m_1 + m_2$, $n := n_1 + n_2$. Let $M = \begin{bmatrix} A & B \\ W & D \end{bmatrix} \in \mathbb{Z}^{m \times n}$ be totally Δ -modular, where $A \in \mathbb{Z}^{m_1 \times n_1}$, $B \in \mathbb{Z}^{m_1 \times n_2}$, $W \in \mathbb{Z}^{m_2 \times n_1}$ and $D \in \mathbb{Z}^{m_2 \times n_2}$. Assume A has at most two non-zeros per row. Let $c \in \mathbb{Z}^n$ and $b \in \mathbb{Z}^m$. Solve (IP).

We reduce Problem 2 to Problem 3 below, in which the main technical difficulty for this project lies.

Problem 3. Fix $\Delta, k \in \mathbb{N}$. Let $A \in \{0, \pm 1\}^{m \times n}$, $W \in \mathbb{Z}^{k \times n}$, let $M = \begin{bmatrix} A \\ W \end{bmatrix}$ be totally Δ -modular and assume A has two non-zeros per row. Let $c, \ell, u \in \mathbb{Z}^n$, $b \in \mathbb{Z}^m$ and $d \in \mathbb{Z}^k$. Solve $\min \{c^\top x : Ax \leq b, Wx = d, \ell \leq x \leq u, x \in \mathbb{Z}^n\}$.

In the (k -dimensional) *partially ordered knapsack problem*, we are given an integer $k \in \mathbb{N}$, a set of elements V with an underlying partial order, a profit function $p : V \rightarrow \mathbb{Z}$, k weight-functions $W_i : V \rightarrow \mathbb{Z}$ for $i \in [k]$, and k budgets $b_i \in \mathbb{Z}$ for $i \in [k]$. The goal is to find the maximum profit subset of V that respects the budgets with respect to the weight functions and adheres to the partial order as precedence constraints, i.e., in order to choose any element, we need to choose all its predecessors with respect to the partial order. This problem is known to be NP-complete even if there is only one weight function with bounded entries, and the partial order is bipartite [19]. It can be formulated as an IP on a totally unimodular constraint matrix that has two non-zeros per row, and k extra constraints. In the following, we refer to this formulation as the respective IP. This makes it an interesting problem to consider in the context of this work. Note that the Hasse diagram of the partial order induces a directed graph G on V .

1.1 Contribution

Our main result is a strongly polynomial-time algorithm for IPs as defined in Problem 2, see Theorem 4. By combining our structural results with a proximity result by

Aprile et al. [2], our proof technique implies an FPT-algorithm for Problem 3 if A is totally unimodular, see Corollary 5 and in particular for the k -dimensional partially ordered knapsack problem, see Corollary 6. For this problem, only few non-trivial polynomially solvable cases are known [21]. A proof for Corollary 5 has previously been given in the author's thesis [20].

Proofs for the statements are deferred to Section 6.

Theorem 4. *There is a strongly polynomial-time algorithm to solve Problem 2.*

Corollary 5. *There is an FPT-algorithm to solve Problem 3 with respect to the largest subdeterminant if A is totally unimodular.*

Corollary 6. *There is an FPT-algorithm for the k -dimensional partially ordered knapsack problem with parameter*

$$\Delta = \max\left\{ \sum_{v \in V(H)} W_i(v) : i \in [k], H \text{ connected subgraph of } G \right\}.$$

Note that for $k = 1$, Δ in Corollary 6 corresponds to the largest subdeterminant of the corresponding constraint matrix, see [20, Corollary 4.24]. For constant k , the subdeterminants of the respective IP are bounded by a function in k and Δ .

Our results contribute to the research on IPs with bounded subdeterminants in the following ways. We combine ideas from two previous technical papers [16,2] and push the boundary of IPs with bounded subdeterminants in terms of polynomial-time algorithms. In particular, we relate the subdeterminants of matrices that have at most two non-zeros after removing one row to a novel invariant of weighted (signed) graphs, called the *alternating weight of a tree*, see Proposition 11. We show that a bound on the alternating weight of a tree in an integer-weighted graph gives a structural restriction on the structure of non-zero weights within the graph. More specifically, given an instance of Problem 3, we obtain a weighted graph and show that graphs corresponding to matrices with bounded subdeterminants admit a structured decomposition, see Theorem 16. We exploit this decomposition in a dynamic program that allows us to solve Problem 2, see Theorem 4.

We remark that it is unclear whether a constant proximity and decomposition result similar to the one in [2] holds in our case. Therefore, in Section 5 we crucially need to exploit the fact that we work in a different graph class within the context of this work, and provide a substantially stronger decomposition. In particular, Aprile et al. [2] show that graphs associated with their instances admit a decomposition such that the bags of the tree-decomposition either contain a *small* number of vertices, correspond to a leaf of the tree and contain a *small* number of roots, or are *almost embeddable in a surface of bounded genus*. We can show that the latter type is not necessary in graphs occurring in the context of this work. Finally, our structural results are largely self-contained and in particular do not make use of the graph minor structure theorem for K_t -minor free graphs as required in [16,2].

1.2 Background and related work

The main focus of our work lies in integer programming with bounded subdeterminants, see Figure 1 for an overview of recent results. There has been a large number

of exciting results in the last years that consider the question of polynomial solvability, e.g., the aforementioned papers [3,16,25,2], if the matrix is $\{a, b, c\}$ -modular [17], if the problem corresponds to the total matching problem [15], or for restricted subdeterminants over a polynomial ring [6]. Further, there is a randomized polynomial-time algorithm for feasibility of IPs on constraint matrices that are strictly Δ -modular for $\Delta \in \{3, 4\}$ [25,24].




Fig. 1. Recent results on IPs with bounded subdeterminants: bimodular matrices [3], $\{a, b, c\}$ -modular matrices [17], two non-zeros per row (2 non-zeros) [16], nearly totally unimodular matrices, the cographic case (cographic + k) [2], and this work (2 non-zeros + k , shaded in green).

Secondly, we work on IPs over a class of constraint matrices *with a constant number of extra rows*. Such problems appear naturally within combinatorial optimization or operations research, where the extra rows can play the role of budget or exactness constraints. Since already the knapsack problem is NP-hard, which only has one budget-constraint, one typically needs bounds on the entries or subdeterminants to design polynomial-time algorithms for such problems. We specifically mention the exact matching problem [26], integer programming on few constraints [13,1], the exact matroid basis problem [23,12], integer programming over TU matrices with few extra constraints [2], or the partially ordered knapsack problem [21].

One important problem in the class of IPs with bounded subdeterminants and two non-zeros per row is the maximum stable set problem with bounded *odd cycle packing number*, i.e., there is a bound on the maximum number of disjoint odd cycles. This problem has an efficient algorithm due to Fiorini et al. [16], which we use in order to solve our base case. Two exemplary problems that are covered by Problem 2 are the budgeted stable set problem studied by Dhahan [9], and the partially ordered knapsack problem, where we further restrict to bounded subdeterminants.

1.3 Overview

We begin by setting up definitions that are used throughout the paper in Section 2. In Section 3, we reduce Problem 2 to Problem 3. We proceed to exploit the bound on the subdeterminants in order to obtain bounds on the parameters of a related graph, as well as a structural description of such graphs in Sections 4 and 5. We conclude the proof of our main result in Section 6 by providing a dynamic program. In the final Section 7, we mention related problems and directions for further research.

2 Preliminaries

Throughout the paper, we use the notation $[k] := \{1, \dots, k\}$. Given an integer matrix $A \in \mathbb{Z}^{m \times n}$, we denote its largest subdeterminant by

$$\Delta(A) := \max\{|\det(B)| : B \text{ is a submatrix of } A\}.$$

We refer to Diestel [10] for more background on the notions from graph theory that we use. We proceed to give an overview of the most important tools and definitions we use in the remainder of the paper.

Given a graph $G = (V, E)$, and a subset of vertices $U \subseteq V$, we define the *neighborhood* $N(U) \subseteq V - U$ of U in G , such that for every $v \in N(U)$, there is an edge $vv' \in E$ with $v' \in U$. Given an edge $e = vv'$ of G , we denote by G/e the graph obtained by *contracting* an edge e , where v and v' are replaced by one vertex v_e that is adjacent to all vertices in $N(\{v, v'\})$. We say for graphs G and H that G contains an *H -minor*, if we can obtain a graph isomorphic to H by deleting vertices, deleting edges, and contracting edges of G . We call a graph G a *subdivision* of a graph H , if we can obtain a graph isomorphic to G by replacing some of the edges of H by paths.

Given a graph G , we define a *tree-decomposition* of G as a tuple (T, \mathcal{B}) , such that T is a tree, and a collection of bags $\mathcal{B} = (B_t)_{t \in V(T)}$ where $B_t \subseteq V(G)$ for each $t \in V(T)$ is a subset of the vertices of G such that the following properties are satisfied:

- $\bigcup_{t \in V(T)} B_t = V(G)$,
- for each edge $vv' \in E(G)$, there is a vertex $t \in V(T)$ such that $\{v, v'\} \subseteq B_t$, and
- for each $v \in V(G)$, the set $\{t \in V(T) : v \in B_t\}$ induces a connected graph in T .

If T is a star, we also call (T, \mathcal{B}) a *star-decomposition*. We define the *adhesion-size* of a tree-decomposition as $\max\{|B_t \cap B_{t'}| : tt' \in E(T)\}$. Note that we allow G to not be connected, and B_t to be empty for some $t \in V(T)$.

3 Reduction

Theorem 7 (Cook, Gerards, Schrijver, & Tardos [7]). *There exists a function $f_7 : \mathbb{Z}^2 \rightarrow \mathbb{Z}$ with the following properties. Fix $\Delta \in \mathbb{N}$ and consider an integer program as in (IP). If M is totally Δ -modular, (IP) is feasible, and x^* is an optimal solution of the linear relaxation, then there exists an optimal solution z^* of (IP) with $\|x^* - z^*\|_\infty \leq f_7(n, \Delta)$.*

Cook, Gerards, Schrijver, & Tardos [7] show that $f_7(n, \Delta) \leq n\Delta$. This was recently improved by Celaya, Kuhlmann, Paat & Weismantel [5] to $f_7(n, \Delta) < \frac{4n+2}{9}\Delta$. If the constraint matrix has a special structure, it is possible to give a bound not depending on the dimension of the problem, see e.g. Aliev, Henk & Oertel [1], or Aprile et al. [2]. In fact, it is not clear that the dimension should appear in the proximity function at all.

The following statement relates the two problems from the introduction. Clearly, any instance of Problem 3 is also an instance of Problem 2. But in fact, there is a polynomial-time reduction from Problem 2 to Problem 3. Note that we need to solve a polynomial number of instances of Problem 3 in order to solve Problem 2.

Proposition 8. *If there is a polynomial-time algorithm to solve Problem 3, then there is a polynomial-time algorithm for Problem 2.*

The proof of the statement is based on the proximity result by Cook et al. [7], see Theorem 7. This theorem allows us to guess on a constant number of variables in any totally Δ -modular instance of (IP). Fiorini et al. [16, Lemma 4] show that A has at most $\log_2(\Delta)$ columns with entries larger than 1 in absolute value. Thus, we can efficiently enumerate all possible assignments to the corresponding variables. Therefore, we can solve a polynomial number of instances of Problem 3 in order to solve an instance of Problem 2.

In addition, we recenter the problem around $\mathbf{0}_n$, i.e., given a solution $x^* \in \mathbb{R}^n$ of the linear relaxation, we redefine our variables $x' := x - \lfloor x^* \rfloor$. By Theorem 7 there is an optimal solution z^* to the integer program, such that $\|z^*\|_\infty \leq f_7(n, \Delta)$. Thus, we can assume without loss of generality that the lower and upper bounds have bounded entries, i.e., $\ell, u \in [-f_7(n, \Delta), f_7(n, \Delta)]^n \cap \mathbb{Z}^n$.

For a detailed description of the reduction, we refer to Fiorini et al. [16], see Lemma 4 and the following discussion.

4 Subdeterminants

Let $M = \begin{bmatrix} A \\ w^\top \end{bmatrix}$, where $A \in \{0, \pm 1\}^{m \times n}$ with at most two non-zeros per row, and $w \in \mathbb{Z}^n$. In this section, we show a relationship between the subdeterminants of M , and the structure of a corresponding graph.

Let $G = (V, E)$ be a graph, and $\sigma : E \rightarrow \{-1, +1\}$ be a signing of its edges. We call the triple $S = (V, E, \sigma)$ a *signed graph*. We call a subgraph S' of S *odd* if $\prod_{e \in E(S')} \sigma(e) = -1$, and *even* otherwise. Observe that if $\sigma(e) = -1$ for all $e \in E$, then the notion of an *odd cycle* coincides with the notion of an odd cycle for unsigned graphs. The *odd cycle packing number* of a signed graph S is denoted by $\text{ocp}(S)$, and is defined as the maximum number of vertex-disjoint odd cycles. For a signed graph S , we define an edge-vertex incidence matrix $M \in \{0, \pm 1\}^{E \times V}$ in the following way. Given an edge $e = vv'$, $M_{e,v}M_{e,v'} = -\sigma(e)$, i.e., if e is odd, then $M_{e,v} = +1$ and $M_{e,v'} = +1$, or $M_{e,v} = -1$ and $M_{e,v'} = -1$, and if e is even, then $M_{e,v} = +1$ and $M_{e,v'} = -1$, or $M_{e,v} = -1$ and $M_{e,v'} = +1$. For v'' not incident to e , we set $M_{e,v''} = 0$. Note that given a signed graph, its edge-vertex incidence matrix is not necessarily unique, but any $\{0, \pm 1\}$ -matrix with two non-zeros per row is the edge-vertex incidence matrix of a signed graph, and thus has a unique associated signed graph.

The following result is well-known.

Proposition 9 (Grossman, Kulkarni, & Schochetman [18]). *Let $A \in \{0, \pm 1\}^{m \times n}$ be the edge-vertex incidence matrix of a signed graph S . Then $2^{\text{ocp}(S)} = \Delta(A)$.*

Let S be a signed graph, denote the set of subgraphs of S that are trees by $\mathcal{T}(S)$ and let $T \in \mathcal{T}(S)$. For v, v' vertices of T , we denote the edges of the unique $v - v'$ -path in T by $P_T(v, v')$. Note that T induces a partition on $V(T)$ as follows. We define $X_T, Y_T \subseteq V(T)$, such that $X_T \cap Y_T = \emptyset$, $X_T \cup Y_T = V(T)$, any even edge of T is in X_T or Y_T , and any odd edge of T crosses between X_T and Y_T . Hence, $P_T(v, v')$ is even if and only if v and v' are both in X_T or in Y_T . This partition can be found greedily.

Definition 10. *Let $S = (V, E, \sigma)$ be a signed graph and $w : V \rightarrow \mathbb{Z}$ be a weight function on the vertices of S . Let $T \in \mathcal{T}(S)$. We define the alternating weight of a tree as*

$$\text{alt}(T, w) := \left| \sum_{v \in X_T} w(v) - \sum_{v \in Y_T} w(v) \right|.$$

We define $\text{alt}(S, w) := \max_{T \in \mathcal{T}(S)} \text{alt}(T, w)$ to be the maximum alternating weight of a tree in S .

Note that $\text{alt}(T, w)$, and $\text{alt}(S, w)$ are well-defined, since the bipartition is unique up to exchanging the sides, and we take the absolute value. The alternating weight of a tree appears naturally in the context of subdeterminants.

Proposition 11. *Let $A \in \{0, \pm 1\}^{m \times n}$ be the edge-vertex incidence matrix of a signed graph $S = (V, E, \sigma)$ and $w : V \rightarrow \mathbb{Z}$ be a weight function on the vertices of S . Then $\text{alt}(S, w) \leq \Delta\left(\begin{bmatrix} A \\ w^\top \end{bmatrix}\right)$.*

Proof. We show that any tree T of S gives rise to a subdeterminant of absolute value $\text{alt}(T, w)$ within the matrix $\begin{bmatrix} A \\ w^\top \end{bmatrix}$. We prove this by induction on the number of vertices of T . Given a tree T , we denote by M the square submatrix of $\begin{bmatrix} A \\ w^\top \end{bmatrix}$, where the rows are indexed by $E(T)$ and w , and the columns are indexed by $V(T)$.

If T has just a single vertex v_1 then $\text{alt}(T, w) = |w(v_1)| = |\det(M)|$. Assume that T has $t \geq 2$ vertices. Then T has a leaf v_1 with an edge $e = v_1 v_2$ to some vertex $v_2 \in V(T)$.

Recall for the following that for an edge $e = (v, v')$, $\sigma(e) = -M_{e,v} M_{e,v'}$. In M , we subtract the row indexed by e scaled with a factor of $w(v_1) \cdot M_{e,v_1}$ from the row indexed by w to obtain M' . To be precise, we define $M'_{w,v_1} := 0$, $M'_{w,v_2} := w(v_2) + \sigma(e) \cdot w(v_1)$, and $M'_{i,j} := M_{i,j}$ otherwise, for $i \in E(T) \cup \{w\}$, and $j \in V(T)$. Since M' is obtained from an elementary row operation on M , we observe that $\det(M') = \det(M)$. In M' , the column indexed by v_1 has exactly one non-zero entry of absolute value one, so we can remove the row indexed by e and the column indexed by v_1 to obtain a matrix of size $(t-1) \times (t-1)$ while preserving the determinant of M' in absolute value.

We define $T' := T - \{v_1\}$, and $w' : V(T') \rightarrow \mathbb{Z}$, where $w'(v_2) := w(v_2) + \sigma(e) \cdot w(v_1)$, and $w'(v) := w(v)$ for $v \in V(T') - \{v_2\}$. By the induction hypothesis, it follows that

$$|\det(M)| = |\det(M')| = \left| \sum_{v \in A_{T'}} w'(v) - \sum_{v \in B_{T'}} w'(v) \right| = \left| \sum_{v \in X_T} w(v) - \sum_{v \in Y_T} w(v) \right|.$$

□

We remark that Proposition 11 also follows from the *max-circuit imbalance* of an associated matrix, see [14, Proposition 3.2] for a relation between subdeterminants and the imbalance of circuits. In contrast, we are not aware of a proof of Proposition 9 based solely on the structure of circuits of the matrix.

5 Structure

Given a graph $G = (V, E)$ and k weight-functions $w_1, \dots, w_k : V \rightarrow \mathbb{Z}$, we define an associated set of terminals $R := \{v \in V \mid \exists i \in [k] : w_i(v) \neq 0\}$ as the set of vertices whose weight is non-zero with respect to at least one of the weight functions. For an integer $r \in \mathbb{N}$, we call G *r-decorated* if G is a tree, and at least r of its leaves are terminals.

Lemma 12. *Let $S = (V, E, \sigma)$ be a signed graph, $k, r \in \mathbb{N}$, and let $w_1, \dots, w_k : V \rightarrow \mathbb{Z}$ be k weight-functions on the vertices. Let $G := (V, E)$, and $R := \{v \in V \mid \exists i \in [k] : w_i(v) \neq 0\}$. If G has an r -decorated tree as a subgraph, then there is some $i \in [k]$ such that $\text{alt}(S, w_i) \geq \frac{r}{2k}$.*

Proof. Let T be an r -decorated tree in G with respect to R . Let $L(T)$ denote the set of leaves of T , and define the set of leaves which are terminals as $L' := L(T) \cap R$. Note that $|L'| \geq r$. By the pigeon-hole principle, there is some $i \in [k]$ such that $w_i(v) \neq 0$ for at least $\frac{r}{k}$ vertices in L' .

Fix a bipartition X_T, Y_T of T in order to compute the alternating weight of T . We denote the set of leaves with a positive contribution to $\text{alt}(T, w_i)$ by $L^+ := \{v \in L \cap X_T \mid w_i(v) > 0\} \cup \{v \in L \cap Y_T \mid w_i(v) < 0\}$, and the set of leaves with a negative contribution to $\text{alt}(T, w_i)$ by $L^- := \{v \in L \cap X_T \mid w_i(v) < 0\} \cup \{v \in L \cap Y_T \mid w_i(v) > 0\}$. Clearly, $T^+ := T \setminus L^-$ and $T^- := T \setminus L^+$ are trees, and $|\text{alt}(T^+, w_i) - \text{alt}(T^-, w_i)| \geq \frac{r}{k}$. Thus,

$$\text{alt}(S, w_i) \geq \max\{|\text{alt}(T^+, w_i)|, |\text{alt}(T^-, w_i)|\} \geq \frac{r}{2k}.$$

□

Theorem 13 (Ding [11, Lemma 3.1]). *Let $G = (V, E)$ be a graph with some set of terminals $R \subseteq V$, and $r \in \mathbb{N}$. There exists a function $f_{13}(r) \in O(r^2)$ with the following properties. If G has no r -decorated tree as a subgraph, then there is a star-decomposition (T, \mathcal{B}) of G (with t_0 the center of T), such that*

- $|B_{t_0} \cap R| \leq f_{13}(r)$,
- $|B_{t_0} \cap B_t| = 2$ for all $t \in V(T) \setminus \{t_0\}$, and both vertices in $B_{t_0} \cap B_t$ are in R ,
- $B_t \cap B_{t'} = \emptyset$ for all $t, t' \in V(T)$ leaves of T ,
- for all $t \in V(T) \setminus \{t_0\}$ it holds that $G[B_t]$ is connected, and each $r \in (B_t \setminus B_{t_0}) \cap R$ is a cut vertex in $G[B_t]$ separating the two vertices in $B_{t_0} \cap B_t$.

Note that this implies that the maximum degree of T is bounded from above by $f_{13}(r)/2$. We further remark that all steps in the proof of [11, Lemma 3.1] are constructive, so one can find the above decomposition in polynomial time.

In the proof of [11, Lemma 3.1], the concrete order of $f_{13}(r)$ is shown to be in $O(s(r)^2)$, where $s(r)$ denotes the minimal number such that every connected graph excluding a $K_{1,r}$ -minor is a subdivision of a graph on $s(r)$ vertices.

Lemma 14. *Let $r \in \mathbb{N}$, and G be a connected graph without $K_{1,r}$ -minor. There exists a function $s_{14}(r) \in O(r)$, such that G is a subdivision of a graph on at most $s_{14}(r)$ vertices.*

Proof. Fix $r \in \mathbb{N}$, and let G be a connected graph without $K_{1,r}$ -minor. Then G does not contain a tree with more than r leaves as a subgraph.

We first claim that given a tree T , the number of leaves is given by

$$\#(\text{leaves of } T) = 2 + \sum_{v \in V(T), \deg_T(v) \geq 3} (\deg(v) - 2).$$

From the handshake lemma and the fact that in a tree, $|E(T)| = |V(T)| - 1$, we know that $2 + \sum_{v \in V(T)} (\deg(v) - 2) = 0$. By splitting the sum into vertices of degree less than 2 and more than 2, we obtain that

$$2 + \sum_{v \in V(T), \deg_T(v) \geq 3} (\deg(v) - 2) = \sum_{v \in V(T), \deg_T(v) = 1} (\deg(v)),$$

which implies the claim.

Let $p \in \mathbb{N}$, and consider a subgraph of G that is a vertex-disjoint packing of K_{1,r_i} with $r_i \geq 3$ for all $i \in [p]$. We claim that

$$\sum_{i \in [p]} r_i \leq 3r, \tag{1}$$

so in particular $p \leq r$. To see this, assume to the contrary that there is a packing of K_{1,r_i} such that $r_i \geq 3$ for all $i \in [p]$ and $\sum_{i \in [p]} r_i > 3r$. Extend the packing to a spanning tree T in G . By the formula for the number of leaves of a tree, we have

$$\#(\text{leaves of } T) = 2 + \sum_{v \in V(T), \deg_T(v) \geq 3} (\deg(v) - 2) \geq 2 + \sum_{i \in [p]} (r_i - 2) > r,$$

a contradiction. The last inequality holds, since each $r_i \geq 3$, and $\sum_{i \in [p]} r_i > 3r$.

Choose an inclusion-wise maximal set of vertices $N_1 = \{v_1, \dots, v_p\}$ of degree at least 3 in G , such that $d(v_i, v_j) \geq 3$ for $i, j \in [p]$ with $i \neq j$. By (1), $|N_1| \leq r$, and $\sum_{i \in [p]} \deg(v_i) \leq 3r$. We define $N_2 := N(N_1)$, and observe that $|N_2| \leq 3r$. We further define $N_3 := N(N_2) \setminus N_1$. Assume that $|N_3| > 9r$. We construct a subset $N^* \subseteq N_2$ greedily, by iteratively choosing the vertex with the largest number of neighbors in N_3 , adding it to N^* and removing these neighbors from N_3 . We stop the process when $N(N^*) \cap N_3 > 3r$. By the pigeonhole principle, each vertex $v \in N^*$ has at least 3 unique neighbors in N_3 (similarly, the first one has at least 4 neighbors). Thus, $|N^*| \leq r$, a contradiction to (1). Since N_1 is maximal, all vertices of degree at least 3 must be contained in $N_1 \cup N_2 \cup N_3$. Further, there is at most r vertices of degree 1 in G (otherwise every spanning tree has more than r leaves).

Thus, the number of vertices in G with degree not equal to 2 is in $O(r)$, which concludes the proof. \square

We proceed to modify the decomposition from Theorem 13 in order to obtain a graph decomposition that allows us to employ a dynamic programming framework in order to solve Problem 3.

Lemma 15. *Let G be a connected graph with set of terminals $R = \{v_0, \dots, v_\alpha\}$ for some $\alpha \in \mathbb{N}$. Assume that for all $i \in [\alpha - 1]$, v_i is a cut vertex of G separating v_0 from v_α .*

Then there is an ordering of R , such that every $v_0 - v_\alpha$ -path in G visits all terminals in sequence with respect to the ordering. Further, any component of $G - R$ attaches to at most two terminals. These terminals must be consecutive with respect to the ordering.

Proof. We first claim that any $v_0 - v_\alpha$ -path contains each terminal in R . Assume otherwise, then there is some $i \in [\alpha - 1]$, such that v_i does not separate v_0 from v_α , a contradiction.

If there are two paths with a different ordering of terminals, then there is some $v_i \in R$, such that its position is different in the orderings. Consider the connected components of $G - v_i$, and denote the components containing v_0 and v_α by C_0 and C_α respectively. By the first claim, every $v \in R - v_i$ is in C_0 or C_α . Since v_i is a cut vertex, every $v_0 - v_\alpha$ -path first traverses C_0 and then C_α , a contradiction to the assumption that the position of v_i is different.

Therefore, we can assume that every $v_0 - v_\alpha$ -path visits all terminals in the same ordering. Now assume for a contradiction that there is a component C of $G - R$ that attaches to v_i and v_j for $i < j \in \{0, \dots, \alpha\}$ where $i + 1 \neq j$. Then, we can find a $v_0 - v_\alpha$ -path avoiding v_{i+1} by combining a $v_0 - v_i$ -path, a $v_i - v_j$ -path in C , and a $v_j - v_\alpha$ -path. This contradicts the first claim. \square




Fig. 2. A depiction of the structure of graphs in Lemma 15. The red vertices are terminals. The collections of components K_i (dashed) and L_j (dotted) are used in the proof of Theorem 16.

Theorem 16. *Let $r \in \mathbb{N}$ and G be a graph with set of terminals R . There is a function $f_{16}(r) \in O(r^2)$ with the following properties. If G has no r -decorated tree as a subgraph, then there is a tree-decomposition (T, \mathcal{B}) of G with adhesion-size and degree bounded by $f_{16}(r)$, such that each bag B_t for $t \in V(T)$ is of at least one of the following two types:*

- (i) $|B_t| \leq f_{16}(r)$
- (ii) t is a leaf of T with neighbor t' , and $B_t \cap R \subseteq B_{t'}$

Further, we can compute such a decomposition in polynomial time.

Proof. Let $f_{16} := \max\{3, f_{13}(r)/2 + 2, f_{13}(r)\}$. We first assume that G is a connected graph. Then, we can apply Theorem 13 and obtain a decomposition (T', \mathcal{B}') of G . We proceed to describe how to modify such a decomposition in order to obtain a decomposition (T, \mathcal{B}) that fulfills the conditions of the lemma.

For $k \geq 1$, we define a k -*comb* as a tree on $2k$ vertices $c_1, \dots, c_k, d_1, \dots, d_k$, consisting of a path c_1, \dots, c_k , and edges $c_i d_i$ for $i \in [k]$. We first describe how we modify T' in order to obtain T . Let w_0 be the center vertex of T' . For any leaf w of T' , we define $\lambda_w := |B'_w \cap R| - 1$. We replace each w by a λ_w -comb on the vertex set $\{c_1^w, \dots, c_{\lambda_w}^w, d_1^w, \dots, d_{\lambda_w}^w\}$, such that there is an edge $w_0 c_1^w$. Finally, we attach a vertex t_{-1} to the center vertex w_0 , and denote the resulting tree by T with center vertex t_0 .

We set $B_{t_{-1}} := B'_{w_0}$ and $B_{t_0} := B'_{w_0} \cap R$. Observe that B_{t_0} has type (i), $B_{t_{-1}}$ has type (ii), and their intersection is bounded by $f_{13}(r)$.

For any leaf w of T' , we denote the two terminals in $B'_w \cap B'_{w_0}$ by $v_0(w)$ and $v_{\lambda_w}(w)$. By Lemma 15 applied to B'_w , we obtain an ordering $v_0(w), v_1(w), \dots, v_{\lambda_w}(w)$ of the terminals, such that any $v_0(w) - v_{\lambda_w}(w)$ -path visits the terminals in this ordering. We set $B_{c_i^w} := \{v_{i-1}(w), v_i(w), v_{\lambda_w}(w)\}$ for $i \in [\lambda_w]$. Observe that each such bag has at most 3 elements and is of type (i).

It remains to describe the bags $B_{d_i^w}$ for $i \in [\lambda_w]$. By Lemma 15, each component of $G[B'_w] \setminus R$ attaches to at most two consecutive terminals in $v_0(w), v_1(w), \dots, v_{\lambda_w}(w)$. For $i \in [\lambda_w]$, we denote by K_i the union of all components that attach to $v_{i-1}(w)$ and $v_i(w)$. For $j \in \{0, \dots, \lambda_w\}$, we denote by L_j the union of all components that attach only to $v_j(w)$. We define $B_{d_1^w} := \{v_0(w), v_1(w), v_{\lambda_w}(w)\} \cup K_1 \cup L_0 \cup L_1$ and $B_{d_i^w} := \{v_{i-1}(w), v_i(w), v_{\lambda_w}(w)\} \cup K_i \cup L_i$ for $i \in \{2, \dots, \lambda_w\}$. Observe that all d_i^w are leaves of T , and any terminal that they contain is also present in c_i^w for all $i \in [\lambda_w]$. Therefore, each such bag is of type (ii) and the decomposition (T, \mathcal{B}) is of the desired form.

If G is not connected, we consider the connected components G_1, \dots, G_γ of G for some $\gamma \in \mathbb{N}$. By the previous arguments, we obtain a tree-decomposition (T_i, \mathcal{B}_i) for each $i \in [\gamma]$. We combine the tree-decompositions into one central decomposition in the following way.

We construct the tree T as a γ -comb on the vertex set $\{c_1, \dots, c_\gamma, d_1, \dots, d_\gamma\}$, where d_i is replaced by T_i , such that c_i attaches to the central vertex of T_i . This increases the maximum degree of the tree-decomposition by at most one. The collection of bags \mathcal{B} is obtained by copying the \mathcal{B}_i for $i \in [\gamma]$. Note that in this way, the bags corresponding to c_i for $i \in [\gamma]$ remain empty. \square

6 Dynamic Program

In this section, we describe how to exploit the structural results from Sections 4 and 5 into an efficient algorithm for Problem 3. We propose a *bottom-up* dynamic program on the tree-decomposition given by Theorem 16, which follows the ideas of the dynamic program presented in [2]. We define a set of *rooted instances* corresponding to subtrees of the tree-decomposition. In each of these subtrees, we compute optimal solutions for the possible assignments on the right hand sides of the extra rows as well as the possible assignments on variables in the intersection with the parent in the tree-decomposition.

In bags of type (ii), we fix all assignments on the terminals of the subgraph, and solve the remaining integer program with at most two non-zeros per row. Bags of type (i) have bounded size. Hence, we can enumerate all possible assignments to the corresponding variables. Since the degree of the tree-decomposition is bounded,

we can also combine the optimal solutions in the corresponding subtrees efficiently. Note that by Theorem 7 and Theorem 16, the number of possible assignments is polynomial in n and Δ .

In the context of the dynamic program, we work with *rooted tree-decompositions*. A rooted tree-decomposition of a graph G is a tree-decomposition (T, \mathcal{B}) with the additional property that some vertex $t_0 \in V(T)$ is identified as the root of T . In a rooted tree-decomposition (T, \mathcal{B}) with root t_0 , any vertex t of T induces a *rooted subtree*, i.e., the tree induced on the vertices $t' \in V(T)$, such that the unique $t' - t_0$ -path in T contains t . We denote this rooted subtree by $T(t)$, and its associated collection of vertices of G by $B(t) := \{v \in V(G) \mid \exists t' \in V(T(t)) : v \in B_{t'}\}$. Further, for any vertex of a rooted tree-decomposition $t \in V(T)$, we define its *adhesion* as $\text{adh}(t) := \emptyset$ if $t = t_0$, and $\text{adh}(t) := B_t \cap B_{t'}$ for t' being the unique parent of t in the rooted tree-decomposition otherwise.

We begin by recalling the setup for Problem 3. Let $\Delta, k \in \mathbb{N}$, $A \in \{0, \pm 1\}^{m \times n}$, $W \in \mathbb{Z}^{k \times n}$, such that A has two non-zeros per row. Let $M := \begin{bmatrix} A \\ W \end{bmatrix}$ be totally Δ -modular, $c, \ell, u \in \mathbb{Z}^n$, $b \in \mathbb{Z}^m$, and $d \in \mathbb{Z}^k$, which together define an instance of Problem 3. Let $S = (V, E, \sigma)$ be a signed graph with corresponding incidence matrix A , and denote $G := (V, E)$ the unsigned graph with set of terminals $R \subseteq V$, where $v \in V$ is a terminal if the column vector of W corresponding to v is non-zero. By Proposition 11 and Lemma 12, we conclude that G has no $(2k\Delta + 1)$ -decorated tree as a subgraph.

Let (T, \mathcal{B}) be the tree-decomposition of G given by Theorem 16 with respect to the set of terminals R . We turn (T, \mathcal{B}) into a rooted tree-decomposition by identifying the root as c_1 (in the notation of the proof of Theorem 16), i.e., the first vertex on the path of the comb that combines the decompositions of the components.

Further, we define assignment spaces for the right hand side and the variables respectively: $\mathcal{D} := [-n\Delta f_7(n, \Delta), n\Delta f_7(n, \Delta)]^k \cap \mathbb{Z}^k$, and $\mathcal{X}_t := [-f_7(n, \Delta), f_7(n, \Delta)]^{\text{adh}(t)} \cap \mathbb{Z}^{\text{adh}(t)}$.

Definition 17. *Given a vertex of the tree-decomposition $t \in V(T)$, an assignment on the adhesion to the parent $\chi_t \in \mathcal{X}_t$, and a target for the extra constraints $d_t \in \mathcal{D}$, we define a rooted instance of Problem 3. We denote by A_t the restriction of A to the columns corresponding to vertices of $B(t)$ and to the rows corresponding to edges induced on $G[B(t)]$. Similarly, we denote by W_t, c_t, ℓ_t , and u_t the restriction of W, c, ℓ and u to the vertices of $B(t)$, where $W_t(i, v) = 0$ and $c_t(i, v) = 0$ for $v \in \text{adh}(t)$ and $i \in [k]$. Finally, we denote by b_t the restriction of b to the edges induced on $G[B(t)]$.*

The rooted problem is defined as

$$\min \{c_t^\top x : A_t x \leq b_t, W_t x = d_t, \ell_t \leq x \leq u_t, x(v) = \chi_t(v) \forall v \in \text{adh}(t), x \in \mathbb{Z}^{B(t)}\}.$$

We denote an instance of the rooted problem by $I := I(t, \chi_t, d_t)$, and its optimal value by $\text{OPT}(I)$. By restricting to the vertices and edges in a single bag B_t instead of a rooted subtree $B(t)$, we define a local instance $I^0(t, \chi_t, d_t)$ with optimal value $F_t^0(\chi_t, d_t)$. We denote any optimal solution of $I^0(t, \chi_t, d_t)$ by $x_t^0(\chi_t, d_t)$.

Note that $I(t_0, \emptyset, d)$ is the original instance of Problem 3. Thus, in order to solve Problem 3 it is sufficient to give a recursive algorithm on how to solve rooted problems. Further, for any $t \in V(T)$, there is $(2n\Delta f_7(n, \Delta) + 1)^k \cdot (2f_7(n, \Delta) + 1)^{O((k\Delta)^2)}$ different possible assignments for rooted instances, which is polynomial for fixed $\Delta, k \in \mathbb{N}$.

Definition 18. Let $I(t, \chi_t, d_t)$ be a rooted instance corresponding to vertex $t \in V(T)$. Let t_1, \dots, t_q denote the children of t in T (where $q = 0$ if t is a leaf). We define

$$F_t(\chi_t, d_t) := \min \left\{ F_t^0(\chi_t, d_t^0) + \sum_{i \in [q]} (F_{t_i}(\chi_{t_i}, d_{t_i})) : \chi_{t_i} \in \mathcal{X}_{t_i} \forall i \in [q], \chi_t(v) = \chi_{t_i}(v) \right. \\ \left. \forall i \in [q], v \in B_t \cap B_{t_i}, d_t^0, d_{t_1}, \dots, d_{t_q} \in \mathcal{D}, d_t^0 + \sum_{i \in [q]} d_{t_i} = d_t \right\}.$$

We denote by $x_t(\chi_t, d_t)$ the composition of $x_{t_i}^0(\chi_{t_i}, d_{t_i})$ with $x_{t_i}(\chi_{t_i}, d_{t_i})$ for $i \in [q]$.

Proof (Proof of Theorem 4). By Proposition 8, it suffices to give a polynomial-time algorithm for instances of Problem 3. By Theorem 16 we obtain a tree-decomposition (T, \mathcal{B}) . We show that $F_t(\chi_t, d_t) = \text{OPT}(I(t, \chi_t, d_t))$ for all $t \in T$, all $\chi_t \in \mathcal{X}_t$, and $d_t \in \mathcal{D}$. We proceed by induction on the distance of a vertex $t \in T$ from the root t_0 , starting with the largest distance.

The claimed algorithm corresponds to computing $F_{t_0}(\emptyset, d)$ as in Definition 18.

By definition, the statement is true for any $t \in V(T) - t_0$, where t is a leaf. Assume that t is not a leaf. Then, t has q children t_1, \dots, t_q , such that for each $i \in [q]$, and each $\chi_{t_i} \in \mathcal{X}_{t_i}$ and $d_{t_i} \in \mathcal{D}$, it holds by induction that $F_{t_i}(\chi_{t_i}, d_{t_i}) = \text{OPT}(I(t_i, \chi_{t_i}, d_{t_i}))$. Recall from Section 3, that $\ell, u \in [-f_7(n, \Delta), f_7(n, \Delta)]^n \cap \mathbb{Z}^n$. Consider an optimal solution $x_t^*(\chi_t, d_t) \in [-f_7(n, \Delta), f_7(n, \Delta)]^n \cap \mathbb{Z}^n$ of $I(t, \chi_t, d_t)$. For $i \in [q]$, we can decompose $x_t^*(\chi_t, d_t)$ into solutions for $I_{t_i}(\chi_{t_i}, d_{t_i})$, where χ_{t_i} and d_{t_i} are induced by the corresponding assignments of $x_t^*(\chi_t, d_t)$. By definition of the spaces \mathcal{X}_{t_i} for $i \in [q]$ and \mathcal{D} , we can see that the corresponding optimal values have been pre-computed. Further, the dynamic program explores all valid assignments for variables in B_t . Thus, $c_t^\top x_t^*(\chi_t, d_t) = c_t^\top x_t(\chi_t, d_t)$, certifying optimality.

We conclude the proof with an analysis of the runtime. The total size of the tree-decomposition is linear in n , i.e., $|V(T)| \in O(n)$. For each vertex $t \in V(T)$, we obtain at most $(2n\Delta f_7(n, \Delta) + 1)^k \cdot (2f_7(n, \Delta) + 1)^{O((k\Delta)^2)}$ relevant rooted instances. Each local instance can be solved in polynomial time. For bags of type (i), this corresponds to enumerate all possible assignments within the proximity bound. For bags of type (ii), this corresponds to solving an integer program with bounded subdeterminants and two non-zeros per row, see Theorem 1. Solving a rooted instance corresponds to solving a local instance and combining it with a polynomial number of rooted instances, which can again be done in polynomial time.

Proof (Proof of Corollary 5). The statement follows from Theorem 4 and its proof with the following additional observations.

Since A is totally unimodular, we can solve the local problem for bags of type (ii) such that the exponent of the running time does not depend on Δ , see e.g. Tardos [28]. Further, here the proximity function $f_7(n, \Delta)$ does not depend on the dimension of the problem n , but just on Δ and k , see Aprile et al. [2, Theorem 4].

7 Conclusion

We study IPs with bounded subdeterminants and give a strongly polynomial-time algorithm to solve a new structured class of IPs. We achieve our results by combin-

ing structural results with a dynamic programming approach. Within the dynamic program, we use the algorithm to solve integer programs with bounded subdeterminants and two non-zeros per row [16] in a blackbox fashion. In particular, for our main result, the proximity result by Cook et al. [7] can be used directly. This is in contrast to other recent papers, that make use of auxiliary proximity or solution decomposition results, see e.g. [3,13,12,2,16]. Still, this leaves open, whether one can prove a stronger proximity result for Problem 3 or its generalizations. Such a proximity result could open the door to more general results on the solvability of totally Δ -modular IPs.

In the work by Fiorini et al. [16], the majority of the technical difficulty lies in constraint matrices with bounded subdeterminants and two non-zeros per *row*. They also solve the case with two non-zeros per *column*. We do not extend this part of their result to additional rows and columns. Note that such integer programs can model a version of the *exact matching* problem [26] with a structural restriction. In the exact matching problem, we are given a graph with red and blue edges, and $k \in \mathbb{N}$. The goal is to decide whether there exists a perfect matching with exactly k red edges. This problem is not known to be NP-hard, but no efficient deterministic algorithm is known. Subproblems involving the exact matching problem have proven to be complex to solve deterministically in the literature on subdeterminants, see [25,24,2].

Finally, we remark on a structural question for matrices with bounded subdeterminants. In many recent papers, it has been helpful to exploit a relation between such matrices and certain graph classes, see [3,16,25,24,2]. We ask whether this connection can be made more precise, and if the question of solving integer programs with bounded subdeterminants can be solved by analyzing the right graph classes. One indication for such a relation can be found in recent work on the number of disjoint columns in matrices with bounded subdeterminants, see [22,4].

Acknowledgements

The author thanks Manuel Aprile, Samuel Fiorini, Caleb McFarland, Stefan Weltge and Yelena Yuditsky for helpful discussions. The author acknowledges funding by the ERC Consolidator Grant 615640-ForEFront and the *Fonds de la Recherche Scientifique* - FNRS through research project BD-DELTA (PDR 20222190, 2021–24).

In addition, we would like to thank the anonymous reviewers for their careful reading and their helpful comments which greatly helped to improve the paper.

References

1. Aliev, I., Henk, M., Oertel, T.: Distances to lattice points in knapsack polyhedra. *Mathematical Programming* **182**(1), 175–198 (2020)
2. Aprile, M., Fiorini, S., Joret, G., Kober, S., Seweryn, M.T., Weltge, S., Yuditsky, Y.: Integer programs with nearly totally unimodular matrices: the cographic case. *ACM-SIAM Symposium on Discrete Algorithms (SODA25)* (2025)
3. Artmann, S., Weismantel, R., Zenklusen, R.: A strongly polynomial algorithm for bimodular integer linear programming. In: *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, pp. 1206–1219 (2017)

4. Averkov, G., Schymura, M.: On the maximal number of columns of a Δ -modular integer matrix: bounds and computations. *Mathematical Programming* pp. 1–29 (2023)
5. Celaya, M., Kuhlmann, S., Paat, J., Weismantel, R.: Proximity and flatness bounds for linear integer optimization. *Mathematics of Operations Research* (2023)
6. Celaya, M., Kuhlmann, S., Weismantel, R.: On matrices over a polynomial ring with restricted subdeterminants. In: *International Conference on Integer Programming and Combinatorial Optimization*. pp. 43–56. Springer (2024)
7. Cook, W., Gerards, A.M., Schrijver, A., Tardos, É.: Sensitivity theorems in integer linear programming. *Mathematical Programming* **34**(3), 251–264 (1986)
8. Dadush, D., Koh, Z.K., Natura, B., Olver, N., Vég, L.A.: A strongly polynomial algorithm for linear programs with at most two nonzero entries per row or column. In: *Proceedings of the 56th Annual ACM Symposium on Theory of Computing*. pp. 1561–1572 (2024)
9. Dhahan, J.: The maximum weight stable set problem with a budget constraint. Master's thesis, Simon Fraser University (2020)
10. Diestel, R.: *Graph Theory*, Graduate Texts in Mathematics, vol. 173. Springer, Heidelberg; New York, fourth edn. (2010)
11. Ding, G.: Graphs without large $K_{2,n}$ -minors. arXiv:1702.01355 (2017)
12. Eisenbrand, F., Rohwedder, L., Węgrzycki, K.: Sensitivity, proximity and FPT algorithms for exact matroid problems. In: *2024 IEEE 65th Annual Symposium on Foundations of Computer Science (FOCS)*. pp. 1610–1620. IEEE (2024)
13. Eisenbrand, F., Weismantel, R.: Proximity results and faster algorithms for integer programming using the Steinitz lemma. *ACM Transactions on Algorithms (TALG)* **16**(1), 1–14 (2019)
14. Ekbatani, F., Natura, B., Vég, L.A.: Circuit imbalance measures and linear programming. *Surveys in Combinatorics* pp. 64–114 (2022)
15. Ferrarini, L., Fiorini, S., Kober, S., Yuditsky, Y.: Total matching and subdeterminants. In: *International Symposium on Combinatorial Optimization*. pp. 192–204. Springer (2024)
16. Fiorini, S., Joret, G., Weltge, S., Yuditsky, Y.: Integer programs with bounded subdeterminants and two nonzeros per row. *Journal of the ACM* **72**(1), 1–50 (2025)
17. Glanzer, C., Stallknecht, I., Weismantel, R.: Notes on $\{a, b, c\}$ -modular matrices. *Vietnam Journal of Mathematics* **50**(2), 469–485 (2022)
18. Grossman, J.W., Kulkarni, D.M., Schochetman, I.E.: On the minors of an incidence matrix and its Smith normal form. *Linear Algebra and its Applications* **218**, 213–224 (1995)
19. Johnson, D.S., Niemi, K.: On knapsacks, partitions, and a new dynamic programming technique for trees. *Mathematics of Operations Research* **8**(1), 1–14 (1983)
20. Kober, S.A.: Three perspectives on integer programming: practical and theoretical applications, and the case of bounded subdeterminants. Ph.D. thesis, Technische Universität München (2023)
21. Kolliopoulos, S.G., Steiner, G.: Partially ordered knapsack and applications to scheduling. *Discrete Applied Mathematics* **155**(8), 889–897 (2007)
22. Lee, J., Paat, J., Stallknecht, I., Xu, L.: Polynomial upper bounds on the number of differing columns of Δ -modular integer programs. *Mathematics of Operations Research* (2022)
23. Liu, S., Xu, C.: On the congruency-constrained matroid base. In: *International Conference on Integer Programming and Combinatorial Optimization*. pp. 280–293. Springer (2024)
24. Nägele, M., Nöbel, C., Santiago, R., Zenklusen, R.: Advances on strictly Δ -modular IPs. *Mathematical Programming* pp. 1–30 (2024)
25. Nägele, M., Santiago, R., Zenklusen, R.: Congruency-constrained TU problems beyond the bimodular case. *Mathematics of Operations Research* **49**(3), 1303–1348 (2024)
26. Papadimitriou, C.H., Yannakakis, M.: The complexity of restricted spanning tree problems. *Journal of the ACM (JACM)* **29**(2), 285–309 (1982)

27. Shevchenko, V.: Qualitative topics in integer linear programming. American Mathematical Soc. (1996), original manuscript in russian
28. Tardos, É.: A strongly polynomial algorithm to solve combinatorial linear programs. Operations Research **34**(2), 250–256 (1986)