

Minimizing breaks by minimizing odd cycle transversals *

Koichi Fujii [†] Tomomi Matsui [†]

November 26, 2024

Abstract

Constructing a suitable schedule for sports competitions is a crucial issue in sports scheduling. The round-robin tournament is a competition adopted in many professional sports. For most round-robin tournaments, it is considered undesirable that a team plays consecutive away or home matches; such an occurrence is called a break. Accordingly, it is preferable to reduce the number of breaks in a tournament. A common approach is first to construct a schedule and then determine a home-away assignment based on the given schedule to minimize the number of breaks (first-schedule-then-break). In this study, we concentrate on the problem that arises in the second stage of the first-schedule-then-break approach, namely, the break minimization problem (BMP). We prove that this problem can be reduced to an odd cycle transversal problem, the well-studied graph problem. These results lead to a new approximation algorithm for the BMP.

Keywords: sports scheduling; tournament scheduling; round robin; odd cycle transversal.

1. Introduction

Constructing a suitable schedule for sports competitions is a crucial issue in sports scheduling. A (single) *round-robin tournament* (RRT) is a simple sports competition, where each team plays against every other team once. The RRT is a competition framework adopted in many professional sports such as soccer and basketball, especially in Europe. In this study, we consider an RRT schedule with the following properties:

*This work was supported by JSPS KAKENHI Grant Number JP20K04973.

[†]Department of Industrial Engineering and Economics, Tokyo Institute of Technology

- Each team plays one match in each *slot*, that is the day when a match is held.
- Each team has a home stadium, and each match is held at the home stadium of one of the two playing teams. When a team plays a match at the home of the opponent, we state that the team plays away.
- Each team plays each other team exactly once at home or away.

It is considered undesirable that a team plays consecutive away or home matches to reduce player’s burden; such an occurrence is called a *break*.

Accordingly, the number of breaks in a tournament should be reduced. To construct a tournament schedule with fewer breaks, the following two decomposition approaches are widely used: (i) first-break-then-schedule [3, 7, 15, 18, 25–27] and (ii) first-schedule-then-break [16–18, 20, 22, 24]. In the first-break-then-schedule approach, an home-away assignment (HA-assignment) is generated in the first stage, which assigns a home game or an away game to each team in each slot. In the second stage, a timetable consistent with the generated HA-assignment is found, if it exists. Contrarily, in the first-schedule-then-break approach, a timetable is constructed at the first stage. The second stage determines the home and away teams of each match. In this approach, the home advantage is further determined in the second stage. For detailed information, refer to [21]. In this study, we focus on a problem occurring in the second stage of the first-schedule-then-break approach. This problem is commonly called the break minimization problem (BMP), which determines an HA-assignment minimizing the number of breaks.

The research on the BMP, introduced by the seminal work of de Werra [4], has steadily evolved. By $B_{\min}(\tau)$, we denote the minimum total number of breaks over all possible HA-assignments for a given timetable τ of an RRT with $2n$ teams. For the lower bound of $B_{\min}(\tau)$, de Werra [4] demonstrated that any schedule of a round-robin tournament has at least $2n - 2$ breaks. For the upper bound, Miyashiro and Matsui [16] proposed an algorithm for finding an HA-assignment satisfying the condition that the number of breaks is less than or equal to $n(n - 1)$ for a given timetable. Post and Woeginger [20] revised this analysis and improved the upper bound of the number of breaks from $n(n - 1)$ to $(n - 1)^2$ if the number of teams $2n$ is not a multiple of 4.

Elf, Jünger, and Rinaldi [5] conjectured that the problem of finding $B_{\min}(\tau)$ is NP-hard, which has not been proven yet to the best of our knowledge. Miyashiro and Matsui [17] formulated BMP as MAX RES CUT. They also applied an approximation algorithm for MAX RES CUT based on the

semi-definite programming relaxation, which was proposed by Goemans and Williamson [6].

In this paper, we consider the odd cycle transversal (OCT) problem, which is the problem of deleting vertices to transform a graph into a bipartite graph.

After the breakthrough paper of Reed, Smith, and Vetta [23], various parameterized algorithms have been investigated [9–14]. Not only are there theoretical results, but there are also computational results for both the exact and heuristic approaches [2,8]. We will explore the relationship between the BMP and OCT problems and consequently propose a novel approximation algorithm of the BMP.

2. Preliminaries

2.1 Break Minimization Problem

In this section, we introduce some notations and formal definitions. We use the following notations and symbols throughout this paper:

- $2n$: number of teams, where $n \geq 2$,
- $T = \{1, 2, \dots, 2n\}$: set of teams,
- $S = \{1, 2, \dots, 2n - 1\}$: set of slots.

A schedule of an RRT is described as a pair of a timetable and an HA-assignment, which is defined below. In this study, we assume that a timetable τ of an RRT is a matrix whose rows and columns are indexed by T and S , respectively. An element $\tau(t, s)$ denotes the opponent that plays against team t at slot s . A timetable τ (of an RRT schedule) should satisfy the following conditions: (i) a row of τ indexed by team $t \in T$ is a permutation of teams in $T \setminus \{t\}$, and (ii) $\tau(\tau(t, s), s) = t$ ($\forall (t, s) \in T \times S$). For any pair of different teams $t_1, t_2 \in T$, $s(\{t_1, t_2\})$ indicates a slot in which teams t_1 and t_2 have a match. We denote a subset $\{1, 2, \dots, 2n - 2\} \subset S$ as S^- . Table 1 presents a timetable for an RRT schedule of four teams.

A team is defined to be at *home* in slot s if the team plays a match at its home stadium in s ; otherwise, it is *away* in s . An *HA-assignment* is a matrix $\mathbf{Z} = (z_{t,s})$ in which the components consist of \mathbf{H} and \mathbf{A} , whose rows and columns are indexed by T and S , respectively. A value of $z_{t,s}$ is equal to \mathbf{H} if team t plays a match in slot s at home; otherwise, it is equal to \mathbf{A} . For a given timetable τ , \mathbf{Z} is *consistent* with τ when $\{z_{t,s}, z_{\tau(t,s),s}\} = \{\mathbf{H}, \mathbf{A}\}$ holds

Table 1: Timetable τ

Slot	1	2	3
team 1	2	3	4
team 2	1	4	3
team 3	4	1	2
team 4	3	2	1

Table 2: HA-assignment \mathbf{Z}

Slot	1	2	3
team 1	H	A	H
team 2	A	A	H
team 3	H	H	A
team 4	A	H	A

Table 3: Partial HA-assignment

Slot	1	2	3
team 1	H	A	H
team 2	*	A	H
team 3	*	H	A
team 4	A	H	A

for each $(t, s) \in T \times S$. Table 2 presents an example of an HA-assignment that is consistent with the timetable presented in Table 1.

Given an HA-assignment \mathbf{Z} , we say that team t has a break at slot $s \in S \setminus \{1\}$ if $z_{t,s-1} = z_{t,s}$ holds. For example, in Table 2, team 2 has a break at slot 2, as there are consecutive away matches ($z_{2,1} = z_{2,2} = \text{A}$). The number of breaks in an HA-assignment is defined as the total number of breaks belonging to all teams.

Following the definitions introduced above, we are now ready to present a formal definition of the BMP:

Break Minimization Problem: *Given a timetable τ , the BMP finds an HA-assignment consistent with τ that minimizes the number of breaks.*

A *partial HA-assignment* is a $T \times S$ matrix $\mathbf{Z} = (z_{t,s})$ with the components consisting of $\{\text{H}, \text{A}, *\}$. For a given timetable τ , a partial HA-assignment \mathbf{Z} is *consistent* with τ when

$$(z_{t,s}, z_{\tau(t,s),s}) \in \{(\text{H}, \text{A}), (\text{A}, \text{H}), (\text{H}, *), (\text{A}, *), (*, \text{H}), (*, \text{A}), (*, *)\}$$

holds for each $(t, s) \in T \times S$. Table 3 presents an example of an HA-assignment that is consistent with the timetable presented in Table 1. Suppose that we have a partial HA-assignment \mathbf{Z} consistent with a given timetable τ . We can construct an HA-assignment $\tilde{\mathbf{Z}}$ consistent with τ by assigning H or A to *-components of \mathbf{Z} so that $\tilde{z}_{t,s} = z_{t,s}$ for $z_{t,s} \in \{\text{H}, \text{A}\}$ and $(\tilde{z}_{t,s}, \tilde{z}_{\tau(t,s),s}) \in \{(\text{H}, \text{A}), (\text{A}, \text{H})\}$ are satisfied.

2.2 Odd Cycle Transversal

Given an undirected graph $G = (V, E)$, a vertex subset $V' \subseteq V$ is called an *OCT* if and only if V' has a nonempty intersection with every odd cycle in G . In other words, the induced subgraph $G[V \setminus V']$ is a bipartite graph. The minimum OCT problem finds an OCT with the smallest cardinality.

Given an OCT V' of $G = (V, E)$, we can construct a map $\alpha : V \rightarrow (0, 1, 2)$ (where $(0, 1, 2)$ is an ordered triplet of 3-set $\{0, 1, 2\}$) satisfying that $\alpha^{-1}(0) = V'$ and both $\alpha^{-1}(1)$ and $\alpha^{-1}(2)$ are independent sets of G . Here, we note that this construction is not unique. Conversely, if we have a map $\alpha : V \rightarrow (0, 1, 2)$ satisfying the condition that

$$\text{both } \alpha^{-1}(1) \text{ and } \alpha^{-1}(2) \text{ are independent sets of } G, \quad (1)$$

then $\alpha^{-1}(0)$ becomes an OCT. We call such a map an *OCT-map*. Throughout this paper, we represent an OCT by an OCT-map $\alpha : V \rightarrow (0, 1, 2)$.

3. Main theorem

To describe our main theorem, we consider an auxiliary graph of a given timetable τ . It is an undirected graph $G(\tau) = (\hat{V}, \hat{E})$ with vertex-set $\hat{V} = T \times S^-$ and edge-set \hat{E} defined by $\hat{E} = \hat{E}_H \cup \hat{E}_0 \cup \hat{E}_1$ where

$$\begin{aligned} \hat{E}_H &= \{(t, s-1), (t, s)\} \mid t \in T, s \in S^- \setminus \{1\}, \\ \hat{E}_0 &= \bigcup_{s \in S^-} \{(t_1, s), (t_2, s)\} \mid t_1, t_2 \in T, s(\{t_1, t_2\}) = s\}, \text{ and} \\ \hat{E}_1 &= \bigcup_{s \in S^- \setminus \{1\}} \{(t_1, s-1), (t_2, s-1)\} \mid t_1, t_2 \in T, s(\{t_1, t_2\}) = s\}. \end{aligned}$$

Figure 3 depicts the auxiliary graph corresponding to the timetable in Table 1.

The following is our main theorem.

Theorem 3.1. *Let τ be a given timetable and $B_{\min}(\tau)$ the optimal value of the BMP defined by τ . Then, $B_{\min}(\tau)$ is equal to the size of the minimum OCT of the auxiliary graph $G(\tau)$.*

First, we show that an HA-assignment consistent with a given timetable τ gives an OCT-map of auxiliary graph $G(\tau)$. Let \mathbf{Z} be an HA-assignment consistent with τ . We construct a map $\alpha_{\mathbf{Z}} : \hat{V} \rightarrow (0, 1, 2)$ by

$$\alpha_{\mathbf{Z}}(t, s) = \begin{cases} 1 & (\text{if } z_{t,s} = \mathbf{H} \text{ and } z_{t,s+1} = \mathbf{A}), \\ 2 & (\text{if } z_{t,s} = \mathbf{A} \text{ and } z_{t,s+1} = \mathbf{H}), \\ 0 & (\text{otherwise}). \end{cases}$$




Figure 1: Auxiliary graph

Lemma 3.2. *The map $\alpha_{\mathbf{Z}}$ is an OCT-map, where $|\alpha_{\mathbf{Z}}^{-1}(0)|$, the size of the OCT, is equal to $B_{\min}(\tau)$.*

Proof. We first show that $\alpha_{\mathbf{Z}}^{-1}(1)$ is a independent set of $G(\tau)$. Let us assume that there exists an edge (v, w) whose endpoints belong to $\alpha_{\mathbf{Z}}^{-1}(1)$. The edge (v, w) belongs to \hat{E}_H, \hat{E}_0 , or \hat{E}_1 . If it belongs to \hat{E}_H , then there exists a $(t, s) \in \hat{V}$ such that $(v, w) = \{(t, s - 1), (t, s)\}$. Then, $z_{t,s} = \mathbf{A}$ follows from $\alpha_{\mathbf{Z}}(v) = 1$ and $z_{t,s} = \mathbf{H}$ follows from $\alpha_{\mathbf{Z}}(w) = 1$. This is a contradiction. If it belongs to \hat{E}_0 , then there exists $(s, t_1, t_2) \in S^- \times T \times T$ such that $(v, w) = \{(t_1, s), (t_2, s)\}$. Then, $z_{t_1,s} = \mathbf{H}$ follows from $\alpha_{\mathbf{Z}}(v) = 1$ and $z_{t_2,s} = \mathbf{H}$ follows from $\alpha_{\mathbf{Z}}(w) = 1$. This is also a contradiction. The same occurs for the case belonging to \hat{E}_1 . Therefore, $\alpha_{\mathbf{Z}}^{-1}(1)$ is a independent set. Similarly, $\alpha_{\mathbf{Z}}^{-1}(2)$ is a independent set, and $\alpha_{\mathbf{Z}}$ is an OCT-map of $G(\tau)$ from the definition. The size of the OCT, $|\alpha_{\mathbf{Z}}^{-1}(0)|$, is equal to $B_{\min}(\tau)$. \square

Table 4 indicates the OCT-map $\alpha_{\mathbf{Z}}$ which is obtained from the HA-assignment \mathbf{Z} in Table 3.

Next, we discuss a procedure to construct a partial HA-assignment from an OCT-map of $G(\tau)$. Let $\alpha : \hat{V} \rightarrow (0, 1, 2)$ be an OCT-map of $G(\tau) = (\hat{V}, \hat{E})$, which satisfies that both $\alpha^{-1}(1)$ and $\alpha^{-1}(2)$ are independent sets of

Table 4: OCT map $\alpha_{\mathbf{z}}$

Slot	1	2
team 1	1	2
team 2	0	2
team 3	0	1
team 4	2	1

Table 5: OCT map α

Slot	1	2
team 1	2	0
team 2	0	0
team 3	0	1
team 4	2	0

Table 6: Partial assignment

Slot	1	2	3
team 1	A	H	*
team 2	*	*	*
team 3	*	H	A
team 4	A	H	*

$G(\tau)$. The above property implies that each edge $\{(t, s-1), (t, s)\} \in \hat{E}_H$ satisfies that $(\alpha(t, s-1), \alpha(t, s)) \notin \{(1, 1), (2, 2)\}$. Thus, we can uniquely construct a partial HA-assignment \mathbf{Z}^α defined by

$$z_{t,s}^\alpha = \begin{cases} \text{H} & (\text{if } \alpha(t, s) = 1 \text{ or } \alpha(t, s-1) = 2), \\ \text{A} & (\text{if } \alpha(t, s) = 2 \text{ or } \alpha(t, s-1) = 1), \\ * & (\text{otherwise}). \end{cases}$$

When we apply the above procedure to the OCT-map in Table 4, we obtain the partial HA-assignment in Table 3.

A partial HA-assignment obtained from an OCT-map of $G(\tau)$ is not always consistent with the given timetable τ . For example, Table 5 gives an OCT-map α of $G(\tau)$ defined by timetable τ in Table 1. However, the partial HA-assignment \mathbf{Z}^α , presented in Table 6, is not consistent with τ in Table 1. This inconsistency arises because the pair of teams 1 and 3 play matches at each of their home venues in slot 2, which does not satisfy the properties of RRT schedules.

We will show that we can obtain a consistent partial HA-assignment by repairing α . To facilitate the description of the proof, we introduce some notations. Let t_1 and t_2 be teams that have a game in slot s , that is $t_2 = \tau(t_1, s)$. Here, we assume $s \geq 2$ and $s \in S^-$. Then, the following relations hold:

$$\begin{aligned} \{(t_1, s), (t_2, s)\} &\in \hat{E}_0, \{(t_1, s-1), (t_2, s-1)\} \in \hat{E}_1, \\ \{(t_1, s-1), (t_1, s)\}, \{(t_2, s-1), (t_2, s)\} &\in \hat{E}_H. \end{aligned}$$

Thus, the sequence $((t_1, s-1), (t_1, s), (t_2, s), (t_2, s-1))$ forms a 4-cycle on $G(\tau)$. We call such a 4-cycle a *rectangular cycle*. We denote the rectangular cycle corresponding to a pair of teams (t_1, t_2) as $C(t_1, t_2)$. A map α is *inconsistent* on a rectangular cycle $C(t_1, t_2)$ if the following holds:

$$\begin{pmatrix} \alpha(t_1, s-1) & \alpha(t_1, s) \\ \alpha(t_2, s-1) & \alpha(t_2, s) \end{pmatrix} \in \left\{ \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 2 \\ 1 & 0 \end{pmatrix} \right\};$$

otherwise, it is *consistent* on $C(t_1, t_2)$.

A map α is *locally bipartite* on a rectangular cycle $C(t_1, t_2)$ if and only if $\alpha^{-1}(1) \cap C(t_1, t_2)$ and $\alpha^{-1}(2) \cap C(t_1, t_2)$ are independent sets on $C(t_1, t_2)$.

Lemma 3.3. *Let τ be a given timetable and $G(\tau) = (\hat{V}, \hat{E})$ be the corresponding auxiliary graph. Let $\alpha : \hat{V} \rightarrow (0, 1, 2)$ be an OCT-map of $G(\tau)$. Assume that α is consistent on every rectangular cycle of $G(\tau)$. Then, the corresponding partial HA-assignment \mathbf{Z}^α is consistent with τ .*

Proof. Assume that \mathbf{Z}^α is inconsistent with τ . Then, there is an inconsistent pair $(z_{t,s}^\alpha, z_{\tau(t,s),s}^\alpha)$. If $s \geq 2$, then α becomes inconsistent on rectangular cycle $C(t, \tau(t, s))$, which contradicts the assumption of the lemma. If $s = 1$, then $z_{t,1}^\alpha = z_{\tau(t,1),1}^\alpha \neq *$ holds, which implies that $\alpha(t, 1) = \alpha(\tau(t, 1), 1) \neq 0$. It contradicts the assumption that $\alpha^{-1}(1)$ and $\alpha^{-1}(2)$ are independent sets. \square

The following lemma gives a procedure to construct a consistent partial HA-assignment from a given OCT-map.

Lemma 3.4. *Let τ be a given timetable and $G(\tau) = (\hat{V}, \hat{E})$ be the corresponding auxiliary graph. If we have an OCT-map $\alpha : \hat{V} \rightarrow (0, 1, 2)$ of $G(\tau)$, then there exists an OCT-map $\tilde{\alpha} : \hat{V} \rightarrow (0, 1, 2)$ of $G(\tau)$ satisfying that (1) $|\alpha^{-1}(0)| = |\tilde{\alpha}^{-1}(0)|$ and (2) the partial HA-assignment $\mathbf{Z}^{\tilde{\alpha}}$ is consistent with τ .*

Proof. Assume that the partial HA-assignment \mathbf{Z}^α is not consistent with the given timetable τ . Let s^* be the earliest slot of inconsistent rectangular cycles;

$$s^* = \min \left\{ s \in S^- \mid \begin{array}{l} \exists (t_1, t_2) \in T \times T, s = s(t_1, t_2) \\ \alpha \text{ is inconsistent on } C(t_1, t_2) \end{array} \right\}.$$

We note that $s^* \geq 2$ holds. We denote (t_1^*, t_2^*) by a pair of teams where $s^* = s(t_1^*, t_2^*)$ and α is inconsistent on $C(t_1^*, t_2^*)$. From definition,

$$(\alpha(t_1^*, s^* - 1), \alpha(t_2^*, s^*)) \in \{(1, 2), (2, 1)\} \quad (\text{see Table 7}).$$

We denote $\tau(t_2^*, s^* - 1)$ by t_3^* . In this proof, we assume that $(\alpha(t_1^*, s^* - 1), \alpha(t_2^*, s^*)) = (1, 2)$ without loss of generality.

Table 7: Inconsistent vertex (t_1^*, s^*)

Slot	$s^* - 1$	s^*
team t_1^*	$\alpha(t_1^*, s^* - 1) \neq 0$	0
team t_2^*	0	$\alpha(t_2^*, s^*) \neq 0$

If $s^* = 2$, then vertex $(t_2^*, s^* - 1)$ has three neighbors $\{(t_1^*, s^* - 1), (t_2^*, s^*), (t_3^*, s^* - 1)\} \subseteq \hat{V}$. When $s^* > 2$, vertex $(t_2^*, s^* - 1)$ has four neighbors $\{(t_1^*, s^* - 1), (t_2^*, s^*), (t_3^*, s^* - 1), (t_2^*, s^* - 2)\}$.

Now we introduce a pair of maps α_1 and α_2 (see Tables 8 and 9) defined by

$$\alpha_1(t, s) = \begin{cases} 0 & (\text{if } (t, s) = (t_1^*, s^* - 1)), \\ \alpha(t_1^*, s^* - 1) & (\text{if } (t, s) = (t_2^*, s^* - 1)), \\ \alpha(t, s) & (\text{otherwise}), \end{cases}$$

$$\alpha_2(t, s) = \begin{cases} \alpha(t_2^*, s^*) & (\text{if } (t, s) = (t_2^*, s^* - 1)), \\ 0 & (\text{if } (t, s) = (t_2^*, s^*)), \\ \alpha(t, s) & (\text{otherwise}). \end{cases}$$

Table 8: Map α_1

Slot	$s^* - 1$	s^*
team t_1^*	0	0
team t_2^*	$\alpha(t_1^*, s^* - 1) \neq 0$	$\alpha(t_2^*, s^*) \neq 0$

Table 9: Map α_2

Slot	$s^* - 1$	s^*
team t_1^*	$\alpha(t_1^*, s^* - 1) \neq 0$	0
team t_2^*	$\alpha(t_2^*, s^*) \neq 0$	0

Figure 3 depicts three rectangular cycles connected to $C(t_1^*, t_2^*)$. Both α_1 or α_2 are locally bipartite on $C(t_1^*, t_2^*)$, $C(t_1^*, \tau(t_1^*, s^* - 1))$ and $C(t_2^*, \tau(t_2^*, s^* + 1))$. Both maps are also consistent on them. For $C(t_2^*, \tau(t_2^*, s^* - 1))$, we denote $\{\alpha(t_2^*, s^* - 2), \alpha(t_3^*, s^* - 1), \alpha(t_3^*, s^* - 2)\}$ as $\{(u, v, w)\}$. Figure 3 exhibits the nodes corresponding to $\{(u, v, w)\}$.




Figure 2: Rectangular cycles connected to $C(t_1^*, t_2^*)$.

To obtain the desired OCT-map $\tilde{\alpha}$, we update α to α_\bullet so that α_\bullet satisfies that (a) α_\bullet is an OCT-map, (b) the number of rectangular cycles where α_\bullet is inconsistent decreases by 1, and (c) $|\alpha^{-1}(0)| = |\alpha_\bullet^{-1}(0)|$ holds. For $s^* = 2$, both α_1 and α_2 satisfy those properties. For $s^* > 2$, we have to choose either α_1 or α_2 carefully. The decision on which map to choose from follows

$$\alpha_\bullet = \begin{cases} \alpha_{3-u} & (\text{if } u \neq 0), \\ \alpha_{3-v} & (\text{if } v \neq 0), \\ \alpha_w & (\text{if } w \neq 0), \\ \alpha_1 & \text{otherwise.} \end{cases}$$

First, we demonstrate that the choice is well-defined. It suffices to show it for the following four cases:

(a) $u, v \neq 0, w = 0$; given the minimality of s^* , it follows that $u = v$. Therefore, $\alpha_{3-u} = \alpha_{3-v}$ holds.

(b) $u, w \neq 0, v = 0$; as α is an OCT-map, $u \neq w$ holds. Therefore, $\alpha_{3-u} = \alpha_w$ holds.

(c) $v, w \neq 0, u = 0$; the proof is the same as that for case (b).

(d) $u, v, w \neq 0$; as α is an OCT-map, $u \neq w$ and $u = v$ holds. Therefore, $\alpha_{3-u} = \alpha_{3-v} = \alpha_w$ holds.

Therefore, the choice is well-defined. Next, we will show that the chosen α_\bullet is locally bipartite on $C(t_2^*, \tau(t_2^*, s-1))$. This follows from the conditions $\alpha_\bullet(t_2^*, s-1) \neq \alpha_\bullet(t_2^*, s-2)$ and $\alpha_\bullet(t_2^*, s-1) \neq \alpha_\bullet(t_3^*, s-1)$. As α_\bullet is locally bipartite on every rectangular cycle, it is an OCT-map.

Finally, we will show that the chosen α_\bullet is consistent on $C(t_2^*, \tau(t_2^*, s-1))$. If it is inconsistent, then $\alpha_\bullet(t_2^*, s-1) \neq 0, \alpha_\bullet(t_3^*, s-2) \neq 0, \alpha_\bullet(t_3^*, s-1) \neq \alpha_\bullet(t_3^*, s-2)$ must hold. Meanwhile, $\alpha_\bullet(t_3^*, s-2) = \alpha_\bullet(t_3^*, s-2)$ must hold if $\alpha_\bullet(t_3^*, s-2) = w \neq 0$ holds. This is a contradiction. Therefore, α_\bullet is consistent on $C(t_2^*, \tau(t_2^*, s-1))$.

As a result, α_\bullet satisfies that (a) α_\bullet is an OCT-map, (b) the number of rectangular cycles where α_\bullet is inconsistent decreases by 1, and (c) $|\alpha^{-1}(0)| = |\alpha_\bullet^{-1}(0)|$ holds. Considering the procedure to update α by α_\bullet repeatedly, it is bounded above by the number of rectangular cycles where α is inconsistent. After this finite procedure, we obtain an OCT-map $\tilde{\alpha}$ that is consistent on every rectangular cycle. The corresponding partial HA-assignment $\mathbf{Z}^{\tilde{\alpha}}$ becomes consistent with τ from Lemma 3.4. This proves the lemma. \square

The above proof of the lemma gives a procedure to construct a desired OCT-map $\tilde{\alpha}$ satisfying that (1) $|\alpha^{-1}(0)| = |\tilde{\alpha}^{-1}(0)|$ and (2) the partial HA-

assignment $Z^{\hat{\alpha}}$ is consistent with the given timetable τ . We note that it is done in polynomial time. Now, we are ready to prove the main theorem.

Proof of Theorem 3.1. Let Z^* be an optimal HA-assignment of the BMP defined by τ . From Lemma 3.2, OCT-map α_{Z^*} of $G(\tau)$ satisfies that the size of the corresponding OCT $|\alpha_{Z^*}^{-1}(0)|$ is equal to $B_{\min}(\tau)$. Thus, the size of the minimum OCT of $G(\tau)$, denoted by $O_{\min}(\tau)$, satisfies that $O_{\min}(\tau) \leq B_{\min}(\tau)$.

Let α_* be an OCT-map corresponding to the minimum OCT of $G(\tau)$. Lemma 3.4 implies that there exists an OCT-map $\tilde{\alpha}$ of $G(\tau)$ that satisfies $|\alpha_*^{-1}(0)| = |\tilde{\alpha}^{-1}(0)|$ and the partial HA-assignment $Z^{\hat{\alpha}}$ is consistent with τ . Let \hat{Z} be an HA-assignment consistent with τ , which is obtained from $Z^{\hat{\alpha}}$ by assigning H or A to $*$ -components, as described in Section 2.1. Then, the number of breaks in \hat{Z} is less than or equal to $O_{\min}(\tau)$, which implies that $B_{\min}(\tau) \leq O_{\min}(\tau)$. \square

Agarwal, Charikar, Makarychev and Makarychev [1] proposed $O(\sqrt{\log v})$ -approximation algorithm for the edge bipartization problem defined on a graph with v vertices. The minimum edge bipartization problem finds the smallest edge subset E' such that the graph G' obtained by deleting E' is bipartite. As the degree of the auxiliary graph $G(\tau)$ is bounded by 4, the approximation algorithm of the OCT problem on $G(\tau)$ can be constructed by the approximation algorithm of the edge bipartization problem on $G(\tau)$ with the same approximation ratio. Consequently, we obtain the following result by applying it to the BMP.

Corollary 3.5. *There is a randomized polynomial-time $O(\sqrt{\log n})$ -approximation algorithm for BMP.*

4. Example

In this section, we present an example of a timetable with $n = 4$. Table 10 indicates an example of a timetable. We obtain the optimal OCT-map α by solving the OCT problem of the auxiliary graph $G(\tau)$. Following the integer linear programming formulation described in [8], we solved it using the Nuorium Optimizer version 26.1.0 [19], an integer linear programming solver.

We can anticipate an inconsistent rectangular cycle $C(3, 7)$. It can be recovered by the procedure described in Lemma 3.4. Table 12 indicates the updated OCT-map $\tilde{\alpha}$ and Table 13 indicates the optimal HA-assignment obtained by the OCT-map.

Table 10: Timetable of $n = 4$

Slot	1	2	3	4	5	6	7
1	4	8	5	3	2	6	7
2	3	7	4	8	1	5	6
3	2	6	8	1	7	4	5
4	1	5	2	7	6	3	8
5	7	4	1	6	8	2	3
6	8	3	7	5	4	1	2
7	5	2	6	4	3	8	1
8	6	1	3	2	5	7	4

Table 11: Optimal OCT-map α

Slot	1	2	3	4	5	6
1	2	1	2	1	2	1
2	2	1	0	2	0	1
3	1	2	1	2	0	1
4	1	2	1	2	1	2
5	2	0	1	2	1	2
6	2	1	2	1	0	2
7	1	2	0	0	1	2
8	1	0	2	1	2	1

Table 12: Updated OCT-map $\tilde{\alpha}$

Slot	1	2	3	4	5	6
1	2	1	2	1	2	1
2	2	1	0	2	0	1
3	1	2	1	2	0	1
4	1	2	1	2	1	2
5	2	0	1	2	1	2
6	2	1	2	1	0	2
7	1	2	0	1	0	2
8	1	0	2	1	2	1

Table 13: Optimal HA-assignment

Slot	1	2	3	4	5	6	7
1	A	H	A	H	A	H	A
2	A	H	A	A	H	H	A
3	H	A	H	A	H	H	A
4	H	A	H	A	H	A	H
5	A	H	H	A	H	A	H
6	A	H	A	H	A	A	H
7	H	A	A	H	A	A	H
8	H	A	A	H	A	H	A

References

- [1] Amit Agarwal, Moses Charikar, Konstantin Makarychev, and Yury Makarychev. $O(\sqrt{\log n})$ approximation algorithms for min uncut, min 2CNF deletion, and directed cut problems. In *Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*, pages 573–581, 2005.
- [2] Takuya Akiba and Yoichi Iwata. Branch-and-reduce exponential/FPT algorithms in practice: A case study of vertex cover. *Theoretical Computer Science*, 609:211–225, 2016.
- [3] Dirk Briskorn. Feasibility of home-away-pattern sets for round robin tournaments. *Oper. Res. Lett.*, 36(3):283–284, 2008.
- [4] D. de Werra. Scheduling in sports. *Studies on Graphs and Discrete Programming*, 11:381–395, 1981.
- [5] Matthias Elf, Michael Jünger, and Giovanni Rinaldi. Minimizing breaks by maximizing cuts. *Oper. Res. Lett.*, 31(5):343–349, 2003.
- [6] Michel X Goemans and David P Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *J. ACM*, 42(6):1115–1145, 1995.

- [7] Martin Henz. Scheduling a major college basketball conference—revisited. *Oper. Res.*, 49(1):163–168, 2001.
- [8] Falk Hüffner. Algorithm engineering for optimal graph bipartization. *Journal of Graph Algorithms and Applications*, 13(2):77–98, 2009.
- [9] Yoichi Iwata, Keigo Oka, and Yuichi Yoshida. Linear-time FPT algorithms via network flow. In *Proceedings of the twenty-fifth annual ACM-SIAM symposium on Discrete algorithms*, pages 1749–1761. SIAM, 2014.
- [10] Bart MP Jansen and Stefan Kratsch. On polynomial kernels for structural parameterizations of odd cycle transversal. In *International Symposium on Parameterized and Exact Computation*, pages 132–144. Springer, 2011.
- [11] Ken-ichi Kawarabayashi and Bruce Reed. An (almost) linear time algorithm for odd cycles transversal. In *Proceedings of the twenty-first annual ACM-SIAM symposium on Discrete Algorithms*, pages 365–378. SIAM, 2010.
- [12] Sudeshna Kolay, Pranabendu Misra, MS Ramanujan, and Saket Saurabh. Faster graph bipartization. *J. Comput. System Sci.*, 109:45–55, 2020.
- [13] Daniel Lokshantov, NS Narayanaswamy, Venkatesh Raman, MS Ramanujan, and Saket Saurabh. Faster parameterized algorithms using linear programming. *ACM Trans. Algorithms*, 11(2):1–31, 2014.
- [14] Daniel Lokshantov, Saket Saurabh, and Somnath Sikdar. Simpler parameterized algorithm for OCT. In *IWOCA 2009*, volume 5874 of *Lecture Notes in Computer Science*, pages 380–384. Springer, 2009.
- [15] Ryuhei Miyashiro, Hideya Iwasaki, and Tomomi Matsui. Characterizing feasible pattern sets with a minimum number of breaks. In *PATAT 2002*, volume 2740 of *Lecture Notes in Computer Science*, pages 78–99. Springer, 2002.
- [16] Ryuhei Miyashiro and Tomomi Matsui. A polynomial-time algorithm to find an equitable home-away assignment. *Oper. Res. Lett.*, 33(3):235–241, 2005.
- [17] Ryuhei Miyashiro and Tomomi Matsui. Semidefinite programming based approaches to the break minimization problem. *Comput. Oper. Res.*, 33(7):1975–1982, 2006.
- [18] George L Nemhauser and Michael A Trick. Scheduling a major college basketball conference. *Oper. Res.*, 46(1):1–8, 1998.
- [19] NTT DATA Mathematical Systems Inc. Nuorium Optimizer. <https://www.msi.co.jp/solution/nuopt/index.html>. Accessed: 2024-09-05.
- [20] Gerhard Post and Gerhard J Woeginger. Sports tournaments, home-away assignments, and the break minimization problem. *Discrete Optim.*, 3(2):165–173, 2006.
- [21] Rasmus V. Rasmussen. Scheduling a triple round robin tournament for the best Danish soccer league. *European J. Oper. Res.*, 185(2):795–810, 2008.
- [22] Rasmus V Rasmussen and Michael A Trick. The timetable constrained distance minimization problem. In *CPAIOR 2006*, volume 3990 of *Lecture Notes in Computer Science*, pages 167–181. Springer, 2006.
- [23] Bruce Reed, Kaleigh Smith, and Adrian Vetta. Finding odd cycle transversals. *Oper. Res. Lett.*, 32(4):299–301, 2004.
- [24] Michael A Trick. A schedule-then-break approach to sports timetabling. In *PATAT 2000*, volume 2079 of *Lecture Notes in Computer Science*, pages 242–253. Springer, 2000.
- [25] David Van Bulck and Dries Goossens. On the complexity of pattern feasibility problems in time-relaxed sports timetabling. *Oper. Res. Lett.*, 48(4):452–459, 2020.
- [26] David Van Bulck and Dries Goossens. Optimizing rest times and differences in games played: an iterative two-phase approach. *J. Sched.*, 25(3):261–271, 2022.
- [27] Lishun Zeng and Shinji Mizuno. On the separation in 2-period double round robin tournaments with minimum breaks. *Comput. Oper. Res.*, 39(7):1692–1700, 2012.