

On Two Conservative HDG Schemes for Nonlinear Klein-Gordon Equation

Shipra Gupta ^{*}and Amiya Kumar Pani ^{†‡}and Sangita Yadav [§]

Abstract

In this article, a hybridizable discontinuous Galerkin (HDG) method is proposed and analyzed for the Klein-Gordon equation with local Lipschitz-type non-linearity. *A priori* error estimates are derived, and it is proved that approximations of the flux and the displacement converge with order $O(h^{k+1})$, where h is the discretizing parameter and k is the degree of the piecewise polynomials to approximate both flux and displacement variables. After post-processing of the semi-discrete solution, it is shown that the post-processed solution converges with order $O(h^{k+2})$ for $k \geq 1$. Moreover, a second-order conservative finite difference scheme is applied to discretize in time and it is proved that the discrete energy is conserved with optimal error estimates for the completely discrete method. To avoid solving a nonlinear system of algebraic equations at each time step, a non-conservative scheme is proposed, and its error analysis is also briefly established. Moreover, another variant of the HDG scheme is analyzed, and error estimates are established. Finally, some numerical experiments are conducted to confirm our theoretical findings.

AMS subject classifications. 65N12, 65N22

1 Introduction

This paper focuses on two HDG methods for the following non-linear Klein-Gordon equation:

$$\left. \begin{aligned} u_{tt} - \Delta u + f(u) &= 0 && \text{in } \Omega \times (0, T], \\ u &= 0 && \text{on } \partial\Omega \times (0, T], \\ u|_{t=0} &= u_0, \quad u_t|_{t=0} = u_1 && \text{in } \Omega, \end{aligned} \right\} \quad (1.1)$$

where Ω is a convex polygonal domain in \mathbb{R}^2 with boundary $\partial\Omega$, $f(u)$ is a given nonlinear function, and u_0, u_1 are also given functions. When $f(s) = s^3 - s$, the problem (1.1), popularly known as the Klein-Gordon equation, occurs as a mathematical model in relativistic quantum mechanics [23] and also in order-disorder transition in solid mechanics [16]. Moreover, when $f(s) = \sin s$, (1.1) is called sine-Gordon equation and has a wide range of applications in the propagation of fluxons in Josephson junctions between two superconductors, the motion of a rigid pendulum coupled to a stretched wire, and fluid motion stability. For the existence of a unique weak solution of problem (1.1), we refer to [20, 30].

Earlier literature on numerical methods on finite difference methods for the nonlinear Klein-Gordon equation can be found in [1, 17, 18, 31] and references therein. Tourigny [30] has applied a conforming finite element method in spatial direction combined with product approximation for the nonlinear term and has discussed the optimal rate of convergence for the semi-discrete case. In the context of discontinuous Galerkin (LDG) schemes applied to the two-dimensional sine-Gordon equation on Cartesian grids, Baccouch [2] has derived error estimates and superconvergence results.

Now, we briefly discuss the HDG methods, which is the theme of this paper. The HDG methods have been introduced in [9, 10] in the context of steady-state diffusion problem, their convergence and superconvergence properties in [12] as well as their efficiently implemented in [22]. These schemes provide

^{*}Department of Mathematics, Birla Institute of Technology and Science, Pilani, Pilani Campus, Vidya Vihar, Pilani Rajasthan 333031 India (p20190427@pilani.bits-pilani.ac.in)

[†]Corresponding author

[‡]Department of Mathematics, Birla Institute of Technology and Science, Pilani, K.K. Birla Goa Campus, Zuarinagar, Goa-403726, India. (amiyap@goa.bits-pilani.ac.in)

[§]Department of Mathematics, Birla Institute of Technology and Science, Pilani, Pilani Campus, Vidya Vihar, Pilani Rajasthan 333031 India (sangita.yadav@pilani.bits-pilani.ac.in)

approximations that are more accurate than the ones given by any other DG methods for second-order elliptic problems [6]. The researchers in [5, 15, 27, 28] have addressed the nonlinear source term using the HDG method for semilinear elliptic and parabolic problems, essentially with globally Lipschitz non-linearity. With the help of the interpolatory HDG approach, the nonlinear component has been dealt with efficiently, and the optimal convergence rate has been proved. For a class of DG methods and with the aid of appropriate post-processing of the semi-discrete potential, Pani and Yadav [32] have derived superconvergence for nonlinear parabolic problems. Concerning a second-order linear wave equation, Cockburn *et al.* [8] have devised the energy-conservative HDG methods and have derived uniform in-time estimates along with the superconvergence result of the post-processed solution. For a recent survey on HDG methods, see [7].

In most of the papers discussed earlier on numerical approximations to the Klein-Gordon equation, except [30], the authors have derived their results for a global Lipschitz non-linearity. Since the non-linearity present in the problem (1.1) is of polynomial type, and in fact locally Lipschitz type, as a first step, we derive *a priori* bounds, which helps to deal with the nonlinearity in the discrete problem. This study builds on two conservative HDG techniques for the nonlinear Klein-Gordon equation with odd-degree non-linearity and achieves discrete energy conservation. The major contributions of this paper are summarized below.

- *A priori* bound in L^p norm for the discrete displacement variable in the semidiscrete HDG method is derived, and it is exploited to deal with the local Lipschitz non-linearity in the discrete system.
- Convergence of order $O(h^{k+1})$, is obtained when piece-wise polynomials of degree k approximate both displacement and the flux variables. Subsequently, using a variant of Baker's technique, the optimal rate of convergence for the displacement is established under a less smooth regularity assumption on the exact solution.
- Low-cost local postprocessing for the displacement has helped us to improve $L^\infty(L^2)$ norm convergence, which is of order $k + 2$.
- Fully discrete conservative and non-conservative schemes are proposed using the discrete-time Galerkin method, and corresponding error estimates are established.
- A variant of the HDG method based on [25] is analyzed, and error estimates are derived.
- The HDG approach has been numerically tested with various degrees of polynomials. Numerical results show that the displacement and flux converge with optimal order, while the post-processed solution achieves superconvergence. These results confirm our theoretical findings.

This paper is structured as follows. Section 2 introduces the semi-discrete HDG approach, provides some preliminary results and assumptions, and shows that the method is conservative. Section 3 focuses on a priori error analysis for approximating the HDG solution. The post-processing is covered in Section 4. Section 5 deals with fully discrete conservative and non-conservative schemes and their error analyses. Section 6 demonstrates the numerical results and concluding remarks.

Throughout this paper, whenever there is no confusion, the standard Sobolev spaces $H^m(\Omega)$ will be denoted by H^m with norm $\|\cdot\|_{H^m}$. Moreover, C denotes a generic positive constant, which may vary from step to step.

2 HDG Formulation

For HDG formulation, the original problem (1.1) is first rewritten in mixed form as a system of first-order differential equations by adding an auxiliary flux variable \mathbf{q} . This results in:

$$\mathbf{q} = \nabla u \text{ in } \Omega \times (0, T].$$

Then, (1.1) becomes

$$u_{tt} - \nabla \cdot \mathbf{q} + f(u) = 0, \text{ in } \Omega \times (0, T].$$

2.1 Discretization

Let \mathcal{T}_h be a shape regular triangulation of $\bar{\Omega}$ with $h_K = \text{diam}(K) \forall K \in \mathcal{T}_h$ and $h = \max_{K \in \mathcal{T}_h} h_K$. Let \mathcal{E}_h be the set of the faces E of elements K of the triangulation \mathcal{T}_h , with $\mathcal{P}_k(K)$ as the space of polynomials of degree $\leq k$ on K . Set

$$\begin{aligned} W_h &= \{w \in L^2(\Omega) : w|_K \in \mathcal{P}_k(K) \forall K \in \mathcal{T}_h\}, \\ \mathbf{V}_h &= \{\mathbf{v} \in [L^2(\Omega)]^2 : \mathbf{v}|_K \in [\mathcal{P}_k(K)]^2 \forall K \in \mathcal{T}_h\}, \\ M_h &= \{\mu \in L^2(\mathcal{E}_h) : \mu|_E \in \mathcal{P}_k(E) \forall E \in \mathcal{E}_h\}. \end{aligned}$$

Define

$$L^2(\mathcal{T}_h) = \prod_{K \in \mathcal{T}_h} L^2(K), \quad L^2(\mathcal{E}_h) = \prod_{E \in \mathcal{E}_h} L^2(E),$$

with inner-product and norm, respectively, as

$$\begin{aligned} (\mathbf{f}, \mathbf{g}) &= \sum_{K \in \mathcal{T}_h} (\mathbf{f}, \mathbf{g})_K, \quad (\phi, \psi) = \sum_{K \in \mathcal{T}_h} (\phi, \psi)_K, \quad \langle \phi, \psi \rangle = \sum_{K \in \mathcal{T}_h} \langle \phi, \psi \rangle_{\partial K}, \\ \langle \phi, \psi \rangle_{\mathcal{E}_h/\Gamma_\partial} &= \sum_{E \in \mathcal{E}_h/\Gamma_\partial} \int_E \phi \psi \quad \text{and} \quad \langle \phi, \psi \rangle_{\Gamma_\partial} = \sum_{E \in \Gamma_\partial} \int_E \phi \psi, \end{aligned}$$

where,

$$(\mathbf{f}, \mathbf{g})_K = \int_K \mathbf{f} \cdot \mathbf{g}, \quad (\phi, \psi)_K = \int_K \phi \psi, \quad \langle \phi, \psi \rangle_{\partial K} = \int_{\partial K} \phi \psi.$$

For vector valued square integrable space $\mathbf{L}^2(\mathcal{T}_h) = (L^2(\mathcal{T}_h))^d$, the inner-product and norm are similarly defined.

Now, the HDG semi-discrete formulation is for $t \in (0, T]$ to seek $(u_h(t), \mathbf{q}_h(t), \hat{u}_h(t)) \in W_h \times \mathbf{V}_h \times M_h$ such that

$$(\mathbf{q}_h, \mathbf{v}_h) + (u_h, \nabla \cdot \mathbf{v}_h) - \langle \hat{u}_h, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0 \quad \forall \mathbf{v}_h \in \mathbf{V}_h, \quad (2.1a)$$

$$(u_{htt}, w_h) + (\mathbf{q}_h, \nabla w_h) - \langle \hat{\mathbf{q}}_h \cdot \mathbf{n}, w_h \rangle + (f(u_h), w_h) = 0 \quad \forall w_h \in W_h, \quad (2.1b)$$

$$\langle \hat{\mathbf{q}}_h \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h/\Gamma_\partial} = 0 \quad \forall \mu_1 \in M_h, \quad (2.1c)$$

$$\langle \hat{u}_h, \mu_2 \rangle_{\Gamma_\partial} = 0 \quad \forall \mu_2 \in M_h, \quad (2.1d)$$

$$\hat{\mathbf{q}}_h \cdot \mathbf{n} = \mathbf{q}_h \cdot \mathbf{n} - \tau(u_h - \hat{u}_h) \quad \text{on } \mathcal{E}_h \quad (2.1e)$$

with initial conditions:

$$u_h(0) = u_{h0} \in W_h \quad \text{and} \quad u_{ht}(0) = u_{h1} \in W_h, \quad (2.2)$$

where u_{h0} and u_{h1} are to be defined later with the following approximation property:

$$\|u_0 - u_{h0}\| + \|u_1 - u_{h1}\| = O(h^{k+1}). \quad (2.3)$$

Here, $\Gamma_\partial = \mathcal{E}_h \cap \partial\Omega$ is the collection of exterior edges of the domain Ω , $\tau : \mathcal{E}_h \rightarrow \mathbb{R}^+$ is constant on each face of $K \in \mathcal{T}_h$ and \mathbf{n} is the unit outward normal to element K . We define the faces norm as

$$\|\psi - \hat{\psi}\|_+^2 = \sum_{E \in \mathcal{E}_h} \|\tau^{1/2}(\psi - \hat{\psi})\|_{L^2(E)}^2$$

2.2 HDG Projections

Now introduce HDG projections $\Pi_W : L^2(K) \rightarrow \mathcal{P}_k(K)$ and $\mathbf{\Pi}_V : [L^2(K)]^2 \rightarrow [\mathcal{P}_k(K)]^2$ such that for any $u \in L^2(K)$ and $\mathbf{q} \in [L^2(K)]^2$ define $(\Pi_W u, \mathbf{\Pi}_V \mathbf{q})$ as

$$(\mathbf{\Pi}_V \mathbf{q}, \mathbf{v})_K = (\mathbf{q}, \mathbf{v})_K \quad \text{for all } \mathbf{v} \in [\mathcal{P}_{k-1}(K)]^2, \quad (2.4a)$$

$$(\Pi_W u, w)_K = (u, w)_K \quad \text{for all } w \in \mathcal{P}_{k-1}(K), \quad (2.4b)$$

$$\langle \mathbf{\Pi}_V \mathbf{q} \cdot \mathbf{n} - \tau \Pi_W u, \mu \rangle_E = \langle \mathbf{q} \cdot \mathbf{n} - \tau u, \mu \rangle_E \quad \text{for all } \mu \in \mathcal{P}_k(E). \quad (2.4c)$$

This projection $(\mathbf{\Pi}_V, \Pi_W)$ as shown in the Appendix A of [11] is well defined and has reasonable approximation properties, which are stated below in terms of a Lemma 2.1.

Based on these projections, we define the quantities

$$\vartheta_u = \Pi_W u - u, \quad \vartheta_{\mathbf{q}} = \mathbf{\Pi}_V \mathbf{q} - \mathbf{q}, \quad \widehat{\vartheta}_u = \mathcal{P}u - u, \quad (2.5)$$

where \mathcal{P} is the L^2 projection onto M_h .

Lemma 2.1. Suppose $k \geq 0$, $\tau|_{\partial K}$ is non-negative and $\tau_K^{\max} := \max \tau|_{\partial K} > 0$. Then, the system (2.4) is uniquely solvable for $(\mathbf{\Pi}_V, \Pi_W)$. Furthermore, there is constant C independent of K and τ such that

$$\begin{aligned} \|\vartheta_{\mathbf{q}}\|_{[L^2(K)]^2} &\leq Ch_K^{l_q+1} |\mathbf{q}|_{H^{l_q+1}(K)} + Ch_K^{l_u+1} \tau_K^* |u|_{H^{l_u+1}(K)}, \\ \|\vartheta_u\|_{L^2(K)} &\leq Ch_K^{l_u+1} |u|_{H^{l_u+1}(K)} + C \frac{h_K^{l_q+1}}{\tau_K^{\max}} |\nabla \cdot \mathbf{q}|_{H^{l_q}(K)}, \end{aligned}$$

for l_u, l_q in $[0, k]$, here $\tau_K^* := \max \tau|_{\partial K/E^*}$, E^* is a face of K at which $\tau|_{\partial K}$ is maximum. Hence, if τ_K^* and $\frac{1}{\tau_K^{\max}}$ are uniformly bounded for all $K \in \mathcal{T}_h$, then the projection converges with order $k+1$ for both variables provided the solution is smooth enough.

2.3 Energy Conservation Property

This subsection deals with the discrete conservation property. Set $F(s) = \frac{1}{4}(1-s^2)^2$ and observe that, $F(s) \geq 0$ and $F'(s) = f(s)$ for all $s \in \mathbb{R}$. Below, we present the energy conservation property.

Theorem 2.1. The HDG approximation $(u_h(t), \mathbf{q}_h(t), \widehat{u}_h(t)) \in W_h \times \mathbf{V}_h \times M_h$ given by (2.1a)-(2.1d) satisfies the following discrete conservation property

$$\mathcal{E}(t) = \mathcal{E}(0), \quad \forall t \in [0, T] \quad (2.6)$$

where

$$\mathcal{E}(t) = \frac{1}{2} (\|u_{ht}(t)\|^2 + \|\mathbf{q}_h(t)\|^2 + \|u_h(t) - \widehat{u}_h(t)\|_+^2 + 2(F(u_h(t)), 1)).$$

Proof. Differentiate (2.1a) with respect to t . Then, choose $\mathbf{v}_h = \mathbf{q}_h$ in the resulting equation and $w_h = u_{ht}$ in (2.1b). On adding these two resulting equations, we arrive at

$$\frac{1}{2} \frac{d}{dt} (\|u_{ht}\|^2 + \|\mathbf{q}_h\|^2 + 2(F(u_h), 1)) + (u_{ht}, \nabla \cdot \mathbf{q}_h) + (\mathbf{q}_h, \nabla u_{ht}) - \langle \widehat{u}_{ht}, \mathbf{q}_h \cdot \mathbf{n} \rangle - \langle \widehat{\mathbf{q}}_h \cdot \mathbf{n}, u_{ht} \rangle = 0. \quad (2.7)$$

Now, the use of integration by parts shows

$$(u_{ht}, \nabla \cdot \mathbf{q}_h) + (\mathbf{q}_h, \nabla u_{ht}) = \langle u_{ht}, \mathbf{q}_h \cdot \mathbf{n} \rangle. \quad (2.8)$$

An application of (2.1c) with (2.1d) and (2.8) yields

$$(u_{ht}, \nabla \cdot \mathbf{q}_h) + (\mathbf{q}_h, \nabla u_{ht}) - \langle \widehat{u}_{ht}, \mathbf{q}_h \cdot \mathbf{n} \rangle - \langle \widehat{\mathbf{q}}_h \cdot \mathbf{n}, u_{ht} \rangle = \frac{1}{2} \frac{d}{dt} \|u_h - \widehat{u}_h\|_+^2. \quad (2.9)$$

On substitution of (2.9) in (2.7), we obtain

$$\frac{d}{dt} \mathcal{E}(t) = 0.$$

After integrating from 0 to t , we obtain that $\mathcal{E}(t) = \mathcal{E}(0) \quad \forall t \in (0, T]$. This completes the proof of (2.6). \square

The next corollary is a consequence of the discrete energy conservation property. Note that from (2.3), it follows for a positive constant C that

$$\|u_{h0}\| \leq C \|u_0\| \quad \text{and} \quad \|u_{h1}\| \leq C \|u_1\|. \quad (2.10)$$

Corollary 2.1. There is a positive constant C such that for $t > 0$,

$$\|u_{ht}(t)\|^2 + \|\mathbf{q}_h(t)\|^2 + \|u_h - \widehat{u}_h\|_+^2 + \|u_h(t)\|_{L^4}^4 \leq C.$$

Proof. From (2.6), it is easy to note using (2.10) that

$$\begin{aligned} \|u_{ht}(t)\|^2 + \|\mathbf{q}_h(t)\|^2 + \|u_h - \widehat{u}_h\|_+^2 + \|u_h(t)\|_{L^4}^4 &\leq \mathcal{E}(t) = \mathcal{E}(0) \\ &\leq \|u_{ht}(0)\|^2 + \|\mathbf{q}_h(0)\|^2 + \|u_h(0) - \widehat{u}_h(0)\|_+^2 \\ &\quad + \frac{1}{2}((1 - u_h^2(0))^2, 1) \\ &\leq C \left(\|u_1\|^2 + \|\nabla u_0\|^2 + |\Omega| + \|u_0\|_{L^4}^4 \right). \end{aligned}$$

This completes the rest of the proof. \square

3 Semidiscrete Error Estimates

3.1 L^p Estimate for the Discrete solution

Lemma 3.1. For given $w_h \in W_h$, $\widehat{w}_h \in M_h$ and $\mathbf{y} \in (L^2(\Omega))^2$, let $\mathbf{z} \in (L^2(\Omega))^2$ satisfy

$$(\mathbf{z}, \mathbf{v}_h) + (w_h, \nabla \cdot \mathbf{v}_h) - \langle \widehat{w}_h, \mathbf{v}_h \cdot \mathbf{n} \rangle = (\mathbf{y}, \mathbf{v}_h) \quad (3.1)$$

for all $\mathbf{v}_h \in \mathbf{V}_h$. Then, there exists a positive constant C independent of h such that for $p \in [2, \infty)$

$$\|w_h\|_{L^p} \leq C \left(\|\mathbf{z}\| + \|w_h - \widehat{w}_h\|_+ + \|\mathbf{y}\| \right),$$

and for $p = \infty$

$$\|w_h\|_{L^\infty} \leq C \log \frac{1}{h} \left(\|\mathbf{z}\| + \|w_h - \widehat{w}_h\|_+ + \|\mathbf{y}\| \right),$$

Proof. For $g \in L^q(\Omega)$, $\frac{1}{p} + \frac{1}{q} = 1$, let $\psi \in W^{2,q}(\Omega)$ be a solution of the elliptic problem

$$-\Delta \psi = g \text{ in } \Omega \text{ and } \psi = 0 \text{ on } \partial\Omega$$

satisfying the regularity

$$\|\psi\|_{W^{2,q}} \leq C \|g\|_{L^q}.$$

With $\Psi = \nabla \psi$, and using the commutative property of HDG projection (from [11]), it follows that

$$\begin{aligned} (w_h, g) &= -(w_h, \nabla \cdot \Psi) = -(w_h, \nabla \cdot \mathbf{\Pi}_V \Psi) + \langle w_h, \tau(\mathbf{\Pi}_W \psi - \psi) \rangle \\ &= (\mathbf{z}, \mathbf{\Pi}_V \Psi) - \langle \widehat{w}_h, \mathbf{\Pi}_V \Psi \cdot \mathbf{n} \rangle + \langle w_h, \tau(\mathbf{\Pi}_W \psi - \psi) \rangle - (\mathbf{y}, \mathbf{\Pi}_V \Psi) \quad [\text{using (3.1)}] \\ &= (\mathbf{z}, \mathbf{\Pi}_V \Psi) + \langle \tau(w_h - \widehat{w}_h), \mathbf{\Pi}_W \psi - \psi \rangle - \langle \widehat{w}_h, \mathbf{\Pi}_V \Psi \cdot \mathbf{n} - \tau(\mathbf{\Pi}_W \psi - \psi) \rangle - (\mathbf{y}, \mathbf{\Pi}_V \Psi) \\ &= (\mathbf{z}, \mathbf{\Pi}_V \Psi) + \langle \tau(w_h - \widehat{w}_h), \mathbf{\Pi}_W \psi - \psi \rangle - \langle \widehat{w}_h, \Psi \cdot \mathbf{n} \rangle - (\mathbf{y}, \mathbf{\Pi}_V \Psi). \quad [\text{using (2.4c)}] \end{aligned}$$

As \widehat{w}_h is single valued, $\Psi \in H(\text{div}, \Omega)$ and $\widehat{w}_h = 0$ on $\partial\Omega$, $\langle \widehat{w}_h, \Psi \cdot \mathbf{n} \rangle = 0$. Thus, an application of Theorem 2.1 implies

$$\begin{aligned} (w_h, g) &= (\mathbf{z}, \mathbf{\Pi}_V \Psi) + \langle \tau(w_h - \widehat{w}_h), \mathbf{\Pi}_W \psi - \psi \rangle - (\mathbf{y}, \mathbf{\Pi}_V \Psi) \\ &\leq \|\mathbf{z}\| \|\Psi\|_1 + C (h_k^{3/2} \tau^{-1/2}) \|w_h - \widehat{w}_h\|_+ \|\psi\|_2 + \|\mathbf{y}\| \|\Psi\|_1 \\ &\leq C (\|\mathbf{z}\| + \|w_h - \widehat{w}_h\|_+ + \|\mathbf{y}\|) \|\psi\|_{2,q} \\ &\leq C (\|\mathbf{z}\| + \|w_h - \widehat{w}_h\|_+ + \|\mathbf{y}\|) \|g\|_{L^q}. \end{aligned}$$

This implies

$$\|w_h\|_{L^p} = \sup_{\substack{g \in L^q \\ g \neq 0}} \frac{(w_h, g)}{\|g\|_{L^q}} \leq C (\|\mathbf{z}\| + \|w_h - \widehat{w}_h\|_+ + \|\mathbf{y}\|).$$

Now a use of Sobolev inequality when the dimension $d = 2$ (see, [19, page 47])

$$\|w_h\|_{L^\infty} \leq C \log \frac{1}{h} \|w_h\|_{L^p}$$

now completes the rest of the proof. \square

3.2 *A priori* Error Estimates

This subsection deals with *a priori* error estimates of the semi-discrete method. Decompose, $\mathbf{e}_u = \theta_u - \vartheta_u$, $\mathbf{e}_q = \theta_q - \vartheta_q$, $\widehat{\mathbf{e}}_u = \widehat{\theta}_u - \widehat{\vartheta}_u$, where $\theta_u = \Pi_W u - u_h$, $\theta_q = \Pi_{\mathbf{V}} \mathbf{q} - \mathbf{q}_h$, $\widehat{\theta}_u = \mathcal{P}u - \widehat{u}_h$. Since the estimates of ϑ_u , ϑ_q , $\widehat{\vartheta}_u$ are known, it is enough to estimate θ_u , θ_q and $\widehat{\theta}_u$. Using the HDG projections ((2.4a)-(2.4c)) and ((2.1a)-(2.1c)), equations in θ_u , θ_q and $\widehat{\theta}_u$ are written for all $(\mathbf{v}_h, w_h, \mu_1, \mu_2) \in \mathbf{V}_h \times W_h \times M_h \times M_h$ as

$$(\theta_q, \mathbf{v}_h) + (\theta_u, \nabla \cdot \mathbf{v}_h) - \langle \widehat{\theta}_u, \mathbf{v}_h \cdot \mathbf{n} \rangle = (\vartheta_q, \mathbf{v}_h), \quad (3.2a)$$

$$(\theta_{u_{tt}}, w_h) + (\theta_q, \nabla w_h) - \langle \widehat{\theta}_q \cdot \mathbf{n}, w_h \rangle + (f(u) - f(u_h), w_h) = (\vartheta_{u_{tt}}, w_h), \quad (3.2b)$$

$$\langle \widehat{\theta}_q \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h \setminus \Gamma_\partial} = 0, \quad (3.2c)$$

$$\langle \widehat{\theta}_u, \mu_2 \rangle_{\Gamma_\partial} = 0, \quad (3.2d)$$

where the numerical trace for flux is defined on \mathcal{E}_h as

$$\widehat{\theta}_q \cdot \mathbf{n} = \theta_q \cdot \mathbf{n} - \tau(\theta_u - \widehat{\theta}_u). \quad (3.3)$$

Following [8], we now define initial approximations via the HDG formulation of the associated elliptic equation at $t = 0$, i.e., seeking $(u_h(0), \mathbf{q}_h(0), \widehat{u}_h(0)) \in W_h \times \mathbf{V}_h \times M_h$ as the solution of

$$(\mathbf{q}_h(0), \mathbf{v}_h) + (u_h(0), \nabla \cdot \mathbf{v}_h) - \langle \widehat{u}_h(0), \mathbf{v}_h \cdot \mathbf{n} \rangle = 0 \quad \forall \mathbf{v}_h \in \mathbf{V}_h, \quad (3.4a)$$

$$(\mathbf{q}_h(0), \nabla w_h) - \langle \widehat{\mathbf{q}}_h(0) \cdot \mathbf{n}, w_h \rangle = (\Delta u_0, w_h) \quad \forall w_h \in W_h, \quad (3.4b)$$

$$\langle \widehat{\mathbf{q}}_h(0) \cdot \mathbf{n}, \mu \rangle_{\mathcal{E}_h \setminus \Gamma_\partial} = 0 \quad \forall \mu \in M_h, \quad (3.4c)$$

$$\langle \widehat{u}_h(0), \mu \rangle_{\Gamma_\partial} = 0 \quad \forall \mu \in M_h, \quad (3.4d)$$

$$\widehat{\mathbf{q}}_h(0) \cdot \mathbf{n} = \mathbf{q}_h(0) \cdot \mathbf{n} - \tau(u_h(0) - \widehat{u}_h(0)) \quad \text{on } \mathcal{E}_h. \quad (3.4e)$$

Moreover, the initial approximation $u_{ht}(0)$ of the initial velocity is obtained by differentiating the HDG projection (2.4) concerning time and evaluating at $t = 0$ using $(\Pi_{\mathbf{V}}(\nabla u_1), \Pi_W u_1)$ of $(\nabla u_1, u_1)$.

For our subsequent analysis, we use the following notation. For a Banach space X with norm $\|\cdot\|_X$, we define $W^{m,p}(0, T; X)$, $1 \leq p \leq \infty$ as the space of all strongly measurable and Bochner integrable functions $\phi : (0, T) \rightarrow X$ such that the corresponding norm $\|\phi\|_{W^{m,p}(0, T; X)}$ is finite, that is, for $p \in [1, \infty)$

$$\|\phi\|_{W^{m,p}(0, T; X)} := \left(\sum_{j=0}^m \int_0^T \left\| \frac{d^j \phi}{dt^j}(s) \right\|_X^p ds \right)^{1/p}$$

and for $p = \infty$

$$\|\phi\|_{W^{m,\infty}(0, T; X)} := \max_{0 \leq j \leq m} \operatorname{ess\,sup}_{0 < t < T} \left\| \frac{d^j \phi}{dt^j}(t) \right\|_X.$$

The following Lemma is useful for our subsequent use.

Lemma 3.2. Let u_{h0} be defined by HDG elliptic problem (3.4) and $u_{h1} = \Pi_W u_1$. Then, there holds

$$\left(\|\theta_q(0)\|^2 + \|(\theta_u - \widehat{\theta}_u)(0)\|_+^2 \right)^{1/2} \leq C h^{k+1} \|u_0\|_{H^{k+2}}. \quad (3.5)$$

Moreover, an additional elliptic $H^2 \cap H_0^1$ -regularity result of the associated adjoint problem in a duality argument as in [11, Lemma 4.1] shows

$$\|\theta_u(0)\| \leq C h^{k+2} \|u_0\|_{H^{k+2}}. \quad (3.6)$$

Proof. Using HDG projections (2.4) at $t = 0$ and (3.4), we now arrive at error equations at $t = 0$ as

$$(\theta_q(0), \mathbf{v}_h) + (\theta_u(0), \nabla \cdot \mathbf{v}_h) - \langle \widehat{\theta}_u(0), \mathbf{v}_h \cdot \mathbf{n} \rangle = 0 \quad \forall \mathbf{v}_h \in \mathbf{V}_h, \quad (3.7a)$$

$$(\theta_q(0), \nabla w_h) - \langle \widehat{\theta}_q(0) \cdot \mathbf{n}, w_h \rangle = 0 \quad \forall w_h \in W_h, \quad (3.7b)$$

$$\langle \widehat{\theta}_q(0) \cdot \mathbf{n}, \mu \rangle_{\mathcal{E}_h \setminus \Gamma_\partial} = 0 \quad \forall \mu \in M_h, \quad (3.7c)$$

$$\langle \widehat{\theta}_u(0), \mu \rangle_{\Gamma_\partial} = 0 \quad \forall \mu \in M_h, \quad (3.7d)$$

$$\widehat{\theta}_q(0) \cdot \mathbf{n} = \theta_q(0) \cdot \mathbf{n} - \tau(\theta_u(0) - \widehat{\theta}_u(0)) \quad \text{on } \mathcal{E}_h. \quad (3.7e)$$

Now, a use of arguments in [11] (see also [8, Lemma 3.6], completes estimate (3.5). In order to prove (3.6), we appeal to the Aubin-Nitsche duality argument as in [11, Lemma 4.1, Theorem 4.1] to conclude the remaining part of the proof. \square

By using the semidiscrete formulation (3.2), below, we estimate $\|\theta_{u_t}\|$ and $\|\theta_{\mathbf{q}}\|$.

Lemma 3.3. With u_{h0} defined by HDG elliptic problem (3.4), $u_{h1} = \Pi_W u_1$, and $u \in W^{2,1}(0, T; H^{k+2})$, there exists a positive constant C independent of h such that

$$\left(\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 \right)^{1/2} \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{2,1}(0, T; H^{k+2})} \right). \quad (3.8)$$

Moreover, there holds for $p \in [2, \infty)$

$$\|\theta_u(t)\|_{L^p} \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{2,1}(0, T; H^{k+2})} \right), \quad (3.9)$$

and for $p = \infty$

$$\|\theta_u(t)\|_{L^\infty} \leq C \log \frac{1}{h} h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{2,1}(0, T; H^{k+2})} \right). \quad (3.10)$$

Proof. For estimating $\|\theta_{u_t}\|$, differentiate (3.2a) with respect to t and then, choose $\mathbf{v}_h := \theta_{\mathbf{q}}$ and $w_h := \theta_{u_t}$ in (3.2b) to obtain

$$\frac{1}{2} \frac{d}{dt} \left(\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 \right) + (f(u) - f(u_h), \theta_{u_t}) = (\vartheta_{\mathbf{q}t}, \theta_{\mathbf{q}}) + (\vartheta_{u_{tt}}, \theta_{u_t}). \quad (3.11)$$

With the help of the Cauchy-Schwarz inequality, it follows that

$$\frac{d}{dt} \left(\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 \right) \leq \left(\|f(u) - f(u_h)\| + \|\vartheta_{\mathbf{q}t}\| + \|\vartheta_{u_{tt}}\| \right) \left(\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 \right)^{1/2},$$

and hence,

$$\frac{d}{dt} \left(\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 \right)^{1/2} \leq \|f(u) - f(u_h)\| + \|\vartheta_{\mathbf{q}t}\| + \|\vartheta_{u_{tt}}\|. \quad (3.12)$$

With the help of HDG projection, we split the nonlinear terms into two parts. Since $f(u) = u^3 - u$, using the mean value theorem, for some $\lambda \in (0, 1)$ and the Cauchy-Schwarz inequality

$$\begin{aligned} \|f(u) - f(\Pi_W u)\| &\leq C \|1 + (\Pi_W u)^2 + u^2\|_{L^\infty} \|\vartheta_u\| \\ &\leq C \|\vartheta_u\|. \end{aligned} \quad (3.13)$$

Here, we have used boundedness of $\Pi_W u \in L^\infty(\Omega)$ as $u \in L^\infty(\Omega)$ is bounded and using an inverse estimate and Theorem-2.1. Now, a use of Holder's inequality with Lemma 3.1 shows

$$\begin{aligned} \|f(\Pi_W u) - f(u_h)\|^2 &\leq C \int_{\Omega} (1 + |\Pi_W u|^4 + |u_h|^4) |\theta_u|^2 ds \\ &\leq C \left((1 + \|\Pi_W u\|_{L^6}^4 + \|u_h\|_{L^6}^4) \|\theta_u\|_{L^6}^2 \right) \\ &\leq C \left((1 + \|\mathbf{q}_h\|^4 + \|u_h - \widehat{u}_h\|_+^4) (\|\theta_{\mathbf{q}}\|^2 + \|\theta_h - \widehat{\theta}_h\|_+ + \|\vartheta_{\mathbf{q}}\|^2) \right). \end{aligned}$$

Now, a direct consequence of Corollary 2.1 shows that $1 + \|\mathbf{q}_h\|^4 + \|u_h - \widehat{u}_h\|_+^4 \leq C$. Hence,

$$\|f(\Pi_W u) - f(u_h)\|^2 \leq C (\|\theta_{\mathbf{q}}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 + \|\vartheta_{\mathbf{q}}\|^2). \quad (3.14)$$

On combining (3.13) and (3.14), there holds

$$\|f(u) - f(u_h)\|^2 \leq C (\|\vartheta_u\|^2 + \|\theta_{\mathbf{q}}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 + \|\vartheta_{\mathbf{q}}\|^2). \quad (3.15)$$

A use of inequality (3.15) in (3.12) yields

$$\begin{aligned} \frac{d}{dt} \left(\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 \right)^{1/2} &\leq C (\|\vartheta_u\| + \|\vartheta_{\mathbf{q}}\| + \|\vartheta_{\mathbf{q}t}\| + \|\vartheta_{u_{tt}}\| \\ &\quad + \|\theta_{\mathbf{q}}\| + \|\theta_u - \widehat{\theta}_u\|_+). \end{aligned}$$

After integrating the above inequality and noting that $\theta_{u_t}(0) = 0$, we use Grönwall's lemma to arrive at

$$\begin{aligned} \left(\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_+^2 \right)^{1/2} &\leq \left(\left(\|\theta_{\mathbf{q}}(0)\|^2 + \|(\theta_u - \widehat{\theta}_u)(0)\|_+^2 \right)^{1/2} \right. \\ &\quad \left. + \int_0^t (\|\vartheta_u\| + \|\vartheta_{\mathbf{q}}\| + \|\vartheta_{\mathbf{q}t}\| + \|\vartheta_{u_{tt}}\|) ds \right) \exp(Ct). \end{aligned} \quad (3.16)$$

A use of Lemma 2.1 with Lemma 3.2 completes the estimate (3.8).

A use of Lemma 3.1 for the error equation (3.2a) shows that for $p \in [2, \infty)$

$$\|\theta_u(t)\|_{L^p} \leq C \left(\|\theta_{\mathbf{q}}\| + \|\theta_u - \widehat{\theta}_u\|_+ + \|\vartheta_{\mathbf{q}}\| \right), \quad (3.17)$$

and for $p = \infty$

$$\|\theta_u(t)\|_{L^\infty} \leq C \log \frac{1}{h} \left(\|\theta_{\mathbf{q}}\| + \|\theta_u - \widehat{\theta}_u\|_+ + \|\vartheta_{\mathbf{q}}\| \right). \quad (3.18)$$

Then, a use of (3.8) completes the rest of the proof. \square

Lemma 3.4. Under the assumptions in Lemma 3.3 and with additional assumption $u_{ttt} \in L^1(0, T; H^{k+2})$, there holds for a positive constant C independent of h :

$$\left(\|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_{tt}}\|^2 + \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_+^2 \right)^{1/2} \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{W^{3,1}(0, T; H^{k+2})} \right). \quad (3.19)$$

Proof. In order to prove (3.19), now differentiate (3.2a) twice with respect to time and differentiate (3.2b)-(3.2d) and then choose $\mathbf{v}_h := \theta_{\mathbf{q}_t}$ and $w_h := \theta_{u_{tt}}$ to obtain

$$\frac{1}{2} \frac{d}{dt} \left(\|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_{tt}}\|^2 + \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_+^2 \right) + (f_t(u) - f_t(u_h), \theta_{u_{tt}}) = (\vartheta_{\mathbf{q}_{tt}}, \theta_{\mathbf{q}_t}) + (\vartheta_{u_{ttt}}, \theta_{u_{tt}}).$$

A use of the Cauchy-Schwarz inequality shows

$$\frac{d}{dt} \left(\|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_{tt}}\|^2 + \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_+^2 \right)^{1/2} \leq \left(\|f_t(u) - f_t(u_h)\| + \|(\vartheta_{\mathbf{q}_{tt}})\| + \|(\vartheta_{u_{ttt}})\| \right)$$

Integrating from 0 to t for $t \in (0, T]$, we arrive at

$$\begin{aligned} \left(\|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_{tt}}\|^2 + \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_+^2 \right)^{1/2} &\leq \left(\|\theta_{\mathbf{q}_t}(0)\|^2 + \|\theta_{u_{tt}}(0)\|^2 + \|(\theta_{u_t} - \widehat{\theta}_{u_t})(0)\|_+^2 \right)^{1/2} \\ &+ C \int_0^t \left(\|\vartheta_{\mathbf{q}_{tt}}\| + \|\vartheta_{u_{ttt}}\| + \|f_t(u) - f_t(u_h)\| \right) ds. \end{aligned} \quad (3.20)$$

Since $f(u) = u^3 - u$, there holds

$$\begin{aligned} f_t(u) - f_t(\Pi_W u) &= 3(u^2 u_t - (\Pi_W u)^2 \Pi_W u_t) - (u_t - \Pi_W u_t) \\ &= (3u^2 - 1)(u_t - \Pi_W u_t) + 3\Pi_W u_t(u^2 - (\Pi_W u)^2) \end{aligned}$$

and using the boundedness of $\Pi_W u$ and $\Pi_W u_t$ in $L^\infty(\Omega)$, it follows that

$$\begin{aligned} \|f_t(u) - f_t(\Pi_W u)\| &\leq \|3u^2 + 1\|_{L^\infty} \|u_t - \Pi_W u_t\| + \|3\Pi_W u_t\|_{L^\infty} \|u + \Pi_W u\|_{L^\infty} \|u - \Pi_W u\| \\ &\leq C \left(\|\vartheta_{u_t}\| + \|\vartheta_u\| \right). \end{aligned} \quad (3.21)$$

Similarly, we arrive with replacing u_{ht} by $\theta_{u_t} + \Pi_W u_t$ at

$$\begin{aligned} \|f_t(\Pi_W u) - f_t(u_h)\| &\leq \|(3(\Pi_W u)^2 - 1)\theta_{u_t}\| + 3\|(\theta_{u_t} + \Pi_W u_t)(\Pi_W u + u_h)\theta_u\| \\ &\leq \|3(\Pi_W u)^2 + 1\|_{L^\infty} \|\theta_{u_t}\| + 3\|\Pi_W u + u_h\|_{L^\infty} (\|\theta_{u_t}\| \|\theta_u\|_{L^\infty} + \|\Pi_W u_t\|_{L^\infty} \|\theta_u\|) \\ &\leq C \left((1 + \|\theta_u\|_{L^\infty}) \|\theta_{u_t}\| + \|\theta_u\| \right). \end{aligned} \quad (3.22)$$

Therefore, adding (3.21) with (3.22), it follows using Lemma 3.1 and inequality (3.18) that

$$\begin{aligned} \|f_t(u) - f_t(u_h)\| &\leq C \left(1 + \log \frac{1}{h} \left(\|\theta_{\mathbf{q}}\| + \|\theta_u - \widehat{\theta}_u\|_+ + \|\vartheta_{\mathbf{q}}\| \right) \right) \int_0^t \left(\|\theta_{\mathbf{q}}\| + \|\theta_u - \widehat{\theta}_u\|_+ + \|\vartheta_{\mathbf{q}}\| \right) ds \\ &+ C \left(\|\theta_{\mathbf{q}}\| + \|\theta_u - \widehat{\theta}_u\|_+ + \|\vartheta_{\mathbf{q}}\| \right) + \|\vartheta_{u_t}\| + \|\vartheta_u\|. \end{aligned} \quad (3.23)$$

Now, it remains to estimate $\|\theta_{u_{tt}}(0)\|$. We rewrite (3.2b) at $t = 0$ using (3.7b) to obtain

$$(\theta_{u_{tt}}(0), w_h) + (f(u)(0) - f(u_h(0)), w_h) = (\vartheta_{u_{tt}}(0), w_h), \quad (3.24)$$

Note that

$$\|(f(u) - f(u_h))(0)\|^2 \leq C (\|\vartheta_u(0)\|^2 + \|\vartheta_{\mathbf{q}}(0)\|^2). \quad (3.25)$$

Finally, set $w_h = \theta_{u_{tt}}(0)$ in (3.24), and use the bound (3.25) to arrive at

$$\|\theta_{u_{tt}}(0)\| \leq C(\|\vartheta_u(0)\| + \|\vartheta_{\mathbf{q}}(0)\|) + \|\vartheta_{u_{tt}}(0)\|, \quad (3.26)$$

Similarly, for $\theta_{\mathbf{q}_t}(0)$, differentiating the first error equation with respect to time then write the resulting equation at $t = 0$ and set $\mathbf{v}_h = \theta_{\mathbf{q}_t}(0)$, to obtain

$$(\theta_{\mathbf{q}_t}(0), \theta_{\mathbf{q}_t}(0)) + (\theta_{u_t}(0), \nabla \cdot \theta_{\mathbf{q}_t}(0)) - \langle \widehat{\theta}_{u_t}(0), \theta_{\mathbf{q}_t}(0) \cdot \mathbf{n} \rangle = (\vartheta_{\mathbf{q}_t}(0), \theta_{\mathbf{q}_t}(0)). \quad (3.27)$$

Here $(\theta_{u_t}(0), \nabla \cdot \theta_{\mathbf{q}_t}(0))$ is zero, as we choose $u_{ht}(0) = \Pi_W u_1$. Now, a use of (3.7c) with (3.7d) and (3.7e) at $t = 0$ in (3.27) shows after simplification

$$\|\theta_{\mathbf{q}_t}(0)\|^2 + \|(\theta_{u_t} - \widehat{\theta}_{u_t})(0)\|_+^2 \leq C \|\vartheta_{\mathbf{q}_t}(0)\|^2. \quad (3.28)$$

On combining (3.21), (3.23), (3.26) and (3.28) in (3.20), an application of Lemmas 2.1 and 3.3 completes the rest of the proof. \square

Theorem 3.1. Let $u \in L^\infty(0, T; H^{k+2})$, $u_t \in L^\infty(0, T; H^{k+2})$ and $u_{tt} \in L^1(0, T; H^{k+2})$. With assumptions in Lemma 3.3, the following estimates holds true for all $h = \max_{K \in \mathcal{T}_h} h_K$ and for all $t \in (0, T]$:

$$\|(u_t - u_{ht})(t)\| + \|(\mathbf{q} - \mathbf{q}_h)(t)\| \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{2,1}(0,T;H^{k+2})} + \|u\|_{W^{1,2}(0,T;H^{k+2})} \right). \quad (3.29)$$

If in addition $u_{tt} \in L^\infty(0, T; H^{k+2})$ and $u_{ttt} \in L^2(0, T; H^{k+2}(\Omega))$, then

$$\|(u_{tt} - u_{htt})(t)\| + \|(\mathbf{q}_t - \mathbf{q}_{ht})(t)\| \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{W^{3,1}(0,T;H^{k+2})} + \|u\|_{W^{2,2}(0,T;H^{k+2})} \right). \quad (3.30)$$

Proof. With the help of HDG projections and triangle inequality, we write

$$\|(\mathbf{q} - \mathbf{q}_h)(t)\| \leq \|(\mathbf{q} - \Pi_{\mathbf{V}} \mathbf{q})(t)\| + \|(\Pi_{\mathbf{V}} \mathbf{q} - \mathbf{q}_h)(t)\|.$$

Now, from Theorem-2.1 and Lemma-3.3, the result follows, completing the rest of the proof. \square

As a consequence of (3.29) and using the fact that $\phi(t) = \phi(0) + \int_0^t \phi(s) ds$, we obtain

$$\|(u - u_h)(t)\| \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{2,1}(0,T;H^{k+2})} + \|u\|_{W^{1,2}(0,T;H^{k+2})} \right), \quad (3.31)$$

under higher regularity : $u \in L^\infty(0, T; H^{k+2})$, $u_t \in L^\infty(0, T; H^{k+2})$ and $u_{tt} \in L^2(0, T; H^{k+2})$.

Below, we discuss using a variant of Baker's nonstandard energy technique, see [3] and [29] a direct proof of the optimal order of convergence in $L^\infty(0, T; L^2)$ norm with reduced regularity results on the exact solution.

For any $\tilde{\phi}$, define $\tilde{\phi}(s) = \int_0^s \phi(z) dz$ for $0 \leq s \leq t$.

Lemma 3.5. With u_{h0} defined by HDG elliptic problem (3.4), $u_{h1} = \Pi_W u_1$, let $u \in W^{1,2}(0, T; H^{k+2})$. Then, here exists a positive constant C independent of h such that

$$\left(\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\theta_u\|^2 + \|\tilde{\theta}_u - \widehat{\theta}_u\|_+^2 \right)^{1/2} \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{1,2}(0,T;H^{k+2})} \right).$$

Proof. Integrating (3.2b) from 0 to s and choosing $w_h := \theta_u$ and $\mathbf{v}_h := \tilde{\theta}_{\mathbf{q}}$ in (3.2a) and adding, we obtain

$$(\theta_{u_s}, \theta_u) + (\theta_{\mathbf{q}}, \tilde{\theta}_{\mathbf{q}}) + \langle \tau(\tilde{\theta}_u - \widehat{\theta}_u), \theta_u - \widehat{\theta}_u \rangle = (\vartheta_{\mathbf{q}}, \tilde{\theta}_{\mathbf{q}}) + (\vartheta_{u_s}, \theta_u) - \left(\int_0^s (f(u) - f(u_h)) dz, \theta_u \right). \quad (3.32)$$

Now, rewrite (3.32) as

$$\frac{1}{2} \frac{d}{ds} \left(\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\theta_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2 \right) = (\vartheta_{\mathbf{q}}, \tilde{\theta}_{\mathbf{q}}) + (\vartheta_{u_s}, \theta_u) - \left(\int_0^s (f(u) - f(u_h)) dz, \theta_u \right). \quad (3.33)$$

A use of the Cauchy-Schwarz and Young's inequalities in (3.33), we arrive at

$$\begin{aligned} \frac{1}{2} \frac{d}{ds} \left(\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\theta_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2 \right) &\leq C \left(\|\vartheta_{\mathbf{q}}\|^2 + \|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\vartheta_{u_s}\|^2 + \|\theta_u\|^2 \right. \\ &\quad \left. + \left\| \int_0^s (f(u) - f(u_h)) dz \right\|^2 \right). \end{aligned} \quad (3.34)$$

Now, with the help of HDG projection and equation (3.13), we obtain

$$\left\| \int_0^s (f(u) - f(\Pi_W u)) dz \right\|^2 \leq C \int_0^s (\|1 + (\Pi_W u)^2 + u^2\|_{L^\infty}^2 \|\vartheta_u\|^2) dz. \quad (3.35)$$

Also, using the mean value theorem, for some $\lambda \in (0, 1)$,

$$\begin{aligned} f(\Pi_W u) - f(u_h) &= (3((1-\lambda)\Pi_W u + \lambda u_h)^2 - 1)(u_h - \Pi_W u), \\ &= (3((1-\lambda)\Pi_W u + \lambda u_h)^2 - 1) \frac{d}{ds} \tilde{\theta}_u, \\ &= \frac{d}{ds} ((3((1-\lambda)\Pi_W u + \lambda u_h)^2 - 1) \tilde{\theta}_u) \\ &\quad - 6((1-\lambda)\Pi_W u + \lambda u_h)((1-\lambda)\Pi_W u_s + \lambda u_{hs}) \tilde{\theta}_u. \end{aligned}$$

Hence,

$$\begin{aligned} \left\| \int_0^s (f(\Pi_W u) - f(u_h)) dz \right\|^2 &\leq C \left((1 + \|\Pi_W u\|_{L^6}^4 + \|u_h\|_{L^6}^4) \|\tilde{\theta}_u\|_{L^6}^2 \right. \\ &\quad \left. + \int_0^s (1 + \|\Pi_W u\|_{L^6}^2 + \|u_h\|_{L^6}^2) (1 + \|\Pi_W u_s\|_{L^6}^2 + \|u_{hs}\|_{L^6}^2) \|\tilde{\theta}_u\|_{L^6}^2 dz \right). \end{aligned}$$

If we consider the functions $u, u_s \in L^\infty(\Omega) \subset L^6(\Omega)$ and apply the L^p estimates from Lemma-3.1 and use the second bound, derived from Theorem-3.1, we obtain

$$\begin{aligned} \left\| \int_0^s (f(\Pi_W u) - f(u_h)) dz \right\|^2 &\leq C \left(\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\tilde{\vartheta}_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2 \right. \\ &\quad \left. + \int_0^s (\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\tilde{\vartheta}_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2) dz \right). \end{aligned} \quad (3.36)$$

Combine expression (3.35) and (3.36), and then put in the (3.34) to arrive at

$$\begin{aligned} \frac{d}{ds} \left(\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\theta_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2 \right) &\leq C \left(\|\vartheta_{\mathbf{q}}\|^2 + \|\vartheta_{u_s}\|^2 + \|\theta_u\|^2 + \|\vartheta_u\|^2 + \|\tilde{\theta}_{\mathbf{q}}\|^2 \right. \\ &\quad \left. + \|\tilde{\vartheta}_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2 + \int_0^s (\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\tilde{\vartheta}_u\|^2 \right. \\ &\quad \left. + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2) dz \right). \end{aligned} \quad (3.37)$$

Integration of the above-resulting equation (3.37) yields

$$\begin{aligned} \|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\theta_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2 &\leq C \left(\|\theta_u(0)\|^2 + \int_0^t (\|\vartheta_{\mathbf{q}}\|^2 + \|\vartheta_{u_s}\|^2 + \|\theta_u\|^2 + \|\vartheta_u\|^2 \right. \\ &\quad \left. + \|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\tilde{\vartheta}_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2) ds + \int_0^t \int_0^s (\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\tilde{\vartheta}_u\|^2 + \|\tilde{\theta}_u - \tilde{\theta}_u\|_+^2) dz ds \right). \end{aligned} \quad (3.38)$$

Solving the double integral term in (3.38), we can write it,

$$\begin{aligned} \|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\theta_u\|^2 + \|\tilde{\theta}_u - \tilde{\tilde{\theta}}_u\|_+^2 &\leq C(1+t) \left(\|\theta_u(0)\|^2 + \int_0^t \left(\|\vartheta_{\mathbf{q}}\|^2 + \|\vartheta_{u_s}\|^2 + \|\vartheta_u\|^2 \right) ds \right) \\ &+ C(1+t) \int_0^t \left(\|\tilde{\theta}_{\mathbf{q}}\|^2 + \|\theta_u\|^2 + \|\tilde{\theta}_u - \tilde{\tilde{\theta}}_u\|_+^2 \right) ds. \end{aligned}$$

Finally, an application of the Grönwall's lemma with projection estimates from Lemma 2.1 completes the rest of the proof. \square

Theorem 3.2. With u_{h0} defined by HDG elliptic problem (3.4), $u_{h1} = \Pi_W u_1$, let $u \in L^\infty(0, T; H^{k+2})$ and $u_t \in L^2(0, T; H^{k+1})$. Then, there exists a positive constant C independent of h such that for all $h = \max_{K \in \mathcal{T}_h} h_K$ and for all $t \in (0, T]$, the following estimate holds:

$$\|(u - u_h)(t)\| + \|(\tilde{\mathbf{q}} - \tilde{\mathbf{q}}_h)(t)\| \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u\|_{L^\infty(0, T; H^{k+2})} + \|u_t\|_{L^2(0, T; H^{k+2})} \right).$$

Proof. With the help of HDG projections, we can write $\|u - u_h\| \leq \|u - \Pi_W u\| + \|\Pi_W u - u_h\|$. Now from Theorem-2.1 and Lemma-3.5, the result follows, and this completes the proof. \square

4 Post-processing

This section deals with local post-processing of the semi-discrete approximation of the displacement variable, which provides a super-convergence result when compared with the exact solution.

For HDG approaches, Cockburn *et al.* devised element-by-element post-processing of the scalar variable for elliptic problems and the velocity variable in the Stokes problem in [12,14]. He also demonstrated how to use this local postprocessing to achieve a close approximation to the original scalar unknown that also converges with order $O(h^{k+2})$ for $k \geq 1$ for the acoustic wave problem [13]. Inspired by their approach, we seek u_h^* in the space

$$W_h = \{w \in L^2(\Omega) : w|_K \in \mathcal{P}_{k+1}(K) \forall K \in \mathcal{T}_h\},$$

such that, in each element $K \in \mathcal{T}_h$, it satisfies:

$$\left. \begin{aligned} (\nabla u_h^*, \nabla w) &= (\mathbf{q}_h, \nabla w) \quad \forall w \in \mathcal{P}_{k+1}(K), \\ (u_h^*, 1) &= (u_h, 1). \end{aligned} \right\} \quad (4.1)$$

4.1 The BDM Projection

For our subsequent analysis, we define the BDM projection, introduced earlier by Brezzi, Douglas, and Marini [4] and denoted by $\mathbf{\Pi}^{BDM}$ is an operator that maps a vector-valued function $\boldsymbol{\varphi} \in \mathbf{H}^1(\Omega)$ onto BDM finite dimensional space \mathbf{V}_h satisfying:

$$(\mathbf{\Pi}^{BDM} \boldsymbol{\varphi}, v)_K = (\boldsymbol{\varphi}, v)_K, \quad \forall v \in [\mathcal{P}_k(K)]^d, \quad (4.2a)$$

$$(\mathbf{\Pi}^{BDM} \boldsymbol{\varphi}, p)_E = \langle \boldsymbol{\varphi}, p \rangle_E, \quad \forall p \in \mathcal{P}_k(E), \quad \forall E \in \partial K \quad (4.2b)$$

with property,

$$(\nabla \cdot \mathbf{\Pi}^{BDM} \boldsymbol{\varphi}, v)_K = (\nabla \cdot \boldsymbol{\varphi}, v)_K \quad \forall v \in \mathcal{P}_k(K), \quad (4.3)$$

and

$$\|\mathbf{\Pi}^{BDM} \boldsymbol{\varphi} - \boldsymbol{\varphi}\| \leq C h \|\boldsymbol{\varphi}\|_{H^1(K)} \quad \forall \boldsymbol{\varphi} \in H^1(K). \quad (4.4)$$

The main theorem of this section is given below.

Theorem 4.1. Under the assumptions of Theorem 3.1, there holds for $k \geq 1$

$$\|u - u_h^*\| \leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{W^{3,1}(0, T; H^{k+2})} + \|u\|_{W^{0,2}(0, T; H^{k+2})} \right).$$

To prove this theorem, we first prove a superconvergence of $P_{k-1}\theta_u$ with the help of the duality argument.

4.2 The dual problem

Now, we introduce a dual problem of the wave equation as follows. We define the function $\phi \in H_0^1(\Omega)$ to be the solution of the wave equation and for any given function $\Theta \in L^2(\Omega)$, ϕ satisfying

$$\phi_{ss}(s) - \Delta\phi(s) + f_u(u)\phi(s) = 0, \quad \text{in } \Omega \times [0, t], \quad (4.5a)$$

$$\phi = 0, \quad \text{on } \partial\Omega \times [0, t], \quad (4.5b)$$

$$\phi(t) = 0, \quad \text{in } \Omega, \quad (4.5c)$$

$$\phi_s(t) = \Theta, \quad \text{in } \Omega. \quad (4.5d)$$

Proposition 4.1. With $u \in L^\infty(\Omega)$, there is a positive constant C such that the following holds:

$$\|\phi\|_{L^\infty(0,t;L^2(\Omega))} + \|\phi\|_{L^\infty(0,t;H^1(\Omega))} + \|\phi_t\|_{L^\infty(0,t;L^2(\Omega))} \leq C\|\Theta\|. \quad (4.6)$$

Moreover,

$$\|\underline{\phi}\|_{L^\infty(0,t;H^2(\Omega))} \leq C\|\Theta\|, \quad (4.7)$$

where $\|\underline{\phi}(s)\| = \int_s^t \phi(z) dz$.

Proof. By change of variable $s \rightarrow t - s$ and with $\bar{\phi}(s) = \phi(t - s)$, the problem (4.5) is transformed into a forward problem in $\bar{\phi}$.

$$\bar{\phi}_{ss}(s) - \Delta\bar{\phi}(s) + f_u(\bar{u})\bar{\phi}(s) = 0, \quad \text{in } \Omega \times [0, t], \quad (4.8a)$$

$$\bar{\phi} = 0, \quad \text{on } \partial\Omega \times [0, t], \quad (4.8b)$$

$$\bar{\phi}(0) = 0, \quad \text{in } \Omega, \quad (4.8c)$$

$$\bar{\phi}_s(0) = -\Theta, \quad \text{in } \Omega. \quad (4.8d)$$

Now, the standard energy argument along with the boundedness of f_u and Grönwall's inequality, shows

$$\|\bar{\phi}\|_{L^\infty(0,t;L^2(\Omega))} + \|\bar{\phi}\|_{L^\infty(0,t;H^1(\Omega))} + \|\bar{\phi}_t\|_{L^\infty(0,t;L^2(\Omega))} \leq C\|\Theta\|. \quad (4.9)$$

and on integrating concerning time, use of (4.9) yields in a straightforward way the estimate

$$\|\bar{\underline{\phi}}\|_{L^\infty(0,t;H^2(\Omega))} \leq C\|\Theta\|, \quad (4.10)$$

where $\|\bar{\underline{\phi}}(s)\| = \int_s^t \bar{\phi}(z) dz$. Then an application of inverse transformation $t - s \rightarrow s$ and with $\bar{\phi}(t - s) = \phi(s)$ completes the rest of the proof. \square

Next, in order to prove the Theorem 4.1, we need to find the estimate of $P_{k-1}\theta_u$

4.3 Estimate of $P_{k-1}\theta_u$

Lemma 4.1. Under the assumptions of Theorem 3.1, there holds for $k \geq 1$

$$\|P_{k-1}\theta_u(t)\| \leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{W^{3,1}(0,T;H^{k+2})} \right). \quad (4.11)$$

Proof. On multiplying $P_{k-1}\theta_u$ to (4.5a) and integrating over Ω , we note that

$$\int_0^t \frac{d}{ds} \left((\phi_s, P_{k-1}\theta_u) - (\phi, P_{k-1}\theta_{u_s}) \right) = \int_0^t \left(-(\phi, P_{k-1}\theta_{u_{ss}}) + (\Delta\phi, P_{k-1}\theta_u) - (f_u(u)\phi, P_{k-1}\theta_u) \right) ds.$$

Using equation (4.5d) from the dual problem, the property of projection and the property of BDM-projection (4.3), gives

$$\begin{aligned} (\Theta, P_{k-1}\theta_u) &= (\phi_s(0), P_{k-1}\theta_u(0)) - (\phi(0), P_{k-1}\theta_{u_s}(0)) + \int_0^t \left(-(\phi, P_{k-1}\theta_{u_{ss}}) \right. \\ &\quad \left. + (\nabla \cdot (\mathbf{\Pi}^{BDM} \nabla \phi), \theta_u) - (f_u(u)\phi, P_{k-1}\theta_u) \right) ds. \end{aligned}$$

Since, $u_{ht}(0) = \mathbf{\Pi} \mathbf{v} u_1$, the second term on the right-hand side become zero. For the second last term under the time integral, use of (3.2a) shows

$$(\nabla \cdot (\mathbf{\Pi}^{BDM} \nabla \phi), \theta_u) = (\mathbf{\Pi} \mathbf{v} \mathbf{q} - \mathbf{q}, \mathbf{\Pi}^{BDM} \nabla \phi) - (\theta_{\mathbf{q}}, \mathbf{\Pi}^{BDM} \nabla \phi) + \langle \widehat{\theta}_u, \mathbf{\Pi}^{BDM} \nabla \phi \cdot \mathbf{n} \rangle.$$

Here, the boundary term becomes zero as $\widehat{\theta}_u = 0$ on $\partial\Omega$ and $\mathbf{\Pi}^{BDM} \nabla \phi \in H(\text{div}, \Omega)$. Thus, we arrive at

$$\begin{aligned} (\Theta, P_{k-1} \theta_u) &= (\phi_s(0), P_{k-1} \theta_u(0)) + \int_0^t \left(-(\phi, P_{k-1} \theta_{u_{ss}}) + (\mathbf{q}_h - \mathbf{q}, \mathbf{\Pi}^{BDM} \nabla \phi) - (f_u(u) \phi, P_{k-1} \theta_u) \right) ds, \\ &= (\phi_s(0), P_{k-1} \theta_u(0)) + \int_0^t \left(-(P_{k-1} \phi, \theta_{u_{ss}}) + (\mathbf{q}_h - \mathbf{q}, \mathbf{\Pi}^{BDM} \nabla \phi - \nabla I_h \phi) \right) ds \\ &\quad - \int_0^t \left((\theta_{\mathbf{q}}, \nabla I_h \phi) - (f_u(u) \phi, P_{k-1} \theta_u) \right) ds, \end{aligned}$$

where I_h indicates the interpolation operator from $L^2(\Omega)$ into $W_h \cap H_0^1(\mathcal{T}_h)$. Using the second error equation (3.2b) for the second last term in the above equation, we arrive at

$$\begin{aligned} (\Theta, P_{k-1} \theta_u) &= (\phi_s(0), P_{k-1} \theta_u(0)) + \int_0^t \left(-(P_{k-1} \phi, \theta_{u_{ss}}) + (\mathbf{q}_h - \mathbf{q}, \mathbf{\Pi}^{BDM} \nabla \phi - \nabla I_h \phi) + (\theta_{u_{ss}}, I_h \phi) \right. \\ &\quad \left. + (f(u) - f(u_h), I_h \phi) - (\mathbf{\Pi}_W u_{ss} - u_{ss}, I_h \phi) - (f_u(u) \phi, P_{k-1} \theta_u) \right) ds. \end{aligned}$$

Here, as $\langle \widehat{\theta}_{\mathbf{q}}, \mu \rangle = 0$ on $\mathcal{E}_h / \Gamma_{\partial}$ and $I_h \phi = 0$ on $\partial\Omega$, we have used $\langle \widehat{\theta}_{\mathbf{q}} \cdot \mathbf{n}, I_h \phi \rangle = 0$,

$$\begin{aligned} (\Theta, P_{k-1} \theta_u) &= (\phi_s(0), P_{k-1} \theta_u(0)) + \int_0^t \left((\mathbf{q}_h - \mathbf{q}, \mathbf{\Pi}^{BDM} \nabla \phi - \nabla I_h \phi) + (u_{ss} - u_{hss}, I_h \phi - P_{k-1} \phi) \right) \\ &\quad + \int_0^t \left((f(u) - f(u_h), I_h \phi) - (f_u(u) \phi, P_{k-1} \theta_u) \right) ds. \end{aligned} \quad (4.12)$$

For the first term in integration, use the identity,

$$\int_0^t h(z) g(z) dz = h(0) \underline{g}(0) + \int_0^t h_z(z) \underline{g}(z) dz, \quad (4.13)$$

and utilize Taylor's series expansion to deal with the term $f(u) - f(u_h)$ on the right-hand side of (4.12) to arrive with $\mathbf{e}_{\mathbf{q}s} = \mathbf{q}_s - \mathbf{q}_{hs}$ at,

$$\begin{aligned} (\Theta, P_{k-1} \theta_u) &= (\phi_s(0), P_{k-1} \theta_u(0)) - (\mathbf{e}_{\mathbf{q}}(0), \mathbf{\Pi}^{BDM} \nabla \underline{\phi}(0) - \nabla I_h \underline{\phi}(0)) - \int_0^t \left((\mathbf{e}_{\mathbf{q}s}, \mathbf{\Pi}^{BDM} \nabla \underline{\phi} - \nabla I_h \underline{\phi}) \right. \\ &\quad \left. + (u_{ss} - u_{hss}, I_h \phi - P_{k-1} \phi) + (f_u(u)(u_h - u), I_h \phi - \phi) + (f_u(u)(u_h - u), \phi) \right. \\ &\quad \left. + (f_{uu}(u)(u_h - u)^2, I_h \phi) - (f_u(u) \phi, P_{k-1} \theta_u) \right) ds. \end{aligned} \quad (4.14)$$

A use of the property of P_{k-1} in (4.14) yields

$$\begin{aligned} (\Theta, P_{k-1} \theta_u) &= (\phi_s(0), P_{k-1} \theta_u(0)) - (\mathbf{e}_{\mathbf{q}}(0), \mathbf{\Pi}^{BDM} \nabla \underline{\phi}(0) - \nabla I_h \underline{\phi}(0)) - \int_0^t \left((\mathbf{e}_{\mathbf{q}s}, \mathbf{\Pi}^{BDM} \nabla \underline{\phi} - \nabla I_h \underline{\phi}) \right. \\ &\quad \left. + (u_{ss} - u_{hss}, I_h \phi - P_{k-1} \phi) + (f_u(u)(u_h - u), I_h \phi - \phi) + (f_{uu}(u)(u_h - u)^2, I_h \phi) \right. \\ &\quad \left. - (f_u(u) \phi - P_{k-1}(f_u(u) \phi), P_{k-1} \theta_u + \theta_u) + (f_u(u) \phi - P_{k-1}(f_u(u) \phi), \vartheta_u) \right) ds \\ &= J_0 + J_1 + \int_0^t (J_2 + J_3 + J_4 + J_5 + J_6 + J_7) ds. \end{aligned} \quad (4.15)$$

A straightforward applications of the Cauchy-Schwarz inequality with (3.5)-(3.6) elliptic regularity results yield

$$\begin{aligned} |J_0| &\leq \|\phi_s(0)\| \|\theta_u(0)\| \leq h^{k+2} \|u_0\|_{H^{k+2}} \|\phi_s(0)\| \\ |J_1| &\leq \|\mathbf{\Pi}^{BDM} \nabla \underline{\phi}(0) - \nabla I_h \underline{\phi}(0)\| \|\mathbf{q}_h(0) - \mathbf{q}(0)\| \leq C h^{k+2} \|\underline{\phi}(0)\|_{H^2}, \end{aligned}$$

A use of Theorem 3.1 with approximation property in BDM projection (4.4) and in interpolant shows

$$\begin{aligned} |J_2| &\leq \left(\|\mathbf{\Pi}^{BDM} \nabla \underline{\phi} - \nabla \underline{\phi}\| + \|\nabla(\underline{\phi} - I_h \underline{\phi})\| \right) \|\mathbf{q}_{ht} - \mathbf{q}_t\| \\ &\leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{W^{3,1}(0,T;H^{k+2})} \right) \|\underline{\phi}\|_{H^2}. \end{aligned}$$

Similarly an application of Theorem 3.1 with approximation property in BDM projection (4.4), L^2 projection and interpolant shows

$$\begin{aligned} |J_3| &\leq \|u_{tt} - u_{htt}\| (\|I_h \phi - \phi\| + \|\phi - P_{k-1} \phi\|) \leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{W^{3,1}(0,T;H^{k+2})} \right) \|\phi\|_{H^1}, \\ |J_4| &\leq \|u - u_h\| \|I_h \phi - \phi\| \leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{2,1}(0,T;H^{k+2})} \right) \|\phi\|_{H^1}, \\ |J_5| &\leq \|u - u_h\|^2 \|I_h \phi\|_{L^\infty} \leq C h^{2k+2} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{2,1}(0,T;H^{k+2})} \right) \|\phi\|_{H^1}, \\ |J_6 + J_7| &\leq C h \|\phi\|_{H^1} (\|\theta_u\| + \|\vartheta_u\|) \leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u\|_{W^{2,1}(0,T;H^{k+2})} \right) \|\phi\|_{H^1}. \end{aligned}$$

Substitute all above estimate J_0, \dots, J_7 in (4.15), a use of Proposition 4.1 yields

$$\begin{aligned} (\Theta, P_{k-1} \theta_u(t)) &\leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{W^{3,1}(0,T;H^{k+2})} \right) \left(\|\underline{\phi}(0)\|_{H^2} + \sup_{s \in (0,T)} \|\phi_s(s)\| \right. \\ &\quad \left. + \int_0^t (\|\phi\|_{H^1} + \|\underline{\phi}\|_{H^2}) ds \right), \\ &\leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{W^{3,1}(0,T;H^{k+2})} \right) \|\Theta\|. \end{aligned}$$

Since

$$\|P_{k-1} \theta_u(t)\| = \sup_{\Theta \in L^2(\Omega)} \frac{(\Theta, P_{k-1} \theta_u(t))}{\|\Theta\|},$$

this concludes the rest of the proof. \square

4.4 Proof of Theorem 4.1

For finding the estimate of $\|u - u_h^*\|$, we rewrite

$$\|u - u_h^*\|_K \leq \|u - P_{k+1} u\|_K + \|P_{k+1} u - u_h^*\|_K.$$

For $w \in H^1(K)$ and $k \geq 1$, we observe that

$$\|w\|_K \leq \|P_0 w\|_K + \|P_0 w - w\|_K \leq \|P_0 w\|_K + Ch \|\nabla w\|_K. \quad (4.16)$$

Now, set $w = P_{k+1} u - u_h^*$ and note that

$$\|P_0 w\|_K^2 = (P_{k+1} u - u_h^*, P_0 w)_K = (u - u_h^*, P_0 w)_K.$$

Now, with the help of the HDG projection, we rewrite

$$\|P_0 w\|_K^2 = (\theta_u, P_0 w)_K = (P_{k-1} \theta_u, P_0 w)_K \Rightarrow \|P_0 w\|_K \leq \|P_{k-1} \theta_u\|_K. \quad (4.17)$$

For any $v \in P_{k+1}(K)$, we note that

$$\begin{aligned} (\nabla w, \nabla v)_K &= (\nabla(P_{k+1} u - u_h^*), \nabla v)_K, \\ &= (\nabla(P_{k+1} u - u), \nabla v)_K + (\nabla(u - u_h^*), \nabla v)_K, \\ &= (\nabla(P_{k+1} u - u), \nabla v)_K + (\mathbf{q} - \mathbf{q}_h), \nabla v)_K, \end{aligned} \quad (4.18)$$

by the first equation in (4.1). Hence, substituting the estimates (4.17) and (4.18) in (4.16), we obtain that

$$\|u - u_h^*\|_K \leq \|u - P_{k+1} u\|_K + \|P_{k-1} \theta_u\|_K + Ch (\|\nabla(u - P_{k+1} u)\|_K + \|\mathbf{q} - \mathbf{q}_h\|_K).$$

Now squaring both sides and then using the estimate of $\|\mathbf{q} - \mathbf{q}_h\|_K$ from Theorem 3.1 and $\|P_{k-1} \theta_u\|_K$ from (4.11), we complete the proof of the Theorem 4.1.

5 Completely Discrete Schemes

This section focuses on conservative and non-conservative schemes with error analyses.

5.1 A Fully Discrete Conservative Scheme

Let Δt be the time step, and $t_n = n\Delta t$, where $1 \leq n \leq N$ and $N\Delta t = 1$. We now define some notations for our subsequent use:

$$\begin{aligned}\varphi^n &= \varphi(t_n), & \varphi^{n+1/2} &= \frac{1}{2}(\varphi^{n+1} + \varphi^n), & \partial_t^* \varphi^{1/2} &= \frac{(\varphi^1 - \varphi^0)}{\Delta t} \\ \bar{\partial}_t \varphi^n &= \frac{(\varphi^n - \varphi^{n-1})}{\Delta t}, & \partial_t \varphi^n &= \frac{(\varphi^{n+1} - \varphi^n)}{\Delta t}, & \delta_t \varphi^n &= \frac{\varphi^{n+1} - \varphi^{n-1}}{2\Delta t} \\ \mathcal{A}\varphi^n &= \frac{1}{2}(\varphi^{n+1} + \varphi^{n-1}), & \partial_t^2 \varphi &= \frac{1}{(\Delta t)^2}(\varphi^{n+1} - 2\varphi^n + \varphi^{n-1}) = \partial_t \bar{\partial}_t \varphi^n\end{aligned}$$

The HDG fully discretized formulation related to HDG method (2.1a)-(2.1e) reads as: Seek $(U^n, \mathbf{Q}^n, \widehat{U}^n) \in W_h \times \mathbf{V}_h \times M_h$ such that

$$\frac{2}{\Delta t}(\partial_t^* U^{1/2}, w_h) + (\mathbf{Q}^{1/2}, \nabla w_h) - \langle \widehat{\mathbf{Q}}^{1/2} \cdot \mathbf{n}, w_h \rangle + (\mathcal{F}(U^1, U^0), w_h) - \frac{2}{\Delta t}(u_1, w_h) = 0, \quad (5.1a)$$

$$(\partial_t^2 U^n, w_h) + (\mathcal{A}\mathbf{Q}^n, \nabla w_h) - \langle \widehat{\mathbf{Q}}^n \cdot \mathbf{n}, w_h \rangle + (\mathcal{F}(U^{n+1}, U^{n-1}), w_h) = 0, \quad \forall n \geq 1, \quad (5.1b)$$

$$(\mathbf{Q}^n, \mathbf{v}_h) + (U^n, \nabla \cdot \mathbf{v}_h) - \langle \widehat{U}^n, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0, \quad \forall n \geq 0, \quad (5.1c)$$

$$\langle \widehat{\mathbf{Q}}^n \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h/\Gamma_\partial} = 0, \quad \forall n \geq 0, \quad (5.1d)$$

$$\langle \widehat{U}^n, \mu_2 \rangle_{\Gamma_\partial} = 0, \quad \forall n \geq 0, \quad (5.1e)$$

for all $(w_h, \mathbf{v}_h, \mu_1, \mu_2) \in W_h \times \mathbf{V}_h \times M_h \times M_h$, where,

$$\widehat{\mathbf{Q}}^n \cdot \mathbf{n} = \mathbf{Q}^n \cdot \mathbf{n} - \tau(U^n - \widehat{U}^n) \quad \text{on } \mathcal{E}_h, \quad (5.2)$$

and the non-linear term is defined as:

$$\mathcal{F}(U^{n+1}, U^{n-1}) = \begin{cases} \frac{F(U^{n+1}) - F(U^{n-1})}{U^{n+1} - U^{n-1}}, & U^{n+1} \neq U^{n-1} \\ F'(U^{n+1}), & U^{n+1} = U^{n-1} \end{cases}$$

with $F(U) = \frac{1}{4}(1 - U^2)^2$. Given initial values U^0 , \mathbf{Q}^0 , and \widehat{U}^0 , we can determine U^1 , \mathbf{Q}^1 , and \widehat{U}^1 using equations (5.1a) and (5.1c).

5.1.1 Discrete energy conservative property.

Below, we establish the following discrete conservation law for the energy of the fully discrete scheme (5.1).

Theorem 5.1. The fully discrete HDG approximation $(U^{n+1}, \mathbf{Q}^{n+1}, \widehat{U}^{n+1}) \in W_h \times \mathbf{V}_h \times M_h$ given by (5.1b)-(5.1d) satisfies the following fully discrete energy conservation property for $n \geq 1$,

$$\mathbb{E}^{n+1/2} = \mathbb{E}^{3/2}, \quad (5.3)$$

where,

$$\begin{aligned}\mathbb{E}^{n+1/2} &= \|\partial_t U^n\|^2 + \frac{1}{2} \left(\|\mathbf{Q}^n\|^2 + \|\mathbf{Q}^{n+1}\|^2 + \|\tau^{1/2}(U^n - \widehat{U}^n)\|^2 + \|U^{n+1} - \widehat{U}^{n+1}\|_+^2 \right. \\ &\quad \left. + 2(F(U^n), 1) + 2(F(U^{n+1}), 1) \right).\end{aligned}$$

Proof. Taking the difference between two levels, i.e., between $t = t_{n+1}$ and $t = t_{n-1}$ and then dividing by $2\Delta t$ in (5.1c), we find that

$$(\delta_t \mathbf{Q}^n, \mathbf{v}_h) + (\delta_t U^n, \nabla \cdot \mathbf{v}_h) - \langle \delta_t \widehat{U}^n, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0 \quad (5.4)$$

Now, choose $\mathbf{v}_h = \mathcal{A}\mathbf{Q}^n$, and $w_h = \delta_t U^n$ in equations (5.1b) and (5.4) respectively, then add resulting equations

$$\begin{aligned} & (\partial_t^2 U^n, \delta_t U^n) + (\delta_t \mathbf{Q}^n, \mathcal{A}\mathbf{Q}^n) + (\delta_t U^n, \nabla \cdot \mathcal{A}\mathbf{Q}^n) - \langle \delta_t \widehat{U}^n, \mathcal{A}\mathbf{Q}^n \cdot \mathbf{n} \rangle + (\mathcal{A}\mathbf{Q}^n, \nabla \delta_t U^n) \\ & \quad - \langle \mathcal{A}\widehat{\mathbf{Q}}^n \cdot \mathbf{n}, \delta_t U^n \rangle + (\mathcal{F}(U^{n+1}, U^{n-1}), \delta_t U^n) = 0. \end{aligned}$$

An application of (5.1d) with (5.1e) and (5.2) yields

$$(\partial_t^2 U^n, \delta_t U^n) + (\delta_t \mathbf{Q}^n, \mathcal{A}\mathbf{Q}^n) + \tau \langle \delta_t U^n - \delta_t \widehat{U}^n, \mathcal{A}U^n - \mathcal{A}\widehat{U}^n \rangle + (\mathcal{F}(U^{n+1}, U^{n-1}), \delta_t U^n) = 0,$$

and hence, we arrive at

$$\begin{aligned} & \frac{1}{2\Delta t} \left(\|\partial_t U^n\|^2 - \|\partial_t U^{n-1}\|^2 + \frac{1}{2} \left(\|\mathbf{Q}^{n+1}\|^2 - \|\mathbf{Q}^{n-1}\|^2 + \|U^{n+1} - \widehat{U}^{n+1}\|_+^2 \right. \right. \\ & \quad \left. \left. - \|U^{n-1} - \widehat{U}^{n-1}\|_+^2 \right) + (\mathcal{F}(U^{n+1}, U^{n-1}), \delta_t U^n) \right) = 0. \end{aligned}$$

From the definition of δ_t , it follows that

$$\begin{aligned} (\mathcal{F}(U^{n+1}, U^{n-1}), \delta_t U^n) &= \left(\frac{F(U^{n+1}) - F(U^{n-1})}{U^{n+1} - U^{n-1}}, \frac{U^{n+1} - U^{n-1}}{2\Delta t} \right) \\ &= \frac{1}{2\Delta t} (F(U^{n+1}) - F(U^{n-1}), 1). \end{aligned}$$

Altogether, we obtain

$$\begin{aligned} \|\partial_t U^n\|^2 + \frac{1}{2} (\|\mathbf{Q}^{n+1}\|^2 + \|U^{n+1} - \widehat{U}^{n+1}\|_+^2) + (F(U^{n+1}), 1) &= \|\partial_t U^{n-1}\|^2 \\ &+ \frac{1}{2} (\|\mathbf{Q}^{n-1}\|^2 + \|U^{n-1} - \widehat{U}^{n-1}\|_+^2) + (F(U^{n-1}), 1). \end{aligned} \quad (5.5)$$

By adding

$$\frac{1}{2} \left(\|\mathbf{Q}^n\|^2 + \|U^n - \widehat{U}^n\|_+^2 + 2(F(U^n), 1) \right)$$

on both sides of (5.37c), we derive the energy conservation property:

$$E^{n+1/2} = E^{n-1/2} = \dots = E^{3/2}.$$

Solving equations (5.1a) and (5.1c) at $n = 0$ and add

$$\frac{1}{2} \left(\|\mathbf{Q}^0\|^2 + \|U^0 - \widehat{U}^0\|_+^2 + 2(F(U^0), 1) \right)$$

on both sides to arrive at

$$\begin{aligned} 2\|\partial_t U^0\|^2 + \frac{1}{2} \left(\|\mathbf{Q}^0\|^2 + \|\mathbf{Q}^1\|^2 + \|U^0 - \widehat{U}^0\|_+^2 + \|U^1 - \widehat{U}^1\|_+^2 \right) + (F(U^0), 1) \\ + (F(U^1), 1) = \|\mathbf{Q}^0\|^2 + \|U^0 - \widehat{U}^0\|_+^2 + 2(F(U^0), 1) + 2(u_1, \partial_t U^0). \end{aligned} \quad (5.6)$$

From (5.6), we obtain the relation

$$\mathbb{E}^{3/2} = -\|\partial_t U^0\|^2 + \|\mathbf{Q}^0\|^2 + \|U^0 - \widehat{U}^0\|_+^2 + 2(F(U^0), 1) + 2(u_1, \partial_t U^0).$$

With the initial energy at $n = 0$ given by

$$\mathcal{E}^0 = 2\|\partial_t U^0\|^2 + \frac{1}{2} \left(\|\mathbf{Q}^0\|^2 + \|\mathbf{Q}^1\|^2 + \|U^0 - \widehat{U}^0\|_+^2 + \|U^1 - \widehat{U}^1\|_+^2 \right) + (F(U^0), 1) + (F(U^1), 1),$$

it is observe that $\mathbb{E}^{3/2} \leq \mathcal{E}^0$ and this concludes the rest of the proof. \square

The next lemma helps us to deal with uniqueness and error estimates.

Lemma 5.1. For $\Phi_1, \Phi_2, \Psi_1, \Psi_2 \in L^6(\Omega)$, there holds

$$\|\mathcal{F}(\Phi_1, \Psi_1) - \mathcal{F}(\Phi_2, \Psi_2)\|^2 \leq C \left(\|\Phi_1 - \Phi_2\|_{L^6}^2 + \|\Psi_1 - \Psi_2\|_{L^6}^2 \right) \left(\|\Phi_1\|_{L^6}^2 + \|\Psi_1\|_{L^6}^2 + \|\Phi_2\|_{L^6}^2 + \|\Psi_2\|_{L^6}^2 + 1 \right).$$

Proof. From the definition of $\mathcal{F}(U, V)$ with $F(U) = \frac{1}{4}(1 - U^2)^2$, it follows that

$$\mathcal{F}(\Phi_1, \Psi_1) = \frac{1}{4}(\Phi_1 + \Psi_1)(\Phi_1^2 + \Psi_1^2 - 2),$$

and therefore,

$$\begin{aligned} \mathcal{F}(\Phi_1, \Psi_1) - \mathcal{F}(\Phi_2, \Psi_2) &= \frac{1}{4}(\Phi_1^3 - \Phi_2^3 + \Psi_1^3 - \Psi_2^3 + \Phi_1\Psi_1^2 - \Phi_2\Psi_2^2 + \Phi_1^2\Psi_1 - \Phi_2^2\Psi_2 \\ &\quad - 2(\Phi_1 - \Phi_2) - 2(\Psi_1 - \Psi_2)) \end{aligned}$$

Adding and subtracting $\Phi_2\Psi_1^2$ and $\Phi_1^2\Psi_2$, we obtain

$$\begin{aligned} \mathcal{F}(\Phi_1, \Psi_1) - \mathcal{F}(\Phi_2, \Psi_2) &= \frac{1}{4}\left((\Phi_1^3 - \Phi_2^3) + \Psi_1^2(\Phi_1 - \Phi_2) + \Psi_2(\Phi_1^2 - \Phi_2^2) - 2(\Phi_1 - \Phi_2) \right. \\ &\quad \left. + (\Psi_1^3 - \Psi_2^3) + \Phi_1^2(\Psi_1 - \Psi_2) + \Phi_2(\Psi_1^2 - \Psi_2^2) - 2(\Psi_1 - \Psi_2)\right) \end{aligned} \quad (5.7)$$

A use of the Cauchy-Schwarz inequality with the property of L^p bounds yields

$$\begin{aligned} \|\mathcal{F}(\Phi_1, \Psi_1) - \mathcal{F}(\Phi_2, \Psi_2)\|^2 &\leq \frac{1}{16}\left\|\left((\Phi_1 - \Phi_2) + (\Psi_1 - \Psi_2)\right)(\Phi_1 + \Psi_1 + \Phi_2 + \Psi_2 + 2)\right\|^2 \\ &\leq C\left(\|\Phi_1 - \Phi_2\|_{L^6}^2 + \|\Psi_1 - \Psi_2\|_{L^6}^2\right)\left(\|\Phi_1\|_{L^6}^2 \right. \\ &\quad \left. + \|\Psi_1\|_{L^6}^2 + \|\Phi_2\|_{L^6}^2 + \|\Psi_2\|_{L^6}^2 + 1\right), \end{aligned} \quad (5.8)$$

and this completes the rest of the proof. \square

5.2 Well-posedness of Discrete Problem

The discrete system 5.1 gives rise to a system of nonlinear algebraic equations, requiring a discussion of its well-posedness using the following variant of the Brouwer fixed point theorem.

Lemma 5.2 (The Brouwer fixed point theorem [21]). Let X be a finite-dimensional Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$. Further let $\mathcal{H} : X \rightarrow X$ be a continuous map such that $(\mathcal{H}(x), x) > 0 \forall x \in X$ with $\|x\|_X = R > 0$. Then, there exist $x^* \in X$ with $\|x^*\|_X < R$ such that $\mathcal{H}(x^*) = 0$.

Theorem 5.2 (Existence and Uniqueness Result). Given $(U^m, \mathbf{Q}^m, \widehat{U}^m) \in W_h \times \mathbf{V}_h \times M_h$ for $m = 0, 1, \dots, n$, there exists a unique discrete solution triplet $(U^{n+1}, \mathbf{Q}^{n+1}, \widehat{U}^{n+1}) \in W_h \times \mathbf{V}_h \times M_h$ to (5.1).

Proof. In order to apply Lemma 5.2, now re-write system (5.1) as

$$\left(\frac{2AU^n - 2U^n}{\Delta t^2}, w_h\right) + (\mathcal{A}\mathbf{Q}^n, \nabla w_h) - \langle \mathcal{A}\widehat{\mathbf{Q}}^n \cdot \mathbf{n}, w_h \rangle + (\mathcal{F}(2AU^n - U^{n-1}), U^{n-1}), w_h) = 0, \quad (5.9a)$$

$$(\mathcal{A}\mathbf{Q}^n, \mathbf{v}_h) + (AU^n, \nabla \cdot \mathbf{v}_h) - \langle \mathcal{A}\widehat{U}^n, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0, \quad (5.9b)$$

$$\langle \mathcal{A}\widehat{\mathbf{Q}}^n \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h/\Gamma_\theta} = 0, \quad (5.9c)$$

$$\langle \mathcal{A}\widehat{U}^n, \mu_2 \rangle_{\Gamma_\theta} = 0, \quad (5.9d)$$

and

$$\mathcal{A}\widehat{\mathbf{Q}}^n \cdot \mathbf{n} = \mathcal{A}\mathbf{Q}^n \cdot \mathbf{n} - \tau(AU^n - \mathcal{A}\widehat{U}^n) \text{ on } \mathcal{E}_h, \quad (5.10)$$

With $W_1 = AU^n$, $\mathbf{W}_{22} = \mathcal{A}\mathbf{Q}^n$ and $\widehat{W}_1 = \mathcal{A}\widehat{U}^n$, we need to discuss first the existence of solution triplet $\mathcal{W} = (W_1, \mathbf{W}_{22}, \widehat{W}_1) \in W_h \times \mathbf{V}_h \times M_h$ using Lemma 5.2. Now a use of the flux equation (5.10) with (5.9c) and (5.9d) shows with $x = \mathcal{W}$ and $(\mathcal{H}(x), x)$ as

$$\begin{aligned} (\mathcal{H}(\mathcal{W}), \mathcal{W}) &:= \left(\frac{2W_1 - 2U^n}{\Delta t^2}, W_1\right) + (\mathbf{W}_{22}, \mathbf{W}_{22}) + (\mathbf{W}_{22}, \nabla W_1) - \langle \mathbf{W}_{22} \cdot \mathbf{n}, W_1 \rangle + \langle \tau(W_1 - \widehat{W}_1), W_1 \rangle \\ &\quad + (\mathcal{F}(2W_1 - U^{n-1}), U^{n-1}), W_1) + (W_1, \nabla \cdot \mathbf{W}_{22}) - \langle \widehat{W}_1, \mathbf{W}_{22} \cdot \mathbf{n} \rangle, \end{aligned}$$

and hence, using integration by parts, we obtain in a standard way

$$\begin{aligned} (\mathcal{H}(\mathcal{W}), \mathcal{W}) &:= \|W_1\|^2 - (U^n, W_1) + \frac{(\Delta t)^2}{2}(\|\mathbf{W}_{22}\|^2 + \|(W_1 - \widehat{W}_1)\|_+^2) \\ &\quad - (U^n, W_1) + \frac{(\Delta t)^2}{2}(\mathcal{F}(2W_1 - U^{n-1}), U^{n-1}), W_1) - \frac{\Delta t^2}{2}\langle \widehat{W}_1, \mathbf{W}_{22} \cdot \mathbf{n} \rangle. \end{aligned}$$

Using the definition of $\mathcal{F}(U, V)$, the non linear term can be rewritten as

$$(\mathcal{F}(2W_1 - U^{n-1}, U^{n-1}), W_1) = \left(\frac{1}{4}(2W_1((2W_1 - U^{n-1})^2 + U^{n-1}^2 - 2)), W_1\right).$$

and hence, with

$$\|\mathcal{W}\|^2 = \|W_1\|^2 + \frac{(\Delta t)^2}{2} \left(\|\mathbf{W}_{22}\|^2 + \|(W_1 - \widehat{W}_1)\|_+^2 \right)$$

we now derive for $0 < \Delta t < 1$

$$\begin{aligned} (\mathcal{H}(\mathcal{W}), \mathcal{W}) &= \|\mathcal{W}\|^2 - (U^n, W_1) - \frac{(\Delta t)^2}{2} \|W_1\|^2 + \frac{(\Delta t)^2}{4} (W_1((2W_1 - U^{n-1})^2 + U^{n-1}^2), W_1), \\ &\geq \|\mathcal{W}\|^2 - \frac{(\Delta t)^2}{2} \|W_1\|^2 - \|U^n\| \|W_1\| \\ &\geq \left(1 - \frac{(\Delta t)^2}{2}\right) \|\mathcal{W}\| - \|U^n\| \|\mathcal{W}\| \\ &\geq \left(\frac{1}{2} \|\mathcal{W}\| - \|U^n\|\right) \|\mathcal{W}\| \end{aligned}$$

Therefore, the map $(\mathcal{H}(\mathcal{W}), \mathcal{W}) > 0$ for all $\|\mathcal{W}\| = 2\|U^n\| + 1 = R > 0$ and now from Lemma 5.2, there exists $\mathcal{W}^* = (U^{n+1}, \mathbf{Q}^{n+1}, \widehat{U}^{n+1}) \in W_h \times \mathbf{V}_h \times M_h$ with $\|(U^{n+1}, \mathbf{Q}^{n+1}, \widehat{U}^{n+1})\| < R$ such that $\mathcal{H}(\mathcal{W}^*) = 0$. This completes the existence of solution to the discrete system (5.1).

For uniqueness of the discrete solutions, let $(U_1^{n+1}, \mathbf{Q}_1^{n+1}, \widehat{U}_1^{n+1}), (U_2^{n+1}, \mathbf{Q}_2^{n+1}, \widehat{U}_2^{n+1}) \in W_h \times \mathbf{V}_h \times M_h$ be two arbitrary solutions of (5.1). With

$$W_{12}^{n+1} = U_1^{n+1} - U_2^{n+1}, \quad \mathbf{W}_{22}^{n+1} = \mathbf{Q}_1^{n+1} - \mathbf{Q}_2^{n+1} \quad \text{and} \quad \widehat{W}_{12}^{n+1} = \widehat{U}_1^{n+1} - \widehat{U}_2^{n+1}, \quad (5.11)$$

taking difference appropriately, it follows that

$$(\partial_t^2 W_{12}^n, w_h) + (\mathcal{A} \mathbf{W}_{22}^n, \nabla w_h) - \langle \mathcal{A} \widehat{\mathbf{W}}_{22}^n \cdot \mathbf{n}, w_h \rangle + (\mathcal{F}(U_1^{n+1}, U_1^{n-1}) - \mathcal{F}(U_2^{n+1}, U_2^{n-1}), w_h) = 0, \quad (5.12a)$$

$$(\mathbf{W}_{22}^{n+1}, \mathbf{v}_h) + (W_{12}^{n+1}, \nabla \cdot \mathbf{v}_h) - \langle \widehat{W}_{12}^{n+1}, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0, \quad (5.12b)$$

$$\langle \widehat{\mathbf{W}}_{22}^{n+1} \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h / \Gamma_\partial} = 0, \quad (5.12c)$$

$$\langle \widehat{W}_{12}^{n+1} \cdot \mathbf{n}, \mu_2 \rangle_{\Gamma_\partial} = 0, \quad (5.12d)$$

for all $(w_h, \mathbf{v}_h, \mu_1, \mu_2) \in W_h \times \mathbf{V}_h \times M_h \times M_h$, where,

$$\widehat{\mathbf{W}}_{22}^{n+1} \cdot \mathbf{n} = \mathbf{W}_{22}^{n+1} \cdot \mathbf{n} - \tau(W_{12}^{n+1} - \widehat{W}_{12}^{n+1}). \quad (5.13)$$

Now, choose $w_h = \delta_t W_{12}^n$ in (5.12a). In (5.12b), taking difference between between $t = t_{n+1}$ and $t = t_{n-1}$ and dividing by $2\Delta t$, set $\mathbf{v}_h = \mathcal{A} \mathbf{W}_{22}^n$. Then, add the resulting equations to arrive at

$$\begin{aligned} \frac{1}{2} \bar{\partial}_t \|\partial_t W_{12}^n\|^2 + \delta_t \left(\|\mathbf{W}_{22}^n\|^2 + \|W_{12}^n - \widehat{W}_{12}^{n+1}\|_+^2 \right) \\ + (\mathcal{F}(U_1^{n+1}, U_1^{n-1}) - \mathcal{F}(U_2^{n+1}, U_2^{n-1}), \delta_t W_{12}^n) = 0. \end{aligned} \quad (5.14)$$

Multiplying by $2\Delta t$, a use of the Cauchy-Schwarz with Young's inequality shows

$$\begin{aligned} \|\partial_t W_{12}^n\|^2 - \|\partial_t W_{12}^{n-1}\|^2 + \frac{1}{2} \left(\|\mathbf{W}_{22}^{n+1}\|^2 - \|\mathbf{W}_{22}^{n-1}\|^2 + \|W_{12}^{n+1} - \widehat{W}_{12}^{n+1}\|_+^2 - \|W_{12}^{n-1} - \widehat{W}_{12}^{n-1}\|_+^2 \right) \\ \leq \Delta t \left((\mathcal{F}(U_1^{n+1}, U_1^{n-1}) - \mathcal{F}(U_2^{n+1}, U_2^{n-1}))^2 + \frac{\Delta t}{4} (\|\partial_t W_{12}^n\|^2 + \|\partial_t W_{12}^{n-1}\|^2) \right). \end{aligned} \quad (5.15)$$

For the nonlinear term on the right hand side of (5.14), we note as a consequence of Theorem 5.1 the boundedness of solution in L^6 norm that

$$\begin{aligned} \|\mathcal{F}(U_1^{n+1}, U_1^{n-1}) - \mathcal{F}(U_2^{n+1}, U_2^{n-1})\|^2 &\leq C \left((\|U_1^{n+1}\|_{L^6}^4 + \|U_1^{n-1}\|_{L^6}^4 + \|U_2^{n+1}\|_{L^6}^4 \right. \\ &\quad \left. + \|U_2^{n-1}\|_{L^6}^4 + 2^4) (\|\mathbf{W}_{12}^{n+1}\|_{L^6}^2 + \|\mathbf{W}_{12}^{n-1}\|_{L^6}^2) \right), \\ &\leq C \left(\|\mathbf{W}_{22}^{n+1}\|^2 + \|\mathbf{W}_{22}^{n-1}\|^2 + \|W_{12}^{n+1} - \widehat{W}_{12}^{n+1}\|_+^2 + \|W_{12}^{n-1} - \widehat{W}_{12}^{n-1}\|_+^2 \right). \end{aligned} \quad (5.16)$$

Substituting (5.16) in (5.15), and summing up the resulting inequality from $n = 1$ to m , we arrive at

$$\begin{aligned} \|\partial_t W_{12}^m\|^2 + \frac{1}{2} \left(\|\mathbf{W}_{22}^{m+1}\|^2 + \|W_{12}^{m+1} - \widehat{W}_{12}^{m+1}\|_+^2 \right) &\leq \|\partial_t W_{12}^0\|^2 + \frac{1}{2} \left(\|\mathbf{W}_{22}^0\|^2 + \|W_{12}^0 - \widehat{W}_{12}^0\|_+^2 \right) \\ &+ \|\mathbf{W}_{22}^1\|^2 + \|W_{12}^1 - \widehat{W}_{12}^1\|_+^2 + \Delta t \sum_{n=0}^m \left(C \left(\|\mathbf{W}_{22}^{n+1}\|^2 + \|\mathbf{W}_{22}^{n-1}\|^2 + \|W_{12}^{n+1} - \widehat{W}_{12}^{n+1}\|_+^2 \right) \right. \\ &\left. + \|W_{12}^{n-1} - \widehat{W}_{12}^{n-1}\|_+^2 + \frac{1}{4} \left(\|\partial_t W_{12}^n\|^2 + \|\partial_t W_{12}^{n-1}\|^2 \right) \right). \end{aligned} \quad (5.17)$$

With $\widehat{W}_{12}^0 = 0$, $\mathbf{W}_{22}^0 = 0$ and $W_{12}^0 - \widehat{W}_{12}^0 = 0$, a use of the kickback argument with an application of the discrete Grönwall's lemma yields

$$\begin{aligned} \left(1 - \frac{\Delta t}{4}\right) \|\partial_t W_{12}^n\|^2 + \left(\frac{1}{2} - C\Delta t\right) \left(\|\mathbf{W}_{22}^{n+1}\|^2 + \|W_{12}^{n+1} - \widehat{W}_{12}^{n+1}\|_+^2 \right) &\leq C \left(\|\partial_t W_{12}^0\|^2 \right. \\ &\left. + \|\mathbf{W}_{22}^1\|^2 + \|W_{12}^1 - \widehat{W}_{12}^1\|_+^2 \right). \end{aligned} \quad (5.18)$$

Choose Δt so that $2C\Delta t < 1$ and $0 < \Delta t < 1$, and to complete the remaining part, we need to obtain an estimate for $n = 1$. Proceed similarly to arrive at

$$\frac{2}{(\Delta t)^2} (W_{12}^1, w_h) - \left(\frac{1}{2} \mathbf{W}_{22}^1, \nabla w_h\right) + \left\langle \frac{1}{2} \widehat{W}_{12}^1 \cdot \mathbf{n}, w_h \right\rangle + (\mathcal{F}(U_1^1, U_1^0) - \mathcal{F}(U_2^1, U_2^0), w_h) = 0, \quad (5.19a)$$

$$(\mathbf{W}_{22}^1, \mathbf{v}_h) + (W_{12}^1, \nabla \cdot \mathbf{v}_h) - \langle \widehat{W}_{12}^1, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0, \quad (5.19b)$$

$$\langle \widehat{W}_{22}^1 \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h/\Gamma_\partial} = 0, \quad (5.19c)$$

$$\langle \widehat{W}_{12}^1, \mu_2 \rangle_{\Gamma_\partial} = 0, \quad (5.19d)$$

for all $(w_h, \mathbf{v}_h, \mu_1, \mu_2) \in W_h \times \mathbf{V}_h \times M_h \times M_h$, where

$$\widehat{W}_{22}^1 \cdot \mathbf{n} = \mathbf{W}_{22}^1 \cdot \mathbf{n} - \tau(W_{12}^1 - \widehat{W}_{12}^1). \quad (5.20)$$

Now, in (5.19a), choosing $w_h = W_{12}^1/(\Delta t)$ and in (5.19b) set $\mathbf{v}_h = \mathbf{W}_{22}^1/(\Delta t)$. Then, add these two resulting equations and then multiply by Δt to find that

$$2\|W_{12}^1/(\Delta t)\|^2 + \frac{1}{2} \left(\|\mathbf{W}_{22}^1\|^2 + \|W_{12}^1 - \widehat{W}_{12}^1\|_+^2 \right) = -\Delta t \left(\mathcal{F}(U_1^1, U_1^0) - \mathcal{F}(U_2^1, U_2^0), W_{12}^1/(\Delta t) \right). \quad (5.21)$$

For the nonlinear term, using Lemma 5.1, then Lemma 3.1 and Theorem 5.1 gives

$$2\|W_{12}^1/(\Delta t)\|^2 + \frac{1}{2} \left(\|\mathbf{W}_{22}^1\|^2 + \|W_{12}^1 - \widehat{W}_{12}^1\|_+^2 \right) \leq C \Delta t \left(\|\mathbf{W}_{22}^1\|^2 + \|W_{12}^1 - \widehat{W}_{12}^1\|_\tau^2 \right) + \|W_{12}^1/(\Delta t)\|^2. \quad (5.22)$$

Hence, the use of a kickback argument shows

$$\|W_{12}^1/(\Delta t)\|^2 + \left(\frac{1}{2} - \Delta t C\right) \left(\|\mathbf{W}_{22}^1\|^2 + \|W_{12}^1 - \widehat{W}_{12}^1\|_+^2 \right) \leq 0. \quad (5.23)$$

By choosing Δt small so that $(\frac{1}{2} - C\Delta t) > 0$, there holds $U_1^1 = U_2^1$, $\mathbf{Q}_1^1 = \mathbf{Q}_2^1$, $\widehat{U}_1^1 = \widehat{U}_2^1$. On substitution in (5.18), this completes the rest of the proof. \square

5.3 Error analysis

Since the error estimates for the semidiscrete scheme are derived in Section 3, we now split the error in the completely discrete scheme (5.1) as

$$u(t_n) - U^n = u(t_n) - u_h(t_n) + \xi_u^n \quad \text{and} \quad \mathbf{q}(t_n) - \mathbf{Q}^n = \mathbf{q}(t_n) - \mathbf{q}_h(t_n) + \xi_q^n.$$

Now, it remains to estimate $\|\xi_u^n\|$ and $\|\xi_q^n\|$. Since $U^0 = u_h(0)$ and $\mathbf{Q}^0 = \mathbf{q}_h(0)$ are the solutions of problem 3.4, it follows that $\xi_u(0) = 0$, $\xi_q(0) = 0$.

With the help of (3.2) and (5.1), we now obtain

$$\begin{aligned} \frac{2}{\Delta t}(\partial_t^* \xi_u^{1/2}, w_h) + (\xi_q^{1/2}, \nabla w_h) - \langle \widehat{\xi}_q^{1/2} \cdot \mathbf{n}, w_h \rangle + (f(u_h(t_{1/2})) - \mathcal{F}(U^1, U^0), w_h) \\ + \frac{2}{\Delta}(u_1, w_h) = \left(\frac{2}{\Delta} \partial_t^* u_h(t_{1/2}) - u_{htt}(t_{1/2}), w_h\right), \end{aligned} \quad (5.24a)$$

$$(\xi_q^{1/2}, \mathbf{v}_h) + (\xi_u^{1/2}, \nabla \cdot \mathbf{v}_h) - \langle \widehat{\xi}_u^{1/2}, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0, \quad (5.24b)$$

$$\begin{aligned} (\partial_t^2 \xi_u^n, w_h) + (\mathcal{A} \xi_q^n, \nabla w_h) - \langle \mathcal{A} \widehat{\xi}_q^n \cdot \mathbf{n}, w_h \rangle + (\mathcal{A} f(u_h(t_n)) - \mathcal{F}(U^{n+1}, U^{n-1}), w_h) \\ = (\partial_t^2 u_h(t_n) - \mathcal{A} u_{htt}(t_n), w_h), \end{aligned} \quad (5.24c)$$

$$(\xi_q^n, \mathbf{v}_h) + (\xi_u^n, \nabla \cdot \mathbf{v}_h) - \langle \widehat{\xi}_u^n, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0, \quad (5.24d)$$

Below, we discuss the main theorem on error analysis of the fully discrete scheme.

Theorem 5.3. Let $\{U^n, Q^n, \widehat{U}^n\}$, $n \geq 0$ be defined by (5.1a)-(5.1e). Then, there is a positive constant C independent of h and k such that for all $J \geq 1$

$$\max_{0 \leq n \leq J} (\|\partial_t(u(t^{n+1}) - U^{n+1})\| + \|\mathbf{q}(t^{n+1}) - \mathbf{Q}^{n+1}\|) \leq C (h^{k+1} + (\Delta t)^2), \quad (5.25)$$

$$\max_{0 \leq n \leq J} \|u(t^{n+1}) - U^{n+1}\| \leq C (h^{k+1} + (\Delta t)^2). \quad (5.26)$$

Proof. Choose $w_h = \delta_t \xi_u^n$ in (5.24c) and in (5.24d) taking difference between two levels, i.e., between $t = t_{n+1}$ and $t = t_{n-1}$ and then dividing by $2\Delta t$, then choose $\mathbf{v}_h = \mathcal{A} \xi_q^n$, we add the resulting equations.

$$\begin{aligned} \frac{1}{2} \overline{\partial}_t \|\partial_t \xi_u^n\|^2 + \delta_t (\|\xi_q^n\|^2 + \|\xi_u^n - \widehat{\xi}_u^n\|_+^2) = (\partial_t^2 u_h(t_n) - \mathcal{A} u_{htt}(t_n), \delta_t \xi_u^n) \\ - (\mathcal{A} f(u_h(t_n)) - \mathcal{F}(U^{n+1}, U^{n-1}), \delta_t \xi_u^n). \end{aligned} \quad (5.27)$$

side of (5.27) We can easily deal with the first term on the right-hand side in (5.27) using Taylor's series expansion

$$\begin{aligned} \partial_t^2 u_h(t_n) - \mathcal{A} u_{htt}(t_n) &= \partial_t^2 (u_h(t_n) - u(t_n)) + \partial_t^2 u(t_n) - \mathcal{A} u_{htt}(t_n), \\ &= -(\partial_t^2 (u(t_n) - u_h(t_n))) + (\mathcal{A} u_{tt}(t_n) - \mathcal{A} u_{htt}(t_n)) + \frac{1}{(\Delta t)^2} R_1^* - \frac{1}{2} R_2^*, \end{aligned}$$

where,

$$R_1^* = \frac{1}{6} \int_{-\Delta t}^{\Delta t} (\Delta t - |\mu|)^3 u_{tttt}(t_n + \mu) d\mu, \quad R_2^* = \int_{-\Delta t}^{\Delta t} (\Delta t - |\mu|) u_{tttt}(t_n + \mu) d\mu.$$

A use of Theorem 3.1 with Lemma 3.3 and following the argument given in [24] to estimate R_1^* and R_2^* shows

$$\begin{aligned} \|\partial_t^2 u_h(t_n) - \mathcal{A} u_{htt}(t_n)\|^2 &\leq C h^{2(k+1)} \|u_{tt}\|_{L^\infty(H^{k+1})}^2 + \frac{1}{(\Delta t)^4} \|R_1^*\|^2 + \frac{1}{4} \|R_2^*\|^2 \\ &\leq C (h^{2(k+1)} \|u_{tt}\|_{L^\infty(H^{k+1})}^2 + (\Delta t)^3 \int_{t_{n-1}}^{t_{n+1}} \|u_{tttt}(s)\|^2 ds). \end{aligned} \quad (5.28)$$

Therefore, we need to estimate the nonlinear term and now, we rewrite it as

$$\begin{aligned} \mathcal{A} f(u_h(t_n)) - \mathcal{F}(U^{n+1}, U^{n-1}) &= \mathcal{A} F'(u_h(t_n)) - \mathcal{F}(U^{n+1}, U^{n-1}), \\ &= \mathcal{I}_1 + \mathcal{I}_2, \end{aligned} \quad (5.29)$$

where

$$\mathcal{I}_1 = \mathcal{A} F'(u_h(t_n)) - \mathcal{F}(u_h(t_{n+1}), u_h(t_{n-1})), \quad \mathcal{I}_2 = \mathcal{F}(u_h(t_{n+1}), u_h(t_{n-1})) - \mathcal{F}(U^{n+1}, U^{n-1}).$$

Again, an application of the Taylor series expansion shows

$$\mathcal{I}_1 = \frac{1}{2} (R_1 + R_2) - \frac{1}{u_h(t_{n+1}) - u_h(t_{n-1})} (R_1 - R_2),$$

where, the remainder R_1 and R_2 are given by

$$\begin{aligned} R_1 &= (u_h(t_{n+1}) - u_h(t_n))^2 \int_0^1 F''(u_h(t_n) + \lambda(u_h(t_{n+1}) - u_h(t_n)))(1 - \lambda)d\lambda, \\ R_2 &= (u_h(t_{n-1}) - u_h(t_n))^2 \int_0^1 F''(u_h(t_n) + \lambda(u_h^{n-1} - u_h(t_n)))(1 - \lambda)d\lambda. \end{aligned}$$

After substituting the value of F'' and integrating in \mathcal{I}_1 , we use Theorem 3.2 to derive

$$\|\mathcal{I}_1\| \leq C (h^{k+1} + (\Delta t)^2).$$

For the estimate of \mathcal{I}_2 , a use of Lemma 5.1 with Lemma 3.1 and Theorem 5.1 yields

$$\begin{aligned} \|\mathcal{I}_2\|^2 &\leq C (\|\xi_u^{n-1}\|_{L^6}^2 + \|\xi_u^{n+1}\|_{L^6}^2)(\|u_h(t_{n+1})\|_{L^6}^4 + \|u_h(t_{n-1})\|_{L^6}^4 + \|U^{n+1}\|_{L^6}^4 + \|U^{n-1}\|_{L^6}^4 + 1) \\ &\leq C (\|\xi_q^{n+1}\|^2 + \|\xi_q^{n-1}\|^2 + \|\xi_u^{n+1} - \widehat{\xi}_u^{n+1}\|_+^2 + \|\xi_u^{n-1} - \widehat{\xi}_u^{n-1}\|_+^2). \end{aligned}$$

Substitute the estimates of \mathcal{I}_1 and \mathcal{I}_2 in (5.29) to arrive at

$$\begin{aligned} \|\mathcal{A}f(u_h(t_n)) - \mathcal{F}(U^{n+1}, U^{n-1})\|^2 &\leq C \left((h^{k+1} + (\Delta t)^2)^2 + \|\xi_q^{n+1}\|^2 + \|\xi_q^{n-1}\|^2 \right. \\ &\quad \left. + \|\xi_u^{n+1} - \widehat{\xi}_u^{n+1}\|_+^2 + \|\xi_u^{n-1} - \widehat{\xi}_u^{n-1}\|_+^2 \right). \end{aligned} \quad (5.30)$$

Substitute (5.28) and (5.30) in (5.27), multiply by $2\Delta t$ and sum up from $n = 1$ to m with $U^0 = u_h(0)$, $Q^0 = \mathbf{q}_h(0)$ and $\widehat{U}^0 = \widehat{u}_h(0)$, to obtain

$$\begin{aligned} \|\partial_t \xi_u^m\|^2 + \frac{1}{2} (\|\xi_q^{m+1}\|^2 + \|\xi_u^{m+1} - \widehat{\xi}_u^{m+1}\|_+^2) &\leq \frac{1}{2} (\|\xi_q^1\|^2 + \|\xi_u^1 - \widehat{\xi}_u^1\|_+^2) + C (h^{2(k+1)} + (\Delta t)^4) \\ &\quad + \frac{1}{4} (\|\partial_t \xi_u^n\|^2 + \|\partial_t \xi_u^{n-1}\|^2) + C \left(\Delta t \sum_{n=1}^m (\|\xi_q^{n+1}\|^2 + \|\xi_q^{n-1}\|^2 \right. \\ &\quad \left. + \|\xi_u^{n+1} - \widehat{\xi}_u^{n+1}\|_+^2 + \|\xi_u^{n-1} - \widehat{\xi}_u^{n-1}\|_+^2) \right). \end{aligned} \quad (5.31)$$

Using the kickback argument, we arrive at

$$\begin{aligned} (1 - \frac{\Delta t}{4}) \|\partial_t \xi_u^m\|^2 + (\frac{1}{2} - C\Delta t) (\|\xi_q^{m+1}\|^2 + \|\xi_u^{m+1} - \widehat{\xi}_u^{m+1}\|_+^2) &\leq \frac{1}{2} (\|\xi_q^1\|^2 + \|\xi_u^1 - \widehat{\xi}_u^1\|_+^2) \\ &\quad + C (h^{2(k+1)} + (\Delta t)^4) + C \Delta t \sum_{n=1}^m (\|\partial_t \xi_u^{n-1}\|^2 + \|\xi_q^n\|^2 + \|\xi_u^n - \widehat{\xi}_u^n\|_+^2). \end{aligned} \quad (5.32)$$

Now, with the help of (5.24a) and (5.24b), we plan to bound $\|\xi_q^1\|^2 + \|\xi_u^1 - \widehat{\xi}_u^1\|_+^2$. Choose $w_h = \partial_t^* \xi_u^{1/2}$ in (5.24a) and $\mathbf{v}_h = \xi_q^{1/2}$ in (5.24b), after adding resulting equation, we obtain

$$\begin{aligned} \frac{2}{\Delta t} \|\partial_t^* \xi_u^{1/2}\|^2 + \frac{1}{2\Delta t} (\|\xi_q^1\|^2 - \|\xi_q^0\|^2 + \|\xi_u^1 - \widehat{\xi}_u^1\|_+^2 - \|\xi_u^0 - \widehat{\xi}_u^0\|_+^2) \\ = (\frac{2}{\Delta t} (\partial_t^* u_h(t_{1/2}) - U^0) - u_{htt}(t_{1/2}), \partial_t^* \xi_u^{1/2}) - (f(u_h(t_{1/2})) - \mathcal{F}(U^1, U^0), \partial_t^* \xi_u^{1/2}). \end{aligned} \quad (5.33)$$

An application of the Taylor's series expansion in (5.33) shows

$$\|\xi_u^1\|^2 + \|\xi_q^1\|^2 + \|\xi_u^1 - \widehat{\xi}_u^1\|_+^2 \leq C (h^{2k+2} + (\Delta t)^4). \quad (5.34)$$

On substitution (5.34) in (5.32) and choosing $\Delta t > 0$ small sothat $(2C\Delta t - 1) > 0$, an application of Grönwall's lemma yields

$$\max_{0 \leq m \leq J} (\|\partial_t \xi_u^m\|^2 + \|\xi_q^{m+1}\|^2 + \|\xi_u^{m+1} - \widehat{\xi}_u^{m+1}\|_+^2) \leq C (h^{2k+2} + (\Delta t)^4). \quad (5.35)$$

For the estimate (5.26), note using Lemma 3.1 that

$$\|\xi_u^{m+1}\|_{L^p} \leq C \left(\|\xi_q^{m+1}\| + \|\xi_u^{m+1} - \widehat{\xi}_u^{m+1}\|_+ \right). \quad (5.36)$$

A use of (5.35) in (5.36) for $p = 2$ with Lemma 3.2 and triangle inequality completes the rest of the proof. \square

5.4 A Fully Discrete Non-Conservative Scheme

This subsection is focused on the fully discrete, non-conservative HDG scheme. We approximate the nonlinear term in system (5.1) by replacing $\mathcal{F}(U^{n+1}, U^{n-1})$ simply by $f(U^n)$. The advantage of this scheme is that at each time step, a linear system of algebraic equations has to be solved.

The HDG fully discretized formulation reads as: for each $n \geq 0$, we seek $(U^n, \mathbf{Q}^n, \widehat{U}^n) \in W_h \times \mathbf{V}_h \times M_h$ such that

$$\frac{2}{\Delta t}(\partial_t^* U^{1/2}, w_h) + (\mathbf{Q}^{1/2}, \nabla w_h) - \langle \widehat{\mathbf{Q}}^{1/2} \cdot \mathbf{n}, w_h \rangle + (f(U^0), w_h) - \frac{2}{\Delta t}(u_1, w_h) = 0 \quad (5.37a)$$

$$(\partial_t^2 U^n, w_h) + (\mathcal{A}\mathbf{Q}^n, \nabla w_h) - \langle \mathcal{A}\widehat{\mathbf{Q}}^n \cdot \mathbf{n}, w_h \rangle + (f(U^n), w_h) = 0 \quad \forall n \geq 1, \quad (5.37b)$$

$$(\mathbf{Q}^n, \mathbf{v}_h) + (U^n, \nabla \cdot \mathbf{v}_h) - \langle \widehat{U}^n, \mathbf{v} \cdot \mathbf{n} \rangle = 0 \quad \forall n \geq 0, \quad (5.37c)$$

$$\langle \widehat{\mathbf{Q}}^n \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h/\Gamma_\partial} = 0 \quad \forall n \geq 0, \quad (5.37d)$$

$$\langle \widehat{U}^n, \mu_2 \rangle_{\Gamma_\partial} = 0 \quad \forall n \geq 0, \quad (5.37e)$$

for all $(w_h, \mathbf{v}_h, \mu_1, \mu_2) \in W_h \times \mathbf{V}_h \times M_h \times M_h$, where,

$$\widehat{\mathbf{Q}}^n \cdot \mathbf{n} = \mathbf{Q}^n \cdot \mathbf{n} - \tau(U^n - \widehat{U}^n) \quad \text{on } \mathcal{E}_h, \quad (5.38)$$

Theorem 5.4. Assume that the solution triplet $\{U^n, \mathbf{Q}^n, \widehat{U}^n\}$, $n \geq 0$ is defined by the system (5.37). Then there is a constant C independent of h and k such that for all $J \geq 1$,

$$\max_{0 \leq n \leq J} (\|\partial_t(u(t^{n+1}) - U^{n+1})\| + \|\mathbf{q}(t^{n+1}) - \mathbf{Q}^{n+1}\|) \leq C(h^{k+1} + (\Delta t)^2), \quad (5.39)$$

$$\max_{0 \leq n \leq J} (\|u(t^{n+1}) - U^{n+1}\|) \leq C(h^{k+1} + (\Delta t)^2). \quad (5.40)$$

Proof. Similar to what is done in Theorem 5.3, we arrive at

$$\begin{aligned} & \frac{1}{2\Delta t} \left(\|\partial_t \xi_u^n\|^2 - \|\partial_t \xi_u^{n-1}\|^2 + \frac{1}{2} (\|\xi_q^{n+1}\|^2 - \|\xi_q^{n-1}\|^2 + \|\xi_u^{n+1} - \widehat{\xi}_u^{n+1}\|_+^2 - \|\xi_u^{n-1} - \widehat{\xi}_u^{n-1}\|_+^2) \right) \\ & = (\partial_t^2 u_h(t_n) - Au_{htt}(t_n), \delta_t \xi_u^n) - (\mathcal{A}f(u_h(t_n)) - f(U^n), \delta_t \xi_u^n) \end{aligned} \quad (5.41)$$

Only the nonlinear term on the right-hand side of the equation (5.41) needs to be handled. Therefore, by applying Taylor's expansion, we arrive at

$$|(\mathcal{A}f(u_h(t_n)) - f(U^n), \delta_t \xi_u^n)| \leq \|f(u_h(t_n)) - f(U^n) + \frac{1}{2}(r_1 + r_2)\| \|\delta_t \xi_u^n\|,$$

where,

$$\begin{aligned} r_1 &= (u_h(t_{n+1}) - u_h(t_n))^2 \int_0^1 f'(u_h(t_n) + \lambda(u_h(t_{n+1}) - u_h(t_n)))(1 - \lambda) d\lambda, \\ r_2 &= (u_h(t_{n-1}) - u_h(t_n))^2 \int_0^1 f'(u_h(t_n) + \lambda(u_h(t_{n-1}) - u_h(t_n)))(1 - \lambda) d\lambda. \end{aligned}$$

Note from Lemma 5.1 that

$$\|f(u_h(t_n)) - f(U^n)\|^2 \leq C \|\xi_u^n\|_{L^6}^2 (\|u_h(t_n)\|_{L^6}^4 + \|U^n\|_{L^6}^4 + 1),$$

and using the L^6 estimates of $u_h(t_n), U^n$ at n th level with an application of energy arguments, Lemma 3.1 and the Young's inequality yield,

$$|(\mathcal{A}f(u_h(t_n)) - f(U^n), \delta_t \xi_u^n)| \leq C(h^{2k+2} + (\Delta t)^4 + \|\xi_q^n\|^2 + \|\xi_u^n - \widehat{\xi}_u^n\|_+^2 + \|\delta_t \xi_u^n\|^2).$$

Putting this inequality in (5.41), then multiplying by $2\Delta t$ and summing from $n = 1$ to J , it follows that

$$\begin{aligned} & (1 - \frac{\Delta t}{4}) \|\partial_t \xi_u^J\|^2 + (\frac{1}{2} - C\Delta t) (\|\xi_q^{J+1}\|^2 + \|\xi_u^{J+1} - \widehat{\xi}_u^{J+1}\|_+^2) \leq C(\Delta t) \left(\sum_{n=1}^J (h^{2k+2} + (\Delta t)^2) \right. \\ & \left. + \|\partial_t^2 u_h(t_n) - u_{htt}(t_n)\|^2 + \|\partial_t \xi_u^n\|^2 + \|\xi_q^n\|^2 + \|\xi_u^n - \widehat{\xi}_u^n\|_+^2 + \|\xi_q^1\|^2 + \|\xi_u^1 - \widehat{\xi}_u^1\|_+^2 \right). \end{aligned}$$

Now, for $\Delta t > 0$ small, $1 - 2C\Delta t > 0$. The second term on the right-hand side of (5.41) is one that we previously dealt with in the Theorem 5.3. The desired outcome is now obtained by utilizing Grönwall's inequality. The last inequality (5.40) in this theorem can be proved similarly as in the conservative case. This concludes the rest of the proof. \square

6 One Variation of the HDG method

This section deals with a novel HDG method proposed by Qiu and Shi in [25], a variation of the HDG method. It differs from the existing method in that polynomials of degree $k + 1$ and k , with $k \geq 0$, approximate the displacement and its gradient, respectively. As a result, the convergence of order $O(h^{k+2})$ for the displacement in $L^\infty(L^2)$ -norm is derived, while the convergence of order $O(h^{k+1})$ is shown for the gradient in $L^\infty(L^2)$ -norm.

In this section, we indicate only the differences in the proof of the error analysis. With \mathbf{V}_h and M_h as in Section 2, the space W_h is now changed to

$$W_h = \{w \in L^2(\Omega) : w|_K \in \mathcal{P}_{k+1}(K) \ \forall K \in \mathcal{T}_h\}$$

The new variation of the HDG formulation is to seek $(u_h(t), \mathbf{q}_h(t), \hat{u}_h(t)) \in W_h \times \mathbf{V}_h \times M_h$ such that for $t \in (0, T]$

$$(\mathbf{q}_h, \mathbf{v}_h) + (u_h, \nabla \cdot \mathbf{v}_h) - \langle \hat{u}_h, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0 \quad \forall \mathbf{v}_h \in \mathbf{V}_h, \quad (6.1a)$$

$$(u_{htt}, w_h) + (\mathbf{q}_h, \nabla w_h) - \langle \hat{\mathbf{q}}_h \cdot \mathbf{n}, w_h \rangle + (f(u_h), w_h) = 0 \quad \forall w_h \in W_h, \quad (6.1b)$$

$$\langle \hat{\mathbf{q}}_h \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h/\Gamma_\partial} = 0 \quad \forall \mu_1 \in M_h, \quad (6.1c)$$

$$\langle \hat{u}_h, \mu_2 \rangle_{\Gamma_\partial} = 0 \quad \forall \mu_2 \in M_h, \quad (6.1d)$$

where the new flux is defined as

$$\hat{\mathbf{q}}_h \cdot \mathbf{n} = \mathbf{q}_h \cdot \mathbf{n} + \frac{\tau}{h_K} (\mathcal{P}u_h - \hat{u}_h) \text{ on } \mathcal{E}_h. \quad (6.2)$$

Here, \mathcal{P} is the L^2 projection onto M_h .

Energy conservation is the same as in Theorem 2.1 with Corollary 2.1.

Theorem 6.1. Let $u_0, u_1 \in L^\infty(0, T; H^{k+2})$ and $u \in L^\infty(0, T; H^{k+2} \cap H_0^1)$, $u_t \in L^2(0, T; H^{k+2})$. Then for all $h = \max_{K \in \mathcal{T}_h} h_K$ and for all $t \in (0, T]$, the following estimates hold:

$$\|(u_t - u_{ht})(t)\| + \|(\mathbf{q} - \mathbf{q}_h)(t)\| + \|\nabla(u - u_h)(t)\| \leq C h^{k+1} \left(\|u_0\|_{H^{k+2}} + \|u\|_{L^\infty(0, T; H^{k+2})} + \|u_t\|_{L^2(0, T; H^{k+2})} \right). \quad (6.3)$$

with the additional assumptions $u_{tt} \in L^2(0, T; H^{k+2})$,

$$\|(u_{tt} - u_{htt})(t)\| + \|(\mathbf{q}_t - \mathbf{q}_{ht})(t)\| \leq C h^{k+1} \left(\|u_1\|_{H^{k+2}} + \|u_t\|_{L^\infty(0, T; H^{k+2})} + \|u_{tt}\|_{L^2(0, T; H^{k+2})} \right). \quad (6.4)$$

Moreover,

$$\|(u - u_h)(t)\| \leq C h^{k+2} \left(\|u_0\|_{H^{k+2}} + \|u_1\|_{H^{k+2}} + \|u\|_{L^\infty(0, T; H^{k+2})} + \|u_{tt}\|_{L^2(0, T; H^{k+2})} \right). \quad (6.5)$$

Proof of this Theorem is given in the next two subsections.

6.1 Error Estimates

We redefine the projections Π_W and Π_V as the standard L^2 orthogonal projection onto W_h and \mathbf{V}_h , respectively, and followings are the properties of these projections [25]:

$$\left. \begin{aligned} \|\vartheta_u\| &\leq Ch^{l_u} \|u\|_{H^{l_u}(\mathcal{T}_h)}, & \forall 0 \leq l_u \leq k+2, \\ \|\hat{\vartheta}_u\|_{\mathcal{E}_h} &\leq Ch^{l_u-1/2} \|\mathbf{q}\|_{H^{l_u}(\mathcal{T}_h)}, & \forall 1 \leq l_u \leq k+1, \\ \|\vartheta_u\|_E &\leq Ch^{l_u-1/2} \|u\|_{H^{l_u}(K)}, & \forall 1 \leq l_u \leq k+2, \\ \|\vartheta_{\mathbf{q}}\| &\leq Ch^{l_q} \|\mathbf{q}\|_{H^{l_q}(\mathcal{T}_h)}, & \forall 0 \leq l_q \leq k+1, \\ \|\hat{\vartheta}_{\mathbf{q}}\|_E &\leq Ch^{l_q-1/2} \|\mathbf{q}\|_{H^{l_q}(K)}, & \forall 1 \leq l_q \leq k+1, \\ \|\vartheta_{\mathbf{q}}\|_E &\leq Ch^{l_q-1/2} \|\mathbf{q}\|_{H^{l_q}(K)}, & \forall 1 \leq l_q \leq k+1, \\ \|w\|_E &\leq Ch^{1/2} \|w\|_K, & \forall w \in W_h. \end{aligned} \right\} \quad (6.6)$$

Here, E denotes the face of an element K . Also, define

$$\|\psi\|^2 = \sum_{K \in \mathcal{T}_h} \|\psi\|_{L^2(K)}^2, \quad \|\psi - \widehat{\psi}\|_{\partial\mathcal{T}_h}^2 = \sum_{K \in \mathcal{T}_h} \left\| \left(\frac{\tau}{h_K} \right)^{1/2} (\mathcal{P}\psi - \widehat{\psi}) \right\|_{L^2(\partial K)}^2.$$

Using L^2 projections $(\Pi_W, \Pi_V, \mathcal{P})$, error equations in θ_u , θ_q and $\widehat{\theta}_u$ are written for all $(\mathbf{v}_h, w_h, \mu_1, \mu_2) \in \mathbf{V}_h \times W_h \times M_h \times M_h$ as

$$(\theta_q, \mathbf{v}_h) + (\theta_u, \nabla \cdot \mathbf{v}_h) - \langle \widehat{\theta}_u, \mathbf{v}_h \cdot \mathbf{n} \rangle = 0, \quad (6.7a)$$

$$(\theta_{u_{tt}}, w_h) + (\theta_q, \nabla w_h) - \langle (\mathbf{q} - \widehat{\mathbf{q}}_h) \cdot \mathbf{n}, w_h \rangle + (f(u) - f(u_h), w_h) = 0, \quad (6.7b)$$

$$\langle (\mathbf{q} - \widehat{\mathbf{q}}_h) \cdot \mathbf{n}, \mu_1 \rangle_{\mathcal{E}_h/\Gamma_\partial} = 0, \quad (6.7c)$$

$$\langle \widehat{\theta}_u, \mu_2 \rangle_{\Gamma_\partial} = 0, \quad (6.7d)$$

where the numerical trace for flux is defined as

$$(\mathbf{q} - \widehat{\mathbf{q}}_h) \cdot \mathbf{n} = (\theta_q - \vartheta_q) \cdot \mathbf{n} - \frac{\tau}{h_K} (\mathcal{P}\theta_u - \widehat{\theta}_u) + \frac{\tau}{h_K} \mathcal{P}\vartheta_u. \quad (6.8)$$

With the help of the error equations (6.7), we derive the estimates of θ_q . Differentiating equation (6.7a) with respect to time and set $\mathbf{v}_h = \theta_q$ and choose $w_h = \theta_{u_t}$ in (6.7b), then adding these resulting equations to obtain

$$\frac{1}{2} \frac{d}{dt} (\|\theta_q\|^2 + \|\theta_{u_t}\|^2) + \langle \theta_q \cdot \mathbf{n}, \theta_{u_t} - \widehat{\theta}_{u_t} \rangle - \langle (\mathbf{q} - \widehat{\mathbf{q}}_h) \cdot \mathbf{n}, \theta_{u_t} \rangle + (f(u) - f(u_h), \theta_{u_t}) = 0. \quad (6.9)$$

Adding the zero-term $\langle (\mathbf{q} - \widehat{\mathbf{q}}_h) \cdot \mathbf{n}, \widehat{\theta}_{u_t} \rangle$, a use of (6.8) shows

$$\frac{1}{2} \frac{d}{dt} (\|\theta_q\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}^2) + \left\langle \vartheta_q \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \theta_{u_t} - \widehat{\theta}_{u_t} \right\rangle + (f(u) - f(u_h), \theta_{u_t}) = 0.$$

On integrating from 0 to t , it follows that

$$\|\theta_q\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}^2 = - \int_0^t \left(\left\langle \vartheta_q \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \theta_{u_t} - \widehat{\theta}_{u_t} \right\rangle + (f(u) - f(u_h), \theta_{u_t}) \right) ds. \quad (6.10)$$

For estimating the right-hand side terms in (6.10), we first rewrite the first term as

$$\begin{aligned} \int_0^t \left(\left\langle \vartheta_q \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \theta_{u_t} - \widehat{\theta}_{u_t} \right\rangle \right) ds &= \int_0^t \left(\frac{d}{ds} \left\langle \vartheta_q \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \theta_u - \widehat{\theta}_u \right\rangle \right. \\ &\quad \left. - \left\langle \vartheta_{q_t} \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_{u_t}, \theta_u - \widehat{\theta}_u \right\rangle \right) ds, \\ &= \left\langle \vartheta_q \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \theta_u - \widehat{\theta}_u \right\rangle (t) - \left\langle \vartheta_q \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \theta_u - \widehat{\theta}_u \right\rangle (0) \\ &\quad - \int_0^t \left\langle \vartheta_{q_t} \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_{u_t}, \theta_u - \widehat{\theta}_u \right\rangle ds. \end{aligned}$$

Using the properties of L^2 projection \mathcal{P} , the Cauchy-Schwarz inequality, and the trace's inequality, we arrive at

$$\begin{aligned} \langle \vartheta_q \cdot \mathbf{n}, \theta_u - \widehat{\theta}_u \rangle &\leq h^{1/2} \|\vartheta_q \cdot \mathbf{n}\|_{\partial\mathcal{T}_h} \|h_K^{-1/2} (\theta_u - \widehat{\theta}_u)\|_{\partial\mathcal{T}_h} \\ &\leq C h^{l_q} \|\mathbf{q}\|_{H^{l_q}(\mathcal{T}_h)} \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}, \quad 0 \leq l_q \leq k+1 \end{aligned} \quad (6.11)$$

Similarly,

$$\begin{aligned} \left\langle \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \theta_u - \widehat{\theta}_u \right\rangle &\leq \|\tau h_K^{-1/2} \vartheta_u\|_{\partial\mathcal{T}_h} \|h_K^{-1/2} (\theta_u - \widehat{\theta}_u)\|_{\partial\mathcal{T}_h} \\ &\leq C h^{l_u-1} \|u\|_{H^{l_u}(\mathcal{T}_h)} \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}, \quad 1 \leq l_u \leq k+2 \end{aligned} \quad (6.12)$$

The non-linear term is dealt with similarly as done in Section 3. Hence, we find that

$$\|f(u) - f(u_h)\|^2 \leq C (\|\vartheta_u\|^2 + \|\theta_{\mathbf{q}}\|^2 + \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}^2). \quad (6.13)$$

Substituting (6.11), (6.12) and (6.13) in (6.10), and using the Young's inequality with kickback argument, we arrive at

$$\begin{aligned} \|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}^2 &\leq C \left(h^{2l_q} (\|\mathbf{q}\|_{H^{l_q}(\mathcal{T}_h)}^2 + \|\nabla u_0\|_{H^{l_q}(\mathcal{T}_h)}^2) + h^{2l_u-2} (\|u\|_{H^{l_u}(\mathcal{T}_h)}^2 + \|u_0\|_{H^{l_u}(\mathcal{T}_h)}^2) \right) \\ &\quad + C \int_0^t (h^{2l_u-2} \|u_t\|_{H^{l_u}(\mathcal{T}_h)}^2 + h^{2l_q} \|\mathbf{q}_t\|_{H^{l_q}(\mathcal{T}_h)}^2 + h^{2l_u} \|u\|_{H^{l_u}(\mathcal{T}_h)}^2) ds \\ &\quad + C \int_0^t (\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}^2) ds. \end{aligned} \quad (6.14)$$

Then, an application of Grönwall's inequality $1 \leq l_u \leq k+2$ and $0 \leq l_q \leq k+1$, yields

$$\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}^2 \leq C h^{2(k+1)} \left(\|u_0\|_{H^{k+2}}^2 + \|u\|_{H^{k+2}}^2 + \int_0^t (\|u_t\|_{H^{k+2}}^2 + \|\mathbf{q}_t\|_{H^{k+1}}^2 + \|u\|_{H^{k+2}}^2) ds \right). \quad (6.15)$$

Next, a use of the estimate of θ_u from article [26, Lemma 3.2] shows

$$\begin{aligned} \|\theta_u\|^2 &\leq C (\|\nabla\theta_u\|^2 + \|h_K^{-1/2}(\theta_u - \widehat{\theta}_u)\|_{\partial\mathcal{T}_h}^2), \\ &\leq C (\|\theta_{\mathbf{q}}\|^2 + \|h_K^{-1/2}(\mathcal{P}\theta_u - \widehat{\theta}_u)\|_{\partial\mathcal{T}_h}^2), \\ &\leq C (\|\theta_{\mathbf{q}}\|^2 + \|\theta_{u_t}\|^2 + \|\theta_u - \widehat{\theta}_u\|_{\partial\mathcal{T}_h}^2) \end{aligned}$$

Hence, we derive the optimal estimates of $\|\theta_u\|$ from (6.15). A use of Lemma 3.2 and Theorem 2.1 in (6.15) completes the proof of (6.3).

In order to prove (6.4), differentiate (6.7a) twice with respect to time and differentiate (6.7b)-(6.7d) and then choose $\mathbf{v}_h := \theta_{\mathbf{q}_t}$ and $w_h := \theta_{u_{tt}}$ to obtain

$$\frac{1}{2} \frac{d}{dt} (\|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_{tt}}\|^2) + \langle \theta_{\mathbf{q}_t} \cdot \mathbf{n}, \theta_{u_{tt}} - \widehat{\theta}_{u_{tt}} \rangle - \langle (\mathbf{q}_t - \widehat{\mathbf{q}}_{ht}) \cdot \mathbf{n}, \theta_{u_{tt}} \rangle + (f(u_t) - f(u_{ht}), \theta_{u_{tt}}) = 0. \quad (6.16)$$

Use of (6.8), after differentiating with respect to time shows

$$\frac{1}{2} \frac{d}{dt} (\|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_{tt}}\|^2 + \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_{\partial\mathcal{T}_h}^2) + \left\langle \vartheta_{\mathbf{q}_t} \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_{u_t}, \theta_{u_{tt}} - \widehat{\theta}_{u_{tt}} \right\rangle + (f(u_t) - f(u_{ht}), \theta_{u_{tt}}) = 0.$$

Proceed similarly as done in proving (6.3),

$$\begin{aligned} \int_0^t \left(\left\langle \vartheta_{\mathbf{q}_t} \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_{u_t}, \theta_{u_{tt}} - \widehat{\theta}_{u_{tt}} \right\rangle \right) ds &= \left\langle \vartheta_{\mathbf{q}_t} \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_{u_t}, \theta_{u_t} - \widehat{\theta}_{u_t} \right\rangle (t) \\ &\quad - \left\langle \vartheta_{\mathbf{q}_t} \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_{u_t}, \theta_{u_t} - \widehat{\theta}_{u_t} \right\rangle (0) \\ &\quad - \int_0^t \left\langle \vartheta_{\mathbf{q}_{tt}} \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_{u_{tt}}, \theta_{u_t} - \widehat{\theta}_{u_t} \right\rangle ds. \end{aligned}$$

Hence

$$\begin{aligned} \left| \left\langle \vartheta_{\mathbf{q}_t} \cdot \mathbf{n} + \frac{\tau}{h_K} \mathcal{P}\vartheta_{u_t}, \theta_{u_t} - \widehat{\theta}_{u_t} \right\rangle \right| &\leq C (h^{l_q} \|\mathbf{q}_t\|_{H^{l_q}(\mathcal{T}_h)} + h^{l_u-1} \|u_t\|_{H^{l_u}(\mathcal{T}_h)}) \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_{\partial\mathcal{T}_h}, \\ &\quad \forall 0 \leq l_q \leq k+1, 1 \leq l_u \leq k+2. \end{aligned}$$

The non-linear term is dealt with similarly to Section 3. Thus, using the Young's inequality with kickback argument,

$$\begin{aligned} \|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_{tt}}\|^2 + \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_{\partial\mathcal{T}_h}^2 &\leq C \left(h^{2l_q} (\|\nabla u_1\|_{H^{l_q}(\mathcal{T}_h)}^2 + \|\mathbf{q}_t\|_{H^{l_q}(\mathcal{T}_h)}^2) + h^{2l_u-2} (\|u_1\|_{H^{l_u}(\mathcal{T}_h)}^2 + \|u_t\|_{H^{l_u}(\mathcal{T}_h)}^2) \right) \\ &\quad + \int_0^t (h^{2l_u-2} \|u_{tt}\|_{H^{l_u}(\mathcal{T}_h)}^2 + h^{2l_q} \|\mathbf{q}_{tt}\|_{H^{l_q}(\mathcal{T}_h)}^2 + \|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_{\partial\mathcal{T}_h}^2 \\ &\quad + \|\theta_{u_{tt}}\|^2 + \|\vartheta_{u_t}\|^2) ds. \end{aligned}$$

Then, an application of Grönwall's inequality $1 \leq l_u \leq k+2$ and $0 \leq l_q \leq k+1$, yields

$$\|\theta_{\mathbf{q}_t}\|^2 + \|\theta_{u_{tt}}\|^2 + \|\theta_{u_t} - \widehat{\theta}_{u_t}\|_{\partial\mathcal{T}_h}^2 \leq C h^{2(k+1)} (\|u_1\|_{H^{k+2}}^2 + \|u_t\|_{H^{k+2}}^2 + \int_0^t (\|u_{tt}\|_{H^{k+2}}^2 + \|\mathbf{q}_{tt}\|_{H^{k+1}}^2) ds). \quad (6.17)$$

This completes the rest of the proof.

6.2 Super-convergence Results

This section derives the third result (6.5) of the Theorem 6.1. We use the same dual problem (4.5) defined in Section 4 with the same regularity results (4.6) and (4.7). Multiplying by θ_u to (4.5a) and integrating over Ω , then split the time derivative and use (4.5b)-(4.5d) to find that

$$(\Theta, \theta_u) = (\phi_s(0), \theta_u(0)) - (\phi(0), \theta_{u_s}(0)) + \int_0^t \left(-(\phi, \theta_{u_{ss}}) + (\nabla \cdot (\nabla \phi), \theta_u) - (f_u(u)\phi, \theta_u) \right) ds.$$

Here, we set $u_h(0) = \Pi_W u_0$ and $u_{ht}(0) = \Pi_W u_1$, which makes the first and second terms on the right-hand side zero. Using the L^2 projection and integration by parts yields

$$(\Theta, \theta_u) = \int_0^t \left(-(\phi, \theta_{u_{ss}}) + (\nabla \cdot (\mathbf{\Pi}_V \nabla \phi), \theta_u) + \langle (\nabla \phi - \mathbf{\Pi}_V \nabla \phi) \cdot \mathbf{n}, \theta_u \rangle - (f_u(u)\phi, \theta_u) \right) ds.$$

Using (6.7a) with $\mathbf{v} = \mathbf{\Pi}_V \nabla \phi$, we obtain

$$(\Theta, \theta_u) = \int_0^t \left(-(\phi, \theta_{u_{ss}}) - (\theta_{\mathbf{q}}, \mathbf{\Pi}_V \nabla \phi) + \langle \widehat{\theta}_u, \mathbf{\Pi}_V \nabla \phi \rangle + \langle (\nabla \phi - \mathbf{\Pi}_V \nabla \phi) \cdot \mathbf{n}, \theta_u \rangle - (f_u(u)\phi, \theta_u) \right) ds.$$

The second term on the right-hand side can be rewritten as

$$(\theta_{\mathbf{q}}, \mathbf{\Pi}_V \nabla \phi) = (\theta_{\mathbf{q}}, \mathbf{\Pi}_V \nabla \phi - \nabla \Pi_W \phi + \nabla \Pi_W \phi) = (\theta_{\mathbf{q}}, \nabla \Pi_W \phi) + (\theta_{\mathbf{q}}, \nabla \phi - \nabla \Pi_W \phi).$$

Hence, using (6.7b) for the term $(\theta_{\mathbf{q}}, \nabla \Pi_W \phi)$, we arrive at

$$\begin{aligned} (\Theta, \theta_u) &= \int_0^t \left(-(\phi, \theta_{u_{ss}}) + (\theta_{u_{ss}}, \Pi_W \phi) - \langle (\mathbf{q} - \widehat{\mathbf{q}}_h) \cdot \mathbf{n}, \Pi_W \phi \rangle + (f(u) - f(u_h), \Pi_W \phi) \right. \\ &\quad \left. - (\theta_{\mathbf{q}}, \nabla \phi - \nabla \Pi_W \phi) + \langle \widehat{\theta}_u, \mathbf{\Pi}_V \nabla \phi \rangle + \langle (\nabla \phi - \mathbf{\Pi}_V \nabla \phi) \cdot \mathbf{n}, \theta_u \rangle - (f_u(u)\phi, \theta_u) \right) ds, \\ &= Q_1 + Q_2 + Q_3 + Q_4 + Q_5 + Q_6 + Q_7 + Q_8. \end{aligned}$$

Substituting the identity (4.13) in terms Q_3 and Q_5 , a use of (6.7c) with properties of L^2 projection shows

$$\begin{aligned} \int_0^t Q_3 ds &= \langle (\mathbf{q} - \widehat{\mathbf{q}}_h)(0) \cdot \mathbf{n}, \mathcal{P}\phi(0) - \Pi_W \phi(0) \rangle \\ &\quad + \langle (\theta_{\mathbf{q}} - \vartheta_{\mathbf{q}}) \cdot \mathbf{n} - \frac{\tau}{h_K} (\mathcal{P}\theta_u - \widehat{\theta}_u) + \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \mathcal{P}\phi - \Pi_W \phi \rangle. \end{aligned}$$

And

$$\int_0^t Q_5 ds = ((\theta_{\mathbf{q}})(0), \nabla \phi(0) - \nabla \Pi_W \phi(0)) + \int_0^t (\theta_{\mathbf{q}_s}, \nabla \phi - \nabla \Pi_W \phi) ds.$$

With the help of the Cauchy Schwarz inequality, the trace inequality, and the properties of L^2 projections from (6.6), we now arrive at

$$\begin{aligned} |\langle (\theta_{\mathbf{q}} - \vartheta_{\mathbf{q}}) \cdot \mathbf{n}, \mathcal{P}\phi - \Pi_W \phi \rangle| &\leq Ch^{-1/2} (\|\theta_{\mathbf{q}} \cdot \mathbf{n}\| + \|\vartheta_{\mathbf{q}} \cdot \mathbf{n}\|) \|\phi - \Pi_W \phi\|, \\ &\leq Ch (\|\theta_{\mathbf{q}} \cdot \mathbf{n}\| + \|\vartheta_{\mathbf{q}} \cdot \mathbf{n}\|) \|\phi\|_2, \end{aligned} \quad (6.18)$$

similar proceed for the other two terms,

$$\left| \left\langle \frac{\tau}{h_K} \mathcal{P}\vartheta_u, \mathcal{P}\phi - \Pi_W \phi \right\rangle \right| \leq Ch^{-3/2} \|\vartheta_u\| \|\phi - \Pi_W \phi\| \leq Ch^{k+2} \|u\|_{H^{k+2}} \|\phi\|_2, \quad (6.19)$$

similarly

$$\left| \left\langle \frac{\tau}{h_K} (\mathcal{P}\theta_u - \widehat{\theta}_u) \right\rangle \right| \leq Ch^{k+2} \|u\|_{H^{k+2}} \|\underline{\phi}\|_2. \quad (6.20)$$

Combining (6.18), (6.19) and (6.20) gives the estimate for Q_3 . Estimate (6.17), help us to find the bound for Q_5 . Q_6 and Q_7 can be combined by adding the zero term $\langle \widehat{\theta}_u, \nabla \phi \cdot \mathbf{n} \rangle$ as $\widehat{\theta}_u$ is single-valued across the interior faces and us the identity (4.13), therefore,

$$Q_6 + Q_7 = \langle (\theta_u - \widehat{\theta}_u)(0), (\nabla \underline{\phi}(0) - \mathbf{\Pi}_V \nabla \underline{\phi}(0)) \cdot \mathbf{n} \rangle + \int_0^t \langle \theta_{u_s} - \widehat{\theta}_{u_s}, (\nabla \underline{\phi} - \mathbf{\Pi}_V \nabla \underline{\phi}) \cdot \mathbf{n} \rangle$$

$$\begin{aligned} |\langle \theta_{u_s} - \widehat{\theta}_{u_s}, (\nabla \underline{\phi} - \mathbf{\Pi}_V \nabla \underline{\phi}) \cdot \mathbf{n} \rangle| &\leq \|\theta_{u_s} - \widehat{\theta}_{u_s}\| \|\nabla(\phi - I_h \phi)\| \\ &\leq C h^{k+2} (\|u_1\|_{H^{k+2}}^2 + \|u_t\|_{H^{k+2}}^2 + \int_0^t \|u_{tt}\|_{H^{k+2}}^2 ds) \|\underline{\phi}\|_{H^2}. \end{aligned}$$

For the nonlinear terms, we rewrite as

$$Q_4 + Q_8 = (f_u(u)\theta_u, \Pi_W \phi - \phi) + (f_u(u)\vartheta_u, \Pi_W \phi) + (f_{uu}(u)(u - u_h)^2, \Pi_W \phi).$$

To complete the derivation, we proceed by bounding each term systematically. Using the estimates of $\theta_{u_{ss}}$ from (6.4), we bound Q_1 and Q_2 . For Q_5 , proceed similarly as done in subsection 4.2. At last, combining all the terms, applying regularity results (4.6) and (4.7) with the properties of L^2 projections, we derive the required result. This completes the rest of the proof.

7 Numerical Experiments

This section illustrates three numerical examples to highlight our proposed methods' convergence, superconvergence, and energy conservation properties.

Example-1: In this example, we consider the domain $\Omega = [0, 1] \times [0, 1]$ with homogeneous Dirichlet boundary conditions. The source terms $f_1(x, y, t)$, and the initial conditions are chosen so that the exact solution $u(x, y, t) = t^2 \sin(\pi x) \sin(\pi y)$ for the nonlinear Klein-Gordon equation

$$\begin{aligned} u_{tt} - \Delta u + u^3 - u &= f_1(x, y, t) && \text{in } [0, 1] \times [0, 1] \times (0, T], \\ u|_{t=0} &= u_0, \quad u_t|_{t=0} = u_1 && \text{in } [0, 1] \times [0, 1]. \end{aligned}$$

The second-order discrete-time Galerkin scheme is applied for the time discretization and the space discretization; we choose polynomial degrees $k = 1, 2, 3$ with mesh parameters $h = 1/2^m$, for $m = 1, 2, 3, 4$. The time step is chosen to be $\Delta t = h^{(k+1)/2}$ with final time $T = 1$. Table 1 confirms our numerical results with our theoretical findings.

Example-2: With the same problem and domain as in Example 1 and the chosen source term $f_2(x, y, t)$, and the initial conditions so that the exact solution $u = \exp(2t^2) \sin(\pi x) \sin(\pi y)$. In Table 2, we observe that numerical orders of convergence and superconvergence results are the same as have been predicted by our theoretical results.

Example-3: Next, we consider $u = \tanh(\frac{x}{\sqrt{3}} - t)$, can be the solution of nonlinear traveling wave. With the same problem and domain as mentioned in Example 1 and the chosen source term $f_2(x, y, t)$, with the initial conditions, we obtain the superconvergence for higher degree polynomials, see Table 3).

Example-4: We consider the following nonlinear wave equation in the unit square: $\Omega = [0, 1] \times [0, 1]$ with source terms $f_1(x, y, t) = 0$, with the given initial conditions,

$$\begin{aligned} u_{tt} - \Delta u + u^3 - u &= 0 && \text{in } [0, 1] \times [0, 1] \times (0, T], \\ u|_{t=0} &= 20x^2(1-x)^2y^2(1-y)^2, \quad u_t|_{t=0} = 2 \sin(2\pi x) \sin(2\pi y) && \text{in } [0, 1] \times [0, 1]. \end{aligned}$$

For this example, we do not have the explicit exact solution; therefore, we use energy conservation property. We choose polynomial degrees $k = 1$ with mesh parameters $h = 1/2^m$ for $m = 1, 2, 3, 4$. The time step is chosen to be $\Delta t = 0.1$ and $n = 1/\Delta t$ with final time $T = 1$. It is observed that our computational results in Table 4 are conforming to the energy conservation for $n \geq 1$ as has been observed theoretically.

k	m	$\ u - u_h\ $		$\ \mathbf{q} - \mathbf{q}_h\ $		$\ u - u_h^*\ $	
		Error	E.O.C	Error	E.O.C	Error	E.O.C
1	1	1.83e-1		5.44e-1		4.50e-2	
	2	6.46e-2	1.5101	1.47e-1	1.8912	5.20e-3	3.1017
	3	1.34e-2	2.2697	2.98e-2	2.2994	5.00e-4	3.2694
	4	3.40e-3	1.9631	7.40e-3	2.0021	1.00e-4	3.0221
2	1	4.21e-2		7.74e-2		5.00e-3	
	2	4.80e-3	3.1226	9.90e-3	2.9640	3.00e-4	4.0556
	3	6.00e-4	3.0169	1.20e-3	3.0688	1.00e-5	4.0999
	4	1.00e-4	3.0158	1.00e-4	3.0253	1.00e-6	4.0116
3	1	8.60e-3		2.11e-2		1.30e-3	
	2	5.00e-4	4.1152	1.20e-3	4.1710	1.00e-4	5.1703
	3	1.00e-5	4.0331	1.00e-4	4.0807	1.00e-5	5.0574
	4	1.00e-6	3.9941	1.00e-5	4.0138	1.00e-6	5.0117

Table 1: The L_2 -Errors and Estimated Orders of Convergence for Example 1.

k	m	$\ u - u_h\ $		$\ \mathbf{q} - \mathbf{q}_h\ $		$\ u - u_h^*\ $	
		Error	E.O.C	Error	E.O.C	Error	E.O.C
1	1	6.77e-1		5.23e-0		6.65e-1	
	2	1.83e-1	1.7470	1.64e-0	1.6729	5.54e-2	2.7870
	3	9.18e-2	2.0836	5.43e-1	1.7944	1.18e-2	3.0439
	4	5.65e-3	1.9906	1.61e-1	1.9502	7.93e-3	2.9804
2	1	6.52e-1		1.08e-0		9.04e-2	
	2	3.57e-2	4.1904	1.89e-1	2.5145	1.30e-2	3.7951
	3	3.60e-3	3.3099	3.06e-2	2.6259	8.00e-4	4.0171
	4	5.00e-4	2.9537	6.30e-3	2.8868	1.00e-4	4.1590
3	1	6.28e-2		5.33e-1		5.91e-2	
	2	3.09e-3	4.0203	2.54e-2	4.3924	2.30e-3	4.8996
	3	2.00e-4	4.1077	1.10e-3	4.5548	1.00e-4	4.9844
	4	1.39e-5	4.0046	9.12e-5	4.5665	2.79e-6	4.9810

Table 2: The L_2 -Errors and Estimated Orders of Convergence for Example 2.

k	m	$\ u - u_h\ $		$\ \mathbf{q} - \mathbf{q}_h\ $		$\ u - u_h^*\ $	
		Error	E.O.C	Error	E.O.C	Error	E.O.C
1	1	4.44e-2		2.95e-2		8.89e-4	
	2	6.60e-3	2.4571	6.70e-3	2.1305	1.60e-4	2.8746
	3	1.20e-3	2.4239	1.80e-3	1.9218	2.00e-4	2.4726
	4	1.00e-4	2.2456	7.00e-4	2.0021	1.00e-5	3.0256
2	1	6.50e-3		1.59e-2		3.50e-4	
	2	3.50e-3	4.3343	1.00e-3	3.9765	1.09e-5	4.3122
	3	2.00e-4	3.9873	1.00e-4	2.7893	1.00e-5	3.9956
	4	1.00e-4	3.4589	1.00e-5	3.0253	1.00e-6	4.0116
3	1	8.60e-3		7.60e-3		1.60e-3	
	2	1.60e-4	5.1964	3.20e-4	5.1847	1.30e-5	5.1916
	3	1.00e-5	5.0087	1.00e-4	5.0003	1.00e-5	5.0079
	4	1.00e-6	5.0415	1.00e-5	5.0012	1.00e-6	5.0117

Table 3: The L_2 -Errors and Estimated Orders of Convergence for Example 3.

n	$m = 1$	$m = 2$	$m = 3$	$m = 4$
1	0.6402	0.0968	0.4813	0.3576
2	1.3878e-17	3.5534e-15	1.7765e-14	3.1972e-14
3	1.3878e-17	1.7767e-15	1.7765e-14	5.3300e-15
4	1.3878e-17	5.3297e-15	1.1421e-14	3.5500e-15
5	1.3878e-17	7.1052e-15	1.0660e-14	5.5329e-14
6	2.7756e-17	8.8823e-15	2.4876e-14	3.3730e-14
7	2.7756e-17	0.0000e-0	1.0660e-14	5.5070e-14
8	4.1633e-17	3.5537e-15	3.5534e-14	9.2375e-14
9	4.1633e-17	3.5527e-15	2.1324e-14	3.7340e-15
10	1.3878e-17	0.0000e-0	1.4213e-14	8.5278e-14

Table 4: Error Energy $|\mathbb{E}^{n+1/2} - \mathbb{E}^{1/2}|$ for Example 4.

8 Summary and Conclusion

In this study, we have developed and analyzed the HDG approach for solving the nonlinear Klein-Gordon equation. The nonlinear function being odd degree polynomials is locally Lipschitz. HDG projection was used to derive the error estimates for the semi-discrete case. Further, element-by-element post-processing has been proposed. It has been demonstrated that the post-processed discrete displacement has super-convergence property, which is of order $k + 2, k \geq 1$. In contrast, the discrete displacement and its gradient have optimal rates of convergence, that is, of order $k + 1, k \geq 0$, when piecewise polynomials of degree $k \geq 1$ are used to approximate both the displacement and its flux variables. To discretize in time, a second-order conservative finite difference scheme is used. It is shown that discrete energy is conserved for $n \geq 1$, and a priori error estimates are derived. A non-conservative scheme is also suggested, which computes the discrete solutions via a linear system of algebraic equations at each time level, and its optimal error estimates are briefly addressed. A variant of the HDG method is also analyzed, along with error analysis in brief. Finally, several computational experiments are conducted, the results of which confirm our theoretical findings. The present analysis can be easily extended to problems with odd degree polynomial nonlinearity or to sine Gordon equation.

References

- [1] T. Achouri. Finite difference schemes for the two-dimensional semilinear wave equation. *Numerical Methods for Partial Differential Equations*, 35(1):200–221, 2019.

- [2] M. Baccouch. Optimal error estimates of the local discontinuous Galerkin method for the two-dimensional sine-Gordon equation on cartesian grids. *International Journal of Numerical Analysis & Modeling*, 16(3):436–462, 2019.
- [3] G. A. Baker. Error estimates for finite element methods for second order hyperbolic equations. *SIAM journal on numerical analysis*, 13(4):564–576, 1976.
- [4] F. Brezzi, J. Douglas, and L. D. Marini. Two families of mixed finite elements for second order elliptic problems. *Numerische Mathematik*, 47:217–235, 1985.
- [5] G. Chen, B. Cockburn, J. Singler, and Y. Zhang. Superconvergent interpolatory HDG methods for reaction diffusion equations. *Journal of Scientific Computing*, 81(3):2188–2212, 2019.
- [6] B. Cockburn. The hybridizable discontinuous Galerkin methods. In *Proceedings of the International Congress of Mathematicians 2010, (volume 4) Vol. I: Plenary Lectures and Ceremonies Vols. II–IV: Invited Lectures*, pages 2749–2775. World Scientific, 2010.
- [7] B. Cockburn. Hybridizable discontinuous Galerkin methods for second-order elliptic problems: overview, a new result and open problems. *Japan Journal of Industrial and Applied Mathematics*, 40(3):1637–1676, 2023.
- [8] B. Cockburn, Z. Fu, A. Hungria, L. Ji, M. A. Sanchez, and F.J. Sayas. Stormer-Numerov HDG methods for acoustic waves. *Journal of Scientific Computing*, 75(2):597–624, 2018.
- [9] B. Cockburn and J. Gopalakrishnan. The derivation of hybridizable discontinuous Galerkin methods for stokes flow. *SIAM Journal on Numerical Analysis*, 47(2):1092–1125, 2009.
- [10] B. Cockburn, J. Gopalakrishnan, and R. Lazarov. Unified hybridization of discontinuous Galerkin, mixed, and continuous Galerkin methods for second order elliptic problems. *SIAM Journal on Numerical Analysis*, 47(2):1319–1365, 2009.
- [11] B. Cockburn, J. Gopalakrishnan, and F.-J. Sayas. A projection-based error analysis of HDG methods. *Mathematics of Computation*, 79(271):1351–1367, 2010.
- [12] B. Cockburn, W. Qiu, and K. Shi. Conditions for superconvergence of HDG methods for second-order elliptic problems. *Mathematics of Computation*, 81(279):1327–1353, 2012.
- [13] B. Cockburn and V. Quenneville-Bélair. Uniform-in-time superconvergence of the HDG methods for the acoustic wave equation. *Mathematics of Computation*, 83(285):65–85, 2014.
- [14] B. Cockburn and K. Shi. Conditions for superconvergence of HDG methods for Stokes flow. *Mathematics of Computation*, 82(282):651–671, 2013.
- [15] B. Cockburn, J. R. Singler, and Y. Zhang. Interpolatory HDG method for parabolic semilinear PDEs. *Journal of Scientific Computing*, 79(3):1777–1800, 2019.
- [16] J. F. Currie, J. A. Krumhansl, A. R. Bishop, and S. E. Trullinger. Statistical mechanics of one-dimensional solitary-wave-bearing scalar fields: Exact results and ideal-gas phenomenology. *Physical Review B*, 22(2):477, 1980.
- [17] D. B. Duncan. Symplectic finite difference approximations of the nonlinear Klein-Gordon equation. *SIAM Journal on numerical analysis*, 34(5):1742–1760, 1997.
- [18] S. Jiménez and L. Vázquez. Analysis of four numerical schemes for a nonlinear Klein-Gordon equation. *Applied Mathematics and Computation*, 35(1):61–94, 1990.
- [19] C. Johnson and V. Thomée. Error estimates for some mixed finite element methods for parabolic type problems. *RAIRO. Analyse numérique*, 15(1):41–78, 1981.
- [20] L. Kapitanski. Global and unique weak solutions of nonlinear wave equations. *Mathematical Research Letters*, 1(2):211–223, 1994.
- [21] S. Kesavan. *Topics in Functional Analysis and Applications*. New Age International Pvt Limited Publishers, 2015.

- [22] R. M. Kirby, S. J. Sherwin, and B. Cockburn. To HDG or to CG: A comparative study. *Journal of Scientific Computing*, 51(1):183–212, 2012.
- [23] H. Kragh. Equation with the many fathers. The Klein–Gordon equation in 1926. *American Journal of Physics*, 52(11):1024–1033, 1984.
- [24] A. K. Pani and J. Y. Yuan. Mixed finite element method for a strongly damped wave equation. *Numerical Methods for Partial Differential Equations: An International Journal*, 17(2):105–119, 2001.
- [25] W. Qiu, J. Shen, and K. Shi. An HDG method for linear elasticity with strong symmetric stresses. *Mathematics of Computation*, 87(309):69–93, 2018.
- [26] W. Qiu and K. Shi. An HDG method for convection diffusion equation. *Journal of Scientific Computing*, 66(1):346–357, 2016.
- [27] N. Sánchez, T. Sánchez-Vizuet, and M. Solano. Error analysis of an unfitted HDG method for a class of non-linear elliptic problems. *Journal of Scientific Computing*, 90(3):1–28, 2022.
- [28] N. Sánchez, T. Sánchez-Vizuet, and M. E. Solano. A priori and a posteriori error analysis of an unfitted HDG method for semi-linear elliptic problems. *Numerische Mathematik*, 148(4):919–958, 2021.
- [29] R. K. Sinha and A. K. Pani. The effect of spatial quadrature on finite element galerkin approximations to hyperbolic integro-differential equations. *Numerical functional analysis and optimization*, 19(9-10):1129–1153, 1998.
- [30] Y. Tourigny. Product approximation for nonlinear Klein-Gordon equations. *IMA journal of Numerical Analysis*, 10(3):449–462, 1990.
- [31] Y. Wong, Q. Chang, and L. Gong. An initial-boundary value problem of a nonlinear Klein-Gordon equation. *Applied Mathematics and Computation*, 84(1):77–93, 1997.
- [32] S. Yadav and A. K. Pani. Superconvergent discontinuous Galerkin methods for nonlinear parabolic initial and boundary value problems. *Journal of Numerical Mathematics*, 27(3):183–202, 2019.