Stability properties of gradient flow dynamics for the symmetric low-rank matrix factorization problem

Hesameddin Mohammadi, Mohammad Tinati, Stephen Tu, Mahdi Soltanolkotabi, and Mihailo R. Jovanović

Abstract—The symmetric low-rank matrix factorization serves as a building block in many learning tasks, including matrix recovery and training of neural networks. However, despite a flurry of recent research, the dynamics of its training via non-convex factorized gradient-descent-type methods is not fully understood especially in the over-parameterized regime where the fitted rank is higher than the true rank of the target matrix. To overcome this challenge, we characterize equilibrium points of the gradient flow dynamics and examine their local and global stability properties. To facilitate a precise global analysis, we introduce a nonlinear change of variables that brings the dynamics into a cascade connection of three subsystems whose structure is simpler than the structure of the original system. We demonstrate that the Schur complement to a principal eigenspace of the target matrix is governed by an autonomous system that is decoupled from the rest of the dynamics. In the over-parameterized regime, we show that this Schur complement vanishes at an O(1/t) rate, thereby capturing the slow dynamics that arises from excess parameters. We utilize a Lyapunov-based approach to establish exponential convergence of the other two subsystems. By decoupling the fast and slow parts of the dynamics, we offer new insight into the shape of the trajectories associated with local search algorithms and provide a complete characterization of the equilibrium points and their global stability properties. Such an analysis via nonlinear control techniques may prove useful in several related overparameterized problems.

I. INTRODUCTION

Training of massive over-parameterized models that contain many more parameters than training data via first-order algorithms such as (stochastic) gradient descent is a work-horse of modern machine learning. Since these learning problems are typically non-convex, it is not clear if first-order methods converge to a global optimum. Furthermore, because of overparameterization, even when global convergence does occur, the computed solution may not have good generalization properties.

Recent years have witnessed a surge of research efforts aimed at demystifying the complexities of optimization and generalization in such problems. A number of papers demonstrate local convergence of search techniques starting from carefully designed spectral initializations [1]-[9]. Another set of publications focuses on showing that the optimization landscape is favorable, i.e., that all local minima are global, with saddle points exhibiting strict negative curvature [10]. In these, algorithms like trust region methods, cubic regularization [11], [12], and stochastic gradient techniques [13]-[16] are employed to find global optima. There is a growing consensus that a deeper, more detailed analysis of gradient descent trajectories is needed beyond merely examining the optimization landscape [17] or specialized initialization techniques. While a number of papers have started to address this challenge, they typically require very intricate and specialized proofs [18]-[21] that often are not able to fully characterize the trajectory from various initialization points or precisely capture the speed of convergence.

In this letter, we use a control-theoretic approach to examine the behavior of gradient flow dynamics for solving the symmetric low-rank matrix factorization problem; the rank-one case was addressed in [22]. This problem serves as a building block for several non-convex learning tasks, including matrix completion [23], [24], and deep neural networks [25], and it has received significant attention in the literature [26]–[28]. Although the optimization landscape of the matrix factorization is benign in the sense that all local minima are globally optimal [29], the behavior of first-order algorithms in solving this problem is not yet fully understood. In addition, establishing global convergence rates for gradientbased methods applied to a non-convex optimization problem is of interest on its own.

Our control-theoretic approach provides guarantees and offers insight into an important class of non-convex optimization problems that are of interest in machine learning. Compared to existing literature [9], [19], [20] and more recent granular convergence guarantees [30], our approach reveals novel hidden structures and it serves as an important first step towards simpler convergence proofs for commonly-used variants of gradient descent. For example, stochastic gradient descent is still stateof-the-art in most vision applications and examining gradient descent is a well-established first step towards the analysis of these more complex algorithms. We take a step toward providing insight into the global behavior of such algorithms by exploiting the structural properties of the underlying dynamics.

In our analysis, we exploit the lifted coordinates that bring the system into a Riccati-like set of dynamical systems. Our main contribution lies in introducing a novel change of variables that brings the underlying dynamics into a cascade connection of three subsystems. We show that one of these subsystems is governed by autonomous dynamics which is decoupled from the rest of the system. In the over-parameterized regime, this subsystem is associated with the excess parameters; our analysis reveals its stability with an O(1/t) asymptotic convergence rate, which captures the slow dynamics. For the other two subsystems, we utilize a Lyapunov-based approach to establish their exponential convergence under mild assumptions.

Our analysis offers new insight into the shape of the trajectories of first-order algorithms in the presence of overparametrization. It also provides a complete characterization of their behavior by decoupling fast and slow dynamical components. In contrast to many existing works, we present our results in their most general form, without making any assumptions about the initialization or the structure of the target matrix. This generality enables our approach to be applied in a wider variety of settings, from small random initializations to large over-parameterized scenarios.

The rest of this letter is as follows. In section II, we

formulate the symmetric low-rank matrix factorization problem and introduce the corresponding gradient flow dynamics. In section III, we characterize the equilibrium points and examine their local stability properties. Additionally, we introduce the signal/noise decomposition of the optimization variable, which paves the way for global stability analysis. In Section IV, we introduce a novel change of variables that transforms the original problem into a cascade connection of three subsystems, thereby facilitating the proof of global convergence. We also show that matrix recovery is impossible over certain invariant manifolds. Finally, we demonstrate that the optimal point is globally asymptotically stable if the initial condition lies outside the invariant manifold. In Section VI, we offer concluding remarks, and in appendices, we provide proofs.

II. PROBLEM FORMULATION

We study the matrix factorization problem,

$$\underset{X}{\text{minimize }} f(X) := (1/4) \| X X^T - M \|_F^2 \qquad (1)$$

where $M \in \mathbb{R}^{n \times n}$ is a given symmetric matrix, $X \in \mathbb{R}^{n \times r}$ is the optimization variable, and $\|\cdot\|_F$ is the Frobenius norm. This is a non-convex optimization problem [19], and our objective is to find a matrix X such that XX^T approximates M. The gradient flow dynamics associated with (1) are given by

$$\dot{X} = -\nabla_X f(X) = \left(M - XX^T\right) X.$$
 (2)

Let $M = V\Lambda V^T$ be the eigenvalue decomposition of the matrix M where V is the orthogonal matrix of eigenvectors and Λ is the diagonal matrix of eigenvalues of M with

$$\Lambda = \operatorname{diag}(\Lambda_1, -\Lambda_2). \tag{3}$$

Here, $\Lambda_1 \in \mathbb{R}^{r^* \times r^*}$ and $-\Lambda_2 \in \mathbb{R}^{(n-r^*) \times (n-r^*)}$ are diagonal matrices of positive $\{\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_{r^*} > 0\}$ and non-positive eigenvalues of M, respectively.

The positive definite part $V \operatorname{diag}(\Lambda_1, 0) V^T$ of M can be written as $X^* X^{*T}$ for some $X^* \in \mathbb{R}^{n \times r}$ if and only if $r \ge r^*$. We refer to the regimes with $r = r^*$ and $r > r^*$ as the exact and over-parameterized, respectively, indicating presence of excess parameters in the latter. To simplify the analysis, we define a new variable $Z := V^T X \in \mathbb{R}^{n \times r}$ and rewrite system (2) as

$$\dot{Z} = (\Lambda - ZZ^T)Z. \tag{4}$$

By analyzing local and global stability properties of the equilibrium points of system (4), we establish guarantees for the convergence of gradient flow dynamics (2) to the optimal value of non-convex optimization problem (1). We also introduce a novel approach to characterize the convergence rate and offer insights into behavior of the dynamics in the over-parameterized regime and robustness of the recovery process.

III. LOCAL STABILITY ANALYSIS

Let $\overline{Z}\overline{Z}^T$ with $\overline{Z} \in \mathbb{R}^{n \times r}$ be a low rank decomposition of the matrix of positive eigenvalues of M, i.e., $\operatorname{diag}(\Lambda_1, 0) = \overline{Z}\overline{Z}^T$. Then, globally optimal solutions to non-convex problem (1) are parameterized by $Z = \overline{Z}G$, where $G \in \mathbb{R}^{r \times r}$ is an arbitrary unitary matrix. To resolve ambiguity caused by this lack of uniqueness, we introduce the lifted matrix $P = ZZ^T \in \mathbb{R}^{n \times n}$ and analyze properties of system (4) without the need to explicitly consider G. It is straightforward to verify that the matrix $P(t) := Z(t)Z^T(t)$ represents the unique solution of

$$\dot{P} = (\Lambda - P)P + P(\Lambda - P).$$
(5)

In what follows, we utilize the analysis of system (5) to deduce convergence properties of system (4).

A. Equilibrium points

We first identify the equilibrium points of system (5) and characterize their local stability properties. The set of equilibrium points of this system is determined by

$$\mathcal{H} := \left\{ \operatorname{diag}(\bar{P}_1, 0) \mid \bar{P}_1 \succeq 0, \ \bar{P}_1 \Lambda_1 = \bar{P}_1^2 \right\}$$
(6)

where $\bar{P}_1 \in \mathbb{R}^{r^* \times r^*}$; see Appendix A for the proof. Two trivial members of the equilibrium set \mathcal{H} are given by $\bar{P}_1 = 0$ and $\bar{P}_1 = \Lambda_1$. Lemma 1 provides an alternative characterization of the set \mathcal{H} which we exploit in our subsequent analysis.

Lemma 1: The positive semidefinite matrix $\overline{P} = \text{diag}(\overline{P}_1, 0)$ with $\overline{P}_1 \notin \{0, \Lambda_1\}$ is an equilibrium point of system (5) if and only if,

$$\bar{P}_1 = U_{\mathbf{i}} D_{\mathbf{i}} U_{\mathbf{i}}^T. \tag{7}$$

Here, $\Lambda_1 = \begin{bmatrix} U_{\mathbf{i}} & U_{\mathbf{o}} \end{bmatrix} \operatorname{diag}(D_{\mathbf{i}}, D_{\mathbf{o}}) \begin{bmatrix} U_{\mathbf{i}} & U_{\mathbf{o}} \end{bmatrix}^T$ provides a unitary similarity transformation of Λ_1 , $D_{\mathbf{i}} \in \mathbb{R}^{l \times l}$ and $D_{\mathbf{o}} \in \mathbb{R}^{(r^*-l) \times (r^*-l)}$ are diagonal matrices, $U_{\mathbf{i}} \in \mathbb{R}^{r^* \times l}$ and $U_{\mathbf{o}} \in \mathbb{R}^{r^* \times (r^*-l)}$ with $0 < l < r^*$ form a unitary matrix $U = \begin{bmatrix} U_{\mathbf{i}} & U_{\mathbf{o}} \end{bmatrix}$, and $\Lambda_1 - \bar{P}_1 = U_{\mathbf{o}} D_{\mathbf{o}} U_{\mathbf{o}}^T$.

Proof: See Appendix A.

The diagonal matrices $D_{\mathbf{i}} = U_{\mathbf{i}}^T \Lambda_1 U_{\mathbf{i}}$ and $D_{\mathbf{o}} = U_{\mathbf{o}}^T \Lambda_1 U_{\mathbf{o}}$ partition the eigenvalues of Λ_1 , i.e., $\operatorname{eig}(\Lambda_1) = \operatorname{eig}(D_{\mathbf{i}}) \cup$ $\operatorname{eig}(D_{\mathbf{o}})$, into those that are in the spectrum of \overline{P} and those that are not. If Λ_1 has distinct eigenvalues, then U can only be a permutation matrix and system (5) has exactly 2^{r^*} equilibrium points. In this case, the corresponding matrices \overline{P}_1 are diagonal with entries determined by either 0 or the eigenvalues λ_k of Λ_1 . If M has a repeated positive eigenvalue, there is a lack of uniqueness of the eigenvectors of Λ_1 , and thereby infinitely many matrices \overline{P}_1 of the form (7).

Example 1: For $\Lambda_1 = \text{diag}(3, 2, 1)$, apart from $\overline{P}_1 = \Lambda_1$ and $\overline{P}_1 = 0$, there are six additional members of the equilibrium set \mathcal{H} , i.e., $\overline{P}_1 \in \{\text{diag}(3, 0, 0), \text{diag}(0, 2, 0), \text{diag}(0, 0, 1), \text{diag}(3, 2, 0), \text{diag}(3, 0, 1), \text{diag}(0, 2, 1)\}.$

Remark 1: Lemma 1 implies that $\bar{P}_1 = \text{diag}(\Lambda_1, 0)$ is the only equilibrium point with rank r^* . Since the remaining nonzero equilibrium points are given by $U_i D_i U_i^T$ for some diagonal $D_i \in \mathbb{R}^{l \times l}$ with $l < r^*$, their rank is smaller than r^* .

B. Local stability properties

To examine local stability properties, we linearize system (5) around its equilibrium point $\bar{P} \in \mathcal{H}$. Substituting $P = \bar{P} + \epsilon \tilde{P}$ to (5) and keeping $O(\epsilon)$ terms leads to the linearized dynamics,

$$\dot{\tilde{P}} = (\Lambda - 2\bar{P})\tilde{P} + \tilde{P}(\Lambda - 2\bar{P})$$
(8)

which allows us to characterize stability properties for $\bar{P}_1 \neq \Lambda_1$.

Proposition 1: Any equilibrium point $\overline{P} = \text{diag}(\overline{P}_1, 0) \in \mathcal{H}$ of system (5) with $\overline{P}_1 \neq \Lambda_1$ is unstable. *Proof:* Let \overline{P}_1 be parameterized by (D_i, D_o, U_i, U_o) , as described in Lemma 1. Now, consider the mode $q := u^T \tilde{P}u$, where $u := [u_k^T \ 0]^T \in \mathbb{R}^n$ and u_k is the kth column of U_o . For linearized system (8), we have

$$\dot{q} = 2u^T (-\bar{P} + (\Lambda - \bar{P}))\tilde{P}u = 2u^T (\Lambda - \bar{P})\tilde{P}u = 2\lambda q$$

where $\lambda > 0$ is an eigenvalue of Λ_1 that corresponds to the *k*th diagonal entry of D_0 . Since U_i and U_0 are mutually orthogonal, the second equality follows from $u^T \bar{P} = u_k^T U_i D_i U_i^T = 0$, and the last equality follows from the second equation in (7). Thus, q is an unstable mode of the linearized system, and \bar{P} is an unstable equilibrium point of nonlinear system (5). The analysis for $\bar{P} = 0$ is similar and is omitted for brevity.

Lemma 2 proves that $\operatorname{diag}(\Lambda_1, 0)$ is an isolated equilibrium point by establishing a lower bound on the distance between $\operatorname{diag}(\Lambda_1, 0)$ and all other equilibria of system (5).

Lemma 2: For any equilibrium point $\bar{P} \neq \text{diag}(\Lambda_1, 0)$ of system (5), we have $\|\bar{P} - \text{diag}(\Lambda_1, 0)\|_2 \ge \lambda_{r^*}$, where $\lambda_r^* > 0$ is the smallest eigenvalue of Λ_1 and $\|\cdot\|_2$ is the spectral norm.

Proof: The result is trivial for $\overline{P} = 0$. For $\overline{P} \neq 0$, Lemma 1 implies that $\|\overline{P} - \operatorname{diag}(\Lambda_1, 0)\|_2 = \|\overline{P}_1 - \Lambda_1\|_2 = \|D_0\|_2 \ge \lambda_{r^*}$, where the inequality follows because the diagonal matrix D_0 contains a nonempty subset of the eigenvalues of Λ_1 .

Using Lemma 2 in conjunction with a Lyapunov-based argument, we prove local asymptotic stability of the global minimum $\bar{P} = \text{diag}(\Lambda_1, 0)$ of optimization problem (1).

Proposition 2: The isolated equilibrium point $\overline{P} = \text{diag}(\Lambda_1, 0)$ of system (5) is locally asymptotically stable.

Proposition 2 proves local asymptotic stability of diag($\Lambda_1, 0$) using the objective function in (1) as a Lyapunov function candidate, i.e., $V_F(P) = f(X)$. However, this approach does not provide insight into global stability properties because of the presence of additional equilibrium points.

C. Signal/noise decomposition

To prove global asymptotic stability of $\overline{P} = \text{diag}(\Lambda_1, 0)$, we decompose Z into "signal" Z_1 and "noise" Z_2 components,

$$Z^T = \begin{bmatrix} Z_1^T & Z_2^T \end{bmatrix}$$
(9)

where $Z_1 \in \mathbb{R}^{r^* \times r}$ captures the part of Z that is desired to get aligned with the positive definite low-rank structure, i.e, $\lim_{t\to\infty} Z_1(t)Z_1^T(t) = \Lambda_1$, and $Z_2 \in \mathbb{R}^{(n-r^*)\times r}$ is supposed to vanish asymptotically. This decomposition allows us to isolate contribution of "noise", thereby facilitating analysis of the stability properties of the gradient flow dynamics.

Using decomposition (9) of Z we can rewrite system (4) as

$$\dot{Z}_1 = (\Lambda_1 - Z_1 Z_1^T) Z_1 - Z_1 Z_2^T Z_2$$
 (10a)

$$\dot{Z}_2 = -(\Lambda_2 + Z_2 Z_2^T) Z_2 - Z_2 Z_1^T Z_1$$
 (10b)

and by partitioning P conformably with the partition of Z,

$$P = \begin{bmatrix} P_1 & P_0 \\ P_0^T & P_2 \end{bmatrix} := \begin{bmatrix} Z_1 Z_1^T & Z_1 Z_2^T \\ Z_2 Z_1^T & Z_2 Z_2^T \end{bmatrix} = Z Z^T$$

the corresponding lifted system (5) can be written as,

$$P_1 = P_1 \Lambda_1 + \Lambda_1 P_1 - 2(P_1^2 + P_0 P_0^T)$$
(11a)

$$\dot{P}_0 = \Lambda_1 P_0 - P_0 \Lambda_2 - 2(P_1 P_0 + P_0 P_2)$$
(11b)

$$\dot{P}_2 = -P_2\Lambda_2 - \Lambda_2P_2 - 2(P_0^T P_0 + P_2^2).$$
 (11c)

Here, P_1 and P_2 represent the "signal" and "noise" components of P, and P_0 accounts for the coupling between Z_1 and Z_2 . We note that for $r^* = n$ the matrices Z_2 , P_2 , and P_0 disappear.

D. The evolution of the "noise" component P_2

We next examine the evolution of the noise component $P_2 := Z_2 Z_2^T$ and identify the conditions under which system (5) converges to the stable equilibrium point $\overline{P} = \text{diag}(\Lambda_1, 0)$. We show that $P_2(t)$ decays with time and that it asymptotically vanishes as the system approaches the stable equilibrium point.

For any initial condition, we first prove that the spectral norm of the noise component P_2 ,

$$V_N(P_2) := ||P_2||_2 = ||Z_2||_2^2$$
 (12)

converges to zero with a polynomial rate.

Lemma 3: Along the trajectories of system (5), the spectral norm of the noise component $P_2(t)$ satisfies

$$V_N(P_2(t)) \leq 2V_N(P_2(\tau))/(1+t-\tau), \ \forall t \geq \tau > 0.$$

Proof: See Appendix B.

Lemma 3 implies that $V_N(P_2(t))$ is a decreasing function of time that converges to zero with O(1/t) rate. While we demonstrate in Section IV that this result can be improved to exponential convergence for $r = r^*$, we next establish its sharpness for the over-parameterized regime.

Sharpness of Lemma 3 in the over-parameterized regime: If $r > r^*$, we demonstrate through an example that the rate O(1/t) in Lemma 3 cannot be improved. In particular, for $\Lambda_2 = 0$, we identify an invariant manifold of system (5) over which the magnitude of the noise component $P_2(t)$ decreases with O(1/t) rate. In particular, let $Z^T(0) = [Z_1^T(0) Z_2^T(0)]$ be such that $P_1(0) = \Lambda_1$, $P_0(0) = 0$, and $P_2(0) \neq 0$; e.g., this can be achieved with $Z_1(0) = [\Lambda_1^{1/2} \quad 0_{r^* \times (r-r^*)}]$ where the rows of $Z_2(0) \neq 0$ belong to the orthogonal complement of the row space of $Z_1(0)$, which is nonempty because $r > r^*$. It is now easy to verify that $\dot{P}_1 = 0$ and $\dot{P}_0 = 0$, thereby implying $P_1(t) = \Lambda_1$ and $P_0(t) = 0$ for all $t \geq 0$. In this case, system (5) simplifies to $\dot{P}_2 = -2P_2^2$, the eigenvectors of P_2 remain unchanged, and each nonzero eigenvalue $\mu > 0$ of P_2 satisfies $\mu(t) = 2\mu(0)/(1 + t)$.

IV. MAIN RESULT: GLOBAL STABILITY ANALYSIS

Our main theoretical result establishes global convergence of the gradient flow dynamics to $\operatorname{diag}(\Lambda_1, 0)$ if and only if the initial condition P(0) satisfies $\operatorname{rank}(P_1(0)) = r^*$. This condition guarantees recovery of the optimal low-rank positive semi-definite factorization of the target matrix $M = M^T$.

Theorem 1: For any initial condition P(0) of system (5) with $P_1(0) \succ 0$, let the time t be large enough for the exponentially decaying term in function

$$l(t) := \left(\|P_1^{-1}(0) - \Lambda_1^{-1}\|_2 + 2t \|P_1^{-1}(0)P_0(0)\|_2^2 \right) e^{-2\lambda_{r^\star} t}$$

to dominate the linearly growing term such that $l(t) \le 1/(2\lambda_1)$. is governed by the following dynamical system Then, $P_1(t) \succ 0$ for all times and the solution to (5) satisfies

$$\begin{aligned} \|P_1(t) - \Lambda_1\|_2 &\leq 2\,\lambda_1^2\,l(t) \\ \|P_0(t)\|_2 &\leq \|P_1^{-1}(0)\,P_0(0)\|_2\left(\lambda_1 + 2\,\lambda_1^2\,l(t)\right)\,\mathrm{e}^{-\lambda_{r^\star}t}. \end{aligned}$$

Proof: See Appendix C.

Theorem 1 establishes exponential asymptotic convergence of $P_1(t)$ to Λ_1 and of $P_0(t)$ to zero at a rate determined by the smallest positive eigenvalue λ_{r^*} of Λ_1 , provided that the initial condition satisfies $P_1(0) \succ 0$. In conjunction with Lemma 3, this proves global convergence to the optimal solution.

Remark 1 (Exponential convergence for the $r = r^*$): For the exact parameterization with $r = r^*$, the matrix Z_1 remains invertible under the conditions of Theorem 1 and it satisfies $P_1^{-1} = Z_1^{-T} Z_1^{-1}$. This allows us to write

$$P_2 = Z_2 Z_2^T = Z_2 Z_1^T P_1^{-1} Z_1 Z_2^T = P_0^T P_1^{-1} P_0^T$$

Combining this equality with Theorem 1, it is straightforward to verify that $P_2(t)$ vanishes at the exponential rate $-2\lambda_r$.

Corollary 1: For any full row rank matrix $Z_1(0) \in \mathbb{R}^{r^* \times r}$ and arbitrary $Z_2(0) \in \mathbb{R}^{(n-r^*) \times r}$, there exists a point $Z^* = [Z_1^{*T} \ 0]^T \in \mathcal{Z}^*$ and scalars $c_i > 0$ such that system (4) with the initial condition $Z(0) = \begin{bmatrix} Z_1^T(0) & Z_2^T(0) \end{bmatrix}^T$ satisfies

$$||Z_1(t) - Z_1^{\star}||_2 \le c_1 \mathrm{e}^{-\lambda_r \star t}, ||Z_2(t)|| \le c_2/\sqrt{t}$$

where $\mathcal{Z}^{\star} := \{ Z^{\star} \in \mathbb{R}^{n \times r} | Z^{\star} Z^{\star T} = \operatorname{diag}(\Lambda_1, 0) \}$ is the set of optimal solutions of problem (1) in the $Z = V^T X$ coordinates. Furthermore, for exact parameterization $r = r^{\star}$, the bound on $Z_2(t)$ improves to $||Z_2(t)||_2 \le c_2 e^{-\lambda_r \star t}$.

Remark 2: The condition $P_1(0) \succ 0$ in Theorem 1 holds if and only if the matrix $X^{\star T}X(0)$ has rank r^{\star} , where $X^{\star} \in$ $\mathbb{R}^{n \times r}$ is an optimal solution to problem (1) of rank r^* , and $X(0) \in \mathbb{R}^{n \times r}$ is the initial condition of (2). This can be interpreted as the matrices X(0) and X^* being well-aligned.

The key technical result that allows us to prove Theorem 1 is a novel nonlinear change of variables that we introduce in Section IV-A. In Section IV-B, we show that for any rank deficient initial condition $P_1(0)$, the trajectory P(t) of system (5) belongs to an invariant subspace that does not contain stable equilibrium point diag($\Lambda_1, 0$) and, in Section IV-C, we provide convergence guarantees in the new set of coordinates.

A. Change of variables

We next introduce a nonlinear change of variables that simplifies the analysis of system (5) and facilitates the proof of global convergence to $\bar{P} = \text{diag}(\Lambda_1, 0)$ for a full-rank $P_1(0)$.

Proposition 3: Let $P_1(t)$ be invertible for some t > 0. The evolution of the matrices,

$$\begin{aligned}
H_1 &:= P_1^{-1} &\in \mathbb{R}^{r^* \times r^*} \\
H_0 &:= P_1^{-1} P_0 &\in \mathbb{R}^{r^* \times (n-r^*)} \\
H_2 &:= P_2 - P_0^T P_1^{-1} P_0 \in \mathbb{R}^{(n-r^*) \times (n-r^*)}
\end{aligned} \tag{13}$$

$$H_1 = -\Lambda_1 H_1 - H_1 \Lambda_1 + 2(I + H_0 H_0^I)$$
(14a)

$$H_0 = -\Lambda_1 H_0 - H_0 (\Lambda_2 + 2H_2)$$
(14b)

$$H_2 = -\Lambda_2 H_2 - H_2 \Lambda_2 - 2H_2^2.$$
(14c)

Proof: See Appendix C.




Figure 1. Block diagram of system (14) illustrating its cascade structure.

In the H-coordinates, we have a cascade connection of three subsystems; see Fig. 1 for an illustration. While the H_2 dynamics are not influenced by the evolution of $H_1(t)$ and $H_0(t), H_2(t)$ enters as a coefficient into the H_0 -dynamics and $H_0(t)H_0^T(t)$ enters as an additive input into the H_1 -dynamics. It is also worth noting that the matrix $H_0(t)$ which satisfies Sylvester equation (14b) evolves independently of $H_1(0)$.

Remark 3: The matrix H_1 is determined by the inverse of $P_1, H_2 = P/P_1$ is the Schur complement of the matrix P with respect to P_1 , and H_0 in (13) is the best matrix that transforms the signal component Z_1 to the noise component Z_2 , i.e.,

$$H_0 = \operatorname*{argmin}_{E \in \mathbb{R}^{r^* \times (n-r^*)}} \|Z_2 - E^T Z_1\|_F^2.$$

B. Invariant subspaces

We next show that the null space of $P_1(t)$ does not change with time if $P_1(0)$ is singular. In this case, system (5) is restricted to a subspace and convergence to the full-rank solution $\bar{P}_1 = \Lambda_1$ is not possible. In other words, initial conditions with rank-deficient $P_1(0)$ lead to incomplete recovery. To show this, we note that for any fixed v with $v^T P_1 v = 0$ we can write

$$v^T \dot{P}_1 v = 2v^T (P_1 \Lambda_1 - (P_1^2 + P_0 P_0^T)) v = 0.$$

This expression follows from the fact that such v also belongs to the null space of Z_1^T and, hence, $P_0^T v = 0$. This implies that $v^T P_1(t)v$ remains equal to zero for all $t \ge 0$.

C. Convergence guarantees in the H-coordinates

Cascade connection (14) in transformed coordinates (13) allows us to examine stability properties of each H-component. For H_2 -subsystem (14c), we show global asymptotic stability of the origin with a worst-case convergence rate O(1/t), which is achieved when $\Lambda_2 = 0$. We also establish that $H_1(t)$ and $H_0(t)$ converge exponentially to Λ_1^{-1} and 0, respectively. Theorem 2 provides analytical expressions for $H_1(t)$ and $H_0(t)$ that satisfy (14) in terms of $H_0(t)$ and $H_2(t)$, respectively, and it proves the aforementioned convergence rates.

Theorem 2: For any full rank initial condition $H_1(0) \succ 0$, the solution $H_1(t)$ to (14) is given by

$$H_{1}(t) = \Lambda_{1}^{-1} + e^{-\Lambda_{1}t} (H_{1}(0) - \Lambda_{1}^{-1}) e^{-\Lambda_{1}t} + 2 \int_{0}^{t} e^{\Lambda_{1}(\tau-t)} H_{0}(\tau) H_{0}^{T}(\tau) e^{\Lambda_{1}(\tau-t)} d\tau$$

and it satisfies $H_1(t) \succ 0$. Moreover, for any $t \ge 0$, we have

$$\begin{aligned} \|\dot{H}_{1}(t)\|_{2} &\leq \left(\|\dot{H}_{1}(0)\|_{2} + 2t\|H_{0}(0)\|_{2}^{2}\right) e^{-2\lambda_{r}\star t} \\ \|H_{0}(t)\|_{2} &\leq \|H_{0}(0)\|_{2} e^{-\lambda_{r}\star t} \\ \|H_{2}(t)\|_{2} &\leq 2\|H_{2}(0)\|_{2}/(1+t) \end{aligned}$$

where $\tilde{H}_1(t) := H_1(t) - \Lambda_1^{-1}$.

Pre

Theorem 2 demonstrates that starting from any initial condition H(0) with $H_1(0) \succ 0$, $H_1(t)$ and $H_0(t)$ converge exponentially to Λ_1^{-1} and 0 with the respective rates $2\lambda_{r^*}$ and λ_{r^*} . Note that the upper bound on $||H_1(t) - \Lambda_1^{-1}||_2$ involves algebraic growth for small t, i.e., $\log(t)/t \leq \lambda_r$, but the exponential decay eventually dominates. In Remark 4, we provide examples to demonstrate that this algebraic growth in the upper bound on $||H_1 - \Lambda_1^{-1}||_2$ cannot be eliminated.

Exact parameterization: We now specialize Theorem 2 to $r = r^*$. In this case, the matrix $P_1 = Z_1 Z_1^T$ is invertible if and only if $det(Z_1) \neq 0$. Thus, change of variables (13) satisfies

$$H_1 = P_1^{-1} = Z_1^{-T} Z_1^{-1}, \ H_0 = P_1^{-1} P_0 = Z_1^{-T} Z_2^{T}$$

and the Schur complement $H_2 = P/P_1$ vanishes,

 $H_2 = P_2 - P_0^T P_1^{-1} P_0 = Z_2 Z_2^T - Z_2 Z_1^T (Z_1^{-T} Z_1^{-1}) Z_1 Z_2^T = 0.$

Corollary 2: For $r = r^*$, the solution of (14) is given by

$$\dot{H}_1(t) = e^{-\Lambda_1 t} (\dot{H}_1(0) + 2H_0(0)\Psi(t)H_0^T(0))e^{-\Lambda_1 t} H_0(t) = e^{-\Lambda_1 t} H_0(0)e^{-\Lambda_2 t}, \quad H_2(t) = 0$$

where $\tilde{H}_1(t) := H_1(t) - \Lambda_1^{-1}$ and $\Psi(t) := \int_0^t e^{-2\Lambda_2 \tau} d\tau$.

Proof: Since $H_2 = P/P_1 = 0$ for $r = r^*$, the H_0 -dynamics in (14) simplify to $\dot{H}_0 = -\Lambda_1 H_0 - H_0 \Lambda_2$ which leads to the analytical solution for $H_0(t)$ in Corollary 2. Combining this result with Theorem 2 yields the analytical solution for $H_1(t)$ and completes the proof.

Remark 4: For $\Lambda_2 = 0$, the matrix $\Psi(t)$ in Proposition 2 becomes equal to tI, thereby showing that the algebraic growth in the upper bound established in Theorem 2 is inevitable.

Remark 5: While changes of coordinates (3) and (13) are obtained by decomposing the optimization variable Z into $Z_1 \in \mathbb{R}^{i \times r}$ and $Z_2 \in \mathbb{R}^{(n-i) \times r}$ with $i = r^*$, and $\Lambda = \text{diag}(\Lambda_1, -\Lambda_2)$ into its positive and non-positive diagonal blocks, it is easy to verify that a similar change of variables can be introduced for any $i \in \{1, \ldots, n-1\}$. In addition, the H_2 -dynamics in (14c) have a similar structure to the original system (5) that governs the P-dynamics. Now, let $\Lambda_1 = \text{diag}(\hat{\lambda}_1 I, \ldots, \hat{\lambda}_k I)$ be the partitioning of Λ_1 based on its k distinct eigenvalues $\hat{\lambda}_1 > \cdots > \hat{\lambda}_k > 0$. Utilizing the above facts, we can start from the principle eigenspace of Λ associated with $\hat{\lambda}_1$ and successively employ k changes of variables similar to (13) to decouple individual eigenspaces and bring system (5) into a cascade connection of k + 2 subsystems,

$$\hat{H}_{i,1} = -2\hat{\lambda}_i \hat{H}_{i,1} + 2(I + \hat{H}_{i,0} \hat{H}_{i,0}^T)$$
(15a)

$$\hat{H}_{i,0} = -\hat{\lambda}_i \hat{H}_{i,0} + \hat{H}_{i,0} \hat{\Lambda}_i - 2\hat{H}_{i,0} \hat{H}_{i+1,1}$$
(15b)

$$\hat{H}_2 = -\Lambda_2 \hat{H}_2 - \hat{H}_2 \Lambda_2 - 2\hat{H}_2^2.$$
(15c)

for i = 1, ..., k. Here, the pair $(\hat{H}_{i,1}, \hat{H}_{i,0})$ corresponds to the distinct eigenvalue $\hat{\lambda}_i > 0$, $\hat{H}_{k+1,i} := \hat{H}_2$ corresponds to non-positive eigenvalues of Λ , and $\hat{\Lambda}_i := \text{diag}(\hat{\lambda}_i I, ..., \hat{\lambda}_k I, -\Lambda_2)$ is a lower diagonal block of Λ ; see Fig. 2 for an illustration.

Furthermore, as we demonstrate in Appendix D, the autonomous system in (15c) satisfies $\hat{H}_2(t) = H_2(t)$, where H_2 is the original Schur complement in Theorem 2. Finally, using similar arguments as in the proof of Theorem 2, it is straightforward to show under the conditions of Theorem 2,

$$\begin{aligned} \|\hat{H}_{i,1}(t) - \hat{\lambda}_i^{-1}I\|_2 &\leq \hat{c}_{i,1} e^{-2\lambda_i t} \\ \|\hat{H}_{i,0}(t)\|_2 &\leq \hat{c}_{i,0} e^{-(\hat{\lambda}_i - \hat{\lambda}_{i+1})t} \\ \|\hat{H}_2(t)\|_2 &\leq \hat{c}_2/(1+t) \end{aligned}$$

where $\hat{c}_{i,1}$, $\hat{c}_{i,0}$, and \hat{c}_2 are positive scalars that depend on the initial condition, and $\hat{\lambda}_{k+1} := 0$. This decomposition demonstrates the impact of gaps between eigenvalues of Λ_1 on the convergence behavior of system (5).




Figure 2. Block diagram of the 2k + 1 subsystems in (15).

Remark 6: When $r = r^*$ and $\Lambda_2 = 0$, we can introduce a new variable $W = H_0 H_0^T$ to reduce (14) to a stable LTI system with stability margin λ_{r^*} ,

$$\dot{W} = -\Lambda_1 W - W \Lambda_1, \ \dot{H}_1 = -\Lambda_1 H_1 - H_1 \Lambda_1 + 2I + 2W.$$

V. COMPUTATIONAL EXPERIMENTS

Herein, we provide an example to demonstrate the merits of our theoretical results. We set n = 10, $r^* = 4$, r = 8, $\Lambda_1 = \text{diag}(4, 3, 2, 1)$, and $\Lambda_2 = 0$. The black, blue, green, red, and yellow curves mark trajectories of gradient flow dynamics (2) with random initial conditions around the equilibrium points $\bar{P}_1 = \text{diag}(0, 0, 0, 0)$, $\bar{P}_1 = \text{diag}(4, 0, 0, 0)$, $\bar{P}_1 = \text{diag}(4, 3, 0, 0)$, and $\bar{P}_1 = \text{diag}(4, 3, 2, 0)$, and a normal random initialization, respectively. Figure 3a illustrates the objective function in (1). Figures 3b and 3c demonstrate the exponential convergence of $P_1(t)$ to Λ_1 and of $P_0(t)$ to 0, respectively. While at early stages, we observe a transient behavior, for large enough t, the error converges exponentially. Finally, Figure 3d shows the sub-linear convergence resulting from over-parameterization. We observe that the convergence curves confirm the results established in Theorem 1.

VI. CONCLUDING REMARKS

We have examined the gradient flow dynamics for overparameterized symmetric low-rank matrix factorization problem under the most general conditions. Our proof is based on a novel nonlinear coordinate transformation that converts the original problem into a cascade connection of three subsystems. In spite of the lack of convexity, we proved that this system globally converges to the stable equilibrium point if and only if the initialization is well-aligned with the optimal solution. We




Figure 3. Results obtained using simulation of gradient flow dynamics (2) for a problem with n = 10, $r^* = 4$, r = 8, $\Lambda_1 = \text{diag}(4,3,2,1)$, and $\Lambda_2 = 0$. Colors mark trajectories with random initial conditions where $P_1(0)$ is selected: around the equilibrium points $\bar{P}_1 = \text{diag}(0,0,0,0)$ (black), $\bar{P}_1 = \text{diag}(4,0,0,0)$ (blue), $\bar{P}_1 = \text{diag}(4,3,0,0)$ (green), $\bar{P}_1 = \text{diag}(4,3,2,0)$ (red); with Gaussian distribution (yellow).

used signal/noise decomposition to show that the subsystem associated with the signal component exponentially converges to the target low-rank matrix. Our analysis also reveals that the Schur complement associated with excess parameters vanishes with O(1/t) rate, thereby demonstrating that overparameterization inevitably decelerates the algorithm.

Potential future directions include the extension of our results to (i) the asymmetric setup aimed at computing a low-rank factorization of a rectangular matrix $M \in \mathbb{R}^{m \times n}$; (ii) the matrix sensing problem where only some linear measurements of the target matrix are observed; and (iii) the optimization of structured problems using gradient flow dynamics [31], [32].

REFERENCES

- E. J. Candes, X. Li, and M. Soltanolkotabi, "Phase retrieval via Wirtinger flow: Theory and algorithms," *IEEE Trans. Inform. Theory*, vol. 61, no. 4, pp. 1985–2007, 2015.
- [2] Y. Chen and E. J. Candes, "Solving random quadratic systems of equations is nearly as easy as solving linear systems," *Comm. Pure Appl. Math.*, vol. 70, no. 5, pp. 822–883, 2017.
- [3] C. Ma, K. Wang, Y. Chi, and Y. Chen, "Implicit regularization in nonconvex statistical estimation: Gradient descent converges linearly for phase retrieval and matrix completion," in *Proc. Int. Conf. Mach. Learn.*, pp. 3345–3354, 2018.
- [4] S. Tu, R. Boczar, M. Simchowitz, M. Soltanolkotabi, and B. Recht, "Low-rank solutions of linear matrix equations via Procrustes Flow," in *Proc. Int. Conf. Mach. Learn.*, pp. 964–973, 2016.
- [5] X. Li, S. Ling, T. Strohmer, and K. Wei, "Rapid, robust, and reliable blind deconvolution via nonconvex optimization," *Appl. Comput. Harmon. Anal.*, vol. 47, no. 3, pp. 893–934, 2019.
- [6] S. Ling and T. Strohmer, "Regularized gradient descent: A non-convex recipe for fast joint blind deconvolution and demixing," *Inf. Inference*, vol. 8, no. 1, pp. 1–49, 2019.
- [7] P. Netrapalli, P. Jain, and S. Sanghavi, "Phase retrieval using alternating minimization," in *Proc. Adv. Neural Inf. Process. Syst.*, vol. 26, 2013.
- [8] I. Waldspurger, "Phase retrieval with random Gaussian sensing vectors by alternating projections," *IEEE Trans. Inform. Theory*, vol. 64, no. 5, pp. 3301–3312, 2018.
- [9] D. Stöger and Y. Zhu, "Non-convex matrix sensing: Breaking the quadratic rank barrier in the sample complexity," arXiv preprint arXiv:2408.13276, 2024.
- [10] J. Sun, Q. Qu, and J. Wright, "When are nonconvex problems not scary?," arXiv preprint arXiv:1510.06096, 2015.
- [11] Y. Nesterov and B. T. Polyak, "Cubic regularization of Newton method and its global performance," *Math. Program.*, vol. 108, no. 1, pp. 177–205, 2006.
- [12] J. Nocedal and S. J. Wright, "Trust-region methods," *Numerical optimization*, pp. 66–100, 2006.
- [13] C. Jin, R. Ge, P. Netrapalli, S. M. Kakade, and M. I. Jordan, "How to escape saddle points efficiently," in *Proc. Int. Conf. Mach. Learn.*, pp. 1724–1732, 2017.
- [14] R. Ge, F. Huang, C. Jin, and Y. Yuan, "Escaping from saddle points—online stochastic gradient for tensor decomposition," in *Proc. Conf. Learn. Theory*, pp. 797–842, 2015.

- [15] M. Raginsky, A. Rakhlin, and M. Telgarsky, "Non-convex learning via stochastic gradient Langevin dynamics: A nonasymptotic analysis," in *Proc. Conf. Learn. Theory*, pp. 1674–1703, 2017.
- [16] Y. Zhang, P. Liang, and M. Charikar, "A hitting time analysis of stochastic gradient Langevin dynamics," in *Proc. Conf. Learn. Theory*, pp. 1980– 2022, 2017.
- [17] S. Arora, N. Cohen, W. Hu, and Y. Luo, "Implicit regularization in deep matrix factorization," in *Advances in Neural Information Processing Systems*, vol. 32, 2019.
- [18] Y. Chen, Y. Chi, J. Fan, and C. Ma, "Gradient descent with random initialization: Fast global convergence for nonconvex phase retrieval," *Math. Program.*, vol. 176, pp. 5–37, 2019.
- [19] D. Stöger and M. Soltanolkotabi, "Small random initialization is akin to spectral learning: Optimization and generalization guarantees for overparameterized low-rank matrix reconstruction," in *Proc. Adv. Neural Inf. Process. Syst.*, vol. 34, pp. 23831–23843, 2021.
- [20] M. Soltanolkotabi, D. Stöger, and C. Xie, "Implicit balancing and regularization: Generalization and convergence guarantees for overparameterized asymmetric matrix sensing," in *Proc. Conf. Learn. Theory*, pp. 5140–5142, 2023.
- [21] T. Ye and S. S. Du, "Global convergence of gradient descent for asymmetric low-rank matrix factorization," in *Proc. Adv. Neural Inf. Process. Syst.*, vol. 34, pp. 1429–1439, 2021.
- [22] H. Mohammadi, M. Razaviyayn, and M. R. Jovanović, "On the stability of gradient flow dynamics for a rank-one matrix approximation problem," in *Proc. Amer. Control Conf.*, pp. 4533–4538, 2018.
- [23] R. Ge, J. D. Lee, and T. Ma, "Matrix completion has no spurious local minimum," 2016.
- [24] K. Kawaguchi, "Deep learning without poor local minima," in Proc. Adv. Neural Inf. Process. Syst., vol. 29, 2016.
- [25] S. Arora, S. Du, W. Hu, Z. Li, and R. Wang, "Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks," in *Proc. Int. Conf. Mach. Learn.*, pp. 322–332, 2019.
- [26] E. Oja, "Simplified neuron model as a principal component analyzer," J. Math. Biol., vol. 15, pp. 267–273, 1982.
- [27] E. Oja, "Neural networks, principal components, and subspaces," Int. J. Neural Syst., vol. 1, no. 01, pp. 61–68, 1989.
- [28] U. Helmke and J. B. Moore, *Optimization and Dynamical Systems*. Springer Science & Business Media, 2012.
- [29] N. Srebro and T. Jaakkola, "Weighted low-rank approximations," in Proc. Int. Conf. Mach. Learn., pp. 720–727, 2003.
- [30] S. Tarmoun, G. Franca, B. D. Haeffele, and R. Vidal, "Understanding the dynamics of gradient flow in overparameterized linear models," in *Proc. Int. Conf. Mach. Learn.*, pp. 10153–10161, 2021.
- [31] F. Scott, R. Conejeros, and V. S. Vassiliadis, "Constrained nlp via gradient flow penalty continuation: Towards self-tuning robust penalty schemes," *Comput. Chem. Eng.*, vol. 101, pp. 243–258, 2017.
- [32] E. A. del Rio-Chanona, C. Bakker, F. Fiorelli, M. Paraskevopoulos, F. Scott, R. Conejeros, and V. S. Vassiliadis, "On the solution of differential-algebraic equations through gradient flow embedding," *Comput. Chem. Eng.*, vol. 103, pp. 165–175, 2017.
- [33] C. A. Desoer and M. Vidyasagar, Feedback Systems: Input-Output Properties. Philadelphia, PA, USA: SIAM, 2009.
- [34] G. B. Folland, Real Analysis: Modern Techniques and Their Applications, vol. 40. John Wiley & Sons, 1999.
- [35] D. E. Crabtree and E. V. Haynsworth, "An identity for the Schur complement of a matrix," *Proc. Am. Math. Soc.*, vol. 22, no. 2, pp. 364– 366, 1969.

APPENDIX

A. Characterization of equilibrium points

It is easy to verify that any $\bar{P} \in \mathcal{H}$ is an equilibrium point of system (5). To show the converse, let the pair $(\bar{P} \succeq 0, \Lambda)$ make the right-hand side of (11) vanish,

$$0 = \bar{P}_1 \Lambda_1 + \Lambda_1 \bar{P}_1 - 2(\bar{P}_1^2 + \bar{P}_0 \bar{P}_0^T)$$
(16a)

$$0 = \Lambda_1 \bar{P}_0 - \bar{P}_0 \Lambda_2 - 2(\bar{P}_1 \bar{P}_0 + \bar{P}_0 \bar{P}_2) \tag{16b}$$

$$0 = -\bar{P}_2\Lambda_2 - \Lambda_2\bar{P}_2 - 2(\bar{P}_0^T\bar{P}_0 + \bar{P}_2^2).$$
(16c)

Since $P \succeq 0$, we have $\bar{P}_2 \succeq 0$. To show that $\bar{P}_2 = 0$, let us assume that \bar{P}_2 has a positive eigenvalue $\mu > 0$ with the associated eigenvector p. Pre and post multiplying (16c) by p^T and p, respectively, and rearranging terms yields,

$$\mu p^T \Lambda_2 p = -\|\bar{P}_0 p\|_2^2 - \mu^2 \|p\|_2^2 < 0.$$
(17)

This contradicts positive semi-definiteness of Λ_2 and implies that $\bar{P}_2 = 0$. Furthermore, substitution of $\bar{P}_2 = 0$ to (16c) gives $\bar{P}_0 = 0$ and equation (16a) simplifies to,

$$\bar{P}_1\Lambda_1 + \Lambda_1\bar{P}_1 = 2\bar{P}_1^2.$$
(18)

Now, consider the eigenvalue decomposition of \bar{P}_1 , $\bar{P}_1 = \bar{V}\bar{\Sigma}\bar{V}^T = \bar{V}_i\bar{\Sigma}_i\bar{V}_i^T \neq 0$, where $\bar{\Sigma} = \text{diag}(\bar{\Sigma}_i, 0) \in \mathbb{R}^{r^* \times r^*}$ with $\bar{\Sigma}_i \in \mathbb{R}^{l \times l}$ containing the nonzero eigenvalues and, $\bar{V} := [\bar{V}_i \ \bar{V}_o]$ with $\bar{V}_i \in \mathbb{R}^{r^* \times l}$ containing the corresponding orthonormal eigenvectors as its columns. Equation (18) yields

$$A\Sigma + \Sigma A = 2\Sigma^2 \tag{19a}$$

where $A := \bar{V}^T \Lambda_1 \bar{V}$.

If $l = r^*$, i.e., if $\overline{\Sigma}$ is full rank, (19a) implies that A is also diagonal and, hence, $A = \overline{\Sigma}$. From the definition of A and \overline{V} , it follows that $\overline{P}_1 = \Lambda_1$ is an equilibrium point.

To address the case with $l \neq r^{\star}$, let us partition A

$$A = \begin{bmatrix} A_{\mathbf{i}} & A_{\mathbf{a}} \\ A_{\mathbf{a}}^T & A_{\mathbf{o}} \end{bmatrix}$$
(19b)

with $A_i \in \mathbb{R}^{l \times l}$. Using (19a), we observe that

$$\begin{bmatrix} A_{\mathbf{i}}\bar{\Sigma}_{i}+\bar{\Sigma}_{i}A_{\mathbf{i}} & \bar{\Sigma}_{i}A_{\mathbf{a}} \\ A_{\mathbf{a}}^{T}\bar{\Sigma}_{i} & 0 \end{bmatrix} = \begin{bmatrix} 2\bar{\Sigma}_{i}^{2} & 0 \\ 0 & 0 \end{bmatrix}.$$
 (19c)

Thus, $A_i = \overline{\Sigma}_i$ and $A_a = 0$. Substituting this to $A := \overline{V}^T \Lambda_1 \overline{V}$ yields

$$\Lambda_1 = \bar{V} \begin{bmatrix} \bar{\Sigma}_i & 0\\ 0 & A_o \end{bmatrix} \bar{V}^T = \bar{P}_1 + \bar{V}_o A_o \bar{V}_o^T \qquad (19d)$$

Hence, $\Lambda_1 - \bar{P}_1 = \bar{V}_{\mathbf{o}} A_{\mathbf{o}} \bar{V}_{\mathbf{o}}^T$. Since $\bar{V}_{\mathbf{i}}$ and $\bar{V}_{\mathbf{o}}$ are mutually orthogonal, i.e., $\bar{V}_{\mathbf{i}}^T \bar{V}_{\mathbf{o}} = 0$, we have $\bar{P}_1(\Lambda_1 - \bar{P}_1) = 0$ and \mathcal{H} is the set of equilibrium points of (5).

1) Proof of Lemma 1: First, we show that the existence of such a matrix V and diagonal matrices D_i and D_o is sufficient for $\overline{P} \in \mathcal{H}$. We observe that

$$P_{1}\Lambda_{1} = P_{1}(P_{1} + \Lambda_{1} - P_{1}) = \bar{P}_{1}^{2} + P_{1}(\Lambda_{1} - P_{1})$$
$$= \bar{P}_{1}^{2} + \bar{P}_{1}V_{o}D_{o}V_{o}^{T} \stackrel{(a)}{=} \bar{P}_{1}^{2}$$

where (a) follow from the fact that V_i and V_o are orthogonal. To show the necessity, first, let $A_o = U_o D_o U_o^T$ be the eigenvalue

decomposition of A_{o} . We can write (19d) as

$$\bar{V}_{\mathbf{i}}\bar{\Sigma}_{\mathbf{i}}\bar{V}_{\mathbf{i}}^{T} + \bar{V}_{\mathbf{o}}A_{\mathbf{o}}\bar{V}_{\mathbf{o}}^{T} = \bar{V}_{\mathbf{i}}\bar{\Sigma}_{\mathbf{i}}\bar{V}_{\mathbf{i}}^{T} + \bar{V}_{\mathbf{o}}U_{\mathbf{o}}D_{\mathbf{o}}U_{\mathbf{o}}^{T}\bar{V}_{\mathbf{o}}^{T}.$$

Let us define

$$V_{\mathbf{i}} := V_{\mathbf{i}}, D_{\mathbf{i}} := \Sigma_{\mathbf{i}}, V_{\mathbf{o}} := V_{\mathbf{o}}U_{\mathbf{o}}$$

We observe that V_{o} is a unitary matrix and orthogonal to V_{i} . We will show that such a $(V_{i}, D_{i}, V_{o}, D_{o})$ satisfies the conditions in Lemma 1. Based on the definition, we have

$$\bar{P}_1 = V_{\mathbf{i}} D_{\mathbf{i}} V_{\mathbf{i}}^T, \quad \Lambda_1 - \bar{P}_1 = V_{\mathbf{o}} D_{\mathbf{o}} V_{\mathbf{o}}^T.$$

We also know that $\operatorname{eig}(D_{\mathbf{i}}) \cup \operatorname{eig}(A_{\mathbf{o}}) = \operatorname{eig}(\Lambda_1)$, and $\operatorname{eig}(D_{\mathbf{o}}) = \operatorname{eig}(A_{\mathbf{o}})$, which proves the necessity. Finally, suppose that the eigenvalues of Λ_1 are all distinct. We know that

$$\Lambda_1 = \begin{bmatrix} V_{\mathbf{i}} & V_{\mathbf{o}} \end{bmatrix} \begin{bmatrix} D_{\mathbf{i}} & 0 \\ 0 & D_{\mathbf{o}} \end{bmatrix} \begin{bmatrix} V_{\mathbf{i}}^T \\ V_{\mathbf{o}}^T \end{bmatrix}.$$

In this case, both Λ_1 and diag (D_i, D_o) are diagonal matrices with distinct values. Hence, the matrix $\begin{bmatrix} V_i & V_o \end{bmatrix}$ has to be a permutation matrix as the eigenvalues are unique. This implies that diag (D_i, D_o) can only be a permuted version of Λ_1 . Thus, there is a one-to-one correspondence between the equilibrium points and the 2^{r^*} subsets of positive eigenvalues. This completes the proof.

B. Local convergence results

1) Proof of Proposition 2: We consider the objective function in optimization problem (1) as a Lyapunov function candidate,

$$V_F(P) = (1/4) ||P - \Lambda||_F^2.$$

The derivative of V_F along the trajectories of (5) satisfies

$$\dot{V}_F = (1/4) \operatorname{trace} \left[(P - \Lambda) \dot{P} \right] + (1/4) \operatorname{trace} \left[\dot{P} (P - \Lambda) \right]$$

$$\stackrel{(a)}{=} (1/2) \operatorname{trace} \left[(P - \Lambda) ((\Lambda - P)P + P(\Lambda - P)) \right]$$

$$\stackrel{(b)}{=} -\operatorname{trace} \left[(P - \Lambda)^2 P \right] = - \| (P - \Lambda) P^{\frac{1}{2}} \|_F^2$$

where (a) and (b) follow by the cyclic property of the matrix trace. Stability of $\overline{P} = \operatorname{diag}(\Lambda_1, 0)$ in the sense of Lyapunov follows from $\dot{V}_F \leq 0$ for all $P \succeq 0$. To show local asymptotic stability, we note that $\dot{V}_F(P) = 0$ if and only if $(P-\Lambda)P^{\frac{1}{2}} = 0$. Thus, $(P - \Lambda)P = 0$ and $\dot{V}_F(P) = 0$ only at equilibria of system (5). Since $\operatorname{diag}(\Lambda_1, 0)$ is an isolated equilibrium point, $\dot{V}_F(P)$ is negative over the open neighborhood of $\operatorname{diag}(\Lambda_1, 0)$, $\{P \mid ||P - \operatorname{diag}(\Lambda_1, 0)||_2 < \lambda_{r^*}\}$, and $\dot{V}_F(P) = 0$ if and only if $P = \operatorname{diag}(\Lambda_1, 0)$. Hence, $\overline{P} = \operatorname{diag}(\Lambda_1, 0)$ is a locally asymptotically stable equilibrium point of system (5).

2) Proof of Lemma 3: Let $(\lambda(t), w(t)) \in (\mathbb{R}_+, \mathbb{R}^{n-r^*})$ be the principal eigenpair of the matrix $P_2(t)$ with $w^T(t)w(t) = 1$. Since

$$\dot{w}^{T}(t)P_{2}(t)w(t) + w^{T}(t)P_{2}(t)\dot{w}(t) = \lambda(t) \frac{\mathrm{d}(w^{T}(t)w(t))}{\mathrm{d}t} = 0$$

the derivative of $V_N(t) := \lambda(t) = w^T(t)P_2(t)w(t)$ along the the spectral norm $||H_0||_2 = u^T H_0 v$ satisfies solutions of (5) satisfies

$$\begin{split} \dot{V}_N &= w^T \dot{P}_2 w + \dot{w}^T P_2 w + w^T P_2 \dot{w} = w^T \dot{P}_2 w \\ &= -w^T (P_2 \Lambda_2 + \Lambda_2 P_2 + 2(P_0^T P_0 + P_2^2)) w \\ &= -2V_N^2 - 2 \|\Lambda_2^{\frac{1}{2}} w\|_2^2 V_N - 2\|P_0 w\|_2^2 \leq -2V_N^2. \end{split}$$

For $x(t_0) > 0$, $x(t) = \frac{2x(t_0)}{1+t-t_0}$ solves $\dot{x} = -2x^2$ and the result follows from comparison principle.

C. Global convergence results

1) Proof of Proposition 3: We can write

$$\dot{H}_1 = d(P_1^{-1})/dt = -H_1\dot{P}_1H_1 = -H_1 \left(P_1\Lambda_1 + \Lambda_1P_1 - 2(P_1^2 + P_0P_0^T)\right)H_1 = -H_1 \left(H_1^{-1}\Lambda_1 + \Lambda_1H_1^{-1} - 2(H_1^{-2} + P_0P_0^T)\right)H_1 = -\Lambda_1H_1 - H_1\Lambda_1 + 2I + 2H_1P_0P_0^TH_1 = -\Lambda_1H_1 - H_1\Lambda_1 + 2I + 2H_0H_0^T$$

where the last equality follows from $H_1P_0 = H_0$. Similarly, for the H_0 -dynamics we have

$$H_{0} = d(H_{1}P_{0})/dt$$

$$= H_{1}\dot{P}_{0} + \dot{H}_{1}P_{0}$$

$$= H_{1} (\Lambda_{1}P_{0} - P_{0}\Lambda_{2} - 2(P_{1}P_{0} + P_{0}P_{2}))$$

$$+ (-\Lambda_{1}H_{1} - H_{1}\Lambda_{1} + 2I + 2H_{0}H_{0}^{T})P_{0}$$

$$= -\Lambda_{1}H_{1}P_{0} - H_{1}P_{0}\Lambda_{2} - 2(H_{1}P_{0}P_{2} - H_{0}H_{0}^{T}P_{0})$$

$$= -\Lambda_{1}H_{0} - H_{0}\Lambda_{2} - 2H_{0}(P_{2} - P_{0}^{T}P_{1}^{-1}P_{0})$$

$$= -\Lambda_{1}H_{0} - H_{0}\Lambda_{2} - 2H_{0}H_{2}.$$

Finally, the H_2 -dynamics are given by

$$\begin{aligned} H_2 &= \mathrm{d}(P_2 - P_0^T H_1 P_0)/\mathrm{d}t \\ &= \dot{P}_2 - \dot{P}_0^T H_1 P_0 - P_0^T H_1 \dot{P}_0 - P_0^T \dot{H}_1 P_0 \\ &= -P_2 \Lambda_2 - \Lambda_2 P_2 - 2(P_0^T P_0 + P_2^2) \\ &- (\Lambda_1 P_0 - P_0 \Lambda_2 - 2(P_1 P_0 + P_0 P_2))^T H_1 P_0 \\ &- P_0^T H_1 (\Lambda_1 P_0 - P_0 \Lambda_2 - 2(P_1 P_0 + P_0 P_2)) \\ &- P_0^T (-\Lambda_1 H_1 - H_1 \Lambda_1 + 2I + 2H_0 H_0^T) P_0 \\ &= -\Lambda_2 (P_2 - P_0^T H_1 P_0) - (P_2 - P_0^T H_1 P_0) \Lambda_2 \\ &- 2(P_2 - P_0^T H_1 P_0)^2 \\ &= -\Lambda_2 H_2 - H_2 \Lambda_2 - 2H_2^2 \end{aligned}$$

which completes the proof.

2) Proof of Theorem 2: We first present a technical result. The next lemma establishes the exponential decay of H_0 .

Lemma 4: For the matrix H_0 governed by (14), the derivative of the spectral norm satisfies

$$\frac{\mathrm{d} \|H_0\|_2}{\mathrm{d} t} \leq -\lambda_{r^{\star}} \|H_0\|_2$$

Proof: Let u(t) and v(t) be the principal left and right singular vectors of $H_0(t)$ with ||u(t)|| = ||v(t)|| = 1. Then,

$$\frac{\|H_0\|_2}{dt} = u^T \dot{H}_0 v = u^T (-\Lambda_1 H_0 - H_0 (2P/P_1 + \Lambda_2)) v = -(u^T \Lambda_1 u + v^T (2P/P_1 + \Lambda_2) v) \|H_0\|_2 \leq -\lambda_{r^\star} \|H_0\|_2$$

Here, the first equality is a well-known property of the derivative of singular vectors, the inequality follows from the fact that $u^T \Lambda_1 u > \lambda_{r^*}$ and $2P/P_1 + \Lambda_2 \succeq 0$. This proves the first inequality in Lemma 4.

We are now ready to prove Theorem 2.

The shifted matrix variable $\tilde{H}_1 = H_1 - \Lambda_1^{-1}$ brings equation (14) for H_1 to

$$\dot{\tilde{H}}_1 = -\Lambda_1 \tilde{H}_1 - \tilde{H}_1 \Lambda_1 + 2H_0 H_0^T.$$

This system is linear in \tilde{H}_1 , and it is driven by the exogenous input $2H_0H_0^T$. The state transition operator is determined by $\dot{M} \rightarrow e^{-\Lambda_1 t} \dot{M} e^{-\Lambda_1 t}$, the variation of constants formula yields

$$\tilde{H}_1(t) = e^{-\Lambda_1 t} \tilde{H}_1(0) e^{-\Lambda_1 t} + 2\Phi(t)$$

where the forced response is determined by

$$\Phi(t) := \int_0^t e^{-\Lambda_1(t-\tau)} H_0(\tau) H_0^T(\tau) e^{-\Lambda_1(t-\tau)} d\tau.$$
 (20)

Substituting $\tilde{H}_1 = H_1 - \Lambda_1^{-1}$ in the above equation yields the expression in Theorem 2 for $H_1(t)$. The norm of this forced response can be bounded by

$$\begin{split} \|\Phi(t)\|_{2} &\stackrel{(a)}{\leq} \int_{0}^{t} \|\mathrm{e}^{-\Lambda_{1}(t-\tau)}H_{0}(\tau)H_{0}(\tau)^{T}\mathrm{e}^{-\Lambda_{1}(t-\tau)}\|_{2}\,\mathrm{d}\tau\\ &\stackrel{(b)}{\leq} \int_{0}^{t} \|\mathrm{e}^{-\Lambda_{1}(t-\tau)}\|_{2}^{2}\,\|H_{0}(\tau)\|_{2}^{2}\,\mathrm{d}\tau\\ &= \int_{0}^{t} \mathrm{e}^{-2\lambda_{r^{\star}}(t-\tau)}\,\|H_{0}(\tau)\|_{2}^{2}\,\mathrm{d}\tau \end{split}$$
(21)

where (a) follows from the triangle inequality and (b) follows from the sub-multiplicative property of the spectral norm.

To complete the proof, we note that if $H_1(t) \succ 0$ then $P_1(t) = H_1^{-1}$ and $P_0 = H_1^{-1}H_0$. In addition, the Schur complement P/P_1 exists which together with the exponential decay rate of $H_0(t)$ established in Lemma 4 yields

$$||H_0(t)||_2 \le e^{-\lambda_r \star t} ||H_0(0)||_2.$$

By combining this inequality with (21), we obtain

$$\|\Phi(t)\|_{2} \leq \int_{0}^{t} e^{-2\lambda_{r} \star t} d\tau \|H_{0}(0)\|_{2}^{2} \leq t e^{-2\lambda_{r} \star t} \|H_{0}(0)\|_{2}^{2}$$

Finally, to derive the upper bound on $\|\tilde{H}_1(t)\|_2$, we write

$$\begin{split} \|\tilde{H}_{1}(t)\|_{2} & \stackrel{(a)}{\leq} \|\mathrm{e}^{-\Lambda_{1}t}\tilde{H}_{1}(0)\mathrm{e}^{-\Lambda_{1}t}\|_{2} + 2\|\Phi(t)\|_{2} \\ & \stackrel{(b)}{\leq} \left(\|\tilde{H}_{1}(0)\|_{2} + 2t\|H_{0}(0)\|_{2}^{2}\right)\mathrm{e}^{-2\lambda_{r}\star t} \end{split}$$

where (a) follow from triangle inequality and (b) follows from combining the above aforementioned facts. This completes the proof of Theorem 2.

3) Proof of Theorem 1: We first present a lemma that we use to establish an upper bound on the error $||P_1 - \Lambda_1||_2$ as a function of $||P_1^{-1} - \Lambda_1^{-1}||_2$.

Lemma 5: Let the matrices $A \succ 0$ and B be such that

$$||A - B||_2 < \sigma := \sigma_{\min}(A).$$

Then, the matrix B is invertible, and it satisfies

$$||A^{-1} - B^{-1}||_2 \le \frac{||A - B||_2}{\sigma (\sigma - ||A - B||_2)}$$

Proof: For
$$Q := I - BA^{-1} = (A - B)A^{-1}$$
, we have
 $\|Q\|_2 \le \|A - B\|_2 \|A^{-1}\|_2 = \|A - B\|_2 / \sigma < 1.$ (22)

Thus, the matrix $BA^{-1} = I - Q$ and consequently B are both invertible by the small gain theorem [33]. In addition,

$$B^{-1} - A^{-1} = A^{-1}(AB^{-1} - I)$$

= $A^{-1}((I - Q)^{-1} - I) = A^{-1}\sum_{k=1}^{\infty} Q^k$

Thus, we can use the triangle inequality and the submultiplicative property of the spectral norm to obtain

$$\begin{split} \|B^{-1} - A^{-1}\|_{2} &\leq \sigma^{-1} \sum_{k=1}^{\infty} \|Q^{k}\|_{2} \\ &\stackrel{(a)}{\leq} \sigma^{-1} \sum_{k=1}^{\infty} \sigma^{-k} \|A - B\|_{2}^{k} = \frac{\|A - B\|_{2}}{\sigma \left(\sigma - \|A - B\|_{2}\right)} \end{split}$$

where (a) follows from (22). This completes the proof.

To prove the upper bound for $||P_1(t) - \Lambda_1||_2$, we note that the function l(t) can be written in the *H*-coordinates as

$$l(t) = \left(\|P_1^{-1}(0) - \Lambda_1^{-1}\|_2 + 2t \|P_1^{-1}(0)P_0(0)\|_2^2 \right) e^{-2\lambda_{r^*}t} = \left(\|\tilde{H}_1(0)\|_2 + 2t \|H_0(0)\|_2^2 \right) e^{-2\lambda_{r^*}t}.$$

Thus, we have

$$\begin{aligned} \|P_1^{-1}(t) - \Lambda_1^{-1}\|_2 &= \|\tilde{H}_1(t)\|_2 \le l(t) \\ &\le 1/(2\lambda_1) < 1/\lambda_1 = \sigma_{\min}(\Lambda_1^{-1}) \end{aligned} (23)$$

where the first inequality follows from Theorem 2 and the second inequality holds by assumption as stated in Theorem 1. The bound in (23) allows us to apply Lemma 5 with $A := \Lambda_1^{-1}$ and $B := P_1^{-1}(t)$ to obtain

$$\begin{aligned} \|P_{1}(t) - \Lambda_{1}\|_{2} &\leq \frac{\|P_{1}^{-1}(t) - \Lambda_{1}^{-1}\|_{2}}{\lambda_{1}^{-1}(\lambda_{1}^{-1} - \|P_{1}^{-1}(t) - \Lambda_{1}^{-1}\|_{2})} \\ &\leq 2\lambda_{1}^{2}\|P_{1}^{-1}(t) - \Lambda_{1}^{-1}\|_{2} \leq 2\lambda_{1}^{2}l(t) \end{aligned}$$
(24)

which establishes the desired upper bound for $||P_1(t) - \Lambda_1||_2$.

Furthermore, the upper bound on $||P_0(t)||_2$ is given by

$$\begin{aligned} \|P_{0}(t)\|_{2} & \stackrel{(a)}{\leq} \|P_{1}(t)\|_{2} \|P_{1}^{-1}(t)P_{0}(t)\|_{2} \\ & \stackrel{(b)}{\leq} (\|\Lambda_{1}\|_{2} + \|P_{1}(t) - \Lambda_{1}\|_{2}) \|P_{1}^{-1}(t)P_{0}(t)\|_{2} \\ & \stackrel{(c)}{\leq} (\lambda_{1} + 2\lambda_{1}^{2}l(t)) \|P_{1}^{-1}(0)P_{0}(0)\|_{2} e^{-\lambda_{r} \star t}. \end{aligned}$$

where (a) follows from the sub-multiplicative property of the spectral norm, (b) follows from triangle inequality, and (c)

follows from combining equation (24) and the fact that

$$|P_1^{-1}(t)P_0(t)||_2 = ||H_0(t)||_2 \le e^{-\lambda_r \star t} ||H_0(0)||_2$$

established in Theorem 2.

Finally, the inequality $P_1(t) \succ 0$ can be verified by noting that that $P_1^{-1}(t) = H_1(t) \succ 0$ exists and is bounded according to Theorem 2. This completes the proof.

4) Proof of Corollary 1: We begin by noting that the condition on Z(0) is equivalent to the condition $P_1(0) \succ 0$ in Theorem 1. Thus, the convergence results in Theorem 1 hold. For some $c_1 > 0$, Lemma 3 implies,

$$||P_2(t)||_2 = ||Z_2(t)||_2^2 \le c_1/t$$

and thus $||Z_2(t)||_2 \le \sqrt{c_1/t}$. Moreover, for $r = r^*$ we can use the exponential convergence of $P_2(t)$ established in Remark 1 to conclude that $||Z_2(t)||_2 \le c_2 e^{-\lambda_r \star t}$, for some $c_2 > 0$.

We next prove the convergence of $Z_1(t)$. Applying the triangle inequality and the submultiplicity of the spectral norm to equation (10a) yields

$$\begin{aligned} \|\dot{Z}_1\|_F &\leq \|\Lambda_1 - Z_1 Z_1^T\|_2 \|Z_1\|_F + \|Z_1 Z_2^T\|_2 \|Z_2\|_F \\ &= \|\Lambda_1 - P_1\|_2 \|Z_1\|_F + \|P_0\|_2 \|Z_2\|_F. \end{aligned} (25)$$

For any positive scalars $t_1 < t_2$, we have

$$\|\int_{t_1}^{t_2} \dot{Z}_1(s) \,\mathrm{d}s\|_F \leq \int_{t_1}^{t_2} \|\dot{Z}_1(s)\|_F \,\mathrm{d}s$$

$$\leq \int_{t_1}^{t_2} \left(\|\Lambda_1 - P_1(s)\|_2 \|Z_1(s)\|_F + \|P_0(s)\|_2 \|Z_2(s)\|_F\right) \,\mathrm{d}s \quad (26)$$

where the first inequality follows from the triangle inequality and the second follows from equation (25). By Theorem 1, we have $\|\Lambda_1 - P_1(t)\|_2 \leq c_3 e^{-2\lambda_r \star t}$ and $\|P_0(t)\|_2 \leq c_4 e^{-\lambda_r \star t}$ for scalars $c_3, c_4 > 0$. Moreover, using convergence of $P(t) = Z(t)Z^T(t)$ shown in Theorem 1, it is straightforward to verify that $\|Z(t)\|_F^2 = \|Z_1(t)\|_F^2 + \|Z_2(t)\|_F^2 \leq c^2$ for a large enough c > 0. Thus, for $t \gg 1$ we have

$$\|\int_{t}^{\infty} \dot{Z}_{1}(s) \,\mathrm{d}s\|_{F} \leq c \int_{t}^{\infty} \left(\|\Lambda_{1} - P_{1}(s)\|_{2} \right)$$
(27)

$$+ \|P_0(s)\|_2 ds$$
 (28)

$$\leq c' \int_{t}^{\infty} \left(e^{-\lambda_r \star s} + e^{-2\lambda_r \star s} \right) ds \quad (29)$$

$$\leq c'' \mathrm{e}^{-\lambda_r \star t} \tag{30}$$

where c', c'' > 0 are large enough scalars. Since $Z_1(t)$ is a bounded function with finite Frobenius norm of integral of \dot{Z} , it converges to a matrix Z_1^* [34]. In addition, since $Z(0)Z^T(0)$ satisfies the condition of Theorem 1, $P_1(t) = Z_1(t)Z_1^T(t)$ converges to Λ_1 and thereby $Z_1^*Z_1^{*T} = \Lambda_1$. To prove the convergence rate of $Z_1(t)$, note that equation (27) yields

$$||Z_1(t) - Z_1^{\star}||_F = ||\int_t^{\infty} \dot{Z}_1(s) \mathrm{d}s||_F \leq c_5 \mathrm{e}^{-\lambda_r \star t}.$$

for some $c_5 > 0$. This completes the proof.

D. Expanded decomposition

Let us formalize the strategy sketched in Remark 5 by defining the recursive equations

$$U_{i+1} = U_i / \hat{P}_{i,1} = \hat{P}_{i,2} - \hat{P}_{i,0}^T \hat{P}_{i,1}^{-1} \hat{P}_{i,0} \in \mathbb{R}^{m_i \times m_i}$$
(31)

for i = 1..., k with the initialization $U_1 = P \in \mathbb{R}^{n \times n}$. Here, the matrices $\hat{P}_{i,j}$ are obtained by the block decomposition

$$U_{i} = \begin{bmatrix} \hat{P}_{i,1} & \hat{P}_{i,0} \\ \hat{P}_{i,0}^{T} & \hat{P}_{i,2} \end{bmatrix} \in \mathbb{R}^{m_{i-1} \times m_{i-1}}$$
(32)

and they satisfy $\hat{P}_{i,1} \in \mathbb{R}^{n_i \times n_i}$, $\hat{P}_{i,0} \in \mathbb{R}^{n_i \times m_i}$, and $\hat{P}_{i,2} \in \mathbb{R}^{m_i \times m_i}$, where n_i is the dimension of the *i*th principal subspace of Λ_1 associated with the eigenvalue $\hat{\lambda}_i$ and $m_i = n - (n_1 + \cdots + n_i)$ with $m_0 := n$. The change of variables

$$\hat{H}_{i,1} := \hat{P}_{i,1}^{-1} \in \mathbb{R}^{n_i \times n_i}
\hat{H}_{i,0} := \hat{P}_{i,1}^{-1} \hat{P}_{i,0} \in \mathbb{R}^{n_i \times m_i}
\hat{H}_{i,2} := U_{i+1} \in \mathbb{R}^{m_i \times m_i}$$
(33)

yields the cascade system (15).

For the set of equations in (31) to be well defined, the matrices $\hat{P}_{i,1}$ need to be invertible. A necessary and sufficient condition for the invertibility of $\hat{P}_{i,1}$ is that P_1 is invertible.

Proposition 4: System (31) satisfies

$$\det(P_1) = \prod_{i=1}^{k} \det(\hat{P}_{i,1})$$
(34)

Furthermore, we have

$$H_2 = U_{k+1} = \hat{H}_{k,2} \tag{35}$$

where $H_2 = P/P_1$ is the Schur complement of P_1 in P.

Proposition 4 establishes the equivalence between the invertibility of matrices $\hat{P}_{i,1}$ and P_1 . To prove this result, we next present two key properties of the Schur complement.

Lemma 6: For any symmetric block matrix

$$X = \left[\begin{array}{cc} X_1 & X_0 \\ X_0^T & X \end{array} \right]$$

with invertible X, we have the LU factorization

$$X = \begin{bmatrix} I & 0 \\ X_0^T X_1^{-1} & I \end{bmatrix} \begin{bmatrix} X_1 & 0 \\ 0 & X/X_1 \end{bmatrix} \begin{bmatrix} I & X_1^{-1} X_0 \\ 0 & I \end{bmatrix}$$

Lemma 7: Consider the symmetric block matrices

$$X = \begin{bmatrix} X_1 & X_0 \\ X_0^T & X_2 \end{bmatrix}, \quad X_1 := \begin{bmatrix} X_1 & \bar{X}_0 \\ \bar{X}_0^T & \bar{X}_2 \end{bmatrix}$$

For invertible X_1 , we have the identity

$$X/X_1 = Y/Y_1$$

where $Y := \begin{bmatrix} Y_1 & Y_0 \\ Y_0^T & Y_2 \end{bmatrix} = X/\bar{X}_1$ is the Schur complement of \bar{X}_1 in X and its block dimensions are conformable with X.

Proof: We first show that $Y_1 = X_1/\bar{X}_1$. We have

$$Y = \begin{bmatrix} Y_1 & Y_0 \\ Y_0^T & Y_2 \end{bmatrix} := X/\bar{X}_1$$
$$= \begin{bmatrix} \bar{X}_2 & S^T \\ S & X_2 \end{bmatrix} - \begin{bmatrix} \bar{X}_0^T \\ R^T \end{bmatrix} \bar{X}_1^{-1} \begin{bmatrix} \bar{X}_0 & R \end{bmatrix}$$
$$= \begin{bmatrix} \bar{X}_2 - \bar{X}_0^T \bar{X}_1^{-1} \bar{X}_0 & S^T - \bar{X}_0^T \bar{X}_1^{-1} R \\ S - R^T \bar{X}_1^{-1} \bar{X}_0 & X_2 - R^T \bar{X}_1^{-1} R \end{bmatrix}$$

where $X_0^T = \begin{bmatrix} R^T & S^T \end{bmatrix}$. Matching the 11-block in the above equation yields $Y_1 = X_1/\bar{X}_1$.

We can now write

$$X/X_1 = (X/X_1)/(X_1/X_1) = Y/Y_1$$

where the first equality is the quotient identity [35] and the second one follows from the definition of Y. This completes the proof.

1) Proof of Proposition 4: We use induction to show that

$$P/\hat{P}_i = U_{i+1} \tag{36}$$

where $\hat{P}_i \in \mathbb{R}^{(n-m_i)\times(n-m_i)}$ is the 11-block of P. Equation (36) for i = 1 follows from (31) and the fact that $\hat{P}_1 = \hat{P}_{1,1}$. To prove the case i + 1, we write

$$P/P_{i+1} = (P/P_i)/((P/P_i))_{11}$$

= $U_{i+1}/(U_{i+1})_{11}$
= $U_{i+1}/\hat{P}_{i+1,1} = U_{i+2}$

where we use $(\cdot)_{11}$ to denote the 11-block of the size n_{i+1} . Here, the first equality follows from Lemma 7 with X = P, $X_1 = \hat{P}_{i+1}$, and $\bar{X}_1 = \hat{P}_i$, the second equality is the inductive hypothesis, the third equality follows from (32), and the last equality follows from (31). This completes the proof of (36). This equation for i = k allows us to write

$$H_2 = P/P_1 = P/\hat{P}_k = U_{k+1} \stackrel{(a)}{=} \hat{H}_{k,2} =: \hat{H}_2.$$

where (a) follows from the definition in (33). This completes the proof of (35)

To prove (34), we can recursively apply Lemma 6 to the matrices U_i starting from $U_1 = P$ to form an LU decomposition $P = \Phi Y \Psi$, where Φ and Ψ are lower and upper diagonal matrices with 1 on the main diagonal, respectively, and

$$Y = \text{diag}(P_{1,1}, \cdots, P_{k,1}, H_{k,2})$$

Now, since $det(\Phi) = det(\Psi) = 1$, we obtain that

$$\det(P) = \det(Y)$$

= $\det(\hat{H}_{k,2}) \prod_{i=1}^{k} \det(\hat{P}_{i,1}) = \det(P/P_1) \prod_{i=1}^{k} \det(\hat{P}_{i,1}).$

where the last equality follows from (35). Combining this equation with $\det(P) = \det(P_1) \det(P/P_1)$ completes the proof of (34).