THE BOOLEAN SPECTRUM OF A GROTHENDIECK CATEGORY

HENNING KRAUSE

ABSTRACT. A notion of support for objects in any Grothendieck category is introduced. This is based on the spectral category of a Grothendieck category and uses its Boolean lattice of localising subcategories. The support provides a classification of all subcategories that are closed under arbitrary coproducts, subobjects, and essential extensions. There is also a notion of exact support which classifies certain thick subcategories. As an application, the coproduct decompositions of objects are described in terms of Boolean lattices. Also, for any ring Crawley-Boevey's correspondence between definable subcategories of modules and closed subsets of the Ziegler spectrum is extended.

1. Introduction

The spectrum $\operatorname{Sp} \mathcal{A}$ of a Grothendieck category \mathcal{A} is the set of isomorphism classes of indecomposable injective objects. This definition is due to Gabriel, and he uses this spectrum when he reconstructs a noetherian scheme from its category of quasi-coherent sheaves [12]. A finer invariant is introduced in work of Gabriel and Oberst [13]. The spectral category $\operatorname{Spec} \mathcal{A}$ is again a Grothendieck category with the additional property that every short exact sequence is split exact. There is a canonical functor $\mathcal{A} \to \operatorname{Spec} \mathcal{A}$ which identifies $\operatorname{Sp} \mathcal{A}$ with the isomorphism classes of simple objects in $\operatorname{Spec} \mathcal{A}$.

In this work we study the lattice $L(\operatorname{Spec} A)$ of localising subcategories of $\operatorname{Spec} A$, which one may think of as an intermediate between $\operatorname{Sp} A$ and $\operatorname{Spec} A$. This is actually a complete Boolean lattice; so it identifies via Stone duality with the Boolean lattice of clopen sets of its spectrum

$$\operatorname{Spc} A := \operatorname{Spec}(\mathbf{L}(\operatorname{Spec} A))$$

which we call the Boolean spectrum of A. There is a canonical embedding

$$\operatorname{Sp} A \hookrightarrow \operatorname{Spc} A$$

which is a bijection if and only if Spec \mathcal{A} is *discrete*, which means that every object decomposes into a coproduct of simple objects. This happens, for example, when \mathcal{A} is locally noetherian.

To each object $X \in \mathcal{A}$ we assign its support $\operatorname{supp}(X) \subseteq \operatorname{Spc} \mathcal{A}$, and this sets up a bijection

$$(1.1) \mathcal{C} \longmapsto \operatorname{supp}(\mathcal{C}) = \bigvee_{X \in \mathcal{C}} \operatorname{supp}(X)$$

between the essentially closed subcategories of \mathcal{A} and the clopen subsets of $\operatorname{Spc} \mathcal{A}$ (Theorem 3.3), where a full subcategory is called *essentially closed* if it is closed under arbitrary coproducts, subobjects, and essential extensions. Let $\langle X \rangle$ denote

Date: December 10, 2024.

 $^{2020\} Mathematics\ Subject\ Classification.\ 18E10\ (primary),\ 16D70,\ 16E50,\ 18E40,\ 18E45$ (secondary).

Key words and phrases. Boolean lattice, definable subcategory, Grothendieck category, spectral category, support, Ziegler spectrum.

the smallest essentially closed subcategory containing X. The support enjoys the following universal property, which is the analogue of a property of support for tensor triangulated categories due to Balmer [1]. We do not assume any monoidal structure, but our analogue seems reasonable given that one has the identity $\langle X \otimes Y \rangle = \langle X \rangle \cap \langle Y \rangle$ in the tensor triangulated setting.

Theorem (Theorem 3.8). The map supp: $Ob A \to Clop(Spc A)$ has the following properties:

- (S0) $supp(X) = \emptyset \text{ for } X = 0.$
- (S1) $\operatorname{supp}(X) = \operatorname{Spc} A$ for every generator X of A.
- $(S \vee) \operatorname{supp}(X) \cup \operatorname{supp}(Y) = \operatorname{supp}(X \oplus Y) \text{ for all objects } X, Y.$
- $(\mathrm{S}\wedge)\ \mathrm{supp}(X)\cap\mathrm{supp}(Y)=\mathrm{supp}(Z)\ \textit{for all objects}\ X,Y,Z\ \textit{with}\ \langle X\rangle\cap\langle Y\rangle=\langle Z\rangle.$

Moreover, for any pair (T, σ) which consists of a topological space T and a map $\sigma \colon \operatorname{Ob} \mathcal{A} \to \operatorname{Clop}(T)$ satisfying the analogue of the above conditions, there exists a unique continuous map $f \colon T \to \operatorname{Spc} \mathcal{A}$ such that $\sigma(X) = f^{-1}(\operatorname{supp}(X))$ for all X.

The local structure of an object can be described in terms of its support. More precisely, we introduce for each injective object X a complete Boolean lattice $\mathbf{D}(X)$ which is given by the clopen subsets of $\mathrm{supp}(X)$; its elements parameterise the direct summands of X up to an appropriate equivalence relation and there is a canonical isomorphism

$$\mathbf{D}(X) \xrightarrow{\sim} \mathbf{B}(\operatorname{End}(X)/J(\operatorname{End}(X))),$$

where $\mathbf{B}(\Lambda)$ denotes the lattice of central idempotents of a ring Λ (Theorem 4.5). The ring $\operatorname{End}(X)/J(\operatorname{End}(X))$ is von Neumann regular; so an elaborated theory of decompositions and types applies [23, 29]. It is interesting to note that any complete Boolean lattice can be realised as $\mathbf{D}(X)$ for some injective object X and some appropriate Grothendieck category.

The support of an object has the useful property that $\operatorname{supp}(X) = \emptyset$ implies X = 0. This one cannot expect when using $\operatorname{Sp} \mathcal{A}$ as target for a support function, because $\operatorname{Sp} \mathcal{A}$ may be empty. Another option for defining support would be to use as target the frame $\mathbf{L}(\mathcal{A})$ of localising subcategories of \mathcal{A} . However, its set of points may be empty as well [37]. The following result shows that $\operatorname{Spc} \mathcal{A}$ is an invariant which encompasses both $\operatorname{Sp} \mathcal{A}$ and $\mathbf{L}(\mathcal{A})$.

Theorem (Theorem 3.16). Taking a localising subcategory to its support induces an embedding of frames

$$\mathbf{L}(\mathcal{A}) \hookrightarrow \mathbf{L}(\operatorname{Spec} \mathcal{A}) \xrightarrow{\sim} \operatorname{Clop}(\operatorname{Spc} \mathcal{A}).$$

For example, when $\mathcal{A} = \operatorname{Mod} \Lambda$ equals the module category of a commutative noetherian ring Λ , then $\operatorname{Sp} \mathcal{A} \xrightarrow{\sim} \operatorname{Spc} \mathcal{A}$ identifies with the set of prime ideals $\operatorname{Spec} \Lambda$, and the above correspondence (1.1) extends Gabriel's bijection between the localising subcategories of $\operatorname{Mod} \Lambda$ and the specialisation closed subsets of $\operatorname{Spec} \Lambda$, which is one of the ingredients for the reconstruction of a noetherian scheme [12].

The work of Gabriel and his use of the spectrum of indecomposable injectives has been a great inspiration for non-commutative algebraic geometry; see for example [3, 22, 35] where several possible definitions of a spectrum of an abelian category are discussed. Here we follow a somewhat different route. For not necessarily commutative rings there is a spectrum which receives much attention and provides another motivation for this work.

The Ziegler spectrum of a ring Λ is given by the isomorphism classes of indecomposable pure-injective Λ -modules, together with a topology introduced by Ziegler in model-theoretic terms [40]. Let Ind Λ denote the Ziegler spectrum of Λ . This identifies with the spectrum of a locally coherent Grothendieck category; so the above theory applies. To be more precise, the module category Mod Λ embeds into

its purity category $\mathbf{P}(\operatorname{Mod}\Lambda)$, and then $\operatorname{Ind}\Lambda$ identifies with $\operatorname{Sp}\mathbf{P}(\operatorname{Mod}\Lambda)$. The Boolean spectrum provides an extension

$$(1.2) \qquad \operatorname{Ind} \Lambda \xrightarrow{\sim} \operatorname{Sp} \mathbf{P}(\operatorname{Mod} \Lambda) \hookrightarrow \operatorname{Spc} \mathbf{P}(\operatorname{Mod} \Lambda) =: \operatorname{PSpc} \Lambda$$

and we propose to study this as an invariant of Λ .

The Ziegler spectrum has been used in various contexts, for instance when studying representations of Artin algebras, but also in more general categorical settings, including derived and stable categories; see [32] for an encyclopedic survey. An important feature of this spectrum is Crawley-Boevey's fundamental correspondence

$$\mathcal{C} \longmapsto \mathcal{C} \cap \operatorname{Ind} \Lambda$$

between definable subcategories of Mod Λ and Ziegler closed subsets of Ind Λ ; see [7]. In order to illustrate the use of the Boolean spectrum PSpc Λ we extend this correspondence as follows.

Theorem (Theorem 6.6). The assignment $\mathcal{C} \mapsto \operatorname{supp}(\mathcal{C})$ induces an inclusion preserving bijection between

- (1) the set of subcategories of Mod Λ that are closed under arbitrary direct sums, pure subobjects, and pure-injective envelopes, and
- (2) the clopen subsets of $PSpc \Lambda$.

In particular, the embedding (1.2) identifies $\mathfrak{C} \cap \operatorname{Ind} \Lambda$ with $\operatorname{supp}(\mathfrak{C}) \cap \operatorname{Ind} \Lambda$ when \mathfrak{C} is definable.

Returning to the above notion of support $X \mapsto \operatorname{supp}(X)$ for an arbitrary Grothendieck category, we mention that there is also an exact version $X \mapsto \operatorname{supp}_{\operatorname{ex}}(X)$ such that for each exact sequence $0 \to X_1 \to X_2 \to X_3 \to 0$ one has

$$\operatorname{supp}_{\mathrm{ex}}(X_i) \subseteq \operatorname{supp}_{\mathrm{ex}}(X_j) \cup \operatorname{supp}_{\mathrm{ex}}(X_k) \quad \text{when} \quad \{i, j, k\} = \{1, 2, 3\}.$$

This is in line with Neeman's celebrated classification of localising subcategories for the derived category of a commutative noetherian ring [30], because it agrees with his notion of support for a complex when specialised to a module. The exact support provides a classification of certain thick subcategories (Theorem 5.5), which is an analogue of the bijective correspondence (1.1) and again not requiring any noetherianess assumption. This generalises a classification for modules over a commutative noetherian ring from [38] and another more recent classification for locally noetherian categories [39].

Let us give a quick outline of this paper. In §2 we recall the basic facts about spectral categories and introduce the Boolean spectrum of a Grothendieck category. In §3 we set up the support for Grothendieck categories and establish the classification of essentially closed subcategories. A first application is given in §4, where coproduct decompositions of objects are studied in terms of Boolean lattices. The following §5 provides the derived analogue of the results from §3, which leads to the exact support. The final §6 discusses the applications for module categories, but in the more general setting of exactly definable categories.

2. Spectral Grothendieck categories

Let \mathcal{A} be a Grothendieck category. We recall the notion of its spectral category from the work of Gabriel and Oberst [13]. Then we study the lattice of localising subcategories and use this to define a space, which we call the Boolean spectrum of \mathcal{A} .

Spectral categories. We write Spec \mathcal{A} for the spectral category of \mathcal{A} which is obtained from \mathcal{A} by formally inverting all essential monomorphisms; see [13] or [36, V.7]. The category Spec \mathcal{A} is again a Grothendieck category in which all exact sequences are split exact. A Grothendieck category with this property is called spectral. The canonical functor

$$P: \mathcal{A} \longrightarrow \operatorname{Spec} \mathcal{A} := \mathcal{A}[\operatorname{Ess}^{-1}]$$

maps any injective envelope $X \to E(X)$ in \mathcal{A} to an isomorphism and it identifies the indecomposable injective objects in \mathcal{A} with the simple objects in Spec \mathcal{A} . These properties are clear from the definition. Also, P is left exact and preserves all coproducts since the class of essential monomorphism is closed under all coproducts. More precisely, for any directed union $X = \sum_{\alpha} X_{\alpha}$ in \mathcal{A} we have $P(X) = \sum_{\alpha} P(X_{\alpha})$. We write Sp \mathcal{A} for the set of isomorphism classes of indecomposable injective objects in \mathcal{A} , and P induces a bijection Sp $\mathcal{A} \xrightarrow{\sim} \operatorname{Sp}(\operatorname{Spec} \mathcal{A})$.

Let $\operatorname{Inj} \mathcal{A}$ denote the full subcategory of injective objects in \mathcal{A} . For later use we note that P induces an equivalence

$$(2.1) \qquad (\operatorname{Inj} \mathcal{A}) / \operatorname{Rad}(\operatorname{Inj} \mathcal{A}) \xrightarrow{\sim} \operatorname{Spec} \mathcal{A},$$

where Rad C denotes the Jacobson radical of an additive category C; see [27, Proposition 2.5.9]. Recall that Rad C is an ideal of morphisms that is given by subgroups

$$Rad(X, Y) \subseteq Hom(X, Y)$$

for each pair of objects X, Y such that $\operatorname{Rad}(X, X)$ equals the Jacobson radical of the endomorphism ring $\operatorname{End}(X)$. A useful consequence of (2.1) is that for any pair of objects $X, Y \in \mathcal{A}$ we have

$$(2.2) P(X) \cong P(Y) \Longleftrightarrow E(X) \cong E(Y)$$

where E(X) denotes an injective envelope of X; see also [13, Satz 1.5].

The following lemma shows that the spectral category controls the decompositions of injective objects.

Lemma 2.1. Let $X \in \mathcal{A}$ be an injective object and $P(X) = Y_1 \oplus Y_2$ a decomposition. Then there exists a decomposition $X = X_1 \oplus X_2$ such that $Y_i = P(X_i)$ for i = 1, 2.

Proof. There is a decomposition into idempotent endomorphisms $\mathrm{id}_{P(X)} = \phi_1 + \phi_2$ such that $Y_i = \mathrm{Ker}\,\phi_i$. Using (2.1) we decompose $\mathrm{id}_X = \psi_1 + \psi_2$ such that $P(\psi_i) = \phi_i$. Now set $X_i = E(\mathrm{Ker}\,\psi_i) \subseteq X$ and apply (2.2).

Spectral categories and their structure are closely related to (modules over) self-injective von Neumann regular rings, because the spectral categories are precisely (up to equivalence) the categories of non-singular injective modules over such rings. This is explained in [13, 33], and for the structure of such rings we refer to [15]. There is an elaborated decomposition theory for operator algebras which uses certain types and goes back to Murray, von Neumann, and Kaplansky [23, 29]. The analogous decomposition theory for spectral categories is developed in [16, 33]. For some further structure of spectral categories, see [10].

Localising subcategories. Let \mathcal{A} be a Grothendieck category. A full subcategory of \mathcal{A} is *localising* if it is closed under subobjects, quotients, extensions, and arbitrary coproducts. The localising subcategories of \mathcal{A} form a set, because for a fixed generator $G \in \mathcal{A}$ any localising subcategory $\mathcal{U} \subseteq \mathcal{A}$ is determined by the set

¹Given a spectral category and a generator G, the endomorphism ring of G is self-injective von Neumann regular, and the equivalence is induced by Hom(G, -).

of quotients G/G' which belong to \mathcal{U} . The localising subcategories are partially ordered by inclusion and closed under arbitrary intersections; so they form a complete lattice which we denote by $\mathbf{L}(A)$.

For a localising subcategory $\mathcal{U} \subseteq \mathcal{A}$ and an object $X \in \mathcal{A}$ let $t_{\mathcal{U}}(X) \subseteq X$ denote the maximal subobject in \mathcal{U} . The following lemma provides a way of describing the join in $\mathbf{L}(\mathcal{A})$.

Lemma 2.2. Let (\mathcal{U}_{α}) be a family in $\mathbf{L}(\mathcal{A})$ and set $\mathcal{U} = \bigvee_{\alpha} \mathcal{U}_{\alpha}$. For any $X \in \mathcal{A}$ we have

$$\sum_{\alpha} t_{\mathfrak{U}_{\alpha}}(X) = t_{\mathfrak{U}}(X).$$

In particular,

$$t_{\mathfrak{U}}(X) = 0 \qquad \iff \qquad t_{\mathfrak{U}_{\alpha}}(X) = 0 \quad \text{for all} \quad \alpha.$$

Proof. We begin with the second assertion. Let $X \to E(X)$ denote an injective envelope. Then $t_{\mathcal{U}}(X) = 0$ if and only if $t_{\mathcal{U}}(E(X)) = 0$. So we may assume that X is injective. But then the assertion is clear, since $t_{\mathcal{U}}(X) = 0$ if and only if $\operatorname{Hom}(U,X) = 0$ for all $U \in \mathcal{U}$, and the kernel of $\operatorname{Hom}(-,X)$ is a localising subcategory.

Now consider the subobjects

$$X' := \sum_{\alpha} t_{\mathfrak{U}_{\alpha}}(X) \subseteq t_{\mathfrak{U}}(X) \subseteq X$$

If X is in \mathcal{U} , then $t_{\mathcal{U}_{\alpha}}(X/X') = 0$ for all α , and therefore the first part of the proof yields $t_{\mathcal{U}}(X/X') = 0$. Thus X' = X.

A complete lattice is a *frame* if finite meets distribute over arbitrary joins, and it is a *coframe* if finite joins distribute over arbitrary meets. It is well known that the localising subcategories of a Grothendieck form a frame; see for instance [2] (in a more general categorical context) or [14] (for module categories). We refer to [37] for a proof that uses an explicit construction of the join in the lattice of localising subcategories.

Proposition 2.3. Let A be a Grothendieck category. For an element U and a family (V_{α}) in L(A) we have

$$\mathcal{U} \wedge \left(\bigvee_{\alpha} \mathcal{V}_{\alpha}\right) = \bigvee_{\alpha} (\mathcal{U} \wedge \mathcal{V}_{\alpha}).$$

Thus the lattice $\mathbf{L}(A)$ is a frame.

Proof. The inclusion \supseteq is automatic; so we need to show \subseteq . Let X be an object in $\mathcal{U} \wedge (\bigvee_{\alpha} \mathcal{V}_{\alpha})$. Then Lemma 2.2 implies $X = \sum_{\alpha} t_{\mathcal{V}_{\alpha}}(X)$, and therefore X belongs to $\bigvee_{\alpha} (\mathcal{U} \wedge \mathcal{V}_{\alpha})$ since $t_{\mathcal{V}_{\alpha}}(X) \in \mathcal{U} \wedge \mathcal{V}_{\alpha}$ for each α .

Proposition 2.4. Let A be a spectral Grothendieck category. For an element U and a family (V_{α}) in $\mathbf{L}(A)$ we have

$$\mathcal{U} \vee \left(\bigwedge_{\alpha} \mathcal{V}_{\alpha} \right) = \bigwedge_{\alpha} (\mathcal{U} \vee \mathcal{V}_{\alpha}).$$

Thus the lattice L(A) is a coframe.

Proof. The inclusion \subseteq is automatic; so we need to show \supseteq . Any object X can be written as $X = t_{\mathcal{U}}(X) \oplus X/t_{\mathcal{U}}(X)$. A simple calculation shows that $X \in \mathcal{U} \vee \mathcal{V}_{\alpha}$ if and only if $X/t_{\mathcal{U}}(X) \in \mathcal{V}_{\alpha}$. Thus $X \in \bigwedge_{\alpha} (\mathcal{U} \vee \mathcal{V}_{\alpha})$ implies $X/t_{\mathcal{U}}(X) \in \bigwedge_{\alpha} \mathcal{V}_{\alpha}$, and therefore $X \in \mathcal{U} \vee (\bigwedge_{\alpha} \mathcal{V}_{\alpha})$.

Discrete versus continuous. Let \mathcal{A} be a spectral category. The category \mathcal{A} is called *discrete* if every object is a coproduct of simple objects. Any object $X \in \mathcal{A}$ admits a decomposition

$$X = \operatorname{soc}(X) \oplus \operatorname{rad}(X),$$

where the $socle\ soc(X)$ is the sum of all simple subobjects and the $radical\ rad(X)$ is the intersection of all maximal subobjects. We set

$$\mathcal{A}_{d} := \{X \in \mathcal{A} \mid \operatorname{soc}(X) = X\}$$
 and $\mathcal{A}_{c} := \{X \in \mathcal{A} \mid \operatorname{rad}(X) = X\}.$

This yields a decomposition $\mathcal{A} = \mathcal{A}_d \times \mathcal{A}_c$ of spectral Grothendieck categories such that \mathcal{A}_d is discrete and $(\mathcal{A}_c)_d = 0 = (\mathcal{A}_d)_c$; see [13, §3].

Lemma 2.5. Let \mathcal{A} be spectral. The assignment $\mathcal{U} \mapsto \mathcal{U} \cap \operatorname{Sp} \mathcal{A}$ induces a lattice isomorphism between $\mathbf{L}(\mathcal{A}_d)$ and the lattice of subsets of $\operatorname{Sp} \mathcal{A}$.

Proof. The inverse map sends a subset $\mathcal{V} \subseteq \operatorname{Sp} \mathcal{A}$ to the subcategory of \mathcal{A} consisting of all coproducts of objects in \mathcal{V} .

Example 2.6. Let \mathcal{A} be a locally noetherian Grothendieck category. Then its spectral category Spec \mathcal{A} is discrete [12, IV.2]. Thus in this case $\mathbf{L}(\operatorname{Spec} \mathcal{A})$ identifies with the lattice of subsets of $\operatorname{Sp} \mathcal{A}$.

The Boolean spectrum. Let \mathcal{A} be a Grothendieck category. For a subcategory $\mathcal{C} \subseteq \mathcal{A}$ we set

$$\mathfrak{C}^{\perp} := \{ X \in \mathcal{A} \mid \operatorname{Hom}(C, X) = 0 \text{ for all } C \in \mathfrak{C} \},$$

$${}^{\perp}\mathfrak{C} := \{ X \in \mathcal{A} \mid \operatorname{Hom}(X, C) = 0 \text{ for all } C \in \mathfrak{C} \}.$$

Localising subcategories of \mathcal{A} are in bijective correspondence to hereditary torsion pairs, by taking a subcategory \mathcal{C} to the pair $(\mathcal{C}, \mathcal{C}^{\perp})$; see [36, VI.3]. Torsion pairs for spectral categories admit the following elementary description.

Lemma 2.7. Let A be a spectral Grothendieck category. A pair $(\mathfrak{C}, \mathfrak{D})$ of full subcategories is a torsion pair if and only if

- (1) C and D are closed under direct summands,
- (2) $\mathcal{C} \cap \mathcal{D} = 0$, and
- (3) $A = \{C \oplus D \mid C \in \mathcal{C}, D \in \mathcal{D}\}.$

Proof. Clearly, for any torsion pair $(\mathcal{C}, \mathcal{D})$ the conditions (1)–(3) hold. Conversely, these conditions imply that $\mathcal{C}^{\perp} = \mathcal{D}$ and $\mathcal{C} = {}^{\perp}\mathcal{D}$. Thus $(\mathcal{C}, \mathcal{D})$ is a torsion pair. \square

We record the above in lattice-theoretical terms. Recall that a bounded lattice L is complemented if for each $x \in L$ there exists a complement $y \in L$, so $x \wedge y = 0$ and $x \vee y = 1$. Such a complement is unique when L is distributive, and then we write $\neg x$ for the complement of x. A complemented distributive lattice is called Boolean lattice.

Proposition 2.8. Let A be a spectral Grothendieck category. Then the lattice L(A) is a complete Boolean lattice.

Proof. First observe that all torsion pairs are hereditary since \mathcal{A} is spectral. Now let $(\mathcal{C}, \mathcal{D})$ be a torsion pair. Then it follows from Lemma 2.7 that $(\mathcal{D}, \mathcal{C})$ is again a torsion pair. Thus $\mathcal{D} = \mathcal{C}^{\perp}$ is a complement for $\mathcal{C} \in \mathbf{L}(\mathcal{A})$.

Let us collect some basic facts from Stone duality; for details and proofs see [9, 21]. Let $A = (A, \vee, \wedge)$ be a Boolean lattice. We may identify A with the corresponding Boolean ring $(A, +, \cdot)$, with operations given by

$$x + y := (x \land \neg y) \lor (y \land \neg x)$$
 and $x \cdot y := x \land y$,

and keeping in mind that the category of Boolean lattices is isomorphic to the category of Boolean rings. Let $\operatorname{Spec}(A)$ denote the spectrum of prime ideals of A with the Zariski topology. For a topological space T we write $\operatorname{Clop}(T)$ for its Boolean lattice of clopen subsets. This yields an adjoint pair of contravariant functors

$$\textbf{Top} \xleftarrow{\frac{\text{Clop} = \text{Hom}(-, \mathbf{2})}{\text{Spec} = \text{Hom}(-, \mathbf{2})}} \textbf{BRing}$$

where $\mathbf{2} = \{0, 1\}$ is either viewed as topological space with the discrete topology, or $\mathbf{2} = \mathbb{F}_2$ is viewed as a Boolean ring. This means there is a natural bijection

which sends $\phi \colon A \to \operatorname{Clop}(T)$ to the map $T \to \operatorname{Spec}(A)$ given by

$$p \longmapsto \{x \in A \mid p \notin \phi(x)\}$$
 for $p \in T$.

Stone duality implies that for any Boolean ring A the unit of the adjunction

$$(2.4) A \longrightarrow \operatorname{Clop}(\operatorname{Spec}(A)), \quad x \mapsto \{\mathfrak{p} \in \operatorname{Spec}(A) \mid x \notin \mathfrak{p}\}\$$

is an isomorphism.

Definition 2.9. The Boolean spectrum of a Grothendieck category A is the space

$$\operatorname{Spc} A := \operatorname{Spec}(\mathbf{L}(\operatorname{Spec} A)).$$

For a Grothendieck category \mathcal{A} the decomposition $\operatorname{Spec} \mathcal{A} = (\operatorname{Spec} \mathcal{A})_d \times (\operatorname{Spec} \mathcal{A})_c$ implies that $\operatorname{Spc} \mathcal{A}$ equals the disjoint union

(2.5)
$$\operatorname{Spc} A = \operatorname{Spec}(\mathbf{L}((\operatorname{Spec} A)_{d})) \sqcup \operatorname{Spec}(\mathbf{L}((\operatorname{Spec} A)_{c}))$$

where

$$\operatorname{Spec}(\mathbf{L}((\operatorname{Spec} \mathcal{A})_{\operatorname{d}})) \cong \operatorname{Sp} \mathcal{A}$$

carries the discrete topology. In particular, we see that the set $\operatorname{Sp} \mathcal{A}$ of isoclasses of indecomposable injective objects embeds canonically into $\operatorname{Spc} \mathcal{A}$; cf. Lemma 2.5.

Central idempotents. For an additive category \mathcal{A} let $Z(\mathcal{A})$ denote its *centre* which is the ring of natural transformations $\mathrm{id}_{\mathcal{A}} \to \mathrm{id}_{\mathcal{A}}$. For a ring Λ let $\mathbf{B}(\Lambda)$ denote the lattice of central idempotents, with operations given by

$$x \lor y := x + y - x \cdot y$$
 and $x \land y := x \cdot y$.

Let \mathcal{A} be a spectral Grothendieck category. Any torsion pair $(\mathcal{C}, \mathcal{D})$ gives rise to a functorial decomposition $X = t_{\mathcal{C}}(X) \oplus t_{\mathcal{D}}(X)$ for each object $X \in \mathcal{A}$, which amounts to an idempotent $e_{\mathcal{C}} \colon \mathrm{id}_{\mathcal{A}} \to \mathrm{id}_{\mathcal{A}}$ such that the image of $(e_{\mathcal{C}})_X$ equals $t_{\mathcal{C}}(X)$. Conversely, any idempotent $e \in Z(\mathcal{A})$ yields a torsion pair $(\mathcal{C}, \mathcal{D})$ if we set $\mathcal{C} = \{X \in \mathcal{A} \mid \operatorname{Im} e_X = X\}$ and $\mathcal{D} = \{X \in \mathcal{A} \mid \operatorname{Ker} e_X = X\}$. Clearly, these operations are mutually inverse to each other, and this yields another description of $\mathbf{L}(\mathcal{A})$.

Lemma 2.10. Let A be a spectral Grothendieck category. Then the assignment $\mathfrak{C} \mapsto e_{\mathfrak{C}}$ induces a lattice isomorphism

$$\mathbf{L}(\mathcal{A}) \xrightarrow{\sim} \mathbf{B}(Z(\mathcal{A})).$$

Moreover, when $G \in \mathcal{A}$ is a generator with $\Lambda = \operatorname{End}(G)$, then

$$x \longmapsto \{X \in \mathcal{A} \mid \operatorname{Hom}(G,X)x = \operatorname{Hom}(G,X)\}$$

induces a lattice isomorphism

$$\mathbf{B}(\Lambda) \xrightarrow{\sim} \mathbf{L}(\mathcal{A}).$$

Proof. The first assertion is clear from the above discussion. Let \mathcal{A}' denote the category of non-singular injective right modules over Λ . The functor Hom(G,-) induces an equivalence $\mathcal{A} \xrightarrow{\sim} \mathcal{A}'$ and an isomorphism $Z(\mathcal{A}) \xrightarrow{\sim} Z(\mathcal{A}')$. On the other hand, multiplication with a central element induces an isomorphism $Z(\Lambda) \xrightarrow{\sim} Z(\mathcal{A}')$. Thus the second assertion follows from the first.

I am grateful to Ken Goodearl for pointing out the following converse to Proposition 2.8.

Proposition 2.11 (Goodearl). Every complete Boolean lattice is isomorphic to L(A) for some spectral Grothendieck category A.

Proof. We begin with the following observation. Let Λ be a right self-injective von Neumann regular ring and let \mathcal{A} denote the corresponding spectral category consisting of the non-singular injective Λ -modules. Then it follows from Lemma 2.10 that the assignment $x \mapsto \{M \in \mathcal{A} \mid Mx = M\}$ induces a lattice isomorphism $\mathbf{B}(\Lambda) \xrightarrow{\sim} \mathbf{L}(\mathcal{A})$.

Now let L be a complete Boolean lattice and view this as a Boolean ring. Then L is von Neumann regular with $\mathbf{B}(L) = L$, and the ring is self-injective because L is complete. Thus for \mathcal{A} the category of non-singular injective L-modules, we have that $L \cong \mathbf{L}(\mathcal{A})$.

The proof shows that the realisation of a complete Boolean lattice is canonical. For a ring Λ let $\operatorname{Inj}_{\operatorname{ns}} \Lambda$ denote the category of non-singular injective right modules over Λ . Then for any complete Boolean lattice L there is a canonical isomorphism

$$L \xrightarrow{\sim} \mathbf{L}(\operatorname{Inj}_{\operatorname{ns}} L).$$

The above result suggests a procedure for constructing explicit examples of objects with specific decomposition properties. For instance, we obtain examples of superdecomposable objects. Recall that an object in an abelian category is *superdecomposable* if there are no indecomposable direct summands.

Example 2.12 (Goodearl). Let Λ be an infinite direct product of copies of $\mathbb{Z}/2$ and let $\bar{\Lambda}$ denote the maximal quotient ring of $\Lambda/\operatorname{soc}(\Lambda)$. Then $\bar{\Lambda}$ is injective as a $\bar{\Lambda}$ -module and superdecomposable by construction, since $\operatorname{soc}(\bar{\Lambda}) = 0$. Moreover, $\mathbf{B}(\bar{\Lambda}) = \bar{\Lambda}$ for its Boolean lattice of direct summands.

The ring $\bar{\Lambda}$ has characteristic 2 and is of type I, but further examples of different characteristic and different type can be constructed via methods from Propositions 5-3.11 and 5-3.12 in [17].

For general Grothendieck categories the decompositions of objects will be discussed in more detail in §4.

Coproduct decompositions. We provide an explicit description of the join in the lattice L(A) when A is spectral. We use the following property of a Boolean lattice.

Lemma 2.13. Let A be a complete Boolean lattice and $x = \bigvee_{\alpha} x_{\alpha}$ an element. Then there are elements $y_{\alpha} \leq x_{\alpha}$ such that $x = \bigvee_{\alpha} y_{\alpha}$ and $y_{\alpha} \wedge y_{\beta} = 0$ for all $\alpha \neq \beta$.

Proof. We consider the set of families (y_{α}) of objects in A such that $y_{\alpha} \leq x_{\alpha}$ and $y_{\alpha} \wedge y_{\beta} = 0$ for all $\alpha \neq \beta$. This set is partially ordered via $(y_{\alpha}) \leq (y'_{\alpha})$ if $y_{\alpha} \leq y'_{\alpha}$ for all α . An application of Zorn's lemma yields a maximal element and it is easily checked that $x = \bigvee_{\alpha} y_{\alpha}$ for such maximal (y_{α}) .

Proposition 2.14. Let A be a spectral Grothendieck category. For a family (U_{α}) in $\mathbf{L}(A)$ we have

$$\bigvee_{\alpha} \mathfrak{U}_{\alpha} = \left\{ \coprod_{\alpha} X_{\alpha} \mid X_{\alpha} \in \mathfrak{U}_{\alpha} \text{ for all } \alpha \right\}.$$

Proof. Set $\mathcal{U} = \bigvee_{\alpha} \mathcal{U}_{\alpha}$. We appy Lemma 2.13 and obtain a decomposition $\mathcal{U} = \bigvee_{\alpha} \mathcal{V}_{\alpha}$ such that $\mathcal{V}_{\alpha} \subseteq \mathcal{U}_{\alpha}$ and $\mathcal{V}_{\alpha} \cap \mathcal{V}_{\beta} = 0$ for all $\alpha \neq \beta$. Given $X \in \mathcal{U}$ we have $X = \sum_{\alpha} t_{\mathcal{V}_{\alpha}}(X)$ by Lemma 2.2. Moreover, $t_{\mathcal{V}_{\beta}}(X) \cap \sum_{\alpha \neq \beta} t_{\mathcal{V}_{\alpha}}(X) = 0$ for all β since $\mathcal{V}_{\beta} \wedge \left(\bigvee_{\alpha \neq \beta} \mathcal{V}_{\alpha}\right) = 0$. Thus $X = \coprod_{\alpha} t_{\mathcal{V}_{\alpha}}(X)$.

Functoriality. We briefly discuss the functoriality of the assignment $\mathcal{A} \mapsto \operatorname{Spec} \mathcal{A}$. An exact and coproduct preserving functor $\mathcal{A} \to \mathcal{B}$ between Grothendieck categories admits a right adjoint which restricts to an additive functor $\operatorname{Inj} \mathcal{B} \to \operatorname{Inj} \mathcal{A}$. Using (2.1) one obtains an exact functor $\operatorname{Spec} \mathcal{B} \to \operatorname{Spec} \mathcal{A}$, at least when $\mathcal{A} \to \mathcal{B}$ is the inclusion of a localising subcategory or a quotient functor.

Now let $\mathcal{A}' \subseteq \mathcal{A}$ be a localising subcategory and set $\mathcal{A}'' = \mathcal{A}/\mathcal{A}'$.

Lemma 2.15. The exact sequence $A' \rightarrow A \rightarrow A''$ induces an exact sequence

$$\operatorname{Spec} A'' \longrightarrow \operatorname{Spec} A \twoheadrightarrow \operatorname{Spec} A'.$$

These functors admit right adjoints and yield decompositions

$$\operatorname{Spec} A = \operatorname{Spec} A' \times \operatorname{Spec} A''$$
 and $\operatorname{Spc} A = \operatorname{Spc} A' \sqcup \operatorname{Spc} A''$.

Proof. The functor $\operatorname{Spec} \mathcal{A} \to \operatorname{Spec} \mathcal{A}'$ is induced by $t_{\mathcal{A}'} \colon \mathcal{A} \to \mathcal{A}'$. It is easily checked that $\operatorname{Spec} \mathcal{A}'' \to \operatorname{Spec} \mathcal{A}$ identifies $\operatorname{Spec} \mathcal{A}''$ with the kernel of $\operatorname{Spec} \mathcal{A} \to \operatorname{Spec} \mathcal{A}'$, which is closed under arbitrary coproducts. The inclusion $\mathcal{A}' \to \mathcal{A}$ preserves essential extensions and induces therefore a functor $\operatorname{Spec} \mathcal{A}' \to \operatorname{Spec} \mathcal{A}$ which is a right inverse for $\operatorname{Spec} \mathcal{A} \to \operatorname{Spec} \mathcal{A}'$. Thus $\operatorname{Spec} \mathcal{A} \to \operatorname{Spec} \mathcal{A}'$ is a localisation functor, and this yields the decomposition $\operatorname{Spec} \mathcal{A} = \operatorname{Spec} \mathcal{A}' \times \operatorname{Spec} \mathcal{A}''$. The functor from spectral categories to spaces given by $\mathcal{C} \mapsto \operatorname{Spec} \mathbf{L}(\mathcal{C})$ is contravariant and takes products to coproducts.

Monoidal structure. Let \mathcal{A} be a spectral Grothendieck category. Any choice of a generator $G \in \mathcal{A}$ induces a tensor product $-\otimes_G -: \mathcal{A} \times \mathcal{A} \to \mathcal{A}$ as follows. We write \mathcal{G} for the full subcategory given by the direct summands of finite direct sums of copies of G. Let $-\otimes_G -: \mathcal{G} \times \mathcal{G} \to \mathcal{G}$ be the unique additive functor such that $C \otimes_G G = C = G \otimes_G C$ for all $C \in \mathcal{G}$. An arbitrary object $X \in \mathcal{A}$ can be written canonically as (filtered) colimit $X = \operatorname{colim}_{C \to X} C$ where $C \to X$ runs through all morphisms from objects in \mathcal{G} . Then set

$$X \otimes_G Y := \underset{\substack{C \to X \\ D \to Y}}{\operatorname{colim}} C \otimes_G D.$$

It is clear that the $X \otimes_G$ – is exact and preserves all coproducts. Also, any localising subcategory is a tensor ideal. Let $\langle X \rangle$ denote the localising subcategory generated by X.

Lemma 2.16. For a pair of objects X, Y in A we have $\langle X \rangle \cap \langle Y \rangle = \langle X \otimes_G Y \rangle$.

Proof. We have $t_{\langle X \rangle} = t_{\langle X \rangle}(G) \otimes_G -$. Thus for an object $M \in \langle X \rangle \cap \langle Y \rangle$ we have

$$M \cong t_{\langle X \rangle}(G) \otimes_G t_{\langle Y \rangle}(G) \otimes_G M \in \langle t_{\langle X \rangle}(G) \otimes_G t_{\langle Y \rangle}(G) \rangle \subseteq \langle X \otimes_G Y \rangle.$$

The other inclusion is clear.

3. Essentially closed subcategories

We fix a Grothendieck category \mathcal{A} and introduce a class of subcategories which is controlled by its spectral category. This yields a notion of support for the objects of \mathcal{A} . To simplify notation we set

$$\mathbf{S}(\mathcal{A}) := \mathbf{L}(\operatorname{Spec} \mathcal{A}) \xrightarrow{\sim} \operatorname{Clop}(\operatorname{Spc} \mathcal{A}).$$

Essentially closed subcategories and support. We begin with the main definition of this section; the terminology is justified by Theorem 3.3, because we look at the operations that are preserved by the functor $P: \mathcal{A} \to \operatorname{Spec} \mathcal{A}$ inverting all essential monomorphisms.

Definition 3.1. A full subcategory of \mathcal{A} is called *essentially closed* if it is closed under arbitrary coproducts, subobjects, and essential extensions.

The essentially closed subcategories are partially ordered by inclusion and closed under arbitrary intersections; so they form a complete lattice. When $\mathcal A$ is spectral, then its essentially closed subcategories are precisely its localising subcategories. For a class $\mathcal X\subseteq\mathcal A$ let $\langle\mathcal X\rangle$ denote the smallest essentially closed subcategory of $\mathcal A$ that contains $\mathcal X$.

Lemma 3.2. Let \mathcal{A} be a Grothendieck category.

- (1) If $U \subseteq \operatorname{Spec} A$ is localising, then $P^{-1}(U) \subseteq A$ is essentially closed.
- (2) If $\mathcal{C} \subseteq \mathcal{A}$ is essentially closed, then $P(\mathcal{C}) \subseteq \operatorname{Spec} \mathcal{A}$ is localising.

Proof. (1) Let $X \subseteq Y$ be a subobject in \mathcal{A} with $P(Y) \in \mathcal{U}$. Then $P(X) \in \mathcal{U}$ since P is left exact. Thus $P^{-1}(\mathcal{U})$ is closed under subobjects. Analogously, $P^{-1}(\mathcal{U})$ is closed under coproducts since P preserves coproducts, and it is closed under essential extensions as P maps those to isomorphisms.

(2) First observe that the subcategory $P(\mathcal{C})$ is replete, so given a pair of objects $X,Y\in\mathcal{A}$ such that $P(X)\cong P(Y)$, then $X\in\mathcal{C}$ implies $Y\in\mathcal{C}$. This follows from (2.2), because when X and Y have isomorphic injective envelopes, then $X\in\mathcal{C}$ implies $Y\in\mathcal{C}$. Next we show that $P(\mathcal{C})$ is closed under coproducts and subobjects. Clearly, $P(\mathcal{C})$ is closed under coproducts since P preserves coproducts. Any subobject $Y\subseteq P(X)$ of an object in $P(\mathcal{C})$ is actually a direct summand of P(X), and we may assume that X is injective. Using the equivalence (2.1), we have $Y\cong \operatorname{Ker} P(\phi)\cong P(\operatorname{Ker}\phi)$ for some morphism $\phi\colon X\to X$. Thus $P(\mathcal{C})$ is closed under subobjects.

For an object $X \in \mathcal{A}$ its *support* is by definition

(3.1)
$$\operatorname{supp}(X) := \inf\{\mathcal{U} \in \mathbf{S}(\mathcal{A}) \mid P(X) \in \mathcal{U}\} = \langle P(X) \rangle.$$

For a class $\mathfrak{X} \subseteq \mathcal{A}$ we set $\operatorname{supp}(\mathfrak{X}) := \bigvee_{X \in \mathfrak{X}} \operatorname{supp}(X)$ and note that

(3.2)
$$\operatorname{supp}(\mathfrak{X}) = \bigvee_{X \in \mathfrak{X}} \langle P(X) \rangle = \langle P(\mathfrak{X}) \rangle.$$

Theorem 3.3. Let A be a Grothendieck category. The assignment

$$\mathcal{C} \longmapsto \operatorname{supp}(\mathcal{C}) = P(\mathcal{C})$$

induces a lattice isomorphism

$$\{\mathfrak{C} \subseteq \mathcal{A} \mid \mathfrak{C} \text{ essentially closed}\} \xrightarrow{\sim} \mathbf{S}(\mathcal{A}).$$

The inverse map is given by $\mathcal{U} \mapsto P^{-1}(\mathcal{U})$.

Proof. The functor P equals the identity on objects. From this and Lemma 3.2 it follows that $\mathcal{C} \mapsto P(\mathcal{C})$ and $\mathcal{U} \mapsto P^{-1}(\mathcal{U})$ provide mutually inverse bijections between the essentially closed subcategories of \mathcal{A} and $\operatorname{Spec} \mathcal{A}$, respectively. The equality $\operatorname{supp}(\mathcal{C}) = P(\mathcal{C})$ follows from (3.2).

In this generality the theorem seems to be new. Special cases were studied before; see the discussion in Remark 5.7 when we establish another correspondence for subcategories of \mathcal{A} . Essentially closed subcategories were introduced for module categories by Dauns under the name 'saturated class' [8]. He showed that they form a Boolean lattice, though his proof is different from ours and does not use the spectral category. Saturated classes were used in [8] to develop a decomposition theory for modules, analogous to the one for von Neumann regular rings. Proposition 4.1 gives some flavour of such decomposition results.

Corollary 3.4. Every essentially closed subcategory of A is of the form $\langle X \rangle$ for some object $X \in A$.

Proof. The assertion is clear when \mathcal{A} is spectral, because any localising subcategory is a Grothendieck category and contains therefore a generator. When \mathcal{A} is arbitrary and $\mathcal{C} \subseteq \mathcal{A}$ is essentially closed, we can choose $X \in \mathcal{A}$ with $P(\mathcal{C}) = \langle P(X) \rangle$. This implies $\mathcal{C} = \langle X \rangle$.

Remark 3.5. The complement of an essentially closed subcategory $\mathcal{C} \subseteq \mathcal{A}$ admits an easy description: it is the essentially closed subcategory consisting of all objects $X \in \mathcal{A}$ such that any subobject of X is zero when it belongs to \mathcal{C} .

Support data. The notion of a support datum for tensor triangulated categories was introduced by Balmer in [1]. The following is the analogue for Grothendieck categories, where the intersection of essentially closed subcategories plays the role of the tensor product.²

Definition 3.6. A support datum on a Grothendieck category \mathcal{A} is a pair (T, σ) consisting of a topological space T and a map $\sigma \colon \mathrm{Ob}\,\mathcal{A} \to \mathrm{Clop}(T)$ such that

- (S0) $\sigma(X) = \emptyset$ for X = 0,
- (S1) $\sigma(X) = T$ for every generator X of \mathcal{A} ,
- (SV) $\sigma(X) \cup \sigma(Y) = \sigma(X \oplus Y)$ for all objects X, Y in \mathcal{A} , and
- $(S \land) \ \sigma(X) \cap \sigma(Y) = \sigma(Z) \text{ for all objects } X, Y, Z \text{ in } \mathcal{A} \text{ with } \langle X \rangle \cap \langle Y \rangle = \langle Z \rangle.$

A morphism $(T, \sigma) \to (T', \sigma')$ is given by a continuous map $f: T \to T'$ such that $\sigma(X) = f^{-1}(\sigma'(X))$ for all $X \in \mathcal{A}$.

It is convenient to reformulate the notion of a support datum as follows.

Lemma 3.7. Let (T, σ) be a support datum. Given objects X, Y in A, the condition $\langle X \rangle \subseteq \langle Y \rangle$ implies $\sigma(X) \subseteq \sigma(Y)$. Therefore the assignment $\langle X \rangle \mapsto \sigma(X)$ induces a map

$$\{\mathfrak{C} \subseteq \mathcal{A} \mid \mathfrak{C} \ essentially \ closed\} \longrightarrow \operatorname{Clop}(T)$$

which is a lattice homomorphism.

Proof. From condition $(S \land)$ it follows that $\langle X \rangle \subseteq \langle Y \rangle$ implies $\sigma(X) \subseteq \sigma(Y)$. From Corollary 3.4 we know that every essentially closed subcategory of \mathcal{A} is generated by a single object. Thus the assignment $\langle X \rangle \mapsto \sigma(X)$ yields a well defined map $\{\mathcal{C} \subseteq \mathcal{A} \mid \mathcal{C} \text{ essentially closed}\} \to \operatorname{Clop}(T)$. Conditions (S0)– $(S \land)$ imply that this is a lattice homomorphism.

By slight abuse of notation and using the isomorphism (2.4) we write supp for the composite

$$\mathrm{Ob}\,\mathcal{A} \xrightarrow{\mathrm{supp}} \mathbf{S}(\mathcal{A}) \xrightarrow{\sim} \mathrm{Clop}(\mathrm{Spec}(\mathbf{S}(\mathcal{A}))) = \mathrm{Clop}(\mathrm{Spc}\,\mathcal{A}).$$

²This is reasonable since $\langle X \otimes Y \rangle = \langle X \rangle \cap \langle Y \rangle$ for objects X, Y in a tensor triangulated category, where $\langle X \rangle$ denotes the radical thick tensor ideal generated by X. See also Lemma 2.16 for the case of a spectral category.

Theorem 3.8. The pair (Spc A, supp) is a support datum on A. Moreover, for any support datum (T, σ) there is a unique continuous map $f: T \to \operatorname{Spc} A$ such that $\sigma(X) = f^{-1}(\operatorname{supp}(X))$ for all $X \in A$. Explicitly, the map f is given by the adjunction (2.3) and therefore

$$f(p) = \{ \langle X \rangle \in \mathbf{S}(\mathcal{A}) \mid p \notin \sigma(X) \}$$
 for $p \in T$.

Proof. It is clear that (Spc \mathcal{A} , supp) is a support datum. Each support datum (T,σ) induces a homomorphism $\{\mathfrak{C}\subseteq\mathcal{A}\mid \mathfrak{C} \text{ essentially closed}\}\to \mathrm{Clop}(T)$ by Lemma 3.7. The composite with the isomorphism $\mathbf{S}(\mathcal{A})\stackrel{\sim}{\to} \{\mathfrak{C}\subseteq\mathcal{A}\mid \mathfrak{C} \text{ essentially closed}\}$ from Theorem 3.3 yields a lattice homomorphism, which corresponds via the adjunction (2.3) to a continuous map $T\to\mathrm{Spec}(\mathbf{S}(\mathcal{A}))=\mathrm{Spc}\,\mathcal{A}$ with the above explicit description.

The clopen subsets of $\operatorname{Spc} \mathcal{A}$ form a complete lattice. This reflects the existence of arbitrary coproducts in \mathcal{A} , but it need not hold for an arbitrary space.

Proposition 3.9. Let (X_{α}) be a family of objects in A. Then we have

$$\operatorname{supp}\left(\coprod_{\alpha} X_{\alpha}\right) = \bigvee_{\alpha} \operatorname{supp}(X_{\alpha}).$$

Proof. The assertion follows from the equality $\langle \coprod_{\alpha} X_{\alpha} \rangle = \bigvee_{\alpha} \langle X_{\alpha} \rangle$.

We end our discussion of support with a simple observation which demonstrates the benefit of the extension $\operatorname{Sp} \mathcal{A} \hookrightarrow \operatorname{Spc} \mathcal{A}$.

Remark 3.10. For $X \in \mathcal{A}$ we have $\operatorname{supp}(X) \neq \emptyset$ when $X \neq 0$. This property cannot be expected for a map $\operatorname{Ob} \mathcal{A} \to \operatorname{Sp} \mathcal{A}$ since $\operatorname{Sp} \mathcal{A}$ may be empty.

Localising subcategories. Next we show that the functor $\mathcal{A} \to \operatorname{Spec} \mathcal{A}$ induces an embedding for the frame of localising subcategories. To this end we consider essentially closed subcategories that are closed under arbitrary products because they arise naturally from localising subcategories. The following lemma explains a tight connection.

Lemma 3.11. For a full subcategory $\mathbb{D} \subseteq \mathcal{A}$ the following are equivalent.

- (1) The subcategory \mathfrak{D} is essentially closed and closed under arbitrary products.
- (2) There is a localising subcategory $\mathfrak{C} \subseteq \mathcal{A}$ such that $\mathfrak{C}^{\perp} = \mathfrak{D}$.

Proof. (1) \Rightarrow (2): We set $\mathcal{C} := {}^{\perp}\mathcal{D}$ and this is localising since \mathcal{D} is closed under injective envelopes, so $\mathcal{C} = {}^{\perp}(\mathcal{D} \cap \operatorname{Inj} \mathcal{A})$. It remains to note that $\mathcal{C}^{\perp} = \mathcal{D}$ where one uses that \mathcal{D} is closed under arbitrary products; see [36, Proposition VI.2.2].

$$(2) \Rightarrow (1)$$
: This is clear.

We continue with a sequence of technical lemmas.

Lemma 3.12. Let (\mathfrak{C}_{α}) be a finite family of essentially closed subcategories and $X \in \mathcal{A}$ an injective object. Then X belongs to $\bigvee_{\alpha} \mathfrak{C}_{\alpha}$ if and only if there is a decomposition $X = \coprod_{\alpha} X_{\alpha}$ such that $X_{\alpha} \in \mathfrak{C}_{\alpha}$ for all α .

Proof. Consider $\mathcal{C} = \mathcal{C}_1 \vee \mathcal{C}_2$. If $X_1 \in \mathcal{C}_1$ and $X_2 \in \mathcal{C}_2$, then $X_1 \oplus X_2 \in \mathcal{C}_1 \vee \mathcal{C}_2$. Now assume $X \in \mathcal{C}_1 \vee \mathcal{C}_2$. Set Y = P(X) and $Y_1 = t_{P(\mathcal{C}_1)}(Y)$. Then $Y_2 = Y/Y_1$ is in $P(\mathcal{C}_2)$, and therefore $Y = Y_1 \oplus Y_2$ with $Y_i \in P(\mathcal{C}_i)$. Using Lemma 2.1 we find $X_i \in \mathcal{C}_i$ such that $X = X_1 \oplus X_2$. See also Proposition 4.1.

Lemma 3.13. Let (\mathcal{C}_{α}) be a family of essentially closed subcategories and suppose that each \mathcal{C}_{α} is closed under arbitrary products. Then $\bigwedge_{\alpha} \mathcal{C}_{\alpha}$ is closed under arbitrary products, and $\bigvee_{\alpha} \mathcal{C}_{\alpha}$ is closed under arbitrary products provided the family is finite.

Proof. For the lattice of essentially closed subcategories the meet is given by taking intersections, and it is clear that being closed under products is preserved when taking arbitrary intersections. Now suppose the family (\mathcal{C}_{α}) is finite. Observe that an essentially closed subcategory $\mathcal{C} \subseteq \mathcal{A}$ is closed under products if and only if $\mathcal{C} \cap \operatorname{Inj} \mathcal{A}$ is closed under products. For $\mathcal{C} = \bigvee_{\alpha} \mathcal{C}_{\alpha}$ we note that $\mathcal{C} \cap \operatorname{Inj} \mathcal{A}$ consists of all objects $\coprod_{\alpha} X_{\alpha}$ with $X_{\alpha} \in \mathcal{C}_{\alpha} \cap \operatorname{Inj} \mathcal{A}$ for all α , by Lemma 3.12. For an arbitrary family of objects (X_i) in $\mathcal{C} \cap \operatorname{Inj} \mathcal{A}$, the product

$$\prod_i X_i = \prod_i \left(\coprod_lpha X_{i,lpha}
ight) \cong \coprod_lpha \left(\prod_i X_{i,lpha}
ight)$$

belongs to \mathcal{C} since $\prod_i X_{i,\alpha}$ belongs to \mathcal{C}_{α} for all α .

Lemma 3.14. Let $\mathfrak{C} \subseteq \mathcal{A}$ be localising. Then every injective object $X \in \mathcal{A}$ admits a decomposition $X = X' \oplus X''$ such that $t_{\mathfrak{C}}(X) \to X'$ is an injective envelope and $t_{\mathfrak{C}}(X'') = 0$.

Proof. The inclusion $t_{\mathcal{C}}(X) \to X$ induces a monomorphism $E(t_{\mathcal{C}}(X)) \to X$. One takes for X' its image and for X'' its cokernel.

Lemma 3.15. Let $C \subseteq A$ be localising. Then the essential extensions of objects in C form an essentially closed subcategory $\overline{C} \subseteq A$, and the assignment

$$(\mathcal{C}, \mathcal{D}) \mapsto (P(\bar{\mathcal{C}}), P(\mathcal{D}))$$

yields a map between the hereditary torsion pairs for A and Spec A, respectively.

Proof. It is easily checked that the essential extensions of objects in \mathbb{C} form an essentially closed subcategory. Now let (\mathbb{C}, \mathbb{D}) be a hereditary torsion pair for A. From Lemmas 2.7 and 3.14 it follows that $(P(\bar{\mathbb{C}}), P(\mathbb{D}))$ is a hereditary torsion pair for Spec A.

Theorem 3.16. Let A be a Grothendieck category. There is a pair of maps

$$\mathbf{L}(\mathcal{A}) \xrightarrow{\sigma} \mathbf{L}(\operatorname{Spec} \mathcal{A})$$

given by $\sigma(\mathcal{C}) = \operatorname{supp}(\mathcal{C})$ and $\tau(\mathcal{U}) = {}^{\perp}(P^{-1}(\mathcal{U}^{\perp}))$ such that $\tau \sigma = \operatorname{id}$. Moreover, σ preserves arbitrary joins and finite meets.

Proof. First observe that τ is well defined, since $P^{-1}(\mathcal{U}^{\perp})$ is an essentially closed subcategory, and therefore $^{\perp}(P^{-1}(\mathcal{U}^{\perp}))$ is a localising subcategory of \mathcal{A} .

Let $\mathcal{C} \subseteq \mathcal{A}$ be localising and $(\mathcal{C}, \mathcal{D})$ the corresponding hereditary torsion pair. Then $(P(\bar{\mathcal{C}}), P(\mathcal{D}))$ is a hereditary torsion pair for Spec \mathcal{A} by Lemma 3.15, with $\mathcal{U} := \sigma(\mathcal{C}) = P(\bar{\mathcal{C}})$. Thus $\mathcal{U}^{\perp} = P(\mathcal{D})$ and then $P^{-1}(\mathcal{U}^{\perp}) = P^{-1}P(\mathcal{D}) = \mathcal{D}$. It follows that $\tau(\mathcal{U}) = \mathcal{C}$.

The assignment $\mathcal{C} \mapsto \mathcal{C}^{\perp}$ identifies the localising subcategories of \mathcal{A} with the essentially closed subcategories that are closed under products, by Lemma 3.11. This map reverses inclusions and we compose it with the isomorphism from Theorem 3.3. Then it follows from Lemma 3.13 that the image of σ is closed under arbitrary joins and finite meets.

I am grateful to Sira Gratz and Greg Stevenson for suggesting the following couple of remarks.

Given a frame L, its *points* are the frame morphisms $L \to \mathbf{2}$, where $\mathbf{2}$ denotes the frame with two elements $\{0 \le 1\}$. The frame L has *enough points* if for all $x \not\le y$ in L there exists a point $p: L \to \mathbf{2}$ such that p(x) = 1 and p(y) = 0.

Remark 3.17. There are examples showing that the frame $\mathbf{L}(\mathcal{A})$ may have no points [37]. This is in contrast to Proposition 2.8 which shows that the bigger frame $\mathbf{L}(\operatorname{Spec} \mathcal{A})$ has enough points.

Remark 3.18. Suppose that Spec \mathcal{A} is discrete. For example, let \mathcal{A} be locally noetherian. Then $\mathbf{L}(\mathcal{A})$ has enough points, so it is a spatial frame. This is easily seen, because $\mathbf{L}(\operatorname{Spec}\mathcal{A})$ identifies with the set of subsets of $\operatorname{Sp}\mathcal{A}$. For $I \in \operatorname{Sp}\mathcal{A}$ consider the point $p_I : \mathbf{L}(\mathcal{A}) \xrightarrow{\sigma} \mathbf{L}(\operatorname{Spec}\mathcal{A}) \xrightarrow{\pi_I} \mathbf{2}$, where $\pi_I(\mathcal{U}) = 1$ iff $I \in \mathcal{U}$. Given a pair of localising subcategories $\mathcal{C} \not\subseteq \mathcal{D}$ of \mathcal{A} , one chooses $I \in \operatorname{Sp}\mathcal{A} \cap (\langle \mathcal{C} \rangle \setminus \langle \mathcal{D} \rangle)$ and then $p_I(\mathcal{C}) = 1$ and $p_I(\mathcal{D}) = 0$.

4. Coproduct decompositions

Decompositions of objects in a Grothendieck have been studied in great detail in the past. For instance, the theorems of Azumaya, Krull, Remak, and Schmidt treat the decompositions into indecomposable objects with local endomorphism rings.³ For arbitrary objects the decomposition theory is far more complex. In this section we discuss a Boolean lattice which controls the decompositions, at least for injective objects. There is probably no efficient way to describe all decompositions. So we focus on decompositions up to a certain equivalence relation, which reflects the decompositions of essentially closed subcategories.

We fix a Grothendieck category \mathcal{A} and recall that $P: \mathcal{A} \to \operatorname{Spec} \mathcal{A}$ denotes the functor that inverts all essential monomorphisms.

Coproduct decompositions via essentially closed subcategories. A coproduct decomposition of an object in \mathcal{A} induces via the assignment $X \mapsto \langle X \rangle$ a decomposition in the lattice of essentially closed subcategories. This correspondence works in both directions as follows.

Proposition 4.1. Let A be a Grothendieck category and $X \in A$ an object.

- (1) A decomposition $X = \coprod_{\alpha} X_{\alpha}$ in A induces a decomposition $\langle X \rangle = \bigvee_{\alpha} \langle X_{\alpha} \rangle$ into essentially closed subcategories.
- (2) Suppose that X is injective and let $\langle X \rangle = \bigvee_{\alpha} \mathbb{C}_{\alpha}$ be a decomposition into essentially closed subcategories. Then there are direct summands $X_{\alpha} \subseteq X$ such that $X = E(\coprod_{\alpha} X_{\alpha})$ and $X_{\alpha} \in \mathbb{C}_{\alpha}$ for all α . Moreover, $\langle X_{\alpha} \rangle = \mathbb{C}_{\alpha}$ for all α provided that $\mathbb{C}_{\alpha} \wedge \mathbb{C}_{\beta} = 0$ for all $\alpha \neq \beta$.

Proof. (1) is clear from the definitions. For (2) we may assume that $\mathcal{C}_{\alpha} \wedge \mathcal{C}_{\beta} = 0$ for all $\alpha \neq \beta$ by replacing \mathcal{C}_{α} with a smaller subcategory if needed; see Lemma 2.13. Suppose first that \mathcal{A} is spectral. Then there is an essentially unique decomposition $X = \coprod_{\alpha} X_{\alpha}$ in \mathcal{A} with $\langle X_{\alpha} \rangle = \mathcal{C}_{\alpha}$ for all α , by choosing $X_{\alpha} = t_{\mathcal{C}_{\alpha}}(X)$. This follows from Lemma 2.2; see also Proposition 2.14. When \mathcal{A} is arbitrary and X is injective we obtain a decomposition of X by decomposing P(X) in Spec \mathcal{A} and applying Theorem 3.3. More precisely, for each α we apply Lemma 2.1 and choose a direct summand $X_{\alpha} \subseteq X$ such that $P(X_{\alpha}) = t_{P(\mathcal{C}_{\alpha})}(P(X))$. Then the equality $X = E(\coprod_{\alpha} X_{\alpha})$ follows from (2.2).

Remark 4.2. The identity $X = E(\coprod_{\alpha} X_{\alpha})$ in Proposition 4.1 simplifies to $X = \coprod_{\alpha} X_{\alpha}$ when the index set is finite or when \mathcal{A} is locally noetherian.

Remark 4.3. In part (2) of the above proposition, the assumption on X to be injective cannot be removed. Consider for instance an indecomposable object X of length 3 with $soc(X) = S_1 \oplus S_2$ such that S_1 and S_2 are non-isomorphic simple

³Splittings or decompositions have been considered since ancient times. The Quran reports about a miraculous splitting of the moon [Quran 54:1-2], and the Bible about the splitting of the Red Sea [Exodus 14:21-22].

objects. Then $\langle X \rangle = \langle S_1 \rangle \vee \langle S_2 \rangle$ and $\langle S_1 \rangle \wedge \langle S_2 \rangle = 0$, but there is no decomposition $X = X_1 \oplus X_2$ with $\langle X_i \rangle = \langle S_i \rangle$.

The lattice of decompositions. We introduce an equivalence relation on the direct summands of an injective object in order to provide an efficient description of all decompositions.

Definition 4.4. Let X be an injective object in \mathcal{A} . Given a pair of direct summands $U \subseteq X$ and $V \subseteq X$, we say they are *essentially equivalent* if $\langle U \rangle = \langle V \rangle$. Let $\mathbf{D}(X)$ denote the set of essential equivalence classes of direct summands of X, with partial order given by $U \leq V$ if $\langle U \rangle \subseteq \langle V \rangle$.

The following result establishes a connection to the decomposition theory for von Neumann regular rings since $\operatorname{End}(X)/J(\operatorname{End}(X))$ has this property. In particular, the elaborated theory of types applies [23, 29].

Theorem 4.5. Given an injective object X in A, the partially ordered set $\mathbf{D}(X)$ of direct summands is a complete Boolean lattice. Moreover, there is a canonical lattice isomorphism

$$\mathbf{D}(X) \xrightarrow{\sim} \mathbf{B}(\mathrm{End}(X)/J(\mathrm{End}(X))).$$

Proof. We claim that the assignment $U\mapsto \langle P(U)\rangle=P(\langle U\rangle)$ induces a lattice isomorphism

(4.1)
$$\mathbf{D}(X) \xrightarrow{\sim} \{ \mathcal{U} \in \mathbf{S}(\mathcal{A}) \mid \mathcal{U} \subseteq \operatorname{supp}(X) \}.$$

The inverse map sends a localising subcategory $\mathcal{U} \subseteq \operatorname{Spec} \mathcal{A}$ to the equivalence class of a direct summand $U \subseteq X$ such that $P(U) = t_{\mathcal{U}}(P(X))$; for the existence of U see Lemma 2.1. Using Theorem 3.3 it is easily checked that these assignments are mutually inverse to each other and preserving the partial order. Thus $\mathbf{D}(X)$ is a Boolean lattice, since $\mathbf{S}(\mathcal{A})$ has this property by Proposition 2.8. Moreover, the above argument shows that $P \colon \mathcal{A} \to \operatorname{Spec} \mathcal{A}$ induces a lattice isomorphism

$$\mathbf{D}(X) \xrightarrow{\sim} \mathbf{D}(P(X)).$$

For the isomorphism with the lattice of central idempotents we consider the Grothendieck category

$$A_X := \langle P(X) \rangle \subseteq \operatorname{Spec} A$$

which is generated by P(X). The functor P induces an isomorphism

$$\operatorname{End}(X)/J(\operatorname{End}(X)) \xrightarrow{\sim} \operatorname{End}(P(X))$$

by (2.1). This yields the following sequence of isomorphisms

$$\mathbf{D}(X) \xrightarrow{\sim} \mathbf{D}(P(X)) \xrightarrow{\sim} \mathbf{L}(\mathcal{A}_X) \xrightarrow{\sim} \mathbf{B}(\mathrm{End}(X)/J(\mathrm{End}(X))),$$

where the first is (4.2), the second is (4.1), and the last comes from Lemma 2.10. \Box

Remark 4.6. The Boolean structure of $\mathbf{D}(X)$ implies that each equivalence class contains a summand $U\subseteq X$ that is maximal in the sense that $\langle U\rangle\cap\langle X/U\rangle=0$. An equivalent condition is that for any pair of monomorphisms $U\leftarrow V\to X/U$ we have V=0.

5. Cohomologically stable subcategories

An essentially closed subcategory of a Grothendieck category is always closed under subobjects and extensions, but it need not be closed under quotients. For that reason we consider another class of subcategories which is also controlled by its spectral category. This leads to a notion of exact support for any Grothendieck category.

Exact support. Let \mathcal{A} be an exact category. Thus \mathcal{A} is an additive category together with a distinguished class of short exact sequences. Recall that a full subcategory $\mathcal{C} \subseteq \mathcal{A}$ is *thick* if \mathcal{C} is closed under direct summands and the *two out of three condition* holds: any short exact sequence belongs to \mathcal{C} if two of its three terms are in \mathcal{C} .

A map $\sigma \colon \operatorname{Ob} \mathcal{A} \to L$ with values in a join-semilattice L is called exact if

- (1) for all objects $X, Y \in \mathcal{A}$ one has $\sigma(X \oplus Y) = \sigma(X) \vee \sigma(Y)$, and
- (2) for each exact sequence $0 \to X_1 \to X_2 \to X_3 \to 0$ in \mathcal{A} one has

$$\sigma(X_i) \le \sigma(X_j) \lor \sigma(X_k)$$
 when $\{i, j, k\} = \{1, 2, 3\}.$

It is easily checked that σ is exact if and only if for each $u \in L$

$$\mathcal{A}_u := \{ X \in \mathcal{A} \mid \sigma(X) \le u \}$$

is a thick subcategory of A.

Now let A be a Grothendieck category. We consider again the Boolean lattice

$$\mathbf{S}(\mathcal{A}) := \mathbf{L}(\operatorname{Spec} \mathcal{A}) \xrightarrow{\sim} \operatorname{Clop}(\operatorname{Spc} \mathcal{A}).$$

Also, we consider the derived category $\mathbf{D}(A)$ and the right derived functor

$$\mathbf{R}P \colon \mathbf{D}(\mathcal{A}) \longrightarrow \mathbf{D}(\operatorname{Spec} \mathcal{A}).$$

For an object $X \in \mathcal{A}$ its exact support is by definition

(5.1)
$$\operatorname{supp}_{\operatorname{av}}(X) := \inf \{ \mathcal{U} \in \mathbf{S}(\mathcal{A}) \mid \mathbf{R}P(X) \in \mathbf{D}(\mathcal{U}) \} \in \mathbf{S}(\mathcal{A}),$$

where we identify

$$\mathbf{D}(\mathcal{U}) = \{ Y \in \mathbf{D}(\operatorname{Spec} \mathcal{A}) \mid H^n(Y) \in \mathcal{U} \text{ for all } n \in \mathbb{Z} \}.$$

For
$$\mathfrak{X} \subseteq \mathcal{A}$$
 we set $\operatorname{supp}_{\operatorname{ex}}(\mathfrak{X}) := \bigvee_{X \in \mathfrak{X}} \operatorname{supp}_{\operatorname{ex}}(X)$.

Lemma 5.1. Let $X \in \mathcal{A}$ be an object and $X \to I$ a minimal injective resolution. Then $\operatorname{supp}_{\operatorname{ex}}(X)$ equals the localising subcategory of $\operatorname{Spec} \mathcal{A}$ consisting of all direct summands of coproducts of objects in $\{P(I^n) \mid n \in \mathbb{Z}\}$.

Proof. The functor P sends each differential $I^n \to I^{n+1}$ to zero because they are radical morphisms; see [27, Proposition 4.3.18]. Thus $H^n(\mathbf{R}P(X)) \cong P(I^n)$ for each $n \in \mathbb{Z}$.

Let $\mathcal{U} \in \mathbf{S}(\mathcal{A})$. We set

$$\operatorname{Inj}_{\mathcal{U}} \mathcal{A} := \{ I \in \operatorname{Inj} \mathcal{A} \mid P(I) \in \mathcal{U} \}$$

and $\mathcal{A}_{\mathcal{U}}$ denotes the full subcategory of objects $X \in \mathcal{A}$ that admit an injective resolution $X \to I$ with $I^n \in \operatorname{Inj}_{\mathcal{U}} \mathcal{A}$ for all $n \in \mathbb{Z}$, so

$$\mathcal{A}_{\mathcal{U}} = \{ X \in \mathcal{A} \mid \mathbf{R}P(X) \in \mathbf{D}(\mathcal{U}) \} = \{ X \in \mathcal{A} \mid \mathrm{supp}_{\mathrm{ex}}(X) \subseteq \mathcal{U} \}.$$

Lemma 5.2. For each $\mathcal{U} \in \mathbf{S}(A)$ the subcategory $A_{\mathcal{U}}$ is thick, and the function

$$\operatorname{supp}_{\operatorname{ex}} \colon \operatorname{Ob} \mathcal{A} \longrightarrow \mathbf{S}(\mathcal{A})$$

is exact.

Proof. Each exact sequence $0 \to X \to Y \to Z \to 0$ in \mathcal{A} gives rise to an exact triangle $X \to Y \to Z \to X[1]$ in $\mathbf{D}(\mathcal{A})$. The right derived functor $\mathbf{R}P$ is exact and $\mathbf{D}(\mathcal{U})$ is a triangulated subcategory which is closed under direct summands. Then it follows that $\mathcal{A}_{\mathcal{U}}$ is thick. We have already seen that thickness of $\mathcal{A}_{\mathcal{U}}$ for each \mathcal{U} implies the exactness of the corresponding support function.

Cohomologically stable subcategories. The following definition is the derived analogue of Definition 3.1.

Definition 5.3. We call a thick subcategory $\mathcal{C} \subseteq \mathcal{A}$ cohomologically stable if

- (1) for every family (X_{α}) of objects in \mathcal{C} the injective envelope of $\coprod_{\alpha} X_{\alpha}$ belongs to \mathcal{C} , and
- (2) for every exact sequence $X^0 \to X^1 \to X^2 \to \cdots$ with all X^n in \mathcal{C} the kernel of $X^0 \to X^1$ belongs to \mathcal{C} .

The cohomologically stable subcategories are partially ordered by inclusion and closed under arbitrary intersections; so they form a complete lattice. When $\mathcal A$ is spectral, then its cohomologically stable subcategories are precisely its localising subcategories. For an object $X \in \mathcal A$ let $\langle X \rangle_{\mathrm{ex}}$ denote the smallest cohomologically stable subcategory of $\mathcal A$ that contains X.

Lemma 5.4. For $\mathcal{U} \in \mathbf{S}(A)$ the subcategory $A_{\mathcal{U}}$ is cohomologically stable.

Proof. The category $\mathcal{A}_{\mathcal{U}}$ is thick by Lemma 5.2. Let (X_{α}) be a family of objects in $\mathcal{A}_{\mathcal{U}}$ and choose injective envelopes $X_{\alpha} \to I_{\alpha}$. Then $I_{\alpha} \in \operatorname{Inj}_{\mathcal{U}} \mathcal{A}$ for all α , and therefore $E(\coprod_{\alpha} X_{\alpha}) \cong E(\coprod_{\alpha} I_{\alpha})$ belongs to $\operatorname{Inj}_{\mathcal{U}} \mathcal{A}$. Now let

$$X^0 \to X^1 \to X^2 \to \cdots$$

be exact with all X^n in $\mathcal{A}_{\mathcal{U}}$. We set $Z^n = \operatorname{Ker}(X^n \to X^{n+1})$ and need to show that Z^0 belongs to $\mathcal{A}_{\mathcal{U}}$. Choose an injective envelope $X^0 \to I^0$. The cokernel $C = \operatorname{Coker}(Z^0 \to I^0)$ fits into an exact sequence $0 \to Z^1 \to C \to C' \to 0$ with $C' = \operatorname{Coker}(X^0 \to I^0)$ in $\mathcal{A}_{\mathcal{U}}$. We form the pushout

and the object \bar{X}^1 belongs to $\mathcal{A}_{\mathcal{U}}$. Thus we obtain an exact sequence

$$I^0 \to \bar{X}^1 \to X^2 \to X^3 \to \cdots$$

with all terms in $\mathcal{A}_{\mathcal{U}}$ and $Z^0 = \operatorname{Ker}(I^0 \to \bar{X}^1)$. Proceeding by induction we obtain an exact sequence

$$I^0 \rightarrow I^1 \rightarrow I^2 \rightarrow I^3 \rightarrow \cdots$$

with all terms in $\operatorname{Inj}_{\mathcal{U}} \mathcal{A}$ and $Z^0 = \operatorname{Ker}(I^0 \to I^1)$. Thus Z^0 belongs to $\mathcal{A}_{\mathcal{U}}$.

Theorem 5.5. Let A be a Grothendieck category. The assignment $\mathcal{C} \mapsto \operatorname{supp}_{\operatorname{ex}}(\mathcal{C})$ induces a lattice isomorphism

$$\{\mathfrak{C} \subseteq \mathcal{A} \mid \mathfrak{C} \ cohomologically \ stable\} \xrightarrow{\sim} \mathbf{S}(\mathcal{A}).$$

The inverse map is given by $\mathcal{U} \mapsto \mathcal{A}_{\mathcal{U}}$.

Proof. Lemma 5.4 shows that the assignment $\mathcal{U} \mapsto \mathcal{A}_{\mathcal{U}}$ is well defined. Using the description of support in Lemma 5.1, it is clear that $\operatorname{supp}_{\operatorname{ex}}(\mathcal{A}_{\mathcal{U}}) = \mathcal{U}$. Now let $\mathcal{C} \subseteq \mathcal{A}$ be a cohomologically stable subcategory and set $\mathcal{U} = \operatorname{supp}_{\operatorname{ex}}(\mathcal{C})$. We claim that $\operatorname{Inj}_{\mathcal{U}} \mathcal{A} \subseteq \mathcal{C}$. When $X \to I$ is a minimal injective resolution of an object in $X \in \mathcal{C}$, then the definition of cohomologically stable implies that $I^n \in \mathcal{C}$ for all $n \in \mathbb{Z}$. Moreover, cohomologically stable implies that any object $I \in \operatorname{Inj} \mathcal{A}$ belongs to \mathcal{C} when P(I) arises as a direct summand of a coproduct of objects of the form $P(I^n)$. Here we use again that P preserves coproducts. This yields the claim, and the equality $\mathcal{A}_{\mathcal{U}} = \mathcal{C}$ then follows.

Remark 5.6. When \mathcal{A} is locally noetherian, then a cohomologically stable subcategory is closed under all coproducts. This follows from the fact that any coproduct of injectives is again injective. Moreover, in this case $\mathbf{S}(\mathcal{A})$ identifies with the lattice of subsets of $\operatorname{Sp} \mathcal{A}$; see Example 2.6.

Remark 5.7. In this generality Theorem 5.5 seems to be new. For some special cases it was already known that subcategories of \mathcal{A} are determined by subsets of $\operatorname{Sp} \mathcal{A}$.

- (1) When \mathcal{A} equals the category of modules over a commutative noetherian ring Λ , then $\mathbf{S}(\mathcal{A})$ identifies with the lattice of subsets of the prime ideal spectrum of Λ ; see Example 2.6. In this case the localising subcategories of \mathcal{A} are cohomologically stable and correspond to the specialisation closed subsets of Spec Λ . This goes back to work of Gabriel in [12]; it was extended by Takahashi to a correspondence for all cohomologically stable subcategories in [38] and more recently in [28], using Neeman's classification of localising subcategories of the derived category $\mathbf{D}(\mathcal{A})$ from [30]. A generalisation of Gabriel's correspondence for locally noetherian Grothen-dieck categories was established in [39].
- (2) For a locally coherent Grothendieck category \mathcal{A} , the localising subcategories of finite type are classified in terms of Sp \mathcal{A} ; see [19, 24]. This is somewhat orthogonal to Gabriel's correspondence. More precisely, a localising subcategory $\mathcal{C} \subseteq \mathcal{A}$ is of *finite type* when the corresponding subcategory of torsion-free objects

$$\mathcal{C}^{\perp} = \{ X \in \mathcal{A} \mid \text{Hom}(C, X) = 0 \text{ for all } C \in \mathcal{C} \}$$

is closed under filtered colimits. The subcategory \mathcal{C}^{\perp} is an essentially closed subcategory, and both $\mathcal{C}^{\perp} \cap \operatorname{Sp} \mathcal{A}$ and $P(\mathcal{C}^{\perp}) \in \mathbf{S}(\mathcal{A})$ determine \mathcal{C} , while \mathcal{C} need not be cohomologically stable; cf. Theorem 3.16.

Support data. The following is the analogue of Definition 3.6. The only difference appears in the last condition $(E \land)$ where the intersection is based on cohomologically stable subcategories, whereas in condition $(S \land)$ essentially closed subcategories are used.

Definition 5.8. An exact support datum on a Grothendieck category \mathcal{A} is a pair (T, σ) consisting of a topological space T and a map $\sigma \colon \mathrm{Ob}\,\mathcal{A} \to \mathrm{Clop}(T)$ such that

- (E0) $\sigma(X) = \emptyset$ for X = 0,
- (E1) $\sigma(X) = T$ for every generator X of \mathcal{A} ,
- $(E\vee)$ $\sigma(X)\cup\sigma(Y)=\sigma(X\oplus Y)$ for all objects X,Y in \mathcal{A} , and
- $(E \wedge) \ \sigma(X) \cap \sigma(Y) = \sigma(Z) \text{ for all objects } X, Y, Z \text{ in } \mathcal{A} \text{ with } \langle X \rangle_{\mathrm{ex}} \cap \langle Y \rangle_{\mathrm{ex}} = \langle Z \rangle_{\mathrm{ex}}.$

A morphism $(T, \sigma) \to (T', \sigma')$ is given by a continuous map $f: T \to T'$ such that $\sigma(X) = f^{-1}(\sigma'(X))$ for all $X \in \mathcal{A}$.

The following is the exact analogue of Theorem 3.8; it has the same proof.

Theorem 5.9. The pair ($\operatorname{Spc} A, \operatorname{supp}_{\operatorname{ex}}$) is an exact support datum on A. Moreover, for any exact support datum (T,σ) there is a unique continuous map $f:T\to \operatorname{Spc} A$ such that $\sigma(X)=f^{-1}(\operatorname{supp}_{\operatorname{ex}}(X))$ for all $X\in A$. Explicitly, the map f is given by the adjunction (2.3) and therefore

$$f(p) = \{ \langle X \rangle \in \mathbf{S}(\mathcal{A}) \mid p \notin \sigma(X) \} \quad \text{for } p \in T.$$

The computation of $\operatorname{supp}_{\operatorname{ex}}$ in Lemma 5.1 shows that for each object $X \in \mathcal{A}$ we have an inclusion

$$supp(X) \subseteq supp_{ex}(X)$$
.

There is an equality when A is spectral, but small examples show that it is a proper inclusion in general.

Example 5.10. Let $\mathcal{A} = \operatorname{Mod} \Lambda$ be the module category of $\Lambda = \begin{bmatrix} k & k \\ 0 & k \end{bmatrix}$ where k is a field. Then there are two indecomposable injective Λ -modules and $\operatorname{Sp} \mathcal{A} = \{I_1, I_2\} = \operatorname{Spc} \mathcal{A}$. There is a short exact sequence $0 \to S_1 \to I_1 \to I_2 \to 0$ where $S_1 = \operatorname{soc}(I_1)$, and therefore

$$supp(S_1) = \{I_1\}$$
 and $supp_{ex}(S_1) = \{I_1, I_2\}.$

Stable localising subcategories. Let \mathcal{A} be a Grothendieck category. A localising subcategory is called stable if it is closed under essential extensions, equivalently if it is closed under injective envelopes. The stable localising subcategories are partially ordered by inclusion and closed under all intersections; so they form a complete lattice.

Given an element $\mathcal{U} \in \mathbf{S}(\mathcal{A})$, we have always the inclusion $\mathcal{A}_{\mathcal{U}} \subseteq P^{-1}(\mathcal{U})$. So one may ask when the equality $\mathcal{A}_{\mathcal{U}} = P^{-1}(\mathcal{U})$ holds.

Lemma 5.11. For a full subcategory $\mathfrak{C} \subseteq \mathcal{A}$ the following are equivalent.

- (1) C is stable localising
- (2) C is essentially closed and cohomologically stable.
- (3) There exists $\mathcal{U} \in \mathbf{S}(\mathcal{A})$ such that $\mathcal{C} = \mathcal{A}_{\mathcal{U}} = P^{-1}(\mathcal{U})$.

Proof. If $\mathfrak{C} \subseteq \mathcal{A}$ is stable localising, then $\mathfrak{C} = P^{-1}(\mathfrak{U})$ for some $\mathfrak{U} \in \mathbf{S}(\mathcal{A})$. Also, we have $\mathfrak{C} = \mathcal{A}_{\mathfrak{U}}$ since \mathfrak{C} is cohomologically stable. On the other hand, when $\mathcal{A}_{\mathfrak{U}} = P^{-1}(\mathfrak{U})$, then this subcategory is clearly stable localising.

Proposition 5.12. Let A be a Grothendieck category. The assignment

$$\mathcal{C} \longmapsto \operatorname{supp}(\mathcal{C}) = P(\mathcal{C})$$

induces an embedding

$$\{\mathcal{C} \subseteq \mathcal{A} \mid \mathcal{C} \text{ stable localising}\} \hookrightarrow \mathbf{S}(\mathcal{A})$$

which preserves arbitrary joins and meets.

Proof. We apply Lemma 5.11. Each stable localising subcategory is essentially closed. Thus the embedding follows from Theorem 3.3. The stable localising subcategories of \mathcal{A} are closed under arbitrary intersections. So arbitrary meets are preserved. For a family $(\mathcal{C}_{\alpha})_{\alpha \in A}$ of stable localising subcategories with $\mathcal{U}_{\alpha} = P(\mathcal{C}_{\alpha})$ we consider the join $\mathcal{U} = \bigvee_{\alpha} \mathcal{U}_{\alpha}$ and need to show that $P^{-1}(\mathcal{U})$ is a stable localising subcategory. The subcategory is essentially closed. So it suffices to show that for any epimorphism $\phi \colon I \to X$ with I injective and $P(I) \in \mathcal{U}$ we have that $P(X) \in \mathcal{U}$. We begin with the finite case and may assume $A = \{\alpha, \beta\}$. Let $I = I_{\alpha} \oplus I_{\beta}$ with $I_{\alpha} \in \mathcal{C}_{\alpha}$ and $I_{\beta} \in \mathcal{C}_{\beta}$. This decomposition induces an exact sequence $0 \to X_{\alpha} \to X \to X_{\beta} \to 0$ where $X_{\alpha} = \phi(I_{\alpha})$ and X_{β} is a quotient of I_{β} . Since \mathcal{C}_{α} and \mathcal{C}_{β} are closed under quotients, it follows that $P(X) \in \mathcal{U}$. Writing A as directed union of its finite subsets, we may assume that $\mathcal{U} = \bigvee_{\alpha} \mathcal{U}_{\alpha}$ is directed. Then $I = \sum_{\alpha} I_{\alpha}$ is the directed union of injectives $I_{\alpha} \in \mathcal{C}_{\alpha}$ by Lemma 2.2. This implies $X = \sum_{\alpha} X_{\alpha}$, where $X_{\alpha} = \phi(I_{\alpha})$. We have $P(X) = \sum_{\alpha} P(X_{\alpha})$, and this belongs to \mathcal{U} since $P(X_{\alpha}) \in \mathcal{U}_{\alpha}$ for all α .

Corollary 5.13. The stable localising subcategories of a Grothendieck category form a frame and a coframe.

Proof. This follows from the above theorem since $\mathbf{S}(A)$ is a frame and a coframe, by Propositions 2.3 and 2.4.

Example 5.14. Let $\mathcal{A} = \operatorname{Mod} \Lambda$ be the module category of a ring Λ . When Λ is commutative noetherian, then all localising subcategories are stable. This fails when Λ is not commutative. For $\Lambda = \begin{bmatrix} k & k \\ 0 & k \end{bmatrix}$ and k a field, there are two proper localising subcategories and one of them is not stable.

Derived categories. For a Grothendieck category \mathcal{A} we have already considered the right derived functor of the canonical functor $P: \mathcal{A} \to \operatorname{Spec} \mathcal{A}$. The restriction $\operatorname{Inj} \mathcal{A} \to (\operatorname{Inj} \mathcal{A}) / \operatorname{Rad}(\operatorname{Inj} \mathcal{A}) \xrightarrow{\sim} \operatorname{Spec} \mathcal{A}$ induces an exact functor

$$\mathbf{K}(\operatorname{Inj} \mathcal{A}) \longrightarrow \mathbf{K}(\operatorname{Spec} \mathcal{A}) \xrightarrow{\sim} \mathbf{D}(\operatorname{Spec} \mathcal{A})$$

where $\mathbf{K}(\operatorname{Inj} \mathcal{A})$ denotes the category of complexes in $\operatorname{Inj} \mathcal{A}$ up to homotopy. Its computation for any $X \in \mathbf{K}(\operatorname{Inj} \mathcal{A})$ is very explicit. By removing non-zero contractible summands we may assume that the complex X is homotopically minimal so that all differentials are radical morphisms; see [27, Proposition 4.3.18]. Then P(X) has zero differentials.

For example, when \mathcal{A} is locally noetherian the triangulated category $\mathbf{K}(\operatorname{Inj}\mathcal{A})$ is compactly generated with category of compacts equivalent to $\mathbf{D}^{\mathrm{b}}(\operatorname{Noeth}\mathcal{A})$, where $\operatorname{Noeth}\mathcal{A}$ denotes the full subcategory of noetherian objects in \mathcal{A} ; see [26]. Moreover, in this case the functor $\mathbf{K}(\operatorname{Inj}\mathcal{A}) \to \mathbf{D}(\operatorname{Spec}\mathcal{A})$ preserves all coproducts.

Given a triangulated category \mathcal{T} with coproducts, let $\mathbf{L}(\mathcal{T})$ denote the lattice of localising subcategories. The inclusion $i \colon \operatorname{Spec} \mathcal{A} \to \mathbf{D}(\operatorname{Spec} \mathcal{A})$ induces a lattice isomorphism

$$\mathbf{L}(\mathbf{D}(\operatorname{Spec} \mathcal{A})) \xrightarrow{\sim} \mathbf{L}(\operatorname{Spec} \mathcal{A})$$

by taking $\mathcal{U} \subseteq \mathbf{D}(\operatorname{Spec} \mathcal{A})$ to $i^{-1}(\mathcal{U})$. The inverse map sends $\mathcal{V} \subseteq \operatorname{Spec} \mathcal{A}$ to $\mathbf{D}(\mathcal{V})$ where we identify

$$\mathbf{D}(\mathcal{V}) = \{ X \in \mathbf{D}(\operatorname{Spec} \mathcal{A}) \mid H^n(X) \in \mathcal{V} \text{ for all } n \in \mathbb{Z} \}.$$

This yields a notion of support for objects $X \in \mathbf{K}(\operatorname{Inj} \mathcal{A})$ with $\operatorname{supp}(X) \subseteq \operatorname{Spc} \mathcal{A}$, since $\mathbf{L}(\operatorname{Spec} \mathcal{A}) \xrightarrow{\sim} \operatorname{Clop}(\operatorname{Spc} \mathcal{A})$. Details and properties of this are left to the interested reader.

6. Exactly definable categories

There are interesting classes of rings where the Boolean spectrum of its module category is a finite set with the discrete topology. For instance this includes all artinian rings. In such a case it is more interesting to study the pure-injective modules and the corresponding pure spectral category, as the class of injective modules is rather small. The notion of purity for modules goes back to Cohn [5] and was later generalised by Crawley-Boevey in the context of locally finitely presented additive categories [6]. A key idea is to identify the relevant additive category $\mathcal C$ with the full subcategory of fp-injective objects in a much larger locally coherent Grothendieck category $\mathbf P(\mathcal C)$, the purity category of $\mathcal C$. Following [25] such categories $\mathcal C$ are called *exactly definable*, because they identify with the category of exact functors $\mathcal A \to \mathbf A \mathbf b$ from a small abelian category $\mathcal A$ to the category of abelian groups.

Exactly definable categories. Let \mathcal{A} be a Grothendieck category that is *locally coherent*. Thus \mathcal{A} has a generating set of finitely presented objects and the full subcategory $\operatorname{fp} \mathcal{A}$ of finitely presented objects is abelian. An object $X \in \mathcal{A}$ is *fp-injective* if $\operatorname{Ext}^1(F,X) = 0$ for all $F \in \operatorname{fp} \mathcal{A}$. Let $\mathcal{C} := \operatorname{Fpinj} \mathcal{A}$ denote the full subcategory of fp-injective objects in \mathcal{A} . This is an exact subcategory of \mathcal{A} that is closed under filtered colimits and products. In fact, the exact structure of \mathcal{C} is intrinsic, because it is the smallest exact structure on \mathcal{C} such that the exact

sequences are closed under filtered colimits.⁴ The exact sequences in \mathcal{C} are called *pure-exact*, and \mathcal{C} admits pure-injective envelopes which identify with the injective envelopes in the ambient category \mathcal{A} .

Definition 6.1. A category that is equivalent to one of the form Fpinj \mathcal{A} for a locally coherent category \mathcal{A} is called *exactly definable*; it has filtered colimits, products and an intrinsic exact structure given by the pure-exact sequences.

Let \mathcal{C} be an exactly definable category, given by a locally coherent category \mathcal{A} . Then \mathcal{A} is uniquely determined by \mathcal{C} , because the category of pure-injective objects in \mathcal{C} identifies with the category of injective objects in \mathcal{A} , and any abelian category with enough injective objects is uniquely determined by its category of injective objects. We write $\mathbf{P}(\mathcal{C})$ for \mathcal{A} and call it the *purity category* of \mathcal{C} .

Example 6.2. Let Λ be a ring and let $\operatorname{mod} \Lambda$ denote its category of finitely presented modules. The category $(\operatorname{mod}(\Lambda^{\operatorname{op}}), \operatorname{Ab})$ of additive functors $\operatorname{mod}(\Lambda^{\operatorname{op}}) \to \operatorname{Ab}$ is a locally coherent Grothendieck category, and the assignment

$$X \longmapsto X \otimes_{\Lambda} -|_{\operatorname{mod}(\Lambda^{\operatorname{op}})}$$

identifies $\operatorname{Mod} \Lambda$ with the full subcategory of fp-injective objects in $(\operatorname{mod}(\Lambda^{\operatorname{op}}), \operatorname{Ab})$. Thus $\operatorname{Mod} \Lambda$ is an exactly definable category and

$$\mathbf{P}(\operatorname{Mod}\Lambda) = (\operatorname{mod}(\Lambda^{\operatorname{op}}), \operatorname{Ab}).$$

This goes back to Gruson and Jensen [18]; see also [6, §3] or [27, 12.4].

Definable subcategories. Fix an exactly definable category \mathcal{C} , given by a locally coherent category \mathcal{A} . Following Crawley-Boevey [7] a full subcategory $\mathcal{D} \subseteq \mathcal{C}$ is called *definable* if it is closed under filtered colimits, products, and pure subobjects. The terminology is justified by the fact that definable subcategories of \mathcal{C} correspond to Serre subcategories of fp \mathcal{A} : we have

$$\mathcal{D} = \{ X \in \mathcal{C} \mid \text{Hom}(F, X) = 0 \text{ for all } F \in \mathcal{S} \}$$

for some Serre subcategory $S \subseteq \text{fp} \mathcal{A}$; see [27, 12.2] for details. From the above description it follows that \mathcal{D} is a thick subcategory. Moreover, \mathcal{D} is closed under pure-injective envelopes; see [27, Proposition 12.2.8]. We generalise the definition of a definable subcategory in two directions, following the Definitions 3.1 and 5.3.

Definition 6.3. A full subcategory $\mathcal{D} \subseteq \mathcal{C}$ is called *pure-essentially closed* if it is closed under arbitrary coproducts, pure subobjects, and pure-injective envelopes.

Definition 6.4. A thick subcategory $\mathcal{D} \subseteq \mathcal{C}$ is called *pure-cohomologically stable* if

- (1) for every family (X_{α}) of objects in \mathcal{D} the pure-injective envelope of $\coprod_{\alpha} X_{\alpha}$ belongs to \mathcal{D} , and
- (2) for every pure-exact sequence $X^0 \to X^1 \to X^2 \to \cdots$ with all X^n in \mathcal{D} the kernel of $X^0 \to X^1$ belongs to \mathcal{D} .

For an exact category we recall that a sequence of composable morphisms (ϕ^n) is exact if each ϕ^n is the composite $\phi^n = \iota_n \pi_n$ of an admissible epimorphism (deflation) followed by an admissible monomorphism (inflation) so that each pair (ι_n, π_{n+1}) is a conflation.

 $^{^4}$ I am grateful to Kevin Schlegel for suggesting the argument: For any object $X \in \mathcal{C}$ there is a reduced power $\bar{X} = \prod_I X/\mathcal{F}$ (given by an appropriate ultrafilter \mathcal{F} on some sufficiently large set I) which is injective in \mathcal{A} ; see [32, 4.2.5]. The canonical morphism $X \to \bar{X}$ is by construction a filtered colimit of split monos, and it factors through any mono $X \to Y$ since \bar{X} is injective. Thus any mono $X \to Y$ is an admissible mono with respect to any exact structure that is closed under filtered colimits.

Lemma 6.5. Any definable subcategory of an exactly definable category is pureessentially closed and pure-cohomologically stable.

Proof. This follows from the defining properties of a definable subcategory and the additional properties that have already been mentioned. Specifically, one uses that a definable subcategory is thick and closed under pure-injective envelopes. \Box

The pure-essentially closed and the pure-cohomologically stable subcategories are partially ordered by inclusion and closed under arbitrary intersections; so they form complete lattices.

The Ziegler topology. The class of pure-injective modules has been studied in seminal work of Ziegler on the model theory of modules [40]. In particular, he introduced for any ring a topology on the set of isoclasses of indecomposable pure-injective modules.

Let Ind \mathcal{C} denote the set of isoclasses of indecomposable pure-injective objects in \mathcal{C} ; this identifies with Sp $\mathbf{P}(\mathcal{C})$. The assignment

$$(6.1) \mathcal{D} \longmapsto \mathcal{D} \cap \operatorname{Ind} \mathfrak{C}$$

identifies the definable subcategories of \mathcal{C} with the Ziegler closed subsets of Ind \mathcal{C} . The inverse map sends a Ziegler closed subset \mathcal{U} to the definable subcategory generated by \mathcal{U} , which consists of all pure subobjects of products of objects in \mathcal{U} . This correspondence is due to Crawley-Boevey for module categories [7, 2.5] and we refer to [27, 12.2] for the general version.

The Boolean spectrum. Let PInj \mathcal{C} denote the full subcategory of pure-injective objects in \mathcal{C} ; it identifies with Inj $\mathbf{P}(\mathcal{C})$ via the embedding $\mathcal{C} \to \mathbf{P}(\mathcal{C})$. The *pure spectral category* is the quotient

$$PSpec \mathcal{C} := (PInj \mathcal{C}) / Rad(PInj \mathcal{C}) \xrightarrow{\sim} Spec \mathbf{P}(\mathcal{C})$$

which is a spectral Grothendieck category because of the equivalence from (2.1). We consider the lattice of localising subcategories

$$\mathbf{L}(\operatorname{PSpec} \mathfrak{C}) \xrightarrow{\sim} \mathbf{S}(\mathbf{P}(\mathfrak{C})).$$

This is a Boolean lattice by Proposition 2.8 and its spectrum

$$PSpc \mathcal{C} := Spec(\mathbf{L}(PSpec \mathcal{C}))$$

can be written as the disjoint union of a discrete and a continuous part as in (2.5). This yields a canonical embedding Ind $\mathcal{C} \hookrightarrow \operatorname{PSpc} \mathcal{C}$ which identifies Ind \mathcal{C} with the discrete part. We endow Ind \mathcal{C} with the Ziegler topology and write $\operatorname{Cl}(\operatorname{Ind} \mathcal{C})$ for its lattice of closed subsets.

In the following we provide two possible extensions of Crawley-Boevey's correspondence (6.1) for definable subcategories, which amounts to a lattice isomorphism

$$\{\mathcal{D} \subseteq \mathcal{C} \mid \mathcal{D} \text{ definable}\} \xrightarrow{\sim} \text{Cl}(\text{Ind }\mathcal{C}).$$

We view \mathcal{C} as a full subcategory of the purity category $\mathbf{P}(\mathcal{C})$ so that $\mathrm{supp}(X)$ and $\mathrm{supp}_{\mathrm{ex}}(X)$ for objects $X \in \mathcal{C}$ are defined via (3.1) and (5.1). Moreover, these are viewed as subsets of PSpc \mathcal{C} via the isomorphism

(6.2)
$$\mathbf{S}(\mathbf{P}(\mathcal{C})) \xrightarrow{\sim} \mathbf{L}(\mathrm{PSpec}\,\mathcal{C}) \xrightarrow{(2.4)} \mathrm{Clop}(\mathrm{PSpc}\,\mathcal{C}).$$

Theorem 6.6. Let \mathcal{C} be an exactly definable category. The assignment $\mathcal{D} \mapsto \text{supp}(\mathcal{D})$ induces a lattice isomorphism

$$\{\mathcal{D} \subseteq \mathcal{C} \mid \mathcal{D} \text{ pure-essentially closed}\} \xrightarrow{\sim} \operatorname{Clop}(\operatorname{PSpc} \mathcal{C}).$$

Moreover, the embedding $\operatorname{Ind} \mathcal{C} \hookrightarrow \operatorname{PSpc} \mathcal{C}$ identifies $\mathcal{D} \cap \operatorname{Ind} \mathcal{C}$ with $\operatorname{supp}(\mathcal{D}) \cap \operatorname{Ind} \mathcal{C}$ when $\mathcal{D} \subseteq \mathcal{C}$ is definable.

Proof. We view $\mathfrak{C} \subseteq \mathbf{P}(\mathfrak{C})$ as a full subcategory. Taking an essentially closed subcategory $\mathfrak{U} \subseteq \mathbf{P}(\mathfrak{C})$ to $\mathfrak{U} \cap \mathfrak{C}$ induces a bijection between the essentially closed subcategories of $\mathbf{P}(\mathfrak{C})$ and the pure-essentially closed subcategories of \mathfrak{C} . This follows from the fact that any essentially closed subcategory $\mathfrak{U} \subseteq \mathbf{P}(\mathfrak{C})$ is determined by $\mathfrak{U} \cap \operatorname{Inj} \mathbf{P}(\mathfrak{C})$, keeping in mind that $\operatorname{PInj} \mathfrak{C} = \operatorname{Inj} \mathbf{P}(\mathfrak{C})$. Given this correspondence the isomorphism is obtained by composing the isomorphisms from Theorem 3.3 and (6.2). The additional assertion about definable subcategories follows by unraveling the construction of the embedding $\operatorname{Ind} \mathfrak{C} \hookrightarrow \operatorname{PSpc} \mathfrak{C}$.

Theorem 6.7. Let \mathfrak{C} be an exactly definable category. The assignment $\mathfrak{D} \mapsto \sup_{\mathbf{e}_{\mathbf{x}}}(\mathfrak{D})$ induces a lattice isomorphism

$$\{\mathcal{D} \subseteq \mathcal{C} \mid \mathcal{D} \text{ pure-cohomologically stable}\} \xrightarrow{\sim} \operatorname{Clop}(\operatorname{PSpc} \mathcal{C}).$$

Moreover, the embedding Ind $\mathcal{C} \hookrightarrow \operatorname{PSpc} \mathcal{C}$ identifies $\mathcal{D} \cap \operatorname{Ind} \mathcal{C}$ with $\operatorname{supp}_{\operatorname{ex}}(\mathcal{D}) \cap \operatorname{Ind} \mathcal{C}$ when $\mathcal{D} \subseteq \mathcal{C}$ is definable.

Proof. Adapt the proof of Theorem 6.6 by using Theorem 5.5 instead of Theorem 3.3.

Remark 6.8. For every definable subcategory $\mathcal{D} \subseteq \mathcal{C}$ we have $\operatorname{supp}(\mathcal{D}) = \operatorname{supp}_{\operatorname{ex}}(\mathcal{D})$.

Module categories. Let Λ be a ring and consider its module category $\mathcal{C} = \operatorname{Mod} \Lambda$ which is exactly definable. We use the obvious notation and set $\operatorname{PInj} \Lambda = \operatorname{PInj} \mathcal{C}$, $\operatorname{Ind} \Lambda = \operatorname{Ind} \mathcal{C}$, and $\operatorname{PSpc} \Lambda = \operatorname{PSpc} \mathcal{C}$. The books of Prest [31] and also Jensen and Lenzing [20] discuss the basic structure of the *pure spectral category*

$$\operatorname{PSpec} \Lambda = (\operatorname{PInj} \Lambda) / \operatorname{Rad}(\operatorname{PInj} \Lambda).$$

For instance, it is shown that any pure-injective module X admits an essentially unique decomposition $X = X_d \oplus X_c$ into a *discrete* and a *continuous* part [20, Corollary 8.28]. This reflects the decomposition

$$PSpec \Lambda = (PSpec \Lambda)_d \times (PSpec \Lambda)_c$$

of the spectral category. For a more detailed treatment of decompositions of pure-injective modules we refer to [11]. Unfortunately, not much seems to be known about the continuous part of the spectral category, except that one knows for various classes of rings when the continuous part vanishes. On the other hand, superdecomposable modules have been constructed more or less explicitly for several rings, but these constructions do not provide much insight into the ways they decompose; see [34] for a survey. In any case, the discussion in §4 explains how the Boolean lattice $\mathbf{L}(\mathrm{PSpec}\,\Lambda)$ controls the decompositions of pure-injective Λ -modules; see also Example 2.12.

We characterise the absence of superdecomposable modules and may ask which Boolean lattices arise from representations of finite dimensional algebras, following ideas from [4].

Proposition 6.9. We have $X = X_d$ for every pure-injective Λ -module X if and only if the canonical embedding $\operatorname{Ind} \Lambda \hookrightarrow \operatorname{PSpc} \Lambda$ is a bijection.

Proof. This follows from the above decomposition of the pure spectral category PSpec Λ and the corresponding decomposition of the Boolean spectrum (2.5). \square

Problem 6.10. Let Λ be a finite dimensional algebra of strictly wild representation type and L a complete Boolean lattice. Is there a pure-injective Λ -module X such that its lattice of direct summands $\mathbf{D}(X)$ is isomorphic to L?

The following result complements our extension of Crawley-Boevey's correspondence for definable subcategories.

Proposition 6.11. For an Artin algebra Λ the following are equivalent.

- (1) The embedding $Cl(\operatorname{Ind}\Lambda) \to Clop(\operatorname{PSpc}\Lambda)$ is a bijection.
- (2) Every pure-essentially closed subcategory of Mod Λ is definable.
- (3) Every pure-cohomologically stable subcategory of Mod Λ is definable.
- (4) The Ziegler spectrum Ind Λ is a discrete space.
- (5) The algebra Λ is of finite representation type.

Proof. (1) \Leftrightarrow (2): This follows from Theorem 6.6.

- $(1) \Leftrightarrow (3)$: This follows from Theorem 6.7.
- $(2) \Rightarrow (4)$: Every subset of PSpc Λ contained in Ind Λ is clopen, by Lemma 2.5. It follows from Theorem 6.6 that every subset of Ind Λ is Ziegler closed.
- $(4) \Rightarrow (5)$: The space Ind Λ is finite when it is discrete, because Ind Λ is quasi-compact. For an Artin algebra, every indecomposable finitely presented module is pure-injective, so Λ is of finite representation type.
- $(5) \Rightarrow (2)$: When Λ is of finite representation type, then every Λ -module is endofinite, and in particular pure-injective [27, Theorem 13.2.10]. Let \mathcal{D} be a pure-essentially closed subcategory. Then it follows that \mathcal{D} is closed under pure subobjects and pure quotients. In particular, \mathcal{D} is closed under filtered colimits because any filtered colimit of objects in \mathcal{D} is a pure quotient of a coproduct of objects in \mathcal{D} . Finite representation type implies that every Λ -module decomposes into a coproduct of indecomposables, and every product of objects in \mathcal{D} decomposes into indecomposables from \mathcal{D} , because only finitely many indecomposables are involved. This follows again from endofiniteness. Thus \mathcal{D} is closed under products. \square

We close our discussion of module categories with some remarks.

Remark 6.12. Beyond finite representation type the correspondences for pure-essentially closed and pure-cohomologically stable subcategories (Theorems 6.6 and 6.7) seem to be new. In particular, Proposition 6.11 shows that the extension of the Ziegler spectrum yields additional information whenever the ring is not of finite representation type. This is illustrated by looking at the category of abelian groups $Ab = \text{Mod } \mathbb{Z}$. Consider the Prüfer group $\mathbb{Z}(p^{\infty})$ given by a prime p. The direct sums of copies of $\mathbb{Z}(p^{\infty})$ form a subcategory of Ab which is pure-essentially closed and pure-cohomologically stable, but not definable, because it is not closed under products. More precisely, the rationals \mathbb{Q} arise as direct summand of a product of copies of $\mathbb{Z}(p^{\infty})$, and then $\{\mathbb{Z}(p^{\infty}), \mathbb{Q}\}$ is the Ziegler closure of $\mathbb{Z}(p^{\infty})$.

The Ziegler spectrum plus all definable subcategories are well understood for various classes of rings, including the rings of finite representation type, commutative Dedekind domains, or tame hereditary algebras; see [32] for a survey. For these rings it seems reasonable to expect similar descriptions for the pure-essentially closed and the pure-cohomologically stable subcategories. Note that in these examples Ind $\Lambda = \operatorname{PSpc} \Lambda$; so the relevant subcategories are parameterised by all subsets of Ind Λ .

Acknowledgements. It is a pleasure to thank Greg Stevenson for several discussions and helpful comments on this work. Also, I am grateful to Ken Goodearl and Jan Šťovíček for their valuable input. The work was supported by the Deutsche Forschungsgemeinschaft (Project-ID 491392403 – TRR 358).

References

- P. Balmer, The spectrum of prime ideals in tensor triangulated categories, J. Reine Angew. Math. 588 (2005), 149–168.
- [2] F. Borceux and G. M. Kelly, On locales of localizations, J. Pure Appl. Algebra 46 (1987), no. 1, 1–34.
- [3] M. Brandenburg, Rosenberg's reconstruction theorem, Expo. Math. 36 (2018), no. 1, 98–117.

- [4] S. Brenner, Decomposition properties of some small diagrams of modules, in Symposia Mathematica, Vol. XIII (Convegno di Gruppi Abeliani & Convegno di Gruppi e loro Rappresentazioni, INDAM, Rome, 1972), 127–141, Academic Press, London, 1974.
- [5] P. M. Cohn, On the free product of associative rings, Math. Z. 71 (1959), 380–398.
- [6] W. Crawley-Boevey, Locally finitely presented additive categories, Comm. Algebra 22 (1994), no. 5, 1641–1674.
- [7] W. Crawley-Boevey, Infinite-dimensional modules in the representation theory of finite-dimensional algebras, in Algebras and modules, I (Trondheim, 1996), 29–54, CMS Conf. Proc., 23, Amer. Math. Soc., Providence, RI, 1998.
- [8] J. Dauns, Module types, Rocky Mountain J. Math. 27 (1997), no. 2, 503–557.
- [9] M. A. Dickmann, N. Schwartz and M. Tressl, Spectral spaces, New Mathematical Monographs, 35, Cambridge Univ. Press, Cambridge, 2019.
- [10] A. Facchini, Spectral categories and varieties of preadditive categories, J. Pure Appl. Algebra 29 (1983), no. 3, 219–239.
- [11] A. Facchini, Decompositions of algebraically compact modules, Pacific J. Math. 116 (1985), no. 1, 25–37.
- [12] P. Gabriel, Des catégories abéliennes, Bull. Soc. Math. France 90 (1962), 323-448.
- [13] P. Gabriel and U. Oberst, Spektralkategorien und reguläre Ringe im von-Neumannschen Sinn, Math. Z. 92 (1966), 389–395.
- [14] J. S. Golan, Torsion theories, Pitman Monographs and Surveys in Pure and Applied Mathematics, 29, Longman Sci. Tech., Harlow, 1986.
- [15] K. R. Goodearl, von Neumann regular rings, Monographs and Studies in Mathematics, 4, Pitman, Boston, MA, 1979.
- [16] K. R. Goodearl and A. K. Boyle, Dimension theory for nonsingular injective modules, Mem. Amer. Math. Soc. 7 (1976), no. 177, viii+112 pp.
- [17] K. R. Goodearl and F. Wehrung, The complete dimension theory of partially ordered systems with equivalence and orthogonality, Mem. Amer. Math. Soc. 176 (2005), no. 831, vii+117 pp.
- [18] L. Gruson and C. U. Jensen, Dimensions cohomologiques reliées aux foncteurs \(\begin{array}{lim}\)(i), in Paul Dubreil and Marie-Paule Malliavin Algebra Seminar, 33rd Year (Paris, 1980), 234–294, Lecture Notes in Math., 867, Springer, Berlin, 1981.
- [19] I. Herzog, The Ziegler spectrum of a locally coherent Grothendieck category, Proc. London Math. Soc. (3) 74 (1997), no. 3, 503–558.
- [20] C. U. Jensen and H. Lenzing, Model-theoretic algebra with particular emphasis on fields, rings, modules, Algebra, Logic and Applications, 2, Gordon and Breach, New York, 1989.
- [21] P. T. Johnstone, Stone spaces, Cambridge Stud. Adv. Math., vol. 3, Cambridge Univ. Press, Cambridge, 1982.
- [22] R. Kanda, Classification of categorical subspaces of locally Noetherian schemes, Doc. Math. 20 (2015), 1403–1465.
- [23] I. Kaplansky, Projections in Banach algebras, Ann. of Math. (2) 53 (1951), 235-249.
- [24] H. Krause, The spectrum of a locally coherent category, J. Pure Appl. Algebra 114 (1997), no. 3, 259–271.
- [25] H. Krause, Exactly definable categories, J. Algebra 201 (1998), no. 2, 456-492.
- [26] H. Krause, The stable derived category of a Noetherian scheme, Compos. Math. 141 (2005), no. 5, 1128–1162.
- [27] H. Krause, Homological theory of representations, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 195, Cambridge, 2022.
- [28] H. Matsui and R. Takahashi, Filtrations in module categories, derived categories, and prime spectra, Int. Math. Res. Not. IMRN 2022, no. 5, 3457–3492.
- [29] F. J. Murray and J. von Neumann, On rings of operators, Ann. of Math. (2) 37 (1936), no. 1, 116–229.
- [30] A. Neeman, The chromatic tower for D(R), Topology **31** (1992), no. 3, 519–532.
- [31] M. Prest, Model theory and modules, London Mathematical Society Lecture Note Series, 130, Cambridge Univ. Press, Cambridge, 1988.
- [32] M. Prest, Purity, spectra and localisation, Encyclopedia of Mathematics and its Applications, 121, Cambridge Univ. Press, Cambridge, 2009.
- [33] J.-E. Roos, Locally distributive spectral categories and strongly regular rings, in Reports of the Midwest Category Seminar, 156–181, Lecture Notes in Math., No. 47, Springer, Berlin, 1967.
- [34] C. M. Ringel, Infinite length modules. Some examples as introduction, in *Infinite length modules (Bielefeld, 1998)*, 1–73, Trends Math, Birkhäuser, Basel, 2000.
- [35] A. L. Rosenberg, Noncommutative algebraic geometry and representations of quantized algebras, Mathematics and its Applications, 330, Kluwer Acad. Publ., Dordrecht, 1995.

- [36] B. T. Stenström, Rings of quotients, Die Grundlehren der mathematischen Wissenschaften, Band 217, Springer, New York, 1975.
- $[37] \ \ {\rm G.\ Stevenson,\ Localizing\ subcategories\ of\ Grothendieck\ categories,\ Preprint,\ 2024.}$
- [38] R. Takahashi, On localizing subcategories of derived categories, J. Math. Kyoto Univ. 49 (2009), no. 4, 771–783.
- [39] K. Wu and X. Ma, A classification of some thick subcategories in locally noetherian Grothendieck categories, Glasg. Math. J. 66 (2024), no. 1, 175–182.
- [40] M. Ziegler, Model theory of modules, Ann. Pure Appl. Logic 26 (1984), no. 2, 149–213.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, D-33501 BIELEFELD, GERMANY $Email\ address$: hkrause@math.uni-bielefeld.de