Broadcasting in Heterogeneous Tree Networks with Edge Weight Uncertainty

Cheng-Hsiao Tsou* Ching-Chi Lin[†] Gen-Huey Chen*

Abstract

A broadcasting problem in heterogeneous tree networks with edge weight uncertainty under the postal model is considered in this paper. The broadcasting problem asks for a minmax-regret broadcast center, which minimizes the worst-case loss in the objective function. Due to the presence of edge weight uncertainty, it is not easy to attack the broadcasting problem. An $O(n \log n \log \log n)$ -time algorithm is proposed for solving the broadcasting problem.

Keywords: algorithm, broadcasting, edge weight uncertainty, heterogeneous tree network, minmax-regret, optimization problem.

^{*}Department of Computer Science and Information Engineering, National Taiwan University, Taipei 10617, Taiwan. Email: {f97922063, ghchen}@csie.ntu.edu.tw.

[†]Department of Computer Science and Engineering, National Taiwan Ocean University, Keelung 20224, Taiwan. Corresponding author. Email: lincc@mail.ntou.edu.tw.

1 Introduction

We consider the broadcasting problem in heterogeneous tree networks T = (V(T), E(T)) with edge weight uncertainty, where the edge weight $w_{u,v}$ can take any value from $[w_{u,v}^-, w_{u,v}^+]$ with unknown distribution. A heterogeneous network interconnects computers and other devices that can employ diverse operating systems and communication protocols. Representing such a network as a graph denoted as G proves advantageous, wherein the set of network nodes corresponds to V(G), representing the vertices in G, while the set of communication links corresponds to E(G), signifying the edges within the graph. For each link $(u, v) \in E(G)$, there exists a positive weight $w_{u,v}$ that signifies the time required to transmit a message across the link.

In response to the exact communication service requirements of a network, a variety of well-suited communication models have been devised [20, 21, 32, 33]. Among these communication models, the postal model, as explored in [9–11], is well-suited for characterizing packet-switching techniques. In the postal model, the process involves a sender u connecting with a receiver v and subsequently transmitting a message to v. A sender u is not permitted to connect with two or more receivers simultaneously. However, u can connect with a receiver, while transmitting a message to another receiver. Throughout this paper, we use ρ to denote the time consumed for link connection, where ρ is a positive constant. Let's consider a scenario where a sender u intends to broadcast a message to its k neighbors v_1, v_2, \ldots, v_k . Node v_1 will establish a connection with u at time unit ρ and receive the message at time unit $\rho + w_{u,v_1}$, ν_2 will connect with ν at time unit ν and receive the message at time unit ν and finally, ν will connect with ν at time unit ν and receive the message at time unit ν and finally, ν will connect with ν at time unit ν and receive the message at time unit ν and finally, ν will connect with ν at time unit ν and receive the message at time unit ν and finally, ν will connect with ν at

In recent years, broadcasting problems [1, 9–11, 13, 26, 32, 35] have gained considerable attention due to the rapid advancements in multimedia and network technologies. When utilizing the postal model, these problems can be viewed as extensions of center problems (with $\rho = 0$). The objective of broadcasting problems is to identify one or more nodes, referred to as broadcast centers, from which a message can be efficiently disseminated to other nodes with minimal time delay. Specifically, Su et al. [33] introduced a linear-time

algorithm for finding one broadcast center on a tree network.

Previous research on broadcasting problems typically assumed that the value of $w_{u,v}$ is deterministic. However, in practice, considering the uncertainty of $w_{u,v}$ is essential, as the available bandwidth of a communication link can vary over time. Thus, it is valuable to account for $w_{u,v}$ as uncertain, falling within a range $[w_{u,v}^-, w_{u,v}^+]$, where $w_{u,v}^-$ and $w_{u,v}^+$ are non-negative real numbers. The concept of uncertainty has been previously introduced in various optimization problems, such as facility location [5–8, 12, 16, 27, 29, 37, 38], resource allocation [2–4, 15], and job scheduling [17, 22, 28]. There was an approach, named minmax regret [5–8, 25, 36, 38, 39], proposed for solving them. Given an optimization problem with link weight uncertain, the minmax regret approach tries to minimize the worst-case loss to the objective function, without specifying the probability distribution of link weights. For a comprehensive discussion of the minmax regret approach, please refer to [25, 30].

In this paper, under the postal model, we investigate the broadcasting problem in which a single broadcast center is designated, and all edge weights are uncertain. When the underlying network has a random topology, this problem is known as NP-hard, even if all edge weights are deterministic [32]. The broadcasting problem considered was inspired by the 1-center problem. Note that the broadcasting problem becomes the 1-center problem when $\rho=0$. The 1-center problem is polynomial-time solvable on a random network with deterministic edge weights [18]. Clearly, the problem to be investigated is more intractable than the 1-center problem. With $\rho>0$, each sender has to determine an optimal broadcast sequence for its neighbors so as to minimize the broadcasting time to the whole network. Further, with uncertain edge weights, such an optimal broadcast sequence is more difficult to find, because it relies on the values of edge weights.

In their work [14], Burkard and Dollani addressed uncertain edge weights in the 1-center problem (with $\rho = 0$) and successfully solved it on a tree network in $O(n \log n)$ time. In this paper, also on a tree network with uncertain edge weights, we extend Burkard and Dollani's work by solving the broadcasting problem (with $\rho > 0$) with one broadcast center in $O(n \log n \log \log n)$ time. In comparison, a multiplier of $O(\log \log n)$

is incurred, as a consequence of the inclusion of $\rho > 0$. In Table 1, we show the best results thus far for the two problems on both a random network with deterministic edge weights and a tree network with uncertain edge weights.

Table 1: Comparison between the 1-center problem and the broadcasting problem with one broadcast center.

	value of ρ	on a random network with deterministic edge wights	on a tree network with uncertain edge wights
1-center problem	0	$O(n^2 \log n + nm) \ [18]$	$O(n\log n)$ [14]
broadcasting problem with one broadcast center	a positive constant	NP-hard [32]	$\frac{O(n\log n\log\log n)}{\text{(this paper)}}$

The execution of the proposed algorithm is iterative and based on the prune-andsearch strategy. To begin, the algorithm was executed on the tree T. Then, a centroid x of T, a worst-case scenario, and a broadcast center κ under the worst-case scenario were found, which takes $O(n \log \log n)$ time. As a consequence of Lemma 7, a minmax-regret broadcast center κ^* of T is either x or located in the subtree of T-x that contains κ . If $\kappa^* \neq x$, then the algorithm execution continues on the subtree containing κ . Since at most $O(\log n)$ iterations are executed, the overall time complexity is $O(n \log n \log \log n)$.

The most challenging aspect of the proposed algorithm is conducting a worst-case scenario search in $O(n \log \log n)$ time, given the potentially infinite number of scenarios. To achieve this, we successfully reduced the search space to a maximum of n-1 scenarios through a series of scenario transformations. Through each transformation, we pinned down a particular edge weight, leading to the reduction of the search space. Finally, a worst-case scenario could be found in $O(n \log \log n)$ time, by the aid of a preprocessing and some dedicated data structures.

The rest of this paper is organized as follows. In Section 2, some notations and definitions, together with some properties, are introduced. In Section 3, an $O(n \log n \log \log n)$ -time algorithm is proposed for solving the broadcasting problem, with one broadcast center and uncertain edge weights, on a tree network. The correctness of the algorithm is verified in Section 4, and the time complexity is analyzed in Section 5. Finally, this paper concludes with some remarks in Section 6.

2 Preliminaries

Let T be a tree and $w_{u,v}$ be the weight of $(u,v) \in E(T)$. We suppose that n = |V(T)| and the value of $w_{u,v}$ is randomly generated from $[w_{u,v}^-, w_{u,v}^+]$, where $w_{u,v}^-$ and $w_{u,v}^+$ are two non-negative real numbers. Define C to be a Cartesian product of all n-1 intervals $[w_{u,v}^-, w_{u,v}^+]$ for T. For example, refer to the tree of Figure 1, where $[w_{u,v_1}^-, w_{u,v_1}^+] = [2, 6]$ and $[w_{u,v_2}^-, w_{u,v_2}^+] = [1, 4]$. We have $C = \{(g, h) \mid 2 \le g \le 6 \text{ and } 1 \le h \le 4\}$ or $C = \{(g, h) \mid 1 \le g \le 4 \text{ and } 2 \le h \le 6\}$, where g and h are two real numbers. Throughout this paper, we assume that C is predetermined. Also, notice that for the notations introduced in this section, if they are defined with respect to T, we remove T from them for simplicity (so, we use C, instead of C_T or C(T)).

$$\overset{v_1}{\bigcirc} \overset{[2,6]}{\bigcirc} \overset{u}{\bigcirc} \overset{[1,4]}{\bigcirc} \overset{v_2}{\bigcirc}$$

Figure 1: A tree T.

Clearly, C is an infinite set. A tuple $s = (s_1, s_2, \ldots, s_{n-1})$ is called a *scenario* of T, if $s \in C$. For the example of Figure 1, if $C = \{(g, h) \mid 2 \leq g \leq 6 \text{ and } 1 \leq h \leq 4\}$, then (5, 2) is a scenario of T, but (2, 5) is not. In fact, a scenario of T represents a feasible weight assignment of all edges in E(T), in the sequence specified by C. Suppose $s \in C$. Define $w_{u,v}^s$ to be the weight of $(u, v) \in E(T)$, $b_time^s(u, G)$ to be the minimum time required for u to broadcast a message to all other vertices of G, and B_Ctr^s to be the set of broadcast centers of T, all under the scenario s. That is, $B_Ctr^s = \{u \mid b_time^s(u, T) \leq b_time^s(v, T)$ for all $v \in V(T)\}$.

Further, for any $x, y \in V(T)$, we define $r_{x,y}^s = b_time^s(x,T) - b_time^s(y,T)$ and $max_r(x) = \max\{r_{x,y}^s \mid y \in V(T) \text{ and } s \in C\}$, where $r_{x,y}^s$ is called the relative regret of x with respect to y and $max_r(x)$ is called the maximum regret of x. Intuitively, $r_{x,y}^s$ represents the time lost, if x was chosen, instead of y, to broadcast a message over T under the scenario s. Notice that if $max_r(x) = r_{x,y'}^{s'}$ for some $y' \in V(T)$ and $s' \in C$, then $y' \in B_Ctr^{s'}$, and s' is called a worst-case scenario of T with respect to x. It is possible that there are multiple worst-case scenarios of T with respect to x. The minmax-

regret broadcasting problem on T with edge weight uncertainty is to find a minmax-regret broadcast center $\kappa^* \in V(T)$ such that its maximum regret is minimized, i.e., $max_r(\kappa^*) \le max_r(v)$ for all $v \in V(T)$.

Suppose $x, y \in V(T)$. Removing x from T will result in some subtrees of T. We use $B_{x,y}$ to denote the subtree that contains y, and let $\bar{B}_{x,y} = T - B_{x,y}$ (refer to Figure 2). We have $B_{x,y} = \bar{B}_{y,x}$, if $(x,y) \in E(T)$.




Figure 2: $B_{x,y}$ and $\bar{B}_{x,y}$.

Lemma 1 ([33]) Suppose $s \in C$ and $(u, v) \in E(T)$. If $b_time^s(u, \bar{B}_{u,v}) \leq b_time^s(v, \bar{B}_{v,u})$, then the following hold:

- $b_time^s(u,T) = \rho + w_{u,v}^s + b_time^s(v,\bar{B}_{v,u});$
- $b_time^s(v,T) \le b_time^s(u,T)$.

A tree is a *star*, if it has one or two vertices, or has exactly one vertex whose degree is greater than one. The latter is called the *center* of the star. For the one-vertex star, the vertex is the center. For the two-vertex star, either vertex can be the center.

Lemma 2 ([33]) The subgraph of T induced by B_Ctr^s is a star.

Define $N_T(u) = \{v \mid (u,v) \in E(T)\}$ to be the set of neighboring vertices of u in T. A vertex $\hat{\kappa} \in B_Ctr^s$ is a *prime broadcast center* of T under the scenario s if and only if $b_time^s(\hat{\kappa}, \bar{B}_{\hat{\kappa},u}) \geq b_time^s(u, \bar{B}_{u,\hat{\kappa}})$ for all $u \in N_T(\hat{\kappa})$. Further, according to Lemma 1, if $\hat{\kappa}$ is a prime broadcast center of T under the scenario s, then $b_time^s(\hat{\kappa}, T) \leq b_time^s(u, T)$ for all $u \in N_T(\hat{\kappa})$. **Lemma 3** A prime broadcast center of T always exists under any scenario of T.

Proof. Let $s \in C$ be an arbitrary scenario of T. When |V(T)| = 1, it is easy to see that the only vertex is a prime broadcast center of T. So, we assume $|V(T)| \geq 2$. According to Lemma 2, the subgraph of T induced by B_Ctr^s is a star. Let κ be the center of the star. If κ is not a prime broadcast center of T, then there exists $x \in N_T(\kappa)$ such that $b_time^s(\kappa, \bar{B}_{\kappa,x}) < b_time^s(x, \bar{B}_{x,\kappa})$. It follows, as a consequence of Lemma 1, that $b_time^s(x,T) \leq b_time^s(\kappa,T)$, which implies $x \in B_Ctr^s$. Clearly, x is a prime broadcast center of T, if $N_T(x) - {\kappa} = \emptyset$. Otherwise, for each $v \in N_T(x) - {\kappa}$, we have $v \notin B_Ctr^s$, i.e., $b_time^s(x,T) < b_time^s(v,T)$ (as a consequence of κ being the center of the star). Again, as a consequence of Lemma 1, we have $b_time^s(x,\bar{B}_{x,v}) > b_time^s(v,\bar{B}_{v,x})$. Therefore, x is a prime broadcast center of T.

Suppose $x, y \in V(T)$. Let $P_{x,y}$ be the set of edges contained in the x-to-y path of T, $d_{x,y} = |P_{x,y}|$, and $\tilde{w}_{x,y}^s$ be the total weight of all edges in $P_{x,y}$ under the scenario s, i.e. $\tilde{w}_{x,y}^s = \sum_{(u,v) \in P_{x,y}} w_{u,v}^s$. The following lemma shows a relation between $b_time^s(x,T)$ and $b_time^s(\hat{\kappa}, \bar{B}_{\hat{\kappa},x})$, where $\hat{\kappa}$ is a prime broadcast center of T under the scenario s.

Lemma 4 Suppose that $s \in C$ and $\hat{\kappa}$ is a prime broadcast center of T under s. If $x \in V(T) - \{\hat{\kappa}\}$, then $b_time^s(x,T) = d_{x,\hat{\kappa}} \cdot \rho + \tilde{w}^s_{x,\hat{\kappa}} + b_time^s(\hat{\kappa}, \bar{B}_{\hat{\kappa},x})$.

Proof. Suppose that (u_0, u_1, \ldots, u_h) is the x-to- $\hat{\kappa}$ path of T, where $u_0 = x$ and $u_h = \hat{\kappa}$. Refer to Figure 3. For $0 \le i \le h-1$, $\bar{B}_{u_i,u_{i+1}}$ is a subgraph of $\bar{B}_{u_{h-1},\hat{\kappa}}$, and $\bar{B}_{\hat{\kappa},u_{h-1}}$ is a subgraph of \bar{B}_{u_{i+1},u_i} . Now that $b_time^s(\hat{\kappa}, \bar{B}_{\hat{\kappa},u_{h-1}}) \ge b_time^s(u_{h-1}, \bar{B}_{u_{h-1},\hat{\kappa}})$, we have $b_time^s(u_i, \bar{B}_{u_i,u_{i+1}}) \le b_time^s(u_{i+1}, \bar{B}_{u_{i+1},u_i})$ for $0 \le i \le h-1$. Then, according to Lemma 1, we have

$$b_time^s(u_i, T) = \rho + w_{u_i, u_{i+1}}^s + b_time^s(u_{i+1}, \bar{B}_{u_{i+1}, u_i}) \text{ for } 0 \le i \le h - 1.$$
 (1)

Observe $\{(u_i, u_{i+1})\} \cup E(\bar{B}_{u_{i+1}, u_i}) \subseteq E(\bar{B}_{u_i, u_{i-1}}) \subset E(T)$ for $1 \leq i \leq h-1$. It implies $\rho + w^s_{u_i, u_{i+1}} + b_time^s(u_{i+1}, \bar{B}_{u_{i+1}, u_i}) \leq b_time^s(u_i, \bar{B}_{u_i, u_{i-1}}) \leq b_time^s(u_i, T)$ for $1 \leq i \leq h-1$, where the two equalities hold, as a consequence of (1). That is, we have

 $b_time^s(u_i, \bar{B}_{u_i, u_{i-1}}) = \rho + w^s_{u_i, u_{i+1}} + b_time^s(u_{i+1}, \bar{B}_{u_{i+1}, u_i}) \text{ for } 1 \leq i \leq h-1. \text{ It follows}$ that $b_time^s(x, T) = d_{x,\hat{\kappa}} \cdot \rho + \tilde{w}^s_{x,\hat{\kappa}} + b_time^s(\hat{\kappa}, \bar{B}_{\hat{\kappa}, u_{h-1}}), \text{ where } \bar{B}_{\hat{\kappa}, u_{h-1}} = \bar{B}_{\hat{\kappa}, x}.$




Figure 3: proof of Lemma 4.

Also notice that $d_{x,\hat{\kappa}} \cdot \rho + \tilde{w}^s_{x,\hat{\kappa}}$ in Lemma 4 is the minimum time requirement for x to broadcast a message along the x-to- $\hat{\kappa}$ path. Actually, Lemma 4 reveals that $b_time^s(x,T)$ can be calculated as the minimum time requirement for x to broadcast a message over a subgraph, i.e., the union of the x-to- $\hat{\kappa}$ path and $\bar{B}_{\hat{\kappa},x}$, of T. Moreover, the proof of Lemma 4 shows that an optimal transmission sequence for $b_time^s(x,T)$ has each u_i connected to u_{i+1} immediately after u_i has received the message. In general, for any $x, y \in V(T)$ and $x \neq y$, if y is not a prime broadcast center of T under the scenario s, then we have $b_time^s(x,T) \geq d_{x,y} \cdot \rho + \tilde{w}^s_{x,y} + b_time^s(y,\bar{B}_{y,x})$. Therefore, we have the following lemma for $x \neq y$.

Lemma 5 Suppose $s \in C$ and $x, y \in V(T)$. If $x \neq y$, then $b_time^s(x, T) \geq d_{x,y} \cdot \rho + \tilde{w}^s_{x,y} + b_time^s(y, \bar{B}_{y,x})$.

Lemma 6 Suppose $s \in C$. A prime broadcast center $\hat{\kappa}$ of T under s is a center of the star induced by B- Ctr^s .

Proof. If $\hat{\kappa}$ is not a center of the star induced by B_Ctr^s (refer to Lemma 2), then there exist two broadcast centers κ_1 and κ_2 in B_Ctr^s such that $(\hat{\kappa}, \kappa_1, \kappa_2)$ is a path in T. According to Lemma 4, we have $b_time^s(\kappa_1, T) = d_{\kappa_1, \hat{\kappa}} \cdot \rho + \tilde{w}^s_{\kappa_1, \hat{\kappa}} + b_time^s(\hat{\kappa}, \bar{B}_{\hat{\kappa}, \kappa_1})$ and $b_time^s(\kappa_2, T) = d_{\kappa_2, \hat{\kappa}} \cdot \rho + \tilde{w}^s_{\kappa_2, \hat{\kappa}} + b_time^s(\hat{\kappa}, \bar{B}_{\hat{\kappa}, \kappa_2})$. Since $P_{\kappa_1, \hat{\kappa}} \subset P_{\kappa_2, \hat{\kappa}}$ and $\bar{B}_{\hat{\kappa}, \kappa_1} = \bar{B}_{\hat{\kappa}, \kappa_2}$, we have $b_time^s(\kappa_1, T) < b_time^s(\kappa_2, T)$, a contradiction.

3 Finding a minmax-regret broadcast center

In this section, assuming that T is with edge weight uncertainty, we intend to find a minmax-regret broadcast center on T under the postal model. It was shown in [33] that given any $x \in V(T)$ and $s \in C$, both B_Ctr^s and $b_time^s(x,T)$ can be determined in O(n) time. Suppose that a worst-case scenario s' of T with respect to $u \in V(T)$ is available, i.e., $max_r(u) = r_{u,v'}^{s'}$ for some $v' \in V(T)$. Since $v' \in B_Ctr^{s'}$, we can compute $max_r(u)$ in O(n) time, using the results of [33]. Further, a minmax-regret broadcast center of T can be obtained, if $max_r(u)$ is computed for all $u \in V(T)$.

In subsequent discussion, we use $\ddot{s}(u)$ to denote some worst-case scenario of T with respect to u. If Q(n) is the time requirement for finding $\ddot{s}(u)$, then the approach above takes $O(n^2 + n \cdot Q(n))$ time to find a minmax-regret broadcast center of T, because it examines all n vertices of T. In this section, a more efficient approach to finding a minmax-regret broadcast center of T is suggested. Inspired by the following lemma, the new approach can result in an $O(n \log n + Q(n) \cdot \log n)$ time algorithm.

Lemma 7 Suppose $x \in V(T)$. If $x \in B_Ctr^{\ddot{s}(x)}$, then x is a minmax-regret broadcast center of T. Otherwise, a minmax-regret broadcast center of T can be found in $V(B_{x,\kappa}) \cup \{x\}$, where $\kappa \in B_Ctr^{\ddot{s}(x)}$.

Proof. Clearly, when $x \in B_Ctr^{\check{s}(x)}$, we have $max_r(x) = 0$, implying that x is a minmax-regret broadcast center of T, because $max_r(v) \geq 0$ for every $v \in V(T)$. Then, we assume $x \not\in B_Ctr^{\check{s}(x)}$. It suffices to show that for every $x' \in V(T) - V(B_{x,\kappa}) - \{x\}$, we have $max_r(x') > max_r(x)$. Suppose that $\hat{\kappa}$ is a prime broadcast center of T under $\check{s}(x)$. According to Lemma 4, we have $b_time^{\check{s}(x)}(x',T) = d_{x',\hat{\kappa}} \cdot \rho + \tilde{w}^{\check{s}(x)}_{x',\hat{\kappa}} + b_time^{\check{s}(x)}(\hat{\kappa},\bar{B}_{\hat{\kappa},x'})$ and $b_time^{\check{s}(x)}(x,T) = d_{x,\hat{\kappa}} \cdot \rho + \tilde{w}^{\check{s}(x)}_{x,\hat{\kappa}} + b_time^{\check{s}(x)}(\hat{\kappa},\bar{B}_{\hat{\kappa},x})$. As a consequence of Lemma 2, we have $\hat{\kappa} \in B_Ctr^{\check{s}(x)} \subseteq V(B_{x,\kappa})$. Since $P_{x,\hat{\kappa}} \subset P_{x',\hat{\kappa}}$, we have $d_{x',\hat{\kappa}} \cdot \rho + \tilde{w}^{\check{s}(x)}_{x',\hat{\kappa}} > d_{x,\hat{\kappa}} \cdot \rho + \tilde{w}^{\check{s}(x)}_{x,\hat{\kappa}}$. It follows that we have $b_time^{\check{s}(x)}(x',T) > b_time^{\check{s}(x)}(x,T)$, because $\bar{B}_{\hat{\kappa},x'} = \bar{B}_{\hat{\kappa},x}$. So, we have $max_r(x') \geq r^{\check{s}(x)}_{x',\hat{\kappa}} = b_time^{\check{s}(x)}(x',T) - b_time^{\check{s}(x)}(\hat{\kappa},T) > b_time^{\check{s}(x)}(x,T) - b_time^{\check{s}(x)}(\hat{\kappa},T) = r^{\check{s}(x)}_{x,\hat{\kappa}} = max_r(x)$.

With Lemma 7, a minmax-regret broadcast center of T can be found by the pruneand-search strategy. The search starts on T. A vertex z is first determined such that the largest open z-branch has minimum size, i.e., $\max_{y \in V(T) - \{z\}} \{|B_{z,y}|\} \le \max_{y \in V(T) - \{v\}} \{|B_{v,y}|\}$ for every $v \in V(T)$. The vertex z thus found is called a *centroid* of T. If $z \in B \mathcal{L}tr^{\ddot{s}(z)}$, then z is a minmax-regret broadcast center of T and the search terminates.

Otherwise, a broadcast center $\kappa \in B_Ctr^{\ddot{s}(z)}$ of T under $\ddot{s}(z)$ is found, and the search continues on the subtree of T induced by $V(B_{z,\kappa}) \cup \{z\}$. The resulting subtree is denoted by T'. A centroid z' of T' is then determined. Again, if $z' \in B_Ctr^{\ddot{s}(z')}$, then the search terminates with z' being a minmax-regret broadcast center of T. If $z' \notin B_Ctr^{\ddot{s}(z')}$, then some $\kappa' \in B_Ctr^{\ddot{s}(z')}$ is found and the search continues on the subtree of T' induced by $V(T') \cap \{V(B_{z',\kappa'}) \cup \{z'\}\}$.

The search will proceed until z is a minmax-regret broadcast center of T or the size of the induced subtree is small enough. For the latter case, a minmax-regret broadcast center of T can be found with an additional O(n + Q(n)) time. The detailed execution is expressed as Algorithm 1 below.

Algorithm 1 Determining a minmax-regret broadcast center of a tree

```
Input: a tree T.
Output: a minmax-regret broadcast center of T.
 1: T' \leftarrow T;
 2: while |V(T')| \ge 3 do
         determine a centroid z of T';
 3:
         find \ddot{s}(z) and B L Ctr^{\ddot{s}(z)};
 4:
         if z \in B L Ctr^{\ddot{s}(z)} then
 5:
             return z;
 6:
         else
 7:
             find \kappa \in B L Ctr^{\ddot{s}(z)};
 8:
             T' \leftarrow \text{the subtree of } T' \text{ induced by } V(T') \cap \{V(B_{z,\kappa}) \cup \{z\}\};
 9:
         end if
10:
11: end while
12: determine v \in V(T') whose max_r(v) is minimum;
13: return v.
```

It was shown in [19,23] that a centroid z of T can be found in O(n) time. Besides, the largest open z-branch contains at most $\lfloor \frac{n}{2} \rfloor$ vertices. However, it is not easy to find $\ddot{s}(z)$, because there are an infinite number of scenarios. Suppose $V(T) = \{v_1, v_2, \dots, v_n\}$.

For $x \in V(T)$, we would like to construct a finite set $\ddot{S}(x) = \bigcup_{1 \leq \ell \leq n} \Psi_{\ell}(x)$ of scenarios such that a worst-case scenario of T with respect to x can be found in some $\Psi_{\ell}(x)$, i.e., $\ddot{S}(x) \in \Psi_{\ell}(x)$. The construction of $\ddot{S}(x)$ is described below.

In the rest of this paper, we assume $x = v_n$, without losing generality. We set $\Psi_n(x) = \{c\}$, where $c \in C$ is an arbitrary scenario of T. Define α_{x,v_i} to be the base scenario of T with respect to x and v_i , where $1 \le i \le n-1$, which satisfies the following:

- $w_{a,b}^{\alpha_{x,v_i}} = w_{a,b}^+$, if $(a,b) \in P(x,v_i) \cup E(\bar{B}_{v_i,x})$;
- $w_{a,b}^{\alpha_{x,v_i}} = w_{a,b}^-$, else.

Refer to Figure 4, where an illustrative example is shown with $\rho = 1$. In Figure 4(b), we use $\overline{7}$ and $\underline{5}$, for example, to denote $w_{a,b}^+$ and $w_{a,b}^-$, respectively. For any $s \in C$, we have $d_{x,v_i} \cdot \rho + \tilde{w}_{x,v_i}^s + b_time^s(v_i, \bar{B}_{v_i,x}) \leq d_{x,v_i} \cdot \rho + \tilde{w}_{x,v_i}^{\alpha_{x,v_i}} + b_time^{\alpha_{x,v_i}}(v_i, \bar{B}_{v_i,x})$. To construct $\Psi_i(x)$, we need to create new scenarios, which are modifications of α_{x,v_i} . The edge weights under α_{x,v_i} are to be changed according to a predefined sequence that is suggested by the following lemma.




Figure 4: An illustrative example with $\rho = 1$. (a) A tree T. (b) The base scenario α_{x,v_i} .

Lemma 8 [33] Suppose $x \in V(T)$, $s \in C$, and $N_T(x) = \{u_1, u_2, \ldots, u_h\}$, where $h = |N_T(x)|$. If $w_{x,u_k}^s + b$ -time^s $(u_k, \bar{B}_{u_k,x})$ is nonincreasing as k increases from 1 to h, then u_1, u_2, \ldots, u_h is an optimal sequence (with respect to minimum broadcast time) for x to

broadcast a message to $N_T(x)$ under s. That is, $b_time^s(x,T) = \max\{k \cdot \rho + w_{x,u_k}^s + b_time^s(u_k, \bar{B}_{u_k,x}) \mid 1 \le k \le h\}.$

Let $h_i = |N_{\bar{B}_{v_i,x}}(v_i)|$. If $h_i = 0$, then we set $\Psi_i(x) = \emptyset$. Otherwise, the vertices of $N_{\bar{B}_{v_i,x}}(v_i)$ are arranged as $u_{i,1}, u_{i,2}, \ldots, u_{i,h_i}$ so that $w_{v_i,u_{i,k}}^{\alpha_{x,v_i}} + b_time^{\alpha_{x,v_i}}(u_{i,k}, \bar{B}_{u_{i,k},v_i})$ is nonincreasing as k increases from 1 to h_i . For example, refer to Figure 4(b) again, where $w_{v_i,u_{i,k}}^{\alpha_{x,v_i}} + b_time^{\alpha_{x,v_i}}(u_{i,k}, \bar{B}_{u_{i,k},v_i}) = 13$, 12, 7, as k = 1, 2, 3, respectively. Then, we set $\Psi_i(x) = \{\beta_{x,v_i}^1, \beta_{x,v_i}^2, \ldots, \beta_{x,v_i}^{h_i}\}$, where each β_{x,v_i}^j , $1 \leq j \leq h_i$, is a scenario of T with respect to x and v_i that satisfies the following:

- $w_{a,b}^{\beta_{x,v_i}^j} = w_{a,b}^-$, if $(a,b) \in \bigcup_{j < k \le h_i} (\{(v_i, u_{i,k})\} + E(\bar{B}_{u_{i,k},v_i}));$
- $w_{a,b}^{\beta_{x,v_i}^j} = w_{a,b}^{\alpha_{x,v_i}}$, else.

Refer to Figure 5, where β_{x,v_i}^1 and β_{x,v_i}^2 are illustrated for the example of Figure 4. Compared with α_{x,v_i} , the changes (i.e., $w_{a,b}^{\beta_{x,v_i}^j} = w_{a,b}^-$) for β_{x,v_i}^j are based on the sequence of $u_{i,1}, u_{i,2}, \ldots, u_{i,h_i}$. That is, β_{x,v_i}^1 changes the weights of all edges in $(\{(v_i, u_{i,2})\} + E(\bar{B}_{u_{i,3},v_i})) \cup (\{(v_i, u_{i,3})\} + E(\bar{B}_{u_{i,3},v_i}))$, β_{x,v_i}^2 changes the weights of all edges in $(\{(v_i, u_{i,3})\} + E(\bar{B}_{u_{i,3},v_i}))$, and β_{x,v_i}^3 remains the same as α_{x,v_i} . For this example, if v_i is a prime broadcast center of T under some worst-case scenario of T with respect to x, then β_{x,v_i}^1 is a worst-case scenario of T with respect to x (i.e., $\beta_{x,v_i}^1 = \ddot{s}(x)$), as explained below.




Figure 5: Two instances of β_{x,v_i}^j . (a) β_{x,v_i}^1 . (b) β_{x,v_i}^2 .

According to Lemma 4, we have $b_time^{\ddot{s}(x)}(x,T) = d_{x,v_i} \cdot \rho + \tilde{w}_{x,v_i}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(v_i, \bar{B}_{v_i,x}),$ i.e., $b_time^{\ddot{s}(x)}(x,T) - b_time^{\ddot{s}(x)}(v_i, \bar{B}_{v_i,x}) = d_{x,v_i} \cdot \rho + \tilde{w}_{x,v_i}^{\ddot{s}(x)}.$ Further, we have $d_{x,v_i} \cdot \rho + \tilde{w}_{x,v_i}^{\ddot{s}(x)} \leq d_{x,v_i} \cdot 1 + \tilde{w}_{x,v_i}^{\beta_{x,v_i}^1} = 2 + 4 = 6,$ as a consequence of β_{x,v_i}^1 and α_{x,v_i} . Since $r_{x,v_i}^{\ddot{s}(x)} = b_time^{\ddot{s}(x)}(x,T) - b_time^{\ddot{s}(x)}(v_i,T) \leq b_time^{\ddot{s}(x)}(x,T) - b_time^{\ddot{s}(x)}(v_i,\bar{B}_{v_i,x}),$ we have $r_{x,v_i}^{\ddot{s}(x)} \leq d_{x,v_i} \cdot \rho + \tilde{w}_{x,v_i}^{\beta_{x,v_i}^1} = 6.$ On the other hand, we have $r_{x,v_i}^{\beta_{x,v_i}^1} = b_time^{\beta_{x,v_i}^1}(x,T) - b_time^{\beta_{x,v_i}^1}(v_i,T) = 20 - 14 = 6 \geq r_{x,v_i}^{\ddot{s}(x)}.$ Now that v_i is a prime broadcast center of T under $\ddot{s}(x)$, we have $r_{x,v_i}^{\ddot{s}(x)} = max_r(x)$. So, we have $r_{x,v_i}^{\beta_{x,v_i}^1} = max_r(x)$, implying that β_{x,v_i}^1 is a worst-case scenario of T with respect to x (i.e., $\beta_{x,v_i}^1 = \ddot{s}(x)$).

In general, if $\ddot{s}(x) \notin \Psi_n(x)$, then there exists $\beta^j_{x,v_i} = \ddot{s}(x)$ for some $1 \leq i < n$ and $1 \leq j \leq h_i$, where v_i is a prime broadcast center of T under some worst-case scenario of T with respect to x. After $\ddot{S}(x)$ (= $\bigcup_{1 \leq \ell \leq n} \Psi_{\ell}(x)$) is constructed, $\ddot{s}(x)$ can be determined from $\ddot{S}(x)$, as a consequence that $\max\{r^s_{x,y} \mid s \in \ddot{S}(x) \text{ and } y \in B_Ctr^s\}$ occurs when $s = \ddot{s}(x)$. The detailed execution for finding $\ddot{s}(x)$ is expressed as Algorithm 2 below.

Algorithm 2 Finding $\ddot{s}(x)$

```
Input: a tree T with vertices v_1, v_2, \ldots, v_n and a vertex x = v_n of T.
Output: \ddot{s}(x).
  1: set \Psi_n(x) = \{c\}, where c is an arbitrary scenario of T;
  2: for each v_i \in V(T) - \{x\} do
            construct \alpha_{x,v_i};
 3:
 4:
            h_i \leftarrow |N_{\bar{B}_{v_i,x}}(v_i)|;
            if h_i = 0 then
 5:
                 set \Psi_i(x) = \emptyset;
  6:
  7:
            else
                 arrange the vertices of N_{\bar{B}_{v_i,x}}(v_i) as u_{i,1}, u_{i,2}, \ldots, u_{i,h_i} so that w_{v_i,u_{i,k}}^{\alpha_{x,v_i}}+
 8:
                 b\_time^{\alpha_{x,v_i}}(u_{i,k},\bar{B}_{u_{i,k},v_i}) is nonincreasing as k increases from 1 to h_i;
                 construct \beta_{x,v_i}^1, \beta_{x,v_i}^2, \dots, \beta_{x,v_i}^{h_i};
 9:
                 set \Psi_i(x) = \{\beta_{x,v_i}^1, \beta_{x,v_i}^2, \dots, \beta_{x,v_i}^{h_i}\};
10:
11:
            end if
12: end for
13: determine s' \in \bigcup_{1 \le \ell \le n} \Psi_{\ell}(x) so that r_{x,y'}^{s'} = \max\{r_{x,y}^s \mid s \in \bigcup_{1 \le \ell \le n} \Psi_{\ell}(x) \text{ and } t \in \mathcal{S}_{\ell}(x) \}
      y \in B\_Ctr^s} for some y' \in B\_Ctr^{s'};
14: return s'.
```

The main result of this paper is stated as the following theorem, whose proof is shown in the following two sections.

Theorem 9 A minmax-regret broadcast center on a tree of n vertices with edge weight uncertainty can be found in $O(n \log n \log \log n)$ time.

4 Correctness

Clearly, with Lemma 7, a minmax-regret broadcast center κ^* of T can be found by Algorithm 1, provided $\ddot{s}(x)$ can be determined for any $x \in V(T)$. So, it suffices to determine $\ddot{s}(x)$, i.e., to verify Algorithm 2, in order to prove the correctness of Algorithm 1. Recall that we have $\Psi_n(x) = \{c\}$, where $c \in C$. If x is a prime broadcast center of T under $\ddot{s}(x)$, then we have $\max_x r(x) = 0$. Since $r_{x,y}^c \geq 0 = \max_x r(x)$ for any $y \in B_*Ctr^c$, we have $c = \ddot{s}(x) \in \Psi_n(x)$. In the rest of this section, we assume that x is not a prime broadcast center of T under any worst-case scenario of T with respect to x (which implies $|V(T)| \geq 3$). Besides, unless specified particularly, whenever a worst-case scenario c is mentioned, it means that c is a worst-case scenario of T with respect to x.

In order to identify a worst-case scenario in some $\Psi_{\ell}(x)$, where $1 \leq \ell < n$, we perform a series of scenario transformations, starting with a worst-case scenario (it does exist, although its edge weights are unknown), denoted by $\ddot{s}(x)$. First, we transform $\ddot{s}(x)$ into another worst-case scenario by specifying some edge weights. This transformation can be realized by the following fact whose proof will be presented in Section 4.1.

Fact 1 Suppose $x, y \in V(T)$ and y is a prime broadcast center of T under $\ddot{s}(x)$. Let s be a scenario of T that satisfies $w_{a,b}^s = w_{a,b}^{\alpha_{x,y}}$ if $(a,b) \in P_{x,y} \cup E(B_{y,x})$, and $w_{a,b}^s = w_{a,b}^{\ddot{s}(x)}$ else. Then, s is a worst-case scenario of T with respect to x. Besides, we have $y \in B$ -Ctr^s.

According to Fact 1, $\ddot{s}(x)$ is transformed into s, which is also a worst-case scenario. The weights of edges $(a,b) \in P_{x,y} \cup E(B_{y,x})$ are specified under s. However, y is not necessarily a prime broadcast center under s. The following fact finds out a worst-case scenario and an associated prime broadcast center.

Fact 2 Suppose $x \in V(T)$. There exist $\ddot{s}(x)$ and $y \in V(T)$ satisfying $w_{a,b}^{\ddot{s}(x)} = w_{a,b}^{\alpha_{x,y}}$ for each $(a,b) \in P_{x,y} \cup E(B_{y,x})$. Besides, y is a prime broadcast center of T under $\ddot{s}(x)$.

Fact 2 will be proved in Section 4.2, where a transformation from s (the worst-case scenario obtained by Fact 1) to another worst-case scenario is shown and an associated

prime broadcast center is found. To simplify notations, we use $\ddot{s}(x)$ and y to represent the resulting worst-case scenario and prime broadcast center, respectively. They may differ from those used in Fact 1.

Finally, we transform $\ddot{s}(x)$ into another worst-case scenario by specifying the weights of the other edges, i.e., all edges $(a,b) \in E(\bar{B}_{y,x})$ (= $E(T) - \{P_{x,y} \cup E(B_{y,x})\}$). The resulting scenario is identical with some $\beta_{x,y}^j$ ($\in \Psi_{\ell}(x)$, as $y = v_{\ell}$), and thus Algorithm 2 is verified. The transformation, which depends on two cases, can be carried out by the following two facts, where the notations μ , q_{0,τ_0} , h, and t^* are introduced in subsequent paragraphs ($h = h_{\ell}$ and $\beta_{x,y}^h$, $\beta_{x,y}^{t^*} \in \Psi_{\ell}(x)$, as $y = v_{\ell}$). Their proofs, which will be presented in Section 4.3 and Section 4.4, show two transformations from $\ddot{s}(x)$ to $\beta_{x,y}^h$ and $\beta_{x,y}^{t^*}$, respectively.

Fact 3 If $w_{y,\mu}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(\mu, B_{y,x}) \ge w_{y,q_{0,\tau_0}}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(q_{0,\tau_0}, \bar{B}_{q_{0,\tau_0},y})$, then $\beta_{x,y}^h$ is a worst-case scenario of T with respect to x.

Fact 4 If $w_{y,\mu}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(\mu, B_{y,x}) < w_{y,q_{0,\tau_{0}}}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(q_{0,\tau_{0}}, \bar{B}_{q_{0,\tau_{0}},y})$, then $\beta_{x,y}^{t^{*}}$ is a worst-case scenario of T with respect to x.

We use μ to denote the neighbor of y in $B_{y,x}$, and u_1, u_2, \ldots, u_h to denote the neighbors of y in $\bar{B}_{y,x}$ such that $w_{y,u_k}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(u_k, \bar{B}_{u_k,y})$ is nonincreasing as k increases from 1 to k. Refer to Figure 6. Let $\gamma_{x,y}^0, \gamma_{x,y}^1, \gamma_{x,y}^2, \ldots, \gamma_{x,y}^h$ denote k+1 scenarios of k with respect to k and k where k0 is k2 in k3. We satisfies the following:

•
$$w_{a,b}^{\gamma_{x,y}^{j}} = w_{a,b}^{\alpha_{x,y}}$$
, if $(a,b) \in \bigcup_{1 \le k \le j} (\{(y,u_k)\} + E(\bar{B}_{u_k,y}))$;

•
$$w_{a,b}^{\gamma_{x,y}^{j}} = w_{a,b}^{\ddot{s}(x)}$$
, else.

Since the weights of edges in $\{(y,\mu)\} \cup E(B_{y,x})$, i.e., the subtree at the right-bottom of Figure 6, were specified by Fact 2, only the weights of edges in $\bigcup_{j< k \leq h} (\{(y,u_k)\} + E(\bar{B}_{u_k,y}))$ are not specified under $\gamma_{x,y}^j$, where $\gamma_{x,y}^j$ and $\beta_{x,y}^j$ differ (refer to Figure 6).

Let $Q_0 = (q_{0,1}, q_{0,2}, \dots, q_{0,h})$ be an arrangement of u_1, u_2, \dots, u_h such that $w_{y,q_{0,k}}^{\gamma_{x,y}^0} + b_time^{\gamma_{x,y}^0}(q_{0,k}, \bar{B}_{q_{0,k},y})$ is nonincreasing as k increases from 1 to k. According to Lemma 8,




Figure 6: Comparison between $\gamma_{x,y}^j$ and $\beta_{x,y}^j$.

 Q_0 is an optimal sequence for y to broadcast a message to $N_{\bar{B}_{y,x}}(y)$ under $\gamma_{x,y}^0$. Moreover, for $1 \leq j \leq h$, let $Q_j = (q_{j,1}, q_{j,2}, \ldots, q_{j,h})$ be the sequence obtained from $Q_{j-1} = (q_{j-1,1}, q_{j-1,2}, \ldots, q_{j-1,h})$ by cyclically shifting $q_{j-1,j}, q_{j-1,j+1}, \ldots, q_{j-1,\ell}$ one position toward the right, where $q_{j-1,\ell} = u_j$ is assumed and $j \leq \ell$. That is, we have $(q_{j,1}, \ldots, q_{j,j-1}) = (q_{j-1,1}, \ldots, q_{j-1,j-1}), (q_{j,j}, q_{j,j+1}, q_{j,j+2}, \ldots, q_{j,\ell}) = (q_{j-1,\ell}, q_{j-1,j}, q_{j-1,j+1}, \ldots, q_{j-1,\ell-1}),$ and $(q_{j,\ell+1}, \ldots, q_{j,h}) = (q_{j-1,\ell+1}, \ldots, q_{j-1,h})$. It is not difficult to see that the first j vertices in Q_j are u_1, u_2, \ldots, u_j .

Also notice that $w_{y,q_{j,k}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(q_{j,k}, \bar{B}_{q_{j,k},y})$ is nonincreasing as k increases from 1 to h, as a consequence of $\gamma_{x,y}^{j}$ (refer to Figure 6). According to Lemma 8, Q_{j} is an optimal sequence for y to broadcast a message to $N_{\bar{B}_{y,x}}(y)$ under $\gamma_{x,y}^{j}$. Let $q_{j,\tau_{j}}$ be the first vertex in Q_{j} with $b_time^{\gamma_{x,y}^{j}}(y, \bar{B}_{y,x}) = \tau_{j} \cdot \rho + w_{y,q_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(q_{j,\tau_{j}}, \bar{B}_{q_{j,\tau_{j}},y})$, where the right-hand side is the broadcast time induced by $\bar{B}_{q_{j,\tau_{j}},y}$ ($q_{0,\tau_{0}}$ has the same definition as $q_{j,\tau_{j}}$).

Define Q_{t^*} to be the first sequence in Q_1, Q_2, \ldots, Q_h that has $q_{t^*,\tau_{t^*}} = q_{t^*,t^*}$. The proof of Fact 4 shows that $\gamma_{x,y}^{t^*}$ is a worst-case scenario and consequently, $\beta_{x,y}^{t^*}$ is a worst-case scenario (recall that $\gamma_{x,y}^{t^*}$ and $\beta_{x,y}^{t^*}$ differ in the weights of edges in $\bigcup_{t^* < k \le h} (\{(y, u_k)\} + E(\bar{B}_{u_k,y}))$. Below the existence of Q_{t^*} is guaranteed, by showing that Q_{τ_h} has $q_{\tau_h,\tau_h} = q_{\tau_h,\tau_h}$ (hence, $t^* \le \tau_h$).

Since Q_{τ_h} is an optimal sequence for y to broadcast a message to $N_{\bar{B}y,x}(y)$ under $\gamma_{x,y}^{\tau_h}$ and $u_{\tau_h}(=q_{\tau_h,\tau_h})$ is the τ_h -th vertex in Q_{τ_h} , we have $b_time^{\gamma_{x,y}^{\tau_h}}(y,\bar{B}_{y,x}) \geq \tau_h \cdot \rho + w_{y,q_{\tau_h,\tau_h}}^{\gamma_{\tau_h}} + b_time^{\gamma_{x,y}^{\tau_h}}(q_{\tau_h,\tau_h},\bar{B}_{q_{\tau_h,\tau_h},y})$, i.e., the broadcast time induced by $\bar{B}_{q_{\tau_h,\tau_h},y}$. Besides, we have $q_{\tau_h,\tau_h} = u_{\tau_h} = q_{h,\tau_h}$, because u_{τ_h} is also the τ_h -th vertex in Q_h . Since $\gamma_{x,y}^{\tau_h}$ and $\gamma_{x,y}^{h}$ have the same weights of those edges in $\{(y,u_{\tau_h})\} + E(\bar{B}_{u_{\tau_h},y})$, we have $w_{y,u_{\tau_h}}^{\gamma_{x,y}^{\tau_h}} + b_time^{\gamma_{x,y}^{\tau_h}}(u_{\tau_h},\bar{B}_{u_{\tau_h},y})$. Consequently, we have

$$\begin{array}{ll} b_time^{\gamma_{x,y}^{\tau_h}}(y,\bar{B}_{y,x}) & \geq & \tau_h \cdot \rho + w_{y,q_{\tau_h,\tau_h}}^{\gamma_{x,y}^{\tau_h}} + b_time^{\gamma_{x,y}^{\tau_h}}(q_{\tau_h,\tau_h},\bar{B}_{q_{\tau_h,\tau_h},y}) \\ & = & \tau_h \cdot \rho + w_{y,q_{h,\tau_h}}^{\gamma_{x,y}^{h}} + b_time^{\gamma_{x,y}^{h}}(q_{h,\tau_h},\bar{B}_{q_{h,\tau_h},y}) \\ & = & b_time^{\gamma_{x,y}^{h}}(y,\bar{B}_{y,x}). \end{array}$$

On the other hand, since $\tau_h \leq h$, we have $b_time^{\gamma_{x,y}^{\tau_h}}(y, \bar{B}_{y,x}) \leq b_time^{\gamma_{x,y}^{h}}(y, \bar{B}_{y,x})$ (refer to Figure 6). Therefore, $b_time^{\gamma_{x,y}^{\tau_h}}(y, \bar{B}_{y,x}) = b_time^{\gamma_{x,y}^{h}}(y, \bar{B}_{y,x})$ holds, which implies $b_time^{\gamma_{x,y}^{\tau_h}}(y, \bar{B}_{y,x}) = \tau_h \cdot \rho + w_{y,q\tau_h,\tau_h}^{\gamma_{x,y}^{\tau_h}} + b_time^{\gamma_{x,y}^{\tau_h}}(q_{\tau_h,\tau_h}, \bar{B}_{q\tau_h,\tau_h,y})$. Moreover, we have

$$b_time^{\gamma_{x,y}^{\tau_h}}(y, \bar{B}_{y,x}) = b_time^{\gamma_{x,y}^{h}}(y, \bar{B}_{y,x})$$

$$> k \cdot \rho + w_{y,q_{h,k}}^{\gamma_{x,y}^{h}} + b_time^{\gamma_{x,y}^{h}}(q_{h,k}, \bar{B}_{q_{h,k},y})$$

$$= k \cdot \rho + w_{y,q_{h,k}}^{\gamma_{x,y}^{\tau_h}} + b_time^{\gamma_{x,y}^{\tau_h}}(q_{\tau_h,k}, \bar{B}_{q_{\tau_h,k},y})$$

for all $1 \leq k < \tau_h$. It follows that we have $q_{\tau_h, \tau_{\tau_h}} = q_{\tau_h, \tau_h}$.

In the rest of this section, we show the proofs of Fact 1, Fact 2, Fact 3, and Fact 4.

4.1 Proof of Fact 1

In order to prove Fact 1, it suffices to show $r_{x,y}^s \geq r_{x,y}^{\ddot{s}(x)}$. As explained below, we first claim

$$\text{b_time}^s(y,T) \leq \text{b_time}^{\ddot{s}(x)}(y,T) + \tilde{w}^s_{x,y} - \tilde{w}^{\ddot{s}(x)}_{x,y}.$$

Without loss of generality, suppose $N_T(y) = \{v_1, v_2, \dots, v_h\}$, where $w_{y,v_k}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(v_k, \bar{B}_{v_k,y})$ is nonincreasing as k increases from 1 to h. According to Lemma 8, (v_1, v_2, \dots, v_h) is an optimal sequence for y to broadcast a message to $N_T(y)$ under $\ddot{s}(x)$. Besides, we have $b_time^{\ddot{s}(x)}(y, T) = \max\{k \cdot \rho + w_{y,v_k}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(v_k, \bar{B}_{v_k,y}) \mid 1 \leq k \leq h\}$. Since $b_time^s(y, T) \leq \max\{k \cdot \rho + w_{y,v_k}^s + b_time^s(v_k, \bar{B}_{v_k,y}) \mid 1 \leq k \leq h\}$, the claim is true provided $w_{y,v_k}^s + b_time^s(v_k, \bar{B}_{v_k,y}) \leq (w_{y,v_k}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(v_k, \bar{B}_{v_k,y})) + \tilde{w}_{x,y}^s - \tilde{w}_{x,y}^{\ddot{s}(x)}$ for $1 \leq k \leq k$

h. When $x \notin V(\bar{B}_{v_k,y})$, we have $w_{y,v_k}^s + b \pm time^s(v_k, \bar{B}_{v_k,y}) = w_{y,v_k}^{\bar{s}(x)} + b \pm time^{\bar{s}(x)}(v_k, \bar{B}_{v_k,y})$ and $\tilde{w}_{x,y}^s = \tilde{w}_{x,y}^{\alpha_{x,y}} \geq \tilde{w}_{x,y}^{\bar{s}(x)}$, as a consequence of the scenario s. The claim then follows.

So, we assume $x \in V(\bar{B}_{v_k,y})$. Let s' be a scenario of T, which has $w_{a,b}^{s'} = w_{a,b}^{\alpha_{x,y}}$ if $(a,b) \in P_{x,y}$, and $w_{a,b}^{s'} = w_{a,b}^{\bar{s}(x)}$ else. Then, we have $b_time^s(v_k, \bar{B}_{v_k,y}) \leq b_time^{s'}(v_k, \bar{B}_{v_k,y})$. Since s' and $\ddot{s}(x)$ differ in the edge weights of $P_{x,y}$, we have $b_time^{s'}(v_k, \bar{B}_{v_k,y}) \leq b_time^{s'}(v_k, \bar{B}_{v_k,y}) + \tilde{w}_{v_k,x}^{s'} - \tilde{w}_{v_k,x}^{\bar{s}(x)}$ (the right-hand side is an upper bound on the time requirement for v_k to broadcast over $\bar{B}_{v_k,y}$ under s' using the optimal broadcast sequence with respect to $b_time^{\ddot{s}(x)}(v_k, \bar{B}_{v_k,y})$). Further, since $\tilde{w}_{v_k,x}^{s'} = \tilde{w}_{v_k,x}^{s}$, $\tilde{w}_{x,y}^{s} = w_{y,v_k}^{s} + \tilde{w}_{v_k,x}^{s}$, and $\tilde{w}_{x,y}^{\ddot{s}(x)} = w_{y,v_k}^{\ddot{s}(x)} + \tilde{w}_{v_k,x}^{\ddot{s}(x)}$, the claim follows, because

$$w_{y,v_{k}}^{s} + b_time^{s}(v_{k}, \bar{B}_{v_{k},y})$$

$$\leq w_{y,v_{k}}^{s} + b_time^{\ddot{s}(x)}(v_{k}, \bar{B}_{v_{k},y}) + \tilde{w}_{v_{k},x}^{s} - \tilde{w}_{v_{k},x}^{\ddot{s}(x)}$$

$$= w_{y,v_{k}}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(v_{k}, \bar{B}_{v_{k},y}) + \tilde{w}_{x,y}^{s} - \tilde{w}_{x,y}^{\ddot{s}(x)}.$$

Then, the lemma holds, as a consequence of

$$\begin{split} r_{x,y}^s &= b_time^s(x,T) - b_time^s(y,T) \\ &\geq (d_{x,y} \cdot \rho + \tilde{w}_{x,y}^s + b_time^s(y,\bar{B}_{y,x})) - (b_time^{\ddot{s}(x)}(y,T) + \tilde{w}_{x,y}^s - \tilde{w}_{x,y}^{\ddot{s}(x)}) \\ &\quad \text{(by Lemma 5 and the claim)} \\ &= d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(y,\bar{B}_{y,x}) - b_time^{\ddot{s}(x)}(y,T) \\ &\quad (\because b_time^s(y,\bar{B}_{y,x}) = b_time^{\ddot{s}(x)}(y,\bar{B}_{y,x})) \\ &= b_time^{\ddot{s}(x)}(x,T) - b_time^{\ddot{s}(x)}(y,T) \quad \text{(by Lemma 4)} \\ &= r_{x,y}^{\ddot{s}(x)}. \end{split}$$

4.2 Proof of Fact 2

Let s_1 be the worst-case scenario and $y_1 \in B_-Ctr^{s_1}$, both obtained by Fact 1. If y_1 is a prime broadcast center of T under s_1 , then s_1 is a desired $\ddot{s}(x)$. Otherwise, it is implied by Lemma 6 that the prime broadcast center, say $\hat{\kappa}_1$, of T under s_1 is neighboring to y_1 in T. If $\hat{\kappa}_1 \ (\neq x)$ is on the x-to- y_1 path, then we have $P_{x,\hat{\kappa}_1} \cup E(B_{\hat{\kappa}_1,x}) \subset P_{x,y_1} \cup E(B_{y_1,x})$. Besides, we have $w_{a,b}^{s_1} = w_{a,b}^{\alpha_{x,y_1}} = w_{a,b}^{\alpha_{x,\hat{\kappa}_1}}$ for each $(a,b) \in P_{x,\hat{\kappa}_1} \cup E(B_{\hat{\kappa}_1,x})$, implying that s_1 is a desired $\ddot{s}(x)$.

If $\hat{\kappa}_1$ is not on the x-to- y_1 path, i.e., $d_{x,\hat{\kappa}_1} = d_{x,y_1} + 1$, then according to Fact 1, there is another worst-case scenario s_2 , which has $w_{a,b}^{s_2} = w_{a,b}^{\alpha_{x,\hat{\kappa}_1}}$ if $(a,b) \in P_{x,\hat{\kappa}_1} \cup E(B_{\hat{\kappa}_1,x})$, and $w_{a,b}^{s_2} = w_{a,b}^{s_1}$ else, where $\hat{\kappa}_1 \in B_Ctr^{s_2}$. Similarly, it can be concluded that either s_2 is a desired $\ddot{s}(x)$ or $d_{x,\hat{\kappa}_2} = d_{x,\hat{\kappa}_1} + 1$ holds, where $\hat{\kappa}_2$ is a prime broadcast center of T under s_2 . Thus, a desired $\ddot{s}(x)$ will be obtained eventually, if the above process is done repeatedly.

4.3 Proof of Fact 3

Notice that $\ddot{s}(x)$ and y used in Fact 3 are inherited from Fact 2. The notations μ , h, $\gamma_{x,y}^0$, Q_0 , and τ_0 , which will be used below, were defined earlier (after Fact 4) in this section. Since $\beta_{x,y}^h = \alpha_{x,y}$, it suffices to show $r_{x,y}^{\alpha_{x,y}} \geq r_{x,y}^{\ddot{s}(x)}$, in order to prove Fact 3.

Let Q' be the sequence that is obtained from Q_0 by inserting μ between $q_{0,p-1}$ and $q_{0,p}$, where $w_{y,q_{0,p-1}}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(q_{0,p-1}, \bar{B}_{q_{0,p-1},y}) > w_{y,\mu}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(\mu, B_{y,x}) \geq w_{y,q_{0,p}}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(q_{0,p}, \bar{B}_{q_{0,p},y})$. That is, we have $Q' = (q'_1, \ldots, q'_{p-1}, q'_p, q'_{p+1}, \ldots, q'_{h+1}) = (q_{0,1}, \ldots, q_{0,p-1}, \mu, q_{0,p}, q_{0,p+1}, \ldots, q_{0,h})$. According to Lemma 8, Q' is an optimal sequence for y to broadcast a message to $N_T(y)$ under $\ddot{s}(x)$. Since $q'_p = \mu$ and $q'_{\tau_0+1} = q_{0,\tau_0}$, we have

$$b_time^{\ddot{s}(x)}(y,T) = \max\{p \cdot \rho + w_{y,\mu}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(\mu, B_{y,x}),$$

$$(1 + \tau_0) \cdot \rho + w_{y,q_0,\tau_0}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(q_{0,\tau_0}, \bar{B}_{q_{0,\tau_0},y})\}$$

$$= \max\{p \cdot \rho + w_{y,\mu}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(\mu, B_{y,x}), \rho + b_time^{\ddot{s}(x)}(y, \bar{B}_{y,x})\}.$$
(2)

Recall that $Q_h = (q_{h,1}, q_{h,2}, \dots, q_{h,h})$ is an optimal sequence for y to broadcast a message to $N_{\bar{B}_{y,x}}(y)$ under $\gamma_{x,y}^h = (-\alpha_{x,y})$. Similarly, inserting μ between $q_{h,p-1}$ and $q_{h,p}$, we have $b_time^{\alpha_{x,y}}(y,T) \le \max\{p \cdot \rho + w_{y,\mu}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(\mu,B_{y,x}), \rho + b_time^{\alpha_{x,y}}(y,\bar{B}_{y,x})\}$. If $p \cdot \rho + w_{y,\mu}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(\mu,B_{y,x}) \ge \rho + b_time^{\alpha_{x,y}}(y,\bar{B}_{y,x})$, then $b_time^{\alpha_{x,y}}(y,T) \le p \cdot \rho + w_{y,\mu}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(\mu,B_{y,x})$. Since $w_{y,\mu}^{\bar{s}(x)} + b_time^{\bar{s}(x)}(\mu,B_{y,x}) = w_{y,\mu}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(\mu,B_{y,x})$ (as a consequence of Fact 2), we have

$$b_time^{\ddot{s}(x)}(y,T) \geq p \cdot \rho + w_{y,\mu}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(\mu, B_{y,x})$$
$$= p \cdot \rho + w_{y,\mu}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(\mu, B_{y,x})$$
$$\geq b_time^{\alpha_{x,y}}(y,T)$$

Further, since $b_time^{\alpha_{x,y}}(x,\bar{B}_{y,x}) \geq b_time^{\ddot{s}(x)}(x,\bar{B}_{y,x})$, we have

$$r_{x,y}^{\alpha_{x,y}} = b_time^{\alpha_{x,y}}(x,T) - b_time^{\alpha_{x,y}}(y,T)$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(y,\bar{B}_{y,x}) - b_time^{\alpha_{x,y}}(y,T) \quad \text{(by Lemma 5)}$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\bar{s}(x)} + b_time^{\bar{s}(x)}(y,\bar{B}_{y,x}) - b_time^{\alpha_{x,y}}(y,T)$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\bar{s}(x)} + b_time^{\bar{s}(x)}(y,\bar{B}_{y,x}) - b_time^{\bar{s}(x)}(y,T)$$

$$= b_time^{\bar{s}(x)}(x,T) - b_time^{\bar{s}(x)}(y,T) \quad \text{(by Lemma 4)}$$

$$= r_{x,y}^{\bar{s}(x)}. \quad (3)$$

If $p \cdot \rho + w_{y,\mu}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(\mu, B_{y,x}) < \rho + b_time^{\alpha_{x,y}}(y, \bar{B}_{y,x})$, then $b_time^{\alpha_{x,y}}(y, T) \le \rho + b_time^{\alpha_{x,y}}(y, \bar{B}_{y,x})$, and hence

$$r_{x,y}^{\alpha_{x,y}} = b_time^{\alpha_{x,y}}(x,T) - b_time^{\alpha_{x,y}}(y,T)$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(y,\bar{B}_{y,x}) - b_time^{\alpha_{x,y}}(y,T) \quad \text{(by Lemma 5)}$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\alpha_{x,y}} + b_time^{\alpha_{x,y}}(y,\bar{B}_{y,x}) - (\rho + b_time^{\alpha_{x,y}}(y,\bar{B}_{y,x}))$$

$$= d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\alpha_{x,y}} - \rho$$

$$= d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\tilde{s}(x)} - \rho \quad \text{(by Fact 2)}. \tag{4}$$

Moreover, we have $b_time^{\ddot{s}(x)}(x,T) = d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(y,\bar{B}_{y,x})$ by Lemma 4. Since $b_time^{\ddot{s}(x)}(y,T) \ge \rho + b_time^{\ddot{s}(x)}(y,\bar{B}_{y,x})$ (according to (2)), we have

$$r_{x,y}^{\ddot{s}(x)} = b_time^{\ddot{s}(x)}(x,T) - b_time^{\ddot{s}(x)}(y,T)$$

$$\leq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(y,\bar{B}_{y,x}) - (\rho + b_time^{\ddot{s}(x)}(y,\bar{B}_{y,x}))$$

$$= d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\ddot{s}(x)} - \rho$$

$$\leq r_{x,y}^{\alpha_{x,y}} \text{ (according to (4))}.$$

4.4 Proof of Fact 4

Notice that $\ddot{s}(x)$ and y used in Fact 4 are inherited from Fact 2 and the notations μ , h, t^* , $\gamma_{x,y}^j$, Q_j , and τ_j , which will be used below, were defined earlier in this section. In order to prove Fact 4, three lemmas are introduced first.

Lemma 10 $w_{y,q_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(q_{j,\tau_{j}}, \bar{B}_{q_{j,\tau_{j}},y})$ is nondecreasing as j increases from 0 to t^{*} .

Proof. It is equivalent to show

 $w_{y,q_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b \text{-}time^{\gamma_{x,y}^{j}}(q_{j,\tau_{j}}, \bar{B}_{q_{j,\tau_{j}},y}) \leq w_{y,q_{j+1},\tau_{j+1}}^{\gamma_{x,y}^{j+1}} + b \text{-}time^{\gamma_{x,y}^{j+1}}(q_{j+1,\tau_{j+1}}, \bar{B}_{q_{j+1},\tau_{j+1},y})$ for $0 \leq j < t^{*}$. We first claim that Q_{j} has $q_{j,\tau_{j}} \notin \{q_{j,1}, q_{j,2}, \dots, q_{j,j}\}$ under $\gamma_{x,y}^{j}$ for the following reasons. Clearly, the claim is true for j = 0. For $1 \leq j < t^{*}$, it suffices to show $b \text{-}time^{\gamma_{x,y}^{j}}(y, \bar{B}_{y,x}) > k \cdot \rho + w_{y,q_{j,k}}^{\gamma_{x,y}^{j}} + b \text{-}time^{\gamma_{x,y}^{j}}(q_{j,k}, \bar{B}_{q_{j,k},y})$ for all $1 \leq k \leq j$. As a consequence of $\gamma_{x,y}^{j}$ and $\gamma_{x,y}^{k}$, we have $b \text{-}time^{\gamma_{x,y}^{j}}(y, \bar{B}_{y,x}) \geq b \text{-}time^{\gamma_{x,y}^{k}}(y, \bar{B}_{y,x})$ and $w_{y,q_{j,k}}^{\gamma_{x,y}^{j}} + b \text{-}time^{\gamma_{x,y}^{j}}(q_{k,k}, \bar{B}_{q_{k,k},y})$. Besides, since $k < t^{*}$, we have $b \text{-}time^{\gamma_{x,y}^{k}}(y, \bar{B}_{y,x}) > k \cdot \rho + w_{y,q_{k,k}}^{\gamma_{x,y}^{k}} + b \text{-}time^{\gamma_{x,y}^{k}}(q_{k,k}, \bar{B}_{q_{k,k},y})$. Thus, the claim follows.

Suppose $q_{j,\ell} = u_{j+1}$, where $j < \ell \le h$. We first assume $\tau_j < \ell$. For $1 \le k \le h$ and $k \ne j+1$, since $w_{y,q_{j+1,k}}^{\gamma_{x,y}^{j+1}} = w_{y,q_{j+1,k}}^{\gamma_{x,y}^{j+1}}$ and $b_time^{\gamma_{x,y}^{j}}(q_{j+1,k}, \bar{B}_{q_{j+1,k},y}) = b_time^{\gamma_{x,y}^{j+1}}(q_{j+1,k}, \bar{B}_{q_{j+1,k},y})$, we have

$$\begin{split} k \cdot \rho + w_{y,q_{j+1,k}}^{\gamma_{x,y}^{j+1}} + b_time^{\gamma_{x,y}^{j+1}} \big(q_{j+1,k}, \bar{B}_{q_{j+1,k},y} \big) \\ &= k \cdot \rho + w_{y,q_{j+1,k}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}} \big(q_{j+1,k}, \bar{B}_{q_{j+1,k},y} \big) \\ &\leq \rho + \max\{ (k-1) \cdot \rho + w_{y,q_{j,k-1}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}} \big(q_{j,k-1}, \bar{B}_{q_{j,k-1},y} \big), \\ &\qquad \qquad (k \cdot \rho + w_{y,q_{j,k}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}} \big(q_{j,k}, \bar{B}_{q_{j,k},y} \big) \big) - \rho \} \quad (\because q_{j+1,k} \in \{q_{j,k-1}, q_{j,k}\}) \\ &\leq \rho + \big(\tau_{j} \cdot \rho + w_{y,q_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}} \big(q_{j,\tau_{j}}, \bar{B}_{q_{j,\tau_{j}},y} \big) \big) \\ &= (\tau_{j} + 1) \cdot \rho + w_{y,q_{j+1,\tau_{j}+1}}^{\gamma_{x,j}^{j+1}} + b_time^{\gamma_{x,y}^{j+1}} \big(q_{j+1,\tau_{j}+1}, \bar{B}_{q_{j+1,\tau_{j}+1},y} \big), \end{split}$$

where the last equality holds, because the claim implies $\tau_j + 1 > j + 1$ and hence $q_{j+1,\tau_j+1} = q_{j,\tau_j}$.

Consequently, we have $q_{j+1,\tau_{j+1}} \in \{q_{j+1,\tau_{j+1}}, q_{j+1,j+1}\}$. If $q_{j+1,\tau_{j+1}} = q_{j+1,\tau_{j+1}} \ (= q_{j,\tau_{j}})$, then $w_{y,q_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}} (q_{j,\tau_{j}}, \bar{B}_{q_{j,\tau_{j}},y}) = w_{y,q_{j+1,\tau_{j+1}}}^{\gamma_{x,y}^{j+1}} + b_time^{\gamma_{x,y}^{j+1}} (q_{j+1,\tau_{j+1}}, \bar{B}_{q_{j+1,\tau_{j+1}},y})$. If $q_{j+1,\tau_{j+1}} = q_{j+1,j+1}$, then

$$\begin{split} & w_{y,q_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(q_{j,\tau_{j}},\bar{B}_{q_{j,\tau_{j}},y}) \\ &= w_{y,q_{j+1,\tau_{j}+1}}^{\gamma_{x,y}^{j+1}} + b_time^{\gamma_{x,y}^{j+1}}(q_{j+1,\tau_{j}+1},\bar{B}_{q_{j+1,\tau_{j}+1},y}) \\ &\leq w_{y,q_{j+1,\tau_{j}+1}}^{\gamma_{x,y}^{j+1}} + b_time^{\gamma_{x,y}^{j+1}}(q_{j+1,\tau_{j+1}},\bar{B}_{q_{j+1,\tau_{j+1}},y}) \\ & \text{(recall that } w_{y,q_{j+1,k}}^{\gamma_{x,y}^{j+1}} + b_time^{\gamma_{x,y}^{j+1}}(q_{j+1,k},\bar{B}_{q_{j+1,k},y}) \text{ is nonincreasing as} \\ & k \text{ increases from 1 to } h). \end{split}$$

Next we assume $\tau_j \geq \ell$. Similarly, we have $q_{j+1,\tau_{j+1}} \in \{q_{j+1,j+1}, q_{j+1,j+2}, \dots, q_{j+1,\ell}\} \cup \{q_{j+1,\tau_j}\}$, i.e., $\tau_{j+1} \leq \tau_j$. Since $b_time^{\gamma_{x,y}^j}(y, \bar{B}_{y,x}) \leq b_time^{\gamma_{x,y}^{j+1}}(y, \bar{B}_{y,x})$, we have $w_{y,q_{j,\tau_j}}^{\gamma_{x,y}^j} + b_time^{\gamma_{x,y}^j}(q_{j,\tau_j}, \bar{B}_{q_{j,\tau_j},y}) \leq w_{y,q_{j+1,\tau_{j+1}}}^{\gamma_{x,y}^{j+1}} + b_time^{\gamma_{x,y}^{j+1}}(q_{j+1,\tau_{j+1}}, \bar{B}_{q_{j+1,\tau_{j+1}},y})$.

Lemma 11 If $w_{y,\mu}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(\mu, B_{y,x}) < w_{y,q_{0,\tau_{0}}}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(q_{0,\tau_{0}}, \bar{B}_{q_{0,\tau_{0}},y})$, then $b_time^{\gamma_{x,y}^{\dot{j}}}(y,T) - b_time^{\gamma_{x,y}^{\dot{j}}}(y,\bar{B}_{y,x})$ is nonincreasing as j increases from 0 to t^{*} .

Proof. It is equivalent to show

$$b_time^{\gamma_{x,y}^{j-1}}(y,T) - b_time^{\gamma_{x,y}^{j-1}}(y,\bar{B}_{y,x}) \ge b_time^{\gamma_{x,y}^{j}}(y,T) - b_time^{\gamma_{x,y}^{j}}(y,\bar{B}_{y,x})$$

for $1 \leq j \leq t^*$. First, we define $\hat{Q}_j = (\hat{q}_{j,1}, \hat{q}_{j,2}, \dots, \hat{q}_{j,h+1})$ as follows. Initially, let \hat{Q}_0 be obtained from Q_0 by inserting μ between $q_{0,p-1}$ and $q_{0,p}$, where $w_{y,q_{0,p-1}}^{\gamma_{x,y}^0} + b_time^{\gamma_{x,y}^0}(q_{0,p-1}, \bar{B}_{q_{0,p-1},y}) \geq w_{y,\mu}^{\gamma_{x,y}^0} + b_time^{\gamma_{x,y}^0}(\mu, B_{y,x}) > w_{y,q_{0,p}}^{\gamma_{x,y}^0} + b_time^{\gamma_{x,y}^0}(q_{0,p}, \bar{B}_{q_{0,p},y}).$

Then, for $1 \leq j \leq t^*$, \hat{Q}_j is obtained from \hat{Q}_{j-1} by cyclically shifting $\hat{q}_{j-1,c}$, $\hat{q}_{j-1,c+1}$, ..., $\hat{q}_{j-1,\ell}$ one position toward the right, where $\hat{q}_{j-1,\ell} = u_j$ is assumed, c = j if $w_{y,\hat{q}_{j-1,\ell}}^{\gamma_{x,y}^j} + b \pm time^{\gamma_{x,y}^j} (\hat{q}_{j-1,\ell}, \bar{B}_{\hat{q}_{j-1,\ell},y}) \geq w_{y,\mu}^{\gamma_{x,y}^j} + b \pm time^{\gamma_{x,y}^j} (\mu, B_{y,x})$, and c = j+1 else. If we remove μ from \hat{Q}_j , then Q_j results. Since $w_{y,\hat{q}_{j,k}}^{\gamma_{x,y}^j} + b \pm time^{\gamma_{x,y}^j} (\hat{q}_{j,k}, \bar{B}_{\hat{q}_{j,k},y})$ is nonincreasing as k increases from 1 to k+1, Lemma 8 guarantees that \hat{Q}_j is an optimal sequence for k0 broadcast a message to k1 under k2, under k3.

Notice that we have

$$(\hat{q}_{i,1}, \hat{q}_{i,2}, \dots, \hat{q}_{i,\tau_i}) = (q_{i,1}, q_{i,2}, \dots, q_{i,\tau_i}), \tag{5}$$

because

$$w_{y,\mu}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(\mu, B_{y,x})$$

$$= w_{y,\mu}^{\tilde{s}(x)} + b_time^{\tilde{s}(x)}(\mu, B_{y,x}) \quad \text{(as a consequence of } \gamma_{x,y}^{j})$$

$$< w_{y,q_{0,\tau_{0}}}^{\gamma_{x,y}^{0}} + b_time^{\gamma_{x,y}^{0}}(q_{0,\tau_{0}}, \bar{B}_{q_{0,\tau_{0}},y}) \quad (\because \ddot{s}(x) = \gamma_{x,y}^{0})$$

$$\leq w_{y,q_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(q_{j,\tau_{j}}, \bar{B}_{q_{j,\tau_{j}},y}) \quad \text{(by Lemma 10)}.$$
(6)

As a consequence of (5), we have

$$b_time^{\gamma_{x,y}^{j}}(y,T) \geq \tau_{j} \cdot \rho + w_{y,\hat{q}_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(\hat{q}_{j,\tau_{j}}, \bar{B}_{\hat{q}_{j,\tau_{j}},y})$$
$$> k \cdot \rho + w_{y,\hat{q}_{j,k}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(\hat{q}_{j,k}, \bar{B}_{\hat{q}_{j,k},y})$$

for $1 \leq k < \tau_j$. Let \hat{q}_{j,ϱ_j} be the first vertex in \hat{Q}_j with $b_time^{\gamma_{x,y}^j}(y,T) = \varrho_j \cdot \rho + w_{y,\hat{q}_{j,\varrho_j}}^{\gamma_{x,y}^j} + b_time^{\gamma_{x,y}^j}(\hat{q}_{j,\varrho_j},\bar{B}_{\hat{q}_{j,\varrho_j},y})$. Clearly, we have $\varrho_j \geq \tau_j$. When $\varrho_j = \tau_j$, the lemma follows, because $b_time^{\gamma_{x,y}^j}(y,T) = b_time^{\gamma_{x,y}^j}(y,\bar{B}_{y,x})$ and $b_time^{\gamma_{x,y}^{j-1}}(y,T) \geq b_time^{\gamma_{x,y}^{j-1}}(y,\bar{B}_{y,x})$. In subsequent discussion, we assume $\varrho_j > \tau_j$.

Suppose $q_{j-1,\ell'}=u_j$, where $j\leq \ell'\leq h$. We first consider $\tau_{j-1}<\ell'$. Since $q_{j-1,\tau_{j-1}}=q_{j,\tau_{j-1}+1}$, we have

$$b_time^{\gamma_{x,y}^{j}}(y,\bar{B}_{y,x}) \geq (\tau_{j-1}+1) \cdot \rho + w_{y,q_{j,\tau_{j-1}+1}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(q_{j,\tau_{j-1}+1},\bar{B}_{q_{j,\tau_{j-1}+1},y})$$

$$= \rho + (\tau_{j-1} \cdot \rho + w_{y,q_{j-1,\tau_{j-1}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(q_{j-1,\tau_{j-1}},\bar{B}_{q_{j-1,\tau_{j-1}},y}))$$

$$= \rho + (\tau_{j-1} \cdot \rho + w_{y,q_{j-1,\tau_{j-1}}}^{\gamma_{x,y}^{j-1}} + b_time^{\gamma_{x,y}^{j-1}}(q_{j-1,\tau_{j-1}},\bar{B}_{q_{j-1,\tau_{j-1}},y}))$$

$$= \rho + b_time^{\gamma_{x,y}^{j-1}}(y,\bar{B}_{y,x}). \tag{7}$$

Since $\varrho_j > \tau_j \geq j$, we have $\hat{q}_{j,\varrho_j} \neq q_{j,j} = u_j$ (as a consequence of (5)), which implies $\hat{q}_{j,\varrho_j} \in \{\hat{q}_{j-1,\varrho_j-1}, \hat{q}_{j-1,\varrho_j}\}$. Hence, we have

$$b_time^{\gamma_{x,y}^{j}}(y,T) = \varrho_{j} \cdot \rho + w_{y,\hat{q}_{j,\varrho_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(\hat{q}_{j,\varrho_{j}}, \bar{B}_{\hat{q}_{j,\varrho_{j}},y})$$

$$= \rho + \max\{(\varrho_{j} - 1) \cdot \rho + w_{y,\hat{q}_{j-1,\varrho_{j}-1}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(\hat{q}_{j-1,\varrho_{j}-1}, \bar{B}_{\hat{q}_{j-1,\varrho_{j}-1},y}),$$

$$(\varrho_{j} \cdot \rho + w_{y,\hat{q}_{j-1,\varrho_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(\hat{q}_{j-1,\varrho_{j}}, \bar{B}_{\hat{q}_{j-1,\varrho_{j}-1},y})) - \rho\}$$

$$= \rho + \max\{(\varrho_{j} - 1) \cdot \rho + w_{y,\hat{q}_{j-1,\varrho_{j}-1}}^{\gamma_{x,y}^{j-1}} + b_time^{\gamma_{x,y}^{j-1}}(\hat{q}_{j-1,\varrho_{j}-1}, \bar{B}_{\hat{q}_{j-1,\varrho_{j}-1},y}),$$

$$(\varrho_{j} \cdot \rho + w_{y,\hat{q}_{j-1,\varrho_{j}}}^{\gamma_{x,y}^{j-1}} + b_time^{\gamma_{x,y}^{j-1}}(\hat{q}_{j-1,\varrho_{j}}, \bar{B}_{\hat{q}_{j-1,\varrho_{j}},y})) - \rho\}$$

$$\leq \rho + b_time^{\gamma_{x,y}^{j-1}}(y,T). \tag{8}$$

Thus, the lemma follows, as a consequence of (7) and (8).

Next we consider $\tau_{j-1} \geq \ell'$. Suppose $\hat{q}_{j,\eta} = \mu$. We have $\varrho_j \geq \eta$, because for $1 \leq k < \eta$,

$$\begin{split} k \cdot \rho + w_{y,\hat{q}_{j,k}}^{\gamma_{x,y}^{j}} + b \text{-}time^{\gamma_{x,y}^{j}} (\hat{q}_{j,k}, \bar{B}_{\hat{q}_{j,k},y}) \\ &= k \cdot \rho + w_{y,q_{j,k}}^{\gamma_{x,y}^{j}} + b \text{-}time^{\gamma_{x,y}^{j}} (q_{j,k}, \bar{B}_{q_{j,k},y}) \\ &\leq \tau_{j} \cdot \rho + w_{y,q_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b \text{-}time^{\gamma_{x,y}^{j}} (q_{j,\tau_{j}}, \bar{B}_{q_{j,\tau_{j}},y}) \\ &= \tau_{j} \cdot \rho + w_{y,\hat{q}_{j,\tau_{j}}}^{\gamma_{x,y}^{j}} + b \text{-}time^{\gamma_{x,y}^{j}} (\hat{q}_{j,\tau_{j}}, \bar{B}_{\hat{q}_{j,\tau_{j}},y}) \\ &< \varrho_{j} \cdot \rho + w_{y,\hat{q}_{j,\varrho_{j}}}^{\gamma_{x,y}^{j}} + b \text{-}time^{\gamma_{x,y}^{j}} (\hat{q}_{j,\varrho_{j}}, \bar{B}_{\hat{q}_{j,\varrho_{j}},y}). \end{split}$$

Besides, $u_{j} (= q_{j-1,\ell'})$ precedes μ in \hat{Q}_{j-1} , because $w_{y,\mu}^{\gamma_{x,y}^{j-1}} + b_time^{\gamma_{x,y}^{j-1}} (\mu, B_{y,x}) < w_{y,q_{j-1},\tau_{j-1}}^{\gamma_{x,y}^{j-1}} + b_time^{\gamma_{x,y}^{j-1}} (q_{j-1,\ell'}, \bar{B}_{q_{j-1},\ell',j-1}) \le w_{y,q_{j-1},\ell'}^{\gamma_{x,y}^{j-1}} + b_time^{\gamma_{x,y}^{j-1}} (q_{j-1,\ell'}, \bar{B}_{q_{j-1},\ell',y})$, where the relation < holds with arguments similar to (6). So, we have $\hat{q}_{j,\varrho_{j}} = \hat{q}_{j-1,\varrho_{j}}$, and hence

$$\begin{array}{lll} b_time^{\gamma_{x,y}^{j}}(y,T) & = & \varrho_{j}\cdot\rho + w_{y,\hat{q}_{j,\varrho_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(\hat{q}_{j,\varrho_{j}},\bar{B}_{\hat{q}_{j,\varrho_{j}},y}) \\ & = & \varrho_{j}\cdot\rho + w_{y,\hat{q}_{j-1,\varrho_{j}}}^{\gamma_{x,y}^{j}} + b_time^{\gamma_{x,y}^{j}}(\hat{q}_{j-1,\varrho_{j}},\bar{B}_{\hat{q}_{j-1,\varrho_{j}},y}) \end{array}$$

$$= \varrho_{j} \cdot \rho + w_{y,\hat{q}_{j-1},\varrho_{j}}^{\gamma_{x,y}^{j-1}} + b time^{\gamma_{x,y}^{j-1}} (\hat{q}_{j-1,\varrho_{j}}, \bar{B}_{\hat{q}_{j-1},\varrho_{j}}, y)$$

$$\leq b time^{\gamma_{x,y}^{j-1}} (y, T).$$

On the other hand, since $b_time^{\gamma_{x,y}^j}(y,T) \ge b_time^{\gamma_{x,y}^{j-1}}(y,T)$, we have $b_time^{\gamma_{x,y}^j}(y,T) = b_time^{\gamma_{x,y}^{j-1}}(y,T)$. The lemma then follows, because $b_time^{\gamma_{x,y}^j}(y,\bar{B}_{y,x}) \ge b_time^{\gamma_{x,y}^{j-1}}(y,\bar{B}_{y,x})$.

Lemma 12 If $w_{y,\mu}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(\mu, B_{y,x}) < w_{y,q_0,\tau_0}^{\ddot{s}(x)} + b_time^{\ddot{s}(x)}(q_{0,\tau_0}, \bar{B}_{q_0,\tau_0,y})$, then for $0 \leq j \leq t^*$, $\gamma_{x,y}^j$ is a worst-case scenario of T with respect to x. Besides, we have $\max_{x} T(x) = d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\gamma_{x,y}^j} + b_time^{\gamma_{x,y}^j}(y, \bar{B}_{y,x}) - b_time^{\gamma_{x,y}^j}(y, T)$.

Proof. This lemma can be proved by induction on j. Recall that we have $\gamma_{x,y}^0 = \ddot{s}(x)$, where y is a prime broadcast center of T under $\ddot{s}(x)$. The lemma holds for j = 0, as a consequence of Lemma 4. Suppose that the lemma holds for $j = k \geq 0$. When j = k + 1, we have

$$r_{x,y}^{\gamma_{x,y}^{k+1}} = b_time^{\gamma_{x,y}^{k+1}}(x,T) - b_time^{\gamma_{x,y}^{k+1}}(y,T)$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\gamma_{x,y}^{k+1}} + b_time^{\gamma_{x,y}^{k+1}}(y,\bar{B}_{y,x}) - b_time^{\gamma_{x,y}^{k+1}}(y,T) \quad \text{(by Lemma 5)}$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\gamma_{x,y}^{k+1}} + b_time^{\gamma_{x,y}^{k}}(y,\bar{B}_{y,x}) - b_time^{\gamma_{x,y}^{k}}(y,T) \quad \text{(by Lemma 11)}$$

$$= d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\gamma_{x,y}^{k}} + b_time^{\gamma_{x,y}^{k}}(y,\bar{B}_{y,x}) - b_time^{\gamma_{x,y}^{k}}(y,T)$$

$$= max_r(x) \quad \text{(by the induction hypothesis)}.$$

Now it is ready to prove Fact 4. First observe that Fact 4 and Lemma 12 have the same sufficient condition, but different necessary conditions. In the following, we show that the necessary condition (i.e., $\beta_{x,y}^{t^*}$ is a worst-case scenario) of Fact 4 is derivable from the necessary conditions (i.e., $\gamma_{x,y}^{t^*}$ is a worst-case scenario and $\max_x r(x) = d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\gamma_{x,y}^{t^*}} + b_t time^{\gamma_{x,y}^{t^*}}(y, \bar{B}_{y,x}) - b_t time^{\gamma_{x,y}^{t^*}}(y, T)$) of Lemma 12. It suffices to show $r_{x,y}^{\beta_{x,y}^{t^*}} \geq \max_x r(x)$, in order to show $\beta_{x,y}^{t^*}$ a worst-case scenario.

Let $Q'' = (q''_1, q''_2, \dots, q''_h)$ be an arrangement of u_1, u_2, \dots, u_h , which has $q''_k = u_k$ $(= q_{t^*,k})$ for $1 \le k \le t^*$ and $w^{\beta^{t^*}_{x,y}}_{y,q''_k} + b_time^{\beta^{t^*}_{x,y}}(q''_k, \bar{B}_{q''_k,y})$ nonincreasing as k increases from $t^* + 1$ to h. Since $\gamma^{t^*}_{x,y}$ and $\beta^{t^*}_{x,y}$ differ in the weights of edges in $\bigcup_{t^* < k \le h} (\{(y, u_k)\} + E(\bar{B}_{u_k,y})), w^{\beta^{t^*}_{x,y}}_{y,q''_k} + b_time^{\beta^{t^*}_{x,y}}(q''_k, \bar{B}_{q''_k,y})$ is also nonincreasing as k increases from 1 to t^*

(refer to Figure 6). Actually, it is nonincreasing as k increases from 1 to h, because $w_{y,q''_{t*}}^{\beta^{t*}_{x,y}} + b_time^{\beta^{t*}_{x,y}}(q''_{t*}, \bar{B}_{q''_{t*},y}) \ge w_{y,q''_{t*}+1}^{\beta^{t*}_{x,y}} + b_time^{\beta^{t*}_{x,y}}(q''_{t*+1}, \bar{B}_{q''_{t*+1},y})$, as explained below.

Suppose $q''_{t^*+1} = q_{t^*,p}$, where $t^* . We have$

$$\begin{split} & w_{y,q_{t^*+1}^{t^*}}^{\beta_{x,y}^{t^*}} + b_time^{\beta_{x,y}^{t^*}}(q_{t^*+1}'', \bar{B}_{q_{t^*+1}',y}) \\ & \leq w_{y,q_{t^*,p}}^{\gamma_{x,y}^{t^*}} + b_time^{\gamma_{x,y}^{t^*}}(q_{t^*,p}, \bar{B}_{q_{t^*,p},y}) \quad \text{(refer to Figure 6)} \\ & \leq w_{y,q_{t^*,t^*}}^{\gamma_{x,y}^{t^*}} + b_time^{\gamma_{x,y}^{t^*}}(q_{t^*,t^*}, \bar{B}_{q_{t^*,t^*},y}) \\ & = w_{y,q_{t^*}'}^{\beta_{x,y}^{t^*}} + b_time^{\beta_{x,y}^{t^*}}(q_{t^*}', \bar{B}_{q_{t^*,y}'}). \end{split}$$

Then, according to Lemma 8, Q'' is an optimal sequence for y to broadcast a message to $N_{\bar{B}_{y,x}}(y)$ under $\beta_{x,y}^{t^*}$. So, we have

$$\begin{array}{ll} b_time^{\beta^{t^*}_{x,y}}(y,\bar{B}_{y,x}) & \geq t^* \cdot \rho + w^{\beta^{t^*}_{x,y}}_{y,q''^*_{t^*}} + b_time^{\beta^{t^*}_{x,y}}(q''_{t^*},\bar{B}_{q''_{t^*},y}) \\ & = t^* \cdot \rho + w^{\gamma^{t^*}_{x,y}}_{y,q_{t^*,t^*}} + b_time^{\gamma^{t^*}_{x,y}}(q_{t^*,t^*},\bar{B}_{q_{t^*,t^*},y}) \\ & = b_time^{\gamma^{t^*}_{x,y}}(y,\bar{B}_{y,x}). \end{array}$$

On the other hand, since $b_time^{\beta_{x,y}^{t^*}}(y, \bar{B}_{y,x}) \leq b_time^{\gamma_{x,y}^{t^*}}(y, \bar{B}_{y,x})$, we have $b_time^{\beta_{x,y}^{t^*}}(y, \bar{B}_{y,x}) = b_time^{\gamma_{x,y}^{t^*}}(y, \bar{B}_{y,x})$. Further, since $\tilde{w}_{x,y}^{\beta_{x,y}^{t^*}} = \tilde{w}_{x,y}^{\gamma_{x,y}^{t^*}}$ and $b_time^{\beta_{x,y}^{t^*}}(y, T) \leq b_time^{\gamma_{x,y}^{t^*}}(y, T)$ (refer to Figure 6), we have

$$r_{x,y}^{\beta_{x,y}^{t^*}} = b_time^{\beta_{x,y}^{t^*}}(x,T) - b_time^{\beta_{x,y}^{t^*}}(y,T)$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\beta_{x,y}^{t^*}} + b_time^{\beta_{x,y}^{t^*}}(y,\bar{B}_{y,x}) - b_time^{\beta_{x,y}^{t^*}}(y,T) \quad \text{(by Lemma 5)}$$

$$= d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\gamma_{x,y}^{t^*}} + b_time^{\gamma_{x,y}^{t^*}}(y,\bar{B}_{y,x}) - b_time^{\beta_{x,y}^{t^*}}(y,T)$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\gamma_{x,y}^{t^*}} + b_time^{\gamma_{x,y}^{t^*}}(y,\bar{B}_{y,x}) - b_time^{\gamma_{x,y}^{t^*}}(y,T)$$

$$= max_r(x) \quad \text{(by Lemma 12)}. \tag{9}$$

5 Time complexity

It is known that given a scenario $s \in C$ and a vertex $v \in V(T)$, B_Ctr^s and $b_time^s(v,T)$ can be determined in O(n) time [33]. Since there are at most n-1 scenarios in $\Psi_1(x) \cup \Psi_2(x) \cup \ldots \cup \Psi_n(x)$, it takes $O(n^2)$ time for Algorithm 2 to find $\ddot{s}(x)$. In this section, we show that the time complexity of Algorithm 2 can be reduced to $O(n \log \log n)$, provided an $O(n \log n)$ time preprocessing is made. Consequently, Algorithm 1 requires $O(n \log n) + O(n \log \log n) \cdot O(\log n) = O(n \log \log n)$ time. We assume below that x

is not a prime broadcast center of T under $\ddot{s}(x)$, i.e., $\ddot{s}(x) \in \Psi_1(x) \cup \Psi_2(x) \cup \ldots \cup \Psi_{n-1}(x)$ (refer to the first paragraph of Section 4). The following lemma is useful to find $\ddot{s}(x)$.

Lemma 13 Suppose $x \in V(T)$. If $d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\beta_{x,y}^{j}} + b_time^{\beta_{x,y}^{j}}(y, \bar{B}_{y,x}) - b_time^{\beta_{x,y}^{j}}(y, T)$ has maximum value as y = y' and j = j', then we have $\beta_{x,y'}^{j'} = \ddot{s}(x)$.

Proof. The notations μ , q_{0,τ_0} , h, and t^* used in this proof inherit from Fact 3 and Fact 4. First we claim that there exists $\beta^j_{x,y}$, satisfying $\beta^j_{x,y} = \ddot{s}(x)$ and $\max_x r(x) = d_{x,y} \cdot \rho + \tilde{w}^{\beta^j_{x,y}}_{x,y} + b_x time^{\beta^j_{x,y}}(y, \bar{B}_{y,x}) - b_x time^{\beta^j_{x,y}}(y, T)$, as explained below. If $w^{\ddot{s}(x)}_{y,\mu} + b_x time^{\ddot{s}(x)}(\mu, B_{y,x}) \geq w^{\ddot{s}(x)}_{y,q_0,\tau_0} + b_x time^{\ddot{s}(x)}(q_{0,\tau_0}, \bar{B}_{q_{0,\tau_0},y})$, then we have $r^{\alpha_{x,y}}_{x,y} \geq d_{x,y} \cdot \rho + \tilde{w}^{\alpha_{x,y}}_{x,y} + b_x time^{\alpha_{x,y}}(y, \bar{B}_{y,x}) - b_x time^{\alpha_{x,y}}(y, T) \geq r^{\ddot{s}(x)}_{x,y}$, as a consequence of (3) and (4). Also, since $r^{\ddot{s}(x)}_{x,y} = \max_x r(x) \geq r^{\alpha_{x,y}}_{x,y}$, we have $\max_x r(x) = r^{\alpha_{x,y}}_{x,y} = d_{x,y} \cdot \rho + \tilde{w}^{\alpha_{x,y}}_{x,y} + b_x time^{\alpha_{x,y}}(y, \bar{B}_{y,x}) - b_x time^{\alpha_{x,y}}(y, T)$ (recall that $\alpha_{x,y} = \beta^h_{x,y}$). If $w^{\ddot{s}(x)}_{y,\mu} + b_x time^{\ddot{s}(x)}(\mu, B_{y,x}) < w^{\ddot{s}(x)}_{y,q_{0,\tau_0}} + b_x time^{\ddot{s}(x)}(q_{0,\tau_0}, \bar{B}_{q_{0,\tau_0},y})$, then we have $\max_x r(x) = r^{\beta^t_{x,y}}_{x,y} = d_{x,y} \cdot \rho + \tilde{w}^{\beta^t_{x,y}}_{x,y} + b_x time^{\beta^t_{x,y}}(y, \bar{B}_{y,x}) - b_x time^{\beta^t_{x,y}}(y, T)$, similarly, as a consequence of (9).

It follows that we have

$$r_{x,y'}^{\beta_{x,y'}^{j'}} = b_time^{\beta_{x,y'}^{j'}}(x,T) - b_time^{\beta_{x,y'}^{j'}}(y',T)$$

$$\geq d_{x,y'} \cdot \rho + \tilde{w}_{x,y'}^{\beta_{x,y'}^{j'}} + b_time^{\beta_{x,y'}^{j'}}(y',\bar{B}_{y',x}) - b_time^{\beta_{x,y'}^{j'}}(y',T) \quad \text{(by Lemma 5)}$$

$$\geq d_{x,y} \cdot \rho + \tilde{w}_{x,y}^{\beta_{x,y}^{j}} + b_time^{\beta_{x,y}^{j}}(y,\bar{B}_{y,x}) - b_time^{\beta_{x,y}^{j}}(y,T)$$

$$= max_r(x),$$

which implies $\beta_{x,y'}^{j'} = \ddot{s}(x)$.

With Lemma 13, it suffices to find a scenario β_{x,v_i}^j that can maximize the value of $d_{x,v_i} \cdot \rho + \tilde{w}_{x,v_i}^{\beta_{x,v_i}^j} + b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x}) - b_time^{\beta_{x,v_i}^j}(v_i, T)$, in order to determine $\ddot{s}(x)$. By a depth-first traversal of T, d_{x,v_i} and $\tilde{w}_{x,v_i}^{\alpha_{x,v_i}}$ ($=\tilde{w}_{x,v_i}^{\beta_{x,v_i}^j}$ for all $1 \leq j \leq h_i$) for all $1 \leq i < n$ can be determined in O(n) time. On the other hand, as shown below, $b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x})$ (and $b_time^{\beta_{x,v_i}^j}(v_i, T)$ similarly) for all $1 \leq i < n$ and $1 \leq j \leq h_i$ can be determined in $O(n \log \log n)$ time. Therefore, $\ddot{s}(x)$ can be determined in $O(n \log \log n)$ time.

We suppose $N_{\bar{B}_{v_i,x}}(v_i) = \{u_{i,1}, u_{i,2}, \dots, u_{i,h_i}\}$ in the rest of this section. The following fact will be shown in Section 5.2.

Fact 5 With an $O(n \log n)$ -time preprocessing, b_time $\beta_{x,v_i}^j(u_{i,k}, \bar{B}_{u_{i,k},v_i})$ can be determined in constant time, where $1 \leq j \leq h_i$ and $1 \leq k \leq h_i$.

With Fact 5, $b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x})$ can be determined in additional $O(h_i)$ time, as described below. Suppose $s \in C$, and let bucket[1], bucket[2], ..., $bucket[h_i]$ be h_i buckets. We assume, without loss of generality, that $u_{i,1}$ can maximize the value of $w_{v_i,u_{i,k}}^s + b_time^s(u_{i,k}, \bar{B}_{u_{i,k},v_i})$. Like [33], we insert $u_{i,k}$ into $bucket[\ell]$, if $\ell - 1 \leq (w_{v_i,u_{i,1}}^s + b_time^s(u_{i,1}, \bar{B}_{u_{i,1},v_i})) - (w_{v_i,u_{i,k}}^s + b_time^s(u_{i,k}, \bar{B}_{u_{i,k},v_i})) < \ell$, where $1 \leq k \leq h_i$ and $1 \leq \ell \leq h_i$. Define $acc(\ell) = \sum_{1 \leq r \leq \ell} |bucket[r]|$, which is the total number of vertices contained in bucket[1], bucket[2], ..., $bucket[\ell]$, and $min_v(\ell) = min\{w_{v_i,u_{i,k}}^s + b_time^s(u_{i,k}, \bar{B}_{u_{i,k},v_i}) | u_{i,k} \in bucket[\ell]\}$. Previously, $acc(\ell)$ and $min_v(\ell)$ were successfully used in [33] to reduce the time requirement for determining an optimal broadcast sequence.

Suppose $u_{i,k} \in bucket[\ell]$, satisfying $min_v(\ell) = w_{v_i,u_{i,k}}^s + b_time^s(u_{i,k}, \bar{B}_{u_{i,k},v_i})$, where $1 \leq \ell \leq h_i$. Then, as a consequence of Lemma 8, $u_{i,k}$ is the $acc(\ell)$ -th vertex in some optimal sequence for v_i to broadcast a message to $N_{\bar{B}_{v_i,x}}(v_i)$. Also, for any other $u_{i,k'} \in bucket[\ell]$, we have $u_{i,k'}$ preceding $u_{i,k}$ in the optimal sequence, implying that the broadcast time (i.e., $acc(\ell) \cdot \rho + w_{v_i,u_{i,k}}^s + b_time^s(u_{i,k}, \bar{B}_{u_{i,k},v_i})$) induced by $\bar{B}_{u_{i,k},v_i}$ is at least as much as the broadcast time induced by $\bar{B}_{u_{i,k'},v_i}$. Thus, we have $b_time^s(v_i, \bar{B}_{v_i,x}) = \max\{min_v(\ell) + acc(\ell) \cdot \rho \mid 1 \leq \ell \leq h_i\}$, which can be computed in $O(h_i)$ time.

It follows that $b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x})$ for all $1 \leq j \leq h_i$ can be determined in additional $O(h_i^2)$ time. As elaborated below, the time complexity can be reduced to $O(h_i \log \log h_i)$ (hence, it takes $O(n \log \log n)$ time to determine $b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x})$ for all $1 \leq i < n$ and $1 \leq j \leq h_i$). The following fact will be shown in Section 5.2 as well.

Fact 6 With an $O(n \log n)$ -time preprocessing, a vertex ordering $(u_{i,1}, u_{i,2}, \ldots, u_{i,h_i})$ of $N_{\bar{B}_{v_i,x}}(v_i)$ can be determined in $O(h_i)$ time such that $w_{v_i,u_{i,k}}^{\beta_{x_i,v_i}^h} + b_time^{\beta_{x_i,v_i}^h}(u_{i,k}, \bar{B}_{u_{i,k},v_i})$ is nonincreasing as k increases from 1 to h_i .

With Fact 6, $(w_{v_i,u_{i,1}}^{\beta_{x,v_i}^h} + b_time^{\beta_{x,v_i}^{h_i}}(u_{i,1}, \bar{B}_{u_{i,1},v_i})) - (w_{v_i,u_{i,k}}^{\beta_{x,v_i}^h} + b_time^{\beta_{x,v_i}^{h_i}}(u_{i,k}, \bar{B}_{u_{i,k},v_i}))$ is nondecreasing as k increases from 1 to h_i . Under $\beta_{x,v_i}^{h_i}$, we have $u_{i,1} \in bucket[1]$. Moreover, there exists $1 \leq h_i' \leq h_i$ such that $u_{i,1}, u_{i,2}, \ldots, u_{i,h_i'}$ are contained in

buckets, while $u_{i,h'_i+1}, u_{i,h'_i+2}, \ldots, u_{i,h_i}$ are not contained in buckets. Recall that when $h'_i+1 \leq j \leq h_i$ and $1 \leq k \leq h'_i$, we have $w_{v_i,u_i,k}^{\beta_{h'_i}}+b_time^{\beta_{x,v_i}^{h'_i}}(u_{i,k}, \bar{B}_{u_{i,k},v_i}) = w_{v_i,u_{i,k}}^{\beta_{x,v_i}^j}+b_time^{\beta_{x,v_i}^{h'_i}}(u_{i,k}, \bar{B}_{u_{i,k},v_i})$ (refer to Figure 6), implying that $u_{i,k}$ is contained in the same bucket under $\beta_{x,v_i}^{h'_i}$ and β_{x,v_i}^j . Similarly, when $h'_i+1 \leq j \leq h_i$ and $h'_i+1 \leq k \leq h_i$, we have $w_{v_i,u_{i,k}}^{\beta_{x'_i}^h}+b_time^{\beta_{x,v_i}^h}(u_{i,k}, \bar{B}_{u_{i,k},v_i}) \leq w_{v_i,u_{i,k}}^{\beta_{x,v_i}^h}+b_time^{\beta_{x,v_i}^h}(u_{i,k}, \bar{B}_{u_{i,k},v_i}) \leq w_{v_i,u_{i,k}}^{\beta_{x'_i}^h}+b_time^{\beta_{x,v_i}^h}(u_{i,k}, \bar{B}_{u_{i,k},v_i})$ implying that $u_{i,k}$ is not contained in any bucket under $\beta_{x,v_i}^{h'_i}$ and β_{x,v_i}^j . Consequently, $\beta_{x,v_i}^{h'_i}$ and β_{x,v_i}^j can induce the same values of $min_v(\ell)$ and $acc(\ell)$ for all $1 \leq \ell \leq h_i$, i.e., $b_time^{\beta_{x,v_i}^{h'_i}}(v_i, \bar{B}_{v_i,x}) = b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x})$.

So, we only need to compute $b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x})$ for all $1 \leq j \leq h_i'$. Suppose $u_{i,j} \in bucket[\tau_j]$ under β_{x,v_i}^j . Let pred(j) (succ(j)) be a list of non-empty buckets such that $bucket[\ell]$ is included in pred(j) (succ(j)) if and only if $bucket[\ell]$ is not empty, $\ell < \tau_j$ ($\ell > \tau_j$), and $min_v(\ell) + acc(\ell) \cdot \rho \geq min_v(t) + acc(t) \cdot \rho$ for all $1 \leq t \leq \ell$ ($\ell \leq t \leq h_i$). Assume $pred(j) = (bucket[\pi_{j,1}], bucket[\pi_{j,2}], \ldots, bucket[\pi_{j,p_j}])$ and $succ(j) = (bucket[\sigma_{j,1}], bucket[\sigma_{j,2}], \ldots, bucket[\sigma_{j,2}], \ldots, bucket[\sigma_{j,q_j}])$, where $\pi_{j,1} < \pi_{j,2} < \ldots < \pi_{j,p_j}$ and $\sigma_{j,1} < \sigma_{j,2} < \ldots < \sigma_{j,q_j}$. Then, we have $b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x}) = \max\{min_v(\pi_{j,p_j}) + acc(\pi_{j,p_j}) \cdot \rho, min_v(\tau_j) + acc(\tau_j) \cdot \rho, min_v(\sigma_{j,1}) + acc(\sigma_{j,1}) \cdot \rho\}$. Hence, it suffices to find the values of $min_v(\ell) + acc(\ell) \cdot \rho$ for all $\ell \in \{\pi_{j,p_j}, \tau_j, \sigma_{j,1}\}$, in order to determine $b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x})$. The use of pred(j) and succ(j) (together with $acc(\ell)$ and $min_v(\ell)$) is crucial for reducing the overall time complexity to $O(h_i \log \log h_i)$.

First, with Fact 5, additional $O(h_i)$ time is enough to determine $b_time^{\beta_{x,v_i}^{h'_i}}(v_i, \bar{B}_{v_i,x})$. Under $\beta_{x,v_i}^{h'_i}$, it takes $O(h_i)$ time to find the buckets (and hence $\tau_{h'_i}$) where $u_{i,1}, u_{i,2}, \ldots, u_{i,h'_i}$ reside and determine the values of $min_v(\ell)$ and $acc(\ell)$ for all $1 \leq \ell \leq h_i$. Another $O(h_i)$ time is required to find $pred(h'_i)$ and $succ(h'_i)$ (and hence $\pi_{h'_i,p_{h'_i}}$ and $\sigma_{h'_i,1}$). Then, it requires $O(h_i \log \log h_i)$ time to determine $b_time^{\beta_{x,v_i}^j}(v_i, \bar{B}_{v_i,x})$, sequentially, for $j = h'_i - 1, h'_i - 2, \ldots, 1$, as explained below.

Suppose $u_{i,j+1} \in bucket[\hat{\ell}]$ under β_{x,v_i}^j . We first prove the following lemma.

Lemma 14 β_{x,v_i}^j and β_{x,v_i}^{j+1} induce the same bucket[ℓ] for all $\ell \in \{1, 2, ..., h_i\} - \{\tau_{j+1}, \hat{\ell}\}$. Proof. Recall that when $1 \leq j \leq h'_i - 1$, β_{x,v_i}^j and β_{x,v_i}^{j+1} differ in the weights of edges in $\{(v_i, u_{i,j+1})\} + E(\bar{B}_{u_{i,j+1},v_i})$ (refer to Figure 6). It implies that among $u_{i,1}, u_{i,2}, ..., u_{i,h'_i}$, only $u_{i,j+1}$ resides in different buckets under β_{x,v_i}^j and β_{x,v_i}^{j+1} . The lemma then follows, because $u_{i,j+1} \in bucket[\tau_{j+1}]$ under β_{x,v_i}^{j+1} .

The proof above also implies that $u_{i,j}$ resides in the same bucket under $\beta_{x,v_i}^j, \beta_{x,v_i}^{j+1}, \ldots, \beta_{x,v_i}^{h'_i}$. Hence, τ_j can be determined in constant time. With Fact 5, the value of $w_{v_i,u_{i,j+1}}^{\beta_{x,v_i}^j} + b_time^{\beta_{x,v_i}^j}(u_{i,j+1}, \bar{B}_{u_{i,j+1},v_i})$ can be obtained, and hence $bucket[\hat{\ell}]$ can be determined, also in constant time. According to Lemma 14, it takes constant time to obtain $min_v(1)$, $min_v(2), \ldots, min_v(h_i), |bucket[1]|, |bucket[2]|, \ldots, |bucket[h_i]|$ under β_{x,v_i}^j from $min_v(1)$, $min_v(2), \ldots, min_v(h_i), |bucket[1]|, |bucket[2]|, \ldots, |bucket[h_i]|$ under β_{x,v_i}^j .

Since $w_{v_i,u_{i,j}}^{\beta_{x_i,v_i}^h} + b_time^{\beta_{x_i,v_i}^{h'_i}}(u_{i,j}, \bar{B}_{u_{i,j},v_i}) \geq w_{v_i,u_{i,j+1}}^{\beta_{x_i,v_i}^h} + b_time^{\beta_{x_i,v_i}^{h'_i}}(u_{i,j+1}, \bar{B}_{u_{i,j+1},v_i}),$ $u_{i,j} \in bucket[\tau_j]$ under β_{x,v_i}^j and $\beta_{x,v_i}^{h'_i}$, and $u_{i,j+1} \in bucket[\tau_{j+1}]$ under β_{x,v_i}^{j+1} and $\beta_{x,v_i}^{h'_i}$, we have $\tau_j \leq \tau_{j+1}$. Also, since $w_{v_i,u_{i,j+1}}^{\beta_{x_i,v_i}^j} + b_time^{\beta_{x,v_i}^{j+1}}(u_{i,j+1}, \bar{B}_{u_{i,j+1},v_i}) \geq w_{v_i,u_{i,j+1}}^{\beta_{x_i,v_i}^j} + b_time^{\beta_{x,v_i}^j}(u_{i,j+1}, \bar{B}_{u_{i,j+1},v_i}) \geq w_{v_i,u_{i,j+1}}^{\beta_{x_i,v_i}^j} + b_time^{\beta_{x,v_i}^j}(u_{i,j+1}, \bar{B}_{u_{i,j+1},v_i})$, we have $\tau_{j+1} \leq \hat{\ell}$. So, by Lemma 14, β_{x,v_i}^j and β_{x,v_i}^{j+1} induce the same $bucket[\ell]$ for all $1 \leq \ell \leq \tau_j - 1$, i.e., the same $acc(\tau_j - 1)$. In this way, $acc(\tau_j - 1)$ remains the same under β_{x,v_i}^j , β_{x,v_i}^{j+1} , ..., $\beta_{x,v_i}^{h'_i}$, and hence $acc(\tau_j)$ (= $acc(\tau_j - 1) + |bucket[\tau_j]|$ under β_{x,v_i}^j can be determined in constant time. Therefore, it takes $O(h_i)$ time to determine the values of $min_v(\tau_j) + acc(\tau_j) \cdot \rho$ under β_{x,v_i}^j for all $1 \leq j \leq h'_i - 1$.

Also, as a consequence of $\tau_j \leq \tau_{j+1}$ and β_{x,v_i}^j , β_{x,v_i}^{j+1} inducing the same $bucket[\ell]$ for all $1 \leq \ell \leq \tau_j - 1$, pred(j) is a sublist of pred(j+1) for all $1 \leq j \leq h'_i - 1$. Since $\pi_{h'_i,p_{h'_i}}$ is available, $O(h_i)$ time is enough to determine π_{j,p_j} for all $1 \leq j \leq h'_i - 1$. Therefore, it takes $O(h_i)$ time to determine the values of $min_-v(\pi_{j,p_j}) + acc(\pi_{j,p_j}) \cdot \rho$ under β_{x,v_i}^j for all $1 \leq j \leq h'_i - 1$. The computation of $min_-v(\sigma_{j,1}) + acc(\sigma_{j,1}) \cdot \rho$ under β_{x,v_i}^j for all $1 \leq j \leq h'_i - 1$, which is detailed in Section 5.1, takes $O(h_i \log \log h_i)$ time.

5.1 Computation of $min_{-}v(\sigma_{j,1}) + acc(\sigma_{j,1}) \cdot \rho$ under β_{x,v_i}^j

Without loss of generality, we suppose $h'_i = h_i$. First we claim that $succ(j) = (bucket[\sigma_{j,1}], bucket[\sigma_{j,2}], \ldots, bucket[\sigma_{j,q_j}])$ for all $1 \leq j < h_i$ can be constructed in $O(h_i \log \log h_i)$ time, which will be verified later. It was shown earlier that the values of $min_v(\ell)$ under β^j_{x,v_i} for all $1 \leq \ell \leq h_i$ and $1 \leq j < h_i$ can be determined in $O(h_i)$ time. On the other hand, we show in the next paragraph that the values of $acc(\sigma_{j,1})$ under β^j_{x,v_i} for all $1 \leq j < h_i$

can be determined in additional $O(h_i)$ time.

Define $\Delta_{j,t} = (acc(\sigma_{j,t}) - acc(\sigma_{j,t-1}) \text{ under } \beta_{x,v_i}^j) - (acc(\sigma_{j,t}) - acc(\sigma_{j,t-1}) \text{ under } \beta_{x,v_i}^{h_i}),$ where $1 \leq t \leq q_j$ and $\sigma_{j,0}$ is set to τ_j . The values of $\Delta_{j,t}$ for all $1 \leq t \leq q_j$ can be obtained, while constructing succ(j). If succ(j) is empty, then $\sigma_{j,t}$ and $\Delta_{j,t}$ are not defined. When t = 1, we have

 $\Delta_{j,1} = (acc(\sigma_{j,1}) - acc(\tau_j) \text{ under } \beta_{x,v_i}^j) - (acc(\sigma_{j,1}) - acc(\tau_j) \text{ under } \beta_{x,v_i}^{h_i}).$ (10) Now that the values of $acc(\tau_j)$ under β_{x,v_i}^j for all $1 \leq j < h_i$ and $acc(\ell)$ under $\beta_{x,v_i}^{h_i}$ for all $1 \leq \ell \leq h_i$ can be determined in $O(h_i)$ time (shown earlier), the values of $acc(\sigma_{j,1})$ under β_{x,v_i}^j for all $1 \leq j < h_i$ can be determined in additional $O(h_i)$ time according to (10).

Next we show the construction of succ(j) and $\Delta_{j,t}$ $(1 \leq t \leq q_j)$ from succ(j+1) and $\Delta_{j+1,t}$ $(1 \leq t \leq q_{j+1})$, where $1 \leq j < h_i$. We use the van Emde Boas priority queue [34] to store the indices, i.e., $\sigma_{j,t}$'s, of succ(j) for all $1 \leq j \leq h_i$. Initially, succ(j) is empty for all $1 \leq j \leq h_i$ (and hence $\Delta_{j,t}$ is not defined for all $1 \leq t \leq q_j$). The construction comprises four sequential stages, i.e., Stage 1 to Stage 4, which can find all buckets $bucket[\ell] \in succ(j)$ for $\ell \in \{\ell + 1, \ell + 2, \dots, h_i\}, \{\ell\}, \{\tau_{j+1} + 1, \tau_{j+1} + 2, \dots, \ell - 1\},$ and $\{\tau_j + 1, \tau_j + 2, \dots, \tau_{j+1}\}$, respectively, where $\tau_{j+1} \leq \ell \leq h_i$. If $\ell = h_i$, then Stage 1 is omitted and the construction starts with Stage 2. If $\ell \leq \tau_{j+1} + 1$, then Stage 3 is omitted (Stage 2 is also omitted as $\ell = \tau_{j+1}$). If $\tau_{j+1} = \tau_j$, then Stage 4 is omitted.

As a consequence of Lemma 14, when $\hat{\ell} > \tau_{j+1}$, we have

$$acc(c) \text{ under } \beta_{x,v_i}^j = \begin{cases} acc(c) - 1 \text{ under } \beta_{x,v_i}^{j+1}, & \text{if } \tau_{j+1} \le c \le \hat{\ell} - 1; \\ acc(c) \text{ under } \beta_{x,v_i}^{j+1}, & \text{if } \hat{\ell} \le c \le h_i. \end{cases}$$

$$(11)$$

Stage 1. $\ell \in \{\hat{\ell}+1, \hat{\ell}+2, \ldots, h_i\} = L_1$. As a consequence of Lemma 14, the values of $\min_{v}(\ell)$ and $acc(\ell)$ remain the same under β_{x,v_i}^j and β_{x,v_i}^{j+1} , which implies $bucket[\ell] \in succ(j)$ if and only if $bucket[\ell] \in succ(j+1)$. Moreover, those buckets $bucket[\ell] \in succ(j)$, if they exist, constitute a sublist $(bucket[\sigma_{j+1,\ell}], bucket[\sigma_{j+1,\ell+1}], \ldots, bucket[\sigma_{j+1,q_{j+1}}])$ of succ(j+1), where $bucket[\sigma_{j+1,\ell}]$ is the bucket that immediately succeeds $bucket[\ell]$ in succ(j+1), i.e., $\sigma_{j+1,\ell} = \min\{\sigma_{j+1,\ell} \mid \sigma_{j+1,\ell} > \ell \text{ and } 0 \le t \le q_{j+1}\}$. If succ(j+1) is empty or $\sigma_{j+1,\ell} \le \ell$ for all $1 \le t \le q_{j+1}$, then succ(j) remains empty and $\Delta_{j,\ell}$ for all $1 \le t \le q_j$ remain not defined.

Otherwise, by the aid of the van Emde Boas priority queue, the value of $\sigma_{j+1,\hat{t}}$ can be determined in $O(\log\log h_i)$ time. Then, $bucket[\sigma_{j+1,\hat{t}}]$ can be found in succ(j+1) and hence the sublist of succ(j+1) can be added to succ(j) in constant time. That is, we have $(bucket[\sigma_{j,\hat{t}}], bucket[\sigma_{j,\hat{t}+1}], \ldots, bucket[\sigma_{j,q_j}]) = (bucket[\sigma_{j+1,\hat{t}}], bucket[\sigma_{j+1,\hat{t}+1}], \ldots, bucket[\sigma_{j+1,q_{j+1}}])$, where $q_j - \ddot{t} = q_{j+1} - \hat{t}$. Moreover, we have $acc(\sigma_{j,\hat{t}}), acc(\sigma_{j,\hat{t}+1}), \ldots, acc(\sigma_{j,q_j})$ under β_{x,v_i}^j equal to $acc(\sigma_{j+1,\hat{t}}), acc(\sigma_{j+1,\hat{t}+1}), \ldots, acc(\sigma_{j+1,q_{j+1}})$ under β_{x,v_i}^{j+1} , respectively, which implies $\Delta_{j,\hat{t}+1}, \Delta_{j,\hat{t}+2}, \ldots, \Delta_{j,q_j}$ equal to $\Delta_{j+1,\hat{t}+1}, \Delta_{j+1,\hat{t}+2}, \ldots, \Delta_{j+1,q_{j+1}}$, respectively. We also set $\Delta_{j,\hat{t}} = \Delta_{j+1,\hat{t}}$ temporarily. It may help the construction of succ(j) in subsequent stages.

Stage 2. $\ell \in \{\hat{\ell}\} = L_2$, where $\hat{\ell} > \tau_{j+1}$. We use $bucket[\sigma_{j+1,\check{\ell}}]$ to denote the bucket that immediately precedes $bucket[\hat{\ell}]$ in $\{bucket[\tau_{j+1}]\} \cup succ(j+1)$, i.e., $\sigma_{j+1,\check{\ell}} = \max\{\sigma_{j+1,t} \mid \sigma_{j+1,t} < \hat{\ell} \text{ and } 0 \leq t \leq q_{j+1}\}$, where $\sigma_{j+1,0} = \tau_{j+1}$. By the aid of the van Emde Boas priority queue, $\sigma_{j+1,\check{\ell}}$ can be found in $O(\log\log h_i)$ time. If succ(j) remains empty after Stage 1, then $bucket[\hat{\ell}+1]$, $bucket[\hat{\ell}+2]$, ..., $bucket[h_i]$ under β_{x,v_i}^j are all empty. Since $bucket[\hat{\ell}]$ is not empty under β_{x,v_i}^j , we have $\hat{\ell} = \sigma_{j,q_j}$ and add $bucket[\hat{\ell}]$ to succ(j). We set $\Delta_{j,q_j} = 1 + \Delta_{j+1,q_{j+1}}$ if $bucket[\hat{\ell}] \in succ(j+1)$, and $1 - (acc(\sigma_{j,q_j}) - acc(\sigma_{j+1,\check{\ell}})$ under $\beta_{x,v_i}^{h_i}$ if $bucket[\hat{\ell}] \not\in succ(j+1)$. Let $\nabla = (acc(\sigma_{j,q_j}) - acc(\sigma_{j+1,\check{\ell}})$ under $\beta_{x,v_i}^j) - (acc(\sigma_{j,q_j}) - acc(\sigma_{j+1,\check{\ell}})$ under $\beta_{x,v_i}^{h_i}$). For both cases, we have $\Delta_{j,q_j} = \nabla$, as explained below.

According to (11), we have $\nabla = 1 + (acc(\sigma_{j,q_j}) - acc(\sigma_{j+1,\check{t}})$ under $\beta_{x,v_i}^{j+1}) - (acc(\sigma_{j,q_j}) - acc(\sigma_{j+1,\check{t}}))$ under $\beta_{x,v_i}^{h_i}$. When $bucket[\hat{\ell}] \in succ(j+1)$, we have $(\sigma_{j,q_j} =) \hat{\ell} = \sigma_{j+1,q_{j+1}}$ and $\sigma_{j+1,\check{t}} = \sigma_{j+1,q_{j+1}-1}$, i.e., $\nabla = 1 + \Delta_{j+1,q_{j+1}}$. When $bucket[\hat{\ell}] \not\in succ(j+1)$, we have $\sigma_{j+1,\check{t}} = \tau_{j+1}$ if succ(j+1) is empty, and $\sigma_{j+1,q_{j+1}}$ otherwise. It follows that we have $bucket[\sigma_{j+1,\check{t}}+1]$, $bucket[\sigma_{j+1,\check{t}}+2]$, ..., $bucket[\hat{\ell}]$ under β_{x,v_i}^{j+1} all empty, and so $acc(\sigma_{j+1,\check{t}}) = acc(\hat{\ell})$ (= $acc(\sigma_{j,q_j})$) under β_{x,v_i}^{j+1} , i.e., $\nabla = 1 - (acc(\sigma_{j,q_j}) - acc(\sigma_{j+1,\check{t}}))$ under $\beta_{x,v_i}^{h_i}$. Also notice that the value of Δ_{j,q_j} above is temporary.

In the rest of Stage 2, we consider the situation that Stage 1 is carried out and succ(j) is not empty after Stage 1. Then we have $succ(j) = (bucket[\sigma_{j,\vec{t}}], bucket[\sigma_{j,\vec{t}+1}], \ldots, bucket[\sigma_{j,q_j}])$ after Stage 1. We claim $bucket[\hat{\ell}] \in succ(j)$ if and only if $min_v(\hat{\ell})$ under $\beta_{x,v_i}^j + acc(\sigma_{j+1,\hat{t}-1}) \cdot \rho$ under $\beta_{x,v_i}^{h_i} \geq min_v(\sigma_{j+1,\hat{t}})$ under $\beta_{x,v_i}^j + acc(\sigma_{j+1,\hat{t}}) \cdot \rho$ under

 $\beta_{x,v_i}^{h_i} + \Delta_{j+1,\hat{t}} \cdot \rho$, as explained below. It follows that whether $bucket[\hat{\ell}] \in succ(j)$ or not can be determined in constant time.

The inequality of the if-and-only-if statement can be written as $\min_{v}(\hat{\ell})$ under $\beta_{x,v_i}^j + acc(\sigma_{j+1,\hat{\ell}-1}) \cdot \rho$ under $\beta_{x,v_i}^{j+1} \geq \min_{v}(\sigma_{j+1,\hat{\ell}})$ under $\beta_{x,v_i}^j + acc(\sigma_{j+1,\hat{\ell}}) \cdot \rho$ under β_{x,v_i}^{j+1} . If $bucket[\hat{\ell}] \in succ(j+1)$, then we have $\hat{\ell} = \sigma_{j+1,\hat{\ell}-1}$. Since $bucket[\hat{\ell}]$ is not empty and $\hat{\ell} > \tau_{j+1} \ (\geq \tau_j)$, we have $bucket[\hat{\ell}] \in succ(j)$ if and only if $\min_{v}(\hat{\ell}) + acc(\hat{\ell}) \cdot \rho \geq \min_{v}(\sigma_{j+1,\hat{\ell}}) + acc(\sigma_{j+1,\hat{\ell}}) \cdot \rho$ under β_{x,v_i}^j . The latter inequality is equivalent to the inequality above, as a consequence of (11).

Then we assume $bucket[\hat{\ell}] \not\in succ(j+1)$, which implies $\hat{\ell} > \sigma_{j+1,\hat{t}-1}$. If $bucket[\hat{\ell}] \not\in succ(j)$, then we have $min_v(\hat{\ell}) + acc(\hat{\ell}) \cdot \rho < min_v(\sigma_{j+1,\hat{t}}) + acc(\sigma_{j+1,\hat{t}}) \cdot \rho$ under β_{x,v_i}^j , from which the inequality $min_v(\hat{\ell})$ under $\beta_{x,v_i}^j + acc(\sigma_{j+1,\hat{t}-1}) \cdot \rho$ under $\beta_{x,v_i}^{j+1} < min_v(\sigma_{j+1,\hat{t}})$ under $\beta_{x,v_i}^j + acc(\sigma_{j+1,\hat{t}}) \cdot \rho$ under β_{x,v_i}^{j+1} can be derived, as a consequence of (11) and $acc(\hat{\ell}) \geq acc(\sigma_{j+1,\hat{t}-1})$ under β_{x,v_i}^{j+1} . If $bucket[\hat{\ell}] \in succ(j)$, then we have $min_v(\hat{\ell}) + acc(\hat{\ell}) \cdot \rho \geq min_v(\sigma_{j+1,\hat{t}}) + acc(\sigma_{j+1,\hat{t}}) \cdot \rho$ under β_{x,v_i}^j , from which the inequality $min_v(\hat{\ell})$ under $\beta_{x,v_i}^j + acc(\sigma_{j+1,\hat{t}-1}) \cdot \rho$ under $\beta_{x,v_i}^{j+1} \geq min_v(\sigma_{j+1,\hat{t}})$ under $\beta_{x,v_i}^j + acc(\sigma_{j+1,\hat{t}}) \cdot \rho$ under β_{x,v_i}^{j+1} can be derived, because $acc(\hat{\ell}) = acc(\sigma_{j+1,\hat{t}-1})$ under β_{x,v_i}^j . The latter equality is verified below.

To prove $acc(\hat{\ell}) = acc(\sigma_{j+1,\hat{t}-1})$ under β_{x,v_i}^{j+1} , it suffices to show that $bucket[\sigma_{j+1,\hat{t}-1}+1]$, $bucket[\sigma_{j+1,\hat{t}-1}+2],\ldots,bucket[\hat{\ell}]$ under β_{x,v_i}^{j+1} are all empty. Suppose to the contrary that there exists $\sigma_{j+1,\hat{t}-1}+1 \leq \ell' \leq \hat{\ell}$ such that under β_{x,v_i}^{j+1} , $bucket[\ell']$ is not empty and $bucket[\ell'+1]$, $bucket[\ell'+2],\ldots,bucket[\hat{\ell}]$ are all empty. Then we have $acc(\ell')=acc(\hat{\ell})$ under β_{x,v_i}^{j+1} . As a consequence of (11), we have $min_v(\ell')+acc(\ell')\cdot\rho$ under $\beta_{x,v_i}^{j+1}\geq min_v(\hat{\ell})+acc(\hat{\ell})\cdot\rho$ under β_{x,v_i}^{j} . Further, since $min_v(\sigma_{j+1,\hat{t}})+acc(\sigma_{j+1,\hat{t}})\cdot\rho>min_v(\hat{\ell})+acc(\hat{\ell})\cdot\rho$ under β_{x,v_i}^{j+1} , we have $min_v(\sigma_{j+1,\hat{t}})+acc(\sigma_{j+1,\hat{t}})+acc(\sigma_{j+1,\hat{t}})\cdot\rho$ under β_{x,v_i}^{j} . Then, by Lemma 14, the inequality $min_v(\sigma_{j+1,\hat{t}})+acc(\sigma_{j+1,\hat{t}})\cdot\rho$ under $\beta_{x,v_i}^{j}>min_v(\hat{\ell})+acc(\hat{\ell})\cdot\rho$ under $\beta_{x,v_i}^{j}>min_v(\hat{\ell})+ac$

Next, the computations of succ(j) and $\Delta_{j,t}$ are described. If $bucket[\hat{\ell}] \in succ(j)$, then we add $bucket[\hat{\ell}]$ to the front of succ(j), i.e., $\sigma_{j,\tilde{t}-1} = \hat{\ell}$. As a consequence of (11), $\sigma_{j,\tilde{t}} = \sigma_{j+1,\hat{t}}$, and $acc(\sigma_{j,\tilde{t}-1}) = acc(\hat{\ell}) = acc(\sigma_{j+1,\hat{t}-1})$ under β_{x,v_i}^{j+1} , we have $\Delta_{j,\tilde{t}} = (acc(\sigma_{j,\tilde{t}}) - acc(\sigma_{j,\tilde{t}}))$

 $acc(\sigma_{j,\tilde{t}-1})$ under $\beta^{j}_{x,v_{i}}$) $-(acc(\sigma_{j,\tilde{t}}) - acc(\sigma_{j,\tilde{t}-1}))$ under $\beta^{h_{i}}_{x,v_{i}}$) $= \Delta_{j+1,\hat{t}} - (acc(\sigma_{j+1,\hat{t}-1}) - acc(\sigma_{j,\tilde{t}-1}))$ under $\beta^{h_{i}}_{x,v_{i}}$), which can be computed in constant time. In addition, we set $\Delta_{j,\tilde{t}-1} = 1 + \Delta_{j+1,\hat{t}-1}$ if $\hat{\ell} = \sigma_{j+1,\hat{t}-1}$, and $1 - (acc(\sigma_{j,\tilde{t}-1}) - acc(\sigma_{j+1,\tilde{t}}))$ under $\beta^{h_{i}}_{x,v_{i}}$ if $\hat{\ell} > \sigma_{j+1,\hat{t}-1}$. Let $\nabla' = (acc(\sigma_{j,\tilde{t}-1}) - acc(\sigma_{j+1,\tilde{t}}))$ under $\beta^{j}_{x,v_{i}} - (acc(\sigma_{j,\tilde{t}-1}) - acc(\sigma_{j+1,\tilde{t}}))$ under $\beta^{h_{i}}_{x,v_{i}}$. For both cases, we have $\Delta_{j,\tilde{t}-1} = \nabla'$, as a consequence of (11) and the fact that we have $\sigma_{j+1,\tilde{t}} = \sigma_{j+1,\hat{t}-2}$ if $\hat{\ell} = \sigma_{j+1,\hat{t}-1}$, and $\sigma_{j+1,\hat{t}-1}$ if $\hat{\ell} > \sigma_{j+1,\hat{t}-1}$.

If $bucket[\hat{\ell}] \not\in succ(j)$, then succ(j) remains unchanged. We set $\Delta_{j,\ddot{t}} = 1 + \Delta_{j+1,\hat{t}-1} + \Delta_{j+1,\hat{t}}$ if $\hat{\ell} = \sigma_{j+1,\hat{t}-1}$, and $1 + \Delta_{j+1,\hat{t}}$ if $\hat{\ell} > \sigma_{j+1,\hat{t}-1}$. Let $\nabla'' = (acc(\sigma_{j,\ddot{t}}) - acc(\sigma_{j+1,\breve{t}}))$ under $\beta_{x,v_i}^j - (acc(\sigma_{j,\ddot{t}}) - acc(\sigma_{j+1,\breve{t}}))$ under $\beta_{x,v_i}^{h_i}$. For both cases, we have $\Delta_{j,\ddot{t}} = \nabla''$, as a consequence of (11) and $\sigma_{j,\ddot{t}} = \sigma_{j+1,\hat{t}}$. The values of $\Delta_{j,\ddot{t}-1}$ (as $bucket[\hat{\ell}] \in succ(j)$) and $\Delta_{j,\ddot{t}}$ (as $bucket[\hat{\ell}] \not\in succ(j)$) above are temporary.

Recall that we use the van Emde Boas priority queue to store the indices, i.e., $\sigma_{j+1,t}$'s, of succ(j+1), and construct succ(j) from succ(j+1). We need to insert $\hat{\ell}$ into the queue if $bucket[\hat{\ell}] \not\in succ(j+1)$ and $bucket[\hat{\ell}] \in succ(j)$, and delete $\hat{\ell}$ from the queue if $bucket[\hat{\ell}] \in succ(j+1)$ and $bucket[\hat{\ell}] \not\in succ(j)$. Both take $O(\log \log h_i)$ time.

Stage 3. $\ell \in \{\tau_{j+1} + 1, \tau_{j+1} + 2, \dots, \hat{\ell} - 1\} = L_3$, where $\hat{\ell} > \tau_{j+1} + 1$. We first claim that if $bucket[\ell] \not\in succ(j+1)$, then $bucket[\ell] \not\in succ(j)$, as explained below. Suppose $bucket[\ell] \not\in succ(j+1)$. If $bucket[\ell]$ is empty under β_{x,v_i}^{j+1} , then $bucket[\ell]$ is also empty under β_{x,v_i}^j , according to Lemma 14. Otherwise, there exists some $bucket[\ell']$, where $\ell < \ell'$, satisfying $min_v(\ell) + acc(\ell) \cdot \rho < min_v(\ell') + acc(\ell') \cdot \rho$ under β_{x,v_i}^{j+1} . As a consequence of Lemma 14 and (11), we have $(min_v(\ell) + acc(\ell) \cdot \rho$ under $\beta_{x,v_i}^j) = (min_v(\ell) + (acc(\ell) \cdot \rho) + acc(\ell') \cdot \rho$ under $\beta_{x,v_i}^j)$. Similarly, we have $(min_v(\ell') + acc(\ell') \cdot \rho) + acc(\ell') \cdot \rho$ under $\beta_{x,v_i}^j) = (min_v(\ell') + acc(\ell') \cdot \rho$ under $\beta_{x,v_i}^j)$ if $\ell' \in L_1$, and $\sum (min_v(\ell') + (acc(\ell') - 1) \cdot \rho) + acc(\ell') \cdot \rho$ under $\beta_{x,v_i}^j)$ if $\ell' \in L_2 \cup L_3$. It follows that we have $min_v(\ell) + acc(\ell) \cdot \rho < min_v(\ell') + acc(\ell') \cdot \rho$ under β_{x,v_i}^j , i.e., $bucket[\ell] \not\in succ(j)$.

So, we only need to decide whether $bucket[\ell] \in succ(j)$ or not for each $bucket[\ell] \in succ(j+1)$. All $bucket[\ell]$'s in succ(j+1), if they exist, constitute a sublist, i.e., $(bucket[\sigma_{j+1,1}], bucket[\sigma_{j+1,2}], \ldots, bucket[\sigma_{j+1,\check{\ell}}])$, of succ(j+1). Suppose that $bucket[\sigma_{j,\eta}]$ is the first bucket in succ(j) after Stage 2. For $1 \le t \le \check{t}$, since $\sigma_{j+1,t} > \tau_{j+1}$ ($\ge \tau_j$), we

have $bucket[\sigma_{j+1,t}] \in succ(j)$ if and only if $min_{-}v(\sigma_{j+1,t}) + acc(\sigma_{j+1,t}) \cdot \rho \geq min_{-}v(\sigma_{j+1,\eta}) + acc(\sigma_{j+1,\eta}) \cdot \rho$ under β_{x,v_i}^j , as a consequence of Lemma 14. Let $\overline{\nabla}_t = (acc(\sigma_{j,\eta}) - acc(\sigma_{j+1,t}))$ under $\beta_{x,v_i}^j = (acc(\sigma_{j,\eta}) - acc(\sigma_{j+1,t}))$ under $\beta_{x,v_i}^j = (acc(\sigma_{j,\eta}) - acc(\sigma_{j+1,t}))$ under $\beta_{x,v_i}^i = (acc(\sigma_{$

Recall that $\Delta_{j,\eta}$ (= ∇ or ∇' or ∇'') was set to $(acc(\sigma_{j,\eta}) - acc(\sigma_{j+1,\check{t}}))$ under β_{x,v_i}^j ($acc(\sigma_{j,\eta}) - acc(\sigma_{j+1,\check{t}})$) under β_{x,v_i}^h after Stage 2, which is identical with $\overline{\nabla}_{\check{t}}$. As a consequence of (11), we have $\overline{\nabla}_{\check{t}-1} = (acc(\sigma_{j,\eta}) - acc(\sigma_{j+1,\check{t}}))$ under β_{x,v_i}^j ($acc(\sigma_{j,\eta}) - acc(\sigma_{j+1,\check{t}})$) under β_{x,v_i}^h ($acc(\sigma_{j+1,\check{t}}) - acc(\sigma_{j+1,\check{t}-1})$) under β_{x,v_i}^h ($acc(\sigma_{j+1,\check{t}-1}) - acc(\sigma_{j+1,\check{t}-1})$) und

If $bucket[\sigma_{j+1,t}] \not\in succ(j)$ for all $1 \leq t \leq \check{t}$, then succ(j) remains unchanged and we set $\Delta_{j,\eta} = \overline{\nabla}_1 + \Delta_{j+1,1}$. Otherwise, suppose $bucket[\sigma_{j+1,t'}] \in succ(j)$ and $bucket[\sigma_{j+1,t}] \not\in succ(j)$ for all $t' < t \leq \check{t}$. Then, as a consequence of Lemma 14, we have $bucket[\sigma_{j+1,1}]$, $bucket[\sigma_{j+1,2}], \ldots, bucket[\sigma_{j+1,t'-1}]$ all contained in succ(j). They can be added in constant time to the front of succ(j), i.e., $(bucket[\sigma_{j,t''}], bucket[\sigma_{j,t''+1}], \ldots, bucket[\sigma_{j,\eta-1}]) = (bucket[\sigma_{j+1,1}], bucket[\sigma_{j+1,2}], \ldots, bucket[\sigma_{j+1,t'}])$, where $\eta - t'' = t'$. Moreover, as a consequence of (11), we have $(\Delta_{j,t''+1}, \Delta_{j,t''+2}, \ldots, \Delta_{j,\eta-1}) = (\Delta_{j+1,2}, \Delta_{j+1,3}, \ldots, \Delta_{j+1,t'})$. We also set $\Delta_{j,t''} = \Delta_{j+1,1}$ and $\Delta_{j,\eta} = \overline{\nabla}_{t'}$. It follows that we have $\Delta_{j,\eta} = (acc(\sigma_{j,\eta}) - acc(\sigma_{j+1,t'}))$ under $\beta_{x,v_i}^i) - (acc(\sigma_{j,\eta}) - acc(\sigma_{j,\eta-1}))$ under $\beta_{x,v_i}^i) - (acc(\sigma_{j,\eta}) - acc(\sigma_{j,\eta-1}))$ under $\beta_{x,v_i}^i) - (acc(\sigma_{j,\eta}) - acc(\sigma_{j,\eta-1}))$ under $\beta_{x,v_i}^i)$.

Let $bucket[\sigma_{j,\eta'}]$ be the first bucket in the current succ(j), and $\nabla''' = (acc(\sigma_{j,\eta'}) - acc(\tau_{j+1}))$ under $\beta_{x,v_i}^j - (acc(\sigma_{j,\eta'}) - acc(\tau_{j+1}))$ under $\beta_{x,v_i}^{h_i}$, where $\eta' = \eta$ if $bucket[\sigma_{j+1,t}] \notin succ(j)$ for all $1 \le t \le \check{t}$, and $\eta' = t''$ else. As a consequence of (10) and (11), we have $\Delta_{j+1,1} = (acc(\sigma_{j+1,1}) - acc(\tau_{j+1}))$ under $\beta_{x,v_i}^j - (acc(\sigma_{j+1,1}) - acc(\tau_{j+1}))$ under $\beta_{x,v_i}^{h_i} - acc(\tau_{j+1})$ under β_{x,v_i}^{h

Recall that we have to delete $\sigma_{j+1,t}$ from the van Emde Boas priority queue, if

 $bucket[\sigma_{j+1,t}] \notin succ(j)$. If there are δ_j buckets in the sublist $(bucket[\sigma_{j+1,1}], bucket[\sigma_{j+1,2}], \ldots, bucket[\sigma_{j+1,\check{t}}])$ of succ(j+1) that are not in succ(j), then it requires $O(\delta_j \log \log h_i)$ time to delete these buckets from the queue.

Stage 4. $\ell \in \{\tau_j + 1, \tau_j + 2, \dots, \tau_{j+1}\} = L_4$. Since $u_{i,j}$ resides in the same bucket under β_{x,v_i}^j and $\beta_{x,v_i}^{h_i}$ (refer to the proof of Lemma 14), we have $u_{i,j} \in bucket[\tau_j]$ and $u_{i,j+1} \in bucket[\tau_{j+1}]$ under $\beta_{x,v_i}^{h_i}$. It means that $bucket[\ell']$ is empty under $\beta_{x,v_i}^{h_i}$ for all $\ell' \in L_4 - \{\tau_{j+1}\}$. Again, as a consequence of Lemma 14, β_{x,v_i}^j and $\beta_{x,v_i}^{h_i}$ induce the same $bucket[\ell']$ for all $\ell' \in L_4 - \{\tau_{j+1}\}$, implying that $bucket[\ell']$ is empty under β_{x,v_i}^j for all $\ell' \in L_4 - \{\tau_{j+1}\}$. So, we only need to decide whether $bucket[\tau_{j+1}] \in succ(j)$ or not. Also notice that we have $acc(\tau_{j+1}) = acc(\tau_j) + |bucket[\tau_{j+1}]|$ under β_{x,v_i}^j .

If $bucket[\tau_{j+1}]$ is empty under β_{x,v_i}^j , then we have $bucket[\tau_{j+1}] \not\in succ(j)$. In this case, $\Delta_{j,1}$ was previously set to $\Delta_{j,\bar{t}}$ (in Stage 1) or $\Delta_{j,\eta}$ (= ∇ or ∇' or ∇'' in Stage 2) or $\Delta_{j,\eta'}$ (= ∇''' in Stage 3), and these previous values are all equal to $(acc(\sigma_{j,1}) - acc(\tau_{j+1}))$ under $\beta_{x,v_i}^j - (acc(\sigma_{j,1}) - acc(\tau_{j+1}))$ under $\beta_{x,v_i}^k - (acc(\tau_{j+1}))$ is not empty under $\beta_{x,v_i}^j - (acc(\tau_{j+1}))$ is not empty under $\beta_{x,v_i}^j - (acc(\tau_{j+1}))$

If succ(j) is empty at the end of Stage 3, then $bucket[\tau_{j+1}+1]$, $bucket[\tau_{j+1}+2]$, ..., $bucket[h_i]$ are all empty under β_{x,v_i}^j . In this case, there is only one bucket, i.e., $bucket[\tau_{j+1}]$, in succ(j). We set $\Delta_{j,1} = |bucket[\tau_{j+1}]|$ under $\beta_{x,v_i}^j - (acc(\sigma_{j,1}) - acc(\tau_j))$ under $\beta_{x,v_i}^{h_i}$, which can be computed in constant time. The value is identical with the value of (10), because $\sigma_{j,1} = \tau_{j+1}$ and $acc(\tau_{j+1}) = acc(\tau_j) + |bucket[\tau_{j+1}]|$ under β_{x,v_i}^j .

Otherwise, let $bucket[\sigma_{j,\eta''}]$ be the first bucket in succ(j). In this case, $\Delta_{j,\eta''}$ was set to $\Delta_{j,\bar{t}}$ (in Stage 1) or $\Delta_{j,\eta}$ (= ∇ or ∇' or ∇'' in Stage 2) or $\Delta_{j,\eta'}$ (= ∇''' in Stage 3), and they are all equal to $(acc(\sigma_{j,\eta''}) - acc(\tau_{j+1}) \text{ under } \beta_{x,v_i}^j) - (acc(\sigma_{j,\eta''}) - acc(\tau_{j+1}) \text{ under } \beta_{x,v_i}^k)$. We have $bucket[\tau_{j+1}] \in succ(j)$ if and only if $min_v(\tau_{j+1}) + acc(\tau_{j+1}) \cdot \rho \geq min_v(\sigma_{j,\eta''}) + acc(\sigma_{j,\eta''}) \cdot \rho$ under β_{x,v_i}^j . The inequality can be written as $min_v(\tau_{j+1})$ under $\beta_{x,v_i}^j + acc(\tau_{j+1}) \cdot \rho$ under $\beta_{x,v_i}^{h_i} \geq \Delta_{j,\eta''} \cdot \rho + min_v(\sigma_{j,\eta''})$ under $\beta_{x,v_i}^j + acc(\sigma_{j,\eta''}) \cdot \rho$ under $\beta_{x,v_i}^{h_i}$, which can be determined in constant time. Further, if $bucket[\tau_{j+1}] \not\in succ(j)$,

then succ(j) remains unchanged, i.e., $\eta'' = 1$. The final value of $\Delta_{j,1}$ can be obtained in constant time by adding $(|bucket[\tau_{j+1}]| \text{ under } \beta_{x,v_i}^j - (acc(\tau_{j+1}) - acc(\tau_j) \text{ under } \beta_{x,v_i}^{h_i}))$ to the previous value of $\Delta_{j,1}$, because $acc(\tau_{j+1}) = acc(\tau_j) + |bucket[\tau_{j+1}]|$ under β_{x,v_i}^j .

If $bucket[\tau_{j+1}] \in succ(j)$, then we have $\sigma_{j,1} = \tau_{j+1}$ and $\eta'' = 2$. We add $bucket[\tau_{j+1}]$ to the front of succ(j), and set $\Delta_{j,1} = |bucket[\tau_{j+1}]|$ under $\beta_{x,v_i}^j - (acc(\sigma_{j,1}) - acc(\tau_j))$ under $\beta_{x,v_i}^{h_i}$, which is the same as the situation that succ(j) is empty at the end of Stage 3. The previous value of $\Delta_{j,2}$ is identical with its final value. If Stage 4 is omitted (as $\tau_{j+1} = \tau_j$), then succ(j) obtained after Stage 3 is what we desire and the previous value of $\Delta_{j,1}$ is identical with its final value.

The overall time complexity for constructing $succ(h_i)$, $succ(h_i-1)$, ..., succ(1) is computed as $\sum_{1 \leq j \leq h_i} O(\delta_j \log \log h_i) = O((\sum_{1 \leq j \leq h_i} \delta_j) \cdot \log \log h_i)$, where δ_j is the number of buckets in the sublist $(bucket[\sigma_{j+1,1}], bucket[\sigma_{j+1,2}], \ldots, bucket[\sigma_{j+1,\check{\ell}}])$ of succ(j+1) that are not in succ(j). Since $succ(h_i)$ is empty initially and at most two buckets, i.e., $bucket[\hat{\ell}]$ and $bucket[\tau_{j+1}]$, are newly added to each succ(j) (refer to Stage 2 and Stage 4), we have $\sum_{1 \leq j \leq h_i} \delta_j \leq 2(h_i - 1)$. Therefore, the overall time complexity is $O(h_i \log \log h_i)$.

5.2 Proofs of Fact 5 and Fact 6

We first prove Fact 5 below. Let s^+ (s^-) denote the scenario of T which has $w_{u,v}^{s^+} = w_{u,v}^+$ ($w_{u,v}^{s^-} = w_{u,v}^-$) for all edges $(u,v) \in E(T)$. Then, we have $b_time^{\beta_{x,v_i}^j}(u_{i,k}, \bar{B}_{u_{i,k},v_i}) = b_time^{s^+}(u_{i,k}, \bar{B}_{u_{i,k},v_i})$ if $k \leq j$, and $b_time^{s^-}(u_{i,k}, \bar{B}_{u_{i,k},v_i})$ else (refer to Figure 6). The required preprocessing is to determine $b_time^{s^+}(u, \bar{B}_{u,v})$ and $b_time^{s^-}(u, \bar{B}_{u,v})$ for all edges $(u,v) \in E(T)$. Clearly, after the preprocessing is done, each $b_time^{\beta_{x,v_i}^j}(u_{i,k}, \bar{B}_{u_{i,k},v_i})$ can be determined in constant time. In the following, we show that the preprocessing can be completed in $O(n \log n)$ time. Only the scenario s^+ is considered; the scenario s^- can be treated all the same.

As shown in [33], O(n) time is sufficient to find a broadcast center $\kappa \in B_Ctr^{s^+}$ of T and compute $b_time^{s^+}(u, \bar{B}_{u,\kappa})$ for all $u \in V(T) - \{\kappa\}$ under s^+ . Let $\eta(u)$ denote the neighbor of u in $B_{u,\kappa}$. With the work of [33], we only need to determine $b_time^{s^+}(\eta(u), \bar{B}_{\eta(u),u})$ for all $u \in V(T) - \{\kappa\}$, in order to complete the preprocessing, which depends on the

value of $d_{\kappa,u}$.

For the vertices u with $d_{\kappa,u}=1$, it takes $O(h\log h)$ time to determine an arrangement (q_1,q_2,\ldots,q_h) of $N_T(\kappa)$ such that $w_{q_j,\kappa}^{s^+}+b_time^{s^+}(q_j,\bar{B}_{q_j,\kappa})$ is nonincreasing as j increases from 1 to h, where $h=|N_T(\kappa)|$. We need to compute $b_time^{s^+}(\kappa,\bar{B}_{\kappa,q_j})$ for all $1\leq j\leq h$. When j=1, by Lemma 8, (q_2,q_3,\ldots,q_h) is an optimal sequence for κ to broadcast a message to $N_{\bar{B}_{\kappa,q_1}}(\kappa)$ under s^+ . Besides, we have $b_time^{s^+}(\kappa,\bar{B}_{\kappa,q_1})=\max\{(i-1)\cdot\rho+w_{\kappa,q_i}^{s^+}+b_time^{s^+}(q_i,\bar{B}_{q_i,\kappa})\mid 1\leq i\leq h\}$, computable in additional O(h) time.

When j > 1, q_1, q_2, \ldots, q_h are inserted into buckets bucket[1], bucket[2], \ldots , bucket[h] as before, and $min_v(\ell)$, $acc(\ell)$ for all $1 \le \ell \le h$ are determined. Both require additional O(h) time. Suppose $q_j \in bucket[z_j]$, and let $\lambda(q_j) = \min\{w_{\kappa,q_i}^{s^+} + b_time^{s^+}(q_i, \bar{B}_{q_i,\kappa}) \mid q_i \in bucket[z_j] - \{q_j\}\}$. Also, we define $\pi^-(\ell) = \max\{0, min_v(t) + acc(t) \cdot \rho \mid 1 \le t < \ell\}$ and $\pi^+(\ell') = \max\{0, min_v(t) + acc(t) \cdot \rho \mid \ell' < t \le h - 1\}$, where $1 \le \ell \le h$ and $1 \le \ell' \le h - 1$. Additional O(h) time is sufficient to compute $\lambda(q_j), \pi^-(\ell)$, and $\pi^+(\ell')$ for all $1 \le j \le h$, $1 \le \ell \le h$, and $1 \le \ell' \le h - 1$.

Notice that after inserting the vertices of $N_{\bar{B}\kappa,q_j}(\kappa)$ (= $N_T(\kappa)$ - $\{q_j\}$) into buckets $bucket'[1], bucket'[2], \ldots, bucket'[h-1]$, we have $b_time^{s^+}(\kappa, \bar{B}_{\kappa,q_j}) = \max\{min_v'(\ell) + acc'(\ell) \cdot \rho \mid 1 \leq \ell \leq h-1\}$ (refer to the two paragraphs immediately after Fact 5), where $min_v'(\ell)$ and $acc'(\ell)$ are defined on $bucket'[1], bucket'[2], \ldots, bucket'[h-1]$. Clearly, we have $min_v'(\ell) = min_v(\ell)$ if $\ell \neq z_j$; $min_v'(\ell) = \lambda(q_j)$ if $\ell = z_j$; $acc'(\ell) = acc(\ell)$ if $\ell < z_j$; $acc'(\ell) = acc(\ell) - 1$ if $\ell \geq z_j$. Hence, $b_time^{s^+}(\kappa, \bar{B}_{\kappa,q_j})$ can be obtained in constant time by finding the maximum of $\max\{min_v(\ell) + acc(\ell) \cdot \rho \mid 1 \leq \ell < z_j\}$ (= $\pi^-(z_j)$), $\lambda(q_j) + (acc(z_j) - 1) \cdot \rho$, and $\max\{min_v(\ell) + (acc(\ell) - 1) \cdot \rho \mid z_j < \ell \leq h-1\}$ (= $\pi^+(z_j) - 1$). Consequently, it takes O(h) time to determine $b_time^{s^+}(\kappa, \bar{B}_{\kappa,q_j})$ for all $1 < j \leq h$.

For the vertices u with $d_{\kappa,u}=2$, we need to compute $b_time^{s^+}(\eta(u), \bar{B}_{\eta(u),u})$, where $\eta(u) \in \{q_1, q_2, \dots, q_h\}$. Let us consider those vertices u with $\eta(u) = q_j$, where $1 \leq j \leq h$. Since $b_time^{s^+}(\kappa, \bar{B}_{\kappa,q_j})$ and $b_time^{s^+}(u, \bar{B}_{u,q_j})$ (= $b_time^{s^+}(u, \bar{B}_{u,\kappa})$) are available, we can determine an arrangement $(p_1, p_2, \dots, p_{h_j})$ of $N_T(q_j)$ such that $w_{p_k,q_j}^{s^+} + b_time^{s^+}(p_k, \bar{B}_{p_k,q_j})$ is nonincreasing as k increases from 1 to h_j in $O(h_j \log h_j)$ time, where $h_j = |N_T(q_j)|$. Then it takes additional $O(h_j)$ time to determine $b_time^{s^+}(q_j, \bar{B}_{q_j,p_k})$ for all $1 \leq k \leq h_j$.

It follows that $\sum_{1 \leq j \leq h} O(h_j \log h_j)$ time is sufficient to determine $b_time^{s^+}(\eta(u), \bar{B}_{\eta(u),u})$ for all u having $\eta(u) \in \{q_1, q_2, \dots, q_h\}$, where $h_j = |N_T(q_j)|$.

For the other vertices u (i.e, $d_{\kappa,u} > 2$), $b_time^{s^+}(\eta(u), \bar{B}_{\eta(u),u})$ can be obtained, similarly, in the sequence of $d_{\kappa,u} = 3, 4, \ldots$, by the aid of $b_time^{s^+}(\eta(\eta(u)), \bar{B}_{\eta(\eta(u)),\eta(u)})$ and $b_time^{s^+}(u, \bar{B}_{u,\eta(u)})$ (= $b_time^{s^+}(u, \bar{B}_{u,\kappa})$), where $b_time^{s^+}(\eta(\eta(u)), \bar{B}_{\eta(\eta(u)),\eta(u)})$ is available (because $d_{\kappa,\eta(u)} = d_{\kappa,u} - 1$). For each $\eta(u)$, $O(|N_T(\eta(u))|\log|N_T(\eta(u))|)$ time is spent to determine an arrangement $(g_1, g_2, \ldots, g_{|N_T(\eta(u))|})$ of $N_T(\eta(u))$ such that $w_{g_k,\eta(u)}^{s^+} + b_time^{s^+}(g_k, \bar{B}_{g_k,\eta(u)})$ is nonincreasing as k increases from 1 to $|N_T(\eta(u))|$. Then, $b_time^{s^+}(\eta(u), \bar{B}_{\eta(u),g_k})$ for all $1 \le k \le |N_T(\eta(u))|$ can be obtained in additional $O(|N_T(\eta(u))|)$ time.

According to the discussion above, the time required to determine $b_time^{s^+}(\eta(u), \bar{B}_{\eta(u),u})$ for all $u \in V(T) - \{\kappa\}$ is computed as the summation of $O(|N_T(v)|\log |N_T(v)|)$ for all $v \in \{\eta(u) \mid u \in V(T) - \{\kappa\}\}$, which is bounded by $O(n \log n)$. This completes the proof of Fact 5.

Next we present a proof of Fact 6. The required preprocessing is to determine a vertex ordering (u_1, u_2, \ldots, u_h) of $N_T(v)$ for each $v \in V(T)$ such that $w_{v,u_k}^{s^+} + b_time^{s^+}(u_k, \bar{B}_{u_k,v})$ is nonincreasing as k increases from 1 to h. Recall that in the proof of Fact 5 above, $b_time^{s^+}(u, \bar{B}_{u,v})$ for all edges $(u, v) \in E(T)$ are determined. It follows that the preprocessing can be done in $O(n \log n)$ time. Moreover, since $w_{v_i,u_{i,k}}^{\beta_{n_i}^+} + b_time^{\beta_{x,v_i}^{h_i}}(u_{i,k}, \bar{B}_{u_{i,k},v_i}) = w_{v_i,u_{i,k}}^{s^+} + b_time^{s^+}(u_{i,k}, \bar{B}_{u_{i,k},v_i})$, the vertex ordering, i.e., $(u_{i,1}, u_{i,2}, \ldots, u_{i,h_i})$, of $N_{\bar{B}_{v_i,x}}(v_i)$ can be obtained in $O(h_i)$ time by deleting the neighbor of v_i in $B_{v_i,x}$ from the vertex ordering of $N_T(v_i)$.

6 Conclusion

In this paper, a broadcasting problem with edge weight uncertainty was treated on heterogeneous tree networks under the postal model. All previous broadcasting problems (refer to [9–11, 13, 24, 31–33, 35]) assumed deterministic edge weights. One challenging problem encountered in this paper is to determine a worst-case scenario from an infinite number of candidates. To find the worst-case scenario, we first restricted it to a finite set

of scenarios, and then search the finite set for it by the prune-and-search strategy.

The following results were obtained in this paper. For each vertex x of a tree T, there are at most n-1 candidates for the worst-case scenario $\ddot{s}(x)$, where n is the number of vertices in T. Besides, $\ddot{s}(x)$ (and the maximum regret of x) and a minmax-regret broadcast center of T (and its maximum regret) can be found in $O(n \log \log n)$ time and $O(n \log n \log \log n)$ time, respectively. These results may be useful to the researchers who are interested in broadcasting problems.

It is still unknown if $O(n \log n \log \log n)$ time is the best result for finding a minmaxregret broadcast center of T. It is not easy to reduce the time complexity to $O(n \log n)$. Instead, it is more likely to derive a lower bound or make a limited improvement on the time complexity of the problem.

References

- [1] T. Albouy, D. Frey, M. Raynal, and F. Taïani. Asynchronous Byzantine reliable broadcast with a message adversary. *Theoret. Comput. Sci.*, 978:114110, 2023.
- [2] G. P. Arumugam, J. Augustine, M. J. Golin, and P. Srikanthan. Minmax regret k-sink location on a dynamic path network with uniform capacities. *Algorithmica*, 81(9):3534–3585, 2019.
- [3] I. Averbakh. On the complexity of a class of combinatorial optimization problems with uncertainty. *Mathematical Programming*, 90(2, Ser. A):263–272, 2001.
- [4] I. Averbakh. Minmax regret linear resource allocation problems. *Operations Research Letters*, 32(2):174–180, 2004.
- [5] I. Averbakh and S. Bereg. Facility location problems with uncertainty on the plane. *Discrete Optimization*, 2(1):3–34, 2005.
- [6] I. Averbakh and O. Berman. Minimax regret p-center location on a network with demand uncertainty. Location Science, 5(4):247–254, 1997.
- [7] I. Averbakh and O. Berman. Algorithms for the robust 1-center problem on a tree. European Journal of Operational Research, 123(2):292–302, 2000.
- [8] I. Averbakh and O. Berman. Minmax regret median location on a network under uncertainty. *INFORMS Journal on Computing*, 12(2):104–110, 2000.
- [9] A. Bar-Noy, S. Guha, J. Naor, and B. Schieber. Message multicasting in heterogeneous networks. *SIAM Journal on Computing*, 30(2):347–358, 2000.
- [10] A. Bar-Noy and S. Kipnis. Designing broadcasting algorithms in the postal model for message-passing systems. *Mathematical Systems Theory*, 27(5):431–452, 1994.
- [11] A. Bar-Noy and S. Kipnis. Multiple message broadcasting in the postal model. *Networks*, 29(1):1–10, 1997.

- [12] B. Bhattacharya, T. Kameda, and Z. Song. A linear time algorithm for computing minmax regret 1-median on a tree network. *Algorithmica*, 70(1):2–21, 2014.
- [13] L. Bortolussi, J. Hillston, and M. Loreti. Fluid approximation of broadcasting systems. *Theoret. Comput. Sci.*, 816:221–248, 2020.
- [14] R. E. Burkard and H. Dollani. A note on the robust 1-center problem on trees. *Annals of Operations Research*, 110(1–4):69–82, 2002.
- [15] E. Conde. On the complexity of the continuous unbounded knapsack problem with uncertain coefficients. *Operations Research Letters*, 33(5):481–485, 2005.
- [16] T. T. Dam, T. A. Ta, and T. Mai. Robust maximum capture facility location under random utility maximization models. *European J. Oper. Res.*, 310(3):1128–1150, 2023.
- [17] M. Drwal and J. Józefczyk. Robust min-max regret scheduling to minimize the weighted number of late jobs with interval processing times. *Ann. Oper. Res.*, 284(1):263–282, 2020.
- [18] D. Dvir and G. Y. Handler. The absolute center of a network. *Networks*, 43(2):109–118, 2004
- [19] A. J. Goldman. Optimal center location in simple networks. *Transportation Science*, 5(2):212–221, 1971.
- [20] H. A. Harutyunyan, A. L. Liestman, K. Makino, and T. C. Shermer. Nonadaptive broadcasting in trees. *Networks*, 57(2):157–168, 2011.
- [21] H. A. Harutyunyan, A. L. Liestman, and B. Shao. A linear algorithm for finding the k-broadcast center of a tree. Networks, 53(3):287–292, 2009.
- [22] C.-L. Hsu, W.-C. Lin, L. Duan, J.-R. Liao, C.-C. Wu, and J.-H. Chen. A robust two-machine flow-shop scheduling model with scenario-dependent processing times. *Discrete Dyn. Nat. Soc.*, 2020.
- [23] A. N. C. Kang and D. A. Ault. Some properties of a centroid of a free tree. *Information Processing Letters*, 4(1):18–20, 1975/76.
- [24] S. Khuller, Y.-A. Kim, and Y.-C. Wan. On generalized gossiping and broadcasting. *Journal of Algorithms*, 59(2):81–106, 2006.
- [25] P. Kouvelis and G. Yu. Robust discrete optimization and its applications, volume 14 of Non-convex Optimization and its Applications, pages xvi+356. Kluwer Academic Publishers, Dordrecht, 1997.
- [26] D. R. Kowalski, M. A. Mosteiro, and K. Zaki. Dynamic multiple-message broadcast: bounding throughput in the affectance model. *Theory Comput. Syst.*, 67(4):825–854, 2023.
- [27] M. Labbé, J.-F. Thisse, and R. E. Wendell. Sensitivity analysis in minisum facility location problems. *Operations Research*, 39(6):961–969, 1991.
- [28] W. Liao and Y. Fu. Min-max regret criterion-based robust model for the permutation flow-shop scheduling problem. *Eng. Optim.*, 52(4):687–700, 2020.
- [29] P. B. Mirchandani and A. R. Odoni. Locations of medians on stochastic networks. *Transportation Science*, 13(2):85–97, 1979.

- [30] Y. Nikulin. Robustness in combinatorial optimization and scheduling theory: An extended annotated bibliography. Technical report, Christian-Albrechts-Universitat zu Kiel, Institut für Betriebswirtschaftslehre, 01 2004.
- [31] D. Richards and A. L. Liestman. Generalizations of broadcasting and gossiping. *Networks*, 18(2):125–138, 1988.
- [32] P. J. Slater, E. J. Cockayne, and S. T. Hedetniemi. Information dissemination in trees. SIAM J. Comput., 10(4):692–701, 1981.
- [33] Y.-H. Su, C.-C. Lin, and D. T. Lee. Broadcasting in weighted trees under the postal model. *Theoret. Comput. Sci.*, 621:73–81, 2016.
- [34] P. van Emde Boas, R. Kaas, and E. Zijlstra. Design and implementation of an efficient priority queue. *Theory of Computing Systems*, 10(1):99–127, 1976.
- [35] M. Čevnik and J. Žerovnik. Broadcasting on cactus graphs. J. Comb. Optim., 33(1):292–316, 2017.
- [36] B.-F. Wang. Efficient algorithms for the minmax regret path center problem with length constraint on trees. *Theoret. Comput. Sci.*, 918:18–31, 2022.
- [37] B.-F. Wang, J.-H. Ye, and C.-Y. Li. An improved algorithm for the minmax regret path center problem on trees. *J. Comput. System Sci.*, 114:36–47, 2020.
- [38] J.-H. Ye and B.-F. Wang. On the minmax regret path median problem on trees. *J. Comput. System Sci.*, 81(7):1159–1170, 2015.
- [39] H.-I. Yu, T.-C. Lin, and B.-F. Wang. Improved algorithms for the minmax-regret 1-center and 1-median problems. *ACM Transactions on Algorithms*, 4(3):Art. 36, 27, 2008.