

Equivalent of Non-Trivial Riemann Zeros Condition in Matrix Representation and Particle Anti-Particle Symmetry Breaking

Chee Kian Yap
Queensland University of Technology

ABSTRACT. A Hermitian matrix can be formed for a large enough cutoff N on the number of terms in the Riemann zeta function. At zeros, the matrix trace equals the negative of the off-diagonal sum. Riemann's hypothesis is true if and only if:

$$\sum_i^N \frac{1}{i} = - \sum_{i \neq j} \frac{(-1)^{i+1} (-1)^{j+1}}{i^{\frac{1}{2}} j^{\frac{1}{2}} e^{\sqrt{-1} t \ln\left(\frac{i}{j}\right)}}$$

This equation suggests a relationship between the diagonal and off-diagonal elements of the matrix formed. By forming a matrix with Riemann non-trivial zero, the quantum field vacuum loop level diagram can be opened exactly by coupling the matrix with an anti-particle annihilation operator, which leads to asymmetry between matter and anti-matter.

1 Introduction

The Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. The hypothesis was proposed by Bernhard Riemann and is one of the oldest unsolved problems in pure mathematics. Riemann zeta function is a complex function that takes complex variable $s = \sigma + it$ and sends it to complex number. The Riemann zeta function is related to the distribution of prime numbers p for $\text{Real}(s) > 1$:

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_p \frac{1}{1-p^{-s}} \quad (1)$$

There is no bound for the counting of primes, so it is necessary to use divergence series to represent the collection of prime numbers. One such divergence series is Riemann zeta at $s = 1$; it is the pole of the function and corresponds to the Harmonic series.

According to Riemann's hypothesis, apart from trivial zeros, all fall on that vertical critical line with $\text{Real}(s) = \frac{1}{2}$ on the complex plane, where the sea level of the function landscape lies. At $\text{Real}(s) = \frac{1}{2}$ with large $t = \text{Im}(s)$, the Riemann function was shown to be bounded,^{1 2} guaranteeing that a finite value of an element of a meaningful matrix can be formed with the Riemann function near non-trivial zero, which is demonstrated in section 2. If we allow t to go to infinity, the number of zeros bounded by t will be infinite as well,³ the zeros go on and go, akin to the counting of an infinite number of primes. This study shows that Riemann zero is directly connected to the Harmonic series. Therefore, to justify why the real part of zero must be at $\frac{1}{2}$, it is to make the trace or diagonal sum of our proposed matrix to be a harmonic series given as:

$$\text{Tr}(\bar{\zeta}_i (s^*) \zeta_j (s^*)) = \sum_{n=1}^{\infty} \frac{1}{n} \quad (2)$$

the gamma function is intimately related to the Riemann zeta function, which is involved in calculating the vacuum loop diagram in quantum field theory. In the supersymmetry theory, it assumes matter and antimatter are treated equally. Matter is like the mirror image of anti-matter. They are not equivalent as left and right are switched, there is no continuous transformation between particle and anti-particle, and they have different creation/annihilation operators.

However, we find the world is made up mostly of matter, how the asymmetry came about is unknown. Here we show by coupling the matrix with an anti-particle annihilation operator in the loop integral, this chiral symmetry is broken in a vacuum.

2 Methods

For region in $0 < \text{Real}(s) < 1$, Riemann zeta function is given by:

$$\zeta(s) = \frac{1}{1-2^{1-s}} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s} \quad (3)$$

A matrix can be formed by taking each term in the function:

$$P_{ij}(s) = \bar{\zeta}_i(s) \zeta_j(s) \quad (4)$$

$P_{ij}(s)$ is analogous to density matrix in the formalism of quantum mechanics, it is also a Hermitian matrix. The bar in Eq. (4) indicates complex conjugation. Suppose the Riemann hypothesis is true for large enough N , at location of zeros:

$$\sum_i^N \zeta_i \left(s = \frac{1}{2} + \sqrt{-1}t \right) \cong 0 \quad (5)$$

COROLLARY 1 *Assume the Riemann Hypothesis. Then at zero*

$$\sum_{i,j}^N P_{ij} \left(s = \frac{1}{2} + \sqrt{-1}t \right) \cong 0 \quad (6)$$

COROLLARY 2 *Assume the Riemann Hypothesis. Then at zero*

$$\text{Tr}(P_{ij}) = \sum_i^N \frac{1}{i} = - \sum_{i \neq j} \frac{(-1)^{i+1} (-1)^{j+1}}{i^{\frac{1}{2}} j^{\frac{1}{2}} e^{\sqrt{-1}t \ln\left(\frac{j}{i}\right)}} \quad (7)$$

Corollary 1 is true by self-evident as it involves the multiplication of Riemann zero at each row i .

We shall prove the corollary 2 instead in the following.

Proof.

We find the trace of the matrix contains Harmonic series in the following:

$$\begin{aligned} \text{Tr}(P_{ij}) &= \sum_i^N \frac{1}{\left(1-2^{1-\frac{1}{2}-\sqrt{-1}t}\right)^* \left(i^{\frac{1}{2}+\sqrt{-1}t}\right)^*} \cdot \frac{1}{1-2^{1-\frac{1}{2}-\sqrt{-1}t}} \frac{(-1)^{i+1}}{i^{\frac{1}{2}+\sqrt{-1}t}} \\ &= \sum_i^N \frac{1}{\left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)^* \left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)} \frac{1}{ie^{(-\sqrt{-1}t+\sqrt{-1}t)\ln i}} \end{aligned} \quad (8)$$

$$\text{Tr}(P_{ij}) = \sum_i^N \frac{1}{\left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)^* \left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)} \frac{1}{i} \quad (9)$$

For the off-diagonal sum:

$$\begin{aligned} \sum_{i \neq j}^N P_{ij} &= \sum_{i \neq j} \frac{1}{\left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)^* \left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)} \frac{(-1)^{i+1}(-1)^{j+1}}{i^{\frac{1}{2}} j^{\frac{1}{2}} e^{-\sqrt{-1}t \ln i} e^{\sqrt{-1}t \ln j}} \\ \sum_{i \neq j}^N P_{ij} &= \sum_{i \neq j} \frac{1}{\left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)^* \left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)} \frac{(-1)^{i+1}(-1)^{j+1}}{i^{\frac{1}{2}} j^{\frac{1}{2}} e^{\sqrt{-1}t \ln \left(\frac{j}{i}\right)}} \end{aligned} \quad (10)$$

Because of Corollary 1:

$$\text{Tr}(P_{ij}) = -\sum_{i \neq j} P_{ij} \quad (11)$$

$$\begin{aligned} \sum_i^N \frac{1}{\left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)^* \left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)} \frac{1}{i} &= -\sum_{i \neq j} \frac{1}{\left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)^* \left(1-2^{\frac{1}{2}-\sqrt{-1}t}\right)} \frac{(-1)^{i+1}(-1)^{j+1}}{i^{\frac{1}{2}} j^{\frac{1}{2}} e^{\sqrt{-1}t \ln \left(\frac{j}{i}\right)}} \\ \sum_i^N \frac{1}{i} &= -\sum_{i \neq j} \frac{(-1)^{i+1}(-1)^{j+1}}{i^{\frac{1}{2}} j^{\frac{1}{2}} e^{\sqrt{-1}t \ln \left(\frac{j}{i}\right)}} \end{aligned} \quad (12)$$

Proved.

3 Riemann Matrix Coupling with Annihilation Operator

For time development $\Delta t = x_0 - y_0, x_0 > y_0$, The propagator or Green function for a Klein-Gordon particle is given as:

$$\int \frac{d^4 p}{(2\pi)^4} \frac{G(\Delta t; x-y)}{p^2 - m^2} \quad (13)$$

It has an incoming leg of y and an outgoing leg of x , p is four-momentum, and the propagator is a bare propagator that does not involve interactions of any form. Feynman diagrams without legs are possible with virtual particles. Virtual particles appear and annihilate in pairs of particles and anti-particles in a quantum vacuum; the ground state of quantum fields is empty and void. There are no tree-level diagrams in a vacuum when no operator is acting on a vacuum; only virtual

particles/quanta contribute to vacuum amplitudes in the form of loop integral; loop integrals are also called bubble Feynman diagrams. The first order one loop integral of the vacuum diagram is given as:

$$\int \frac{d^4p}{(2\pi)^4} a_y(p) \frac{G(\Delta t; y-x)}{p^2-m^2} a_x^\dagger(p) a_x(p) \frac{G(\Delta t; x-y)}{p^2-m^2} a_y^\dagger(p) \quad (14)$$

with $a_x^\dagger(p)$, $a_y^\dagger(p)$ are creation operators acting at space points x and y , respectively. By quantum relation:

$$a_x^\dagger(p) a_x(p) = 1 \quad (15)$$

$$\int \frac{d^4p}{(2\pi)^4} a_y(p) \frac{G(\Delta t; y-x)}{p^2-m^2} \frac{G(\Delta t; x-y)}{p^2-m^2} a_y^\dagger(p) \quad (16)$$

It involved joining the end vertex of a particle with the start vertex of the anti-particle propagator and vice versa so that the legs are cancelled, and given as:

$$\int \frac{d^4p}{(2\pi)^4} a_y(p) \frac{1}{p^2-m^2} \frac{1}{p^2-m^2} a_y^\dagger(p) \quad (17)$$

The formula⁴ can be applied involving gamma function:

$$= \frac{2\pi^2}{\Gamma(2)} \int_0^\infty dp p^3 a_y(p) \frac{1}{p^2-m^2} \frac{1}{p^2-m^2} a_y^\dagger(p) \quad (18)$$

Relation of gamma function with Riemann zeta function ζ provided by Hadamard gives:

$$= \frac{2\zeta(2)e^{-2\log(2\pi)+2+\gamma}}{\prod_\rho (1-\frac{2}{\rho})e^{\frac{2}{\rho}}} 2\pi^2 \int_0^\infty dp p^3 a_y(p) \frac{1}{p^2-m^2} \frac{1}{p^2-m^2} a_y^\dagger(p) \quad (19)$$

$$= Ae^\gamma \int_0^\infty dp p^3 a_y(p) \frac{1}{p^2-m^2} \frac{1}{p^2-m^2} a_y^\dagger(p) \quad (20)$$

Here, γ is Euler-Mascheroni constant, the expression can be further derived as:

$$B \int_0^\infty dp p^3 e^{\sum_{k=1}^\infty \frac{1}{k}} a_y(p) \frac{1}{p^2-m^2} \frac{1}{p^2-m^2} a_y^\dagger(p) \quad (21)$$

$$= B \int_0^\infty dp p^3 e^{\sum_{k=1}^\infty \frac{1}{k}} a_y(p) \frac{G(\Delta t; y-x)}{p^2-m^2} \frac{G(\Delta t; x-y)}{p^2-m^2} a_y^\dagger(p) \quad (22)$$

$$= B \int_0^\infty dpp^3 e^{\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*))} a_y(p) \frac{G(\Delta t; y-x)}{p^2-m^2} \frac{G(\Delta t; x-y)}{p^2-m^2} a^\dagger_y(p) \quad (23)$$

The state of anti-particle can be expressed as:

$$\langle p| = a_y(p) \frac{G(\Delta t; y-x)}{p^2-m^2} \quad (24)$$

$$\langle p|a^\dagger(p) = \langle 0| \quad (25)$$

$$\langle p|a^\dagger(p)a^\dagger(p) = \langle 0|a^\dagger(p) = 0 \quad (26)$$

where $a^\dagger(p)$ is the annihilation operator of an anti-particle in momentum space. By coupling the Riemann matrix with the anti-particle, the annihilation operator is given as:

$$\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*)) \rightarrow \text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*))a^\dagger(p) \quad (27)$$

The transformation acting as the ruling out of Riemann zeros, we have,

$$\langle p|e^{\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*))a^\dagger(p)} = \langle p| + \langle p|\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*))a^\dagger(p) + 0 \quad (28)$$

$$\langle p|e^{\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*))a^\dagger(p)} = \langle p| + \langle 0|\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*)) \quad (29)$$

The vacuum diagram reads, after the introduction of the annihilation operator as:

$$B \int_0^\infty dpp^3 a_y(p) \frac{G(\Delta t; y-x)}{p^2-m^2} e^{\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*))a^\dagger(p)} \frac{G(\Delta t; x-y)}{p^2-m^2} a^\dagger_y(p) \quad (30)$$

The introduced annihilation operator in the integral annihilates all copies of anti-particle distributed over the momentum space p and evaluated as:

$$B \int_0^\infty dpp^3 a_y(p) \frac{G(\Delta t; y-x)}{p^2-m^2} e^{\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*))a^\dagger(p)} \frac{G(\Delta t; x-y)}{p^2-m^2} a^\dagger_y(p) \quad (31)$$

$$= B \int_0^\infty dpp^3 a_y(p) \frac{G(\Delta t; y-x)}{p^2-m^2} \frac{G(\Delta t; x-y)}{p^2-m^2} a^\dagger_y(p) + \langle 0|\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*)) \int_0^\infty dpp^3 \frac{G(\Delta t; x-y)}{p^2-m^2} a^\dagger_y(p) \quad (32)$$

$$= B \int_0^\infty dpp^3 a_y(p) \frac{1}{(p^2-m^2)^2} a^\dagger_y(p) + \langle 0|\text{Tr}(\bar{\zeta}_i(s^*)\zeta_j(s^*)) \int_0^\infty dpp^3 \frac{G(\Delta t; x-y)}{p^2-m^2} a^\dagger_y(p) \quad (33)$$

Therefore, for each bubble in the expansion series of the vacuum partition function, we have:

$$\langle 0 \rangle_{bubble}^n \rightarrow \langle 0 \rangle_{bubble}^n + \langle G(x, y) \rangle_{legs}^n .$$

n is the order of perturbation.

4 Conclusion

To form a Harmonic series in the diagonal of the proposed Riemann matrix, the real part of the non-trivial Riemann zero must be $\frac{1}{2}$. Chiral symmetry is broken by coupling the matrix with an anti-particle annihilation/lowering operator, as matter and anti-matter are not treated equally.

References

- (1) Chandee, V.; Soundararajan, K. Bounding $|\zeta(\frac{1}{2}+it)|$ on the Riemann hypothesis. *The Bulletin of the London Mathematical Society* **2011**, *43* (2), 243-250. DOI: 10.1112/blms/bdq095.
- (2) Carneiro, E.; Chandee, V. Bounding $\zeta(s)$ in the critical strip. *Journal of number theory* **2011**, *131* (3), 363-384. DOI: 10.1016/j.jnt.2010.08.002.
- (3) Berndt, B. C. The Number of Zeros for $\zeta(k)(s)$. *Journal of the London Mathematical Society* **1970**, *2* (Part-4), 577-580. DOI: 10.1112/jlms/2.Part_4.577.
- (4) Peskin, M. E.; Schroeder, D. V. *An introduction to quantum field theory*; Addison-Wesley, 1995.