

A Multi-Agent Dual Dialogue System to Support Mental Health Care Providers

Onno P. Kampman¹, Ye Sheng Phang¹, Stanley Han¹, Michael Xing¹,
Xinyi Hong¹, Hazirah Hoosainsah¹, Caleb Tan¹, Genta Indra Winata^{2*},
Skyler Wang³, Creighton Heaukulani¹, Janice Huiqin Weng¹, Robert JT Morris¹

¹MOH Office for Healthcare Transformation, Singapore, ²Capital One, ³McGill University

Correspondence: onno.kampman@moht.com.sg

Abstract

We introduce a general-purpose, human-in-the-loop dual dialogue system to support mental health care professionals. The system, co-designed with care providers, is conceptualized to assist them in interacting with care seekers rather than functioning as a fully automated dialogue system solution. The AI assistant within the system reduces the cognitive load of mental health care providers by proposing responses, analyzing conversations to extract pertinent themes, summarizing dialogues, and recommending localized relevant content and internet-based cognitive behavioral therapy exercises. These functionalities are achieved through a multi-agent system design, where each specialized, supportive agent is characterized by a large language model. In evaluating the multi-agent system, we focused specifically on the proposal of responses to emotionally distressed care seekers. We found that the proposed responses matched a reasonable human quality in demonstrating empathy, showing its appropriateness for augmenting the work of mental health care providers.

1 Introduction

Mental healthcare systems globally are experiencing increased pressure due to rising demands and shortages of trained professionals. This situation has resulted in overall inadequate care and support. Waiting lists for talk therapy, one of the main interventions for mental health conditions, are long, and therapy sessions can be prohibitively expensive for many. Furthermore, mental health care professionals frequently experience burnout, exhibiting symptoms such as emotional exhaustion and depersonalization (O'Connor et al., 2018).

Digital technology can play a crucial role in addressing these needs and alleviating the pressure on the mental healthcare system. This paper proposes a *dual dialogue system* to support therapists

in conversations with their clients. To clarify, we refer to all mental health professionals as therapists throughout this paper. However, support systems may also include other professionals, such as counselors, psychologists, or general practitioners, as well as paraprofessionals, such as peer supporters or trained volunteers. The proposed system is designed for general purposes with sufficient flexibility to be adapted to various domains and contexts.

Dual dialogue systems (Figure 1) can be distinguished from standard *dialogue* systems because they do not directly interact with users (or *clients* in a therapeutic context) but rather facilitate conversation between two humans. They can be differentiated from *trilogue* or *triad* systems (Lai et al., 2023; Lee et al., 2023) because they are not conversations between three agents; there is no communication between the client and the AI assistant. A system that augments human-to-human interaction instead of facilitating human-to-AI interaction in a therapeutic setting may be more suitable for real-world deployment, as it acknowledges and respects the importance of the interpersonal relationship between a care provider and a care seeker.

In our system, the AI assistant analyzes conversations, proposes responses, recommends psychoeducational resources and self-help exercises, and evaluates and summarizes conversations for a therapist. Importantly, the system is not intended to detect or diagnose mental illness (Zhang et al., 2022) but to facilitate conversation by supporting therapists. Some potential beneficial areas may include suggestions on what is troubling the user, an instant summary of lengthy or previous sessions, or the immediate generation of resources tailored to the client's needs.

Below, we first discuss the background and context of the proposed system. The initial focus is on its application within the Singaporean mental health ecosystem. Being specific is valuable, as it

*The work was done outside Capital One.




Figure 1: Schematic of dual dialogue system design.

allows us to discuss a real-world application and collect grounded evidence. However, we stress that the system may be generalized to different nations and contexts. We then detail the requirements of the system, its design principles, and the functionalities included. Subsequently, we describe system evaluation strategies and present a response-quality evaluation study that includes both machine and human raters. Finally, we conclude with the system’s limitations and ethical considerations.

2 Background

2.1 AI and Chatbots in Mental Health

Technology-enabled assisting approaches, such as chatbots and dialogue systems, have been proposed as ways to support and augment therapeutic labor to scale and improve mental health support (Althoff et al., 2016; Bendig et al., 2019; Caceres Najarro et al., 2023; Cameron et al., 2017; Hua et al., 2024; Jin et al., 2023; Lai et al., 2023; Lee et al., 2020; Li et al., 2023; Maddela et al., 2023; Malgaroli et al., 2023; Vaidyam et al., 2019; van Heerden et al., 2023). Such approaches may be particularly useful in cultures in which seeking professional help is still stigmatized or otherwise inaccessible. More specifically, we consider agents in such systems as *role-playing* mental health professionals (Shanahan et al., 2023).

Therapeutic chatbots are not a new phenomenon, with a long heritage stretching back to the first chatbot ever developed, ELIZA, which mimicked conversations with a Rogerian psychotherapist (Weizenbaum, 1966). However, recent advances in large language models (LLMs) have led to increased interest in and hopes for developing systems with real-world applicability.

Previous studies have explored such systems (Brocki et al., 2023; Caceres Najarro et al., 2023; Winata et al., 2017). Some have been brought to market as commercial products, such as

Woebot (Fitzpatrick et al., 2017) and *Wysa* (Inkster et al., 2018). Other studies have demonstrated the efficacy of fully automated internet-based cognitive behavioral therapy (iCBT) delivery (Kwek et al., 2024; Lu et al., 2021; Linardon et al., 2024). However, mere access to self-help resources and online CBT exercises has not been a panacea for improving care (Torous and Blease, 2024). Having a human in the loop remains important for effective care and safety requirements, as most users prefer a more safeguarded experience. Moreover, standalone and self-help apps have limited utility if they are not well-integrated into healthcare systems.

Recent studies evaluating the efficacy of LLMs in offering mental health intervention have shown promising results. When evaluated against human responses, responses by LLMs such as GPT-4 were found to demonstrate more empathy (Luo et al., 2024), active listening, and helpfulness when responding to relationship or general health-related questions (Vowels, 2024; Young et al., 2024). Another study indicated that GPT-4, when prompted to act as a therapist, exhibits competency, empathy, and therapeutic capacity when delivering single-session therapy to individuals seeking assistance with relationship challenges (Vowels et al., 2024). All of this evidence suggests the potential for deploying LLMs to assist human therapists in their work.

2.2 Mental Health in Singapore

In Singapore, the Covid-2019 pandemic has brought mental health to the fore, becoming a key priority for the government (Yeo et al., 2024). This has led to the establishment of national mental health support platforms, such as *mindline.sg*¹ (Weng et al., 2024; Yoon et al., 2024) and *MindSG*².

From this former initiative, a public Reddit-inspired forum aimed at younger Singaporeans, a demographic found to be particularly vulnerable (Subramaniam et al., 2019), was born. Users on this forum can anonymously discuss topics, provide peer support (Griffiths et al., 2012), and ask questions to professional therapists on the platform (Phang et al., 2023). Various other support systems and touchpoints exist in Singapore, including crisis helplines, educational content providers, community outreach programs, and private (physical and online) support providers. However, like

¹<https://mindline.sg/>

²<https://www.healthhub.sg/programmes/mindsg/>




Figure 2: Schematic of system architecture of dual dialogue system.

many other nations, Singapore experiences a shortage of mental health professionals. In 2023, there were 298 registered psychiatrists (Singapore Medical Council, 2023) and 691 registered psychologists (Singapore Psychological Society, 2024). This translates to 5.1 and 11.9 licensed psychiatrists and psychologists per 100,000 population, respectively. The shortage of psychiatrists and psychologists results in a median waiting time of 45 days and 42 days for a subsidized appointment, respectively (Ministry of Health Singapore, 2023). Like many other nations, such an understaffed mental health workforce is often overwhelmed by workload and experiences burnout (Yang et al., 2024).

3 System Description

3.1 System Design and Requirements

Our system is co-designed and co-developed with mental health stakeholders in Singapore. Ethical considerations in the system design are important due to the sensitivity of personal data in the mental health domain. System requirements must be such that trust is maintained for all users of the system. We envisioned a dual dialogue system without interaction between the client and the AI assistant to minimize the risk of any LLM hallucinations directly affecting clients. Further, we wanted the client interface to be as familiar as possible, so we drew design inspiration from popular messaging apps (Telegram or WhatsApp). On the therapist’s side, the system is designed to be general-purpose and modular; therapists may opt to use only parts of the system. In contrast to dialogue systems that directly interact with a user, our dual dialogue system centers on therapists driving the conversation.

Natural language understanding in the AI assistant service is managed through a collection of LLM agents. In the context of our system, an “LLM agent” is defined as a specialized LLM-based module designed to perform a distinct task within the broader architecture of a given mental health support system. Each agent employs customized prompts to elicit specific types of information or responses (Priyadarshana et al., 2024), ensuring that its functionality is precisely aligned with its designated role. Further, LLM agents are distinguished by their access to different types of information. While some may tap into external databases or resources (e.g., a curated library of mental health materials), others operate solely based on the input provided by therapist and client interactions without needing external information. This modular and specialized approach allows for a scalable, adaptable system in which each agent contributes uniquely to the holistic support framework. The therapist is considered the “orchestrator” of the system agents.

Because empathy is a particularly pertinent ingredient to help achieve therapeutic outcomes (Browne et al., 2021) and promote therapeutic change (Rogers, 1957), it is important for our dual dialogue system to embody this quality. Already, empathy in chatbots has been a popular topic of study and an important requirement for existing conversational systems (Fung et al., 2016; Ma et al., 2020; Zhou et al., 2018). Empathy fosters a more supportive and warm environment and aids in the development of emotional connection, allowing users to feel more supported, validated, cared for, and understood both emotionally and cognitively (Decker et al., 2013), ultimately facili-




Figure 3: The therapist’s front-end user interface showcases the two concurrent conversations and AI features, including "Propose response," "Recommend resources," and open-ended chat.

tating client openness towards therapeutic change. Furthermore, studies have shown that empathetic responses can drive therapeutic impacts such as developing a deeper awareness and acceptance towards personal emotions and needs and gaining better emotional regulation strategies (Irrarázaval and Kalawski, 2022). Commonly used LLMs already appear to be empathetic, likely through its reinforcement learning from human feedback (RLHF) fine-tuning, but we will evaluate this further in Section 4. Traditionally, emotionally intelligent dialogue systems would include “affective computing” modules, but such approaches have become less crucial in the age of generative AI (Amin et al., 2023).

3.2 System Architecture

As shown in Figure 2, the back-end is separated from the therapist-facing front-end and the client-facing messaging platform.

The back-end infrastructure was constructed using a microservice architecture. This architecture enables scalable interaction between the front-end and containerized backend services through well-defined API calls. The back-end service was designed to deliver AI functionalities, such as proposing tailored responses, suggesting relevant

resources, analyzing and summarizing conversations, and supporting open-ended chats with the AI assistant. All of these rely heavily on the LLM model for its text generation and chat capabilities. For each specific function, we crafted customized template prompts based on specific mental health needs and adjusted the model’s initialization settings, such as whether to use retrieval-augmented generation (RAG) and the optimal maximum output token length. Our interaction with the LLM service API is facilitated by LangChain (Chase, 2022) for its extensive compatibility and strong community support. Alternatives that support RAG and could act as a drop-in replacement of LangChain include LlamaIndex (Liu, 2022) and Haystack (Deepset, 2019).

The therapist-facing front-end was built using the React JavaScript library. The UI (see Figure 3) was developed and co-designed alongside therapists to ensure the interface meets their needs and expectations. It primarily consists of two chat spaces in which mental health professionals converse, allocating the space on the left for their clients and that on the right for the AI assistant. The chat space on the left follows the design of a conventional chat interface, where the chats are listed in a side panel, and the current conversation

with the client appears as a scrolling list of message bubbles. Our system only considers text modality, but multimedia input can be included through automatic speech recognition (ASR) and multi-modal models. The chat space on the right is where the care provider interacts with the AI assistant. The system is designed to be flexible with a modular interface. Therapists may only use functions of their choosing, depending on their preferred workflow and needs. Each LLM model is tasked with one separate functionality through the multi-agent system design. The available AI functionalities were designed based on feedback from professional therapists and include the following:

1. **Propose Response:** This feature processes the entire conversation with the client to suggest an appropriate response, which can be refined before sending. Suggesting response templates in this manner may improve service quality by reducing the reply time of therapists. It relies on prompt engineering to foster engagement and promote therapeutic dialogue, focusing on eliciting clients' feelings and facilitating a deeper exploration of their experiences.
2. **Recommend Resources.** This feature retrieves the most pertinent self-help resources for care seekers using a RAG framework. Resources are pulled from the *mindline.sg* database, including over 600 articles, videos, chatbot-guided exercises inspired by cognitive behavioral theory (CBT), and in-person opportunities for follow-up support. These curated resources cater to a broad spectrum of self-help needs. This capability significantly reduces the time therapists would otherwise spend sifting through the website to find appropriate and relevant resources. RAG also helps to avoid LLM hallucination (Ji et al., 2023b) by limiting the options of what is retrieved to a relatively small set of curated content. Specifically, we converted the descriptions of resources available on *mindline.sg* into vector formats using the GPT-based `text-embedding-ada-002` embedding model from OpenAI's Embeddings API. The use of commercial APIs here was considered ethical because no personally sensitive data was included. Subsequently, we employed the Facebook AI Similarity Search (FAISS) library for efficient similarity search

and clustering within this high-dimensional vector space. FAISS, designed for fast nearest-neighbor retrieval in large datasets, allows our system to rapidly identify and cluster resource descriptions based on their semantic similarity to the content of the conversation (Douze et al., 2024).

3. **Analyze Conversation.** This feature analyzes the entire conversation with the client to extract key themes, intents, and issues that emerge during the dialogue. This analysis is focused on helping the therapist find effective leverage points for psychological first aid, offering guidance on therapeutic techniques, and/or providing constructive feedback on potential instances of ineffective or inappropriate advice. By distilling the essence of conversations and suggesting follow-up actions, therapists are equipped with a potentially deeper and more nuanced understanding of clients' concerns and the therapeutic process.
4. **Summarize Conversation.** This feature summarizes the entire conversation transcript with the client to quickly bring a therapist up to speed with the matters discussed. This is especially relevant when one therapist takes over a client from another or when a client reaches out again after an extended period of inactivity.
5. **Empathetic Rewrite.** This feature takes a draft reply from the therapist and rewrites it in an empathetic manner. This injection of empathy follows a set of guidelines that should be adopted in the response, including engagement in active listening, usage of non-judgmental language, and validation of the client's emotions and perspectives. The conversation with the client is also included as context in the system prompt so that the rewritten response would fit the flow and style of the conversation.
6. **Open-ended Chat.** This functionality loads the conversation into an LLM, from which the therapist can have an open-ended chat about the conversation with the client.

4 System Evaluation

This section describes the research methodology, study design, and corresponding empirical results

TES facet	Response Generation Source			
	Human	GPT-4o	Llama 3 70B	Llama 3 8B
concern	4.48 ± 1.53	4.00 ± 1.66 (-1.06, 0.29)	5.20 ± 1.53 (1.66, 0.10)	5.16 ± 0.99 (1.87, 0.07)
resonate	4.12 ± 1.48	4.08 ± 1.61 (-0.09, 0.93)	4.84 ± 1.49 (1.72, 0.09)	5.08 ± 1.29 (2.45, 0.02)
warmth	4.52 ± 1.81	3.80 ± 1.87 (-1.38, 0.17)	5.04 ± 1.65 (1.06, 0.29)	5.28 ± 1.46 (1.64, 0.11)
attuned	4.08 ± 1.63	4.00 ± 1.58 (-0.18, 0.86)	5.08 ± 1.63 (2.17, 0.04)	4.80 ± 1.44 (1.65, 0.10)
cognitive	4.36 ± 1.58	3.84 ± 1.28 (-1.28, 0.21)	4.60 ± 1.83 (0.50, 0.62)	5.00 ± 1.08 (1.67, 0.10)
understanding	4.44 ± 1.83	3.84 ± 1.62 (-1.23, 0.23)	4.76 ± 1.79 (0.63, 0.54)	5.00 ± 1.55 (1.17, 0.25)
acceptance	4.80 ± 1.66	4.20 ± 1.71 (-1.26, 0.21)	5.32 ± 1.44 (1.19, 0.24)	5.44 ± 1.36 (1.49, 0.14)
average	4.40 ± 1.65	3.97 ± 1.62	4.98 ± 1.62	5.11 ± 1.31

Table 1: Human rater perceived empathy scores (TES item ratings) for each response generation source. The mean ± standard deviation and *t*-test results are shown between machine-generated responses and (professional) human-written responses (*t*-value and *p*-value in brackets). Significant differences ($p < 0.05$) are bolded.

TES facet	Response Generation Source			
	Human	GPT-4o	Llama 3 70B	Llama 3 8B
concern	5.40 ± 1.26	5.48 ± 0.51 (0.29, 0.77)	5.64 ± 0.57 (0.87, 0.34)	5.60 ± 0.58 (0.72, 0.47)
resonate	3.96 ± 0.93	4.48 ± 0.51 (2.44, 0.02)	4.32 ± 0.56 (1.65, 0.10)	4.12 ± 0.44 (0.77, 0.47)
warmth	5.48 ± 1.08	5.60 ± 0.50 (0.50, 0.62)	5.84 ± 0.37 (1.57, 0.12)	5.72 ± 0.61 (0.96, 0.34)
attuned	4.00 ± 1.15	4.36 ± 0.49 (1.44, 0.16)	4.48 ± 0.51 (1.90, 0.06)	4.32 ± 0.63 (1.22, 0.23)
cognitive	4.76 ± 0.88	5.16 ± 0.47 (2.00, 0.05)	5.08 ± 0.57 (1.53, 0.13)	5.04 ± 0.45 (1.41, 0.16)
understanding	4.12 ± 1.20	4.40 ± 0.50 (1.08, 0.29)	4.56 ± 0.51 (1.69, 0.10)	4.44 ± 0.51 (1.23, 0.23)
acceptance	5.16 ± 1.18	5.44 ± 0.51 (1.09, 0.28)	5.56 ± 0.51 (1.56, 0.13)	5.56 ± 0.51 (1.56, 0.13)
average	4.70 ± 1.10	4.99 ± 0.50	5.07 ± 0.51	4.97 ± 0.53

Table 2: Machine (GPT-4o) rated perceived empathy scores (TES item ratings) for each response generation source. The mean ± standard deviation and *t*-test results are shown between machine-generated responses and (professional) human-written responses (*t*-value and *p*-value in brackets). Significant differences ($p < 0.05$) are bolded.

and discusses them in the context of our system evaluation. The proposed multi-agent system poses difficulty for a full system evaluation due to the absence of a clear performance benchmark and the presence of multiple functionalities that serve various purposes (Jin et al., 2023; Qiu et al., 2023; Rashkin et al., 2018). Moreover, we present a general-purpose system, which requires further evaluation in its specific application. In this evaluation, we focus on evaluating the *empathy* of the "Propose response" functionality, a key value proposition in helping mental health professionals and key for therapeutic effectiveness.

4.1 Study Design

We randomly sampled 100 query-response pairs from the local, anonymous mental health forum previously discussed. The responses to those queries were written by professional therapists. To evaluate the quality of responses from LLMs, we used a state-of-the-art commercial model, GPT-4o (Achiam et al., 2023), and a small and large version of a state-of-the-art open source model, Llama 3 with 8 billion and 70 billion param-

eters, to propose responses for 25 queries each, randomly sampled from the 100 queries, alongside 25 responses from the human therapists. These responses were then blindly scored against an adaptation of the Therapist Empathy Scale (TES) (Decker et al., 2013), which consists of nine rating items: "concern", "expressiveness" (unused), "resonate or capture client feelings", "warmth", "attuned to client's inner world", "understanding cognitive framework", "understanding feelings/inner experience", "acceptance of feelings/inner experiences", and "responsiveness" (unused). This evaluation excludes the items "expressiveness" and "responsiveness" because they are largely irrelevant to non-verbal exchanges. Each item is rated on a 7-point Likert scale based on the extent to which the item is demonstrated in the response, from 1 (not at all) to 7 (extremely). This adapted version of the TES can be found in Table 4.

We consider human evaluation the ground truth, but we are also interested in investigating whether machines can evaluate responses, which would allow for scaling up evaluation efforts. Therefore, responses were evaluated both by human raters and

TES facet	Response Generation Source			
	Human	GPT-4o	Llama 3 70B	Llama 3 8B
concern	2.33 (0.029)	4.39 (0.000)	1.59 (0.126)	2.53 (0.018)
resonate	-0.48 (0.637)	1.11 (0.278)	-1.87 (0.073)	-3.59 (0.001)
warmth	2.75 (0.011)	5.06 (0.000)	2.49 (0.020)	1.55 (0.134)
attuned	-0.24 (0.811)	1.14 (0.265)	-1.90 (0.070)	-1.63 (0.117)
cognitive	1.19 (0.246)	5.28 (0.000)	1.33 (0.196)	0.19 (0.852)
understanding	-0.90 (0.376)	1.80 (0.085)	-0.58 (0.569)	-1.90 (0.070)
acceptance	0.91 (0.372)	3.72 (0.001)	0.90 (0.376)	0.44 (0.664)

Table 3: Comparison between human and machine (GPT-4o) rating. Paired t -test of empathy scores (TES item ratings) between ratings by GPT-4o (r_1) and ratings by human raters (r_2). Shown are the corresponding t -statistics and p -values: negative t -statistic values indicate that human raters perceived the responses as more empathetic. Significant differences ($p < 0.05$) are bolded.

by GPT-4o. We recruited ten human raters (three males and seven females), ages 21 to 30. Each human rater was randomly assigned 10 different query-response pairs from the 100 randomly sampled query-response pairs and tasked with evaluating each response according to the TES. All human raters had existing mental health or psychological first aid-related training, with some being experienced social workers or counselors-in-training. Ethical considerations were addressed by briefing all human raters on the nature and purpose of the research and the risks of generative AI. In addition to human evaluation, we prompted GPT-4o to score each response using the adapted TES. This was achieved through prompt engineering, where the LLM was tasked to return a score for each facet.

4.2 Results and Discussion

Table 1 shows the means and standard deviations of the human-rated TES scores for each response-generating source. The p -values derived from t -tests on the rating scores between the human-generated responses and each of the machine-generated responses are included. The rating score distributions are shown in Figure 6. Few significant differences are found between the human- and machine-generated responses. In these evaluations by human raters, Llama 3 8B performed significantly better in the item resonate ($p = 0.02$), while Llama 3 70B performed significantly better in the item attuned ($p = 0.04$) as compared to respective facets of empathy in human professional written responses. This highlights an LLM’s comparable ability to capture emotion, thought, tone, and nuance underlying a statement or query, and respond to an individual in distress in a manner that signals active listening and care.

GPT-4o was also used as a source of evaluation

to validate our findings and explore alternative and more scalable evaluation methods. Table 2 shows the means and standard deviations of the GPT-4o-rated TES rating scores for each response source, as well as the p -values derived from t -tests on the rating scores between the human-generated responses and each of the machine-generated responses. The rating score distributions are shown in Figure 5. Again, we do not find many significant differences between human- and machine-generated responses. In these evaluations by GPT-4o, we only found that responses from GPT-4o performed significantly better in the item resonate ($p = 0.02$). The consistently lower standard deviation among LLM responses may be due to the standardized format of LLM responses in demonstrating empathy and the fixed preference of GPT-4o as the sole model for LLM evaluation. Results from GPT-4o evaluation roughly resonate with results from human evaluation.

In both the AI and human evaluations, LLM responses generally had similar TES ratings to professional responses. LLMs only showed a significant improvement in a few rating items of the TES but lacked any significant overall difference when compared to professional responses. This highlights the overall competency of LLMs in demonstrating a level of empathy comparable to that of a human therapist. Our results align with previous research on an LLM’s capacity for empathy in a therapeutic context. In the specific context of relationship issues, Vowels (2024) found that when evaluating empathy in responses from GPT-4 and relationship experts, results from a sample of 20 human evaluators showed that LLMs performed similarly to or even better than relationship experts. We stress that our goal is not to show that these LLMs are better than professional therapists, as achieving

therapeutic outcomes goes greatly beyond the ability to demonstrate empathy (Browne et al., 2021). Rather, we show that LLMs can propose responses of similar empathetic quality and may be used to augment human-provided care.

4.3 Comparing LLM and Human Evaluation

The current reliance on human evaluation is time-consuming and does not scale well, especially since underlying LLMs may be frequently updated. Reliable LLM evaluation may automate the assessing the domain of empathy. To determine the reliability of LLM evaluation, we explored the similarity between results from LLM and human evaluations.

In human evaluation (Table 1), GPT-4o-generated responses showed the lowest average mean ($M = 3.97$), but in LLM evaluation (Table 2), GPT-4o-generated responses reflected an average mean ($M = 4.99$) higher than professional responses. One explanation for the disparity between GPT-4o's performance between human and GPT-4o evaluations may be due to the presence of a dataset or AI bias (Srinivasan and Chander, 2021). When using an LLM model to evaluate a response generated by the same LLM model, the output from both the response and the evaluation comes from the same model trained in the same manner and on the same dataset. This likely increases the evaluation model's preference towards responses generated by the same model due to the better alignment to the data it was trained on. Therefore, this results in a higher rating in GPT-4o responses when rated by GPT-4o. In addition, a common feedback that the AI responses were easily identifiable was heard from a few human evaluators after their evaluations. This is likely due to the distinct point-form format that GPT-4o uses compared to Llama 3 responses, which use a more elaborative and lengthy format that lacks the point-form structure when responding to most mental health queries. Given that awareness towards the use of AI use in mental health may lead to prejudice or bias against the AI response (Jain et al., 2024), human evaluators may have given the easily identified AI response (GPT-4o responses) a lower rating as compared to responses that had less resemblance to an AI response (human or Llama responses).

A second observed difference between human and LLM evaluation lies in the difference in the reported standard deviations. Across all response sources, the overall standard deviation in

human evaluation was higher than the overall standard deviation in GPT-4o-generated evaluation. Compared to the GPT-4o evaluation, this increased inconsistency among human raters is likely due to the subjectivity of human interpretations and preferences towards psychological interventions (Williams et al., 2016). Just as how no single therapist can suit every client, each individual has slight preferences.

Table 3 shows the statistical test results between human and GPT-4o evaluation for each of the four response generation methods. The GPT-4o evaluation found human responses relatively more empathetic in eight facets across the generation sources, but particularly finds GPT-4o responses more empathetic than humans judged them to be. These results highlight the differences between LLM and human evaluation. LLMs may not be suitable for drop-in replacement of human evaluation efforts. However, the reliability of LLM evaluation and its adherence to human preferences may be further improved through prompt engineering or fine-tuning.

5 Conclusion

We presented a general-purpose dual dialogue system to support mental health care providers in their conversations with clients. The system is designed to assist in crafting responses, recommending resources, and addressing key emotional struggles of care seekers. It can also reduce the cognitive load and fatigue of care providers. This system does not aim to replace mental health professionals or promote a thoughtless regurgitation of LLM-generated responses to care seekers. The human-in-the-loop design allows professional therapists to incorporate the uniqueness of their personality, personal qualities, and style as a therapist (Smith, 2003), and ensure the quality and safety of responses with their professional judgment. Our evaluation study demonstrated that LLMs are comparable to professional therapists in generating empathetic responses.

Limitations

As a system demonstration, we acknowledge several limitations of both the current form of our system and its future applicability.

First, using a commercial API such as GPT-4o raises privacy concerns, so this system cannot be easily expanded to private conversations. We intend to explore the use of open-source models such

as Llama 3 (Touvron et al., 2023) and evaluate how this changes the system's performance and acceptability. Our evaluation study indicates that Llama 3 may be a viable alternative to GPT-4o.

Second, challenges in LLM applications include interpretability, hallucinations (Ji et al., 2023a), privacy, bias, and clinical effectiveness (Chung et al., 2023). Hallucinations can be obvious, such as nonsensical language or switching to another language, or more subtle, such as returning seemingly correct but erroneous information. LLMs have also been shown to perpetuate stereotypes of mental disorders such as schizophrenia (King, 2022).

Third, engagement remains an issue in this space. We already know that mental health apps fail to consistently engage users. However, tools for clinicians are often not engaging either. For the current system to be embraced and adopted, we intend to conduct further usability studies (Greenhalgh et al., 2017).

Fourth, while we have specifically framed this system as a human-in-the-loop system, we acknowledge that, in practice, this distinction may fade. When the proposed responses are good, and care professionals are overwhelmed, the proposed responses may not be validated by humans anymore as an unintended consequence (an effect known as *automation bias*). This reintroduces some of the risks of a system that directly interacts with clients. This phenomenon has been observed in self-driving cars, where drivers start to trust the system and stop paying attention to such a degree that they can no longer be considered a robust safety agent in the loop.

Fifth, our results' generalizability is limited by our small sample size of 100 query responses and ten human evaluators. The human evaluators, all between 21 and 30 and 70 percent female, may skew our results. Furthermore, the sole focus on empathy will not be sufficient to test the utility of the system at large. Therefore, an evaluation with a larger sample to ensure the generalizability of our results and a system-as-a-whole evaluation is needed.

Ethics Statement

The dual dialogue system discussed may have far-reaching societal impacts. It may represent, at least in part, a redeployment of resources towards technology that could otherwise have been used to train more mental health professionals. Moreover, rapid

proliferation of such systems could destabilize professional norms and institutional practices in unsustainable ways (Wang et al., 2024). However, we reiterate that our system is designed to *support* therapists, helping them to better help their clients rather than to replace them.

Current state-of-the-art LLMs exhibit markedly stronger performance in English than in other languages. In diverse and multilingual societies such as Singapore, a system that only works well in English is likely to be inadequate for many users. It may even be the case that, given the nature of existing service provision, it is even more important to serve users in languages other than English (Love-*nia et al.*, 2024). We hope that future evaluations can explore system performance in other languages in greater detail and consider the evaluation of models explicitly prioritizing multilinguality (Üstün et al., 2024).

LLMs are not value-neutral and have been shown to encode specific social, cultural, and political viewpoints, often aligned with North American values (Santurkar et al., 2023; Arora et al., 2023). Mental health conditions and the seeking of mental health support are often stigmatized in various ways around the world. For an LLM-based mental health tool to be effective, it may need to understand the specific socio-cultural context of its users, which may engender additional fine-tuning by incorporating diverse and representative viewpoints.

As this work is a demonstration of an early system, it has not yet been extensively evaluated with real human participants in (one of) its applications, which includes the system operators and the downstream end users. All conversation extracts used in our analysis and included in the figures throughout this work were taken from public forum responses. Evaluating the performance and acceptability of this system with real users seeking mental health support would require a thorough ethics review by an institutional review board.

Acknowledgments

We thank Samuel J. Bell for his helpful feedback on our manuscript. We also thank our human raters: Kellie Sim, Lee Li Hwee, Zuyuan Cheong, and Allyson Grace Chai Y. J. for their contributions. We thank Claire Seah for diagram design.

References

- Josh Achiam, Steven Adler, Sandhini Agarwal, Lama Ahmad, Ilge Akkaya, Florencia Leoni Aleman, Diogo Almeida, Janko Altmenschmidt, Sam Altman, Shyamal Anadkat, et al. 2023. GPT-4 technical report. *arXiv preprint arXiv:2303.08774*.
- Tim Althoff, Kevin Clark, and Jure Leskovec. 2016. Large-scale analysis of counseling conversations: An application of natural language processing to mental health. *Transactions of the Association for Computational Linguistics*, 4:463–476.
- Mostafa M. Amin, Erik Cambria, and Björn W. Schuller. 2023. Will Affective Computing Emerge from Foundation Models and General AI? A First Evaluation on ChatGPT. *arXiv preprint*.
- Arnav Arora, Lucie-aimée Kaffee, and Isabelle Augenstein. 2023. Probing Pre-Trained Language Models for Cross-Cultural Differences in Values. In *Proceedings of the First Workshop on Cross-Cultural Considerations in NLP (C3NLP)*, pages 114–130.
- Eileen Bendig, Benjamin Erb, Lea Schulze-Thuesing, and Harald Baumeister. 2019. The next generation: Chatbots in clinical psychology and psychotherapy to foster mental health – a scoping review. *Verhaltenstherapie*, 32:64 – 76.
- Lennart Brocki, George C. Dyer, Anna Gładka, and Neo Christopher Chung. 2023. Deep learning mental health dialogue system. *2023 IEEE International Conference on Big Data and Smart Computing (Big-Comp)*, pages 395–398.
- Julia Browne, Corinne Cather, and Kim T. Mueser. 2021. Common factors in psychotherapy.
- Lismer Andres Caceres Najarro, Yonggeon Lee, Kobiljon E. Toshnazarov, Yoonhyung Jang, Hyungsook Kim, and Youngtae Noh. 2023. WMGPT: Towards 24/7 online prime counseling with ChatGPT. *Adjunct Proceedings of the 2023 ACM International Joint Conference on Pervasive and Ubiquitous Computing & the 2023 ACM International Symposium on Wearable Computing*.
- Gillian Cameron, David Cameron, Gavin Megaw, Raymond R. Bond, Maurice D. Mulvenna, Siobhan O’neill, Chérie Armour, and Michael F. McTear. 2017. Towards a chatbot for digital counselling. In *British Computer Society Conference on Human-Computer Interaction*.
- Harrison Chase. 2022. LangChain.
- Neo Christopher Chung, George C. Dyer, and Lennart Brocki. 2023. Challenges of large language models for mental health counseling. *ArXiv*, abs/2311.13857.
- Suzanne E. Decker, Charla Nich, Kathleen M. Carroll, and Steve Martino. 2013. Development of the therapist empathy scale. *Behavioural and Cognitive Psychotherapy*, 42:339 – 354.
- Deepset. 2019. Haystack.
- Matthijs Douze, Alexandr Guzhva, Chengqi Deng, Jeff Johnson, Gergely Szilvasy, Pierre-Emmanuel Mazaré, Maria Lomeli, Lucas Hosseini, and Hervé Jégou. 2024. The faiss library. *arXiv preprint arXiv:2401.08281*.
- Kathleen Kara Fitzpatrick, Alison M Darcy, and Molly Vierhile. 2017. Delivering Cognitive Behavior Therapy to Young Adults With Symptoms of Depression and Anxiety Using a Fully Automated Conversational Agent (Woebot): A Randomized Controlled Trial. *JMIR Mental Health*, 4.
- Pascale Fung, Dario Bertero, Yan Wan, Anik Dey, Ricky Ho Yin Chan, Farhad Bin Siddique, Yang Yang, Chien-Sheng Wu, and Ruixi Lin. 2016. Towards empathetic human-robot interactions. In *Conference on Intelligent Text Processing and Computational Linguistics*.
- Trisha Greenhalgh, Joseph P. Wherton, Chrysanthi Pappoutsis, Jennifer Lynch, Gemma Hughes, Christine A’Court, Susan Hinder, Nick Fahy, R. N. Procter, and Sara E. Shaw. 2017. Beyond adoption: A new framework for theorizing and evaluating nonadoption, abandonment, and challenges to the scale-up, spread, and sustainability of health and care technologies. *Journal of Medical Internet Research*, 19.
- Kathleen Margaret Griffiths, Andrew J Mackinnon, Dimity Ann Crisp, Helen Christensen, Kylie Bennett, and Louise M. Farrer. 2012. The effectiveness of an online support group for members of the community with depression: A randomised controlled trial. *PLoS ONE*, 7.
- Yining Hua, Fenglin Liu, Kailai Yang, Zehan Li, Yi-han Sheu, Peilin Zhou, Lauren V Moran, Sophia Ananiadou, and Andrew Beam. 2024. Large Language Models in Mental Health Care: A Scoping Review. *arXiv preprint arXiv:2401.02984*.
- Becky Inkster, Shubhankar Sarda, and Vinod Subramanian. 2018. An Empathy-Driven, Conversational Artificial Intelligence Agent (Wysa) for Digital Mental Well-Being: Real-World Data Evaluation Mixed-Methods Study. *JMIR mHealth and uHealth*, 6.
- Leonor Irarrázaval and Juan Pablo Kalawski. 2022. Phenomenological considerations on empathy and emotions in psychotherapy. *Frontiers in Psychology*, 13.
- Gagan Jain, Samridhi Pareek, and Per Carlbring. 2024. Revealing the source: How awareness alters perceptions of AI and human-generated mental health responses. *Internet Interventions*, 36:100745.
- Shaoxiong Ji, Tianlin Zhang, Kailai Yang, Sophia Ananiadou, and Erik Cambria. 2023a. Rethinking large language models in mental health applications. *ArXiv*, abs/2311.11267.

- Ziwei Ji, Nayeon Lee, Rita Frieske, Tiezheng Yu, Dan Su, Yan Xu, Etsuko Ishii, Ye Jin Bang, Andrea Madotto, and Pascale Fung. 2023b. Survey of hallucination in natural language generation. *ACM Computing Surveys*, 55(12):1–38.
- Haoan Jin, Siyuan Chen, Mengyue Wu, and Kenny Q Zhu. 2023. PsyEval: A Comprehensive Large Language Model Evaluation Benchmark for Mental Health. *arXiv preprint arXiv:2311.09189*.
- Morgan King. 2022. Harmful biases in artificial intelligence. *The Lancet Psychiatry*.
- Tammie Rong Rong Kwek, Jackki Hoon Eng Yim, Erik Andersson, Oliver Suendermann, Mythily Subramaniam, Elna Yadin, Janhavi Vaingankar, and Bhanu Gupta. 2024. Effectiveness and acceptability of Internet-based Cognitive Behavioral Therapy for individuals with Obsessive Compulsive Disorder in Singapore. *Journal of Behavioral and Cognitive Therapy*, 34(2):100487.
- Tin Lai, Yukun Shi, Zicong Du, Jiajie Wu, Ken Fu, Yichao Dou, and Ziqi Wang. 2023. Psy-LLM: Scaling up Global Mental Health Psychological Services with AI-based Large Language Models. *ArXiv*, abs/2307.11991.
- Peter Lee, Carey Goldberg, and Isaac Kohane. 2023. *The AI revolution in medicine: GPT-4 and beyond*. Pearson.
- Yi-Chieh Lee, Naomi Yamashita, Yun Huang, and Wai-Tat Fu. 2020. "I Hear You, I Feel You": Encouraging Deep Self-disclosure through a Chatbot. *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems*.
- Han Li, Renwen Zhang, Yi-Chieh Lee, Robert E Kraut, and David C Mohr. 2023. Systematic review and meta-analysis of AI-based conversational agents for promoting mental health and well-being. *NPJ Digital Medicine*, 6(1):236.
- Jake Linardon, John B Torous, Joseph Firth, Pim Cuijpers, Mariel Messer, and Matthew Fuller-Tyszkiewicz. 2024. Current evidence on the efficacy of mental health smartphone apps for symptoms of depression and anxiety: A meta-analysis of 176 randomized controlled trials. *World Psychiatry*, 23.
- Jerry Liu. 2022. *LlamaIndex*.
- Holy Lovenia, Rahmad Mahendra, Salsabil Maulana Akbar, Lester James V. Miranda, Jennifer Santoso, Elyanah Aco, Akhdan Fadhilah, Jonibek Mansurov, Joseph Marvin Imperial, Onno P. Kampman, Joel Ruben Antony Moniz, Muhammad Ravi Shulthan Habibi, Frederikus Hudi, Railey Montalan, Ryan Ignatius, Joanito Agili Lopo, William Nixon, Börje F. Karlsson, James Jaya, Ryandito Diandaru, Yuze Gao, Patrick Amadeus, Bin Wang, Jan Christian Blaise Cruz, Chenxi Whitehouse, Ivan Halim Parmonangan, Maria Khelli, Wenyu Zhang, Lucky Susanto, Reynard Adha Ryanda, Sonny Lazuardi Hermawan, Dan John Velasco, Muhammad Dehan Al Kautsar, Willy Fitra Hendria, Yasmin Moslem, Noah Flynn, Muhammad Farid Adilazuarda, Haochen Li, Johannes Lee, R. Damanhuri, Shuo Sun, Muhammad Reza Qorib, Amirbek Djanibekov, Wei Qi Leong, Quyet V. Do, Niklas Muennighoff, Tanrada Pansuwan, Ilham Firdausi Putra, Yan Xu, Ngee Chia Tai, Ayu Purwarianti, Sebastian Ruder, William Tjhi, Peerat Limkonchotiwat, Alham Fikri Aji, Sedrick Keh, Genta Indra Winata, Ruochoen Zhang, Fajri Koto, Zheng-Xin Yong, and Samuel Cahyawijaya. 2024. SEACrowd: A Multilingual Multimodal Data Hub and Benchmark Suite for Southeast Asian Languages. *Proceedings of EMNLP*.
- Sharon HX Lu, Hanita A Assudani, Tammie RR Kwek, Shaun WH Ng, Trisha EL Teoh, and Geoffrey CY Tan. 2021. A randomised controlled trial of clinician-guided internet-based cognitive behavioural therapy for depressed patients in singapore. *Frontiers in Psychology*, 12.
- Man Luo, Christopher J. Warren, Lu Cheng, Haidar M. Abdul-Muhsin, and Imon Banerjee. 2024. Assessing empathy in large language models with real-world physician-patient interactions. *arXiv preprint*.
- Yukun Ma, Khanh Linh Nguyen, Frank Xing, and E. Cambria. 2020. A survey on empathetic dialogue systems. *Inf. Fusion*, 64:50–70.
- Mounica Maddela, Megan Ung, Jing Xu, Andrea Madotto, Heather Foran, and Y-Lan Boureau. 2023. Training models to generate, recognize, and reframe unhelpful thoughts. *arXiv preprint arXiv:2307.02768*.
- Matteo Malgaroli, Thomas D. Hull, Jamie M. Zech, and Tim Althoff. 2023. Natural language processing for mental health interventions: A systematic review and research framework. *Translational Psychiatry*, 13.
- Ministry of Health Singapore. 2023. Waiting time for appointments for mental health treatment at public healthcare institutions. <https://www.moh.gov.sg/news-highlights/details/waiting-time-for-appointments-for-mental-health-treatment-at-public-healthcare-institutions>.
- Karen O'Connor, Deirdre Muller Neff, and Steve Pitman. 2018. Burnout in mental health professionals: A systematic review and meta-analysis of prevalence and determinants. *European psychiatry: The journal of the Association of European Psychiatrists*, 53:74–99.
- Ye Sheng Phang, Creighton Heaukulani, Wijaya Martanto, Robert Morris, Mian Mian Tong, and Roger Ho. 2023. Perceptions of a Digital Mental Health Platform Among Participants With Depressive Disorder, Anxiety Disorder, and Other Clinically Diagnosed Mental Disorders in Singapore: Usability and Acceptability Study. *JMIR Human Factors*, 10:e42167.

- Y.H.P.P. Priyadarshana, Ashala Senanayake, Zilu Liang, and Ian Piumarta. 2024. Prompt engineering for digital mental health: A short review. *Frontiers in Digital Health*, 6:1410947.
- Huachuan Qiu, Tong Zhao, Anqi Li, Shuai Zhang, Hongliang He, and Zhenzhong Lan. 2023. A benchmark for understanding dialogue safety in mental health support. In *Natural Language Processing and Chinese Computing*.
- Hannah Rashkin, Eric Michael Smith, Margaret Li, and Y-Lan Boureau. 2018. Towards empathetic open-domain conversation models: A new benchmark and dataset. In *Annual Meeting of the Association for Computational Linguistics*.
- Carl R. Rogers. 1957. The necessary and sufficient conditions of therapeutic personality change. *Journal of Consulting Psychology*, 21:95–103.
- Shibani Santurkar, Esin Durmus, Faisal Ladhak, Cino Lee, Percy Liang, and Tatsunori Hashimoto. 2023. Whose Opinions Do Language Models Reflect? Preprint, arxiv:2303.17548.
- Murray Shanahan, Kyle McDonell, and Laria Reynolds. 2023. Role play with large language models. *Nature*, 623:493–498.
- Singapore Medical Council. 2023. Annual report 2023. [https://www.healthprofessionals.gov.sg/docs/librariesprovider2/publications-newsroom/smc-annual-reports/smc_annual-report-2023-03sept-pdf-a-\(1\).pdf?sfvrsn=b379c36a_1](https://www.healthprofessionals.gov.sg/docs/librariesprovider2/publications-newsroom/smc-annual-reports/smc_annual-report-2023-03sept-pdf-a-(1).pdf?sfvrsn=b379c36a_1).
- Singapore Psychological Society. 2024. Directory of sps members. <https://singaporepsychologicalsociety.org/members-directory/>.
- Edward WL Smith. 2003. *The person of the therapist*. McFarland.
- Ramya Srinivasan and Ajay Chander. 2021. Biases in ai systems. *Communications of the ACM*, 64(8):44–49.
- Mythily Subramaniam, Edimansyah Abidin, Janhavi Ajit Vaingankar, Saleha Shafie, Boon Yiang Chua, Rajeswari Sambasivam, Y. J. Zhang, Shazana Shahwan, S. M. Chang, Hong Choon Chua, Swapna Verma, Lyn James, Kian Woon Kwok, Derrick Mk Heng, and Siow Ann Chong. 2019. Tracking the mental health of a nation: Prevalence and correlates of mental disorders in the second Singapore mental health study. *Epidemiology and Psychiatric Sciences*, 29.
- John B. Torous and Charlotte Blease. 2024. Generative artificial intelligence in mental health care: Potential benefits and current challenges. *World Psychiatry*, 23.
- Hugo Touvron, Louis Martin, Kevin R. Stone, Peter Albert, Amjad Almahairi, Yasmine Babaei, Nikolay Bashlykov, Soumya Batra, Prajjwal Bhargava, Shruti Bhosale, Daniel M. Bikel, Lukas Blecher, Cristian Cantón Ferrer, Moya Chen, Guillem Cucurull, David Esiobu, Jude Fernandes, Jeremy Fu, Wenyin Fu, Brian Fuller, Cynthia Gao, Vedanuj Goswami, Naman Goyal, Anthony S. Hartshorn, Saghar Hosseini, Rui Hou, Hakan Inan, Marcin Kardas, Viktor Kerkez, Madian Khabsa, Isabel M. Kloumann, A. V. Korenev, Punit Singh Koura, Marie-Anne Lachaux, Thibaut Lavril, Jenya Lee, Diana Liskovich, Yinghai Lu, Yuning Mao, Xavier Martinet, Todor Mihaylov, Pushkar Mishra, Igor Molybog, Yixin Nie, Andrew Poulton, Jeremy Reizenstein, Rashi Rungta, Kalyan Saladi, Alan Schelten, Ruan Silva, Eric Michael Smith, R. Subramanian, Xia Tan, Binh Tang, Ross Taylor, Adina Williams, Jian Xiang Kuan, Puxin Xu, Zhengxu Yan, Iliyan Zarov, Yuchen Zhang, Angela Fan, Melanie Kambadur, Sharan Narang, Aurelien Rodriguez, Robert Stojnic, Sergey Edunov, and Thomas Scialom. 2023. Llama 2: Open foundation and fine-tuned chat models. *ArXiv*, abs/2307.09288.
- Ahmet Üstün, Viraat Aryabumi, Zheng-Xin Yong, Wei-Yin Ko, Daniel D’souza, Gbemileke Onilude, Neel Bhandari, Shivalika Singh, Hui-Lee Ooi, Amr Kayid, Freddie Vargus, Phil Blunsom, Shayne Longpre, Niklas Muennighoff, Marzieh Fadaee, Julia Kreutzer, and Sara Hooker. 2024. Aya Model: An Instruction Finetuned Open-Access Multilingual Language Model. Preprint, arxiv:2402.07827.
- Aditya Nrusimha Vaidyam, Hannah Wisniewski, John D. Halamka, Matcheri S. Kashavan, and John B Torous. 2019. Chatbots and conversational agents in mental health: A review of the psychiatric landscape. *The Canadian Journal of Psychiatry*, 64:456 – 464.
- Alastair C. van Heerden, Julia R Pozuelo, and Brandon A. Kohrt. 2023. Global mental health services and the impact of artificial intelligence-powered large language models. *JAMA psychiatry*.
- Laura M. Vowels. 2024. Are chatbots the new relationship experts? Insights from three studies. *Computers in Human Behavior: Artificial Humans*, 2.
- Laura M. Vowels, Rachel R.R. Francois-Walcott, and Joëlle Darwiche. 2024. AI in Relationship Counselling: Evaluating ChatGPT’s Therapeutic Capabilities in Providing Relationship Advice. *Computers in Human Behavior: Artificial Humans*, 2.
- Skyler Wang, Ned Cooper, and Margaret Eby. 2024. From human-centered to social-centered artificial intelligence: Assessing ChatGPT’s impact through disruptive events. *Big Data & Society*, 11(4).
- Joseph Weizenbaum. 1966. ELIZA - A computer program for the study of natural language communication between man and machine. *Communications of the ACM*, 9(1):36–45.
- Janice Huiqin Weng, Yanyan Hu, Creighton Heaukulani, Clarence Tan, Julian Kuiyu Chang, Ye Sheng Phang, Priyanka Rajendram, Weng Mooi Tan, Wai Chiong Loke, and Robert JT Morris. 2024. Mental Wellness

Self-Care in Singapore With mindline.sg: A Tutorial on the Development of a Digital Mental Health Platform for Behavior Change. *Journal of Medical Internet Research*, 26.

Ryan Williams, Lorna Farquharson, Lucy Palmer, Paul Bassett, Jeremy Clarke, David M. Clark, and Mike J. Crawford. 2016. Patient preference in psychological treatment and associations with self-reported outcome: National cross-sectional survey in England and Wales. *BMC Psychiatry*, 16.

Genta Indra Winata, Onno P. Kampman, Yang Yang, Anik Dey, and Pascale Fung. 2017. Nora the empathetic psychologist. In *INTERSPEECH*, pages 3437–3438.

Suyi Yang, Germaine Ke Jia Tan, Kang Sim, Lucas Jun Hao Lim, Benjamin Yong Qiang Tan, Abhiram Kanneganti, Shirley Beng Suat Ooi, and Lue Ping Ong. 2024. [Stress and burnout amongst mental health professionals in Singapore during Covid-19 endemicity](#). *PLOS ONE*, 19(1).

Pearlie Mei En Yeo, Vicky Mengqi Qin, Chin-Siang Ang, Michael Chia, Ringo Moon-Ho Ho, Andy Hau Yan Ho, and Josip Car. 2024. [Prevalence and correlates of depressive symptoms among matriculated university students in singapore during covid-19 pandemic: Findings from a repeated cross-sectional analysis](#). *BMC Public Health*, 24.

Sungwon Yoon, Hendra Goh, Xinyi Casuarine Low, Janice Huiqin Weng, and Creighton Heaukulani. 2024. User perceptions and utilisation of features of an ai-enabled workplace digital mental wellness platform ‘mindline at work’. *BMJ Health & Care Informatics*, 31(1).

Jordyn Young, Laala M Jawara, Diep N Nguyen, Brian Daly, Jina Huh-Yoo, and Afsaneh Razi. 2024. [The Role of AI in Peer Support for Young People: A Study of Preferences for Human- and AI-Generated Responses](#). In *Proceedings of the 2024 CHI Conference on Human Factors in Computing Systems*.

Tianlin Zhang, Annika M Schoene, Shaoxiong Ji, and Sophia Ananiadou. 2022. Natural language processing applied to mental illness detection: A narrative review. *NPJ digital medicine*, 5(1):1–13.

Li Zhou, Jianfeng Gao, Di Li, and Harry Shum. 2018. [The Design and Implementation of XiaoIce, an Empathetic Social Chatbot](#). *Computational Linguistics*, 46:53–93.

A Example Conversations

Figure 4 shows several examples of conversational functionalities. We highlight several system capabilities. For the **Propose Response** functionality, the AI assistant acknowledges the client’s emotions and stresses the importance of a social support network. For the **Recommend Resources**

functionality, it briefly empathizes with the client, and then recommends a relevant sleep-related exercise, while also explaining why such exercises may be helpful. For the **Analyze Conversation** functionality, the AI assistant extracts relevant themes from the client’s message and discusses potential therapeutic directions, such as learning more about managing work-related stressors, encouraging open communication with their partner, and exploring self-esteem issues that may underlie the symptoms discussed. While any trained therapist will understand these insights, this functionality can prove useful when meeting a new client with a chat history or when the therapist has not received extensive training. For the **Summarize Conversation** functionality, the AI assistant condenses an exchange between client and therapist in a short paragraph by extracting the main themes and issues and discussing focal points of the healing process.

<p style="text-align: center;">Propose response</p> <p>Client:</p> <p>I'm feeling incredibly sorrowful and isolated after the passing of my cat, and I'm unsure how to cope with these emotions.</p> <p>AI Assistant:</p> <p>I'm sorry to hear about the loss of your cat. It's understandable to feel sad and lonely during this time. Grieving the loss of a pet can be incredibly difficult because they are not just pets; they are family members. It's important to allow yourself to feel these emotions and not to rush the grieving process. You might find it helpful to remember the good times you had with your cat and perhaps celebrate their life in a way that feels right to you. Additionally, talking about your feelings with friends, family, or a professional can provide comfort.</p>	<p style="text-align: center;">Recommend resources</p> <p>Client:</p> <p>Hey there, it's 4 am and I'm wide awake, feeling absolutely terrible.</p> <p>AI Assistant:</p> <p>I'm sorry to hear that you're having trouble sleeping and feeling awful. It's important to address both your immediate discomfort and the underlying issues that might be causing your insomnia. For tonight, I recommend trying a guided sleep meditation or a sleep story to help your mind relax and drift off to sleep. Mindline.sg offers a tool called Wysa, an emotionally intelligent AI chatbot, which provides a guided sleep meditation called "Drift to Sleep." This meditation focuses on developing positive feelings before bedtime, which might help you find a more peaceful state of mind. Additionally, consider exploring resources on improving sleep hygiene and creating a bedtime routine that encourages relaxation and sleep. Remember, it's also beneficial to consult with a professional if your sleep issues persist, as they can offer personalized advice and support.</p> <p>Resource:</p> <ul style="list-style-type: none"> - https://www.mindline.sg/workspace/resource-group/practising-self-care/resource-sub-group/getting-better-sleep
<p style="text-align: center;">Analyze conversation</p> <p>Client:</p> <p>Feeling overwhelmed, I struggle to cope with the immense pressure at work, recognizing that some of it is self-inflicted. Consequently, finding any happiness becomes challenging. Despite my daily hard work, I fail to see results, leading to a decline in my confidence. Unfortunately, I vent my frustrations on my boyfriend, making our weekend encounters strained and difficult. The sense of incompetence and worthlessness weighs heavily on me.</p> <p>AI Assistant:</p> <p>The client is experiencing significant stress due to work pressure, some of which they recognize as self-imposed. This stress is affecting their happiness, self-esteem, and relationships, particularly with their boyfriend. They feel incompetent and struggle with coping mechanisms.</p> <p>Themes and issues discussed include work-related stress, self-imposed pressure, impact on personal relationships, and feelings of incompetence. Potential solutions could involve strategies for managing tough work situations, such as recognizing and resolving conflicts, communicating effectively with supervisors about workload, practicing self-care, and building a support network. Encouraging the client to engage in open communication with their partner about their feelings and stressors could also be beneficial. It might be helpful to explore the underlying reasons for the self-imposed pressure and work on building self-esteem and confidence through positive self-talk and setting realistic expectations.</p>	<p style="text-align: center;">Summarize conversation</p> <p>Client:</p> <p>After our breakup last year, my ex and I recently rekindled things. I find myself drawn back to her, feeling a strong need for her presence in my life. However, I'm unsure if I can handle just being friends. Despite our honest conversations and her recent clarification that she doesn't want more, I've found myself spiraling. Since she made her feelings clear, I've been isolating myself and emotionally numbing. I've lost my appetite, struggle to sleep at night, and find it hard to even drink.</p> <p>Therapist:</p> <p>Thank you for opening up about your struggles with romantic relationships. Dealing with a breakup can be devastating, but finding a way to rekindle things must have felt like a second chance. I want you to know that your thoughts and feelings are valid given the circumstances. However, it's concerning to see you feeling overwhelmed and neglecting self-care.</p> <p>Humans naturally crave connections, especially in romantic relationships where we find comfort, passion, and shared values. There's something significant in that relationship that keeps drawing you back. Perhaps it's worth reflecting on what those needs are and exploring if they can be fulfilled in other ways.</p> <p>Consider listing down the things you value in a relationship, as well as your deal-breakers. This exercise can help you understand what you're looking for and how you can contribute to those aspects. Additionally, think about how you can attract the kind of people you want to be around by embodying those values yourself.</p> <p>I also wonder about the underlying thoughts that lead you to isolate and "punish" yourself. Have you noticed this pattern of behavior before, and how did you cope with it in the past? While these behaviors may not be helpful in the long run, it's essential to be kinder to yourself and shift to a more constructive perspective. Do you have someone you trust that you can talk to and seek support from? Remember, you deserve and can receive the necessary support. Take care, and I hope to hear from you soon.</p> <p>AI Assistant:</p> <p>The conversation between the therapist and the client revolves around the client's struggle with a recent rekindling with an ex-partner, leading to feelings of isolation, self-neglect, and emotional turmoil. The therapist acknowledges the client's feelings and suggests introspection to understand the underlying needs that the relationship fulfilled, and to explore if these needs can be met in other ways. The therapist encourages the client to identify values and deal-breakers in relationships, and to consider how they can attract the kind of people they want to be around. Additionally, the therapist prompts the client to reflect on their pattern of isolating and 'punishing' themselves, suggesting a shift towards more helpful behaviors and self-compassion. The therapist also emphasizes the importance of seeking support from trusted individuals.</p>

Figure 4: Examples of functionalities, including proposing responses, extracting relevant resources, and analyzing and summarizing conversations.

Item	Label	Description
Concern	concern	A therapist conveys concern by showing a regard for and interest in the client. The therapist seems engaged and involved with the client and attentive to what the client has said. The therapist's voice has a soft resonance that supports and enhances the client's concerned expressions.
Expressiveness	<i>N/A (unused in a text-based environment)</i>	A therapist's voice demonstrates expressiveness when the therapist speaks with energy and varies the pitch of his or her voice to accommodate the mood or disposition of the client.
Resonate or capture client feelings	resonate	A therapist resonates with or captures the intensity of the client's feelings when he or she speaks with a tone and emphasis that matches the client's emotional state or pitches words or phrases that underscore how the client feels.
Warmth	warmth	A therapist resonates with or captures the intensity of the client's feelings when he or she speaks with a tone and emphasis that matches the client's emotional state or pitches words or phrases that underscore how the client feels.
Attuned to the client's inner world	attuned	A client's inner world is defined as the client's feelings, perceptions, memories, meanings, bodily sensations, and core values. A therapist is attuned to a client's inner world when they provide moment-to-moment verbal acknowledgment of the client's expressions. These acknowledgments suit, agree with, or support the client's mood and reflections. The therapist is attentive to nuances of meaning and feeling conveyed in a client's statements beyond surface content and shows a genuine understanding of the client's inner world.
Understanding cognitive framework	cognitive	A therapist demonstrates an understanding of the client's cognitive framework and meanings when he or she clearly follows what the client has said and accurately reflects this understanding to the client. In short, the therapist and client are on the same page. The therapist is careful to provide ample opportunities for the client to state his or her views in order to permit the fullest and most accurate understanding of the client. The interaction conveys that the therapist values knowing what the client means or intends by his or her statements without predetermination or judgment.
Understanding feelings/inner experience	understanding	A therapist conveys an understanding of a client's feelings and inner experience when they show a sensitive appreciation and gentle caring for the client's emotional state. A therapist provides ample opportunities for the client to explore his or her emotional reactions. The therapist accurately reflects how the client feels by appropriately labeling feeling states with words (e.g., anger, sadness, frustration, etc.) or metaphors (e.g., "It's as if you are pent up and feel about to explode") to clarify and crystallize for the client what they are experiencing emotionally.
Acceptance of feelings/inner experiences	acceptance	A therapist shows acceptance of the client's feelings and inner experience when he or she validates the client's experience and reflects the client's feelings without judgment or a dismissive attitude. The therapist is unconditionally open to and respectful of how the client feels. The therapist's stance is one of genuineness and honesty instead of seemingly feigning concern or appearing inauthentic.
Responsiveness	<i>N/A (unused in a text-based environment)</i>	A therapist shows responsiveness to the client by adjusting his or her responses to the client's statements or nonverbal communications during the conversation. The therapist follows the client's lead in the conversation instead of trying to steer the discussion to the therapist's agenda or interests.

Table 4: Adapted TES rating items and their descriptions (Decker et al., 2013).











Figure 5: Histograms of rating items for responses generated by GPT-4o (first row), Llama 3 70b (second), Llama 3 8b (third), and professional therapists (last), as evaluated by GPT-4o.

Figure 6: Histograms of rating items for responses generated by GPT-4o (first row), Llama 3 70b (second), Llama 3 8b (third), and professional therapists (last), as evaluated by human raters.