

Bhirkuti's Test of Bias Acceptance: Examining in Psychometric Simulations

Aneel Bhusal, Todd D. Little (Texas Tech University)

Abstract

*This study introduces **Bhirkuti's Test of Bias Acceptance**, a systematic graphical framework for evaluating bias and determining its acceptability under varying experimental conditions. Absolute Relative Bias (ARB), while useful for understanding bias, is sensitive to outliers and population parameter magnitudes, often overstating bias for small values and understating it for larger ones. Similarly, Relative Efficiency (RE) can be influenced by variance differences and outliers, occasionally producing counterintuitive values exceeding 100%, which complicates interpretation. By addressing the limitations of traditional metrics such as Absolute Relative Bias (ARB) and Relative Efficiency (RE), the proposed graphical methodology framework leverages ridgeline plots and standardized estimate to provide a comprehensive visualization of parameter estimate distributions. Ridgeline plots done this way offer a robust alternative by visualizing full distributions, highlighting variability, trends, outliers, descriptives and facilitating more informed decision-making. This study employs multivariate Latent Growth Models (LGM) and Monte Carlo simulations to examine the performance of growth curve modeling under planned missing data designs, focusing on parameter estimate recovery and efficiency. By combining innovative visualization techniques with rigorous simulation methods, **Bhirkuti's Test of Bias Acceptance** provides two methods of versatile and interpretable toolset for advancing quantitative research in bias evaluation and efficiency assessment.*

Absolute Relative Bias (ARB)

Absolute Relative Bias (ARB) is the measure determined by taking the difference between the true population parameter (θ) and the average model-estimated parameter ($\hat{\theta}$) across all successfully converged simulations (refer to Equation 1). This difference is then divided by the population parameter, and the absolute value of the resulting proportion is calculated. The resulting ARB value indicates the proportional deviation of the estimated parameter from the true population value. Bias levels exceeding 10% are regarded as unacceptable, as supported by prior research (Muthén et al., 1987; Garnier-Villarreal et al., 2014; Lang et al., 2020; McArdle & Epstein, 1987; Moore et al., 2020). ARB plays a vital role in quantifying the degree of bias introduced into parameter estimates under various experimental conditions. It helps evaluate the feasibility of using a specific study design and provides insights into conditions that may cause parameter estimates to be biased. However, ARB is influenced by the size of the population parameter, potentially overestimating bias for smaller values and underestimating it for larger ones.

$$ARB_{\theta} = \frac{|\hat{\theta} - \theta|}{\theta} * 100\% \quad (1)$$

One limitation of Absolute Relative Bias (ARB) is its sensitivity to statistical outliers and the magnitude of the population parameter. Outliers in the model-estimated parameters ($\hat{\theta}$) can disproportionately influence the ARB calculation, skewing the results and potentially


exaggerating or minimizing the perceived bias. This sensitivity is particularly problematic in scenarios where parameter estimates vary widely, as outliers may not represent the typical behavior of the data. Additionally, ARB depends on the magnitude of the true population parameter (θ), which can lead to distortions in bias interpretation. Specifically, for smaller population parameter values, even minor deviations in estimates can result in a disproportionately large ARB, overstating the degree of bias. Conversely, for larger population parameter values, similar deviations might appear relatively small, leading to an underestimation of bias. These limitations underscore the importance of interpreting ARB within the context of the data and considering complementary methods to account for its susceptibility to outliers and parameter scaling effects.

Latent growth curve models

This study employs a two-construct multivariate Latent Growth Model (LGM) based on Rhemtulla et al. (2014) to examine the joint evolution of Bullying (B) and Homophobic Teasing (H) across five equally spaced time points. The model replicates the structure and population parameter values from Figure 11.7 in Little (2024; see Figure 3.1) to ensure consistency and facilitate comparisons. A key feature is the inclusion of correlated latent intercepts and slopes for both constructs, highlighting their interdependencies. Simulated data generated under this framework is analyzed using LGCM techniques to estimate growth trajectories, construct relationships, and factor loadings, with results compared to the original parameters. Variations may arise due to sampling error or simulation limitations. Each construct uses three indicators per time point, adhering to the just-identified construct principle to enable precise evaluation under constrained conditions (Little, 2024).

Figure 1

Proposed Growth Curve Model for Simulation



Note: From Figure 11.7, in *Longitudinal Structural Equation Modeling* (p. 398), by T. D. Little, 2024, Guilford Publications. Copyright 2024 by T. D. Little. Reprinted with permission.

Simple Wave Missing Design (SWMD-6)

Graham et al. (2001) introduced a simple wave missing design comprising six distinct participant groups. Among these, five groups exhibit systematic wave-level missingness, while Group 1 follows a complete data collection pattern across all waves. This group is particularly beneficial for ensuring convergence when using Full Information Maximum Likelihood (FIML) estimation to fit longitudinal models. The study underscores the importance of strategically designing data collection patterns to enhance convergence and accuracy in FIML-based longitudinal modeling.

Table 1
Simple Wave Missing Design with Six Groups (SWMD-6)

Random Group	Occasion Of Measurement				
	Time 1	Time 2	Time 3	Time 4	Time 5
1	√	√	√	√	√
2	√	√	√	√	X
3	√	√	√	X	√
4	√	√	X	√	√
5	√	X	√	√	√
6	X	√	√	√	√

Note: Within each time occasion, √ = data present, and X = data missing; (Graham et al., 2001; Little, 2024)

Full-Information Maximum Likelihood (FIML)

Full-Information Maximum Likelihood (FIML) is a statistical approach for handling missing data by estimating parameters directly from the available data, without imputing missing values. Rather than predicting or filling in the missing values, FIML uses the observed data to estimate overall patterns through a process called maximum likelihood estimation. This method computes a likelihood function for each case, even when some values are missing, and adjusts these calculations to account for the missing data, resulting in accurate overall estimates (Enders & Bandalos, 2001). FIML is most effective when the missing data mechanism is either Missing at Random (MAR) or Missing Completely at Random (MCAR), where it provides unbiased parameter estimates. However, if the data are Missing Not at Random (MNAR), the method may yield biased results. Enders (2022) notes that for large sample sizes, FIML and Multiple Imputation (MI) tend to produce similar outcomes, particularly when a sufficient number of imputations are used (e.g., 100 or more).

Simulation Design and Conditions

This study examines the performance of an alternative measure through simulated complete datasets and Graham’s wave planned missing data design, employing Full Information Maximum Likelihood (FIML) to analyze data generated from a proposed growth model. The model investigates the relationships between initial levels (intercepts) and rates of change (slopes) for two constructs, Bullying and Homophobic Teasing, measured across five time points. Population values and theoretical frameworks are drawn from Espelage et al. (2012), Rhemtulla et al. (2014), and Little (2024). The initial slope correlation between the constructs was set at 0.55, following Rhemtulla et al. (2014), but varied across simulation conditions. Other statistical properties, including means, variances, and covariances, were held constant to ensure analytical consistency.

The constructs were modeled with 10 latent variables (five time points for each construct), each measured by three observed indicators. Consistent with Meredith's (1993) guidelines, measurement invariance was achieved by holding factor loadings and intercepts constant over time, while allowing residual variances to vary and covary across time points. This ensured stable measurement while accommodating temporal changes in residuals. Slope loadings were fixed at 0 for the first time point, 1.0 for the second, and freely estimated for subsequent points. Latent intercepts were given fixed loadings of 1.0 across all time points, allowing the model to capture the observed rates of change. Population slope values for Bullying were 1.32, -0.83, and -0.64 for the third, fourth, and fifth time points, respectively, while for Homophobic Teasing, they were 1.45, 0.68, and -0.01. These benchmarks guided model estimation but were adjusted based on observed data for improved accuracy.


Simulated data were generated for 24 conditions by crossing three levels of latent slope correlation ($\rho_{s1,s2} = 0.1, 0.3, 0.55$) with eight sample sizes ($n=40, 60, 80, 100, 300, 500, 800, 1000$). Each condition was repeated 5,000 times to ensure robust parameter estimates. FIML was applied to estimate slope correlations under these conditions, with a focus on its ability to recover parameters accurately in planned missingness scenarios with SWMD-6. The study does not address unplanned missingness, as its primary focus is on leveraging ridgeline plots and standardized estimate standardization to provide a comprehensive visualization of parameter estimate distributions. It is designed to evaluate planned missing data designs exclusively to assess the performance of the proposed alternative measure to better understand the bias.

All simulations and analyses were conducted in R (R Core Team, 2024) using the RStudio integrated development environment (RStudio Team, 2024). Growth Curve Modeling was performed with the lavaan package (Rosseel, 2012), which facilitated the specification and estimation of growth curve models for both complete and planned missing datasets. A nested iteration structure automated the simulation process, varying experimental conditions and extracting key parameters. The estimates produced will be stored in a data frame for post-simulation analysis and will be used to evaluate Absolute Relative Bias (ARB), Relative Efficiency (RE), and the proposed alternative graphical metric. This analysis leverages ridgeline plots and standardized estimate standardization to provide a comprehensive visualization of parameter estimate distributions.

Exploring Alternatives to Absolute Relative Bias (ARB)

While Absolute Relative Bias (ARB) has been widely used to assess the proportional deviation of parameter estimates from their true population values, its limitations have prompted the exploration of alternative metrics (Muthén et al., 1987; Garnier-Villarreal et al., 2014; Lang et al., 2020; McArdle & Epstein, 1987; Moore et al., 2020). However, ARB's sensitivity to outliers and its dependency on the magnitude of the population parameter can distort the evaluation of bias. Specifically, smaller population parameter values can lead to inflated bias estimates, while larger values may understate the true bias which can be seen in Figure 2.

Figure 2




These findings highlight the importance of larger sample sizes and larger correlations in reducing parameter bias, although none of the methods achieved negligible bias levels as defined by the 10% threshold. While the absolute relative bias metric helps make bias comparable across parameters, changes in population value magnitude will have different bias values for the same absolute difference which overstates the degree of bias for small population values. The population parameter for slope correlation was set at .1, .3, and .55. This value is used in the denominator of the standardized bias computation. Consequently, an absolute bias of just .005 translates into 5%, 1.6%, and .9% absolute relative bias, respectively. While the ARB metric helps to make bias comparable across parameters, it can overstate the degree of bias for small population values as for .1 value. This characteristic of ARB occurs because changes in population estimate magnitude result in different bias values for the same absolute difference, making the bias seem more significant when the population magnitude value is smaller. To address these issues, alternative approaches aim to provide more robust and interpretable measures of bias, especially in complex modeling scenarios.

Relative Efficiency (RE)

Relative Efficiency (RE), which evaluates the efficiency of the planned missing data design compared to a complete dataset. RE is calculated by dividing the variance of parameter estimates in the complete data by the variance of parameter estimates in the incomplete data (see Equation 3). RE values range from 0 to 1, where values closer to 1 indicate minimal efficiency loss due to missing data. An RE of 0.80, for instance, suggests that the incomplete data estimates perform as efficiently as if the complete data had only 80% of its original sample size (Wu et al., 2016; Garnier-Villarreal et al., 2014; Rhemtulla et al., 2014).

$$RE_{\hat{\theta}} = \frac{var(\hat{\theta})_{complete}}{var(\hat{\theta})_{incomplete}} * 100\% \quad (2)$$

Figure 3



Note: Red line is at y=100.


This measure is instrumental in determining how much efficiency is retained under planned missingness conditions and whether the design remains viable (Wu et al., 2016). Efficient designs typically achieve RE values of 0.90 or higher across all parameters, although the acceptable threshold may vary depending on the study's tolerance for efficiency loss (Wu et al., 2016). However, RE is sensitive to variance, and extreme values or outliers may distort its accuracy, leading to potentially misleading efficiency estimates.

Relative Efficiency (RE) values exceeding 100% can complicate interpretation, as they suggest the evaluated condition produces lower variance in parameter estimates than the reference condition. This challenges the assumption that the reference condition, such as complete data, represents the standard for minimal variance. Such values often result from methodological artifacts, like unequal sample distributions or differences in admissible solutions, rather than genuine improvements in efficiency. Since RE is typically interpreted as the proportion of retained information relative to the reference condition, values over 100% may misrepresent efficiency and should be contextualized carefully to avoid misleading conclusions. The complexity of interpreting ARB and RE highlights the need for a balanced, nuanced approach to decision-making. We provide a systematic graphical visual approach in making the decision-making.

Visual Distribution of Parameter Estimates

Ridgeline plots serve as a powerful visualization tool for examining the distribution of parameter estimates across multiple groups (Wilke, 2021). In this study, they are utilized to analyze estimates generated from Latent Growth Models (LGM) via Monte Carlo simulations, providing valuable insights into variations under different experimental conditions. These plots effectively depict trends, patterns, outliers and descriptives, enabling researchers to assess the consistency and reliability of parameter estimates. Unlike traditional metrics such as Absolute Relative Bias (ARB) and Relative Efficiency (RE), which can be sensitive to outliers or parameter magnitudes, ridgeline plots visualize the full distribution, offering a more robust and intuitive perspective. By directly presenting data distributions, ridgeline plots help address limitations of conventional summary metrics, facilitating more informed decision-making about the performance of the proposed design and missing data techniques. Furthermore, the ability to compare descriptive statistics—such as means, medians, and variances—within the plots enhances the capacity for making nuanced judgments, providing a clearer understanding of systematic effects and the underlying data-generating processes. This approach combines visual accessibility with analytical depth, supporting robust evaluation and interpretation of the simulation results.

Figure 4




Note: Dots are the distribution means and bars show ± 1 SD from the mean. The vertical line represents the population value. Distribution of estimate for 24 conditions. “V:” variance of the estimate produced and “M:” mean of the estimate produced.

The figure 4 illustrates the distribution of estimated slope-slope correlations across 24 conditions, varying by sample size and population correlation values ($\rho=0.1,0.3,0.55$). An FIML with SWMD-6 can be considered acceptable if the population value (indicated by the vertical line) passes through the mean of the distribution (marked by dots), as this reflects unbiased estimation. A deviation of up to 10% from the population value is often deemed acceptable. However, this threshold introduces inconsistencies across different population values, as 10% of $\rho=0.1,0.3$, and 0.55 corresponds to varying levels of tolerance, which may result in problematic evaluations. This inconsistency can be resolved with two methods:

Method A: Using ridgeline plots to examine the distribution of absolute relative bias (ARB) across multiple groups for all parameter estimates. This approach visualizes the distribution of bias obtained from Equation 1 for each individual estimate within each group, providing insights into bias variations across different conditions.

Figure 5




Note: Dots are the distribution means. Red dotted vertical line represents at $x=0$. Blue dashed vertical line represents $x=-0.1$ and $x=0.1$. Distribution of standardized estimate for 24 conditions. “M:” mean value of the estimate.

Method B: Standardized estimate similarly to Z-score, calculated as

$$\text{Standardized estimate} = \frac{\text{Estimate Produced} - \text{Population Value}}{\text{Standard Deviation of Estimate}}$$


In simulation studies with 5000 iterations per condition, this large number of replications is often considered the gold standard. It ensures stable and reliable estimates by minimizing random variability and providing a robust basis for evaluating performance metrics. Standardized estimate calculated under these conditions reflects an accurate measure of bias and consistency across varying population values and conditions, making it a reliable tool for assessing the quality of the estimation method and comparison robust.

Figure 6




Note: Dots are the distribution means. Red dotted vertical line represents at $x=0$. Blue dashed vertical line represents $x=-0.1$ and $x=0.1$. Distribution of standardized estimate for 24 conditions. “V:” variance of the estimate produced and “M:” mean of the estimate produced.

Figure 7



Note: Red dotted vertical line represents at $x=0$. Blue dashed vertical line represents $x=-0.1$ and $x=0.1$. Distribution of standardized estimate for 24 conditions. “M:” mean value of the estimate.

Figure 8



Note: Note: Red dotted vertical line represents at $x=0$. Blue dashed vertical line represents $x=-0.1$ and $x=0.1$. Distribution of standardized estimate for 24 conditions. “M:” mean value of the estimate.

In Figure 7 and 8, the red dashed line at $x = 0$ represents no bias, while the dotted lines at $x = 0.1$ and $x = -0.1$ delineate the acceptable bias range, allowing for a 10% margin. This method addresses limitations associated with the sensitivity of Absolute Relative Bias (ARB) while looking at as percentage and variances to outliers, particularly in relation to the magnitude of population parameters. Bias is considered acceptable if any measure of central tendency, such as the mean, median, or mode, falls within or touches the acceptable range. While the mean is the recommended measure, the median and mode can also offer valuable insights into the distribution of estimates depending on the context of the study. Both method standardized estimate and ARB estimate are used to ensure comparability across groups by placing all distributions on a common scale. This approach provides consistency across all parameters being tested and preserves the relative distribution and structure of the data. However, it can introduce visual distortions: groups with smaller original variances appear broader after standardization due to stretching, while groups with larger variances appear narrower due to compression. While standardization enhances the ability to directly compare distributions across groups, it can alter the visual spread of the data. This method strikes a balance between maintaining consistency across parameters and improving the interpretability of bias and variance estimates, while acknowledging the inherent trade-offs in visual representation.

Conclusion

Bhirkuti's Test of Bias Acceptance points out the limitations of traditional metrics such as Absolute Relative Bias (ARB) and Relative Efficiency (RE) in evaluating bias and efficiency across experimental conditions. ARB, while useful for standardizing bias, often overemphasizes discrepancies for small population parameters and underrepresents them for larger ones, potentially skewing interpretations. Similarly, Relative Efficiency (RE), while effective in quantifying the efficiency of planned missing data designs, can be influenced by differences in variances, outliers, or methodological artifacts. These factors may result in counterintuitive values exceeding 100%, complicating interpretation, particularly when dealing with variances expressed with decimal precision.

In contrast, ridgeline plots provide a more robust and comprehensive alternative. By visualizing the full distribution of parameter estimates, these plots highlight variability, trends, and outliers across conditions. This approach addresses the sensitivity of ARB to population parameter magnitudes and the variance distortions that can misrepresent RE. Additionally, ridgeline plots allow researchers to directly assess the acceptability of bias within predefined thresholds (e.g., $\pm 10\%$), offering a clearer perspective on central tendencies like the mean, median, or mode while acknowledging the impact of data structure and variance behavior.

Standardization estimate or ARB estimate further enhances interpretability by placing distributions on a common scale. Although these methods can visually distort the perceived spread of data (e.g., broadening smaller variances and compressing larger ones), they strike a balance between consistency and comparability, making cross-group evaluations more meaningful. By combining ridgeline plots with these standardization techniques, researchers gain a nuanced toolset for analyzing parameter bias and efficiency. This approach avoids oversimplifying the data by using single numbers like ARB and RE. Instead, it helps to better understand patterns and unusual results in different experimental conditions better.

Ultimately, ridgeline plots empower more informed decision-making and enhance the interpretability of complex data, addressing the challenges posed by traditional metrics and offering a versatile solution for evaluating parameter performance under varying conditions. This methodological test for evaluating bias and determining its acceptability is formally termed **Bhirkuti's Test of Bias Acceptance**, providing a systematic graphical framework for assessing the reliability of parameter estimates under varying conditions. Although this test is performed within the context of psychometric simulation studies, it holds significant potential for application in other research study domains where bias is critical in simulation. Its robust methodology and adaptability make it a valuable tool for analyzing and interpreting data across diverse fields, broadening its impact beyond psychometrics.

Additional Information

Analysis of additional parameters and additional information is available on request to Aneel Bhusal abhusal@ttu.edu.

References

- Carsey, T. M., & Harden, J. J. (2013). *Monte Carlo simulation and resampling methods for social science*. Sage Publications.
- Dempster, A. P., Laird, N. M., & Rubin, D. B. (1977). Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1), 1–38. <http://www.jstor.org/stable/2984875>
- Enders, C. K. (2022). *Applied missing data analysis* (2nd ed.). The Guilford Press.
- Enders, C. K., & Bandalos, D. L. (2001). The relative performance of full information maximum likelihood estimation for missing data in structural equation models. *Structural Equation Modeling*, 8(3), 430–457. https://doi.org/10.1207/S15328007SEM0803_5
- Garnier-Villareal, M., Rhemtulla, M., & Little, T. D. (2014). Two-method planned missing designs for longitudinal research. *International Journal of Behavioral Development*, 38(5), 411–422. <https://doi.org/10.1177/0165025414542711>
- Gower, J. C. (1966). Some distance properties of latent root and vector methods used in multivariate analysis. *Biometrika*, 53(3/4), 325–338. <https://doi.org/10.2307/2333639>
- Graham, J. W., Taylor, B. J., & Cumsille, P. E. (2001). Planned missing-data designs in analysis of change. In L. M. Collins & A. G. Sayer (Eds.), *New methods for the analysis of change* (pp. 335–353). American Psychological Association. <https://doi.org/10.1037/10409-011>
- Graham, J. W., Taylor, B. J., Olchowski, A. E., & Cumsille, P. E. (2006). Planned missing data designs in psychological research. *Psychological Methods*, 11, 323–343. <https://doi.org/10.1037/1082-989X.11.3.323>
- Hertzog, C., Lindenberger, U., Ghisletta, P., & Von Oertzen, T. (2006). On the power of multivariate latent growth curve models to detect correlated change. *Psychological Methods*, 11(3), 244–252. <https://doi.org/10.1037/1082-989X.11.3.244>
- Jia, F., Moore, W. G., Kinai, R., Crowe, K. S., Schoemann, A. M., & Little, T. D. (2014). Planned missing data designs with small sample sizes: How small is too small? *International Journal of Behavioral Development*, 38(5), 435–452. <https://doi.org/10.1177/0165025414531095>
- Lang, K. M., Moore, E. W. G., & Grandfield, E. M. (2020). A novel item-allocation procedure for the three-form planned missing data design. *MethodsX*, 7, 100941. <https://doi.org/10.1016/j.mex.2020.100941>

- Little, R. J. A., & Rubin, D. B. (2019). *Statistical analysis with missing data* (3rd ed.). John Wiley & Sons.
- Little, T. D. (2024). *Longitudinal structural equation modeling* (2nd ed.). The Guilford Press.
- Little, T. D., & Rhemtulla, M. (2013). Planned missing data designs for developmental researchers. *Child Development Perspectives*, 7(4), 199–204. <https://doi.org/10.1111/cdep.12043>
- McArdle, J. J., & Epstein, D. (1987). Latent growth curves within developmental structural equation models. *Child Development*, 58(1), 110–133. <https://doi.org/10.2307/1130295>
- Meredith, W., & Tisak, J. (1990). Latent curve analysis. *Psychometrika*, 55(1), 107–122. <https://doi.org/10.1007/BF02294746>
- Mistler, S. A., & Enders, C. K. (2012). Planned missing data designs for developmental research. In B. Laursen, T. D. Little, & N. A. Card (Eds.), *Handbook of developmental research methods* (pp. 742–754). The Guilford Press.
- Moore, E. W. G., Lang, K. M., & Grandfield, E. M. (2020). Maximizing data quality and shortening survey time: Three-form planned missing data survey design. *Psychology of Sport and Exercise*, 51, 101701. <https://doi.org/10.1016/j.psychsport.2020.101701>
- R Core Team. (2024). *R: A language and environment for statistical computing* (Version 4.4.1) [Software]. R Foundation for Statistical Computing. <https://www.r-project.org/>
- Rhemtulla, M., Jia, F., & Little, T. D. (2014). Planned missing designs to optimize the efficiency of latent growth parameter estimates. *International Journal of Behavioral Development*, 38(5), 423–434. <https://doi.org/10.1177/0165025413514324>
- Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2), 1–36. <https://doi.org/10.18637/jss.v048.i02>
- RStudio Team. (2024). *RStudio: Integrated Development Environment for R* (Version 2024.4.2.764) [Software]. Posit Software, PBC. <https://posit.co/>
- Rubin, D. B. (1987). *Multiple imputation for nonresponse in surveys*. John Wiley & Sons.
- Savalei, V., & Kolenikov, S. (2008). Constrained versus unconstrained estimation in structural equation modeling. *Psychological Methods*, 13(2), 150–170. <https://doi.org/10.1037/1082-989X.13.2.150>
- Wilke, C. O. (2021). *ggridges: Ridgeline plots in 'ggplot2'* (Version 0.5.4) [Computer software]. <https://CRAN.R-project.org/package=ggridges>
- Wu, W., Jia, F., Rhemtulla, M., & Little, T. D. (2016). Search for efficient complete and planned missing data designs for analysis of change. *Behavior Research Methods*, 48(3), 1047–1061. <https://doi.org/10.3758/s13428-015-0629-5>