

GENERALIZED FAST KRASNOSEL'KII–MANN METHOD WITH PRECONDITIONERS

RADU I. BOȚ*, ENIS CHENCHENE*, AND JALAL M. FADILI†

ABSTRACT. We study accelerated Krasnosel'kii–Mann-type methods with preconditioners in both continuous and discrete time. From a continuous time model, we derive a generalized fast Krasnosel'kii–Mann method, providing a new yet simple proof of convergence that allows for unprecedented flexibility in parameter tuning. Our analysis unifies inertial and anchoring acceleration mechanisms and offers a broad range of parameter choices, which prove beneficial in practice. Additionally, we extend our analysis to the case where preconditioners are allowed to be degenerate, unifying the treatment of various splitting methods and establishing the weak convergence of shadow sequences.

Keywords: Inertial Accelerations, Nesterov Acceleration, Halpern Method, Fast-Krasnosel'kii–Mann Method, Degenerate Proximal Point Algorithm

1. INTRODUCTION

Let $T: H \rightarrow H$ be a nonexpansive operator on the real Hilbert space H . Our objective is to solve the fixed-point problem:

$$(1.1) \quad \text{Find } x \in H \text{ such that } x = T(x),$$

with fast methods in continuous and discrete time, and apply these results to splitting methods providing convergence of the so-called shadow sequences (to be made precise later).

If $T = I - \nabla f$, where $f: H \rightarrow \mathbb{R}$ is a convex and smooth function whose set of minimizers is nonempty, (1.1) is equivalent to minimizing f . The most popular accelerated method to minimize f in discrete time is the celebrated Nesterov's algorithm [36]. Su, Boyd, and Candès in [42] later discovered that Nesterov's algorithm can be understood as a specific finite differences discretization of the following ODE

$$(1.2) \quad \ddot{x}(t) + \frac{\alpha}{t} \dot{x}(t) + \nabla f(x(t)) = 0, \quad \text{for } t \geq t_0 > 0$$

with $\alpha = 3$ and initial data $(x(t_0), \dot{x}(t_0)) = (x_0, v_0) \in H^2$. Although the convergence of $x(t)$ to a minimizer of f remains an open problem for $\alpha = 3$ (except in one dimension where convergence holds [6]), the trajectories of (1.2) satisfy $f(x(t)) - \min f = O(t^{-2})$ as $t \rightarrow +\infty$, which is an improvement over the $o(t^{-1})$ rate of the standard gradient flow. Later, in [7, 23], it has been shown that replacing the critical parameter $\alpha = 3$ with any $\alpha > 3$ ensures the weak convergence of $x(t)$ to a minimizer of f , and improves the rate of convergence for the function values from $O(t^{-2})$ to $o(t^{-2})$.

However, these inertial acceleration methods often suffer from significant oscillations. This issue was addressed by Attouch, Peypouquet, and Redont [9], and later by Attouch, Chbani, Fadili, and Riahi [5, 4], who proposed to couple the asymptotically vanishing viscous damping term in (1.2) to an additional *geometric Hessian-driven damping term*, previously introduced in [2, 8], resulting in the system

$$(1.3) \quad \ddot{x}(t) + \frac{\alpha}{t} \dot{x}(t) + \beta(t) \frac{d}{dt} \nabla f(x(t)) + \nabla f(x(t)) = 0, \quad \text{for } t \geq t_0,$$

which, for $\alpha > 3$ and appropriate coefficient $\beta(t) > 0$, has the important benefit of retaining the $o(t^{-2})$ rate on the objective, while ensuring the weak convergence of trajectories, and attenuating oscillations.

The research of RIB and EC has been supported by the Austrian Science Fund (FWF), projects W 1260 and P 34922-N, respectively.

The Hessian-driven damping proves advantageous not only for optimization but also when ∇f is replaced by any monotone and Lipschitz continuous operator $Q: H \rightarrow H$:

$$(1.4) \quad \ddot{x}(t) + \frac{\alpha}{t}\dot{x}(t) + \beta(t)\frac{d}{dt}Q(x(t)) + b(t)Q(x(t)) = 0, \quad \text{for } t \geq t_0.$$

For instance, to solve (1.1), it is natural to consider $Q := I - T$, which is indeed monotone and Lipschitz, and the equilibria of (1.4) are solutions to (1.1).

Although the system in (1.3) generates fast converging trajectories for a range of admissible parameters (cf. [4, Theorem 2.1]), when one turns to the general operator setting (1.4) the situation becomes significantly more delicate. A notable contribution in this direction was made by BoȚ, Csetnek and Nguyen in [14], who studied (1.4) with $b(t) := \frac{1}{2}(\dot{\beta}(t) + \frac{\alpha}{t}\beta(t))$, the so-called fast Optimistic Gradient Descent Ascent (Fast-OGDA) system. In particular, they showed that when β is constant and $\alpha > 2$, this dynamical system generates trajectories weakly converging to a zero of Q (provided that the set of zeros of Q is non-empty) with a convergence rate of $\|Q(x(t))\| = o(t^{-1})$ as $t \rightarrow +\infty$. Therefore, (1.4) can be seen as an accelerated version of the first order system

$$(1.5) \quad \dot{x}(t) + Q(x(t)) = 0, \quad \text{for } t \geq t_0.$$

Indeed, when Q is a cocoercive operator, it is known that each solution trajectory generated by (1.5) weakly converges to a zero of Q with a convergence rate of $\|Q(x(t))\| = o(t^{-\frac{1}{2}})$ as $t \rightarrow +\infty$ [3].

It turns out that the specific choice $b(t) := \frac{1}{2}(\dot{\beta}(t) + \frac{\alpha}{t}\beta(t))$ in (1.4) considered by [14] is ultimately due to the impossibility of exploiting function values in the Lyapunov analysis. This immediately leads to the natural question:

Can Fast-OGDA in [14] be extended to accommodate more general parameters b , enabling new fast and flexible numerical algorithms for solving (1.1)?

In this paper, we show that is possible to consider the more general and flexible choice

$$(1.6) \quad b(t) := \eta\dot{\beta}(t) + \alpha_\eta\frac{\beta(t)}{t}, \quad \text{for } \eta \in (0, 1) \quad \text{and} \quad \alpha_\eta := \eta + (1 - \eta)(\alpha - 1),$$

which recovers [14] by setting $\eta = \frac{1}{2}$. Our convergence guarantees for a general parameter $\eta \in (0, 1)$ cover those in [14]: in particular, we establish that, e.g., if $\alpha > 2$ and β is constant, we have weak convergence of the trajectories to zeros of Q with a convergence rate of $\|Q(x(t))\| = o(t^{-1})$ as $t \rightarrow +\infty$.

In the recent work [16], BoȚ and Nguyen have further highlighted that the Fast-OGDA system is in turn related to a seemingly completely different mechanism: the Tikhonov regularization, also known as anchoring mechanism. Specifically, for the choice $\beta(t) := \beta_0 t^{\alpha-2}$ with $\beta_0 > 0$, it follows from [16, Proposition 3.1 and Remark 3.3] that the dynamical system (1.4), is equivalent to:

$$(1.7) \quad \dot{x}(t) + \beta(t)Q(x(t)) + \frac{\alpha - 1}{t}(x(t) - v) = 0, \quad \text{for } t \geq t_0,$$

with anchor point

$$v := x_0 + \frac{t_0}{\alpha - 1}(\dot{x}(t_0) + \beta(t_0)Q(x_0)).$$

(1.7) is nothing but the Tikhonov regularization of the monotone operator flow (1.5), see [21, 39, 26], that in fact not only generates trajectories that converge strongly to the projection of v onto $\text{zer } Q$, but also exhibits the rate $O(t^{-(\alpha-1)})$ as $t \rightarrow +\infty$. The system (1.7) can also be recognized as the continuous time model of the celebrated Halpern method which is a classical method in fixed-point theory [16, 43]. A simplified version of (1.7) was studied in [43] in finite-dimensional spaces. In the discrete time setting, Sabach and Shtern in [41] and independently Lieder in [33], proved that Halpern's method achieves the convergence rate of $O(k^{-1})$ as $k \rightarrow +\infty$ on the norm of the fixed-point residual. Interestingly, this cannot be improved to $o(k^{-1})$ in normed spaces, see [27]. The connection between Halpern's method and momentum-type algorithms have been observed previously in [27, 46, 45].

The $o(k^{-1})$ convergence rate on the norm of the fixed-point residual is in fact attained by Algorithm 1 that we propose with $\alpha > 2$. The latter is a generalization of the so-called Fast-Krasnosel'kiĭ–Mann method (Fast-KM) in [15], with general parameters $\eta \in (0, 1)$ and $\sigma > 0$. Algorithm 1 allows us to

Algorithm 1: (Generalized) Fast Krasnosel’kiĭ–Mann Method

Pick $x^0, x^1 \in H$, $\eta \in (0, 1)$, $\sigma > 0$, $\alpha \geq 2$;

Set $\alpha_\eta := \eta + (1 - \eta)(\alpha - 1)$;

for $k = 1, 2, \dots$ **do**

$$\left[\begin{array}{l} x^{k+1} = \left(1 - \frac{\alpha_\eta}{k + \sigma}\right) x^k + \frac{\alpha_\eta}{k + \sigma} T(x^k) + \left(1 - \frac{\alpha}{k + \sigma}\right) (T(x^k) - T(x^{k-1})) \end{array} \right.$$

solve (1.1), and can be understood as a discrete time counterpart of (1.4) with b as in (1.6), $\beta \equiv 1$, and $Q = I - T$. We strongly rely on this important connection to derive a simple convergence proof that matches almost exactly the continuous time approach, and allows us to cover the case of $\alpha = 2$, for which the method reduces to the Optimal Halpern Method [33]. This is a consequence of the first significant difference with the approach in [15], as Algorithm 1 is obtained with a *semi-implicit* discretization of (1.4), while Algorithm 2.3 in [15] is obtained through an explicit discretization, thus requiring a more involved analysis. Numerical experiments also show that considering $\eta < \frac{1}{2}$ does indeed have a beneficial effect on the convergence performance, surpassing the baseline method in several considered instances.

Since Algorithm 1 allows us to find fixed-points of nonexpansive operators, it emerges as a unified algorithmic framework to design accelerations for *operator splitting methods*, such as primal-dual [24], forward-backward [11], Douglas–Rachford [29, 34], Davis–Yin [28, 38], and several other extensions. Following the recent insights in [32, 17, 18], these methods can be studied as fixed-point iterations with respect to nonexpansive mappings in a renormed Hilbert space, see Section 4. In this context, the metric might actually be *degenerate*, i.e., only inducing a Hilbert seminorm, and thus the analysis requires a careful adaptation. We do so by delineating a *degenerate* Opial Lemma, a fundamental ingredient that allows us to adapt the convergence analysis of Algorithm 1 almost effortlessly to this degenerate setting. We apply this result to accelerate the convergence of a recent extension of the Douglas–Rachford splitting method [34, 18], ensuring the weak convergence of the so-called *shadow* sequences in an infinite-dimensional Hilbert space setting.

1.1. Contributions. Our main contributions are summarized as follows:

- We introduce a generalization of the Fast-OGDA system introduced by Boţ, Nguyen and Csetnek in [14] with b as in (1.6).
- We introduce Algorithm 1 as a discretization of the proposed continuous time model and propose a new convergence proof that matches almost exactly the continuous time approach. The method generalizes [15] integrating two degrees of freedom: it consider a general parameter $\sigma > 0$ and a general parameter $\eta \in (0, 1)$. Further, we cover the convergence analysis also with $\alpha = \sigma = 2$, which corresponds to the Optimal Halpern method [46], hence providing a unified analysis of *inertial* and *anchoring* (Tikhonov) acceleration mechanisms.
- We establish an Opial-type lemma with degenerate preconditioners, and adapt our convergence proof to this setting. We further discuss how this allows us to systematically obtain convergence results for several splitting methods, establishing the weak convergence of the so-called *shadow sequences* in infinite dimensional Hilbert spaces, thus closing a gap in [15].
- We present numerical experiments that demonstrate that the Fast-KM method with $\eta < \frac{1}{2}$ seemingly outperforms the baseline choice with $\eta = \frac{1}{2}$.

1.2. Notation and Preliminaries. In this section, we fix the notation and introduce some preliminary results on nonexpansive operators in Hilbert spaces and their connection to monotone operators; see [10] for a comprehensive account. An operator $T: H \rightarrow H$ is said to be *nonexpansive* if

$$(1.8) \quad \|T(x) - T(x')\| \leq \|x - x'\|, \quad \text{for all } x, x' \in H.$$

The set of *fixed-points* of T is the defined by $\text{Fix } T := \{x \in H \mid x = T(x)\}$. In this paper, we always assume that $\text{Fix } T \neq \emptyset$. If T is nonexpansive, the operator $T_s := sT + (1 - s)I$ for $s \in (0, 1)$ is said to

be *s-averaged*. $T_{1/2}$ is called firmly nonexpansive. It follows from [10, Remark 4.34(iv)] that $I - T_s$ is $\frac{1}{2s}$ -cocoercive, i.e., it satisfies the following inequality:

$$(1.9) \quad \langle x - x', (I - T_s)(x) - (I - T_s)(x') \rangle \geq \frac{1}{2s} \|(I - T_s)(x) - (I - T_s)(x')\|^2, \quad \text{for } x, x' \in H.$$

In particular, $I - T_s$ is a *monotone* operator. A set-valued mapping $A: H \rightarrow 2^H$ is *monotone* if it obeys

$$(1.10) \quad \langle a - a', u - u' \rangle \geq 0, \quad \text{for all } (u, a), (u', a') \in \text{gra } A,$$

where the set $\text{gra } A \subseteq H \times H$ is the *graph* of A and is defined by $\text{gra } A := \{(u, a) \mid a \in A(u)\}$. A monotone operator is called *maximal monotone* if its graph is maximal (with respect to the inclusions of sets) among the class of monotone operators. The inverse of a set-valued mapping A is denoted by A^{-1} and defined by $\text{gra } A^{-1} := \{(a, u) \mid (u, a) \in \text{gra } A\}$. The *resolvent* of A is the mapping $J_A := (I + A)^{-1}$, where I denotes the identity operator on H . From [10, Proposition 23.8], A is maximal monotone if and only if J_A is a single-valued mapping from H to H , which is 1-cocoercive, or simply *cocoercive* (or firmly nonexpansive). Moreover, it is clear that the operator $T := 2J_A - I$ is nonexpansive. Conversely, from [10, Corollary 23.9], if T is nonexpansive, there exists a maximal monotone operator A such that $T = 2J_A - I$. Furthermore, fixed-points of T correspond to *zeros* of A , i.e., elements of $\text{Fix } T = \text{zer } A := \{u \in H \mid 0 \in A(u)\}$. We also denote \rightharpoonup as weak convergence and \rightarrow as strong convergence.

2. CONTINUOUS TIME SYSTEM AND ANALYSIS

In this section, we introduce a general dynamical system that extends Fast-OGDA from [14]. Let $Q: H \rightarrow H$ be a monotone and Lipschitz continuous operator¹ and consider the dynamical system:

$$(2.1) \quad \ddot{x}(t) + \frac{\alpha}{t} \dot{x}(t) + \beta(t) \frac{d}{dt} Q(x(t)) + b(t) Q(x(t)) = 0, \quad t \geq t_0 > 0,$$

where b is defined for all $t \geq t_0$ by

$$(2.2) \quad b(t) := \eta \dot{\beta}(t) + \alpha_\eta \frac{\beta(t)}{t} = \eta \dot{\beta}(t) + (\eta + (1 - \eta)(\alpha - 1)) \frac{\beta(t)}{t},$$

with the additional parameter $\eta \in (0, 1)$. The system (2.1) recovers the choice in [14] by setting $\eta = \frac{1}{2}$, for which $\alpha_\eta = \frac{1}{2}\alpha$, and can be leveraged to approach solutions to (1.1) by setting $Q = I - T$, since indeed:

$$\text{zer } Q = \text{zer}(I - T) = \text{Fix } T.$$

We will need that (2.1) admits at least one strong global solution for every starting point $x(0) = x_0 \in H$, i.e., such that x and \dot{x} are locally absolutely continuous on $[t_0, +\infty)$, and x satisfies (2.1) *almost everywhere* (a.e.) on $[t_0, +\infty)$. In turn, this will entail that for every locally absolutely continuous trajectory $t \mapsto x(t)$ the trajectory $t \mapsto Q(x(t))$ is also locally absolutely continuous. In fact, using [31, Proposition 6.2.1], Lipschitz continuity of Q and mild conditions on $\beta(t)$, it can be shown that the system (2.1) has a unique strong solution trajectory. In fact, Lipschitz continuity of Q can even be weakened to Lipschitz continuous over the bounded subsets of H to show the existence and uniqueness of a maximal solution. The passage to a global solution would use a classical contradiction argument and boundedness of the trajectory that we will show in our Lyapunov analysis.

To study the second order dynamical system (2.1), we introduce, for $x^* \in \text{zer } Q \neq \emptyset$, the energy function $E^\lambda: [t_0, +\infty) \rightarrow \mathbb{R}_+$ that will play a fundamental role in our Lyapunov analysis,

$$(2.3) \quad \begin{aligned} E^\lambda(t) &:= \delta(t) \langle Q(x(t)), x(t) - x^* \rangle + \frac{\xi(t)}{2} \|Q(x(t))\|^2 + \frac{c}{2} \|x(t) - x^*\|^2 + \frac{1}{2} \|v^\lambda(t)\|^2, \\ v^\lambda(t) &:= \lambda(x(t) - x^*) + t(\dot{x}(t) + \eta\beta(t)Q(x(t))), \end{aligned}$$

for some constant $\lambda \geq 0$, and parameters given by

$$(2.4) \quad \delta(t) := (1 - \eta)\lambda\beta(t)t, \quad \xi(t) := (1 - \eta)\eta\beta(t)^2 t^2, \quad c := \lambda(\alpha - 1 - \lambda), \quad \text{for } t \geq t_0.$$

¹Note that the Lipschitz continuity of Q is only useful to establish the existence and uniqueness of a strong solution to (2.1) which in turn entails the almost everywhere differentiability of $t \mapsto Q(x(t))$.

This choice of the parameters allows us to establish the following result, whose proof is postponed to Appendix B.

Lemma 2.1. *Let $x^* \in \text{zer } Q \neq \emptyset$, $x: [t_0, +\infty) \rightarrow H$ be a solution trajectory to (2.1), and E^λ the energy function defined in (2.3). Then, for almost all $t \geq t_0$, we have*

$$(2.5) \quad \begin{aligned} \frac{d}{dt} E^\lambda(t) &= \lambda t(1-\eta) \left(\dot{\beta}(t) + \frac{2-\alpha}{t} \beta(t) \right) \langle Q(x(t)), x(t) - x^* \rangle - \frac{t}{2} (\alpha - 1 - \lambda) \|\dot{x}(t) + \beta(t)Q(x(t))\|^2 \\ &\quad + \beta(t)t^2 \left((1-\eta)\eta \left(\dot{\beta}(t) + \frac{2-\alpha}{t} \beta(t) \right) + \frac{1}{2} (\alpha - 1 - \lambda) \frac{\beta(t)}{t} \right) \|Q(x(t))\|^2 \\ &\quad - (1-\eta)t^2 \beta(t) \left\langle \dot{x}(t), \frac{d}{dt} Q(x(t)) \right\rangle - \frac{t}{2} (\alpha - 1 - \lambda) \|\dot{x}(t)\|^2. \end{aligned}$$

Lemma 2.1 naturally leads to consider the following *growth condition* for $t \mapsto \beta(t)$:

$$(2.6) \quad 0 \leq \frac{\dot{\beta}(t)t}{\beta(t)} \leq \alpha - 2 - \varepsilon, \quad \text{for } \varepsilon \in [0, \alpha - 2], \text{ and all } t \geq t_0,$$

which with a suitable rearrangement, writes equivalently as

$$(2.7) \quad \dot{\beta}(t) + \frac{2-\alpha}{t} \beta(t) \leq -\varepsilon \frac{\beta(t)}{t}, \quad \text{for all } t \geq t_0.$$

This makes the coefficient in front of the term $\|Q(x(t))\|^2$ negative and bounded away from zero at least for λ sufficiently close to $\alpha - 1$, hence leading to the desired energy decrease. This observation is crucial to establish the following convergence result.

Theorem 2.2. *Let $Q: H \rightarrow H$ be a monotone and Lipschitz continuous operator with $\text{zer } Q \neq \emptyset$. Let $x^* \in \text{zer } Q$, $x: [t_0, +\infty) \rightarrow H$ be a solution trajectory to (2.1) for some $\eta \in (0, 1)$ and $\alpha \geq 2$. Suppose that (2.6) holds with $\varepsilon \geq 0$. Then, it holds:*

$$(2.8) \quad \langle Q(x(t)), x(t) - x^* \rangle = O\left(\frac{1}{t\beta(t)}\right) \quad \text{and} \quad \|Q(x(t))\| = O\left(\frac{1}{t\beta(t)}\right) \quad \text{as } t \rightarrow +\infty.$$

Additionally, if $\alpha > 2$ and $\varepsilon > 0$, then the following holds:

(i) *We have the following integral bounds:*

$$\int_{t_0}^{\infty} \beta(t) \langle Q(x(t)), x(t) - x^* \rangle dt < +\infty, \quad \int_{t_0}^{\infty} t\beta(t)^2 \|Q(x(t))\|^2 dt < +\infty, \quad \int_{t_0}^{\infty} t \|\dot{x}(t)\|^2 dt < +\infty.$$

(ii) *$x(t)$ converges weakly to a zero of Q as $t \rightarrow +\infty$.*

(iii) *The following fast rates of convergence hold:*

$$\langle Q(x(t)), x(t) - x^* \rangle = o\left(\frac{1}{t\beta(t)}\right), \quad \|Q(x(t))\| = o\left(\frac{1}{t\beta(t)}\right) \quad \text{and} \quad \|\dot{x}(t)\| = o\left(\frac{1}{t}\right) \quad \text{as } t \rightarrow +\infty.$$

Proof. To lighten notation, we abbreviate (t) with t as a subscript. Additionally, we write Q_t instead of $Q(x(t))$ and ∂_t instead of $\frac{d}{dt}$. Setting $\lambda := \alpha - 1$ in (2.5), discarding the negative terms, we get using (2.7) and monotonicity of Q , that for almost all $t \geq t_0$:

$$\partial_t E_t^{\alpha-1} \leq -\varepsilon(\alpha-1)(1-\eta)\beta_t \langle Q_t, x_t - x^* \rangle - (1-\eta)t^2 \beta_t \langle \dot{x}_t, \partial_t Q_t \rangle \leq 0.$$

Thus, $E_t^{\alpha-1}$ is nonincreasing and since all the terms in (2.3) are nonnegative, we get

$$(\alpha-1)(1-\eta)t\beta_t \langle Q_t, x_t - x^* \rangle + \frac{\eta(1-\eta)}{2} \beta_t^2 t^2 \|Q_t\|^2 \leq E_{t_0}^{\alpha-1}, \quad \text{for all } t \geq t_0,$$

hence (2.8).

Let us now turn to claims (i)-(iii). Suppose that (2.6) holds for some $\varepsilon > 0$. Thus, (2.5) and monotonicity of Q yield for all $0 \leq \lambda \leq \alpha - 1$ and almost all $t \geq t_0$:

$$\partial_t E_t^\lambda \leq -\varepsilon\lambda(1-\eta)\beta_t \langle Q_t, x_t - x^* \rangle - \frac{t}{2} (\alpha - 1 - \lambda) \|\dot{x}_t\|^2 + \beta_t^2 t \left(-(1-\eta)\eta\varepsilon + \frac{1}{2} (\alpha - 1 - \lambda) \right) \|Q_t\|^2.$$

Let $\bar{\lambda} := \alpha - 1 - \varepsilon\eta(1 - \eta)$. Observe that $\bar{\lambda} \in [3(\alpha - 2)/4, \alpha - 1]$. We then have

$$0 \leq \alpha - 1 - \lambda \leq \varepsilon\eta(1 - \eta), \quad \text{for all } \lambda \in [\bar{\lambda}, \alpha - 1].$$

It then follows that for all $\lambda \in [\bar{\lambda}, \alpha - 1]$ and almost all $t \geq t_0$:

$$(2.9) \quad \partial_t E_t^\lambda \leq -\lambda(1 - \eta)\varepsilon\beta_t \langle Q_t, x_t - x^* \rangle - \frac{t}{2}(\alpha - 1 - \lambda)\|\dot{x}_t\|^2 - \frac{1}{2}\beta_t^2 t(1 - \eta)\|Q_t\|^2.$$

Integrating this inequality and using that the energy function E^λ is nonnegative, we get (i).

To show (ii), we start by expressing $E_t^\lambda - E_t^{\alpha-1}$ which reads

$$E_t^\lambda - E_t^{\alpha-1} = -(1 - \eta)(\alpha - 1 - \lambda)\beta_t t \langle Q_t, x_t - x^* \rangle + \frac{\lambda}{2}(\alpha - 1 - \lambda)\|x_t - x^*\|^2 + \frac{1}{2}\|v_t^\lambda\|^2 - \frac{1}{2}\|v_t^{\alpha-1}\|^2.$$

Using the three point identity, we get

$$\begin{aligned} \frac{1}{2}\|v_t^\lambda\|^2 - \frac{1}{2}\|v_t^{\alpha-1}\|^2 &= -\frac{1}{2}\|v_t^\lambda - v_t^{\alpha-1}\|^2 + \langle v_t^\lambda, v_t^\lambda - v_t^{\alpha-1} \rangle \\ &= -\frac{(\alpha - 1 - \lambda)(\alpha - 1 + \lambda)}{2}\|x_t - x^*\|^2 - (\alpha - 1 - \lambda)t \langle x_t - x^*, \dot{x}_t + \eta\beta_t Q_t \rangle. \end{aligned}$$

Plugging this above we arrive at

$$(2.10) \quad \begin{aligned} E_t^\lambda - E_t^{\alpha-1} &= -(1 - \eta)(\alpha - 1 - \lambda)\beta_t t \langle Q_t, x_t - x^* \rangle \\ &\quad - \frac{(\alpha - 1)(\alpha - 1 - \lambda)}{2}\|x_t - x^*\|^2 - (\alpha - 1 - \lambda)t \langle x_t - x^*, \dot{x}_t + \eta\beta_t Q_t \rangle \\ &= -(\alpha - 1 - \lambda)\beta_t t \langle Q_t, x_t - x^* \rangle \\ &\quad - \frac{(\alpha - 1)(\alpha - 1 - \lambda)}{2}\|x_t - x^*\|^2 - (\alpha - 1 - \lambda)t \langle x_t - x^*, \dot{x}_t \rangle. \end{aligned}$$

Therefore, we get for almost all $t \geq t_0$

$$(2.11) \quad E_t^\lambda - E_t^{\alpha-1} = -(\alpha - 1 - \lambda) \left[(\alpha - 1)q(t) + t\dot{q}(t) - (\alpha - 1)I(t) \right],$$

where we have denoted:

$$q(t) := \frac{1}{2}\|x_t - x^*\|^2 + I(t), \quad \text{and} \quad I(t) := \int_{t_0}^t \beta_s \langle Q_s, x_s - x^* \rangle ds.$$

Note that $I(t)$ admits a limit as $t \rightarrow +\infty$ by (i). Moreover, from (2.9) we also get that for each $\lambda \in [\bar{\lambda}, \alpha - 1]$ the limit $\lim_{t \rightarrow +\infty} E_t^\lambda$ exists. In particular the limit of $E_t^\lambda - E_t^{\alpha-1}$ exists. Hence, we deduce from (2.11) that $\lim_{t \rightarrow +\infty} q(t) + \frac{t}{\alpha-1}\dot{q}(t)$ exists. This implies, thanks to [9, Lemma A.2] and since $\alpha > 2$, that also $\lim_{t \rightarrow +\infty} q(t)$ exists. In view of the definition of $q(t)$, $I(t)$ admitting a limit, we get that $\lim_{t \rightarrow +\infty} \|x_t - x^*\|$ exists. $x(t)$ is in particular bounded, and one can extract a weakly convergent subsequence. Using (2.8) and that $t \mapsto \beta(t)$ is nondecreasing, it is immediate to see that the weak cluster point of each such weakly converging subsequence is a zero of Q . We now invoke the standard Opial's lemma to get (ii).

Let us now establish the little- o rates in (iii). To do so, we first note that, since the limit of $t \mapsto q(t)$ exists, from (2.11) we get:

$$\lim_{t \rightarrow +\infty} t\dot{q}(t) = \lim_{t \rightarrow +\infty} t \langle \dot{x}_t + \beta_t Q_t, x_t - x^* \rangle \in \mathbb{R}.$$

Therefore, since for almost all $t \geq t_0$

$$\begin{aligned} E_t^{\alpha-1} &= \frac{1}{2}\|(\alpha - 1)(x_t - x^*) + t(\dot{x}_t + \eta\beta_t Q_t)\|^2 \\ &\quad + (1 - \eta)(\alpha - 1)\beta_t t \langle Q_t, x_t - x^* \rangle + \frac{\eta}{2}(1 - \eta)\beta_t^2 t^2 \|Q_t\|^2 \\ &= \frac{1}{2}(\alpha - 1)^2 \|x_t - x^*\|^2 + (\alpha - 1)t \langle x_t - x^*, \dot{x}_t + \beta_t Q_t \rangle + \frac{t^2}{2}\|\dot{x}_t + \eta\beta_t Q_t\|^2 + \frac{\eta}{2}(1 - \eta)\beta_t^2 t^2 \|Q_t\|^2, \end{aligned}$$

we deduce that

$$0 \leq \ell := \lim_{t \rightarrow +\infty} t^2 (\|\dot{x}_t + \eta\beta_t Q_t\|^2 + \eta(1 - \eta)\|\beta_t Q_t\|^2)$$

exists. Suppose that $\ell > 0$. Then, there exists $s \geq t_0$ such that

$$\begin{aligned} & \int_{t_0}^{+\infty} t (\|\dot{x}_t + \eta\beta_t Q_t\|^2 + \eta(1-\eta)\|\beta_t Q_t\|^2) dt \\ & \geq \int_s^{+\infty} t^2 (\|\dot{x}_t + \eta\beta_t Q_t\|^2 + \eta(1-\eta)\|\beta_t Q_t\|^2) t^{-1} dt \geq \int_s^{+\infty} \frac{\ell}{2t} dt = +\infty, \end{aligned}$$

leading to a contradiction with the last two integral bounds in (i). This shows that $\ell = 0$ and combining this with Jensen's inequality proves the last two little- o rates in (iii). For the first rate, we have

$$\begin{aligned} & \lim_{t \rightarrow +\infty} t\beta_t \langle Q_t, x_t - x^* \rangle \\ & = \lim_{t \rightarrow +\infty} (t \langle \dot{x}_t + \beta_t Q_t, x_t - x^* \rangle - t \langle \dot{x}_t, x_t - x^* \rangle) = \lim_{t \rightarrow +\infty} t \langle \dot{x}_t + \beta_t Q_t, x_t - x^* \rangle \in \mathbb{R}, \end{aligned}$$

where we used that $t\|\dot{x}_t\| \rightarrow 0$ and x_t is bounded in the last inequality. Arguing with a contradiction argument as above, using the first integral bound in (i), yields the first little- o rate. \square

Remark 2.3 (Connection to Tikhonov Regularization). *It is instructive to inspect the edge case in Theorem 2.2 with $\varepsilon \equiv 0$, i.e., when*

$$(2.12) \quad \dot{\beta}(t) + \frac{\beta(t)}{t}(2-\alpha) = 0, \quad \text{i.e.,} \quad \beta(t) = \beta_0 t^{\alpha-2}, \quad \text{for } \beta_0 > 0.$$

One can check that in this case (as well as when $\alpha = 1$ and any $\varepsilon \geq 0$) we have $b(t) = \eta\dot{\beta}(t) + \alpha\eta\frac{\beta(t)}{t} = \dot{\beta}(t) + \frac{\beta(t)}{t}$ and thus (2.1) reads as

$$(2.13) \quad \ddot{x}(t) + \frac{\alpha}{t}\dot{x}(t) + \beta(t)\frac{d}{dt}Q(x(t)) + \left(\dot{\beta}(t) + \frac{1}{t}\beta(t)\right)Q(x(t)) = 0, \quad t \geq t_0.$$

Multiplying (2.13) by t we shall write:

$$(2.14) \quad \begin{aligned} 0 & = t\ddot{x}(t) + \dot{x}(t) + t\beta(t)\frac{d}{dt}Q(x(t)) + \left(t\dot{\beta}(t) + \beta(t)\right)Q(x(t)) + (\alpha-1)\dot{x}(t) = 0 \\ & = \frac{d}{dt} \left[t(\dot{x}(t) + \beta(t)Q(x(t))) \right] + (\alpha-1)\dot{x}(t). \end{aligned}$$

Therefore, a solution to (2.13) also solves

$$(2.15) \quad \dot{x}(t) + \beta_0 t^{\alpha-2}Q(x(t)) + \frac{\alpha-1}{t}(x(t) - v) = 0, \quad t \geq t_0,$$

where $v := t_0(\dot{x}(t_0) + \beta(t_0)Q(x(t_0))) + (\alpha-1)x(t_0)$. This reveals two interesting properties. First, if $\alpha = 1$ and $\varepsilon \geq 0$, the system is fundamentally equivalent to a first-order system with perturbation and time-scaling (see [3]). Secondly, if $\alpha > 1$ and $\varepsilon = 0$, e.g., $\alpha = 2$ (which yields $\beta(t) = \beta_0$), we are in the setting of Tikhonov regularization and continuous time version of Halpern method: The system (2.15) generates trajectories that converge strongly to the projection of v onto the set of zeros of Q , see, e.g., [26, 16].

3. CONVERGENCE ANALYSIS OF ALGORITHM 1

In this section, we present the convergence analysis of Algorithm 1. Let us first show that the latter can indeed be understood as a discretization of (2.1). Let $(x^k)_{k \in \mathbb{N}}$ be the sequence generated by Algorithm 1, $Q := I - T$ and, for all $k \geq 0$, set

$$(3.1) \quad z^{k+1} := T(x^k), \quad \text{and} \quad Q^{k+1} := Q(x^k) = (I - T)(x^k) = x^k - z^{k+1}.$$

By definition of Q and Algorithm 1, we have for all $k \geq 1$:

$$\begin{aligned}
(3.2) \quad z^{k+2} + Q^{k+2} &= x^{k+1} \\
&= \left(1 - \frac{\alpha_\eta}{k + \sigma}\right) x^k + \frac{\alpha_\eta}{k + \sigma} T(x^k) + \left(1 - \frac{\alpha}{k + \sigma}\right) (T(x^k) - T(x^{k-1})) \\
&= \left(1 - \frac{\alpha_\eta}{k + \sigma}\right) (Q^{k+1} + z^{k+1}) + \frac{\alpha_\eta}{k + \sigma} z^{k+1} + \left(1 - \frac{\alpha}{k + \sigma}\right) (z^{k+1} - z^k) \\
&= z^{k+1} + \left(1 - \frac{\alpha}{k + \sigma}\right) (z^{k+1} - z^k) + \left(1 - \frac{\alpha_\eta}{k + \sigma}\right) Q^{k+1}.
\end{aligned}$$

Now, bringing every term to the left hand-side, we get for all $k \geq 1$:

$$(3.3) \quad z^{k+2} - 2z^{k+1} + z^k + \frac{\alpha}{k + \sigma} (z^{k+1} - z^k) + (Q^{k+2} - Q^{k+1}) + \frac{\alpha_\eta}{k + \sigma} Q^{k+1} = 0.$$

Denoting by $t_k := (k - 1) + \sigma$, we get for all $k \geq 1$:

$$(3.4) \quad (z^{k+2} - 2z^{k+1} + z^k) + \frac{\alpha}{t_{k+1}} (z^{k+1} - z^k) + (Q^{k+2} - Q^{k+1}) + \frac{\alpha_\eta}{t_{k+1}} Q^{k+1} = 0,$$

which, when $z^{k+1} \simeq x^k$ (asymptotically the case), can be understood as an implicit finite-difference discretization of (2.1) with $\beta \equiv 1$, step-size 1, and $Q := I - T$, which is Lipschitz continuous and monotone. The main difference with the implicit discretization of the Fast-OGDA system proposed in [14] is that the latter does not lead to an easily implementable numerical method as it involves computing the resolvent of Q which is hardly available. Note that an explicit scheme is also proposed in [14] which is very different from ours.

Remark 3.1 (Connection to [15]). *As we mentioned in the introduction, Algorithm 1 is a generalization of the scheme (2.8) in [15]. The latter can be recovered by setting $\sigma = \alpha$ and $\eta = \frac{1}{2}$, which gives for all $k \geq 1$:*

$$(3.5) \quad x^{k+1} = \left(1 - \frac{\alpha}{2(k + \alpha)}\right) x^k + \frac{\alpha}{2(k + \alpha)} T(x^k) + \left(1 - \frac{\alpha}{k + \alpha}\right) (T(x^k) - T(x^{k-1})).$$

This is also a special case of [15, Algorithm 2.1], which can be recovered by introducing a step-size $s \in (0, 1]$ and considering $T_s := (1 - s)I + sT$, which is now s -averaged and has the same fixed-points as T . Indeed, substituting T by T_s in (3.5) we get for all $k \geq 1$:

$$(3.6) \quad x^{k+1} = \left(1 - \frac{s\alpha}{2(k + \alpha)}\right) x^k + \frac{(1 - s)k}{k + \alpha} (x^k - x^{k-1}) + \frac{s\alpha}{2(k + \alpha)} T(x^k) + \frac{sk}{k + \alpha} (T(x^k) - T(x^{k-1})).$$

Conversely, if T is θ -averaged, we shall consider (3.6) with $s \in (0, 1/\theta]$, which is equivalent to applying (3.5) with T replaced by $sT + (1 - s)I$. Since there is no loss of generality, in the following, we provide the analysis for general nonexpansive operators.

Remark 3.2 (Connection to Halpern-type Methods). *Since β is constant here, $\alpha = 2$ corresponds to the edge case in the growth condition (2.6), i.e., $\varepsilon = 0$, and we have a similar behaviour as that described in Remark 3.1: The method reduces to the Optimal Halpern Method (OHM). In fact, for $\alpha = 2$, $\sigma = \alpha$ and any $\eta \in (0, 1)$, Algorithm 1 reads as:*

$$(3.7) \quad x^{k+1} := \left(1 - \frac{1}{k + 2}\right) x^k + \frac{1}{k + 2} T(x^k) + \frac{k}{k + 2} (T(x^k) - T(x^{k-1})),$$

which coincides with the momentum form of OHM, see [46]. It is interesting to observe that in the edge case $\alpha = 2$ we lose the little- o rates, and the convergence of iterates is in fact in the strong topology, but is established with completely different techniques.

To analyze Algorithm 1, we denote finite differences with the Δ symbol, i.e., for a sequence $(a^k)_{k \in \mathbb{N}}$, we define $\Delta a_k := a^{k+1} - a^k$ for all $k \geq 0$. Second order differences are denoted by Δ^2 , i.e., $\Delta^2 a_k = \Delta(\Delta a)_k := \Delta a_{k+1} - \Delta a_k$. These notations will prove handy in our analysis since we follow closely the

continuous time Lyapunov analysis. In this sense, it is convenient to recall some discrete time calculus rule listed in Appendix A.

3.1. Lyapunov Energy. Inspired by the continuous time approach, we introduce an energy function that matches almost exactly² (2.3). Specifically, we consider for all $k \geq 0$ the sequences $z^{k+1} := T(x^k)$ and $Q^{k+1} := Q(x^k)$, where $Q := I - T$, and for all $k \geq 2$, $\lambda \in (0, \alpha - 1]$, and any $z^* \in \text{zer } Q = \text{Fix } T$, we consider the energy function:

$$(3.8) \quad \begin{aligned} E_k^\lambda &:= \delta_k \langle Q^k, z^k - z^* \rangle + \frac{\delta_k}{2} \|Q^k\|^2 + \frac{\xi_k}{2} \|Q^k\|^2 + \frac{c}{2} \|z^{k-1} - z^*\|^2 + \frac{1}{2} \|v^k\|^2, \\ v^k &:= \lambda(z^{k-1} - z^*) + t_{k-1} (\Delta z_{k-1} + \eta Q^k), \end{aligned}$$

where $t_k := (k-1) + \sigma$, $\delta_k := (1-\eta)\lambda t_{k-1}$, $\xi_k := (1-\eta)\eta t_{k-1}^2$, and $c := \lambda(\alpha - 1 - \lambda)$. Note that the only substantial difference with (2.3) is that the first inner product $\langle Q^k, z^k - z^* \rangle$ is no longer nonnegative, since $Q^k = Q(x^{k-1})$, i.e. Q is evaluated at x^{k-1} rather than at z^k . The second term allows us to make E_k^λ nonnegative. Indeed, observing that $Q/2$ is firmly nonexpansive, it follows from [10, Proposition 4.4] that for every $x, y \in H$:

$$(3.9) \quad \begin{aligned} \langle Q(x) - Q(y), T(x) - T(y) \rangle &= \langle Q(x) - Q(y), x - y \rangle - \|Q(x) - Q(y)\|^2 \\ &\geq -\frac{1}{2} \|Q(x) - Q(y)\|^2. \end{aligned}$$

Using this with $x = x^k$ and $y = z^* \in \text{Fix } T$, hence $Q(z^*) = 0$, we get that $\langle Q^k, z^k - z^* \rangle + \frac{1}{2} \|Q^k\|^2 \geq 0$ for all $k \geq 2$.

3.2. Energy Decrease. We start with the following preliminary result, which represents the discrete time counterpart of Lemma 2.1.

Lemma 3.3. *Let $T: H \rightarrow H$ be a nonexpansive operator with $\text{Fix } T \neq \emptyset$. Let $(x^k)_{k \in \mathbb{N}}$ be a sequence generated by Algorithm 1 for some $\eta \in (0, 1)$, $\sigma > 0$, and $\alpha \geq 2$. Then there exists $\underline{\lambda} < \alpha - 1$ such that for all $\lambda \in (\underline{\lambda}, \alpha - 1]$ and all $k \geq 2$ such that $t_k \geq 2\lambda$ it holds:*

$$(3.10) \quad \begin{aligned} \Delta E_k^\lambda &\leq -\lambda(1-\eta)(\alpha-2) \left(\langle x^{k-1} - T(x^{k-1}), T(x^{k-1}) - z^* \rangle + \frac{1}{2} \|x^{k-1} - T(x^{k-1})\|^2 \right) \\ &\quad \left[-\eta(1-\eta)(\alpha-2)t_k - \frac{1-\eta}{2} (\eta + (1-\eta)(\alpha-1)^2 - \lambda(\alpha-1)) \right] \|x^{k-1} - T(x^{k-1})\|^2 \\ &\quad + (\alpha-1-\lambda) \frac{t_k - (2(1-\eta)\alpha_\eta + (1-\eta + \alpha_\eta))}{2} \|x^{k-1} - T(x^{k-1})\|^2 \\ &\quad - (\alpha-1-\lambda) \frac{t_k - (\alpha+2 + (1-\eta)(\alpha-1))}{2} \|T(x^{k-1}) - T(x^{k-2})\|^2. \end{aligned}$$

Observe that the first term on the right hand side is negative thanks to (3.9). The proof of Lemma 3.3, which is postponed to Appendix B, follows exactly the continuous time approach with $Q := I - T$. We are now ready to establish the discrete counterpart of Theorem 2.2.

Theorem 3.4. *Let $T: H \rightarrow H$ be a nonexpansive operator with $\text{Fix } T \neq \emptyset$. Let $(x^k)_{k \in \mathbb{N}}$ be a sequence generated by Algorithm 1 for some $\eta \in (0, 1)$, $\sigma > 0$, and $\alpha \geq 2$. Then:*

(i) *We have $\|x^k - T(x^k)\| = O(k^{-1})$ as $k \rightarrow +\infty$.*

Suppose now that $\alpha > 2$. Then:

(ii) *We have the summability properties:*

$$\begin{aligned} \sum_{k \in \mathbb{N}} \left(\langle x^k - T(x^k), T(x^k) - z^* \rangle + \frac{1}{2} \|x^k - T(x^k)\|^2 \right) &< +\infty, \\ \sum_{k \in \mathbb{N}} k \|x^k - T(x^k)\|^2 &< +\infty, \quad \text{and} \quad \sum_{k \in \mathbb{N}} k \|x^k - x^{k-1}\|^2 < +\infty. \end{aligned}$$

²Yet another difference with prior work and in particular with [14, 5] is that the term in v^k is scaled differently.

- (iii) The sequence $(x^k)_{k \in \mathbb{N}}$ converges weakly to a fixed point of T as $k \rightarrow +\infty$.
(iv) We have the following fast rates of convergence:

$$\|x^k - x^{k-1}\| = o(k^{-1}) \quad \text{and} \quad \|x^k - T(x^k)\| = o(k^{-1}) \quad \text{as } k \rightarrow +\infty.$$

Proof. As in the proof of Lemma 3.3, it is convenient to introduce the additional $(z^k)_{k \in \mathbb{N}}$. We recall that the latter is defined by $z^{k+1} := T(x^k)$ and thus satisfies (3.4) with $Q^{k+1} := Q(x^k) := (I - T)(x^k) = x^k - z^{k+1}$ for all $k \geq 0$.

- (i) When $\alpha = 2$, we set $\lambda = 1$, which also yields

$$\eta + (1 - \eta)(\alpha - 1)^2 - \lambda(\alpha - 1) = 0,$$

hence $\Delta E_k^1 \leq 0$ for all $k \geq 2$. For $\alpha > 2$, we set $\lambda = \alpha - 1$, and (3.10) becomes

$$(3.11) \quad \Delta E_k^{\alpha-1} \leq -(1 - \eta)(\alpha - 1)(\alpha - 2) \left(\langle Q^k, z^k - z^* \rangle + \frac{1}{2} \|Q^k\|^2 \right) \\ + \left[-\eta(1 - \eta)(\alpha - 2)t_k + \frac{\eta(1 - \eta)}{2} ((\alpha - 1)^2 - 1) \right] \|Q^k\|^2.$$

Thus, there exists $\bar{k} \geq 2$ large enough, i.e. $t_{\bar{k}} \geq \frac{(\alpha-1)^2-1}{2(\alpha-2)}$, such that for all $k \geq \bar{k}$ we have $\Delta E_k^{\alpha-1} \leq 0$. Therefore, for all $k \geq \bar{k}$, we have $\frac{\xi_{k+1}}{2} \|Q^{k+1}\|^2 \leq E_{k+1}^1 \leq E_{\bar{k}}^1$ whence we get the claim recalling the expressions of ξ_k and t_k .

Suppose from now on that $\alpha > 2$.

- (ii) Setting again $\lambda = \alpha - 1$ and embarking from (3.11), we have after summing on both sides that

$$(1 - \eta)(\alpha - 1)(\alpha - 2) \sum_{i=\bar{k}}^k \left(\langle Q^i, z^i - z^* \rangle + \frac{1}{2} \|Q^i\|^2 \right) + \eta(1 - \eta)(\alpha - 2) \sum_{i=\bar{k}}^k (t_i - t_{\bar{k}}) \|Q^i\|^2 \\ \leq E_{\bar{k}}^{\alpha-1} - E_{k+1}^{\alpha-1} \leq E_{\bar{k}}^{\alpha-1} < +\infty.$$

Passing to the limit as $k \rightarrow +\infty$ we get the first two summability claims. For the last one, we set $\lambda = \alpha - 1 - \varepsilon$, for $\varepsilon > 0$ to be chosen small enough. Then, discarding the first negative term in (3.10), we have

$$(3.12) \quad \Delta E_k^{\alpha-1-\varepsilon} \leq \left[-\left(\eta(1 - \eta)(\alpha - 2) - \frac{\varepsilon}{2} \right) t_k + \frac{\eta(1 - \eta)}{2} ((\alpha - 1)^2 - 1) \right] \|Q^k\|^2 \\ - \varepsilon \frac{t_k - (\alpha + 2 + (1 - \eta)(\alpha - 1))}{2} \|\Delta z_{k-1}\|^2.$$

Therefore, for $0 < \varepsilon < \eta(1 - \eta)(\alpha - 2)$, there exists $\tilde{k} \geq 2$ large enough such that for all $k \geq \tilde{k}$,

$$\Delta E_k^{\alpha-1-\varepsilon} \leq -\frac{\varepsilon}{2} (t_k - t_{\tilde{k}}) \|\Delta z_{k-1}\|^2,$$

where we have discarded the first negative term in (3.12). Summing again we get $\sum_{k \in \mathbb{N}} k \|\Delta z_{k-1}\|^2 < +\infty$. Observe now that since $Q^{k+1} = x^k - z^{k+1}$ by definition, we have

$$\|\Delta x_{k-1}\|^2 = \|x^k - x^{k-1}\|^2 = \|\Delta z_k + \Delta Q_k\|^2 \leq 2\|\Delta z_k\|^2 + 2\|Q^{k+1}\|^2 + \|Q^k\|^2.$$

Multiplying by k , summing and using the summability claims proved above, we conclude.

- (iii) We argue as in the continuous time setting and set $\lambda = \alpha - 1 - \varepsilon$, for $\varepsilon > 0$ small enough as before. We will emphasize the dependency of $(v^k)_{k \in \mathbb{N}}$ on λ by writing explicitly $(v_\lambda^k)_{k \in \mathbb{N}}$. Let $k \geq 2$. We have

$$E_k^\lambda - E_k^{\alpha-1} = -(1 - \eta)(\alpha - 1 - \lambda)t_{k-1} \left(\langle Q^k, z^k - z^* \rangle + \frac{1}{2} \|Q^k\|^2 \right) \\ + \frac{\lambda}{2} (\alpha - 1 - \lambda) \|z^{k-1} - z^*\|^2 + \underbrace{\frac{1}{2} \|v_\lambda^k\|^2 - \frac{1}{2} \|v_{\alpha-1}^k\|^2}_{(I)}.$$

Let us now compute (I) separately. By the three point identity, we get

$$(I) := \frac{1}{2}\|v_\lambda^k\|^2 - \frac{1}{2}\|v_{\alpha-1}^k\|^2 = \frac{\lambda^2 - (\alpha-1)^2}{2}\|z^{k-1} - z^*\|^2 - t_{k-1}(\alpha-1-\lambda)\langle \Delta z_{k-1} + \eta Q^k, z^{k-1} - z^* \rangle.$$

Therefore,

$$\begin{aligned} E_k^\lambda - E_k^{\alpha-1} &= -(\alpha-1-\lambda)\left[t_{k-1}\langle Q^k, z^k - z^* - \eta\Delta z_{k-1} \rangle + \frac{1-\eta}{2}t_{k-1}\|Q^k\|^2\right. \\ &\quad \left. + \frac{(\alpha-1)}{2}\|z^{k-1} - z^*\|^2 + t_{k-1}\langle \Delta z_{k-1}, z^{k-1} - z^* \rangle\right]. \end{aligned}$$

Therefore, this can be written as

$$(3.13) \quad \begin{aligned} E_k^\lambda - E_k^{\alpha-1} &= -(\alpha-1-\lambda)\left[(\alpha-1)q_k + t_{k-1}\Delta q_k - (\alpha-1)I_k\right. \\ &\quad \left. - \eta t_{k-1}\langle Q^k, \Delta z_{k-1} \rangle - \frac{t_{k-1}}{2}(\|\Delta z_{k-1}\|^2 + \eta\|Q^k\|^2)\right], \end{aligned}$$

where we have denoted

$$q_k := \frac{1}{2}\|z^{k-1} - z^*\|^2 + I_k, \quad \text{and} \quad I_k := \sum_{j=1}^{k-1} \left(\langle Q^j, z^j - z^* \rangle + \frac{1}{2}\|Q^j\|^2 \right).$$

Recall that I_k is nonnegative thanks to (3.9), hence so is q_k . As argued in the proof of (i), $\Delta E_k^{\alpha-1} \leq 0$ for all k large enough and thus $\frac{c}{2}\|z_k - z^*\|^2 \leq E_k^{\alpha-1} < +\infty$, i.e. $(z^k)_{k \in \mathbb{N}}$ is a bounded sequence. It then follows from the last two summability properties of (ii) that the terms terms in the second line of (3.13) converge to 0. (ii) also gives that I_k admit a limit. Moreover, E_k^λ and E_k^α are nonnegative and decreasing for k large enough as argued above. Therefore, we deduce that

$$(3.14) \quad \lim_{k \rightarrow +\infty} q_k + \frac{t_{k-1}}{\alpha-1}\Delta q_k = \lim_{k \rightarrow +\infty} (E_k^\lambda - E_k^{\alpha-1}) \in \mathbb{R}.$$

We are then in position to apply [4, Lemma A.2] to (3.14) coupled with the fact that $\frac{t_{k-1}}{\alpha-1} = \frac{k-2+\sigma}{\alpha-1}$ (recall that $\alpha-1 > 1$ by assumption), to infer that $(q_k)_{k \in \mathbb{N}}$ admits a limit. Using again that I_k has a limit by (ii), we get that $(\|z^k - z^*\|)_{k \in \mathbb{N}}$ converges. Since $\|Q^k\| = \|x^{k-1} - T(x^{k-1})\| = \|x^{k-1} - z^k\| \rightarrow 0$ by (i), we have $\lim_{k \rightarrow +\infty} \|x^{k-1} - z^*\|^2 = \lim_{k \rightarrow +\infty} \|z^k - z^*\|^2$ and thus the limit of $\|x^k - z^*\|$ exists. The weak convergence claim of $(x_k)_{k \in \mathbb{N}}$ then follows from the standard Opial's lemma.

(iv) Let us first note that since the limit of $(q^k)_{k \in \mathbb{N}}$ exists (see the proof of (iii)), (3.14) yields

$$(3.15) \quad \begin{aligned} \mathbb{R} \ni \lim_{k \rightarrow +\infty} t_{k-1}\Delta q_k &= \lim_{k \rightarrow +\infty} t_{k-1} \left[\langle Q^k, z^k - z^* \rangle + \frac{1}{2}\|Q^k\|^2 \right. \\ &\quad \left. + \langle \Delta z_{k-1}, z^k - z^* \rangle - \frac{1}{2}\|\Delta z_{k-1}\|^2 \right]. \end{aligned}$$

Now, from (3.8), we have

$$\begin{aligned} E_k^{\alpha-1} &= \frac{1}{2}\|(\alpha-1)(z^{k-1} - z^*) + t_{k-1}(\Delta z_{k-1} + \eta Q^k)\|^2 \\ &\quad + (1-\eta)(\alpha-1)t_{k-1} \left(\langle Q^k, z^k - z^* \rangle + \frac{1}{2}\|Q^k\|^2 \right) + \frac{(1-\eta)\eta t_{k-1}^2}{2}\|Q^k\|^2 \\ &= \frac{(\alpha-1)^2}{2}\|z^{k-1} - z^*\|^2 + \frac{t_{k-1}^2}{2}\|\Delta z_{k-1} + \eta Q^k\|^2 + \frac{1}{2}(1-\eta)\eta t_{k-1}^2\|Q^k\|^2 \\ &\quad - \frac{1}{2}(\alpha-1)\eta t_{k-1}\|Q^k\|^2 - (\alpha-1)\eta t_{k-1}\langle Q^k, \Delta z_{k-1} \rangle \\ &\quad + (\alpha-1)t_{k-1} \left[\langle Q^k, z^k - z^* \rangle + \frac{1}{2}\|Q^k\|^2 + \langle \Delta z_{k-1}, z^k - z^* \rangle - \frac{1}{2}\|\Delta z_{k-1}\|^2 \right]. \end{aligned}$$

Combining (3.15) (hence the last line has a limit), (ii) (hence the second line converges to 0), and that $\lim_{k \rightarrow \infty} \|z^{k-1} - z^*\|^2$ and $\lim_{k \rightarrow \infty} E_k^{\alpha-1}$ exist as proved above, we deduce that

$$\lim_{t \rightarrow +\infty} t_{k-1}^2 \left(\frac{1}{2} \|\Delta z_{k-1} + \eta Q^k\|^2 + \frac{1}{2} (1 - \eta) \eta \|Q^k\|^2 \right) \in \mathbb{R}.$$

Observe now that the summability claims of (ii) already give us that

$$(3.16) \quad \liminf_{k \rightarrow +\infty} k^2 \|\Delta z_k\|^2 = 0, \quad \text{and} \quad \liminf_{k \rightarrow +\infty} k^2 \|Q^k\|^2 = 0.$$

Thus these are actually limits. To transfer the first rate from z^k to x^k , we use again the relation $x^k = Q^{k+1} + z^{k+1}$. This concludes the proof. \square

4. APPLICATION TO SPLITTING METHODS WITH PRECONDITIONERS

The goal here is apply our results to operator splitting methods for solving monotone inclusion problems, i.e. finding a zero of one or sum of maximally monotone operators. We follow the algorithmic framework introduced in [17]. For a maximal monotone operator $A: H \rightarrow 2^H$, we introduce a *preconditioner* M , i.e., a self-adjoint positive semidefinite bounded linear operator $M: H \rightarrow H$, and consider the resolvent:

$$(4.1) \quad J_{M^{-1}A} := (I + M^{-1}A)^{-1} = (M + A)^{-1}M.$$

It is worth noting that without further hypothesis, $J_{M^{-1}A}$ is not necessarily single-valued nor defined everywhere on H . On the other hand, this representation is helpful in practice, and evaluating $J_{M^{-1}A}$ can be facilitated by the right choice of M . As first remarked by He and Yuan in [32], this is in fact the case for primal-dual-type methods. Later, by allowing the preconditioner M to be *degenerate*, i.e., with $\ker M \neq \emptyset$, Bredies and Sun in [19, 20], and later [17] recognized that the class of methods representable as iterations with respect to degenerate preconditioners can be greatly broadened to include methods such as the Douglas–Rachford Splitting (DRS) [34] and its extensions [18].

4.1. Preliminaries on Degenerate Proximal Point. The notion of monotonicity and cocoercivity can be readily extended to a *preconditioned* setting, even with a degenerate preconditioner $M: H \rightarrow H$. In the degenerate case, M induces a Hilbertian semi-inner product $\langle u, v \rangle_M := \langle u, Mv \rangle$ and associated seminorm $\|u\|_M := \sqrt{\langle u, u \rangle_M}$ for all $u, v \in H$. For a maximal monotone operator A and a preconditioner M we shall consider the operator $M^{-1}A$ as a composition of the multi-valued operators M^{-1} (pre-image) and A , defined as

$$(4.2) \quad M^{-1}A(u) := \bigcup_{a \in A(u)} M^{-1}(a), \quad \text{for all } u \in H.$$

It is easy to check that $M^{-1}A$ is monotone with respect to the semi-inner product induced by M . We then say that $M^{-1}A$ is *M-monotone*. Likewise, its resolvent $J_{M^{-1}A}$ is *M-cocoercive* in the sense that

$$(4.3) \quad \langle v - v', u - u' \rangle_M \geq \|v - v'\|_M^2, \quad \text{for all } (u, v), (u', v') \in \text{gra } J_{M^{-1}A}.$$

Due to the degeneracy of the preconditioner, the operator $J_{M^{-1}A}$ is not necessarily single-valued and could even have empty domain. To avoid such pathological cases, we suppose $\text{dom } J_{M^{-1}A} = H$. A further hypothesis we shall require is that the operator

$$(4.4) \quad (M + A)^{-1} \text{ is Lipschitz over } \text{Im } M,$$

which makes $J_{M^{-1}A}$ a well-defined Lipschitz operator on H , since indeed $J_{M^{-1}A} = (M + A)^{-1}M$.

An appealing property of preconditioned resolvents with positive semidefinite preconditioners is that evaluating $J_{M^{-1}A}$ automatically produces a quotient space operation. Suppose that M has closed range and consider a factorization of M as $M = CC^*$, with $C: D \rightarrow H$ one-to-one from the Hilbert space D to H . We say that CC^* is an *onto decomposition* of M . In this case, one can prove that C^* is onto [17, Proposition 2.3] and the operator $C^* \triangleright A := (C^* A^{-1} C)^{-1}$ is maximal monotone [17, Theorem 2.13]. In fact, if M has closed range, the latter is an equivalence in the spirit of Minty's Theorem, see [25,

Theorem 3.5.4]. The resolvents of $M^{-1}A$ and of $C^* \triangleright A$ are tightly connected: If M is a preconditioner with closed range and $M = CC^*$ is onto, then

$$(4.5) \quad C^* J_{M^{-1}A}(u) = J_{C^* \triangleright A}(C^* u), \quad \text{for all } u \in H.$$

This is particularly useful as it reveals that the evaluation of $J_{M^{-1}A}$ is somewhat equivalent to evaluating $J_{C^* \triangleright A}$, which is a resolvent of a maximal monotone operator; see [25, Theorem 3.3.3] for a proof. While the preconditioner M can be degenerate, if M has closed range, in view of (4.5), fixed-point iterations for $J_{M^{-1}A}$ are in fact closely related to standard proximal point algorithms associated to maximal monotone operators. However, the convergence proof of the so-called shadow-sequences in infinite-dimensional spaces does not follow directly from the classical analysis. For clarity, we present a prototypical example:

Example 4.1 (Douglas–Rachford [20, 17]). *Let $A_1, A_2: H \rightarrow 2^H$ be two maximal monotone operators on H and assume that $\text{zer}(A_1 + A_2) \neq \emptyset$. Following [17], the DRS method for finding a zero of $A_1 + A_2$ can be understood as iterating $J_{M^{-1}A}$ with the operators:*

$$(4.6) \quad A := \begin{bmatrix} A_1 & I & -I \\ -I & A_2 & I \\ I & -I & 0 \end{bmatrix}, \quad \text{and} \quad M := \begin{bmatrix} I & -I & I \\ -I & I & -I \\ I & -I & I \end{bmatrix} = \begin{bmatrix} I \\ -I \\ I \end{bmatrix} \underbrace{\begin{bmatrix} I & -I & I \end{bmatrix}}_{:=C^*}.$$

Indeed, since $M + A$ is block lower-triangular, evaluating $J_{M^{-1}A} = (M + A)^{-1}M$ only requires accessing the resolvents of A_1 and A_2 , we refer to [17] for the detailed computation. If $u^k := (x_1^k, x_2^k, v^k) \in H^3$ is such that $u^{k+1} := J_{M^{-1}A}(u^k)$ the DRS method can be obtained via the substitution $w^k := x_1^k - x_2^k + v^k$, which yields for all $k \in \mathbb{N}$:

$$(4.7) \quad \begin{cases} x_1^{k+1} = J_{A_1}(w^k), \\ x_2^{k+1} = J_{A_2}(2x_1^{k+1} - w^k), \\ w^{k+1} = w^k + x_2^{k+1} - x_1^{k+1}. \end{cases}$$

This corresponds to taking $w^k = C^* u^k$, which defines an iteration with respect to the maximal monotone operator $C^* \triangleright A$; see Section 4.1. It follows that $(w^k)_{k \in \mathbb{N}}$ weakly converges to some $w^* \in \text{zer } C^* \triangleright A$, and hence $x_1^* = J_{A_1}(w^*) \in \text{zer}(A_1 + A_2)$. The asymptotic behaviour of the so-called solution estimates $(x_1^{k+1})_{k \in \mathbb{N}}$ and $(x_2^{k+1})_{k \in \mathbb{N}}$ (a.k.a. shadow sequences), in infinite dimensional Hilbert spaces, which is related to the asymptotic behaviour of $(J_{M^{-1}A}(u^k))_{k \in \mathbb{N}}$, remained an open problem for almost 30 years until Svaiter proposed a convergence proof in [44]. The analysis of iterations associated to (4.1) with a degenerate preconditioner allows to recover this result from general principles applicable to a broad range of algorithms; see [17].

4.2. A Degenerate Opial Lemma. The following lemma captures the argument that allows us to prove convergence of several method with degenerate preconditioners. The mechanism underlying this result can be traced back to [17].

Lemma 4.2 (Opial in the Degenerate Setting). *Let A be a maximal monotone operator with $\text{zer } A \neq \emptyset$, and M a self-adjoint positive semidefinite preconditioner such that $(M + A)^{-1}$ is Lipschitz over $\text{Im } M$ and $J_{M^{-1}A}$ has full domain. Let $(u^k)_{k \in \mathbb{N}}$ be a sequence, and suppose that:*

- (i) *For all $u^* \in \text{Fix } J_{M^{-1}A}$ the limit $\lim_{k \rightarrow +\infty} \|u^k - u^*\|_M^2$ exists;*
- (ii) *It holds $\lim_{k \rightarrow +\infty} \|u^k - J_{M^{-1}A}(u^k)\|_M = 0$.*

Then $J_{M^{-1}A}(u^k) \rightharpoonup u^ \in \text{zer } A$ as $k \rightarrow +\infty$. If, additionally, M has closed range and $M = CC^*$ is onto, then for $w^k := C^* u^k$:*

$$w^k \rightharpoonup w^* \in \text{zer } C^* \triangleright A, \quad \text{and} \quad (A + M)^{-1}C(w^k) = J_{M^{-1}A}(u^k) \rightharpoonup u^* \in \text{zer } A, \quad \text{as } k \rightarrow +\infty.$$

Proof. For brevity, let us denote by $v^k := J_{M^{-1}A}(u^k)$ for all $k \in \mathbb{N}$. From (i), we deduce that the sequence $(Mu^k)_{k \in \mathbb{N}}$ is bounded. Therefore, since $(M + A)^{-1}$ is Lipschitz and $J_{M^{-1}A} = (M + A)^{-1}M$, which is then single-valued everywhere on H , also $(v^k)_{k \in \mathbb{N}}$ is bounded. Since $M(u^k - v^k) \in A(v^k)$ for all $k \in \mathbb{N}$, $\lim_{k \rightarrow +\infty} \|M(u^k - v^k)\| = 0$ from point (ii), and A is maximal monotone (in particular its graph is weak-to-strong sequentially closed [10, Proposition 20.38]), and in turn every weak cluster point

of $(v^k)_{k \in \mathbb{N}}$ belongs to $\text{zer } A$. Now we show that, in fact, it admits only one weak cluster point and thus $(v^k)_{k \in \mathbb{N}}$ converges weakly to it. Let $(v^{k_j})_{j \in \mathbb{N}}$ and $(v^{k_i})_{i \in \mathbb{N}}$ be two weakly converging subsequences and $v^*, v^{**} \in H$ their respective weak cluster points. Note that, from points (i) and (ii), for all $u^* \in \text{zer } A$, $\lim_{k \rightarrow +\infty} \|v^k - u^*\|_M = \lim_{k \rightarrow +\infty} \|u^k - u^*\|_M$, which we denote by $\ell(u^*)$. Suppose by contradiction that $\|v^* - v^{**}\|_M \neq 0$. Thus,

$$(4.8) \quad \begin{aligned} \ell(v^{**}) &= \lim_{i \rightarrow +\infty} \|v^{k_i} - v^{**}\|_M^2 = \lim_{i \rightarrow +\infty} (\|v^{k_i} - v^*\|_M^2 + 2\langle v^{k_i} - v^*, v^* - v^{**} \rangle_M + \|v^* - v^{**}\|_M^2) \\ &= \|v^* - v^{**}\|_M^2 + \lim_{i \rightarrow +\infty} \|v^{k_i} - v^*\|_M^2 > \ell(v^*), \end{aligned}$$

where we have used that $v^{k_i} \rightharpoonup v^*$. Reversing the role of v^* and v^{**} , we also get $\ell(v^{**}) < \ell(v^*)$, hence the desired contradiction, which yields $Mv^* = Mv^{**}$. Hence, $J_{M^{-1}A}(v^*) = J_{M^{-1}A}(v^{**})$, and, since $v^*, v^{**} \in \text{zer } A = \text{Fix } J_{M^{-1}A}$, we obtain $v^* = v^{**}$. Recalling (4.5), the second statement follows easily from the first. \square

Naturally, a similar results holds in continuous time. We now describe how this lemma can be utilized to prove convergence of several splitting methods in a unified way. We opted to present the adaptation of Theorem 3.4, i.e., the one for algorithmic schemes. The same can be done for Theorem 2.2, i.e., for continuous-time dynamics, but we leave this to the reader.

Theorem 4.3. *Let A be a maximal monotone operator with $\text{zer } A \neq \emptyset$, and M a self-adjoint positive semidefinite preconditioner such that $(M + A)^{-1}$ is Lipschitz over $\text{Im } M$ and $J_{M^{-1}A}$ has full domain. Let $(u^k)_{k \in \mathbb{N}}$ be the sequence³ generated by Algorithm 1 with $T = (1 - s)I + sJ_{M^{-1}A}$, $s \in (0, 2]$, $\eta \in (0, 1)$, $\sigma > 0$, and $\alpha > 2$. Let $u^* \in \text{zer } A$. Then:*

(i) *We have*

$$\sum_{k \in \mathbb{N}} k \|u^k - J_{M^{-1}A}(u^k)\|_M^2 < \infty, \quad \sum_{k \in \mathbb{N}} k \|u^k - u^{k-1}\|_M^2 < +\infty.$$

(ii) *$(J_{M^{-1}A}(u^k))_{k \in \mathbb{N}}$ converges weakly to some element in $\text{zer } A$ as $k \rightarrow +\infty$.*

(iii) *We have the following fast rates of convergence:*

$$\|u^k - u^{k-1}\|_M = o(k^{-1}) \quad \text{and} \quad \|u^k - J_{M^{-1}A}(u^k)\|_M = o(k^{-1}) \quad \text{as } k \rightarrow +\infty.$$

Suppose furthermore that M has closed range and that $M = CC^*$ is onto with $C: D \rightarrow H$. Then, $w^k := C^*u^k$ satisfies (i)-(iii) with $M^{-1}A$ replaced by $C^* \triangleright A$ and seminorms replaced by norms (in D), and $w^k \rightharpoonup C^*u^*$ as $k \rightarrow +\infty$.

Proof. The proof follows exactly the same lines of Theorem 3.4 by replacing every norm by M -seminorms, every inner product with M -semi-inner products and setting $Q = s(I - J_{M^{-1}A})$. In fact, the only property utilized is the nonexpansiveness of T , which in this context translates as the M -nonexpansiveness of $sJ_{M^{-1}A} + (1 - s)I$ for $s \in (0, 2]$ see [17, Remark 2.7]. Specifically, (i) can be proven by following the arguments in the proof Theorem 3.4(ii). This gives that $\lim_{k \rightarrow +\infty} \|u^k - J_{M^{-1}A}(u^k)\|_M = s^{-1} \lim_{k \rightarrow +\infty} \|Q(u^k)\|_M = 0$. Regarding (ii), following the proof of statement Theorem 3.4(iii), we get that $\lim_{k \rightarrow +\infty} \|u^k - u^*\|_M^2$ exists for any $u^* \in \text{zer } A$. We are now in the setting of the degenerate Opial lemma (Lemma 4.2), which yields (ii). Finally, statement (iii) is a consequence of Theorem 3.4(iv). \square

Remark 4.4. *Theorem 4.3, via Lemma 4.2, allows us to extend our results of Section 3 without much effort to degenerate preconditioned proximal point methods, hence obtaining the weak convergence of the sequence*

$$(4.9) \quad J_{M^{-1}A}(u^k) = (M + A)^{-1}(Mu^k) = (M + A)^{-1}(Cw^k), \quad w^k := C^*u^k,$$

to a zero of A . It is important to note, however, that this result does not say anything regarding the asymptotic convergence of the sequence $(u^k)_{k \in \mathbb{N}}$ itself, which requires dedicated analysis. In certain instances such as in [17, Theorem 2.9], one can prove that the stronger statement $\lim_{k \rightarrow 0} \|u^k - J_{M^{-1}A}(u^k)\| = 0$,

³We denote by $(u^k)_{k \in \mathbb{N}}$ the sequence generated by Algorithm 1 to avoid confusion with $x^k = (x_1^k, \dots, x_N^k)$ generated by Algorithm 2 discussed in Section 4.3.

i.e., in the norm and not only the M -seminorm. Unfortunately, the proof of this property does not seem to follow from general principles. For a general setting such as ours, this remains an open problem.

Remark 4.5. If $\alpha = 2$, we are in the setting of the OHM method (cf. Remark 3.2), which produces a sequence that converges strongly. In this case, if M has closed range, there is indeed no need to rely on a different analysis with degenerate preconditioners since the map $(M + A)^{-1}C$ is Lipschitz, and the convergence of $(u^k)_{k \in \mathbb{N}}$ would follow from that of $(w^k)_{k \in \mathbb{N}}$.

4.3. Application to Graph Douglas–Rachford Splitting. In this section, we show the application of Theorem 4.3 to a slight extension of the graph-DRS method [18] discussed in [25].

Consider the monotone inclusion problem:

$$(4.10) \quad \text{Find } x \in H : \quad 0 \in \sum_{i=1}^N A_i(x),$$

where $A_1, \dots, A_N: H \rightarrow 2^H$ are *resolvent-friendly* maximal monotone operators, i.e., such that $J_{\sigma A_i}$ is easy to evaluate for each i and each $\sigma > 0$. To solve (4.10), we can employ the graph-DRS method introduced in [18]. Defining a graph-DRS operator only requires choosing two real matrices:

$$(4.11) \quad Z, \bar{Z} \in \mathbb{R}^{N \times (N-1)}, \quad \text{with} \quad \ker Z^\top = \text{span}\{\mathbf{1}\} \subset \ker \bar{Z}^\top, \quad \mathbf{1} := (1, \dots, 1)^\top.$$

Then, denoting by $L := ZZ^\top$, $\bar{L} := \bar{Z}\bar{Z}^\top$ and $d := \text{diag}(L + \bar{L})$, and for $\tau > 0$, the update rule reads as:

$$(4.12) \quad \begin{cases} x_i^+ := J_{\frac{\tau}{d_i} A_i} \left(-\frac{2}{d_i} \sum_{h=1}^{i-1} (L_{hi} + \bar{L}_{hi}) x_h^+ + \frac{1}{d_i} \sum_{j=1}^{N-1} Z_{ij} w_j \right), \\ w_j^+ := w_j - \sum_{i=1}^N Z_{ij} x_i^+. \end{cases}$$

This method generalizes a number of splitting methods. The baseline DRS solves (4.10) with $N = 2$, and can be obtained from (4.12) choosing $Z^\top = [1, -1]$ and $\bar{Z}^\top = [0, 0]$. The graph-DRS method [18], by choosing Z as an onto decomposition of the graph laplacian of some directed connected graph $\mathcal{G} = (\mathcal{N}, \mathcal{E})$ with nodes $\mathcal{N} := \{1, \dots, N\}$ and edges \mathcal{E} such that $(i, j) \in \mathcal{E}$ if and only if $i < j$, and \bar{Z} as the onto decomposition of the graph laplacian of a (non-necessarily connected) graph on \mathcal{N} . This in turn encompasses, the Ryu’s splitting method [40], the sequential and parallel extensions of DRS [17], the Malitsky–Tam splitting [35] and several other methods.

The method can be understood in the framework of degenerate proximal point methods with respect to the operators

$$(4.13) \quad \mathbf{A} := \begin{bmatrix} \mathbf{A}_L & -\mathbf{Z} \\ \mathbf{Z} & \mathbf{0} \end{bmatrix}, \quad \mathbf{M} := \begin{bmatrix} \mathbf{L} & \mathbf{Z} \\ \mathbf{Z}^* & \mathbf{I} \end{bmatrix} = \begin{bmatrix} \mathbf{Z} \\ \mathbf{I} \end{bmatrix} \underbrace{\begin{bmatrix} \mathbf{Z}^* & \mathbf{I} \end{bmatrix}}_{=: \mathbf{C}^*},$$

where $\mathbf{Z} := Z \otimes I$, $\mathbf{L} := \mathbf{Z}\mathbf{Z}^* = L \otimes I$, \mathbf{I} and $\mathbf{0}$ are, respectively, the identity map and the zero operator on H^{N-1} , $\mathbf{A}_L := \tau \text{diag}(A_1, \dots, A_N) + \mathbf{S} + \mathbf{P}$, $\mathbf{S} := S \otimes I$ and $\mathbf{P} := P \otimes I$. S and P are respectively, skew-symmetric and lower-triangular matrices, defined by

$$(4.14) \quad S_{ij} := \begin{cases} -L_{ij} & \text{if } i < j, \\ L_{ij} & \text{if } i > j, \\ 0 & \text{else,} \end{cases} \quad \text{and} \quad P_{ij} := \begin{cases} \bar{L}_{ij} & \text{if } i = j, \\ 2\bar{L}_{ij} & \text{if } i > j, \\ 0 & \text{else.} \end{cases}$$

One can in fact check that, denoting by $w^+ := (w_1^+, \dots, w_{N-1}^+)$, and $w := (w_1, \dots, w_{N-1})$, we have $w^+ = J_{\mathbf{C}^* \triangleright \mathbf{A}}(w)$ and $x^+ := (x_1^+, \dots, x_N^+)$ are the first N components of $(\mathbf{M} + \mathbf{A})^{-1}(\mathbf{C}w)$, since indeed $x^+ := (\mathbf{L} + \mathbf{A}_L)^{-1}(\mathbf{Z}w)$. Furthermore, $w \in \text{Fix } J_{\mathbf{C}^* \triangleright \mathbf{A}}$ if and only if $x^* = x_1 = \dots = x_N$ and x^* solves (4.10). Fixed-points of $J_{\mathbf{C}^* \triangleright \mathbf{A}}$ are thus related to consensus on the components of x . This can also be quantified via the fixed-point residual $\|w^k - J_{\mathbf{C}^* \triangleright \mathbf{A}}(w^k)\|$. Indeed, it can be shown from [18,

Algorithm 2: Fast Douglas–Rachford Method for N operators

Pick $\eta \in (0, 1)$, $\sigma > 0$, $\alpha \geq 2$, two step-sizes $\tau > 0$ and $s \in (0, 2]$, and Z, \bar{Z} such that (4.11) hold ;

Set $\alpha_\eta := \eta + (1 - \eta)(\alpha - 1)$ and $d = \text{diag}(L + \bar{L})$;

Initialize $w^0, w^1 \in H^{N-1}$;

for $k = 1, 2, \dots$ **do**

for $i = 1, \dots, N$ **do**

$$x_i^{k+1} := J_{\frac{\tau}{d_i} A_i} \left(-\frac{2}{d_i} \sum_{h=1}^{i-1} (L_{hi} + \bar{L}_{hi}) x_h^{k+1} + \frac{1}{d_i} \sum_{j=1}^{N-1} Z_{ij} w_j^k \right)$$

for $j = 1, \dots, N-1$ **do**

$$w_j^{k+1} = w_j^k + \left(1 - \frac{\alpha}{k + \sigma}\right) (w_j^k - w_j^{k-1}) - \frac{s\alpha_\eta}{k + \sigma} \sum_{i=1}^N Z_{ij} x_i^{k+1} + s \left(1 - \frac{\alpha}{k + \sigma}\right) \sum_{i=1}^N Z_{ij} (x_i^{k+1} - x_i^k)$$

Proposition 3.8] or [25, Lemma 5.2.1] that this residual controls a natural dispersion functional between the components of x , referred to as *variance*:

$$(4.15) \quad \text{Var}(x) := \frac{1}{N} \sum_{i=1}^N \|x_i - \bar{x}\|^2 \leq \frac{1}{\lambda_1 N} \|w - J_{\mathcal{C}^* \triangleright \mathbf{A}}(w)\|^2,$$

where $\bar{x} := \frac{1}{N} \sum_{i=1}^N x_i$, λ_1 is the smallest nonzero eigenvalue of L . In plain words, the smaller the residual the closer the components of x are to consensus. In the other hand, it can be shown, see [25, Lemma 5.2.2], that the variance (4.15) provides an upper-bound on the norm of the elements in A_i defined implicitly through the evaluations of their resolvents, i.e., on

$$(4.16) \quad a_i^{k+1} := -\frac{2}{\tau} \sum_{h=1}^{i-1} (L_{hi} + \bar{L}_{hi}) x_h^{k+1} - \frac{d_i}{\tau} x_i^{k+1} + \frac{1}{\tau} \sum_{j=1}^{N-1} Z_{ij} w_j^k \in A(x_i^{k+1}), \quad k \in \mathbb{N},$$

for all $i \in \{1, \dots, N\}$.

Plugging $T := sJ_{\mathcal{M}^{-1}\mathbf{A}} + (1-s)I$ for $s \in (0, 2]$, where \mathbf{A} and \mathcal{M} are defined in (4.13), into Algorithm 1, we get the method described in Algorithm 2 whose convergence proof is an adaptation of Theorem 4.3.

Proposition 4.6. *Let $A_1, \dots, A_N: H \rightarrow 2^H$ maximal monotone operators such that $\text{zer}(A_1, \dots, A_N) \neq \emptyset$ and let $(w_j^k)_{k \in \mathbb{N}}$ and $(x_i^k)_{k \in \mathbb{N}}$ be the iterates generated by Algorithm 2 with $\alpha > 2$, $\sigma > 0$, $\eta \in (0, 1)$ and $s \in (0, 2]$, for $j \in \{1, \dots, N-1\}$ and $i \in \{1, \dots, N\}$. Then:*

(i) $(w_j^k)_{k \in \mathbb{N}}$ converges weakly to some $w_j^* \in H$ for all $j \in \{1, \dots, N-1\}$,

(ii) $(x_i^k)_{k \in \mathbb{N}}$ converges weakly to some $x^* \in \text{zer}(A_1 + \dots + A_N)$ (independent of i) with rate

$$\text{Var}(x^{k+1}) := o(k^{-2}), \quad \text{and} \quad \|a_1^{k+1} + \dots + a_N^{k+1}\| = o(k^{-1}), \quad \text{as } k \rightarrow +\infty.$$

(iii) Suppose $A_i = \partial f_i$ for all $i \in \{1, \dots, N\}$ and fix an index $j \in \{1, \dots, N\}$. Suppose f_i is locally Lipschitz continuous for all $j \neq i$. Let $f := f_1 + \dots + f_N$, then

$$f(x_j^{k+1}) - \inf f = o(k^{-1}) \quad \text{as } k \rightarrow +\infty.$$

Furthermore, if each f_i is locally Lipschitz, the same holds for $f(\bar{x}^{k+1}) - \inf f$.

Proof. Since x^{k+1} coincides with the block-vector containing the first N components of $(\mathcal{M} + \mathbf{A})^{-1}(\mathcal{C}w^k)$, (i) and (ii) follow directly from Theorem 4.3, (4.15), and the same argument as in [25, Lemma 5.2.2]. Statement (iii) follows by using the same argument as in [18, Proposition 3.9], which basically consists in showing $f(x_j^{k+1}) - \inf f$ can be bounded by the fixed-point residual. \square

5. NUMERICAL EXPERIMENTS

In this section, we present our numerical experiments, which were performed in Python on a 12th-Gen. Intel(R) Core(TM) i7-1255U, 1.70–4.70 GHz laptop with 16 Gb of RAM. The code is available for reproducibility at <https://github.com/echnen/fast-km>.

5.1. **Toy problems.** In these first numerical experiments, we consider two main instances of T :

- We consider an optimization problem that can be written as

$$\min_{x \in \mathbb{R}^d} f_1(x) + \cdots + f_N(x),$$

for $f_1, \dots, f_N: \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ proper, convex and lower-semicontinuous that are also *prox-friendly*, i.e., such that $\text{prox}_{\tau f_i}$ can be easily computed for each $\tau > 0$. In this case, we take T as given in (4.12) and thus use Algorithm 2 initialized with $w^0 = w^1 = 0$.

- As in [15], for dimension $d \in \mathbb{N}$ even, we consider $T: \mathbb{R}^d \rightarrow \mathbb{R}^d$ defined by $T(x) := J_{\tau M}(x)$, where $\tau = 10^{-1}$ and M is the skew-symmetric (hence monotone) matrix given block-wise by:

$$M := \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix},$$

where I is now the identity operator in $\mathbb{R}^{d/2}$ and 0 represents the zero operator in $\mathbb{R}^{d/2}$. In this case, we use Algorithm 1 initialized with $x^0 = x^1 = 1$.

5.2. **The influence of parameters.** As we introduced two degrees of flexibility in Algorithm 1 and Algorithm 2 consisting in the parameter $\eta \in (0, 1)$ and $\sigma > 0$, we first numerically investigate the influence of these two parameters separately. We consider the two instances of T in Section 5.1. In the first one, we set $d = 2$, $N = 2$ and take:

$$f_1(x) := 10^{-3}|x|, \quad \text{and} \quad f_2(x) := \frac{1}{2} \text{dist}^2(x, \mathbb{B}), \quad x \in \mathbb{R}^2,$$

where \mathbb{B} is the unit euclidean ball centered at $(1, 1)^\top$. The two proximity operators are easily accessed through, soft-thresholding and averaged projection onto \mathbb{B} , respectively; see [10, Example 14.5 and Proposition 24.8(vii)]. In the second one, we set $d = 10$.

5.2.1. *Influence of η .* In this section, we perform a systematic comparison between $\eta = \frac{1}{2}$, which was introduced (when $\sigma = \alpha$) in [15] and the choices $\eta \in (0, 1)$. Specifically, we consider $\alpha = 2, 4, 16, 32$ and two different choices of σ , $\sigma = 2$ and $\sigma = 16$. Note in particular that the Fast-KM method in [15] is considered with $\alpha = 32$. When $\alpha = 2$ the methods with different values of η all coincide (since $\alpha_\eta = 1$ for all $\eta \in (0, 1)$) and the resulting method is OHM, see Remark 3.2. This is reflected in our experiments. We plot the results in Figure 1 and Figure 2.

Our numerical experiments show that Algorithm 1 is particularly competitive against the baseline choice with $\eta = \frac{1}{2}$. From Figure 1, we can indeed observe an improved convergence behavior when $\eta < \frac{1}{2}$, which is not reflected in Figure 2. The nonincrease property of $E_k^{\alpha-1}$ (see Lemma 3.3 and Theorem 3.4) provides us with interesting insights supporting this numerical evidence. Consider (3.10) and set $\lambda = \alpha - 1$. Let us also emphasize here the dependency of E_k^λ on η by writing $E_k^{\lambda, \eta}$. Recalling that the first term in (3.10) is negative, simple algebraic operations lead to:

$$(5.1) \quad \Delta E_k^{\alpha-1, \eta} \leq -\eta(1-\eta)(\alpha-2) \left[t_k - \frac{\alpha}{2} \right] \|x^{k-1} - T(x^{k-1})\|^2, \quad \text{for all } k \geq 2 \text{ such that } t_k \geq 2(\alpha-1).$$

Therefore, if $t_k \geq 2(\alpha-1)$ for all $k \geq 2$ (hence $t_k \geq \alpha/2$ as well since $\alpha \geq 2$), which is the case if $\sigma \geq 2\alpha-3$, it holds $\Delta E_k^{\alpha-1, \eta} \leq 0$ for all $k \geq 2$. Observe that in our examples, T is firmly nonexpansive, and is thus the resolvent of the maximal monotone operator $A := T^{-1} + I$. We then have $Q(x) \in A(T(x)) = A(z)$. In turn, the term $\langle Q^k, z^k - z^* \rangle$ in (3.8) is nonnegative since $Q^k \in A(z^k) = A(T(x^{k-1}))$. To emphasize this fact, let us rename here Q^k as A^{k-1} . We then obtain for all $k \geq 2$:

$$(5.2) \quad \langle A^{k-1}, T(x^{k-1}) - z^* \rangle \leq \frac{E_2^{\alpha-1, \eta}}{(1-\eta)(\alpha-1)t_{k-1}}, \quad \text{and} \quad \frac{1}{2} \|x^{k-1} - T(x^{k-1})\|^2 \leq \frac{E_2^{\alpha-1, \eta}}{(1-\eta)\eta t_{k-1}^2},$$




FIGURE 1. Results of Section 5.2.1 with the first instance of T described in Section 5.1: Comparison of different choices of $\eta \in (0, 1)$, with corresponding colors indicated in the colorbar. $\eta = \frac{1}{2}$ corresponds to the baseline case considered in [15] and is highlighted in red.

If $x^0 = x^1$, the energy function at $k = 2$ reads as

$$\begin{aligned}
 (5.3) \quad E_2^{\alpha-1, \eta} &= (1 - \eta)(\alpha - 1)t_1 \left(\langle A^0, T(x^0) - z^* \rangle + \frac{1}{2} \|x^0 - T(x^0)\|^2 \right) \\
 &\quad + \frac{(1 - \eta)\eta t_1^2}{2} \|x^0 - T(x^0)\|^2 + \frac{1}{2} \|(\alpha - 1)(T(x^0) - z^*) + t_1 \eta (x^0 - T(x^0))\|^2.
 \end{aligned}$$

By developing the square in the last norm in (5.3), it is easy to observe that $E_2^{\alpha-1, \eta}$ has a linear dependency on η , i.e., $E_2^{\alpha-1, \eta} = c_0 + c_1 \eta$ for some $c_0 \geq 0$ and $c_1 \in \mathbb{R}$ such that $c_1 \geq -c_0$, only depending on x^0 , on the operator T , on α and σ . The optimal rate for the fixed-point residual is then attained for η minimizing $\eta \mapsto \frac{c_0 + c_1 \eta}{\eta(1 - \eta)}$, which is greater or smaller than $\frac{1}{2}$ depending only on c_0 and c_1 . On the other hand, the optimal rate for the inner product $\langle A^{k-1}, T(x^{k-1}) - z^* \rangle$ is attained by minimizing $\eta \mapsto \frac{c_0 + c_1 \eta}{1 - \eta}$, which always suggests setting $\eta \simeq 0$ (since $\eta = 0$ is not admissible).

If A is a linear skew-symmetric operator, as in our second example, the inner product $\langle A^{k-1}, T(x^{k-1}) - z^* \rangle$ is always zero, suggesting to fix η as the minimizer of $\eta \mapsto \frac{c_0 + c_1 \eta}{\eta(1 - \eta)}$, which in our case is in fact $\eta \simeq \frac{1}{2}$, at least fixing $\sigma = 2\alpha - 3$. In other cases, such as when T is a resolvent of a primal-dual operator A , like in our first example, the inner product on the left hand-side of (5.2) can become more relevant as it is often related to primal-dual gaps, hence suggesting $\eta \simeq 0$. We leave a more detailed analysis on this intriguing trade-off between the two rates in (5.2) for a future work.

5.2.2. *The influence of σ .* Keeping in mind (3.4), the parameter σ is related to the starting time t_0 in the continuous dynamics (2.1). Recall in fact that the discrete time is defined by $t_k = k - 1 + \sigma$ so $t_0 = \sigma - 1$. We test 20 different choices of σ spaced evenly from 1 to 100 and take the four choices $\alpha = 2, 4, 16, 32$. We put particular emphasis on the choice $\sigma = \alpha$, which was introduced in [15]. We consider and only the operator T corresponding to the second instance of Section 5.1. The results are depicted in Figure 3. We refrain from displaying the results for the first instance of T in Section 5.1 since the results therein




FIGURE 2. Results of Section 5.2.1 with T as described the second instance of T described in Section 5.1: Comparison of different choices of $\eta \in (0, 1)$, with corresponding colors indicated in the colorbar. $\eta = \frac{1}{2}$ corresponds to the baseline case considered in [15] and is highlighted in red.

are almost indistinguishable. Regarding the tuning of the parameter σ , our experiments suggest indeed setting $\sigma = \alpha$.

5.3. Cooling Heuristics. Motivated by the insights of Sections 5.2.1, and 5.2.2, we propose a “cooling” heuristics⁴, which consists in gradually increasing the parameter α along the iterations until a certain upper bound α_{\max} is reached:

$$(5.4) \quad \alpha_0 < \alpha_1 < \dots < \alpha_{\lfloor \text{maxit}/2 \rfloor} = \alpha_{\max}.$$

Specifically, we consider two different strategies: a *linear* and a *logarithmic* parameter scaling. In either case, we set $\alpha_0 > 0$, $\alpha_{\max} = 100\alpha_0$, and we fix a maximum number of iterations maxit . Then, in first case, at each iteration we increase α with a fixed step-size, in such a way that α_{\max} is reached for $k = \lfloor \text{maxit}/2 \rfloor$. In the second case, the increase is defined in a logarithmic scale. We consider $\alpha = 2, 4, 16, 32$, and the operator T as given in the second choice in Section 5.1. For each α , we compare the performance of the vanilla Krasnosel’kiĭ–Mann (KM) method, Fast-KM with $\eta = \frac{1}{2}$, and Fast-KM with linear and logarithmic scaling in the cooling heuristics and $\eta = \frac{1}{2}$, $\alpha_0 = \alpha$, abbreviated (fKM-LNC) and (fKM-LGC), respectively. Recall that $\alpha_{\max} = 100\alpha_0$. The results are shown in Figure 4. A notable insight from these experiment is the beneficial convergence speed of Algorithm 1 with $\alpha = 32$, but without incurring into the undesirable initial instability state; see [1] who thoroughly studied the choice of α for the ODE (1.4) when $\beta(t) \equiv 0$, $b(t) \equiv 1$ and Q is the gradient of a convex function.

5.4. Application to Realistic Problems. In this section, we test Algorithm 1 and its application in Algorithm 2 to solve optimization problems arising from realistic settings. We start with an application to Optimal Transport (OT), and in particular, to the *Beckmann minimal flow formulation* of the Wasserstein 1 distance, where DRS proved particularly effective [37, 12, 22]. In the second experiment, we consider the geometric median problem, a standard nonsmooth benchmark especially for distributed optimization.

⁴The name is borrowed from Consensus Based Optimization literature [30].




FIGURE 3. Results of Section 5.2.2: Comparison between different choices of $\sigma \in (0, 100)$. $\sigma = \alpha$ corresponds to the baseline case considered in [15] and is highlighted in red.




FIGURE 4. Results of Section 5.3: Testing Cooling Heuristics.

In these two experiments, we compare the performance of i) the vanilla Krasnosel'kiĭ–Mann (KM) method, ii) Algorithm 1 with $\eta = \frac{1}{2}$, iii) Algorithm 1 with $\eta = 0.1$ and iii) Algorithm 1 with a linear cooling heuristics as introduced in Section 5.3. The results are reported in Figure 5.

5.4.1. Optimal Transport. In this experiment, we tackle a high dimensional OT problem, which is usually solved in its primal form using the celebrated Sinkhorn Algorithm [13], which actually solves entropic-regularized versions of the OT problem.

Consider a square grid $\Omega := \{(i, j) \mid i, j \in \{1, \dots, p\}\}$, with $p \in \mathbb{N}$, and two discrete probability measures μ , and ν on Ω , i.e., $\mu, \nu \in \mathbb{R}_+^n$ with $n = p^2$ such that $\sum_{i=1}^n \mu_i = \sum_{i=1}^n \nu_i = 1$. Denoted by div the discrete divergence operator with Neumann boundary conditions on Ω defined through finite




FIGURE 5. Results of Section 5.4. Left: Results of Section 5.4.1 (Optimal Transport). Right: Results of Section 5.4.2 (Geometric Median).

differences (see, e.g., [22]), we aim at solving the following convex problem:

$$(5.5) \quad \min_{\sigma \in \mathbb{R}^{n \times 2}} \|\sigma\|_{2,1}, \quad \text{s.t.:} \quad \text{div } \sigma = \mu - \nu,$$

where $\|\sigma\|_{2,1}$ is the *group-lasso* penalty, i.e., $\|\sigma\|_{2,1} := \sum_{i=1}^n \|\sigma_i\|$, and for each $i \in \{1, \dots, n\}$, $\sigma_i \in \mathbb{R}^2$. The problem can be addressed by applying the DRS method with $f_1(\sigma) := \|\sigma\|_{2,1}$, and $f_2(\sigma) := \iota_{\mathcal{C}}(\sigma)$, where $\mathcal{C} := \{\sigma \in \mathbb{R}^{n \times 2} \mid \text{div } \sigma = \mu - \nu\}$. These two functions are indeed *prox-friendly*. prox_{f_1} is the group soft-thresholding operator (see [10, Example 14.5]), while prox_{f_2} coincides with the projection onto the constraint set \mathcal{C} , which requires solving a Poisson equation on Ω . This can be efficiently solved either by using dedicated sparse solvers or by FFT. We apply Algorithm 2 to solve (5.5). As can be seen from Figure 5, Algorithm 2 emerges here as a great practical alternative: the method outperformed the standard approach with KM, and confirmed once again that a choice of η small is indeed beneficial in practice, especially when combined with the cooling heuristic.

5.4.2. Solving the Geometric Median Problem. We now turn to computing the geometric median of N sample points in \mathbb{R}^d . Given a sample $\{x_i\}_i$ of N points in \mathbb{R}^d we aim at solving the following convex problem

$$(5.6) \quad \min_{x \in \mathbb{R}^d} \sum_{i=1}^N \|x - x_i\|.$$

This problem can be shown to be strictly convex if $\{x_i\}_i$ are not collinear and hence admits a unique solution. To solve (5.6), we consider the graph-DRS operator in (4.12) splitting (5.6) in N terms. All the terms $x \mapsto \|x - x_i\|$ are in fact prox-friendly with proximity operators that can be computed by simply applying the soft-thresholding operator suitably shifted by x_i . To define T in (4.12) and apply Algorithm 2, we consider $\bar{Z} = 0$ and build an arbitrary Z such that (4.11) holds. The same conclusions drawn from the OT experiment in Section 5.4.2 remain true here.

6. CONCLUSION

In this paper, we introduced a generalized version of the Fast-KM method proposed in [15], incorporating two additional degrees of flexibility through the parameters $\eta \in (0, 1)$ and $\sigma > 0$. We provided a new convergence proof based on a semi-implicit discretization of a continuous time model. Notably, our approach encompasses the case $\alpha = 2$, which corresponds to the Optimal Halpern Method, thereby unifying the convergence analysis of these two acceleration mechanisms. We demonstrated how these results can be applied to operator splitting methods potentially with preconditioners, offering critical convergence

guarantees for the so-called shadow sequences. Furthermore, our numerical experiments showcased the benefits afforded by the added flexibility of the method. This paper opens up new intriguing questions, among which: i) is the concept of Hessian-driven damping truly fundamental in the operator setting ? ii) Can the restrictions on η be removed by setting, e.g., $\eta = 0$ or $\eta = 1$? We leave this for future research.

APPENDICES

A. Discrete Calculus Rules. For three sequences $(a_k)_{k \in \mathbb{N}}$, $(b_k)_{k \in \mathbb{N}}$, $(c_k)_{k \in \mathbb{N}}$, we have:

$$(6.1) \quad \begin{aligned} \Delta(ab)_k &= \Delta a_k b^k + a^{k+1} \Delta b_k, \\ \Delta(abc)_k &= \Delta a_k b^k c^k + a^{k+1} \Delta b_k c^k + a^{k+1} b^{k+1} \Delta c_k, \\ \Delta a_k^2 &= 2(a_{k+1} - a_k) a_k + (\Delta a_k)^2 = 2(a_{k+1} - a_k) a_{k+1} - (\Delta a_k)^2. \end{aligned}$$

B. Proof of the main technical lemmas.

Proof of Lemma 2.1. For the sake of notation, we also abbreviate $E_t^\lambda := E^\lambda(t)$, $\dot{x}_t := \dot{x}(t)$, $v_t^\lambda := v^\lambda(t)$, $\delta_t := \delta(t)$, and $\xi_t := \xi(t)$. Additionally, instead of writing $\frac{d}{dt}$ we write ∂_t . Using Leibniz rule, we get for almost all $t \geq t_0$:

$$(6.2) \quad \begin{aligned} \partial_t E_t^\lambda &= \dot{\delta}_t \langle Q_t, x_t - x^* \rangle + \delta_t \langle \partial_t Q_t, x_t - x^* \rangle + \delta_t \langle Q_t, \dot{x}_t \rangle \\ &\quad + \frac{\dot{\xi}_t}{2} \|Q_t\|^2 + \xi_t \langle \partial_t Q_t, Q_t \rangle + c \langle x_t - x^*, \dot{x}_t \rangle + \langle v_t, \dot{v}_t \rangle. \end{aligned}$$

Let us compute the last inner product $\langle v_t^\lambda, \dot{v}_t^\lambda \rangle$. For \dot{v}_t we have from the constitute equation (2.1) that for almost all $t \geq t_0$:

$$\begin{aligned} \dot{v}_t^\lambda &= (\lambda + 1) \dot{x}_t + \eta \beta_t Q_t + t \left(-\frac{\alpha}{t} \dot{x}_t - \beta_t \partial_t Q_t - b_t Q_t + \eta \dot{\beta}_t Q_t + \eta \beta_t \partial_t Q_t \right) \\ &= (\lambda + 1 - \alpha) \dot{x}_t + \eta \beta_t Q_t - t(1 - \eta) \beta_t \partial_t Q_t + t(\eta \dot{\beta}_t - b_t) Q_t \\ &= -(\alpha - 1 - \lambda) \dot{x}_t - t \left(b_t - \eta \dot{\beta}_t - \eta \frac{\beta_t}{t} \right) Q_t - t \beta_t (1 - \eta) \partial_t Q_t. \end{aligned}$$

Denoting by $w_t := b_t - \eta \dot{\beta}_t - \eta \frac{\beta_t}{t}$, the inner products reads for almost all $t \geq t_0$ as

$$(6.3) \quad \begin{aligned} \langle v_t^\lambda, \dot{v}_t^\lambda \rangle &= -\lambda(\alpha - 1 - \lambda) \langle x_t - x^*, \dot{x}_t \rangle - (\alpha - 1 - \lambda) t \|\dot{x}_t\|^2 - \eta(\alpha - 1 - \lambda) t \beta_t \langle Q_t, \dot{x}_t \rangle \\ &\quad - \lambda t w_t \langle x_t - x^*, Q_t \rangle - t^2 w_t \langle \dot{x}_t, Q_t \rangle - t^2 \beta_t w_t \eta \|Q_t\|^2 \\ &\quad (1 - \eta) t \lambda \beta_t \langle x_t - x^*, \partial_t Q_t \rangle - (1 - \eta) t^2 \beta_t \langle \dot{x}_t, \partial_t Q_t \rangle - t^2 \beta_t^2 (1 - \eta) \eta \langle Q_t, \partial_t Q_t \rangle. \end{aligned}$$

Plugging (6.3) into (6.2), and canceling out the appropriate terms, we obtain for almost all $t \geq t_0$ that

$$(6.4) \quad \begin{aligned} \partial_t E_t^\lambda &= \left(\dot{\delta}_t - t \lambda w_t \right) \langle Q_t, x_t - x^* \rangle - (1 - \eta) t^2 \beta_t \langle \dot{x}_t, \partial_t Q_t \rangle \\ &\quad + \left(\delta_t - \eta(\alpha - 1 - \lambda) t \beta_t - t^2 w_t \right) \langle Q_t, \dot{x}_t \rangle + \left(\frac{\dot{\xi}_t}{2} - \eta t^2 \beta_t w_t \right) \|Q_t\|^2 - (\alpha - 1 - \lambda) t \|\dot{x}_t\|^2. \end{aligned}$$

Let us study the quadratic form in the second line of (6.4) separately⁵. Plugging in the definitions of δ_t , ξ_t in (2.4) and of b in (2.2), we get, first, $w_t = (\alpha - 1)(1 - \eta) \frac{\beta_t}{t}$, hence,

$$\begin{aligned} \delta_t - \eta(\alpha - 1 - \lambda) t \beta_t - t^2 w_t &= -t \beta_t (\alpha - 1 - \lambda), \\ \frac{\dot{\xi}_t}{2} - \eta t^2 \beta_t w_t &= \beta_t t^2 (1 - \eta) \eta \left(\dot{\beta}_t + \frac{(2 - \alpha)}{t} \beta_t \right), \end{aligned}$$

⁵The coefficient b_t in (2.2) is in fact the only parameter that (i) does not depend on λ (artificial parameter introduced in the Lyapunov analysis), and (ii) can make this quadratic form negative semidefinite for all $\lambda \in (0, \alpha - 1]$. To see this, consider $\lambda = \alpha - 1$. Then, δ_t must be equal to $t^2 w_t$, which yields (2.2). This is the crucial difference to the case $Q = \nabla f$ in [5], where the blue term does not appear.

so the quadratic form simplifies to:

$$\begin{aligned}
& (\delta_t - \eta(\alpha - 1 - \lambda)t\beta_t - t^2w_t) \langle Q_t, \dot{x}_t \rangle + \left(\frac{\dot{\xi}_t}{2} - \eta t^2 \beta_t w_t \right) \|Q_t\|^2 - (\alpha - 1 - \lambda)t \|\dot{x}_t\|^2 \\
&= -t\beta_t(\alpha - 1 - \lambda) \langle Q_t, \dot{x}_t \rangle + \beta_t t^2(1 - \eta)\eta \left(\dot{\beta}_t + \frac{(2 - \alpha)}{t} \beta_t \right) \|Q_t\|^2 - (\alpha - 1 - \lambda)t \|\dot{x}_t\|^2 \\
&= -\frac{t}{2}(\alpha - 1 - \lambda) \|\dot{x}_t + \beta_t Q_t\|^2 - \frac{t}{2}(\alpha - 1 - \lambda) \|\dot{x}_t\|^2 \\
&\quad + \beta_t t^2 \left((1 - \eta)\eta \left(\dot{\beta}_t + \frac{\beta_t}{t}(2 - \alpha) \right) + \frac{1}{2}(\alpha - 1 - \lambda) \frac{\beta_t}{t} \right) \|Q_t\|^2.
\end{aligned}$$

Plugging this and the definition of ξ_t and of w_t into (6.4), we get the claim. \square

Proof of Lemma 3.3. Let $k \geq 2$. Apply the discrete Leibniz rule (see Appendix A) to get:

$$\begin{aligned}
(6.5) \quad \Delta E_k^\lambda &= \Delta \delta_k \langle Q^k, z^k - z^* \rangle + \delta_{k+1} \langle \Delta Q_k, z^k - z^* \rangle + \delta_{k+1} \langle Q^{k+1}, \Delta z_k \rangle + \Delta \left(\frac{\delta}{2} \|Q\|^2 \right)_k \\
&\quad + \frac{1}{2} \Delta \xi_k \|Q^k\|^2 + \xi_{k+1} \langle \Delta Q_k, Q^{k+1} \rangle - \frac{1}{2} \xi_{k+1} \|\Delta Q_k\|^2 \\
&\quad + c \langle \Delta z_{k-1}, z^k - z^* \rangle - \frac{c}{2} \|\Delta z_{k-1}\|^2 \\
&\quad + \langle \Delta v_k, v^{k+1} \rangle - \frac{1}{2} \|\Delta v_k\|^2.
\end{aligned}$$

Following the continuous time approach, we first write Δv_k only in terms of $z^{k-1} - z^*$, Q^k and ΔQ_k , then compute $\langle \Delta v_k, v^{k+1} \rangle$ and finally plug everything into ΔE_k^λ . This will give the necessary cancellations. Using that $\Delta t_k = 1$ and the update of Algorithm 1 (see (3.4)), we get

$$\begin{aligned}
(6.6) \quad \Delta v_k &= \lambda \Delta z_{k-1} + \Delta t_{k-1} (\Delta z_{k-1} + \eta Q^k) + t_k (\Delta^2 z_{k-1} + \eta \Delta Q_k) \\
&= \lambda \Delta z_{k-1} + (\Delta z_{k-1} + \eta Q^k) + t_k \left(-\frac{\alpha}{t_k} \Delta z_{k-1} - \Delta Q_k - \frac{\alpha \eta}{t_k} Q^k + \eta \Delta Q_k \right) \\
&= -(\alpha - 1 - \lambda) \Delta z_{k-1} - (\alpha_\eta - \eta) Q^k - t_k (1 - \eta) \Delta Q_k \\
&= -(\alpha - 1 - \lambda) \Delta z_{k-1} - (1 - \eta)(\alpha - 1) Q^k - t_k (1 - \eta) \Delta Q_k.
\end{aligned}$$

Let us now evaluate $\langle \Delta v_k, v^{k+1} \rangle$, we get:

$$\begin{aligned}
(6.7) \quad \langle \Delta v_k, v^{k+1} \rangle &= -\lambda(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, z^k - z^* \rangle - t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta z_k \rangle - (\alpha - 1 - \lambda)t_k \eta \langle \Delta z_{k-1}, Q^{k+1} \rangle \\
&\quad - \lambda(1 - \eta)(\alpha - 1) \langle Q^k, z^k - z^* \rangle - t_k(1 - \eta)(\alpha - 1) \langle Q^k, \Delta z_k \rangle - t_k \eta(1 - \eta)(\alpha - 1) \langle Q^k, Q^{k+1} \rangle \\
&\quad - \lambda t_k(1 - \eta) \langle \Delta Q_k, z^k - z^* \rangle - t_k^2(1 - \eta) \langle \Delta Q_k, \Delta z_k \rangle - \eta(1 - \eta)t_k^2 \langle \Delta Q_k, Q^{k+1} \rangle.
\end{aligned}$$

Let us now inspect $\|\Delta v_k\|^2$:

$$\begin{aligned}
(6.8) \quad -\frac{1}{2} \|\Delta v_k\|^2 &= -\frac{1}{2}(1 - \eta)^2(\alpha - 1)^2 \|Q^k\|^2 - \frac{t_k^2}{2}(1 - \eta)^2 \|\Delta Q_k\|^2 - \frac{1}{2}(\alpha - 1 - \lambda)^2 \|\Delta z_{k-1}\|^2 \\
&\quad - (\alpha - 1)(1 - \eta)^2 t_k \langle Q^k, \Delta Q_k \rangle - (1 - \eta)(\alpha - 1 - \lambda)(\alpha - 1) \langle Q^k, \Delta z_{k-1} \rangle \\
&\quad - (1 - \eta)t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta Q_k \rangle.
\end{aligned}$$

Plugging (6.7) and (6.8) into (6.5) and rearranging suitably we get

$$\begin{aligned}
(6.9) \quad \Delta E_k^\lambda &= -\lambda(1 - \eta)(\alpha - 2) \langle Q^k, z^k - z^* \rangle - t_k^2(1 - \eta) \langle \Delta Q_k, \Delta z_k \rangle \\
&\quad - (1 - \eta)t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta Q_k \rangle
\end{aligned}$$

$$(6.10) \quad + \frac{1}{2} \eta(1 - \eta) \Delta t_{k-1}^2 \|Q^k\|^2 - t_k \eta(1 - \eta)(\alpha - 1) \langle Q^k, Q^{k+1} \rangle$$

$$\begin{aligned}
(6.11) \quad & - \frac{(1-\eta)^2}{2}(\alpha-1)^2\|Q^k\|^2 - (\alpha-1)(1-\eta)^2t_k \langle Q^k, \Delta Q_k \rangle \\
& + \lambda(1-\eta)t_k \langle Q^{k+1}, \Delta z_k \rangle - (\alpha-1-\lambda)t_k\eta \langle \Delta z_{k-1}, Q^{k+1} \rangle - t_k(1-\eta)(\alpha-1) \langle Q^k, \Delta z_k \rangle \\
& - (1-\eta)(\alpha-1-\lambda)(\alpha-1) \langle Q^k, \Delta z_{k-1} \rangle
\end{aligned}$$

$$\begin{aligned}
(6.12) \quad & - t_k(\alpha-1-\lambda) \langle \Delta z_{k-1}, \Delta z_k \rangle \\
& - \frac{1}{2}(\alpha-1-\lambda)^2\|\Delta z_{k-1}\|^2
\end{aligned}$$

$$(6.13) \quad - \frac{1}{2}(1-\eta)t_k^2\|\Delta Q_k\|^2 - \frac{\lambda}{2}(\alpha-1-\lambda)\|\Delta z_{k-1}\|^2 + \Delta \left(\frac{\delta}{2}\|Q\|^2 \right)_k,$$

where in the last line we have used $\eta(1-\eta) + (1-\eta)^2 = (1-\eta)$. To deal with the various nonnegative off-sets in the inner products (6.10), (6.11) and (6.12), we make a discrete version of the the continuous time quadratic form appear and control the remainder with Young's inequality. This gives

$$\begin{aligned}
(6.10) \quad & := \frac{1}{2}\eta(1-\eta)\Delta t_{k-1}^2\|Q^k\|^2 - t_k\eta(1-\eta)(\alpha-1) \langle Q^k, Q^{k+1} \rangle \\
& - \frac{(1-\eta)^2}{2}(\alpha-1)^2\|Q^k\|^2 - (\alpha-1)(1-\eta)^2t_k \langle Q^k, \Delta Q_k \rangle \\
& = (1-\eta) \left(\frac{\eta}{2}\Delta t_{k-1}^2 - \eta t_k(\alpha-1) - \frac{(1-\eta)}{2}(\alpha-1)^2 \right) \|Q^k\|^2 - t_k(\alpha-1)(1-\eta) \langle Q^k, \Delta Q_k \rangle \\
& = - \underbrace{\left[\eta(1-\eta) \left((\alpha-2)t_k + \frac{1}{2} \right) + \frac{1}{2}(1-\eta)^2(\alpha-1)^2 \right]}_{(Q.1)} \|Q^k\|^2 - t_k(\alpha-1)(1-\eta) \langle Q^k, \Delta Q_k \rangle,
\end{aligned}$$

where in the coefficient that multiplies $\|Q^k\|^2$ we have used, by the definition of t_k ,

$$(6.14) \quad \frac{1}{2}\Delta t_{k-1}^2 = \frac{1}{2}((k-1+\sigma)^2 - (k-2+\sigma)^2) = t_k - \frac{1}{2},$$

hence $\frac{1}{2}\Delta t_{k-1}^2 - t_k(\alpha-1) = -\frac{1}{2} - t_k(\alpha-2)$. The first term of our quadratic form involving $\|Q^k\|^2$ appears. We now force $\langle Q^k, \Delta z_{k-1} \rangle$ to appear. This comes from the fact that

$$\begin{aligned}
(6.11) \quad & := \lambda(1-\eta)t_k \langle Q^{k+1}, \Delta z_k \rangle - (\alpha-1-\lambda)t_k\eta \langle \Delta z_{k-1}, Q^{k+1} \rangle - t_k(1-\eta)(\alpha-1) \langle Q^k, \Delta z_k \rangle \\
& - (1-\eta)(\alpha-1-\lambda)(\alpha-1) \langle Q^k, \Delta z_{k-1} \rangle \\
& = \lambda(1-\eta)t_k \langle Q^k, \Delta z_{k-1} \rangle - (\alpha-1-\lambda)t_k\eta \langle Q^k, \Delta z_{k-1} \rangle - t_k(1-\eta)(\alpha-1) \langle Q^k, \Delta z_{k-1} \rangle \\
& + \lambda(1-\eta)t_k \langle Q^{k+1}, \Delta^2 z_{k-1} \rangle - (\alpha-1-\lambda)t_k\eta \langle \Delta Q_k, \Delta z_{k-1} \rangle - t_k(1-\eta)(\alpha-1) \langle Q^k, \Delta^2 z_{k-1} \rangle \\
& + \lambda(1-\eta)t_k \langle \Delta Q_k, \Delta z_{k-1} \rangle - (1-\eta)(\alpha-1-\lambda)(\alpha-1) \langle Q^k, \Delta z_{k-1} \rangle \\
& = \underbrace{-(\alpha-1-\lambda)t_k \langle Q^k, \Delta z_{k-1} \rangle - t_k(1-\eta)(\alpha-1-\lambda) \langle Q^k, \Delta^2 z_{k-1} \rangle}_{(Q.2)} \\
& + \lambda(1-\eta)t_k \langle \Delta Q_k, \Delta^2 z_{k-1} \rangle - (\alpha-1-\lambda)t_k\eta \langle \Delta Q_k, \Delta z_{k-1} \rangle \\
& + \lambda(1-\eta)t_k \langle \Delta Q_k, \Delta z_{k-1} \rangle - (1-\eta)(\alpha-1-\lambda)(\alpha-1) \langle Q^k, \Delta z_{k-1} \rangle \\
& = (Q.2) \\
& \underbrace{-t_k(1-\eta)(\alpha-1-\lambda) \langle Q^k, \Delta^2 z_{k-1} \rangle}_{(II)} + \lambda(1-\eta)t_k \langle \Delta Q_k, \Delta z_k \rangle - (\alpha-1-\lambda)t_k\eta \langle \Delta Q_k, \Delta z_{k-1} \rangle \\
& \underbrace{-(1-\eta)(\alpha-1-\lambda)(\alpha-1) \langle Q^k, \Delta z_{k-1} \rangle}_{(N.1)}.
\end{aligned}$$

We will handle (II) separately later. To conclude, we make the term in $\|\Delta z_{k-1}\|^2$ appear. Starting with (6.12) we have:

$$(6.12) := -t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta z_k \rangle - \frac{1}{2}(\alpha - 1 - \lambda)^2 \|\Delta z_{k-1}\|^2 \\ = \underbrace{-t_k(\alpha - 1 - \lambda) \|\Delta z_{k-1}\|^2}_{(Q.3)} - \underbrace{t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta^2 z_{k-1} \rangle}_{(III)} - \underbrace{\frac{1}{2}(\alpha - 1 - \lambda)^2 \|\Delta z_{k-1}\|^2}_{(N.2)}.$$

Once again, we will deal with (III) later. Gathering all the terms together, we get

$$(6.15) \quad \Delta E_k^\lambda = -\lambda(1 - \eta)(\alpha - 2) \langle Q^k, z^k - z^* \rangle - t_k^2(1 - \eta) \langle \Delta Q_k, \Delta z_k \rangle \\ - (1 - \eta)t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta Q_k \rangle \\ + (Q.1) - t_k(\alpha - 1)(1 - \eta) \langle Q^k, \Delta Q_k \rangle \\ + (Q.2) + (II) + \lambda(1 - \eta)t_k \langle \Delta Q_k, \Delta z_k \rangle - (\alpha - 1 - \lambda)t_k\eta \langle \Delta Q_k, \Delta z_{k-1} \rangle + (N.1) \\ + (Q.3) + (III) + (N.2) \\ - \frac{1}{2}(1 - \eta)t_k^2 \|\Delta Q_k\|^2 - \frac{\lambda}{2}(\alpha - 1 - \lambda) \|\Delta z_{k-1}\|^2 + \Delta \left(\frac{\delta}{2} \|Q\|^2 \right)_k \\ = -\lambda(1 - \eta)(\alpha - 2) \langle Q^k, z^k - z^* \rangle - (t_k^2 - \lambda t_k)(1 - \eta) \langle \Delta Q_k, \Delta z_k \rangle \\ - t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta Q_k \rangle \\ + (Q.1) - t_k(\alpha - 1)(1 - \eta) \langle Q^k, \Delta Q_k \rangle \\ + (Q.2) + (II) + (N.1) \\ + (Q.3) + (III) + (N.2) \\ - \frac{1}{2}(1 - \eta)t_k^2 \|\Delta Q_k\|^2 - \frac{\lambda}{2}(\alpha - 1 - \lambda) \|\Delta z_{k-1}\|^2 + \Delta \left(\frac{\delta}{2} \|Q\|^2 \right)_k,$$

Let us now collect all the terms in (Q.2) and (Q.3) and argue analogously to the continuous-time setting to get

$$(6.16) \quad (Q.2) + (Q.3) = -(\alpha - 1 - \lambda)t_k \langle Q^k, \Delta z_{k-1} \rangle - t_k(\alpha - 1 - \lambda) \|\Delta z_{k-1}\|^2 \\ = -(\alpha - 1 - \lambda)t_k \left(\langle Q^k, \Delta z_{k-1} \rangle + \|\Delta z_{k-1}\|^2 \right) \\ = -(\alpha - 1 - \lambda)t_k \left(\frac{1}{2} \|Q^k + \Delta z_{k-1}\|^2 + \frac{1}{2} \|\Delta z_{k-1}\|^2 - \frac{1}{2} \|Q^k\|^2 \right) \\ = (\alpha - 1 - \lambda) \frac{t_k}{2} \|Q^k\|^2 - (\alpha - 1 - \lambda) \frac{t_k}{2} \|\Delta z_{k-1}\|^2 - (\alpha - 1 - \lambda) \frac{t_k}{2} \|Q^k + \Delta z_{k-1}\|^2.$$

Note that the positive term $(\alpha - 1 - \lambda) \frac{t_k}{2}$ in the last display is not a concern when $\alpha > 2$ since we shall take λ arbitrarily close to $\alpha - 1$ in our Lyapunov analysis in such a way that when combined with (Q.1), the coefficient in front of $\|Q^k\|^2$ will be negative. The case $\alpha = 2$ will be treated with $\lambda = \alpha - 1$. We now deal with (II) and (III), while the terms labeled with a capital ‘‘N’’ will be handled properly later.

We have

$$(6.17) \quad (II) = -(\alpha - 1 - \lambda)t_k(1 - \eta) \langle Q^k, \Delta^2 z_{k-1} \rangle \\ = -(\alpha - 1 - \lambda)t_k(1 - \eta) \left\langle Q^k, -\frac{\alpha}{t_k} \Delta z_{k-1} - \Delta Q_k - \frac{\alpha\eta}{t_k} Q^k \right\rangle \\ = (\alpha - 1 - \lambda)(1 - \eta)t_k \langle Q^k, \Delta Q_k \rangle + \underbrace{(\alpha - 1 - \lambda) [\alpha(1 - \eta) \langle Q^k, \Delta z_{k-1} \rangle + (1 - \eta)\alpha_\eta \|Q^k\|^2]}_{(N.3)}.$$

As far as (III) is concerned, we have

$$\begin{aligned}
(III) &= -t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta^2 z_{k-1} \rangle \\
(6.18) \quad &= -t_k(\alpha - 1 - \lambda) \left\langle \Delta z_{k-1}, -\frac{\alpha}{t_k} \Delta z_{k-1} - \Delta Q_k - \frac{\alpha \eta}{t_k} Q^k \right\rangle \\
&= t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta Q_k \rangle + \underbrace{\alpha(\alpha - 1 - \lambda) \|\Delta z_{k-1}\|^2 + \alpha \eta (\alpha - 1 - \lambda) \langle \Delta z_{k-1}, Q^k \rangle}_{(N.4)}.
\end{aligned}$$

Plugging now (II) and (III) into (6.15) we get

$$\begin{aligned}
\Delta E_k^\lambda &= -\lambda(1 - \eta)(\alpha - 2) \langle Q^k, z^k - z^* \rangle - (t_k^2 - \lambda t_k)(1 - \eta) \langle \Delta Q_k, \Delta z_k \rangle \\
&\quad + (Q.1) - t_k \lambda (1 - \eta) \langle Q^k, \Delta Q_k \rangle \\
&\quad + (Q.2) + (N.1) + (N.3) \\
&\quad + (Q.3) + (N.2) + (N.4) \\
&\quad - \frac{1}{2}(1 - \eta)t_k^2 \|\Delta Q_k\|^2 - \underbrace{\frac{\lambda}{2}(\alpha - 1 - \lambda) \|\Delta z_{k-1}\|^2}_{(N.5)} + \Delta \left(\frac{\delta}{2} \|Q\|^2 \right)_k,
\end{aligned}$$

where the term $t_k(\alpha - 1 - \lambda) \langle \Delta z_{k-1}, \Delta Q_k \rangle$ canceled out with the corresponding one in (III), and we simplified the coefficient that multiplies $\langle Q^k, \Delta Q_k \rangle$ using the contribution from (II). Let us also define for brevity: $(N) := (N.1) + (N.2) + (N.3) + (N.4) + (N.5)$. We now make use of (3.9) to infer that

$$(6.19) \quad \langle \Delta Q_k, \Delta z_k \rangle \geq -\frac{1}{2} \|\Delta Q_k\|^2.$$

This allows us to get for k large enough (such that $t_k \geq 2\lambda$)

$$\begin{aligned}
\Delta E_k^\lambda &\leq -\lambda(1 - \eta)(\alpha - 2) \left(\langle Q^k, z^k - z^* \rangle + \frac{1}{2} \|Q^k\|^2 \right) + \frac{\lambda}{2}(1 - \eta)(\alpha - 2) \|Q^k\|^2 \\
(6.20) \quad &\quad - \frac{t_k}{2} \lambda (1 - \eta) \|\Delta Q_k\|^2 + \Delta \left(\frac{\delta}{2} \|Q\|^2 \right)_k \\
&\quad + (Q.1) + (Q.2) + (Q.3) \underbrace{-t_k \lambda (1 - \eta) \langle Q^k, \Delta Q_k \rangle}_{(IV)} + (N).
\end{aligned}$$

Using the polarization identity on (IV) and recalling that $\delta_k = t_{k-1} \lambda (1 - \eta)$, we get:

$$\begin{aligned}
(IV) &= -t_k \lambda (1 - \eta) \langle Q^k, \Delta Q_k \rangle \\
&= -t_k \lambda (1 - \eta) \left(\frac{1}{2} \|Q^{k+1}\|^2 - \frac{1}{2} \|Q^k\|^2 - \frac{1}{2} \|\Delta Q_k\|^2 \right) \\
(6.21) \quad &= -\frac{1}{2} t_k \lambda (1 - \eta) \|Q^{k+1}\|^2 + \frac{1}{2} t_{k-1} \lambda (1 - \eta) \|Q^k\|^2 + \frac{\lambda}{2} (1 - \eta) \|Q^k\|^2 + \frac{t_k}{2} \lambda (1 - \eta) \|\Delta Q_k\|^2 \\
&= -\Delta \left(\frac{\delta}{2} \|Q\|^2 \right)_k + \frac{\lambda}{2} (1 - \eta) \|Q^k\|^2 + \frac{t_k}{2} \lambda (1 - \eta) \|\Delta Q_k\|^2.
\end{aligned}$$

Inserting (IV) and (6.16) into (6.20), simplifying and grouping terms yields (6.22)

$$\begin{aligned}
\Delta E_k^\lambda &\leq -\lambda(1-\eta)(\alpha-2) \left(\langle Q^k, z^k - z^* \rangle + \frac{1}{2} \|Q^k\|^2 \right) \\
&\quad + \left[\frac{\lambda}{2}(1-\eta)(\alpha-2) + \frac{\lambda}{2}(1-\eta) \right] \|Q^k\|^2 \\
&\quad - \left[\eta(1-\eta) \left((\alpha-2)t_k + \frac{1}{2} \right) + \frac{1}{2}(1-\eta)^2(\alpha-1)^2 \right] \|Q^k\|^2 + (N) \\
&\quad + (\alpha-1-\lambda) \frac{t_k}{2} \|Q^k\|^2 - (\alpha-1-\lambda) \frac{t_k}{2} \|\Delta z_{k-1}\|^2 - (\alpha-1-\lambda) \frac{t_k}{2} \|Q^k + \Delta z_{k-1}\|^2 \\
&= -\frac{1-\eta}{2} \left(\eta + (1-\eta)(\alpha-1)^2 - \lambda(\alpha-1) \right) \|Q^k\|^2 - \left(\eta(1-\eta)(\alpha-2) - \frac{\alpha-1-\lambda}{2} \right) t_k \|Q^k\|^2 \\
&\quad - (\alpha-1-\lambda) \frac{t_k}{2} \|\Delta z_{k-1}\|^2 - (\alpha-1-\lambda) \frac{t_k}{2} \|Q^k + \Delta z_{k-1}\|^2 + (N).
\end{aligned}$$

Now, invoking the Young's inequality on all the inner products appearing in (N), we obtain (6.23)

$$\begin{aligned}
(N) &= -(1-\eta)(\alpha-1-\lambda)(\alpha-1) \langle Q^k, \Delta z_{k-1} \rangle - \frac{1}{2}(\alpha-1-\lambda)^2 \|\Delta z_{k-1}\|^2 \\
&\quad + (\alpha-1-\lambda) \left[\alpha(1-\eta) \langle Q^k, \Delta z_{k-1} \rangle + (1-\eta)\alpha_\eta \|Q^k\|^2 \right] \\
&\quad + \alpha(\alpha-1-\lambda) \|\Delta z_{k-1}\|^2 + \alpha_\eta(\alpha-1-\lambda) \langle \Delta z_{k-1}, Q^k \rangle - \frac{\lambda}{2}(\alpha-1-\lambda) \|\Delta z_{k-1}\|^2 \\
&= (\alpha-1-\lambda) \left((1-\eta)\alpha_\eta \|Q^k\|^2 + (1-\eta+\alpha_\eta) \langle Q^k, \Delta z_{k-1} \rangle + \frac{\alpha+1}{2} \|\Delta z_{k-1}\|^2 \right) \\
&\leq (\alpha-1-\lambda) \left((1-\eta)\alpha_\eta + \frac{1-\eta+\alpha_\eta}{2} \right) \|Q^k\|^2 + (\alpha-1-\lambda) \frac{\alpha+2+(1-\eta)(\alpha-1)}{2} \|\Delta z_{k-1}\|^2.
\end{aligned}$$

Inserting (6.23) into (6.22) and rearranging we get the claim. \square

REFERENCES

- [1] S. Adly and H. Attouch. "Complexity analysis based on tuning the viscosity parameter of the Su-Boyd-Candès inertial gradient dynamics". In: *Set-valued and Variational Analysis* (2024).
- [2] F. Alvarez, H. Attouch, J. Bolte, and P. Redont. "A second-order gradient-like dissipative dynamical system with Hessian-driven damping: Application to optimization and mechanics". In: *Journal de Mathématiques Pures et Appliquées* 81.8 (2002), pp. 747–779.
- [3] H. Attouch, R. I. Boş, and D.-K. Nguyen. "Fast convex optimization via time scale and averaging of the steepest descent". In: *Mathematics of Operations Research* (2024).
- [4] H. Attouch, Z. Chbani, J. Fadili, and H. Riahi. "Convergence of iterates for first-order optimization algorithms with inertia and Hessian driven damping". In: *Optimization* 72.5 (2023), pp. 1199–1238.
- [5] H. Attouch, Z. Chbani, J. Fadili, and H. Riahi. "First-order optimization algorithms via inertial systems with Hessian driven damping". In: *Mathematical Programming* 193.1 (2022), pp. 113–155.
- [6] H. Attouch, Z. Chbani, J. Fadili, and H. Riahi. "Rate of convergence of the Nesterov accelerated gradient method in the subcritical case $\alpha \leq 3$ ". In: *ESAIM: Control, Optimisation and Calculus of Variations* 25 (2019), p. 2.
- [7] H. Attouch, Z. Chbani, J. Peypouquet, and P. Redont. "Fast convergence of inertial dynamics and algorithms with asymptotic vanishing viscosity". In: *Mathematical Programming* 168.1 (2018), pp. 123–175.
- [8] H. Attouch, P.-E. Maingé, and P. Redont. "A second-order differential system with Hessian-driven damping; Application to non-elastic shock laws". In: *Differential Equations and Applications* 4 (2012), pp. 27–65.
- [9] H. Attouch, J. Peypouquet, and P. Redont. "Fast convex optimization via inertial dynamics with Hessian driven damping". In: *Journal of Differential Equations* 261.10 (2016), pp. 5734–5783.
- [10] H. H. Bauschke and P. L. Combettes. *Convex analysis and monotone operator theory in Hilbert spaces*. Second. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer, Cham, 2017.

- [11] A. Beck and M. Teboulle. “A Fast Iterative Shrinkage-Thresholding algorithm for linear inverse problems”. In: *SIAM Journal On Imaging Sciences* 2 (2009), pp. 183–202.
- [12] J.-D. Benamou and G. Carlier. “Augmented Lagrangian Methods for Transport Optimization, Mean Field Games and Degenerate Elliptic Equations”. In: *Journal of Optimization Theory and Applications* 167.1 (2015), pp. 1–26.
- [13] J.-D. Benamou, G. Carlier, M. Cuturi, L. Nenna, and G. Peyré. “Iterative Bregman Projections for Regularized Transportation Problems”. In: *SIAM Journal on Scientific Computing* 37.2 (2015), A1111–A1138.
- [14] R. I. Boţ, E. R. Csetnek, and D.-K. Nguyen. “Fast Optimistic Gradient Descent Ascent (OGDA) Method in Continuous and Discrete Time”. In: *Foundations of Computational Mathematics* (2023).
- [15] R. I. Boţ and D.-K. Nguyen. “Fast Krasnoselskii–Mann Algorithm with a Convergence Rate of the Fixed Point Iteration of $o\left(\frac{1}{k}\right)$ ”. In: *SIAM Journal on Numerical Analysis* 61.6 (2023), pp. 2813–2843.
- [16] R. I. Boţ and D.-K. Nguyen. *Tikhonov regularization of monotone operator flows not only ensures strong convergence of the trajectories but also speeds up the vanishing of the residuals*. arXiv:2406.00852. 2024.
- [17] K. Bredies, E. Chenchene, D. A. Lorenz, and E. Naldi. “Degenerate Preconditioned Proximal Point Algorithms”. In: *SIAM Journal on Optimization* 32.3 (2022), pp. 2376–2401.
- [18] K. Bredies, E. Chenchene, and E. Naldi. “Graph and Distributed Extensions of the Douglas–Rachford Method”. In: *SIAM Journal on Optimization* 34.2 (2024), pp. 1569–1594.
- [19] K. Bredies and H. Sun. “A proximal point analysis of the preconditioned alternating direction method of multipliers”. In: *Journal of Optimization Theory and Applications* 173.3 (2017), pp. 878–907.
- [20] K. Bredies and H. Sun. “Preconditioned Douglas–Rachford splitting methods for convex-concave saddle-point problems”. In: *SIAM Journal on Numerical Analysis* 53.1 (2015), pp. 421–444.
- [21] F. E. Browder. “Nonlinear Operators and Nonlinear Equations of Evolution in Banach Spaces”. In: *Proceedings of Symposia in Pure Mathematics*. Vol. 18, part 2. Providence, RI: American Mathematical Society, 1976.
- [22] G. Carlier, E. Chenchene, and K. Eichinger. “Wasserstein Medians: Robustness, PDE Characterization, and Numerics”. In: *SIAM Journal on Mathematical Analysis* 56.5 (2024), pp. 6483–6520.
- [23] A. Chambolle and C. Dossal. “On the Convergence of the Iterates of the “Fast Iterative Shrinkage/Thresholding Algorithm””. In: *Journal of Optimization Theory and Applications* 166.3 (2015), pp. 968–982.
- [24] A. Chambolle and T. Pock. “A First-Order Primal-Dual Algorithm for Convex Problems with Applications to Imaging”. In: *Journal of Mathematical Imaging and Vision* 40.1 (2011), pp. 120–145.
- [25] E. Chenchene. “Splitting methods in nonsmooth convex degenerate optimization”. Dissertation (PhD). Graz: University of Graz, 2023.
- [26] R. Cominetti, J. Peypouquet, and S. Sorin. “Strong asymptotic convergence of evolution equations governed by maximal monotone operators with Tikhonov regularization”. In: *Journal of Differential Equations* 245.12 (2008), pp. 3753–3763.
- [27] J. P. Contreras and R. Cominetti. “Optimal error bounds for non-expansive fixed-point iterations in normed spaces”. In: *Mathematical Programming* 199.1 (2023), pp. 343–374.
- [28] D. Davis and W. Yin. “A three-operator splitting scheme and its optimization applications”. In: *Set-Valued and Variational Analysis. Theory and Applications* 25.4 (2017), pp. 829–858.
- [29] J. Douglas Jr. and H. H. Rachford Jr. “On the numerical solution of heat conduction problems in two and three space variables”. In: *Transactions of the American Mathematical Society* 82 (1956), pp. 421–439.
- [30] M. Fornasier, H. Huang, L. Pareschi, and P. Sünnen. “Anisotropic Diffusion in Consensus-Based Optimization on the Sphere”. In: *SIAM Journal on Optimization* 32.3 (2022), pp. 1984–2012.
- [31] A. Haraux. *Systèmes dynamiques dissipatifs et applications*. Vol. 17. Recherches en Mathématiques Appliquées. Paris: Masson, 1991.
- [32] B. He and X. Yuan. “Convergence Analysis of Primal-Dual Algorithms for a Saddle-Point Problem: From Contraction Perspective”. In: *SIAM Journal on Imaging Sciences* 5 (2012), pp. 119–149.
- [33] F. Lieder. “On the convergence rate of the Halpern-iteration”. In: *Optimization Letters* 15.2 (2021), pp. 405–418.
- [34] P. L. Lions and B. Mercier. “Splitting Algorithms for the Sum of Two Nonlinear Operators”. In: *SIAM Journal on Numerical Analysis* 16.6 (1979), pp. 964–979.
- [35] Y. Malitsky and M. K. Tam. “Resolvent splitting for sums of monotone operators with minimal lifting”. In: *Mathematical Programming* 201.1 (2023), pp. 231–262.
- [36] Y. Nesterov. “A method for solving the convex programming problem with convergence rate $O(1/k^2)$ ”. In: *Proceedings of the USSR Academy of Sciences* 269 (1983), pp. 543–547.

- [37] N. Papadakis, G. Peyré, and E. Oudet. “Optimal transport with proximal splitting”. In: *SIAM Journal on Imaging Sciences* 7.1 (2014), pp. 212–238.
- [38] H. Raguet, J. Fadili, and G. Peyré. “A generalized forward-backward splitting”. In: *SIAM Journal on Imaging Sciences* 6.3 (2013), pp. 1199–1226.
- [39] S. Reich. “Nonlinear evolution equations and nonlinear ergodic theorems”. In: *Nonlinear Analysis: Theory, Methods & Applications* 1.4 (1977), pp. 319–330.
- [40] E. K. Ryu. “Uniqueness of DRS as the 2 operator resolvent-splitting and impossibility of 3 operator resolvent-splitting”. In: *Mathematical Programming* 182.1 (2020), pp. 233–273.
- [41] S. Sabach and S. Shtern. “A First Order Method for Solving Convex Bilevel Optimization Problems”. In: *SIAM Journal on Optimization* 27.2 (2017), pp. 640–660.
- [42] W. Su, S. Boyd, and E. J. Candès. “A Differential Equation for Modeling Nesterov’s Accelerated Gradient Method: Theory and Insights”. In: *Journal of Machine Learning Research* 17.153 (2016), pp. 1–43.
- [43] J. Suh, J. Park, and E. K. Ryu. “Continuous-time Analysis of Anchor Acceleration”. In: *Advances in Neural Information Processing Systems*. Vol. 36. 2023, pp. 32782–32866.
- [44] B. F. Svaiter. “On weak convergence of the Douglas-Rachford method”. In: *SIAM Journal on Control and Optimization* 49.1 (2011), pp. 280–287.
- [45] Q. Tran-Dinh. “From Halpern’s fixed-point iterations to Nesterov’s accelerated interpretations for root-finding problems”. In: *Computational Optimization and Applications* 87 (2024), pp. 181–218.
- [46] T. Yoon, J. Kim, J. J. Suh, and E. K. Ryu. “Optimal Acceleration for Minimax and Fixed-Point Problems is Not Unique”. In: *Proceedings of the 41st International Conference on Machine Learning*. Vol. 235. Proceedings of Machine Learning Research. PMLR, 2024, pp. 57244–57314.

*FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, 1090 VIENNA, AUSTRIA
Email address: radu.bot@univie.ac.at, enis.chenchene@univie.ac.at

†NORMANDIE UNIV, ENSICAEN, CNRS, GREYC, FRANCE.
Email address: Jalal.Fadili@ensicaen.fr