

On the Unknowable Limits to Prediction

Jiani Yan^{1,2,3,*} and Charles Rahal^{1,4,*}

¹Leverhulme Centre for Demographic Science, University of Oxford

²Department of Sociology, University of Oxford

³Wolfson College, University of Oxford

⁴Nuffield College, University of Oxford

*Both authors contributed jointly to this work.

December 2, 2024

Keywords: *Computational Science, Machine Learning, Philosophy of Prediction*

arXiv:2411.19223v1 [cs.LG] 28 Nov 2024

For Correspondence: Jiani Yan and Charles Rahal, Leverhulme Centre for Demographic Science, Demographic Science Unit, University of Oxford, OX1 1JD, United Kingdom. Tel: 01865 286170. Email: jiani.yan@sociology.ox.ac.uk and charles.rahal@demography.ox.ac.uk. **Acknowledgements:** Funding is gratefully acknowledged from the Leverhulme Trust (Grant RC-2018-003) for the Leverhulme Centre for Demographic Science, Nuffield College, and the Economic and Social Research Council. Comments gratefully received from Kyla Chasalow, Mark Verhagen, Ridhi Kashyap, Ben Domingue, Michael Biggs, Sander Wagner, Haohao Lei and Edith Darin. We are especially grateful to comments received from Nicholas Irons.

“We may regard the present state of the universe as the effect of its past and the cause of its future. An intellect which at a certain moment would know all forces that set nature in motion, and all positions of all items of which nature is composed, if this intellect were also vast enough to submit these data to analysis, . . . nothing would be uncertain.”

– Laplace (1814), *Essai philosophique sur les probabilités*

The classic dichotomisation of prediction error into statically defined ‘reducible’ and ‘irreducible’ terms is unable to capture the intricacies of progressive research design where specific types of *truly* reducible error are being rapidly eliminated at differential speeds. This is especially true of prediction in social systems where questions regarding the ‘predictability’ of outcomes are gradually reaching the fore (Rahal et al., 2024). Canonical work which convenes common predictive tasks involving conventional social surveys (Salganik et al., 2020) shows low power, but rapidly emerging computational approaches which utilize non-standard administrative data and information contained within things such as children’s autobiographical essays – often combined with generative Artificial Intelligence and distributed computing (Wolfram et al., 2022; Savcicens et al., 2024) – begin to allow high accuracy. We propose an enhanced framework which builds upon existing work (Hüllermeier and Waegeman, 2021) in a way which we believe helpfully decomposes various types of truly reducible ‘epistemic’ error which are in theory eliminable (resulting from a lack of knowledge), and isolates residual, ‘aleatoric’ error (inherent randomness which can *never* be modeled) that research endeavor will not be able to capture in the limit of human progress. We further emphasize the impossibility of claiming whether things are ‘predictable’, not least in the context of open, dynamically evolving social systems.

Decomposing prediction error into reducible and irreducible terms is not new.¹ However, statements regarding ‘predictability’ and ‘irreducibility’ require the reinforcement of an important quantifier; they are entirely conditional on information sets. For example, the insightful mixed-methods work of Lundberg et al. (2024) begins its abstract with ‘Why are some life outcomes difficult to predict?’. In that case, they are difficult to predict based on information which is currently measurable, and the ‘task’ at hand. An outcome which is difficult to predict with one feature set may not be ‘unpredictable’ in an aleatoric sense: life trajectories may be trivial to predict in the future. More reflexive terminology is essential, lest conclusions be drawn that there is no space for future positive policy interventions based on ever increasingly accurate predictions which prevent negative outcomes. It is impossible to know whether we have eliminated all reducible epistemic error to approach the practical ‘ceiling’ of accuracy, make statements about how predictable outcomes will become, or understand where we are in the process. Predictability of an outcome can only be asserted when all epistemic errors are eliminated: when ‘unmeasured’ information is measured and constructed perfectly, features are constructed perfectly, and when functional forms are able to be exactly recovered.

Consider Equation 1 which better conceptually decomposes error when predicting within contexts under the assumption of stationarity, where y_{true} represents the true outcome, measured after constructing on the phenomenon of interest:

$$\begin{aligned}
 y_{\text{true}} = & \underbrace{f^*(\mathbf{x}_{\text{true}})}_{\text{Predictive Ceiling}} + \underbrace{\varepsilon}_{\text{Aleatoric Error}} \\
 & + \underbrace{[f^*(\mathbf{x}_{\text{true}}) - f(\mathbf{x}_{\text{true}} | y_{\text{true}})]}_{\text{Model approximation gain (Epistemic)}} \\
 & + \underbrace{[f(\mathbf{x}_{\text{true}} | y_{\text{true}}) - f(\mathbf{x}_{\text{true}} | y_{\text{observed}})]}_{\text{Measurement gain from } y \text{ (Epistemic)}} \\
 & + \underbrace{[f(\mathbf{x}_{\text{true}} | y_{\text{observed}}) - f(\mathbf{x}_{\text{observed}} | y_{\text{observed}})]}_{\text{Measurement gain from } \mathbf{x} \text{ (Epistemic)}} \\
 & + \underbrace{y_{\text{predicted}}}_{\text{Current Prediction}} + \underbrace{\varepsilon}_{\text{Irreducible (Aleatoric)}} .
 \end{aligned} \tag{1}$$

where \mathbf{x}_{true} (from a possibly uncountably infinite collection of features in the ‘universe’) and y_{true} are ‘enhanced’

¹See, for example, Sections 2.9 and 7.3 of Hastie et al. (2009); a landmark instruction manual in the field. This foundational text uses the term ‘reducible’ to taxonomically classify errors into a type that can be eliminated, and the term ‘irreducible’ to denote errors which cannot be eliminated *given current information on target variables and feature sets*.




Figure 1: Decomposed Learning Curves: The functional form of learning curves can vary depending on the outcomes and features being predicted, and different amounts of truly irreducible error may exist for different outcomes (see Supplementary Information Figure S1). Panel ‘a.’ represents a baseline scenario which positions specific studies against how well information on target and features has been measured (and how well such information has been mapped into learning algorithms), while Panels ‘b.’ and ‘c.’ represent changes (and gains and changes in predictive performance) with regards to construct validity of y and x respectively. Note: ‘obs.’ denotes ‘observed’ variables as per Equation 1.

information sets over and above $x_{observed}$ and $y_{observed}$ which could tautologically reduce what may currently have been termed ‘irreducible’.² Predictive accuracy could also improve through better construct validity (O’Leary-Kelly and Vokurka, 1998); the extent to which the model’s predictions or features accurately capture the theoretical construct or concept the model is intended to measure or represent.³ Incorporation of previously unmeasured information (from $x_{observed}$ to x_{true}) or better measurement of existing features (from $x_{observed}$ and $y_{observed}$ to x_{true} and y_{true}) will also reduce epistemic error, as will reductions in model approximation error (from f to f^*).⁴ To illustrate how these different forms of error reduction work in practice, we visualise our generalized framework in Figure 1. This demonstrates how learning curves evolve as different sources of epistemic uncertainty are reduced (Figure 1a.) and targets and features are constructed differently (Figure 1b.-c.).⁵

However, the existence of aleatoric uncertainty means we cannot appreciate a practical limit to prediction, because that itself cannot be measured. We have no reasonable expectation of what data will be collectable, or how powerful deep learning algorithms may become. It is only recently that the concept of measuring epigenetic data – let alone interpreting it at scale – has become viable due to advances in molecular biology. Aleatoric error may be large and nontrivial, with suggestions that predictive performance may be bounded

²For class labeling tasks (such as binary classification), our conceptual framework remains viable at the aggregate level with appropriately defined model evaluation metrics: we leave this – as well as extensions to things such as probability distributions, entropy, and stochastic processes adapted to filtration – for further work.

³Note: better construction of the target and features may not necessarily involve the collection of additional information, but simply a re-framing of a variable or new utilization of existing information, as depicted in Figures 1b.-c.

⁴Kindly see our Supplementary Information for a fully delineated decomposition of Equation 1.

⁵Supplementary Information Figure S1a. shows the case where aleatoric error may be low and the speed of improvement relative to information gain is high with large gains from measurement error possible. Supplementary Information Figure S1b. shows the case where aleatoric error may be high, but learning is relatively slower and only minimal gains from reducing learning error are possible.

well below deterministic precision. Some data may be potentially impossible to collect, and some outcomes may be genuinely stochastic. The ‘time to event’ will also affect how large aleatoric error is; it is easier to predict how somebody will vote at the next election than how the same person will vote in three election’s time. Similarly, re-framing the outcome from ‘when will somebody have a child’ to ‘whether somebody will have a child’ will also affect the size of aleatoric error. We may be able to better approximate ‘perfect prediction’ in some easily constructed and perfectly measurable outcomes (e.g., height) than others (e.g., behavioral traits). Research is required to perturb information sets across individual prediction exercises (i.e., comparing better measurement and construct through inclusion/omission), illuminating how quickly each type of epistemic uncertainty is reduced. Work on systematically indexing predictive accuracy is also of utility, as are designs which seek to maximize external validity. Emerging qualitative approaches which explore predictive errors as a function of pivotal moment’s in people’s lives (Lundberg et al., 2024) are also immensely promising, and represent a more nuanced form of information. Collecting more detailed information also necessitates careful discussions around privacy, compliance, and ethical issues. However, until we have reason to believe that we have eliminated error through better measurement and construction of features and outputs – in addition to the elimination of learning error – we should only tentatively discuss how predictable things are with current data and technology, and not yet uniformly infer a ‘failure to predict’ or an ‘unpredictability of life outcomes’ (Garip, 2020, p. 8235).

References

- Garip, F. (2020). *Proceedings of the National Academy of Sciences* 117(15), 8234–8235.
- Hastie, T., R. Tibshirani, J. H. Friedman, and J. H. Friedman (2009). Volume 2. Springer.
- Hüllermeier, E. and W. Waegeman (2021). *Machine learning* 110(3), 457–506.
- Laplace, P.-S. (1814). *Essai philosophique sur les probabilités*. Paris: Courcier.
- Lundberg, I., R. Brown-Weinstock, S. Clampet-Lundquist, S. Pachman, T. J. Nelson, V. Yang, K. Edin, and M. J. Salganik (2024). *Proceedings of the National Academy of Sciences* 121(24), e2322973121.
- O’Leary-Kelly, S. W. and R. J. Vokurka (1998). *Journal of operations management* 16(4), 387–405.
- Rahal, C., M. Verhagen, and D. Kirk (2024). *AI & SOCIETY* 39(2), 799–801.
- Salganik, M. J., I. Lundberg, A. T. Kindel, C. E. Ahearn, K. Al-Ghoneim, A. Almaatouq, D. M. Altschul, J. E. Brand, N. B. Carnegie, R. J. Compton, et al. (2020). *Proceedings of the National Academy of Sciences* 117(15), 8398–8403.
- Savcisen, G., T. Eliassi-Rad, L. K. Hansen, L. H. Mortensen, L. Lilleholt, A. Rogers, I. Zettler, and S. Lehmann (2024). *Nature Computational Science* 4(1), 43–56.
- Wolfram, T., F. C. Tropf, and C. Rahal (2022), *SocArXiv*, 10.31235/osf.io/a8ht9.

Supplementary Information

SI.1 Delineating Epistemic and Aleatoric Errors

In predictive modeling, researchers often aim to predict a phenomenon or concept of interest; they aim to construct a concrete and measurable representation of this as y_{true} . However, due to measurement errors that arise from factors such as subjective assessment, accounting inaccuracies, or sampling bias, researchers are often forced to conceptualize this phenomenon which they want to measure as something similar to y_{observed} . Such measurement errors exist in both feature and target variables; it is essential to understand how these errors propagate through the model and affect predictive accuracy. This supplementary section presents a mathematical framework to link y_{true} with y_{observed} , highlighting epistemic (reducible) and aleatoric (irreducible) errors in prediction, and then equates it to the familiar bias-variance trade-off.

SI.1.1 Problem Setup

Consider the following definition of the ‘true’ target and features, the observable target and features, and the model which binds them as follows.

1. True Target and Features

Allow us to equate the relationship between the ‘true’ target variable and feature set as:

$$y_{\text{true}} := f^*(\mathbf{x}_{\text{true}}) + \varepsilon, \quad (\text{S2})$$

where:

- y_{true} is the true outcome, measured *after* constructing on the phenomenon of interest.
- \mathbf{x}_{true} represents the best possible feature set both in terms of quantity and in quality; a perfectly chosen, constructed and measured feature set.
- $f^*(\cdot)$ is the conceptually true underlying function which maps \mathbf{x}_{true} onto y_{true} .
- ε is aleatoric error (inherent randomness) for y_{true} , with $\mathbb{E}[\varepsilon] = 0$ and variance $\text{Var}(\varepsilon) = \sigma_\varepsilon^2$. It is uncorrelated with $f^*(\mathbf{x}_{\text{true}})$. A proof of this is available upon request.

2. Observed Target and Features

Next, allow us to introduce measurement error in y_{true} and \mathbf{x}_{true} by relating them to what we can observe:

$$y_{\text{observed}} := y_{\text{true}} + \delta_y = f^*(\mathbf{x}_{\text{true}}) + \varepsilon + \delta_y, \quad (\text{S3})$$

$$\mathbf{x}_{\text{observed}} := \mathbf{x}_{\text{true}} + \delta_{\mathbf{x}}, \quad (\text{S4})$$

where:

- δ_y is measurement error in y , with $\mathbb{E}[\delta_y] = \mu_{\delta_y}$ and $\text{Var}(\varepsilon) = \sigma_{\delta_y}^2$.
- $\delta_{\mathbf{x}}$ is measurement error in \mathbf{x} , with $\mathbb{E}[\delta_{\mathbf{x}}] = \mu_{\delta_{\mathbf{x}}}$ and variance $\text{Var}(\varepsilon) = \sigma_{\delta_{\mathbf{x}}}^2$. $\mathbf{x}_{\text{observed}}$ is a vector with $1 \times K$ dimensions, with each $x_k \in \mathbf{X}$ for $k=\{1, 2, \dots, K\}$ as each one of the predictors. Note: $\mathbf{x}_{\text{observed}}$ can have different dimensions to \mathbf{x}_{true} .

As y_{observed} is observed based on a constructed concept, δ_y is a function of how well y_{true} is measured. $\delta_{\mathbf{x}}$ constitutes measurement and construct error both at the single predictor and aggregate levels. At the level of an individual feature (\mathbf{x}_k), it regards how well the feature is constructed and measured, and at the aggregate level, how well the feature set itself is measured (as a function of individual features \mathbf{x}_k and the set \mathbf{X} as a whole). Construct validity of $x_k \in \mathbf{X}$ can be achieved with the same amount of underlying information (see Figure 1c). For instance, if x_k is calculated by averaging the response of multiple items, construct validity can be improved by finding optimal weights by which to combine the individual predictors. $\delta_{\mathbf{x}}$ can be reduced through improved construction of each $x_k \in \mathbf{X}$; increased sample representativeness, and conceptualizing the feature set so that it contains the most relevant set of predictors. In general, we have assumed that y_{observed} captures all aleatoric errors in y_{true} ; changing δ_y does not impact the level of aleatoric error ε . A discussion of the violation of this condition can be found in [SI.1.5](#)

3. Predictive Model

The predictive model is estimated using observable training data $\mathbf{x}_{\text{observed}}$ and y_{observed} :

$$y_{\text{pred}} := f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}}), \quad (\text{S5})$$

where f is the estimated model trained on $\{\mathbf{x}_{\text{observed}}, y_{\text{observed}}\}$. The notation $f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}})$ emphasizes that f depends on y_{observed} in training processes. The difference between the conceptually true underlying function f^* and the currently selected predictive function $f()$ on \mathbf{x}_{true} and y_{true} is defined as follows:

$$f(\mathbf{x}_{\text{true}} \mid y_{\text{true}}) := f^*(\mathbf{x}_{\text{true}}) + \delta_f, \quad (\text{S6})$$

where δ_f is defined based on the optimal \mathbf{x}_{true} and y_{true} , and $\mathbb{E}[\delta_f] = \mu_f$, $\text{Var}[\delta_f] = \sigma_{\delta_f}$. The difference between $f(\mathbf{x}_{\text{true}} \mid y_{\text{true}})$ and y_{pred} attributes to improvement in δ_x and δ_y ; further details can be found in Equation S7.

SI.1.2 Decomposing y_{true}

Following these aforementioned definitions, we can decompose the output of a predictive machine which predicts y_{true} as perfectly as possible, and in the process link it to existent predictions generated by $f()$ to create y_{pred} as follows:

$$\begin{aligned} y_{\text{true}} &= f^*(\mathbf{x}_{\text{true}}) + \varepsilon \\ &= [f^*(\mathbf{x}_{\text{true}}) - f(\mathbf{x}_{\text{true}} \mid y_{\text{true}})] \\ &\quad + [f(\mathbf{x}_{\text{true}} \mid y_{\text{true}}) - f(\mathbf{x}_{\text{true}} \mid y_{\text{observed}})] \\ &\quad + [f(\mathbf{x}_{\text{true}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}})] \\ &\quad + f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}}) + \varepsilon \\ &= \underbrace{-\delta_f}_{\substack{\text{Model approximation gain} \\ \text{(Epistemic)}}} \\ &\quad + \underbrace{[f(\mathbf{x}_{\text{true}} \mid y_{\text{true}}) - f(\mathbf{x}_{\text{true}} \mid y_{\text{true}} + \delta_y)]}_{\substack{\text{Measurement gain from y} \\ \text{(Epistemic)}}} \\ &\quad + \underbrace{[f(\mathbf{x}_{\text{true}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{true}} + \delta_x \mid y_{\text{observed}})]}_{\substack{\text{Measurement gain from X} \\ \text{(Epistemic)}}} \\ &\quad + \underbrace{y_{\text{predicted}}}_{\text{Current Prediction}} + \underbrace{\varepsilon}_{\substack{\text{Irreducible} \\ \text{(Aleatoric)}}}. \end{aligned} \quad (\text{S7})$$

Therefore, we can achieve improved predictive performance in comparison to existing predictions based on $\mathbf{x}_{\text{observed}}$, y_{observed} , and $f()$. This is made possible through opportunities to eliminate measurement and construction error in $\mathbf{x}_{\text{observed}}$ (i.e., reducing δ_x), measurement in y_{observed} (i.e., reducing δ_y), and better model approximation $f()$ (i.e., reducing δ_f). When $\delta_x = \delta_y = \delta_f = 0$, we reach a scenario where the difference between y_{true} and $y_{\text{predicted}}$ is ε : pure aleatoric error. This essentially represents the ‘upper bound’ of predictive accuracy, occasionally referred to as a ‘predictive ceiling’ (Garip, 2020).

SI.1.3 Error Decomposition

For any individual observation, the total prediction error (denoted below as *Error*) is defined as the difference between the predicted and true outcome:

$$\begin{aligned}
Error &:= y_{\text{pred}} - y_{\text{true}} \\
&= f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}}) - y_{\text{true}} \\
&= [f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{true}} \mid y_{\text{observed}})] \\
&\quad + [f(\mathbf{x}_{\text{true}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{true}} \mid y_{\text{true}})] \\
&\quad + [f(\mathbf{x}_{\text{true}} \mid y_{\text{true}}) - f^*(\mathbf{x}_{\text{true}})] + [f^*(\mathbf{x}_{\text{true}}) - y_{\text{true}}].
\end{aligned} \tag{S8}$$

Substituting for Equations S2-S4, we obtain:

$$\begin{aligned}
Error &= \underbrace{[f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{observed}} - \delta_{\mathbf{x}} \mid y_{\text{observed}})]}_{\substack{\text{Error due to measurement error in } X \\ \text{(Epistemic)}}} \\
&\quad + \underbrace{[f(\mathbf{x}_{\text{true}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{true}} \mid (y_{\text{observed}} - \delta_y))]}_{\substack{\text{Error due to measurement error in } y \text{ affecting model estimation} \\ \text{(Epistemic)}}} \\
&\quad + \underbrace{\delta_f}_{\substack{\text{Model approximation error} \\ \text{(Epistemic)}}} + \underbrace{\varepsilon}_{\substack{\text{Irreducible} \\ \text{(Aleatoric)}}}.
\end{aligned} \tag{S9}$$

It then logically follows that predicting the target variable as accurately as can be done in practice (see Section SI.1.1) results in the case where all epistemic errors are eliminated, and only aleatoric error remains: $Error = y_{\text{pred}} - y_{\text{true}} = \varepsilon$.

SI.1.4 Relationship to Existing Frameworks

Our framework mainly decomposes errors in a conceptually simple way (i.e., up until now, we have been dealing with scalar target outcome observations). We next develop the aggregate level decomposition of the expected sum of squared errors and relate our framework to the bias, variance, and error frameworks of canonical textbooks (e.g., James et al., 2013; Hastie et al., 2009) as well as papers (Pedro, 2000; James, 2003) in the field of statistical learning. Suppose we have a sample with N observations, for each $y_{\text{true},i} \in \mathbf{y}_{\text{true}}$ and $y_{\text{pred},i} \in \mathbf{y}_{\text{predicted}}$ for $i = \{1, 2, \dots, N\}$. Equation S9 holds as follows:

$$\begin{aligned}
Error &= y_{\text{pred}} - y_{\text{true}} \\
&= \underbrace{[f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{observed}} - \delta_{\mathbf{x}} \mid y_{\text{observed}})]}_{\substack{\text{Error due to measurement error in } X \\ \text{(Epistemic)}}} \\
&\quad + \underbrace{[f(\mathbf{x}_{\text{true}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{true}} \mid (y_{\text{observed}} - \delta_y))]}_{\substack{\text{Error due to measurement error in } y \text{ affecting model estimation} \\ \text{(Epistemic)}}} \\
&\quad + \underbrace{\delta_f}_{\substack{\text{Model approximation error} \\ \text{(Epistemic)}}} + \underbrace{\varepsilon}_{\substack{\text{Irreducible} \\ \text{(Aleatoric)}}} \\
&= \delta_{\mathbf{x}}^{\text{error}} + \delta_y^{\text{error}} + \delta_f + \varepsilon,
\end{aligned} \tag{S10}$$

where:

- $\delta_{\mathbf{x}}^{\text{error}} := f(\mathbf{x}_{\text{observed}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{observed}} - \delta_{\mathbf{x}} \mid y_{\text{observed}})$; the difference in predicted values owing to $\delta_{\mathbf{x}}$. $\mathbb{E}[\delta_{\mathbf{x}}^{\text{error}}] = \mu_{\mathbf{x}}^{\text{error}}$ and variance $\text{Var}(\delta_{\mathbf{x}}^{\text{error}}) = \sigma_{\delta_{\mathbf{x}}^{\text{error}}}^2$. As $\delta_{\mathbf{x}}^{\text{error}}$ is expressed as the difference between the fitted values, the calculation of $\text{Var}(\delta_{\mathbf{x}}^{\text{error}})$ depends on the specific form of f , the distribution of $\mathbf{x}_{\text{observed}}$ and $\mathbf{x}_{\text{observed}} - \delta_{\mathbf{x}}$, and their dependencies.
- $\delta_y^{\text{error}} := f(\mathbf{x}_{\text{true}} \mid y_{\text{observed}}) - f(\mathbf{x}_{\text{true}} \mid (y_{\text{observed}} - \delta_y))$; the difference in predicted values owing to δ_y . $\mathbb{E}[\delta_y^{\text{error}}] = \mu_y^{\text{error}}$ and variance $\text{Var}(\delta_y^{\text{error}}) = \sigma_{\delta_y^{\text{error}}}^2$. Similarly, the calculation of $\text{Var}(\delta_y^{\text{error}})$ depends on the specific form of f , the distribution of y_{observed} and $y_{\text{observed}} - \delta_y$, and their dependencies.

- δ_f is defined as before in Equation S5.

Therefore,

$$\begin{aligned}\mathbb{E}[Error] &= \mathbb{E}[\delta_{\mathbf{x}}^{\text{error}} + \delta_y^{\text{error}} + \delta_f + \varepsilon] \\ &= \mu_{\mathbf{x}}^{\text{error}} + \mu_y^{\text{error}} + \mu_{\delta_f}.\end{aligned}\tag{S11}$$

Since ε is uncorrelated with other sources of error:

$$\begin{aligned}\text{Var}(Error) &= \text{Var}[(\delta_{\mathbf{x}}^{\text{error}} + \delta_y^{\text{error}} + \delta_f + \varepsilon)] \\ &= \text{Var}[(\delta_{\mathbf{x}}^{\text{error}} + \delta_y^{\text{error}} + \delta_f)] + \text{Var}(\varepsilon).\end{aligned}\tag{S12}$$

With Equations S11-S12 we can write our squared error loss as:

$$\begin{aligned}\mathbb{E}[(y_{\text{true}} - y_{\text{pred}})^2] &= \mathbb{E}[(y_{\text{true}} - y_{\text{pred}})]^2 + \text{Var}[(y_{\text{true}} - y_{\text{pred}})] \\ &= \mathbb{E}[Error]^2 + \text{Var}(Error) \\ &= \underbrace{(\mu_{\mathbf{x}}^{\text{error}} + \mu_y^{\text{error}} + \mu_{\delta_f})^2}_{\text{Bias}} + \underbrace{\text{Var}[(\delta_{\mathbf{x}}^{\text{error}} + \delta_y^{\text{error}} + \delta_f)]}_{\text{Variance}} + \underbrace{\text{Var}(\varepsilon)}_{\text{Aleatoric Error}},\end{aligned}\tag{S13}$$

which results in the familiar Irreducible Error + Bias² + Variance framework akin to Equation 7.9 of [Hastie et al. \(2009\)](#). For simplicity, we don't expand $\text{Var}[(\delta_{\mathbf{x}}^{\text{error}} + \delta_y^{\text{error}} + \delta_f)]$, of which the full expansion contains the covariance terms between each of the δ terms.

SI.1.5 Partially captured aleatoric error in observed data

In Equations S3-S4, we assume that y_{observed} captures all aleatoric error in y_{true} . However, since y_{observed} is measured after constructing y_{true} , this assumption may not hold in all situations. For example, observed data may be a biased sub-sample of the whole population. With such observed data, we can only optimize epistemic errors ($\delta_f, \delta_{\mathbf{x}}$ and δ_y) that are inherent in training data. In this case – even when all epistemic error is reduced – $\varepsilon_{y_{\text{observed}}}$ cannot represent the true ε_y . Consider a true target variable which is measurable across a population (y_{true}), but y_{observed} only covers a non-representative 70% of the population. In this case, $\varepsilon_{y_{\text{observed}}}$ only captures the ε_y of the observed sample, and lacks the ability to capture the component which would be inherent in the remaining 30% of the population. The representativeness of the error term $\varepsilon_{y_{\text{observed}}}$ depends on how well y_{observed} represents true reality (y_{true}). It also affects assumptions we need to make in terms of the bias-variance decomposition as per Equations S12-S13. If $\varepsilon_{y_{\text{observed}}} \neq \varepsilon_y$, the covariance between $\varepsilon_{y_{\text{observed}}}$ and $\delta_f, \delta_y^{\text{error}}$ and $\delta_{\mathbf{x}}^{\text{error}}$ are no longer zero. The $\text{Var}(Error)$ term must also incorporate $\text{Cov}(\varepsilon_{y_{\text{observed}}}, \delta_f), \text{Cov}(\varepsilon_{y_{\text{observed}}}, \delta_y^{\text{error}}), \text{Cov}(\varepsilon_{y_{\text{observed}}}, \delta_{\mathbf{x}}^{\text{error}})$. Therefore, guaranteeing the representativeness of y_{observed} to y_{true} (which also affects the representativeness of $\mathbf{x}_{\text{observed}}$ to \mathbf{x}_{true} from a measurement perspective) is a necessary step before considering epistemic errors in the form of δ_f, δ_y , and $\delta_{\mathbf{x}}$, as it interrogates whether the questions we are answering are really what we meant to ask.

SII Supplementary Figures




Figure S1: Decomposed Learning Curves: Two complimentary example. Panel 'a.' represents a target variable with relatively little aleatoric error, and a rapid learning rate with large gains from eliminating measurement error. Panel 'b.' represents a target variable with relatively higher aleatoric error, a slower learning rate, and relatively fewer gains from improving measurement. Note: there is no requirement for such learning curves to be continuous functions. Dashed horizontal red lines denote what is commonly known as the 'predictive ceiling' (no prediction error). Golden arrows denote the rate at which reducible error is reducing. Dashed blue lines represents a less accurate learning algorithm, and the solid blue line represents the case where learning error is entirely eliminated.

Supplementary References

- Garip, F. (2020). What failure to predict life outcomes can teach us. *Proceedings of the National Academy of Sciences* 117(15), 8234–8235.
- Hastie, T., R. Tibshirani, J. H. Friedman, and J. H. Friedman (2009). The elements of statistical learning Volume 2. Springer.
- James, G. (2003). Variance and bias for general loss functions. *Machine learning* 51, 115–135.
- James, G., D. Witten, T. Hastie, R. Tibshirani, et al. (2013). An introduction to statistical learning Volume 112. Springer.
- Pedro, D. (2000). A unified bias-variance decomposition and its applications. In *17th International conference on machine learning*, pp. 231–238.