Better Approximation for Weighted k-Matroid Intersection

Neta Singer * Theophile Thiery *[†]

Abstract

We consider the problem of finding an independent set of maximum weight simultaneously contained in k matroids over a common ground set. This k-matroid intersection problem appears naturally in many contexts, for example in generalizing graph and hypergraph matching problems. In this paper, we provide a $(k + 1)/(2 \ln 2)$ -approximation algorithm for the weighted k-matroid intersection problem. This is the first improvement over the longstanding (k - 1)-guarantee of Lee, Sviridenko and Vondrák (2009). Along the way, we also give the first improvement over GREEDY for the more general weighted matroid k-parity problem.

Our key innovation lies in a randomized reduction in which we solve almost unweighted instances iteratively. This perspective allows us to use insights from the unweighted problem for which Lee, Sviridenko, and Vondrák have designed a k/2-approximation algorithm. We analyze this procedure by constructing refined matroid exchanges and leveraging randomness to avoid bad local minima.

^{*}Ecole Polytechnique Fédérale de Lausanne (neta.singer@epfl.ch, theophile.thiery@epfl.ch)

[†]Supported by the Swiss State Secretariat for Education, Research and Innovation (SERI) under contract number MB22.00054

Contents

1	Introduction 1			
	1.1 Our local search algorithm	2		
	1.2 Construction of matroid exchanges	3		
	1.3 Computing the approximation ratio	3		
2	Preliminaries	4		
3	Sliding Local-Search Algorithm	5		
	3.1 INTERVAL LOCAL-SEARCH	5		
	3.2 Sliding Local-Search	6		
4	A better-than-k approximation guarantee	6		
	4.1 Approach and Definitions	7		
	4.2 Proof of Lemma 2	8		
	4.3 Escaping local-optima using randomness	10		
	4.4 Obtaining an $\Omega(k)$ approximation ratio	11		
	4.5 Conclusion and Open Questions	11		
Α	Omitted Proofs	15		
	A.1 Proof of Lemma 1	15		
	A.2 Proof of Lemma 3	15		
	A.3 Proof of Theorem 5 and Theorem 1	16		
В	Improved $(k+1)/(2\ln 2)$ approximation ratio	17		
	B.1 Improved Approximation Ratio	18		
С	Proof of Matroid Properties	20		
	C.1 Proof of Theorem 2	20		
	C.2 Proof of Theorem 3	21		
D	Runtime and probability guarantee of Algorithm 2	21		
	D.1 Polynomial runtime	21		
	D.2 Last interval has negligible weight – Proof of Proposition 1	22		
	D.3 Probability of Algorithm 2 succeeding	22		
Е	Reduction and Simplifying Assumptions	23		
	E.1 Reduction from k-Matroid Intersection to Matroid k-Parity	23		
	E.2 Disjointness Assumption	23		

1 Introduction

k-Matroid Intersection is a fundamental question in combinatorial optimization where the goal is to find a maximum weight independent set in the intersection of k matroids defined on a common ground set. This problem generalizes many classical optimization tasks, including finding maximum matchings in bipartite graphs and k-partite hypergraphs. Beyond its versatile formulation, it provides a rich mathematical environment capturing the success of popular approximation algorithms heuristics. For k = 1, the problem is solvable in polynomial time using GREEDY. Moreover, any independence system (or downward closed family of sets) for which GREEDY is optimal is a matroid [Rad57, Gal68]. For k = 2, Edmonds' celebrated polynomial-time algorithm for matroid intersection, generalizing the Hungarian Method, finds the largest independent set in the intersection of two matroids [Edm03].

As soon as the intersection is taken over 3 or more matroids, no algorithm is known to have optimal approximation guarantee. For $k \ge 3$, k-Matroid Intersection was recently shown to be APX-hard to approximate within a factor k/12 unless **NP** \subseteq **BPP** [LST24], improving on the bound of [HSS06]. Currently, the best approximation ratio for the *unweighted* problem stands at k/2 [LSV10]. Notably, this result does not extend to settings with arbitrary weights. In fact, for weighted k-Matroid Intersection Lee, Sviridenko and Vondrák obtained a tight k – 1-approximation algorithm, which remains the best known to date [LSV09].

Since Lee, Sviridenko and Vondrák's result, most algorithmic research has been devoted to the study of special cases of weighted *k*-Matroid Intersection. A notable case is *k*-Set Packing (technically *k*-Dimensional Matching), which consists of finding a maximum weight matching in a *k*-partite hypergraph. After a series of work [AH98, CH99, Ber00, BK03, Neu21, Neu22, TW23, Neu23], the state-of-the-art approximation algorithm for the weighted problem is equal to $\frac{k}{2.006}$ [Neu23]. Despite great algorithmic progress, known algorithms for *k*-Set Packing don't appear to extend naturally to the more general *k*-Matroid Intersection. The simplest method accomplishing a (k+1)/2 approximation factor is due to Berman [Ber00]. This method performs a local-search procedure on an instance with squared-weights and is used and refined in [BK03, Neu21, Neu22, TW23, Neu23]. However, their analyses rely on properties that are only true for the intersection of partition matroids. This situation highlights our limited understanding of *k*-Matroid Intersection beyond *k*-Set Packing and the need for a clean and universal algorithm for the more general problem. Our main result is a novel algorithm for weighted *k*-Matroid Intersection, which is the first to improve upon the k - 1 approximation algorithm of [LSV09].

Theorem 1. There is $\frac{k+1}{2 \ln(2)}$ -approximation algorithm for weighted k-Matroid Intersection.

Our algorithm is the first approximation algorithm asymptotically improving over GREEDY for weighted k-Matroid Intersection. This improvement over GREEDY is substantial since the ratio $\frac{1}{2\ln(2)} \simeq 0.722$ improves the leading constant to more than halfway between GREEDY and the state-of-the-art ratio of k/2 for the unweighted setting. Our algorithm applies the standard local-search algorithm in an iterative fashion and thus also provides an alternative method to the w^2 -algorithm of Berman [Ber00] for weighted k-Set Packing. We, in fact, design a $\frac{k+1}{2\ln(2)}$ -approximation algorithm for a more general

problem: weighted Matroid k-Parity. In this problem, we are given a collection of weighted hyperedges each incident to k vertices, and we aim to find the largest weight hypergraph matching whose groundset is independent in a matroid over the vertices of the hypergraph. k-Matroid Intersection is a special case of this problem. Matroid (2-)Parity was proposed by Lawler [Law01] to generalize non-bipartite matching and 2-Matroid Intersection. It is solvable in polynomial time for *linear* matroids as shown by Lovász in the unweighted setting [Lov80] and by Iwata and Kobayashi with general weights [IK17]. With an arbitrary matroid, the problem requires exponentially many calls to the independence oracle [JK82]. This problem has a wide variety of applications across different areas. A non-exhaustive list of applications of this problem include Path Packing, Minimum Pinning Set, Maximum Genus Embedding, and Maximum Planar Subgraph which have connection to approximation algorithms, combinatorial rigidity and electric ciruits. We refer the reader to [CLL14] for an in-depth literature review of the problem. For *unweighted* Matroid k-Parity with $k \ge 3$, Lee, Vondrák and Sviridenko [LSV10] designed a $\frac{k}{2}$ approximation algorithm. Meanwhile, GREEDY is still the best known algorithm for handling the problem with general hyperedge weights, giving a k-approximation factor. Our contribution is thus to also give the first improvement over GREEDY for weighted Matroid k-Parity. This answers the question by Lee, Vondrák and Sviridenko [LSV10] who asked about the existence of improved local search algorithms for the weighted version of Matroid k-Parity.

1.1 Our local search algorithm

The vast majority of approximation algorithms for k-Matroid Intersection and k-Set Packing, including ours, use a local-search heuristic as a subroutine. Local-search procedures repeatedly improve a solution A by adding some small number of elements $S \subseteq V \setminus A$ and removing *conflicting* elements $T \subseteq A$, while preserving the independence of the solution. The pair (S,T) is called a *swap*. The algorithm terminates when there is no *improving swap*. The absence of improving swaps then allows to compare the value of the algorithm's output A with the value of an optimal solution. The approximation ratio is dictated by the size of the local search swaps. For example, if A is obtained using GREEDY then there is no improving swap of size 1, and implies that GREEDY is a k-approximation algorithm. Prior improvements over the factor k were obtained by looking at larger swaps as in [LSV10, LSV09] for k-Matroid Intersection and [HS89, Cyg13, Neu21, TW23] for k-Set Packing.

However, for weighted *k*-Matroid Intersection the standard local-search algorithm fails even with large swaps and yields a *tight* k - 1-approximation in the worst-case [AH98, LSV09]. Our algorithm instead attempts to apply unweighted algorithms that are known to perform well in a black-box manner. It partitions the instance into several weight classes, where each class is an almost unweighted instance. It then performs local-search on each weight classes, we are able to leverage Lee, Sviridenko and Vondrák's k/2-approximation algorithm for the unweighted problem [LSV10] and avoid the k - 1 local-optima. Crucially, this iterative procedure allows us to construct non-improving swaps based on the currently processed weight class. This approach differs from known algorithms which apply various local search methods over the entire instance directly. Beyond our result, our algorithm is a blueprint for solving a weighted maximization problem by decomposing it in many unweighted instances. Using our framework, it seems very likely that an improved local-search algorithms

rithm for unweighted *k*-Matroid Intersection will yield an improved approximation for the weighted problem surpassing our guarantees. Note that for unweighted *k*-Set Packing, there is a (k + 1)/3-approximation algorithm [Cyg13]. We hope that our work will resonate with other problems where weighted variants are challenging.

1.2 Construction of matroid exchanges

The bulk of our work consists of defining a collection of non-improving swaps between our current solution A and the optimal solution O. By processing the instance greedily over decreasing weight classes, we are able to understand the algorithm as follows. After processing the i^{th} weight class, any element of O in the i^{th} weight class conflicts with the items that were already added to A. The optimal elements in the i^{th} weight class are necessarily conflicting as otherwise we could have added them to A to improve the value of our solution. Thus, each optimal element is either conflicting before or while processing the i^{th} weight class. We can therefore split the optimal elements contained in the i^{th} interval into two sets: (1) the elements that conflict within their own weight class, and (2) the elements that conflict with our choices from the previous i - 1 weight classes.

In matroid terminology, each iteration i defines a large swap between the vertices added to the current solution and conflicting vertices of O. Using Rota's exchange property, we refine this swap by constructing two disjoint swaps of smaller size between A and just the vertices from O in the i^{th} weight-class, and A and just the vertices from O outside the i^{th} weight-class. The existence of these two swaps allows us in the first case to apply Lee, Sviridenko and Vondrák [LSV10] analysis and recover their guarantees. This involves another refinement of the large constructed swap that we break this into many small swaps. We point out that our analysis is conceptually different than that for k-Set Packing. We start from the existence of a large matroid exchange that we gradually refine to obtain small swaps that our algorithm can find. This top-to-bottom approach diverges from other local-search analyses which typically have a bottom-to-top approach. In fact, a fundamental step in the analysis of Berman's algorithm [Ber00] lies in the ability to combine two swaps (S_1, T_1) and (S_2, T_2) to obtain a larger swap $(S_1 \cup S_2, T_1 \cup T_2)$. Berman's analysis [Ber00] heavily relies on this fact to group optimal elements by their *individual* conflicts with edges in A to get a larger swap, and the same is used for [CH99]. Unfortunately, for general matroids the union of two swaps is not necessarily a feasible swap. One such example is the graphic matroid on the complete graph with 4 vertices, K_4 . The techniques used in [LSV10] are also not directly applicable, as their algorithm can informally be understood as solving 2-Matroid Intersection and greedily extending the solution. This essentially reduces the number of matroid dependencies by 1 to obtain a k - 1-factor.

1.3 Computing the approximation ratio

The final approximation ratio depends on the following two factors: (1) the fraction of the optimal elements conflicting only within their weight class, and (2) the fraction of optimal elements which conflict with a prior but adjacent weight class. As explained, the contribution of (1) can essentially be reduced to the unweighted analysis of Lee, Sviridenko and Vondrák [LSV10]. Given that the weights within a weight class differ at most by a factor of $(1 - \varepsilon)$ for some $\varepsilon > 0$, we obtain a $\frac{k}{2(1-\varepsilon)}$ -approximation of the optimal elements conflicting only within their weight class. The difficulty then

lies in understanding the effect of optimal elements that conflict with a previous weight class. This event is *bad* if it behaves like GREEDY in that an element in the current solution (from the previous interval) conflicts with up to *k* optimal elements (in the next interval), all of which have roughly the same weight. This suggests that a careful weight class partitioning is needed. We use a randomized weight class partitioning to avoid the greedy bottleneck. Instead of choosing the *i*th weight class as $[W(1-\varepsilon)^i, W(1-\varepsilon)^{i-1}]$, where *W* is the largest weight of the instance we randomly shift all intervals by a small factor. Using this randomized procedure, we argue that, with high probability, an optimal element is far from being in an anterior weight class. This shows that even if *o* was conflicting with some element in a prior weight class the weight difference is large enough to improve over the factor *k* that GREEDY attains (with high probability). We achieve an improved approximation guarantee by choosing the parameter ε that balances these two factors.

2 Preliminaries

For any integer $k \in \mathbb{N}$, we use the convention $[k] = \{1, ..., k\}$. We assume that the reader is familiar with basic matroid definitions (for a good survey, see [S⁺03]). We define the *k*-matroid intersection and matroid *k*-parity problems. All of our results will hold for a more general problem, Matroid *k*-Parity, which is the main focus of this paper.

k-Matroid Intersection: Given k matroids $\{\mathcal{M}_i = (V, \mathcal{I}_i)\}_{i \in [k]}$ defined on the same groundset V and a linear weight function $w : V \to \mathbb{R}_{\geq 0}$, the goal is to find a set $S \subseteq V$ of maximum weight independent in each matroid, i.e. such that $S \in \mathcal{I}_i$ for all $i \in [k]$.

Matroid *k*-Parity: We are given an edge-weighted hypergraph G = (V, E) on a set of vertices V with linear weight function $w : E \to \mathbb{R}_{\geq 0}$ and a matroid $\mathcal{M} = (V, \mathcal{I})$ defined on V. Each hyperedge $e \in E$ has size at most k. The goal is to find a maximum weight collection of disjoint hyperedges $M \subseteq E$ such that the vertices incident to M are independent in \mathcal{I} . We assume access to a matroid independence oracle, which for any set $S \subseteq V$ answers whether $S \in \mathcal{I}$. Throughout the document, we will denote by e(V') the set of edges incident to $V' \subseteq V$ with $V' \in \mathcal{I}$, and conversely let v(E') be the set of vertices contained in $E' \subseteq E$. We say that $A \subseteq E$ is a *feasible solution* if $v(A) \in \mathcal{I}$.

Observe that if \mathcal{M} is the *free matroid* (where $\mathcal{I} = 2^V$), then the problem is known as *k*-Set Packing. As shown by Lee, Sviridenko and Vondrák [LSV10] finding a maximum weight independent set in the intersection of *k*-matroids can be cast as a matroid *k*-parity problem in which all the edges $e \in E$ are disjoint. In fact without loss of generality, we may assume that the input graph G = (V, E) has the property that each vertex *v* belongs to a unique hyperedge. Repeating [LSV10], we provide a proof of these two facts in Appendix E for completeness. As Matroid *k*-Parity generalizes *k*-Matroid Intersection, we provide all algorithms and analyses for the Matroid *k*-Parity. It will also be helpful to define a vertex-weighted function.

Definition 1. Let $A \subseteq E$ be a collection of disjoint hyperedges, we define $c : v(A) \to \mathbb{R}_{\geq 0}$ where $c_v = w_e$ where $v \in e$, and $c(S) = \sum_{v \in S} c_v$ for all $S \subseteq v(A)$. Thus, c_v is the cost of the hyperedge $e \in A$ covering v. This function is well defined since A is a collection of disjoint hyperedges.

Our analyis will use the following two theorems (proved in Appendix C) about matroid exchanges.

Theorem 2. Let $\mathcal{M} = (V, \mathcal{I})$ be a matroid and S, T be independent sets of \mathcal{M} with $|S| \leq |T|$. For any partition S_1, \ldots, S_m of S, there are disjoint sets $T_1, \ldots, T_m \subseteq T$ such that $|S_i| = |T_i|$ and

$$S_i \cup (T \setminus T_i) \in \mathcal{I}.$$

In particular, if |S| = |T|, then the sets T_i partition T.

This is a slightly more general version of a Rota-exchange where S and T are not necessarily bases of \mathcal{M} . The second proposition, which is an almost immediate corollary, will be useful in our analysis. It states that we can construct exchanges in a top-to-bottom fashion. We can use the existence of a large matroid exchange to repartition into disjoint smaller matroid exchanges.

Theorem 3. Let $\mathcal{M} = (V, \mathcal{I})$ be a matroid and S, T be independent sets. Suppose that there is $N_S \subseteq T$ such that $|N_S| = |S|$ and $S \cup (T \setminus N_S) \in \mathcal{I}$. Then, for any partition of S into S_1, \ldots, S_m there is a partition $\{N_{S_i}\}_{i \in [m]}$ of N_S such that $|N_{S_i}| = |S_i|$ and $S_i \cup (T \setminus N_{S_i}) \in \mathcal{I}$.

Other Notations: We denote by $W \triangleq \max_{e \in E, v(e) \in \mathcal{I}} w(e)$ the maximum weight of any feasible edge and denote an interval by $I \subseteq [0, W]$. For simplicity, we overload the notation of I and also let I refer to the set of edges $e \in E$ where $w_e \in I$. Thus, given a set of edges $R \subseteq E$, we let $R \cap I \triangleq \{e \in R : w_e \in I\}$. An interval is then a weight class. Finally, given a collection of intervals $\{I_j\}_{j=1}^m$ with $m \in \mathbb{N}$ and a set of edges $R \subseteq E$, we will denote $R_{\leq i} = \bigcup_{j=1}^i R \cap I_j, R_{\geq i} = \bigcup_{j=i}^m R \cap I_j$.

3 Sliding Local-Search Algorithm

We describe our main algorithm SLIDING LOCAL-SEARCH (Algorithm 2), abbreviated by SLIDINGLS. SLIDINGLS iteratively runs a local-search algorithm over disjoint intervals of bounded-length in order of decreasing weights.

3.1 INTERVAL LOCAL-SEARCH

INTERVAL LOCAL-SEARCH (Algorithm 1) is a standard local-search algorithm which attempts to improve the current solution by finding improving (A, I, I)-swaps where $I \subseteq [0, W]$.

Definition 2 ((A, I, J)-swap). Let $A \subseteq E$ be any feasible solution, and $I, J \subseteq [0, W]$ be two intervals. A pair (S, N) is a (A, I, J)-swap if $S \subseteq (E \setminus A) \cap I$ and $|S| \leq 2$, $N \subseteq A \cap J$ and $|N| \leq 2k$, and $(A \setminus N) \cup S$ is a feasible solution. We further say that (S, T) is *improving* if w(S) > w(N).

We recall that without loss of generality (see Appendix E.2) all edges of our *k*-Matroid Parity instance are disjoint. Thus, performing an (A, I, J)-swap necessarily yields a feasible collection of disjoint hyperedges.

Algorithm 1: Interval Local-Search (A, I)			
Input: Feasible solution $A \subseteq E$, and interval $I \subseteq [0, W]$			
while There exists (S, N) that is an improving (A, I, I) -swap do			
$A \leftarrow A ackslash N \cup S$ // Improve the solution			
return A			

3.2 SLIDING LOCAL-SEARCH

SLIDING LOCAL-SEARCH iteratively calls INTERVAL LOCAL-SEARCH on geometrically decreasing bounded length intervals (except the last one). Formally, the interval [0, W] is partitioned into L + 1 disjoint intervals $\{I_j\}_{j=1}^{L+1}$, for some $L \in \mathbb{N}$. Each interval $I_j \subseteq [0, W]$ for $j = 1, \ldots, L + 1$ is defined by two markers. We let $I_j = (m_j, m_{j-1}]$ where $m_j \in [0, W]$ is the j^{th} -marker. The size of each interval is controlled by a parameter $\varepsilon > 0$, which will be optimized later, such that $\frac{m_{i-1}}{m_i} = 1 - \varepsilon$. Instead of deterministically choosing $m_i = W(1 - \varepsilon)^i$, SLIDINGLS chooses a random threshold $\tau \sim \mathcal{U}[0, \varepsilon)$, where $\mathcal{U}[a, b)$ is the uniform distribution between a and b, to shift the marker placement and lets $m_i = W(1 - \tau)(1 - \varepsilon)^{i-1}$.

Definition 3 (Markers). Each interval I_j is defined by two markers m_{j-1} and m_j such that $I_j = (m_j, m_{j-1}]$ for $j \in [L]$ and $I_{L+1} = [m_{L+1}, m_L]$. For some τ sampled from $\mathcal{U}[0, \varepsilon)$, the markers are chosen as: $m_j = W \cdot (1 - \varepsilon)^{j-1} (1 - \tau)$ for $j = 0, 1, \ldots, L$ and $m_{L+1} = 0$.

SLIDINGLS then calls INTERVAL LOCAL-SEARCH on I_1 , then I_2 and so on. More precisely, it runs INTERVAL LOCAL-SEARCH on $I_1 = (m_1, m_0]$ to find an initial solution $A_{\leq 1} \subseteq I_1$. It then seeks to improve the solution using edges of the next interval. It runs INTERVAL LOCAL-SEARCH $(A_{\leq 1}, I_2)$ to obtain solution $A_{\leq 2} \subseteq I_1 \cup I_2$, where $I_2 = (m_2, m_1]$. Since, INTERVAL LOCAL-SEARCH attempts to find improving $(A_{\leq 2}, I_2, I_2)$ -swap, the edges in $A_{\leq 1} \subseteq I_1$ aren't discarded when processing I_2 . This implies that $A_{\leq 1} \subseteq A_{\leq 2}$. We denote by $A_{\leq i+1}$ the output of INTERVAL LOCAL-SEARCH $(A_{\leq i}, I_{i+1})$. Similarly, we have that $A_{\leq i} \subseteq A_{\leq i+1}$.

Remark 1. When SLIDINGLS terminates, there is no improving $(A_{\langle i}, I_i, I_i)$ -swap for all $i \in [L + 1]$.

In Appendix D, we prove, using standard weight-scaling techniques [LSV09, LSV10, Ber00] that SLIDINGLS terminates in polynomial time as long as k is constant. We note this is also needed in [Ber00, Cyg13, LSV10]. Additionally, broadly inspired by streaming algorithms such as [BMKK14, KMZ⁺19], we prove that we can "discard" edges in the last interval.

Proposition 1. Let $L \triangleq \lceil -\log_{1-\varepsilon}(|E|\delta^{-1}) \rceil + 1$. Let O be the optimal solution and let A be the output of SLIDINGLS, and $O_{\leq L} = \{o \in O : w_o \geq m_L\}$. Then, $w(O \setminus O_{\leq L}) \leq \delta w(O)$.

Input: Parameters δ , ε that quantify the number and the size of the intervals				
Let $1 - \tau \sim \mathcal{U}(1 - \varepsilon, 1]$	// Random marker placement			
Let $L \triangleq \left\lceil -\log_{1-\varepsilon}(E \delta^{-1}) \right\rceil + 1$	<pre>// Number of intervals</pre>			
Let $W = \max_{e \in E, v(e) \in \mathcal{I}} w(e)$				
Let $m_0 = W(1-\tau)/(1-\varepsilon)$, $m_i = W(1-\tau)(1-\varepsilon)^{i-1}$ for $i \in [L]$, and $m_{L+1} = 0$				
for $i \leftarrow 0$ to L by 1 do				
$A \leftarrow \text{Interval Local-Search}(A, (m_{i+1}, m_i])$				
return A				

4 A better-than-*k* approximation guarantee

In this section, we give a simplified analysis of Algorithm 2 and obtain a $\frac{9}{10}(k+1)$ -approximation algorithm. In Appendix B, we show that the approximation factor can be improved to $(k+1)/(2\ln(2))$. Formally,

Theorem 4. For any $k \ge 3$, there is a randomized $\frac{9}{10}(k+1)$ -approximation algorithm for weighted Matroid *k*-Parity.

This is the first result that improves over GREEDY which yields a k-approximation for Matroid k-Parity and over the k - 1-approximation algorithm of Lee, Sviridenko and Vondrák [LSV09] for k-Matroid Intersection. We show in Appendix D.3 that this guarantee holds with high probability. To analyze the approximation guarantee of Algorithm 2, we will construct (A, I, J)-swaps (Definition 2) in Lemma 2. In Section 4.1, we detail our approach, give key definitions and state our main technical lemma (Lemma 2). We prove Lemma 2 in Section 4.2 using the local optimality of our solution within intervals. In Section 4.3, we analyze the contribution of the swaps across different intervals.

4.1 Approach and Definitions

To find a collection of local swaps between A and O we think of our algorithm as the following process. Before the start of iteration i, our algorithm maintains a solution $v(A_{\leq i-1})$, where $v(A_0) = \emptyset$ and a set $T_{i-1} \subseteq v(O)$ such that $v(A_{\leq i-1}) \cup (v(O) \setminus T_{i-1})$ is independent. The set T_{i-1} are the vertices *conflicting* with $v(A_{i-1})$. At the end of iteration i, we have extended $v(A_{\leq i-1})$ by finding a set $v(A_i)$. This set further blocks new vertices – the set $T_i \setminus T_{i-1}$. Lemma 1 makes this construction precise. This, however, doesn't define local swaps yet. We repeatedly apply Rota's exchange (Theorem 2 and Theorem 3) to define the set of optimal edges that are blocked at any iteration, and then construct local swaps in Lemma 2.

Lemma 1. There exists a nested sequence of sets $\emptyset = T_0 \subseteq T_1 \subseteq \ldots \subseteq T_{L+1} \subseteq v(O)$ such that $v(A_{\leq i}) \cup (v(O) \setminus T_i) \in \mathcal{I}$ and $|T_i \setminus T_{i-1}| = |v(A_i)|$. Moreover, for any $i \in [L+1]$, and any $o \in O \cap I_i$ such that $w_o > 0$, there exists some $j \leq i$ such that $v(o) \cap T_j \neq \emptyset$.

Lemma 1 shows the existence of a set of vertices $T_i \setminus T_{i-1} \subseteq v(O)$ which *conflict* with $v(A_i)$ at the i^{th} iteration and thus have to be discarded. The proof is in Appendix A.1. This implies that some optimal edges conflict with the solution $A_{\leq i}$ and we define these edges as follows.

Definition 4. Let $O^{(i)} \triangleq \{o \in O : v(o) \cap T_i \neq \emptyset, \text{ and } v(o) \cap T_{i-1} = \emptyset\}$ be the edges of O which *conflict* with A for the first time when processing interval $i \in [L+1]$.

We will construct swaps for edges in $O^{(i)}$. In particular, it will be convenient to expand Definition 2 for a partial solution $A_{\leq j}$ at a j^{th} iteration of the algorithm.

Definition 5 (Local swap). Let A be any feasible solution, and let (S, N) be a pair of sets. We say that (S, N) is a *local swap* if there exists $i \in [L+1]$ and $j \leq i$ such that (S, N) is a $(A_{\leq j}, I_i, I_j)$ -swap.

4.1.1 Recovering the Unweighted Guarantees

To leverage the power of local-search in almost unweighted sub-instances, consider the edges of $O^{(i)} \cap I_i$ that lie in I_i and are blocked for the first time at the *i*th iteration of Algorithm 2. We will prove that these sets behave like an unweighted *k*-matroid parity instance. As their weights differ by only a small factor $1 - \varepsilon$, this implies that the local-search gives a nearly k/2-approximation using [LSV10]. For every $i \in [L]$, we define the unweighted portion of the instance as follows,

- O_s⁽ⁱ⁾ = {o ∈ O⁽ⁱ⁾ ∩ I_i: |v(o) ∩ T_i| = 1} is the set of blocked edges in the ith interval with a single vertex conflict, and let O_s = U_{i∈[L]}O_s⁽ⁱ⁾ be the set of singles.
- O⁽ⁱ⁾_d = {o ∈ O⁽ⁱ⁾ ∩ I_i: |v(o) ∩ T_i| ≥ 2} is the set of blocked edges in the ith interval with a double vertex conflict, and let O_d = ∪_{i∈[L]} O⁽ⁱ⁾_d be the set of doubles.

Note that these sets are only defined up to the L^{th} interval. Equipped with the above definitions, we state our main result that constructs local swaps $\{(o, e(N_o))\}_{o \in O}$ between A and O at each iteration of Algorithm 2. Each swap is non-improving and satisfies a nice partition property such that each edge $a \in A$ isn't part of more than k swaps. Additionally, every $a \in A$ appears in the local swap of at most 1 *single* optimal edge. We also show that for every $o \in O_d$, its weight is much smaller than the weight of its neighborhood. This will allow us to obtain two straightforward corollaries. The first one says that Algorithm 2 is always a k-approximation. The second bounds the total weight of O_s (similarly to [LSV10]).

Lemma 2. For any random choice of τ , there exists a collection $\{(o, e(N_o))\}_{o \in O}$ of local swaps such that

- **1.** for all $i \in [L+1]$, and every $o \in O^{(i)}$, we have $\emptyset \neq N_o \subseteq v(A_i)$ and $w_o \leq c(N_o)$,
- **2.** each $v \in v(A)$ appears in at most one of the sets $\{N_o\}_{o \in O}$,
- **3.** for every $o \in O_s$, $c(N_o) = w(e(N_o))$ and for $o' \neq o \in O_s$, we have $e(N_o) \cap e(N'_o) = \emptyset$,
- 4. for every $o \in O_d$, we have that $|N_o| \ge 2$ and $w_o \le \frac{1}{2(1-\varepsilon)}c(N_o)$.

Corollary 1. Let $\{(o, e(N_o))\}_{o \in O}$ be the collection from Lemma 2, then $\sum_{o \in O} c(N_o) \leq kw(A)$. In particular, we have that $w(O) \leq kw(A)$.

Proof of Corollary 1. By Lemma 2, each $v \in v(A)$ appears at most once in one set N_o for $o \in O$. Since every $a \in A$ contains at most k vertices of v(A) we get that,

$$\sum_{o \in O} c(N_o) = \sum_{a \in A} w_a \sum_{o \in O} |v(a) \cap N_o| \le \sum_{a \in A} w_a \cdot k = k \cdot w(A).$$

Applying Property 1 of Lemma 2 and the previous bound, we get that: $w(O) \leq \sum_{o \in O} c(N_o) \leq kw(A)$.

Corollary 2. By the properties 1 and 3 of Lemma 2, we have $w(O_s) \le w(A)$.

4.2 Proof of Lemma 2

Proof of Lemma 2. Let $\{T_i\}_{i=1}^{L+1}$ be the sequence of sets obtained from Lemma 1. We apply Rota exchange properties (Theorem 2) on \mathcal{M}_i , the matroid contracted on $v(A_{\leq i-1}) \cup (v(O) \setminus T_i)$, to obtain local swaps $\{(o, e(N_o))\}_{o \in O}$. By Lemma 1, the sets $T_i \setminus T_{i-1}$ and $v(A_i)$ are independent in \mathcal{M}_i and $|v(A_i)| = |T_i \setminus T_{i-1}|$. We partition $T_i \setminus T_{i-1}$ as follows: let $B_s = v(O_s^{(i)}) \cap T_i$ and $B' = v(O^{(i)} \setminus O_s^{(i)}) \cap T_i$ and $B'' = (T_i \setminus T_{i-1}) \setminus (B_s \cup B')$. Note that this is a partition of $T_i \setminus T_{i-1}$ by definition of $O^{(i)}$. Here B_s is simply empty when i = L + 1. By Theorem 2, there exist disjoint sets $N_s, N' \subseteq v(A_i)$ such that $v(A_i)\setminus N_s \cup B_s$ and $v(A_i)\setminus N' \cup B'$ are independent in \mathcal{M}_i and $|N_s| = |B_s|$, and |N'| = |B'|respectively. Property 3 follows from the following two claims which we now prove. Let $i \in [L]$, then

- 1. For all $a \in A_i$, we have that $|v(a) \cap N_s| \leq 1$,
- 2. $N_s = \bigsqcup_{o \in O_s^{(i)}} N_o$, and $|N_o| = 1$ for $o \in O_s^{(i)}$.

We start with the first claim. Suppose conversely that there exists an edge $a \in A_i$ such that $|v(a) \cap N_s| \geq 2$. This implies that $|v(A_i) \setminus (N_s \cup v(a)) \cup B_s| = |v(A_i)| - (|N_s| + |v(a)| - |N_s \cap v(a)|) + |B_s| \geq |v(A_i - a)| + 2$ since $|N_s| = |B_s|$. Because $v(A_i) \setminus (N_s \cup v(a)) \cup B_s$ and $v(A_i - a)$ are independent in \mathcal{M}_i , we can find an extension $b, b' \in B_s$ such that $v(A_i - a) \cup \{b, b'\} \in \mathcal{I}_i$. Observe that the vertices b, b' belong to distinct hyperedges in $O_s^{(i)}$, which we denote by $e, e' \in O_s^{(i)}$, respectively. Since $v(A_i - a) \cup \{b, b'\}$ is independent in \mathcal{M}_i , we have that $(v(A_{\leq i} - a) \cup \{b, b'\} \cup (v(O) \setminus \mathcal{T}_i) \in \mathcal{I}$. In particular, $A_{\leq i} \setminus \{a\} \cup \{e, e'\}$ is feasible since $\{e, e'\} \subseteq \{b, b'\} \cup (v(O) \setminus \mathcal{T}_i)$ by definition of $O^{(i)}$. We prove that this yields an improving $(A_{\leq i}, I_i)$ -swap as long as $\varepsilon \in (0, 0.5)$ and $i \in [L]$. We have that $a \in A_i \subseteq I_i$ and $e, e' \in O_s^{(i)} \subseteq I_i$. Since $I_i = (m_i, m_{i-1}]$, we have

$$w_e + w'_e - w_a > 2m_i - m_{i-1} = 2(1 - \varepsilon)m_{i-1} - m_{i-1} = m_{i-1}(1 - 2\varepsilon) > 0,$$

where we use $m_i = (1 - \varepsilon)m_{i-1}$ for $i \in [L]$, contradicting the absence of improving $(A_{\leq i}, I_i)$ -swaps.

We prove the next claim together with Property 1 and 2. For every $i \in [L + 1]$, we partition B_s into $\{v(o) \cap B_s\}_{o \in O_s^{(i)}}$ and B' into $\{v(o) \cap B'\}_{o \in O^{(i)} \setminus O_s^{(i)}}$. Applying Theorem 3, we get that there are collections of disjoints sets $\{N_o\}_{o \in O_s^{(i)}}$ and $\{N'_o\}_{o \in O^{(i)} \setminus O_s^{(i)}}$ partitioning N_s and N' such that $v(A_i) \setminus N_o \cup (v(o) \cap B_s) \in \mathcal{I}_i$ for all $o \in O_s^{(i)}$ and $v(A_i) \setminus N'_o \cup (v(o) \cap B') \in \mathcal{I}_i$ for all $o \in O^{(i)} \setminus O_s^{(i)}$, respectively. Theorem 3 further says that $|N'_o| = |v(o) \cap B'| \ge 1$ and $|N_o| = |v(o) \cap B_s| = 1$, where the second equality uses that $|v(o) \cap T_i| = 1$ for all $o \in O_s^{(i)}$. We prove that $\{(o, e(N_o))\}_{o \in O_s^{(i)} \setminus O_s^{(i)}}$ are local swaps. Fix $i \in [L+1]$ and let $o \in O_s^{(i)}$ together with its pair $(o, e(N_o))$. The proof is identical for $o \in O^{(i)} \setminus O_s^{(i)}$. We prove that the swap is independent:

$$v(A_{\leq i}) \setminus \{v(e(N_o))\} \cup v(o) \subseteq v(A_{\leq i}) \setminus N_o \cup v(o)$$
$$\subseteq \left(v(A_i) \setminus N_o \cup (v(o) \cap T_i)\right) \cup \left(v(A_{\leq i-1}) \cup (v(O) \setminus T_i)\right)$$
$$\subseteq \left(v(A_i) \setminus N_o \cup (v(o) \cap (T_i \setminus T_{i-1}))\right) \cup \left(v(A_{\leq i-1}) \cup (v(O) \setminus T_i)\right)$$

where we used in the last containment that $v(o) \cap T_{i-1} = \emptyset$ for all $o \in O^{(i)}$. The last term is independent since $v(o) \cap B_s = v(o) \cap T_i \setminus T_{i-1}$ and $v(A_i) \setminus N_o \cup (v(o) \cap B_s)$ is independent in \mathcal{M}_i . We have that $(o, e(N_o))$ is, in fact, a $(A_{\leq i}, I_\ell, I_i)$ -swap for some $\ell \geq i$, since $o \in I_\ell$ by the second part of Lemma 1. We now prove Property 1. Let $v \in N_o$ (which exists since $|N_o| \geq 1$) and let a be the edge incident to v. If $i = \ell$, then the property follows from the absence of $(A_{\leq i}, I_i)$ -swap, which implies that $w_o \leq w(e(N_o)) \leq c(N_o)$. Then, suppose $i < \ell$, we have that $w_o \leq m_{\ell-1} \leq m_i < w_a \leq c(N_o)$ by definition of the markers. Property 2 follows from the fact that that $\{v(A_i)\}_{i \in [L+1]}$ are disjoint sets of vertices and that the sets N_o are also disjoint.

Lastly, we show these local swaps satisfy Property 4 for $o \in O_d$. Let $i \in [L]$ be any index and fix $o \in O_d^{(i)}$. By construction of the local swaps $(o, e(N'_o))$, we have $|N'_o| = |v(o) \cap B'| = |v(o) \cap T_i \setminus T_{i-1}| \ge 2$. Let $v_1, v_2 \in N_o$ be two distinct vertices. Further, let $a_1, a_2 \subseteq e(N_o) \subseteq A_i$ be the edges incident to v_1 and v_2 . It is possible that $a_1 = a_2$. Since $a_j \in A_i$, we have that $m_{i-1} \ge w_{a_j} > m_i$ for j = 1, 2. Additionally, we have that $o \in O_d^{(i)}$, so o must lie in I_i . This means that $w_o \le m_{i-1}$. Therefore,

$$w_o = \frac{w_o}{c(v_1) + c(v_2)} \left(c(v_1) + c(v_2) \right) \le \frac{m_{i-1}}{2m_i} c(N_o) = \frac{1}{2(1-\varepsilon)} c(N_o).$$

4.3 Escaping local-optima using randomness

By Property 3 and 4 of Lemma 2 we are able to analyze the absence of improving swap in a given interval. Unfortunately, the optimal edges in the *i*th interval might already be blocked after processing an anterior interval j < i. This poses problems as an edge in $a \in A_{i-1}$ may block up to k edges in $O \cap I_i$ that differ in weight only slightly. If this happens, then SLIDINGLS has the same guarantees as GREEDY. Importantly, this situation can be avoided by slightly shifting the *i*th-marker. We use the random marker placement to show that such events occur with low probability. This motivates the following definitions. Let $\gamma \in (0, 1)$ be a fixed constant.

Definition 6. For each $o \in O$, we define by m_o the closest marker to w_o such that $m_o \ge w_o$.

We partition the remaining edges of $O \setminus (O_s \cup O_d)$ into those that are close to m_o and

- $O_b = \{o \in O \setminus (O_s \cup O_d) : w_o \ge (1 + \gamma)^{-1} m_o\}$ is the set of blocked edges whose weight is close to m_o . These edges are *bad* for the analysis.
- O_g = {o ∈ O\(O_s ∪ O_d) : w_o < (1 + γ)⁻¹m_o} is the set of blocked edges whose weight is far from m_o. These edges are good for the analysis as shown in Lemma 3 proved in Appendix A.2.

Lemma 3. Let $o \in O_g \cap O_{\leq L}$ and $(o, e(N_o))$ be its local swap given by Lemma 2. Then,

$$(1+\gamma)w_o \le c(N_o).$$

Motivated by Lemma 3, our objective is then to estimate the expected weight of O_b . In Lemma 4, we use the random threshold to argue that the expected weight is small.

Lemma 4. Let $\varepsilon > 0$ be the parameter from Algorithm 2. For any $o \in O$ and $\gamma \in [0, (1 - \varepsilon)^{-1} - 1)$, we have that

$$\mathbb{P}_{\tau} \left[o \in O_b \right] \le \mathbb{P}_{\tau} \left[w_o \ge (1+\gamma)^{-1} m_o \right] \le \frac{\gamma}{\varepsilon (1+\gamma)}.$$

In particular, the following holds: $\mathbb{E}\left[w(O_b)\right] \leq \frac{\gamma}{\varepsilon(1+\gamma)}w(O)$.

The first inequality follows from the definition of O_b . Thus, Lemma 4 tells that it is sufficient to compute the probability that an edge o lands close to m_o to bound the contribution of O_b .

Proof of Lemma 4. Fix any $o \in O$. We start by proving the lemma when $w_o \in [1 - \varepsilon, 1] \cdot W$ and split the proof into two computations depending on the value of w_o . Suppose that $w_o \in [1 - \varepsilon, \frac{1}{1+\gamma}] \cdot W$. Then, the only marker for which $w_o \ge (1 + \gamma)^{-1} m_o$ is if $m_o = m_1$. Thus,

$$\mathbb{P}_{\tau}\left[w_o \ge (1+\gamma)^{-1} m_o\right] = \mathbb{P}_{\tau}\left[m_1 \in [w_o, w_o(1+\gamma)]\right] = \frac{1}{\varepsilon \cdot W} \cdot (w_o(1+\gamma) - w_o) \le \frac{\gamma}{\varepsilon(1+\gamma)}$$

In the second inequality, we used that $m_1 = (1 - \tau)W \sim \mathcal{U}(1 - \varepsilon, 1] \cdot W$ and that $[w_o, w_o(1 + \gamma)] \subseteq [1 - \varepsilon, 1] \cdot W$. The last inequality uses that $w_o \leq \frac{W}{1+\gamma}$. Suppose alternatively that $w_o \in [\frac{1}{1+\gamma}, 1] \cdot W$. Then, we have:

$$\mathbb{P}_{\tau} \left[w_o \ge (1+\gamma)^{-1} m_o \right] = \mathbb{P}_{\tau} \left[(m_1 \in [w_o, 1]) \lor (m_0 \in [1, (1+\gamma)w_o]) \right]$$
$$= \mathbb{P}_{\tau} \left[m_1 \in [w_o, 1] \right] + \mathbb{P}_{\tau} \left[m_1 \in [1-\varepsilon, (1-\varepsilon)(1+\gamma)w_o] \right]$$
$$= \frac{1}{\varepsilon \cdot W} \cdot \left(\varepsilon + w_o \left((1-\varepsilon)(1+\gamma) - 1 \right) \right)$$
$$\leq \frac{1}{\varepsilon} \cdot \frac{\gamma}{1+\gamma}.$$

In the second equality, we use that events are disjoint from each other and that $m_0(1-\varepsilon) = m_1$. The third equality uses that $m_1 \sim \mathcal{U}(1-\varepsilon, 1] \cdot W$. In the last inequality, we observe that $1+\gamma \leq (1-\varepsilon)^{-1}$ so $(1-\varepsilon)(1+\gamma) - 1 \leq 0$. Substituting $w_o \geq (1+\gamma)^{-1}W$ into the last inequality yields the desired result. This proves the result when $w_o \in [1-\varepsilon, 1] \cdot W$.

Suppose now that $w_o \in [(1-\varepsilon)^i, (1-\varepsilon)^{i-1}] \cdot W$ for some $i \in [L+1]$, we reduce this case to the previous computation. Indeed, the probability that $w_o \in [\frac{1}{1+\gamma}m_j, m_j]$ for some j is equal to the probability that the point $\frac{w_o}{(1-\varepsilon)^{i-1}}$ is in $[\frac{1}{(1+\gamma)}m_{j-i+1}]$. Note that $\frac{w_o}{(1-\varepsilon)^{i-1}} \in [1-\varepsilon, 1] \cdot W$, thus, by the previous computation, we get that o is in O_b with probability at most $\frac{\gamma}{\varepsilon(1+\gamma)}$. Lastly, suppose that $w_o \in [0, (1-\varepsilon)^{L+1}] \cdot W$. Then as $m_L = W(1-\tau)(1-\varepsilon)^{L-1}$, we have that $(1+\gamma)^{-1}m_o = (1+\gamma)^{-1}m_L \geq W(1-\varepsilon)^{L+1} \geq w_o$, where we used that $(1-\tau) \geq 1-\varepsilon$ and $(1+\gamma)^{-1} \geq 1-\varepsilon$. This implies that $o \in O_g$.

4.4 Obtaining an $\Omega(k)$ approximation ratio

We combine the improvements due to the local-search analysis of Lemma 2 and to the random marker analysis of Lemma 4 to show that Algorithm 2 is an $\Omega(k)$ -approximation algorithm for weighted *k*-Matroid Intersection and weighted *k*-Matroid Parity. The full proof is in Appendix A.3.

Theorem 5. Let $\varepsilon \in (0, 1/2)$, $\delta \in (0, 1)$ be the parameters of Algorithm 2, and let $\gamma \in (0, 1)$ be a constant such that $\gamma \leq (1 - \varepsilon)^{-1} - 1$ and $\gamma \leq 2(1 - \varepsilon) - 1$. Let *A* be the output of Algorithm 2. Then,

$$w(O) \le \frac{k+\gamma}{1+\gamma\left(1-\frac{\gamma}{\varepsilon(1+\gamma)}\right)-\delta} \cdot \mathbb{E}\left[w(A)\right].$$

Setting $\varepsilon = 0.3873$, $\gamma = 0.2253$ and δ small enough we get that $w(O) \leq \frac{9(k+1)}{10} \cdot \mathbb{E}[w(A)]$.

4.5 Conclusion and Open Questions

We study weighted k-matroid intersection and its generalization as weighted matroid k-parity. In contrast to special cases, such as k-set packing, both problems are not as well understood. Prior

to this work, GREEDY was the state-of-the-art algorithm with a k-approximation. We significantly improve over this bound and provide a $\frac{k+1}{2\ln(2)} \simeq 0.722(k+1)$ -approximation which largely reduces the gap between the unweighted and weighted setting. Our algorithm is conceptually simple and provides an alternative to the current algorithms for special cases such as Berman's algorithm. We iteratively apply a weighted local-search over random intervals which prevents the algorithm from reaching certain bad local optima. Beyond an improved constant, our algorithm offers a blueprint for transforming algorithms over the unweighted setting into algorithms over the weighted setting. This may be of independent interest and could lead to improvements for other problems beyond Matroid k-Parity.

Open Questions: It remains interesting to close the gap between the unweighted and the weighted setting, even in the simpler case of *k*-Set Packing for which there is still a gap [Cyg13, Neu23]. On the hardness front, as mentioned in [LST24], better hardness of approximation for *k*-CSP might improve the k/12-hardness of *k*-Dimensional Matching which would extend to *k*-Matroid Intersection. Finally, a clear direction for future improvements is to investigate whether there exists a $\frac{k+1}{3}$ -approximation for unweighted *k*-matroid intersection similarly to unweighted *k*-set packing or give a separation.

Acknowledgments

The authors are grateful to Ola Svensson for the many helpful discussions and comments.

References

- [AH98] Esther M. Arkin and Refael Hassin. On local search for weighted *k*-set packing. *Math. Oper. Res.*, 23(3):640–648, 1998. (Pages 1 and 2)
- [Ber00] Piotr Berman. A d/2 approximation for maximum weight independent set in d-claw free graphs. In *Scandinavian Workshop on Algorithm Theory*, pages 214–219. Springer, 2000. (Pages 1, 3, 6, and 21)
- [BK03] Piotr Berman and Piotr Krysta. Optimizing misdirection. In Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms, January 12-14, 2003, Baltimore, Maryland, USA, pages 192–201. ACM/SIAM, 2003. (Page 1)
- [BMKK14] Ashwinkumar Badanidiyuru, Baharan Mirzasoleiman, Amin Karbasi, and Andreas Krause. Streaming submodular maximization: massive data summarization on the fly. In Sofus A. Macskassy, Claudia Perlich, Jure Leskovec, Wei Wang, and Rayid Ghani, editors, *The 20th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, KDD '14, New York, NY, USA - August 24 - 27, 2014*, pages 671–680. ACM, 2014. (Page 6)
- [CH99] Barun Chandra and Magnús M. Halldórsson. Greedy local improvement and weighted set packing approximation. In Robert Endre Tarjan and Tandy J. Warnow, editors, Proceedings of the Tenth Annual ACM-SIAM Symposium on Discrete Algorithms, 17-19 January 1999, Baltimore, Maryland, USA, pages 169–176. ACM/SIAM, 1999. (Pages 1 and 3)

- [CLL14] Ho Yee Cheung, Lap Chi Lau, and Kai Man Leung. Algebraic algorithms for linear matroid parity problems. *ACM Transactions on Algorithms (TALG)*, 10(3):1–26, 2014. (Page 2)
- [Cyg13] Marek Cygan. Improved approximation for 3-dimensional matching via bounded pathwidth local search. In 54th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2013, 26-29 October, 2013, Berkeley, CA, USA, pages 509–518. IEEE Computer Society, 2013. (Pages 2, 3, 6, and 12)
- [Edm03] Jack Edmonds. *Submodular Functions, Matroids, and Certain Polyhedra*, pages 11–26. Springer Berlin Heidelberg, Berlin, Heidelberg, 2003. (Page 1)
- [Gal68] David Gale. Optimal assignments in an ordered set: an application of matroid theory. *Journal of Combinatorial Theory*, 4(2):176–180, 1968. (Page 1)
- [HS89] Cor A. J. Hurkens and Alexander Schrijver. On the size of systems of sets every t of which have an sdr, with an application to the worst-case ratio of heuristics for packing problems. *SIAM J. Discret. Math.*, 2(1):68–72, 1989. (Page 2)
- [HSS06] Elad Hazan, Shmuel Safra, and Oded Schwartz. On the complexity of approximating *k*-set packing. *Comput. Complex.*, 15(1):20–39, 2006. (Page 1)
- [IK17] Satoru Iwata and Yusuke Kobayashi. A weighted linear matroid parity algorithm. In Hamed Hatami, Pierre McKenzie, and Valerie King, editors, Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2017, Montreal, QC, Canada, June 19-23, 2017, pages 264–276. ACM, 2017. (Page 2)
- [JK82] Per M. Jensen and Bernhard Korte. Complexity of matroid property algorithms. *SIAM J. Comput.*, 11(1):184–190, 1982. (Page 2)
- [KMZ⁺19] Ehsan Kazemi, Marko Mitrovic, Morteza Zadimoghaddam, Silvio Lattanzi, and Amin Karbasi. Submodular streaming in all its glory: Tight approximation, minimum memory and low adaptive complexity. In *International Conference on Machine Learning*, pages 3311–3320. PMLR, 2019. (Page 6)
- [Las15] Michal Lason. List coloring of matroids and base exchange properties. *Eur. J. Comb.*, 49:265–268, 2015. (Page 20)
- [Law01] Eugene L Lawler. *Combinatorial optimization: networks and matroids*. Courier Corporation, 2001. (Page 2)
- [Lov80] László Lovász. Matroid matching and some applications. *J. Comb. Theory B*, 28(2):208–236, 1980. (Page 2)
- [LST24] Euiwoong Lee, Ola Svensson, and Theophile Thiery. Asymptotically optimal hardness for *k*-set packing and *k*-matroid intersection. *arXiv preprint arXiv:2409.17831*, 2024. (Pages 1 and 12)

- [LSV09] Jon Lee, Maxim Sviridenko, and Jan Vondrák. Submodular maximization over multiple matroids via generalized exchange properties. In Irit Dinur, Klaus Jansen, Joseph Naor, and José D. P. Rolim, editors, *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, 12th International Workshop, APPROX 2009, and 13th International Workshop, RANDOM 2009, Berkeley, CA, USA, August 21-23, 2009. Proceedings, volume 5687 of Lecture Notes in Computer Science*, pages 244–257. Springer, 2009. (Pages 1, 2, 6, 7, and 21)
- [LSV10] Jon Lee, Maxim Sviridenko, and Jan Vondrák. Matroid matching: the power of local search. In Leonard J. Schulman, editor, *Proceedings of the 42nd ACM Symposium on Theory of Computing, STOC 2010, Cambridge, Massachusetts, USA, 5-8 June 2010*, pages 369–378. ACM, 2010. (Pages 1, 2, 3, 4, 6, 7, 8, 21, and 23)
- [Neu21] Meike Neuwohner. An improved approximation algorithm for the maximum weight independent set problem in d-claw free graphs. In Markus Bläser and Benjamin Monmege, editors, 38th International Symposium on Theoretical Aspects of Computer Science, STACS 2021, March 16-19, 2021, Saarbrücken, Germany (Virtual Conference), volume 187 of LIPIcs, pages 53:1–53:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021. (Pages 1 and 2)
- [Neu22] Meike Neuwohner. The limits of local search for weighted k-set packing. In Karen I. Aardal and Laura Sanità, editors, *Integer Programming and Combinatorial Optimization* 23rd International Conference, IPCO 2022, Eindhoven, The Netherlands, June 27-29, 2022, Proceedings, volume 13265 of Lecture Notes in Computer Science, pages 415–428. Springer, 2022. (Page 1)
- [Neu23] Meike Neuwohner. Passing the limits of pure local search for weighted k-set packing. In Nikhil Bansal and Viswanath Nagarajan, editors, *Proceedings of the 2023 ACM-SIAM Symposium on Discrete Algorithms, SODA 2023, Florence, Italy, January 22-25, 2023,* pages 1090–1137. SIAM, 2023. (Pages 1 and 12)
- [Rad57] Richard Rado. Note on independence functions. Proceedings of the London Mathematical Society, 3(1):300–320, 1957. (Page 1)
- [S⁺03] Alexander Schrijver et al. Combinatorial optimization: polyhedra and efficiency, volume 24. Springer, 2003. (Pages 4 and 23)
- [TW23] Theophile Thiery and Justin Ward. An improved approximation for maximum weighted k-set packing. In Nikhil Bansal and Viswanath Nagarajan, editors, *Proceedings of the* 2023 ACM-SIAM Symposium on Discrete Algorithms, SODA 2023, Florence, Italy, January 22-25, 2023, pages 1138–1162. SIAM, 2023. (Pages 1 and 2)

A Omitted Proofs

For completeness, we provide the missing proofs from Section 4.

A.1 Proof of Lemma 1

Lemma 1. There exists a nested sequence of sets $\emptyset = T_0 \subseteq T_1 \subseteq \ldots \subseteq T_{L+1} \subseteq v(O)$ such that $v(A_{\leq i}) \cup (v(O) \setminus T_i) \in \mathcal{I}$ and $|T_i \setminus T_{i-1}| = |v(A_i)|$. Moreover, for any $i \in [L+1]$, and any $o \in O \cap I_i$ such that $w_o > 0$, there exists some $j \leq i$ such that $v(o) \cap T_i \neq \emptyset$.

Without loss of generality, we may assume that $|v(O)| \ge |v(A)|$ by adding disjoint edges of weight equal to 0, each incident to k new dummy vertices independent with everything else. We prove the statement by induction on i. Let $v(A_1)$ be the vertices incident to A_1 and apply Theorem 2 to find a set $T_1 \subseteq v(O)$ such that $v(A_1) \cup (v(O) \setminus T_1) \in \mathcal{I}$ and $|T_1| = |v(A_1)|$. This concludes the proof for i = 1.

By the induction hypothesis, we have a collection $T_0 \subseteq \ldots \subseteq T_{i-1} \subseteq v(O)$ such that $v(A_{\leq i-1}) \cup (v(O) \setminus T_{i-1}) \in \mathcal{I}$ and $|T_{i-1} \setminus T_{i-2}| = |v(A_{i-1})|$. Let $\mathcal{M}' = (V, \mathcal{I}')$ be the matroid \mathcal{M} contracted on $v(A_{\leq i-1})$. Observe that $v(O) \setminus T_{i-1}$ and $v(A_i)$ are independent in \mathcal{M}' . Moreover, by the induction hypothesis $0 \geq |v(A_{\leq i})| - |v(O)| = |v(A_{\leq i-1})| + |v(A_i)| - |v(O) \setminus T_{i-1}| - |T_{i-1}| = |v(A_i)| - |v(O) \setminus T_{i-1}|$. So $|v(O) \setminus T_{i-1}| \geq |v(A_i)|$. Thus, applying Theorem 2 to $v(A_i)$ and $v(O) \setminus T_{i-1}$ in \mathcal{M}' , we get that there exists a set $T_i \setminus T_{i-1} \subseteq v(O) \setminus T_{i-1}$ such that $v(A_i) \cup (v(O) \setminus T_{i-1}) \setminus (T_i \setminus T_{i-1}) \in \mathcal{I}'$ and that $|T_i \setminus T_{i-1}| = |v(A_i)|$. We finish the induction by uncontracting the matroid.

It remains to prove the second part of the statement. Let $o \in O \cap I_i$ be such that $w_o > 0$ for some $i \in [L+1]$. Suppose by contradiction that $v(o) \cap T_j = \emptyset$ for all $j \leq i$. This implies that:

$$v(A_{\leq i}) \cup v(o) \subseteq v(A_{\leq i}) \cup v(o) \cup (v(O) \setminus T_i) = v(A_{\leq i}) \cup (v(O) \setminus T_i) \in \mathcal{I},$$

where we used the first part of the lemma to claim the independence of the right-hand side. So v(o) can be added to $v(A_{\leq i})$ without violating the independence constraint. Since $w_o > 0$, the pair $(\{o\}, \emptyset)$ is an improving $(A_{\leq i}, I_i, I_i)$ -swap contradicting the termination of Algorithm 2.

A.2 Proof of Lemma 3

Lemma 3. Let $o \in O_q \cap O_{\leq L}$ and $(o, e(N_o))$ be its local swap given by Lemma 2. Then,

$$(1+\gamma)w_o \le c(N_o).$$

Let $o \in O_g \cap O_{\leq L}$ and let $(o, e(N_o))$ be its local swap from Lemma 2. Let $j \in [L]$ be the index such that $o \in I_j$. Since $o \notin O_s \cup O_d$, there exists an index i < j such that $o \in O^{(i)} \setminus (O_d^{(i)} \cup O_s^{(i)})$. In particular, by Property 1 of Lemma 2, we have that $\emptyset \neq e(N_o) \subseteq A_i$. This implies that $m_o = m_{j-1} \leq m_i$ so $w_o \leq (1+\gamma)^{-1}m_o \leq (1+\gamma)^{-1}m_i < (1+\gamma)^{-1}c(N_o)$.

A.3 Proof of Theorem 5 and Theorem 1

Theorem 5. Let $\varepsilon \in (0, 1/2)$, $\delta \in (0, 1)$ be the parameters of Algorithm 2, and let $\gamma \in (0, 1)$ be a constant such that $\gamma \leq (1 - \varepsilon)^{-1} - 1$ and $\gamma \leq 2(1 - \varepsilon) - 1$. Let *A* be the output of Algorithm 2. Then,

$$w(O) \le \frac{k+\gamma}{1+\gamma\left(1-\frac{\gamma}{\varepsilon(1+\gamma)}\right)-\delta} \cdot \mathbb{E}\left[w(A)\right].$$

Setting $\varepsilon = 0.3873$, $\gamma = 0.2253$ and δ small enough we get that $w(O) \leq \frac{9(k+1)}{10} \cdot \mathbb{E}[w(A)]$.

Recall that $O_{\leq L} \triangleq \{o \in O : w_o \geq m_L\}$. The proof follows from the fact that the probability of being in O_b is small (Lemma 4). As a consequence,

$$\left(1 - \frac{\gamma}{\varepsilon(1+\gamma)}\right) w(O) \leq \mathbb{E} \left[w(O_s \cup O_d \cup O_g)\right]$$

$$\leq \mathbb{E} \left[w(O_d)\right] + \mathbb{E} \left[w(O_g \cap O_{\leq L})\right] + \mathbb{E} \left[w(O_g \setminus O_{\leq L})\right] + \mathbb{E} \left[w(A)\right]$$

$$\leq \mathbb{E} \left[w(O_d)\right] + \mathbb{E} \left[w(O_g \cap O_{\leq L})\right] + \mathbb{E} \left[w(O \setminus O_{\leq L})\right] + \mathbb{E} \left[w(A)\right]$$

$$\leq \mathbb{E} \left[w(O_d)\right] + \mathbb{E} \left[w(O_g \cap O_{\leq L})\right] + \delta w(O) + \mathbb{E} \left[w(A)\right],$$

where the first inequality comes from Lemma 4 and the fact that $O = O_g \sqcup O_b \sqcup O_s \sqcup O_d$. The second inequality uses that $w(O_s) \le w(A)$ (Corollary 2). The last inequality uses Proposition 1. Next, recall that $(1 + \gamma)w_o \le c(N_o)$ for every $o \in O_d \cup (O_g \cap O_{\le L})$ whenever $(1 + \gamma) \le 2(1 - \varepsilon)$ by Lemma 2 and Lemma 3. Then by Proposition 1,

$$w(O) + \left(\gamma \left(1 - \frac{\gamma}{\varepsilon(1+\gamma)}\right) - \delta\right) w(O) \le w(O) + \gamma \left(\mathbb{E}\left[w(O_d)\right] + \mathbb{E}\left[w(O_g \cap O_{\le L})\right] + \mathbb{E}\left[w(A)\right]\right)$$
$$= (1+\gamma) \left(\mathbb{E}\left[w(O_d)\right] + \mathbb{E}\left[w(O_g \cap O_{\le L})\right]\right)$$
$$+ \gamma \cdot \mathbb{E}\left[w(A)\right] + \mathbb{E}\left[w(O_b \cup O_s \cup (O_g \setminus O_{\le L})\right]$$
$$\le \sum_{o \in O} \mathbb{E}\left[c(N_o)\right] + \gamma \mathbb{E}\left[w(A)\right]$$
$$\le k \cdot \mathbb{E}\left[w(A)\right] + \gamma \mathbb{E}\left[w(A)\right],$$

where the last inequality uses Corollary 1. Rearranging the terms yields the desired result.

Corollary 3. Let *A* be the output of Algorithm 2. Then,

$$w(O) \le \frac{k + 0.2253}{1.118} \cdot \mathbb{E}\left[w(A)\right]$$

Proof of Corollary 3. Substituting the following values for $\varepsilon = 0.3873$ and $\gamma = 0.2253$ and $\delta = .0001$ we obtain the desired result. We check that $\gamma \leq (1-\varepsilon)^{-1} - 1$ and that $\gamma \leq 2(1-\varepsilon) - 1$. We have that $(1-\varepsilon)^{-1} - 1 > 0.6322$ and $2(1-\varepsilon) - 1 > 0.2254$. On the other hand, $1+\gamma \left(1-\frac{\gamma}{\varepsilon(1+\gamma)}\right) - \delta \geq 1.118$.

Note that as $1/1.118 \le \frac{9}{10}$ Corollary 3 guarantees the approximation factor Theorem 1.

B Improved $(k+1)/(2\ln 2)$ approximation ratio

In Section 4.3, we defined the sets O_b and O_g as a function of γ measuring the distance between w_o and the closest marker of greater weight m_o . We have split $O \setminus (O_s \cup O_d)$ into the set of $o \in O$ such that $w_o \ge (1 + \gamma^{-1})m_o$ and those for which $w_o < (1 + \gamma)^{-1}m_o$. However, an optimal edge is only bad if it is very close to the marker, otherwise a small factor improvement can still be made in the analysis. Given this, we analyze the probability that any o falls within any prescribed range from m_o . This motivates the definition of the following sets. Let $m \in \mathbb{N}$ and let $\delta_j = \frac{\varepsilon_j}{m-\varepsilon_j}$ for $j = 0, \ldots, m$.

Definition 7. For $j \in [m]$, we define $R_j \triangleq \{o \in O : w_o \in ((1 + \delta_j)^{-1}, (1 + \delta_{j-1})^{-1}] \cdot m_o\}$ as the set of optimal edges in a factor between δ_j and δ_{j-1} close to their closest larger weight marker.

Remark 2. Observe that for any $o \in O_{\leq L}$, there exists an index $j \in [m]$ such that $o \in R_j$. This is by definition of δ as $\delta_0 = 0$ and $\delta_m = \frac{\varepsilon}{1-\varepsilon}$. In particular, we get $(1 + \delta_m)^{-1} = 1 - \varepsilon$ which is equal to multiplicative gap between two consecutive markers.

Observe that the set R_j is a random set due to the random choice to τ . We show that we can derive stronger swap properties than those derived in Lemma 2 and Lemma 4.

Lemma 5. Let $\{(o, e(N_o))\}_{o \in O}$ be the local swaps given by Lemma 2. Then, for any $j \in [m]$ and any $o \in (O_{\leq L} \setminus O_s) \cap R_j$, we have

$$(1+\delta_{j-1})w_o \le c(N_o).$$

Proof of Lemma 5. Suppose first that $o \in O_d^{(i)} \cap R_j$ for some $i \in [L]$ and let $(o, e(N_o))$ be its local swap given by Lemma 2. We have that $|N_o| \ge 2$ for all $o \in O_d^{(i)}$. Let v_1, v_2 be two distinct vertices, and let $a_1, a_2 \subseteq e(N_o) \subseteq A_i$ be the edges incident to v_1 and v_2 . It is possible that $a_1 = a_2$. Since $a_j \in A_i$, we have that $m_{i-1} \ge w_{a_j} > m_i$ for j = 1, 2. Additionally, we have that $o \in O_d^{(i)}$, so o must lie in I_i and $m_o = m_{i-1}$. This means that $w_o \le (1 + \delta_{j-1})^{-1}m_{i-1}$, where we used the definition of R_j . Therefore,

$$w_o = \frac{w_o}{c(v_1) + c(v_2)} \left(c(v_1) + c(v_2) \right) \le \frac{(1 + \delta_{j-1})^{-1} m_{i-1}}{2m_i} c(N_o) = \frac{1}{2(1 - \varepsilon)(1 + \delta_{j-1})} c(N_o),$$

where the second equality follows from $m_i/m_{i-1} = (1 - \varepsilon)$ for $i \in [L]$. This proves that Lemma 5 holds for $o \in O_d \cap R_j$ using that $(2(1 - \varepsilon))^{-1} \leq 1$ for $\varepsilon \in (0, 1/2)$.

Suppose that $o \in (O_{\leq L} \setminus (O_d \cup O_s)) \cap R_j$ and let $(o, e(N_o))$ be its local swap given by Lemma 2. Let $\ell \in [L]$ be the index such that $o \in I_\ell$. Since $o \notin O_s \cup O_d$, there exists an index $i < \ell$ such that $o \in O^{(i)} \setminus (O_d^{(i)} \cup O_s^{(i)})$. In particular, by Lemma 2, we have that $\emptyset \neq e(N_o) \subseteq A_i$. This implies that $m_o = m_{\ell-1} \leq m_i$ so $w_o \leq (1 + \delta_{j-1})^{-1} m_o \leq (1 + \delta_{j-1})^{-1} m_i < (1 + \delta_{j-1})^{-1} c(N_o)$.

Then by Lemma 4, we get the following result:

Corollary 4. Let $\varepsilon > 0$ be the parameter from Algorithm 2. For any $o \in O$, we have that

$$\mathbb{P}\left[w_o \ge (1+\delta_j)^{-1} \cdot m_o\right] \le \frac{\delta_j}{\varepsilon(1+\delta_j)} = \frac{\frac{\varepsilon_j}{m-\varepsilon_j}}{\varepsilon\left(1+\frac{\varepsilon_j}{m-\varepsilon_j}\right)} = \frac{j}{m}$$

B.1 Improved Approximation Ratio

Compared to the 9(k + 1)/10 approximation that we have shown so far, we obtain an improved approximation ratio arbitrarily close to $(k + 1)/(2 \ln(2))$.

Theorem 6. For any $\eta > 0$, there exists a choice of parameters $\varepsilon \in (0,1)$ and $\delta \in (0,1)$ for Algorithm 2 such that its output A satisfies:

$$w(O) \le \frac{k+1}{2\ln(2) - \eta} \cdot \mathbb{E}[w(A)].$$

Proof of Theorem 6. For any $\varepsilon > 0$, define $\delta \triangleq \frac{\varepsilon(1-\varepsilon)\eta}{-\ln(1-\varepsilon)}$. For each $o \in O$, let $(o, e(N_o))$ be the local swap defined by Lemma 2. Then, either $o \in O_s$ or $o \in (O_{\leq L} \setminus O_s) \cap R_j$ for some $j \in [m]$ or $o \in O \setminus O_{\leq L}$. In the first and last case, we have that $w_o \leq c(N_o)$. In the second case Lemma 5 shows that $(1 + \delta_{j-1})w_o \leq c(N_o)$. By Remark 2 and any outcome τ , we have that

$$w(O) = w(O_s) + \sum_{j=1}^{m} w((O_{\leq L} \setminus O_s) \cap R_j) + w(O \setminus O_{\leq L})$$

$$\leq \sum_{o \in O} c(N_o) - \sum_{j=1}^{m} \delta_{j-1} w((O_{\leq L} \setminus O_s) \cap R_j)$$

$$\leq kw(A) - \sum_{j=1}^{m} \delta_{j-1} w((O_{\leq L} \setminus O_s) \cap R_j),$$
(B.1)

where in the last inequality, we used that $\sum_{o \in O} c(N_o) \leq kw(A)$ by Corollary 1. Our proof continues by estimating the rightmost term. Let $\bar{O} \triangleq \{o \in O : w_o \geq (1 - \varepsilon)^{L-1} \cdot W\}$. Since $\delta_j \leq 1$ for $\varepsilon \in (0, 1/2)$, we have that:

$$\sum_{j=1}^{m} \delta_{j-1} w((O_{\leq L} \setminus O_s) \cap R_j) \ge \sum_{j=1}^{m} \delta_{j-1} w(\bar{O} \cap R_j) - w(O_s)$$
$$\ge \sum_{j=1}^{m} \delta_{j-1} w(\bar{O} \cap R_j) - w(A),$$
(B.2)

where we used Corollary 2 in the last inequality. Let $p_j = \mathbb{P}[o \in R_j]$ be the probability that $o \in R_j$. Note that this probability depends only on the outcome τ that determines the placement of the markers and is therefore equal for every $o \in \bar{O}$. Taking the expectation we get that, $\sum_{j=1}^{m} \delta_{j-1} \mathbb{E} \left[w(\bar{O} \cap R_j) \right] = w(\bar{O}) \cdot \sum_{j=1}^{m} p_j \cdot \delta_{j-1}$. We now lower bound the quantity $\sum_{j=1}^{m} p_j \cdot \delta_{j-1}$. By Corollary 4,

$$\sum_{i=j}^{m} p_i = 1 - \mathbb{P}\left[w_o \ge (1+\delta_{j-1})^{-1} m_o\right] \ge 1 - \frac{j-1}{m} \quad \text{for all } j \in [m].$$

We claim that $\sum_{j=1}^{m} p_j \cdot \delta_{j-1} \ge \sum_{j=1}^{m} \frac{\delta_{j-1}}{m}$ whenever (p_1, \ldots, p_m) satisfy $\sum_{i=j}^{m} p_i \ge 1 - \frac{j-1}{m}$ for all $j \in [m]$. Suppose otherwise that there exists some $p' \in [0, 1]^m$ that achieves the minimum value

of $\sum_{j=1}^{m} p_j \cdot \delta_{j-1}$ and that is not $(1/m, \dots, 1/m)$. Let ℓ be the largest index where $p'_{\ell} \neq 1/m$. Then $p'_{\ell} > 1/m$ otherwise $\sum_{i=\ell}^{m} p'_i < 1 - \frac{\ell-1}{m}$. Moreover, there exists an index q such that $p'_q < 1/m$ since otherwise the summed value cannot be less than $\sum_{j=1}^{m} \frac{\delta_{j-1}}{m}$. Let q be the largest such index and observe that $q < \ell$ as otherwise $\sum_{j=q}^{m} p'_j < 1 - \frac{q-1}{m}$ contradicting the constraint. Let p'' be the vector identical to p' except $p''_q = p'_q + \lambda$ and $p''_{\ell} = p'_{\ell} - \lambda$ for some small enough $\lambda > 0$. Then since $\delta_{\ell} \ge \delta_q$,

we have that summed values are strictly smaller for p'' and that p'' still satisfies $\sum_{i=j}^{m} p''_i \ge 1 - \frac{j-1}{m}$ for all $j \in [m]$. This is a contradiction to the minimality of p'. This implies that:

$$\sum_{j=1}^{m} p_{j} \delta_{j-1} \geq \sum_{j=1}^{m} \frac{\delta_{j-1}}{m}$$

$$= \frac{1}{m} \sum_{j=0}^{m-1} \frac{\varepsilon_{j}}{m - \varepsilon_{j}}$$

$$\geq \frac{1}{m} \int_{0}^{m-1} \frac{\varepsilon_{x}}{m - \varepsilon_{x}} dx$$

$$= \frac{-1}{m} \cdot \left[\frac{m \ln(m - \varepsilon_{x})}{\varepsilon} + x \right]_{0}^{m-1}$$

$$= \frac{-1}{m} \left(\frac{m}{\varepsilon} \ln \left(\frac{m - \varepsilon(m - 1)}{m} \right) + m - 1 \right)$$

$$= -\frac{\ln(1 - \varepsilon + \varepsilon/m)}{\varepsilon} - 1 + \frac{1}{m}$$
(B.4)

Taking the expectation of Equation (B.2) and substituting Equation (B.4) and Equation (B.2) into Equation (B.1), we obtain the following result

$$w(O) \le (k+1)\mathbb{E}\left[w(A)\right] - w(\bar{O}) \cdot \left(-\frac{\ln(1-\varepsilon+\varepsilon/m))}{\varepsilon} - 1 + \frac{1}{m}\right)$$
(B.5)

Similar to the proof of Proposition 1, observe that: $w(O\setminus\bar{O}) \leq |E| (1-\varepsilon)^{L-1}W \leq \frac{\delta}{1-\varepsilon}w(O)$. Thus, $w(\bar{O}) \geq (1-\frac{\delta}{1-\varepsilon})w(O)$. Substituting the above bound, using that $-\frac{\ln(1-x)}{x} \geq 1$ for all $x \in (0,1)$, and using the definition of δ , we get that, as $m \to \infty$,

$$\begin{split} w(O) + w(\bar{O}) \cdot \left(-\frac{\ln(1-\varepsilon)}{\varepsilon} - 1 \right) &\geq w(O) \left(1 + \left(1 - \frac{\delta}{1-\varepsilon} \right) \left(-\frac{\ln(1-\varepsilon)}{\varepsilon} - 1 \right) \right) \\ &\geq w(O) \left(\frac{-\ln(1-\varepsilon)}{\varepsilon} - \eta \right) \\ &\geq w(O) \left(\frac{-\ln(1-\varepsilon) - \eta}{\varepsilon} \right), \end{split}$$

Substituting the above bound in Equation (B.5), we get that:

$$w(O) \le \frac{\varepsilon}{-\ln(1-\varepsilon)-\eta}(k+1)\mathbb{E}[w(A)].$$

Taking the limit as $\varepsilon \to 1/2$ yields the desired result.

C Proof of Matroid Properties

In this section, we prove Theorem 2 and Theorem 3. Both theorems are based on the following known result about matroid exchanges:

Theorem 7. [Proposition 6, [Las15]] Let $\mathcal{M} = (V, \mathcal{I})$ be a matroid and A, B be bases of \mathcal{M} . Then, for any partition A_1, \ldots, A_m of A, there exists a partition B_1, \ldots, B_m of B such for each $i \in [m]$, we have:

$$A_i \cup (B \setminus B_i)$$
 is a basis of \mathcal{M} .

C.1 Proof of Theorem 2

We extend the previous Theorem 7 to handle the case where A and B aren't necessarily bases.

Theorem 2. Let $\mathcal{M} = (V, \mathcal{I})$ be a matroid and S, T be independent sets of \mathcal{M} with $|S| \leq |T|$. For any partition S_1, \ldots, S_m of S, there are disjoint sets $T_1, \ldots, T_m \subseteq T$ such that $|S_i| = |T_i|$ and

$$S_i \cup (T \setminus T_i) \in \mathcal{I}.$$

In particular, if |S| = |T|, then the sets T_i partition T.

Proof of Theorem 2. Suppose first that |S| = |T|. If S is a basis then we can apply Theorem 7. Suppose then, that S, and T aren't bases of \mathcal{M} . Let $\mathcal{M}' = \mathcal{M}_{|S\cup T}$ be the matroid restricted to $S \cup T$. More precisely, $\mathcal{M}' = \{A \subseteq S \cup T : A \in \mathcal{I}\}$. Let B be any basis of \mathcal{M}' . By matroid extension property, there exists $D \subseteq T \setminus S$ and $D' \subseteq S \setminus T$ such that $S \cup D$ and $T \cup D'$ are basis of \mathcal{M} . Let $\mathcal{M}'' = \{R \subseteq (S \cup T) \setminus (D \cup D') : R \cup (D \cup D') \in \mathcal{M}'\}$ be the matroid \mathcal{M}' contracted on $D \cup D'$. For $R \subseteq S \cup T$, the rank of R is equal to

$$\operatorname{rank}_{\mathcal{M}''}(R) = \operatorname{rank}_{\mathcal{M}'}(R \cup (D \cup D')) - \operatorname{rank}_{\mathcal{M}'}(D \cup D')$$

Let $n = \operatorname{rank}(\mathcal{M}')$, then $n = |S \cup D| = |T \cup D'|$, and let $m = \operatorname{rank}(\mathcal{M}'')$. Since $D \cup D' \subseteq T \cup D' \in \mathcal{I}$, we have that $\operatorname{rank}_{\mathcal{M}'}(D \cup D') = 2d$ where |D| = d. The equation above implies that m = n - 2d, and observe that $\operatorname{rank}_{\mathcal{M}''}(S) = \operatorname{rank}_{\mathcal{M}'}(S \cup D) - \operatorname{rank}_{\mathcal{M}'}(D \cup D') = n - 2d = m$. This computation shows that $S \setminus D'$ is a base of \mathcal{M}'' . An identical computation shows that $T \setminus D$ is a base of \mathcal{M}'' .

Constructing the swaps: We will apply Theorem 7 to $S \setminus (D \cup D')$ and $T \setminus (D \cup D')$ in \mathcal{M}'' . Before doing so, we define an arbitrary bijection $\pi : D' \to D$ between D' and D. This is well defined as |D| = |D'|. Using Theorem 7 with starting partition $S'_i = S_i \setminus D'$ of $S \setminus D'$, we find a partition

 $T'_1, \ldots, T'_m \subseteq T \setminus D$ such that $|S'_i| = |T'_i|$ and $((T \setminus D) \setminus T'_i) \cup S'_i \in \mathcal{I}''$. We extend these swaps to \mathcal{M}' and define $T_i = T'_i \cup \pi(S_i \cap D')$. Clearly, we have $|T_i| = |S_i|$. We prove that $(T \setminus T_i) \cup S_i \in \mathcal{I}$. Indeed,

$$T \setminus T_i \cup S_i = \left((T \setminus T_i \cup S_i) \cap (D \cup D') \right) \cup \left((T \setminus T_i \cup S_i) \setminus (D \cup D') \right).$$

Observe that the second term $((T \setminus T_i \cup S_i) \setminus (D \cup D')) \subseteq ((T \setminus D) \setminus T'_i) \cup S'_i \in \mathcal{I}''$ so is independent on the matroid contracted on $D \cup D'$. Additionally, we have that the first term $((T \setminus T_i \cup S_i) \cap (D \cup D')) \subseteq D \cup D'$. Combining both observations implies that: $T \setminus T_i \cup S_i \in \mathcal{I}'$ and so is independent in \mathcal{I} .

It remains to prove the theorem whenever $|S| \leq |T|$. Since T has larger size than S, there exists a set $T' \subseteq T \setminus S$ such that $S \cup (T \setminus T') \in \mathcal{I}$ and $|S \cup (T \setminus T')| = |T|$. In particular, |T'| = |S|. Moreover, both T' and S are independent in the matroid $\mathcal{M}' = (V, \mathcal{I}')$ contracted on $T \setminus T'$. By the first part of the statement, for any partition S_1, \ldots, S_m of S, there is a partition T'_1, \ldots, T'_m such that $|S_i| = |T'_i|$ and such that $(T' \setminus T'_i) \cup S_i \in \mathcal{I}'$. Therefore, we have that $(T \setminus T') \cup (T' \setminus T'_i) \cup S_i = T \setminus T'_i \cup S_i \in \mathcal{I}$. This concludes the proof.

C.2 Proof of Theorem 3

Theorem 3. Let $\mathcal{M} = (V, \mathcal{I})$ be a matroid and S, T be independent sets. Suppose that there is $N_S \subseteq T$ such that $|N_S| = |S|$ and $S \cup (T \setminus N_S) \in \mathcal{I}$. Then, for any partition of S into S_1, \ldots, S_m there is a partition $\{N_{S_i}\}_{i \in [m]}$ of N_S such that $|N_{S_i}| = |S_i|$ and $S_i \cup (T \setminus N_{S_i}) \in \mathcal{I}$.

Proof of Theorem 3. We contract \mathcal{M} on $T \setminus N_S$ and observe that both S and N_S are independent in the contracted matroid. By Theorem 2, we have that there exists a partition $\{N_{S_i}\}_{i \in [m]}$ such that $|N_{S_i}| = |S_i|$ and $N_S \setminus N_{S_i} \cup S_i$ is independent in the contracted matroid. In particular, we have that $T \setminus N_S \cup (N_S \setminus N_{S_i} \cup S_i) = T \setminus N_{S_i} \cup S_i \in \mathcal{I}$.

D Runtime and probability guarantee of Algorithm 2

D.1 Polynomial runtime

Let n = |E|. SLIDING LOCAL-SEARCH calls INTERVAL LOCAL-SEARCH L times. Hence, SLIDING LOCAL-SEARCH runs in $O(L \cdot |LS|)$ -time, where |LS| is the run time of INTERVAL LOCAL-SEARCH. The latter algorithm runs in polynomial time. Finding an (A, I, I)-swap uses $O(n^{O(k)})$ calls to the independence matroid oracle. Indeed, for all possible subsets of size 2, i.e., $O(n^2)$ of them, we verify the existence of a set of size 2k that we can discard by checking $O(n^{2k})$ possible sets. Although the number of improving (A, I, I)-swaps can be exponential, standard weight-scaling techniques [LSV09, LSV10, Ber00] can be used to reduce the number of iterations to ensure polynomial-time convergence and termination.

D.1.1 Weight rescaling

Suppose that A is a locally-optimal solution with respect to the improvement considered in Algorithm 1. Let $W \triangleq \max_{e \in E, v(e) \in \mathcal{I}} w_e$ be the maximum weight in our instance, let $\varepsilon > 0$ be an arbitrarily small constant. We apply Algorithm 2 on the instance where we rescale the weights of the instance as follows. For each $e \in E$, we define $\tilde{w}(e) = \lfloor Mw(e) \rfloor$, where $M \triangleq \frac{n}{\varepsilon W}$.

First, we prove that the weight-scaling doesn't affect the approximation guarantee of Algorithm 2. Let *A* be the solution output by Algorithm 2, let *O* be the optimal solution and let $\alpha > 0$ be the approximation factor, such that $\alpha w(A) \ge w(O)$. By definition of \tilde{w} , we have that $Mw(a) \ge \tilde{w}(a) \ge Mw(a) - 1$. Combining the above properties, we get that

$$\alpha Mw(A) \ge \alpha \tilde{w}(A) \ge \tilde{w}(O) \ge Mw(O) - |O| \ge Mw(O) - n$$

Dividing through by M, using that $M = \frac{n}{\varepsilon W}$ and that $W \leq w(O)$, we get that

$$\alpha w(A) \ge w(O) - \frac{n}{M} \ge w(O) - \varepsilon W \ge (1 - \varepsilon)w(O)$$

This proves that the weight-scaling doesn't affect the approximation guarantee of Algorithm 2. It remains to prove that the technique makes the local-search run in polynomial-time. Here, we use the fact that $\tilde{w}(e)$ is an integer. Indeed, the optimal solution has value at most $\tilde{w}(O) \leq$

 $|O| M \cdot W = \frac{n^2}{\varepsilon}$. Thus, there is at most $n^2 \varepsilon^{-1}$ improving (A, I, I)-swaps.

D.2 Last interval has negligible weight – Proof of Proposition 1

Proposition 1. Let $L \triangleq \lceil -\log_{1-\varepsilon}(|E|\delta^{-1})\rceil + 1$. Let O be the optimal solution and let A be the output of SLIDINGLS, and $O_{\leq L} = \{o \in O : w_o \geq m_L\}$. Then, $w(O \setminus O_{\leq L}) \leq \delta w(O)$.

Using that $L = \left\lceil -\log_{1-\varepsilon} \left(|E| \, \delta^{-1} \right) \right\rceil + 1$, we show that $w(O \setminus O_{\leq L})$ is small.

$$w(O \setminus O_{\leq L}) \leq |O \setminus O_{\leq L}| \cdot m_L \leq |E| (1 - \varepsilon)^{-\log_{1-\varepsilon}(|E|\delta^{-1})} (1 - \tau)W \leq \delta W \leq \delta w(O),$$

where we used that $w(O) \ge W$ since W is the largest weight of an independent edge and $(1-\tau) \le 1$ for all outcomes τ .

D.3 Probability of Algorithm 2 succeeding

We prove that SLIDING LOCAL-SEARCH succeeds with high-probability. Let *A* be the solution returned by SLIDINGLS, and let α be the approximation factor such that $\mathbb{E}[A] \ge (\alpha/k) \cdot w(O)$. We show that $w(A) \ge (\alpha/k) \cdot w(O)$ with high probability. This follows from a reverse Markov inequality.

Theorem 8. Let *X* be a random variable such that $\mathbb{P}[X \le a] = 1$ for some constant *a*. Then, for any $d < \mathbb{E}[X]$, the following holds:

$$\mathbb{P}\left[X > d\right] \ge \frac{\mathbb{E}\left[X\right] - d}{a - d}.$$

Applying Theorem 8 with a = w(O) and $d = \alpha(1 - \xi)/kw(O)$ for some small parameter ξ , we get that

$$\mathbb{P}\left[X > \alpha(1-\xi)/k \cdot w(O)\right] \ge \frac{\alpha\xi}{k - \alpha(1-\xi)}$$

Running $O(k/\xi)$ independent copies of SLIDING LOCAL-SEARCH and taking the best solution ensures that the expected guarantee of SLIDING LOCAL-SEARCH holds with probability arbitrarily close to 1.

E Reduction and Simplifying Assumptions

E.1 Reduction from *k*-Matroid Intersection to Matroid *k*-Parity

For completeness, we provide the reduction from k-Matroid Intersection to Matroid k-Parity as shown by Lee, Sviridenko and Vondrák [LSV10]. Let Π be an instance of k-Matroid Intersection. We are given k-matroid $\mathcal{M}_1 = (V, \mathcal{I}_1), \ldots, \mathcal{M}_k = (V, \mathcal{I}_k)$ on the same groundset V. We would like to construct an instance $\Pi' = (G = (V', E), \mathcal{M})$ of Matroid k-Parity.

The set of vertices V' consists of k copies of V, which we denote by V_1, \ldots, V_k . For each copy, we define the matroid $\mathcal{M}'_i = (V_i, \mathcal{I}_i)$ that is the matroid \mathcal{M}_i on the i^{th} copy of V. Our hypergraph G = (V', E) then contains n = |V| parallel (disjoint) hyperedges on the k copies of the same element from V. Observe that a collection of hyperedges such that, for all $i \in [k]$, the set of incident vertices in the i^{th} copy is independent defines an independent set in the intersection of k matroids and vice-versa.

It remains to prove that this property can be encoded as a single matroid. In fact, we simply define \mathcal{M} as the *union* of the matroid $\mathcal{M}'_1, \ldots, \mathcal{M}'_k$. More precisely, we let $\mathcal{M} = \mathcal{M}'_1 \vee \ldots \vee \mathcal{M}'_k$ defined on the groundset V' so that $\mathcal{I} = \{I_1 \cup \ldots \cup I_k : I_j \in \mathcal{I}_j \text{ for all } j \in [k]\}$. In contrast to matroid intersection, \mathcal{M} is in fact a matroid (See [S⁺03] for a proof).

E.2 Disjointness Assumption

In this section, we show that, given an instance of k-Matroid Parity $\Pi = (G = (V, E), \mathcal{M})$, we may assume that each vertex belongs to a unique hyperedge. This assumption is proved in [LSV10]. We recall the proof here. Let Π be our k-Matroid Parity instance. We create a novel instance $\Pi' = (G' = (V', E'), \mathcal{M}')$ equivalent to Π . For each vertex $v \in V$, create $n_v \in V'$ copies of v, where n_v is the degree of v in the hypergraph G. We replace each hyperedge in G by a collection of distinct copies of its elements, so that the new hyperedges are disjoint. It remains to define \mathcal{M}' . A set $S' \subseteq V'$ is independent in \mathcal{M}' if S contains at most 1 copy of each vertex V and the respective set $S \subseteq V$ is independent in \mathcal{M} . It is simple to check that \mathcal{M}' is a matroid.