

Optimality of Gerver's Sofa


Jineon Baek¹

December 2, 2024

¹Department of Mathematics, Yonsei University, Seoul, Korea. jineon@yonsei.ac.kr

Abstract

We resolve the *moving sofa problem* by showing that Gerver's construction with 18 curve sections attains the maximum area $2.2195\dots$.



Contents

Abstract	ii
1 Moving Sofa Problem	1
1.1 Introduction	1
1.2 Monotone Sofas and Caps	4
1.2.1 Monotone Sofa	5
1.2.2 Cap and Niche	7
1.3 Balancing Argument of Gerver	8
1.3.1 Balancing Argument	8
1.3.2 Logical Gap	10
1.4 Balanced Maximum Sofas and Caps	11
1.4.1 Limit of Maximum Polygon Caps	11
1.4.2 Balancedness of Maximum Polygon Cap	12
1.5 Rotation Angle of Balanced Maximum Sofas	13
1.5.1 Statement	13
1.5.2 Proof Outline	15
1.6 Surface Area Measure	16
1.7 Injectivity Condition	17
1.7.1 Statement	17
1.7.2 A Differential Inequality	18
1.7.3 Solving the Differential Inequality	20
1.8 Optimality of Gerver's Sofa	21
1.8.1 Definition of Q	21
1.8.2 Quadraticity of Q	23
1.8.3 Optimality of Q at Gerver's Sofa	24
2 Monotone Sofas and Caps	29
2.1 Planar Convex Body	29
2.2 Supporting Hallway	32
2.3 Monotone Sofa	34
2.4 Cap and Niche	37
2.5 Cap Contains Niche	38
3 Balanced Maximum Sofas and Caps	43
3.1 Simple Nef Polygon	43
3.2 Polygon Cap and Niche	45
3.3 Extensions of Polygon Cap Space	47

3.4	Maximum Polygon Cap	49
3.5	Balanced Maximum Sofa	54
4	Rotation Angle of Balanced Maximum Sofas	57
4.1	Horizontal Side Lengths	57
4.2	Right Rotation Angle	58
5	Surface Area Measure	63
5.1	Lebesgue–Stieltjes Measure	63
5.2	Differential Gauss–Minkowski Theorem	65
6	Injectivity Condition	67
6.1	Statement	67
6.2	Arm Lengths	68
6.3	Inequality on Maximum Polygon Caps	70
6.4	Inequality on Balanced Maximum Caps	74
6.5	Bounding Arm Lengths	77
7	Convex Domain and Convex Curves	80
7.1	Convex Domain	80
7.2	Curve Area Functional	82
7.3	Convex Curve	85
7.4	Mamikon’s Theorem	88
8	Optimality of Gerber’s Sofa	90
8.1	Domain of \mathcal{Q}	90
8.2	Definition of \mathcal{Q}	94
8.3	Concavity of \mathcal{Q}	97
8.4	Gerber’s Sofa	100
8.5	Directional Derivative of \mathcal{Q}	106
A	Table of Symbols	111

Chapter 1

Moving Sofa Problem

1.1 Introduction

Moving a large couch through a narrow hallway requires a well-planned pivoting. The *moving sofa problem* is asked in a two-dimensional idealization of such a situation:

What is the largest area α_{\max} of a connected planar shape that can move around the right-angled corner of a hallway with unit width?

Such a movable shape is called a *moving sofa* that we define precisely as below.

Definition 1.1.1. Define the *hallway* L as the union $L := H_L \cup V_L$ of its *horizontal side* $H_L := (-\infty, 1] \times [0, 1]$ and *vertical side* $V_L := [0, 1] \times (-\infty, 1]$.

Definition 1.1.2. A *moving sofa* S is any translation¹ of a nonempty, connected, and closed² subset of H_L that can be moved inside L by a continuous rigid motion to a subset of V_L .³

The moving sofa problem combines the two objectives of *motion planning* and *area maximization*. Despite numerous works on each subject, the problem has remained open since the initial publication by Leo Moser in 1966 [Mos66].

Definition 1.1.3. Denote the area (Borel measure) of a Borel measurable $X \subseteq \mathbb{R}^2$ as $|X|$.

The best bounds known so far on the maximum area α_{\max} of a moving sofa are summarized as

$$|G| = 2.2195 \dots \leq \alpha_{\max} \leq 2.37. \quad (1.1)$$

The lower bound comes from Gerver's sofa G of area $|G| = 2.2195 \dots$ constructed in 1992 [Ger92] (see Figure 1.1). The upper bound comes from a computer-assisted approach of Kallus and Romik in 2018 [KR18].

¹We allow arbitrary translation of a moving sofa S to locate it at any position we want, even outside the hallway L . Only a translation of S needs to be inside the horizontal side H_L , navigate its way inside L , and end at the vertical side V_L .

²Taking the closure of S does not hurt the movability.

³Recall that the *special Euclidean group* $\text{SE}(2)$ is the Lie group of all sign-preserving isometries of \mathbb{R}^2 . The movability can be stated formally as follows: there is a continuous curve $\Phi_t \in \text{SE}(2)$ parametrized by $t \in [0, 1]$, such that Φ_0 is a translation, $\Phi_0(S) \subseteq H_L$, $\Phi_t(S) \subseteq L$ for all $t \in [0, 1]$, and $\Phi_1(S) \subseteq V_L$.




Figure 1.1: Gerver's sofa G . The ticks denote the endpoints of 18 analytic curves and segments constituting the boundary of G [Rom18]. The supporting hallways L_t containing G are depicted as grey in the right side.

There were many evidences supporting that Gerver's sofa G attains the maximum area $\alpha_{\max} = |G|$. Gerver proved that a maximum-area moving sofa satisfies a certain local optimality condition (Theorem 1 of [Ger92]), and showed that his sofa G also satisfies the same condition (Theorem 2 of [Ger92]). Local optimality of G was further explored in [Rom18] and [Den24], and many numerical experiments also supported $\alpha_{\max} = |G|$ [Gib14; Bat22; Len+24].

We show that Gerver's sofa G indeed attains the maximum area. The proof does not require computer assistance, except for numerical computations that can be done on a scientific calculator.

Theorem 1.1.1. *Gerver's sofa G attains the maximum area α_{\max} of a moving sofa.*

The problem is difficult because there is no universal formula for the area that works for all possible moving sofas. To address this, we prove a property called the *injectivity condition* for a maximum-area moving sofa S_{\max} . For each moving sofa S satisfying the condition, we will define a larger shape R that resembles the shape of Gerver's sofa (Figure 1.2). The area $\mathcal{Q}(S)$ of R is then an upper bound of the area of S , and $\mathcal{Q}(S)$ matches the exact area of S if it is Gerver's sofa G . Injectivity condition of S ensures that the boundary of region R forms a Jordan curve, allowing us to compute $\mathcal{Q}(S)$ by using Green's theorem.




Figure 1.2: A moving sofa S (light yellow) is enclosed by a slightly larger region R (bold lines) of area $\mathcal{Q}(S)$ with a shape similar to Gerver's sofa. Three convex bodies K , B , and D represent different parts of R (bold and thin lines). K is a superset of R , and B , D are subsets of R .

The upper bound $\mathcal{Q}(S)$ of the area of a moving sofa S is then maximized with respect to S as follows. We use Brunn-Minkowski theory to express \mathcal{Q} as a quadratic functional on

the space \mathcal{L} of tuples (K, B, D) of convex bodies (Figure 1.2). We use Mamikon's theorem to establish the global concavity of \mathcal{Q} on \mathcal{L} (Figure 1.13). We use the local optimality equations on Gerver's sofa G by Romik [Rom18] to show that $S = G$ locally maximizes $\mathcal{Q}(S)$. Because \mathcal{Q} is concave, G also maximizes \mathcal{Q} globally. As the upper bound \mathcal{Q} matches the area at G , the sofa G also maximizes the area globally, establishing Theorem 1.1.1.

The full proof of Theorem 1.1.1 is divided into three main steps. Step 1 restricts the possible shapes of a maximum-area moving sofa S_{\max} . Step 2 establishes the injectivity condition for S_{\max} . Step 3 constructs the upper bound $\mathcal{Q}(S)$ for the area of a moving sofa S satisfying the injectivity condition, and maximizes $\mathcal{Q}(S)$ with respect to S .

1. Reduce the possible shapes of S_{\max} .
 - (a) S_{\max} is *monotone* (Section 1.2, Chapter 2).
 - (b) S_{\max} is *balanced* (Section 1.4, Chapter 3).
 - (c) S_{\max} have *rotation angle* $\pi/2$ (Section 1.5, Chapter 4).
2. Show that S_{\max} satisfies the *injectivity condition* (Section 1.7, Chapter 6).
3. Establish the upper bound \mathcal{Q} of sofa area with injectivity condition (Section 1.8, Chapter 8).
 - (a) Define the convex domain \mathcal{L} of \mathcal{Q} (Section 1.8.1, Section 8.1).
 - (b) Define a quadratic functional \mathcal{Q} on \mathcal{L} and show that it is an upper bound of sofa area (Section 1.8.2, Section 8.2).
 - (c) Show that \mathcal{Q} is concave on \mathcal{L} (Section 1.8.3, Section 8.3).
 - (d) Show that Gerver's sofa is a local (and thus global) optimum of \mathcal{Q} (Section 1.8.3, Section 8.5).

Step 1-(a) narrows down the possible shapes of S_{\max} to a *monotone sofa*, a convex body with a dent carved out by the inner corner of the supporting hallways (Figure 1.4). Step 1-(b) reprove an important local optimality condition by Gerver that the side lengths of S_{\max} should balance each other (Theorem 1.3.1). As the original proof by Gerver has a logical gap that does not address the connectedness of a moving sofa, we introduce new ideas and rework the proof carefully. Step 1-(c) uses previous steps and elementary geometry to show that S_{\max} rotates the full right angle in its movement.

Step 2 proves the *injectivity condition* on S_{\max} which is the key for establishing the upper bound \mathcal{Q} later. It states that the trajectory of the inner corner $(0, 0)$ of L does not make self-loops in the perspective (frame of reference) of the moving sofa (Figure 1.9). To prove this condition for S_{\max} , we establish a new differential inequality on S_{\max} (Equation (1.9)) heavily inspired by an ODE of Romik that balance the differential sides of Gerver's sofa (Equation (1.8)).

Step 3-(a) extends the space of all moving sofas S with injectivity condition to a collection \mathcal{L} of tuples (K, B, D) of convex bodies, so that each S maps to $(K, B, D) \in \mathcal{L}$ one-to-one (but not necessarily onto). The convex bodies describe different parts of the region R enclosing S (Figure 1.2). Step 3-(b) defines the upper bound \mathcal{Q} on the extended domain \mathcal{L} . We follow the boundary of R and express its area \mathcal{Q} using Green's theorem and the quadratic area expressions on K, B , and D from Brunn-Minkowski theory. We use injectivity condition and Jordan curve theorem to rigorously show that $\mathcal{Q}(K, B, D)$ is an upper bound of the area of S .

Step 3-(c) uses Mamikon’s theorem to establish the concavity of \mathcal{Q} on \mathcal{L} (Figure 1.13). Step 3-(d) calculates the directional derivative of \mathcal{Q} at the convex bodies $(K, B, D) \in \mathcal{L}$ arising from Gerber’s sofa G . The local optimality ODEs on G by Romik [Rom18] are used to show that the directional derivative is always non-positive. This implies that G is a local optimum of \mathcal{Q} in \mathcal{L} . The concavity of \mathcal{Q} on \mathcal{L} implies that G is also a global optimum of \mathcal{Q} in \mathcal{L} . As the value of \mathcal{Q} at G matches the area, the sofa G also globally maximizes the area, completing the proof of Theorem 1.1.1.

Chapter 2 to Chapter 8 provide the full details of the proof of Theorem 1.1.1. Given the large volume, Section 1.2 to Section 1.8 overviews each chapter and explains its motivation. Readers are strongly encouraged to start with the overview sections to understand the core idea hidden in the details.

The notations and definitions used in the overviews will be often simpler than the ones used in the full proof. That is, the definitions made in this Chapter 1 starting Section 1.2 are specific to this chapter alone. Starting from Chapter 2, all notations and definitions will be redefined for consistency in the detailed proofs. We always assume the plane with x - and y -coordinates, and the variables x and y are always associated with these coordinates.

Acknowledgements

The author thanks Dan Romik for his thorough support and encouragement that greatly helped the research process. His feedback on the presentation significantly improved the clarity of this work. His package `MovingSofas.nb`⁴ helped making the intricate details of the problem much more accessible to the author. The package was also used to generate figures of Gerber’s sofa in this work.

Acknowledgment is extended to Joseph Gerber and Thomas Hales for their interest in this work and their in-depth discussions. The author also appreciates David Speyer’s efforts in understanding the details and help in refining the presentation. The author thanks Michael Zieve and Joonkyung Lee for their mentorship and valuable advice.

Thanks are also due to Martin Strauss, Jeffrey Lagarias and Alexander Barvinok for their interest, help, and advice during the early stages of the research, as well as to Rolf Schneider for his suggestions on the proof of Theorem 5.2.2. The author acknowledges Hyunuk Nam, Seewoo Lee, Changki Yun, Jaemin Choi, Yeonghyeon Kim, Joonhyung Shin, Yugeun Shim, and Seungwon Park for their interest, discussions, and encouragement.

A prior version of the proof of Theorem 1.1.1 was computer-assisted. Although the software developed for this purpose⁵ does not appear in the final proof, it played an important role in shaping the intuition and strategy behind the full proof. The author thanks an anonymous mentor and Hyunuk Nam for their discussions that helped the development of the software.

This research was supported by the National Research Foundation of Korea (NRF) under grant MSIT NRF-2022R1C1C1010300. The author also acknowledges support from the Korea Foundation for Advanced Studies during the completion of this research.

1.2 Monotone Sofas and Caps

Summary: This section is an overview of Chapter 2. We show that a moving sofa S of maximum area can be assumed to be a *monotone sofa*, which is an intersection of the *supporting hallways* L_t of S (Section 1.2.1). A monotone sofa

⁴<https://www.math.ucdavis.edu/~romik/data/uploads/software/movingsofas-v1.3.nb>

⁵<https://github.com/jcpaik/sofa-designer>

S is equal to its *cap* $K := \mathcal{C}(S)$, a convex body, subtracted by the *niche* $\mathcal{N}(K)$ determined by cap K . Thus, the monotone sofa S can be identified with its cap K , and the moving sofa problem becomes the maximization of the *sofa area functional* $\mathcal{A}(K) = |K| - |\mathcal{N}(K)|$ with respect to the cap K (Section 1.2.2).

1.2.1 Monotone Sofa

A fundamental idea of Gerver [Ger92] is to see a moving sofa S as the intersection of rotating hallways. Look at the movement of S inside the hallway L in perspective of S . Then S is fixed in our frame of reference and L rotates and translates around S while containing S inside (bottom of Figure 1.3). So S is a common subset of the rotating hallways (right side of Figure 1.1).




Figure 1.3: The movement of a moving sofa in the perspective of hallway (top) and sofa (bottom).

We will make the details of this idea precise. First, define the angle ω that S rotates inside L .

Definition 1.2.1. The *rotation angle* ω of a moving sofa S is the *clockwise* angle that it rotates as it moves from H_L to V_L inside L .⁶

Define the unit-width strips H and V_ω .

Definition 1.2.2. Let $R_t : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote the rotation of \mathbb{R}^2 around the origin by the counterclockwise angle of $t \in \mathbb{R}$.

Definition 1.2.3. Define the horizontal strip $H := \mathbb{R} \times [0, 1]$, vertical strip $V := [0, 1] \times \mathbb{R}$, and its rotation V_ω around the origin by a counterclockwise angle $\omega \in \mathbb{R}$.

⁶This is the angular difference between the two rigid motions Φ_0 and Φ_1 sending S to H_L and V_L respectively.

Gerver showed that we can assume $\omega \in (0, \pi/2]$ for the moving sofa problem (see Theorem 1.5.1 for details). Let S be any moving sofa with rotation angle $\omega \in (0, \pi/2]$. Without loss of generality, we will always translate S and put it in the *standard position* defined as below. Recall that a line *supports* S if it contains a point of S but does not separate any two points of S .

Definition 1.2.4. A moving sofa S with rotation angle $\omega \in (0, \pi/2]$ is in *standard position* if the upper sides $y = 1$ of H and $x \cos \omega + y \sin \omega = 1$ of V_ω support S from above.

Proposition 1.2.1. *For any moving sofa S with rotation angle $\omega \in (0, \pi/2]$, there is a translation of S in standard position which is (i) unique if $\omega < \pi/2$, or (ii) unique up to horizontal translations if $\omega = \pi/2$.*

Proof. Observe that the lines $y = 1$ and $x \cos \omega + y \sin \omega = 1$ intersect properly if $\omega < \pi/2$, and overlaps if $\omega = \pi/2$. \square

A moving sofa S put in standard position is a common subset of H , V_ω and rotating hallways L_t parametrized by its counterclockwise angle $t \in [0, \omega]$.

Proposition 1.2.2. *Fix an arbitrary moving sofa S with rotation angle $\omega \in (0, \pi/2]$ in standard position. Then S is contained in each of the following sets.*

1. *The horizontal strip H .*
2. *For every angle $t \in [0, \omega]$, the rotating hallway L_t which is a translation of $R_t(L)$.*
3. *The rotated vertical strip $V_\omega = R_\omega(V)$.*

Proof. The initial position of S at L is contained in $H_L \subset H$. So the width of S measured along the y -axis is at most one. Because S is in standard position, the line $y = 1$ supports S from above and we have $S \subseteq H$.

The sofa S is rotated clockwise by ω after its movement in L . By the intermediate value theorem, for every $t \in [0, \omega]$ there is a moment in the movement where a copy of S is rotated clockwise by t inside L . See this in the frame of reference of S to conclude that $S \subset L_t$ for some translation L_t of $R_t(L)$.

The final position of S at L is contained in $V_L \subset V$. Look at this in the frame of reference of S . Then S is in a translation of V_ω , so the width of S measured along the direction $(\cos \omega, \sin \omega)$ is at most one. Because S is in standard position, the line $x \cos \omega + y \sin \omega = 1$ is a supporting line above S , and we have $S \subseteq V_\omega$. \square

By Proposition 1.2.2, any moving sofa S with rotation angle $\omega \in (0, \pi/2]$ in standard position is contained in the intersection

$$\mathcal{I} := H \cap V_\omega \cap \bigcap_{t \in [0, \omega]} L_t. \quad (1.2)$$

of two strips H , V_ω and the hallways L_t each rotated counterclockwise by $t \in [0, \omega]$ and translated. So we have $S \subseteq \mathcal{I}$, and it is natural to identify a maximum-area moving sofa S with the intersection \mathcal{I} and maximize \mathcal{I} by fixing H , V_ω and translating the hallways L_t for each $t \in [0, \omega]$. All known derivations of Gerver's sofa G [Ger92; Rom18; Den24] follow this approach.

However, recall that a moving sofa S is defined as a *connected* set (e.g. page 267 of [Ger92]). So the connectedness of \mathcal{I} in Equation (1.2) is necessary to identify a maximum-area S with the intersection \mathcal{I} . But it has not been rigorously established in the existing works that uses the idea $S = \mathcal{I}$ [Ger92; Rom18; KR18].⁷ Also, Proposition 1.2.2 does not yet imply that the hallways L_t should move continuously with respect to t .

(See the right side of Figure 1.4) To resolve these issues, we let each rotated hallway L_t in the Equation (1.2) be the *supporting hallway* of angle t making contact with S . We first give names to the different parts of L_t for further discussions.

Definition 1.2.5. (See Figure 2.2) Let L_t be the hallway rotated counterclockwise by $t \in [0, \omega]$ in Equation (1.2). Let $\mathbf{x}(t)$ be the *inner corner* of L_t corresponding to the point $(0, 0)$ of L . Let $\mathbf{y}(t)$ be the *outer corner* of L_t corresponding to the point $(1, 1)$ of L . Let $a(t)$ and $c(t)$ be the right and left *outer walls* of L_t respectively, corresponding to the walls $x = 1$ and $y = 1$ of L . Let $b(t)$ and $d(t)$ be the right and left *inner walls* of L_t respectively, corresponding to the walls $x = 0$ and $y = 0$ of L .

Starting from any hallway L_t of counterclockwise angle t containing S , the *supporting hallway* is obtained by pushing L_t in the directions of $-(\cos t, \sin t)$ and $-(-\sin t, \cos t)$ continuously, until the two outer walls $a(t)$ and $c(t)$ of L_t makes contact with S . As this move only pulls the inner walls $b(t)$ and $d(t)$ of L_t away from S , the new supporting hallway L_t still contains S and now moves continuously with respect to t .

After letting each L_t be the supporting hallways of S , the intersection \mathcal{I} in Equation (1.2) is now completely determined by S , so that we will denote it as $\mathcal{I}(S)$. We show that this $\mathcal{I}(S)$ is always connected for any moving sofa S of rotation angle $\omega \in (0, \pi/2]$ (Theorem 2.3.6). By looking at $\mathcal{I}(S) \subseteq L_t$ in the frame of reference of L_t , the intersection $\mathcal{I}(S)$ also admits a continuous movement inside L . So $\mathcal{I}(S)$ is a moving sofa containing S (Theorem 2.3.2).

Define a *monotone sofa* as the intersection $\mathcal{I}(S)$ of supporting hallways arising from some moving sofa S . Then we can always assume that a maximum-area sofa S is monotone by taking the intersection $\mathcal{I}(S)$ and making it larger. In particular, Gerver's sofa G is a monotone sofa because G is the intersection of supporting hallways (Figure 1.1). We also show that for any monotone sofa S , taking the intersection again does not enlarge the set and $S = \mathcal{I}(S)$ itself is the intersection of supporting hallways L_t of S (Theorem 2.4.4),

1.2.2 Cap and Niche

Let S be a monotone sofa with rotation angle $\omega \in (0, \pi/2]$. The outer walls $a(t)$ and $c(t)$ of the supporting hallways L_t of S form the supporting lines of a convex body $K := \mathcal{C}(S)$ that we call the *cap* of S . Define the *parallelogram* $P_\omega := H \cap V_\omega$. Then the cap $K = \mathcal{C}(S)$ is

$$\mathcal{C}(S) := P_\omega \cap \bigcap_{t \in [0, \omega]} Q_t^+ \quad (1.3)$$

where Q_t^+ is the closed convex cone with vertex $\mathbf{y}(t)$ bounded from above by the outer walls $a(t)$, $c(t)$ of L_t . Because S was in standard position (Definition 1.2.4), the cap K is inscribed in the parallelogram P_ω and makes contact with all four sides of P_ω ((1) of Definition 2.4.1).

⁷Gerver requires a moving sofa S to be connected (Page 267 of [Ger92]). The proof of Theorem 1 in [Ger92] then defines a subcollection \mathcal{T} of intersections \mathcal{I} in Equation (1.2) and uses compactness to find a set $T \in \mathcal{T}$ of maximum area. However, Gerver does not show in his proof that the set T should be connected, which is a logical gap not trivial to fix. In [Rom18], Romik assumes the equality $S = \mathcal{I}$ (Equation 8, p319) to give a streamlined derivation of Gerver's sofa, but does not rigorously prove $S = \mathcal{I}$ for a maximum-area S . In [KR18], Kallus and Romik require S to be connected and choose the largest-area connected component S of \mathcal{I} , allowing the possibility of $S \neq \mathcal{I}$.




Figure 1.4: The movement of a monotone sofa S with rotation angle $\omega = \pi/2$ in perspective of the hallway (left) and the sofa (right).

The monotone sofa S is obtained from the cap K by subtracting the *niche* $\mathcal{N}(K)$ of cap K , the union of all the triangular regions carved out by the inner walls $b(t)$, $d(t)$ of L_t . Explicitly, define the *fan*

$$F_\omega := \{(x, y) : y \geq 0, x \cos \omega + y \sin \omega \geq 0\}$$

bounded from below by the bottom sides of the parallelogram P_ω . Then the niche $\mathcal{N}(K)$ is

$$\mathcal{N}(K) = F_\omega \cap \bigcup_{t \in [0, \omega]} Q_t^- \quad (1.4)$$

where Q_t^- is the open convex cone with vertex $\mathbf{x}(t)$ bounded from above by the inner walls $b(t)$ and $d(t)$ of L_t . We can derive $S = K \setminus \mathcal{N}(K)$ from the equality $L_t = Q_t^+ \setminus Q_t^-$ (Theorem 2.4.2). Note that L_t and Q_t^- can be recovered from the supporting lines of cap K (Lemma 2.3.5), so the niche $\mathcal{N}(K)$ is indeed determined by K .

Because a monotone sofa $S = K \setminus \mathcal{N}(K)$ is completely determined by its cap $K := \mathcal{C}(S)$, we will identify S with its cap K . We will prove $\mathcal{N}(K) \subset K$ using elementary geometry (Theorem 2.5.9). Then the area $|K| - |\mathcal{N}(K)|$ of S can be understood in terms of the cap and niche separately. We will define \mathcal{K}_ω^c as the space of all caps with rotation angle $\omega \in (0, \pi/2]$. Now the moving sofa problem becomes the maximization of the *sofa area functional* $\mathcal{A}_\omega(K) := |K| - |\mathcal{N}(K)|$ on $K \in \mathcal{K}_\omega^c$.

1.3 Balancing Argument of Gerber

Summary: This section reviews an important theorem of Gerber, stating that there is a maximum-area moving sofa which is a limit of polygons with opposite sides of the same length (Section 1.3.1). We argue that the balancing argument of Gerber, while holds the essence of the proof, has a subtle logical gap that does not take account of the connectedness of a moving sofa (Section 1.3.2).

1.3.1 Balancing Argument

Call a polygon P *balanced* if, for any two parallel lines l^+ and l^- of distance one on the plane, the total length of all edges of P in one line l^+ is equal to that of the other line l^- . Theorem

1 in [Ger92] by Gerver states that there exist a maximum-area moving sofa S_ω that can be approximated sufficiently close by balanced polygons S_Θ .

We copy the full statement of the theorem as appears exactly in Gerver's paper [Ger92] (footnote ours). We will rephrase the theorem in our words, so the reader may skim it for first read.

Theorem 1.3.1. (Theorem 1 in [Ger92]) *There exists a real number γ , $\pi/3 \leq \gamma \leq \pi/2$, and a region S , such that S can move around the corner of H ,⁸ rotating through an angle of $-\gamma$ in the process,⁹ such that no region of greater area can move around the corner, and such that for arbitrarily large n , S can be approximated arbitrarily closely by a polygonal region P_n with the following properties:¹⁰ The boundary of P_n is a balanced polygon. P_n is the intersection of $n+1$ sets H_α (where $\alpha = k\gamma/n$ and $0 \leq k \leq n$). H_0 is the half-strip¹¹ $x \leq 1, 0 \leq y \leq 1$. H_γ is a translation of the half strip¹² $y \leq 1, 0 \leq x \leq 1$ rotated by angle γ . For $0 < \alpha < \gamma$, H_α is a translation of H rotated by angle¹³ γ .*

We now explain the Theorem 1.3.1 and its proof by Gerver in our words. Fix the rotation angle $\omega \in (0, \pi/2)$. As described in Section 1.2, a maximum-area moving sofa S is the connected intersection

$$\mathcal{I} := H \cap V_\omega \cap \bigcap_{t \in [0, \omega]} L_t$$

of two unit-width strips H, V_ω and hallways L_t of counterclockwise angle t . Discretize the problem by taking a finite nonempty subset Θ of $(0, \omega)$ and the polygon intersection

$$S_\Theta := H \cap V_\omega \cap \bigcap_{t \in \Theta} L_t \quad (1.5)$$

instead. The approximated problem now is to maximize the area of S_Θ by translating the hallways L_t each rotated counterclockwise by $t \in \Theta$.

(See Figure 1.5) Gerver's main idea in [Ger92] is that each maximum-area polygon S_Θ in Equation (1.5) should be balanced. Theorem 1.3.1 states that, as $n \rightarrow \infty$ and the angle set $\Theta = \Theta_n := \{i\omega/n : 1 \leq i < n\}$ gets denser in $[0, \omega]$, the balanced polygons S_Θ should converge to some maximum-area moving sofa S_ω . For the proof of Theorem 1.3.1, Gerver uses the following *balancing argument* to show that each S_Θ is indeed balanced,¹⁴ and use compactness to show that such S_Θ 's converge to some maximum-area sofa S_ω .

Balancing Argument: Assume for the sake of contradiction that a maximum-area polygon S_Θ in Equation (1.5) is not balanced. Take any pair of two parallel lines l^+ and l^- of distance one, so that the total side lengths s^+ and s^- of S_Θ respectively on the lines l^+ and l^- are not equal. Then all sides of S_Θ on l^\pm are contributed by exactly one of $X = H, V_\omega$ or L_t . Let $\pm v$ be the normal unit vectors of parallel lines l^\pm respectively, directing outwards from each other. If $s^+ > s^-$

⁸This H in [Ger92] is the hallway L in our paper.

⁹This γ in [Ger92] is the rotation angle ω in our paper. His proof of the bound $\pi/3 \leq \gamma \leq \pi/2$ is factored out separately as Theorem 1.5.1.

¹⁰This P_n in [Ger92] is the polygon S_{Θ_n} in our description (Equation (1.5)).

¹¹This H_0 in [Ger92] is the horizontal side H_L of L in our paper.

¹²This H_γ in [Ger92] is the vertical side V_L of L in our paper rotated counterclockwise by γ .

¹³This H_α in [Ger92] is the rotating hallway L_α containing S in our paper. The proof of Theorem 1 in [Ger92] actually takes each L_α as the supporting hallway of angle α , using the support functions $p(\alpha)$ and $q(\alpha)$ of S .

¹⁴This balancing argument on S_Θ (or P_n in [Ger92]) is done in the second paragraph of page 273 in [Ger92].

(resp. $s^- > s^+$), translate X slightly by ϵv (resp. $-\epsilon v$) for sufficiently small $\epsilon > 0$. If we pushed either $X = H$ or V_ω , translate the whole S_Θ with H, V_ω, L_t together, to put H and V_ω back to their initial positions. We just increased the area of S_Θ by $\epsilon|s^+ - s^-| + o(\epsilon) > 0$ by translating the hallways L_t , contradicting the maximality of S_Θ .




Figure 1.5: A maximum-area polygon intersection S_Θ should have balanced side lengths (left). By taking the angle set Θ denser in $[0, \omega]$, the polygon S_Θ converges to a maximum-area monotone sofa with balanced side lengths (right).

1.3.2 Logical Gap

The balancing argument of Gerber, while holds great importance and contains the gist of the proof of Theorem 1.3.1, has a subtle logical gap that does not address the connectedness of moving sofas.

In the first paragraph of [Ger92], he defines a moving sofa as a *connected* planar region. However, neither the connectedness of the polygons S_Θ , nor the limiting shape S_ω of S_Θ are established in the proof of Theorem 1 in [Ger92].¹⁵ To fill this gap in Gerber's proof, it is natural to simply assume that each maximum-area polygon S_Θ is taken *among* connected intersections.¹⁶ However, this will not work because the balancing argument on S_Θ may break the connectedness of S_Θ . See the following example.

(See Figure 1.6) Take the angle set $\Theta = \{\pi/6, \pi/3\}$ and rotation angle $\omega = \pi/2$. Define the unit vector $\mathbf{u} := (\cos \pi/6, \sin \pi/6)$. Take a sufficiently small positive real number $c > 0$. Take the hallways $L_{\pi/6}, L_{\pi/3}$ with angles in Θ and inner corners $\mathbf{x}(\pi/6) = (0, 1) - c\mathbf{u}$, $\mathbf{x}(\pi/3) = (-0.9, 0.98)$ respectively. The intersection S_Θ in Equation (1.5) is not balanced, as the side of S_Θ with normal angle \mathbf{u} is larger than the side with opposite normal angle $-\mathbf{u}$ for all $c \geq 0$ (depicted green). The balancing argument will now push $L_{\pi/6}$ in the positive direction of \mathbf{u} , decreasing c as long as $c \geq 0$. But as c becomes negative, the intersection S_Θ becomes disconnected.

Thus, while the balancing argument of Gerber can guarantee the balancedness of a maximum-area S_Θ , it cannot guarantee the connectedness of S_Θ . In the example above, it is actually possible to preserve the connectedness of S_Θ by carefully choosing another pair of edges to balance. However, such an extra consideration is not also made in [Ger92]. The next Section 1.4 provides a strategy that circumvents this issue.

¹⁵In comparison, a lot of work in Chapter 2 and Chapter 3 are done to ensure the connectedness of the intersection \mathcal{I} or S_Θ that we find.

¹⁶The other option is to allow each maximum-area polygon S_Θ to be disconnected, but then proving that its limit S_ω is connected would require completely new ideas.




Figure 1.6: Balancing argument breaks the connectivity of a polygon intersection S_Θ .

1.4 Balanced Maximum Sofas and Caps

Summary: This section is an overview of Chapter 3. We rework the proof of Theorem 1.3.1 by Gerver, taking account of the connectedness of moving sofas. We show the existence of a *balanced maximum sofa*, a monotone sofa of the maximum area that can be approximated sufficiently close by balanced polygons.

1.4.1 Limit of Maximum Polygon Caps

Our goal now is to bridge the gap discussed in Section 1.3.2 and show that the *connected* polygon intersection

$$S_\Theta := H \cap V_\omega \cap \bigcap_{t \in \Theta} L_t \tag{1.6}$$

of maximum area is balanced. Recall that the strips H and V_ω are fixed, and each hallway L_t of counterclockwise angle $t \in \Theta$ can translate freely.

We will first write the polygon $S_\Theta = K \setminus \mathcal{N}_\Theta(K)$ as the difference of the *polygon cap* $K := \mathcal{C}_\Theta(K)$ and the *polygon niche* $\mathcal{N}_\Theta(K)$, analogous to the cap K and niche $\mathcal{N}(K)$ of a monotone sofa S in Section 1.2. Explicitly, the polygon cap K is defined as

$$\mathcal{C}_\Theta(K) := P_\omega \cap \bigcap_{t \in \Theta} Q_t^+$$

following the Equation (1.3) of caps, and the polygon niche $\mathcal{N}_\Theta(K)$ is defined as

$$\mathcal{N}_\Theta(K) := F_\omega \cap \bigcup_{t \in \Theta} Q_t^-$$

following the Equation (1.4) of niche. From $L_t = Q_t^+ \setminus Q_t^-$, we can also obtain $S_\Theta = K \setminus \mathcal{N}_\Theta(K)$ back.

Instead of maximizing S_Θ directly, we will maximize the *polygon area functional* $\mathcal{A}_\Theta(K) := |K| - |\mathcal{N}_\Theta(K)|$ with respect to the polygon cap K , where we allow the polygon sofa $S_\Theta = K \setminus \mathcal{N}_\Theta(K)$ to be disconnected. For an example, we allow the case $c = -0.05$ in Figure 1.6 where $\mathcal{N}_\Theta(K) \not\subset K$. Call such a maximizer K_Θ of $\mathcal{A}_\Theta(K)$ a *maximum polygon cap*.

We will show in Section 3.4 that the side lengths of maximum polygon cap K_Θ and niche $\mathcal{N}_\Theta(K_\Theta)$ balance each other (Theorem 3.4.9) and that $\mathcal{N}_\Theta(K_\Theta) \subset K_\Theta$ (Theorem 3.4.10). This is the technical part of the proof that we outline in the next Section 1.4.2. By taking the angle set Θ denser in $[0, \omega]$, the maximum polygon caps K_Θ converge to some cap K_ω with rotation angle ω that we call as the *balanced maximum cap* (Definition 3.5.2).

As the maximum polygon caps K_Θ converge to a balanced maximum cap K_ω , that $\mathcal{N}_\Theta(K_\Theta) \subset K_\Theta$ implies $\mathcal{N}(K_\omega) \subseteq K_\omega$ too, so that the set $S_\omega := K_\omega \setminus \mathcal{N}(K_\omega)$ is connected and forms a monotone sofa.¹⁷ We call such S_ω a *balanced maximum sofa* (Definition 3.5.3). As each K_Θ is a maximizer of \mathcal{A}_Θ , the limit K_ω is also a maximizer of \mathcal{A}_ω , and the area $\mathcal{A}_\omega(K_\omega) = |K_\omega| - |\mathcal{N}(K_\omega)|$ of a balanced maximum sofa S_ω achieves the maximum area among all monotone sofas of rotation angle ω .

1.4.2 Balancedness of Maximum Polygon Cap

Now we overview the technical proof that the side lengths of a maximum polygon cap K_Θ and its polygon niche $\mathcal{N}_\Theta(K_\Theta)$ are balanced. That is, for any unit vector v , the total length of all sides of K_Θ and $\mathcal{N}_\Theta(K_\Theta)$ with normal angle v is equal to that of normal angle $-v$ (Definition 3.4.5; see Figure 3.1). We will also obtain $\mathcal{N}_\Theta(K_\Theta) \subset K_\Theta$ as a consequence. We omit many details that can be found in the full Chapter 3.

We extend the space \mathcal{K}_Θ^c of all polygon caps K with angle set Θ using the *support function* of K .

Definition 1.4.1. Define $u_t := (\cos t, \sin t)$ and $v_t := (-\sin t, \cos t)$.

Definition 1.4.2. For any planar convex body K (a compact, convex subset of \mathbb{R}^2), define the *support function* $h_K(t) := \sup \{u_t \cdot p : p \in K\}$.

The support function $h_K(t)$ of K is the signed distance from the origin $(0, 0)$ to the supporting line of K with normal vector u_t outwards from K . Let $\Theta^\circ = \Theta \cup (\Theta + \pi/2) \cup \{\omega, \pi/2\}$. We embed the space \mathcal{K}_Θ^c of all polygon caps K with the angle set Θ to the space \mathcal{H}_Θ of all functions $h : \Theta^\circ \rightarrow \mathbb{R}$ by taking the support function h_K and restricting it to Θ° . This embedding allows to see \mathcal{H}_Θ as an extension of \mathcal{K}_Θ^c .

We will extend the polygon area functional $\mathcal{A}_\Theta(K)$ on $K \in \mathcal{K}_\Theta^c$ to the larger space $h \in \mathcal{H}_\Theta$. To do so, we write the cap K and niche $\mathcal{N}_\Theta(K)$ as the *Nef polygons* obtained from boolean set operations on half-planes. For any $t \in S^1$ and $h \in \mathbb{R}$, define the closed half-planes $H_\pm(t, h)$ and the open half-planes $H_\pm^\circ(t, h)$ with the boundary $l(t, h)$ as the following.

$$\begin{aligned} H_-(t, h) &:= \{p \in \mathbb{R}^2 : p \cdot u_t \leq h\} & H_-^\circ(t, h) &:= \{p \in \mathbb{R}^2 : p \cdot u_t < h\} \\ H_+(t, h) &:= \{p \in \mathbb{R}^2 : p \cdot u_t \geq h\} & H_+^\circ(t, h) &:= \{p \in \mathbb{R}^2 : p \cdot u_t > h\} \end{aligned}$$

¹⁷Theorem 2.5.9 shows that for any cap K , we have $\mathcal{N}(K) \subset K$ if and only if the set $K \setminus \mathcal{N}(K)$ is connected.

Let $h := h_K$ be the support function of K . Then we can then write the cap

$$K = \bigcap_{t \in \{\omega, \pi/2\}} (H_-(t, h(t)) \cap H_+(t, h(t) - 1)) \cap \bigcap_{t \in \Theta \cup (\Theta + \pi/2)} H_-(t, h(t))$$

and and the niche

$$\begin{aligned} \mathcal{N}_\Theta(K) = & \bigcap_{t \in \{\omega, \pi/2\}} H_+(t, h(t) - 1) \cap \\ & \bigcup_{t \in \Theta} (H_-^\circ(t, h(t) - 1) \cap H_-^\circ(t + \pi/2, h(t + \pi/2) - 1)) \end{aligned}$$

purely as boolean operations on the half-planes $H_\pm(t, h(t))$ and $H_\pm(t, h(t) - 1)$ determined by $h = h_K$ (Definition 3.3.3). This amounts to saying that K and $\mathcal{N}_\Theta(K)$ are the *Nef polygons* determined by such half-planes. The Nef polygon formulas $\mathcal{C}_\Theta(h)$ and $\mathcal{N}_\Theta(h)$ of K and $\mathcal{N}_\Theta(K)$ respectively in $h = h_K$ generalizes to all $h \in \mathcal{H}_\Theta$. Now the polygon area $\mathcal{A}_\Theta(K)$ extends to

$$\mathcal{A}_\Theta(h) := |\mathcal{C}_\Theta(h)| - |\mathcal{N}_\Theta(h)|$$

over all $h \in \mathcal{H}_\Theta$.

To prove that a maximum polygon cap K_Θ is balanced, we use the method of contradiction and assume that K_Θ is not balanced. Let $h := h_{K_\Theta}$ so that $\mathcal{A}_\Theta(K_\Theta) = \mathcal{A}_\Theta(h)$. Lemma 3.4.6 carefully chooses the angle $t \in \Theta^\circ$ so that the balancing move on unbalanced sides of $K_\Theta = \mathcal{C}_\Theta(h)$ and $\mathcal{N}_\Theta(K_\Theta) = \mathcal{N}_\Theta(h)$ always moves a hallway in a *positive* direction of u_t . So the move increases the value of $h(t)$ by a sufficiently small $\epsilon > 0$ and makes $\mathcal{A}_\Theta(h)$ slightly larger. Let $h^+ \in \mathcal{H}_\Theta$ be the incremented function so that $\mathcal{A}_\Theta(h^+) > \mathcal{A}_\Theta(h)$.

Our way of choosing the angle t guarantees that the convex polygon $K^+ := \mathcal{C}_\Theta(h^+)$ obtained back from h^+ is always a *translation* of some polygon cap $K^0 \in \mathcal{K}_\Theta^c$ (Lemma 3.4.8). By translating K^+ back to K^0 , we conclude $\mathcal{A}_\Theta(h^+) = \mathcal{A}_\Theta(K^0)$ and thus

$$\mathcal{A}_\Theta(K_\Theta) = \mathcal{A}_\Theta(h) < \mathcal{A}_\Theta(h^+) = \mathcal{A}_\Theta(K^0),$$

reaching contradiction with the maximality of K_Θ . This is the sketch of the rigorous proof of Theorem 3.4.9.

We can then use the balancedness of K_Θ and $\mathcal{N}_\Theta(K_\Theta)$ to show that $\mathcal{N}_\Theta(K_\Theta) \subset K_\Theta$. See Figure 3.1. Balancedness essentially implies that the *polyline* \mathbf{p}_{K_Θ} obtained from the bottom sides of $S_\Theta = K_\Theta \setminus \mathcal{N}_\Theta(K_\Theta)$ is a ‘permutation’ of the upper sides of polygon cap K_Θ . This implies that the polyline \mathbf{p}_{K_Θ} should be contained inside K_Θ , so that $\mathcal{N}_\Theta(K_\Theta) \subset K_\Theta$. This is the essential idea behind Theorem 3.4.10 that rigorously proves $\mathcal{N}_\Theta(K_\Theta) \subset K_\Theta$.

1.5 Rotation Angle of Balanced Maximum Sofas

Summary: This section is an overview of Chapter 4. We show that the balanced maximum sofa found in the previous step admits a movement with rotation angle $\pi/2$.

1.5.1 Statement

Recall that the *rotation angle* ω of a moving sofa S is the clockwise angle that S rotates during its movement inside L (Definition 2.3.3). It can be strictly less than the angle $\pi/2$ of

the hallway L . For example, the square $S := [0, 1]^2$ have rotation angle $\omega = 0$ as it can be moved inside L by only translation.

Gerver showed that there is a maximum-area moving sofa S_{\max} with rotation angle $\pi/3 \leq \omega \leq \pi/2$ (Theorem 1.3.1). His argument, reproduced in the Theorem 1.5.1 below, actually proves a slightly improved lower bound $\omega \geq \sec^{-1}(2.2) = 62.96 \dots^\circ$.

Theorem 1.5.1. (Modification of page 271 of [Ger92]) *Let S be any moving sofa of area ≥ 2.2 . Then S admits a movement in L with rotation angle $\omega \in [\sec^{-1}(2.2), \pi/2]$.*

Proof. Assume any movement of S inside L with rotation angle $\omega \in \mathbb{R}$. Without loss of generality, we can assume that S is in its initial position at $H_L \subseteq H$.

First assume $\omega \leq -\pi/4$. By the intermediate value theorem, there is a moment where S is rotated clockwise by $-\pi/4 \in [\omega, 0]$ (or, counterclockwise by $\pi/4$) inside L during its assumed movement. Looking at this in the perspective of S , the sofa S is contained in a hallway L' rotated clockwise by $\pi/4$ and translated. The intersection $H \cap L'$ containing S have area $\sqrt{2} = 1.4142 \dots$ and we get contradiction as $|S| \geq 2.2$.

Now assume $|\omega| < \sec^{-1}(2.2)$. The sofa S is rotated clockwise by ω in its final position at $V_L \subseteq V$. Look at this in perspective of S , then S is contained in a translation of V_ω . The intersection of H and a translation of V_ω is a parallelogram of area $\sec(\omega) < 2.2$. This contradicts $|S| \geq 2.2$.

So we should have $\sec^{-1}(2.2) \leq \omega$ because $\sec^{-1}(2.2) = 62.96 \dots^\circ > \pi/4$. We finish the proof by assuming $\omega > \pi/2$ and finding another movement of S in L with rotation angle $\pi/2$. By the intermediate value theorem, there is a moment in the movement of S with rotation angle ω , where S is rotated clockwise by $\pi/2 \in [0, \omega]$ in L . Call the position of S at this moment $S_{\pi/2}$. Instead of following the rest of the movement of S , translate $S_{\pi/2}$ horizontally in the positive direction of the x -axis until it makes contact with the outer wall $x = 1$ of L . Since S was initially in H_L , the width of $S_{\pi/2}$ measured along the x -axis is at most one. So after the horizontal translation, $S_{\pi/2}$ will lie completely inside the destination V_L , finishing a full moment of S with rotation angle $\pi/2$. \square

By Theorem 1.5.1 and Gerver's sofa of area $|G| = 2.2195 \dots > 2.2$, we can assume the rotation angle $\omega \in [\sec^{-1}(2.2), \pi/2]$ of a maximum-area moving sofa. Chapter 4 proves the equality $\omega = \pi/2$ for balanced maximum sofas.

Theorem 1.5.2. *Let S_ω be an arbitrary balanced maximum sofa with area ≥ 2.2 and rotation angle $\omega \in [\sec^{-1}(2.2), \pi/2]$. Then a rotated copy of S_ω admits a movement inside L with rotation angle $\omega = \pi/2$.*

To prove Theorem 1.5.2, we will show that (a rotated copy of) S_ω can rotate an extra angle of $\pi/2 - \omega$ inside H_L before its movement with angle ω .

The main step is to show that a triangular region Δ_ω is disjoint from S_ω (Theorem 4.2.5). Recall that the cap of the monotone sofa S_ω is inscribed in the parallelogram P_ω of width 1 with the lower-left corner $O := (0, 0)$ and upper-right corner $o_\omega := (\tan(\pi/4 - \omega/2), 1)$, making an angle of $\omega + \pi/2$ at both corners (see Equation (1.3) and the left of Figure 1.7). The region Δ_ω is then defined as the triangular region near O formed by three vertices O , $o_\omega - (0, 1)$, and $o_\omega - (\cos \omega, \sin \omega)$.

Once we show the main step that $S_\omega \subseteq P_\omega$ is disjoint from Δ_ω , we obtain enough room to rotate S_ω counterclockwise by an angle of $\pi/2 - \omega$ inside the horizontal side H_L (see the right of Figure 1.7). Follow this rotation of S_ω in reverse so that it rotates clockwise by $\pi/2 - \omega$. Then follow the original movement of S_ω in L with rotation angle ω . We have just found the movement of a rotated copy of S_ω with full rotation angle $\pi/2$, proving Theorem 1.5.2.




Figure 1.7: The moving sofa S_ω of maximum area with a fixed rotation angle ω is inscribed in the parallelogram P_ω and disjoint from the triangular region Δ_ω (left). So it can rotate counterclockwise by the angle of $\pi/2 - \omega$ inside the horizontal side H_L (right).

1.5.2 Proof Outline

We now outline the proof of the main step that S_ω is disjoint from Δ_ω (Theorem 4.2.5). We use the balancedness of S_ω established in Section 1.4. In particular, the horizontal sides of S_ω should be equal in their length (the blue sides of Figure 1.5 and Figure 1.8).




Figure 1.8: Proof of Theorem 1.5.2. We find two points $q_0, q_1 \in S_\omega$ sufficiently away from the origin. Then we take a supporting hallway L_t containing S_ω and thus the two points $q_0, q_1 \in S_\omega$ (dashed), so that L_t is disjoint from Δ_ω .

(See Figure 1.8) Let K be the cap of the monotone sofa S_ω . We will find two points $q_0, q_1 \in S_\omega$ on the upper boundary of K sufficiently away from O . Have the right endpoint q_0 of K on the x -axis sufficiently away from the origin O by the distance $d_{\omega, \min}$ (Definition 4.2.2), by using $|K| > 2.2$ and reflecting K along the line passing through O and o_ω if necessary. Take a right triangle with the right-angled vertex q_0 and side 1 (green), to find a lower bound g (blue) of the horizontal side length of S_ω on the line $y = 0$. By the balancedness of S_ω , the side length of S_ω on the line $y = 1$ is also bounded from below by g . Define $q_1 \in K$ as the point on the line $y = 1$ exactly g away from the endpoint o_ω on this side.

Now that we found two points $q_0, q_1 \in S_\omega$, we take the supporting hallway L_t of S_ω angle $t = \pi/2 - \omega \in (0, \omega)$. Using that L_t contains the two points $q_0, q_1 \in S_\omega$ sufficiently away from O , technical calculations show that the region Δ_ω must be enclosed by the inner walls of L_t as in Figure 1.8. So Δ_ω must be disjoint with L_t and thus also with S_ω as desired.

1.6 Surface Area Measure

Summary: This section is an overview of Chapter 5. A planar convex body K does not necessarily have a differentiable boundary. Using the Lebesgue–Stieltjes measure and Brunn–Minkowski theory, we prove an equality that allows us to use the *surface area measure* σ_K of K as a weak derivative of the boundary of K .

(See Figure 2.1) A *planar convex body* K is a nonempty, compact, and convex subset of \mathbb{R}^2 . For any planar convex body K and angle t , define $l_K(t)$ as the supporting line of K with normal vector u_t directing outwards from K . Define the *edge* $e_K(t)$ of K as the intersection $e_K(t) := K \cap l_K(t)$.

Let S^1 be the circle taken as \mathbb{R} modulo 2π . The *surface area measure* σ_K of K is a measure on S^1 that describes the length of the edges $e_K(t)$ of a planar convex body K in terms of the angle t . For any Borel subset X of S^1 , the value $\sigma_K(X)$ is equal to the one-dimensional length of the set $\bigcup_{t \in X} e_K(t)$. We give two examples.

1. If K is the rectangle $[-1, 1] \times [0, 1]$, then σ_K measures the side lengths of K . That is, $\sigma_K(\{t\})$ is equal to 1 if $t = 0, \pi$, and equal to 2 if $t = \pi/2, 3\pi/2$. The measure σ_K on S^1 is zero outside the finite set $\{0, \pi/2, \pi, 3\pi/2\}$ of normal angles of K .

In general, if K is a polygon, then σ_K at the singleton $\{t\}$ of angle t will measure the side length of the edge of K with normal vector u_t .

2. If K is the semicircle $\{(x, y) : x^2 + y^2 \leq 1, y \geq 0\}$ of radius one above the x -axis, then σ_K measures the *differential* side lengths of K . That is, σ_K restricted to $[0, \pi]$ is the usual Borel measure on $[0, \pi]$. The value $\sigma_K(\{3\pi/2\})$ is equal to 2. The measure σ_K is zero on $(\pi, 2\pi) \setminus \{3\pi/2\}$.

In general, if K has a smooth boundary and each edge $e_K(t)$ is a single point on the boundary with curvature $\kappa(t) > 0$, then the density of the measure σ_K at the point $e_K(t)$ is the *radius of curvature* $R(t) := 1/\kappa(t)$.

The measure σ_K is very useful in our analysis of K . Recall that the support function $h_K(t) := \sup\{p \cdot u_t : p \in K\}$ is the signed distance that the edge $e_K(t)$ makes from the origin. The area $|K|$ of K can be expressed using σ_K and h_K as

$$|K| = \frac{1}{2} \int_{t \in S^1} h_K(t) \sigma_K(dt).$$

The measure σ_K also acts as a ‘weak derivative’ of a possibly non-differentiable boundary of K . Recall that $v_t := (-\sin t, \cos t)$. For each angle t , define the *vertices* $v_K^-(t)$ and $v_K^+(t)$ of K as the endpoints of the edge $e_K(t)$ that are furthestmost in the direction of $-v_t$ and v_t respectively. Note that $v_K^+(t)$ depends on K but v_t does not. Then the equality

$$dv_K^+(t) = v_t \sigma_K \tag{1.7}$$

holds, whose meaning we elaborate as below.

Let $I := [a, b]$ be a closed interval. For any right-continuous $f : I \rightarrow \mathbb{R}$ of bounded variation, let df denote the *Lebesgue–Stieltjes measure* of f which is the unique Borel measure on I such that $df(\{a\}) = 0$ and $df((c, d]) = f(d) - f(c)$ for any $(c, d] \subset I$.

The Lebesgue–Stieltjes measure df acts as a rigorous justification of the differential df . We can state informal calculations of differentials like $d(t^2) = 2t dt$ rigorously as the equality

$d(t^2) = 2t dt$ of measures, where the variable t parametrizes the interval I . Note that the Lebesgue–Stieltjes measure dt is from the function $g(t) = t$ on $t \in I$, so that dt denotes the usual Borel measure of I . Correspondingly, the measure $2t dt$ have density function $2t$ on $t \in I$. So the value of measure $2t dt$ on $(c, d]$ is $\int_c^d 2t dt = d^2 - c^2$, which is equal to that of $d(t^2)$.

It turns out that the vertex $v_K^+ : S^1 \rightarrow \mathbb{R}^2$ as a function of angle $t \in S^1$ is right-continuous and of bounded variation. So using the notion above, the left-hand side $dv_K^+(t)$ of Equation (1.7) makes sense as a pair of Lebesgue–Stieltjes measures of the x and y -coordinates of $v_K^+(t)$. The right-hand side of Equation (1.7) is the pair $(-\sin t \cdot \sigma_K, \cos t \cdot \sigma_K)$ of measures on S^1 , where $-\sin t \cdot \sigma_K$ is the measure σ_K on S^1 multiplied pointwise with the measurable function $-\sin t$ on $t \in S^1$.

Intuitively, Equation (1.7) states that the differential of v_K^+ is the vector with direction v_t and side length σ_K at $t \in S^1$. The equality will be used frequently in later parts.

1.7 Injectivity Condition

Summary: This section is an overview of Chapter 6. We show the key property, called the *injectivity condition*, on any balanced maximum sofa S of rotation angle $\pi/2$. The main idea is to prove (Section 1.7.2) and solve for (Section 1.7.3) a differential inequality on S that compares the differential side lengths of S (Equation (1.9)). The inequality is inspired by an ODE of Romik [Rom18] that balances the differential side lengths of moving sofas (Equation (1.8)).

1.7.1 Statement

Recall from Section 1.2 and Equation (1.3) that a monotone sofa S with rotation angle $\omega = \pi/2$ is the intersection

$$S = H \cap \bigcap_{t \in [0, \pi/2]} L_t$$

of the strip H and supporting hallways L_t of S (the vertical strip V_ω overlaps with H as $\omega = \pi/2$). Recall that $\mathbf{x}(t)$ is the inner corner of L_t . The curve $\mathbf{x} : [0, \pi/2] \rightarrow \mathbb{R}^2$, called the *rotation path* of S by Romik [Rom18], determines L_t , the monotone sofa S , and its area $\alpha(\mathbf{x})$ completely. Gerver’s sofa G is derived so that any local perturbation of the rotation path $\mathbf{x} := \mathbf{x}_G$ of G does not increase the area $\alpha(\mathbf{x})$ [Ger92; Rom18; Den24].

A major obstacle in showing the global optimality of G is that there is no manageable formula of the area $\alpha(\mathbf{x})$ of the sofa in terms of the rotation path $\mathbf{x} : [0, \pi/2] \rightarrow \mathbb{R}^2$. All known derivations of G assumes a specific shape of G to find a workable formula of $\alpha(\mathbf{x})$ [Ger92; Rom18; Den24]. We prove the following condition to overcome this obstacle. Recall that $u_t = (\cos t, \sin t)$ and $v_t = (-\sin t, \cos t)$.

Theorem 1.7.1. (*Injectivity condition; abridged*) *The rotation path $\mathbf{x} : [0, \pi/2] \rightarrow \mathbb{R}^2$ of any balanced maximum sofa S is continuously differentiable, and $\mathbf{x}'(t) \cdot u_t < 0$ and $\mathbf{x}'(t) \cdot v_t > 0$ for all $t \in (0, \pi/2)$.*

Theorem 1.7.1 is an abridged version that captures the essence of the full statement (Theorem 6.1.1). We call it the *injectivity condition* as it implies that the rotation path $\mathbf{x} : [0, \pi/2] \rightarrow \mathbb{R}^2$ does not self-intersect (Figure 1.9). Assuming Theorem 1.7.1, we have

$$\mathbf{x}'(t) \cdot (1, 0) = \cos t (\mathbf{x}'(t) \cdot u_t) - \sin t (\mathbf{x}'(t) \cdot v_t) < 0$$

for all $t \in (0, \pi/2)$ so the x -coordinate of $\mathbf{x}(t)$ strictly decreases as t increases. Thus the trajectory of $\mathbf{x}(t)$ forms a Jordan arc, and the area enclosed by $\mathbf{x}(t)$ can be expressed using Green's theorem.




Figure 1.9: Injectivity condition on a monotone sofa $S = K \setminus \mathcal{N}(K)$ with cap K implies that the inner corner $\mathbf{x}(t)$ of the supporting hallways L_t , as a curve over $t \in [0, \pi/2]$, does not self-intersect.

1.7.2 A Differential Inequality

Assume an arbitrary balanced maximum sofa S with rotation angle $\pi/2$ and supporting hallways L_t for $t \in [0, \pi/2]$. In [Rom18], Romik introduced a set of ODEs that balance the differential side lengths of S .

Definition 1.7.1. (See the figures below Theorem 8.4.2) For any angle t where the wall $a(t)$ (resp. $b(t)$, $c(t)$, and $d(t)$) of the hallway L_t is tangent to S , define $\mathbf{A}(t)$ (resp. $\mathbf{B}(t)$, $\mathbf{C}(t)$, and $\mathbf{D}(t)$) as the corresponding point of tangency.

The balancing ODEs by Romik are in terms of the four curves $\mathbf{A}(t)$, $\mathbf{B}(t)$, $\mathbf{C}(t)$, $\mathbf{D}(t)$ in Definition 1.7.1 and the inner corner $\mathbf{x}(t)$ of L_t . In particular, he derives Gerver's sofa $S = G$ by parametrizing the boundary with the five curve segments (Figure 8.3), each on a different interval of t , and solving for the ODEs on the curve segments.

Assume for now that all five curves are well-defined and continuously differentiable on their respective domain. This be alleviated in the full proof at Chapter 6. As S is a monotone sofa, the points $\mathbf{A}(t)$ and $\mathbf{C}(t)$ are well-defined over all $t \in (0, \pi/2)$ and extends naturally to $t \in [0, \pi/2]$ by taking limits. If the point $\mathbf{B}(t)$ is well-defined (that is, if S makes contact with the wall $b(t)$ of L_t), then since $\mathbf{A}(t)$ and $\mathbf{B}(t)$ are points of tangency of parallel lines $a(t)$ and $b(t)$ of distance one, we have $\mathbf{B}(t) = \mathbf{A}(t) - u_t$ (Theorem 1 of [Rom18]). Likewise, we have $\mathbf{D}(t) = \mathbf{B}(t) - v_t$ if $\mathbf{D}(t)$ is well-defined.

(See the right side of Figure 1.5) Assume that for some right angle t and its neighborhood, the supporting hallway L_t makes contact with S at three points $\mathbf{A}(t)$, $\mathbf{B}(t)$, and $\mathbf{x}(t)$. Recall that S is the limit of balanced polygons S_Θ (Section 1.4.1). As S_Θ converges to S , the three balanced sides (green) of S_Θ on the walls $a(t)$ and $b(t)$ becomes the differential sides of G contributed by three points $\mathbf{A}(t)$, $\mathbf{B}(t)$, and $\mathbf{x}(t)$. So their lengths balance each other as

$$\langle \mathbf{A}'(t), v_t \rangle = \langle -\mathbf{B}'(t), v_t \rangle + \langle \mathbf{x}'(t), v_t \rangle \quad (1.8)$$

which is one of the ODEs by Romik (Equation (20) of [Rom18]). See the equations following Theorem 8.4.2 for many other examples.

Equation (1.8) is very useful but depends on the assumption that S makes contact with L_t at three points $\mathbf{A}(t), \mathbf{B}(t), \mathbf{x}(t)$. So we (essentially) prove the following weaker *inequality* that works for any S regardless of whether it makes contact with L_t at $\mathbf{B}(t)$ or $\mathbf{x}(t)$.

$$\langle \mathbf{A}'(t), v_t \rangle \leq \max(\langle -\mathbf{B}'(t), v_t \rangle, 0) + |\langle \mathbf{x}'(t), v_t \rangle| \quad (1.9)$$

Even if S does not make contact with L_t on the line $b(t)$, the point $\mathbf{B}(t)$ extends naturally over all $t \in [0, \pi/2]$ by letting $\mathbf{B}(t) := \mathbf{A}(t) - u_t$.

We sketch the idea behind Equation (1.9). Our description here is only a rough sketch of the ideas and we hide the details of magnitude analysis. See the proof of Theorem 6.3.3 for full details.

Take the maximum polygon sofa S_Θ that approximates S . Take three adjacent angles $t - \delta, t, t + \delta$ from the finite angle set Θ . It turns out that the side of S_Θ on the line $a(t)$ is of magnitude $\delta \langle \mathbf{A}'(t), v_t \rangle + O(\delta^2)$, so after dividing by δ , converges to $\langle \mathbf{A}'(t), v_t \rangle$ as $\delta \rightarrow 0$ and Θ gets denser in $[0, \pi/2]$. This is the left-hand side of Equation (1.9).

Let $\vec{b}(t)$ be the half-line on $b(t)$ from $\mathbf{x}(t)$ that represents the right inner wall of L_t starting with $\mathbf{x}(t)$. We now overestimate all sides of S_Θ on the half-line $\vec{b}(t)$. Define R as the union of three closed half-planes $H^d(s)$, each of angle $s = t - \delta, t, t + \delta$ bounded from below by the left inner wall $d(s)$. We will overestimate the sides of S_Θ on $\vec{b}(t) \cap R$ and $\vec{b}(t) \setminus R$ respectively. Adding two estimates below and sending $\delta \rightarrow 0$ will give the right-hand side of Equation (1.9).

- The length of set $\vec{b}(t) \cap R$ is of magnitude $\leq \delta |\langle \mathbf{x}'(t), v_t \rangle| + O(\delta^2)$.

To see this, observe that the point $\mathbf{x}(t)$ is on the boundary $d(t)$ of $H^d(t)$ and is away from the boundary $d(t \pm \delta)$ of $H^d(t \pm \delta)$ by the signed distance $\mp \delta \langle \mathbf{x}'(t), v_t \rangle + O(\delta^2)$ along the direction v_t . Exact verification is done in Lemma 6.3.2.

- The sides of S_Θ on $\vec{b}(t) \setminus R$ are of magnitude $\leq \delta \max(\langle -\mathbf{B}'(t), v_t \rangle, 0) + O(\delta^2)$.

To see this, observe that for each angle $s = t - \delta, t, t + \delta$, the set S_Θ is disjoint from the inner quadrant Q_s^- of L_t bounded from above by $b(s)$ and $d(s)$. So for each s , the set $S_\Theta \setminus R$ is contained in the closed half-plane $H^b(s)$ bounded from below by the line $b(s)$. Now the sides of S_Θ on $\vec{b}(t) \setminus R$ is contained in the segment of $H^b(t - \delta) \cap H^b(t) \cap H^b(t + \delta)$ on the line $b(t)$. This segment is contributed by the lines $b(t)$ and $b(t \pm \delta)$ and is of the claimed magnitude.

In the actual proof prested in Chapter 6, Equation (1.9) is not stated as-is and formulated quite differently as

$$\sigma_K \leq k_0(g(t)) dt \quad (1.10)$$

in Theorem 6.4.3 where

$$k_0(x) := \max(|x - 1|, (|x - 1| + 1)/2).$$

This is to ensure that the inequality works for general S that may have the contact point $\mathbf{A}(t)$ that is not differentiable in t , or have more than one contact points with outer wall $a(t)$. So the actual proof proceeds with Equation (1.10) that works for any K , but it essentially follows the idea behind the proof of Equation (1.9) sketched above.

We derive Equation (1.10) from Equation (1.9) as below. This explains why the inequalities are more or less equivalent and why the function k_0 is involved. First use $\mathbf{B}(t) = \mathbf{A}(t) - u_t$ and $u'_t = v_t$ and write

$$\langle -\mathbf{B}'(t), v_t \rangle = -\langle \mathbf{A}'(t), v_t \rangle + 1.$$

Then by letting $\alpha := \langle \mathbf{x}'(t), v_t \rangle$, Equation (1.9) implies

$$\langle \mathbf{A}'(t), v_t \rangle \leq \max(|\alpha|, (|\alpha| + 1)/2)$$

in both cases $\langle \mathbf{A}'(t), v_t \rangle < 1$ and $\langle \mathbf{A}'(t), v_t \rangle \geq 1$. Using Equation (1.7), the left-hand side is the differential side length at $\mathbf{A}(t)$ equal to the density of σ_K . It turns out that the value $\alpha = \langle \mathbf{x}'(t), v_t \rangle$ in right-hand side is equal to $g(t) - 1$ where $g(t)$ is the *arm length* that will be defined soon. Substituting both sides, we get Equation (1.10).

1.7.3 Solving the Differential Inequality

We now sketch the argument that solves the Equation (1.10) and proves the injectivity condition.

Recall that $\mathbf{y}(t)$ is the outer corner of L_t corresponding to $(1, 1)$ of L (Definition 1.2.5). For each $t \in [0, \pi/2]$, the *arm lengths* $f(t)$ and $g(t)$ measure the distance from outer corner $\mathbf{y}(t)$ to $\mathbf{A}(t)$ and $\mathbf{C}(t)$ respectively.¹⁸ A computation (Theorem 6.2.3) shows that

$$\mathbf{x}'(t) = -(f(t) - 1)u_t + (g(t) - 1)v_t$$

so that proving $f(t), g(t) > 1$ on $t \in (0, \pi/2)$ is sufficient for establishing the injectivity condition (Theorem 1.7.1).

We will express Equation (1.10) purely in terms of arm lengths. The derivative of $f(t)$ is

$$f'(t) = g(t) - \langle \mathbf{A}'(t), v_t \rangle$$

because $\langle \mathbf{y}'(t), v_t \rangle = g(t)$ and $f(t) = \langle \mathbf{y}(t) - \mathbf{A}(t), v_t \rangle$ (Theorem 6.2.5). As the side length $\langle \mathbf{A}'(t), v_t \rangle$ corresponds to σ_K at t , Equation (1.10) is equivalent to

$$f'(t) \geq g(t) - k_0(g(t)) = m_0(g(t)) \quad (1.11)$$

where

$$m_0(x) := x - k_0(x) = x - \max(|x - 1|, (|x - 1| + 1)/2)$$

is monotonically increasing. This is done rigorously in Theorem 6.5.1.

We now use Equation (1.11) to iteratively obtain better lower bounds $f_0(t), f_1(t), \dots$ of $f(t)$ on $t \in [0, \pi/2]$. Let $f_0(t) := 0$ so that $f_0(t)$ is a trivial lower bound of $f(t)$. The same argument on S reflected along the y -axis shows that $f_0(\pi/2 - t)$ is a lower bound of $g(t)$. We have $f(0) = 1$ because the point $\mathbf{A}(0)$ should be on the x -axis. Equation (1.11) implies that

$$f(t) \geq 1 + \int_0^t m_0(g(u)) du \geq 1 + \int_0^t m_0(f_0(\pi/2 - u)) du.$$

We just obtained a new lower bound of $f(t)$ in the right-hand side. By letting

$$f_1(t) := \max \left(f_0(t), 1 + \int_0^t m_0(f_0(\pi/2 - u)) du \right) \quad (1.12)$$

¹⁸The actual definition of arm lengths (Definition 6.2.1) have signs $f_K^\pm(t)$ and $g_K^\pm(t)$ in superscript as we cannot guarantee that the cap K meets the line $a(t)$ at a single point $\mathbf{A}(t)$. This sketch assumes that K meets $a(t)$ at a single point, so that $f(t) = f_K^\pm(t)$ (and the same for g).

we obtain a better lower bound $f_1(t)$ of $f(t)$. A symmetric argument also shows that $g(t) \geq f_1(\pi/2 - t)$. Further iterations of Equation (1.12) will give monotonically increasing lower bounds f_2, f_3, \dots of f . Somewhat magically, eleven iterations of this improvement gives $f(t) \geq f_{11}(t) > 1$, proving the injectivity condition (Figure 1.10). Detailed computations are done in Section 6.5.




Figure 1.10: The arm length $f(t)$ of Gerver's sofa G and the lower bounds $f_0(t), f_1(t), \dots$ of $f(t)$. Numerical computations show that three iterations are sufficient to give $f(t) \geq f_3(t) > 1$. But to minimize computer assistance, we do more iterations and show $f_i(t) \geq (i-1)/12$ for $i \leq 10$ in Lemma 6.5.3, which is sufficient to prove the injectivity hypothesis.

1.8 Optimality of Gerver's Sofa

Summary: This section is an overview of Chapter 8 that proves the main Theorem 1.1.1. The previous Chapter 7 prepares a minimal theoretical framework needed to execute the ideas below in Chapter 8.

We establish an upper bound $\mathcal{Q}(S)$ of the area of any monotone sofa S satisfying the injectivity condition. To do so, we construct a region R enclosing S so that $R = S$ if S is Gerver's sofa G . The upper bound \mathcal{Q} is then defined as the area of R (Section 1.8.1). We define a convex space \mathcal{L} of tuples (K, B, D) of convex bodies, so that each sofa S embeds one-to-one to a tuple $(K, B, D) \in \mathcal{L}$ and \mathcal{Q} is a quadratic functional on \mathcal{L} via Brunn-Minkowski theory (Section 1.8.2). The concavity of \mathcal{Q} on \mathcal{L} is established using Mamikon's theorem, and the local optimality of \mathcal{Q} at G is established using the local optimality ODEs on G by Romik (Section 1.8.3). As G is a local optimum of a globally concave \mathcal{Q} , it is also a global optimum of \mathcal{Q} and thus the area.

1.8.1 Definition of \mathcal{Q}

(See Figure 1.1) The niche of Gerver's sofa G has a characteristic shape made of one 'core' colored blue and the two 'tails' colored red. Assuming that a maximum-area sofa S_{\max} follows the same shape, the derivation $S_{\max} = G$ is essentially done in the existing works establishing

the local optimality of G [Ger92; Rom18; Den24]. So the difficulty of the moving sofa problem lies in showing that S_{\max} indeed follows the same shape as G .

We circumvent the difficulty by defining a *larger* region R contains S_{\max} and have the desired shape of one core and two tails. Then the upper bound \mathcal{Q} of the area of a moving sofa is simply defined as the area of R . Take any monotone sofa $S = K \setminus \mathcal{N}(K)$ of rotation angle $\pi/2$ with cap K , satisfying the injectivity condition. Recall that $\mathbf{x}(t)$, $b(t)$, and $d(t)$ are respectively the inner corner, right inner wall, and left inner wall of supporting hallway L_t with angle t . We construct R as follows.

(See Figure 1.11) There is a specific angle $\varphi \in [0.039, 0.040]$ such that the rotation path $\mathbf{x}_G : [0, \pi/2] \rightarrow \mathbb{R}$ of Gerver's sofa draws the core portion of the niche at the interval $[\varphi, \pi/2 - \varphi]$. Using the same angles, cut the cap K and niche $\mathcal{N}(K)$ of arbitrary monotone sofa S into three parts, using the lines $b(\varphi)$ and $d(\pi/2 - \varphi)$ passing through $\mathbf{x}(\varphi)$ and $\mathbf{x}(\pi/2 - \varphi)$ of S respectively. Let H^R (resp. H^L) be the half-planes bounded from left by $b(\varphi)$ (resp. right by $d(\pi/2 - \varphi)$). Let H^M be the region $\mathbb{R}^2 \setminus H^R \setminus H^L$. Then the sets H^R, H^M, H^L partition the plane into three parts.¹⁹

The injectivity condition on S (Theorem 1.7.1) implies that the rotation path $\mathbf{x}(t)$ should be in H^R , H^M , and H^L as t is in the interval $[0, \varphi]$, $[\varphi, \pi/2 - \varphi]$, and $[\pi/2 - \varphi, \pi/2]$ respectively (Lemma 8.1.6). Using this, we take a subset N' of the niche $\mathcal{N}(K)$ as follows. In the region H^R , take only the region swept out by the right inner wall $b(t)$ as $t \in [\varphi, \pi/2]$, where $\mathbf{x}(t)$ is outside H^R (colored blue). In the region H^M , take only the region bounded by the lines $b(\varphi)$, $d(\pi/2 - \varphi)$, $y = 0$, and the inner corner $\mathbf{x}(t)$ restricted to $[\varphi, \pi/2 - \varphi]$ (colored green). In the region H^L , take only the region swept out by the left inner wall $d(t)$ as $t \in [0, \pi/2 - \varphi]$, where $\mathbf{x}(t)$ is outside H^L (colored red). The injectivity condition guarantees that the final region N' is a subset of the niche $\mathcal{N}(K)$.




Figure 1.11: The overestimated region R is obtained by taking the region N' (blue in H^R , green in H^M , red in H^L) away from the cap K of monotone sofa S .

The region R is now simply defined as $K \setminus N'$. While R may not be a moving sofa, the ‘niche’ N' of R consists of one core and two tails like that of G does. For a general monotone sofa S , the endpoints of the core and two tails of N' does not match each other, so we simply connect them by the line segments each on $b(\varphi)$ and $d(\pi/2 - \varphi)$. As Gerver's sofa G is constructed by design to have the matching endpoints of core and tails, the region R is equal to S if $S = G$ (Theorem 8.4.6).

¹⁹The half-planes H^R and H^L do overlap technically, but it does not matter as the region of overlap $H^R \cap H^L$ is disjoint from the sets K and $\mathcal{N}(K)$ that we divide (Lemma 8.1.4).

1.8.2 Quadraticity of \mathcal{Q}

The collection \mathcal{K} of all planar convex bodies form a *convex domain* with the barycentric operation $c_\lambda(K_1, K_2) := (1 - \lambda)K_1 + \lambda K_2$. Here,

$$aK := \{ap : p \in K\}$$

is the *dilation* of a convex body K by $a \geq 0$, and

$$K_1 + K_2 := \{x_1 + x_2 \in \mathbb{R}^2 : x_1 \in K_1, x_2 \in K_2\}$$

is the *Minkowski sum* of convex bodies K_1, K_2 . The operations satisfy necessary properties (commutative, associative, and distributive) that makes \mathcal{K} an abstract convex cone.

Many values on convex body K , including the support function $h_K(t) := v_K^\perp(t) \cdot u_t$, vertex $v_K^\perp(t)$, and surface area measure σ_K are convex-linear in K . Correspondingly, the area

$$|K| = \frac{1}{2} \int_{t \in S^1} h_K(t) \sigma_K(t)$$

of $K \in \mathcal{K}$ is a *quadratic* functional on the convex domain \mathcal{K} . This notion of a quadratic functional on a (abstract) convex domain is established rigorously in Section 7.1.

In the previous Section 1.8.1, we defined an overestimation R of a monotone sofa S . We will now define the three convex bodies K , B , and D from S that represents different parts of the region R . This step is very important. While the area of R does not have a quadratic expression involving K only, it does have a quadratic expression involving K, B , and D (Definition 8.2.2) that is amenable to further analysis (see also Remark 8.1.2).

(Compare Figure 1.11 with Figure 1.2) Again, let S be any monotone sofa of rotation angle $\pi/2$ satisfying the injectivity condition. The convex body K is the (usual) cap of S . Convex bodies B and D are the portions of the region R in the half-planes H^R and H^L respectively. Put precisely, B (resp. D) is the cap K intersected with the closed half-planes bounded from below by $b(t)$ for all $t \in [\varphi, \pi/2]$ (resp. $d(t)$ for all $t \in [0, \pi/2 - \varphi]$). This defines the convex bodies K, B , and D from S .

The three convex bodies satisfy certain linear constraints (Lemma 8.1.7). An example is

$$h_K(t) + h_B(\pi + t) \leq 1 \tag{1.13}$$

for every $t \in [\varphi^R, \pi/2]$, which holds because B is bounded from below by the line $b(t)$ which is distance one away from the the supporting line $a(t)$ of K . With this, define \mathcal{L} as the collection of all tuples (K, B, D) of convex bodies satisfying such constraints (Definition 8.1.3). The collection of all monotone sofas S with rotation angle $\pi/2$ now embeds to a subset of \mathcal{L} by constructing the convex bodies (K, B, D) from S as above. The space \mathcal{L} is a convex domain with pairwise barycentric operation

$$c_\lambda((K_1, B_1, D_1), (K_2, B_2, D_2)) := (c_\lambda(K_1, K_2), c_\lambda(B_1, B_2), c_\lambda(D_1, D_2)).$$

We now define the upper bound $\mathcal{Q}(K, B, D)$ of the area of a monotone sofa S as a quadratic functional on $(K, B, D) \in \mathcal{L}$ (Definition 8.2.2). Recall that \mathcal{Q} is equal to $|K| - |N'|$ where $|N'|$ is the area of the underestimated niche in Figure 1.11. Using injectivity condition, we essentially²⁰ show that the boundary γ of N' is a Jordan curve. Take γ counterclockwise,

²⁰The actual proof takes three Jordan curves, two bounding the red and blue regions of Figure 1.11 (Lemma 8.2.2) and one bounding the green region of Figure 1.11 (Lemma 8.2.3).

then by Green's theorem the region N' have area

$$\mathcal{J}(\gamma) := \frac{1}{2} \int_a^b \gamma(t) \times \gamma'(t) dt$$

which we call the *curve area functional* on $\gamma : [a, b] \rightarrow \mathbb{R}^2$. So we have $\mathcal{Q} = |K| - \mathcal{J}(\gamma)$ in particular.

For a convex body $X = B$ or D , define the segment $\mathbf{u}_X^{a,b}$ of the boundary of X as the union of all edges of X with normal vectors u_t of angle $t \in (a, b)$. We further express $\mathcal{J}(\gamma)$ as a quadratic term on K, B , and D by breaking the boundary γ of N' into the following five segments.

1. The segment $\mathbf{d}_D := \mathbf{u}_D^{3\pi/2, 3\pi/2+\varphi^L}$ of the boundary of D , representing the left tail of N' .
2. The line segment connecting the right end Y_D of the left tail \mathbf{d}_D , to the left end $\mathbf{x}_K^L := \mathbf{x}_K(\pi/2 - \varphi)$ of the core \mathbf{x}_K .
3. The rotation path $\mathbf{x}_K : [\varphi, \pi/2 - \varphi] \rightarrow \mathbb{R}^2$ of cap K reversed in direction, representing the core of N' .
4. The line segment connecting the right end $\mathbf{x}_K^R := \mathbf{x}_K(\varphi)$ of the core \mathbf{x}_K , to the left end X_B of the right tail \mathbf{b}_B .
5. The segment $\mathbf{b}_B := \mathbf{u}_B^{\pi+\varphi^R, 3\pi/2}$ of the boundary of B , representing the right tail of N' .

Each segment corresponds to each term in the appearing order of the Definition 8.2.2 of $\mathcal{Q}(K, B, D) = |K| - \mathcal{J}(\gamma) =$

$$|K| + \mathcal{J}(\mathbf{d}_D) + \mathcal{J}(Y_D, \mathbf{x}_K^L) - \mathcal{J}(\mathbf{x}_K|_{[\varphi, \pi/2-\varphi]}) + \mathcal{J}(\mathbf{x}_K^R, X_B) + \mathcal{J}(\mathbf{b}_B)$$

where $\mathcal{J}(p, q) := (x_p y_q - x_q y_p)/2$ is the curve area functional of the segment from $p = (x_p, y_p)$ to $q = (x_q, y_q)$.

We now argue that \mathcal{Q} is quadratic in K, B , and D . The area $|K|$ is quadratic in K as seen above. The quadraticity of the core term $\mathcal{J}(\mathbf{x}_K|_{[\varphi, \pi/2-\varphi]})$ comes from linearity of \mathbf{x}_K in K . Theorem 7.3.2 computes

$$\mathcal{J}(\mathbf{u}_X^{a,b}) = \frac{1}{2} \int_{t \in (a,b)} h_K(t) \sigma_K(dt)$$

which is quadratic in K . This establishes the quadraticity of two tail terms $\mathcal{J}(\mathbf{d}_D)$ and $\mathcal{J}(\mathbf{b}_B)$. The terms on two line segments come from bilinearity of $\mathcal{J}(p, q) := (x_p y_q - x_q y_p)/2$ in $p, q \in \mathbb{R}^2$.

1.8.3 Optimality of \mathcal{Q} at Gerver's Sofa

(See Figure 1.12) Let K be any convex body. Take an interval $[a, b]$ of length $\leq 2\pi$. For each angle $t \in [a, b]$, assume a tangent segment s_t of K with length $\alpha(t)$ and one endpoint $v_K^+(t)$ on K , making an angle of t from the y -axis. *Mamikon's theorem* states that the region swept out by the segments s_t over all $t \in [a, b]$ is exactly $\frac{1}{2} \int_a^b \alpha(t)^2 dt$.

(See Figure 1.13) Mamikon's theorem is used to show that \mathcal{Q} is globally concave on \mathcal{L} . The idea is to attach multiple 'Mamikon regions' (grey) to the region R of area \mathcal{Q} . The length




Figure 1.12: Mamikon's theorem.

$\alpha(t)$ of each tangent segment in grey with angle t turns out to be linear in \mathcal{L} . So the area $\frac{1}{2} \int_a^b \alpha(t)^2 dt$ of each Mamikon region is convex and quadratic in \mathcal{L} . We show in Lemma 8.3.7 that the total area of R and all Mamikon regions (bounded by bold lines) is linear in K . So the area \mathcal{Q} of R is a linear functional (bold lines) subtracted by convex quadratic functionals (grey regions), which is concave. Section 8.3 rigorously checks the full details.

To establish the main Theorem 1.1.1, it suffices to show that the tuple $(K, B, D) \in \mathcal{L}$ arising from Gerver's sofa G is a maximizer of \mathcal{Q} . Assuming this, recall that a balanced maximum sofa S^* attaining the maximum area also satisfies the injectivity condition (Theorem 1.7.1). So the maximum-area S^* also corresponds to another tuple $(K^*, B^*, D^*) \in \mathcal{L}$ of convex bodies as described in Section 1.8.2. Because the region R of Gerver's sofa G matches with G , we have $\mathcal{Q}(K, B, D) = |R| = |G|$. By the optimality of \mathcal{Q} at (K, B, D) , and that $\mathcal{Q}(K^*, B^*, D^*)$ is an upper bound of the area $|S^*|$, we have

$$\mathcal{Q}(K, B, D) \geq \mathcal{Q}(K^*, B^*, D^*) \geq |S^*|.$$

So we have $|G| \geq |S^*|$, proving the main Theorem 1.1.1.

We now show that \mathcal{Q} is maximized at the point $(K, B, D) \in \mathcal{L}$ from Gerver's sofa G . Choose an arbitrary $(K^*, B^*, D^*) \in \mathcal{L}$. The directional derivative of \mathcal{Q} at (K, B, D) in the direction towards (K^*, B^*, D^*) is defined as

$$\begin{aligned} & D\mathcal{Q}(K, B, D; K^*, B^*, D^*) \\ & := \left. \frac{d}{d\lambda} \right|_{\lambda=0} \mathcal{Q}(c_\lambda((K, B, D), (K^*, B^*, D^*))) \end{aligned} \tag{1.14}$$

where $\lambda \in [0, 1]$ interpolates between (K, B, D) and (K^*, B^*, D^*) . If the value is ≤ 0 regardless of the choice of $(K^*, B^*, D^*) \in \mathcal{L}$, then (K, B, D) indeed achieves the maximum value of concave and quadratic \mathcal{Q} as desired; this can be shown by quadraticity of \mathcal{Q} (Theorem 7.1.5).

So it remains to compute Equation (1.14) and show that it is non-positive. Assuming that (K^*, B^*, D^*) is close enough to (K, B, D) , the value of $D\mathcal{Q}$ is approximately the rate of change of \mathcal{Q} along the interval $\lambda \in [0, 1]$, so

$$D\mathcal{Q}(K, B, D; K^*, B^*, D^*) \simeq \mathcal{Q}(K^*, B^*, D^*) - \mathcal{Q}(K, B, D)$$




Figure 1.13: Mamikon's theorem applied to the upper bound Q of sofa area. As Mamikon regions in grey are added to the region R with area Q , the resulting shape bounded by bold lines have an area linear in K .

and we use them interchangeably here for ease of explanation (we do not use this in full calculation). Instead of computing the full DQ for general $(K^*, B^*, D^*) \in \mathcal{L}$ as in Section 8.5, we take two representative cases of (K^*, B^*, D^*) and illustrate how DQ is computed.

Recall that the boundary of G is parametrized by the contact points $\mathbf{A}(t)$, $\mathbf{B}(t)$, $\mathbf{C}(t)$, $\mathbf{D}(t)$, and $\mathbf{x}(t)$ that G makes with supporting hallways L_t (Figure 8.3). In both representative cases of (K^*, B^*, D^*) , fix a particular angle $t \in (0, \pi/2)$ so that G meets L_t at three points $\mathbf{A}(t)$, $\mathbf{B}(t)$, and $\mathbf{x}(t)$ as in the right side of Figure 1.5. Also, fix a sufficient small $\delta > 0$ and let $I := [t, t + \delta]$. Take an arbitrary $\epsilon > 0$ that is sufficiently small relative to δ . We now assume the first case of $(K^*, B^*, D^*) \in \mathcal{L}$.

Case 1: Recall that $G = H \cap \bigcap_{s \in [0, \pi/2]} L_s$ is the intersection of the horizontal strip H and supporting hallways L_s . Translate each L_s to $L_s^* := L_s + \epsilon u_t$ by ϵu_t for any $s \in I = [t, t + \delta]$ and fix $L_s^* := L_s$ for any other $s \notin I$. Take the new sofa $G^* := H \cap \bigcap_{t \in [0, \pi/2]} L_s^*$ which is a slight perturbation of G . Assume the case where the convex bodies $(K^*, B^*, D^*) \in \mathcal{L}$ come from G^* as described in Section 1.8.2.

In this Case 1, the new sofa G^* is obtained from G by the following changes in region. We ignore second-order or smaller terms of δ and ϵ in length.

1. Adding a rectangle of approximate base $\delta \langle \mathbf{A}'(t), u_t \rangle$ and height ϵ near the point $\mathbf{A}(t)$.
2. Removing a rectangle of approximate base $\delta \langle -\mathbf{B}'(t), u_t \rangle$ and height ϵ near the point $\mathbf{B}(t)$.
3. Removing a parallelogram with approximate sides of vector $\delta \mathbf{x}'(t)$ and ϵu_t near the point $\mathbf{x}(t)$.

So the area change for Case 1 is approximately

$$\begin{aligned} D\mathcal{Q}(K, B, D; K^*, B^*, D^*) &\simeq |G^*| - |G| \\ &\simeq \delta (\epsilon \langle \mathbf{A}'(t), u_t \rangle - \epsilon \langle -\mathbf{B}'(t), u_t \rangle - \epsilon \langle \mathbf{x}'(t), u_t \rangle). \end{aligned} \quad (1.15)$$

Romik's ODE Equation (1.8) on G now balance the differential side lengths and make the value of $D\mathcal{Q}$ from (K, B, D) to (K^*, B^*, D^*) equal to zero. In fact, this is essentially how Romik derived his ODE for G in [Rom18].

Now we assume a slightly more general Case 2 where the edges of K and B along the angle t can move in the direction of $\pm u_t$ independently.

Case 2. Take $\delta' > 0$ sufficiently smaller than δ so that $U := (t - \delta', t + \delta + \delta')$ is a sufficiently close neighborhood of $I = [t, t + \delta]$. Take ϵ_K and $\epsilon_B > 0$ sufficiently smaller than δ and δ' . Take any $(K^*, B^*, D^*) \in \mathcal{L}$ so that the followings are true.

- $h_{K^*}(s) = h_K(s) + \epsilon_K$ for $s \in I$ and $h_{K^*}(s) = h_K(s)$ for $s \notin U$
- $h_{B^*}(s) = h_B(s) - \epsilon_B$ for $s \in I + \pi = [\pi + t, \pi + t + \delta]$ and $h_{B^*}(s) = h_B(s)$ for $s \notin U + \pi$.

Observe that this Case 2 reduces to the previous Case 1 if we let $\epsilon_K = \epsilon_B = \epsilon$. Recall that for a general $(K^*, B^*, D^*) \in \mathcal{L}$, the value $\mathcal{Q}(K^*, B^*, D^*)$ is equal to $|K^*| - |N'|$ where N' is the region bounded by five curves and lines from K^* , B^* , and D^* as in Section 1.8.2. In this case, the rectangle in (1) near $\mathbf{A}(t)$ now have height ϵ_K , and the rectangle near $\mathbf{B}(t)$ in (2) and parallelogram near $\mathbf{x}(t)$ in (3) have sides of length ϵ_B and ϵ_K respectively. Now the change in the value of \mathcal{Q} is approximately

$$\begin{aligned} D\mathcal{Q}(K, B, D; K^*, B^*, D^*) \\ \simeq \delta (\epsilon_K \langle \mathbf{A}'(t), u_t \rangle - \epsilon_B \langle -\mathbf{B}'(t), u_t \rangle - \epsilon_K \langle \mathbf{x}'(t), u_t \rangle). \end{aligned}$$

Now it takes a bit more than just Equation (1.8) to show $D\mathcal{Q} \leq 1$. Because the hallway L_t meets G at the points $\mathbf{A}(t)$ and $\mathbf{B}(t)$, we have $h_K(t) + h_B(\pi + t) = 1$. The condition in Equation (1.13) to $(K^*, B^*, D^*) \in \mathcal{L}$ implies $h_{K^*}(t) + h_{B^*}(\pi + t) \leq 1$. So we should have $\epsilon_K \leq \epsilon_B$. Now Equation (1.8) with $\langle -\mathbf{B}'(t), u_t \rangle \geq 0$, that the differential side at $\mathbf{B}(t)$ is nondegenerate, imply

$$\begin{aligned} D\mathcal{Q}(K, B, D; K^*, B^*, D^*) &\simeq |G^*| - |G| \\ &\simeq \delta (\epsilon_K \langle \mathbf{A}'(t), u_t \rangle - \epsilon_B \langle -\mathbf{B}'(t), u_t \rangle - \epsilon_K \langle \mathbf{x}'(t), u_t \rangle) \\ &\leq \delta (\epsilon_K \langle \mathbf{A}'(t), u_t \rangle - \epsilon_K \langle -\mathbf{B}'(t), u_t \rangle - \epsilon_K \langle \mathbf{x}'(t), u_t \rangle) = 0. \end{aligned}$$

In the full calculation of $D\mathcal{Q}$, all the differential sides of K, B , and D are simultaneously taken into account. The interval $t \in [0, \pi/2]$ is divided into five intervals I_1, \dots, I_5 , where the set of contact points between G and L_t is fixed for each $t \in I_i$ (Definition 8.4.1; see also the second column of Theorem 8.4.2). For each $t \in I_i$, the differential sides with normal vectors $\pm u_t$ (or $\pm v_t$) contribute to $D\mathcal{Q}$ in a way analogous to the Case 2 above. In particular, Case 2 corresponds to the case $t \in I_4$ and normal vectors $\pm u_t$.

In the full proof of $D\mathcal{Q} \leq 0$ at Gerger's sofa G , we first import a total of 10 ODEs in $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$, and \mathbf{x} that balance all the differential sides of G from [Rom18] (Theorem 8.4.2). Each ODE is effective on an interval $t \in I_i$ and a specific direction $\pm u_t$ or $\pm v_t$. Then the side lengths of G on $\mathbf{A}(t), \mathbf{B}(t), \mathbf{C}(t), \mathbf{D}(t)$, and $\mathbf{x}(t)$ are represented as the surface area measures

of K , B , and D (Proposition 8.4.4) and a measure ι_K (Definition 8.4.6). The 10 ODEs of Romik are translated to equalities of measures on each I_i in Theorem 8.4.5. They are used in the full computation at Theorem 8.5.7 where the contribution by each angle $t \in I_i$ is shown to be nonnegative. The analysis of Case 2 above, in particular, corresponds to the third case in the proof of Theorem 8.5.7.

Chapter 2

Monotone Sofas and Caps

This chapter follows the overview in Section 1.2 and shows that a maximum-area sofa is a *monotone sofa* S which is the *cap* K subtracted by *niche* $\mathcal{N}(K)$.

- Section 2.1 prepares standard notions on a planar convex body K .
- Section 2.2 defines the *supporting hallway* $L_S(t)$ of any moving sofa S making contact with S in the outer walls.
- Section 2.3 defines the intersection $\mathcal{I}(S)$ of unit-width stripes H, V_ω , and the supporting hallways $L_S(t)$ containing S . We show that $\mathcal{I}(S)$ is connected (Theorem 2.3.6) so that it is a larger moving sofa containing S (Theorem 2.3.2). With this, we define a *monotone sofa* as the intersection $\mathcal{I}(S)$ coming from some moving sofa S (Definition 2.3.7).
- Section 2.4 defines the notion of cap K (Definition 2.4.1) and niche $\mathcal{N}(K)$ (Definition 2.4.5), and shows that a monotone sofa S is equal to the cap $K := \mathcal{C}(S)$ subtracted by niche $\mathcal{N}(K)$ (Theorem 2.4.3).
- Section 2.5 shows that for a monotone sofa S , its cap K contains the niche $\mathcal{N}(K)$ as a subset (Theorem 2.5.9). Along the way, many notions on the cap and niche are defined. We define the space of all caps \mathcal{K}_ω^c with rotation angle ω and turn the problem into the maximization of the sofa area functional $\mathcal{A}_\omega(K) := |K| - |\mathcal{N}(K)|$ on caps $K \in \mathcal{K}_\omega^c$.

2.1 Planar Convex Body

Definition 2.1.1. A *planar convex body* K is a nonempty, compact, and convex subset of \mathbb{R}^2 . Define \mathcal{K} as the collection of all planar convex bodies K .

All convex bodies appearing in this work will be planar, so we will omit the word ‘planar’. Many authors also require K° to be nonempty, but we follow [Sch13] and allow K° to be empty. That is, a closed line segment or a point is also a convex body.

We define standard notions on a convex body K . Note that the notions generalize naturally to any nonempty and compact $S \subseteq \mathbb{R}^2$. Here, S^1 is taken as $\mathbb{R}/2\pi\mathbb{Z}$. We will denote an element or interval of S^1 by its representation in \mathbb{R} .

Definition 2.1.2. For any subset X of \mathbb{R}^2 , denote the topological closure, boundary, and interior as \overline{X} , ∂X , and X° respectively.




Figure 2.1: A planar convex body K with its edge, vertices, supporting line, and half-plane.

Definition 2.1.3. For any angle t in S^1 or \mathbb{R} , define the unit vectors $u_t = (\cos t, \sin t)$ and $v_t = (-\sin t, \cos t)$.

Definition 2.1.4. For any $t \in S^1$ and $h \in \mathbb{R}$, define the line $l(t, h)$ with the *normal angle* t and the signed distance h from the origin as

$$l(t, h) = \{p \in \mathbb{R}^2 : p \cdot u_t = h\}.$$

Definition 2.1.5. For any $t \in S^1$ and $h \in \mathbb{R}$, define the *closed half-planes* $H_{\pm}(t, h)$ and the *open half-planes* $H_{\pm}^{\circ}(t, h)$ with the boundary $l(t, h)$ as the following.

$$\begin{aligned} H_-(t, h) &:= \{p \in \mathbb{R}^2 : p \cdot u_t \leq h\} & H_-^{\circ}(t, h) &:= \{p \in \mathbb{R}^2 : p \cdot u_t < h\} \\ H_+(t, h) &:= \{p \in \mathbb{R}^2 : p \cdot u_t \geq h\} & H_+^{\circ}(t, h) &:= \{p \in \mathbb{R}^2 : p \cdot u_t > h\} \end{aligned}$$

We say that $H_-(t, h)$ and $H_-^{\circ}(t, h)$ have the *normal angle* t and *normal vector* u_t .

Consequently, $H_+(t, h)$ and $H_+^{\circ}(t, h)$ have the normal angle $t + \pi$.

Definition 2.1.6. For any nonempty and compact $S \subseteq \mathbb{R}^2$, define its *support function* $h_S : S^1 \rightarrow \mathbb{R}$ as the value $h_S(t) := \sup \{s \cdot u_t : s \in S\}$.

Definition 2.1.7. For any nonempty and compact $S \subseteq \mathbb{R}^2$ and angle $t \in S^1$, define the *supporting line* $l_S(t)$ of S with *normal angle* t as $l_S(t) := l(t, h_S(t))$. Define the *supporting half-plane* $H_S(t)$ of S with *normal angle* t as $H_S(t) := H_-(t, h_S(t))$.

Definition 2.1.8. For any nonempty and compact S and angle t in S^1 or \mathbb{R} , the *width* of S along the direction of angle t (or unit vector u_t) is the distance between the parallel supporting lines $l_S(t)$ and $l_S(t + \pi)$ of S defined as $h_S(t) + h_S(t + \pi)$.

In this work only, we use the following notions of *vertices* and *edges* of a convex body K .

Definition 2.1.9. For any convex body K and $t \in S^1$, define the *edge* $e_K(t)$ of K as the intersection of K with the supporting line $l_K(t)$.

Definition 2.1.10. For any convex body K and $t \in S^1$, let $v_K^+(t)$ and $v_K^-(t)$ be the endpoints of the edge $e_K(t)$ such that $v_K^+(t)$ is positioned farthest in the direction of v_t and $v_K^-(t)$ is positioned farthest in the opposite direction $-v_t$. We call $v_K^\pm(t)$ the *vertices* of K .

It is possible that the edge $e_K(t)$ can be a single point. In such case, the supporting line $l_K(t)$ makes contact with K at the single point $v_K^+(t) = v_K^-(t)$.

Definition 2.1.11. Define \mathcal{H}^1 as the *Hausdorff measure of dimension one* on \mathbb{R}^2 .

That is, if X is a disjoint union of finite line segments in \mathbb{R}^2 , then $\mathcal{H}^1(X)$ is the sum of all lengths of the line segments.

Definition 2.1.12. Denote the *Hausdorff distance* between convex bodies K_1 and K_2 as d_H . This is the supremum norm between h_{K_1} and h_{K_2} (Lemma 1.8.14, page 66 of [Sch13]).

Definition 2.1.13. Denote the *surface area measure* of a convex body K as σ_K . For simplicity, denote $\sigma_K(\{t\})$ as $\sigma_K(t)$.

We refer to page 214 of [Sch13] for a full construction of σ_K . It measures the side lengths of K , which is the defining property of σ_K .

Theorem 2.1.1. (*Theorem 4.2.3 of [Sch13]*) For any Borel subset X of S^1 , the value $\sigma_K(X)$ is the Hausdorff measure \mathcal{H}^1 of the union $\bigcup_{t \in X} e_K(t)$ of edges of K .

So if K is a convex polygon, then σ_K is a discrete measure such that the measure $\sigma_K(t)$ at point t is the length of the edge $e_K(t)$. For another example, assume that K is a smooth convex polygon where for every $t \in S^1$, the tangent line $l_K(t)$ always meets K at a single point $v(t) = v_K^\pm(t)$ which is smooth in $t \in S^1$. Then the distribution function $R : S^1 \rightarrow \mathbb{R}_{\geq 0}$ of σ_K is the radius of curvature $R(t) = \|v'(t)\|$ of ∂K at $v(t)$.

Proposition 2.1.2. For any convex body K and $t \in S^1$, $\sigma_K(t)$ is the length of the edge $e_K(t)$. Consequently, $v_K^+(t) = v_K^-(t) + \sigma_K(t)v_t$.

Proof. Let $X = \{t\}$ in Theorem 2.1.1. □

We add the following definition.

Definition 2.1.14. Let K be any convex body. For every $a, b \in S^1$ such that $b \neq a, a + \pi$, define $v_K(a, b)$ as the intersection $l_K(a) \cap l_K(b)$.

We prove a technical lemma on the limit of vertices of K .

Theorem 2.1.3. Let K be any convex body and $t \in S^1$ be an arbitrary angle. We have the following right limits converging to $v_K^+(t)$. In particular, the vertex $v_K^+(t)$ is right-continuous on $t \in S^1$.

$$\lim_{s \rightarrow t^+} v_K^+(t) = \lim_{s \rightarrow t^+} v_K^-(u) = \lim_{s \rightarrow t^+} v_K(t, s) = v_K^+(t)$$

Similarly, we have the following left limits.

$$\lim_{s \rightarrow t^-} v_K^+(s) = \lim_{s \rightarrow t^-} v_K^-(s) = \lim_{s \rightarrow t^-} v_K(s, t) = v_K^-(t)$$

Proof. We only compute the right limits. Left limits can be shown using a symmetric argument.

Let $\epsilon > 0$ be arbitrary. Let $p = v_K^+(t) + \epsilon v_t$. By the definition of $v_K^+(t)$, the point p is not in K . As $\mathbb{R}^2 \setminus K$ is open, we can take some positive $\epsilon' < \epsilon$ such that the closed line segment s connecting p and $q = p - \epsilon' u_t$ is disjoint from K as well. Define the closed right-angled triangle T with vertices $v_K^+(t)$, p , and q . Take the line l that passes through both q and $v_K^+(t)$. Call the two closed half-planes divided by the line l as H_T and H' , where H_T contains p and H' does not contain p . By definition, H' has normal angle $s \in (t, t + \pi/2)$.

We show that $K \cap H_T \subseteq T$. Observe that the intersection $X := H_K(t) \cap H_T$ is a cone centered at the point $v_K^+(t)$, with the line segment s dividing X into triangle T and an unbounded convex set $X \setminus T$. Now take any point $r \in K \cap H_T$. If $r \notin T$, then since $r \in X \setminus T$ and $v_K^+(t) \in T$ the line connecting $r \in K$ and $v_K^+(t) \in K$ should pass through a point in s . This, however, contradicts that s is disjoint from convex K . So we should have $r \in T$ and thus $K \cap H_T \subseteq T$.

Now take arbitrary $t_0 \in (t, s)$. We show that the edge $e_K(t_0)$ should lie inside T . It suffices to show that any point z in K that attains the maximum value of $z \cdot u_{t_0}$ is in T . Define the fan $F := H_K(t) \cap H'$, so that F is bounded by lines $l_K(t)$ and l with the vertex $v_K^+(t)$. If $z \in F$, it should be that $z = v_K^+(t) \in T$, because $v_K^+(t) \in K$ and $v_K^+(t) \cdot u_{t_0} > z \cdot u_{t_0}$ for every point z in F other than $z = v_K^+(t)$. If $z \in K \setminus F$ on the other hand, we have $K \setminus F = K \setminus H' \subseteq K \cap H_T \subseteq T$ so $z \in T$. This completes the proof of $e_K(t_0) \subseteq T$.

Observe that the triangle T contains $v_K^+(t)$ and has diameter $< 2\epsilon$ because the two perpendicular sides of T containing p have length $\leq \epsilon$. So the endpoints $v_K^+(u)$ and $v_K^-(u)$ of the edge $e_K(t_0) \subseteq T$ are distance at most 2ϵ away from $v_K^+(t)$. This completes the epsilon-delta argument for $\lim_{s \rightarrow t^+} v_K^+(s) = \lim_{s \rightarrow t^+} v_K^-(s) = v_K^+(t)$.

From $e_K(t_0) \subseteq T$ and that the vertex p of T maximizes the value of $z \cdot u_{t_0}$ over all $z \in T$, we get that p is either on $l_K(t_0)$ or outside the half-plane $g_K(t_0)$. On the other hand we have $v_K^+(t) \in g_K(t_0)$. So the line $l_K(t_0)$ passes through the segment connecting p and $v_K^+(t)$, and the intersection $v_K(t, t_0) = l_K(t) \cap l_K(t_0)$ is inside T . This with that the diameter of T is less than 2ϵ proves $\lim_{s \rightarrow t^+} v_K(t, s) = v_K^+(t)$. \square

2.2 Supporting Hallway

In this Section 2.2, we define the *supporting hallways* $L_S(t)$ of a moving sofa S . We first name the parts of the hallway L .

Definition 2.2.1. Let $\mathbf{x}_L = (0, 0)$ and $\mathbf{y}_L = (1, 1)$ be the inner and outer corner of L respectively.

Let a_L and c_L be the lines $x = 1$ and $y = 1$ representing the *outer walls* of L passing through \mathbf{y}_L . Let \vec{b}_L and \vec{d}_L be the half-lines $\{0\} \times (-\infty, 0]$ and $(-\infty, 0] \times \{0\}$ from the inner corner \mathbf{x}_L representing the *inner walls* of L . Let b_L and d_L be the lines $x = 0$ and $y = 0$ extending \vec{b}_L and \vec{d}_L respectively.

Let $Q_L^+ = (-\infty, 1]^2$ be the closed quarter-plane bounded by outer walls a_L and c_L . Let $Q_L^- = (-\infty, 0)^2$ be the open quarter-plane bounded by inner walls \vec{b}_L and \vec{d}_L , so that $L = Q_L^+ \setminus Q_L^-$.

Now we define the supporting hallways on any nonempty and compact subset of \mathbb{R}^2 .

Definition 2.2.2. For any nonempty and compact $S \subset \mathbb{R}^2$ and angle $t \in S^1$, define the rigid




Figure 2.2: The hallway L (Definition 1.1.1, on left), supporting hallway $L_S(t)$ (Definition 2.2.2, on right), and their corresponding parts (Definition 2.2.1, Definition 2.2.3).

transformation $f_{S,t}$ of \mathbb{R}^2 as

$$f_{S,t}(p) := R_t(p) + (h_S(t) - 1)u_t + (h_S(t + \pi/2) - 1)v_t$$

and define the *supporting hallway* $L_S(t) := f_{S,t}(L)$ of S with angle t .

The following Proposition 2.2.1 is the defining property of $L_S(t)$.

Proposition 2.2.1. *For any shape S and angle $t \in S^1$, the supporting hallway $L_S(t)$ is the unique translation of $R_t(L)$ such that the outer walls of $L_S(t)$ corresponding to the outer walls a and c of L are the tangent lines $l_S(t)$ and $l_S(t + \pi/2)$ of S respectively.*

Proof. Let c_1 and c_2 be arbitrary real values. Then $L' = R_t(L) + c_1u_t + c_2v_t$ is an arbitrary rigid transformation of L rotated counterclockwise by t . The outer walls of L' corresponding to the outer walls a and c of L (Definition 2.2.1) are $l(t, c_1+1)$ and $l(t+\pi/2, c_2+1)$ respectively. They match with the tangent lines $l_S(t) = l(t, h_S(t))$ and $l_S(t + \pi/2) = l(t + \pi/2, h_S(t + \pi/2))$ of S if and only if $c_1 = h_S(t) - 1$ and $c_2 = h_S(t + \pi/2) - 1$. That is, if and only if $L' = L_S(t)$. \square

Name the parts of supporting hallway $L_S(t)$ corresponding to the parts of L (Definition 2.2.1) under the transformation $f_{S,t}$ for future use.

Definition 2.2.3. For any shape S and angle $t \in S^1$, let $\mathbf{x}_S(t), \mathbf{y}_S(t), a_S(t), b_S(t), c_S(t), d_S(t), \vec{b}_S(t), \vec{d}_S(t), Q_S^+(t), Q_S^-(t)$ be the parts of $L_S(t)$ corresponding to the parts $\mathbf{x}_L, \mathbf{y}_L, a_L, b_L, c_L, d_L, \vec{b}_L, \vec{d}_L, Q_L^+, Q_L^-$ of L respectively with $f_{S,t}$. That is, for any $? = \mathbf{x}, \mathbf{y}, a, b, c, d, \vec{b}, \vec{d}, Q^+, Q^-$, let $?_S(t) := f_{S,t}(?_L)$.

Proposition 2.2.2. *We have $L_S(t) = Q_S^+(t) \setminus Q_S^-(t)$. Also we can express the parts of $L_S(t)$*

purely in terms of the supporting lines, half-planes, and support function h_S of S as below.

$$\begin{aligned}
\mathbf{x}_S(t) &= (h_S(t) - 1)u_t + (h_S(t + \pi/2) - 1)v_t \\
\mathbf{y}_S(t) &= h_S(t)u_t + h_S(t + \pi/2)v_t \\
a_S(t) &= l_S(t) = l(t, h_S(t)) & b_S(t) &= l(t, h_S(t) - 1) \\
c_S(t) &= l_S(t + \pi/2) = l(t + \pi/2, h_S(t + \pi/2)) \\
d_S(t) &= l(t + \pi/2, h_S(t + \pi/2) - 1) \\
Q_S^+(t) &= H_S(t) \cap H_S(t + \pi/2) = H_-(t, h_S(t)) \cap H_-(t + \pi/2, h_S(t + \pi/2)) \\
Q_S^-(t) &= H_-^\circ(t, h_S(t) - 1) \cap H_-^\circ(t + \pi/2, h_S(t + \pi/2))
\end{aligned}$$

Proof. Follows from Definition 2.2.2 and Proposition 2.2.1. \square

With the following Proposition 2.2.3, we can assume that the rotating hallways L_t in Proposition 1.2.2 containing a moving sofa S are the supporting hallways $L_t = L_S(t)$ of S .

Proposition 2.2.3. *Let S be any nonempty compact set contained in a translation of $R_t(L)$ with angle $t \in S^1$. Then the supporting hallway $L_S(t)$ with angle t also contains S .*

Proof. Assume that a translation L' of $R_t(L)$ contains S . Then $L' = f(L)$ for some rigid transformation f with counterclockwise rotation t . Define $Q'^+ = f(Q_L^+)$ and $Q'^- = f(Q_L^-)$ as the quarter-planes of L' corresponding to that of L . Then $L' = Q'^+ \setminus Q'^-$ and Q'^+ is a convex cone containing S with boundaries of normal angles t and $t + \pi/2$. By Proposition 2.2.2, $Q_S^+(t)$ is the intersection of two supporting half-planes of S with normal angles t and $t + \pi/2$. So we should have $Q_S^+(t) \subseteq Q'^+$. Shifting this by $-u_t - v_t$, we get $Q_S^-(t) \subseteq Q'^-$. Now $S \subset L'$ is disjoint from $Q_S^-(t) \subseteq Q'^-$, and we have $S \subseteq Q_S^+(t) \setminus Q_S^-(t) = L_S(t)$. \square

2.3 Monotone Sofa

We now define the notion of monotone sofas. We first prepare basic definitions.

Definition 2.3.1. Let $R_t : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote the rotation of \mathbb{R}^2 around the origin by the counterclockwise angle of $t \in \mathbb{R}$.

Definition 2.3.2. Define the horizontal strip $H := \mathbb{R} \times [0, 1]$, vertical strip $V := [0, 1] \times \mathbb{R}$, and its rotation $V_\omega := R_\omega(V)$ around the origin by a counterclockwise angle $\omega \in \mathbb{R}$.

Definition 2.3.3. The *rotation angle* ω of a moving sofa S is the clockwise angle that it rotates as it moves from H_L to V_L inside L .

Definition 2.3.4. A moving sofa S with rotation angle $\omega \in (0, \pi/2]$ is in *standard position* if $h_S(\omega) = h_S(\pi/2) = 1$.

We will assume without loss of generality that a moving sofa S can be put in standard position after some translation.

Definition 2.3.5. For any $\omega \in (0, \pi/2]$, define the *parallelogram* $P_\omega := H \cap V_\omega$ with rotation angle ω . Let $O := (0, 0)$ and $o_\omega := (\tan(\pi/4 - \omega/2), 1)$ represent the lower left and upper right vertices of P_ω respectively.

Proposition 2.3.1. *For any moving sofa S with rotation angle $\omega \in (0, \pi/2]$, there is a translation of S in standard position which is (i) unique if $\omega < \pi/2$, or (ii) unique up to horizontal translations if $\omega = \pi/2$. After such a translation, we have $S \subseteq P_\omega$.*

Proof. If $\omega < \pi/2$, there is a unique translation of S making contact with the two supporting lines $l(\omega, 1)$ and $l(\pi/2, 1)$ from below. If $\omega = \pi/2$, there is a unique translation of S making contact with the supporting line $l(\pi/2, 1)$ from below, up to horizontal translations. We have $S \subseteq H$ and $S \subseteq V_\omega$ by the proof of Proposition 1.2.2. \square

Define the intersection \mathcal{I} in Equation (1.2) of Section 1.2 with the supporting hallways $L_t := L_S(t)$.

Definition 2.3.6. Let S be any moving sofa with rotation angle $\omega \in (0, \pi/2]$ in standard position. Define the intersection

$$\mathcal{I}(S) = P_\omega \cap \bigcap_{t \in [0, \omega]} L_S(t).$$

We will establish that $\mathcal{I}(S)$ is a moving sofa containing S .

Theorem 2.3.2. *For any moving sofa S with rotation angle $\omega \in (0, \pi/2]$ in standard position, $\mathcal{I}(S)$ is a moving sofa with the same rotation angle ω in standard position containing S .*

With Theorem 2.3.2, we will call any sofa of form $\mathcal{I}(S)$ a *monotone sofa*, and it suffices to consider monotone sofas for the moving sofa problem.

Definition 2.3.7. Define a *monotone sofa* as the intersection $\mathcal{I}(S)$ of some moving sofa S in standard position.

Proof of Theorem 2.3.2

Proposition 2.3.3. *For any moving sofa S in standard position, $S \subseteq \mathcal{I}(S)$.*

Proof. Assume rotation angle $\omega \in (0, \pi/2]$. We have $S \subseteq P_\omega$ by Proposition 2.3.1. For all $t \in [0, \omega]$, we have $S \subset L_t$ for some hallway L_t rotated counterclockwise by t from (2) of Proposition 1.2.2, and then $S \subseteq L_S(t)$ by Proposition 2.2.3. \square

We prepare the following terminologies.

Definition 2.3.8. Say that a set $X \subseteq \mathbb{R}^2$ is *closed in the direction of vector $v \in \mathbb{R}^2$* if, for any $x \in X$ and $\lambda \geq 0$, we have $x + \lambda v \in X$.

Definition 2.3.9. Any line l of \mathbb{R}^2 divides the plane into two half-planes. Assuming l is not parallel to the y -axis, call the *left side* (resp. *right side*) of l as the closed half-plane with boundary l containing the point $-Nu_0$ (resp. Nu_0) for sufficiently large N .

Definition 2.3.10. Let S be any moving sofa with rotation angle $\omega \in (0, \pi/2]$ in standard position. Define the convex set

$$\mathcal{C}(S) := P_\omega \cap \bigcap_{0 \leq t \leq \omega} Q_S^+(t).$$

We prepare useful lemmas on $\mathcal{C}(S)$.

Proposition 2.3.4. *For any moving sofa S in standard position, $S \subseteq \mathcal{I}(S) \subseteq \mathcal{C}(S)$.*

Proof. By Proposition 2.3.3 and $L_S(t) \subset Q_S^+(t)$ for all $t \in [0, \omega]$. \square

Definition 2.3.11. For any $\omega \in [0, \pi/2]$, define the set $J_\omega := [0, \omega] \cup [\pi/2, \omega + \pi/2]$.

Lemma 2.3.5. *Let S be any moving sofa with rotation angle $\omega \in [0, \pi/2]$ in standard position. Then the support functions h_X of the sets $X = S, \mathcal{I}(S), \mathcal{C}(S)$ are the same on the set J_ω . Consequently, for any $t \in [0, \omega]$, the supporting hallways $L_X(t)$ on the sets $X = S, \mathcal{I}(S), \mathcal{C}(S)$ are the same.*

Proof. We have $S \subseteq \mathcal{I}(S) \subseteq \mathcal{C}(S)$ by Proposition 2.3.4. So it remains to show $h_{\mathcal{C}(S)}(t) \leq h_S(t)$ for every t in J_ω to show that h_X s on J_ω are the same. This follows from Definition 2.3.10 as $\mathcal{C}(S) \subseteq H_S(t)$ for any $t \in J_\omega$. To show that the supporting hallways $L_X(t)$ are the same, observe that $L_X(t)$ depends solely on the values $t, h_X(t)$, and $h_X(t + \pi/2)$ by its Definition 2.2.2. \square

We establish the connectedness of $\mathcal{I}(S)$ which is the hardest part.

Theorem 2.3.6. *For any moving sofa S with rotation angle $\omega \in (0, \pi/2]$ in standard position, $\mathcal{I}(S)$ is connected.*

Proof. Fix an arbitrary point p in $\mathcal{I}(S)$. It suffices to show that $\mathcal{I}(S)$ is connected by finding a line segment s_θ inside $\mathcal{I}(S)$ that connects p to the connected subset S of $\mathcal{I}(S)$. Here $\theta \in [\omega, \pi/2]$ is a value that will be fixed later. Letting $\theta \in [\omega, \pi/2]$ arbitrary as of now, define the line l_θ passing through p in the direction of u_θ and let $s_\theta := l_\theta \cap \mathcal{I}(S)$. Then s_θ is a subset of $\mathcal{I}(S)$ containing p .

Our goal now is to show that s_θ is a line segment that overlaps with S for some $\theta \in [\omega, \pi/2]$. We first show that s_θ is a nonempty line segment. Define the set $X := \bigcup_{0 \leq t \leq \omega} Q_S^-(t)$. By plugging $L_S(t) = Q_S^+(t) \setminus Q_S^-(t)$ in Proposition 2.2.2 to Definition 2.3.6, we have $\mathcal{I}(S) = \mathcal{C}(S) \setminus X$. The set $\mathcal{C}(S)$ is a convex body containing S by Proposition 2.3.1, and the set X is closed in the direction of $-u_\theta$ (Definition 2.3.8) since each $Q_S^-(t)$ is. Now s_θ is a line segment because it is the line segment $l_\theta \cap \mathcal{C}(S)$ subtracted by the half-line $l_\theta \setminus X$. Our goal now is to find some $\theta \in [\omega, \pi/2]$ such that l_θ meets S , so that s_θ connects p to S inside $\mathcal{I}(S)$.

Assume by contradiction that for every $\theta \in [\omega, \pi/2]$ the line l_θ is disjoint from S . Because the line l_θ is disjoint from S for any $\theta \in [\omega, \pi/2]$, the set S is inside the set $Y = \mathbb{R}^2 \setminus \bigcup_{\theta \in [\omega, \pi/2]} l_\theta$. Note that Y has exactly two connected components Y_L and Y_R on the left and the right side of the lines l_θ respectively. We will find a point at each $S \cap Y_L$ and $S \cap Y_R$, reaching the contradiction as S is connected.

By Lemma 2.3.5, we have $l_{\mathcal{I}(S)}(t) = l_S(t)$ for every $t \in J_\omega = [0, \omega] \cup [\pi/2, \omega + \pi/2]$. Because $p \in \mathcal{I}(S)$, the line $l_{\pi/2}$ passing through p is on the left side of $l_S(0)$. So any point of $e_S(0)$ is on the right side of $l_{\pi/2}$, and should be in $S \cap Y_R$. Likewise, as $p \in \mathcal{I}(S)$, the line l_ω passing through p is on the right side of $l_S(\omega + \pi/2)$. So any point of $e_S(\omega + \pi/2)$ is on the left side of l_ω , and should be in $S \cap Y_L$. This establishes the contradiction we wanted, and we finally prove that $\mathcal{I}(S)$ is connected. \square

We are now ready to prove Theorem 2.3.2.

Proof of Theorem 2.3.2. We first show that $S' := \mathcal{I}(S)$ is a moving sofa. As S' is connected by Theorem 2.3.6, it suffices to show that S' can move continuously inside L from H_L to V_L with rotation angle ω . See the movement of $L_S(t) = L_{S'}(t)$ containing S' for $t \in [0, \omega]$ (Lemma 2.3.5) in perspective of the hallway, to find a movement of S' inside L . In particular,

$S' \subseteq H$ so the horizontal side of $L_{S'}(0)$ (corresponding to H_L of L) contains S' . Also, $S' \subseteq V_\omega$ so the vertical side of $L_{S'}(\omega)$ (corresponding to V_L of L) contains S' . Since $h_{S'}$ is continuous, this movement of S' is also continuous.

Because $L_{S'}(\omega)$ is rotated counterclockwise by ω , the sofa S' have rotation angle ω . By Proposition 2.3.3, the sofa S' contains S . The sofa S' is in standard position because $S \subseteq S' \subseteq H \cap V_\omega$ and S is in standard position. \square

2.4 Cap and Niche

We show that any intersection $\mathcal{I}(S)$ of a moving sofa S is equal to the *cap* K of S minus the *niche* $\mathcal{N}(K)$ of K (Theorem 2.4.2). We first define the notion of *cap* as a kind of convex body.

Definition 2.4.1. A *cap* K with *rotation angle* $\omega \in (0, \pi/2]$ is a convex body such that the followings hold.

1. $h_K(\omega) = h_K(\pi/2) = 1$ and $h_K(\omega + \pi) = h_K(3\pi/2) = 0$.
2. K is an intersection of closed half-planes with normal angles in $J_\omega \cup \{\omega + \pi, 3\pi/2\}$.

Definition 2.4.2. Define the *space of caps* \mathcal{K}_ω^c with the *rotation angle* $\omega \in (0, \pi/2]$ as the collection of all caps K with rotation angle ω .

We now show that the convex set $\mathcal{C}(S)$ from a moving sofa S (Definition 2.3.10) is a cap, justifying calling $\mathcal{C}(S)$ *the cap of S* .

Theorem 2.4.1. For any moving sofa S with rotation angle $\omega \in (0, \pi/2]$ in standard position, the set $\mathcal{C}(S)$ in Definition 2.3.10 is a cap with rotation angle ω as in Definition 2.4.1.

Proof. The second condition of Definition 2.4.1 on $\mathcal{C}(S)$ is satisfied by Definition 2.3.10. Since S is in standard position, it suffices to check $h_{\mathcal{C}(S)}(\omega + \pi) = h_{\mathcal{C}(S)}(3\pi/2) = 0$ in the first condition of Definition 2.4.1.

We first prove the case $\omega = \pi/2$. Since $h_S(\pi/2) = 1$ and $S \subseteq \mathcal{C}(S)$ by Proposition 2.3.4, we can take a point $q \in S \cap l(\pi/2, 1) \subseteq \mathcal{C}(S)$. For any $t \in [0, \pi/2]$, each $Q_S^+(t)$ is closed in the direction of $-v_0$ (Definition 2.3.8). This with $q \in \mathcal{C}(S)$ implies that $q - v_0 \in \mathcal{C}(S)$. Note that $q - v_0 \in l(\pi/2, 0)$, so $q - v_0 \in \mathcal{C}(S) \subseteq P_{\pi/2}$ implies $h_{\mathcal{C}(S)}(3\pi/2) = 0$ as we desired.

We now prove the case $\omega < \pi/2$. Since $h_S(\omega) = h_S(\pi/2) = 1$, we can take two points $q_\omega \in S \cap l(\omega, 1)$ and $q_{\pi/2} \in S \cap l(\pi/2, 1)$. Observe that the three points $q_\omega, o_\omega, q_{\pi/2}$ are in monotonically decreasing order of x -coordinates and form an angle of $\omega + \pi/2$. Take any supporting hallway $L_S(t)$ with angle $t \in [0, \omega]$. Then $Q_S^+(t)$ should contain both $q_\omega, q_{\pi/2} \in S$, so we also have $Q_S^+(t) \ni o_\omega$. This implies $o_\omega \in \mathcal{C}(S)$. For any $t \in [0, \omega]$, each $Q_S^+(t)$ is closed in the directions v_0 and u_ω (Definition 2.3.8). So $o_\omega \in \mathcal{C}(S)$ implies $o_\omega - v_0, o_\omega - u_\omega \in \mathcal{C}(S)$. This with $\mathcal{C}(S) \subseteq H \cap V_\omega$ implies $h_{\mathcal{C}(S)}(\omega + \pi) = h_{\mathcal{C}(S)}(3\pi/2) = 0$ as we desired. \square

Definition 2.4.3. With Theorem 2.4.1, call $\mathcal{C}(S)$ *the cap of the moving sofa S* .

We now define the *niche* of a cap. Note that the following fan F_ω contains $H \cap V_\omega$ in particular.

Definition 2.4.4. For any angle $\omega \in [0, \pi/2]$, define the *fan* $F_\omega := H_+(\omega, 0) \cap H_+(\pi/2, 0)$.

Definition 2.4.5. Let $K \in \mathcal{K}_\omega^c$ be arbitrary. Define the *niche* of K as

$$\mathcal{N}(K) := F_\omega \cap \bigcup_{t \in (0, \omega)} Q_K^-(t).$$

Theorem 2.4.2. Let S be a moving sofa with rotation angle $\omega \in (0, \pi/2]$ in standard position. The monotone sofa $\mathcal{I}(S)$ from S is equal to $K \setminus \mathcal{N}(K)$ where $K := \mathcal{C}(S)$ is the cap of S .

Proof. By writing each $L_S(t)$ as $Q_S^+(t) \setminus Q_S^-(t)$, the set $\mathcal{I}(S)$ can be represented as follows.

$$\begin{aligned} \mathcal{I}(S) &= P_\omega \cap \bigcap_{t \in [0, \omega]} L_S(t) \\ &= \left(P_\omega \cap \bigcap_{t \in [0, \omega]} Q_S^+(t) \right) \setminus \left(F_\omega \cap \bigcup_{t \in [0, \omega]} Q_S^-(t) \right) \end{aligned} \quad (2.1)$$

By Lemma 2.3.5 we have $Q_S^-(t) = Q_K^-(t)$. So we have $\mathcal{I}(S) = K \setminus \mathcal{N}(K)$ by the definitions of K and $\mathcal{N}(K)$. \square

A monotone sofa is its cap subtracted by its niche. Note that unlike Theorem 2.4.2 above, the following Theorem 2.4.3 does not depend on another moving sofa.

Theorem 2.4.3. For any monotone sofa S with cap $K := \mathcal{C}(S)$, we have $S = K \setminus \mathcal{N}(K)$.

Proof. Let $S = \mathcal{I}(S')$ for some moving sofa S' in standard position. By Lemma 2.3.5 and that Definition 2.3.10 depends solely on the values of support function on J_ω , the cap $K = \mathcal{C}(S)$ of S is also the cap $\mathcal{C}(S')$ of S' . So by Theorem 2.4.2 we have $S = \mathcal{I}(S') = K \setminus \mathcal{N}(K)$. \square

Taking the intersection $\mathcal{I}(-)$ enlarges any moving sofa to monotone sofas and fixes monotone sofas.

Theorem 2.4.4. For any moving sofa S' in standard position, we have $\mathcal{I}(\mathcal{I}(S')) = \mathcal{I}(S')$. Consequently, the equality $S = \mathcal{I}(S)$ holds if and only if S is a monotone sofa.

Proof. Let $S := \mathcal{I}(S')$ so that S is a monotone sofa with cap K . Then $\mathcal{I}(S) = K \setminus \mathcal{N}(K) = S$ by Theorem 2.4.2 and Theorem 2.4.3. So $\mathcal{I}(\mathcal{I}(S')) = \mathcal{I}(S')$ and the equality $S = \mathcal{I}(S)$ holds for monotone sofa S . On the other hand, any moving sofa S with equality $S = \mathcal{I}(S)$ is immediately a monotone sofa. \square

2.5 Cap Contains Niche

We now define the parts of an arbitrary cap K .

Definition 2.5.1. Let $K \in \mathcal{K}_\omega^c$ be arbitrary. For any $t \in [0, \omega]$, define the *vertices* $A_K^+(t) = v_K^+(t)$, $A_K^-(t) = v_K^-(t)$, $C_K^+(t) = v_K^+(t + \pi/2)$, and $C_K^-(t) = v_K^-(t + \pi/2)$ of K .

Note that the outer wall $a_K(t)$ (resp. $c_K(t)$) of $L_K(t)$ is in contact with the cap K at the vertices $A_K^+(t)$ and $A_K^-(t)$ (resp. $C_K^+(t)$ and $C_K^-(t)$) respectively. We also define the *upper boundary* of a cap K .

Definition 2.5.2. Let $K \in \mathcal{K}_\omega^c$ be arbitrary. Define the *upper boundary* δK of K as the set $\delta K = \bigcup_{t \in [0, \omega + \pi/2]} e_K(t)$.

For any cap K with rotation angle ω , the upper boundary δK is exactly the points of K making contact with the outer walls $a_K(t)$ and $c_K(t)$ of supporting hallways $L_K(t)$ for every $t \in [0, \omega]$. In particular, upper boundary δK is a curve from the right endpoint $A_{\bar{K}}(0)$ to the left endpoint $C_{\bar{K}}^+(\omega)$. We collect some observations on δK .

Proposition 2.5.1. *Let $K \in \mathcal{K}_\omega^c$ be arbitrary. The set δK is the boundary of K in the subset topology of F_ω .*

Proof. Let X be the boundary of K in the subset topology of F_ω . We first show $\delta K \subseteq X$. Take any point z of δK . Then $z \in e_K(t)$ for some $t \in [0, \omega + \pi/2]$. Since K is a planar convex body, for any $\epsilon > 0$ the point $z' = z + \epsilon u_t$ is not in K . This with $z \in K$ implies $z \in X$. We now show $X \subseteq \delta K$. Assume by contrary that there is a point $z \in X \setminus \delta K$. Then for every $t \in [0, \omega + \pi/2]$ we have $z \notin e_K(t)$ so that z is in the interior of $H_K(t)$. So by compactness of $[0, \omega + \pi/2]$, an open ball U of radius ϵ centered at z is contained in the half-space $H_K(t)$ for all $t \in [0, \omega + \pi/2]$. Now $U \cap F_\omega \subseteq K$ and so $z \notin X$, leading to contradiction. \square

Proposition 2.5.2. *For any cap K , its upper boundary δK is connected.*

Proof. Let $\omega \in (0, \pi/2]$ be the rotation angle of K . Let $I := [0, \omega + \pi/2]$. Assume contradictory that $\delta K = \bigcup_{t \in I} e_K(t)$ is disconnected. That is, there are open subsets U, V of \mathbb{R}^2 such that δK is the disjoint union of nonempty subsets $\delta K \cap U$ and $\delta K \cap V$. For any $t \in I$, each $e_K(t) \subset \delta K$ is connected so it should be contained in exactly one of U or V . Now define $I_U := \{t \in I : e_K(t) \subseteq U\}$ and $I_V := \{t \in I : e_K(t) \subseteq V\}$. Then I is a disjoint union of I_U and I_V . By Theorem 2.1.3 and that each $e_K(t)$ is the line segment connecting $v_{\bar{K}}(t)$ to $v_{\bar{K}}^+(t)$, the sets I_U and I_V are open in the subspace topology of I . Since I is connected, either $I_U = I$ or $I_V = I$, and they imply either $\delta K \subset U$ or $\delta K \subset V$, leading to contradiction. \square

We also define some notions on the niche $\mathcal{N}(K)$ of a cap K .

Definition 2.5.3. For any $K \in \mathcal{K}_\omega^c$ and $t \in (0, \omega)$, define the *wedge* $T_K(t) := F_\omega \cap Q_{\bar{K}}^-(t)$ of K with angle t .

Proposition 2.5.3. *For any $K \in \mathcal{K}_\omega^c$, we have $\mathcal{N}(K) = \bigcup_{t \in (0, \omega)} T_K(t)$.*

Proof. Immediate from Definition 2.4.5. \square

The following Definition 2.5.4 defines the left and right endpoints of the wedge $T_K(t)$.

Definition 2.5.4. For any $K \in \mathcal{K}_\omega^c$ and $t \in (0, \omega)$, define $W_K(t)$ as the intersection of lines $b_K(t)$ and $l(\pi/2, 0)$, and define $Z_K(t)$ as the intersection of lines $d_K(t)$ and $l(\omega, 0)$.

Note that if the wedge $T_K(t)$ contains the origin O , then $T_K(t)$ is a quadrilateral with vertices $O, W_K(t), Z_K(t)$, and $\mathbf{x}_K(t)$, and the points $W_K(t)$ and $Z_K(t)$ are the leftmost and rightmost point of $T_K(t)$ respectively.

Definition 2.5.5. For any cap K with rotation angle ω and $t \in (0, \omega)$, define the *right wedge gap* $w_K(t) := (A_{\bar{K}}^-(0) - W_K(t)) \cdot u_0$ with angle t , which is the signed distance from $W_K(t)$ to $A_{\bar{K}}^-(0)$ along the line $l(\pi/2, 0)$ in the direction of u_0 . Likewise, define the *left wedge gap* $z_K(t) = (C_{\bar{K}}^+(\omega) - Z_K(t)) \cdot v_\omega$ with angle t , which is the signed length from $Z_K(t)$ to $C_{\bar{K}}^+(\omega)$ along the line $l(\omega, 0)$ in the direction of v_ω .

We introduce the notion of reflecting a cap K that will reduce symmetric arguments without loss of generality.

Definition 2.5.6. Let $\omega \in (0, \pi/2]$ be arbitrary. Define $M_\omega : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ as the reflection along the line passing through O and o_ω . For any cap K with rotation angle ω , define the *mirror reflection* $K^m := M_\omega(K)$ of K .

Many definitions on K are symmetric along the reflection M_ω .

Proposition 2.5.4. Let $K \in \mathcal{K}_\omega^c$ be arbitrary. The parts of supporting hallway $L_K(t)$, cap K , and niche $\mathcal{N}(K)$ are equivariant under M_ω . That is, for any $t \in [0, \omega]$:

- $?_{K^m}(t) = M_\omega(?_K(\omega - t))$ for $? = L, \mathbf{x}, \mathbf{y}, a, b, c, d, W, Z$.
- $A_{K^m}^\pm(t) = M_\omega(C_{K^m}^\mp(\omega - t))$ and $C_{K^m}^\pm(t) = M_\omega(A_{K^m}^\mp(\omega - t))$.
- $w_{K^m}(t) = z_K(\omega - t)$ and $z_{K^m}(t) = w_K(\omega - t)$.
- $\delta K^m = M_\omega(\delta K)$, $T_{K^m}(t) = M_\omega(T_K(\omega - t))$, and $\mathcal{N}(K^m) = M_\omega(\mathcal{N}(K))$.
- $\sigma_{K^m}(E) = \sigma_K(\omega + \pi/2 - E)$ for any $E \subseteq S^1$.

Proof. Check the symmetry of the definition of each. For the surface area measure σ_{K^m} , use Equation (4.14), page 215 of [Sch13]. \square

Remark 2.5.1. We use the mirror reflection of a cap in Definition 2.5.6 and Proposition 2.5.4 extensively to exploit the symmetry without loss of generality. For example, say we want to show both $w_K(t) > 0$ and $z_K(t) > 0$ for arbitrary $K \in \mathcal{K}_\omega^c$ and $t \in (0, \omega)$. Then it suffices to show the case $w_K(t) > 0$, as the symmetric case $z_K(t) = w_{K^m}(\omega - t) > 0$ follows by Proposition 2.5.4.

We now show that for any monotone sofa S , its cap K contains the niche $\mathcal{N}(K)$ (Theorem 2.5.9). The positivity of $w_K(t)$ and $z_K(t)$ in Theorem 2.5.5 is important in establishing $\mathcal{N}(K) \subset K$.

Definition 2.5.7. Say that a point p_1 is *further than* (resp. *strictly further than*) the point p_2 in the direction of nonzero vector $v \in \mathbb{R}^2$ if $p_1 \cdot v \geq p_2 \cdot v$ (resp. $p_1 \cdot v > p_2 \cdot v$).

Theorem 2.5.5. Let $K \in \mathcal{K}_\omega^c$ be arbitrary. For any angle $t \in (0, \omega)$, we have $w_K(t), z_K(t) > 0$.

Proof. By mirror symmetry (Remark 2.5.1), it suffices to show $w_K(t) > 0$. We need to show that the point $A_{K^-}(0)$ is strictly further than the point $W_K(t)$ in the direction of u_0 (Definition 2.5.7). The point $q := a_K(t) \cap l(\pi/2, 1)$ is strictly further than $W_K(t) = b_K(t) \cap l(\pi/2, 0)$ in the direction of u_0 , because the lines $a_K(t)$ and $b_K(t)$ form the boundary of a unit-width vertical strip rotated counterclockwise by t . The points $q = l_K(t) \cap l_K(\pi/2)$, $A_{K^-}(t), A_{K^-}(0)$ are consecutively further in the direction of u_0 because K is a convex body, completing the proof. \square

Lemma 2.5.6. Fix an arbitrary $K \in \mathcal{K}_\omega^c$ and an angle $t \in (0, \omega)$. If the inner corner $\mathbf{x}_K(t)$ is in K , then the wedge $T_K(t)$ is a subset of K .

Proof. Assume $\mathbf{x}_K(t) \in K$. If $\omega = \pi/2$, then by $\mathbf{x}_K(t) \in K$ the wedge $T_K(t)$ is the triangle with vertices $W_K(t)$, $\mathbf{x}_K(t)$, and $Z_K(t)$ in counterclockwise order. Also $W_K(t)$ is further than $Z_K(t)$ in the direction of u_0 (Definition 2.5.7). This with $w_K(t), z_K(t) > 0$ (Theorem 2.5.5) implies that all the three vertices of $T_K(t)$ are in K .

If $\omega < \pi/2$, we divide the proof into four cases on whether the origin $O = (0, 0)$ lies strictly below the lines $b_K(t)$ and $d_K(t)$ or not respectively.

- If $(0, 0)$ lies on or above both $b_K(t)$ and $d_K(t)$, then the corner $\mathbf{x}_K(t)$ should be outside F_ω° and this contradicts $\mathbf{x}_K(t) \in K$.
- If $(0, 0)$ lies on or above $b_K(t)$ but lies strictly below $d_K(t)$, then $T_K(t)$ is a triangle with vertices $\mathbf{x}_K(t)$, $Z_K(t)$ and the intersection $p := l(\omega, 0) \cap b_K(t)$ in clockwise order, with the point p on the line segment connecting $Z_K(t)$ and $(0, 0)$. As $z_K(t) > 0$ (Theorem 2.5.5) the point $Z_K(t)$ lies in the segment connecting $C_K^+(\omega)$ and the origin $(0, 0)$. So the vertices $\mathbf{x}_K(t)$, $Z_K(t)$, p of $T_K(t)$ are in K , showing $T_K(t) \subseteq K$.
- If $(0, 0)$ lies strictly below $b_K(t)$ but lies on or above $d_K(t)$, apply the previous case to the mirror reflection K^m and reflect back (Remark 2.5.1).
- If $(0, 0)$ lies strictly below both $b_K(t)$ and $d_K(t)$, then $T_K(t)$ is a quadrilateral with vertices $\mathbf{x}_K(t)$, $Z_K(t)$, $W_K(t)$ and $(0, 0)$. As $w_K(t) > 0$ (resp. $z_K(t) > 0$) by Theorem 2.5.5, the point $W_K(t)$ (resp. $Z_K(t)$) is in the line segment connecting $(0, 0)$ and $A_K^-(0)$ (resp. $C_K^+(\omega)$). So all vertices of $T_K(t)$ are in K and $T_K(t) \subseteq K$.

□

Lemma 2.5.7. *For any $K \in \mathcal{K}_\omega^c$, we have $A_K^-(0), C_K^+(\omega) \in K \setminus \mathcal{N}(K)$.*

Proof. We only need to show that $A_K^-(0), C_K^+(\omega)$ are not in $\mathcal{N}(K)$. That is, for any $t \in (0, \omega)$, neither points are in $T_K(t)$. Since $w_K(t) > 0$ by Theorem 2.5.5, the point $A_K^-(0)$ is on the right side of the boundary $b_K(t)$ of $T_K(t)$. So $A_K^-(0) \notin T_K(t)$. Similarly, $z_K(t) > 0$ implies $C_K^+(\omega) \notin T_K(t)$. □

We identify the exact condition where $\mathcal{N}(K) \subseteq K$ for a general cap K .

Theorem 2.5.8. *For any $K \in \mathcal{K}_\omega^c$, the followings are all equivalent.*

1. $\mathcal{N}(K) \subseteq K$
2. $\mathcal{N}(K) \subseteq K \setminus \delta K$
3. For every $t \in (0, \omega)$, either $\mathbf{x}_K(t) \notin F_\omega^\circ$ or $\mathbf{x}_K(t) \in K$.
4. The set $S = K \setminus \mathcal{N}(K)$ is connected.

Proof. (1) and (2) are equivalent because the niche $\mathcal{N}(K)$ is open in the subset topology of F_ω by Definition 2.4.5, and the set $K \setminus \delta K$ is the interior of K in the subset topology of F_ω by Proposition 2.5.1.

(1 \Rightarrow 3) We will prove the contraposition and assume $\mathbf{x}_K(t) \in F_\omega^\circ \setminus K$. Then a neighborhood of $\mathbf{x}_K(t)$ is inside F_ω and disjoint from K , so a subset of $T_K(t)$ is outside K and $\mathcal{N}(K) \not\subseteq K \setminus \delta K$. (3 \Rightarrow 1) If $\mathbf{x}_K(t) \notin F_\omega^\circ$ then $T_K(t)$ is an empty set. If $\mathbf{x}_K(t) \in K$ then by Lemma 2.5.6 we have $T_K(t) \subseteq K$.

(2 \Rightarrow 4) As δK is disjoint from $\mathcal{N}(K)$, we have $\delta K \subseteq S$. We show that S is connected. The set δK is connected by Proposition 2.5.2. Take any point $p \in S$. Take the half-line r starting from p in the upward direction v_0 . Then r touches a point in δK as $p \in K$. Moreover, r is disjoint from $\mathcal{N}(K)$ as the set $\mathcal{N}(K) \cup (\mathbb{R}^2 \setminus F_\omega)$ is closed in the direction $-v_0$ (Definition 2.3.8). Now $r \cap K$ is a line segment inside S connecting arbitrary $p \in S$ to a point in δK . So S is connected.

(4 \Rightarrow 3) Assume by contradiction that $\mathbf{x}_K(t) \in F_\omega^\circ \setminus K$ for some $t \in (0, \omega)$. The ray with initial point $\mathbf{x}_K(t)$ and direction v_0 is disjoint from K , as $F_\omega^\circ \setminus K$ is closed in the direction v_0 . The ray with initial point $\mathbf{x}_K(t)$ and the opposite direction $-v_0$ is not in S , as $\mathbf{x}_K(t)$ is the corner of $Q_K^-(t)$, and $Q_K^-(t)$ is closed in the direction of $-v_0$. So the vertical line l passing through $\mathbf{x}_K(t)$ in the direction of v_0 is disjoint from S .

By $z_K(t) > 0$ and $\mathbf{x}_K(t) \in F_\omega^\circ$, the points $C_K^+(\omega)$, $Z_K(t)$, $\mathbf{x}_K(t)$ are consecutively strictly further in the direction of u_0 . Likewise, by $w_K(t) > 0$ and $\mathbf{x}_K(t) \in F_\omega^\circ$, the points $\mathbf{x}_K(t)$, $W_K(t)$, $A_K^-(0)$ are consecutively strictly further in the direction of u_0 . So the point $C_K^+(\omega)$ lies strictly left to l , and $A_K^-(0)$ lies strictly right to l . By Lemma 2.5.7 we have $C_K^+(\omega), A_K^-(0) \in S$, and this with S disjoint from l contradicts that that S is connected. \square

Theorem 2.5.9. *Let $K \in \mathcal{K}_\omega^c$ be arbitrary. Then K is the cap of a monotone sofa if and only if K contains $\mathcal{N}(K)$.*

Proof. Assume first that K is the cap of a monotone sofa S . We have $S = K \setminus \mathcal{N}(K)$ by Theorem 2.4.3. In particular, $K \setminus \mathcal{N}(K)$ is a moving sofa so it is connected. Since (4) implies (1) in Theorem 2.5.8 we have $\mathcal{N}(K) \subseteq K$.

Now assume that $\mathcal{N}(K)$ contains K . Then the set $S = K \setminus \mathcal{N}(K)$ is connected by (4) of Theorem 2.5.8. We have $S = K \setminus \mathcal{N}(K) = P_\omega \cap \bigcap_{t \in [0, \omega]} L_K(t)$ as in Equation (2.1). Now see the movement of $L_K(t)$ containing S for $t \in [0, \omega]$ in perspective of S to show that S is a moving sofa ($S \subseteq P_\omega$ and $h_K(\omega) = h_K(\pi/2) = 1$ implies that S moves from H_L to V_L). By (2) of Theorem 2.5.8 we have $\delta K \subseteq S \subseteq K$. So h_S and h_K agree on J_ω , and we have $\mathcal{C}(S) = K$. Now $S = K \setminus \mathcal{N}(K) = \mathcal{I}(S)$ by Theorem 2.4.3 and S is monotone by Theorem 2.4.4. \square

Remark 2.5.2. Not all cap K contains its niche $\mathcal{N}(K)$. For example, take a long cap $K = [0, 100] \times [0, 1]$ with rotation angle $\omega = \pi/2$. Then K is very wide and the inner quadrant $Q_K^-(\pi/4)$ of $L_K(\pi/4)$ is outside K , so that $\mathcal{N}(K) \not\subseteq K$. By Theorem 2.5.9, this K can never be the cap of a monotone sofa with rotation angle $\pi/2$.

From now on, we will *always* understand a monotone sofa S with rotation angle ω by its cap $K := \mathcal{C}(S)$ in \mathcal{K}_ω^c . By Theorem 2.4.3, the monotone sofa $S = K \setminus \mathcal{N}(K)$ can be recovered from its cap K . In other words, the collection of all monotone sofas S with the rotation angle $\omega \in (0, \pi/2]$ embeds into \mathcal{K}_ω^c by taking the cap $S \mapsto \mathcal{C}(S)$. That is, the space of caps \mathcal{K}_ω^c extends the space of all monotone sofas with rotation angle ω .

Define the *sofa area functional* \mathcal{A}_ω on \mathcal{K}_ω^c extending the area $|S|$ of a monotone sofa S .

Definition 2.5.8. For any angle $\omega \in (0, \pi/2]$, define the *sofa area functional* $\mathcal{A}_\omega : \mathcal{K}_\omega^c \rightarrow \mathbb{R}$ on the space of caps \mathcal{K}_ω^c as $\mathcal{A}_\omega(K) = |K| - |\mathcal{N}(K)|$.

Theorem 2.5.10. *For any cap $K := \mathcal{C}(S)$ of a monotone sofa S with rotation angle ω , we have $\mathcal{A}_\omega(K) = |S|$.*

Proof. An immediate consequence of Theorem 2.4.3 and Theorem 2.5.9. \square

We will change the moving sofa problem to the maximization of the sofa area functional \mathcal{A}_ω on cap space \mathcal{K}_ω^c . But recall that as in Remark 2.5.2, not all cap $K \in \mathcal{K}_\omega^c$ is the cap $\mathcal{C}(S)$ of a monotone sofa S . So it is not obvious yet that the maximizer K_ω of \mathcal{A}_ω will correspond to a monotone sofa S_ω of maximum area with a fixed rotation angle ω . This will be established later in Theorem 3.5.6.

Chapter 3

Balanced Maximum Sofas and Caps

This chapter follows the overview in Section 1.4 and shows the existence of a *balanced maximum sofa*, a monotone sofa of the maximum area that can be approximated sufficiently close by balanced polygons.

- Section 3.1 rigorously define the notion of *simple Nef polygons* and prove Theorem 3.1.2 that measures the area difference in balancing moves on a simple Nef polygon.
- Section 3.2 builds the notion of *polygon cap* K with finite angle set Θ and *polygon niche* $\mathcal{N}_\Theta(K)$.
- Section 3.3 extends the space \mathcal{K}_Θ^c of all polygon caps with angle set Θ to a larger space \mathcal{H}_Θ of functions $h : \Theta^\circ \rightarrow \mathbb{R}$.
- Section 3.4 defines the *polyline* \mathbf{p}_K of polygon cap K that represents the bottom sides of polygon sofa $S_\Theta := K \setminus \mathcal{N}_\Theta(K)$. Theorem 3.4.9 shows that for a *maximum polygon cap* K that maximizes $|K| - |\mathcal{N}_\Theta(K)|$, the upper sides of K should balance with the sides of polyline \mathbf{p}_K , following the outline in Section 1.4.2. Theorem 3.4.10 shows that $\mathcal{N}_\Theta(K) \subset K$ for a maximum polygon cap K .
- Section 3.5 defines the *balanced maximum cap* K_ω as the limit of maximum polygon caps $K = K_\Theta$ with angle set Θ and rotation angle ω , as Θ gets denser in $[0, \omega]$. The set $S_\omega := K_\omega \setminus \mathcal{N}(K_\omega)$ is then a *balanced maximum sofa* which attains the maximum area among all monotone sofas of rotation angle $\omega \in (0, \pi/2]$.

3.1 Simple Nef Polygon

A *Nef polygon* is a subset of \mathbb{R}^2 that can be obtained by applying a finite number of boolean operations (e.g. complement $\mathbb{R}^2 \setminus X$ of X or intersection $X_1 \cap X_2$ of X_1 and X_2) to open or closed half-planes [Bie95]. For our purpose, we introduce the notion of *simple Nef polygon*. The polygon cap K and niche $\mathcal{N}_\Theta(K)$ that will be defined in the next Section 3.4 are simple Nef polygons. We will establish Theorem 3.1.2 to balance the sides of K and $\mathcal{N}_\Theta(K)$.

Definition 3.1.1. Let *true* and *false* be the *boolean values* denoting the truth value of a predicate (e.g., $1 + 1 = 2$ is *true*, but $1 + 2 = 4$ is *false*). A (n -ary) *boolean function* \mathcal{E} is a function from $\{\text{true}, \text{false}\}^n$ to $\{\text{true}, \text{false}\}$.

Definition 3.1.2. For two boolean values P and Q , write $P \Rightarrow Q$ if and only if either P is false or both P and Q are true. An n -ary boolean function \mathcal{E} is *monotone* if for any boolean values $P_1, P_2, \dots, P_n, Q_1, Q_2, \dots, Q_n$, with $P_i \Rightarrow Q_i$ for all $1 \leq i \leq n$, we have

$$\mathcal{E}(P_1, \dots, P_n) \Rightarrow \mathcal{E}(Q_1, \dots, Q_n).$$

Proposition 3.1.1. Any n -ary boolean function $\mathcal{E}(P_1, \dots, P_n)$ is monotone if it is obtained from variables P_1, \dots, P_n by applying logical conjunctions \wedge (AND) and disjunctions \vee (OR).

Proof. Check that each P_i is itself monotone as an n -ary boolean function on the variables P_1, \dots, P_n . Then observe that any logical conjunction or disjunction of monotone functions are monotone. \square

Definition 3.1.3. For any n -ary boolean function \mathcal{E} and n closed or open half-planes H_1, \dots, H_n of \mathbb{R}^2 , define the set

$$\mathcal{E}(H_1, \dots, H_n) := \{p \in \mathbb{R}^2 : \mathcal{E}(p \in H_1, \dots, p \in H_n) \text{ is true}\}.$$

Call any such set $X := \mathcal{E}(H_1, \dots, H_n)$ a *Nef polygon*.

Definition 3.1.4. Call X a *simple Nef polygon* with *defining half-planes* H_1, H_2, \dots, H_n if $X = \mathcal{E}(H_1, \dots, H_n)$ for a monotone boolean function \mathcal{E} and the half-planes H_1, \dots, H_n have different boundaries l_1, \dots, l_n .

A line l in \mathbb{R}^2 is a Nef polygon since l is the intersection of two closed half-planes H_l^+ and H_l^- with boundary l . However, l is not simple because the half-planes H_l^+ and H_l^- shares the same boundary. The idea of Definition 3.1.4 is that a point $p \in \mathbb{R}^2$ is more likely to be contained in X if p is contained in more defining half-planes H_1, \dots, H_n . Recall that \mathcal{H}^1 is the Hausdorff measure of dimension one on \mathbb{R}^2 measuring the length of finite segments.

Definition 3.1.5. For real-valued expressions f and g that may depend on other parameters, write $f = O(g)$ if and only if there is an absolute constant $C > 0$ that does *not* depend on any parameters such that $|f| \leq Cg$.

If the inequality $|f| \leq Cg$ holds for a constant $C > 0$ that depends *only* on certain parameters (e.g. a and b), write these as the subscripts of O (e.g. $f = O_{a,b}(g)$).

Theorem 3.1.2. Let $X = \mathcal{E}(H_1, \dots, H_n)$ be a simple Nef polygon with monotone boolean function \mathcal{E} and defining half-planes $H_i = H_-(t_i, h_i)$ or $H_-(t_i, h_i)$ of different boundaries $l_i = l(t_i, h_i)$ for $1 \leq i \leq n$. Fix an arbitrary index $1 \leq i \leq n$. There exists a constant $\epsilon = \epsilon(X, i) > 0$ such that the following holds.

Let δ be any real value with $|\delta| \leq \epsilon$. Define

$$X'_\delta = \mathcal{E}(H_1, \dots, H_{i-1}, H'_\delta, H_{i+1}, \dots, H_n)$$

where $H'_\delta := H_-(t_i, h_i + \delta)$ replaces $H_i = H_-(t_i, h_i)$ or, $H'_\delta := H_-(t_i, h_i + \delta)$ replaces $H_i = H_-(t_i, h_i)$. Then we have

$$|X'_\delta| = |X| + \mathcal{H}^1(\partial X \cap l_i) \cdot \delta + O_{X,i}(\delta^2).$$

Proof. The lines l_j for all index $j \neq i$ divide the plane into open polygons R_1, R_2, \dots, R_N with $N \leq 2^{n-1}$. Fix an arbitrary region R_k with $1 \leq k \leq N$. We will show that there exists some $\epsilon_k = \epsilon_k(X, i, k) > 0$ such that

$$|X'_\delta \cap R_k| = |X \cap R_k| + \mathcal{H}^1(\partial X \cap l_i \cap R_k) \cdot \delta + O(\delta^2) \quad (3.1)$$

for any δ with $|\delta| \leq \epsilon_k$. Once Equation (3.1) is shown, we can take $\epsilon > 0$ to be the minimum of ϵ_k over all k and sum Equation (3.1) over all k to complete the proof. Note that the boundary of any region R_k and the line l_i intersect in a finite number of points, so adding $\mathcal{H}^1(\partial X \cap l_i \cap R_k)$ over all k gives $\mathcal{H}^1(\partial X \cap l_i)$.

Take an arbitrary point $p \in R_k$ in an open region R_k . By the definition of R_k , for any index $j \neq i$, the predicate $p \in H_j$ is a constant true or false no matter which $p \in R_k$ we take. Define the restriction

$$\mathcal{F}(Q) := \mathcal{E}(p \in H_1, \dots, p \in H_{i-1}, Q, p \in H_{i+1}, \dots, p \in H_n)$$

of \mathcal{E} to a single boolean variable Q . Given that $p \in R_k$, the predicate $\mathcal{F}(p \in H_i)$ is equivalent to $p \in X$ and the predicate $\mathcal{F}(p \in H'_{i,\delta})$ is equivalent to $p \in X'_\delta$. Since \mathcal{E} is monotone, \mathcal{F} is also monotone and $\mathcal{F}(Q)$ cannot be the negation of Q . We now have three cases.

Case 1 (resp. Case 2): $\mathcal{F}(Q)$ is the constant false (resp. true). In this case, for any $p \in R_k$ the predicates $\mathcal{F}(p \in H_i)$ and $\mathcal{F}(p \in H'_i)$ are false (resp. true) so R_k is disjoint from (resp. contained in) both X and X'_δ . So R_k is disjoint from ∂X and Equation (3.1) holds.

Case 3: $\mathcal{F}(Q)$ is Q . In this case, for any $p \in R_k$ the predicate $p \in H_i$ (resp. $p \in H'_{i,\delta}$) is equivalent to $p \in X$ (resp. $p \in X'_\delta$). Consequently, we have $X \cap R_k = H_i \cap R_k$ and $X'_\delta \cap R_k = H'_\delta \cap R_k$. In particular, $\partial X \cap R_k = l_i \cap R_k$. Now Equation (3.1) becomes

$$|H'_\delta \cap R_k| = |H_i \cap R_k| + \mathcal{H}^1(l_i \cap R_k) \cdot \delta + O(\delta^2). \quad (3.2)$$

Define $g(x) := |H_-(t, x) \cap R_k|$ so that $|H'_\delta \cap R_k| = g(h_i + \delta)$ and $|H_i \cap R_k| = g(h_i)$. Let $f(x) := \mathcal{H}^1(l(t_i, x) \cap R_k)$ then by Cavalieri's principle we have $g(x) = \int_{-\infty}^x f(u) du$. Note that R_k is a convex polygon with edges different from $l_i = l(t_i, h_i)$. So $f(x)$ is Lipschitz near $x = h_i$. Now by approximating $g(x)$ near $x = h_i$, we get the linear estimate $g(h + \delta) = g(h) + f(h)\delta + O(\delta^2)$ which is exactly Equation (3.2). In all three cases, we establish Equation (3.1) and complete the proof. \square

3.2 Polygon Cap and Niche

Following the outline in Section 1.3, we define a polygon cap K and its polygon niche $\mathcal{N}_\Theta(K)$.

Definition 3.2.1. Define an *angle set* Θ with *rotation angle* $\omega \in (0, \pi/2]$ as the pair (ω, X) of ω and a nonempty finite subset X of $(0, \omega)$.

Remark 3.2.1. The angle set Θ is essentially a finite subset X of $(0, \omega)$ that does not forget the value of ω . With an abuse of notation, we will treat the angle set Θ like the subset $X \subseteq (0, \omega)$ by, for example, saying $t \in \Theta$ instead of saying $t \in X$. Unless specified otherwise, the value ω will always denote the rotation angle of Θ .

Definition 3.2.2. For any angle set Θ with rotation angle ω , define Θ° as the set $\Theta \cup (\Theta + \pi/2) \cup \{\omega, \pi/2\}$.

Definition 3.2.3. Define the space \mathcal{K}_Θ^c of *polygon caps with angle set* Θ as the set of caps $K \in \mathcal{K}_\omega^c$ which is the intersection of closed half-planes with normal angles in $\Theta^\circ \cup \{\omega + \pi, 3\pi/2\}$.

Any cap $K \in \mathcal{K}_\omega^c$ can be approximated by the polygon cap $\mathcal{C}_\Theta(K) \in \mathcal{K}_\Theta^c$ with angle set Θ circumscribing K with the edges of normal angles in $\Theta^\circ \cup \{\omega + \pi, 3\pi/2\}$. The denser Θ is in $(0, \omega)$, the better the approximation $\mathcal{C}_\Theta(K)$ of K .

Definition 3.2.4. For any $K \in \mathcal{K}_\omega^c$ and angle set Θ of rotation angle ω , define

$$\mathcal{C}_\Theta(K) = P_\omega \cap \bigcap_{t \in \Theta} Q_K^+(t).$$

Proposition 3.2.1. For any cap $K \in \mathcal{K}_\omega^c$, the set $\mathcal{C}_\Theta(K) \in \mathcal{K}_\Theta^c$ contains K as a subset. With this, call $\mathcal{C}_\Theta(K)$ the polygon cap with angle set Θ approximating K . The map $\mathcal{C}_\Theta : \mathcal{K}_\omega^c \rightarrow \mathcal{K}_\Theta^c$ is a surjective map fixing the elements of $\mathcal{K}_\Theta^c \subset \mathcal{K}_\omega^c$.

Proof. That $K \subseteq \mathcal{C}_\Theta(K)$ comes from $K \subseteq P_\omega$ and $K \subseteq Q_K^+(t) = H_K(t) \cap H_K(t + \pi/2)$. We have $\mathcal{C}_\Theta(K) \in \mathcal{K}_\omega^c$ as $K \subseteq \mathcal{C}_\Theta(K) \subseteq P_\omega$. We have $\mathcal{C}_\Theta(K) \in \mathcal{K}_\Theta^c$ as the formula in Definition 3.2.4 is the intersection of closed half-planes with normal angles in $\Theta^\circ \cup \{\omega + \pi, 3\pi/2\}$.

Now it remains to show that for any $K \in \mathcal{K}_\Theta^c$, we have $\mathcal{C}_\Theta(K) = K$. Since $K \in \mathcal{K}_\Theta^c$ is an intersection of closed half-planes with normal angles in $\Pi := \Theta^\circ \cup \{\omega + \pi, 3\pi/2\}$, we have $K = \bigcap_{t \in \Pi} H_K(t)$. As $K \in \mathcal{K}_\Theta^c \subset \mathcal{K}_\omega^c$, we have $h_K(\omega) = h_K(\pi/2) = 1$ and $h_K(\omega + \pi) = h_K(3\pi/2) = 0$ so

$$K = \bigcap_{t \in \Pi} H_K(t) = P_\omega \cap \bigcap_{t \in \Theta} Q_K^+(t) = \mathcal{C}_\Theta(K)$$

as desired. \square

Definition 3.2.5. For every cap $K \in \mathcal{K}_\omega^c$ and finite nonempty $\Theta \subset (0, \omega)$, define its *polygon niche*

$$\mathcal{N}_\Theta(K) = P_\omega \cap \bigcup_{t \in \Theta} Q_K^-(t)$$

with angle set Θ .

Proposition 3.2.2. For any $K \in \mathcal{K}_\omega^c$, we have $\mathcal{N}_\Theta(K) = \mathcal{N}_\Theta(\mathcal{C}_\Theta(K)) \subseteq \mathcal{N}(K)$.

Proof. Let $K' := \mathcal{C}_\Theta(K)$. The support functions h_X of $X = K, K'$ agree on the set $\Theta \cup (\Theta + \pi/2)$ by Definition 3.2.4. Observe that the Definition 3.2.5 of $\mathcal{N}_\Theta(X)$ only depends on ω and the values of h_X on $\Theta \cup (\Theta + \pi/2)$. So we have $\mathcal{N}_\Theta(K) = \mathcal{N}_\Theta(K')$. That $\mathcal{N}_\Theta(K) \subseteq \mathcal{N}(K)$ follows from their respective definitions. \square

Definition 3.2.6. Let Θ be an angle set with rotation angle ω . Define the *polygon sofa area functional* $\mathcal{A}_\Theta : \mathcal{K}_\omega^c \rightarrow \mathbb{R}$ as $\mathcal{A}_\Theta(K) := |\mathcal{C}_\Theta(K)| - |\mathcal{N}_\Theta(K)|$.

Theorem 3.2.3. For any polygon cap $K \in \mathcal{K}_\Theta^c$ we have $\mathcal{A}_\Theta(K) = |K| - |\mathcal{N}_\Theta(K)|$. For every cap $K \in \mathcal{K}_\omega^c$, we have $\mathcal{A}_\omega(K) \leq \mathcal{A}_\Theta(K)$.

Proof. Consequence of Proposition 3.2.1 and Proposition 3.2.2. \square

Remark 3.2.2. The upper bound $\alpha_{\max} \leq 2.37$ of [KR18] was essentially established by computing the upper bound \mathcal{A}_Θ of $\mathcal{A}_{\pi/2}$ for a specific set Θ of five angles.

3.3 Extensions of Polygon Cap Space

We extend the space \mathcal{K}_Θ^c of polygon caps to the space \mathcal{K}_Θ^t of all *translates* of polygon caps.

Definition 3.3.1. Define the space of *polygon cap translates* \mathcal{K}_Θ^t as the collection of all translations K' of every cap $K \in \mathcal{K}_\Theta^c$ with angle set Θ .

Proposition 3.3.1. *A convex polygon K' is in \mathcal{K}_Θ^t if and only if the followings are all true.*

1. *The widths $h_{K'}(\omega) + h_{K'}(\omega + \pi)$ and $h_{K'}(\pi/2) + h_{K'}(3\pi/2)$ of K' along the angles ω and $\pi/2$ are exactly one.*
2. *K' is a convex polygon with normal angles in the set Θ° .*

Proof. Any translate K' of an arbitrary $K \in \mathcal{K}_\Theta^c$ satisfies (1) and (2) immediately. On the other hand, for any K' satisfying conditions (1) and (2), find a translation K of K' so that $h_K(\omega) = h_K(\pi/2) = 1$ and $K \in \mathcal{K}_\Theta^c$. Since the width of K measured along the angles ω and $\pi/2$ are one, we get $h_K(\omega + \pi) = h_K(3\pi/2) = 0$. \square

Now extend the set of all support functions of $K \in \mathcal{K}_\Theta^t$ to the space \mathcal{H}_Θ of all functions $h : \Theta^\circ \rightarrow \mathbb{R}$.

Definition 3.3.2. Define \mathcal{H}_Θ as the space of all functions $h : \Theta^\circ \rightarrow \mathbb{R}$.

Proposition 3.3.2. *The map $\mathcal{K}_\Theta^t \rightarrow \mathcal{H}_\Theta$ mapping each $K' \in \mathcal{K}_\Theta^t$ to its support function $h_{K'}$ restricted to the domain Θ° is an injection.*

Proof. By (2) of Proposition 3.3.1, the values of $h_{K'}$ on Θ° determine

$$K' = \bigcap_{t \in \Theta^\circ \cup \{\omega + \pi, 3\pi/2\}} H_{K'}(t) \in \mathcal{K}_\Theta^t$$

uniquely. \square

We now have a series of extensions $\mathcal{K}_\Theta^c \subseteq \mathcal{K}_\Theta^t \hookrightarrow \mathcal{H}_\Theta$. We will extend the notion of cap, niche, and the polygon sofa area functional \mathcal{A}_Θ from \mathcal{K}_Θ^c to \mathcal{H}_Θ .

Definition 3.3.3. For any $h \in \mathcal{H}_\Theta$, define its *parallelogram*

$$P_h := \bigcap_{t \in \{\omega, \pi/2\}} H_-(t, h(t)) \cap H_+(t, h(t) - 1)$$

and *cap*

$$\mathcal{C}_\Theta(h) := P_h \cap \bigcap_{t \in \Theta \cup (\Theta + \pi/2)} H_-(t, h(t))$$

and *fan*

$$F_h := \bigcap_{t \in \{\omega, \pi/2\}} H_+(t, h(t) - 1)$$

and *niche*

$$\mathcal{N}_\Theta(h) := F_h \cap \bigcup_{t \in \Theta} (H_-^\circ(t, h(t) - 1) \cap H_-^\circ(t + \pi/2, h(t + \pi/2) - 1))$$

and the *polygon sofa area functional*

$$\mathcal{A}_\Theta(h) := |\mathcal{C}(h)| - |\mathcal{N}_\Theta(h)|.$$

The polygons appearing in Definition 3.3.3 are simple Nef polygons (Definition 3.1.4).

Proposition 3.3.3. *Take arbitrary $h \in \mathcal{H}_\Theta$.*

1. *The sets $\mathcal{C}_\Theta(h)$ is a simple Nef polygon with the defining half-planes $H_-(t, h(t))$ for each $t \in \Theta^\circ$, and $H_+(t, h(t) - 1)$ for each $t \in \{\omega, \pi/2\}$.*
2. *The set $\mathcal{N}_\Theta(h)$ is a simple Nef polygon with the defining half-planes $H_-^\circ(t, h(t) - 1)$ for each $t \in \Theta \cup (\Theta + \pi/2)$, and $H_+(t, h(t) - 1)$ for each $t \in \{\omega, \pi/2\}$.*

Proof. Check that Definition 3.1.4 holds for $\mathcal{C}_\Theta(h)$ and $\mathcal{N}_\Theta(h)$. The boundaries of the defining half-planes differ as they differ in its normal angle or distance from the origin. The formulas of $\mathcal{C}_\Theta(h)$ and $\mathcal{N}_\Theta(h)$ in Definition 3.3.3 are defined from the mentioned half-planes by applying only union and intersection, so by Proposition 3.1.1 the defining boolean function is monotone. \square

We show that the extended notion of cap, niche, and polygon sofa area functional on \mathcal{H}_Θ in Definition 3.3.3 are compatible with that of \mathcal{K}_Θ^c .

Proposition 3.3.4. *For any polygon cap translate $K' \in \mathcal{K}_\Theta^t$ with $h := h_K \in \mathcal{H}_\Theta$ we have $\mathcal{C}_\Theta(h) = K'$.*

Proof. By (2) of Proposition 3.3.1, $K' = \bigcap_{t \in \Theta^\circ \cup \{\omega + \pi, 3\pi/2\}} H_{K'}(t)$. By (1) of Proposition 3.3.1, the intersection matches the Definition 3.3.3 of $\mathcal{C}_\Theta(h_K)$. \square

Proposition 3.3.5. *For any polygon cap $K \in \mathcal{K}_\Theta^c$ with $h := h_K \in \mathcal{H}_\Theta$ we have $\mathcal{N}_\Theta(h) = \mathcal{N}_\Theta(K)$. Consequently, we have $\mathcal{A}_\Theta(h) = \mathcal{A}_\Theta(K)$.*

Proof. That $\mathcal{N}_\Theta(h) = \mathcal{N}_\Theta(K)$ comes from the last equation of Proposition 2.2.2. Then $\mathcal{A}_\Theta(h) = \mathcal{A}_\Theta(K)$ comes from Proposition 3.3.4. \square

Now inherit Definition 3.3.3 from \mathcal{H}_Θ to \mathcal{K}_Θ^t under the extension $\mathcal{K}_\Theta^c \subseteq \mathcal{K}_\Theta^t \hookrightarrow \mathcal{H}_\Theta$ (Definition 3.3.4) and observe that they act equivariantly under translation (Theorem 3.3.6).

Definition 3.3.4. For any cap translate $K' \in \mathcal{K}_\Theta^t$ with support function $h := h_{K'} \in \mathcal{H}_\Theta$, define $\mathcal{N}_\Theta(K') := \mathcal{N}_\Theta(h)$ and $\mathcal{A}_\Theta(K') := \mathcal{A}_\Theta(h')$.

Theorem 3.3.6. *For any translation $K' := K + \mathbf{v} \in \mathcal{K}'_\Theta$ of a polygon cap $K \in \mathcal{K}_\Theta^c$ by $\mathbf{v} \in \mathbb{R}^2$, we have $\mathcal{N}_\Theta(K') = \mathcal{N}_\Theta(K) + \mathbf{v}$ and $\mathcal{A}_\Theta(K') = \mathcal{A}_\Theta(K)$. So with $\mathbf{v} := (0, 0)$, Definition 3.3.4 is a proper extension of \mathcal{N}_Θ and \mathcal{A}_Θ from \mathcal{K}_Θ^c to \mathcal{K}_Θ^t .*

Proof. By Proposition 3.3.3 and Definition 3.3.4, the polygon $\mathcal{N}_\Theta(K')$ is a simple Nef polygon with defining half-planes $H_{K'}(t)$ and $H_{K'}(t) - u_t$ (or their complement and/or interior) for $t \in \Theta^\circ$. For any convex body K and fixed $t \in S^1$, the supporting half-plane $H_K(t)$ is equivariant under the translation of K so that $H_{K+\mathbf{v}}(t) = H_K(t) + \mathbf{v}$. So we have $\mathcal{N}_\Theta(K') = \mathcal{N}_\Theta(K) + \mathbf{v}$. We then have

$$\begin{aligned} \mathcal{A}_\Theta(K') &= |\mathcal{C}_\Theta(h_{K'})| - |\mathcal{N}_\Theta(h_{K'})| = |K'| - |\mathcal{N}_\Theta(K')| \\ &= |K| - |\mathcal{N}_\Theta(K)| = \mathcal{A}_\Theta(K) \end{aligned}$$

by Proposition 3.3.4 and Proposition 3.3.5. \square

Proposition 3.3.7. *Assume that $h^+ \in \mathcal{H}_\Theta$ is taken so that $K^+ := \mathcal{C}_\Theta(h^+)$ is a polygon cap translate in \mathcal{K}_Θ^t .¹ Then $\mathcal{A}_\Theta(h^+) \leq \mathcal{A}_\Theta(K^+)$.*

Proof. By the definition of $K^+ := \mathcal{C}_\Theta(h^+)$, we always have $h_{K^+}(t) \leq h^+(t)$ on $t \in \Theta^\circ$ and $K^+ \subseteq P_{h^+}$. Since K^+ is a polygon cap translate, it has width 1 in the directions of u_ω and $u_{\pi/2}$, so is circumscribed in P_{h^+} . So we have equality $h_{K^+}(t) = h^+(t)$ for $t = \omega, \pi/2$, and $F_{h_{K^+}} = F_{h^+}$ in particular. From this and $h_{K^+}(t) \leq h^+(t)$ on $t \in \Theta^\circ$, we have $\mathcal{N}_\Theta(h_{K^+}) \subseteq \mathcal{N}_\Theta(h^+)$. By Proposition 3.3.4 on K^+ we have $\mathcal{C}_\Theta(h_{K^+}) = K^+ = \mathcal{C}_\Theta(h^+)$. Now

$$\begin{aligned} \mathcal{A}_\Theta(h^+) &= |\mathcal{C}_\Theta(h^+)| - |\mathcal{N}_\Theta(h^+)| \\ &\leq |\mathcal{C}_\Theta(h_{K^+})| - |\mathcal{N}_\Theta(h_{K^+})| = \mathcal{A}_\Theta(h_{K^+}) = \mathcal{A}_\Theta(K^+) \end{aligned}$$

where we use Definition 3.3.4 in the last equality. \square

3.4 Maximum Polygon Cap

We now define the polygon cap K_Θ attaining the maximum value of \mathcal{A}_Θ .

Definition 3.4.1. Call a polygon cap $K_\Theta \in \mathcal{K}_\Theta^c$ a *maximum polygon cap* with angle set Θ if $o_\omega \in K_\Theta$ and $K := K_\Theta$ attains the maximum value of $\mathcal{A}_\Theta : \mathcal{K}_\Theta^c \rightarrow \mathbb{R}$.

Remark 3.4.1. For any rotation angle $\omega < \pi/2$, any cap $K \in K_\Theta$ must contain the point o_ω . So the condition $o_\omega \in K_\Theta$ in Definition 3.4.1 is only relevant when $\omega = \pi/2$, and its purpose is to prevent the cap from sliding horizontally far away.

Lemma 3.4.1. *The mirror reflection of any maximum polygon cap $K_\Theta \in \mathcal{K}_\Theta^c$ is a maximum polygon cap with the angle set $\omega - \Theta$.*

Proof. Observe that \mathcal{A}_Θ and o_ω are preserved under the mirror reflection M_ω . \square

Lemma 3.4.2. *Let $\omega \in (0, \pi/2]$ and $t \in (0, \omega)$ be arbitrary. There exists a constant $c_{\omega, t} > 0$ such that the following holds. Let Θ be any angle set of rotation angle ω containing t . Assume that $K \in \mathcal{K}_\Theta^c$ satisfies $\mathcal{A}_\Theta(K) > 0$. Then the width of K along the direction u_0 is at most $c_{\omega, t}$.*

Proof. If $\omega < \pi/2$, then as $K \subseteq P_\omega$ it suffices to take $c_{\omega, t}$ as the width of P_ω along the direction u_0 . Now assume $\omega = \pi/2$ and fix t . Take any $K \in \mathcal{K}_\Theta^c$ with the angle set Θ containing t , and let d be the width of K along u_0 . Then $|K| \leq d$. The wedge $T_K(t)$ is a right triangle of side $\geq d - \sec t - \csc t$ and the acute angle t . So $|\mathcal{N}_\Theta(K)| \geq |T_K(t)| \geq Q_t(d)$, where $Q_t(d)$ is a quadratic polynomial of d completely determined by t with positive leading coefficient. Now there exists a constant $c_{\pi/2, t} > 0$ so that for any $K \in \mathcal{K}_\Theta^c$ with width $d > c_{\pi/2, t}$ we have $\mathcal{A}_\Theta(K) = |K| - |\mathcal{N}_\Theta(K)| \leq d - Q_t(d) \leq 0$. Take the contraposition to finish the proof. \square

Theorem 3.4.3. *For any angle set Θ , a maximum polygon cap K_Θ exists.*

Proof. We first show that $\mathcal{A}_\Theta(K) = |K| - |\mathcal{N}_\Theta(K)|$ on $K \in \mathcal{K}_\Theta^c$ is continuous with respect to the Hausdorff distance d_H on $K \in \mathcal{K}_\Theta^c$. By Theorem 1.8.20, page 68 of [Sch13], $|K|$ is continuous with respect to d_H . Fix an $K \in \mathcal{K}_\Theta^c$ and take any $K' \in \mathcal{K}_\Theta^c$ sufficiently close to K

¹Note that we do not require $h^+ = h_{K^+}$ here. It is possible for the resulting cap K^+ to have support function strictly less than h^+ after intersection, as the lines obtained from h^+ are not guaranteed to be in convex position.

in d_H . By Proposition 2.5.3, the absolute value of $|\mathcal{N}_\Theta(K')| - |\mathcal{N}_\Theta(K)|$ is at most the sum of the areas of the symmetric difference $\Delta(t)$ between wedges $T_K(t)$ and $T_{K'}(t)$ over all angles $t \in \Theta$. As $K' \rightarrow K$ in d_H , we have $h_{K'}(t) \rightarrow h_K(t)$ and $h_{K'}(t + \pi/2) \rightarrow h_K(t)$ so $|\Delta(t)| \rightarrow 0$. This shows that $|\mathcal{N}_\Theta(K')| \rightarrow |\mathcal{N}_\Theta(K)|$ as $K' \rightarrow K$ in d_H . So $\mathcal{A}_\Theta(K)$ is continuous in K with respect to d_H .

Let \mathcal{B}_Θ be the collection of all $K \in \mathcal{K}_\Theta^c$ such that $\mathcal{A}_\Theta(K) \geq 0$ and $o_\omega \in K$. Let $K_1 \in \mathcal{K}_\Theta^c$ be the polygon cap with the support function $h_{K_1}(t) = 1$ for every $t \in \Theta^\circ$ (this K_1 is equal to $\mathcal{C}_\Theta(K)$ where K is the semicircle of radius 1 and angle $\omega + \pi/2$ centered at O). Then $\mathcal{N}_\Theta(K_1)$ is empty and $o_\omega \in K_1$ so we have $K_1 \in \mathcal{B}_\Theta$ and \mathcal{B}_Θ is nonempty. By Lemma 3.4.2, the width of any member of \mathcal{B}_Θ is bounded by the constant $c_{\omega,t}$. So by the Blaschke selection theorem, the domain \mathcal{B}_Θ is compact in \mathcal{K}_Θ^c . So a maximizer K_Θ of $\mathcal{A}_\Theta(K_\Theta)$ on the compact and nonempty domain \mathcal{B}_Θ exists.

We finally show that $K := K_\Theta$ maximizes $\mathcal{A}_\Theta(K)$ over the domain $K \in \mathcal{K}_\Theta^c$ larger than \mathcal{B}_Θ . Take any other $K' \in \mathcal{K}_\Theta^c$. Our goal is to show that $\mathcal{A}_\Theta(K') \leq \mathcal{A}_\Theta(K_\Theta)$. If $\mathcal{A}_\Theta(K') < 0$, then we have $\mathcal{A}_\Theta(K') < 0 \leq \mathcal{A}_\Theta(K_\Theta)$ since $K_\Theta \in \mathcal{B}_\Theta$ so the proof is done. So assume $\mathcal{A}_\Theta(K') \geq 0$. If $\omega < \pi/2$, then we have $o_\omega \in K'$ so $K' \in \mathcal{B}_\Theta$ and again the proof is done. So assume also $\omega = \pi/2$. Now any horizontal translation of K' is also a cap $K'' \in \mathcal{K}_\Theta^c$ with the same $\mathcal{A}_\Theta(K'') = \mathcal{A}_\Theta(K')$. Find one translate K'' that contains the point $o_\omega = (0, 1)$, then $K'' \in \mathcal{B}_\Theta$ and we have $\mathcal{A}_\Theta(K') = \mathcal{A}_\Theta(K'') \leq \mathcal{A}_\Theta(K_\Theta)$ also completing the proof. \square

Definition 3.4.2. For any $n \geq 1$ and a finite sequence $p_1, p_2, \dots, p_n \in \mathbb{R}^2$ of points on a plane with strictly increasing x -coordinates, call the union of all closed segments connecting adjacent points p_i to p_{i+1} ($1 \leq i < n$) an x -monotone polyline.




Figure 3.1: A polygon cap $K \in \mathcal{K}_\Theta^c$ and polygon niche $\mathcal{N}_\Theta(K)$ with balanced side lengths. Two black points $A_K^-(0)$ and $C_K^+(\omega)$ separate the upper boundary δK of K and the polyline \mathbf{p}_K . Let $\theta_1, \dots, \theta_8$ be the elements of Θ in increasing order. Each side length $\sigma_i := \sigma(\theta_i)$ of δK then balance exactly with a corresponding side length $\tau_i := \tau(\theta_i)$ of \mathbf{p}_K .

Theorem 3.4.4. For any polygon cap $K \in \mathcal{K}_\Theta^c$, the boundary of the closed set $F_\omega \setminus \mathcal{N}_\Theta(K)$ is a disjoint union of the following subsets, from left to right in \mathbb{R}^2 .

1. The open half-line \vec{l}_K from $C_K^+(\omega)$ extending in the direction v_ω but not containing $C_K^+(\omega)$.
2. An x -monotone polyline \mathbf{p}_K from left to right connecting $C_K^+(\omega)$ to $A_K^-(0)$, with each segment of normal angle $t \in \Theta^\circ$.
3. The open half-line \vec{r}_K from $A_K^-(0)$ extending in the direction u_0 but not containing $A_K^-(0)$.

Proof. Let $X := \bigcup_{t \in \Theta} Q_K^-(t)$. Observe that X is a Nef polygon closed in the direction of $-v_0$ (Definition 2.3.8) with edges of normal angles in $\Theta \cup (\Theta + \pi/2)$. So since $F_\omega \setminus \mathcal{N}_\Theta(K) = F_\omega \setminus X$, it suffices to show that \vec{l}_K and \vec{r}_K are disjoint from X . Take any $t \in \Theta$. Since $w_K(t) > 0$ by Theorem 2.5.5, the point $A_K^-(0)$ and thus the half-line \mathbf{r}_K is on the right side of the line $b_K(t)$. So \mathbf{r}_K is disjoint from $Q_K^-(t)$. Likewise, from $z_K(t) > 0$ of Theorem 2.5.5, $Q_K(t)$ is disjoint from $Q_K^-(t)$ as well. \square

Definition 3.4.3. Define the polyline \mathbf{p}_K in Theorem 3.4.4 from $C_K^+(\omega)$ to $A_K^-(0)$ as the *polyline of cap* $K \in \mathcal{K}_\Theta^c$.

Definition 3.4.4. For any cap $K \in \mathcal{K}_\Theta^c$ and angle $t \in \Theta^\diamond$, define $\tau_K(t)$ as the sum of the lengths of all edges in the polyline \mathbf{p}_K with normal angle t .

Lemma 3.4.5. Let $K \in \mathcal{K}_\Theta^c$ be arbitrary.

1. For any $t \in \Theta$, we have

$$\mathcal{H}^1(\partial\mathcal{N}_\Theta(K) \cap b_K(t)) = \mathcal{H}^1(\partial\mathcal{N}_\Theta(K) \cap \vec{b}_K(t)) = \tau_K(t)$$

and

$$\mathcal{H}^1(\partial\mathcal{N}_\Theta(K) \cap d_K(t)) = \mathcal{H}^1(\partial\mathcal{N}_\Theta(K) \cap \vec{d}_K(t)) = \tau_K(t + \pi/2).$$

2. For any $t \in \{\omega, \pi/2\}$, we have

$$\mathcal{H}^1(\partial\mathcal{N}_\Theta(K) \cap l(t, 0)) = \mathcal{H}^1(\mathcal{N}_\Theta(K) \cap l(t, 0)) = \sigma_K(t + \pi) - \tau_K(t).$$

Proof. Define $X := \bigcup_{t \in \Theta} Q_K^-(t)$, a Nef polygon with edges of normal angles in $\Theta \cup (\Theta + \pi/2)$, and recall that $\mathcal{N}_\Theta(K) = F_\omega \cap X$. Define the followings.

- The intersection $M := X \cap \partial F_\omega$ which is open in the subspace topology of ∂F_ω .
- The boundary $N \subseteq F_\omega^\circ$ of $\mathcal{N}_\Theta(K)$ in the subspace topology of F_ω° .
- The intersection $P := \partial X \cap \partial F_\omega$ which is a finite set of points in ∂F_ω .

We check that the boundary $\partial\mathcal{N}_\Theta(K)$ is a disjoint union of M, N, P . Since F_ω° is open, we have $N = \partial\mathcal{N}_\Theta(K) \cap F_\omega^\circ$. Since $M, P \subseteq \partial F_\omega$, and they are disjoint because X is open, it suffices to show that the set $X' := \partial\mathcal{N}_\Theta(K) \setminus N = \partial\mathcal{N}_\Theta(K) \cap \partial F_\omega$ is a union of M and P . The union of M and P is $\overline{X} \cap \partial F_\omega$ so it contains X' , completing the check for $\partial\mathcal{N}_\Theta(K)$.

Let $Y := \partial(F_\omega \setminus \mathcal{N}_\Theta(K)) = \partial(F_\omega \setminus X)$. We check that $Y = N \cup (\partial F_\omega \setminus M)$. We have $Y \cap F_\omega^\circ = N$ as N is also the boundary of $F_\omega \setminus \mathcal{N}_\Theta(K)$ in the subspace topology of F_ω° . It remains to show $Y \cap \partial F_\omega = \partial F_\omega \setminus M$. This is equivalent to showing that ∂F_ω is a disjoint union of $M = X \cap \partial F_\omega$ and $\partial(F_\omega \setminus X) \cap \partial F_\omega$. They are disjoint because X is open. For any $p \in \partial F_\omega \setminus M = \partial F_\omega \setminus X$, because $F_\omega \setminus X$ is closed in the direction v_0 , there is a point $p + \epsilon v_0 \in F_\omega \setminus X$ sufficiently close to p , so $p \in \partial(F_\omega \setminus X)$. This completes the check for Y .

Define $D := K \cap \partial F_\omega$. Then $D = e_K(\omega + \pi) \cup e_K(3\pi/2)$ since K is a cap. By Theorem 3.4.4, the set \mathbf{p}_K is equal to $Y \setminus \vec{l}_K \setminus \vec{r}_K$. Now, since $Y = N \cup (\partial F_\omega \setminus M)$ by the check above, and $D = \partial F_\omega \setminus \mathbf{l}_K \setminus \mathbf{r}_K$, we have $\mathbf{p}_K = N \cup (D \setminus M)$. The portion N of \mathbf{p}_K is contributed by the lines $b_K(t)$ and $d_K(t)$ for angles $t \in \Theta$. The portion $D \setminus M$ of \mathbf{p}_K is contributed by the bottom sides ∂F_ω of fan F_ω .

We prove (1). Take any $t \in \Theta$.

$$\mathcal{H}^1(\partial\mathcal{N}_\Theta(K) \cap b_K(t)) = \mathcal{H}^1(N \cap b_K(t)) = \mathcal{H}^1(\mathbf{p}_K \cap b_K(t)) = \tau_K(t)$$

because $M, P \subseteq \partial\mathcal{N}_\Theta(K)$ are on ∂F_ω and so only intersect $b_K(t)$ at a finite number of points. The value is also equal to $\mathcal{H}^1(\partial\mathcal{N}_\Theta(K) \cap \vec{b}_K(t))$ because the sides of $\mathcal{N}_\Theta(K)$ with normal angle t is only contributed by the half-line $\vec{b}_K(t)$. Use mirror symmetry to show the corresponding equality for $\tau_K(t + \pi/2)$.

We prove (2). Take any $t \in \{\omega, \pi/2\}$.

$$\mathcal{H}^1(\partial\mathcal{N}_\Theta(K) \cap l(t, 0)) = \mathcal{H}^1(M \cap l(t, 0)) = \mathcal{H}^1((D \setminus \mathbf{p}_K) \cap l(t, 0)) = \sigma_K(t + \pi) - \tau_K(t)$$

because $N \subseteq \partial\mathcal{N}_\Theta(K)$ is disjoint from ∂F_ω and $P \subseteq \partial\mathcal{N}_\Theta(K)$ is a finite number of points. \square

Now we define the notion of balancedness on $K \in \mathcal{K}_\Theta^c$.

Definition 3.4.5. (See Figure 3.1) Say that a polygonal cap $K \in \mathcal{K}_\Theta^c$ is *balanced* if and only if for any $t \in \Theta^\circ$, we have $\sigma_K(t) = \tau_K(t)$.

We now show that any maximum polygon cap $K_\Theta \in \mathcal{K}_\Theta^c$ is balanced. The following lemma chooses the right angle $t \in \Theta^\circ$ to balance $K \in \mathcal{K}_\Theta^c$.

Lemma 3.4.6. *Assume that a polygon cap $K \in \mathcal{K}_\Theta^c$ is not balanced. Then there exists an angle $t \in \Theta^\circ$ such that $\sigma_K(t) > \tau_K(t)$.*

Proof. By following the polyline \mathbf{p}_K from right to left, we have the identity

$$C_K^+(\omega) - A_K^-(0) = \sum_{t \in \Theta^\circ} \tau_K(t) v_t.$$

Also, by following the upper boundary δK of K from right to left, we also have the identity

$$C_K^+(\omega) - A_K^-(0) = \sum_{t \in \Theta^\circ} \sigma_K(t) v_t.$$

So we have $\sum_{t \in \Theta^\circ} (\tau_K(t) - \sigma_K(t))(v_t \cdot u_0) = 0$ where $v_t \cdot u_0 < 0$ for all $t \in \Theta^\circ$. If $\sigma_K(t) \leq \tau_K(t)$ for all $t \in \Theta^\circ$, then the equality $\sigma_K(t) = \tau_K(t)$ should hold and K should be balanced. Taking the contraposition concludes the proof. \square

Execute the balancing step by pusing the edge $e_K(t)$ in the positive direction of u_t .

Lemma 3.4.7. *Let $h \in \mathcal{H}_\Theta$ be the support function of some $K \in \mathcal{K}_\Theta^c$. Let $t \in \Theta^\circ$ be arbitrary. Take sufficiently small $\epsilon > 0$ relative to K and t . Define $h^+ \in \mathcal{H}_\Theta$ as $h^+(t) := h(t) + \epsilon$ and $h^+(s) := h(s)$ for all $s \neq t$. Then we have*

$$\mathcal{A}_\Theta(h^+) = \mathcal{A}_\Theta(h) + (\sigma_K(t) - \tau_K(t))\epsilon + O(\epsilon^2).$$

Proof. First consider the case $t \notin \{\omega, \pi/2\}$. We have

$$|\mathcal{C}_\Theta(h^+)| = |\mathcal{C}_\Theta(h)| + \sigma_K(t)\epsilon + O(\epsilon^2)$$

by applying Theorem 3.1.2 to the simple Nef polygon $\mathcal{C}_\Theta(h)$ ((1) of Proposition 3.3.3). We also have

$$|\mathcal{N}_\Theta(h^+)| = |\mathcal{N}_\Theta(h)| + \tau_K(t)\epsilon + O(\epsilon^2)$$

by applying Theorem 3.1.2 to the simple Nef polygon $\mathcal{N}_\Theta(h) = \mathcal{N}_\Theta(K)$ ((2) of Proposition 3.3.3) and using (1) of Lemma 3.4.5. Subtract the two equations above to conclude the proof.

Now consider the case $t \in \{\omega, \pi/2\}$. Recall that by Proposition 3.3.3, the simple Nef polygon $\mathcal{C}_\Theta(h)$ have defining half-planes including $H_-(t, h(t))$ and $H_+(t, h(t) - 1)$. We will apply Theorem 3.1.2 twice to $\mathcal{C}_\Theta(h)$ to get the following.

$$|\mathcal{C}_\Theta(h^+)| = |\mathcal{C}_\Theta(h)| + (\sigma_K(t) - \sigma_K(t + \pi))\epsilon + O(\epsilon^2)$$

Here, we first push $H_-(t, h(t))$ to $H_-(t, h^+(t))$ by ϵ while keeping $H_+(t, h(t) - 1)$ intact. Then, we push $H_+(t, h(t) - 1)$ towards $H_+(t, h^+(t) - 1)$ by ϵ while keeping $H_-(t, h^+(t))$ intact. Since the two half-planes have parallel boundaries of distance 1, the two applications of Theorem 3.1.2 do not interfere with each other. We also have

$$|\mathcal{N}_\Theta(h^+)| = |\mathcal{N}_\Theta(h)| + (\sigma_K(t + \pi) - \tau_K(t))\epsilon + O(\epsilon^2)$$

by applying Theorem 3.1.2 to $\mathcal{N}_\Theta(h)$ and using (2) of Lemma 3.4.5. Subtract the two equations above to conclude the proof. \square

Choosing the right angle $t \in \Theta^\circ$ in Lemma 3.4.6 guarantees that the new function $h^+ \in \mathcal{H}_\Theta$ with $h^+(t) := h(t) + \epsilon$ corresponds to a cap translate K^+ .

Lemma 3.4.8. *Let $K \in \mathcal{K}_\Theta^c$ be arbitrary with the support function $h := h_K \in \mathcal{H}_\Theta$, so that $K = \mathcal{C}_\Theta(h)$ by Proposition 3.3.4. Let $t \in \Theta^\circ$ be arbitrary such that $\sigma_K(t) > 0$.*

Take any $\epsilon > 0$ sufficiently small relative to K and t , and define $h^+ \in \mathcal{H}_\Theta$ as $h^+(t) := h(t) + \epsilon$ and $h^+(s) := h(s)$ on $s \neq t$. Then the intersection $K^+ := \mathcal{C}_\Theta(h^+)$ is a polygon cap translate in \mathcal{K}_Θ^t .

Proof. We first prove the case $t \notin \{\omega, \pi/2\}$. We have $K = \mathcal{C}_\Theta(h) \subseteq \mathcal{C}_\Theta(h^+) = K^+$ by the definition of \mathcal{C}_Θ and K^+ . So $K \subseteq K^+ \subseteq P_\omega$ and $K^+ \in \mathcal{K}_\Theta^c$ as we want.

Now assume $t \in \{\omega, \pi/2\}$. We first show that K^+ is a polygon cap translate. By Proposition 3.3.1 it suffices to show that the width of K^+ is one in the angles ω and $\pi/2$.

Since $\sigma_K(t) > 0$, we have $h_{K^+}(t) = h_K(t) + \epsilon$ for sufficiently small $\epsilon > 0$.² As $\epsilon > 0$, we have $h_{K^+}(t + \pi) = h_K(t + \pi) - \epsilon$ as well. So the width of K^+ along u_t is one. If $\omega = \pi/2$ then we are done. If $\omega < \pi/2$, let t' be the other value than t in $\{\omega, \pi/2\}$. We have $h_{K^+}(t') = h_K(t')$ as $l_K(t)$ moves upwards. Also, as $\sigma_K(t' + \pi) \geq o_\omega \cdot u_0 > 0$, we have $h_{K^+}(t' + \pi) = h_K(t' + \pi)$ as $l_K(t + \pi)$ moves upwards. So the width of K^+ along $u_{t'}$ is also one, completing the proof. \square

Remark 3.4.2. In Lemma 3.4.8, the adjusted function h^+ might not be the supporting function h_{K^+} of K^+ .

Combine the steps above to show the balancedness of a maximum polygon cap.

Theorem 3.4.9. *Any maximum polygon cap $K_\Theta \in \mathcal{K}_\Theta^c$ is balanced.*

Proof. Assume by contradiction that $K := K_\Theta$ is not balanced. Let $h := h_K \in \mathcal{H}_\Theta$ so that $\mathcal{A}_\Theta(K) = \mathcal{A}_\Theta(h)$ by Proposition 3.3.5. Take $t \in \Theta^\circ$ with $\sigma_K(t) > \tau_K(t) \geq 0$ as in Lemma 3.4.6. Now take sufficiently small $\epsilon > 0$ and define $h^+ \in \mathcal{H}_\Theta$ and the polygon cap translate $K^+ := \mathcal{C}_\Theta(h^+) \in \mathcal{K}_\Theta^c$ as in Lemma 3.4.8. Then we have $\mathcal{A}_\Theta(h) < \mathcal{A}_\Theta(h^+)$

²That $\sigma_K(t) > 0$ is extremely crucial here. Otherwise, K^+ might have width less than zero in the angle of either $t = \omega$ or $t = \pi/2$, which is necessary for K^+ to be a polygon cap translate. This is why we choose t according to Lemma 3.4.6.

by Lemma 3.4.7. We also have $\mathcal{A}_\Theta(h^+) \leq \mathcal{A}_\Theta(K^+)$ by Proposition 3.3.7. Since K^+ is a translation of some polygon cap $K_0 \in \mathcal{K}_\Theta^c$, we have $\mathcal{A}_\Theta(K^+) = \mathcal{A}_\Theta(K_0)$. Summing up, we have

$$\mathcal{A}_\Theta(K) = \mathcal{A}_\Theta(h) < \mathcal{A}_\Theta(h^+) \leq \mathcal{A}_\Theta(K^+) = \mathcal{A}_\Theta(K_0)$$

where $K_0 \in \mathcal{K}_\Theta^c$ is a polygon cap, so $K \in \mathcal{K}_\Theta^c$ cannot attain the maximum value of \mathcal{A}_Θ , leading to contradiction. \square

Theorem 3.4.10. *Any maximum polygon cap K with angle set Θ contains its polygon niche $\mathcal{N}_\Theta(K)$.*

Proof. Denote K_Θ simply as K . By Theorem 3.4.4, it suffices to show that any vertex p of the polyline \mathbf{p}_K is contained in K . It suffices to show $p \in H_K(s)$ for any $s \in \Theta \cup \{\omega\}$. Once this is done, apply a corresponding argument to mirror reflection K^m of K (Lemma 3.4.1) and reflect back to conclude $p \in H_K(\omega + \pi/2 - t)$ for any $t \in (\omega - \Theta) \cup \{\omega\}$ as well. Then $p \in F_\omega \cap \bigcap_{t \in \Theta^\circ} H_K(t) = K$, completing the proof.

K is balanced by Theorem 3.4.9. Follow \mathbf{p}_K from right to left, from the right endpoint $A_K^-(0)$ to the point p and stop. By summing up the contribution of edges between $A_K^-(0)$ and p in \mathbf{p}_K , we have

$$A_K^-(0) - p = \sum_{t \in \Theta^\circ} c_t v_t$$

with coefficients $c_t \in [0, \tau_K(t)]$ for every $t \in \Theta^\circ$. Under this constraint on c_t , the value

$$(p - A_K^-(0)) \cdot u_s = \sum_{t \in \Theta^\circ} c_t (v_t \cdot u_s)$$

is maximized when $c_t = \tau_K(t)$ for all $t \leq s$ and $c_t = 0$ otherwise. For such c_t we have the equality

$$\sum_{t \in \Theta^\circ} c_t v_t = A_K^+(s) - A_K^-(0)$$

by using the balancedness $\tau_K(t) = \sigma_K(t)$ and following the upper boundary δK from $A_K^-(0)$ to $A_K^+(s)$. So we have

$$(p - A_K^-(0)) \cdot u_s \leq (A_K^+(s) - A_K^-(0)) \cdot u_s$$

which implies $p \cdot u_s \leq A_K^+(s) \cdot u_s$. This in turn implies $p \in H_K(s)$ as desired. \square

3.5 Balanced Maximum Sofa

We now take the limit of the balanced polygon sofas in Theorem 3.4.9 to find a monotone sofa S_ω .

Definition 3.5.1. Define the *uniform angle set* $\Theta_{\omega,n}$ of n intervals with rotation angle $\omega \in (0, \pi/2]$ as $\Theta_{\omega,n} := \{i/n\omega : 1 \leq i < n\}$.

Definition 3.5.2. For every $\omega \in (0, \pi/2]$, call a cap $K_\omega \in \mathcal{K}_\omega^c$ with the following additional data a *balanced maximum cap* with the rotation angle ω .

1. There exists a strictly increasing sequence $1 < n_1 < n_2 < \dots$ of powers of two.

2. For each $i \geq 1$, there exists a maximum polygon cap K_i with uniform angle set $\Theta_i := \Theta_{\omega, n_i}$.
3. As $i \rightarrow \infty$, the polygon cap K_i converges to K_ω in Hausdorff distance d_H .

Proposition 3.5.1. *The mirror reflection of any balanced maximum cap $K_\omega \in \mathcal{K}_\omega^c$ is also a balanced maximum cap.*

Proof. Apply Lemma 3.4.1 to the maximum polygon caps K_i converging to K_ω . \square

Theorem 3.5.2. *For every $\omega \in (0, \pi/2]$, there exists a balanced maximum cap \mathcal{K}_ω^c with rotation angle ω .*

Proof. For each $i \geq 1$, take some maximum polygon cap K_i with the uniform angle set $\Theta_{\omega, 2^i}$ of 2^i intervals (Theorem 3.4.3). Since $o_\omega \in K_i$ and every K_i is uniformly bounded in diameter $\sqrt{1 + c_{\omega, \omega/2}}$ by Lemma 3.4.2, we can use the Blaschke convergence theorem to find a convex body K that a subsequence of K_1, K_2, \dots converges to in d_H . Checking $K \in \mathcal{K}_\omega^c$ is easy. \square

Lemma 3.5.3. *Let X, Y and X_i, Y_i for all $i \geq 1$ be bounded nonempty subsets of \mathbb{R}^2 . Assume that $X_i \rightarrow X$ and $Y_i \rightarrow Y$ in Hausdorff distance d_H as $i \rightarrow \infty$. Assume also that Y is compact. Then $X_i \subseteq Y_i$ for all $i \geq 1$ implies that $X \subseteq Y$.*

Proof. For two points $p, q \in \mathbb{R}^2$, let $d(p, q)$ denote the Euclidean distance between them. For any point $p \in \mathbb{R}^2$ and $Z \subseteq \mathbb{R}^2$, define $d(p, Z) := \inf_{q \in Z} d(p, q)$. Recall that $d_H(Z_1, Z_2)$ is defined as the maximum of $\sup_{p \in Z_1} d(p, Z_2)$ and $\sup_{q \in Z_2} d(Z_1, q)$. Take any $p \in X$. Then $X_i \rightarrow X$ in d_H implies that $d(p, X_i) \rightarrow 0$. So we can take $p_i \in X_i \subseteq Y_i$ such that $d(p, p_i) \rightarrow 0$ as well. That is, $p_i \rightarrow p$. Because $Y_i \rightarrow Y$ in d_H , we have $d(p_i, Y) \rightarrow 0$. As $p_i \rightarrow p$ and Y is compact, we have $p \in Y$. \square

Theorem 3.5.4. *For any balanced maximum cap $K_\omega \in \mathcal{K}_\omega^c$ we have $\mathcal{N}(K_\omega) \subset K_\omega$.*

Proof. Write K_ω as K to avoid cluttering. Take the maximum polygon cap K_i with the uniform angle set $\Theta_i := \Theta_{\omega, n_i}$ of n_i intervals, where n_i is an increasing powers of two, so that $K_i \rightarrow K$ in d_H . Let $\Theta = \cup_i \Theta_i$ so that Θ is the set of dyadic angles. Note that $\mathcal{N}_{\Theta_1}(K) \subseteq \mathcal{N}_{\Theta_2}(K) \subseteq \dots$ by definition. We also have $\bigcup_j \mathcal{N}_{\Theta_j}(K) = \mathcal{N}(K)$ as the open set $Q_K^-(t)$ changes continuously with respect to t . Fix an arbitrary $j \geq 1$. For any $i \geq j$, we have $\mathcal{N}_{\Theta_j}(K_i) \subseteq \mathcal{N}_{\Theta_i}(K_i) \subseteq K_i$ by Theorem 3.4.10. As $K_i \rightarrow K$ in d_H , we have h_{K_i} converging to h_K uniformly so as $i \rightarrow \infty$ and j is fixed, either $\mathcal{N}_{\Theta_j}(K)$ is empty or $\mathcal{N}_{\Theta_j}(K_i) \rightarrow \mathcal{N}_{\Theta_j}(K)$ in d_H . By Lemma 3.5.3 on $\mathcal{N}_{\Theta_j}(K_i) \subseteq K_i$, we get $\mathcal{N}_{\Theta_j}(K) \subseteq K$. Now take $j \rightarrow \infty$ to conclude $\mathcal{N}(K) \subseteq K$. \square

Theorem 3.5.5. *A balanced maximum cap $K_\omega \in \mathcal{K}_\omega^c$ attains the maximum value of the sofa area functional $\mathcal{A}_\omega : \mathcal{K}_\omega^c \rightarrow \mathbb{R}$.*

Proof. Write $K := K_\omega$ to avoid cluttering, and take K_i and Θ_i as in the proof of Theorem 3.5.4 above. We will show $\mathcal{A}_{\Theta_i}(K_i) \rightarrow \mathcal{A}_\omega(K)$ as $n \rightarrow \infty$. We have $|K_i| \rightarrow |K|$ as $K_i \rightarrow K$ in Hausdorff distance and they are convex bodies (Theorem 1.8.20 of [Sch13]). So it remains to show $|\mathcal{N}_{\Theta_i}(K_i)| \rightarrow |\mathcal{N}(K)|$ as $n \rightarrow \infty$. We always have $\mathcal{A}_{\Theta_i}(K_i) \geq \mathcal{A}_{\Theta_i}(K) \geq \mathcal{A}_\omega(K)$ by Definition 3.4.1 and Theorem 3.2.3. This with $|K_i| \rightarrow |K|$ establishes $\limsup_n |\mathcal{N}_{\Theta_i}(K_i)| \leq |\mathcal{N}(K)|$. On the other hand, fix any $m \geq 1$, then

$$\liminf_n |\mathcal{N}_{\Theta_i}(K_i)| \geq \lim_{n \rightarrow \infty} |\mathcal{N}_{\Theta_m}(K_i)| = |\mathcal{N}_{\Theta_m}(K)|$$

because $h_{K_i} \rightarrow h_K$ uniformly. Now taking $m \rightarrow \infty$ shows $\liminf_n |\mathcal{N}_{\Theta_i}(K_i)| \geq |\mathcal{N}(K)|$, completing the proof.

Let m_ω be the supremum of $\mathcal{A}_\omega : \mathcal{K}_\omega^c \rightarrow \mathbb{R}$. Take any $\epsilon > 0$. There exists some $K' \in \mathcal{K}_\omega^c$ such that $\mathcal{A}_\omega(K') > m_\omega - \epsilon$. Now

$$m_\omega - \epsilon < \mathcal{A}_\omega(K') \leq \mathcal{A}_{\Theta_i}(K') \leq \mathcal{A}_{\Theta_i}(K_i)$$

by maximality of K_i on \mathcal{A}_{Θ_i} . Let $n \rightarrow \infty$ to have $m_\omega - \epsilon \leq \mathcal{A}_\omega(K)$. Take $\epsilon \rightarrow 0$ to have $m_\omega \leq \mathcal{A}_\omega(K)$, so that K attains the supremum of \mathcal{A}_ω as desired. \square

Definition 3.5.3. Call a monotone sofa S_ω with the balanced maximum cap $K_\omega := \mathcal{C}(S_\omega)$ a *balanced maximum sofa*.

Theorem 3.5.6. *A balanced maximum sofa $S_\omega := K_\omega \setminus \mathcal{N}(K_\omega)$ with the balanced maximum cap K_ω exists, and attains the maximum area among all moving sofas of rotation angle $\omega \in (0, \pi/2]$.*

Proof. By Theorem 3.5.2, a balanced maximum cap K_ω exists. Such K_ω is the cap of a monotone sofa S_ω by Theorem 3.5.4 and Theorem 2.5.9. This S_ω attains the maximum area by Theorem 2.5.10 and Theorem 3.5.5. \square

Chapter 4

Rotation Angle of Balanced Maximum Sofas

This chapter proves the Theorem 1.5.2 that any maximum-area moving sofa admits a movement with rotation angle $\omega = \pi/2$. Take any $\omega \in [\sec^{-1}(2.2), \pi/2)$. Section 4.1 compares the horizontal sides of a balanced maximum sofa S_ω . Section 4.2 proves the Theorem 1.5.2 by following the outline in Section 1.5.2.

4.1 Horizontal Side Lengths

We first establish the horizontal side length comparison in Theorem 4.1.4, which is a crucial step in the proof of main Theorem 1.5.2.

Definition 4.1.1. For any cap $K \in \mathcal{K}_\omega^c$, define $w_K^\circ := \inf_{t \in (0, \omega)} w_K(t)$ and $z_K^\circ := \inf_{t \in (0, \omega)} z_K(t)$.

Lemma 4.1.1. Assume that $\omega < \pi/2$ and the caps $K, K' \in \mathcal{K}_\omega^c$ have Hausdorff distance $\epsilon := d_H(K, K')$. Then $|w_K^\circ - w_{K'}^\circ| \leq (1 + \sec \omega)\epsilon$

Proof. Since d_H is the supremum norm between h_K and $h_{K'}$, the distance between $l_K(t)$ and $l_{K'}(t)$ is at most ϵ . So the distance between $W_K(t) = l_K(t) \cap l(\pi/2, 0)$ and $W_{K'}(t)$ is at most $\epsilon \sec \omega$. As $|h_K(0) - h_{K'}(0)| \leq \epsilon$ the distance between $A_K^-(0)$ and $A_{K'}^-(0)$ is at most ϵ . So $w_K(t) = (A_K^-(0) - W_K(t)) \cdot u_0$ and $w_{K'}(t)$ differ by at most $(1 + \sec \omega)\epsilon$. \square

Theorem 4.1.2. For any maximum polygon cap $K \in \mathcal{K}_\Theta^c$ with angle set Θ of rotation angle $\omega < \pi/2$, we have $w_K^\circ \leq \sigma_K(\pi/2)$ and $z_K^\circ \leq \sigma_K(\omega)$.

Proof. By mirror symmetry (Lemma 3.4.1), we only need to show $w_K^\circ \leq \sigma_K(\pi/2)$.

Take any $t \in \Theta$. Define the closed segment s_t of length $w_K(t)$ connecting $W_K(t)$ to $A_K^-(0)$ from left to right. By Theorem 2.5.5, s_t is on the right side of the boundary $b_K(t)$ of the wedge $T_K(t)$. So s_t is disjoint from $T_K(t)$.

Let s be the intersection of the edge $e_K(3\pi/2)$ from O to $A_K^-(0)$ and the segment s_t over all $t \in \Theta$. Since each s_t is disjoint from $T_K(t)$, the segment s is disjoint from $\mathcal{N}_\Theta(K) = \bigcup_{t \in \Theta} T_K(t)$.

We will also check that s have length $\geq w_K^\circ$. Since both $e_K(3\pi/2)$ and s_t have right endpoint $A_K^-(0)$, it suffices to check that both $e_K(3\pi/2)$ and s_t have length $\geq w_K^\circ$. For any

$t \in (0, \omega)$, the point $W_K(t)$ is the intersection of lines $l(t, h_K(t) - 1)$ and $l(\pi/2, 0)$, we have

$$\lim_{t \rightarrow \omega^-} W_K(t) = l(\omega, h_K(\omega) - 1) \cap l(\pi/2, 0) = O.$$

So

$$\lim_{t \rightarrow \omega^-} w_K(t) = \lim_{t \rightarrow \omega^-} (A_K^-(0) - W_K(t)) \cdot u_0 = h_K(0)$$

which is the length of $e_K(3\pi/2)$. So $e_K(3\pi/2)$ have length $\geq w_K^\circ$. For any $t \in \Theta$, the segment s_t have length $w_K(t)$ so it has length $\geq w_K^\circ$. Thus s have length $\geq w_K^\circ$.

Then s is a segment of length $\geq w_K^\circ$ in the edge $e_K(3\pi/2)$ disjoint from $\mathcal{N}_\Theta(K)$. So we have

$$\mathcal{H}^1(s) + \mathcal{H}^1(\mathcal{N}_\Theta(K) \cap e_K(3\pi/2)) \leq \mathcal{H}^1(e_K(3\pi/2))$$

which becomes

$$w_K^\circ + \sigma_K(3\pi/2) - \tau_K(\pi/2) \leq \sigma_K(3\pi/2)$$

by (2) of Lemma 3.4.5. So we have $w_K^\circ \leq \tau_K(\pi/2)$. By the balancedness of K (Theorem 3.4.9), we have $\tau_K(\pi/2) = \sigma_K(\pi/2)$ so the result follows. \square

Theorem 4.1.3. (Theorem 4.2.1, page 212 of [Sch13]) *As $n \rightarrow \infty$ and the convex bodies K_n converge to K in the Hausdorff distance d_H , the surface area measure σ_{K_n} on S^1 converges weakly to σ_K .*

Theorem 4.1.4. *For $\omega < \pi/2$, any balanced maximum cap $K := K_\omega$ with rotation angle ω satisfies $\sigma_K(\pi/2) \geq w_K^\circ$ and $\sigma_K(\omega) \geq z_K^\circ$. Theorem 4.1.3*

Proof. By mirror symmetry (Remark 2.5.1), it suffices to show $\sigma_K(\pi/2) \geq w_K^\circ$. Take the maximum polygon caps K_n converging to K . As $K_n \rightarrow K$ in Hausdorff distance, $w_{K_n}^\circ \rightarrow w_K^\circ$ by Lemma 4.1.1. Because $\sigma_{K_n} \rightarrow \sigma_K$ in weak convergence (Theorem 4.1.3), we have $\limsup_n \sigma_{K_n}(\pi/2) \leq \sigma_K(\pi/2)$. This combined with $w_{K_n}^\circ \leq \sigma_{K_n}(\pi/2)$ (Theorem 4.1.2) prove $w_K^\circ \leq \sigma_K(\pi/2)$ as we want. \square

4.2 Right Rotation Angle

We will prove Theorem 4.2.5 which is the main step for establishing Theorem 1.5.2 that $\omega = \pi/2$. We factor out technical calculations in lemmas, and encourage the reader to jump to Theorem 4.2.5 right away and refer to them only when needed. The outline in Section 1.5.2 and Figure 1.8 should help navigating the proof.

Definition 4.2.1. Say that a right triangle T have *base* b and *angle* θ , if an edge e of T connecting the right-angled vertex to some other vertex B have length b , and the angle at B is equal to θ . We call the length of the edge orthogonal to e the *height* of T .

Proposition 4.2.1. *For any $\omega \in [0, \pi/2)$, define $c_\omega = \tan((\pi/2 - \omega)/2)$. Then $o_\omega - v_0 = c_\omega u_0$ and $o_\omega - v_\omega = c_\omega v_\omega$. Also, the parallelogram P_ω has base $e_{P_\omega}(3\pi/2)$ connecting $O = (0, 0)$ to $l(\pi/2, 0) \cap l(\omega, 1) = (\sec \omega, 0)$, and consequently $c_\omega = \sec \omega - \tan \omega$.*

Proof. The points $O, o_\omega - v_0, o_\omega$ (resp. $O, o_\omega - v_\omega, o_\omega$) forms a right triangle with base one and angle $(\pi/2 - \omega)/2$. Let $P := l(\pi/2, 0) \cap l(\omega, 1)$, then $P = (\sec \omega, 0)$ by computation, and the points $o_\omega, o_\omega - v_0, P$ forms a right triangle with base one and angle ω . This implies $c_\omega = \sec \omega - \tan \omega$. \square

Definition 4.2.2. Let $\omega \in [\sec^{-1}(2.2), \pi/2)$ be arbitrary. Define $d_{\omega, \min}$ as 1.25 if $\omega < \tan^{-1}(2.2)$ and 1.1 otherwise.

Definition 4.2.3. Let $\omega \in [\sec^{-1}(2.2), \pi/2)$ and $d \in [0, \tan \omega]$ be arbitrary. Define the region

$$R_{\omega, d} := P_{\omega} \cap H_{-}(0, d + c_{\omega}) \cap H_{-}(\omega + \pi/2, d + c_{\omega}).$$

Lemma 4.2.2. Let $\omega \in [\sec^{-1}(2.2), \pi/2)$ be arbitrary. Then for $d = d_{\omega, \min}$, the set $R_{\omega, d}$ have area < 2.2 .

Proof. We first prove the case $\omega \geq \tan^{-1}(2.2)$. It suffices to show that the region $R_{\omega, d}$ with $d = 1.1$ in Lemma 4.2.2 have area ≤ 2.2 . Define Q_1 as the convex quadrilateral with vertices $O, o_{\omega} - v_0, o_{\omega}, o_{\omega} - v_{\omega}$. Define Q_2 as the rectangle with vertices $A := o_{\omega} - v_0 + 1.1u_0, B := o_{\omega} + 1.1u_0, o_{\omega}, o_{\omega} - v_0$. Then $|R_{\omega, 1.1}| = |Q_1| + 2|Q_2 \cap R_{\omega}|$.

Check that Q_1 is contained in the right triangle with vertices $o_{\omega}, o_{\omega} - v_0, o_{\omega} - v_0 - \cot \omega \cdot u_0$ of base one, height $\cot \omega$ and angle $\pi/2 - \omega$. So $|Q_1| \leq (\cot \omega)/2$. Check $1.1 \leq \tan \omega$ by calculating $\tan(\sec^{-1}(2.2)) > 1.959 > 1.1$. So $|Q_2 \setminus R_{\omega}|$ is a right-angled triangle with base 1.1 and angle $\pi/2 - \omega$ of area $(1.21 \cot \omega)/2$. Now we have

$$\begin{aligned} |R_{\omega, 1.1}| &= |Q_1| + 2|Q_2 \cap R_{\omega}| \\ &\leq |Q_1| + 2(|Q_2| - |Q_2 \setminus R_{\omega}|) \\ &\leq (\cot \omega)/2 + 2(1.1 - (1.21 \cot \omega)/2) < 2.2 \end{aligned}$$

proving the goal.

Now we prove the case $\omega < \tan^{-1}(2.2)$. We calculate the area of $R_{\omega, d}$ explicitly. The region $R_{\omega, d}$ is the parallelogram P_{ω} of base $\sec \omega$ and height one, subtracted by two right-angled triangles of base $b := \tan \omega - d$ and height $b \cot \omega$. So the area of $R_{\omega, d}$ is $\sec \omega - (\tan \omega - d)^2 \cot \omega$. Now it suffices to show that the area is < 2.2 for $d = d_{\omega, \min} = 1.25$. Since $\omega \in [\sec^{-1}(2.2), \tan^{-1}(2.2)]$, the following estimates hold.

$$\begin{aligned} 2.2 &\leq \sec \omega \leq \sec(\tan^{-1}(2.2)) = \sqrt{146}/5 \\ 2.2 &\geq \tan \omega \geq \tan(\sec^{-1}(2.2)) = \sqrt{96}/5 \end{aligned}$$

Now

$$\begin{aligned} |R_{\omega, 1.25}| &= \sec \omega - (\tan \omega - 1.25)^2 \cot \omega \\ &\leq \sqrt{146}/5 - \left(\sqrt{96}/5 - 1.25\right)^2 / 2.2 \\ &= 2.187736 \dots < 2.2 \end{aligned}$$

completing the proof. □

Lemma 4.2.3. For any constant $d \geq 1$, the functions $(1 - d \cot \omega)^2$ and $\cos^2 \omega$ are convex on $\omega \in [\pi/4, \pi/2]$.

Proof. To see the convexity of $(1 - d \cot \omega)^2$, compute the second derivative

$$\begin{aligned} \frac{d^2}{d\omega^2} (1 - d \cot \omega)^2 &= 2d \csc^2 \omega (d \csc^2 \omega + 2d \cot^2 \omega - 2 \cot \omega) \\ &= 2d \csc^4 \omega (d + 2d \cos^2 \omega - 2 \sin \omega \cos \omega) \\ &= 2d \csc^4 \omega (2d + d \cos(2\omega) - \sin(2\omega)) \end{aligned}$$

which is nonnegative. It is easy to check that $\cos^2 \omega = (1 + \cos(2\omega))/2$ is convex. □

Definition 4.2.4. Let $\omega \in [\sec^{-1}(2.2), \pi/2)$ and $d \in [d_{\omega, \min}, \tan \omega]$ be arbitrary. Define the values $r_y, g \in \mathbb{R}$ and $q_0, q_1 \in \mathbb{R}^2$ depending solely on ω and d as follows.

$$\begin{aligned} r_y &:= 1 - d \cot \omega \\ g &:= \sqrt{1 - r_y^2} \\ q_0 &:= o_\omega - v_0 + du_0 \\ q_1 &:= o_\omega - gu_0 \end{aligned}$$

Lemma 4.2.4. Let $\omega \in [\arctan(2.2), \pi/2)$ and $d \in [d_{\omega, \min}, \tan \omega]$ be arbitrary such that $r_y \geq 0$. Then the following inequalities are true.

1. $(q_0 - (o_\omega - v_0)) \cdot u_{\pi/2-\omega} > 1$
2. $(q_1 - (o_\omega - u_\omega)) \cdot v_{\pi/2-\omega} > 1$

Proof. (1) By the definition of q_0 , the inequality to show is $du_0 \cdot u_{\pi/2-\omega} > 1$. As $u_0 \cdot u_{\pi/2-\omega} = \sin \omega$, we need to check $d_{\omega, \min} \sin \omega > 1$. Depending on whether $\omega < \tan^{-1}(2.2)$ or not, computations

$$1.25 \sin(\sec^{-1}(2.2)) = 1.1134 \dots > 1 \quad 1.1 \sin(\tan^{-1}(2.2)) = 1.0014 \dots > 1$$

prove the result.

(2) By the definition of q_1 , the inequality to show is $(u_\omega - gu_0) \cdot v_{\pi/2-\omega} > 1$. Computing the left-hand side gives $-\cos(2\omega) + g \cos \omega > 1$. Rearranging the terms, we need to prove $g > 2 \cos \omega$ as $(1 + \cos(2\omega))/\cos \omega = 2 \cos \omega$. Both g and $2 \cos \omega$ are nonnegative, so we only need to compare their squares and prove $g^2 = 1 - r_y^2 > 4 \cos^2 \omega$. Since $0 \leq r_y = 1 - d \cot \omega \leq 1 - d_{\omega, \min} \cot \omega$, it suffices to prove

$$1 > (1 - d_{\omega, \min} \cot \omega)^2 + 4 \cos^2 \omega \tag{4.1}$$

on $\omega \in [\arctan(2.2), \pi/2)$.

As $d_{\omega, \min}$ is the constant 1.25 or 1.1 on each interval $\omega \in (\sec^{-1}(2.2), \tan^{-1}(2.2)]$ or $\omega \in (\tan^{-1}(2.2), \pi/2]$, both $(1 - d_{\omega, \min} \cot \omega)^2$ and $4 \cos^2 \omega$ are convex functions of ω on each interval by Lemma 4.2.3. So it suffices to check Equation (4.1) at the four endpoints, which are true by following calculations.

$$\begin{aligned} (1 - 1.25 \cot(\sec^{-1}(2.2)))^2 + 4 \cos^2(\sec^{-1}(2.2)) &< 0.9576 < 1 \\ (1 - 1.25 \cot(\tan^{-1}(2.2)))^2 + 4 \cos^2(\tan^{-1}(2.2)) &< 0.8714 < 1 \\ (1 - 1.1 \cot(\tan^{-1}(2.2)))^2 + 4 \cos^2(\tan^{-1}(2.2)) &< 0.9350 < 1 \\ (1 - 1.1 \cot(\pi/2))^2 + 4 \cos^2(\pi/2) &= 1 \end{aligned}$$

□

Theorem 4.2.5. Let $\omega \in [\sec^{-1}(2.2), \pi/2)$ be arbitrary. Let $K := K_\omega$ be any balanced maximum cap with rotation angle ω in Theorem 3.5.2. Assume also that $\mathcal{A}_\omega(K) \geq 2.2$. Then there exists some $t \in (0, \omega)$ such that the three points $O, o_\omega - v_0, o_\omega - u_\omega$ are in the closure of the quadrant $Q_K^-(t)$.

Proof. We first show that at least one of $h_K(0)$ or $h_K(\omega + \pi/2)$ should be large enough. That is, $h_K(0) \geq d_{\omega, \min} + c_\omega$ or $h_K(\omega + \pi/2) \geq d_{\omega, \min} + c_\omega$ should be true (see Proposition 4.2.1 and Definition 4.2.2). Assume the contrary. Then the cap K is contained in the set $R_{\omega, d}$ with $d = d_{\omega, \min}$ in Definition 4.2.3. But by Lemma 4.2.2 the area of $R_{\omega, d}$ is less than 2.2, contradicting the assumption $|K| \geq \mathcal{A}_\omega(K) \geq 2.2$.

So we have either $h_K(0) \geq d_{\omega, \min} + c_\omega$ or $h_K(\omega + \pi/2) \geq d_{\omega, \min} + c_\omega$. Appeal to the mirror reflection¹ of K and assume without loss of generality that $h_K(0) \geq d_{\omega, \min} + c_\omega$. Now define $d := h_K(0) - c_\omega$ so that $d \geq d_{\omega, \min}$ in particular. We now have the point $A_K^-(0) = q_0$ in K , where $q_0 := o_\omega - v_0 + du_0$ is in Definition 4.2.4. In particular, q_0 is in K .

Define the intersection $r := l_K(0) \cap l_K(\omega)$. Then $r, o_\omega, o_\omega + du_0$ forms a right-angled triangle. Solving for the coordinates of r , we get $r = (d + c_\omega, r_y)$ where $r_y := 1 - d \cot \omega$ is defined in Definition 4.2.4. Since K is a convex subset of P_ω we have $0 \leq r_y \leq 1$. Define $g := \sqrt{1 - r_y^2}$ as in Definition 4.2.4 and the point $s := q_0 - gu_0$, so that the triangle with vertices s, q_0, r is right-angled at q_0 with base g , height r_y and side of length 1.

The main idea is that the point s is not contained in $Q_K^-(t)$ for any $t \in (0, \omega)$, so that $g \leq w_K^\circ$. We prove this rigorously. Take any $t \in (0, \omega)$. Since $r := l_K(0) \cap l_K(\omega)$ and K is a convex body, we have $h_K(t) \leq r \cdot u_t$. So

$$h_K(t) - 1 \leq r \cdot u_t - 1 \leq r \cdot u_t - (r - s) \cdot u_t = s \cdot u_t$$

as $r - s$ is a unit vector. Thus $s \notin H_-^\circ(h_K(t) - 1, t)$ and s is further than $W_K(t) = b_K(t) \cap l(\pi/2, 0)$ in the direction of u_0 . This with $s = q_0 - gu_0$ and $W_K(t) = q_0 - w_K(t)u_0$ implies $g \leq w_K(t)$. Thus we get $g \leq w_K^\circ$.

By Theorem 4.1.4 we have $g \leq w_K^\circ \leq \sigma_K(\pi/2)$. Define $q_1 := o_\omega - gu_0$ as in Definition 4.2.4, then as o_ω is the rightmost vertex of K on the edge $e_K(\pi/2)$ we also have $q_1 \in K$.

We now have two points q_0, q_1 in K , and the set $X = \{q_0, q_1\}$ is a subset of K . From now on, fix the angle $t = \pi/2 - \omega \in (0, \pi/4) \subset (0, \omega)$ and take the quadrant $Q_X^-(t)$. By $X \subset K$ we have $Q_X^-(t) \subseteq Q_K^-(t)$. So the proof is done if we show that the three points $O, o_\omega - v_0, o_\omega - u_\omega$ are in the closure of $Q_X^-(t)$.

We will show that $q_1 - q_0 = -\alpha u_t + \beta v_t$ for real coefficients $\alpha, \beta \geq 0$. By definition of q_0 and q_1 we have $q_1 - q_0 = v_0 - (d + g)u_0$ where $d + g \geq d \geq 1.1$. So the vector $q_1 - q_0$ is in the convex cone generated by $-u_0$ and $v_0 - u_0$. Because the angle $t = \pi/2 - \omega$ is in between 0 and $\pi/4$, both $-u_0$ and $v_0 - u_0$ are in the convex cone generated by $-u_t$ and v_t . So the vector $q_1 - q_0$ is in the convex cone generated by $-u_t$ and v_t , and we have $\alpha, \beta \geq 0$.

Now by $q_1 - q_0 = -\alpha u_t + \beta v_t$ for $\alpha, \beta \geq 0$, and the definition of $L_X(t)$, the point q_0 lies in the outer wall $a_X(t)$ of $L_X(t)$ and q_1 lies in the outer wall $c_X(t)$ of $L_X(t)$. Thus we can write

$$Q_X^-(t) = H_-^\circ(t, q_0 \cdot u_t - 1) \cap H_-^\circ(t + \pi/2, q_1 \cdot v_t - 1). \quad (4.2)$$

We now show that the three points $O, o_\omega - v_0 = c_\omega u_0, o_\omega - u_\omega = c_\omega v_\omega$ (Proposition 4.2.1) are contained in $Q_X^-(t)$. By Equation (4.2) and $c_\omega > 0$, it suffices to show that $c_\omega u_0 \in H_-^\circ(t, q_0 \cdot u_t - 1)$ and $c_\omega v_\omega \in H_-^\circ(t + \pi/2, q_1 \cdot v_t - 1)$, both are true by Lemma 4.2.4. \square

Theorem 1.5.2 is now a consequence of Theorem 4.2.5 as described in Section 1.5.1.

Proof of Theorem 1.5.2. By Theorem 4.2.5, the triangle Δ_ω formed by $O, o_\omega - v_0, o_\omega - u_\omega$ is contained in $\mathcal{N}(K_\omega)$, so is disjoint from S_ω (left of Figure 1.7). Now $S_\omega \subseteq P_\omega \setminus \Delta_\omega$ and

¹This argument depends on the fact that the statement of Theorem 4.2.5 on K and its mirror image K^m is equivalent. In particular, the angle $t \in (0, \omega)$ that we take for K would correspond to the angle $\omega - t$ for K^m .

observe that the set $P_\omega \setminus \Delta_\omega$ have width ≤ 1 for every direction u_t with angle $t \in [\omega, \pi/2]$. So S_ω can rotate counterclockwise by $\pi/2 - \omega$ inside H (right of Figure 1.7). Take S' as a copy of $R_{\pi/2-\omega}(S_\omega)$ translated horizontally to the left inside H_L . First rotate S' clockwise by $\pi/2 - \omega$ inside H_L . Then translate it to the right until it hits the wall $x = 1$ of L , to put it in the initial position of the monotone sofa S_ω . Then follow the original movement of S_ω with the rotation angle ω . We have found a movement of S' with rotation angle $\pi/2$, so $|S_\omega| = |S'| \leq |S_{\pi/2}|$ and this completes the proof. \square

Chapter 5

Surface Area Measure

This chapter proves the equality $dv_K^\pm(t) = v_t \sigma_K$ described in Section 1.6. Section 5.1 defines the Lebesgue–Stieltjes measure, and Section 5.2 proves the equality in Theorem 5.2.2.

5.1 Lebesgue–Stieltjes Measure

All real-valued measurable functions used in this and upcoming chapters will be bounded and defined on some finite interval I of \mathbb{R} . All measures μ used in this and upcoming chapters will be finite signed Borel measures on some finite interval I of \mathbb{R} . That is, for any Borel subset X of I , the value $\mu(X)$ will be real and not $\pm\infty$.

Here, we will define the *Lebesgue–Stieltjes measure* df of a right-continuous $f : I \rightarrow \mathbb{R}$ of bounded variation (Definition 5.1.3). We use the following notations.

Definition 5.1.1. For any bounded measurable function f and finite signed Borel measure μ on a finite interval I of \mathbb{R} , define the scalar multiplication $f\mu$ of f and μ as the measure on I defined as $f\mu(X) := \int_{t \in X} f(t) \mu(dt)$. Note that $f\mu$ is also a finite signed Borel measure.

- If f is a pair (f_1, f_2) of bounded measurable function and μ is a finite signed Borel measure, the notion $f\mu$ denotes the pair $(f_1\mu, f_2\mu)$.
- If f is a bounded measurable function and $\mu = (\mu_1, \mu_2)$ is a pair of finite signed Borel measure, then the notion $f\mu$ denotes $(f\mu_1, f\mu_2)$.
- If both $f = (f_1, f_2)$ and $\mu = (\mu_1, \mu_2)$ are pairs of bounded measurable functions and finite signed Borel measures respectively, then $f \cdot \mu$ denotes $f_1\mu_1 + f_2\mu_2$.

Recall the following standard real analysis definition.

Definition 5.1.2. A function $f : [a, b] \rightarrow \mathbb{R}$ is of *bounded variation* if there is an absolute constant C such that, for any partition $a = t_0 < t_1 < \dots < t_n = b$ of I , the sum $\sum_{i=1}^n |f(t_i) - f(t_{i-1})|$ is bounded from above by C .

We use the notation df to denote the *Lebesgue–Stieltjes measure* of f .

Definition 5.1.3. (Theorem 4.3, page 5 of [RY13]) For any right-continuous $f : I \rightarrow \mathbb{R}$ on interval $I := [a, b]$ of bounded variation, define the *Lebesgue–Stieltjes measure* df on I as the

unique finite signed Borel measure such that $df(\{a\}) = 0$ and $df((a, t]) = f(t) - f(a)$ for all $t \in I$.

If $f(t) = (f_1(t), f_2(t))$ is a pair of such functions, then df denotes the pair (df_1, df_2) of measures on I .

Definition 5.1.4. Fix an interval $I := [a, b]$ parametrized by t . Let $f : I \rightarrow \mathbb{R}$ be arbitrary right-continuous function of bounded variation. Let $g : I \rightarrow \mathbb{R}$ be bounded and measurable. Let X be any Borel subset of I . Define the *Lebesgue–Stieltjes integral* of g on X with respect to f as

$$\int_{t \in X} g(t) df(dt)$$

which is the integral of g on X with respect to the Lebesgue–Stieltjes measure df of f . We also denote the integral as

$$\int_{t \in X} g(t) df(t) \quad \text{or} \quad \int_X g df.$$

The map $f \mapsto df$ is linear like differentials would do.

Proposition 5.1.1. For any right-continuous $f, g : [a, b] \rightarrow \mathbb{R}$ of bounded variation and real values $r, s \in \mathbb{R}$, we have $d(rf + sg) = r df + s dg$ as signed measures on I .

The product rule $d(fg) = g df + f dg$ is more subtle. We need one of f or g to be continuous.

Lemma 5.1.2. (Proposition 4.5, page 6 of [RY13]) For any right-continuous $f, g : [a, b] \rightarrow \mathbb{R}$ of bounded variation, we have

$$\int_{t \in (a, b]} g(t) df(t) + \int_{t \in (a, b]} f(t-) dg(t) = f(b)g(b) - f(a)g(a)$$

where $f(t-)$ denotes the left limit of f at t .

Lemma 5.1.3. For any right-continuous $f, g : [a, b] \rightarrow \mathbb{R}$ of bounded variation, if one of f or g is continuous then the equality $d(fg) = g df + f dg$ holds.

Proof. Assume without loss of generality that f is continuous. It suffices to show that both sides, as measures on $[a, b]$, agree on the subset $(a, x]$ for all $x \in (a, b]$. This is true by Lemma 5.1.2. \square

Finally, we note the following characterization of absolutely continuous functions f .

Proposition 5.1.4. For any right-continuous $f : [a, b] \rightarrow \mathbb{R}$ of bounded variation, the followings are equivalent.

1. The function $f : [a, b] \rightarrow \mathbb{R}$ is absolutely continuous.
2. We have $df = r dt$ for some measurable and bounded $r : [a, b] \rightarrow \mathbb{R}$.

In such a case, we have $f'(t)$ equal to $r(t)$ for almost every $t \in [a, b]$.

Proof. (2 \Rightarrow 1) By Definition 5.1.3, we have $f(t) = f(a) + \int_a^t r(s) ds$, so f should be absolutely continuous. (1 \Rightarrow 2) Let $r : [a, b] \rightarrow \mathbb{R}$ be the derivative of $f(t)$ that exists on almost every $t \in [a, b]$. Then $f(t) = f(a) + \int_a^t r(s) ds$ by absolute continuity of f , so by the uniqueness of df we should have $df = r dt$. \square

5.2 Differential Gauss–Minkowski Theorem

Recall that for any $K \in \mathcal{K}$, the vertex $v_K^+(t)$ is right-continuous with respect to $t \in S^1$ (Theorem 2.1.3). It is also of bounded variation, so the pair $dv_K^+(t)$ of Lebesgue–Stieltjes measures of the x - and y -coordinates of $v_K^+(t)$ exists.

Lemma 5.2.1. *For any $K \in \mathcal{K}$ and any interval $[a, b]$ of S^1 , the function $v_K^+ : [a, b] \rightarrow \mathbb{R}^2$ is of bounded variation.*

Proof. For the interval $t \in [0, \pi/4]$, observe that the x -coordinate (resp. y -coordinate) of $v_K^+(t)$ monotonically decreases (resp. increases) with respect to t , so v_K^+ is of bounded variation on $[0, \pi/4]$. A similar logic can be used to angles $[\pi/4, \pi/2]$, $[\pi/2, 3\pi/4]$, and $[3\pi/4, 2\pi]$. Any larger domain $[a, b]$ of t can be divided into such intervals or their subintervals. \square

Theorem 5.2.2 evaluates $dv_K^+(t)$ in terms of the surface area measure σ_K of K . This will be used frequently.

Theorem 5.2.2. *Let $a, b \in \mathbb{R}$ be arbitrary such that $a < b \leq a + 2\pi$. Let K be any planar convex body. Then the equality*

$$dv_K^+(t) = v_t \sigma_K$$

of pairs of measures on the half-open interval $I := (a, b]$ holds,¹ where $v_K^+(t) : \bar{I} \rightarrow \mathbb{R}^2$ and $v_t : \bar{I} \rightarrow \mathbb{R}^2$ are taken as functions of $t \in \bar{I} = [a, b]$.

Note that the notations $v_K^+(t)$ and v_t denote different things: $v_K^+(t)$ is a vertex of the convex body K , while v_t is the direction $(-\sin t, \cos t)$ independent of K . Integrating Theorem 5.2.2 on any bounded measurable function $p : I \rightarrow \mathbb{R}^2$, we get

$$\int_{t \in I} p(t) \cdot dv_K^+(t) = \int_{t \in I} (p(t) \cdot v_t) \sigma(dt).$$

Proof of Theorem 5.2.2. It suffices to check that the pairs of measures $dv_K^+(t)$ and $u_t \sigma$ agree on the subset $(a, x]$ of I for any $x \in I$. That is, we only need to check

$$v_K^+(x) - v_K^+(a) = \int_{t \in (a, x]} v_t \sigma_K(dt). \quad (5.1)$$

We first show Equation (5.1) for polygon K . For polygon K , the measure σ_K is a discrete measure where each proper edge $e_K(t)$ of K with normal angle t corresponds to a point mass of σ_K concentrated at t with the weight $\sigma_K(\{t\})$ which is the length of $e_K(t)$. So the right-hand side of Equation (5.1) is the sum of all vectors $v_t \sigma_K(\{t\}) = v_K^+(t) - v_K^-(t)$ over all normal angles $t \in (a, x]$ of proper edges of K . The telescopic sum is the left-hand side $v_K^+(x) - v_K^+(a)$ as we want.

Now we prove Equation (5.1) for general convex body K . As in the proof of Theorem 8.3.3, page 466 of [Sch13], we can take a series K_1, K_2, \dots of polygons converging to K in the Hausdorff distance d_H such that $e_{K_n}(a) = e_K(a)$ and $e_{K_n}(x) = e_K(x)$ for all $n \geq 1$. By Theorem 4.1.3, the measure σ_{K_n} on S^1 converges to σ_K weakly as $n \rightarrow \infty$.

For any measure σ on S^1 , define the restriction $\sigma|_A$ of σ to a Borel subset $A \subseteq S^1$ as the measure on S^1 such that $\sigma|_A(X) = \sigma(A \cap X)$ for all Borel subset $X \subseteq S^1$. Define U as the open set $S^1 \setminus \{a, x\}$ of S^1 , and V as the open interval (a, x) of S^1 . Define u_n and u

¹The equality does *not* hold in general on the left endpoint $\{a\}$, as it is (somewhat artificially) defined in Definition 5.1.3 that the Lebesgue–Stieltjes measure is zero on the left endpoint.

as the restriction of σ_{K_n} and σ_K to U , then u_n converges to u weakly as $n \rightarrow \infty$ because $\sigma_{K_n}(\{a\}) = \sigma_K(\{a\})$ and $\sigma_{K_n}(\{x\}) = \sigma_K(\{x\})$.

Define λ_n and λ as the restriction of σ_{K_n} and σ_K to V . We will prove that $\lambda_n \rightarrow \lambda$ weakly as $n \rightarrow \infty$. Take any continuity set $X \subseteq S^1$ of λ so that $\lambda(\partial X) = 0$. By the Portmanteau theorem on finite measures, it suffices to show $\lambda_n(X) \rightarrow \lambda(X)$. Because $\partial(X \cap V) \subseteq (\partial X \cap V) \cup \partial V$, and both $u(\partial X \cap V) = \lambda(\partial X)$ and $u(\partial V)$ are zero, the set $X \cap V$ is a continuity set of u . So $u_n(X \cap V) \rightarrow u(X \cap V)$ and thus $\lambda_n(X) \rightarrow \lambda(X)$ as $n \rightarrow \infty$. This completes the proof that $\lambda_n \rightarrow \lambda$ weakly as $n \rightarrow \infty$.

Now take the limit $n \rightarrow \infty$ to the Equation (5.1) for polygons K_n :

$$v_{K_n}^+(x) - v_{K_n}^+(a) = \int_{t \in (a, x]} v_t \sigma_{K_n}(dt).$$

The left-hand side is equal to $v_K^+(x) - v_K^+(a)$ by $e_{K_n}(a) = e_K(a)$ and $e_{K_n}(x) = e_K(x)$. The right-hand side is equal to

$$(v_{K_n}^+(x) - v_{K_n}^-(x)) + \int_{t \in S^1} v_t \lambda_n(dt)$$

and by $e_{K_n}(x) = e_K(x)$ and the weak convergence $\lambda_n \rightarrow \lambda$, the expression converges to

$$(v_K^+(x) - v_K^-(x)) + \int_{t \in S^1} v_t \lambda(dt) = \int_{t \in (a, x]} v_t \sigma_K(dt)$$

thus completing the proof of Equation (5.1) for general convex body K . \square

Remark 5.2.1. The *Gauss–Minkowski correspondence* maps any planar convex body K , up to translation, bijectively to the Borel measure $\sigma := \sigma_K$ on S^1 satisfying $\int_{t \in S^1} v_t \sigma(dt) = 0$ ([MR14] or Theorem 8.3.1 of [Sch13]). Theorem 5.2.2 can be seen as a differential version of this correspondence, as we can recover it from integrating Theorem 5.2.2.

- By integrating both sides of Theorem 5.2.2 over all $t \in S^1$, we immediately the equality $\int_{t \in S^1} v_t \sigma_K(dt) = 0$ which is one direction of the correspondence.
- For any Borel measure σ on S^1 such that $\int_{t \in S^1} v_t \sigma(dt) = 0$, we can recover K with $\sigma_K = \sigma$ by taking K as the convex hull of the partial integrals $v^-(s) := \int_{(0, s]} v_t \sigma(dt)$ and $v^+(s) := \int_{(0, s]} v_t \sigma(dt)$ of Theorem 5.2.2 for all $s \in (0, 2\pi]$. While we omit the details, it is easy to see that such points are in convex position and $v^\pm(s) = v_K^\pm(s)$, so that $\sigma_K = \sigma$.

Chapter 6

Injectivity Condition

This chapter follows the sketch in Section 1.7 and proves the injectivity condition on any balanced maximum sofa S . The proof is centered around the differential inequality Equation (1.9).

- Section 6.1 states the full injectivity condition.
- Section 6.2 defines the *arm lengths* f_K, g_K of a cap K and makes calculations related to it.
- Section 6.3 establishes a discrete version of Equation (1.9) on maximum polygon sofas (Theorem 6.3.3). The proof follows the sketch in Section 1.7.2.
- Section 6.4 takes limit on the inequality on maximum polygon sofas, and establish the continuous version of Equation (1.9) (Theorem 6.4.3) by taking the limit in the discrete version.

6.1 Statement

We showed in the previous Chapter 4 that we can assume the rotation angle $\omega = \pi/2$ for the moving sofa problem (Theorem 1.5.2). So we will omit ω in the subscript to denote $\omega = \pi/2$.

Definition 6.1.1. Define \mathcal{K}^c as the space of caps $\mathcal{K}_{\pi/2}^c$ with rotation angle $\pi/2$. Define $\mathcal{A} : \mathcal{K}^c \rightarrow \mathbb{R}$ as the sofa area functional $\mathcal{A}_{\pi/2}$ with rotation angle $\pi/2$.

We fully state the main Theorem 6.1.1 of this paper. Recall that a *balanced maximum cap* K is the cap of a *balanced maximum sofa* S with rotation angle $\pi/2$ attaining the maximum area α_{\max} (Theorem 3.5.6). Note that the following Definition 6.1.2 also defines the notions r_K and s_K for K satisfying the injectivity condition.

Definition 6.1.2. Say that a cap $K \in \mathcal{K}^c$ satisfies the *injectivity condition* if the followings are true.

1. There exists measurable functions $r_K, s_K : [0, \pi/2] \rightarrow \mathbb{R}_{\geq 0}$, unique up to measure zero, such that $\sigma_K = r_K(t)dt$ on the interval $t \in [0, \pi/2)$ and $\sigma_K = s_K(t - \pi/2)dt$ on the interval $t \in (\pi/2, \pi]$.

2. The inner corner $\mathbf{x}_K : [0, \pi/2] \rightarrow \mathbb{R}^2$ is continuously differentiable.
3. For all $t \in (0, \pi/2)$, we have $\mathbf{x}'_K(t) \cdot u_t < 0$ and $\mathbf{x}'_K(t) \cdot v_t > 0$.

Theorem 6.1.1. *Any balanced maximum cap $K \in \mathcal{K}^c$ satisfies the injectivity condition.*

We will establish Theorem 6.1.1 in the last Section 6.5 of this chapter. Once this is done, we can also say the following.

Theorem 6.1.2. *The cap $K := \mathcal{C}(G)$ of Gerver's sofa satisfies the injectivity condition.*

Proof. Theorem 2 of [Ger92] explicitly constructs a sequence of maximum polygon sofas converging to Gerver's sofa G . So G is a balanced maximum sofa, and Theorem 6.1.1 proves the claim. \square

Remark 6.1.1. Note that for the cap $K := \mathcal{C}(G)$ of Gerver's sofa, a slightly weaker version $\mathbf{x}'_K(t) \cdot u_t \leq 0$ and $\mathbf{x}'_K(t) \cdot v_t \geq 0$ of (3) of Definition 6.1.2 is already assumed by Romik [Rom18] in order to derive Gerver's sofa G .

Although Romik does not explicitly put the derived G back and verify this starting assumption, the equations determining G as provided in [Rom18] should be sufficient to verify Theorem 6.1.2 independently of Theorem 2 by [Ger92]. In particular, the red graph of Figure 1.10 depicts the numerical values of $f_K(t) = 1 - \mathbf{x}'_K(t) \cdot u_t$ (Proposition 6.4.6), verifying one inequality of (3) of Definition 6.1.2 numerically.

6.2 Arm Lengths

Define the *arm lengths* $f_K^\pm(t)$ and $g_K^\pm(t)$ of supporting hallways of a cap K as the following.

Definition 6.2.1. Let $K \in \mathcal{K}^c$ and $t \in [0, \pi/2]$ be arbitrary. Define

$$f_K^+(t) = (\mathbf{y}_K(t) - A_K^+(t)) \cdot v_t \quad f_K^-(t) = (\mathbf{y}_K - A_K^-(t)) \cdot v_t$$

and

$$g_K^+(t) = (\mathbf{y}_K(t) - C_K^+(t)) \cdot u_t \quad g_K^-(t) = (\mathbf{y}_K(t) - C_K^-(t)) \cdot u_t.$$

Proposition 6.2.1. *Let $K \in \mathcal{K}^c$ and $t \in [0, \pi/2]$ be arbitrary. Then we have $\mathbf{y}_K(t) = A_K^\pm(t) + f_K^\pm(t)v_t$ and $\mathbf{y}_K(t) = C_K^\pm(t) + g_K^\pm(t)u_t$.*

Proof. The point $\mathbf{y}_K(t)$ and the vertices $A_K^\pm(t)$ are on the tangent line $a_K(t)$ in the direction of v_t . Likewise $\mathbf{y}_K(t)$ and $C_K^\pm(t)$ are on the tangent line $c_K(t)$ in the direction of u_t . \square

Proposition 6.2.2. *For any $K \in \mathcal{K}^c$ and $t \in [0, \pi/2]$, we have $f_{K^m}^\pm(t) = g_K^\mp(t)$ and $g_{K^m}^\pm(t) = f_K^\mp(t)$.*

Proof. By Proposition 2.5.4 and Definition 6.2.1. \square

It turns out that the condition (3) of Definition 6.1.2 is equivalent to stating that $f_K^\pm(t) > 1$ and $g_K^\pm(t) > 1$, because the derivative of \mathbf{x}_K can be expressed in terms of the arm lengths of K (Theorem 6.2.3).

Definition 6.2.2. For any function f from interval $I \subseteq \mathbb{R}$ to \mathbb{R} , denote its left (resp. right) derivative at $t \in I$ as $\partial^+ f$ (resp. $\partial^- f$).

Theorem 6.2.3. *For any $K \in \mathcal{K}^c$ and $t \in [0, \pi/2)$, the right derivatives of the outer corner $\mathbf{y}_K(t)$ and inner corner $\mathbf{x}_K(t)$ exists for all $0 \leq t < \pi/2$ and is equal to the following.*

$$\partial^+ \mathbf{y}_K(t) = -f_K^+(t)u_t + g_K^+(t)v_t \quad \partial^+ \mathbf{x}_K(t) = -(f_K^+(t) - 1)u_t + (g_K^+(t) - 1)v_t$$

Likewise, the left derivatives of $\mathbf{y}_K(t)$ and $\mathbf{x}_K(t)$ exists for all $0 < t \leq \pi/2$ and is equal to the following.

$$\partial^- \mathbf{y}_K(t) = -f_K^-(t)u_t + g_K^-(t)v_t \quad \partial^- \mathbf{x}_K(t) = -(f_K^-(t) - 1)u_t + (g_K^-(t) - 1)v_t$$

Proof. Fix an arbitrary cap K and omit the subscript K in the arm lengths $f_K^\pm(t), g_K^\pm(t)$, vertices $\mathbf{y}_K(t), \mathbf{x}_K(t)$ and the tangent line $a_K(t)$. Take any $0 \leq t < \pi/2$ and set $s = t + \delta$ for sufficiently small and arbitrary $\delta > 0$. We evaluate $\partial^+ \mathbf{y}(t) = \lim_{\delta \rightarrow 0^+} (\mathbf{y}(s) - \mathbf{y}(t))/\delta$. Define $A_{t,s} = a(t) \cap a(s)$. Since $A_{t,s}$ is on the lines $a(t)$ and $a(s)$, it satisfies both $A_{t,s} \cdot u_t = \mathbf{y}(t) \cdot u_t$ and $A_{t,s} \cdot u_s = \mathbf{y}(s) \cdot u_s$. Rewrite $u_s = (\cos \delta)u_t + (\sin \delta)v_t$ on the second equation and we have

$$(\cos \delta)A_{t,s} \cdot u_t + (\sin \delta)A_{t,s} \cdot v_t = \cos \delta(\mathbf{y}(s) \cdot u_t) + \sin \delta(\mathbf{y}(s) \cdot v_t).$$

Group by $\cos \delta$ and $\sin \delta$ and substitute $A_{t,s} \cdot u_t$ with $\mathbf{y}(t) \cdot u_t$, then

$$\cos \delta(\mathbf{y}(s) \cdot u_t - \mathbf{y}(t) \cdot u_t) = \sin \delta(A_{t,s}(s) \cdot v_t - \mathbf{y}(s) \cdot v_t).$$

Divide by δ and send $\delta \rightarrow 0^+$. We get the following limit as $A_{t,s} \rightarrow A^+(t)$ (Theorem 2.1.3).

$$\partial^+(\mathbf{y}(t) \cdot u_t) = (A^+(t) - \mathbf{y}(t)) \cdot v_t = -f^+(t)$$

A similar argument can be applied to show $\partial^+(\mathbf{y}(t) \cdot v_t) = g^+(t)$ and thus the first equation of the theorem. The right derivative of $\mathbf{x}_K(t)$ comes from $\mathbf{x}_K(t) = \mathbf{y}_K(t) - u_t - v_t$. A mirror-symmetric argument calculates the left derivative of \mathbf{y}_K and \mathbf{x}_K . \square

Remark 6.2.1. The resulting equation $\partial^+ \mathbf{x}_K(t) = -(f_K^+(t) - 1)u_t + (g_K^+(t) - 1)v_t$ in Theorem 6.2.3 can be interpreted intuitively as the following. Imagine moving the hallway $L_K(t)$ slightly by incrementing t by small $\epsilon > 0$. The wall $c_K(t)$ rotates with the pivot $C_K^+(t)$ as center. So the v_t component $g_K^+(t) - 1$ of the derivative $\partial^+ \mathbf{x}_K(t)$ is the distance from the pivot $C_K^+(t)$ to $\mathbf{x}_K(t)$ measured in the direction of u_t . The u_t component $-(f_K^+(t) - 1)$ of $\partial^+ \mathbf{x}_K(t)$ can be interpreted similarly as the distance from the pivot $A_K^+(t)$ to $\mathbf{x}_K(t)$ along the direction v_t .

We prove some lemmas that compute the arm lengths.

Lemma 6.2.4. *For any $K \in \mathcal{K}^c$ and $t \in [0, \pi/2]$, we have*

$$g_K^+(t) = \int_{u \in (t, t + \pi/2]} \sin(u - t) \sigma_K(du).$$

Proof. Since $\mathbf{y}_K(t), A_K^+(t) \in l_K(t)$, we have

$$g_K^+(t) = (A_K^+(t) - C_K^+(t)) \cdot u_t = -u_t \cdot (v_K^+(t + \pi/2) - v_K^+(t)).$$

Parametrize the interval $(t, t + \pi/2]$ by s , then we have $dv_K^+(s) = v_s \sigma_K$ by Theorem 5.2.2 so

$$\begin{aligned} g_K^+(t) &= -u_t \cdot (v_K^+(t + \pi/2) - v_K^+(t)) \\ &= -u_t \cdot \int_{s \in (t, t + \pi/2]} dv_K^+(t') = -u_t \cdot \int_{s \in (t, t + \pi/2]} v_s \sigma_K(ds) \\ &= \int_{s \in (t, t + \pi/2]} (-u_t \cdot v_s) \sigma_K(ds) \\ &= \int_{s \in (t, t + \pi/2]} \sin(s - t) \sigma_K(ds) \end{aligned}$$

is proved. \square

We calculate the differentiation of arm length $f_K^+(t)$.

Theorem 6.2.5. *For any $K \in \mathcal{K}$, the function $f_K^+(t)$ on $t \in [0, \pi/2]$ is right-continuous and of bounded variation. Moreover, its Lebesgue–Stieltjes measure on the half-open interval $t \in (0, \pi/2]$ is*

$$df_K^+(t) = g_K^+(t) dt - \sigma_K$$

where $f_K^+(t), g_K^+(t), t$ are functions of $t \in [0, \pi/2]$, so that dt denotes the usual Borel measure of $(0, \pi/2]$.

Proof. First evaluate

$$\begin{aligned} dh_K(t) &= d(v_K^+(t) \cdot u_t) = u_t \cdot dv_K^+(t) + v_K^+(t) \cdot du_t \\ &= u_t \cdot (v_t \sigma_K) + (v_K^+(t) \cdot v_t) dt = (v_K^+(t) \cdot v_t) dt \end{aligned}$$

for all $t \in S^1$ using Lemma 5.1.3 and Theorem 5.2.2. Now take the Lebesgue–Stieltjes measure of

$$f_K^+(t) = (\mathbf{y}_K(t) - A_K^+(t)) \cdot v_t = h_K(\pi/2 + t) - v_K^+(t) \cdot v_t$$

for all $t \in (0, \pi/2]$ using Lemma 5.1.3 and Theorem 5.2.2 to get

$$\begin{aligned} df_K^+(t) &= dh_K(\pi/2 + t) - v_t \cdot dv_K^+(t) - v_K^+(t) \cdot dv_t \\ &= (v_K^+(\pi/2 + t) \cdot v_{\pi/2+t}) dt - v_t \cdot (v_t \sigma_K) + v_K^+(t) \cdot u_t dt \\ &= -\sigma_K + (-v_K^+(\pi/2 + t) \cdot u_t + h_K(t)) dt = g_K^+(t) dt - \sigma_K \end{aligned}$$

which completes the proof. \square

6.3 Inequality on Maximum Polygon Caps

In this Section 6.3, we prove Theorem 6.3.3 that bounds the side lengths of a maximum polygon cap K .

Recall that a balanced maximum cap K_∞ defined as the limit of *maximum polygon caps* K_1, K_2, \dots converging to K in Hausdorff distance d_H (Definition 3.5.2). Each maximum polygon cap K_i have angle set $\Theta_{n_i} := \{(\pi/2)j/n_i : 1 \leq j < n_i\}$ where $1 < n_1 < n_2 < \dots$ are increasing powers of 2. To refer each polygon cap K_i more easily, we introduce the following notion.

Definition 6.3.1. For any $n \geq 1$, define the angle set $\Theta_n := \Theta_{\pi/2, n}$ with rotation angle $\pi/2$ so that $\Theta_n = \{(\pi/2)i/n : 1 \leq i < n\}$.

Definition 6.3.2. Say that $K \in \mathcal{K}$ is a *maximum polygon cap* with n steps of step size $\delta := (\pi/2)/n$, if $n > 1$ is a power of two, and K is a maximum polygon cap with the angle set Θ_n .

The main Theorem 3.4.9 of Section 3.5 is that each maximum polygon cap $K = K_i$ is *balanced* in the side lengths of K_i and its polygon niche. Using this balancedness condition, we will establish an upper bound of the surface area measure σ_K of a maximum polygon cap K with n steps of step size $\delta = (\pi/2)/n$ in Theorem 6.3.3. Then we will take the limit $n \rightarrow \infty$ and accordingly $\delta \rightarrow 0$, so that the polygon cap K converges to the balanced maximum cap K_∞ with its surface area measure σ_{K_∞} bounded from above.

Lemma 6.3.1. *Any maximum polygon cap K with n steps have the diameter at most 5. Consequently, the functions f_K^\pm and g_K^\pm are bounded from above by 5.*

Proof. Since $n > 1$ is a power of two by Definition 6.3.2, the angle set Θ_n of K contains the angle $\pi/4$. By Theorem 3.5.4, the cap K satisfies the condition (1) of Theorem 2.5.8. So the cap K also satisfies the condition (3) of Theorem 2.5.8, and the y -coordinate of $\mathbf{x}_K(\pi/4)$ is at most one. Now $K \subseteq H \cap Q_K^+(\pi/4)$ and the polygon $H \cap Q_K^+(\pi/4)$ have diameter at most $2 + 2\sqrt{2} < 5$. So K have the diameter at most 5.

By Definition 6.2.1, for any $t \in [0, \pi/2]$ we have $A_K^\pm(t) - C_K^\pm(t) = -f_K^\pm(t)v_t + g_K^\pm(t)u_t$ and since $A_K^\pm(t), C_K^\pm(t)$ are in K , the values $f_K^\pm(t)$ and $g_K^\pm(t)$ are also bounded by the diameter of K which is at most 5. \square

Definition 6.3.3. For any cap $K \in \mathcal{K}^c$ and angle $t \in [0, \pi/2]$, define

$$H_K^b(t) := H_+(t, h_K(t) - 1)$$

which is the closed half-plane with normal angle $t + \pi$ bounded from below by $b_K(t)$. Likewise, define

$$H_K^d(t) := H_+(t + \pi/2, h_K(t + \pi/2) - 1)$$

which is the closed half-plane with normal angle $t + 3\pi/2$ bounded from below by $d_K(t)$.

Lemma 6.3.2. *For any maximum polygon cap K with the angle set $\Theta := \Theta_n$ of step size $\delta := (\pi/2)/n$, and any angle $t \in \Theta$, we have the following calculations.*

1. $\mathcal{H}^1\left(\vec{b}_K(t) \cap H_K^d(t - \delta)\right) = \tan \delta \cdot \max(0, g_K^-(t) - 1 + \tan(\delta/2))$
2. $\mathcal{H}^1\left(\vec{b}_K(t) \cap H_K^d(t + \delta)\right) = \tan \delta \cdot \max(0, 1 - g_K^+(t) + \tan(\delta/2))$

Proof. (1) See the left side of Figure 6.1. Let p be the intersection (orange) of $d_K(t - \delta)$ and $b_K(t)$. Since $p \in b_K(t)$, there is a unique real value α such that $p + \alpha v_t = \mathbf{x}_K(t)$. As $\vec{b}_K(t)$ is the half-line from $\mathbf{x}_K(t)$ extending in the direction of $-v_t$, we have $\mathcal{H}^1\left(\vec{b}_K(t) \cap H_K^d(t - \delta)\right) = \max(\alpha, 0)$ and it remains to evaluate α . Define q as the intersection of $d_K(t + \delta)$ and $d_K(t)$, depicted green in the figure, and β as the unique real value such that $q + \beta u_t = \mathbf{x}_K(t)$. Since the points $p, q, \mathbf{x}_K(t)$ form a right-angled triangle of angle δ at q , we have $\alpha = \tan \delta \cdot \beta$. It remains to compute β .




Figure 6.1: The hallway $L_K(t)$ depicted in upright position with sides $c_K(t \pm \delta)$ and $d_K(t \pm \delta)$ for the proof of Lemma 6.3.2. The points p, q, r are colored as orange, green, blue respectively.

Let r be the intersection of $c_K(t - \delta)$ and $c_K(t)$, depicted blue in the figure. Since the lines $c_K(t)$ and $d_K(t)$ are parallel of distance one, and so are $c_K(t - \delta)$ and $d_K(t - \delta)$, and the two pairs of lines make an angle of δ , follow the dashed lines in the figure and we get $r = q + v_t + \tan(\delta/2)u_t$. As K have angle set Θ and the angles t and $t + \delta$ are adjacent, we have $r = C_K^-(t)$. By Proposition 6.2.1 we have $r + g_K^-(t)u_t = \mathbf{y}_K(t)$. Summing up, we now have

$$\begin{aligned} \beta &= (\mathbf{x}_K(t) - q) \cdot u_t = (\mathbf{x}_K(t) + v_t + \tan(\delta/2)u_t - r) \cdot u_t \\ &= (\mathbf{x}_K(t) + v_t + \tan(\delta/2)u_t - \mathbf{y}_K(t) + g_K^-(t)u_t) \cdot u_t \\ &= g_K^-(t) + \tan(\delta/2) - 1 \end{aligned}$$

and the result follows from $\mathcal{H}^1(\vec{b}_K(t) \cap H_K^d(t - \delta)) = \max(\alpha, 0) = \tan \delta \cdot \max(0, \beta)$.

(2) The proof is analogous to that of (1). See the right side of Figure 6.1. Let p be the intersection of $d_K(t + \delta)$ and $b_K(t)$. Since $p \in b_K(t)$ there is a unique real value α such that $p + \alpha v_t = \mathbf{x}_K(t)$. As $\vec{b}_K(t)$ is the half-line from $\mathbf{x}_K(t)$ extending in the direction of $-v_t$, we have $\mathcal{H}^1(\vec{b}_K(t) \cap H_K^d(t + \delta)) = \max(\alpha, 0)$ and it remains to evaluate α . Define q as the intersection of $d_K(t + \delta)$ and $d_K(t)$, and β as the unique real value such that $\mathbf{x}_K(t) + \beta u_t = q$. Since the points $p, q, \mathbf{x}_K(t)$ form a right-angled triangle of angle δ at q , we have $\alpha = \tan \delta \cdot \beta$. It remains to compute β .

Let r be the intersection of $c_K(t + \delta)$ and $c_K(t)$. Since the lines $c_K(t)$ and $d_K(t)$ are parallel of distance one, and so are $c_K(t + \delta)$ and $d_K(t + \delta)$, and the two pairs of lines make an angle of δ , we have $r = q + v_t - \tan(\delta/2)u_t$. As K have angle set Θ and the angles t and $t + \delta$ are adjacent, we have $r = C_K^+(t)$. By Proposition 6.2.1 we have $r + g_K^+(t)u_t = \mathbf{y}_K(t)$. Summing up, we now have

$$\begin{aligned} \beta &= (q - \mathbf{x}_K(t)) \cdot u_t = (-\mathbf{x}_K(t) - v_t + \tan(\delta/2)u_t + r) \cdot u_t \\ &= (-\mathbf{x}_K(t) - v_t + \tan(\delta/2)u_t + \mathbf{y}_K(t) - g_K^+(t)u_t) \cdot u_t \\ &= 1 - g_K^+(t) + \tan(\delta/2) \end{aligned}$$

and the result follows from $\mathcal{H}^1(\vec{b}_K(t) \cap H_K^d(t + \delta)) = \max(\alpha, 0) = \tan \delta \cdot \max(0, \beta)$. \square

We introduce the following auxiliary real functions. Note that m_0 is monotonically increasing, and is piecewise linear on the intervals $[0, 1]$, $[1, 2]$, and $[2, \infty)$ with values $m_0(0) = -1$, $m_0(1) = 1/2$, $m_0(x) = 1$ for $x \geq 2$.

Definition 6.3.4. Define $k_0, m_0 : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$ as $k_0(x) := \max(|x - 1|, (|x - 1| + 1)/2)$ and $m_0(x) = x - k_0(x)$.

We now bound the side lengths of a maximum polygon cap using balancedness. This is the most important part of the analysis.

Theorem 6.3.3. *Let $K \in \mathcal{K}_\Theta$ be any maximum polygon cap of n steps with step size $\delta = (\pi/2)/n$ and uniform angle set $\Theta := \Theta_n$. Take any $t \in \{0\} \cup \Theta$. We have the following upper bound of side length $\sigma_K(t)$.*

$$\sigma_K(t) \leq k_0(g_K^+(t)) \cdot \delta + O(\delta^2)$$

Proof. If $t = 0$, then as K have angle set Θ and rotation angle $\omega = \pi/2$, it is an intersection of a finite number of half-planes with normal angles $\neq 0 \in S^1$ (see Definition 3.2.3). So we have $\sigma_K(t) = 0$ and the bound holds trivially. Now assume $t \in \Theta$.

Write $s := \sigma_K(t)$. Because K is balanced by Theorem 3.4.9, we have $s = \tau_K(t)$. By (1) of Lemma 3.4.5, the value $\tau_K(t) = \mathcal{H}^1(X)$ is the length of the set $X := \vec{b}_K(t) \cap \partial \mathcal{N}_\Theta(K)$, the side(s) of the polygon niche $\mathcal{N}_\Theta(K)$ contributed by the half-line $\vec{b}_K(t)$ form $\mathbf{x}_K(t)$. We will bound $\mathcal{H}^1(X)$ from above, thus bounding s .

Define the set $U := \{t - \delta, t, t + \delta\}$ and $R := \bigcup_{u \in U} H_K^d(u)$ as the union of three half-planes. Divide X into $X \cap R$ and $X \setminus R$. We bound $\mathcal{H}^1(X \cap R)$ and $\mathcal{H}^1(X \setminus R)$ from above separately.

We first bound $\mathcal{H}^1(X \cap R)$ from above. As $X \subseteq \vec{b}_K(t)$, we have

$$\mathcal{H}^1(X \cap R) \leq \mathcal{H}^1(\vec{b}_K(t) \cap R) = \max(\vec{b}_K(t) \cap H_K^d(t - \delta), \vec{b}_K(t) \cap H_K^d(t + \delta))$$

and by computations in Lemma 6.3.2 and Lemma 6.3.1, we have

$$\mathcal{H}^1(X \cap R) \leq \delta \cdot \max(0, g_K^-(t) - 1, 1 - g_K^+(t)) + O(\delta^2).$$

Let $M := \max(0, g_K^-(t) - 1, 1 - g_K^+(t))$. We have $g_K^-(t) \leq g_K^+(t)$ by definition. So if $g_K^-(t) \leq 1$ then $M = \max(0, 1 - g_K^+(t)) \leq |1 - g_K^+(t)|$. If $g_K^-(t) > 1$ then we have $M = \max(0, g_K^-(t) - 1) \leq |1 - g_K^+(t)|$. Either way, we have $M \leq |1 - g_K^+(t)|$ and

$$\mathcal{H}^1(X \cap R) \leq |g_K^+(t) - 1| \cdot \delta + O(\delta^2). \quad (6.1)$$

Now we bound $\mathcal{H}^1(X \setminus R)$ from above. Define the closed region $S := \bigcap_{u \in U} H_K^b(u)$. Our intermediate goal is to show $\mathcal{H}^1(X \setminus R) \leq \mathcal{H}^1(b_K(t) \cap S)$. Define $H_0 := H_+(\pi/2, 0)$ as the closed half-plane bounded from below by the line $y = 0$. Let p_0 be the intersection of $b_K(t)$ and the line $y = 0$. Because $\mathcal{N}_\Theta(K)$ is open in the subspace topology of H_0 , we have $X \setminus \{p_0\} \subseteq H_0 \setminus \mathcal{N}_\Theta(K)$. For each $u \in U$, we have $Q_K^-(u) \cup H_K^b(u) \cup H_K^d(u) = \mathbb{R}^2$ so

$$\begin{aligned} X \setminus \{p_0\} &\subseteq H_0 \setminus \mathcal{N}_\Theta(K) \subseteq H_0 \setminus \bigcup_{u \in U} Q_K^-(u) \\ &= H_0 \cap \bigcap_{u \in U} (H_K^b(u) \cup H_K^d(u)) \\ &\subseteq R \cup S. \end{aligned}$$

So $X \setminus \{p_0\} \setminus R \subseteq S$ and we have $\mathcal{H}^1(X \setminus R) \leq \mathcal{H}^1(b_K(t) \cap S)$.

Define the intersections $B_+ := b_K(t) \cap b_K(t + \delta)$ and $B_- := b_K(t) \cap b_K(t - \delta)$. There is a unique real value β such that $B_+ = B_- - \beta v_t$. Elementary geometry shows that the side $b_K(t) \cap S$ of S is empty if $\beta < 0$, or is a finite segment of length β if $\beta \geq 0$. Another elementary geometry shows that $\beta = 2 \tan(\delta/2) - \sigma_K(t)$. So

$$\mathcal{H}^1(X \setminus R) \leq \max(0, 2 \tan(\delta/2) - \sigma_K(t)) = \max(0, \delta - s) + O(\delta^2). \quad (6.2)$$

Recall that $s = \nu_K(t) = \mathcal{H}^1(X)$. We divide the proof into cases on whether $s \leq \delta$ or not. If $s \leq \delta$, then adding Equation (6.1) and Equation (6.2) gives

$$s \leq |g_K^+(t) - 1| \cdot \delta + \delta - s + O(\delta^2)$$

so rearranging gives

$$s \leq (|g_K^+(t) - 1| + 1)/2 \cdot \delta + O(\delta^2) \leq k_0(g_K^+(t)) \cdot \delta + O(\delta^2).$$

On the other hand, if $s > \delta$, then adding Equation (6.1) and Equation (6.2) gives

$$s \leq |g_K^+(t) - 1| \cdot \delta + O(\delta^2) \leq k_0(g_K^+(t)) \cdot \delta + O(\delta^2).$$

Either way, the claimed inequality holds. \square

6.4 Inequality on Balanced Maximum Caps

In this Section 6.4, we prove the inequality $\sigma_K \leq k_0(g_K(t)) dt$ on the interval $t \in [0, \pi/2]$ for balanced maximum caps K in Theorem 6.4.3. This is done by taking the limit of Theorem 6.3.3 on maximum polygon caps. Using this, we prove the inequality $f'_K(t) \geq m_0(g_K(t))$ on the arm lengths of K in Theorem 6.5.1.

Lemma 6.4.1. *Let K be any maximum polygon cap with n steps of step size $\delta := (\pi/2)/n$ and angle set $\Theta := \Theta_n$. For any angle $t \in \{0\} \cup \Theta$ and open interval $I := (t, t + \delta)$, the followings are true.*

1. For any $t' \in I$, we have $g_K^+(t) \geq g_K^+(t') = g_K^-(t') \geq g_K^-(t + \delta)$.
2. Moreover, the gap $g_K^+(t) - g_K^-(t + \delta)$ between the upper and the lower bound is at most 5δ .

Proof. Since the angles t and $t + \delta$ are two adjacent angles of the finite set $\{0, \pi/2\} \cup \Theta$, we have

$$A_K^+(t) = A_K^-(t + \delta) = A_K^+(t') = A_K^-(t')$$

for all $t' \in I$. Call the common point A . Similarly, we have

$$C_K^+(t) = C_K^-(t + \delta) = C_K^+(t') = C_K^-(t')$$

for all $t' \in I$. Call the common point C . Take any $t' \in \bar{I} = [t, t + \delta]$. The lines $a_K(t')$ and $c_K(t')$ meet orthogonally at the point $\mathbf{y}_K(t')$. The line $a_K(t')$ passes through A , and the line $c_K(t')$ passes through C . So by elementary geometry, the trajectory of the point $\mathbf{y}_K(t')$ over all $t' \in \bar{I}$ forms an arc Δ of the circle Γ with the diameter of length ≤ 5 (Lemma 6.3.1) connecting A and C . As the lines $a_K(t)$ and $a_K(t + \delta)$ makes an angle of δ at point A , the arc Δ have central angle 2δ in Γ . The value $g_K^+(t)$ is the distance from C to $\mathbf{y}_K(t)$, and $g_K^-(t + \delta)$ is the distance from C to $\mathbf{y}_K(t + \delta)$. So (1) holds. The arc Δ connecting $\mathbf{y}_K(t)$ to $\mathbf{y}_K(t + \delta)$ have length at most 5δ . So (2) holds by triangle inequality. \square

Lemma 6.4.2. *Assume that a sequence of polygon caps K_n converge to a cap $K \in \mathcal{K}^c$ in Hausdorff distance as $n \rightarrow \infty$. Then we have*

$$\lim_{n \rightarrow \infty} \int_0^{\pi/2} |g_{K_n}^+(t) - g_K^+(t)| dt = 0.$$

Proof. The function $g_{K_n}^+$ is nonnegative and bounded from above by the diameter of K_n . The diameter of K_n converges to the diameter of K . So by dominated convergence theorem, it suffices to show that for almost every $t \in (0, \pi/2)$, we have $g_{K_n}^+(t) \rightarrow g_K^+(t)$ as $n \rightarrow \infty$.

Note that the set of all $t \in (0, \pi/2)$ such that either $\sigma_K(\{t\}) > 0$ or $\sigma_{K_n}(\{t\}) > 0$ for some n is countable; otherwise, it contradicts that the measures σ_{K_n} and σ_K are finite. So we can exclude such measure zero case and assume that $\sigma_K(\{t\}) = 0$ and $\sigma_{K_n}(\{t\}) = 0$ for all $n \geq 1$. By our choice of t , we have $g_{K_n}^-(t) = g_{K_n}^+(t)$ and $g_K^-(t) = g_K^+(t)$. We now show $g_{K_n}^+(t) \rightarrow g_K^+(t)$ as $n \rightarrow \infty$ for such t .

The function $s : S^1 \rightarrow \mathbb{R}$, defined as $s(u) := \sin(u - t)$ for $u \in (t, t + \pi/2]$ and zero otherwise, is upper semicontinuous. By Lemma 6.2.4, the value $g_{K_n}^+(t)$ (resp. $g_K^+(t)$) is the integral of s over σ_{K_n} (resp. σ_K). Because the measure σ_{K_n} converges weakly to σ_K as $n \rightarrow \infty$ (Theorem 4.1.3), by the Portmanteau theorem on finite measures we have $\limsup_n g_{K_n}^+(t) \leq g_K^+(t)$.

Similarly, define $s^- : S^1 \rightarrow \mathbb{R}$ as $s^-(u) := \sin(u - t)$ for $u \in (t, t + \pi/2)$ and zero otherwise. Then s^- is lower semicontinuous. By Lemma 6.2.4 and Proposition 2.1.2, the value $g_{K_n}^-(t)$ (resp. $g_K^-(t)$) is the integral of s^- over σ_{K_n} (resp. σ_K). Now by the Portmanteau theorem on finite measures we have $\liminf_n g_{K_n}^-(t) \geq g_K^-(t)$. Because $g_{K_n}^-(t) = g_{K_n}^+(t)$ and $g_K^-(t) = g_K^+(t)$, we get $g_{K_n}^+(t) \rightarrow g_K^+(t)$ as $n \rightarrow \infty$, completing the proof. \square

We now take the limit of Theorem 6.3.3 to show a corresponding inequality for the balanced maximum cup K .

Theorem 6.4.3. *Let K be any balanced maximum cap. Then on the interval $t \in [0, \pi/2)$, we have $\sigma_K \leq k_0(g_K^+(t)) dt$.*

Proof. Let μ_K be the measure $k_0(g_K^+(t)) dt$ on $t \in [0, \pi/2)$. It suffices to show

$$\sigma_K(I) \leq \mu_K(I) = \int_{t \in I} k_0(g_K^+(t)) dt \quad (6.3)$$

for any open interval $I = (a, b) \subseteq [0, \pi/2)$ and half-open interval $I = [0, b) \subseteq [0, \pi/2)$. Then for any set U open in the subspace topology of $[0, \pi/2)$ is a countable union of such intervals, so we have $\sigma_K(U) \leq \int_{t \in U} k_0(g_K^+(t)) dt$. Since $k_0(g_K^+(t)) dt$ is outer regular, by letting U to converge to arbitrary Borel subset of $[0, \pi/2)$ from above, we have $\sigma_K \leq k_0(g_K^+(t)) dt$ on $[0, \pi/2)$.

By Definition 3.5.2, K is the limit of maximum polygon caps K_n of n steps with step size $\delta = (\pi/2)/n$ as $n \rightarrow \infty$, where $n = n_1, n_2, \dots$ takes values in a strictly increasing powers of two. Let $\Theta := \bigcup_n \Theta_n$ be the union of angle sets Θ_n for powers of two $n = n_1, n_2, \dots$, so that Θ is the set of dyadic angles in $(0, \pi/2)$. As Θ is dense in $[0, \pi/2)$, it suffices to show Equation (6.3) for $I = (a, b)$ and $I = [0, b)$ for $a, b \in \Theta$ with $a < b$ by taking limits.

Take sufficiently large n so that we can assume $a, b \in \Theta_n$. For every $t \in \{0\} \cup \Theta_n$, we have

$$\sigma_{K_n}([t, t + \delta)) = \sigma_{K_n}(t) \leq k_0(g_{K_n}^+(t)) \cdot \delta + O(\delta^2) = \int_t^{t+\delta} k_0(g_{K_n}^+(u)) du + O(\delta^2) \quad (6.4)$$

by Theorem 6.3.3 and Lemma 6.4.1. For the open interval $I := (a, b)$, sum up Equation (6.4) for all $t \in [a, b) \cap \Theta_n$ to get

$$\begin{aligned} \sigma_{K_n}(I) &\leq \sum_{t \in [a, b) \cap \Theta_n} \sigma_{K_n}([t, t + \delta)) = \sum_{t \in [a, b) \cap \Theta_n} \sigma_{K_n}(\{t\}) \\ &\leq \left(\sum_{t \in [a, b) \cap \Theta_n} \int_t^{t+\delta} k_0(g_{K_n}^+(u)) du \right) + O_\delta(\delta) \\ &= \int_a^b k_0(g_{K_n}^+(u)) du + O_\delta(\delta) = \mu_{K_n}(I) + O(\delta). \end{aligned} \quad (6.5)$$

Here we use that the size of $[a, b) \cap \Theta_n$ is $\leq n = O(1/\delta)$.

We will now take $n \rightarrow \infty$ in Equation (6.5), first at the left-hand side. As $K_n \rightarrow K$ in Hausdorff distance, $\sigma_{K_n} \rightarrow \sigma_K$ in the weak convergence of measures (Theorem 4.1.3). Since I is open in S^1 , we have $\sigma_K(I) \leq \liminf_n \sigma_{K_n}(I)$. Now we take $n \rightarrow \infty$ in the right-hand side. By Lemma 6.4.2 and that k_0 is 1-Lipschitz, we have

$$\begin{aligned} \lim_n |\mu_{K_n}(I) - \mu_K(I)| &\leq \lim_n \int_{t \in I} |k_0(g_{K_n}^+(t)) - k_0(g_K^+(t))| dt \\ &\leq \lim_n \int_{t \in I} |g_{K_n}^+(t) - g_K^+(t)| = 0. \end{aligned} \quad (6.6)$$

So by taking $n \rightarrow \infty$ in Equation (6.5), we get $\sigma_K(I) \leq \mu_K(I)$ for $I = (a, b)$ as desired.

Similarly, for the interval $I := [0, b)$, set $a = 0$ and observe that Equation (6.5) still holds. Because K_n and K are caps, if we set $U = (-\pi/2, b)$ then we have $\sigma_{K_n}(I) = \sigma_{K_n}(U)$ and $\sigma_K(I) = \sigma_K(U)$. So we still have $\sigma_K(I) \leq \liminf_n \sigma_{K_n}(I)$. We also have $\lim_n \mu_{K_n}(I) = \mu_K(I)$ by following Equation (6.6). So by taking $n \rightarrow \infty$ in Equation (6.5), we have $\sigma_K(I) \leq \mu_K(I)$ for $I = [0, b)$ as desired. \square

Corollary 6.4.4. *Any balanced maximum cap K satisfies the condition (1) of Definition 6.1.2.*

Proof. For $t \in [0, \pi/2)$, use Theorem 6.4.3 and the Radon-Nikodym theorem to prove the condition (4) and thus (1). For $t \in (\pi/2, \pi]$, use a mirror-symmetric argument (Proposition 3.5.1). \square

Note that the condition (1) of Definition 6.1.2 implies that $A_K^+(t) = A_K^-(t)$ and $f_K^+(t) = f_K^-(t)$ for all $t \in [0, \pi/2)$. In the rest of this paper and the upcoming work, we will denote the common value as $A_K(t)$ and $f_K(t)$ respectively.

Proposition 6.4.5. *Assume that a cap $K \in \mathcal{K}^c$ satisfies the condition (1) of Definition 6.1.2. Then the followings are true.*

1. For any $t \in [0, \pi/2)$, we have $A_K^+(t) = A_K^-(t)$ and $f_K^+(t) = f_K^-(t)$.
2. For any $t \in (0, \pi/2]$, we have $C_K^+(t) = C_K^-(t)$ and $g_K^+(t) = g_K^-(t)$.

Proof. We have $\sigma_K(\{t\}) = 0$ for any $t \in [0, \pi] \setminus \{\pi/2\}$. Now use Proposition 2.1.2. \square

Definition 6.4.1. For any cap $K \in \mathcal{K}^c$ satisfying (1) of Definition 6.1.2, define $A_K, C_K : [0, \pi/2] \rightarrow \mathbb{R}^2$ and $f_K, g_K : [0, \pi/2] \rightarrow \mathbb{R}^2$ as below.

- For any $t \in [0, \pi/2)$, denote the common values in (1) of Proposition 6.4.5 as $A_K(t) := A_K^\pm(t)$ and $f_K(t) = f_K^\pm(t)$.
- Also, define $A_K(\pi/2) := A_K^-(\pi/2)$ and $f_K(\pi/2) := f_K^-(\pi/2)$.
- For any $t \in (0, \pi/2]$, denote the common values in (2) of Proposition 6.4.5 as $C_K(t) := C_K^\pm(t)$ and $g_K(t) = g_K^\pm(t)$.
- Also, define $C_K(0) := C_K^+(0)$ and $g_K(0) := g_K^+(0)$.

Proposition 6.4.6. *For any cap $K \in \mathcal{K}^c$ satisfying the condition (1) of Definition 6.1.2, the followings are true.*

1. *The functions A_K, C_K, f_K, g_K in Definition 6.4.1 are continuous on $[0, \pi/2]$.*
2. *The corners $\mathbf{x}_K, \mathbf{y}_K : [0, \pi/2] \rightarrow \mathbb{R}^2$ are continuously differentiable, and we have $\mathbf{x}'_K(t) = -(f_K(t) - 1)u_t + (g_K(t) - 1)v_t$ and $\mathbf{y}'_K(t) = -f_K(t)u_t + g_K(t)v_t$.*

Proof. (1) is by Theorem 2.1.3. (2) is by Theorem 6.2.3. □

Now it makes sense to use the notations $f_K(t) = f_K^\pm(t)$, $g_K(t) = g_K^\pm(t)$, r_K and s_K for balanced maximum caps K by Corollary 6.4.4 and Definition 6.4.1.

6.5 Bounding Arm Lengths

We state Theorem 6.4.3 purely in terms of arm lengths of K , following Section 1.7.3.

Theorem 6.5.1. *Let K be a balanced maximum cap. Then $f_K : [0, \pi/2] \rightarrow \mathbb{R}$ is absolutely continuous, and we have $f'_K(t) \geq m_0(g_K(t))$ for almost every $t \in [0, \pi/2]$ except measure zero set.*

Proof. By Corollary 6.4.4, we have $\sigma_K = r_K(t)dt$ on the domain $t \in [0, \pi/2)$ for some measurable function $r_K : [0, \pi/2) \rightarrow \mathbb{R}_{\geq 0}$ that is unique up to null set. In particular, by Theorem 6.4.3 we can assume $r_K(t) \leq k_0(g_K(t))$ on $t \in [0, \pi/2)$. Now by Theorem 6.2.5 we have $df_K(t) = (g_K(t) - r_K(t))dt$ on $t \in [0, \pi/2]$ (check the case $t = 0$ separately). So by Proposition 5.1.4, the function $f_K(t)$ is absolutely continuous and its derivative is $f'_K(t) = g_K(t) - r_K(t)$ for almost every $t \in [0, \pi/2]$. Now check

$$g_K(t) - r_K(t) \geq g_K(t) - k_0(g_K(t)) = m_0(g_K(t)).$$

to conclude $f'_K(t) \geq m_0(g_K(t))$. □

Using Theorem 6.5.1, we will iteratively obtain better lower bounds of f_K and g_K for balanced maximum caps K , and show the lower bound $f_K(t), g_K(t) > 1$ in Theorem 6.5.6. Then we use the equation $\mathbf{x}'_K(t) = -(f_K(t) - 1)u_t + (g_K(t) - 1)v_t$ of Proposition 6.4.6 to prove the main Theorem 6.1.1.

Definition 6.5.1. Define the operator \mathcal{F} mapping any continuous function $f : [0, \pi/2] \rightarrow \mathbb{R}$, to another continuous function $\mathcal{F}f : [0, \pi/2] \rightarrow \mathbb{R}$ as

$$\mathcal{F}f(x) = 1 + \int_0^x m_0(f(\pi/2 - u)) du.$$

Definition 6.5.2. Define the continuous functions $f_n : [0, \pi/2] \rightarrow \mathbb{R}$ for all integers $n \geq 0$ as the following.

1. $f_0(x) := 0$ for all $x \in [0, \pi/2]$.
2. $f_{n+1}(x) := \max(f_n(x), \mathcal{F}f_n(x))$ for all $x \in [0, \pi/2]$.

Lemma 6.5.2. *The following holds for any $n \geq 0$. Let $K \in \mathcal{K}$ be any balanced maximum cap. Then we have $f_K(t) \geq f_n(t)$ for all $t \in [0, \pi/2)$ and $g_K(t) \geq f_n(\pi/2 - t)$ for all $t \in (0, \pi/2]$.*

Proof. Induct on n . The base case $n = 0$ holds trivially. Now assume the inductive hypothesis that for any balanced maximum cap K' , we have $f_{K'}(t) \geq f_n(t)$ for all $t \in [0, \pi/2)$ and $g_{K'}(t) \geq f_n(\pi/2 - t)$ for all $t \in (0, \pi/2]$.

Fix an arbitrary balanced maximum cap K . For any $t \in [0, \pi/2)$, we have

$$\begin{aligned} f_K(t) &= f_K(0) + \int_0^t f'_K(u) du \\ &\geq f_K(0) + \int_0^t m_0(g_K(u)) du \\ &\geq f_K(0) + \int_0^t m_0(f_K(\pi/2 - u)) du \end{aligned}$$

by the inductive hypothesis on $K' := K$ and Theorem 6.5.1. This combined with the inductive hypothesis $f_K(t) \geq f_n(t)$ on $K' := K$ implies that $f_K(t) \geq f_{n+1}(t)$.

Next, apply the inductive hypothesis on the mirror image $K' := K^m$. Then we also have $f_{K^m}(t) \geq f_{n+1}(t)$ for all $t \in [0, \pi/2)$. This, by mirror symmetry, is equivalent to the other inequality $g_K(t) \geq f_{n+1}(\pi/2 - t)$ for all $t \in (0, \pi/2]$. \square

Definition 6.5.3. For any constant $c \in [0, 1]$, define the function $j_c : [0, \pi/2] \rightarrow \mathbb{R}$ as $j_c(x) := \max(1 - x, c)$.

Lemma 6.5.3. *Define the constant $d_0 := 1/12$. Then for any $c \in [0, 2/3]$, we have $\mathcal{F}j_c(x) \geq j_{c+d_0}(x)$ on $x \in [0, \pi/2]$.*

Proof. Because m_0 is bounded from below by -1 , it is always guaranteed that $\mathcal{F}j_c(x) \geq 1 - x$. So to show that $\mathcal{F}j_c(x) \geq j_{c+d_0}(x)$ on $x \in [0, 1]$, it suffices to show $\mathcal{F}j_c(x) \geq c + d_0$.

We show this by simply computing $\mathcal{F}j_c(x)$. Since the range of j_c is in $[0, 1]$, we have

$$\begin{aligned} \mathcal{F}j_c(x) &= 1 + \int_0^x m_0(j_c(\pi/2 - u)) du \\ &= 1 - x + \int_0^x \frac{3}{2} j_c(\pi/2 - u) du \end{aligned} \tag{6.7}$$

and it suffices to show that the value is $\geq c + d_0$.

First assume the case $x \leq \pi/2 - 1 + c$. Then $\pi/2 - u \geq \pi/2 - x \geq 1 - c$ in the integral of Equation (6.7) so

$$\begin{aligned} \mathcal{F}j_c(x) &= 1 - x + \frac{3}{2}cx = 1 - (1 - 3/2 \cdot c)x \\ &\geq 1 - (1 - 3/2 \cdot c)(\pi/2 - 1 + c) \\ &> c + d_0. \end{aligned}$$

In the last inequality, the difference is minimized at $c = 7/6 - \pi/4$ with the value $-1/8 + 3\pi/8 - 3\pi^2/32 > 0$.

It remains to check the case $x > \pi/2 - 1 + c$. Then Equation (6.7) evaluates to

$$\begin{aligned} \mathcal{F}j_c(x) &= 1 - x + \frac{3}{2}c(\pi/2 - 1 + c) + \int_{\pi/2 - 1 + c}^x \frac{3}{2}(1 - \pi/2 + u) du \\ &= 1 - x + \frac{3}{2}c(\pi/2 - 1 + c) + \frac{3}{4}((x - \pi/2 + 1)^2 - c^2). \end{aligned}$$

Minimize the quadratic polynomial over $x \in \mathbb{R}$, then we have

$$\mathcal{F}j_c(x) \geq \frac{5}{3} - \frac{\pi}{2} + \frac{3}{4}c(c + \pi - 2)$$

where the equality holds at $x = \pi/2 - 1/3$. We also have

$$\frac{5}{3} - \frac{\pi}{2} + \frac{3}{4}c(c + \pi - 2) > c + \frac{1}{12}$$

because the difference of both sides is minimized at $c = 5/3 - \pi/2$ with the value $-1/2 + 3\pi/4 - 3\pi^2/16 > 0$. This completes the proof. \square

Lemma 6.5.4. *For any functions $f, g : [0, \pi/2] \rightarrow \mathbb{R}_{\geq 0}$ such that $f \leq g$ on $[0, \pi/2]$, we also have $\mathcal{F}f \leq \mathcal{F}g$.*

Proof. The function m_0 is monotonically increasing; now check the Definition 6.5.1 of \mathcal{F} . \square

Lemma 6.5.5. *We have $f_{11}(x) > 1$ on $x \in (0, 1]$.*

Proof. For any integer $1 \leq m \leq 10$, we will show that $f_m \geq j_{(m-1)/12}$. The base case $f_1 = j_0$ is done by simple computation. Assuming the inductive hypothesis $f_m \geq j_{(m-1)/12}$ for $m \leq 9$, the inductive step

$$f_{m+1} \geq \mathcal{F}f_m \geq \mathcal{F}j_{(m-1)/12} \geq j_{m/12}$$

can be done using Lemma 6.5.4 and Lemma 6.5.3. So $f_{10} \geq j_{9/12} > 2/3$ in particular.

Because $m_0(y) > 0$ for all $y > 2/3$, we now have

$$f_{11}(x) \geq \mathcal{F}f_{10}(x) = 1 + \int_0^x m_0(f_{10}(\pi/2 - u)) du > 1$$

for all $x \in (0, \pi/2]$. \square

Theorem 6.5.6. *For any balanced maximum cap K , we have $f_K(t) > 1$ on $t \in (0, \pi/2]$ and $g_K(t) > 1$ on $t \in [0, \pi/2]$.*

Proof. By Lemma 6.5.2 and Lemma 6.5.5. \square

Now we prove the injectivity condition for any balanced maximum cap.

Proof of Theorem 6.1.1. We check conditions (1)-(3) of Definition 6.1.2 for balanced maximum caps K . Condition (1) holds by Corollary 6.4.4. Condition (2) holds by (2) of Proposition 6.4.6. Finally, condition (3) holds because

$$\mathbf{x}'_K(t) = -(f_K(t) - 1)u_t + (g_K(t) - 1)v_t$$

by (2) of Proposition 6.4.6 and Theorem 6.5.6. \square

Chapter 7

Convex Domain and Convex Curves

This section prepares a minimal amount of technology needed for the next Chapter 8.

- Section 7.1 defines the general notion of *convex domain* \mathcal{V} and linear/quadratic functionals on \mathcal{V} . We show that if the directional derivative of a concave quadratic functional f is non-negative at $V \in f$, then f attains the global maximum value at V .
- Section 7.2 reviews the notion of a *Jordan curve* \mathbf{x} that encloses a region R of the plane, then defines the *curve area functional* $\mathcal{J}(\mathbf{x})$ that measures the area of R using Green's theorem. This will work for any continuous $\mathbf{x} : [a, b] \rightarrow \mathbb{R}^2$ of bounded variation.
- Section 7.3 defines a *convex curve* segment $\mathbf{u}_K^{a,b}$ of the boundary of a planar convex body K , and calculate its curve area functional.
- Section 7.4 establishes the *Mamikon's theorem* on a general convex body K that may have a non-differentiable boundary.

7.1 Convex Domain

Definition 7.1.1. A *convex domain* \mathcal{V} is a space with the barycentric operation $c_\lambda : \mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}$ for all $\lambda \in [0, 1]$, such that there is an embedding $e : \mathcal{V} \rightarrow V$ to a convex subspace of a vector space V preserving c_λ . That is, e is injective and $e(c_\lambda(v_1, v_2)) = (1 - \lambda)e(v_1) + \lambda e(v_2)$.

Remark 7.1.1. Although we will not use this notion in our work, Definition 7.1.1 is equivalent to saying that (\mathcal{V}, c_λ) is a *cancellative convex space* (Theorem 2 of [Sto49]).

Definition 7.1.2. Call a function $f : \mathcal{V}_1 \rightarrow \mathcal{V}_2$ between convex domains *convex-linear* if it preserves the barycentric operation c_λ .

The composition $f \circ g$ of two convex-linear maps f, g would also be convex-linear, as it preserves c_λ .

Definition 7.1.3. Call a function $g : \mathcal{V}_1 \times \mathcal{V}_2 \rightarrow \mathcal{V}_3$ *convex-bilinear* if the maps $v \mapsto g(v_1, v)$ and $v \mapsto g(v, v_2)$ are convex-linear for any fixed $v_1 \in \mathcal{V}_1$ and $v_2 \in \mathcal{V}_2$.

Definition 7.1.4. Call $h : \mathcal{V} \rightarrow \mathbb{R}$ a *quadratic functional* on a convex domain \mathcal{V} if $h(K) = g(K, K)$ for some convex-bilinear $g : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{R}$.

Theorem 7.1.1. *The space \mathcal{K} of all planar convex bodies is a convex domain with the barycentric operation*

$$\begin{aligned} c_\lambda(K_1, K_2) &:= (1 - \lambda)K_1 + \lambda K_2 \\ &= \{(1 - \lambda)p_1 + \lambda p_2 : p_1 \in K_1, p_2 \in K_2\} \end{aligned}$$

given by the Minkowski sum of convex bodies.

Proof. By Remark 1.7.7 of [Sch13], the map $K \mapsto h_K$ embeds the space \mathcal{K} to a convex cone of the space of all continuous functions from S^1 to \mathbb{R} , preserving the barycentric operations of each space. \square

The values on convex body $K \in \mathcal{K}$ appearing in Section 2.1 are convex-linear in K .

Theorem 7.1.2. *The following values are convex-linear in $K \in \mathcal{K}$.*

1. Support function h_K
2. For fixed constants $a, b \in \mathbb{R}$ so that $a < b < a + \pi$, the vertices $v_K^\pm(a)$ and $v_K(a, b)$.
3. Surface area measure σ_K

Proof. (1) follows from Theorem 1.7.5 (a) of [Sch13] that $h_{K_1+K_2} = h_{K_1} + h_{K_2}$, and that $h_{aK} = ah_K$ for any $a \geq 0$ which is easy to check. For (2), note that $p := v_K(a, b)$ is the unique solution satisfying the equations $p \cdot u_a = h_K(a)$ and $p \cdot u_b = h_K(b)$. Let U be the constant 2×2 matrix with column vectors u_a and u_b . Then we have $p = U^{-1}[h_K(a), h_K(b)]^T$ convex-linear in K by (1). Use Theorem 2.1.3 to see that $v_K^\pm(a)$, the limit of $v_K(a, b)$ as $b \rightarrow a$ in either direction, is linear too. (3) comes from (2) and $\sigma_K = v_t \cdot dv_K^+(t)$ which follows from Theorem 5.2.2. \square

The area $|K|$ of $K \in \mathcal{K}$ is a quadratic functional.

Theorem 7.1.3. (Remark 5.1.2, page 276 of [Sch13]) *For any $K \in \mathcal{K}$, we have*

$$|K| = \frac{1}{2} \int_{t \in S^1} h_K(t) \sigma_K(dt)$$

which is quadratic in K by Theorem 7.1.2.

Definition 7.1.5. For any quadratic functional $f : \mathcal{V} \rightarrow \mathbb{R}$ on a convex space \mathcal{V} and $K, K' \in \mathcal{V}$, define the *directional derivative*

$$Df(K; K') := \left. \frac{d}{d\lambda} \right|_{\lambda=0} f(c_\lambda(K, K'))$$

of f at K in the direction towards K' .

For any quadratic functional f and a fixed $K \in \mathcal{V}$, the value $Df(K; K')$ is well-defined and always a linear functional of K' .

Lemma 7.1.4. *Let f be a quadratic functional on a convex domain \mathcal{V} , so that $f(K) = h(K, K)$ for a convex-bilinear map $h : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{R}$. Then we have the following for any $K, K' \in \mathcal{V}$.*

$$Df(K; K') = h(K, K') + h(K', K) - 2h(K, K)$$

So the map $Df(K; -) : \mathcal{V} \rightarrow \mathbb{R}$ is always well-defined and a linear functional.

Proof. We have

$$\begin{aligned} f(c_\lambda(K, K')) &= h(c_\lambda(K, K'), c_\lambda(K, K')) \\ &= (1 - \lambda)^2 h(K, K) + \lambda(1 - \lambda)(h(K, K') + h(K', K)) + \lambda^2 h(K', K') \end{aligned} \quad (7.1)$$

by bilinearity of h . Take the derivative at $\lambda = 0$. □

Definition 7.1.6. A functional $f : \mathcal{V} \rightarrow \mathbb{R}$ on a convex domain \mathcal{V} is *concave* (resp. *convex*) if $f(c_\lambda(K_1, K_2)) \geq (1 - \lambda)f(K_1) + \lambda f(K_2)$ (resp. $f(c_\lambda(K_1, K_2)) \leq (1 - \lambda)f(K_1) + \lambda f(K_2)$) for all $K_1, K_2 \in \mathcal{V}$ and $\lambda \in [0, 1]$.

To prove that K maximizes a concave quadratic functional $f(K)$ on \mathcal{V} , we only need to prove that $Df(K; -)$ is a nonpositive linear functional on \mathcal{V} .

Theorem 7.1.5. *For any concave quadratic functional f on a convex domain \mathcal{V} , the value $K \in \mathcal{V}$ maximizes $f(K)$ if and only if the convex-linear functional $Df(K; -)$ is nonpositive.*

Proof. Assume that K is the maximizer of $f(K)$. Then for any $K' \in \mathcal{V}$, the value $f(c_\lambda(K, K'))$ over all $\lambda \in [0, 1]$ is maximized at $\lambda = 0$. So taking the derivative at $\lambda = 0$, we should have $Df(K; K') \leq 0$.

Now assume on the other hand that $K \in \mathcal{V}$ is chosen such that $Df(K; -)$ is always nonpositive. Fix an arbitrary $K' \in \mathcal{V}$. Observe that $f(c_\lambda(K, K'))$ is a polynomial $p(\lambda)$ of $\lambda \in [0, 1]$ by Equation (7.1). Because f is concave, the polynomial $p(\lambda)$ is also concave with respect to λ and the quadratic coefficient of $p(\lambda)$ is nonpositive. The linear coefficient of $p(\lambda)$ is $Df(K; K')$ and this is nonpositive as well. So $p(\lambda)$ is monotonically decreasing with respect to λ and we have $f(K) \geq f(K')$ as desired. □

Definition 7.1.7. For any real-valued functionals $f(V)$ and $g(V)$ on the convex domain $V \in \mathcal{V}$, write $f(V) \equiv_V g(V)$ if and only if $f(V) - g(V)$ is convex-linear in $V \in \mathcal{V}$.

It is easy to check that the relation \equiv_V in Definition 7.1.7 is an equivalence relation on the real-valued functionals on convex domain \mathcal{V} .

Lemma 7.1.6. *Let $h : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{R}$ be a convex-bilinear map on a convex domain \mathcal{V} . Fix constants $c_1, c_2 \in \mathcal{V}$. Define quadratic functionals $f(K) = h(K, K)$ and $g(K) = h(K + c_1, K + c_2)$, then we have $f(K) \equiv_K g(K)$.*

Proof. We have $g(K) - f(K) = h(c_1, K) + h(K, c_2) + h(c_1, c_2)$ which is linear in K . □

7.2 Curve Area Functional

We first recall the notion of Jordan arc and curve following Chapter 8 of [Apo]. See the reference for details.

Definition 7.2.1. A *Jordan arc* Γ is the image of a continuous and injective function $\mathbf{x} : [a, b] \rightarrow \mathbb{R}^2$. We call \mathbf{x} a *parametrization* of Γ . A *Jordan curve* Γ is the image of a continuous and injective function $\mathbf{x} : S^1 \rightarrow \mathbb{R}^2$.

Equivalently, a Jordan curve Γ is the image of a continuous function $\mathbf{x} : [a, b] \rightarrow \mathbb{R}^2$ with $a < b$ which is injective on $[a, b)$ and $\mathbf{x}(a) = \mathbf{x}(b)$. We allow $a = b$ for Jordan arcs, but we require $a \neq b$ for Jordan curves in order for the following famous theorem to hold.

Theorem 7.2.1. (*Jordan curve theorem*) *The complement $\mathbb{R}^2 \setminus \Gamma$ of a Jordan curve Γ is a disjoint union of two connected components U and V where U is bounded and V is unbounded. Moreover, U and V are open and $\partial U = \partial V = \Gamma$. Say that U is the region enclosed by Γ .*

We give orientations to Jordan arcs and curves.

Definition 7.2.2. Let Γ be any Jordan arc. Take any parametrization $\mathbf{x} : [a, b] \rightarrow \mathbb{R}^2$ of Γ and call the points $\mathbf{x}(a)$ and $\mathbf{x}(b)$ the *endpoints* of Γ . Note that the endpoints of Γ are independent of the choice of \mathbf{x} . An *oriented Jordan arc* $\vec{\Gamma}$ is a Jordan arc Γ with one endpoint p marked as the *starting point* and the other endpoint q marked as the *ending point*. A *parametrization* $\mathbf{x} : [a, b] \rightarrow \mathbb{R}^2$ of an oriented Jordan arc $\vec{\Gamma}$ is a parametrization of Γ that respects the order $\mathbf{x}(a) = p$ and $\mathbf{x}(b) = q$ of the endpoints of $\vec{\Gamma}$.

Definition 7.2.3. Let Γ be any Jordan curve. An *oriented Jordan curve* $\vec{\Gamma}$ is the unoriented curve Γ with a *clockwise* or *counterclockwise* direction assigned. A *parametrization* $\mathbf{x} : S^1 \rightarrow \mathbb{R}^2$ of an oriented Jordan curve $\vec{\Gamma}$ is a parametrization of Γ such that, for any point p inside the region enclosed by Γ , the parametrization \mathbf{x} rotates around p in the specified direction.

We now define the *curve area functional* $\mathcal{J}(\mathbf{x})$ of $\mathbf{x} : [a, b] \rightarrow \mathbb{R}^2$, so that if \mathbf{x} parametrizes a counterclockwise Jordan curve, then $\mathcal{J}(\mathbf{x})$ is the area of the region enclosed by \mathbf{x} .

Definition 7.2.4. For two vectors $p = (a, b)$ and $q = (c, d)$ in \mathbb{R}^2 , define their cross product $p \times q := ad - bc \in \mathbb{R}$. For a pair $p = (p_1, p_2)$ of bounded measurable functions and a pair $\mu = (\mu_1, \mu_2)$ of finite signed measures on a set X , define the finite signed measure $p \times \mu := p_1 \mu_2 - p_2 \mu_1$ on X .

Definition 7.2.5. Let $C^{\text{BV}}[a, b]$ be the real vector space of all continuous maps of bounded variation from $[a, b]$ to \mathbb{R}^2 .

Definition 7.2.6. For any $\mathbf{x} \in C^{\text{BV}}[a, b]$, define its *curve area functional* $\mathcal{J}(\mathbf{x})$ as

$$\mathcal{J}(\mathbf{x}) := \frac{1}{2} \int_a^b \mathbf{x}(t) \times d\mathbf{x}(t).$$

Proposition 7.2.2. $\mathcal{J}(\mathbf{x})$ is quadratic on $\mathbf{x} \in C^{\text{BV}}[a, b]$.

Proof. $\mathcal{J}(\mathbf{x}) = \mathcal{B}(\mathbf{x}, \mathbf{x})$ where $\mathcal{B}(\mathbf{x}_1, \mathbf{x}_2) := \frac{1}{2} \int_a^b \mathbf{x}_1(t) \times d\mathbf{x}_2(t)$ is bilinear in $\mathbf{x}_1, \mathbf{x}_2 \in C^{\text{BV}}[a, b]$. \square

Note that the integral in Definition 7.2.6 is on the Lebesgue–Stieltjes measure $d\mathbf{x}(t)$ of \mathbf{x} taken the cross product with $\mathbf{x}(t)$ as in Definition 7.2.4. Writing the coordinates of $\mathbf{x}(t) = (x(t), y(t))$, we can write $d\mathbf{x} = (dx, dy)$ and $\mathcal{J}(\mathbf{x})$ more explicitly as

$$\mathcal{J}(\mathbf{x}) = \frac{1}{2} \int_a^b R_{\pi/2}(\mathbf{x}(t)) \cdot d\mathbf{x}(t) = \frac{1}{2} \int_a^b x(t) dy(t) - y(t) dx(t) \quad (7.2)$$

instead, where $R_{\pi/2}(x, y) = (-y, x)$ is the rotation of \mathbb{R}^2 along the origin by $\pi/2$.

Theorem 7.2.3. *If \mathbf{x} is a rectifiable parametrization of a Jordan curve oriented counterclockwise, then $\mathcal{J}(\mathbf{x})$ is the area of the region enclosed by \mathbf{x} .*

Proof. Apply Green's theorem (Theorem 10.43, page 289 of [Apo]) on the curve \mathbf{x} and vector field $(P, Q) = (-y, x)$. \square

Remark 7.2.1. If \mathbf{x} is not closed (that is, $\mathbf{x}(a) \neq \mathbf{x}(b)$), the sofa area functional $\mathcal{J}(\mathbf{x})$ measures the signed area of the region bounded by the curve \mathbf{x} , and two line segments connecting the origin to $\mathbf{x}(a)$ and $\mathbf{x}(b)$ respectively.

Fix an oriented Jordan arc Γ and take any parametrization \mathbf{x} of Γ . Then the value $\mathcal{J}(\mathbf{x})$ is a line integral on Γ (Equation (7.2)), so the value of $\mathcal{J}(\mathbf{x})$ is independent of the choice of \mathbf{x} . Similarly, any parametrization of an oriented Jordan curve Γ are circular shifts of each other, so the value $\mathcal{J}(\mathbf{x})$ is fixed.

Definition 7.2.7. For any oriented Jordan arc or curve Γ , the value of $\mathcal{J}(\mathbf{x})$ is independent of the choice of parametrization \mathbf{x} of Γ ; call this value $\mathcal{J}(\Gamma)$.

In particular, the curve area functional of the oriented line segment from p to q is the following.

Definition 7.2.8. For any two points $p, q \in \mathbb{R}^2$, define $\mathcal{J}(p, q) := (p \times q)/2$.

Proposition 7.2.4. *The curve area functional of the oriented line segment from point p to q is $\mathcal{J}(p, q)$. Moreover, if there is some $t \in S^1$ and $h \in \mathbb{R}$ so that $p, q \in l(t, h)$ and $q - p = dv_t$ for some $d \in \mathbb{R}$, then $\mathcal{J}(p, q) = hd/2$.*

Proof. Parametrize the line segment from p to q as $\mathbf{x}(t) = p + (q - p)t$ for $t \in [0, 1]$. Then $\mathcal{J}(\mathbf{x})$ evaluates to $(p \times q)/2$ which is $\mathcal{J}(p, q)$. Moreover,

$$p \times q = p \times (q - p) = p \times (dv_t) = s(p \cdot u_t) = hd$$

so $\mathcal{J}(p, q) = hd/2$. \square

Proposition 7.2.5. *If two points $p, q \in \mathbb{R}^2$ and the origin $O = (0, 0)$ are on a common line, then $\mathcal{J}(p, q) = 0$.*

Proof. Set $h = 0$ in Proposition 7.2.4. \square

We will often form a Jordan curve by concatenating multiple Jordan arcs.

Definition 7.2.9. Say that an oriented Jordan arc or curve Γ is the *concatenation* of the Jordan arcs $\Gamma_1, \Gamma_2, \dots, \Gamma_n$ in order, if the ending point of Γ_i matches with the starting point of Γ_{i+1} for all $1 \leq i \leq n - 1$, and Γ is obtained by following $\Gamma_1, \Gamma_2, \dots$ in order.

Proposition 7.2.6. *If Γ is the concatenation of the Jordan arcs $\Gamma_1, \Gamma_2, \dots, \Gamma_n$ in order, then $\mathcal{J}(\Gamma) = \sum_{i=1}^n \mathcal{J}(\Gamma_i)$.*

Proof. Take the parametrization \mathbf{x} of Γ , and integrate Definition 7.2.6 into \square

Proposition 7.2.7. *Let Γ be a Jordan curve which is the concatenation of the Jordan arcs $\Gamma_1, \Gamma_2, \dots, \Gamma_n$ in order. Assume that there is a half-plane H' containing Γ with the boundary l' and normal angle $t \in S^1$ (and thus the normal vector u_t). Assume that some arc Γ_i of Γ is an oriented line segment s of length > 0 on l' in the positive direction of v_t . Then Γ is oriented counterclockwise.*

Proof. Say that Γ_i is the line segment from p to q in l' . Fix the endpoints p and q , and deform Γ_i slightly towards outside H' so that $\Gamma_i \setminus \{p, q\}$ is strictly outside H' . Now take the point $r := (p+q)/2$. The segment $\Gamma \setminus \Gamma_i \cup \{p, q\}$ from q to p is inside H' , so it rotates around the point r in the counterclockwise angle of π . Similarly, the deformed curve Γ_i also rotates around the point r in the counterclockwise angle of π . So the total angle is 2π clockwise, and Γ is oriented counterclockwise. \square

7.3 Convex Curve

Define the following convex curve segment of the boundary of a convex body K .

Definition 7.3.1. For any planar convex body K and $a, b \in \mathbb{R}$ be arbitrary such that $a < b < a + \pi$, define the segment

$$\mathbf{u}_K^{a,b} := \{v_K^+(a)\} \cup \bigcup_{t \in (a,b)} e_K(t) \cup \{v_K^-(b)\}.$$

of the boundary of K .

The goal of this Section 7.3 is to show that $\mathbf{u}_K^{a,b}$ is a rectifiable curve and evaluate its curve area functional.

Lemma 7.3.1. *Assume arbitrary $K \in \mathcal{K}$ and $a, b \in \mathbb{R}$ such that $a < b < a + \pi$. If $v_K^+(a) = v_K^-(b)$, then $v_K(a, b) = v_K^+(a) = v_K^-(b)$ and $\mathbf{u}_K^{a,b}$ is the single point $\{v_K^+(a)\}$. If $v_K^+(a) \neq v_K^-(b)$, then the followings are true.*

1. *The point $v_K(a, b)$ is not on the line l' connecting $v_K^+(a)$ and $v_K^-(b)$.*
2. *The closed half-plane H' with the boundary l' containing $v_K(a, b)$ have normal angle $t' + \pi$ for some $t' \in (a, b)$.*
3. *The intersection $K' := K \cap H'$ is a planar convex body satisfying the followings.*
 - i. *For any $t \in (t' - \pi, a]$, we have $e_{K'}(t) = \{v_K^+(a)\}$.*
 - ii. *For any $t \in (a, b)$, we have $e_{K'}(t) = e_K(t)$.*
 - iii. *For any $t \in [b, t' + \pi)$, we have $e_{K'}(t) = \{v_K^-(b)\}$.*
 - iv. *The edge $e_{K'}(t' + \pi)$ is the line segment from $v_K^-(b)$ to $v_K^+(a)$.*

Proof. Define the closed cone $X := H_K(a) \cap H_K(b)$ with vertex $v_K(a, b)$ containing K . The boundary of X is the union of two half-lines $X \cap l_K(a)$ and $X \cap l_K(b)$, each containing $v_K^+(a)$ and $v_K^-(b)$ respectively, and both meeting at $v_K(a, b)$ with an angle of $b - a \in (0, \pi)$. So we have $v_K(a, b) = v_K^+(a) + \alpha v_a$ for some $\alpha \geq 0$ and $v_K^-(b) = v_K(a, b) + \beta v_b$ for some $\beta \geq 0$.

First assume $v_K(a, b) \in K$. Then since $v_K(a, b) \in K \subseteq X$, we have $v_K^+(a) = v_K^-(b) = v_K(a, b)$ and $e_K(t) = \{v_K(a, b)\}$ for all $t \in (a, b)$. So $v_K^+(a) = v_K^-(b)$ and $\mathbf{u}_K^{a,b}$ degenerates to the single point $v_K^+(a)$ as claimed.

Now assume $v_K(a, b) \notin K$. Then since $v_K^+(a), v_K^-(b) \in K$ but $v_K(a, b) \notin K$, we have $\alpha, \beta > 0$ and the points $v_K^+(a), v_K^-(b), v_K(a, b)$ are not on the same line, showing (1). Also, the vector $v_K^+(a) - v_K^-(b) = \alpha v_a + \beta v_b$ is equal to $\tau v_{t'}$ for some $\tau > 0$ and $t' \in (a, b)$. So (2) holds. Define $T := X \cap H'$, which is the triangle with vertices $v_K^+(a), v_K^-(b), v_K(a, b)$.

Since $e_T(t' + \pi)$ is the segment connecting $v_K^+(a)$ and $v_K^-(b)$, we have $e_T(t' + \pi) \subseteq K' \subseteq T$. This, with that T have normal angles $a, b, t' + \pi$, implies (i), (iii), (iv) of (3) except $t = a, b$. For the case $t = a$ in (i), use the definition of $v_K^+(a)$ to see that the edge $e_K(a)$ only intersects K at the single point $v_K^+(a)$. Handle the case $t = b$ in (iii) similarly.

It remains to show (ii) of (3). It suffices to show that for all $t \in (a, b)$, we have $e_K(t) \subseteq T$ as this implies $e_K(t) \subseteq K' = K \cap T$. To prove $e_K(t) \subseteq T$, we will show that for any $p \in K \setminus T$ we have $p \notin e_K(t)$. If $t \leq t'$, then the point p is in the convex cone $H_K(a) \setminus H'$ with normal angles a and t that does not contain the vertex $v_K^+(a)$. So we have $p \cdot u_t < v_K^+(a) \cdot u_t \leq h_K(t)$ and $p \notin e_K(t)$ as desired. For the case $t \geq t'$, we can do a similar argument using the cone $H_K(b) \setminus H'$ with normal angles b and t . \square

Theorem 7.3.2. *Let $K \in \mathcal{K}$ and $a, b \in \mathbb{R}$ be arbitrary such that $a < b < a + \pi$. Then the set $\mathbf{u}_K^{a,b}$ in Definition 7.3.1 is a rectifiable oriented Jordan arc from $v_K^+(a)$ to $v_K^-(b)$, and its curve area functional is*

$$\mathcal{J}(\mathbf{u}_K^{a,b}) = \frac{1}{2} \int_{t \in (a,b)} h_K(t) \sigma_K(dt)$$

which is quadratic in $K \in \mathcal{K}$.

Proof. If $v_K^+(a) = v_K^-(b)$, then by Lemma 7.3.1 the set $\mathbf{u}_K^{a,b}$ degenerates to the single point $p = v_K^+(a) = v_K^-(b)$. So for all $t \in (a, b)$, the set $e_K(t)$ is equal to p , and by Theorem 2.1.1 the measure σ_K is zero on the interval (a, b) . So the claimed equality holds.

Now assume the case $v_K^+(a) \neq v_K^-(b)$. Define the convex body K' containing $v_K^+(a)$ and $v_K^-(b)$ as in Lemma 7.3.1. Then by (3) of Lemma 7.3.1, the supporting functions $h_K(t)$ and $h_{K'}(t)$ agree on $t \in [a, b]$, and the surface area measures σ_K and $\sigma_{K'}$ agree on (a, b) by Theorem 2.1.1.

If K' has empty interior, then by (iv) of Lemma 7.3.1 K' should be the line segment connecting $v_K^+(a)$ and $v_K^-(b)$. So $\mathbf{u}_K^{a,b}$ is the line segment from $v_K^+(a)$ to $v_K^-(b)$, and by Proposition 7.2.4 we have $\mathcal{J}(v_K^+(a), v_K^-(b)) = h_K(t')\sigma_K(t')/2$, verifying the equality.

Now assume that K' have nonempty interior. It is a known fact that the boundary $\partial K'$ of K' is a rectifiable Jordan curve.¹ By (3) of Lemma 7.3.1, the boundary $\partial K'$ of K' is the disjoint union of $\mathbf{u}_K^{a,b}$ and the open line segment s from $v_K^-(b)$ to $v_K^+(a)$ excluding endpoints. Since s is a bounded open interval of $\partial K' \simeq S^1$, the curve $\mathbf{u}_K^{a,b}$ is an oriented Jordan arc from $v_K^+(a)$ to $v_K^-(b)$ inheriting the parametrization from $\partial K'$. By Theorem 7.2.3 we have

$$|K'| = \mathcal{J}(\mathbf{u}_K^{a,b}) + \mathcal{J}(v_K^-(b), v_K^+(a)). \quad (7.3)$$

On the other hand, by Theorem 7.1.3 we have

$$|K'| = \frac{1}{2} \int_{t \in S^1} h_{K'}(t) \sigma_{K'}(dt).$$

By (3) of Lemma 7.3.1 and Proposition 7.2.4, this evaluates to

$$|K'| = \frac{1}{2} \int_{t \in (a,b)} h_K(t) \sigma_K(dt) + \mathcal{J}(v_K^-(b), v_K^+(a)). \quad (7.4)$$

¹We outline a proof from [Hag15]. Translate K' so that it has $(0, 0)$ in the interior of K' . Identifying S^1 with the unit vectors of \mathbb{R}^2 , define $f : \partial K' \rightarrow S^1$ as the map $\mathbf{v} \mapsto \mathbf{v}/|\mathbf{v}|$. Then the map is continuous. It is bijective by the convexity of K . As a continuous bijection between compact sets, f is a homeomorphism and the inverse $f^{-1} : S^1 \rightarrow \partial K'$ is continuous. Rectifiability of f^{-1} follows from that the x -coordinate (resp. the y -coordinate) of the point $f^{-1}(t)$ is monotonically decreasing in one interval I of S^1 and monotonically increasing in the complement $S^1 \setminus I$.

By comparing Equation (7.3) to Equation (7.4), we get the desired equality. Quadraticity of $\mathcal{J}(\mathbf{u}_K^{a,b})$ comes from Theorem 7.1.2. \square

Lemma 7.3.3. *Fix $a, b \in \mathbb{R}$ such that $a < b < a + \pi$. The bilinear form $\mathcal{B} : \mathcal{K} \times \mathcal{K} \rightarrow \mathbb{R}$ on \mathcal{K} defined as*

$$\mathcal{B}(K_1, K_2) := \frac{1}{2} \int_{t \in (a,b)} h_{K_1}(t) \sigma_{K_2}(dt)$$

can also be expressed as

$$\mathcal{B}(K_1, K_2) = \frac{1}{2} \int_{t \in (a,b)} v_{K_1}^+(t) \times dv_{K_2}^+(t) = \frac{1}{2} \int_{t \in (a,b)} v_{K_1}^-(t) \times dv_{K_2}^+(t).$$

In particular, for any $K \in \mathcal{K}$ we have

$$\mathcal{J}(\mathbf{u}_K^{a,b}) = \frac{1}{2} \int_{t \in (a,b)} v_K^+(t) \times dv_K^+(t).$$

Proof. Using Theorem 5.2.2, check

$$v_{K_1}^{\pm}(t) \times dv_{K_2}(t) = v_{K_1}^{\pm}(t) \times (v_t \sigma_{K_2}) = (v_{K_1}^{\pm}(t) \times v_t) \sigma_{K_2} = h_{K_1}(t) \sigma_{K_2}$$

as pairs of measures on $t \in (a, b)$. Integrate this on $t \in (a, b)$ to check the equalities for $\mathcal{B}(K_1, K_2)$. Now use $\mathcal{J}(\mathbf{u}_K^{a,b}) = \mathcal{B}(K, K)$. \square

Lemma 7.3.4. *Let $K \in \mathcal{K}$ and $a, b, c \in \mathbb{R}$ be arbitrary such that $a < b < c < a + \pi$. Then the oriented Jordan curve $\mathbf{u}_K^{a,c}$ is the concatenation of $\mathbf{u}_K^{a,b}$, $e_K(b)$, $\mathbf{u}_K^{b,c}$ in order.*

Proof. Use that the union of Jordan curves $\mathbf{u}_K^{a,b}$, $e_K(b)$, $\mathbf{u}_K^{b,c}$ is $\mathbf{u}_K^{a,c}$, and the three curves only meet at respective endpoints $v_K^-(b)$ and $v_K^+(b)$. \square

Lemma 7.3.5. *Take any $K \in \mathcal{K}$ and $a, b \in \mathbb{R}$ such that $a < b < a + \pi$. Assume $v_K^+(a) \neq v_K^-(b)$. Define Γ as the closed curve formed by following the segment from $v_K^+(a)$ to $v_K(a, b)$, then the segment from $v_K(a, b)$ to $v_K^-(b)$, then the curve $\mathbf{u}_K^{a,b}$ in reverse direction. Then the followings are true.*

1. Γ is a counterclockwise Jordan curve.
2. The region R enclosed by Γ is contained in the interior of $H_K(a) \cap H_K(b)$.
3. The region R is disjoint from the set $\cap_{t \in [a,b]} H_K(t)$.

Proof. Let s_a be the line segment from $v_K^+(a)$ to $v_K(a, b)$. Let s_b be the line segment from $v_K(a, b)$ to $v_K^-(b)$. By 3-(i) of Lemma 7.3.1 at $t = a$, the segment s_a intersects with $\mathbf{u}_K^{a,b}$ at the single point $v_K^+(a)$. Likewise, by 3-(iii) of Lemma 7.3.1 at $t = b$, the segment s_b intersects with $\mathbf{u}_K^{a,b}$ at the single point $v_K^-(b)$. By (1) of Lemma 7.3.1, the segments s_a and s_b overlap at the single point $v_K(a, b)$. So the closed curve Γ is indeed a Jordan curve. Use Proposition 7.2.7 to decide the orientation of Γ , proving (1).

Because the curves $\mathbf{u}_K^{a,b}$, s_a , s_b are contained in $H_K(a)$ and $H_K(b)$, the region R is contained in $H_K(a) \cap H_K(b)$. As R is open, this proves (2).

Let $X := \cap_{t \in [a,b]} H_K(t)$. Since $\mathbf{u}_K^{a,b}$, s_a , s_b are disjoint from the interior of X , the curve Γ is contained in the region $\mathbb{R}^2 \setminus X^\circ$. Because $\mathbb{R}^2 \setminus X^\circ$ is simply connected, the region R enclosed by Γ is contained in $\mathbb{R}^2 \setminus X^\circ$. Now the open set R is disjoint from X° , so is disjoint from X , showing (3). \square

7.4 Mamikon's Theorem

We prove a generalized version of *Mamikon's theorem* [Mna97] that works for general convex bodies with non-differentiable boundaries.

Definition 7.4.1. Fix an arbitrary $K \in \mathcal{K}$ and $a, b \in \mathbb{R}$ such that $a < b < a + \pi$. Fix some measurable $\alpha : [a, b] \rightarrow \mathbb{R}$. Define the *Mamikon region* of such parameters as the region swept out by the tangent segment from $v_K^+(t)$ to $\mathbf{z}(t) := v_K^+(t) + \alpha(t)v_t$ over all $t \in [a, b]$.

Follow the boundaries of the Mamikon region, and we can define its area as follows.

Definition 7.4.2. Fix an arbitrary $K \in \mathcal{K}$ and $a, b \in \mathbb{R}$ such that $a < b < a + \pi$. For any continuous function $\mathbf{z} : [a, b] \rightarrow \mathbb{R}^2$ of bounded variation such that $\mathbf{z}(t) \in l_K(t)$ for all $t \in [a, b]$, define the expression

$$\mathcal{M}_K(a, b; \mathbf{z}) = \mathcal{J}(v_K^+(a), \mathbf{z}(a)) + \mathcal{J}(\mathbf{z}|_{[a, b]}) + \mathcal{J}(\mathbf{z}(b), v_K^-(b)) - \mathcal{J}(\mathbf{u}_K^{a, b}).$$

The expression $\mathcal{M}_K(a, b; \mathbf{z})$ in Definition 7.4.2 is the area of the Mamikon region bounded by the two curves $\mathbf{u}_K^{a, b}$ and \mathbf{z} . Mamikon's theorem states that the area is equal to $\frac{1}{2} \int_a^b \alpha(t)^2 dt$ (see Figure 7.1).




Figure 7.1: A Mamikon region divided into differential fans of angle dt and radius $\alpha(t)$.

Theorem 7.4.1. (*Mamikon's theorem, generalized*) Assume the notations in Definition 7.4.2. Then the function $\alpha : [a, b] \rightarrow \mathbb{R}$ defined as $\alpha(t) := (\mathbf{z}(t) - v_K^+(t)) \cdot v_t$ is bounded, measurable, and satisfies $\mathbf{z}(t) = v_K^+(t) + \alpha(t)v_t$. Also, we have

$$\mathcal{M}_K(a, b; \mathbf{z}) = \frac{1}{2} \int_a^b \alpha(t)^2 dt.$$

Proof. It is easy to check the claimed conditions on α . We prove the equality on $\mathcal{M}_K(a, b; \mathbf{z})$. Write $\mathbf{v}(t) := v_K^+(t)$ for all $t \in (a, b]$. Note that \mathbf{v} is right-continuous by Theorem 2.1.3 and of bounded variation by Lemma 5.2.1. So the Lebesgue–Stieltjes measure $d(\mathbf{z} \times \mathbf{v})$ on $[a, b]$ is well-defined, and we have the chain of equalities

$$\begin{aligned}
& \mathbf{z} \times d\mathbf{z} - \mathbf{v} \times d\mathbf{v} + d(\mathbf{z} \times \mathbf{v}) \\
&= \mathbf{z} \times d\mathbf{z} - \mathbf{v} \times d\mathbf{v} + (d\mathbf{z} \times \mathbf{v} + \mathbf{z} \times d\mathbf{v}) \\
&= (\mathbf{z} - \mathbf{v}) \times d(\mathbf{z} + \mathbf{v}) \\
&= (\mathbf{z} - \mathbf{v}) \times d(\mathbf{z} - \mathbf{v}) \\
&= \alpha u_t \times d(\alpha u_t) = \alpha u_t \times (u_t d\alpha + \alpha v_t dt) = \alpha^2 dt
\end{aligned}$$

of measures and functions on $t \in (a, b]$. The first equality uses Lemma 5.1.3 and continuity of \mathbf{z} . The second equality is bilinearity and antisymmetry of \times . As we have $d\mathbf{v}(t) = v_t \sigma$ by Theorem 5.2.2 and $\mathbf{z}(t) - \mathbf{v}(t) = \alpha_K(t)v_t$, they are parallel and we get $(\mathbf{z}(t) - \mathbf{v}(t)) \times d\mathbf{v}(t) = 0$ which is used in the third equality. The last chain of equalities are basic calculations.

If we integrate the formula above on the whole interval (a, b) , the terms $\mathbf{z}(t) \times d\mathbf{z}(t)$ and $\mathbf{v}(t) \times d\mathbf{v}(t)$ becomes $2\mathcal{J}(\mathbf{z})$ and $2\mathcal{J}(\mathbf{v})$ by Definition 7.2.6 and the last equality of Lemma 7.3.3 respectively. The measure $d(\mathbf{z}(t) \times \mathbf{v}(t))$ integrates to the difference $2\mathcal{J}(\mathbf{z}(b), v_K^-(b)) - 2\mathcal{J}(\mathbf{z}(a), v_K^+(a))$. So the integral matches twice the Definition 7.4.2, completing the proof. \square

Theorem 7.4.2. *Fix $a, b \in \mathbb{R}$ such that $a < b < a + \pi$. Assume that the function $\mathbf{z}_K \in C^{\text{BV}}[a, b]$ is determined by $K \in \mathcal{K}$ so that (i) $\mathbf{z}_K(t)$ is on the line $l_K(t)$ for all $t \in [a, b]$, and (ii) the map $K \mapsto \mathbf{z}_K$ is convex-linear in K . Then the expression $\mathcal{M}_K(a, b; \mathbf{z}_K)$ is quadratic and convex as a functional on $K \in \mathcal{K}$.*

Proof. Let $\alpha_K(t) := (\mathbf{z}_K(t) - v_K^+(t)) \cdot v_t$, then α_K is convex-linear in K by convex-linearity of \mathbf{z}_K and $v_K^+(t)$ (Theorem 7.1.2). So $\mathcal{M}_K(a, b; \mathbf{z}_K)$ which is $\frac{1}{2} \int_a^b \alpha_K(t)^2 dt$ by Theorem 7.4.1 is quadratic and convex in K . \square

Chapter 8

Optimality of Gerver's Sofa

This chapter proves the optimality of Gerver's sofa (Theorem 1.1.1). Each section of this chapter establishes the following portion of the overview in Section 1.8. Recall that the previous Chapter 7 prepares necessary technical lemmas for this chapter.

- Section 8.1 defines the domain \mathcal{L} of \mathcal{Q} , which is the tuple (K, B, D) of three convex bodies as described in Section 1.8.2, and build the embedding $K \mapsto (K, B_K, D_K)$ from caps $K \in \mathcal{K}^i$ with injectivity condition to a subset of \mathcal{L} .
- Section 8.2 formally defines \mathcal{Q} using quadratic expressions on K, B , and D . The formula of \mathcal{Q} traces the boundary of the overestimated region R described in Section 1.8.1 and Section 1.8.2. We use Jordan curve theorem to rigorously show that $\mathcal{Q}(K, B_K, D_K)$ is an upper bound of sofa area functional $\mathcal{A}(K)$ on $K \in \mathcal{K}^i$.
- Section 8.3 establishes the concavity of \mathcal{Q} on \mathcal{L} using Mamikon's theorem.
- Section 8.4 translates the local optimality conditions of Gerver's sofa G derived by Romik in [Rom18] to equalities of surface area measures of K, B_K , and D_K corresponding to G .
- Section 8.5 calculates the directional derivative of \mathcal{Q} at $(K, B_K, D_K) \in \mathcal{L}$ arising from Gerver's sofa G . The conditions on G in Section 8.4 show that

8.1 Domain of \mathcal{Q}

In this Section 8.1, we define the domain \mathcal{L} of the soon-to-be-established upper bound \mathcal{Q} which extends the collection \mathcal{K}^i of caps satisfying the injectivity condition.

Definition 8.1.1. Define \mathcal{K}^i as the subset of caps $K \in \mathcal{K}^c$ that (i) satisfies the injectivity condition (Definition 6.1.2), and (ii) have area $|K| \geq 2.2$.

Remind that for any $K \in \mathcal{K}^i$, we can talk about the density function r_K and s_K of the surface area measure σ_K on $[0, \pi/2)$ and $(\pi/2, \pi]$ respectively by (1) of Definition 6.1.2. Likewise, the walls $a_K(t)$ and $c_K(t)$ makes contact with K at unique points $A_K(t)$ and $C_K(t)$ respectively (see Definition 6.4.1), and we can talk about the arm lengths $f_K(t) = f_K^\pm(t)$ and $g_K(t) = g_K^\pm(t)$.

Theorem 8.1.1. *The space \mathcal{K}^i is a convex subspace of \mathcal{K}^c containing all balanced maximum caps $K \in \mathcal{K}^c$ and the cap $\mathcal{C}(G)$ of Gerber's sofa.*

Proof. We first show that $\mathcal{K}^i \subset \mathcal{K}^c$ is a convex subset. The Definition 6.1.2 of injectivity condition on K is made of linear constraints in K , and so is preserved under the barycentric operation of \mathcal{K}^c . For any $\lambda \in [0, 1]$, by the Brunn-Minkowski theorem

$$|(1 - \lambda)K_1 + \lambda K_2| \geq |K_1|^{1-\lambda}|K_2|^\lambda$$

we have $|c_\lambda(K_1, K_2)| \geq 2.2$ if $|K_1|, |K_2| \geq 2.2$. This shows that the condition $|K| \geq 2.2$ is also closed under c_λ . So \mathcal{K}^i defines a convex subset of \mathcal{K}^c .

By Theorem 6.1.1 and Theorem 6.1.2, both the balanced maximum caps K and the cap $\mathcal{C}(G)$ of Gerber's sofa satisfy the injectivity condition. It is known that G have area $2.2195 \dots \geq 2.2$, and any balanced maximum cap K attains the maximum value of sofa area functional $\mathcal{A}(K)$, so it should have area $|K| \geq \mathcal{A}(K) \geq \mathcal{A}(G) = 2.2195 \dots \geq 2.2$ again. \square

Define the angle constants determining Gerber's sofa G .

Definition 8.1.2. Define φ and θ as the angles satisfying $0 < \varphi < \theta < \pi/4$ and the Equations 27 to 44 in [Rom18]. Note that $\varphi \in [0.039, 0.040]$ in particular (see Table 1 of [Rom18]). Define the constants $\varphi^R := \varphi$ and $\varphi^L := \pi/2 - \varphi$.

Remark 8.1.1. The angles φ and θ are used in the definition of Gerber's sofa G ; see Section 8.4 for the details. In particular, the blue core of G in Figure 1.1 is the trajectory of the inner corner $\mathbf{x}_G(t)$ on the interval $t \in [\varphi^R, \varphi^L]$.

Definition 8.1.3. Define \mathcal{L} as the space of all tuples (K, B, D) of convex bodies such that the followings are true.

1. $K \in \mathcal{K}^i$, $B \subseteq K$, and $D \subseteq K$.
2. For every $t \in [\varphi^R, \pi/2]$, we have $h_K(t) + h_B(\pi + t) \leq 1$.
3. Equality holds in (2) at $t = \varphi^R, \pi/2$.
4. For every $t \in [0, \varphi^L]$, we have $h_K(\pi/2 + t) + h_D(3\pi/2 + t) \leq 1$.
5. Equality holds in (4) at $t = 0, \varphi^L$.

Proposition 8.1.2. *The space \mathcal{L} is a convex domain with the barycentric operation*

$$c_\lambda((K_1, B_1, D_1), (K_2, B_2, D_2)) := (c_\lambda(K_1, K_2), c_\lambda(B_1, B_2), c_\lambda(D_1, D_2)).$$

Proof. The product $\mathcal{K} \times \mathcal{K} \times \mathcal{K}$ of convex domains \mathcal{K} is naturally a convex domain. Recall that \mathcal{K}^i is a convex subspace of \mathcal{K} . That $B \subseteq K$ (resp. $D \subseteq K$) can be written as linear constraints $h_B(t) \leq h_K(t)$ (resp. $h_D(t) \leq h_K(t)$) over all $t \in S^1$. So the condition (1) of Definition 8.1.3 is preserved under barycentric operations. By (1) of Theorem 7.1.2, the conditions (2) to (5) of Definition 8.1.3 are linear constraints. \square

We now build the injection $\mathcal{K}^i \rightarrow \mathcal{L}$ mapping $K \in \mathcal{K}^i$ to $(K, B_K, D_K) \in \mathcal{L}$. That $(K, B_K, D_K) \in \mathcal{L}$ is not too difficult to show using injectivity condition, but requires book-keeping of many definitions.

Definition 8.1.4. Take arbitrary cap $K \in \mathcal{K}^i$. Define the convex body

$$B_K := K \cap \bigcap_{t \in [\varphi_R, \pi/2]} H_K^b(t).$$

Similarly, define the convex body

$$D_K := K \cap \bigcap_{t \in [0, \varphi_L]} H_K^d(t).$$




Figure 8.1: Diagram for the upper bound $\mathcal{Q}(K, B, D)$ in Definition 8.2.2.

Definition 8.1.5. (See Figure 8.1) For any $K \in \mathcal{K}^c$, define the line $b_K^R := b_K(\varphi^R)$ and the half-plane $\check{H}_K^R := H_K^b(\varphi^R)$ bounded from below by the line b_K^R . Define the points $W_K^R := W_K(\varphi^R)$ and $\mathbf{x}_K^R := \mathbf{x}_K(\varphi^R)$ on the line b_K^R .

Similarly, define the line $d_K^L := d_K(\varphi^L)$ and the half-plane $\check{H}_K^L := H_K^d(\varphi^L)$ bounded from below by the line d_K^L . Define the point $Z_K^L := Z_K(\varphi^L)$ and $\mathbf{x}_K^L := \mathbf{x}_K(\varphi^L)$ on the line d_K^L .

In Definition 8.1.5, the superscripts **L** and **R** denote that we are essentially plugging in the values φ^L and φ^R respectively. The only exceptions are half-planes \check{H}_K^R and \check{H}_K^L , where the accent \check{H} means that we are using the half-planes H_K^b and H_K^d bounding the sofa from below, not the supporting planes H_K bounding K from above.

Definition 8.1.6. For any $K \in \mathcal{K}^i$ with $|K| \geq 2.2$, define the following parallelograms.

$$P_K^R := H \cap H_K(\varphi^R) \cap \check{H}_K^R \quad P_K^L := H \cap H_K(\pi/2 + \varphi^L) \cap \check{H}_K^L$$

Lemma 8.1.3. The parallelogram P_K^R is bounded by the lines $l(\pi/2, 0)$, $l(\pi/2, 1)$, $a_K(\varphi^R)$ and $b_K(\varphi^R)$. It has base $\sec \varphi$, height 1, and angle of $\pi/2 \pm \varphi$ at each vertices. The lower-left corner of P_K^R is W_K^R .

Proof. Recall that the point $W_K^R = W_K(\varphi^R)$ is the intersection $b_K^R \cap l(\pi/2, 0)$ by definition. So $W_K(\varphi^R)$ is the lower-left corner of the parallelogram P_K^R . \square

Lemma 8.1.4. For any $K \in \mathcal{K}^i$, the sets $K \cap \check{H}_K^R$ and $K \cap \check{H}_K^L$ are disjoint.

Proof. Fix an $K \in \mathcal{K}^i$. Because K is in the horizontal strip H and the supporting half-plane $H_K(\varphi^R)$, the set $K \cap \check{H}_K^R$ is in the parallelogram P_K^R in Definition 8.1.6. Likewise, the set $K \cap \check{H}_K^L$ is contained in the parallelogram P_K^L . Now assume by contradictory that $K \cap \check{H}_K^R$ and $K \cap \check{H}_K^L$ intersect. Then P_K^R and P_K^L should intersect as well. The bases of P_K^R and P_K^L on the line $l(\pi/2, 1)$ have length $\sec \varphi$. Note that K is contained in the trapezoid $R := \check{H} \cap H_K(\varphi^R) \cap H_K(\pi/2 + \varphi^L)$ containing P_K^R and P_K^L . Since P_K^R and P_K^L overlaps, the side length of R on the line $l(\pi/2, 1)$ is at most $2\sec \varphi$, and so the base of R on the line $l(\pi/2, 0)$ is at most $2\sec \varphi + 2\tan \varphi = 2.08 \dots < 2.2$ which can be checked by computation. Now we have $|K| \leq |R| < 2.2$ and get contradiction. \square

Lemma 8.1.5. *For any $K \in \mathcal{K}^i$, the points W_K^R, Z_K^L are in the edge $e_K(3\pi/2)$ excluding the endpoints $A_K(0)$ and $C_K(\pi/2)$.*

Proof. By Theorem 2.5.5, the point W_K^R is strictly on the left side of the right endpoint $A_K(0)$ of $e_K(3\pi/2)$. So the base of the parallelogram P_K^R contains $A_K(0)$. Because $|K| \geq 2.2$, the edge $e_K(3\pi/2)$ have length ≥ 2.2 , which is strictly larger than the base $\sec \varphi < 1.1$ of P_K^R . So the point W_K^R is also strictly on the right side of the left endpoint $C_K(\pi/2)$ as well, completing the proof for W_K^R . Use a mirror-symmetric argument for Z_K^L . \square

The following lemma shows that the region \check{H}_K^R cannot distinguish the wedge $T_K(t)$ from the complement of $H_K^b(t)$.

Lemma 8.1.6. *Fix an arbitrary $K \in \mathcal{K}^i$.*

1. *Take any $t \in (\varphi^R, \pi/2]$. Then the point $\mathbf{x}_K(t)$ is outside the half-plane \check{H}_K^R , and we have $\check{H}_K^R \cap Q_K^-(t) = \check{H}_K^R \setminus H_K^b(t)$. So we also have $\check{H}_K^R \cap T_K(t) = \check{H}_K^R \cap H_+(\pi/2, 0) \setminus H_K^b(t)$.*
2. *Take any $t \in [0, \varphi^L)$. The point $\mathbf{x}_K(t)$ is outside the half-plane \check{H}_K^L , and we have $\check{H}_K^L \cap Q_K^-(t) = \check{H}_K^L \setminus H_K^d(t)$. So we also have $\check{H}_K^L \cap T_K(t) = \check{H}_K^L \cap H_+(\pi/2, 0) \setminus H_K^d(t)$.*

Proof. We prove (1). The point $\mathbf{x}_K(\varphi^R)$ is on the boundary b_K^R of \check{H}_K^R by definition. As $K \in \mathcal{K}^i$ satisfies the injectivity condition by definition, it satisfies (3) of Definition 6.1.2 and we have $\mathbf{x}'_K(t) \cdot u_{\varphi^R} < 0$ for all $t \in (\varphi^R, \pi/2)$. By integrating, we have $\mathbf{x}_K(t) \cdot u_{\varphi^R} < \mathbf{x}_K(\varphi^R) \cdot u_{\varphi^R}$ for all $t \in (\varphi^R, \pi/2]$, so we have $\mathbf{x}_K(t) \notin \check{H}_K^R$.

We now show $\check{H}_K^R \cap Q_K^-(t) = \check{H}_K^R \setminus H_K^b(t)$. Recall that the quadrant $Q_K^-(t)$ have half-lines $\vec{b}_K(t)$ and $\vec{d}_K(t)$ as boundary, so we have $Q_K^-(t) = \mathbb{R}^2 \setminus H_K^b(t) \setminus H_K^d(t)$. Take the cone $C := H_K^d(t) \setminus H_K^b(t)$. Then the vertex $\mathbf{x}_K(t)$ of C is outside the half-plane \check{H}_K^R of normal angle φ^R . Also, as $t \in (\varphi^R, \pi/2]$ the spanning directions u_t and $-v_t$ of C are in the positive direction of the normal angle $-u_{\varphi^R}$ of \check{H}_K^R . So C and \check{H}_K^R are disjoint, and we have

$$\check{H}_K^R \cap Q_K^-(t) = \check{H}_K^R \cap (Q_K^-(t) \cup C) = \check{H}_K^R \cap (\mathbb{R}^2 \setminus H_K^b(t)).$$

Intersect above with $H_+(\pi/2, 0)$ to get the second equation of (1). The proof of (2) is a mirror-symmetric argument. \square

Lemma 8.1.7. *For any cap $K \in \mathcal{K}^i$, let $B := B_K$ and $D := D_K$. Then the followings are true.*

1. *For every $t \in [\varphi^R, \pi/2]$, we have $h_K(t) + h_B(\pi + t) \leq 1$.*

2. Equality holds in (1) at $t = \varphi^R, \pi/2$. So $l_B(3\pi/2) = l(\pi/2, 0)$ and $l_B(\pi + \varphi^R) = b_K^R$.
3. For every $t \in [0, \varphi^L]$, we have $h_K(\pi/2 + t) + h_D(3\pi/2 + t) \leq 1$.
4. Equality holds in (3) at $t = 0, \varphi^R$. So $l_D(3\pi/2) = l(\pi/2, 0)$ and $l_D(3\pi/2 + \varphi^L) = d_K^L$.

Proof. By Definition 8.1.4, we have $B \subseteq H_K^b(t) = H_-(\pi + t, 1 - h_K(t))$ for all $t \in [\varphi^R, \pi/2]$, and this completes the proof of (1). We now prove (2).

By Theorem 2.5.5, we have $A_K(0) \in H_K^b(t)$ for all $t \in [\varphi^R, \pi/2]$ and so $A_K(0) \in B_K$. This with $B_K \subseteq K$ implies that $l_B(3\pi/2) = l(\pi/2, 0)$. It remains to prove $l_B(\pi + \varphi^R) = b_K^R$. Since $B \subseteq H_K^b(\varphi^R)$, and b_K^R is the boundary of $\tilde{H}_K^R = H_K^b(\varphi^R)$, it suffices to show that there exists some point $p \in B \cap b_K^R$.

We will show that the upper boundary δK of K and the line b_K^R intersect at a single point p . By Lemma 8.1.5, the line b_K^R passes through W_K^R which is in the edge $e_K(3\pi/2) \setminus \{A_K(0), C_K(\pi/2)\}$ of K . So it should pass through exactly one another point p of the boundary ∂K of K . As $\delta K = \partial K \setminus e_K(3\pi/2) \cup \{A_K(0), C_K(\pi/2)\}$, the point p is in δK and is the unique intersection of δK and b_K^R .

Now take p as the unique intersection of δK and b_K^R . Recall that our goal is to show $p \in B \cap b_K^R$. Since $p \in K \cap b_K^R$, it suffices to show that $p \in H_K^b(t)$ for all $t \in [\varphi^R, \pi/2]$. By (2) of Theorem 2.5.8 and that $p \in \delta K$, the point p is outside the niche $\mathcal{N}(K)$. So p is outside all $Q_K^-(t)$ over $t \in [0, \pi/2]$. By (1) of Lemma 8.1.6, the point p should be inside $H_K^b(t)$ for all $t \in [\varphi^R, \pi/2]$. This completes the goal.

The proofs of (3) and (4) can be done using mirror-symmetric arguments. □

We now make the extension $\mathcal{K}^i \rightarrow \mathcal{L}$.

Theorem 8.1.8. *For any $K \in \mathcal{K}^i$, the triple (K, B_K, D_K) is in \mathcal{L} .*

Proof. We need to check the conditions of Definition 8.1.3 for $B := B_K$ and $D := D_K$. Condition (1) is evident from the definitions. Conditions (2) to (5) are exactly Lemma 8.1.7. □

Remark 8.1.2. Even though both \mathcal{K}^i and \mathcal{L} are convex domains, the injection $\mathcal{K}^i \rightarrow \mathcal{L}$ defined as map $K \mapsto (K, B_K, D_K)$ in Theorem 8.1.8 is *not* convex-linear. The lower left portion of the convex body B_K is the *Aleksandrov body* (or the *Wulff shape*) $\bigcap_{t \in [\varphi^R, \pi/2]} H_+(t, f(t))$ of the function $f(t) := 1 - h_K(\pi + t)$ over $t \in [\varphi^R, \pi/2]$, which is hard to understand in terms of f directly. This makes the injection $\mathcal{K}^i \rightarrow \mathcal{L}$ very important; it ‘irons out’ the sofa area functional $\mathcal{A} : \mathcal{K}^i \rightarrow \mathbb{R}$, which is not quadratic, to a quadratic functional $\mathcal{Q} : \mathcal{L} \rightarrow \mathbb{R}$ (Theorem 8.2.4).

8.2 Definition of \mathcal{Q}

In this Section 8.2, we give the full Definition 8.2.2 of the upper bound $\mathcal{Q} : \mathcal{L} \rightarrow \mathbb{R}$ of area (see Figure 8.1). For the cap $K := \mathcal{C}(G)$ of Gerver’s sofa G , the inner corner $\mathbf{x}_K(t)$ draws a main ‘core’ part of the niche $\mathcal{N}(K)$ at the interval $t \in [\varphi^R, \varphi^L]$. Following this, we break the niche $\mathcal{N}(K)$ into three parts for the construction of the upper bound \mathcal{Q} .

Definition 8.2.1. (See Figure 8.1) For any convex body B , define the convex curve

$$\mathbf{b}_B := \mathbf{u}_B^{\pi + \varphi^R, 3\pi/2}$$

which is from the vertex $X_B := v_B^+(\pi + \varphi^R)$ to the vertex $W_B := v_B^-(3\pi/2)$. Likewise, for any convex body D , define the convex curve

$$\mathbf{d}_D := \mathbf{u}_D^{3\pi/2, 3\pi/2 + \varphi^L}$$

which is from the vertex $Z_D := v_D^+(3\pi/2)$ to the vertex $Y_D := v_D^-(3\pi/2 + \varphi^L)$.

Definition 8.2.2. Define the function $\mathcal{Q} : \mathcal{L} \rightarrow \mathbb{R}$ as $\mathcal{Q}(K, B, D) :=$

$$|K| + \mathcal{J}(\mathbf{d}_D) + \mathcal{J}(Y_D, \mathbf{x}_K^L) - \mathcal{J}(\mathbf{x}_K|_{[\varphi^R, \varphi^L]}) + \mathcal{J}(\mathbf{x}_K^R, X_B) + \mathcal{J}(\mathbf{b}_B).$$

Proposition 8.2.1. \mathcal{Q} is a quadratic functional on \mathcal{L} .

Proof. The term $\mathcal{J}(\mathbf{x}_K|_{[\varphi^R, \varphi^L]})$ is quadratic in K by Proposition 7.2.2 and that $\mathbf{x}_K(t) = (h_K(t) - 1)u_t + (h_K(t + \pi/2) - 1)v_t$ is convex-linear in K (Proposition 2.2.2 and (1) of Theorem 7.1.2). The terms $\mathcal{J}(\mathbf{d}_D)$ and $\mathcal{J}(\mathbf{b}_B)$ are quadratic in D and B by Theorem 7.3.2. The terms $\mathcal{J}(Y_D, \mathbf{x}_K^L)$ and $\mathcal{J}(\mathbf{x}_K^R, X_B)$ are quadratic because X_B, \mathbf{x}_K^R and \mathbf{x}_K^L, Y_D are linear in B, K, D respectively by (2) of Theorem 7.1.2. \square

Now we bound the right and left part of $\mathcal{N}(K)$.

Lemma 8.2.2. Fix an arbitrary $K \in \mathcal{K}^i$ and take $B := B_K$. We have

$$\left| \mathcal{N}(K) \cap \check{H}_K^R \right| \geq \mathcal{J}(X_B, W_K^R) - \mathcal{J}(\mathbf{b}_B)$$

and similarly for $D := D_K$, we have

$$\left| \mathcal{N}(K) \cap \check{H}_K^L \right| \geq \mathcal{J}(Z_K^L, Y_D) - \mathcal{J}(\mathbf{d}_D).$$

Proof. We show the first inequality. The second inequality can be proven by a mirror-symmetric argument.

Recall the Definition 8.2.1 that the curve $\mathbf{b}_B := \mathbf{u}_B^{\pi + \varphi^R, 3\pi/2}$ is the segment of ∂B from the vertex $X_B := v_B^+(\varphi^R + \pi)$ of B to the vertex $W_B := v_B^-(3\pi/2)$ of B . Also, we have $W_K^R = v_B(\pi + \varphi^R, 3\pi/2)$ by (2) of Lemma 8.1.7. Because W_K^R and W_B are on the line $l(\pi/2, 0)$, we have $\mathcal{J}(W_K^R, W_B) = 0$, which we will use implicitly.

If $X_B = W_B$, then by Lemma 7.3.1 the curve \mathbf{b}_B and the points X_B, W_B, W_K^R degenerate to a single point. So the lower bound becomes zero if $X_B = W_B$. Now assume otherwise that $X_B \neq W_B$.

Define the closed curve Γ obtained by following \mathbf{b}_B in the reverse direction, and then the line segments from X_B to W_K^R and from W_K^R to W_B respectively. Then by (1) of Lemma 7.3.5, the curve Γ is a counterclockwise Jordan curve enclosing a region R . By Theorem 7.2.3 and Proposition 7.2.6, the area of R is equal to $\mathcal{J}(X_B, W_K^R) - \mathcal{J}(\mathbf{b}_B)$. So it remains to show that the region R is contained in the set $\mathcal{N}(K) \cap \check{H}_K^R$.

Take any point $p \in R$. Our goal is to show that $p \in \mathcal{N}(K) \cap \check{H}_K^R$. By (2) of Lemma 7.3.5 and (2) of Lemma 8.1.7, we have

$$p \in H_B(\pi + \varphi^R) \cap H_B(3\pi/2) = \check{H}_K^R \cap H_+(\pi/2, 0).$$

Because $B \subseteq K$ by definition, we have $\mathbf{b}_B \subseteq K$. Also by Lemma 8.1.5, we have $W_K^R \subseteq K$ so the closed curve Γ is contained in K . Now $R \subseteq K$ as K is contractible. We also have R disjoint from B by (3) of Lemma 7.3.5. By $p \in R \subseteq K$ and the Definition 8.1.4 of B , we have $p \notin H_K^b(t)$ for some $t \in [\varphi^R, \pi/2]$. By (2) of Lemma 7.3.5, we have $t \neq \varphi^R, \pi/2$. By (1) of Lemma 8.1.6, we now have $p \in \check{H}_K^R \cap T_K(t) \subseteq \check{H}_K^R \cap \mathcal{N}(K)$. \square

We now approximate the middle part of the niche traced out by the core \mathbf{x}_K restricted to the interval $[\varphi^R, \varphi^L]$.

Lemma 8.2.3. *For any $K \in \mathcal{K}^i$, we have*

$$\left| \mathcal{N}(K) \setminus \check{H}_K^R \setminus \check{H}_K^L \right| \geq \mathcal{J}(W_K^R, \mathbf{x}_K^R) + \mathcal{J}(\mathbf{x}_K|_{[\varphi^R, \varphi^L]}) + \mathcal{J}(\mathbf{x}_K^L, Z_K^L).$$

Proof. Denote $\mathbf{x}_K|_{[\varphi^R, \varphi^L]}$, b_K^R , d_K^L simply as \mathbf{x} , b , d . Define $Y := \mathbb{R}^2 \setminus \check{H}_K^R \setminus \check{H}_K^L$ which is the open cone bounded by the lines b and d . The curve $\mathbf{x}(t)$ on $t \in [\varphi^R, \varphi^L]$ starts at b and ends at d by definition. Also, by Lemma 8.1.6, the middle parts $\mathbf{x}(t)$ for $t \in (\varphi^R, \varphi^L)$ are inside the open cone Y .

Take the horizontal line l_h described by the equation $y = -h$ for sufficiently large $h > 0$, so that the trajectory of \mathbf{x} is strictly above l_h . We construct a closed curve Γ which is the concatenation of the following four curves in order.

1. $\mathbf{x} : [\varphi^R, \varphi^L] \rightarrow \mathbb{R}^2$.
2. The line segment s_L from $\mathbf{x}(\varphi^L)$ to $d \cap l_h$.
3. The line segment s_M from $d \cap l_h$ to $b \cap l_h$.
4. The line segment s_R from $b \cap l_h$ to $\mathbf{x}(\varphi^R)$.

Then Γ is a Jordan curve since the interior of \mathbf{x} is inside the open cone Y , and h is taken so that s_M is disjoint from \mathbf{x} . Let G be the open region enclosed by the Jordan curve Γ . Similarly, let R be the closed trapezoid right below the line $l(\pi/2, 0)$ with the vertices $Z_K^L, d \cap l_h, b \cap l_h, W_K^R$ in counterclockwise order. Then the lower bound of the stated inequality is $|G| - |R|$ by following the boundaries of G and R in order and using Theorem 7.2.3 twice.

So it suffices to show $G \setminus R \subseteq \mathcal{N}(K) \cap Y$, as this will imply

$$\left| \mathcal{N}(K) \setminus \check{H}_K^R \setminus \check{H}_K^L \right| = |\mathcal{N}(K) \cap Y| \geq |G| - |R|$$

and proving the theorem. Let H_h be the closed half-plane with boundary l_h above the line l_h . As G is contained in the cone Y and strictly above the line l_h , we have $G \subseteq Y \cap H_h$. On the other hand, by the definition of R we have $Y \cap H_h \setminus H_+(\pi/2, 0) \subseteq R$. Thus we have $G \setminus R \subseteq H_+(\pi/2, 0) \cap Y$. So it remains to show $G \cap H_+(\pi/2, 0) \subseteq \mathcal{N}(K)$.

Take any point $p \in G \cap H_+(\pi/2, 0)$. Let r be the half-line from p in the direction of v_0 , then since $p \in G$ the half-line r intersects Γ at a point $q \neq p$. As $p \in H_+(\pi/2, 0)$, the point q is not in s_M . Thus q is in one of the curves \mathbf{x} , s_L , or s_R . If $q = \mathbf{x}(t)$ for some $t \in [\varphi^R, \varphi^L]$, then $p \in Q_K^-(t)$. If $q \in s_R$, then $p \in Q_K^-(\varphi^R)$. If $q \in s_L$, then $p \in Q_K^-(\varphi^L)$. In any case, we have $p \in \mathcal{N}(K)$, completing the proof. \square

Theorem 8.2.4. *For any $K \in \mathcal{K}^i$, we have $\mathcal{A}(K) \leq \mathcal{Q}(K, B_K, D_K)$.*

Proof. Set $B := B_K$ and $D := D_K$. Add all the inequalities in Lemma 8.2.2 and Lemma 8.2.3. We get

$$\begin{aligned} \mathcal{A}(K) &= |K| - |\mathcal{N}(K)| \\ &= (|K| - |\mathcal{N}(K) \setminus \check{H}_K^R \setminus \check{H}_K^L|) - |\mathcal{N}(K) \cap \check{H}_K^R| - |\mathcal{N}(K) \cap \check{H}_K^L| \\ &\leq |K| - \mathcal{J}(W_K^R, \mathbf{x}_K^R) - \mathcal{J}(\mathbf{x}_K|_{[\varphi^R, \varphi^L]}) - \mathcal{J}(\mathbf{x}_K^L, Z_K^L) \\ &\quad - \mathcal{J}(X_B, W_K^R) + \mathcal{J}(\mathbf{b}_B) - \mathcal{J}(Z_K^L, Y_D) + \mathcal{J}(\mathbf{d}_D) \\ &= \mathcal{Q}(K, B, D) \end{aligned}$$

proving the theorem. □

8.3 Concavity of \mathcal{Q}

We prove the concavity of \mathcal{Q} (Theorem 8.3.8). The main idea is depicted in Figure 8.2. Each grey Mamikon region in the figure have an area that is *convex* in the domain \mathcal{K}^i or \mathcal{L} by Theorem 7.4.2. The function \mathcal{Q} plus the area of such Mamikon regions turns out to be linear in $K \in \mathcal{K}^i$, so the function \mathcal{Q} should be concave.




Figure 8.2: The Mamikon regions used in the proof of Theorem 8.3.8 (compare with Figure 8.1).

We first give a particular parametrization of the supporting line $l_K(t)$ of a convex body K .

Definition 8.3.1. Let $K \in \mathcal{K}$ and $t \in \mathbb{R}$ be arbitrary. Define $\mathbf{I}_K^t : (t - \pi, t] \rightarrow \mathbb{R}^2$ as $\mathbf{I}_K^t(s) = v_K(s, t)$ for $s < t$ and $\mathbf{I}_K^t(t) = v_K^-(t)$.

Theorem 8.3.1. Fix arbitrary $K \in \mathcal{K}$ and $t \in \mathbb{R}$. For any $a, b \in (t - \pi, t]$ such that $a \leq b$, the function \mathbf{I}_K^t restricted to $[a, b]$ is a continuous parametrization of the closed segment in $l_K(t)$ from $\mathbf{I}_K^t(a)$ to $\mathbf{I}_K^t(b)$. Consequently, we have $\mathcal{J}(\mathbf{I}_K^t|_{[a,b]}) = \mathcal{J}(\mathbf{I}_K^t(a), \mathbf{I}_K^t(b))$.

Proof. If $b < t$, then the result follows from the geometric fact that for all $s \in [a, b]$, the intersection $v_K(s, t) = l_K(s) \cap l_K(t)$ is continuous in s and contained in the line segment connecting $v_K(a, t)$ and $v_K(b, t)$. Use the limit $\lim_{b \rightarrow t^-} \mathbf{I}_K^t(b) = \mathbf{I}_K^t(t)$ in Theorem 2.1.3 to extend this to $b = t$. □

Theorem 8.3.2. For fixed $t \in \mathbb{R}$ and $a, b \in (t - \pi, t]$, the value $\mathbf{I}_K^t|_{[a,b]} \in C^{\text{BV}}[a, b]$ is linear in $K \in \mathcal{K}$.

Proof. By (2) of Theorem 7.1.2. □

We give names to the Mamikon regions in Figure 8.2.

Definition 8.3.2. For any $K \in \mathcal{K}^i$, define $\mathcal{S}_K :=$

$$\mathcal{M}_K \left(0, \varphi^{\text{R}}; \mathbf{I}_K^{\pi/2} \right) + \mathcal{M}_K \left(\varphi^{\text{R}}, \varphi^{\text{L}}; \mathbf{y}_K \right) + \mathcal{M}_K \left(\varphi^{\text{L}}, \pi/2; \mathbf{I}_K^{\pi/2 + \varphi^{\text{L}}} \right) + \mathcal{M}_K \left(\pi/2, \pi; \mathbf{I}_K^{\pi} \right).$$

Definition 8.3.3. For convex bodies B and D , define

$$\mathcal{R}_B := \mathcal{M}_B \left(\pi/2 + \varphi^{\text{R}}, 3\pi/2; \mathbf{I}_B^{3\pi/2} \right) \quad \text{and} \quad \mathcal{L}_D := \mathcal{M}_D \left(3\pi/2, 3\pi/2 + \varphi^{\text{L}}; \mathbf{I}_D^{3\pi/2 + \varphi^{\text{L}}} \right).$$

Note that Definition 8.3.2 and Definition 8.3.3 only uses Mamikon areas in Definition 7.4.2. So by Theorem 7.4.2, these are convex and quadratic.

Lemma 8.3.3. The values \mathcal{S}_K , \mathcal{L}_D , \mathcal{R}_B are convex and quadratic as functionals on the convex bodies K, B, D respectively.

Proof. By Theorem 7.4.2, and the linearity of $\mathbf{y}_K(t) = h_K(t)u_t + h_K(\pi/2 + t)v_t$ and $\mathbf{I}_K(t)$ in K (Theorem 7.1.2 and Theorem 8.3.2). □

Definition 8.3.4. For any $K \in \mathcal{K}^i$, define the functional

$$\mathcal{P}_K := |K| + \mathcal{J} \left(Z_K^{\text{L}}, \mathbf{x}_K^{\text{L}} \right) - \mathcal{J} \left(\mathbf{x}_K|_{[\varphi^{\text{R}}, \varphi^{\text{L}}]} \right) + \mathcal{J} \left(\mathbf{x}_K^{\text{R}}, W_K^{\text{R}} \right).$$

Lemma 8.3.4. For any $(K, B, D) \in \mathcal{L}$, we have

$$\mathcal{Q}(K, B, D) = \mathcal{P}_K - \mathcal{R}_B - \mathcal{L}_D.$$

Proof. Unfold Definition 8.2.2 of \mathcal{Q} , Definition 8.3.4 of \mathcal{S}_K , and Definition 8.3.3 of \mathcal{R}_B and \mathcal{L}_D . It remains to verify

$$\begin{aligned} & |K| + \mathcal{J}(\mathbf{d}_D) + \mathcal{J}(Y_D, \mathbf{x}_K^{\text{L}}) - \mathcal{J}(\mathbf{x}_K|_{[\varphi^{\text{R}}, \varphi^{\text{L}}]}) + \mathcal{J}(\mathbf{x}_K^{\text{R}}, X_B) + \mathcal{J}(\mathbf{b}_B) \\ = & |K| + \mathcal{J}(Z_K^{\text{L}}, \mathbf{x}_K^{\text{L}}) - \mathcal{J}(\mathbf{x}_K|_{[\varphi^{\text{R}}, \varphi^{\text{L}}]}) + \mathcal{J}(\mathbf{x}_K^{\text{R}}, W_K^{\text{R}}) \\ & - \mathcal{J}(Z_D, Z_K^{\text{L}}) - \mathcal{J}(Z_K^{\text{L}}, Y_D) + \mathcal{J}(\mathbf{d}_D) \\ & - \mathcal{J}(X_B, W_K^{\text{R}}) - \mathcal{J}(W_K^{\text{R}}, W_B) + \mathcal{J}(\mathbf{b}_B) \end{aligned}$$

which holds because the four points $Z_D, Z_K^{\text{L}}, W_K^{\text{R}}, W_B$ are colinear with O , the three points $Z_K^{\text{L}}, \mathbf{x}_K^{\text{L}}, Y_D$ are colinear, and the three points $W_K^{\text{R}}, \mathbf{x}_K^{\text{R}}, X_B$ are colinear. □

Recall the Definition 7.1.7 that we write $f(K) \equiv_K g(K)$ for functionals f, g on cap K if the difference $f(K) - g(K)$ is linear in K .

Lemma 8.3.5. For any $K \in \mathcal{K}^i$, we have

$$|K| \equiv_K \mathcal{J} \left(\mathbf{u}_K^{0, \varphi^{\text{R}}} \right) + \mathcal{J} \left(\mathbf{u}_K^{\varphi^{\text{R}}, \varphi^{\text{L}}} \right) + \mathcal{J} \left(\mathbf{u}_K^{\varphi^{\text{L}}, \pi/2} \right) + \mathcal{J} \left(\mathbf{u}_K^{\pi/2, \pi} \right).$$

Proof. We first break the value $|K|$ into a sum of convex curve area functionals. By Theorem 7.1.3, we have $|K| = \frac{1}{2} \int_{t \in S^1} h_K(t) \sigma_K(dt)$. As K is a cap with rotation angle $\pi/2$, the measure σ_K is zero on the set $(\pi, 2\pi) \setminus \{3\pi/2\}$ and $h_K(3\pi/2) = 0$. As $K \in \mathcal{K}^i$, we have $\sigma_K(0) = \sigma_K(\pi) = 0$ too by (1) of Definition 6.1.2. So

$$|K| = \frac{1}{2} \int_{t \in (0, \pi)} h_K(t) \sigma_K(dt).$$

This with Theorem 7.3.2 implies $|K| = \mathcal{J}(\mathbf{u}_K^{0, \pi})$. Now use Lemma 7.3.4 multiple times to obtain

$$|K| = \mathcal{J}(\mathbf{u}_K^{0, \varphi^R}) + \mathcal{J}(\mathbf{u}_K^{\varphi^R, \varphi^L}) + \mathcal{J}(\mathbf{u}_K^{\varphi^L, \pi/2}) + \mathcal{J}(A_K(\pi/2), C_K(0)) + \mathcal{J}(\mathbf{u}_K^{\pi/2, \pi}).$$

Here we use that each of $e_K(\varphi^R)$ and $e_K(\varphi^L)$ is a single point, because $K \in \mathcal{K}^i$ and so (1) of Definition 6.1.2 holds. The expression $\mathcal{J}(A_K(\pi/2), C_K(0))$ is equal to $\sigma_K(\{\pi/2\})/2$ by Proposition 2.1.2 and Proposition 7.2.4, so is linear in K . This completes the proof. \square

Lemma 8.3.6. *For any $K \in \mathcal{K}^i$, we have the followings.*

1. $\mathcal{J}(\mathbf{y}_K|_{[\varphi^R, \varphi^L]}) \equiv_K \mathcal{J}(\mathbf{x}_K|_{[\varphi^R, \varphi^L]})$
2. $\mathcal{J}(\mathbf{I}_K^{\pi/2}(\varphi^R), \mathbf{y}_K(\varphi^R)) \equiv_K \mathcal{J}(W_K^R, \mathbf{x}_K^R)$
3. $\mathcal{J}(\mathbf{I}_K^{\pi/2+\varphi^L}(\varphi^L), \mathbf{y}_K(\varphi^L)) \equiv_K \mathcal{J}(Z_K^L, \mathbf{x}_K^L)$

Proof. For (1), we have $\mathbf{y}_K(t) = \mathbf{x}_K(t) + u_t + v_t$. So with $I := [\varphi^R, \varphi^L]$ and $c_t := u_t + v_t$, we have

$$\begin{aligned} \mathcal{J}(\mathbf{y}_K|_I) &= \frac{1}{2} \int_{\varphi^R}^{\varphi^L} \mathbf{y}_K(t) \times d\mathbf{y}_K(t) \\ &= \frac{1}{2} \int_{\varphi^R}^{\varphi^L} (\mathbf{x}_K(t) + c_t) \times d(\mathbf{x}_K(t) + c_t) \\ &= \mathcal{J}(\mathbf{x}_K) + \frac{1}{2} \left(\int_0^\omega c_t \times d\mathbf{x}_K(t) + \int_0^\omega \mathbf{x}_K(t) \times dc_t + \int_0^\omega c_t \times dc_t \right) \end{aligned}$$

and the term in large bracket is convex-linear in K .

For (2), observe that $\mathbf{y}_K(\varphi^R) - \mathbf{x}_K^R = u_{\varphi^R} + v_{\varphi^R}$ is a constant c_1 independent of K . Likewise, the points $\mathbf{I}_K^{\pi/2}(\varphi^R)$ and W_K^R are the vertices of the parallelogram P_K^R , so their difference is a constant $c_2 := (\sec(\varphi^R)(1 - \sin(\varphi^R)), 1)$ independent of K . Now $\mathcal{J}(W_K^R, \mathbf{x}_K^R) \equiv_K \mathcal{J}(W_K^R + c_2, \mathbf{x}_K^R + c_1)$ by bilinearity of \mathcal{J} . Proof of (3) is similar as (2). \square

Lemma 8.3.7. *For any $K \in \mathcal{K}^i$, we have $\mathcal{S}_K \equiv_K -\mathcal{P}_K$.*

Proof. Expand each term $\mathcal{M}_K(-)$ in the Definition 8.3.2 of \mathcal{S}_K using Definition 7.4.2. Then \mathcal{S}_K is equal to the sum of all terms in the matrix below; each row sums up to a single

expression of form $\mathcal{M}_K(-)$. It is easiest to verify this by following the bold boundaries of four Mamikon regions (colored grey) in the upper part of Figure 8.2 from right to left.

$$\begin{array}{llll}
\mathcal{J}\left(A_K(0), \mathbf{1}_K^{\pi/2}(0)\right) & \mathcal{J}\left(\mathbf{1}_K^{\pi/2}(0), \mathbf{1}_K^{\pi/2}(\varphi^R)\right) & \mathcal{J}\left(\mathbf{1}_K^{\pi/2}(\varphi^R), A_K(\varphi^R)\right) & -\mathcal{J}\left(\mathbf{u}_K^{0, \varphi^R}\right) \\
\mathcal{J}\left(A_K(\varphi^R), \mathbf{y}_K(\varphi^R)\right) & \mathcal{J}\left(\mathbf{y}_K|_{[\varphi^R, \varphi^L]}\right) & \mathcal{J}\left(\mathbf{y}_K(\varphi^L), A_K(\varphi^L)\right) & -\mathcal{J}\left(\mathbf{u}_K^{\varphi^R, \varphi^L}\right) \\
\mathcal{J}\left(A_K(\varphi^L), \mathbf{y}_K(\varphi^L)\right) & \mathcal{J}\left(\mathbf{y}_K(\varphi^L), \mathbf{1}_K^{\pi/2+\varphi^L}(\pi/2)\right) & \mathcal{J}\left(\mathbf{1}_K^{\pi/2+\varphi^L}(\pi/2), A_K(\pi/2)\right) & -\mathcal{J}\left(\mathbf{u}_K^{\varphi^L, \pi/2}\right) \\
\mathcal{J}\left(C_K(0), \mathbf{1}_K^\pi(\pi/2)\right) & \mathcal{J}\left(\mathbf{1}_K^\pi(\pi/2), C_K(\pi/2)\right) & & -\mathcal{J}\left(\mathbf{u}_K^{\pi/2, \pi}\right)
\end{array}$$

Call the term in i 'th row and j 'th column, including the signs, as simply J_{ij} . Now check the following calculations.

- $\sum_{i=1}^4 J_{i4} \equiv_K -|K|$ by Lemma 8.3.5.
- $J_{11} = h_K(0)/2 \equiv_K 0$ by (1) of Theorem 7.1.2.
- $J_{12} = \left(\mathbf{1}_K^{\pi/2}(0) - \mathbf{1}_K^{\pi/2}(\varphi^R)\right) \cdot u_0 \equiv_K 0$ by (2) of Theorem 7.1.2.
- $J_{13} + J_{21} = \mathcal{J}\left(\mathbf{1}_K^{\pi/2}(\varphi^R), \mathbf{y}_K(\varphi^R)\right) \equiv_K \mathcal{J}\left(W_K^R, \mathbf{x}_K^R\right)$ by (2) of Lemma 8.3.6.
- $J_{22} = \mathcal{J}\left(\mathbf{y}_K|_{[\varphi^R, \varphi^L]}\right) \equiv_K \mathcal{J}\left(\mathbf{x}_K|_{[\varphi^R, \varphi^L]}\right)$ by (1) of Lemma 8.3.6.
- $J_{23} + J_{31} = 0$.
- $J_{32} \equiv_K \mathcal{J}\left(\mathbf{x}_K^L, Z_K^L\right)$ by (3) of Lemma 8.3.6.
- $J_{33} \equiv_K J_{41} \equiv_K 0$ by (2) of Theorem 7.1.2 and that the points are on $l_K(\pi/2) = l(\pi/2, 1)$.
- $J_{42} = h_K(\pi/2)/2 \equiv_K 0$ by (1) of Theorem 7.1.2.

Add all the calculations in the list above to conclude $\mathcal{S}_K \equiv_K -\mathcal{P}_K$. □

We finally assemble all the lemmas to prove the concavity of \mathcal{Q} .

Theorem 8.3.8. *The functional $\mathcal{Q} : \mathcal{L} \rightarrow \mathbb{R}$ is concave.*

Proof. We need to show that the value $\mathcal{Q}(K, B, D)$ is quadratic and concave on $(K, B, D) \in \mathcal{L}$. By Lemma 8.3.4 and Lemma 8.3.7, we have

$$\mathcal{Q}(K, B, D) = \mathcal{P}_K - \mathcal{R}_B - \mathcal{L}_D \equiv_K -\mathcal{S}_K - \mathcal{R}_B - \mathcal{L}_D.$$

By Lemma 8.3.3, the right-hand side is quadratic and concave on $(K, B, D) \in \mathcal{L}$. □

8.4 Gerver's Sofa

In this Section 8.4, we extract the properties of Gerver's sofa G we need for proving that G is a global optimum. We follow the derivation of G by Romik in Section 4 of [Rom18].




Figure 8.3: Gerver's sofa G is determined by five oriented curves \mathbf{A} , \mathbf{B} , \mathbf{C} , \mathbf{D} , and \mathbf{x} solved in Section 4 of [Rom18].

Boundary Curves of G

In Romik's derivation, Gerver's sofa G is a monotone sofa with the boundary parametrized by five oriented curves \mathbf{A} , \mathbf{B} , \mathbf{C} , \mathbf{D} , and \mathbf{x} (see Figure 8.3).

The five curves are parametrized differently in each of the five intervals I_1, \dots, I_5 in the following Definition 8.4.1.

Definition 8.4.1. Define

$$(t_0, t_1, t_2, t_3, t_4, t_5) := (0, \varphi, \theta, \pi/2 - \theta, \pi/2 - \varphi, \pi/2)$$

so that $0 = t_0 < t_1 < \dots < t_5 = \pi/2$ forms a partition of the interval $[0, \pi/2]$. For each $1 \leq i \leq 5$, define the interval $I_i := [t_{i-1}, t_i]$.

Definition 8.4.2. Define $\mathbf{x} : [0, \pi/2] \rightarrow \mathbb{R}^2$ as the continuously differentiable function satisfying Equations 25 to 44 of [Rom18] as solved in the Section 4 of [Rom18]. In particular, by Equation 25 of [Rom18], for every $1 \leq i \leq 5$ the restriction $\mathbf{x}_i := \mathbf{x}|_{I_i}$ at the i 'th interval I_i is smooth and satisfies the i 'th ODE in Theorem 2 in [Rom18].

Definition 8.4.3. Define the continuous, piecewise smooth curves $\mathbf{A} : [0, \pi/2] \rightarrow \mathbb{R}^2$, $\mathbf{B} : [t_3, t_5] \rightarrow \mathbb{R}^2$, $\mathbf{C} : [0, \pi/2] \rightarrow \mathbb{R}^2$, and $\mathbf{D} : [t_0, t_2] \rightarrow \mathbb{R}^2$ determined by \mathbf{x} in Definition 8.4.2 according to Equations 9 to 12 of [Rom18] respectively.

The boundary of G is traced out by the four curves \mathbf{A} , \mathbf{B} , \mathbf{C} , \mathbf{D} in their full domains and the curve \mathbf{x} restricted to the interval $[t_1, t_4] = [\varphi^R, \varphi^L]$ (see Figure 8.3 and the table below Theorem 8.4.2).

Theorem 8.4.1. Gerver's sofa G is a monotone sofa. Let $K := \mathcal{C}(G)$ be the cap of G . Then the followings are true.

1. The cap K have vertices $A_K(t) = \mathbf{A}(t)$ and $C_K(t) = \mathbf{C}(t)$ and the inner corner $\mathbf{x}_K(t) = \mathbf{x}(t)$ over $t \in [0, \pi/2]$.
2. The niche $\mathcal{N}(K)$ is the region enclosed counterclockwise by the following curves concatenated in order.
 1. The curve $\mathbf{B} : [t_3, t_5] \rightarrow \mathbb{R}^2$ reversed in direction.
 2. The curve \mathbf{x} restricted to $[t_1, t_4] \rightarrow \mathbb{R}^2$.

3. The curve $\mathbf{D} : [t_0, t_2] \rightarrow \mathbb{R}^2$ reversed in direction.
 4. The horizontal line segment on $l(\pi/2, 0)$ from $\mathbf{D}(0)$ to $\mathbf{B}(\pi/2)$.
3. The half-line inner wall $\vec{b}_K(t)$ (resp. $\vec{d}_K(t)$) of the supporting hallway $L_K(t)$ passes through the point $\mathbf{B}(t)$ in the domain $t \in [t_3, t_5]$ of \mathbf{B} (resp. $\mathbf{D}(t)$ in the domain $t \in [t_0, t_2]$ of \mathbf{D}).
 4. The tangent direction $\mathbf{B}'(t)$ of \mathbf{B} is parallel to v_t , and $\mathbf{B}'(t) \cdot v_t < 0$ on the domain $t \in [t_3, t_5]$ of \mathbf{B} . The tangent direction $\mathbf{D}'(t)$ of \mathbf{D} is parallel to u_t , and $\mathbf{D}'(t) \cdot u_t > 0$ on the domain $t \in [t_0, t_2]$ of \mathbf{D} .

Remark 8.4.1. Theorem 8.4.1 summarizes the properties of Gerver's sofa G as described in the remark at the end of [Ger92] and recasts them our terminology. Rigorously speaking, these properties must be established logically *before* asserting that G constitutes a valid moving sofa with five specific stages of movement as claimed in existing works on G [Ger92; Rom18].

The properties are easy to verify numerically and implicitly assumed in [Ger92; Rom18]. Their truth is not entirely self-evident however. Indeed, Gerver observes a subtlety of the construction of G in his remark, that the endpoints $\mathbf{x}(\varphi^R)$ and $\mathbf{x}(\varphi^L)$ of the 'core' of the niche of G come within a distance of only 0.0012 from the inner walls $b_K(t)$ and $d_K(t)$ of the hallway L at $t = \pi/4$. In other words, the construction of G comes within a distance of 0.0012 of 'breaking' near the hallway L_t with $t = \pi/4$. With this, a rigorous symbolic verification of Theorem 8.4.1 would still be worthy.


ODEs of Boundary Curves of G

We recall the concept of *contact points* from [Rom18]. Fix an interval $I_i = [t_{i-1}, t_i]$ and let $t \in I_i$ parametrize I_i . Then Gerver's sofa G with the cap $K := \mathcal{C}(G)$ makes contact with the supporting hallway $L_K(t)$ at the point $\mathbf{A}(t)$ (resp. $\mathbf{B}(t)$, $\mathbf{C}(t)$, $\mathbf{D}(t)$, and $\mathbf{x}(t)$) if and only if the interval I_i is contained in the domain of \mathbf{A} (resp. the domain of \mathbf{B} , \mathbf{C} , \mathbf{D} , or the restricted domain $[t_1, t_4]$ of \mathbf{x}).

As this set of *contact points* $\Gamma(t)$, a subset of $\{\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{x}\}$, that G makes with $L_K(t)$ is determined, a corresponding ODE that balances the differential side lengths on those contact points can be derived. For example, the Equation (1.8) is derived in Section 1.7.2 by assuming $\Gamma(t) = \{\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{x}\}$. Each interval $t \in I_i$ have a corresponding contact points $\Gamma(t)$ fixed in $t \in I_i$ and their respective ODE as follows.

Definition 8.4.4. Denote the dot product $a \cdot b$ of $a, b \in \mathbb{R}^2$ as also $\langle a, b \rangle$.

Theorem 8.4.2. For each $t \in I_i$, the ODEs involving \mathbf{A} , \mathbf{B} , \mathbf{C} , \mathbf{D} , and \mathbf{x} in the last column of the following table holds.

	Interval Contacts	Figure	Equation (Numbering in [Rom18])
(1)-v	$t \in I_1$ $\mathbf{A}, \mathbf{C}, \mathbf{D}$		$\langle \mathbf{A}'(t), v_t \rangle = 0$ (21)

	Interval Contacts	Figure	Equation (Numbering in [Rom18])
(1)-u	$t \in I_1$ A, C, D		$\langle -\mathbf{C}'(t), u_t \rangle = \langle \mathbf{D}'(t), u_t \rangle$ (22)
(2)-v	$t \in I_2$ A, C, D, x		$\langle \mathbf{A}'(t), v_t \rangle = \langle \mathbf{x}'(t), v_t \rangle$ (17)
(2)-u	$t \in I_2$ A, C, D, x		$\langle -\mathbf{C}'(t), u_t \rangle = \langle \mathbf{D}'(t) - \mathbf{x}'(t), u_t \rangle$ (19)
(3)-v	$t \in I_3$ A, C, x		$\langle \mathbf{A}'(t), v_t \rangle = \langle \mathbf{x}'(t), v_t \rangle$ (17)
(3)-u	$t \in I_3$ A, C, x		$\langle -\mathbf{C}'(t), u_t \rangle = \langle -\mathbf{x}'(t), u_t \rangle$ (18)
(4)-v	$t \in I_4$ A, B, C, x		$\langle \mathbf{A}'(t), v_t \rangle = \langle -\mathbf{B}'(t) + \mathbf{x}'(t), v_t \rangle$ (20)
(4)-u	$t \in I_4$ A, B, C, x		$\langle -\mathbf{C}'(t), u_t \rangle = \langle -\mathbf{x}'(t), u_t \rangle$ (18)
(5)-v	$t \in I_5$ A, B, C		$\langle \mathbf{A}'(t), v_t \rangle = \langle -\mathbf{B}'(t), v_t \rangle$ (22) ¹
(5)-u	$t \in I_5$ A, B, C		$\langle -\mathbf{C}'(t), u_t \rangle = 0$ (21) ²

Proof. They are exactly the equations with respective numberings in the proof of Theorem 2 of [Rom18]. □

¹[Rom18] omits exact equations for the interval I_5 as the cases I_1 and I_5 are symmetric to each other.
²See above.

Left, Middle, and Right Parts of G

Theorem 8.4.3. *Let $K := \mathcal{C}(G)$ be the cap of Gerver's sofa. Then the followings are true.*

1. We have $\mathbf{D}(t) = v_{D_K}^\pm(3\pi/2 + t)$ on $t \in (t_0, t_2)$. Likewise, we have $\mathbf{B}(t) = v_{B_K}^\pm(\pi + t)$ on $t \in (t_3, t_5)$.
2. $\mathbf{x}_K^L = Y_{D_K} = \mathbf{D}(t_2) = v_{D_K}^\pm(3\pi/2 + t_2)$ and $\mathbf{d}_{D_K} = \mathbf{D}$ as oriented curves. Likewise, $\mathbf{x}_K^R = X_{B_K} = \mathbf{D}(t_3) = v_{B_K}^\pm(\pi + t_3)$ and $\mathbf{b}_{B_K} = \mathbf{B}$ as oriented curves.
3. We have $h_K(\pi/2 + t) + h_{D_K}(3\pi/2 + t) = 1$ on $t \in [t_0, t_2]$, and $h_K(t) + h_{B_K}(\pi + t) = 1$ on $t \in [t_3, t_5]$.

Proof. Recall that $K \in \mathcal{K}^i$ by Theorem 6.1.2.

We first show (1). First, observe that \mathbf{D} is in the closure of $\mathcal{N}(K)$ by (2) of Theorem 8.4.1, so we have $\mathbf{D} \subseteq K$ by Theorem 2.5.9. By (2) of Theorem 8.4.1, we have the matching endpoints $\mathbf{D}(t_2) = \mathbf{x}_K^L$ of \mathbf{D} and \mathbf{x} , so $\mathbf{D}(t_2) \in d_K^L$. Then by (4) of Theorem 8.4.1, we have $\mathbf{D} \subseteq \check{H}_K^L$. Now \mathbf{D} is a subset of $K \cap \check{H}_K^L$. Second, the curve \mathbf{D} itself is disjoint from $\mathcal{N}(K)$ by (2) of Theorem 8.4.1, so by (2) of Lemma 8.1.6, \mathbf{D} is contained in all $H_K^d(t)$ over all $t \in [0, \varphi^L)$. So we have $\mathbf{D} \subseteq D_K$ by the definition of D_K . Finally, as $\mathbf{D}(t) \in D_K \cap d_K(t)$ for all $t \in [t_0, t_2]$ by (3) of Theorem 8.4.1, and $D_K \subseteq H_K^d(t)$ by definition, the line $d_K(t)$ should be a supporting line of D_K and $\mathbf{D}(t) \in e_{D_K}(3\pi/2 + t)$ over all $t \in [t_0, t_2]$. For each $t \in (t_0, t_2)$, use the continuity of $\mathbf{D}(t) \in e_{D_K}(3\pi/2 + t)$ at t , and take the left and right limit on t , then use Theorem 2.1.3 to conclude $\mathbf{D}(t) = v_{D_K}^\pm(3\pi/2 + t)$. Use a mirror-symmetric argument to prove the coresponding statement of (1) on \mathbf{B} .

We now show (2) using (1). Recall the definitions $\mathbf{d}_{D_K} = \mathbf{u}_{D_K}^{3\pi/2, 3\pi/2 + \varphi^L}$ and $Y_{D_K} = v_{D_K}^-(3\pi/2 + \varphi^L)$. By taking the limits $t \rightarrow t_0^+$ and $t \rightarrow t_2^-$ to $\mathbf{D}(t) = v_{D_K}^\pm(3\pi/2 + t)$ of (1) and using Theorem 2.1.3, we have $\mathbf{D} = \mathbf{u}_{D_K}^{3\pi/2, 3\pi/2 + t_2}$ as oriented curves and $\mathbf{D}(t_2) = v_{D_K}^-(3\pi/2 + t_2)$. We have $\mathbf{D}(t_2) = \mathbf{x}_K^L$ by (2) of Theorem 8.4.1, and the point is on the supporting line $d_K^L = l_{D_K}(3\pi/2 + \varphi^L)$ of D_K by (4) of Lemma 8.1.7. So we have

$$v_{D_K}^-(3\pi/2 + t_2) \in l_{D_K}(3\pi/2 + \varphi^L)$$

and this implies the degeneracy

$$v_{D_K}^-(3\pi/2 + \varphi^L) = v_{D_K}^\pm(3\pi/2 + t_2)$$

and that $\mathbf{u}_{D_K}^{3\pi/2 + t_2, 3\pi/2 + \varphi^L}$ is a single point. Now the statement of (2) for the curve \mathbf{D} and convex body D_K holds. Use a mirror-symmetric argument to prove the corresponding statement of (2) on \mathbf{B} and B_K .

We show (3) using (1) and (2). Let $t \in (t_0, t_2]$ be arbitrary. We have $\mathbf{C}(t) \in c_K(t)$ and $\mathbf{D}(t) \in d_K(t)$ by (3) of Theorem 8.4.1, and the lines are parallel and of distance one. We also have $\mathbf{C}(t) = C_K(t)$ by (1) of Theorem 8.4.1 and $\mathbf{D}(t) = v_{D_K}^\pm(3\pi/2 + t)$ by (1) and (2). So the first equality of (3) holds. Take limits to show the equality for $t = t_0$ too. Again, use a mirror-symmetric argument to prove the second equality of (3) using (1). \square

Surface Area Measures of G

Our next goal is to translate the ODEs of Romik in Theorem 8.4.2 in terms of surface area measure as differential side lengths. We will use the following Definition 8.4.5 for $C = B, D$ to denote the surface area measure and support function of the bottom sides \mathbf{d}_D and \mathbf{b}_B .

Definition 8.4.5. For any convex body C , let $\check{\sigma}_C$ denote the measure on S^1 such that $\check{\sigma}_C(X) = \sigma_C(X + \pi)$ for any Borel subset $X \subseteq S^1$. Likewise, let $\check{h}_C(t) = h_C(t + \pi)$.

Proposition 8.4.4. Let $K := \mathcal{C}(G)$ be the cap of Gerver's sofa. Let $B := B_K$ and $D := D_K$. Then the followings are true.

1. $\langle \mathbf{A}'(t), v_t \rangle dt = \sigma_K$ as measures on $t \in [0, \pi/2]$.
2. $\langle -\mathbf{B}'(t), v_t \rangle dt = \check{\sigma}_B$ as measures on $t \in [t_3, t_5]$.
3. $\langle -\mathbf{C}'(t - \pi/2), u_{t - \pi/2} \rangle dt = \sigma_K$ as measures on $t \in (\pi/2, \pi]$.
4. $\langle \mathbf{D}'(t), u_t \rangle dt = \check{\sigma}_D$ as measures on $t \in (t_0, t_2]$.

Proof. (1) and (3) for $t \in (0, \pi/2)$ and $t \in (\pi/2, \pi)$ respectively follow from (1) of Theorem 8.4.1 and Theorem 5.2.2. (1) and (3) on the singletons $\{0\}$ and $\{\pi\}$ follow from Theorem 6.1.2 and (1) of Definition 6.1.2. (2) and (4) follow from (1) and (2) of Theorem 8.4.3 and Theorem 5.2.2. \square

We also introduce a measure ι_K that captures the differential side lengths contributed by the inner corner \mathbf{x}_K of cap K .

Definition 8.4.6. For any cap $K \in \mathcal{K}^i$, define the function $i_K : (0, \pi] \rightarrow \mathbb{R}$ as $i_K(t) := \langle \mathbf{x}'_K(t), v_t \rangle$ and $i_K(t + \pi/2) := \langle -\mathbf{x}'_K(t), u_t \rangle$ for every $t \in (0, \pi/2]$. Define ι_K as the measure on $[0, \pi]$ derived from the density function i_K . That is, $\iota_K(dt) = i_K(t)dt$ and $\iota_K(\{0\}) = 0$ in particular.

Now we translate the equations in Theorem 8.4.2 to the equations of aforementioned measures.

Definition 8.4.7. For $1 \leq i \leq 5$, define $J_i := [t_{i-1}, t_i)$ where the values t_i are as in Definition 8.4.1. For $6 \leq i \leq 10$, define $J_i := \pi - J_{11-i}$.

The intervals J_1, \dots, J_{10} and the singleton $\{\pi/2\}$ partition the whole interval $[0, \pi]$.

Theorem 8.4.5. Let $K := \mathcal{C}(G)$ be the cap of Gerver's sofa, and let $B := B_K$ and $D := D_K$.

1. $\sigma_K = 0$ on J_1 .
2. $\sigma_K = \iota_K$ on $J_2 \cup J_3$.
3. $\sigma_K = \check{\sigma}_B + \iota_K$ on J_4 .
4. $\sigma_K = \check{\sigma}_B$ on J_5 .
5. $\sigma_K = \check{\sigma}_D$ on J_6 .
6. $\sigma_K = \check{\sigma}_D + \iota_K$ on J_7 .
7. $\sigma_K = \iota_K$ on $J_8 \cup J_9$.
8. $\sigma_K = 0$ on J_{10} .

Proof. Let $1 \leq i \leq 5$ be arbitrary. For the interval J_i (resp. J_{i+5}), convert the Equation (i)-v (resp. Equation (i)-u) of Theorem 8.4.2 on the interval $t \in I_i$ and direction v_t (resp. u_t) using (1) of Theorem 8.4.1, Definition 8.4.6, and Proposition 8.4.4. Check the endpoints carefully. \square

Theorem 8.4.6. *Let $K := \mathcal{C}(G)$ be the cap of Gerver's sofa. Then $\mathcal{A}(K) = \mathcal{Q}(K, B_K, D_K)$.*

Proof. Write $B := B_K$ and $D := D_K$. Start from the expression of $\mathcal{Q}(K, B, D)$ in Definition 8.2.2. By (2) of Theorem 8.4.3, we have $\mathcal{J}(\mathbf{d}_D) = \mathcal{J}(\mathbf{D})$ and $\mathcal{J}(\mathbf{b}_B) = \mathcal{J}(\mathbf{B})$. By (2) of Theorem 8.4.3, we have $\mathcal{J}(Y_D, \mathbf{x}_K^L) = \mathcal{J}(\mathbf{x}_K^R, X_B) = 0$. By integrating the Jordan curve boundary of $\mathcal{N}(K)$ as in (2) of Theorem 8.4.1, we have $|\mathcal{N}(K)| = \mathcal{J}(\mathbf{x}|_{[\varphi^R, \varphi^L]}) - \mathcal{J}(\mathbf{B}) - \mathcal{J}(\mathbf{D})$. So the value of $\mathcal{Q}(K, B, D)$ in Definition 8.2.2 reduces to $|K| - |\mathcal{N}(K)| = \mathcal{A}(K)$. \square

8.5 Directional Derivative of \mathcal{Q}

We calculate the directional derivative $D\mathcal{Q}$ of \mathcal{Q} at the point $(K, B_K, D_K) \in \mathcal{L}$ corresponding to the cap $K := \mathcal{C}(G)$ of Gerver's sofa G . The value turns out to be non-positive (Theorem 8.5.7). Since \mathcal{Q} is concave and quadratic (Theorem 8.3.8), the value $(K, B_K, D_K) \in \mathcal{L}$ is the maximizer of \mathcal{Q} (Theorem 7.1.5).

We first prepare the theorems that calculate the directional derivative on each curve of \mathcal{Q} .

Theorem 8.5.1. *The area $|K|$ of a cap $K \in \mathcal{K}^i$ is quadratic as a functional on \mathcal{K}^i . Moreover, the directional derivative of $f(K) := |K|$ from K to K^* evaluates to*

$$Df(K; K^*) := \int_{t \in [0, \pi]} (h_{K^*}(t) - h_K(t)) \sigma_K(dt).$$

Proof. Recall that the mixed volume $V(K_1, K_2)$ on convex bodies $K_1, K_2 \in \mathcal{K}$ is a symmetric bilinear functional on \mathcal{K} such that $|K| = V(K, K)$ (e.g. Chapter 5 of [Sch13]). So by Lemma 7.1.4, we have

$$Df(K; K^*) = 2V(K^*, K) - 2V(K, K).$$

We use the formula $V(K_1, K_2) = \frac{1}{2} \int_{t \in S^1} h_{K_1}(t) \sigma_{K_2}(dt)$ of mixed volume (Equation (5.19) of [Sch13]). Substitute this in the equation above, and use that for caps $K, K^* \in \mathcal{K}^i$ we have $\sigma_K = \sigma_{K^*} = 0$ on $t \in (\pi, 2\pi) \setminus \{3\pi/2\}$ and $h_K(3\pi/2) = h_{K^*}(3\pi/2) = 0$. \square

Theorem 8.5.2. *Fix $a, b \in \mathbb{R}$ such that $a < b < a + \pi$. Define $f(K) := \mathcal{J}(\mathbf{u}_K^{a,b})$ as a functional on $K \in \mathcal{K}$. Then f is quadratic on \mathcal{K} and its directional derivative from K to K^* evaluates to*

$$Df(K; K^*) = \int_{t \in (a,b)} (h_{K^*}(t) - h_K(t)) \sigma_K(dt) + [\mathcal{J}(v_K^-(b), v_{K^*}^-(b)) - \mathcal{J}(v_K^+(a), v_{K^*}^+(a))].$$

Proof. Write $f(K) = \mathcal{B}(K, K)$ where $\mathcal{B}(K_1, K_2) := \frac{1}{2} \int_{t \in (a,b)} h_{K_1}(t) \sigma_{K_2}(dt)$ is defined as in Lemma 7.3.3. By integration by parts (Lemma 5.1.2) and that the left limit of $v_{K_1}^+$ at t is $v_{K_1}^-(t)$ (Theorem 2.1.3), we get

$$\int_{t \in (a,b)} dv_{K_1}^+(t) \times v_{K_2}^+(t) + \int_{t \in (a,b)} v_{K_1}^-(t) \times dv_{K_2}^+(t) = v_{K_1}^-(b) \times v_{K_2}^-(b) - v_{K_1}^+(a) \times v_{K_2}^+(a).$$

With Lemma 7.3.3 and the anticommutativity $a \times b = -b \times a$ of cross product, the equality becomes

$$\mathcal{B}(K_1, K_2) - \mathcal{B}(K_2, K_1) = \mathcal{J}(v_{K_1}^-(b), v_{K_2}^-(b)) - \mathcal{J}(v_{K_1}^+(a), v_{K_2}^+(a))$$

after dividing by two. Now let $K_1 := K$ and $K_2 := K^*$ then use Lemma 7.1.4 on $f(K) = \mathcal{B}(K, K)$ to get

$$\begin{aligned} Df(K; K^*) &= \mathcal{B}(K, K^*) + \mathcal{B}(K^*, K) - 2\mathcal{B}(K, K) \\ &= 2\mathcal{B}(K^*, K) - 2\mathcal{B}(K, K) + \mathcal{J}(v_K^-(b), v_{K^*}^-(b)) - \mathcal{J}(v_K^+(a), v_{K^*}^+(a)) \end{aligned}$$

and use the definition of \mathcal{B} . □

Theorem 8.5.3. *Treat $\mathbb{R}^2 \times \mathbb{R}^2$ as a convex domain. Then the curve area functional $\mathcal{J}(p, q)$ on $(p, q) \in \mathbb{R}^2 \times \mathbb{R}^2$ is quadratic. For any $(p, q), (p^*, q^*) \in \mathbb{R}^2 \times \mathbb{R}^2$, the directional derivative of $\mathcal{J}(-, -)$ evaluates to*

$$D\mathcal{J}(p, q; p^*, q^*) = \frac{1}{2} ((p^* + q^*) \times (q - p) - 2(p \times q)) + [\mathcal{J}(q, q^*) - \mathcal{J}(p, p^*)]$$

where the first term in the round bracket $(-)$ equates to zero when $p = q$.

Proof. Define the bilinear form $\mathcal{B}((p_1, q_1), (p_2, q_2)) = p_1 \times q_2 + p_2 \times q_1$ on $(p_1, q_1), (p_2, q_2) \in \mathbb{R}^2 \times \mathbb{R}^2$. Apply Lemma 7.1.4 to $4\mathcal{J}(p, q) = \mathcal{B}((p, q), (p, q))$ and use that \mathcal{B} is symmetric to obtain

$$\begin{aligned} 2D\mathcal{J}(p, q; p^*, q^*) &= \mathcal{B}((p^*, q^*), (p, q)) - \mathcal{B}((p, q), (p, q)) \\ &= p^* \times q + p \times q^* - 2(p \times q) \\ &= [(p^* + q^*) \times (q - p) - 2(p \times q)] + q \times q^* - p \times p^* \end{aligned}$$

and divide by two to conclude the proof. □

Theorem 8.5.4. *The curve area functional $\mathcal{J}(\mathbf{x})$ on $\mathbf{x} \in C^{\text{BV}}[a, b]$ is quadratic. Moreover, the directional derivative of \mathcal{J} from \mathbf{x} to \mathbf{x}^* evaluates to*

$$D\mathcal{J}(\mathbf{x}; \mathbf{x}^*) = \int_a^b (\mathbf{x}^*(t) - \mathbf{x}(t)) \times d\mathbf{x}(t) + [\mathcal{J}(\mathbf{x}(b), \mathbf{x}^*(b)) - \mathcal{J}(\mathbf{x}(a), \mathbf{x}^*(a))].$$

Proof. Consider the bilinear form $\mathcal{B}(\mathbf{x}_1, \mathbf{x}_2) = \int_a^b \mathbf{x}_1(t) \times d\mathbf{x}_2(t)$ on $\mathbf{x}_1, \mathbf{x}_2 \in C^{\text{BV}}[a, b]$. Apply Lemma 7.1.4 to $2\mathcal{J}(\mathbf{x}) = \mathcal{B}(\mathbf{x}, \mathbf{x})$ to get

$$2D\mathcal{J}(\mathbf{x}; \mathbf{x}^*) = \mathcal{B}(\mathbf{x}, \mathbf{x}^*) + \mathcal{B}(\mathbf{x}^*, \mathbf{x}) - 2\mathcal{B}(\mathbf{x}, \mathbf{x}). \quad (8.1)$$

With integration by parts (Lemma 5.1.2), we get

$$\int_a^b \mathbf{x}(t) \times d\mathbf{x}^*(t) = \mathbf{x}(b) \times \mathbf{x}^*(b) - \mathbf{x}(a) \times \mathbf{x}^*(a) + \int_a^b \mathbf{x}^*(t) \times d\mathbf{x}(t)$$

or

$$\mathcal{B}(\mathbf{x}, \mathbf{x}^*) = 2\mathcal{J}(\mathbf{x}(b), \mathbf{x}^*(b)) - 2\mathcal{J}(\mathbf{x}(a), \mathbf{x}^*(a)) + \mathcal{B}(\mathbf{x}^*, \mathbf{x}).$$

Plug this back in Equation (8.1) and rearrange the terms to get the desired equality. □

Theorem 8.5.5. *Let $I := [\varphi^R, \varphi^L]$. The value $f(K) := \mathcal{J}(\mathbf{x}_K|_I)$ on caps $K \in \mathcal{K}^i$ is quadratic. Moreover, the directional derivative of $f(K)$ from K to K^* evaluates to*

$$Df(K; K^*) := \langle h_{K^*} - h_K, \iota_K \rangle_{I \cup (I + \pi/2)} + [\mathcal{J}(\mathbf{x}_K^L, \mathbf{x}_{K^*}^L) - \mathcal{J}(\mathbf{x}_K^R, \mathbf{x}_{K^*}^R)].$$

Proof. Apply Theorem 8.5.4 to $\mathbf{x} := \mathbf{x}_K|_I$ and $\mathbf{x}^* := \mathbf{x}_{K^*}|_I$. Then compute

$$\begin{aligned} & \int_{t \in I} (\mathbf{x}^*(t) - \mathbf{x}(t)) \times d\mathbf{x}(t) \\ &= \int_{t \in I} (\mathbf{x}^*(t) - \mathbf{x}(t)) \times (\langle \mathbf{x}'(t), u_t \rangle u_t + \langle \mathbf{x}'(t), v_t \rangle v_t) dt \\ &= \int_{t \in I} \langle \mathbf{x}^*(t) - \mathbf{x}(t), v_t \rangle i_K(\pi/2 + t) dt + \int_{t \in I} \langle \mathbf{x}^*(t) - \mathbf{x}(t), u_t \rangle i_K(t) dt \\ &= \langle h_{K^*} - h_K, \iota_K \rangle_{I \cup (I + \pi/2)} \end{aligned}$$

using the Definition 8.4.6 of i_K and ι_K to finish the proof. \square

Theorem 8.5.6. *Let $I := [\varphi^R, \varphi^L]$. Assume that $(K, B, D) \in \mathcal{L}$ satisfies the constraints $X_B = \mathbf{x}_K^R$ and $Y_D = \mathbf{x}_K^L$. Then the directional derivative of \mathcal{Q} at (K, B, D) in the direction towards arbitrary $(K^*, B^*, D^*) \in \mathcal{L}$ is*

$$\begin{aligned} D\mathcal{Q}(K, B, D; K^*, B^*, D^*) &= \\ & \langle h_{K^*} - h_K, \sigma_K \rangle_{[0, \pi]} - \langle h_{K^*} - h_K, \iota_K \rangle_{I \cup (I + \pi/2)} \\ & + \left\langle \check{h}_{B^*} - \check{h}_B, \check{\sigma}_B \right\rangle_{(\varphi^R, \pi/2)} + \left\langle \check{h}_{D^*} - \check{h}_D, \check{\sigma}_D \right\rangle_{(\pi/2, \pi/2 + \varphi^L)}. \end{aligned}$$

Proof. Apply the following theorems to each term of Definition 8.2.2.

1. Theorem 8.5.1 on $|K|$.
2. Theorem 8.5.2 on $\mathcal{J}(\mathbf{d}_D) = \mathcal{J}(\mathbf{u}_D^{3\pi/2, 3\pi/2 + \varphi^L})$.
3. Theorem 8.5.3 on $\mathcal{J}(Y_D, \mathbf{x}_K^L)$.
4. Theorem 8.5.5 on $-\mathcal{J}(\mathbf{x}_K|_{[\varphi^R, \varphi^L]})$.
5. Theorem 8.5.3 on $\mathcal{J}(\mathbf{x}_K^R, X_B)$.
6. Theorem 8.5.2 on $\mathcal{J}(\mathbf{b}_B) = \mathcal{J}(\mathbf{u}_B^{\pi + \varphi^R, 3\pi/2})$.

Sum up all results from the theorems. The differences appearing in square brackets come from the matching endpoints, so they telescope and zeros in total.³ The remaining terms from (3) and (5) are zero by the assumptions $X_B = \mathbf{x}_K^R$ and $Y_D = \mathbf{x}_K^L$. The remaining terms from (1), (2), (4), and (6) correspond to the right-hand side in the claimed equality. \square

We finally calculate the directional derivative at Gerver's sofa.

³Except for the endpoints W_B, W_{B^*}, Z_D , and Z_{D^*} on the x -axis, but their cross product also zeroes out.

Theorem 8.5.7. *Let $K := \mathcal{C}(G)$ be the cap of Gerber's sofa G . Let $B := B_K$ and $D := D_K$. Take arbitrary $(K^*, B^*, D^*) \in \mathcal{L}$. Then we have*

$$D\mathcal{Q}(K, B, D; K^*, B^*, D^*) \leq 0.$$

Proof. By (2) of Theorem 8.4.3, we can use Theorem 8.5.6 to calculate the directional derivative. Split each term of Theorem 8.5.6 into integrals over each interval J_i in Definition 8.4.7 as the following table. Note that the integrals on singleton $\{\pi/2\}$ is zero because the differences of support functions are zero at $t = \pi/2, 3\pi/2$ ((1) of Definition 2.4.1 and (2) and (4) of Lemma 8.1.7).

	J_1	$J_2 \cup J_3$	J_4	J_5	J_6	J_7	$J_8 \cup J_9$	J_{10}
$\langle h_{K^*} - h_K, \sigma_K \rangle$	0	0	0	0	0	0	0	0
$-\langle h_{K^*} - h_K, \iota_K \rangle$		0	0			0	0	
$\langle \check{h}_{B^*} - \check{h}_B, \check{\sigma}_B \rangle$			0	0				
$\langle \check{h}_{D^*} - \check{h}_D, \check{\sigma}_D \rangle$					0	0		

For each interval, the sum of integrals in Theorem 8.5.6 on the interval J_i is non-positive as follows. For the i 'th interval, we start with the i 'th equality on measures in Theorem 8.4.5.

1. As $\sigma_K = 0$ on J_1 , the sum is zero.
2. As $\sigma_K = \iota_K$ on $J_2 \cup J_3$, the sum is zero.
3. As $\sigma_K = \check{\sigma}_B + \iota_K$ on J_4 , the sum equates to $\langle h_{K^*} - h_K + h_{B^*} - h_B, \check{\sigma}_B \rangle$. We have $h_K + h_B = 1$ on $J_4 \cup J_5$ by (3) of Theorem 8.4.3, and $h_{K^*} + h_{B^*} \leq 1$ on $J_4 \cup J_5$ by (1) of Lemma 8.1.7. This with $\check{\sigma}_B \geq 0$ shows that the sum should be nonnegative.
4. As $\sigma_K = \check{\sigma}_B$ on J_5 , the sum equates to $\langle h_{K^*} - h_K + h_{B^*} - h_B, \check{\sigma}_B \rangle$. Proceed as (3) above.
5. As $\sigma_K = \check{\sigma}_D$ on J_6 , the sum equates to $\langle h_{K^*} - h_K + h_{D^*} - h_D, \check{\sigma}_D \rangle$. Proceed as (6) below.
6. As $\sigma_K = \check{\sigma}_D + \iota_K$ on J_7 , the sum equates to $\langle h_{K^*} - h_K + h_{D^*} - h_D, \check{\sigma}_D \rangle$. We have $h_K + h_D = 1$ on $J_6 \cup J_7$ by (3) of Theorem 8.4.3, and $h_{K^*} + h_{D^*} \leq 1$ on $J_6 \cup J_7$ by (3) of Lemma 8.1.7. This with $\check{\sigma}_D \geq 0$ shows that the sum should be nonnegative.
7. As $\sigma_K = \iota_K$ on $J_8 \cup J_9$, the sum is zero.
8. As $\sigma_K = 0$ on J_{10} , the sum is zero.

Summing up (1)-(8) above, the directional derivative is non-positive as desired. \square

Corollary 8.5.8. *Let $K := \mathcal{C}(G)$ be the cap of Gerber's sofa G . Then the triple $(K, B_K, D_K) \in \mathcal{L}$ attains the maximum area of $\mathcal{Q} : \mathcal{L} \rightarrow \mathbb{R}$.*

Proof. By Theorem 8.5.7 and Theorem 7.1.5. \square

We finally prove the optimality of Gerber's sofa G by assembling the pieces.

Proof of Theorem 1.1.1. By Theorem 1.5.1, Theorem 3.5.6, and Theorem 1.5.2, some balanced maximum sofa S^* with rotation angle $\pi/2$ attains the maximum area of a moving sofa. Let K^* be the cap of S^* , then we have $K^* \in \mathcal{K}^i$ by Theorem 8.1.1, and $(K^*, B_{K^*}, D_{K^*}) \in \mathcal{L}$ by Theorem 8.1.8. We also have $|S^*| = \mathcal{A}(K^*) \leq \mathcal{Q}(K^*, B_{K^*}, D_{K^*})$ by Theorem 2.5.10 and Theorem 8.2.4.

Let $K := \mathcal{C}(G)$ be the cap of Gerver's sofa. By Corollary 8.5.8 and Theorem 8.4.6, we have $\mathcal{Q}(K^*, B_{K^*}, D_{K^*}) \leq \mathcal{Q}(K, B_K, D_K) = |G|$. Summarizing, we have $|S^*| \leq |G|$ and Gerver's sofa G attains the maximum area. \square

Remark 8.5.1. The choice of constants φ^R and φ^L different from $(\varphi^R, \varphi^L) := (\varphi, \pi/2 - \varphi)$ of Gerver's sofa will also give a valid upper bound $\mathcal{Q} : \mathcal{L} \rightarrow \mathbb{R}$ of the sofa area. However, for such a \mathcal{Q} , the maximizer (K, B, D) of \mathcal{Q} does not give a valid moving sofa (that is, not in the embedding $\mathcal{K}^i \rightarrow \mathcal{L}$ of Theorem 8.1.8) because the ends of the tails \mathbf{d}_D , \mathbf{b}_B and core $\mathbf{x}_K|_{[\varphi^R, \varphi^L]}$ do not match. The choice $(\varphi^R, \varphi^L) := (\varphi, \pi/2 - \varphi)$ ensures that the maximizer (K, B, D) of \mathcal{Q} is in the image of $\mathcal{K}^i \rightarrow \mathcal{L}$ and gives back the moving sofa G .

Appendix A

Table of Symbols

Symbol	Meaning	Location
$ X $	Area of $X \subseteq \mathbb{R}^2$	Definition 1.1.3
$\overline{X}, X^\circ, \partial X$	Topological closure, interior, and boundary of $X \subseteq \mathbb{R}^2$	Definition 2.1.2
L, H_L, V_L	The right-angled hallway of unit width and its horizontal/vertical sides	Definition 1.1.1
$\overline{a}_L, \overline{b}_L, c_L, d_L, \overline{b}_L, \overline{d}_L$	Walls of the hallway L	Definition 2.2.1
$\mathbf{x}_L, \mathbf{y}_L$	Inner and outer corner of the hallway L	Definition 2.2.1
S	Moving sofa	Definition 1.1.2
G	Gerver's Sofa	Section 1.1
α_{\max}	Maximum area of a moving sofa	Section 1.1
ω	Rotation angle of a sofa	Definition 2.3.3
H	Horizontal strip $[0, 1] \times \mathbb{R}$	Definition 2.3.2
V, V_ω	Vertical strip $\mathbb{R} \times [0, 1]$, and its counterclockwise rotation by ω at the origin	Definition 2.3.2
P_ω	Parallelogram of rotation angle ω	Definition 2.3.5
O	Origin $(0, 0)$	Definition 2.3.5
o_ω	Upper-right vertex of P_ω	Definition 2.3.5
R_t	Rotation of \mathbb{R}^2 at the origin by a counterclockwise angle t	Definition 2.3.1
u_t, v_t	Orthogonal unit vectors $(\cos t, \sin t)$ and $(-\sin t, \cos t)$ of angle t	Definition 2.1.3
$l(t, h)$	Line with normal angle t and distance h from the origin	Definition 2.1.4
$H_\pm(t, h), H_\pm^\circ(t, h)$	Half-plane with normal angle t and distance h from the origin	Definition 2.1.5
K	A cap or a planar convex body	Definition 2.1.1
\mathcal{K}	Space of all planar convex bodies	Definition 2.1.1
$h_K(t)$	Supporting function of a planar convex body K	Definition 2.1.6
$l_K(t)$	Supporting line of a planar convex body K with normal angle t	Definition 2.1.7

Symbol	Meaning	Location
$H_K(t)$	Supporting half-plane of a planar convex body K with normal angle t	Definition 2.1.7
$v_K^\pm(t)$	Vertices of a planar convex body K	Definition 2.1.10
$v_K(a, b)$	Intersection $l_K(a) \cap l_K(b)$	Definition 2.1.14
$e_K(t)$	Edge (face) of a planar convex body K	Definition 2.1.10
\mathcal{H}^1	Hausdorff measure of dimension one on \mathbb{R}^2	Definition 2.1.11
σ_K	Surface area measure of a planar convex body K	Definition 2.1.13
d_H	Hausdorff distance between two convex bodies	Definition 2.1.12
$L_S(t), L_K(t)$	Supporting hallway of set S or cap K	Definition 2.2.2
$a_K(t), b_K(t), c_K(t), d_K(t), \vec{b}_K(t), \vec{d}_K(t)$	Walls of supporting hallway $L_K(t)$	Definition 2.2.3
$\mathbf{x}_K(t), \mathbf{y}_K(t)$	Inner and outer corner of supporting hallway $L_K(t)$	Definition 2.2.3
\mathcal{K}_ω^c	Space of all caps with rotation angle ω	Definition 2.4.2
\mathcal{K}^c	Space of all caps with rotation angle $\omega = \pi/2$	Definition 6.1.1
\mathcal{K}^i	Space of all caps with rotation angle $\pi/2$ satisfying injectivity condition	Definition 8.1.1
\mathcal{K}_Θ^c	Space of polygon caps with angle set Θ	Definition 3.2.3
\mathcal{K}_Θ^t	Space of all translations of polygon caps with angle set Θ	Definition 3.3.1
\mathcal{H}_Θ	Space of generalized support functions with angle set Θ	Definition 3.3.2
$\mathcal{I}(S)$	The intersection/monotone sofa arising from a moving sofa S in standard position	Definition 2.3.6
$\mathcal{C}(S)$	The cap of a moving sofa	Definition 2.3.10
δK	Upper boundary of a cap K	Definition 2.5.2
$\mathcal{N}(K)$	The niche of a cap	Definition 2.4.5
$T_K(t)$	Wedge of angle t from cap K	Definition 2.5.3
$W_K(t), Z_K(t)$	Left and right endpoints of the wedge $T_K(t)$	Definition 2.5.4
$w_K(t), z_K(t)$	Wedge gap of a cap K	Definition 2.5.5
$A_K^\pm(t), C_K^\pm(t)$	Vertices of a cap K	Definition 2.5.1
Θ	Angle set	Definition 3.2.1
$\Theta_{\omega, n}$	Uniform angle set of n intervals and rotation angle ω	Definition 3.5.1
Θ_n	Uniform angle set of n intervals and rotation angle $\omega = \pi/2$	Definition 6.3.1
τ_K	Edge length of the niche $\mathcal{N}_\Theta(K)$ of polygon cap K	Definition 3.4.4
k_0, m_0	Real-valued functions used in the proof of injectivity condtion.	Definition 6.3.4
$f_K^\pm(t), g_K^\pm(t)$	Arm lengths; distance from $\mathbf{y}_K(t)$ to $A_K^\pm(t)$ and $C_K^\pm(t)$ respectively	Definition 6.2.1
df	Lebesgue–Stieltjes measure of f	Definition 5.1.3
\mathcal{V}, c_λ	Convex domain and its barycentric operation	Definition 7.1.1
Df	Directional derivative of quadratic functional f on a convex domain	Definition 7.1.5
θ, φ	Angle constants determining Gerver’s sofa	Definition 8.1.2
φ^R, φ^L	Constants $\varphi^R := \varphi$ and $\varphi^L := \pi/2 - \varphi$	Definition 8.1.2

Symbol	Meaning	Location
\mathcal{L}	Space of triples (K, B, D) of convex bodies extending \mathcal{K}^1	Definition 8.1.3
\mathcal{Q}	Upper bound of sofa area on the space \mathcal{L}	Definition 8.2.2
$\mathcal{J}(\mathbf{x})$	Curve area functional	Definition 7.2.6
$\mathcal{J}(p, q)$	Curve area functional from point p to q	Definition 7.2.8
$\mathbf{u}_K^{a,b}$	Convex curve segment of the boundary of K	Definition 7.3.1
$\mathcal{M}_K(a, b; \mathbf{z})$	The area of a Mamikon region bounded by $\mathbf{u}_K^{a,b}$ and $\mathbf{z} : [a, b] \rightarrow \mathbb{R}$.	Definition 7.4.2
t_0, \dots, t_5	The angles $0 = t_0 < \dots < t_5 = \pi/2$ used in Gerver's sofa G	Definition 8.4.1
I_1, \dots, I_5	The intervals $I_i = [t_{i-1}, t_i]$	Definition 8.4.1
$\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{x}$	The boundaries of Gerver's sofa G	Definition 8.4.3

Bibliography

- [Apo] Tom M Apostol. *Mathematical Analysis: A Modern Approach To Advanced Calculus*. Second. Pearson Higher Education. ISBN: 978-0-201-00284-3.
- [Bat22] Michał Batsch. “A Numerical Approach for Analysing the Moving Sofa Problem”. In: *Symmetry* 14.7 (July 8, 2022), p. 1409. ISSN: 2073-8994. DOI: [10.3390/sym14071409](https://doi.org/10.3390/sym14071409). URL: <https://www.mdpi.com/2073-8994/14/7/1409> (visited on 11/04/2023).
- [Bie95] Hanspeter Bieri. “Nef Polyhedra: A Brief Introduction”. In: *Geometric Modelling*. Ed. by H. Hagen, G. Farin, and H. Noltemeier. Vienna: Springer Vienna, 1995, pp. 43–60. ISBN: 978-3-7091-7584-2.
- [Den24] Zhipeng Deng. *Solving Moving Sofa Problem Using Calculus of Variations*. July 2, 2024. arXiv: [2407.02587 \[math\]](https://arxiv.org/abs/2407.02587). URL: <http://arxiv.org/abs/2407.02587> (visited on 10/29/2024). Pre-published.
- [Ger92] Joseph L. Gerver. “On Moving a Sofa around a Corner”. In: *Geometriae Dedicata* 42.3 (June 1992), pp. 267–283. ISSN: 0046-5755, 1572-9168. DOI: [10.1007/BF02414066](https://doi.org/10.1007/BF02414066). URL: <http://link.springer.com/10.1007/BF02414066> (visited on 05/16/2023).
- [Gib14] Philip Gibbs. *A Computational Study of Sofas and Cars*. 2014. URL: <https://vixra.org/pdf/1411.0038v2.pdf>. Pre-published.
- [Hag15] Hagen von Eitzen. *Answer to "Connected-ness of the Boundary of Convex Sets, under Additional Assumptions of the Convex Set Being Compact or Bounded"*. Mathematics Stack Exchange. May 25, 2015. URL: <https://math.stackexchange.com/questions/1298125/connected-ness-of-the-boundary-of-convex-sets-in-mathbb-rn-n1-under-a> (visited on 07/30/2024).
- [KR18] Yoav Kallus and Dan Romik. “Improved Upper Bounds in the Moving Sofa Problem”. In: *Advances in Mathematics* 340 (Dec. 2018), pp. 960–982. ISSN: 00018708. DOI: [10.1016/j.aim.2018.10.022](https://doi.org/10.1016/j.aim.2018.10.022). arXiv: [1706.06630 \[math\]](https://arxiv.org/abs/1706.06630). URL: <http://arxiv.org/abs/1706.06630> (visited on 05/16/2023).
- [Len+24] Kuangdai Leng et al. “Deep Learning Evidence for Global Optimality of Gerver’s Sofa”. In: *Symmetry* 16.10 (Oct. 18, 2024), p. 1388. ISSN: 2073-8994. DOI: [10.3390/sym16101388](https://doi.org/10.3390/sym16101388). URL: <https://www.mdpi.com/2073-8994/16/10/1388> (visited on 10/29/2024).
- [Mna97] Mamikon Mnatsakanian. “Annular Rings of Equal Area”. In: *Math Horizons* 5.2 (Nov. 1997), pp. 5–8. ISSN: 1072-4117, 1947-6213. DOI: [10.1080/10724117.1997.11975029](https://doi.org/10.1080/10724117.1997.11975029). URL: <https://www.tandfonline.com/doi/full/10.1080/10724117.1997.11975029> (visited on 06/16/2023).

- [Mos66] Leo Moser. “Problem 66-11, Moving Furniture Through a Hallway”. In: *SIAM Review* 8.3 (1966), pp. 381–381.
- [MR14] Jean-François Marckert and David Renault. “Compact Convex Sets of the Plane and Probability Theory”. In: *ESAIM: Probability and Statistics* 18 (2014), pp. 854–880.
- [Rom18] Dan Romik. “Differential Equations and Exact Solutions in the Moving Sofa Problem”. In: *Experimental Mathematics* 27.3 (July 3, 2018), pp. 316–330. ISSN: 1058-6458, 1944-950X. DOI: [10.1080/10586458.2016.1270858](https://doi.org/10.1080/10586458.2016.1270858). URL: <https://www.tandfonline.com/doi/full/10.1080/10586458.2016.1270858> (visited on 05/16/2023).
- [RY13] Daniel Revuz and Marc Yor. *Continuous Martingales and Brownian Motion*. Vol. 293. Springer Science & Business Media, 2013.
- [Sch13] Rolf Schneider. *Convex Bodies: The Brunn–Minkowski Theory*. 2nd ed. Encyclopedia of Mathematics and Its Applications. Cambridge: Cambridge University Press, 2013. DOI: [10.1017/CB09781139003858](https://doi.org/10.1017/CB09781139003858).
- [Sto49] Marshall Harvey Stone. “Postulates for the Barycentric Calculus”. In: *Annali di Matematica Pura ed Applicata* 29 (1949), pp. 25–30.