## How well behaved is finite dimensional Diffusion Maps embedding?

Wenyu Bo\* Department of Statistics University of Washington Seattle, WA 98195 wbo@uw.edu Marina Meilă Department of Statistics University of Washington Seattle, WA 98195 mmp@stat.washington.edu

## Abstract

Under a set of assumptions on a family of submanifolds  $\subset \mathbb{R}^D$ , we derive a series of geometric properties that remain valid after finite-dimensional and almost isometric Diffusion Maps (DM) embedding , including almost uniform density, finite polynomial approximation and local reach. Leveraging these properties, we establish rigorous bounds on the embedding errors introduced by the DM embedding algorithm is  $O\left(\left(\frac{\log n}{n}\right)^{\frac{1}{8d+16}}\right)$ . These results offer a solid theoretical foundation for understanding the performance and reliability of DM embedding in practical applications.

## 1 Introduction

The Diffusion Maps (DM) embedding[1; 2], a dimensionality reduction technique that captures the geometric structure of data by constructing a diffusion process among data points, is central to manifold learning from samples. [3] showed that in the limit of infinite dimension it is an isometric embedding, while [4] showed that almost isometry can be achieved with a finite number m of eigenfunctions, where this m depends on manifold geometric properties. [5] showed that if isometry is not required, then the sufficient embedding dimension by Laplacian eigenfunctions depends on dimension, injectivity radius, Ricci curvature and volume, thus that it can be arbitrarily larger than the Whitney embedding dimension of 2d.

The DM embedding is widely used for non-linear dimension reduction as the Diffusion Maps algorithm[1; 6; 7; 8], which embeds a sample into m dimensions by the eigenvectors of  $L_n$ , a  $n \times n$  matrix estimator of the Laplace-Beltrami operator  $\Delta$ . For instance, DM is frequently used to analyze high-dimensional single-cell RNA sequencing data, revealing cell differentiation trajectories and underlying biological patterns[9], and it can order cells along their differentiation paths, enabling accurate reconstruction of branching developmental processes[10]. In chemistry, DM can extract the dynamical modes of high-dimensional simulation trajectories, furnishing a kinetically low-dimensional framework[11], it can identify the collective coordinates of rare events in molecular transitions[12]. In Astronomy, DM is applied to estimate galaxy redshifts from photometric data, demonstrating comparable accuracy to existing methods[13].

Stimulated in part by the practical applications of DM, the consistency of finite sample estimators of  $\Delta$ , and of their eigenfunctions has been intensely studied. Many early works discussed the pointwise consistency for empirical operator with smooth functions on M and they derived convergence (with different rates) [14; 15; 16; 17; 18], which provide illuminating results for subsequent studies. Not only  $\Delta$  itself, the consistency of its embedding are also widely studied, they discussed how the

<sup>\*</sup>\_

eigenvectors and eigenvalues of the empirical Laplacian with specific graph (K-NN,  $\epsilon$ -graph) converge to the eigenfunctions and eigenvalues of  $\Delta$  in the different norm  $(L^2, L^{\infty})[19; 20; 21; 22]$ , also there are some other special convergence been studied. For instance, convergence in Lipschitz norm[23], embedding using heat kernel[24; 25]

This paper completes the picture/advances the understanding of the DM embedding in the infinite and finite sample case, by considering geometric properties of a manifold, such as smoothness, injectivity radius, reach, volume, diameter, and examining to what extent, and under the conditions that diffusion time t is small enough and embedding dimension m is large enough, the DM preserves these properties. In other words, if the original manifold  $\mathcal{M}$  is "well behaved", what can be said about its DM embedding  $\varphi(\mathcal{M})$ ?

More specifically, we set a series of geometric assumptions on a family of compact manifolds (Section 2), then we study the smoothness of  $\varphi$  by examining its Sobolev norms (Section 3.1) and use it to implies the existence of local parameterization, then we derived consistent properties that hold uniformly for all manifolds in the family after DM embedding, like sampling densities and reach (Section 3). In Section 4, we identified the correspondence between the embedding via eigenfunctions and the embedding via eigenvectors of the graph matrix and address the convergence between them, then estimate the error between the eigenfunctions and eigenvectors to quantify the noise in the positions of the embedded points.

### 2 Background, challenges and assumptions

#### 2.1 Manifolds and the Diffusion Map

For basic definitions reader should consult [26; 27; 28]. Here we consider sub-manifolds  $M \subset \mathbb{R}^D$  which we will generically call *manifolds*. Note that the ambient dimension D will not appear throughout the paper, only Euclidean distances |x - y| with  $x, y \in \mathbb{R}^D$  will appear, hence D can actually be infinite, and  $\mathbb{R}^D$  could become a Hilbert space.

We also assume that M is a closed manifold with smoothness of class  $C^k$ , as described in Assumption 4. This assumption ensures the existence of a local polynomial expansion up to order k in the neighborhood of each point on M. Such expansions satisfy specific regularity constraints, enabling precise characterization of the local geometry and supporting rigorous analytical derivations. These properties are fundamental for establishing the theoretical results and ensuring consistency in subsequent computations.

In this work, we focus on several key geometric quantities, including the tangent space, reach, Riemannian metric, and geodesic curves. These quantities play critical roles in understanding the geometry of the manifold and in estimating and reconstructing manifolds. While these concepts are broadly defined in the general framework in Riemannian geometry, considering our sampling points are in the submanifolds in  $\mathbb{R}^D$ , we restrict our attention to their specialized formulations for submanifolds embedded in Euclidean space. This setting simplifies their definitions and aligns with the analytical and computational techniques employed in this study. For example, the tangent space is characterized as the hyperplane tangent to M at a given point, and the Riemannian metric is induced from the ambient Euclidean metric.

**Definition 1** (Tangent Space). Let M be a smooth manifold and p be a point of M. If a linear map  $v: C^{\infty}(M) \to \mathbb{R}$  satisfies

$$v(fg) = f(p)vg + g(p)vf \quad \text{for all } f, g \in C^{\infty}(M), \tag{1}$$

then we call v a derivation at p. The set of all derivations at p is called the tangent space to M at p, denoted by  $T_pM$ .

For a manifold  $M \subset \mathbb{R}^D$ , let  $\iota : M \hookrightarrow \mathbb{R}^D$  be the inclusion map. Let  $(U, \mathbf{x})$  be the chart containing p, and  $\hat{U} := \mathbf{x}(U) \subset \mathbb{R}^d$ . We have  $\hat{\iota} := \iota \circ \mathbf{x}^{-1} : \hat{U} \xrightarrow{\mathbf{x}^{-1}} U \xrightarrow{\iota} \mathbb{R}^D$  is a local representation near p by

$$(\mathbf{x}_1(p),\ldots,\mathbf{x}_d(p)) \to (p_1,\ldots,p_D).$$
<sup>(2)</sup>

Then  $\left\{\frac{\partial i}{\partial x_1}(\mathbf{x}(p)), \ldots, \frac{\partial i}{\partial x_d}(\mathbf{x}(p))\right\}$  span a *d*-dimensional subspace at  $\iota(p)$  of  $\mathbb{R}^D$ , which is the common definition of tangent space at p in the Euclidean case.

**Definition 2** (Geodesic Normal Coordinate). Let  $p \in U \subset M$  and  $\exp_p|_V : V \subset T_pM \to U$  be a diffeomorphism, and there is a basis isomorphism B between  $\mathbb{R}^d$  and orthonormal basis  $\{b_i\}$  for  $T_pM$  by  $B(x_1, \ldots, x_d) = \sum_{i=1}^d x_i b_i$ . Then U is the normal neighborhood with normal coordinate  $\left(\exp_p|_V \circ B\right)^{-1}$ .

Since  $d_0(\exp_p)$  is the identity map, thus by inverse function theorem, such neighborhood V always exists, then it is well defined.

**Definition 3** (Riemannian Metric). A Riemannian metric g on a manifold M assigns to every p a inner product  $g_p(\cdot, \cdot)$  on  $T_pM$  which is smooth in the following sense: For a chart  $(U, \mathbf{x})$  containing p, with  $\mathbf{x}^{-1}(x_1, \ldots, x_d) = q \in U$  and  $\frac{\partial}{\partial x_i}(q) = d\mathbf{x}_q^{-1}(0, \ldots, 1, \ldots, 0)$ , then  $\left\langle \frac{\partial}{\partial x_i}(q), \frac{\partial}{\partial x_j}(q) \right\rangle = g_{ij}(x_1, \ldots, x_d)$  is smooth on  $\hat{U}$ .

In our setting, manifolds are embedded into  $\mathbb{R}^D$  via  $\iota$ , then Riemannian metric automatically inherits Euclidean metric, i.e.  $g_p(v, w) = \langle d\iota_p v, d\iota_p w \rangle$ .

Riemannian metric defines a measure on M, called the *Riemannian measure* via the *volume form*. If M is compact, it has a finite volume Vol(M). We can now have other measures on M, which are absolutely continuous w.r.t. the Riemannian measure, through their densities  $f: M \to (0, \infty)$ .

In statistics/manifold learning, it is assumed that we are given an i.i.d. sample  $X_n = \{x_1, \dots, x_n\} \subset M$  from f.

**Definition 4** (Reach[29]). The reach of a subset A of  $\mathbb{R}^n$  is the largest  $\tau$  (possibly  $\infty$ ) such that if  $x \in \mathbb{R}^n$  and the distance from x to A is smaller that  $\tau$ , then A contains a unique point nearest to x. For more detail and properties, refer [30].

The reach of a subset  $A \subset \mathbb{R}^n$  provides a measure of its local geometric regularity. Specifically, if the reach of A is  $\tau > 0$ , then every point within a distance less than  $\tau$  from A has a unique nearest point on A. Geometrically, this implies that A does not exhibit sharp corners, cusps, or regions of high curvature within the specified radius. In particular: For convex subsets of  $\mathbb{R}^n$ , the reach is infinite, reflecting the absence of curvature bounds or sharp features. For smooth submanifolds, the reach is inversely related to its maximum principal curvature. Intuitively, the reach corresponds to the radius of the smallest osculating ball that fits locally around.

Definition 5 (Laplacian-Beltrami Operator).

The Laplacian-Beltrami operator is the linear operator  $\Delta: C^{\infty}(M) \to C^{\infty}(M)$  defined by

$$\Delta f = div(gradf). \tag{3}$$

It is a classical result that the eigenvalues of the Laplace-Beltrami operator  $-\Delta$  on a Riemannian manifold M form a non-decreasing spectrum, i.e.

$$0 \le \lambda_0 \le \lambda_1 \le \lambda_2 \dots, \tag{4}$$

where each eigenvalue is repeated according to its multiplicity. When M is compact, the spectrum of  $-\Delta$  is discrete, and each eigenvalue has finite multiplicities. If M is compact and without boundary, the smallest eigenvalue  $\lambda_0 = 0$ , and its eigenspace consists of constant functions. Furthermore, the regularity theory for elliptic operators ensures that all eigenfunctions are smooth, i.e., they belong to  $C^{\infty}(M)$ .

**The Diffusion map** Let M be a smooth, closed manifold of class  $C_3$  embedded in the (possibly high-dimensional) Euclidean space  $\mathbb{R}^D$  and  $\Delta_M$  be Laplacian operator on M with eigenvalues  $0 = \lambda_0 \leq \lambda_1 \leq \cdots$ . We consider eigenfunctions corresponding these eigenvalues:  $e_0, e_1, \cdots$ , and we normalize them such that  $||e_i||_{L^2} = 1$ . It is easy to check  $e_0 \equiv const$ . Then we have the following embedding theorem.

**Theorem 1.** Let  $\mathcal{M}$  be the set of d dimensional, closed Riemannian manifolds whose Ricci curvature is bounded from below by  $\kappa$ , injectivity radius is bounded from below by  $\iota$ , and the volume is bounded from above by V, we define  $\varphi : \mathcal{M} \subset \mathbb{R}^D \to \mathcal{N} \subset \mathbb{R}^m$  as the following:

$$\varphi(x) = (2t)^{\frac{d+2}{4}} \sqrt{2} (4\pi)^{\frac{d}{4}} (e^{-\lambda_1 t} e_1(x), e^{-\lambda_2 t} e_2(x), \cdots, e^{-\lambda_m t} e_m(x)) \in \mathbb{R}^m,$$
(5)

then there exists a  $t_0 = t_0(d, \kappa, \iota, \epsilon)$  such that for all  $0 < t < t_0$ , there exists a  $N_0(d, \kappa, \iota, \epsilon, V, t)$ such that if  $N > N_0$ , then for all  $M \in \mathcal{M}(d, \kappa, \iota, V)$ , the map above is an embedding of M into  $\mathbb{R}^N$ , and  $1 - \epsilon < \|d\varphi_p v\| < 1 + \epsilon$  where  $\|v\| = 1.[4]$ 

**Lemma 2.** We use the same notation as in the preceding theorem. Then for  $\epsilon' > 0$ , there exists  $N_1 = N_1(d, \kappa, \iota, V, \epsilon', t_0)$  such that when  $N \ge N_1$  and  $t_0 \le t' \le 4$ , we have

$$\|K_N(t', p, \cdot) - K(t', p, \cdot)\| \le \epsilon',\tag{6}$$

where K(t, p, q) is the heat kernel:

$$K(t, p, q) = \sum_{i=0}^{\infty} e^{-\lambda_i t} e_i(p) e_i(q),$$
(7)

and  $K_N(t, p, q)$  is the truncated heat kernel:

$$K_m(t, p, q) = \sum_{i=0}^{m} e^{-\lambda_i t} e_i(p) e_i(q).$$
 (8)

We select  $t = t_0/2$ ,  $t' = t_0$  (if  $t_0 > 4$ , we set t = 2 and t' = 4) and  $m = \max \{N_0, N_1\} + 1$  to make the results above both hold.

In addition,  $\varphi : M \to \varphi(M)$  is homeomorphism, so dim  $M = \dim \varphi(M)$ , and  $\varphi$  is an embedding, also an immersion, thus  $\varphi$  is local diffeomorphism, and  $\varphi$  is bijective, so  $\varphi$  is diffeomorphism. Thus  $\varphi(M)$  is a smooth manifold.[27]

Given a finite sample  $\mathcal{X}_n = \{x_1, \ldots, x_n\}$  the DM algorithm constructs a similarity matrix to measure the pairwise relationships between data points, where the commonly used kernel is the Gaussian kernel, defined as  $k(x_i, x_j) = \exp(-||x_i - x_j||^2/h)$ , where  $\epsilon$  is a scale parameter, and this similarity matrix is then normalized, we denote it as Laplacian graph which is the approximation of Laplacian-Beltrami Operator. By performing eigenvalue decomposition on normalized Laplacian graph, the algorithm extracts the dominant eigenvalues and eigenvectors. These eigenvectors, scaled by their corresponding eigenvalues, define the diffusion coordinates, providing a low-dimensional embedding of the data. This embedding preserves the global geometry of the dataset while emphasizing its intrinsic structure.

In this paper, we will focus mainly on the geometric properties of the DM  $\varphi(M)$ , w.r.t. the original manifold M. The results we obtain will be useful in characterizing the output of the DM algorithm in finite sample settings, and we apply them specifically to the estimation of the tangent subspace  $\{T_{\varphi(x_i)}\varphi(x_i)\}_{i=1}^n$ .

[3] have shown that, in the limit of large m and small t,  $\varphi(M)$  is isometric with M, and that this is possible approximatively with a finite m for manifolds with bounded diameter and Ricci curvature bounded from below. In spite of these seemingly encouraging results, the DM can be highly unstable even for apparently "nice" manifolds. The intuitive explanation is the fact that, even if M is compact and smooth to order k, the local interactions between the manifold curvature and the manifold reach (note that these are not independent quantities) can exert a strong influence of the Laplacian eigenfunctions.

#### 2.2 Assumptions

Thus, in predcting smoothness (w.r.t. Sobolev norms) and geometric properties of  $\varphi(M)$ , one needs to consider  $\tau_M, \iota_M, \kappa_M$  in addition the smoothess of M (made more precise below). We will give the assumptions for the set of manifolds we will perform DM, and then discuss basic properties of M based on these assumptions. We will see all these assumptions intuitively ensure our manifolds have very good shape, which can help us avoid extremely bizarre situations. And we will use them to derive existence of the geometric bound after DM, which are critical in estimating the convergence rate.

Assumption 1 (Curvature). The absolute value of sectional curvature of M is bounded by  $\kappa$ , that is  $|K(u,v)| \leq \kappa$ . which immediately implies Ricci curvature of M is bounded below by  $-\kappa(d-1)$  since  $Ric(v,v) = \frac{1}{d-1} \sum_{i=1}^{d} K(v,x_i)$  where  $\{v, x_2, \ldots, x_d\}$  are orthonormal basis. This assumption will be used in Section 2.1, Section 3.1.1.

Bounded Ricci curvature prevents the submanifold from having extreme geometric variations or "infinite negative curvature" in any direction. This geometric control ensures that local neighborhoods behave predictably, which is crucial for DM that rely on local structure.

Assumption 2 (Reach). The reach  $\tau_M$  of M is bounded below by  $\tau_{min}$ . This assumption will be used in Section 3.4.

Positive minimum reach is crucial because it ensures the manifold doesn't come too close to selfintersecting and has bounded curvature, making it possible to reliably reconstruct the manifold from discrete samples.

**Assumption 3** (Volume). The volume of M is bounded below by  $V_1$  and bounded above by  $V_2$ . This assumption will be used in Section 2.1, Section 3.1.1.

Assumption 4 (Smoothness). For k and  $\mathbf{L} := (L_{\perp}, L_3, \dots, L_k)$ , we assume there exists a local one-to-one parameterization  $\Psi_p$  for all  $p \in M$ :

$$\Psi_p: B_{T_nM}(0, r) \to M \quad \text{by} \quad \Psi_p(v) = p + v + N_p(v) \tag{9}$$

for some  $r\geq \frac{1}{4L_{\perp}}$  with  $N_p(v)\in C^k(B_{T_pM}(0,r),\mathbb{R}^D)$  such that

$$N_p(0) = p, \ d_0 N_p = 0, \ \|d_v^2 N_p\| \le L_\perp, \ \|d_v^i N_p\| \le L_i \text{ for } i = 3, \dots, k$$
(10)

holds for all  $||v|| \leq \frac{1}{4L_1}$ . This assumption will be used in Section 3.2

Smoothness of order k implies manifolds can be approximated locally by multilinear map over tangent space with bounded norm.

In addition, we make the following more technical regularity assumption that will be used in Section 3.1.2, Section 3.4.

Assumption 5 (Christoffel Symbols). In the normal coordinate chart, the derivatives with order not above k-2 of Christoffel symbol (including Christoffel symbol itself) have the uniform upper bound only depending on its order, i.e.

$$\left|\frac{\partial^{l}\Gamma_{ij}^{k}}{\partial x_{i_{1}}\cdots\partial x_{i_{l}}}\right| \leq C(l) \quad \text{for } 1 \leq i, j, k \leq d \text{ and } l \leq k-2$$

$$(11)$$

Christoffel symbol measures the change of a vector along a curve due to curvature, thus bounded Christoffel symbol control the curvature in some manner.

Let  $\mathcal{M}(d, \kappa, K, \tau_{min}, V, k, \mathbf{L}, \Gamma)$  be the set of compact connected submanifolds  $M \subset \mathbb{R}^D$ , with dimension d satisfying Assumption 1–Assumption 5. For simplicity, in the rest of our paper we always assume that the manifold is d-dimensional and the ambient dimension is D, we also use the abbreviation  $\mathcal{M}$  for  $\mathcal{M}(d, \kappa, \tau_{min}, V, k, \mathbf{L}, \Gamma)$ , and sometimes we include some of the parameters above to indicate that the assumptions corresponding to these parameters are satisfied.

We also assume that the sampling density on M does not deviate too much from uniform.

Assumption 6 (Density). Let  $\mathcal{P}_{f_{\min}, f_{\max}}$  denote the set of distribution P with support on  $M \in \mathcal{M}$ , and the density function f of P with respect to Hausdorff measure such that  $0 < f_{\min} \leq f \leq f_{\max} < \infty$ . This assumption will be used in Section 3.3.

All the assumptions above, perhaps with the exception of Assumption 4 and Assumption 5, are generically present in the manifold learning literature.

#### 2.3 Direct Consequences of Assumptions

Here we list some direct results for  $M \in \mathcal{M}$  from assumptions which will also be used in our following proof.

**Corollary 3** (Complete Manifold). Any compact Riemannian manifold is geodesically complete according to Hopf-Rinow theorem.[26]

**Corollary 4** (The injectivity radius). The injectivity radius  $\iota_M$  of M is bounded below by  $\pi \tau_M[30]$ , which implies  $\iota_M$  is bounded by  $\pi \tau_{\min}$  for all  $M \in \mathcal{M}$ . We use it to make sure DM is an embedding.

The injectivity radius bounded from below implies we can find normal coordinate chart with the uniform radius.

**Corollary 5** (Diameter). For  $M \in \mathcal{M}$ , we have

$$\operatorname{diam}(M) \le \frac{C_d}{\tau_M^{d-1} f_{\min}} \le \frac{C_d}{\tau_{\min}^{d-1} f_{\min}}$$
(12)

where  $C_d$  is a constant only depending on d[31]. Thus the diameter of  $M \in \mathcal{M}$  have an uniform upper bound. We will use it in Section 3.1.1.

## **3** Properties of the diffusion maps embedding $\varphi(\mathcal{M})$

In this section, we will discuss the properties of  $\varphi(M)$ , and we will first give some uniform estimate of geometric quantities for all  $M \in \mathcal{M}$ , and we will prove that the family  $\varphi(\mathcal{M}) \in \mathcal{M}(\tau_{\varphi(M)}, \mathbf{L}')$ where  $\mathbf{L}' := (L'_{\perp}, \ldots, L'_k)$  and the density function of  $\varphi(M) \in \varphi(\mathcal{M})$  admits an uniform upper and lower bounds.

#### **3.1** Bounds on $\lambda_m$ and the (higher) derivatives of $\phi$

Estimating and uniformly bounding the k-th derivative of  $\phi$ ,  $||d^k\varphi||$ , plays an important role in controlling other geometric quantities such as  $\tau_{min}$  and proving existence of local ono-to-one parametrization of  $\varphi(\mathcal{M})$ . Recall that k is the smoothness of  $\mathcal{M}$ . Because the *i*-th component of map  $\varphi : \mathcal{M} \to \mathbb{R}^m$  is  $e_i$  multiple of  $e^{-\lambda_i t_0/2}$  which is a bounded scalar when  $t_0$  is fixed, we can ignore it when we estimate the upper bound.

We notice that  $\varphi(M)$  is a d-dimensional manifold embedded in  $\mathbb{R}^m$ , then  $d\varphi_p$  is a map such that

$$d\varphi_p: T_p M \to T_{\varphi(p)}\varphi(M) \hookrightarrow \mathbb{R}^m \tag{13}$$

and we can treat the tangent vector in  $\mathbb{R}^m$  as a tangent vector in  $T_{\varphi}(p)\varphi(M)$  through the natural isomorphism. This isomorphism maps local basis to local basis, thus this map is also isometric. In the following, we will consider  $T_{\varphi(p)}\varphi(M) \cong \mathbb{R}^d$  and  $T_{\varphi(p)}\varphi(M) \subset \mathbb{R}^m$  as equivalent through this mapping.




Figure 1: Isometry

#### **3.1.1** Upper Bound of $\lambda_m$

The most classical result for estimating eigenvalues is Weyl's law:

$$N(\lambda) \sim \frac{\omega(d) Vol(M) \lambda^{\frac{3}{2}}}{(2\pi)^d},\tag{14}$$

where  $N(\lambda)$  is the number of eigenvalues less than or equal to  $\lambda$ . If we ask  $\lambda = \lambda_k$ , the above is equivalent to

$$\lambda_k^{d/2} \sim \frac{(2\pi)^d k}{\omega(n) Vol(M)} \tag{15}$$

Weyl's law provides an asymptotic expression for the eigenvalues of the Laplace-Beltrami operator, offering profound insights into their growth rates. However, in our paper, strict control for eigenvalues is essential. This requires not only asymptotic estimates but also rigorous upper bounds that hold universally. A notable result for the upper bound was established by Li-Yau, who derived explicit upper bounds for the eigenvalues under geometric constraints, which align with the assumptions in our study.

We estimate the upper bound of eigenvalues. If the lower bound of the Ricci curvature  $\kappa_M$  of M is less than 0, we have

$$\lambda_{m} \leq \begin{cases} \frac{(2\beta+1)^{2}}{4}(-\kappa_{M}) + 4(1+2^{\beta})^{2}\pi^{2} \left(\frac{\sinh\sqrt{-\kappa_{M}}\text{diam}}{\sqrt{-\kappa_{M}}\text{diam}}\right)^{\frac{2d-2}{d}} \\ \times \left((m+1)\frac{\alpha(d-1)}{d}\frac{1}{V}\right)^{\frac{2}{d}} \\ \text{when } d = 2(\beta+1), \beta = 0, 1, 2 \dots \\ \frac{(2\beta+2)^{2}}{4}(-\kappa_{M}) + 4(1+\pi^{2})(1+2^{2\beta})^{2} \left(\frac{\sinh\sqrt{-\kappa_{M}}\text{diam}}{\sqrt{-\kappa_{M}}\text{diam}}\right)^{\frac{2d-2}{d}} , \quad (16) \\ \times \left((m+1)\frac{\alpha(d-1)}{d}\frac{1}{V}\right)^{\frac{2}{d}} \\ \text{when } d = 2\beta + 3, \beta = 0, 1, 2 \dots \end{cases}$$

if M has non-negative Ricci curvature  $\kappa_M$ , we have

$$\lambda_m \le (d+4)d^{1-\frac{2}{d}} \left(\frac{m+1}{V}\alpha(d-1)\right)^{\frac{2}{d}},$$
(17)

where d is the dimension of M, V is the volume of M and  $\alpha(n)$  is the volume of  $S^n$  in  $\mathbb{R}^{n+1}$ .[32] Since for  $M \in \mathcal{M}$ , the volume is greater that  $V_1$  and the lower bound of the Ricci curvature is greater than  $\kappa$ , so  $\frac{1}{V} \leq \frac{1}{V_1}$  and  $-\kappa_M \leq -\kappa$ , thus we have

$$\lambda_m \le \begin{cases} C(m, d, V) & \text{if } \kappa_M \text{ is non-negative} \\ C(m, d, \kappa, \text{diam}, V) & \text{if } \kappa_M \text{ is negative} \end{cases}$$
(18)

For  $M \in \mathcal{M}$ , we can control d,  $\kappa$ , diam, V uniformly, so we can find a constant C only depending on m for all  $M \in \mathcal{M}$  such that

$$\lambda_m \le C(m). \tag{19}$$

#### 3.1.2 Estimate of Higher Order Derivatives

In this subsection, we will estimate the  $L^{\infty}$  norm of higher order derivatives for eigenfunctions to control the operator norm of dN' in the next subsection.

One similar result is Bernstein inequality, for eigenfunction  $e_{\lambda}$  of  $\Delta: \Delta e_{\lambda} = \lambda e_{\lambda}$ , we have [33]:

$$\|\nabla e_{\lambda}\|_{L^{\infty}} \le C\lambda^{\frac{1}{2}} \|e_{\lambda}\|_{L^{\infty}} \tag{20}$$

The same author also prove a more general estiamte, from[34], for eigenfunction  $e_{\lambda}$  of  $\Delta$ :  $\Delta e_{\lambda} = \lambda e_{\lambda}$ , we have

$$\|e_{\lambda}\|_{H^{\infty}_{k}} \sim \lambda^{\frac{k}{2}} \|e_{\lambda}\|_{L_{\infty}} \quad \text{as } \lambda \to \infty$$
(21)

where  $\sim$  means the equivalence between two norms, and  $\|\cdot\|_{H^k_{\infty}}$  is Sobolev norm. If we revise this result to have a upper bound (the author says we only need a little effort to do so), we can control  $\|d^k\varphi\|$  since we have had the upper bound of  $\lambda_m$  which only depend on M.

The author also gave a better estimate[35]:

$$\|e_{\lambda}(x)\|_{H^{\infty}_{L}} \le C\lambda^{k/2 + (d-1)/4} \|e_{\lambda}(x)\|_{L^{2}}$$
(22)

however, C may not be an uniform constant for  $M \in \mathcal{M}$ .

To estimate the  $L^{\infty}$  norm of derivatives for eigenfunctions, we first choose the coordinate chart to be normal coordinate chart to make our derivatives well defined. We take derivative of  $e(x) : M \to \mathbb{R}$ in the following sense: Let  $x \in U$  be the normal coordinate chart  $(U, \mathbf{x})$  and  $\hat{U} = \mathbf{x}(U)$ , then  $\hat{e}(x) = e \circ \mathbf{x}^{-1} : \hat{U} \to R$  be the coordinate representation of e(x). We treat  $\partial_i e(x)$  as  $\partial_i \hat{e}(x)$ , and for higher order derivatives as well.

Let  $\gamma$  be a 2*d* multi-index with  $|\gamma| := \sum_i \gamma_i \leq k$ , and  $x, y \in U$  be a geodesic normal coordinate of *M*. We denote  $\sum_{\lambda_j \leq \lambda} e_j(x) e_j(y)$  as  $e(x, y, \lambda)$ . Let  $\gamma = (\gamma_1, \ldots, \gamma_{2d})$ , then the  $\gamma$  order derivative for  $e_j(x) e_j(y)$  is

$$\partial_{x,y}^{\gamma} e_j(x) e_j(y) = \frac{\partial^{|\gamma_1 + \dots + \gamma_d|} e_j(x)}{\partial x_1^{\gamma_1} \cdots \partial x_d^{\gamma_d}} \frac{\partial^{|\gamma_{d+1} + \dots + \gamma_{2d}|} e_j(y)}{\partial y_1^{\gamma_{d+1}} \cdots \partial y_d^{\gamma_{2d}}}$$
(23)

We have [35]

$$|\partial_{x,y}^{\gamma}e(x,y,\lambda)| \le C_{\gamma}(1+\lambda)^{\frac{d+|\gamma|}{2}}.$$
(24)

To have the estimate for derivatives of eigenfunctions, let  $\alpha$  be a d dimension multi-index, and using the inequality above with  $\gamma = (\alpha, \alpha)$ , then we have

$$|\partial^{\alpha} e_{\lambda}(x)|^{2} \leq \sum_{\lambda_{j} \leq \lambda} |\partial^{\alpha} e_{j}(x)|^{2} \leq C_{\alpha} (1+\lambda)^{d/2+|\alpha|}$$
(25)

so

$$\|e_{\lambda}\|_{H_{k}^{\infty}} = \max_{|\alpha| \le k} |\partial^{\alpha} e_{\lambda}(x)| \le \sup_{|\alpha| \le k} \{C_{\alpha}\} (1+\lambda)^{k/2+d/4}$$
(26)

Since  $\lambda_m$  is bounded above by C(m), which implies that  $\|\varphi_j\|_{H^{\infty}_k} \leq C(l)$  for  $\varphi_j : M \to \mathbb{R}$ . Therefore,  $\left|\frac{\partial^l \varphi_j}{\partial x_{i_1} \cdots \partial x_{i_l}}\right| \leq C(l)$  for  $l \leq k$  with respect to normal coordinate.

To estimate the norm of  $d\varphi$ , we can derive its local coordinate representation and compute the covariant derivatives in the normal chart since choosing different chart does not have the influence on the norm. Now we denote  $\varphi$  as *j*-th component of  $\varphi$  out of simplifying the notation and  $(U, \mathbf{x})$  is the normal chart.

For the first order covariant derivative, we have

$$\nabla\varphi(X) = \nabla_X\varphi = X\varphi \tag{27}$$

thus

$$\nabla \varphi = \varphi_i dx^i \quad \text{with } \varphi_i = \partial_i \varphi \tag{28}$$

so for first order covariant derivative, the local coordinate representation is first order derivative of local representation of  $\varphi$ .

For second order covariant derivative, which is also known as covariant Hessian. We have

$$\nabla^2 \varphi(X, Y) = \nabla^2_{Y, X} = \nabla_Y (\nabla_X \varphi) - \nabla_{\nabla_Y X} \varphi = X(Y\varphi) - (\nabla_X Y)\varphi$$
<sup>(29)</sup>

thus in local coordinates,

$$\nabla^2 \varphi = \varphi_{ij} dx^i \otimes dx^j \quad \text{with } \varphi_{ij} = \partial_{ij}^2 \varphi - \Gamma_{ij}^k \partial_k \varphi, \tag{30}$$

with  $i, j, k \in \{1, 2, ..., m\}$ . For the third order coordinate derivative, we have

$$\nabla^3 \varphi(X, Y, Z) = (\nabla_Z \nabla^2 \varphi)(X, Y) \tag{31}$$

$$= Z(\nabla^2 \varphi(X, Y)) - \nabla^2 \varphi(\nabla_Z X, Y) - \nabla^2 \varphi(X, \nabla_Z Y)$$
(32)

$$= Z(X(Y\varphi) - (\nabla_X Y)\varphi) - \nabla_Z X(Y\varphi) + (\nabla_{\nabla_Z X} Y)\varphi - X(\nabla_Z Y\varphi) + (\nabla_X \nabla_Z Y)\varphi$$
(33)

thus in the local coordinate,

$$\nabla^{3}\varphi = \varphi_{ijk}dx^{i} \otimes dx^{j} \otimes dx^{k}$$
(34)  
with 
$$\varphi_{ijk} = \partial^{3}_{ijk}\varphi - \partial_{k}\Gamma^{l}_{ij}\partial_{l}\varphi - \Gamma^{l}_{ij}\partial^{2}_{kl}\varphi - \Gamma^{m}_{ik}\partial^{2}_{mj}\varphi + \Gamma^{l}_{ik}\Gamma^{m}_{lj}\partial_{m}\varphi - \partial_{i}\Gamma^{l}_{jk}\partial_{l}\varphi$$

$$- \Gamma^{l}_{jk}\partial^{2}_{il}\varphi + \Gamma^{m}_{il}\Gamma^{l}_{jk}\partial_{m}\varphi$$
(35)

The local coordinate representation is combined with derivative of Christoffel symbol and third order derivative of  $\varphi$ . And we can prove easily by induction that the local coordinate representation of  $\nabla^l \varphi$  is  $\varphi_{i_1 \dots i_l} dx^{i_1} \otimes \dots \otimes dx^{i_l}$  with  $\varphi_{i_1 \dots i_l}$  can be represented by derivative of  $\varphi$  not higher than order l and derivative of Christoffel symbol not higher than order l-2. To prove this, we only need to notice that

$$\nabla^{l+1}\varphi(X_1,\cdots,X_l,X) = \nabla(\nabla^l\varphi)(X_1,\cdots,X_l,X)$$
(36)

$$= (\nabla_X \nabla^l \varphi)(X_1, \cdots, X_l) \tag{37}$$

$$= X(\nabla^{l}\varphi(X_{1},\cdots,X_{l})) - \sum_{i=1}^{l}\nabla^{l}\varphi(X_{1},\cdots,\nabla_{X}X_{i},\cdots,X_{l}) \quad (38)$$

Since we assume for any  $M \in \mathcal{M}$ , the local coordinate representation in the normal chart of the Christoffel symbols and their derivatives with order not above k - 2 have an uniform bound only depends on its order, and we have claimed derivatives of  $\varphi$  which order is not higher than k have an uniform bound, and we notice that the local representation of  $\nabla^k \varphi$  is combined with them in the same pattern among all  $M \in \mathcal{M}$ . Therefore,  $\varphi_{i_1 \dots i_l}$  is bounded uniformly for all  $l \leq k$ , i.e.

$$|\varphi_{i_1\cdots i_l}| \le A(l) \quad \text{for } l \le k \tag{39}$$

With this control of upper bound, we can estimate the norm of higher derivatives of  $\varphi$ :

$$\|d^{l}\varphi\| = \sup_{\|v_{1}\|=1,\dots,\|v_{l}\|=1} \|\nabla^{l}\varphi(v_{1},\dots,v_{l})\|$$
(40)

$$= \sup_{\|v_1\|=1,\dots,\|v_l\|=1} \|\nabla^l \varphi(v_1 \otimes \dots \otimes v_l)\|$$
(41)

$$= |\varphi_{i_1\cdots i_l} dx^{i_1} \otimes \cdots \otimes dx^{i_l} (v_1^{i_1} \frac{\partial}{\partial x^{i_1}} \otimes \cdots \otimes v_l^{i_l} \frac{\partial}{\partial x^{i_l}})|$$
(42)

$$= |\varphi_{i_1\cdots i_l} v_1^{i_1} \cdots v_l^{i_l}| \tag{43}$$

$$\leq A(l) |\sum_{i_1, i_2, \dots, i_l=1}^{u} v_1^{i_1} \cdots v_l^{i_l}|$$
(44)

$$= A(l) \left| \sum_{i_1=1}^{d} v_1^{i_1} \sum_{i_2,\dots,i_l=1}^{d} v_2^{i_2} \cdots v_l^{i_l} \right|$$
(45)

$$(AM-GM) \le A(l) |\sum_{i_2,\dots,i_l=1}^d v_2^{i_2} \cdots v_l^{i_l} |\sqrt{d}$$
(46)

$$\leq A(l)d^{l/2}.\tag{48}$$

This inequality holds for all  $l \leq k$ . Here  $\varphi: M \to \mathbb{R}$ , thus  $d^l \varphi_j(v_1, \ldots, v_l)$  gives a vector in  $\mathbb{R}$ , and then  $d^l \varphi(v_1, \ldots, v_l) = (d^l \varphi_1(v_1, \ldots, v_l), \ldots, d^l \varphi_m(v_1, \ldots, v_l))$  is a vector in  $\mathbb{R}^m$ , which is also a tangent vector in  $T_{\varphi(p)}\varphi(M)$  through the natural isometry. Since  $d^l \varphi(v_1, \ldots, v_l)$  has m components, it is easy to check the operator norm of  $d^l \varphi$  has the upper bound  $\sqrt{m}A(l)d^{l/2}$ . Therefore,  $||d^l \varphi||$  are bounded uniformly for  $l \leq k$ .

# **3.2** Uniform upper bounds $\{L_{\perp}, L_3, \ldots, L_k\}$ on the higher derivatives of the local parametrization reminder

In this section, we will verify that there exists an common parameter set  $\mathbf{L}' := \{L'_{\perp}, L'_{3}, \ldots, L'_{k}\}$  such that the operator norm of the derivatives of remainder  $N_{\varphi(p)}$  defined in Assumption 4 can be controlled uniformly over  $\varphi(\mathcal{M})$ .

We fix a point  $p' = \varphi(p) \in \varphi(M)$ , for  $v' \in T_{p'}\varphi(M) = d\varphi_p(T_pM)$ , we define

$$\Psi'_{p'}(v') = \varphi(\Psi_p(d(\varphi^{-1})_{p'}(v'))) = p' + v' + N'_{p'}(v')$$
(49)

from definition of  $\Psi_p$ , we know  $\Psi'_{p'}(v')$  and  $N'_{p'}(v')$  are well defined when  $||d(\varphi^{-1})_{p'}(v')|| \leq \frac{1}{4L_+}$ .

We estimate  $||d\varphi_{p'}^{-1}||$ : We can treat  $T_pM$  as a d dimensional subspace of  $\mathbb{R}^D$  and treat  $T_{p'}\varphi(M)$  is a d dimensional subspace of  $\mathbb{R}^m$ , and these inclusions are canonical and isometric, so they are both closed subspaces. And  $d\varphi_p$  is homeomorphism, so graph of  $d\varphi_{p'}^{-1}$  is closed, then  $d\varphi_{p'}^{-1}$  is bounded, i.e.  $||d\varphi_{p'}^{-1}||$  exists.

$$\|d\varphi_{p'}^{-1}\| = \sup\left\{\|d\varphi_{p'}^{-1}v'\| : v' \in Im(d\varphi_p), \|v'\| = 1\right\}$$
(50)

$$= \sup\left\{\frac{\|v\|}{\|d\varphi_{p}v\|} : v' \in Im(d\varphi_{p}), \|v'\| = 1, v = d\varphi_{p'}^{-1}v'\right\}$$
(51)

$$= \sup\left\{\frac{1}{\|d\varphi_{p}v\|} : \|v\| = 1\right\}$$
(52)

$$\leq \frac{1}{1-\epsilon} \tag{53}$$

so when  $||v'|| \le \frac{1}{4L_{\perp}(1-\epsilon)}$ ,  $||d(\varphi^{-1})_{p'}(v')|| \le \frac{1}{4L_{\perp}}$ , then  $\Psi'_{p'}(v')$  and  $N'_{p'}(v')$  are well defined. Now, We check  $N'_{p'}(0) = 0$  and  $d_0 N'_{p'} = 0$ :

$$N'_{p'}(0) + p' = \varphi(\Psi_p(d(\varphi^{-1})_{p'}(0))) = \varphi(\Psi_p(0)) = \varphi(p + 0 + N_p(0)) = \varphi(p) = p'$$
(54)

$$d_0 N'_{p'} = d\varphi_p d_0 \Psi_p d(\varphi)_{p'}^{-1} - I_m = d\varphi_p (I_D + d_0 N_p) d(\varphi)_{p'}^{-1} - I_m = 0$$
(55)

We assume  $||v'|| \leq \frac{1}{4L_{\perp}(1-\epsilon)}$  and denote  $d\varphi_{p'}^{-1}v'$  as v, and  $c(v') = p + v + N_p(v)$ , then for any unit vector  $w' \in T_{p'}\varphi(\mathcal{M})$ , using Faà di Bruno's formula, we have

$$\|d_{v'}^{k}N_{p'}'(w'^{\otimes k})\| = \|\sum_{\pi \in P(k)} d_{c(v')}^{l}\varphi \circ \left(d_{v}^{j_{1}}\Psi_{p}\left(\left\{d_{p'}\varphi^{-1}w'\right\}^{\otimes j_{1}}\right), \dots, d_{v}^{j_{l}}\Psi_{p}\left(\left\{d_{p'}\varphi^{-1}w'\right\}^{\otimes j_{l}}\right)\right)\|$$
(56)

where P(k) is a partition of k with l parts such that  $j_1 + \cdots + j_l = k$ . Since  $\Psi_p(v) = p + v + N_p(v)$ , so  $d_v \Psi_p = I + d_v N_p$ ,  $d_v^i \Psi_p = d_v^i N_p$ , thus

$$|d_v \Psi_p|| = ||I + d_v N_p|| \le 1 + L_\perp ||v||$$
(57)

$$\|d_{v}^{i}\Psi_{p}\| = \|d_{v}^{i}N_{p}\| \le L_{i}$$
(58)

therefore

$$\|d_{v'}^{k}N_{p'}'(w'^{\otimes k})\| \leq \sum_{\pi \in P(k)} \|d^{l}\varphi\| \prod_{i=1}^{l} \|d^{j_{i}}\Psi_{p}\| \|d_{p'}\varphi^{-1}w'\|^{j_{i}}$$
(59)

$$\leq \sum_{\pi \in P(k)} \|d^l \varphi\| \frac{1}{(1-\epsilon)^l} \prod_{i=1}^l \|d^{j_i} \Psi_p\|$$
(60)

We have proved  $||d^j \varphi||$  and  $||d^j_v \Psi_p||$  have the uniform upper bound for  $j = 2, \ldots, k$ , so we can select  $\mathbf{L}' := (L'_{\perp}, \ldots, L'_k)$  as the uniform upper bound of  $||d^2_{v'}N'_{p'}||, \ldots, ||d^k_{v'}N'_{p'}||$ .

#### 3.3 Bounds on the pushforward density

In this subsection, we estimate the pushforward density of the sampling process. By establishing both lower and upper bounds for the pushforward density, we ensure that the sampling over  $\varphi(M)$  remains approximately uniform. This guarantees that there are always some sampling points in any nonzero measure region with nonzero probability, thereby allowing the manifold to be effectively approximated using the sampled points.

We denote  $d\varphi_p : T_p M \to T_{p'} N$  as A, then for  $||v|| = 1, 1 - \epsilon < ||Av|| < 1 + \epsilon$ , which means  $||A|| < 1 + \epsilon$ , so  $\sqrt{\lambda_{\max}(A^T A)} < 1 + \epsilon$ . We know  $A^T A$  is positive-definite symmetric matrix, we consider an unit eigenvector v of it with eigenvalue  $\lambda$ , then

$$\lambda = \|v^T \lambda v\| = \|v^T A^T A v\| = \|Av\|^2 \in ((1 - \epsilon)^2, (1 + \epsilon)^2)$$
(61)

Since  $A^T A$  is symmetric, then  $A^T A = Q \Lambda Q^T$ , where Q is orthogonal matrix and  $\Lambda$  is diagonal matrix whose diagonal elements are eigenvalues of  $A^T A$  and all of them are close to 1.

$$|||\pi_{T_pM}A^TAv|| - ||v||| \le ||\pi_{T_pM}(A^TA - I)v||$$
(62)

$$= \|\pi_{T_pM}Q(\Lambda - I)Q^Tv\|$$
(63)

$$\leq \|Q(\Lambda - I)Q^T v\| \tag{64}$$

$$\leq 3\epsilon$$
 (65)

the last inequality is due to orthogonal matrix does not change norm of vector and diagonal elements in  $\Lambda - I$  are smaller than  $3\epsilon$ .

Thus we have  $1 - 3\epsilon < ||\pi_{T_pM}A^TAv|| < 1 + 3\epsilon$ , so the eigenvalues of  $\pi_{T_pM}A^TA|_T$  is greater than  $1 - 3\epsilon$  and less than  $1 + 3\epsilon$ . Therefore, if

$$\epsilon \le \min\left\{\frac{4^{1/d} - 1}{3}, \frac{1 - \frac{1}{4^{d/1}}}{3}\right\},$$
(66)

then

$$\det(\pi_{T_pM}A^TA|_T) = \prod \lambda_i \in ((1-3\epsilon)^d, (1+3\epsilon)^d) \subset (\frac{1}{4}, 4), \tag{67}$$

which implies

$$\sqrt{\det\left(\pi_{T_pM} \circ d\varphi_p^T \circ d\varphi_p|_{T_pM}\right)} \in (\frac{1}{2}, 2).$$
(68)

According to Theorem A.2, we have the pushforward density of  $\varphi_{\#}P$  is

$$g(p') = f(p) / \sqrt{\det\left(\pi_{T_pM} \circ d\varphi_p^T \circ d\varphi_p|_{T_pM}\right)},\tag{69}$$

where  $p' = \varphi(p)$ . Therefore,  $\frac{f_{min}}{2} < g < 2f_{max}$ .

In our setting, our sampleing is uniform, thus  $f \equiv \frac{1}{Vol(M)}$ , which is bounded from both and above, which implies the density of  $\varphi(\mathcal{M})$  is also bounded from below by  $\min_{M \in \mathcal{M}} \{Vol(M)\}/2$  and bounded above by  $2 \max_{M \in \mathcal{M}} \{Vol(M)\}$ 

#### **3.4** Estimation of $\tau_{min}$

If M has reach  $\tau$ , then at least one of the following cases holds:

- I (Global case) M has a bottleneck, i.e. there exist  $p, q \in M$ , such that  $(p+q)/2 \in \text{Med}(M)$ and  $||p-q|| = 2\tau$ .
- II (Local case) There exists  $p \in M$ , and an arc-length parametrized geodesic  $\gamma$  such that  $\gamma(0) = p$  and  $|\gamma''(0)| = 1/\tau$ .

where  $\operatorname{Med}(M) = \{z \in \mathbb{R}^D : \exists p \neq q \in M, \|z - p\| = \|z - q\| = d_E(z, M)\}, d_E(x, M) \text{ is the distance between } x \text{ and } M.$ 

We denote global reach and local reach as  $\tau_g$  and  $\tau_l$ , respectively. Thus  $\tau = \min \{\tau_g, \tau_l\}$ 




Figure 2: Reach

And we can also define reach  $\tau := \min \{\tau_l, \tau_{wfs}\}$ , where  $\tau_l$  is local reach and  $\tau_{wfs}$  is weak feature size, and we give more details about them. Let  $\Gamma_M(y) = \{x \in M : d_E(y, M) = |x - y|\}$ , then define generalized gradient:

$$\nabla_M(y) := \frac{y - \operatorname{Center}(\Gamma_M(y))}{d_E(y, M)}$$
(70)

where Center(A) is the center of the smallest ball enclosing the bounded subset  $A \subset \mathbb{R}^D$ . We say y is a critical point of  $d_E(\cdot, M)$  if  $\nabla_M(y) = 0$ , then we can define

$$f_{\text{wfs}} := \inf \left\{ d_E(y, M), y \in \mathcal{C} \right\}$$
(71)

where C is the set of critical points.

And we can define  $\tau_l$  easily by

$$\tau_l := \inf_{p \in M} \left\{ \frac{1}{\|II_p\|} \right\}$$
(72)

#### 3.4.1 Local Reach

Local reach is a quantity measuring the "curvature" locally, controlling of local reach will avoid some extremely weird manifold. Now we estimate the lower bound of local reach for  $\varphi(\mathcal{M})$ .

To control the local reach, we will derive the local coordinate representation of geodesic and then use the boundedness of derivative of eigenfunctions to obtain the upper bound of geodesic of  $\varphi(\mathcal{M})$  in  $\mathbb{R}^m$ .

For any  $p \in M$ , we can choose neighborhoods  $U \subset M$  and  $V \subset \varphi(M)$  s.t.  $\varphi(U) = V$ , and  $(U, \mathbf{x})$  is a normal coordinate chart of p, where  $\mathbf{x}^{-1} = \exp_p \circ E$ , E is the isomorphism from  $\mathbb{R}^d$  to  $T_pM$ , and we denote  $\mathbf{x}(U)$  as  $\hat{U} \subset \mathbb{R}^d$  is open.

We have a parametrization of  $U \subset M$ :

$$\mathbf{x}^{-1}: \hat{U} \subset \mathbb{R}^d \to U \subset M \tag{73}$$

$$(x_1, \dots, x_d) \to \mathbf{x}^{-1}(x_1, \dots, x_d) = (\mathbf{x}_1^{-1}(x_1, \dots, x_d), \dots, \mathbf{x}_D^{-1}(x_1, \dots, x_d))$$
 (74)

Using this map, we have a coordinate representation of  $\varphi$ , i.e.  $\hat{\varphi} := \varphi \circ \mathbf{x}^{-1}$  of  $V \subset \varphi(M)$  by  $\hat{U} \xrightarrow{\mathbf{x}^{-1}} U \xrightarrow{\varphi} V \subset \mathbb{R}^m$ :

$$(x_1, \dots, x_d) \xrightarrow{\mathbf{x}^{-1}} \mathbf{x}^{-1}(x_1, \dots, x_d) \xrightarrow{\varphi} (\varphi_1 \circ \mathbf{x}^{-1}, \dots, \varphi_m \circ \mathbf{x}^{-1})(x_1, \dots, x_d)$$
(75)

Let  $r = \hat{\varphi}$  be the parameterization of  $\varphi(M)$ , we compute its k-th tangent vector

$$r_i = \frac{\partial \hat{\varphi}}{\partial x_i} = \left(\frac{\partial \hat{\varphi}_1}{\partial x_i}, \dots, \frac{\partial \hat{\varphi}_m}{\partial x_i}\right) \quad i = 1, \dots, d,$$
(76)

and r's second derivative is

$$r_{ij} = \left(\frac{\partial^2 \hat{\varphi}_1}{\partial x_i \partial x_j}, \dots, \frac{\partial^2 \hat{\varphi}_m}{\partial x_i \partial x_j}\right) \quad i, j = 1, \dots, d.$$
(77)

For  $\varphi(p) \in \varphi(M)$ , we have the orthogonal decomposition:

$$\mathbb{R}^m = T_{\varphi(p)}\varphi(M) \oplus N_{\varphi(p)}\varphi(M),\tag{78}$$

where  $T_{\varphi(p)}\varphi(M)$  is spanned by  $\{r_1, \ldots, r_d\}$  and  $N_{\varphi(p)}\varphi(M)$  is the m-d subspace orthogonal to  $T_{\varphi(p)}\varphi(M)$ . Let  $\pi$  be the orthogonal projection from  $\mathbb{R}^m$  onto  $N_{\varphi(p)}\varphi(M)$ , we can derive the second fundamental form on  $\varphi(M)$ , which is a symmetric 2-tensor field given by

$$II_{ij}dx^{i} \otimes dx^{j} \quad \text{with} \quad II_{ij} = \pi(r_{ij}) \in N_{\varphi(p)}\varphi(M).$$
(79)

Then we can verify that

$$\|II\|_{op} \le d \max\{\pi(r_{ij})\} \le d \max\{|r_{ij}|\}.$$
(80)

We compute the module of  $r_{ij} = \left(\frac{\partial^2 \hat{\varphi}_1}{\partial x_i \partial x_j}, \dots, \frac{\partial^2 \hat{\varphi}_m}{\partial x_i \partial x_j}\right)$  in more detail. We have  $\hat{\varphi}_i = (t_0)^{\frac{d+2}{4}} \sqrt{2} (4\pi)^{\frac{d}{4}} e^{-\lambda_i t_0/2} \hat{e}_i$ , thus

$$\left|\frac{\partial^2 \hat{\varphi}_k}{\partial x_i \partial x_j}\right| = (t_0)^{\frac{d+2}{4}} \sqrt{2} (4\pi)^{\frac{d}{4}} e^{-\lambda_k t_0/2} \left|\frac{\partial^2 \hat{e}_k}{\partial x_i \partial x_j}\right| \tag{81}$$

$$\leq (t_0)^{\frac{d+2}{4}} \sqrt{2} (4\pi)^{\frac{d}{4}} e^{-\lambda_k t_0/2} C_2 (1+\lambda_k)^{1+\frac{d}{4}}, \tag{82}$$

where the inequality is from eq. (26) in Section 3.1.2.

We also have the lower bound of  $\lambda_k$ ,

$$\lambda_k \ge C_1^{1+\operatorname{diam}\sqrt{\kappa}} \operatorname{diam}^{-2} k^{2/d},\tag{83}$$

where  $C_1$  is constant only depending on d.[36].

Therefore,

$$|r_{ij}| \le (t_0)^{\frac{d+2}{4}} \sqrt{2} (4\pi)^{\frac{d}{4}} C_2 \sqrt{\sum_{k=1}^m e^{-\lambda_k t_0} (1+\lambda_k)^{2+\frac{d}{2}}}.$$
(84)

It is easy to see that  $e^{-xt_0}(1+x)^{4+d}$  is increasing when  $x < \frac{4+d}{2t_0} - 1$  and decreasing when  $x > \frac{4+d}{2t_0} - 1$ , thus we can use

$$C_1^{1+\operatorname{diam}\sqrt{\kappa}}\operatorname{diam}^{-2}k^{2/d} \le \lambda_k \le C + C'k^{2/d},\tag{85}$$

which implies  $c_1 k^{2/d} \leq \lambda_k \leq c_2 k^{2/d}$ , where  $c_1, c_2$  are constants depending on  $\kappa$ , diam, V, d. Then we have  $|r_{ij}|$  is bounded from above, thus  $||II||_{op}$  is bounded from above, which implies that  $\tau_{l,\varphi(M)}$  is bounded from below.

## 4 Finite sample error of Diffusion Map $\varphi$

We consider n sample points  $\mathcal{X}_n = \{x_1, \ldots, x_n\} \subset M \in \mathcal{M}$ , we define graph affinity matrix W and the degree matrix D as:

$$W_{ij} = \frac{k_h(x_i, x_j)}{q_h(x_i)q_h(x_j)}, \quad D_{ii} = \sum_{j=1}^n W_{ij},$$
(86)

where  $k_h(x, y) = \exp\left(-\frac{|x-y|^2}{4h^2}\right)$  is the Gaussian kernel and  $q_h(x) = \sum_{i=1}^n k_h(x, x_i)$ .

Then the normalized graph Laplacian  $L_n$  is defined as

$$L_n = \frac{D^{-1}W - I}{h^2}.$$
 (87)

We denote its *i*-th eigenvalue of  $-L_n$  as  $\mu_{i,n,h}$  with corresponding eigenvector  $\tilde{v}_{i,n,h}$  normalized in  $l^2$  norm. It is easy to verify that  $\mu_{0,n,h} = 0$  and  $\tilde{v}_{0,n,h} = \mathbb{1}$ . Let  $\mathbb{N}(i) = |B_h(x_i) \cap \{x_1, \ldots, x_n\}|$  which is the cardinal of points in the *h*-ball of  $x_i$ . Then we define the  $l^2$  norm of  $\tilde{v}$  with respect to inverse estimate density  $1/\hat{p}$  as:

$$\|\tilde{v}\|_{l^{2}(1/\hat{p})} = \sqrt{\frac{|S^{d-1}|h^{d}}{d}} \sum_{i=1}^{n} \frac{\tilde{v}^{2}(i)}{\mathbb{N}(i)},$$
(88)

and we define

$$v_{i,n,h} = \frac{\tilde{v}_{i,n,h}}{\|\tilde{v}_{i,n,h}\|_{l^2(1/\hat{p})}}.$$
(89)

Let  $\Delta$  be the Laplace-Beltrami operator of M, and  $0 = \lambda_0 < \lambda_1 \le \lambda_2 \le \cdots$  be the eigenvalues of  $-\Delta$ . Denote  $e_i$  be the eigenfunctions of  $-\Delta$  corresponding  $\lambda_i$ . Then we have

**Theorem 6.** Let M be a d-dimensional smooth, closed and connected Riemann manifold embedded in  $\mathbb{R}^D$ , f be the smooth probability density function on M with infimum  $f_{min} > 0$  and  $\mathcal{X}_n = \{x_1, \ldots, x_n\}$  be the point cloud sampled following f independently and identically. Suppose eigenvalues of  $\Delta$  are simple. For fixed  $m \in \mathbb{N}$ , denote  $\Gamma_m = \min_{1 \le i \le m} dist(\lambda_i, \sigma(-\Delta) \setminus \{\lambda_i\})$ ,

where  $\sigma(-\Delta)$  is the spectrum of  $-\Delta$ . Suppose

$$h \le \mathcal{K}_1 \min\left(\left(\frac{\min(\Gamma_m, 1)}{\mathcal{K}_2 + \lambda_m^{d/2+5}}\right)^2, \frac{1}{(\mathcal{K}_3 + \lambda_m^{(5d+7)/4})^2}\right),\tag{90}$$

where  $\mathcal{K}_1$  and  $\mathcal{K}_2, \mathcal{K}_3 > 1$  are constants depending on d,  $f_{\min}$ ,  $||f||_{C^2}$ , and the volume, the injectivity radius, the curvature and the second fundamental form of the manifold. Then, when n is sufficiently large so that  $h = h(n) \ge (\frac{\log n}{n})^{\frac{1}{4d+13}}$ , with probability greater than  $1 - n^{-2}$ , for all  $0 \le i < m$ ,

$$|\mu_{i,n,h} - \lambda_i| \le \Omega_1 h^{3/2}.$$
(91)

And when n is sufficiently large so that  $h = h(n) \ge (\frac{\log n}{n})^{\frac{1}{4d+8}}$ , with probability greater than  $1 - n^{-2}$ , there are  $a_i \in \{1, -1\}$  such that for all  $0 \le i < m$ ,

$$\max_{x_j \in \mathcal{X}_n} |a_i v_{i,n,\epsilon}(j) - e_i(x_j)| \le \Omega_2 h^{1/2}.$$
(92)

 $\Omega_1$  depends on d, the the diameter of M,  $f_{\min}$ ,  $||f||_{C^2}$ , and  $\Omega_2$  depends on d, the diameter and the volume of M,  $f_{\min}$ , and  $||f||_{C^2}$ . [25]

**Remark 1.** In our setting, the sampling is uniform, thus  $f \equiv 1/\text{Vol}(M)$ , which implies  $||f||_{C^2} = 1/\text{Vol}(M)$ . In addition, d, diameter, volume, injectivity radius, curvature and second order fundamental form are bounded uniformly, which are compact, thus  $\mathcal{K}_1, \mathcal{K}_2, \mathcal{K}_3, \Omega_1, \Omega_2$  are bounded uniformly for all  $M \in \mathcal{M}$ .

**Remark 2.** In the case when the eigenvalues are not simple, the same proof still works by introducing the eigenprojection.[37]

**Remark 3.** If we choose  $h = (\frac{\log n}{n})^{\frac{1}{4d+13}}$ , then for large enough *n*, we have

$$|\mu_{i,n,h} - \lambda_i| \le \Omega_1 (\frac{\log n}{n})^{\frac{3}{8d+26}}, \quad \text{for } 0 \le i \le m.$$
 (93)

Similarly, if we choose  $h = \left(\frac{\log n}{n}\right)^{\frac{1}{4d+8}}$ , we have

$$\max_{x_j \in \mathcal{X}_n} |a_i v_{i,n,\epsilon}(j) - e_i(x_j)| \le \Omega_2 (\frac{\log n}{n})^{\frac{1}{8d+16}}, \quad \text{for } 0 \le i \le m.$$
(94)

Since  $(\frac{\log n}{n})^{\frac{1}{4d+13}} \ge (\frac{\log n}{n})^{\frac{1}{4d+8}}$ , the inequalities above hold at the same time if we choose  $h = (\frac{\log n}{n})^{\frac{1}{4d+13}}$ .

Now, we introduce our algorithm and estimate the approximation error.

Algorithm 1 Diffusion Map

- 1: Input: point cloud  $\mathcal{X}_n$ , intrinsic dimension d, the lower bound of injectivity radius  $\iota$  (corollary 4), the lower bound of Ricci curvature  $-\kappa$  (Assumption 1), upper bound of volume V (Assumption 3), norm error  $\epsilon$  (eq. (66)).
- 2: Calculate bindwidth  $h = (\frac{\log n}{n})^{\frac{1}{4d+13}}$ , diffusion time according to Theorem 1, Lemma 2, Lemma A.4

$$t = \min\left\{t_0(d,\kappa,\iota,\epsilon), 4, \frac{\iota^2}{4}\right\},\tag{95}$$

and embedding dimension according to Theorem 1,

$$m = N_0(d, \kappa, \iota, \epsilon, V, \frac{t}{2}) + 1 \tag{96}$$

- 3: Construct W and D with bandwidth h according to eq. (87).
- 4: Calculate first *m* eigenvalues and eigenfunctions  $\{\mu_{i,n,h}, \tilde{v}_{i,n,h}\}_{i=1}^{m}$  of  $L_n = \frac{D^{-1}W-I}{h^2}$ .
- 5: For  $1 \le i \le m$ , calculate

$$\mathbb{N}(i) = |B_h(x_i) \cap \{x_1, \dots, x_n\}|.$$
(97)

Calculate

$$\|\tilde{v}\|_{l^2(1/\hat{p})} = \sqrt{\frac{|S^{d-1}|h^d}{d}} \sum_{i=1}^n \frac{\tilde{v}^2(i)}{\mathbb{N}(i)},\tag{98}$$

and normalize

$$v_{i,n,h} = \frac{\tilde{v}_{i,n,h}}{\|\tilde{v}_{i,n,h}\|_{l^2(1/\hat{p})}}.$$
(99)

6: Embed  $x_j$  to  $(t)^{\frac{d+2}{4}}\sqrt{2}(4\pi)^{\frac{d}{4}} \left(e^{-\mu_{i,n,h}t/2}v_{i,n,h}(j)\right)_{i=1}^m$ **Output:** the embedding point clouds  $\left\{(t)^{\frac{d+2}{4}}\sqrt{2}(4\pi)^{\frac{d}{4}}\left(e^{-\mu_{i,n,h}t/2}v_{i,n,h}(j)\right)_{i=1}^m\right\}_{i=1}^n \subset \mathbb{R}^m$ .

We compare this embedding with embedding in Section 2 and estimate the approximation error. To estimate the error term, we consider the error of i-th component, for large enough n, i.e.

$$\left(\frac{\log n}{n}\right)^{\frac{1}{4d+13}} = h \le \mathcal{K}_1 \min\left(\left(\frac{\min(\Gamma_m, 1)}{\mathcal{K}_2 + \lambda_m^{d/2+5}}\right)^2, \frac{1}{(\mathcal{K}_3 + \lambda_m^{(5d+7)/4})^2}\right)$$
(100)

$$\iff n > \frac{1}{\mathcal{K}} \max\left(\left(\frac{\mathcal{K}_2 + \lambda_K^{d/2+5}}{\min(\Gamma_K, 1)}\right)^{8d+26}, (\mathcal{K}_3 + \lambda_K^{(5d+7)/4})^{8d+26}\right),\tag{101}$$

up to a log factor.

Then we have

$$|e^{-\mu_{i,n,h}t/2}v_{i,n,h}(j) - e^{-\lambda_{i}t/2}e_{i}(x_{j})| \le e^{-\mu_{i,n,h}t/2}|v_{i,n,h}(j) - e_{i}(x_{j})| + |e^{-\mu_{i,n,h}t/2} - e^{-\lambda_{i}t/2}|e_{i}(x_{j})|$$
(102)

$$\leq |v_{i,n,h}(j) - e_i(x_j)| + \frac{t}{2} |\mu_{i,n,h} - \lambda_i| ||e_i||_{l^{\infty}}$$
(103)

We also have

$$\|e_i\|_{l^{\infty}} \le \|e_i\|_{L^{\infty}} \le C\lambda_i^{\frac{n-1}{4}},\tag{104}$$

where C only depends on injectivity radius and the absolute value of the sectional curvature[38], which are compact set, thus C has a uniformly upper bound for all  $M \in \mathcal{M}$ . And we have proved  $\lambda_i$  are bounded from above for  $i \leq m$  with fixed m. Consequently,  $||e_i||_{l^{\infty}}$  has the uniformly bound, we denote it as  $C_{\mathcal{M}}$ .

Therefore,

$$|e^{-\mu_{i,n,h}t/2}v_{i,n,h}(j) - e^{-\lambda_{i}t/2}e_{i}(x_{j})| \le \Omega_{2}(\frac{\log n}{n})^{\frac{1}{8d+16}} + \frac{t}{2}C_{\mathcal{M}}\Omega_{1}(\frac{\log n}{n})^{\frac{3}{8d+26}}$$
(105)

$$= \Omega(\frac{\log n}{n})^{\frac{1}{8d+16}},$$
 (106)

hence

$$\left| \left( e^{-\mu_{i,n,h}t/2} \tilde{v}_{i,n,h}(j) \right)_{i=1}^m - \left( e^{-\lambda_i t/2} e_i(x_j) \right)_{i=1}^m \right| \le \sqrt{m} \Omega(\frac{\log n}{n})^{\frac{1}{8d+16}}$$
(107)

#### 4.1 Related Result

In this subsection, we discussed some other convergence results. In the following, we ignore the specific settings like sample size n, bandwidth h and others, we only focus on the convergence rate and norm, when discussing the convergence, we also ignore the constant. We denote the eigenfunctions and eigenvalues of  $\Delta$  as  $e_i$ ,  $\lambda_i$  and denote the eigenvectors and eigenvalues of graph laplacian as  $\mu_i$ ,  $v_i$ , here we will not specify laplacian graph and kernel function.

In [20], we have similar result, the following holds for finite k

$$\frac{1}{n}\sum_{i=1}^{n}(v_k(i) - e_k(x_i))^2 = O\left(\left(\frac{\log n}{n}\right)^{\frac{1}{2d}}\right),\tag{108}$$

$$|\mu_i - \lambda_i| = O\left(\left(\frac{\log n}{n}\right)^{\frac{1}{2d}}\right).$$
(109)

And the author then improved the result to  $O\left(\left(\frac{\log n}{n}\right)^{\frac{1}{d+4}}\right)$  under some other conditions in [39]. This convergence rate is much faster than we use, but the left side describe the average pointwise error. This result has also been improved [23], they obtained the convergence result for  $l^{\infty}$ , Lipschitz norm is  $O\left(\left(\frac{\log n}{n}\right)^{\frac{1}{d+4}}\right)$  for specific manifold, however the constant C in their result depends on M, which is difficult to quantify, so it cannot be applied to the family of manifolds.

In [19], the author derived the convergence rate of eigenvalues is  $O\left(\left(\frac{\log n}{n}\right)^{\frac{1}{d/2+2}}\right)$ , and the convergence rate of eigenvectors is  $O\left(\left(\frac{\log n}{n}\right)^{\frac{1}{d/2+3}}\right)$  under different settings, and it also includes the results we discussed above. In[21], the convergence rate is  $O\left(n^{-\frac{2}{(5d+6)(d+6)}}\right)$  under  $l^{\infty}$  norm, which is a slower than result we use.

## 5 Conclusion

In this paper, we proved that under bounded curvature, bounded from below reach, finite volume, smoothness and regular condition on Christoffel symbols expressed by Assumption 1 - Assumption 5, if the finite dimensional Diffusion Map embedding deviation from isometry is bounded, then the smoothness, local reach and sampling density of the manifold are preserved, uniformly. And the finite sample error is controlled by  $O\left(\left(\frac{\log n}{n}\right)^{\frac{1}{8d+16}}\right)$ . In addition, if we can control the global reach, we can also control the error for the estimate of tangent space.

#### References

[1] R. R. Coifman and S. Lafon, "Diffusion maps," *Applied and Computational Harmonic Analysis*, vol. 21, no. 1, pp. 5–30, 2006. Special Issue: Diffusion Maps and Wavelets.

- [2] M. Belkin and P. Niyogi, "Laplacian eigenmaps for dimensionality reduction and data representation," *Neural computation*, vol. 15, no. 6, pp. 1373–1396, 2003.
- [3] P. Bérard, G. Besson, and S. Gallot, "Embedding riemannian manifolds by their heat kernel," *Geometric & Functional Analysis GAFA*, vol. 4, pp. 373–398, 1994.
- [4] J. W. Portegies, "Embeddings of riemannian manifolds with heat kernels and eigenfunctions," *Communications on Pure and Applied Mathematics*, vol. 69, no. 3, pp. 478–518, 2016.
- [5] J. Bates, "The embedding dimension of laplacian eigenfunction maps," *Applied and Computational Harmonic Analysis*, vol. 37, no. 3, pp. 516–530, 2014.
- [6] S. Lafon and A. B. Lee, "Diffusion maps and coarse-graining: A unified framework for dimensionality reduction, graph partitioning, and data set parameterization," *IEEE transactions on pattern analysis and machine intelligence*, vol. 28, no. 9, pp. 1393–1403, 2006.
- [7] R. R. Coifman, I. G. Kevrekidis, S. Lafon, M. Maggioni, and B. Nadler, "Diffusion maps, reduction coordinates, and low dimensional representation of stochastic systems," *Multiscale Modeling & Simulation*, vol. 7, no. 2, pp. 842–864, 2008.
- [8] C. M. Arvizu and A. R. Messina, "Dimensionality reduction in transient simulations: A diffusion maps approach," *IEEE Transactions on Power Delivery*, vol. 31, no. 5, pp. 2379–2389, 2016.
- [9] L. Haghverdi, F. Buettner, and F. J. Theis, "Diffusion maps for high-dimensional single-cell analysis of differentiation data," *Bioinformatics*, vol. 31, no. 18, pp. 2989–2998, 2015.
- [10] L. Haghverdi, M. Büttner, F. A. Wolf, F. Buettner, and F. J. Theis, "Diffusion pseudotime robustly reconstructs lineage branching," *Nature methods*, vol. 13, no. 10, pp. 845–848, 2016.
- [11] A. L. Ferguson, A. Z. Panagiotopoulos, I. G. Kevrekidis, and P. G. Debenedetti, "Nonlinear dimensionality reduction in molecular simulation: The diffusion map approach," *Chemical Physics Letters*, vol. 509, no. 1-3, pp. 1–11, 2011.
- [12] L. Boninsegna, G. Gobbo, F. Noé, and C. Clementi, "Investigating molecular kinetics by variationally optimized diffusion maps," *Journal of chemical theory and computation*, vol. 11, no. 12, pp. 5947–5960, 2015.
- [13] P. Freeman, J. Newman, A. Lee, J. Richards, and C. Schafer, "Photometric redshift estimation using spectral connectivity analysis," *Monthly Notices of the Royal Astronomical Society*, vol. 398, no. 4, pp. 2012–2021, 2009.
- [14] U. Von Luxburg, M. Belkin, and O. Bousquet, "Consistency of spectral clustering," *The Annals of Statistics*, pp. 555–586, 2008.
- [15] M. Belkin and P. Niyogi, "Towards a theoretical foundation for laplacian-based manifold methods," *Journal of Computer and System Sciences*, vol. 74, no. 8, pp. 1289–1308, 2008.
- [16] E. Giné and V. Koltchinskii, "Empirical graph laplacian approximation of laplace-beltrami operators: large sample results," *Lecture Notes-Monograph Series*, pp. 238–259, 2006.
- [17] D. Ting, L. Huang, and M. I. Jordan, "An analysis of the convergence of graph laplacians," in *Proceedings of the 27th International Conference on International Conference on Machine Learning*, ICML'10, (Madison, WI, USA), p. 1079–1086, Omnipress, 2010.
- [18] M. Hein, J.-Y. Audibert, and U. v. Luxburg, "Graph laplacians and their convergence on random neighborhood graphs.," *Journal of Machine Learning Research*, vol. 8, no. 6, 2007.
- [19] X. Cheng and N. Wu, "Eigen-convergence of gaussian kernelized graph laplacian by manifold heat interpolation," *Applied and Computational Harmonic Analysis*, vol. 61, pp. 132–190, 2022.
- [20] N. García Trillos, M. Gerlach, M. Hein, and D. Slepčev, "Error estimates for spectral convergence of the graph laplacian on random geometric graphs toward the laplace-beltrami operator," *Foundations of Computational Mathematics*, vol. 20, no. 4, pp. 827–887, 2020.

- [21] X. Wang, "Spectral convergence rate of graph laplacian," *arXiv preprint arXiv:1510.08110*, 2015.
- [22] M. Belkin and P. Niyogi, "Convergence of laplacian eigenmaps," Advances in neural information processing systems, vol. 19, 2006.
- [23] J. Calder, N. Garcia Trillos, and M. Lewicka, "Lipschitz regularity of graph laplacians on random data clouds," *SIAM Journal on Mathematical Analysis*, vol. 54, no. 1, pp. 1169–1222, 2022.
- [24] P. W. Jones, M. Maggioni, and R. Schul, "Manifold parametrizations by eigenfunctions of the laplacian and heat kernels," *Proceedings of the National Academy of Sciences*, vol. 105, no. 6, pp. 1803–1808, 2008.
- [25] D. B. Dunson, H.-T. Wu, and N. Wu, "Spectral convergence of graph laplacian and heat kernel reconstruction in l<sup>∞</sup> from random samples," *Applied and Computational Harmonic Analysis*, vol. 55, pp. 282–336, 2021.
- [26] M. P. Do Carmo and J. Flaherty Francis, *Riemannian geometry*, vol. 2. Springer, 1992.
- [27] J. Lee, Introduction to Smooth Manifolds. Graduate Texts in Mathematics, Springer, 2003.
- [28] J. M. Lee, Introduction to Riemannian manifolds, vol. 2. Springer, 2018.
- [29] H. Federer, "Curvature measures," *Transactions of the American Mathematical Society*, vol. 93, no. 3, pp. 418–491, 1959.
- [30] E. Aamari, J. Kim, F. Chazal, B. Michel, A. Rinaldo, and L. Wasserman, "Estimating the reach of a manifold," *Electronic Journal of Statistics*, vol. 13, no. 1, pp. 1359 – 1399, 2019.
- [31] E. Aamari and C. Levrard, "Stability and minimax optimality of tangential delaunay complexes for manifold reconstruction," *Discrete & Computational Geometry*, vol. 59, pp. 923– 971, 2018.
- [32] P. Li and S. Yau, "Estimates of eigenvalues of a compact riemannian manifold," *Proc. Sympos. PureMath.*, vol. 36, 01 1980.
- [33] Y. Shi and B. Xu, "Gradient estimate of an eigenfunction on a compact riemannian manifold without boundary," Annals of Global Analysis and Geometry, vol. 38, 05 2009.
- [34] B. Xu, "Asymptotic behavior of *l*<sup>2</sup>-normalized eigenfunctions of the laplace-beltrami operator on a closed riemannian manifold," *arXiv preprint math/0509061*, 2005.
- [35] B. Xu, "Derivatives of the spectral function and sobolev norms of eigenfunctions on a closed riemannian manifold," *Annals of Global Analysis and Geometry - ANN GLOB ANAL GEOM*, vol. 26, pp. 231–252, 10 2004.
- [36] A. Hassannezhad, G. Kokarev, and I. Polterovich, "Eigenvalue inequalities on riemannian manifolds with a lower ricci curvature bound," *Journal of Spectral Theory*, vol. 6, no. 4, pp. 807– 835, 2016.
- [37] F. Chatelin, Spectral Approximation of Linear Operators. Society for Industrial and Applied Mathematics, 2011.
- [38] H. Donnelly, "Eigenfunctions of the Laplacian on Compact Riemannian Manifolds," Asian Journal of Mathematics, vol. 10, no. 1, pp. 115 – 126, 2006.
- [39] J. Calder and N. Garcia Trillos, "Improved spectral convergence rates for graph laplacians on ε-graphs and k-nn graphs," *Applied and Computational Harmonic Analysis*, vol. 60, pp. 123– 175, 2022.
- [40] H. Federer, Geometric Measure Theory. Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, Springer, 1969.

- [41] P. Li and S. Yau, "On the parabolic kernel of the schödinger operator," Acta Mathematica, vol. 156, pp. 153–201, 07 1986.
- [42] C. B. Croke, "Some isoperimetric inequalities and eigenvalue estimates," in Annales scientifiques de l'École normale supérieure, vol. 13, pp. 419–435, 1980.
- [43] F.-Y. Wang, "Sharp explicit lower bounds of heat kernels," *The Annals of Probability*, vol. 25, no. 4, pp. 1995–2006, 1997.

#### A Appendix / supplemental material

#### A.1 Pushforward Density

Here we list the lemma and theorem for estimating pushforward density.

**Lemma A.1.** (Area Formula) If  $f : \mathbb{R}^m \to \mathbb{R}^n$  is Lipschitzian and  $m \leq n$ , then

$$\int_{A} g(f(x)) J_f(x) d\mathcal{L}^m x = \int_{\mathbb{R}^n} g(y) N(f|A, y) d\mathcal{H}^m y$$

where A is an lebesgue measurable set,  $J_f(x)$  is the Jacobian  $\sqrt{\det(d_x f^T d_x f)}$  and  $g : \mathbb{R}^n \to \mathbb{R}$ and  $N(f|A, y) < \infty$  for  $\mathcal{H}^m$  almost all y.[40]

**Theorem A.2.** If P is distribution on  $\mathcal{M}$  with density f with respect to the d-dimensional Hausdorff measure and  $\varphi$  is a diffeomorphism, then the density g(p') with respect to d-dimensional Hausdorff measure of  $P' := \varphi_{\#}P$  is

$$g(p') = f(p) / \sqrt{\det\left(\pi_{T_pM} \circ d\varphi_p^T \circ d\varphi_p|_{T_pM}\right)}$$

where  $p = \varphi^{-1}(p')$ .

*Proof.* Let  $p \in \mathcal{M}$  be a fixed point and choose r small enough such that exponential map  $\Psi_p : T_p \mathcal{M} \to \mathbb{R}^D$  is an injection onto  $B(p, r) \cap \mathcal{M}$ .

Choose  $A \subset B(p, r) \cap \mathcal{M}$ , by definition of pushforward measure, we have

$$\int_{\varphi(A)} dP' = \int_A dP = \int_A f(y) d\mathcal{H}^d y$$

Since  $\Psi_p$  is an injection, so  $N(\Psi_p|\Psi_p^{-1}(A), y) = 1$  for all  $y \in A$  and 0 otherwise. By Lemma A.1, we have

$$\int_{A} f(y) d\mathcal{H}^{d} y = \int_{\Psi_{p}^{-1}(A)} f(\Psi_{p}(x)) J_{\Psi_{p}}(x) d\mathcal{L}^{d} x$$

And  $\varphi$  is diffeomorphism, so  $h = \varphi \circ \Psi_p$  is also injective, so

$$\int_{\Psi_p^{-1}(A)} f(\Psi_p(x)) J_{\Psi_p}(x) d\mathcal{L}^d x = \int_{\Psi_p^{-1}(A)} f(\varphi^{-1}(h(x))) \frac{J_{\Psi_p}(h^{-1}(h(x)))}{J_{\varphi \circ \Psi_p}(h^{-1}(h(x)))} J_{\varphi \circ \Psi_p}(x)) d\mathcal{L}^d x$$
$$= \int_{\varphi(A)} f(\varphi^{-1}(z)) \frac{J_{\Psi_p}(h^{-1}(z))}{J_{\varphi \circ \Psi_p}(h^{-1}(z))} d\mathcal{H}^d z$$

Thus

$$\int_{\varphi(A)} dP' = \int_{\varphi(A)} f(\varphi^{-1}(z)) \frac{J_{\Psi_p}(h^{-1}(z))}{J_{\varphi \circ \Psi_p}(h^{-1}(z))} d\mathcal{H}^d z$$

 $T_p\mathcal{M}$  is a subspace of  $\mathbb{R}^D$  with dimension d, so we can choose basis of  $T_p\mathcal{M}$  such that all elements in diagonal of transformation matrix are 1. We notice that  $h^{-1}(z) = \Psi_{\varphi^{-1}(z)}^{-1} \circ \varphi^{-1}(z) = 0$ , and  $d_0\Psi_p = I_D + d_0N_p = I_D$  is inclusion map. Thus

$$f(\varphi^{-1}(z))\frac{J_{\Psi_p}(h^{-1}(z))}{J_{\varphi\circ\Psi_p}(h^{-1}(z))} = f(p)/\sqrt{\det\left(\pi_{T_pM} \circ d\varphi_p^T \circ d\varphi_p|_{T_pM}\right)}$$

where  $p = \varphi^{-1}(z)$ .

#### A.2 Heat Kernel Estimate

**Theorem A.3** (Upper Bound for The Heat Kernel). Let M be a complete manifold without boundary. If the Ricci curvature of M is bounded from below by  $-\kappa$  for some constant  $\kappa \ge 0$ , then for any  $1 < \alpha_1 < 2$  and  $0 < \alpha_2 < 1$ , the heat kernel satisfies:

$$K(t,p,q) \le C(\alpha_2)^{\alpha_1} V^{-1/2} (B_p(\sqrt{t})) V^{-1/2} (B_q(\sqrt{t})) \exp\left(C(d)\alpha_2(\alpha_1-1)^{-1} \kappa t - \frac{d(p,q)^2}{(4+\alpha_2)t}\right),$$
(110)

where  $B_x(r)$  is geodesic ball centered at p with radius r,  $C(\alpha_2)$  depends on  $\alpha_2$  with  $C(\alpha_2) \to \infty$ as  $\alpha_2 \to 0.[41]$ 

**Lemma A.4** (Estimate for Volume of Geodesic Ball). Let M be a complete manifold without boundary. then for  $r \le \iota_M/2$ , we have

$$Vol(B_p(r)) \ge C'(d)r^d,\tag{111}$$

where  $C'(d) = \frac{2^{d-1}\omega(d-1)^d}{d^d\omega(d)^{d-1}}$ , and  $\omega(d) = \frac{2\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2})}$  is the volume of the unit d-dimensional sphere [42]. Thus  $C'(d) = \frac{2^d \Gamma(\frac{d}{2})^{d-1}}{d^d \Gamma(\frac{d-1}{2})^d}$ .

**Theorem A.5** (Lower Bound for Heat Kernel). Let M be a complete Riemannian manifold of dimension d with Ricci curvature is greater than  $-\kappa(d-1)$  for some  $\kappa \ge 0$ . For any  $t, \sigma > 0$  and  $p, q \in M$ , we have

$$K(t, p, q) \ge (4\pi t)^{-d/2} \exp\left(-\left(\frac{1}{4t} + \frac{\sigma}{3\sqrt{2t}}\right) d(p, q)^2 - \frac{\beta^2}{4}t - \left(\frac{\beta^2}{4\sigma} + \frac{2d\sigma}{3}\right)\sqrt{2t}\right), \quad (112)$$

where  $\beta = \sqrt{\kappa}(d-1).[43]$ 

#### A.3 Global Reach

**Lemma A.6.** If  $d(p,q) \leq s_0$  where  $s_0 = 2\sqrt{2\tau_l}$ , then p, q cannot satisfy global reach case.

*Proof.* If not, we assume there exists  $p, q \in M$  such that  $d(p,q) \leq s_0$  and p,q satisfy global reach case, i.e.

$$\frac{p+q}{2} \in Med(M),\tag{113}$$

$$\pi(\frac{p+q}{2}) = p, q \text{ and } d_E(p, M) = d_E(q, M) = |p-q|/2$$
 (114)

Since  $d(p,q) \le s_0$ , we have  $d(p,q) \le \frac{3}{2}|p,q|$ .




Figure 3: Local Estimate

The inequality above implies the geodesic  $\gamma(s)$  connecting p and q must be partly inside the ball  $B(\frac{p+q}{2}, \frac{|p-q|}{2})$ , since  $\overline{pq}$  is the diameter of this ball and

$$|\gamma(s)| = d(p,q) \le \frac{3}{2}|p-q| < \frac{\pi}{2}|p-q|,$$
(115)

which implies there exist  $a \in (0, s)$  such that

$$|\gamma(a) - \frac{p+q}{2}| < \frac{|p-q|}{2},\tag{116}$$

which contradicts with  $\varphi(p), \varphi(q)$  satisfy global reach case eq. (114).