

Free-Surface Equatorial Flows with Surface Tension in Spherical Coordinates

Andrei Stan^{1,2}

¹ Department of Mathematics, Babeş-Bolyai University, Cluj-Napoca, 400084, Romania.

² Tiberiu Popoviciu Institute of Numerical Analysis, Romanian Academy, Cluj-Napoca, 400110, Romania.

Contributing authors: andrei.stan@ubbcluj.ro;

Abstract

In this paper, we determine an exact solution to the governing equations in spherical coordinates for an inviscid, incompressible fluid. This solution describes a steady, purely azimuthal equatorial flow with an associated free surface. Using functional analytic techniques, we demonstrate that if a free surface is known beforehand, the variations in pressure needed to achieve this surface implicitly define the shape of the free surface in a unique way.

Keywords: Azimuthal flows, Surface tension, General density, Implicit function theorem, Coriolis force

AMS Subject Classification: 35Q31 , 35Q35 , 35Q86 , 35R35 , 26B10

1 Introduction

This paper focuses on developing an exact solution for the governing equations of geophysical fluid dynamics (GFD) governing inviscid, incompressible, and stratified fluid dynamics in the equatorial region. Specifically, we investigate a steady purely azimuthal flow. Our approach ensures accuracy by consistently employing spherical coordinates, thereby avoiding any simplifications to the geometry within the governing equations. Additionally, within these equations, we incorporate the complexities of Coriolis and centripetal forces, along with the surface tension resulting in highly nonlinear dynamics [1-3].

This study builds upon recent significant advancements in the field. In [1, 4], the authors shown for the first time that exact solutions to the complete governing equations of geophysical fluid dynamics (GFD) can be formulated using spherical coordinates, representing purely azimuthal, depth-varying flows. These solutions are capable of modeling both the equatorial undercurrent (EUC) and the Antarctic Circumpolar Current (ACC). Initially, these solutions describe purely homogeneous fluids without stratification. Subsequently, in works by the authors [7, 8], exact equatorial flow solutions allowing for stratification were developed. These solutions, while relatively simple, feature fluid density varying linearly with depth and remaining independent of latitude. Finally, in [9], the authors established the existence of solutions in the context of general stratification, a paper that the present work aims to build upon and enhance.

In previous works (see, e.g., [6, 9, 10]), the existence and uniqueness of the free surface are guaranteed only for pressure variations close to the pressure required to achieve a flat, undisturbed surface. The novelty of this paper is that if a free surface γ is known beforehand, a similar conclusion holds: sufficiently small variations in pressure from the value required to obtain γ lead to a unique free surface. From a physical point of view, such a result is expected, as small variations in pressure that determine a smooth enough free surface should also result in a similarly smooth free surface.

2 Preliminaries

Throughout this paper we employ spherical coordinates (r, θ, ϕ) , where r represents distance to the sphere's center, θ ranges from 0 to π (with $\frac{\pi}{2} - \theta$ denoting latitude), and ϕ spans from 0 to 2π (longitude). Notably, the North and South poles correspond to $\theta = 0$ and π , respectively, while the Equator lies at $\theta = \frac{\pi}{2}$. Unit vectors in this system, denoted as \mathbf{e}_r , \mathbf{e}_θ , and \mathbf{e}_ϕ , provide directional references, with \mathbf{e}_ϕ from West to East and \mathbf{e}_θ from North to South.

The equations describing the motion of an inviscid and incompressible flow are comprised in the Euler equations and, respectively, mass conservation equation. In spherical coordinates, they are given by

$$\begin{aligned} u_t + uu_r + \frac{v}{r}u_\theta + \frac{w}{r \sin \theta}u_\phi - \frac{1}{r}(v^2 + w^2) &= -\frac{1}{\rho}p_r + F_r \\ v_t + uv_r + \frac{v}{r}v_\theta + \frac{w}{r \sin \theta}v_\phi + \frac{1}{r}(uv - w^2 \cos \theta) &= -\frac{1}{\rho} \frac{1}{r}p_\theta + F_\theta \\ w_t + uw_r + \frac{v}{r}w_\theta + \frac{w}{r \sin \theta}w_\phi + \frac{1}{r}(uw + vw \cot \theta) &= -\frac{1}{\rho} \frac{1}{r \sin \theta}p_\phi + F_\phi, \end{aligned} \quad (1)$$

and

$$\frac{1}{r^2} \frac{\partial}{\partial r}(\rho r^2 u) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta}(\rho v \sin \theta) + \frac{1}{r \sin \theta} \frac{\partial(\rho w)}{\partial \phi} = 0. \quad (2)$$

Here, $\mathbf{u} = u\mathbf{e}_r + v\mathbf{e}_\theta + w\mathbf{e}_\phi$ is the velocity field, $p(r, \theta, \phi)$ represents the pressure, (F_r, F_θ, F_ϕ) the body-force vector, while $\rho = \rho(r, \theta)$ stands for the density distribution.

To ensure an accurate analysis of fluid dynamics in specific locations, it is crucial to consider the effects of Earth's rotation. Additional terms must be integrated into

the Euler equations to account for these effects, namely the Coriolis force $2\boldsymbol{\Omega} \times \mathbf{u}$ and the centripetal acceleration $\boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{r})$, where

$$\boldsymbol{\Omega} = \Omega(\mathbf{e}_r \cos \theta - \mathbf{e}_\theta \sin \theta),$$

Here, $\Omega \approx 7.29 \times 10^{-5}$ rad/s represents the Earth's constant rate of rotation and $\mathbf{r} = r\mathbf{e}_r$. These forces are incorporated into the Euler equations. Consequently, with gravity given by the vector $(-g, 0, 0)$, and the latter two quantities combined,

$$\begin{aligned} 2\boldsymbol{\Omega} \times \mathbf{u} + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{r}) &= 2\Omega(-w \sin \theta \mathbf{e}_r - w \cos \theta \mathbf{e}_\theta + (u \sin \theta + v \cos \theta) \mathbf{e}_\phi) \\ &\quad - r\Omega^2(\sin^2 \theta \mathbf{e}_r + \sin \theta \cos \theta \mathbf{e}_\phi), \end{aligned}$$

the Euler's equations become

$$\begin{aligned} uu_r + \frac{v}{r}u_\theta + \frac{w}{r \sin \theta}u_\phi - \frac{1}{r}(v^2 + w^2) - 2\Omega w \sin \theta - r\Omega^2 \sin^2 \theta &= -\frac{1}{\rho}p_r - g \\ uv_r + \frac{v}{r}v_\theta + \frac{w}{r \sin \theta}v_\phi + \frac{1}{r}(uv - w^2 \cos \theta) - 2\Omega w \cos \theta - r\Omega^2 \sin \theta \cos \theta &= -\frac{1}{\rho}p_\theta \\ uw_r + \frac{v}{r}w_\theta + \frac{w}{r \sin \theta}w_\phi + \frac{1}{r}(uw + vw \cot \theta) + 2\Omega(u \sin \theta + v \cos \theta) &= -\frac{1}{\rho r \sin \theta}p_\phi, \end{aligned} \tag{3}$$

where we considered $u_t = v_t = w_t = 0$. In addition to the mass conservation and Euler equations, the movement of water is influenced by boundary conditions. As the fluid extends infinitely in all horizontal directions, it encounters two boundaries: the rigid flat bed and the water's free surface. On the bed where $r = d(\theta, \phi)$, we apply the kinematic boundary condition,

$$u = \frac{v}{r}d_\theta + \frac{w}{r \sin \theta}d_\phi,$$

while on the free surface where $r = R + h(\theta, z)$, we impose the kinematic boundary condition

$$u = \frac{v}{r}h_\theta + \frac{w}{r \sin \theta}h_\phi.$$

Additionally, on the free surface, we have the dynamic condition

$$p = P(\theta, \phi) + \sigma \nabla \cdot \vec{n}. \tag{4}$$

Here, σ represents the coefficient of surface tension, and \vec{n} is the outward-pointing unit normal vector. The function h is currently unknown and will be determined later.

Next, we recall two well known results from literature used in this paper. We start with the Implicit Function Theorem, which ensures the existence of nontrivial zeros of a C^1 mapping between Banach spaces (see, e.g., [11]).

Theorem 1. *Let X, Y, Z be Banach spaces, $U \subset X \times Y$ an open neighbourhood of a point $(x_0, y_0) \in X \times Y$ and let $f: U \rightarrow Z$ be a continuous functions. Assume that:*

- i) The function f satisfies $f(x_0, y_0) = 0$.
ii) The partial derivative $f_y(x_0, y_0)$ exists and is an linear homeomorphism from Y to Z .

Then, there exists an open neighbourhood U_1 of x_0 and a unique $g: U_1 \rightarrow Y$ continuous function such that $g(x_0) = y_0$ and $f(x, g(x)) = 0$ on U_1 .

The second result concerns the existence and uniqueness of solution for a second order differential equation (see, e.g., [12, 13]).

Theorem 2. *The initial value problem*

$$y'' + p(t)y' + q(t)y = g(t), \quad y(t_0) = y'(t_0) = 0, \quad (5)$$

where p, q , and g are continuous functions on an open interval I that contains the point t_0 , has a unique solution. In addition, the solution $y(t)$ is given by

$$y(t) = y_p + \int_{t_0}^t \frac{g(s) (\Phi_1(t)W_1(s) + \Phi_2(t)W_2(s))}{W(\Phi_1, \Phi_2)(s)} ds,$$

where y_p is a particular solution of (5), $\{\Phi_1, \Phi_2\}$ is a basis for the solutions of the homogeneous equation $y'' + p(t)y' + q(t)y = 0$, W is the Wronskian of the basis $\{\Phi_1, \Phi_2\}$ and W_i ($i=1,2$) is the Wronskian obtained by replacing the i -th column of W with the column vector $(0, 1)$.

3 Main result

In the sequel, we assume an azimuthal flow, i.e., the velocity field $\mathbf{u} = u\mathbf{e}_r + v\mathbf{e}_\theta + w\mathbf{e}_\phi$ satisfies $u = v = 0$ and $w = w(r, \theta)$. Also, we assume the free surface satisfy $h = h(\theta)$ while the flat bed is a fixed real value d . Consequently, the mass conservation equation (2) is automatically satisfied, while the Euler's equations (3) takes the form

$$\begin{cases} -\frac{w^2}{r} - 2\Omega w \sin \theta - r\Omega^2 \sin^2 \theta = -\frac{1}{\rho}p_r - g, \\ -\frac{w^2}{r} \cot \theta - 2\Omega w \cos \theta - r\Omega^2 \sin \theta \cos \theta = -\frac{1}{\rho r}p_\theta, \\ 0 = -\frac{1}{\rho} \frac{1}{r \sin \theta} p_\phi, \end{cases} \quad (6)$$

or equivalently

$$\begin{cases} \frac{\rho(r, \theta)}{r} (w + \Omega r \sin \theta)^2 = p_r + g\rho(r, \theta) \\ \rho(r, \theta) \cot \theta (w + \Omega r \sin \theta)^2 = p_\theta \\ 0 = p_\phi. \end{cases} \quad (7)$$

From the last relation, we obtain the independence of the pressure with respect to ϕ , i.e., $p = p(r, \theta)$. If we denote

$$Z = Z(r, \theta) := \frac{1}{r} (w + \Omega r \sin \theta)^2,$$

and differentiate the first equation with respect to θ and the second one with respect to r , we deduce

$$(\rho Z)_\theta - (\rho r \cot \theta Z)_r = g\rho\theta,$$

which yields

$$-(r \cos \theta)U_r + (\sin \theta)U_\theta = (r \sin \theta)g\rho_\theta(r, \theta), \quad (8)$$

where $U(r, \theta) = \rho(r, \theta) (w + \Omega r \sin \theta)^2$.

Employing the method of characteristics and following steps similar to those in [9, 10], we infer

$$w(r, \theta) = -\Omega r \sin \theta + \frac{1}{\sqrt{\rho(r, \theta)}} \left(F(r \sin \theta) + gr \sin \theta \int_0^{f(\theta)} \rho_\theta(\bar{r}(s), \bar{\theta}(s)) ds \right)^{\frac{1}{2}}, \quad (9)$$

where

$$\bar{r}(s) = \frac{r \sin \theta}{2} (e^s + e^{-s}), \quad \bar{\theta}(s) = \arccos \left(\frac{1 - e^{2s}}{1 + e^{2s}} \right), \quad f(\theta) = \frac{1}{2} \ln \frac{1 - \cos \theta}{1 + \cos \theta},$$

and $F(\xi) := U(\xi, \frac{\pi}{2})$ is an arbitrary smooth function. Note that

$$\bar{\theta}(f(\theta)) = \theta \text{ and } \bar{r}(f(\theta)) = r.$$

To proceed with the determination of the pressure, one sees that

$$\begin{aligned} p_r + g\rho(r, \theta) &= \frac{\rho(r, \theta)}{r} (w + \Omega r \sin \theta)^2 \\ &= \frac{1}{r} \left(F(r \sin \theta) + gr \sin \theta \int_0^{f(\theta)} \rho_\theta(\bar{r}(s), \bar{\theta}(s)) ds \right), \end{aligned} \quad (10)$$

and

$$p_\theta = \cot \theta U(r, \theta) = \cot \theta \left(F(r \sin \theta) + gr \sin \theta \int_0^{f(\theta)} \rho_\theta(\bar{r}(s), \bar{\theta}(s)) ds \right). \quad (11)$$

After integration from a (an arbitrary constant) to r in (10), we infer

$$p(r, \theta) = C(\theta) - g \int_r^a \rho(s, \theta) ds + \int_{a \sin \theta}^{r \sin \theta} \frac{F(s)}{s} ds + \mathcal{F}(s, \theta) dy, \quad (12)$$

where $C(\theta) := p(a, \theta)$ is an arbitrary smooth function and

$$\mathcal{F}(\xi, \theta) = \int_0^{f(\theta)} g\rho_\theta \left(\xi \frac{e^s + e^{-s}}{2}, \bar{\theta}(s) \right) ds.$$

To determine the function C , we differentiate with respect to θ in (12), which gives

$$\begin{aligned} p_\theta &= C'(\theta) - g \int_0^r \rho_\theta(s, \theta) ds + \cot \theta (F(r \sin \theta) - F(a \sin \theta)) \\ &\quad + \mathcal{F}(r \sin \theta) r \cos \theta - \mathcal{F}(a \sin \theta) a \cos \theta + \int_{a \sin \theta}^{r \sin \theta} \mathcal{F}_\theta(s, \theta) ds. \end{aligned}$$

Since

$$\begin{aligned} \mathcal{F}_\theta(s, \theta) &= f'(\theta) g \rho_\theta \left(\frac{\bar{s}(f(\theta))}{\sin \theta}, \bar{\theta}(f(\theta)) \right) \\ &= \frac{g}{\sin \theta} \rho_\theta \left(\frac{s}{\sin \theta}, \theta \right), \end{aligned}$$

we obtain

$$\int_{a \sin \theta}^{r \sin \theta} \mathcal{F}_\theta(s, \theta) ds = \int_a^r g \rho_\theta(s, \theta) ds.$$

Hence, the expression of p_θ becomes

$$p_\theta = C'(\theta) + \cot \theta [F(r \sin \theta) - F(a \sin \theta)] + \mathcal{F}(r \sin \theta) r \cos \theta - \mathcal{F}(a \sin \theta) a \cos \theta. \quad (13)$$

Consequently, from (11) and (13), we deduce

$$C'(\theta) = F(a \sin \theta) \cot \theta + F(a \sin \theta) a \cos \theta. \quad (14)$$

Finally, relations (12) and (14) yields,

$$p(r, \theta) = b - g \int_a^r \rho(s, \theta) ds + \int_{a \sin \theta}^{r \sin \theta} \left(\frac{F(s)}{s} + \mathcal{F}(s, \theta) \right) ds \quad (15)$$

$$+ \int_{\frac{\pi}{2}}^\theta (F(a \sin s) \cot s + \mathcal{F}(a \sin s) a \cos s) ds, \quad (16)$$

where a, b are arbitrary real numbers.

3.1 The dynamic condition

In the sequel, we analyze the dynamic boundary condition (4) on the free surface $r = R + h(\theta)$. Since we assumed an azimuthal flow and h is independent of the depth, relation (4) takes the form

$$p(R + h(\theta), \theta) = P(\theta) + \sigma \nabla \cdot \vec{n}, \quad (17)$$

where P is the known pressure on the surface. Since the implicit equation of the free surface is $H(r, \theta, z) := r - R - h(\theta) = 0$ and the gradient in spherical coordinates has

the representation (see, e.g., [15, Chapter 1]),

$$\nabla = \mathbf{e}_r \partial_r + \mathbf{e}_\theta \frac{1}{r} \partial_\theta + \mathbf{e}_\phi \frac{1}{r \sin \theta} \partial_\phi,$$

we obtain the normal derivative to the surface H as

$$\vec{N} = \mathbf{e}_r - \frac{h_\theta}{r} \mathbf{e}_\theta.$$

Therefore, the pointing unit normal vector is

$$\vec{n} = \frac{\vec{N}}{\|\vec{N}\|} = \frac{r}{\sqrt{r^2 + h_\theta^2}} \cdot \vec{N} = \frac{r}{\sqrt{r^2 + h_\theta^2}} \mathbf{e}_r - \frac{h_\theta}{\sqrt{r^2 + h_\theta^2}} \mathbf{e}_\theta.$$

Whence, given the the divergence in spherical coordinates (see, e.g., [15, Chapter 1]),

$$\nabla \cdot w = \frac{1}{r^2} \partial_r (r^2 w_r) + \frac{1}{r \sin \theta} \partial_\theta (\sin \theta w_\theta) + \frac{1}{r \sin \theta} \partial_\phi w_\phi,$$

one has,

$$\begin{aligned} \nabla \cdot \vec{n} &= \frac{1}{r^2} \partial_r (r^2 n_r) + \frac{1}{r \sin \theta} \partial_\theta (n_\theta \sin \theta) & (18) \\ &= \frac{2r^2 + 3h_\theta^2}{(r^2 + h_\theta^2)^{\frac{3}{2}}} - \frac{r h_{\theta\theta}}{(r^2 + h_\theta^2)^{\frac{3}{2}}} - \frac{\cot \theta h_\theta}{r (r^2 + h_\theta^2)^{\frac{1}{2}}} \\ &= \frac{2(R + h(\theta))^2 + 3h_\theta^2}{((R + h(\theta))^2 + h_\theta^2)^{\frac{3}{2}}} - \frac{(R + h(\theta)) h_{\theta\theta}}{((R + h(\theta))^2 + h_\theta^2)^{\frac{3}{2}}} \\ &\quad - \frac{\cot \theta h_\theta}{(R + h(\theta)) ((R + h(\theta))^2 + h_\theta^2)^{\frac{1}{2}}}. \end{aligned}$$

Finally, from (17) and (18), we find the pressure on the free surface in the form of a Bernoulli type problem

$$\begin{aligned} P(\theta) &= b - g \int_a^{R+h(\theta)} \rho(s, \theta) ds + \int_{a \sin \theta}^{(R+h(\theta)) \sin \theta} \left(\frac{F(s)}{s} + \mathcal{F}(s, \theta) \right) ds & (19) \\ &\quad + \int_{\frac{\pi}{2}}^\theta (F(a \sin s) \cot s + \mathcal{F}(a \sin s) a \cos s) ds \\ &\quad - \sigma \frac{2(R + h(\theta))^2 + 3h_\theta^2}{((R + h(\theta))^2 + h_\theta^2)^{\frac{3}{2}}} + \sigma \frac{h_{\theta\theta} (R + h(\theta))}{((R + h(\theta))^2 + h_\theta^2)^{\frac{3}{2}}} \\ &\quad + \sigma \frac{\cot \theta h_\theta}{(R + h(\theta)) ((R + h(\theta))^2 + h_\theta^2)^{\frac{1}{2}}}. \end{aligned}$$

3.2 The existence of solutions for the free surface

To have a meaningful comparison of the physical quantities, we start by non-dimensionalizing equation (19). For this, we denote with P_{atm} the pressure corresponding to an undisturbed free surface, i.e., the pressure P obtained by setting $h \equiv 0$ in (19). Subsequently, we denote

$$\wp = \frac{P}{P_{atm}} \text{ and } \mathbf{h} = \frac{h}{R}. \quad (20)$$

Whence, relation (19) together with (20), leads us to the functional equation

$$\mathbb{F}(\mathbf{h}, \wp) = 0, \quad (21)$$

where

$$\begin{aligned} \mathbb{F}(\mathbf{h}, \wp) = & -\wp(\theta) - \frac{g}{P_{atm}} \int_a^{R(1+\mathbf{h}(\theta))} \rho(s, \theta) ds \\ & + \frac{1}{P_{atm}} \int_{a \sin \theta}^{R(1+\mathbf{h}(\theta)) \sin \theta} \left(\frac{F(s)}{s} + \mathcal{F}(s, \theta) \right) ds \\ & - \frac{\sigma}{RP_{atm}} \left(\frac{(2(1+\mathbf{h})^2 + 3\mathbf{h}_\theta^2)}{((1+\mathbf{h})^2 + \mathbf{h}_\theta^2)^{\frac{3}{2}}} - \frac{\mathbf{h}_{\theta\theta}(1+\mathbf{h})}{((1+\mathbf{h})^2 + \mathbf{h}_\theta^2)^{\frac{3}{2}}} - \frac{\cot \theta \mathbf{h}_\theta}{(1+\mathbf{h})((1+\mathbf{h})^2 + \mathbf{h}_\theta^2)^{\frac{1}{2}}} \right) \\ & + \varphi(\theta), \end{aligned} \quad (22)$$

and

$$\varphi(\theta) = \frac{b}{P_{atm}} + \frac{1}{P_{atm}} \int_{\pi/2}^{\theta} (F(a \sin s) \cot s + \mathcal{F}(a \sin s) a \cos s) ds.$$

Our goal is to establish a relationship between the pressure variations at the free surface and the shape of the surface, which corresponds to finding a nontrivial solution to the functional equation (21). To achieve this, we assume the existence of a smooth enough "trivial" free surface, denoted by $\bar{\gamma}$, which is derived from specific measurements. The non-dimensionalized pressure \wp_γ required to maintain this shape is obtained by substituting \mathbf{h} with $\gamma = \frac{\bar{\gamma}}{R}$ in (22).

Remark 1. *If $\bar{\gamma} = 0$, then \wp_γ represents the pressure required to maintain a flat, undisturbed free surface.*

Let

$$C_\gamma = \left\{ u \in C^2 \left[\frac{\pi}{2} - \varepsilon, \frac{\pi}{2} + \varepsilon \right] : u \left(\frac{\pi}{2} \right) = \gamma \left(\frac{\pi}{2} \right), u' \left(\frac{\pi}{2} \right) = \gamma' \left(\frac{\pi}{2} \right) \right\},$$

where $\varepsilon = 0.016$ corresponds to a strip approximately 100 km wide around the Equator [5]. Clearly, \mathbb{F} defines a continuously differentiable map

$$\mathbb{F} : C_\gamma \times C \left[\frac{\pi}{2} - \varepsilon, \frac{\pi}{2} + \varepsilon \right] \rightarrow C \left[\frac{\pi}{2} - \varepsilon, \frac{\pi}{2} + \varepsilon \right].$$

In the subsequent, we analyze the Fréchet derivative of \mathbb{F} with respect to the first argument. To this aim, we have

$$\begin{aligned}\mathbb{F}(t\mathbf{h} + \gamma, \wp_\gamma) - \mathbb{F}(\gamma, \wp_\gamma) &= -\frac{g}{P_{atm}} \int_{R(1+\gamma)}^{R(1+t\mathbf{h}+\gamma)} \rho(s, \theta) ds \\ &\quad + \frac{1}{P_{atm}} \int_{R(1+\gamma) \sin \theta}^{R(1+t\mathbf{h}+\gamma) \sin \theta} \left(\frac{F(s)}{s} + \mathcal{F}(s, \theta) \right) ds \\ &\quad - (J(t\mathbf{h} + \gamma) - J(\gamma)),\end{aligned}$$

where

$$J(v) := \frac{\sigma}{RP_{atm}} \left(\frac{2(1+v)^2 + 3v_\theta^2}{((1+v)^2 + v_\theta^2)^{\frac{3}{2}}} - \frac{v_{\theta\theta}(1+v)}{((1+v)^2 + v_\theta^2)^{\frac{3}{2}}} - \frac{\cot \theta v_\theta}{(1+v)((1+v)^2 + v_\theta^2)^{\frac{1}{2}}} \right).$$

Using the mean value theorem, we easily derive

$$\begin{aligned}\lim_{t \rightarrow 0} \frac{1}{t} \left(-\frac{g}{P_{atm}} \int_{R(1+\gamma)}^{R(1+t\mathbf{h}+\gamma)} \rho(s, \theta) ds \right) &+ \frac{1}{P_{atm}} \int_{R(1+\gamma) \sin \theta}^{R(1+t\mathbf{h}+\gamma) \sin \theta} \left(\frac{F(s)}{s} + \mathcal{F}(s, \theta) \right) ds \\ &= \psi(\theta)\mathbf{h},\end{aligned}$$

where

$$\begin{aligned}\psi(\theta) &= -\frac{Rg}{P_{atm}} \rho(R(1+\gamma), \theta) + \frac{R \sin \theta}{P_{atm}} \left(\frac{F(R(1+\gamma) \sin \theta)}{R(1+\gamma) \sin \theta} + \mathcal{F}(R(1+\gamma) \sin \theta, \theta) \right) \\ &= \frac{\rho(R(1+\gamma), \theta)}{P_{atm}} (-gR + (w(R(1+\gamma), \theta) + \Omega R(1+\gamma) \sin \theta)^2).\end{aligned}$$

To compute the derivative of J in γ , we denote

$$\begin{aligned}a(t) &= \frac{2(1+t\mathbf{h}+\gamma)^2 + 3(t\mathbf{h}_\theta + \gamma_\theta)^2}{\left((1+t\mathbf{h}+\gamma)^2 + (t\mathbf{h}_\theta + \gamma_\theta)^2 \right)^{\frac{3}{2}}}, \\ b(t) &= \frac{(t\mathbf{h}_{\theta\theta} + \gamma_{\theta\theta})(1+t\mathbf{h}+\gamma)}{\left((1+t\mathbf{h}+\gamma)^2 + (t\mathbf{h}_\theta + \gamma_\theta)^2 \right)^{\frac{3}{2}}}, \\ c(t) &= \frac{\cot \theta (t\mathbf{h}_\theta + \gamma_\theta)}{(1+t\mathbf{h}+\gamma) \left((1+t\mathbf{h}+\gamma)^2 + (t\mathbf{h}_\theta + \gamma_\theta)^2 \right)^{\frac{1}{2}}}.\end{aligned}$$

Since a, b, c are smooth functions around 0, one has

$$\lim_{t \rightarrow 0} \frac{1}{t} (J(t\mathbf{h} + \gamma) - J(\gamma)) = \frac{\sigma}{RP_{atm}} (a'(0) - b'(0) - c'(0)).$$

Simple computations yield

$$\begin{aligned}
a'(0) &= \frac{4\mathbf{h}(1+\gamma) + 6\mathbf{h}_\theta\gamma_\theta}{((1+\gamma)^2 + \gamma_\theta^2)^{\frac{3}{2}}} - 3 \left(2(1+\gamma)^2 + 3\gamma_\theta^2 \right) \frac{(\mathbf{h}(1+\gamma) + \mathbf{h}_\theta\gamma_\theta)}{((1+\gamma)^2 + \gamma_\theta^2)^{\frac{5}{2}}}, \\
b'(0) &= \frac{\mathbf{h}_\theta\theta(1+\gamma) + \gamma_\theta\theta\mathbf{h}}{((1+\gamma)^2 + \gamma_\theta^2)^{\frac{3}{2}}} - 3\gamma_\theta\theta(1+\gamma) \frac{(\mathbf{h}(1+\gamma) + \mathbf{h}_\theta\gamma_\theta)}{((1+\gamma)^2 + \gamma_\theta^2)^{\frac{5}{2}}}, \\
c'(0) &= \cot\theta \frac{\mathbf{h}_\theta}{(1+\gamma) \left((1+\gamma)^2 + \gamma_\theta^2 \right)^{\frac{1}{2}}} - \cot\theta \frac{\mathbf{h}\gamma_\theta}{(1+\gamma)^2 \left((1+\gamma)^2 + \gamma_\theta^2 \right)^{\frac{1}{2}}} \\
&\quad - \cot\theta \frac{2\gamma_\theta(\mathbf{h}(1+\gamma) + \mathbf{h}_\theta\gamma_\theta)}{(1+\gamma) \left((1+\gamma)^2 + \gamma_\theta^2 \right)^{\frac{3}{2}}}
\end{aligned}$$

Finally,

$$D_{\mathbf{h}}\mathbb{F}(\gamma, \wp_\gamma)(\mathbf{h}) = \tau_1\mathbf{h}_{\theta\theta} + \tau_2\mathbf{h}_\theta + \tau_3\mathbf{h}, \quad (23)$$

where

$$\begin{aligned}
\tau_1 &= \frac{\sigma}{RP_{atm}} \frac{1+\gamma}{((1+\gamma)^2 + \gamma_\theta^2)^{\frac{3}{2}}}, \\
\tau_2 &= \frac{\sigma}{RP_{atm}} \frac{-3\gamma_\theta^3 + 3\gamma_\theta\gamma_{\theta\theta}(1+\gamma)}{((1+\gamma)^2 + \gamma_\theta^2)^{\frac{3}{2}}} + \frac{\sigma}{RP_{atm}} \frac{\cot\theta \left((1+\gamma)^2 - \gamma_\theta^2 \right)}{(1+\gamma) \left((1+\gamma)^2 + \gamma_\theta^2 \right)^{\frac{3}{2}}}, \\
\tau_3 &= \frac{\sigma}{RP_{atm}} \frac{-2(1+\gamma)^3 - 8\gamma_\theta^2(1+\gamma)}{((1+\gamma)^2 + \gamma_\theta^2)^{\frac{3}{2}}} - \frac{\sigma}{RP_{atm}} \frac{\gamma_{\theta\theta}\gamma_\theta^2 - 2\gamma_\theta\theta(1+\gamma)^2}{((1+\gamma)^2 + \gamma_\theta^2)^{\frac{5}{2}}}, \\
&\quad + \frac{\sigma}{RP_{atm}} \cot\theta \frac{-\gamma_\theta^3 + \gamma_\theta(1+\gamma)^2}{(1+\gamma)^2 \left((1+\gamma)^2 + \gamma_\theta^2 \right)^{\frac{3}{2}}} + \psi.
\end{aligned}$$

Theorem 3. *The Fréchet derivative of \mathbb{F} with respect to \mathbf{h} , given in (23), is a linear homeomorphism.*

Proof. Clearly $D_{\mathbf{h}}\mathbb{F}(\gamma, \wp_\gamma)$ is linear and continuous. Thus, if one can prove that it is a bijection, the Inverse Theorem guarantees that its inverse is also continuous (see, e.g., [16], [17]). The bijectivity of $D_{\mathbf{h}}\mathbb{F}(\gamma, \wp_\gamma)$ is equivalent with proving that for any $g \in C\left[\frac{\pi}{2} - \varepsilon, \frac{\pi}{2} + \varepsilon\right]$, the initial value problem

$$\begin{cases} \tau_1 u_{\theta\theta} + \tau_2 u_\theta + \tau_3 u = 0, \\ u\left(\frac{\pi}{2}\right) = \gamma\left(\frac{\pi}{2}\right), \\ u'\left(\frac{\pi}{2}\right) = \gamma'\left(\frac{\pi}{2}\right) \end{cases} \quad (24)$$

has a unique solution. Since $\tau_1 \neq 0$ and τ_2, τ_3 are continuous functions on $\left[\frac{\pi}{2} - \varepsilon, \frac{\pi}{2} + \varepsilon\right]$, we employ Theorem 2 to obtain the conclusion. \square

Our main result of this paper follows from Theorem 1 and Theorem 3, that is, **Theorem 4.** *For any sufficiently small perturbation of \wp from \wp_γ , there exists a unique $\mathbf{h} \in C_\gamma$ such that (21) holds.*

Acknowledgements

The author acknowledges the support provided by the project "Nonlinear Studies of Stratified Oceanic and Atmospheric Flows", funded by the European Union through the Next Generation EU initiative and the Romanian Government under the National Recovery and Resilience Plan for Romania. The project is contracted under number 760040/23.05.2023, cod PNRR-C9-I8-CF 185/22.11.2022, through the Romanian Ministry of Research, Innovation, and Digitalization, within Component 9, Investment I8. Also, the author would like to thank Professor C.I. Martin for valuable discussions and suggestions.

References

- [1] Constantin, A., Johnson, R.S.: An exact, steady, purely azimuthal equatorial flow with a free surface. *J. Phys. Oceanogr.* **46**(6), 1935–1945 (2016). <https://doi.org/10.1175/JPO-D-15-0205.1>
- [2] Cushman-Roisin, B., Beckers, J.-M.: *Introduction to Geophysical Fluid Dynamics: Physical and Numerical Aspects.* Academic Press, Waltham (2011)
- [3] Vallis, G.K.: *Atmospheric and Oceanic Fluid Dynamics.* Cambridge University Press (2006)
- [4] Constantin, A., Johnson, R.S.: An exact, steady, purely azimuthal flow as a model for the Antarctic Circumpolar Current. *J. Phys. Oceanogr.* **46**, 3585–3594 (2016). <https://doi.org/10.1175/JPO-D-16-0121.1>
- [5] Constantin, A., R.S. Johnson, An exact, steady, purely azimuthal equatorial flow with a free surface, *J. Phys. Oceanogr.* **46** (6) (2016) 1935–1945
- [6] Gheorghe, C., Stan, A. Stratified equatorial flows in cylindrical coordinates with surface tension. *Monatsh Math* **205**(2024), 497–509
- [7] Henry, D., Martin, C.I.: Exact, purely azimuthal stratified equatorial flows in cylindrical coordinates. *Dyn. Partial Differ. Equ.* **15**, 337–349 (2018). <http://dx.doi.org/10.4310/DPDE.2018.v15.n4.a4>
- [8] Henry, D., Martin, C.-I.: Free-surface, purely azimuthal equatorial flows in spherical coordinates with stratification. *J. Differential Equations* **266**, 6788–6808 (2019). <https://doi.org/10.1016/j.jde.2018.11.017>
- [9] Henry, D., Martin, C.I.: Exact, Free-Surface Equatorial Flows with General Stratification in Spherical Coordinates. *Arch Rational Mech Anal* **233**, 497–512 (2019).

<https://doi.org/10.1007/s00205-019-01362-z>

- [10] Martin, C., Petruşel, A.: Free surface equatorial flows in spherical coordinates with discontinuous stratification depending on depth and latitude. *Annali di Matematica* 201, 2677–2690 (2022). <https://doi.org/10.1007/s10231-022-01214-w>
- [11] Berger, M.S. *Nonlinearity and Functional Analysis*. Academic Press, New York (1977)
- [12] Logan, J. D.: *A First Course in Differential Equations*. Springer Cham (2015)
- [13] Coddington, E.A.: *An Introduction to Ordinary Differential Equations*. Dover, New York (1961)
- [14] Hsu, H.-C., Martin, C.I.: Free-surface capillary-gravity azimuthal equatorial flows. *Nonlinear Anal.* **144**(10), 1–9 (2016). <https://doi.org/10.1016/j.jde.2018.11.017>
- [15] Griffiths, D.J.: *Introduction to Electrodynamics*, 5th ed. Cambridge University Press (2023).
- [16] Royden, H.L., Fitzpatrick, P.: *Real Analysis*. Prentice Hall, Boston (2010)
- [17] Wilansky, A.: *Modern Methods in Topological Vector Spaces*. Mineola, New York: Dover Publications (2013)