

Constructing optimal treatment length strategies to maximize quality-adjusted lifetimes

Hao Sun

Didi Chuxing Technology Company

Ashkan Ertefaie*

Department of Biostatistics and Computational Biology, University of Rochester

Luke Duttweiler

Department of Biostatistics, Harvard T.H. Chan School of Public Health
and

Brent A. Johnson

Department of Biostatistics and Computational Biology, University of Rochester

December 9, 2024

Abstract

Real-world clinical decision making is a complex process that involves balancing the risks and benefits of treatments. Quality-adjusted lifetime is a composite outcome that combines patient quantity and quality of life, making it an attractive outcome in clinical research. We propose methods for constructing optimal treatment length strategies to maximize this outcome. Existing methods for estimating optimal treatment strategies for survival outcomes cannot be applied to a quality-adjusted lifetime due to induced informative censoring. We propose a weighted estimating equation that adjusts for both confounding and informative censoring. We also propose a non-parametric estimator of the mean counterfactual quality-adjusted lifetime survival curve under a given treatment length strategy, where the weights are estimated using an undersmoothed sieve-based estimator. We show that the estimator is asymptotically linear and provide a data-dependent undersmoothing criterion. We apply our method to obtain the optimal time for percutaneous endoscopic gastrostomy insertion in patients with amyotrophic lateral sclerosis.

Keywords: Amyotrophic lateral sclerosis, highly adaptive lasso, informative censoring, non-parametric IPW estimator, sectional variation norm, undersmoothing.

*The authors gratefully acknowledge *R01DA058996*, *R01DA048764*, *R61NS120240*, and *R33NS120240* from the National Institutes of Health

1 Introduction

1.1 Motivation

In patient care, clinicians often navigate the delicate balance between risks and benefits, particularly in critical care, life-threatening situations, or managing incurable neurodegenerative disorders like Parkinson’s Disease, Alzheimer’s, and ALS. For example, in ALS, as functional decline progresses, patients and providers face the complex decision of whether to pursue a gastric feeding tube (PEG), which can extend survival but may also introduce side effects and reduce quality of life.

ALS is a rare but fatal neurodegenerative disease with no cure ([Zarei et al., 2015](#)). PEG tube insertion provides nutrition when oral intake becomes unsafe, potentially prolonging life. However, it can lead to complications such as infections, reduced mobility, respiratory issues, and discomfort at the insertion site. These risks increase significantly for patients in poor health, underscoring the importance of early intervention ([Mathus-Vliegen et al., 1994](#); [Leigh et al., 2003](#)). Determining the optimal timing and target population for PEG remains an open question, which our article seeks to address using innovative methods.

1.2 Optimal Treatment Strategies with Survival Outcomes

Quality-adjusted lifetime is a composite outcome that integrates patients’ quantity and quality of life ([Goldhirsch et al., 1989](#); [Glasziou et al., 1990](#)). It represents discounted survival time, where the discount factor—a function of time and disease status—ranges from 0 to 1. Consequently, quality-adjusted lifetime is always less than or equal to survival time. Without censoring, patients with poor quality of life contribute to failure sets earlier on the adjusted time scale. However, in the presence of censoring, these patients

may be under-represented in early failure sets due to induced informative censoring, which biases standard survival analysis approaches when quality-adjusted lifetime is the endpoint of interest (Gelber et al., 1989). Zhao and Tsiatis (1997) addressed this by proposing a consistent estimator for the distribution of quality-adjusted lifetime using a weighted estimating equation. The weights are reciprocal to the probability of observing an uncensored quality-adjusted lifetime and are estimated via the Kaplan-Meier method. Alternative approaches include regression-based methods (Fine and Gelber, 2001; Bang and Tsiatis, 2002; Wang and Zhao, 2006), augmented inverse probability-weighted estimators (van der Laan and Hubbard, 1999), and maximum likelihood estimation of the marginal mark-and-time joint distribution (Huang and Louis, 1998).

Precision medicine further emphasizes individualized treatment strategies designed to optimize health outcomes (Robins, 2004; Chakraborty and Moodie, 2013). For survival outcomes, methods include regression-based approaches incorporating Cox’s model (Chen and Tsiatis, 2001; Tian et al., 2014), doubly robust estimators (Hubbard et al., 2000; Robins et al., 2008b; Bai et al., 2013; Zhao et al., 2014; Díaz et al., 2018; Cai and van der Laan, 2020b; Rytgaard et al., 2022), Q-learning extensions for censored outcomes (Goldberg and Kosorok, 2012), and Kaplan-Meier-based survival estimators for treatment strategies (Jiang et al., 2017). These methods, however, are not applicable to settings where the quality-adjusted lifetime is the endpoint of interest.

1.3 Our contribution

We propose an estimating equation approach to estimate an optimal treatment length strategy concerning quality-adjusted survival lifetime. Our method is applicable, for example, in settings where the interest is to construct individualized decision rules for when to stop or initiate therapy. One of the challenges in such cases is the presence of many decision points. While in theory efficient estimators can be constructed, these estimators

will rely on many nuisance parameters and the derivation of the efficient influence function will be cumbersome. Importantly, the resulting estimator can be irregular with large biases when a subset of the nonparametrically estimated nuisance functions fail to be consistent (Van der Laan, 2014; Benkeser et al., 2017). To address this, we propose two inverse probability weighting estimators: one using parametric models for weights and another allowing data-adaptive estimation of weights. The second approach achieves nonparametric efficiency under mild conditions without deriving the efficient influence function (Ertefaie et al., 2023). Our contributions are threefold. First, the proposed method addresses the induced censoring problem after transforming survival time into quality-adjusted lifetime and enables estimating an optimal treatment length strategy. Second, it allows to estimate the nuisance parameters (i.e., weight functions) using an undersmoothed sieve-based (i.e., data adaptive) technique while providing root- n consistency, thereby achieving efficiency in a nonparametric model space. Third, we provide both theoretical and finite sample undersmoothing criteria to attain the best estimator for any given data, which can be of interest in its own right.

Inverse probability weighting estimators with data-adaptive weights often face challenges in achieving asymptotic linearity and root- n convergence. We overcome these issues by using an undersmoothed highly adaptive lasso to estimate the weight functions, yielding an asymptotically linear estimator for the survival curve of quality-adjusted lifetime under a treatment regime. Our method is general and applies to various survival curve functions, including t -year survival probabilities and restricted quality-adjusted lifetime (RQAL). We focus on RQAL in simulation studies due to its relevance to the real data example.

2 Preliminaries

2.1 Notation

Assume that subjects are monitored at K (finite) pre-specified landmarks indexed by $0 = l_0 < l_1 < l_2 < \dots < l_K \leq \tau$, where τ is the end of follow-up or the upper limit of the support in the time-scale. We define the ALS quality-of-life score at time t as $V(t)$ and assume $V(t)$ does not change between two adjacent landmarks. We define $V(s) = 0$ as the absorbing state, which implies that $V(t) = 0$ for $t > s$; in our application, the absorbing state is death and the quality-of-life is assumed to be zero after death. The lifetime T is defined: $T = \inf\{t : V(t) = 0\}$; it is recorded continuously and represents the time it takes for a subject to move from the initial state to the absorbing state. Define \mathcal{Q} as a quality-of-life function mapping the state space $V(t)$ to the interval $[0, 1]$. We further assume the functional form of \mathcal{Q} is known and set to zero at the absorbing state, i.e. $\mathcal{Q}(0) = 0$. With this notation, a subject's quality-adjusted lifetime is defined as

$$U = \int_0^T \mathcal{Q}\{V(t)\} dt. \quad (1)$$

Let $s^*(x) = \inf [s : \int_0^s \mathcal{Q}\{V(t)\} dt \geq x]$ denote the minimum time (on the original scale) required to cross the target quality-adjusted lifetime x . Let Z_0 denote a vector of baseline variables and Z_j denote the updated time-dependent information at the j th landmark. Also, let $A_j \in \{0, 1\}, j = 1, \dots, K$ denote treatment assignment indicator, and we assume the treatment assignment is monotone that is once it is initialized or stopped, the individual will stay at the current status and no longer switch until the end of the study. We assume that the treatment assignment does not occur at baseline, i.e. $A_0 = 0$. Let C denote a continuously recorded censoring time and $\tilde{T} = \min(T, C)$. Let $Y_j = I(\tilde{T} \geq l_j)$ be the at-risk indicator and realizes the value 1 if the subject is still at risk at landmark l_j and 0 otherwise; $Y_0 = 1$ by definition. Observations after censoring or absorbing state are

filled as \emptyset values for simplicity. Our data consists of n independent, identically distributed trajectories $\mathcal{O} = [\{Y_j, Z_j, A_j\}_{j=0}^K, I(T < C), \tilde{T}]$.

Let $T(x) = \min\{T, s^*(x)\}$, then define the censoring indicator at target time x as $\Delta_c(x) = I\{C > T(x)\}$ and $\tilde{T}(x) = \min\{T(x), C\}$. We also define $l(x) = \operatorname{argmax}\{j : \tilde{T}(x) \geq l_j\}$ to denote the last landmark point that an individual was still at risk for a target quality-adjusted lifetime x . We also define landmark based censoring indicators as $C_j(x) \in \{0, 1\}, j = 0, \dots, K$, where $C_j(x) = I\{\Delta_c(x) = 0, \tilde{T}(x) < l_{j+1}\}$. Overbars with a subscript is used for cumulative information, e.g., the covariate history up to the j th landmark is denoted as $\bar{Z}_j = (Z_0, \dots, Z_j)$.

Define the propensity score for treatment assignment $A_j, j = 1, \dots, K$ as

$$h_{a,j}(1 | \bar{a}_{j-1}, \bar{z}_j) = (1 - A_{j-1})P(A_j = 1 | A_{j-1} = 0, Y_j = 1, \bar{Z}_j = \bar{z}_j) + A_{j-1}. \quad (2)$$

Note, $h_{a,j}(1 | \bar{a}_{j-1}, \bar{z}_j) = 1$ if a subject has inserted PEG prior to l_j . We define the censoring mechanism for $C_j(x), j = 1, \dots, K$ as

$$h_{c,j,x}(1 | \bar{a}_j, \bar{z}_j) = \{1 - C_{j-1}(x)\}P\{C_j(x) = 1 | \bar{A}_j = \bar{a}_j, C_{j-1}(x) = 0, Y_j = 1, \bar{Z}_j = \bar{z}_j\} + C_{j-1}(x). \quad (3)$$

Similar to $h_{a,j}(1 | \bar{a}_{j-1}, \bar{z}_j)$ above, we set $h_{c,j,x}(1 | \bar{a}_j, \bar{z}_j) = 1$ if a subject is censored prior to l_j .

2.2 Causal Estimands and Estimating Equations

The goal of our study to construct a dynamic treatment regime for PEG tube insertion that optimizes quality-adjusted lifetime. A treatment length strategy for PEG tube insertion is a sequence of decision rules of whether to undergo surgery for PEG tube insertion given all available information up to that time point. We consider a class of decision rules

$$\mathcal{G} = \{g : g^\eta(A_{j-1}, Z_j) = I[(\eta^\top Z_j)^{1-A_{j-1}} \geq 0], j = 1, \dots, K, \|\eta\| = 1\},$$

where $\|\cdot\|$ is the L_2 norm. Let $\bar{g}_K^\eta = \{g_1^\eta(A_0, Z_1), \dots, g_K^\eta(A_{K-1}, Z_K)\}$. The decision rule $g_j^\eta \equiv g^\eta(A_{j-1}, Z_j)$ triggers PEG tube insertion as soon as the linear score $\eta^\top Z_j$ crosses zero. The power $(1 - A_{j-1})$ in the exponent of the linear score guarantees that the regimes of interest belong to the class of monotone treatment regimes so that patients who have PEG tube inserted do not later have it removed and possibly wait to have it reinserted again at a later date (Strijbos et al., 2017).

Define T^η and U^η as the potential survival time and the corresponding potential quality-adjusted lifetime, respectively, defined by (1) that would have been observed if a patient's lifetime was uncensored during follow-up and had followed the strategy \bar{g}_K^η for PEG tube insertion, i.e., $\{\bar{A}_K = \bar{g}_K^\eta\}$. We define the survivor distribution of quality-adjusted lifetime under the treatment length strategy \bar{g}_K^η , $P(U^\eta > x) = S_U(x; \eta)$, and our target estimand as $f\{S_U(x; \eta)\}$, for some user-defined known functional of $S_U(x; \eta)$. When the goal is to maximize the t -year survival probability (Jiang et al., 2017), $f(x) = x$, and when the target parameter is the restricted mean quality adjusted lifetime, then $f\{S_U(x; \eta)\} = \int_0^{L_U} S_U(x; \eta) dt$ with a user defined upper limit $L_U \leq \tau$. Naturally, estimating $S_U(x; \eta)$ consistently is a principal concern in the proposed method.

Our procedure handles confounding and dependent censoring as two sources of estimation bias. We leverage the inverse probability weighting strategy to adjust for these biases. Let \bar{g}_K^η as $\Delta_a^\eta(x) = \prod_{j \leq l(x)} I\{A_j = g_j^\eta\}$ denote the indicator function for treatment assignment at the target time x under the treatment length strategy \bar{g}_K^η . This indicates whether the observed cumulative treatment assignment $\bar{A}_{l(x)}$ up to landmark $l(x)$ follows the strategy \bar{g}_K^η . Define the probability of following the regime \bar{g}_K^η through landmark $l(x)$ as $H_a(x, \bar{z}_{l(x)}) = \prod_{j \leq l(x)} h_{a,j}(a | \bar{a}_{j-1}, \bar{z}_j)$. We propose the following estimating function for

$S_U(x; \eta)$,

$$\begin{aligned} \mathbb{P}_n \left[\frac{\Delta_a^\eta(x) \Delta_c(x)}{H_a(x) H_c(x)} \{I(U > x) - S_U(x; \eta)\} \right] = \\ \mathbb{P}_n \left[\prod_{j \leq l(x)} \frac{I\{A_j = g_j^\eta\}}{h_{a,j}(A_j | \bar{A}_{j-1}, \bar{Z}_j)} \frac{I(C_j(x) = 0)}{1 - h_{c,j,x}(1 | \bar{Z}_j, \bar{A}_j)} \{I(U > x) - S_U(x; \eta)\} \right], \quad (4) \end{aligned}$$

where $H_a(x) = H_a(x, \bar{z}_{l(x)})$ and $H_c(x) = H_c(x, \bar{a}_{l(x)}, \bar{z}_{l(x)}) := \prod_{j \leq l(x)} \{1 - h_{c,j,x}(1 | \bar{z}_j, \bar{a}_j)\}$ denote the propensity score and the uncensored probability up to landmark $l(x)$. We will discuss the methods to estimate H_c and H_a in Section 3.

Remark 1 (Extension to Quality-adjusted Lifetime). *Mean restricted quality-adjusted lifetime is defined as the integrated quality-adjusted lifetime from zero to L_U , where L_U is a user-defined upper-limit on the quality-adjusted lifetime scale. Hence,*

$$R(\eta) = \int_0^{L_U} S_U(x; \eta) dx. \quad (5)$$

In our application, we set $L_U = 91.12$ which is the 95% quantile of the observed quality-adjusted lifetime.

3 Estimation

3.1 Semiparametric and Nonparametric Estimators

The estimating equation (4) is infeasible due to unknown nuisance functions. We propose two estimators for the quality adjusted lifetime based on (4). The first estimator relies on parametric modeling assumptions for nuisance parameters. Specifically, we define $\hat{S}_U^{par}(x; \eta)$ as a solution of (4) where $h_{a,j}$ and $h_{c,j,x}$ are replaced with their corresponding parametric estimators $\hat{h}_{a,j}^{par}$ and $\hat{h}_{c,j,x}^{par}$. The statistical properties of $\hat{S}_U^{par}(x; \eta)$ depend on the consistency of both of these estimators.

The second estimator relaxes these modeling assumptions and allows one to model the nuisance functions using a nonparametric regression approach (i.e., highly adaptive lasso).

We refer to this estimator as $\hat{S}_U^{npar}(x; \eta)$ with the estimated nuisance functions as $\hat{h}_{a,j}^{npar}$ and $\hat{h}_{c,j,x}^{npar}$. Accordingly, an estimator of mean restricted quality-adjusted lifetime under regime \bar{g}_K^η is

$$\hat{R}^\diamond(\eta) = \int_0^{L_U} \hat{S}_U^\diamond(x; \eta) dx, \quad (6)$$

where $\diamond \in \{par, npar\}$. Accordingly, the optimal dynamic regime to maximize mean restricted quality-adjusted lifetime is therefore $\bar{g}_K^{\hat{\eta}_{opt}^\diamond}$ where $\hat{\eta}_{opt}^\diamond$ is the maximizer of $\hat{R}^\diamond(\eta)$.

3.2 Finite Sample Bias Correction

In Section 4, we show that, under certain conditions, $\hat{R}^\diamond(\eta)$ is consistent for $R(\eta)$ for any given η . However, when the optimal treatment length strategy is estimated using data, $\hat{R}^\diamond(\hat{\eta}_{opt}^\diamond) \geq \hat{R}^\diamond(\eta_{opt}) \rightarrow R(\eta_{opt})$, which means that $\hat{R}^\diamond(\hat{\eta}_{opt}^\diamond) \rightarrow R(\eta_{opt})$ from above as $n \rightarrow \infty$, but $\hat{R}^\diamond(\hat{\eta}_{opt}^\diamond) \geq \hat{R}^\diamond(\eta_{opt})$ with finite sample size and $\hat{R}^\diamond(\hat{\eta}_{opt}^\diamond)$ would slightly overestimate $R(\eta_{opt})$. Such bias has no impact for the estimation of an optimal strategy, but may lead to insufficient coverage rate when the statistical inference of $R(\eta_{opt})$ is of interest. To mitigate this issue, we construct a smooth estimator to approximate the inverse probability weighting estimator (Jiang et al., 2017). Asymptotically, we can replace $g_j^\eta = I\{(\eta^\top Z_j)^{1-A_{j-1}} > 0\}$ with $\Phi\{(\eta^\top Z_j)^{1-A_{j-1}}/\nu\}$, where $\Phi(\cdot)$ is the standard normal distribution function and ν is a bandwidth tending to 0 as $n \rightarrow \infty$. When $\nu \rightarrow 0$, the normal distribution function obviously converges to the indicator function. The indicator function of treatment following $I(A_j = g_j^\eta)$ can thus be replaced by $W_{A_j}^\eta = A_j \times \Phi\{(\eta^\top Z_j)^{1-A_{j-1}}/\nu\} + (1 - A_j) \times [1 - \Phi\{(\eta^\top Z_j)^{1-A_{j-1}}/\nu\}]$, and thus, $\Delta_a^\eta(x) = \Pi_{j \leq l(x)} I(A_j = g_j^\eta)$, a product of indicator function, can be replaced with $\tilde{\Delta}_a^\eta(x) = \Pi_{j \leq l(x)} W_{A_j}^\eta$. Finally, with this modification, we redefine the estimating equation (4) at time x as

$$\mathbb{P}_n \left[\frac{\tilde{\Delta}_a^\eta(x) \Delta_c(x)}{H_a^\diamond(x) H_c^\diamond(x)} \{I(U > x) - S_U(x; \eta)\} \right] \quad (7)$$

and denote the solution of (7) as $\tilde{S}_U^\diamond(x; \eta)$. This is a normal approximation of indicator function which is used to correct the finite sample bias (Yang and Ding, 2017). We recommend and use $\nu = n^{-1/3}sd(\mathcal{Z})/K$, where $\mathcal{Z} = \{\eta^\top Z_j, j = 1, \dots, K\}$, based on our empirical studies. Here, $sd(\mathcal{Z})$ is the sample standard deviation of the linear predictor and K is the number of stages in the study. The extension to restricted quality-adjusted lifetime is straightforward by combining (6) and (7). Accordingly, define $\tilde{\eta}_{opt}^\diamond = \arg \max_\eta \tilde{R}^\diamond(\eta)$ where

$$\tilde{R}^\diamond(\eta) = \int_0^{L_U} \tilde{S}_U^\diamond(x; \eta) dx. \quad (8)$$

3.3 Highly Adaptive Lasso Estimator

The highly adaptive lasso estimator is a data-adaptive modeling technique which is shown to be consistent for the true regression function with a convergence rate faster than $n^{-1/3}$, up to a log factor (Benkeser and Van Der Laan, 2016; Bibaut and van der Laan, 2019). Highly adaptive lasso offers a flexible tool for estimation of nuisance parameters.

Let d_j denote the covariate dimension at time j and $s \subset (1, \dots, d_j)$. The first step to map any subset s of the covariate vector to indicator basis functions. For example, when the covariate is two dimensional $X = (X_1, X_2)$, $\phi_i(x) = \{\phi_{s=X_1,i}(x), \phi_{s=X_2,i}(x), \phi_{s=(X_1, X_2),i}(x)\} = \{I(x_1 \geq X_{i1}), I(x_2 \geq X_{i2}), I(x_1 \geq X_{i1}, x_2 \geq X_{i2})\}, i = 1, \dots, n$, is the collection of the first- and the second-order indicator basis functions. Then, for each j , a regression model is fit (e.g., with logit link) given these set of basis functions denoted as $\phi_{j,s,i}$. Denote the corresponding parameters as $\beta_{j,s,i}$, then the propensity score can be represented as logit $h_{a,j,\beta} = \beta_{j,0}^a + \sum_{s \subset \{1, \dots, d_j\}} \sum_{i=1}^n \beta_{j,s,i}^a \phi_{s,i}$, where $|\beta_{j,0}^a| + \sum_{s \subset \{1, \dots, d_j\}} \sum_{i=1}^n |\beta_{j,s,i}^a|$ is an approximation of the sectional variation norm of logit $h_{a,j}$. The highly adaptive lasso defines a minimum loss based estimator $\hat{\beta}_j^a$ as

$$\hat{\beta}_j^a = \arg \min_{\beta: |\beta_{j,0}^a| + \sum_{s \in \mathcal{T}} \sum_{i=1}^n |\beta_{j,s,i}^a| < \lambda} \mathbb{P}_n L(\text{logit } h_{a,j,\beta}),$$

where $L(\cdot)$ is the negative log-likelihood loss function, \mathbb{P}_n is the empirical average and λ

is the sectional variation norm of logit $h_{a,j}$ which can be obtained using cross-validation (Benkeser and Van Der Laan, 2016). We denote the estimated propensity score at time j as $\hat{h}_{a,j}^{npar} \equiv \hat{h}_{a,j}^{npar}$. Similarly, the parameters associated with the censoring mechanism can be estimated, yielding $h_{c,j,x}^{npar} \equiv \hat{h}_{c,j,x}^{npar}$. Additional details on the highly adaptive lasso are provided in Section S1.4 of the Supplementary Material.

3.4 Undersmoothing in practice

We generalize a data adaptive undersmoothing criterion of Ertefaie et al. (2023) to our longitudinal settings. Define, for $j = 1, 2, \dots, l(x)$,

$$\tilde{\lambda}_{n,j}^a = \arg \min_{\lambda} B^{-1} \sum_{b=1}^B \left[\sum_{(s,i) \in \mathcal{J}_n} \frac{1}{\|\hat{\beta}_{j,\lambda,b}^a\|_{L_1}} \left| \mathbb{P}_{n,b}^1 \tilde{\Omega}_{s,i}(\phi, \hat{h}_{a,j,\lambda,b}) \right| \right], \quad (9)$$

in which $\|\hat{\beta}_{j,\lambda,b}^a\|_{L_1} = |\hat{\beta}_{j,\lambda,0}^a| + \sum_{s \in \{1, \dots, d_j\}} \sum_{i=1}^n |\hat{\beta}_{j,\lambda,s,i}^a|$ is the L_1 -norm of the coefficients $\hat{\beta}_{j,\lambda,s,i}^a$ in the highly adaptive estimator $\hat{h}_{a,j,\lambda}$ for a given λ in fold b , and $\tilde{\Omega}_{s,i}(\phi, h_{a,j,\lambda,b}^{npar}) = \phi_{r,i}(\bar{A}_{j-1}, \bar{Z}_j) \{A_j - \hat{h}_{a,j,\lambda,b}^{npar}(1 | \bar{A}_{j-1}, \bar{Z}_j)\} \{\hat{h}_{a,j,\lambda,b}^{npar}(1 | \bar{A}_{j-1}, \bar{Z}_j)\}^{-1}$.

Undersmoothing can result in a reduced convergence rate for the highly adaptive lasso fit. Our asymptotic linearity proof of $\hat{R}^{npar}(\hat{\eta}^{opt})$ in Theorem 3 requires $\|\hat{h}_{a,j}^{npar} - h_{a,j}\|_2 = o_p(n^{-1/4})$ for all $j = 1, 2, \dots, l(x)$, which is slower than the convergence rate of the cross-validated highly adaptive lasso fit, given by $O_p(n^{-1/3})$. This ensures that a certain degree of undersmoothing can be tolerated without compromising our desired asymptotic properties. Let J represent the number of features included in the fit (i.e., $J = |\mathcal{J}_n|$). As shown by van der Laan (2023), the convergence rate of a highly adaptive lasso is $(J/n)^{1/2}$. To ensure the required $o_p(n^{-1/4})$, we must choose J such that $J < n^{1/2}$, and let $\lambda_{n,J}^a$ be the corresponding L_1 -norm. We propose our criterion as:

$$\check{\lambda}_{n,j}^a = \max(\lambda_{n,J}^a, \tilde{\lambda}_{n,j}^a). \quad (10)$$

In certain instances, the criterion may lead to excessive undersmoothing (i.e., selecting a very small $\tilde{\lambda}_{n,j}^a$), and the max operator aids in mitigating this issue. The same procedure

can be implemented to obtain the undersmoothing parameter $\check{\lambda}_{n,j}^c = \max(\lambda_{n,J}^c, \tilde{\lambda}_{n,j}^c)$ to fit the function $\hat{h}_{c,j,\lambda^c}^{npar}$.

4 Theoretical properties

4.1 Coarsening assumptions

The following assumptions are required for identification of the target parameters using the observed data.

Assumption 1. (*causal inference and censoring assumptions*)

- a. *Consistency:* $I(U > x) = I(U^\eta > x)$ if $\Delta_a^\eta(x)\Delta_c(x) = 1$;
- b. *Sequential randomization assumption (SRA):* given that a patient is still under observation at landmark l_j and given the patient's covariate history \bar{Z}_j, \bar{A}_{j-1} and \bar{C}_{j-1} , A_j doesn't depend on future prognosis; also, given that a patient is still under observation at landmark l_j and given the patient's covariate history \bar{Z}_j, \bar{A}_j and \bar{C}_{j-1} , C_j doesn't depend on future prognosis;
- c. *Strong positivity:* $H_a(x, \bar{z}) > \alpha_1$ and $H_c(x, \bar{a}, \bar{z}) > \alpha_2$ for each a, z and $\alpha_1, \alpha_2 > \epsilon$ with $\epsilon > 0$ for $j = 1, \dots, K$;

Assumption 1a connects the observed outcome with the potential outcomes and validates the use of observed data to estimate the causal estimand. It also states that there is no interference between the observations. Assumption 1b assumes that there is no unmeasured confounding variable over time for treatment assignment and censoring process and is crucial to identify the mean outcome. Assumption 1c ensures that statistical inference of the treatment effect is possible. These assumptions are commonly used in causal inference literature ([Rosenbaum and Rubin, 1983](#); [Robins, 1986](#)).

4.2 Parametric Estimation of the Nuisance Functions

The first two theorems establish the results when the discrete hazard function for treatment assignment (2) and censoring (3) are correctly specified by finite dimensional parametric models, e.g., logistic models.

Theorem 1. *Under conditions (C.1)-(C.5) in Appendix S1, if the functions $h_{a,j}$ and $h_{c,j,x}$, $j = 1, \dots, K$, are correctly specified, for any regime \bar{g}_K^η , we have, as $n \rightarrow \infty$, (1) $\hat{S}_U^{par}(x; \eta) \rightarrow_p S_U(x; \eta)$ for $x \in (0, L_U]$, (2) $n^{1/2}\{\hat{S}_U^{par}(x; \eta) - S_U(x; \eta)\}$ converges to a mean-zero Gaussian process, (3) $n^{1/2}\{\hat{S}_U^{par}(x; \hat{\eta}_{opt}^{par}) - S_U(x; \eta_{opt})\} \rightarrow_d N\{0, \sigma_{\eta_{opt},1}^2(x)\}$, where the expression of $\sigma_{\eta_{opt},1}^2(x)$ can be found in Appendix S1, (4) $n^{1/2}\{\hat{S}_U^{par}(x; \hat{\eta}_{opt}^{par}) - \tilde{S}_U^{par}(x; \tilde{\eta}_{opt}^{par})\} = o_p(1)$.*

Theorem 1 serves as the building block of our main result in Theorem 2.

Theorem 2. *Under conditions (C.1)-(C.5) in Appendix S1, if $h_{a,j}$ and $h_{c,j,x}$, $j = 1, \dots, K$, are correctly specified, for any regime \bar{g}_K^η , we have, as $n \rightarrow \infty$, (1) $\hat{R}^{par}(\eta) \rightarrow_p R(\eta)$, (2) $n^{1/2}\{\hat{R}^{par}(\eta) - R(\eta)\}$ converges to a mean-zero Gaussian process, (3) $n^{1/2}\{\hat{R}^{par}(\hat{\eta}_{opt}^{par}) - R(\eta_{opt})\} \rightarrow_d N(0, \sigma_{\eta_{opt},2}^2)$, where the expression of $\sigma_{\eta_{opt},2}^2$ can be found in Appendix S1, (4) $n^{1/2}\{\hat{R}^{par}(\hat{\eta}_{opt}^{par}) - \tilde{R}^{par}(\tilde{\eta}_{opt}^{par})\} = o_p(1)$.*

4.3 Nonparametric estimator of $S_U(x; \eta)$

To improve the robustness of our approach, we propose a nonparametric estimator where the weight functions are estimated nonparametrically. In general, obtaining an inverse probability weighted estimator with a desired root- n is challenging due to slower rate of convergence of estimated weight functions. We overcome this issue by estimating the nuisance parameters using undersmoothed highly adaptive lasso denoted as \hat{H}_a^{npar} and \hat{H}_c^{npar} . Specifically, we show that (i) under certain assumption listed below $\Psi_{\hat{H}_a^{npar}, \hat{H}_c^{npar}}(x; \eta)$ is an

unbiased estimating equation; and, (ii) the resulting estimator $\hat{S}_U^{npar}(x; \eta)$ is asymptotically linear. The estimated optimal regime at x is defined as $\hat{\eta}_{opt}^{npar} = \operatorname{argmax}_{\eta} \hat{S}_U^{npar}(x; \eta)$.

To improve the performance of our estimator in finite samples, we propose employing cross-fitting for estimating nuisance parameters (Klaassen, 1987; Zheng and van der Laan, 2011; Chernozhukov et al., 2017). The dataset is randomly divided into B mutually exclusive and exhaustive sets, each approximately of size nB^{-1} . The empirical distribution of a training and validation sample is denoted as $\mathbb{P}_{n,b}^0$ and $\mathbb{P}_{n,b}^1$, respectively, for $b = 1, 2, \dots, B$. The resulting estimating equation is

$$\Psi_{H_a^{npar}, H_c^{npar}}(x; \eta) = \frac{1}{B} \sum_{b=1}^B \mathbb{P}_{n,b}^1 \frac{\Delta_a^\eta(x) \Delta_c(x)}{H_{a,b}^{npar}(x) H_{c,b}^{npar}(x)} \{I(U > x) - S_U(x; \eta)\}. \quad (11)$$

The next theorem shows that, when undersmoothed highly adaptive lasso estimator is used to model the nuisance parameters, we can still develop the asymptotic linearity of our estimators.

Assumption 2 (Complexity of nuisance functions). *For each $j = 1, 2, \dots, l(x)$, the functions $Q_j(x) = E\{I(U > x) | \bar{A}_j = \bar{g}_j^\eta, C_{j-1}(x) = 0, \bar{Z}_j\}$, $h_{a,j}$ and $h_{c,j,x}$ are càdlàg with finite sectional variation norm.*

Assumption 2 is rather mild and includes a wide class of true conditional mean models likely encountered in practice. Specifically, it defines the overall smoothness assumption for the true functions, which is considerably less restrictive compared to local smoothness assumptions imposed by Hölder balls (e.g., Robins et al., 2008a, 2017; Mukherjee et al., 2017). This assumption facilitates a rapid convergence rate of $n^{-1/3}$ (up to a log factor) achievable through the highly adaptive lasso, irrespective of the dimensionality (van der Laan, 2023).

Assumption 3 (Undersmoothing). *Let*

$$f_j^a = \frac{Q_j(x)I(\bar{A}_{j-1} = \bar{g}_{j-1}^\eta)I\{C_{j-1}(x) = 0\}\hat{h}_{c,j,x}(0 \mid \bar{z}_s, \bar{a}_s)}{\prod_{s \leq j} \{1 - h_{c,s}(1 \mid \bar{z}_s, \bar{a}_s)\} \prod_{s \leq j} \{h_{a,s}(g_j^\eta \mid \bar{z}_s, \bar{a}_{s-1})\}}$$

$$f_j^c = \frac{Q_j(x)I(\bar{A}_{j-1} = \bar{g}_{j-1}^\eta)I\{C_{j-1}(x) = 0\}I(A_j = g_j^\eta)}{\prod_{s \leq j} \{1 - h_{c,s}(1 \mid \bar{z}_s, \bar{a}_s)\} \prod_{s \leq j} \{h_{a,s}(g_j^\eta \mid \bar{z}_s, \bar{a}_{s-1})\}},$$

where $Q_j(x) = E\{I(U > x) \mid \bar{A}_j = \bar{g}_j^\eta, C_{j-1}(x) = 0, \bar{Z}_j\}$ and $j = 1, 2, \dots, l(x)$. Let $f_{\phi,j}^a$ and $f_{\phi,j}^c$ be the projections of f^a and f^c onto a linear span of basis functions ϕ_k in $L^2(P)$, for ϕ_k satisfying conditions (S12) and (S13) of Lemma S1. Then, $\|f_j^a - f_{\phi,j}^a\|_{2,\mu} = O_p(n^{-1/4})$ and $\|f_j^c - f_{\phi,j}^c\|_{2,\mu} = O_p(n^{-1/4})$ where μ is a σ -finite measure.

Assumption 3 is a crucial condition for maintaining the asymptotic linearity of $\hat{R}^{npar}(\hat{\eta}_{opt}^{npar})$.

It asserts that when the estimated coarsening mechanisms (i.e., $h_{a,j}$ and $h_{c,j,x}$) are appropriately undersmoothed, the generated features can approximate functions f_j^a and f_j^c at a rate of $n^{-1/4}$. Under Assumption 2, f_j^a and f_j^c fall into the class of càdlàg functions with a finite sectional variation norm, and thus can be approximated using the highly adaptive lasso-generated basis functions at a rate of $n^{-1/3}$ (up to a $\log n$ factor). It's noteworthy that the required rate in Assumption 3, $O_p(n^{-1/4})$, is slower than the $O_p(n^{-1/3})$ rate obtained by the highly adaptive lasso estimator, affirming the credibility of the assumption.

In the proof of Theorem 3, we show that our estimator will be asymptotically linear when $\left| \mathbb{P}_n \sum_{j=0}^{l(x)} f_j^a \left\{ I(A_j = g_j^\eta) - \hat{h}_{a,j}^{npar}(g_j^\eta \mid \bar{a}_{j-1}, \bar{z}_j) \right\} \right| = o_p(n^{-1/2})$ and $\left| \mathbb{P}_n \sum_{j=0}^{l(x)} f_j^c \left\{ I(C_j(x) = 0) - \hat{h}_{c,j,x}^{npar}(0 \mid \bar{a}_j, \bar{z}_j) \right\} \right| = o_p(n^{-1/2})$. Lemma S1 in the Supplementary Material provides the theoretical undersmoothing prerequisites for $\hat{h}_{a,j}^{npar}(g_j^\eta \mid \bar{a}_{j-1}, \bar{z}_j)$ and $\hat{h}_{c,j,x}^{npar}(0 \mid \bar{a}_j, \bar{z}_j)$ to satisfy the $o_p(n^{-1/2})$ conditions.

Theorem 3. *Let $L(\cdot)$ be a log-likelihood loss function. Assume that conditions (C.1), (C.2), (C.4) and (C.5) in Appendix S1 hold and both treatment assignment and censoring mechanisms belong to a class of càdlàg functions with finite sectional variation norm. Let $\hat{H}_{a,\lambda_n^a}^{npar}$ and $\hat{H}_{c,\lambda_n^c}^{npar}$ be highly adaptive lasso estimators of H_a and H_c using L_1 norm bounds equal to*

λ_n^a and λ_n^c where λ_n^a and λ_n^c are data dependent parameter chosen such that conditions (S12) and (S13) are satisfied. Then $n^{1/2}\{\hat{S}^{npar}(\hat{\eta}_{opt}^{npar}) - S(\eta_{opt})\} = n^{-1/2} \sum_{i=1}^n IC_{HAL}(o_i) + o_p(1)$, where the influence function $IC_{HAL}(o_i)$ is defined in Appendix S1.

A direct consequence of Theorem 3 is that the RQAL with highly adaptive lasso will be asymptotic linear with $n^{1/2} \left\{ \hat{R}^{npar}(\hat{\eta}_{opt}^{npar}) - R(\eta_{opt}) \right\} = n^{-1/2} \sum_{i=1}^n \int_0^{L_U} IC_{HAL}(o_i; x, \eta_{opt}) dx + o_p(1)$. An important implication of Theorems 2 and 3 is that when the form of the influence functions is unknown, inference is attainable via the standard bootstrap (Cai and van der Laan, 2020a). Alternatively, for $\diamond \in \{par, npar\}$, one may use a conservative variance estimator for $\hat{R}^\diamond(\hat{\eta}_{opt})$ defined as $\hat{\sigma}_{\hat{\eta}_{opt}, \diamond}^2 = n^{-1} \sum_{i=1}^n \left(\int_0^{L_U} \widehat{IC}_{x, \hat{\eta}}^\diamond(o_i) \right)^2$, where $\widehat{IC}_{x, \hat{\eta}}^\diamond(o_i) = W_{x, \hat{\eta}}^\diamond(o_i) \{I(U_i > x) - S_U(x; \hat{\eta})\} / n^{-1} \sum_{i=1}^n W_{x, \hat{\eta}}^\diamond(o_i)$ and $W_{x, \hat{\eta}}^\diamond(o_i) = \Delta_{a, i}^{\hat{\eta}}(x) \Delta_{c, i}(x) \{I(U_i > x) - S_U(x; \hat{\eta})\} / \hat{H}_{a, \lambda_n^a, i}^\diamond(x) \hat{H}_{c, \lambda_n^c, i}^\diamond(x)$.

5 Simulation Results

We conduct simulation studies to investigate the performance of the proposed estimator for restricted quality-adjusted lifetime (RQAL). We estimate the weight functions using logistic regression, highly adaptive lasso, and random forest where we used R packages `randomForest` (Liaw et al., 2002) and `hal9001` (Coyle et al., 2019) with all tuning parameters set to their respective default values. The only exception is the parameter λ in highly adaptive lasso that is specified using the undersmoothing criteria presented in Section 3.4.

Let the end time of the study be L , the gap between two landmarks be G . Let $K = L/G$ be the number of stages in the simulated longitudinal study. Define the true optimal regime as $g^{\eta_{opt}} = \{g_j^{\eta_{opt}} = I(\eta_{opt}^\top Z_j \geq 0)^{1-A_j-1}, j = 1, \dots, K\}$ where $\eta_{opt} = (1, -1, -1)$. For convenience of notation, we define $W_j = I(\bar{A}_j = g^{\eta_{opt}})$ indicating whether the observed treatment assignment \bar{A}_j matches the true optimal regime up to the j th landmark. We generate baseline values $Z_0 = (Z_{10}, Z_{20}, Z_{30})$ independently from `uniform(0.6, 1)` and set

$A_0 = C_0 = 0$. If a patient is still at risk at the j th landmark, we will update the time-dependent information $Z_j = (Z_{1j}, Z_{2j}, Z_{3j})$ where $Z_{1j} = \text{uniform}(0.4 + 0.02 \times K, 1) \times Z_{10}$, $Z_{2j} = \text{uniform}(0.4 + 0.02 \times K, 1) \times Z_{20}$ and $Z_{3j} = \text{uniform}\{0.4 + 0.02 \times K - (1 - W_j) \times 0.1, 1 - (1 - W_j) \times 0.1\} \times Z_{30}$. In our model, Z_{3j} serves the role of the quality of life score assessed at time j . Patients who do not follow the optimal regime $g^{\eta_{opt}}$ will receive a penalty with respect to quality of life score or the hazard rate function for survival, and thus, their expected quality-adjusted lifetime will not be optimal when deviating from η_{opt} .

We assume the data generation process follows the following discrete hazard models: $h_{a,j}(\bar{z}) = \text{logit}^{-1}(\kappa_0 + \kappa_1 \times z_{1j} + \kappa_2 \times z_{2j})$ and $h_{c,j}(\bar{z}, \bar{a}) = \text{logit}^{-1}(\nu_0 + \nu_1 \times z_{1j} + \nu_2 \times z_{2j})$, where $(\kappa_0, \kappa_1, \kappa_2) = (0.5 - 0.1 \times K, -0.5, -0.5)$, $(\nu_0, \nu_1, \nu_2) = (-2 + 0.5 \times \kappa_0, -1, -1)$. We consider two scenarios for the treatment and censoring mechanisms:

- *Scenario 1* (hazard models are correctly specified by logistic models): In this scenario, the covariates Z is observed and can be used to model the discrete hazard functions for treatment assignment and censoring.
- *Scenario 2* (hazard models are mis-specified by logistic models): In this scenario, the transformed covariates $X = \{(Z_1 + Z_2 - 1)^2, (Z_1 - 0.5)^2, Z_3\}$ is observed and can be used model the discrete hazard functions for treatment assignment and censoring.

We generate 500 datasets of sizes 250 and 500 according to each scenario where, for $t \in (jG, (j + 1)G]$ and $j = 0, \dots, K - 1$, the discrete survival hazard model $h_y(t, \bar{a}, \bar{z}) = \text{pr}(T = t | T > t - 1, \bar{A}_j = \bar{a}_j, \bar{C}_j = 0, \bar{Z}_j = \bar{z})$ is $h_y(t, \bar{a}, \bar{z}) = \text{logit}^{-1}\{-5 + t \times (6 \times 0.75)/K - 0.5 \times z_{1j} - 0.5 \times z_{2j} - 0.5 \times a_j \times W_j\}$.

We then the define the utility map as $\mathcal{Q}(Z_{3j}) = Z_{3j} / \max_{k=0, \dots, K} Z_{3k}$ which is always a number between $(0, 1]$. Patients' quality of life scores will be set to zero if they are no longer at-risk. This set up will generate the data with around 15% censoring. To approximate the true optimal restricted quality-adjusted lifetime, we generated 10^5 samples from the

true survival model with treatment assignment according to $g^{\eta_{opt}}$. To estimate the optimal regimes, a genetic optimization algorithm in the R package `rgenoud` (Mebane Jr et al., 2011) was implemented.

When the weight functions are estimated using random forest, the proposed estimator is no longer asymptotically linear, and the theoretical formula for the standard error does not exist. We used the conservative variance estimator presented at the end of Section 4.3.

Remark 2. *To simplify the presentation of different estimators in the tables, we use the notation \check{R} and $\check{\eta}$ to represent the estimated quantities corresponding to the methods specified in each row. For example, in Table 1, $\check{R}(\check{\eta}_{opt})$ in row 2 denotes $\hat{R}^{npar}(\hat{\eta}_{opt}^{npar})$, whereas in row 4, it denotes $\tilde{R}^{par}(\tilde{\eta}_{opt}^{par})$.*

5.1 Correctly Specified Hazard Models

We generate the treatment and censoring variables according to scenario 1 and fit a correctly specified logistic model for the treatment assignment and the censoring mechanism. We consider the following two setups for L and G . In the first set of simulation, we set $L = 60$ and $G = 10$ with the true targwt parameter $R(\eta_{opt}) = 21.04$ weeks. In the second set of simulation, we set $L = 100$ and $G = 4$ with the true $R(\eta_{opt}) = 31.74$ weeks in this case. See Section S1.6 of the Supplementary Material for more details.

Table S3 in Section S1.6 of the Supplementary Material shows the results of the estimated optimal treatment length strategy using the proposed method where weight functions are estimated using logistic regression, highly adaptive lasso and random forest. The highly adaptive lasso and logistic regression perform similarly in terms of estimating RQAL under the estimated optimal regime (i.e., $\check{R}(\check{\eta}_{opt})$). Importantly, the empirical standard errors reported in the parenthesis in the column $\check{R}(\check{\eta}_{opt})$ matches closely the estimated theoretical standard errors (i.e., SE) which confirm our results in Theorem 3. Figure 1 (first row) displays the coverage probabilities where the dashed line represents the nominal coverage




Figure 1: The coverage rates of the estimated RQAL in Scenarios 1 (first row) and 2 (second row) with logistic (LR), highly adaptive lasso (HAL) and random forest (RF) hazard models. IPW is the estimator (6) while BC-IPW is the bias-corrected estimator (8). The number in brackets is the sample size and the dashed line shows the nominal rate of 0.95.

of 0.95. In general, both logistic (circle) and highly adaptive lasso (cross) fits lead to reasonable coverage probabilities. When $K = 25$ and $n = 250$, the estimated theoretical standard errors tend to underestimate empirical standard errors, which partially leads to the insufficient coverage probabilities. This is expected because for larger number of stages, we would need more sample to learn the optimal regime. However, when the sample size is increased to 500, the empirical standard errors match the estimated standard errors and the coverage probabilities are improved considerably. See Section S1.6 of the Supplementary Material for additional discussion and figures.

Table 1: Summary table of simulation studies in Scenario 2. The target parameter is $R(\eta_{\text{opt}}) = 21.04$ when $K = 6$ and $R(\eta_{\text{opt}}) = 31.74$ when $K = 25$. The numbers in brackets are the empirical standard deviation. The notation \check{R} and $\check{\eta}$ denote the estimated quantities corresponding to the methods in each row.

Method	Model	K	n	η_0	η_1	η_2	$\check{R}(\check{\eta}_{\text{opt}})$	SE	CP	$R(\check{\eta}_{\text{opt}})$	MR
IPW	Logit	6	250	1.08(0.19)	1.03(0.32)	0.96(0.15)	21.63(0.57)	0.59	0.82	20.41(0.26)	19.24(7.59)
	HAL			1.01(0.18)	1.07(0.35)	0.99(0.15)	21.26(0.53)	0.55	0.92	20.69(0.33)	10.72(10.40)
	RF			1.04(0.39)	1.08(0.76)	1.00(0.32)	21.75(0.75)	0.69	0.78	20.54(0.52)	16.20(16.04)
BC-IPW	Logit		250	1.07(0.19)	1.06(0.35)	0.95(0.14)	21.46(0.56)	0.59	0.89	20.44(0.25)	19.15(7.40)
	HAL			1.02(0.19)	1.06(0.36)	1.00(0.16)	20.93(0.52)	0.56	0.97	20.70(0.29)	10.46(9.54)
	RF			1.04(0.46)	1.10(0.81)	0.98(0.36)	21.57(0.75)	0.69	0.84	20.51(0.51)	17.09(16.06)
IPW	Logit	6	500	1.06(0.09)	0.99(0.16)	0.96(0.08)	21.49(0.40)	0.42	0.81	20.47(0.15)	17.60(4.55)
	HAL			1.01(0.09)	1.02(0.17)	1.00(0.09)	21.15(0.38)	0.39	0.94	20.84(0.16)	5.81(5.40)
	RF			1.02(0.18)	1.05(0.38)	1.00(0.15)	21.52(0.56)	0.53	0.81	20.77(0.30)	8.92(9.77)
BC-IPW	Logit		500	1.06(0.09)	1.01(0.16)	0.96(0.07)	21.37(0.40)	0.42	0.88	20.51(0.14)	17.28(4.28)
	HAL			1.01(0.09)	1.01(0.16)	1.00(0.08)	20.92(0.37)	0.40	0.95	20.84(0.15)	5.79(4.92)
	RF			1.03(0.20)	1.05(0.37)	1.00(0.16)	21.38(0.56)	0.53	0.86	20.75(0.32)	9.50(10.21)
IPW	Logit	25	250	1.10(0.27)	1.03(0.40)	0.96(0.17)	33.13(1.10)	1.00	0.62	28.45(0.86)	66.84(10.44)
	HAL			1.05(0.37)	1.10(0.65)	0.99(0.27)	32.37(0.83)	0.87	0.85	30.14(0.33)	33.09(24.18)
	RF			1.13(0.78)	1.10(0.89)	1.02(0.28)	33.75(0.89)	0.72	0.30	30.05(1.78)	34.46(24.43)
BC-IPW	Logit		250	1.11(0.29)	1.03(0.43)	0.96(0.18)	32.75(1.08)	1.03	0.74	28.44(0.72)	68.00(10.73)
	HAL			1.07(0.44)	1.06(0.58)	1.01(0.27)	31.59(0.86)	0.92	0.95	30.26(1.06)	32.53(22.61)
	RF			1.11(0.81)	0.96(1.21)	1.05(0.37)	33.46(0.94)	0.77	0.41	30.05(1.35)	36.31(24.60)
IPW	Logit	25	500	1.06(0.10)	1.01(0.17)	0.96(0.09)	32.73(0.87)	0.81	0.70	28.51(0.44)	67.37(8.15)
	HAL			1.02(0.15)	1.03(0.23)	1.00(0.12)	32.08(0.72)	0.69	0.86	30.90(0.77)	18.08(17.11)
	RF			1.05(0.27)	1.03(0.42)	1.02(0.19)	33.31(0.80)	0.72	0.41	30.63(0.94)	25.48(20.44)
BC-IPW	Logit		500	1.06(0.09)	1.01(0.16)	0.96(0.07)	32.44(0.88)	0.84	0.79	28.48(0.41)	68.66(7.68)
	HAL			1.01(0.11)	1.02(0.20)	1.00(0.10)	31.47(0.70)	0.73	0.95	30.95(0.65)	17.20(14.85)
	RF			1.04(0.28)	1.08(0.54)	1.00(0.21)	33.04(0.83)	0.76	0.54	30.56(1.02)	26.84(22.01)

5.2 Misspecified Hazard Models

In Scenario 2, the logistic models include the transformed covariates X instead of Z , and thus, logistic regressions using X are misspecified. We consider two setups for L and G that are similar to those in Section 5.1 and generate 500 datasets of sizes 250 and 500.

Table 1 summarizes the results of Scenario 2. In this table, IPW is the direct estimation using (6), BC-IPW is the bias-correction estimator (8); Logit, logistic regression, HAL, highly adaptive lasso, RF, random forest; K is the number of stages; n is the sample size; η_0, η_1, η_2 are the empirical average of $\check{\eta}_0/\check{\eta}_1, \check{\eta}_1/\check{\eta}_2, \check{\eta}_2/\check{\eta}_0$; $\check{R}(\check{\eta}_{opt})$ is the empirical average of estimated RQAL of the estimated optimal regime. SE is the standard error estimation; CP is the average coverage probability; $R(\check{\eta}_{opt})$ is the average of true RQAL of the estimated optimal regime; MR is the empirical mis-classification rate of the estimated optimal regime. The numbers in brackets are the empirical standard deviation.

The undersmoothed highly adaptive lasso outperforms the logistic regression and random forest based estimators in terms of bias, coverage probabilities and misclassification rates. Specifically, the highly adaptive lasso leads to an estimator with smallest bias in $\hat{R}(\hat{\eta}_{opt}), \tilde{R}(\tilde{\eta}_{opt}), R(\hat{\eta}_{opt})$ and $R(\tilde{\eta}_{opt})$ and the coverage probabilities are close to the nominal rate. The misclassification rate of the highly adaptive lasso is also substantially lower than the other two estimators. Figure 1 (second row) displays the coverage probabilities where the dashed line represents the nominal coverage of 0.95. The random forest results in coverage probabilities as low as 0.3 with $n = 250$ and $K = 25$. The highly adaptive lasso estimator has the coverage probability of 0.95 for the bias corrected estimator for $n = 500$ while the coverage probabilities for the logistic regression and the random forest are 0.77 and 0.54, respectively.

Similar to Section 5.1, we provide additional information about the distribution of the estimators in both $K=6$ and 25 in Section A.4 of the Supplementary Material (Figures S5

and S6). The figures provide insight about the distribution of (a) the true restricted quality-adjusted lifetime under different estimated optimal regimes (i.e., $R(\hat{\eta}_{opt})$ and $R(\tilde{\eta}_{opt})$); (b) the estimated restricted quality-adjusted lifetime under the estimated optimal regime (i.e., $\hat{R}(\hat{\eta}_{opt})$ and $\tilde{R}(\tilde{\eta}_{opt})$); and, (c) the misclassification rates. The plots does not show any evidence of deviating from normal distribution of our estimators, thereby, supporting our theoretical results. The highly adaptive lasso outperforms the logistic regression and the random forest for both sample sized 250 and 500. The importance of using the highly adaptive lasso is more pronounced when there are more stages (i.e., $K=25$ in Figures S6).

In Section A.5 of the Supplementary Material, we provide additional simulation studies to compare the performance of the undersmoothed and the cross-validated highly adaptive lasso. The results show that while the cross-validated based estimator results in severely under-covered confidence intervals (as low as 69%), the coverage of the undersmoothed based estimator is close to the nominal rate.

6 Application: The effect of PEG tube in patients with ALS

6.1 Existing results

Atassi et al. (2011) used marginal structural models to assess the causal effect of PEG on survival time among patients with ALS (Robins et al., 2000). They concluded that PEG placement was associated with a 28% increase in the hazard of death, permanent assisted ventilation (PAV), or tracheostomy after accounting for confounding from vital capacity and functional capability. McDonnell et al. (2017) also found that PEG placement may harm survival with no quality of life related benefit for people with ALS by considering marginal structural models and structural nested models (Van den Berg et al., 2005; Hernán

et al., 2005). There are also a number of studies on elderly subjects which did not find evidence to support PEG feeding (Callahan et al., 2000; Finucane et al., 2007). Despite the lack of convincing evidence on the benefits of PEG tube placement, it is still widely used and considered as the current standard of care (Rabeneck et al., 1997; Rowland and Shneider, 2001; Miller et al., 2009; Andersen et al., 2012).

Most of the existing substantive literature has focused on the average effect of PEG on survival time or quality of life among ALS patients. However, in practice, an alternative clinical endpoint is a combination of the quantity and quality of life (Rabeneck et al., 1997). For example, for a subgroup of patients, PEG insertion may increase the survival time but reduce the quality of life drastically; conversely, delaying PEG insertion may improve quality of life but decrease the survival time. This investigation is motivated by a recent paper by Dr. David Schoenfeld and colleagues (McDonnell et al., 2017) that concluded PEG insertion has no benefit for survival or quality of life outcomes.

6.2 Data source

We use the data from a double-blinded, placebo-controlled randomized clinical trial (NCT00349622) designed to assess ceftriaxone’s effects on survival in patients with ALS (Berry et al., 2013; Cudkowicz et al., 2014). From 2006 to 2012, the ceftriaxone trial enrolled 513 participants with laboratory-supported probable, probable, or definite ALS according to the El Escorial criteria (Brooks et al., 2000). Patients visit the clinic every four weeks by protocol, and may receive PEG based on their disease progression and status. Therefore, PEG insertion is not randomized and its effect can be confounded by patients’ characteristics.

6.3 Our approach

We are using data of 481 samples from a multicenter placebo-controlled RCT (NCT00349622) designed to assess ceftriaxone’s effects on survival and functional decline in patients with

ALS. Per protocol, participants completed assessments at baseline; weeks 1, 2, and 4; and every four weeks thereafter. Dates of PEG placement were recorded. To simplify the analysis, we only consider the first 120 weeks (i.e., at most 30 time points) data, because more than 95% of patients are dead or censored before this time points. Censoring is defined at the time after more than a 17-week gap between the 2 ALS Functional Rating Scale (ALSFRS-R) assessments or more than a 33-week gap between 2 BMI, FVC, grip strength, or ALSSQOL measurements (McDonnell et al., 2017). We set the interval between two assessments as $G = 4$. Thus, we have a $K = 30$ design. Baseline characteristics considered for confounding included sex, age, disease duration, site of symptom onset, respiratory comorbidities (history and prevalence at screening), cardiovascular comorbidities (history and prevalence at screening), use of riluzole, use of ceftriaxone, ALSFRS-R total score and four subscales (bulbar, breathing, fine motor, and gross motor), percent change in body mass index (BMI) from screening, forced vital capacity (FVC), grip strength using handheld dynamometry, and the ALSSQOL. Survival was defined as the time in weeks from screening until death, PAV, or tracheostomy. Quality of life (QOL) was measured using the average total score of the ALS-Specific Quality of Life (ALSSQOL) inventory, a 59-item patient-reported assessment that evaluates perceived QOL concerning negative emotion, interaction with people and the environment, intimacy, religiosity, physical symptoms, and bulbar function. The utility coefficient map is defined as $\mathcal{Q} := \text{ALSSQOL}(t) / \max_t \text{ALSSQOL}(t)$. Here, we focus on maximizing RQAL as the outcome of interest while the restricted value for QAL is set as $L_U = 91.12$ by picking the 95% quantile of the observed QAL. According to experts' advice, two important time-dependent clinical variables, percent change in BMI from screening and FVC, are considered in the treatment regime g^η , which means the PEG insertion is recommended based on the value of the linear combination of percent change of BMI and FVC.

Table S5 in Section S2 of the Supplementary Material shows the estimated coefficients

of the measured variables on the probability of PEG tube insertion and censoring on logit scale. Based on these results, the ALS bulbar and FVC are the most significant variables in deciding for PEG placement such that patients with higher values of bulbar and FVC are less likely to receive PEG. Also, the stage (i.e., number of months since the beginning of the follow-up), site of onset, Ceftriaxone, and ALS total score contribute significantly in hazard of censoring.

To estimate the optimal treatment length strategy, we apply the proposed method using logistic regression and highly adaptive lasso for the hazard models. The estimate nuisance parameters are plugged into the value function (6) and (8) and maximized to obtain the estimated optimal strategy. We find that logistic regression and highly adaptive lasso return similar RQAL estimation under different regimes. The only difference is the effect of BMI reduction in the estimated optimal regimes. The results of estimated optimal strategy $g^{\hat{\eta}_{opt}}$ and $g^{\tilde{\eta}_{opt}}$ are summarized in Table 2. Since these two estimated regimes are nearly identical, we consider only $g^{\hat{\eta}_{opt}}$ for simplicity of representation. The coefficients for FVC and BMI reduction are negative, which means $g^{\hat{\eta}_{opt}}$ tends to recommend PEG when both FVC and BMI reduction decrease (when BMI decreases, this is a negative number, and thus, the larger is the score, the better is the patient), and the direction of these parameters coincides with the experts' knowledge. The FVC variable is on a large scale, which explains the small coefficient.

We compare the RQAL estimation with four other treatment regimes: immediate PEG insertion ($R(1)$), never PEG insertion ($R(0)$), insert the PEG when FVC score is less than 50 ($R(F)$) and insert the PEG when BMI changes -10% or greater ($R(B)$). The latter two treatment regimes are currently used in clinical practice. The estimated regime $g^{\hat{\eta}_{opt}}$ is significantly better than the always/never insertion strategies by looking at the point estimation and 95% confidence intervals of RQAL. Moreover, the estimated RQAL under the estimated treatment regime using highly adaptive lasso is roughly 16 and 11 points




Figure 2: The estimated survival curve of QAL under regime $g^{\tilde{\eta}_{opt}}$ (solid line), never insertion (black dashed line), always insertion (dotted line), FVC regime (red dashed line), and BMI regime (blue dashed line) with logistic hazard models.

higher than the RQAL obtained by the existing treatment rules $\hat{R}(B)$ and $\hat{R}(F)$, respectively. The difference between $\hat{R}(\hat{\eta}_{opt})$ and $\hat{R}(B)$ is also statistically significant. Regardless, the estimated regime provides clinically significant improvement on RQAL compared with both existing rules.

The IPW and Bias-correction estimators are equal for always/never insertion regimes. The reason is because we use $\eta = (1, 0, 0)$ and $(-1, 0, 0)$ as the representative regimes with only intercept, and $sd(\mathcal{Z})$, a scale factor in the bandwidth h , would be 0 under these regimes. With 0 bandwidth, the bias-corrected estimator reduces to the IPW estimator. Another issue is that the standard error estimation of the immediate insertion regime is pretty large. This is expected since few people (2.5%) in this data set start the treatment at stage 1, and this fact results in the high variability of the proposed IPW estimator. This phenomenon can also be seen from Figure 2 about the overall estimated survival curve.

Table 2: Estimated optimal treatment length strategy for PEG insertion.

PS	Est	(Int.)	FVC	BMI Red.	$\hat{R}(\hat{\eta}_{opt})$	$\hat{R}(1)$	$\hat{R}(0)$	$\hat{R}(F)$	$\hat{R}(B)$
Logit	IPW	0.995	-0.011	-0.097	72.24(4.75)	33.07(8.08)	57.30(2.51)	61.26(2.57)	58.51(3.17)
	BC-IPW	0.996	-0.014	-0.084	71.95(4.14)	33.07(8.08)	57.30(2.51)	60.84(2.57)	58.52(3.17)
HAL	IPW	0.998	-0.012	-0.066	76.09(4.29)	23.59(7.08)	60.24(2.16)	65.01(2.53)	60.66(2.39)
	BC-IPW	0.998	-0.012	-0.054	75.05(4.25)	23.59(7.08)	60.24(2.16)	64.70(2.53)	60.66(2.39)

Figure 2 shows the estimated survival curves of QAL under different regimes using logistic regression. For simplicity, we only plot the survival curve of the IPW estimator. The estimated survival curve under the estimated optimal strategy $g^{\hat{\eta}_{opt}}$ is uniformly better than under the other regimes, indicating that $g^{\hat{\eta}_{opt}}$ may lead to improved QAL if used in ALS study. Additionally, the estimated survival curve shows that the regime that no patients insert the PEG leads to better patient outcomes than the regime that all patients insert the PEG immediately, which partially explains why the previous studies by using MSM conclude that PEG insertion is harmful to survival for patients with ALS.

7 Discussion

There are several directions for future work with this interesting problem. The proposed methods directly maximize the estimated restricted quality-adjusted lifetime using a genetic algorithm and it is computationally expensive in multi-stage cases. It is also challenging to handle high-dimensional covariate space in g^n . The extensions of the proposed method to allow for high-dimensional covariates could be considered. Also, the proposed method only search optimal strategies in a restricted functional class, say the linear classifiers with the linear combination of some covariates. This type of regimes are easy to interpret but can be sub-optimal when the actual optimal regime is complex. Thus, the trade-off between interpretability and accuracy of the estimated optimal regime and extending the search

space of optimal regime to more general functional class would be considered.

References

- Andersen, P. M., Abrahams, S., Borasio, G. D., de Carvalho, M., Chio, A., Van Damme, P., Hardiman, O., Kollewe, K., Morrison, K. E. et al. (2012) Efn guidelines on the clinical management of amyotrophic lateral sclerosis (mals)—revised report of an efn task force. *European journal of neurology*, **19**, 360–375.
- Atassi, N., Cudkowicz, M. E. and Schoenfeld, D. A. (2011) Advanced statistical methods to study the effects of gastric tube and non-invasive ventilation on functional decline and survival in amyotrophic lateral sclerosis. *Amyotrophic Lateral Sclerosis*, **12**, 272–277.
- Bai, X., Tsiatis, A. A. and O’Brien, S. M. (2013) Doubly-robust estimators of treatment-specific survival distributions in observational studies with stratified sampling. *Biometrics*, **69**, 830–839.
- Bang, H. and Tsiatis, A. A. (2002) Median regression with censored cost data. *Biometrics*, **58**, 643–649.
- Benkeser, D., Carone, M., Laan, M. V. D. and Gilbert, P. (2017) Doubly robust nonparametric inference on the average treatment effect. *Biometrika*, **104**, 863–880.
- Benkeser, D. and Van Der Laan, M. (2016) The highly adaptive lasso estimator. In *Proceedings of the... International Conference on Data Science and Advanced Analytics. IEEE International Conference on Data Science and Advanced Analytics*, vol. 2016, 689. NIH Public Access.
- Van den Berg, J., Kalmijn, S., Lindeman, E., Veldink, J., De Visser, M., Van der Graaff,

- M., Wokke, J. and Van den Berg, L. (2005) Multidisciplinary als care improves quality of life in patients with als. *Neurology*, **65**, 1264–1267.
- Berry, J. D., Shefner, J. M., Conwit, R., Schoenfeld, D., Keroack, M., Felsenstein, D., Krivickas, L., David, W. S., Vriesendorp, F., Pestronk, A. et al. (2013) Design and initial results of a multi-phase randomized trial of ceftriaxone in amyotrophic lateral sclerosis. *PloS one*, **8**, e61177.
- Bibaut, A. F. and van der Laan, M. J. (2019) Fast rates for empirical risk minimization over $c\backslash adl\backslash ag$ functions with bounded sectional variation norm. *arXiv preprint arXiv:1907.09244*.
- Brooks, B. R., Miller, R. G., Swash, M. and Munsat, T. L. (2000) El escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotrophic lateral sclerosis and other motor neuron disorders*, **1**, 293–299.
- Cai, W. and van der Laan, M. (2020a) Nonparametric bootstrap inference for the targeted highly adaptive least absolute shrinkage and selection operator (lasso) estimator. *The International Journal of Biostatistics*, **16**.
- Cai, W. and van der Laan, M. J. (2020b) One-step targeted maximum likelihood estimation for time-to-event outcomes. *Biometrics*, **76**, 722–733.
- Callahan, C. M., Haag, K. M., Weinberger, M., Tierney, W. M., Buchanan, N. N., Stump, T. E. and Nisi, R. (2000) Outcomes of percutaneous endoscopic gastrostomy among older adults in a community setting. *Journal of the American Geriatrics Society*, **48**, 1048–1054.
- Chakraborty, B. and Moodie, E. (2013) *Statistical methods for dynamic treatment regimes*. Springer.

- Chen, P.-Y. and Tsiatis, A. A. (2001) Causal inference on the difference of the restricted mean lifetime between two groups. *Biometrics*, **57**, 1030–1038.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C. and Newey, W. (2017) Double/debiased/neyman machine learning of treatment effects. *American Economic Review*, **107**, 261–65.
- Coyle, J. R., Hejazi, N. S. and van der Laan, M. J. (2019) *hal9001: The scalable highly adaptive lasso*. URL: <https://doi.org/10.5281/zenodo.3558314>. R package version 0.2.5.
- Cudkowicz, M. E., Titus, S., Kearney, M., Yu, H., Sherman, A., Schoenfeld, D., Hayden, D., Shui, A., Brooks, B., Conwit, R. et al. (2014) Safety and efficacy of ceftriaxone for amyotrophic lateral sclerosis: a multi-stage, randomised, double-blind, placebo-controlled trial. *The Lancet Neurology*, **13**, 1083–1091.
- Díaz, I., Savenkov, O. and Ballman, K. (2018) Targeted learning ensembles for optimal individualized treatment rules with time-to-event outcomes. *Biometrika*, **105**, 723–738.
- Ertefaie, A., Hejazi, N. S. and van der Laan, M. J. (2023) Nonparametric inverse-probability-weighted estimators based on the highly adaptive lasso. *Biometrics*, **79**, 1029–1041.
- Fine, J. P. and Gelber, R. D. (2001) Joint regression analysis of survival and quality-adjusted survival. *Biometrics*, **57**, 376–382.
- Finucane, T. E., Christmas, C. and Leff, B. A. (2007) Tube feeding in dementia: how incentives undermine health care quality and patient safety. *Journal of the American Medical Directors Association*, **8**, 205–208.

- Gelber, R. D., Gelman, R. S. and Goldhirsch, A. (1989) A quality-of-life-oriented endpoint for comparing therapies. *Biometrics*, **45**, 781–795.
- Gill, R. D., van der Laan, M. J. and Wellner, J. A. (1995) Inefficient estimators of the bivariate survival function for three models. In *Annales de l'IHP Probabilités et statistiques*, vol. 31, 545–597.
- Glasziou, P., Simes, R. and Gelber, R. (1990) Quality adjusted survival analysis. *Statistics in medicine*, **9**, 1259–1276.
- Goldberg, Y. and Kosorok, M. R. (2012) Q-learning with censored data. *Annals of statistics*, **40**, 529.
- Goldhirsch, A., Gelber, R. D., Simes, R. J., Glasziou, P. and Coates, A. S. (1989) Costs and benefits of adjuvant therapy in breast cancer: a quality-adjusted survival analysis. *Journal of Clinical Oncology*, **7**, 36–44.
- Heller, G. (2007) Smoothed rank regression with censored data. *Journal of the American Statistical Association*, **102**, 552–559.
- Hernán, M. A., Cole, S. R., Margolick, J., Cohen, M. and Robins, J. M. (2005) Structural accelerated failure time models for survival analysis in studies with time-varying treatments. *Pharmacoepidemiology and drug safety*, **14**, 477–491.
- Huang, Y. and Louis, T. A. (1998) Nonparametric estimation of the joint distribution of survival time and mark variables. **85**, 785.
- Hubbard, A. E., Van Der Laan, M. J. and Robins, J. M. (2000) Nonparametric locally efficient estimation of the treatment specific survival distribution with right censored data and covariates in observational studies. In *Statistical Models in Epidemiology, the Environment, and Clinical Trials*, 135–177. Springer.

- Jiang, R., Lu, W., Song, R. and Davidian, M. (2017) On estimation of optimal treatment regimes for maximizing t-year survival probability. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **79**, 1165–1185.
- Klaassen, C. A. (1987) Consistent estimation of the influence function of locally asymptotically linear estimators. *The Annals of Statistics*, **15**, 1548–1562.
- Kosorok, M. R. (2008) *Introduction to empirical processes and semiparametric inference*. Springer.
- van der Laan, M. (2023) Higher order spline highly adaptive lasso estimators of functional parameters: Pointwise asymptotic normality and uniform convergence rates. *arXiv preprint arXiv:2301.13354*.
- Van der Laan, M. J. (2014) Targeted estimation of nuisance parameters to obtain valid statistical inference. *The international journal of biostatistics*, **10**, 29–57.
- van der Laan, M. J. and Hubbard, A. (1999) Locally efficient estimation of the quality-adjusted lifetime distribution with right-censored data and covariates. *Biometrics*, **55**, 530–536.
- Leigh, P., Abrahams, S., Al-Chalabi, A., Ampong, M., Goldstein, L., Johnson, J., Lyall, R., Moxham, J., Mustafa, N., Rio, A. et al. (2003) The management of motor neurone disease. *Journal of Neurology, Neurosurgery & Psychiatry*, **74**, iv32–iv47.
- Liaw, A., Wiener, M. et al. (2002) Classification and regression by randomforest. *R news*, **2**, 18–22.
- Mathus-Vliegen, L. M., Louwse, L. S., Merkus, M. P., Tytgat, G. N. and de Jong, J. V. (1994) Percutaneous endoscopic gastrostomy in patients with amyotrophic lateral sclerosis and impaired pulmonary function. *Gastrointestinal endoscopy*, **40**, 463–469.

- McDonnell, E., Schoenfeld, D., Paganoni, S. and Atassi, N. (2017) Causal inference methods to study gastric tube use in amyotrophic lateral sclerosis. *Neurology*, **89**, 1483–1489.
- Mebane Jr, W. R., Sekhon, J. S. et al. (2011) Genetic optimization using derivatives: the rgenoud package for r. *Journal of Statistical Software*, **42**, 1–26.
- Miller, R. G., Jackson, C. E., Kasarskis, E. J., England, J., Forshe, D., Johnston, W., Kalra, S., Katz, J., Mitsumoto, H., Rosenfeld, J. et al. (2009) Practice parameter update: the care of the patient with amyotrophic lateral sclerosis: drug, nutritional, and respiratory therapies (an evidence-based review): report of the quality standards subcommittee of the american academy of neurology. *Neurology*, **73**, 1218–1226.
- Mukherjee, R., Newey, W. K. and Robins, J. M. (2017) Semiparametric efficient empirical higher order influence function estimators. *arXiv preprint arXiv:1705.07577*.
- Rabeneck, L., McCullough, L. B. and Wray, N. P. (1997) Ethically justified, clinically comprehensive guidelines for percutaneous endoscopic gastrostomy tube placement. *The Lancet*, **349**, 496–498.
- Robins, J. (1986) A new approach to causal inference in mortality studies with a sustained exposure period—application to control of the healthy worker survivor effect. *Mathematical modelling*, **7**, 1393–1512.
- Robins, J., Li, L., Tchetgen, E., van der Vaart, A. et al. (2008a) Higher order influence functions and minimax estimation of nonlinear functionals. *Probability and statistics: essays in honor of David A. Freedman*, **2**, 335–421.
- Robins, J., Orellana, L. and Rotnitzky, A. (2008b) Estimation and extrapolation of optimal treatment and testing strategies. *Statistics in medicine*, **27**, 4678–4721.

- Robins, J. M. (2004) Optimal structural nested models for optimal sequential decisions. In *Proceedings of the second seattle Symposium in Biostatistics*, 189–326. Springer.
- Robins, J. M., Hernan, M. A. and Brumback, B. (2000) Marginal structural models and causal inference in epidemiology.
- Robins, J. M., Li, L., Mukherjee, R., Tchetgen, E. T. and van der Vaart, A. (2017) Minimax estimation of a functional on a structured high-dimensional model. *The Annals of Statistics*, **45**, 1951–1987.
- Rosenbaum, P. R. and Rubin, D. B. (1983) The central role of the propensity score in observational studies for causal effects. *Biometrika*, **70**, 41–55.
- Rowland, L. P. and Shneider, N. A. (2001) Amyotrophic lateral sclerosis. *New England Journal of Medicine*, **344**, 1688–1700.
- Rytgaard, H. C., Gerds, T. A. and van der Laan, M. J. (2022) Continuous-time targeted minimum loss-based estimation of intervention-specific mean outcomes. *The Annals of Statistics*, **50**, 2469–2491.
- Strijbos, D., Hofstede, J., Keszthelyi, D., Masclee, A. A. and Gilissen, L. P. (2017) Percutaneous endoscopic gastrostomy under conscious sedation in patients with amyotrophic lateral sclerosis is safe: an observational study. *European journal of gastroenterology & hepatology*, **29**, 1303–1308.
- Tian, L., Alizadeh, A. A., Gentles, A. J. and Tibshirani, R. (2014) A simple method for estimating interactions between a treatment and a large number of covariates. *Journal of the American Statistical Association*, **109**, 1517–1532.
- Van der Vaart, A. W. (2000) *Asymptotic statistics*, vol. 3. Cambridge university press.

- Van der Vaart, A. W., Dudoit, S. and van der Laan, M. J. (2006) Oracle inequalities for multi-fold cross validation. *Statistics & Decisions*, **24**, 351–371.
- Vaart, A. W. and Wellner, J. A. (1996) *Weak convergence and empirical processes: with applications to statistics*. Springer.
- van der Laan, M. J. and Dudoit, S. (2003) Unified cross-validation methodology for selection among estimators and a general cross-validated adaptive epsilon-net estimator: Finite sample oracle inequalities and examples.
- Wang, H. and Zhao, H. (2006) Regression analysis of mean quality-adjusted lifetime with censored data. *Biostatistics*, **8**, 368–382.
- Yang, S. and Ding, P. (2017) Asymptotic causal inference with observational studies trimmed by the estimated propensity scores. *arXiv preprint arXiv:1704.00666*.
- Zarei, S., Carr, K., Reiley, L., Diaz, K., Guerra, O., Altamirano, P. F., Pagani, W., Lodin, D., Orozco, G. and Chinae, A. (2015) A comprehensive review of amyotrophic lateral sclerosis. *Surgical neurology international*, **6**.
- Zhang, B., Tsiatis, A. A., Laber, E. B. and Davidian, M. (2012) A robust method for estimating optimal treatment regimes. *Biometrics*, **68**, 1010–1018.
- Zhao, H. and Tsiatis, A. A. (1997) A consistent estimator for the distribution of quality adjusted survival time. *Biometrika*, **84**, 339–348.
- Zhao, Y.-Q., Zeng, D., Laber, E. B., Song, R., Yuan, M. and Kosorok, M. R. (2014) Doubly robust learning for estimating individualized treatment with censored data. *Biometrika*, **102**, 151–168.
- Zheng, W. and van der Laan, M. J. (2011) Cross-validated targeted minimum-loss-based estimation. In *Targeted Learning*, 459–474. Springer.

S1 Appendix

We first list the regularity conditions needed for this section.

Regularity Conditions:

(C.1) The covariates Z_j are bounded.

(C.2) $H_a(x; \bar{z})$ is bounded away from 0 and 1 for all possible values of x and \bar{z} ; $H_c(x; \bar{a}, \bar{z})$ is bounded away from 0 and 1 for all possible values of (x, \bar{a}, \bar{z}) .

(C.3) The treatment assignment model and censoring model are both correctly specified by logistic regression models, e.g., $P(C_j = 1 \mid \bar{Z}_j, C_{j-1} = 0) = 1 / \{1 + \exp(\beta_j^\top \bar{Z}_j)\}$ and $P(A_j = 1 \mid \bar{Z}_j, A_{j-1} = 0) = 1 / \{1 + \exp(\theta_j^\top \bar{Z}_j)\}$.

(C.4) $|\Delta_c(x) / H_a(x, \bar{Z}) H_c(x, \bar{A}, \bar{Z})| < \infty$, $|\Delta_c(x) I(U > x) / H_a(x, \bar{Z}) H_c(x, \bar{A}, \bar{Z})| < \infty$ uniformly for $x \in (0, L_U]$ and η .

(C.5) $n\nu \rightarrow \infty$ and $n\nu^4 \rightarrow 0$ as $n \rightarrow \infty$.

Condition (C.1) is a common assumption of convenience used to bound the score function and influence curves. Condition (C.2) is required to bound the weights H_a and H_c away from zero such that their reciprocals are finite. Condition (C.3) implies that, for $j = 1, \dots, K$, $\hat{\beta}_j$ and $\hat{\theta}_j$ are consistent and asymptotic normal at the nominal \sqrt{n} -rate, i.e., $n^{1/2}(\hat{\beta}_j - \beta_j) = n^{1/2}\mathbb{P}_n\{IC_{j,\beta}(O)\} + o_p(1)$ and $n^{1/2}(\hat{\theta}_j - \theta_j) = n^{1/2}\mathbb{P}_n\{IC_{j,\theta}(O)\} + o_p(1)$. Condition (C.5) gives common conditions on the smoothing parameter

S1.1 Proof of Theorem 1

Proof. In this proof, we omit the superscript ^{'par'} for simplicity, with the understanding that we are focusing on the parametric modeling of the nuisance functions.

For simplicity, we suppress the covariate information in $H_c(x; \bar{a}, \bar{z})$ and $H_a(x; \bar{z})$, and use $H_a(x)$ and $H_c(x)$ instead. We divide the proof into three steps.

Step 1: The consistency and asymptotic normality of $\hat{S}_U(x; \eta)$

We first show the consistency of the direct IPW estimator. Fix a target time x and note that $\Delta_a^\eta(x)I(U > x) = \Delta_a^\eta(x)I(U^\eta > x)$ under Assumption 1a. Thus, $E[\Delta_a^\eta(x)I(U > x)] = S_U(x; \eta)$. Then, by an ordinary Weak Law of Large Numbers,

$$\mathbb{P}_n \left[\frac{\Delta_a^\eta(x)\Delta_c(x)I(U > x)}{\hat{H}_a(x)\hat{H}_c(x)} \right] \rightarrow_p E \left[\frac{\Delta_a^\eta(x)\Delta_c(x)I(U > x)}{H_a(x)H_c(x)} \right] = S_U(x; \eta),$$

uniformly for $x \in (0, L_U]$ as $n \rightarrow \infty$ because of the correct specification of parametric propensity score models in Condition (C.2). Similarly, we have $\mathbb{P}_n[\Delta_a^\eta(x)\Delta_c(x)/\hat{H}_a(x)\hat{H}_c(x)] \rightarrow_p E[\Delta_a^\eta(x)\Delta_c(x)/H_a(x)H_c(x)] = 1$ uniformly for $x \in (0, L_U]$ as $n \rightarrow \infty$. Putting these two results together leads one to conclude that $\hat{S}_U(x; \eta) \rightarrow_p S_U(x; \eta)$ as $n \rightarrow \infty$.

We then show the asymptotic normality of $n^{1/2} \left\{ \hat{S}_U(x; \eta) - S_U(x; \eta) \right\}$. Let $\hat{S}_U(x; \eta, \theta, \beta)$ denote an estimator where the nuisance functions are known. Based on (4) and first order

Taylor expansion, we have

$$\begin{aligned}
& n^{1/2}\{\hat{S}_U(x; \eta) - S_U(x; \eta)\} \\
&= n^{1/2}\{\hat{S}_U(x; \eta, \theta, \beta) - S_U(x; \eta)\} + D_1^\top(x)n^{1/2}(\hat{\theta} - \theta) + D_2^\top(x)n^{1/2}(\hat{\beta} - \beta) + o_p(1) \\
&= n^{-1/2} \frac{\sum_{i=1}^n \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{H_{a,i}(x;\theta)H_{c,i}(x;\beta)} I(U_i > x)}{1/n \sum_{i=1}^n \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{H_{a,i}(x;\theta)H_{c,i}(x;\beta)}} - S_U(x; \eta) \\
&\quad + D_1^\top(x)n^{1/2}(\hat{\theta} - \theta) + D_2^\top(x)n^{1/2}(\hat{\beta} - \beta) + o_p(1) \\
&= n^{-1/2} \frac{\sum_{i=1}^n \left\{ \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{H_{a,i}(x;\theta)H_{c,i}(x;\beta)} I(U_i > x) - \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{H_{a,i}(x;\theta)H_{c,i}(x;\beta)} S_U(x; \eta) \right\}}{1/n \sum_{i=1}^n \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{H_{a,i}(x;\theta)H_{c,i}(x;\beta)}} \\
&\quad + D_1^\top(x)n^{1/2}(\hat{\theta} - \theta) + D_2^\top(x)n^{1/2}(\hat{\beta} - \beta) + o_p(1) \\
&= n^{-1/2} \sum_{i=1}^n \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{H_{a,i}(x;\theta)H_{c,i}(x;\beta)} \{I(U_i > x) - S_U(x; \eta)\} \\
&\quad + D_1^\top(x)n^{1/2}(\hat{\theta} - \theta) + D_2^\top(x)n^{1/2}(\hat{\beta} - \beta) + o_p(1) \\
&= n^{-1/2} \sum_{i=1}^n \{IC_{x;\eta}(o_i) + D_1^\top(x)IC_\theta(o_i) + D_2^\top(x)IC_\beta(o_i)\} + o_p(1) \\
&:= n^{-1/2} \sum_{i=1}^n IC_{PAR}(o_i; x, \eta) + o_p(1),
\end{aligned} \tag{A.2}$$

where $D_1(x) = \lim_{n \rightarrow \infty} \partial \hat{S}_U(x; \eta) / \partial \theta$, $D_2(x) = \lim_{n \rightarrow \infty} \partial \hat{S}_U(x; \eta) / \partial \beta$, $IC_{x;\eta}(o)$ is the influence function under regime g^η at time x when all nuisance parameters are known, and $IC_\beta(o)$ and $IC_\theta(o)$ are part of the parametric models defined in regularity conditions (C.3). $\hat{S}_U(x; \eta) - S_U(x; \eta)$ is thus an asymptotic linear estimator within the range $(0, L_U]$. The first term is composed of bounded probability functions and simple indicator functions, which belong to the Donsker class, and thus $n^{-1/2} \left\{ \hat{S}_U(x; \eta) - S_U(x; \eta) \right\}$ converges to a mean zero Gaussian process by empirical process theory [Van der Vaart \(2000\)](#).

Consequently, a consistent covariance estimation is given by $1/n \sum_{i=1}^n \widehat{IC}_{PAR}(o_i; x, \eta)^2$ with estimated $\hat{\theta}$ and $\hat{\beta}$, where

$$\widehat{IC}_{PAR}(o_i; x, \eta) = \frac{\left\{ \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{H_{a,i}(x;\theta)H_{c,i}(x;\beta)} I(U_i > x) - \hat{S}_U(x; \eta) \right\}}{1/n \sum_{i=1}^n \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{H_{a,i}(x;\theta)H_{c,i}(x;\beta)}} + \hat{D}_1^\top(x) \widehat{IC}_\theta(o_i) + \hat{D}_2^\top(x) \widehat{IC}_\beta(o_i).$$

Step 2: The asymptotic distribution of $\hat{S}_U(x; \hat{\eta}_{opt})$

Let $\hat{\eta}_{opt}$ be the maximizer of $\hat{S}_U(x; \eta)$ and η_{opt} be the maximizer of $S_U(x; \eta)$. We assume $S_U(x; \eta)$ is a smooth function of η . Then by the consistency of $\hat{S}_U(x; \eta)$, $\hat{\eta}_{opt}$ will converge in probability to η_{opt} . Let $W_n(\eta) = n^{1/2} \{ \hat{S}_U(x; \eta) - S_U(x; \eta) \}$. Following the argument in [Zhang et al. \(2012\)](#) and using a stochastic equicontinuity argument, we have $W_n(\hat{\eta}_{opt}) - W_n(\eta_{opt}) \rightarrow 0$ in probability. We can represent

$$\begin{aligned}
& n^{1/2} \{ \hat{S}_U(x; \hat{\eta}_{opt}) - S_U(x; \eta_{opt}) \} \\
&= W_n(\hat{\eta}_{opt}) + n^{1/2} \{ S_U(x; \hat{\eta}_{opt}) - S_U(x; \eta_{opt}) \} \\
&= W_n(\hat{\eta}_{opt}) - W_n(\eta_{opt}) + W_n(\eta_{opt}) + n^{1/2} \{ S_U(x; \hat{\eta}_{opt}) - S_U(x; \eta_{opt}) \} \tag{A.3} \\
&= W_n(\hat{\eta}_{opt}) - W_n(\eta_{opt}) + W_n(\eta_{opt}) + n^{1/2} (\hat{\eta}_{opt} - \eta_{opt}) \frac{\partial}{\partial \eta} S_U(x; \eta) \Big|_{\eta_{opt}} + o_p(1) \\
&= W_n(\eta_{opt}) + o_p(1),
\end{aligned}$$

The last equality follows because $W_n(\hat{\eta}_{opt}) = o_p(1)$ and $\frac{\partial}{\partial \eta} S_U(x; \eta) \Big|_{\eta_{opt}} = 0$ (η_{opt} is the maximizer of the function). Thus, $n^{1/2} \{ \hat{S}_U(x; \hat{\eta}_{opt}) - S_U(x; \eta_{opt}) \}$ converges to the same Gaussian process as $n^{1/2} \{ \hat{S}_U(x; \eta_{opt}) - S_U(x; \eta_{opt}) \}$. That is $\hat{S}_U(x; \hat{\eta}_{opt})$ converges to a Gaussian process with mean $S_U(x; \eta_{opt})$ and variance $E[IC_{PAR}(o; x, \eta_{opt})^2]$ which can be empirically estimated as

$$1/n \sum_{i=1}^n \widehat{IC}_{PAR}(o_i; x, \hat{\eta}_{opt})^2. \tag{A.4}$$

Step 3: The asymptotic equivalence between $\hat{S}_U(x; \hat{\eta}_{opt})$ and $\tilde{S}_U(x; \tilde{\eta}_{opt})$

Define $Q = E \{ \Delta_{a,i}^\eta(x) \Delta_{c,i}(x) / H_{a,i}(x) H_{c,i}(x) \}$ and $\tilde{Q} = E \{ \tilde{\Delta}_{a,i}^\eta(x) \Delta_{c,i}(x) / H_{a,i}(x) H_{c,i}(x) \}$. K is actually 1. Let's first consider the case with one stage.

$\mathbf{K} = 1$

Let's look at the trivial case with stage $\mathbf{K} = 1$. We first show that $n^{1/2}(\tilde{Q} - Q) = o_p(1)$.

Notice that

$$\begin{aligned} & n^{-1/2} \sum_{i=1}^n \frac{\{\Delta_{a,i}^\eta(x) - \tilde{\Delta}_{a,i}^\eta(x)\} \Delta_{c,i}(x)}{\hat{H}_{a,i}(x) \hat{H}_{c,i}(x)} \\ &= n^{-1/2} \sum_{i=1}^n \frac{\{\Delta_{a,i}^\eta(x) - \tilde{\Delta}_{a,i}^\eta(x)\} \Delta_{c,i}(x)}{H_{a,i}(x) H_{c,i}(x)} + o_p(1) \end{aligned} \quad (\text{A.5})$$

The definition of $\Delta_a^\eta(x)$ and $\tilde{\Delta}_a^\eta(x)$ tell us that when $\mathbf{K}=1$, they are the indicator and normal distribution function themselves. Define $L = (A_0, T, \delta)$, $\Xi(Z_0, L) = \Delta_c(x)/H_a(x)H_c(x)$ and $r^\eta = \eta^\top Z_0$. Following the arguments in [Heller \(2007\)](#), we have

$$|(\text{A.5})| \leq M_1 n^{1/2} \sup_{|\eta|=1} \left| \int_l \int_{r^\eta} \{I(r^\eta \geq 0) - \Phi(r^\eta \geq 0)\} \Xi(l) d\hat{F}(r^\eta|l) d\hat{G}(l) \right| \quad (\text{A.6})$$

where M_1 is a finite constant, $\hat{G}(l)$ and $\hat{F}(r^\eta|l)$ are the marginal empirical cumulative distribution functions for l and the conditional empirical cumulative distribution function for r^η , respectively. For simplicity, we omit the superscript η in r^η . [\(A.5\)](#) is thus bounded by $M_1 n^{1/2} \sup_{\|\eta\|=1} U$. Write U as $U_1 + U_2$ where

$$\begin{aligned} U_1 &= \int_l \int_{r^\eta} \{I(r^\eta \geq 0) - \Phi(r^\eta \geq 0)\} \Xi(l) \{d\hat{F}(r^\eta|q) - dF(r^\eta|q)\} d\hat{G}(q), \\ U_2 &= \int_l \int_{r^\eta} \{I(r^\eta \geq 0) - \Phi(r^\eta \geq 0)\} \Xi(l) dF(r^\eta|l) d\hat{G}(l), \end{aligned} \quad (\text{A.7})$$

with $F(r^\eta|l) = \lim_{n \rightarrow \infty} \hat{F}(r^\eta|l)$. By variable transformation $z = r^\eta/\nu$ and integration by parts, we have

$$U_1 = \int_l \int_z \Xi(l) \phi(z) \left\{ \left[\hat{F}(z\nu|l) - F(z\nu|l) \right] - \left[\hat{F}(0|l) - F(0|l) \right] \right\} dz d\hat{G}(q), \quad (\text{A.8})$$

where $\phi(z)$ is the probability density function of standard normal distribution. Under regularity condition [\(C.4\)](#), we apply the results on oscillations of empirical process ([Shorack and Wellner, 2009](#), Theorem 1, p. 542) to equation [\(A.8\)](#) and have

$$n^{1/2} U_1 = O_p \left(\sqrt{\nu \log n \log \left(\frac{1}{\nu \log n} \right)} \right).$$

In addition, by similar arguments and applying second order Taylor expansion of U_2 with respect to ν around 0, we have

$$U_2 = -\nu^2/2 \int_q \int_z \Xi(l)\phi(z)f'(z\nu^*|l)z^2 dz d\hat{G}(l),$$

where $f'(z\nu^*|l) = \partial^2 F(u|l)/\partial u^2$ and ν^* lies between ν and 0. We thus have $n^{1/2}U_2 = O_p(n^{1/2}\nu^2)$. Combine the above results, we have

$$M_1 n^{1/2}U \leq M_1 n^{1/2}(|U_1| + |U_2|) = O_p\left(\sqrt{\nu \log n \log\left(\frac{1}{\nu \log n}\right)} + n^{1/2}\nu^2\right).$$

By condition (C.4)-(C.5), we have $\sup_{|\eta|=1} M_1 n^{1/2}U = o_p(1)$, which means $n^{1/2}(\tilde{Q} - Q) = o_p(1)$.

With this result, let's look at the next step:

$$\begin{aligned} & n^{1/2} \left\{ \hat{S}_U(x; \eta) - \tilde{S}_U(x; \eta) \right\} \\ = & n^{-1/2} \frac{\sum_{i=1}^n \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{\hat{H}_{a,i}(x)\hat{H}_{c,i}(x)} I(U_i > x)}{1/n \sum_{i=1}^n \frac{\Delta_{a,i}^\eta(x)\Delta_{c,i}(x)}{\hat{H}_{a,i}(x)\hat{H}_{c,i}(x)}}} - \frac{\sum_{i=1}^n \frac{\tilde{\Delta}_{a,i}^\eta(x)\Delta_{c,i}(x)}{\hat{H}_{a,i}(x)\hat{H}_{c,i}(x)} I(U_i > x)}{1/n \sum_{i=1}^n \frac{\tilde{\Delta}_{a,i}^\eta(x)\Delta_{c,i}(x)}{\hat{H}_{a,i}(x)\hat{H}_{c,i}(x)}}} \\ & \tag{A.9} \\ = & \frac{1}{n^{1/2}\tilde{Q}Q} \left[\tilde{Q} \sum_{i=1}^n \frac{\{\Delta_{a,i}^\eta(x) - \tilde{\Delta}_{a,i}^\eta(x)\}\Delta_{c,i}(x)I(U_i > x)}{H_{a,i}(x)H_{c,i}(x)} \right] \\ & - \frac{1}{\tilde{Q}Q} \left[(\tilde{Q} - Q)n^{-1/2} \sum_{i=1}^n \frac{\tilde{\Delta}_{a,i}^\eta(x)\Delta_{c,i}(x)I(U_i > x)}{H_{a,i}(x)H_{c,i}(x)} \right] + o_p(1). \end{aligned}$$

The second term of (A.9) is $o_p(1)$ because $n^{1/2}(Q - \tilde{Q}) = o_p(1)$ and condition (C.4)-(C.5). As for the first term, we can show it is $o_p(1)$ by using similar arguments proving $n^{1/2}(Q - \tilde{Q}) = o_p(1)$, and we omit it here. Therefore, we have $n^{1/2} \left\{ \hat{S}_U(x; \hat{\eta}_{opt}) - \tilde{S}_U(x; \tilde{\eta}_{opt}) \right\} = o_p(1)$ uniformly for η .

K = 2

When $K = 2$, $\Delta_a^\eta(x)$ and $\tilde{\Delta}_a^\eta(x)$ can be either the indicator (distribution) function or a product of two indicator (distribution) functions depend on the map of QAL. Let's also start from the equivalence between R and Q . Suppose at target time x , there are two group

of patients who are at stage 1 or stage 2

$$\begin{aligned}
& n^{-1/2} \sum_{i=1}^n \frac{\{\Delta_{a,i}^\eta(x) - \tilde{\Delta}_{a,i}^\eta(x)\} \Delta_{c,i}(x)}{\hat{H}_{a,i}(x) \hat{H}_{c,i}(x)} \\
&= n^{-1/2} \sum_{i \in \text{Stage1}} \frac{\{\Delta_{a,i}^\eta(x) - \tilde{\Delta}_{a,i}^\eta(x)\} \Delta_{c,i}(x)}{H_{a,i}(x) H_{c,i}(x)} + \sum_{i \in \text{Stage2}} \frac{\{\Delta_{a,i}^\eta(x) - \tilde{\Delta}_{a,i}^\eta(x)\} \Delta_{c,i}(x)}{H_{a,i}(x) H_{c,i}(x)} + o_p(1) \\
&= n^{-1/2} \sum_{i=1}^n \frac{\{\Delta_{a,i}^\eta(x) - \tilde{\Delta}_{a,i}^\eta(x)\} \Delta_{c,i}(x) I\{L_i(x) = 1\}}{H_{a,i}(x) H_{c,i}(x)} \\
&\quad + \sum_{i=1}^n \frac{\{\Delta_{a,i}^\eta(x) - \tilde{\Delta}_{a,i}^\eta(x)\} \Delta_{c,i}(x) I\{L_i(x) = 2\}}{H_{a,i}(x) H_{c,i}(x)} + o_p(1) \\
&:= \mathcal{K}_1 + \mathcal{K}_2 + o_p(1),
\end{aligned} \tag{A.10}$$

where $L_i(x)$ is the number of stages the i th patient stays at target time x . Notice that $|(\text{A.10})| \leq |\mathcal{K}_1| + |\mathcal{K}_2|$. We can use similar arguments in the case $K = 1$ to bound the term $|\mathcal{K}_1|$ and we omit the details here. Define $L_1 = (A_0, T_0, \delta_0)$, $L_2 = (L_1, Z_0, A_1, T_1, \delta_1, S)$, $L = (L_1, L_2)$, $K_2(q) = \Delta_c(x) I\{K(x) = 2\} / H_a(x) H_c(x)$ and $r_1^\eta = \eta^\top Z_0$, $r_2^\eta = \eta^\top Z_1$. Following the arguments in [Heller \(2007\)](#), we have

$$\begin{aligned}
|\mathcal{K}_2| &\leq M_2 n^{1/2} \\
&\quad \times \sup_{|\eta|=1} \left| \int_{l_1} \int_{r_1^\eta} \int_{l_2} \int_{r_2^\eta} \{I(r_1^\eta \geq 0) I(r_2^\eta \geq 0) - \Phi(r_1^\eta/\nu) \Phi(r_2^\eta/\nu)\} K_2(q) \right. \\
&\quad \left. \times d\hat{F}(r_2^\eta|l_2) d\hat{G}(l_2|l_1) d\hat{F}(r_1^\eta|l_1) d\hat{G}(l_1) \right|
\end{aligned} \tag{A.11}$$

where M_2 is a finite constant, $\hat{G}(l_1)$ and $\hat{F}(r_1^\eta|l_1)$ are the marginal empirical cumulative distribution functions for l_1 and the conditional empirical cumulative distribution function for r_1^η , respectively at Stage 1 for subjects who belong to Stage 2, while $\hat{F}(r_2^\eta|l_2)$ and $\hat{G}(l_2|l_1)$ are the conditional empirical cumulative distribution function for r_2^η and l_2 at Stage 2 for subjects who stay at stage 2.

Write the absolute integral of the right hand side of (A.11) as $U_1 + U_2$ where

$$\begin{aligned}
U_1 &= \int_{q_1} \int_{r_1^\eta} \int_{l_2} \int_{r_2^\eta} \{I(r_1^\eta \geq 0)I(r_2^\eta \geq 0) - I(r_1^\eta \geq 0)\Phi(r_2^\eta/\nu)\} K_2(l) \\
&\quad \times d\hat{F}(r_2^\eta|l_2)d\hat{G}(l_2|l_1)d\hat{F}(r_1^\eta|l_1)d\hat{G}(l_1) \\
&= \int_{l_1} \int_{r_1^\eta} I(r_1^\eta \geq 0) \int_{l_2} \int_{r_2^\eta} \{I(r_2^\eta \geq 0) - \Phi(r_2^\eta/\nu)\} \\
&\quad \times K_2(q)d\hat{F}(r_2^\eta|l_2)d\hat{G}(l_2|l_1)d\hat{F}(r_1^\eta|l_1)d\hat{G}(l_1) \\
&= \int_{l_1} \int_{r_1^\eta} I(r_1^\eta \geq 0)U_3d\hat{F}(r_1^\eta|l_1)d\hat{G}(l_1), \\
U_2 &= \int_{l_1} \int_{r_1^\eta} \int_{l_2} \int_{r_2^\eta} \{I(r_1^\eta \geq 0)\Phi(r_2^\eta/\nu) - \Phi(r_1^\eta/h)\Phi(r_2^\eta/\nu)\} \\
&\quad \times K_2(q)d\hat{F}(r_2^\eta|l_2)d\hat{G}(l_2|l_1)d\hat{F}(r_1^\eta|l_1)d\hat{G}(l_1) \\
&= \int_{l_1} \int_{r_1^\eta} \{I(r_1^\eta \geq 0) - \Phi(r_1^\eta/\nu)\} \int_{l_2} \int_{r_2^\eta} \Phi(r_2^\eta/\nu)K_2(l) \\
&\quad \times d\hat{F}(r_2^\eta|l_2)d\hat{G}(l_2|l_1)d\hat{F}(r_1^\eta|q_1)d\hat{G}(l_1).
\end{aligned}$$

where $U_3 = \int_{l_2} \int_{r_2^\eta} \{I(r_2^\eta \geq 0) - \Phi(r_2^\eta/\nu)\}K_2(q)d\hat{F}(r_2^\eta|l_2)d\hat{G}(l_2|l_1)$. Using the arguments made in the case $K = 1$, Conditions (C.4)-(C.5), and the boundedness of the term $\int_{l_2} \int_{r_2^\eta} \Phi(r_2^\eta/\nu)K_2(l)d\hat{F}(r_2^\eta|l_2)d\hat{G}(l_2|l_1)$, we can show that $n^{1/2}U_2 = o_p(1)$ and $n^{1/2}U_3 = o_p(1)$. Because r_1^η is bounded, $\int_{l_1} \int_{r_1^\eta} I(r_1^\eta \geq 0)U_3d\hat{F}(r_1^\eta|l_1)d\hat{G}(l_1)$ is also $o_p(1)$. We thus have $n^{1/2}(R - Q) = o_p(1)$.

We then do the expansion in equation (A.9). The second term is obviously $o_p(1)$ due to $n^{1/2}(R - Q) = o_p(1)$. The first term can also be shown to be $o_p(1)$ by repeating the above arguments when $K = 2$. With fixed finite number of stages K , we can do similar things to generalize this result.

Above arguments show that $n^{1/2} \left\{ \hat{S}_U(x; \eta) - \tilde{S}_U(x; \eta) \right\} = o_p(1)$ uniformly in η . Then by Taylor expansion, $n^{1/2} \left\{ \tilde{S}_U(x; \eta_{opt}) - \tilde{S}_U(x; \tilde{\eta}_{opt}) \right\} = o_p(1)$ and $n^{1/2} \left\{ \hat{S}_U(x; \eta_{opt}) - \hat{S}_U(x; \hat{\eta}_{opt}) \right\} = o_p(1)$ and thus $n^{1/2} \left\{ \hat{S}_U(x; \hat{\eta}_{opt}) - \tilde{S}_U(x; \tilde{\eta}_{opt}) \right\} = n^{1/2} \left\{ \hat{S}_U(x; \hat{\eta}_{opt}) - \hat{S}_U(x; \eta_{opt}) + \hat{S}_U(x; \eta_{opt}) - \tilde{S}_U(x; \eta_{opt}) + \tilde{S}_U(x; \eta_{opt}) - \tilde{S}_U(x; \tilde{\eta}_{opt}) \right\} = o_p(1)$ and the proof of property 4 in Theorem 1 completes.

□

S1.2 Proof of Theorem 2

Proof. In this proof, we omit the superscript ^{'par'} for simplicity, with the understanding that we are focusing on the parametric modeling of the nuisance functions.

As shown in Theorem 1, $\hat{S}_U(x; \hat{\eta}_{opt})$ is consistent to $S_U(x; \eta_{opt})$ and

$$n^{1/2} \left\{ \hat{S}_U(x; \hat{\eta}_{opt}) - S_U(x; \eta_{opt}) \right\} = n^{-1/2} \sum_{i=1}^n IC_{PAR}(o_i; x, \eta_{opt}) + o_p(1),$$

where $IC_{PAR}(o; x, \eta_{opt})$ is the influence function when target time is x . For the pre-defined restricted upper limit L_U , the RQAL $R(\eta)$ up to L_U is a continuous mapping of $S(x; \eta)$. By applying continuous mapping theorem, $\hat{R}(\hat{\eta}_{opt}, \hat{\theta}, \hat{\beta})$ is consistent to $R(\eta_{opt})$ when the model for treatment assignment and censoring are correctly specified.

We then show the asymptotic normality. We already have the asymptotic lineality expression of (A.2). With delta method, it is easy to show that

$$\begin{aligned} n^{1/2} \left\{ \hat{R}(\hat{\eta}_{opt}, \hat{\theta}, \hat{\beta}) - R(\eta_{opt}) \right\} &= n^{-1/2} \sum_{i=1}^n \int_0^{L_U} IC_{PAR}(o_i; x, \eta_{opt}) dx + o_p(1) \\ &:= n^{-1/2} \sum_{i=1}^n IC_{2,PAR}(o_i; x, \eta_{opt}) + o_p(1), \end{aligned}$$

and $IC_{2,PAR}(o; x, \eta_{opt})$ is the influence function for RQAL at time x under regime g^η .

Thus, $\hat{R}(\hat{\eta}_{opt}, \hat{\theta}, \hat{\beta})$ is asymptotic normal with mean $R(\eta_{opt})$ and variance $E[IC_{2,PAR}(o; x, \eta_{opt})^2]$ which can be empirically estimated as

$$1/n \sum_{i=1}^n \hat{IC}_{2,PAR}(o_i; x, \hat{\eta}_{opt})^2 = 1/n \sum_{i=1}^n \left\{ \int_0^{L_U} \hat{IC}_{PAR}(o_i; x, \hat{\eta}_{opt}) \right\}^2. \quad (\text{A.12})$$

The asymptotic equivalence between $\hat{S}_U(x; \hat{\eta}_{opt})$ and $\tilde{S}_U(x; \tilde{\eta}_{opt})$ can be shown using the arguments for Theorem 1. □

S1.3 Proof of Theorem 3

Proof. In this proof, we omit the superscript ^{'npars'} for simplicity, with the understanding that we are focusing on the nonparametric modeling of the nuisance functions.

We use the following Lemma to prove the theorem. Recall

$$h_{a,j}(g_j^\eta | \bar{z}) = (1 - A_{j-1})p(A_j = g_j^\eta | A_{j-1} = 0, C_{j-1} = 0, Y_j = 1, \bar{Z}_j = \bar{z}) + A_{j-1}$$

$$h_{c,j}(c | \bar{a}, \bar{z}) = (1 - C_{j-1})p(C_j = c | \bar{A}_j = \bar{a}, C_{j-1} = 0, Y_j = 1, \bar{Z}_j = \bar{z}) + C_{j-1}.$$

Lemma S1 (Theoretical undersmoothing conditions). *Let $\hat{h}_{a,j,\lambda_n^a}$ and $\hat{h}_{c,j,\lambda_n^c}$ denote highly adaptive lasso estimators of $h_{a,j}$ and $h_{c,j}$ with L_1 -norm bounds λ_n^a and λ_n^c chosen such that, for $j = 1, 2, \dots, l(x)$,*

$$\min_{k \in \mathcal{J}_n^a} \left\| P_n \frac{d}{d \logit \hat{h}_{a,j,\lambda_n^a}} L(\logit \hat{h}_{a,j,\lambda_n^a})(\phi_k) \right\| = o_p(n^{-1/2}), \quad (\text{S12})$$

$$\min_{k \in \mathcal{J}_n^c} \left\| P_n \frac{d}{d \logit \hat{h}_{c,j,\lambda_n^c}} L(\logit \hat{h}_{c,j,\lambda_n^c})(\phi_k) \right\| = o_p(n^{-1/2}), \quad (\text{S13})$$

where $L(\cdot)$ is log-likelihood loss and ϕ_k denote the k th basis function (i.e., feature) included in the model. Also, \mathcal{J}_n is a set of indices for the basis functions with nonzero coefficients in the corresponding model. Let $D^a(f^a, \hat{h}_{a,j}) = f^a \cdot (A_j - \hat{h}_{a,j})$, and $D^c(f^c, \hat{h}_{c,j}) = f^c \cdot (C_j - \hat{h}_{c,j})$. The functions f^a and f^c are càdlàg with finite sectional variation norm. Let \tilde{f}_ϕ^a and \tilde{f}_ϕ^c be projections of \tilde{f}^a and \tilde{f}^c onto the linear span of the basis functions ϕ_j in $L^2(P)$, where $\phi_{s,j}$ satisfies condition (S12) and (S13), respectively. Then, we have $P_n D^a(f^a, \hat{h}_{a,j}) = o_p(n^{-1/2})$ and $P_n D^c(f^c, \hat{h}_{c,j}) = o_p(n^{-1/2})$.

Proof. The proof is similar to the proof of Lemma 1 in [Ertefaie et al. \(2023\)](#), and thus is omitted. □

We now return to the proof of Theorem 3. We first show the consistency of the direct inverse probability weighting estimator with highly adaptive lasso. Let $\check{\Psi}_{H_{a,c}}(x; \eta) = \frac{\Delta_d^a(x) \Delta_c(x)}{H_{a,c}(x)} \{I(U > x) - S_U(x; \eta)\}$ and $\Psi_{H_a, H_c}(x; \eta) = \mathbb{P}_n \check{\Psi}_{H_{a,c}}(x; \eta)$ where \mathbb{P}_n is the empirical

average and $H_{a,c,i}(x) = H_{a,i}(x)H_{c,i}(x)$. For simplicity of notation, we will suppress η in our notation such that $\check{\Psi}_{H_{a,c}}(x; \eta) \equiv \check{\Psi}_{H_{a,c}}(x)$. We have

$$\mathbb{P}_n \check{\Psi}_{\hat{H}_{a,c}} - \mathbb{P} \check{\Psi}_{H_{a,c}} = (\mathbb{P}_n - \mathbb{P}) \check{\Psi}_{H_{a,c}} + \mathbb{P}(\check{\Psi}_{\hat{H}_{a,c}} - \check{\Psi}_{H_{a,c}}) + (\mathbb{P}_n - \mathbb{P})(\check{\Psi}_{\hat{H}_{a,c}} - \check{\Psi}_{H_{a,c}}). \quad (\text{A.13})$$

Using empirical process theory and by consistency of the highly adaptive lasso estimate $\check{\Psi}_{\hat{H}_{a,c}}$, the third term will be $o_p(n^{-1/2})$ (Vaart and Wellner, 1996; Kosorok, 2008). The first term is already linear so it will contribute to the influence function. Hence, the asymptotic linearity for our estimator relies on the linearity of the second term. It can be written as

$$\begin{aligned} \mathbb{P}_0(\check{\Psi}_{\hat{H}_{a,c}} - \check{\Psi}_{H_{a,c}}) &= \mathbb{P}_0 \left\{ \frac{\Delta_a^\eta(x) \Delta_c(x) I(U > x)}{\hat{H}_{a,c}(x)} - \frac{\Delta_a^\eta(x) \Delta_c(x) I(U > x)}{H_{a,c}(x)} \right\} \\ &= \mathbb{P}_0 \left(Q(x) \left[-\frac{\hat{H}_{a,c}(x) - H_{a,c}(x)}{H_{a,c}(x)} + \frac{(\hat{H}_{a,c}(x) - H_{a,c}(x))^2}{\hat{H}_{a,c}(x)} \right] \right) \\ &= \mathbb{P}_0 \sum_{j=0}^{l(x)} \left[\frac{Q_j(x) I(\bar{A}_{j-1} = \bar{g}_{j-1}^\eta) I(C_{j-1} = 0)}{\prod_{s \leq j} \{1 - h_{c,s}(1 | \bar{z}_s, \bar{a}_s)\} \prod_{s \leq j} \{h_{a,s}(g_j^\eta | \bar{z}_s, \bar{a}_s)\}} \right. \\ &\quad \left. \left\{ I(A_j = g_j^\eta) I(C_j = 0) - \hat{h}_{a,c,j} \right\} \right] + o_p(n^{1/2}), \end{aligned} \quad (\text{A.14})$$

where $Q_j(x) = E\{I(U > x) | \bar{A}_j = \bar{g}_j^\eta, C_{j-1} = 0, \bar{Z}_j\}$ and $h_{a,c,j} = h_{a,j}(g_j^\eta | \bar{z}) h_{c,j}(0 | \bar{a}, \bar{z})$.

Recall our definitions,

$$h_{a,j}(g_j^\eta | \bar{z}_j) = (1 - A_{j-1})P(A_j = g_j^\eta | A_{j-1} = 0, C_{j-1} = 0, Y_j = 1, \bar{Z}_j = \bar{z}) + A_{j-1},$$

$$h_{c,j}(c_j | \bar{a}_j, \bar{z}_j) = (1 - C_{j-1})P(C_j = c_j | \bar{A}_j = \bar{a}_j, C_{j-1} = 0, Y_j = 1, \bar{Z}_j = \bar{z}_j) + C_{j-1}.$$

The third equality in (A.14) follows from (i) the projection of the terms in the second equality onto the nuisance tangent space at each time point; and (ii) the rate of convergence of the highly adaptive lasso estimate (i.e., $\|\hat{H}_{a,c}(x) - H_{a,c}(x)\|_2 = o_p(n^{-1/4})$). For simplicity of presentation, let

$$D_{CAR}(Q, h_{a,c}) = \sum_{j=0}^{l(x)} \left[\frac{Q_j(x) I(\bar{A}_{j-1} = \bar{g}_{j-1}^\eta) I(C_{j-1} = 0)}{\prod_{s \leq j} \{1 - h_{c,s}(1 | \bar{z}_s, \bar{a}_s)\} \prod_{s \leq j} \{h_{a,s}(g_j^\eta | \bar{z}_s, \bar{a}_s)\}} \left\{ I(A_j = g_j^\eta) I(C_j = 0) - \hat{h}_{a,c,j} \right\} \right]$$

We first rewrite

$$I(A_j = g_j^\eta)I(C_j = 0) - \hat{h}_{a,c,j} = I(A_j = g_j^\eta)\{I(C_j = 0) - \hat{h}_{c,j}(0 | \bar{a}, \bar{z})\} + \hat{h}_{c,j}(0 | \bar{a}, \bar{z})\{I(A_j = g_j^\eta) - \hat{h}_{a,j}(g_j^\eta | \bar{z})\}.$$

Then define

$$f_j^a = \frac{Q_j(x)I(\bar{A}_{j-1} = \bar{g}_{j-1}^\eta)I(C_{j-1} = 0)\hat{h}_{c,j}(0 | \bar{z}_s, \bar{a}_s)}{\prod_{s \leq j} \{1 - h_{c,s}(1 | \bar{z}_s, \bar{a}_s)\} \prod_{s \leq j} \{h_{a,s}(g_j^\eta | \bar{z}_s, \bar{a}_s)\}}$$

$$f_j^c = \frac{Q_j(x)I(\bar{A}_{j-1} = \bar{g}_{j-1}^\eta)I(C_{j-1} = 0)I(A_j = g_j^\eta)}{\prod_{s \leq j} \{1 - h_{c,s}(1 | \bar{z}_s, \bar{a}_s)\} \prod_{s \leq j} \{h_{a,s}(g_j^\eta | \bar{z}_s, \bar{a}_s)\}}.$$

Accordingly,

$$\mathbb{P}_0(\check{\Psi}_{\hat{H}_{a,c}} - \check{\Psi}_{H_{a,c}}) = -(\mathbb{P}_n - \mathbb{P}_0)D_{CAR}(Q, h_{a,c}) + \mathbb{P}_n \sum_{j=0}^{l(x)} f_j^a \left\{ I(A_j = g_j^\eta) - \hat{h}_{a,j}(g_j^\eta | \bar{z}) \right\}$$

$$+ \mathbb{P}_n \sum_{j=0}^{l(x)} f_j^c \left\{ I(C_j = 0) - \hat{h}_{c,j}(0 | \bar{a}_j, \bar{z}_j) \right\} + o_p(n^{1/2}).$$

Gathering all the terms together, we have

$$\mathbb{P}_n \check{\Psi}_{\hat{H}_{a,c}} - \mathbb{P} \check{\Psi}_{H_{a,c}} = (\mathbb{P}_n - \mathbb{P}) \check{\Psi}_{H_{a,c}} - (\mathbb{P}_n - \mathbb{P}_0)D_{CAR}(Q, h_{a,c}) + \mathbb{P}_n \sum_{j=0}^{l(x)} f_j^a \left\{ I(A_j = g_j^\eta) - \hat{h}_{a,j}(g_j^\eta | \bar{z}) \right\}$$

$$+ \mathbb{P}_n \sum_{j=0}^{l(x)} f_j^c \left\{ I(C_j = 0) - \hat{h}_{c,j}(0 | \bar{a}_j, \bar{z}_j) \right\} + o_p(n^{1/2}).$$

(A.15)

Hence, if we undersmooth $\hat{h}_{c,j}(1 | \bar{a}_j, \bar{z}_j)$ and $\hat{h}_{a,j}(1 | \bar{z}_j)$, for each $j = 1, 2, \dots, l(x)$, such that $\left| \mathbb{P}_n \sum_{j=0}^{l(x)} f_j^a \left\{ I(A_j = g_j^\eta) - \hat{h}_{a,j}(g_j^\eta | \bar{z}) \right\} \right| = o_p(n^{-1/2})$ and $\left| \mathbb{P}_n \sum_{j=0}^{l(x)} f_j^c \left\{ I(C_j = 0) - \hat{h}_{c,j}(0 | \bar{a}_j, \bar{z}_j) \right\} \right| = o_p(n^{-1/2})$, the estimator $\mathbb{P}_n \check{\Psi}_{\hat{H}_{a,c}}$ will be asymptotically linear. Subsequently, we have

$$\hat{S}_U(x; \eta) - S_U(x; \eta) = (\mathbb{P}_n - \mathbb{P}) \check{\Psi}_{H_{a,c}} - (\mathbb{P}_n - P_0)D_{CAR}(Q, h_{a,c}) + o_p(n^{1/2}). \quad (\text{S14})$$

Denote $IC_{HAL}(O; x, \eta) = (\mathbb{P}_n - \mathbb{P}) \check{\Psi}_{H_{a,c}} - (\mathbb{P}_n - P_0) \left[Q(x) \frac{\{\Delta_a^\eta(x)\Delta_c(x) - H_{a,c}(x)\}}{H_{a,c}(x)} \right]$. Then, the

RQAL with highly adaptive lasso will be asymptotic linear with

$$n^{1/2} \left\{ \hat{R}(\hat{\eta}_{opt}) - R(\eta_{opt}) \right\} = n^{-1/2} \sum_{i=1}^n \int_0^{L_U} IC_{HAL}(o_i; x, \eta_{opt}) dx + o_p(1). \quad (\text{A.15})$$

Similar arguments in the proof of Theorem 1 can be used to show the asymptotic equivalence between $\hat{S}_U(x; \hat{\eta}_{opt})$ and $\tilde{S}_U(x; \tilde{\eta}_{opt})$.

□

S1.4 The Highly Adaptive Lasso

The basic idea is to model an outcome on a d_j -dimensional covariate vector X by mapping X into a group of indicator basis functions. Let $s \subset (1, \dots, d)$, $\mathcal{T} = \{s : s \subset (1, \dots, d)\}$, and denote $X_i^*(s) = \{X_{il} : l \in s\}$, $i = 1, \dots, n$. For a typical observation x , we generate the basis expansion as $\Phi_n(x) = \{\phi_1(x), \dots, \phi_n(x)\}^\top$, where $\phi_i(x) = \{\phi_{s,i}(x) : s \in \mathcal{T}\}$ and $\phi_{s,i}(x) = \{x \rightarrow I(x^*(s) \geq X_i^*(s))\}$. For example when $X = (X_1, X_2)$, $\phi_i(x) = \{\phi_{1,i}(x), \phi_{2,i}(x), \phi_{12,i}(x)\} = \{I(x_1 \geq X_{i1}), I(x_2 \geq X_{i2}), I(x_1 \geq X_{i1}, x_2 \geq X_{i2})\}$, $i = 1, \dots, n$, is the collection of the first- and the second-order indicator basis functions.

We now discuss the highly adaptive lasso fit of the treatment assignment model. Let $h_{a,j} \in \mathcal{D}_j[0, \tau]$, $j = 1, \dots, K$, where $\mathcal{D}_j[0, \tau]$ be the Banach space of d_j -variate real-valued càdlàg (right-continuous with left-hand limits) functions on a cube $[0, \tau] \in \mathbb{R}^{d_j}$, where τ is the upper bound of all supports and is assumed to be finite. For a given subset $s \subset \{0, 1, \dots, d_j\}$, define $u_s = (u_l : l \in s)$ and u_{-s} as the complement of u_s . Then, $h_{a,j,s} : [0_s, \tau_s] \rightarrow \mathbb{R}$ defined as $h_{a,j,s}(u_s) = h_{a,j,s}(u_s, 0_{-s})$. Thus, $h_{a,j,s}(u_s)$ is a section of function $h_{a,j}$ that sets the components in the complement of s equal to zero and varies only along the variables in u_s .

Following Gill et al. (1995), logit $h_{a,j}$ can be represented as

$$\begin{aligned} \text{logit } h_{a,j}(\bar{z}_j) &:= \text{logit } h_{a,j}(0) + \sum_{s \subset \{1, \dots, d_j\}} \int_{0_s}^{\bar{z}_{j,s}} d\text{logit } h_{a,j,s}(u_s) \\ &= \text{logit } h_{a,j}(0) + \sum_{s \subset \{1, \dots, d_j\}} \int_{0_s}^{\tau_s} I(u_s \leq \bar{z}_{j,s}) d\text{logit } h_{a,j,s}(u_s). \end{aligned} \quad (\text{S15})$$

Let $\phi_{s,i}(c_s) = I(\bar{z}_{j,s,i} \leq c_s)$ where $\bar{z}_{j,s,i}$ are support points of $dh_{a,j,s}$. The representation

(S15) can be approximated $\text{logit } h_{a,j,\beta} = \beta_{j,0}^a + \sum_{s \subset \{1, \dots, d_j\}} \sum_{i=1}^n \beta_{j,s,i}^a \phi_{s,i}$, where $|\beta_{j,0}^a| +$

$\sum_{s \in \{1, \dots, d_j\}} \sum_{i=1}^n |\beta_{j,s,i}^a|$ is an approximation of the sectional variation norm of logit $h_{a,j}$.

The highly adaptive lasso defines a minimum loss based estimator $\hat{\beta}_j^a$ as

$$\hat{\beta}_j^a = \arg \min_{\beta: |\beta_{j,0}^a| + \sum_{s \in \mathcal{T}} \sum_{i=1}^n |\beta_{j,s,i}^a| < \lambda} \mathbb{P}_n L(\text{logit } h_{a,j,\beta}),$$

where $L(\cdot)$ is the negative log-likelihood loss function and \mathbb{P}_n is the empirical average.

We denote the estimated treatment assignment model at time j as $\hat{h}_{a,j,\hat{\beta}^a} \equiv \hat{h}_{a,j}$. The

sectional variation norm λ (i.e., l_1 -norm of the coefficient vector) is unknown and must be

estimated using the observed data. The finite sectional variation norm assumption implies

that there is a positive constant $0 < M < \infty$ such that $0 < \lambda < M$. In practice, λ is

determined using cross-validation which is justified theoretically by the oracle properties

of cross-validation ([van der Laan and Dudoit, 2003](#); [Van der Vaart et al., 2006](#)). Similarly,

we define a minimum loss based estimator $\hat{\beta}_j^c$ as

$$\hat{\beta}_j^c = \arg \min_{\beta: |\beta_{j,0}^c| + \sum_{s \in \mathcal{T}} \sum_{i=1}^n |\beta_{j,s,i}^c| < \lambda} \mathbb{P}_n L(\text{logit } h_{c,j,\beta}).$$

We denote the estimated discrete hazard function for censoring at time j as $\hat{h}_{c,j,x,\hat{\beta}^c} \equiv \hat{h}_{c,j,x}$.

S1.5 Plots for Simulation Section

The first row of these figures shows the distribution of true RQAL of each estimated optimal

regime compared to the true RQAL of the true optimal regime (the dashed line). The

second row shows the estimated RQAL of the estimated optimal regime. The third row

compares the treatment regime recommendation between the true optimal regime and the

estimated optimal regime by computing the mis-classification rate (MR).




Figure S3: The simulation results of the estimated RQAL in scenario 1 with $K = 6$. IPW is the estimator (6) while BC-IPW is the bias-corrected estimator (8). The number in brackets is the sample size.




Figure S4: The simulation results of the estimated RQAL in scenario 1 with $K = 25$. IPW is the estimator (6) while BC-IPW is the bias-corrected estimator (8). The number in brackets is the sample size.




Figure S5: The simulation results of the estimated RQAL in scenario 2 with $K = 6$. IPW is the estimator (6) while BC-IPW is the bias-corrected estimator (8). The number in brackets is the sample size.




Figure S6: The simulation results of the estimated RQAL in scenario 2 with $K = 25$. IPW is the estimator (6) while BC-IPW is the bias-corrected estimator (8). The number in brackets is the sample size.

S1.6 Simulation Results for Scenario 1

In the first set of simulation, we set $L = 60$ and $G = 10$. The mean observed survival time is around 32.4 weeks; the mean quality-adjusted lifetime is around 18.4 weeks. In this setting, $\sim 66\%$ patients initialize the treatment in the follow-up, and $\sim 29\%$ of patients

follow the optimal regime $g^{\eta_{opt}}$ exactly. We choose $L_U = 26$ weeks as the upper bound of restricted quality-adjusted lifetime for identifiability which lies in the 90% quantile of the observed quality-adjusted lifetime and the theoretical $R(\eta_{opt}) = 21.04$ weeks in this case.

In the second set of simulation, we set $L = 100$ and $G = 4$. The mean observed survival time is around 47 weeks; the mean quality-adjusted lifetime is around 25.2 weeks. In this setting, $\sim 55\%$ patients initialize the treatment in the follow-up, and $\sim 20\%$ of patients follow the optimal regime $g^{\eta_{opt}}$ exactly. We choose $L_U = 36$ weeks as the upper bound of restricted quality-adjusted lifetime which lies in the 90% quantile of the observed quality-adjusted lifetime and the true $R(\eta_{opt}) = 31.74$ weeks in this case.

Remark 3. *To simplify the presentation of different estimators in the tables, we use the notation \check{R} and $\check{\eta}$ to represent the estimated quantities corresponding to the methods specified in each row. For example, in Table S3, $\check{R}(\check{\eta}_{opt})$ in row 2 denotes $\hat{R}^{npar}(\hat{\eta}_{opt}^{npar})$, whereas in row 4, it denotes $\tilde{R}^{par}(\tilde{\eta}_{opt}^{par})$.*

Table S3 Specifications: The columns (η_0, η_1, η_2) correspond to the average of estimated parameters used to form the optimal strategy and the numbers in parenthesis are the empirical standard errors. IPW is the direct estimation using (6), BC-IPW is the bias-correction estimator (8); Logit, logistic regression, HAL, undersmoothed highly adaptive lasso, RF, random forest; K is the number of stages; n is the sample size; η_0, η_1, η_2 are the empirical average of $\check{\eta}_0/\check{\eta}_1, \check{\eta}_1/\check{\eta}_2, \check{\eta}_2/\check{\eta}_0$; $\check{R}(\check{\eta}_{opt})$ is the empirical average of estimated RQAL of the estimated optimal regime. MR is the empirical misclassification rate of the estimated optimal regime.

The misclassification rates of both highly adaptive lasso and logistic regression are also very close. Among the methods considered, the random forest demonstrates the poorest performance in both bias and empirical standard errors. Notably, the conservative standard error presented at the end of Section 4.3 underestimates the empirical standard errors of

$\hat{R}(\hat{\eta}_{opt})$, as indicated in parentheses, resulting in significant undercoverage (Figure 1). The misclassification rates are also substantially higher than the other estimators.

Figures S3 and S4 in Section S1.5 provide additional information about the estimators for $K = 6$ and $K = 25$, respectively. Specifically, it provides insight about the distribution of (a) the true restricted quality-adjusted lifetime under different estimated optimal regimes (i.e., $R(\hat{\eta}_{opt})$ and $R(\tilde{\eta}_{opt})$); (b) the estimated restricted quality-adjusted lifetime under the estimated optimal regime (i.e., $\hat{R}(\hat{\eta}_{opt})$ and $\tilde{R}(\tilde{\eta}_{opt})$); and, (c) the misclassification rates. Overall the plots show the superiority of the correctly specified logistic regression and the highly adaptive lasso compared with random forest. Also, there is no evidence of deviating from normal distribution for $\hat{R}(\hat{\eta}_{opt})$, $\tilde{R}(\tilde{\eta}_{opt})$, $R(\hat{\eta}_{opt})$ and $R(\tilde{\eta}_{opt})$ which confirms our theoretical results in Theorems 2 and 3.

S1.7 Additional Simulation Results for Undersmoothed HAL

In this section, we show the necessity of using undersmoothed HAL when modeling nuisance parameters by comparing with cross-validation (CV)-HAL.

S1.7.1 One Stage Study

In this example, we consider a one stage study with Gaussian outcome, and the purpose is to estimate $Y(1)$ which is 0.

- *Outcome model:* $Y = \alpha_1 Z_1 + \alpha_2 Z_2 + \epsilon$.
- *Treatment assignment model:* $\mathbb{P}(A = 1 \mid Z_1, Z_2) = \exp(-\beta_1 Z_1 - \beta_2 Z_2)^{-1}$.

We generate the data as follows:

- (1) Generate $Z_1, Z_2 \sim N(0, 0.5)$, and the error term $\epsilon \sim N(0, 0.5)$.
- (2) Generate the treatment variable A by the treatment assignment model.

Table S3: Summary table of simulation studies in Scenario 1. The target parameter here is $R(\eta_{opt}) = 21.04$ when $K = 6$ and $R(\eta_{opt}) = 31.74$ when $K = 25$; SE is the standard error estimation; CP is the average coverage probability; $R(\hat{\eta}_{opt})$ is the average of true RQAL of the estimated optimal regime. The numbers in brackets are the empirical standard deviation.

	K	n	η_0	η_1	η_2	$\hat{R}(\hat{\eta}_{opt})$	SE	CP	$R(\hat{\eta}_{opt})$	MR	
IPW	6	250	Logit	1.03(0.16)	1.03(0.28)	1.00(0.13)	21.30(0.55)	0.55	0.90	20.74(0.25)	9.12(8.26)
			HAL	1.02(0.16)	1.05(0.32)	0.99(0.14)	21.29(0.58)	0.55	0.90	20.70(0.29)	10.42(9.35)
			RF	1.04(0.30)	1.03(0.51)	1.01(0.24)	21.67(0.75)	0.67	0.80	20.55(0.49)	14.88(15.15)
			Logit	1.02(0.17)	1.06(0.36)	0.99(0.15)	21.14(0.55)	0.55	0.94	20.73(0.27)	9.31(8.68)
			HAL	1.02(0.19)	1.06(0.36)	1.00(0.16)	20.93(0.52)	0.56	0.97	20.70(0.29)	10.46(9.54)
			RF	1.03(0.37)	1.09(0.67)	1.03(0.39)	21.49(0.75)	0.75	0.84	20.55(0.49)	15.71(15.45)
BC-IPW	6	500	Logit	1.00(0.08)	1.02(0.15)	0.99(0.07)	21.17(0.39)	0.39	0.92	20.87(0.14)	4.87(4.69)
			HAL	0.99(0.07)	1.03(0.16)	0.99(0.08)	21.17(0.41)	0.39	0.92	20.85(0.16)	5.39(5.15)
			RF	1.01(0.15)	1.05(0.30)	0.99(0.13)	21.44(0.54)	0.50	0.86	20.76(0.28)	8.42(9.15)
			Logit	1.01(0.08)	1.02(0.17)	1.00(0.08)	21.06(0.39)	0.39	0.95	20.86(0.15)	5.02(5.01)
			HAL	1.01(0.09)	1.01(0.16)	1.00(0.08)	20.92(0.37)	0.40	0.95	20.84(0.15)	5.76(4.86)
			RF	1.01(0.17)	1.07(0.36)	0.99(0.15)	21.31(0.54)	0.51	0.89	20.78(0.28)	8.48(9.15)
IPW	25	250	Logit	1.05(0.29)	1.06(0.45)	1.00(0.19)	32.40(0.90)	0.86	0.82	30.33(1.47)	28.66(21.65)
			HAL	1.05(0.37)	1.10(0.65)	0.99(0.27)	32.37(0.83)	0.88	0.85	30.14(1.39)	33.09(24.18)
			RF	1.10(0.53)	1.18(1.34)	1.02(0.30)	33.74(0.84)	0.72	0.28	29.95(1.74)	35.66(25.54)
			Logit	1.06(0.34)	1.08(0.55)	1.00(0.22)	32.11(0.88)	0.88	0.87	30.34(1.20)	29.62(22.35)
			HAL	1.07(0.44)	1.06(0.58)	1.01(0.27)	31.59(0.86)	0.92	0.95	30.26(1.06)	32.53(22.61)
			RF	1.01(0.84)	1.02(1.76)	1.01(0.39)	33.46(0.90)	0.77	0.39	29.94(1.51)	37.70(26.13)
BC-IPW	25	500	Logit	1.01(0.10)	1.01(0.19)	1.00(0.09)	32.05(0.68)	0.71	0.89	31.01(0.60)	15.92(13.85)
			HAL	1.02(0.14)	1.02(0.23)	1.00(0.12)	32.07(0.71)	0.69	0.86	30.89(0.75)	18.44(16.90)
			RF	1.06(0.30)	1.00(0.39)	1.03(0.19)	33.25(0.77)	0.74	0.44	30.62(0.92)	24.63(20.23)
			Logit	1.01(0.12)	1.02(0.23)	1.00(0.11)	31.83(0.68)	0.72	0.94	30.99(0.65)	16.37(15.01)
			HAL	1.01(0.11)	1.02(0.21)	1.00(0.10)	31.47(0.71)	0.73	0.94	30.93(0.66)	17.64(15.11)
			RF	1.06(0.30)	1.03(0.44)	1.02(0.21)	32.98(0.79)	0.77	0.57	30.65(0.93)	25.02(20.67)

- (3) Generate the outcome variable Y by the outcome model.
- (4) Set the observed covariates as $X_1 = \exp(Z_1/2)$ and $X_2 = (Z_1 + Z_2)^2$.
- (5) The observed data is (X_1, X_2, Y, A) .

Here, $\alpha = (\alpha_1, \alpha_2) = (1, 1)$ and $\beta = (\beta_1, \beta_2) = (-1.5, 1.5)$. The PS model would be mis-specified if we model treatment assignment by logistic regression and the observed covariates. In Table S4, we show the results of IPW estimators by using CV-HAL and undersmooth HAL for the PS model with sample size $n = 500$. The standard error estimation for logistic regression is calculated by sandwich formula.

S1.7.2 Multi Stage Study

In this example, we consider a $K = 6$ stage study which is similar to the example in Section 5 and the outcome of interest is RQAL $R(\eta_{opt})$. The optimal treatment regime is $\eta_{opt} = (1, -1, -1)$. We ignore the censoring part and the estimation of η_{opt} for simplicity, and assume the optimal treatment regime is known. The purpose is to evaluate the performance of the IPW estimator (6) under CV-HAL and undersmooth HAL for the PS model. We still consider the case when PS model is mis-specified by logistic regression with observed covariates. The covariates $Z_j = (Z_{j1}, Z_{j2}, Z_{j3})$ are generated as described in Section 5.

- *Treatment assignment model:* $h_{a,j}(\bar{z}) = \text{logit}^{-1}(\kappa_0 + \kappa_1 \times z_{1j} + \kappa_2 \times z_{2j})$

where $(\kappa_0, \kappa_1, \kappa_2) = (-0.1 \times K, -1, -1)$ and the observed covariates are $X_j = \{X_{j1}, X_{j2}, X_{j3}\} = \{\exp(Z_{j1}/2), (Z_{j1} + Z_{j2})^2, Z_{j3}\}$. We generate the survival time with the following discrete hazard model

$$\begin{aligned}
 h_y(t, \bar{a}, \bar{z}) &= \text{pr}(T = t | T > t - 1, \bar{A}_j = \bar{a}_j, \bar{C}_j = 0, \bar{Z}_j = \bar{z}) \\
 &= \text{logit}^{-1}\{-5 + t \times (6 \times 0.75)/K - 0.5 \times z_{1j} - 0.5 \times z_{2j} - 0.5 \times a_j \times W_j\} \\
 &\text{for } t \in (jG, (j+1)G], \quad j = 0, \dots, K - 1.
 \end{aligned}$$

We then define the utility map as $\mathcal{Q}(Z_{3j}) = Z_{3j} / \max_{k=0, \dots, K} Z_{3k}$ which is always a number between $(0, 1]$. Patients' quality of life scores will reduce to zero if they are no longer at-risk. The upper limit for RQAL is selected as $L_U = 25$ by picking the 90% quantile of observed QAL and $R(\eta_{opt}) = 20.73$ in this case. We use $n = 500$ in this example. See Table S4 of the results.

Table S4: Summary table to compare CV-HAL and undersmoothed HAL. The results are averaged over 500 Monte Carlo data sets. SD is the empirical standard error, SE is the averaged standard error estimation, CP is the averaged coverage probability, and λ is the averaged tuning parameter in l_1 regression.

Example	Method	Bias	SD	SE	CP (%)	λ
One-stage	Logit	0.171	0.065	0.066	11	NA
	CV-HAL	0.085	0.048	0.055	69	0.025
	Under-HAL	0.006	0.054	0.054	94	0.001
Multi-stage	Logit	0.388	0.365	0.361	84	NA
	CV-HAL	0.245	0.365	0.350	85	0.024
	Under-HAL	0.021	0.387	0.364	94	0.007

S2 Additional Tables for the real data example

Table S5 shows the estimated coefficients of the measured variables on the probability of PEG tube insertion and censoring on logit scale. Based on these results, the ALS bulbar and FVC are the most significant variables in deciding for PEG placement such that patients with higher values of bulbar and FVC are less likely to receive PEG. Also, the stage (i.e., number of months since the beginning of the follow-up), site of onset, Ceftriaxone, and ALS total score contribute significantly in hazard of censoring.

Table S5: Summary table of logistic regression for PEG insertion and censoring. Stage represents the number of months since the beginning of the follow-up. Stage² and Stage³ and the square and cubic of Stage.

	PEG insertion			Censoring		
	Est.	S.E.	p-val	Est.	S.E.	p-val
(Intercept)	1.093	1.010	0.279	-5.827	0.907	<0.001
Stage	-0.054	0.098	0.584	0.849	0.131	<0.001
Stage ²	-0.002	0.009	0.781	-0.048	0.009	<0.001
Stage ³	0.000	0.000	0.584	0.001	0.000	<0.001
Sex	-0.099	0.187	0.595	0.048	0.138	0.727
Age	0.013	0.009	0.131	-0.000	0.006	0.984
Race	-0.396	0.404	0.327	0.097	0.265	0.715
Siteofonset	0.480	0.244	0.049	0.576	0.198	0.004
Duration	-0.007	0.014	0.584	0.012	0.009	0.189
Ceftriaxone	-0.340	0.172	0.048	-0.292	0.124	0.019
Riluzole	0.327	0.201	0.105	-0.266	0.136	0.051
BMI	0.002	0.016	0.888	-0.011	0.012	0.361
ALS total	-0.03	0.042	0.446	-0.074	0.029	0.010
ALS bulbar	-0.291	0.061	<0.001	0.065	0.041	0.118
ALS breathing	-0.029	0.059	0.627	0.042	0.041	0.303
ALS gross motor	-0.040	0.070	0.56	-0.023	0.046	0.627
FVC	-0.016	0.004	<0.001	0.004	0.003	0.156
BMI change	-0.012	0.010	0.221	-0.000	0.005	0.967
Grip	-0.004	0.006	0.460	0.003	0.004	0.519
ALSQOL	0.049	0.067	0.465	-0.03	0.046	0.485