High-dimensional classification problems with Barron regular boundaries under margin conditions

Jonathan García * Philipp Petersen [†]

December 11, 2024

Abstract

We prove that a classifier with a Barron-regular decision boundary can be approximated with a rate of high polynomial degree by ReLU neural networks with three hidden layers when a margin condition is assumed. In particular, for strong margin conditions, high-dimensional discontinuous classifiers can be approximated with a rate that is typically only achievable when approximating a low-dimensional smooth function. We demonstrate how these expression rate bounds imply fast-rate learning bounds that are close to n^{-1} where n is the number of samples. In addition, we carry out comprehensive numerical experimentation on binary classification problems with various margins. We study three different dimensions, with the highest dimensional problem corresponding to images from the MNIST data set.

Keywords: Neural Networks, Binary classification, Fourier-analytic Barron space, Barron regular boundaries, Margin condition, Hinge loss. Mathematics Subject Classification: 68T05, 62C20, 41A25, 41A46

1 Introduction

Binary classification is one of the oldest problems in machine learning, and it provides a foundation for understanding multiclass classification [5, 11, 15]. This problem can be reduced to interpreting whether an element \boldsymbol{x} is inside a set Ω or not, and this can be represented mathematically by an indicator function $\mathbb{1}_{\Omega}(\boldsymbol{x})$. Moreover, we define Ω as

$$\Omega := \left\{ \boldsymbol{x} \in [0,1]^d : \eta(\boldsymbol{x}) \ge 1/2 \right\} \quad \text{where} \quad d \in \mathbb{N}_{\ge 2}, \quad \eta(\boldsymbol{x}) := \mu \left(Y = 1 | X = \boldsymbol{x} \right) \tag{1}$$

is the posterior probability, and denote the decision boundary of Ω as $\partial \Omega$. Our goal is to approximate $\mathbb{1}_{\Omega}$ using neural networks, which is a complex problem since $\mathbb{1}_{\Omega}$ is a

^{*}Faculty of Mathematics, University of Vienna, Austria.

jonathan.garcia.rebellon@univie.ac.at

 $^{^\}dagger {\rm Faculty}$ of Mathematics and Research Network Data Science @ Uni Vienna, University of Vienna, Austria.

philipp.petersen@univie.ac.at

discontinuous function, can be very high dimensional, and it is known that approximation of high dimensional discontinuous functions is a serious problem in general [7]. It has also been shown that under classical smoothness assumptions on the decision boundaries $\partial \Omega$ the classification problem admits a curse of dimension [17].

Probably, the most common classification problem is image classification. In this case the dimension depends on the resolution of the image, i.e. the number of pixels used to create it. Typically this leads to a very high dimensional problem, some well known sample sets to train classification models are for example MNIST with images of 28×28 pixels or ImageNet with images of 256×256 pixels.

A classification method that uses a notion of distance to build the decision functions is expected to learn more easily in regions that are not close to the decision boundary. This suggest that learning should be relatively simple for distributions that do not have a lot of mass in the vicinity of the decision boundary. In order to control the mass in the vicinity of the decision boundary the following margin condition (M) is defined [10]. Let μ be a Borel probability measure on $[0, 1]^d$ and \mathcal{D} a joint distribution induced by μ of an i.i.d. sample set

$$S = \{(\boldsymbol{x}_i, y_i)\}_{i=1}^n, \text{ where } (\boldsymbol{x}_i, y_i) \in [0, 1]^d \times \{0, 1\} \text{ and } \mathbb{1}_{\Omega}(\boldsymbol{x}_i) = y_i,$$
 (2)

for all $i \in \{1, ..., n\}$. Then, the margin condition is as follows.

(M) There exist $C, \gamma > 0$, such that for all $\epsilon > 0$,

$$\mu\left(B_{\epsilon}^{*}\right) \leq C\epsilon^{\gamma} \quad \text{where} \quad B_{\epsilon}^{*} := \left\{ \boldsymbol{x} \in [0,1]^{d} : \text{dist}\left(\boldsymbol{x},\partial\Omega\right) \leq \epsilon \right\}$$

is the ball of radius $\epsilon > 0$ around $\partial \Omega$ with respect to Euclidean distance

dist
$$(\boldsymbol{x}, \partial \Omega) := \inf_{\boldsymbol{x'} \in \partial \Omega} \|\boldsymbol{x} - \boldsymbol{x'}\|_2$$

and γ is called the margin exponent.

Furthermore, in [8] it was also shown that by considering the class of measures μ satisfying the following condition, rates independent of the input dimension and appearing in the form of a polynomial factor can be obtained. It is said that μ is tube compatible with parameters $\alpha \in (0, 1]$ and C > 0 if for each measurable function $f : [0, 1]^{d-1} \rightarrow [0, 1]$, each $i \in \{1, \ldots, d\}$ and each $\epsilon \in (0, 1]$, it is satisfied that

$$\mu(T_{f,\epsilon}^{(i)}) \le C\epsilon^{\alpha} \quad \text{where} \quad T_{f,\epsilon}^{(i)} := \left\{ \boldsymbol{x} = (x_1, \dots, x_d) \in [0, 1]^d : |x_i - f(\boldsymbol{x}^{(i)})| \le \epsilon \right\}$$
(3)

and $\boldsymbol{x}^{(i)} := (x_1, \ldots, x_{i-1}, x_{i+1}, \ldots, x_d) \in [0, 1]^{d-1}$. The set $T_{f,\epsilon}^{(i)}$ is called a tube of width ϵ associated to f. Lastly, it was shown in [1, 3, 2] that for a class of functions with finite Fourier moment, the curse of dimensionality can be overcome when the approximation error is measured in the L^2 norm. This influenced further research on the approximation in this class of functions (see e.g. [6, 13]) which is nowadays known as the Barron class. Motivated by the qualities of this class of functions, we will use them throughout this paper. To be more precise, we study how complex the classification learning for deep neural networks (NNs) with the rectified linear unit (ReLU) activation function and using the hinge loss (see Section 1.2) can be, when the margin condition (M) and the tube compatibility (3) are met, and $\partial\Omega$ can be described by functions in Barron class (see Subsection 1.3.1). Under these conditions we obtained that:

- (i) Theorems 2 and 3. It is possible to approximate 1_Ω using a ReLU-NN Φ with three hidden layers and a number of neurons O(N) with N ∈ N, having an error of µ ({1_Ω ≠ Φ}) ≲ N^{-γ/2} for all probability measures µ satisfying the margin condition (M) with margin exponent γ, and the tube compatibility (3). Moreover, when Φ is trained using the hinge loss φ (see (10)), we obtain a learning rate (see Subsection 1.2.1 and Remark 4) of the order of O(n^{-γ/(2+γ)}(1+log n)), where N ≲ n^{γ/(2+γ)}, n = |S| and S is a training set as in (2). Implying that when γ is large, we achieve to approximate 1_Ω (high-dimensional discontinuous function) by Φ, at a rate comparable to that of a low-dimensional smooth function, with a fast learning rate of ≈ n⁻¹(1 + log n).
- (ii) Simulations 2.1. We produce numerical simulations within the framework of our main theorems: we generate samples in $d \in \{3, 50, 784\}$ dimensions to simulate the margin condition (M) applied with a margin exponent γ and varying the number of elements n in various ranges. Then, we conclude as γ and n increase, the test error (see Subsection 1.2.1) decreases at a rate close to that stated in our Theorem 3 and Remark 4, which supports our results.

1.1 Background

In [10, Section 8], the features of SVMs when applied to binary classification were investigated. It was shown that for distributions with noise but low concentration (margin condition (M)) near the decision boundary, the approximation error for Gaussian kernels was relatively small, leading to favorable learning rates. A data-dependent strategy for selecting the regularization and kernel parameters was also studied, introducing the noise exponent to measure the amount of high noise in the labeling process. The learning rates achieved were sometimes as fast as n^{-1} , where n is the number of data points.

Next in [16], it was shown that a convergence rate similar to the rates of [10] can be achieved but using NN classifiers. Particularly, it was proved that the estimated classifier based on NNs with the hinge loss achieves similar fast convergence rates considering the following assumptions on the true classifier:

(I) The noise Tsybakov condition is always met [20]. That is, there exist C > 0 and $q \in [0, \infty]$ such that for any $\epsilon > 0$,

$$\mu\left(\left\{\boldsymbol{x}\in[0,1]^d:|2\eta(\boldsymbol{x})-1|<\epsilon\right\}\right)\leq C\epsilon^q.$$

(II) The decision boundary is Hölder smooth, i.e. $\partial\Omega$ can be described by functions in the space

$$\mathcal{H}^{\ell}(\mathcal{X}) = \left\{ f \in \mathcal{C}^{\lceil \ell - 1 \rceil}(\mathcal{X}) : \max_{\|\boldsymbol{m}\| \leq \lceil \ell - 1 \rceil} \|\partial^{\boldsymbol{m}} f\|_{\infty} + \max_{\|\boldsymbol{m}\| = \lceil \ell - 1 \rceil} [\partial^{\boldsymbol{m}} f]_{\ell - \lceil \ell - 1 \rceil} < \infty \right\},$$

where $\ell > 0$, $\mathcal{C}^m(\mathcal{X})$ is the space of $m \in \mathbb{N}$ times differentiable functions on a set \mathcal{X} whose partial derivatives of order m with $||m|| \leq m$ are continuous;

$$\partial^{\boldsymbol{m}} f = rac{\partial^{\|\boldsymbol{m}\|} f(\boldsymbol{x})}{\partial \boldsymbol{x}^{\boldsymbol{m}}} \quad ext{and} \quad [f]_s = \sup_{\substack{\boldsymbol{x}, \boldsymbol{y} \in \mathcal{X} \\ \boldsymbol{x} \neq \boldsymbol{y}}} rac{|f(\boldsymbol{x}) - f(\boldsymbol{y})|}{\|\boldsymbol{x} - \boldsymbol{y}\|^s}.$$

- (III) The case when the conditional class probability is assumed to be Hölder smooth, meaning that $\eta(\cdot) \in \mathcal{H}^{\ell}(\mathcal{X})$, where η is as in (1).
- (IV) When the margin condition (M) is fulfilled, but with margin exponent $\gamma \in [1, \infty]$.

However, when the margin condition is satisfied, the bound shown in [16] for the learning rate is

$$\left(\frac{\log^3 n}{n}\right)^{\frac{\ell(q+1)}{\ell(q+2)+(d-1)(q+1)/\gamma}}, \quad \text{for which} \quad \lim_{\substack{q,\ell,\gamma \text{ are fixed}}} \frac{\ell(q+1)}{\ell(q+2)+(d-1)(q+1)/\gamma} = 0 \quad (4)$$

and the variables d, q, ℓ, γ are defined in items (I), (II), (IV) above, meaning that in some sense bound (4) is affected by the curse of dimensionality and to avoid it, one must require that $d \leq \gamma$, but this is a rather restrictive condition.

So, our results (i) provide a significant improvement over the bound (4) when (M) is satisfied, since they are not affected by the curse of dimensionality and show faster convergence.

1.2 Neural networks

Neural networks are functions formed by connecting neurons, where the output of one neuron becomes the input to another. Here, a neuron is a function of the form

$$\mathbb{R}^d \ni \boldsymbol{x} \to \sigma(\langle \boldsymbol{w}, \boldsymbol{x} \rangle + b)$$

where $\boldsymbol{w} \in \mathbb{R}^d$ is a weight vector, $b \in \mathbb{R}$ is called bias, and the function σ is referred to as an activation function. There are several types of NNs but in this paper we only work with one of the most common ones, which is the so-called feedforward NN, in this structure, neurons are organized in layers, and the neurons in each layer receive input only from the previous layer. We introduce these NNs as in [19], as follows.

Let $L \in \mathbb{N}, d_0, \ldots, d_{L+1} \in \mathbb{N}$ and $\sigma : \mathbb{R} \to \mathbb{R}$ an activation function. Then, we call a function $\mathbf{\Phi} : \mathbb{R}^{d_0} \to \mathbb{R}^{d_{L+1}}$ a neural network if there exist for $\ell \in \{0, \ldots, L\}$, matrices $\mathbf{W}^{(\ell)} \in \mathbb{R}^{d_{\ell+1} \times d_{\ell}}$ and vectors $\mathbf{b}^{(\ell)} \in \mathbb{R}^{d_{\ell+1}}$ such that for all $\mathbf{x} \in \mathbb{R}^{d_0}$,

$$\boldsymbol{x_0} := \boldsymbol{x}, \qquad \boldsymbol{x}^{(\ell)} := \sigma(\boldsymbol{W}^{(\ell-1)} \boldsymbol{x}^{(\ell-1)} + \boldsymbol{b}^{(\ell-1)}) \quad \text{for all} \quad \ell \in \{1, \dots, L\},$$

and

$$\Phi(x) = x^{(L+1)} := W^{(L)}x^{(L)} + b^{(L)}$$

In addition, we define the following parameters

$$\begin{split} L(\boldsymbol{\Phi}) &:= L + 1 & \text{number of layers,} \\ N(\boldsymbol{\Phi}) &:= \sum_{j=0}^{L+1} d_{\ell} & \text{number of neurons,} \\ \Theta(\boldsymbol{\Phi}) &:= \left(\left(\boldsymbol{W}^{(0)}, \boldsymbol{b}^{(0)} \right), \dots, \left(\boldsymbol{W}^{(L)}, \boldsymbol{b}^{(L)} \right) \right) & \text{parameter tuple,} \\ & \| \cdot \|_{0} & \text{number of non-zero entries of } \cdot, \end{split}$$

$$\begin{split} W(\boldsymbol{\Phi}) &:= \sum_{\ell=0}^{L} \left(\|\boldsymbol{W}^{(\ell)}\|_{0} + \|\boldsymbol{b}^{(\ell)}\|_{0} \right) & \text{number of weights,} \\ W_{\infty}(\boldsymbol{\Phi}) &:= \max_{0 \leq \ell \leq L} \left\{ \max_{0 \leq \ell \leq L} \|\boldsymbol{W}^{(\ell)}\|_{\infty}, \max_{0 \leq \ell \leq L} \|\boldsymbol{b}^{(\ell)}\|_{\infty} \right\} & \text{largest absolute value of } \boldsymbol{\Theta}, \\ d_{0} \text{ and } d_{L+1} & \text{input and output dimensions,} \\ (d_{0}, d_{1}, \dots, d_{L+1}) \in \mathbb{N}^{L+2} & \text{architecture,} \\ L & \text{number of hidden layers.} \end{split}$$

Moreover, when $L = d_2 = 1$, we call Φ a shallow neural network, where

$$\Phi(\boldsymbol{x}) = b^{(1)} + \sum_{i=1}^{d_1} w_i^{(1)} \sigma\left(\left\langle \boldsymbol{w}_i^{(0)}, \boldsymbol{x} \right\rangle + b_i^{(0)}\right)$$
(5)

for some $w_i^{(1)}, b_i^{(0)}, b^{(1)} \in \mathbb{R}$ and $\boldsymbol{w}_i^{(0)} \in \mathbb{R}^{d_0}$ for $i = 1, \dots, d_1$.

In the sequel, we use the ReLU activation function which we denote as $\rho : \mathbb{R} \to \mathbb{R}$ and which is defined as $\rho(x) := \max\{0, x\}$. Let $d \in \mathbb{N}_{\geq 2}$, $N, W \in \mathbb{N}$ and B > 0. We denote by $\mathcal{NN} = \mathcal{NN}(d, N, W, B)$ the set of ReLU-NNs with three hidden layers, *d*-dimensional input and 1-dimensional output, with at most N neurons per layer, with at most W nonzero weights, and with these weights bounded in absolute value by B. Furthermore, we define

 $\mathcal{NN}_* = \mathcal{NN}_*(d, N, W, B) := \left\{ f \in \mathcal{NN} : 0 \le f(\boldsymbol{x}) \le 1 \quad \text{for all} \quad \boldsymbol{x} \in [0, 1]^d \right\}.$ (6)

1.2.1 Loss function and the hinge loss

In supervised learning, determining the optimal parameters of a NN is achieved by minimizing an objective function. This objective is defined based on a collection of inputoutput pairs known as a sample. Concretely, let $S = \{(\boldsymbol{x}_i, y_i)\}_{i=1}^n$ be as in (2), the goal is to find a NN $\boldsymbol{\Phi}$ such that $\boldsymbol{\Phi}(\boldsymbol{x}_i) \approx y_i$ for all $i \in \{1, \ldots, n\}$ in a meaningful sense. For this purpose, a loss function \mathcal{L} is defined which measures the dissimilarity between its inputs, and the so-called empirical risk of $\boldsymbol{\Phi}$ with respect to the sample S and \mathcal{L} , defined as

$$\widehat{\mathcal{E}}_{\mathcal{L},S}(\boldsymbol{\Phi}) = \frac{1}{n} \sum_{i=1}^{n} \mathcal{L}(\boldsymbol{\Phi}(\boldsymbol{x}_i), y_i)$$
(7)

is minimized. Next, after optimization, we want to measure the performance of $\boldsymbol{\Phi}$ on new data $(\boldsymbol{x}_{\text{new}}, y_{\text{new}})$ sampled independently from \mathcal{D} , for this we define the so-called risk (or \mathcal{L} -risk) of $\boldsymbol{\Phi}$, in the form

$$\mathcal{E}_{\mathcal{L}}(\boldsymbol{\Phi}) = \mathbb{E}_{(\boldsymbol{x}_{\text{new}}, y_{\text{new}}) \sim \mathcal{D}} \left[\mathcal{L}(\boldsymbol{\Phi}(\boldsymbol{x}_{\text{new}}), y_{\text{new}}) \right] := \mathbb{E}_{\boldsymbol{x}} \left[\mathcal{L}(\boldsymbol{\Phi}(\boldsymbol{x}), y) \right].$$

If the risk is not much larger than the empirical risk, then we say that the NN Φ has a small generalization error. Otherwise, if the risk is much larger than the empirical risk, then we say that Φ overfits the training data, meaning that it has memorized the training samples, but fails to generalize to new data (see [16, 19]).

In this context, let $d \in \mathbb{N}$; μ be a Borel probability measure on $[0,1]^d$; \mathcal{D} a joint distribution induced by μ of the set S, where \mathcal{D}_X and \mathcal{D}_y are the respective marginal distributions of $X := \{x_i\}_{i=1}^n$ and $y := \{y_i\}_{i=1}^n$;

$$f \in \mathcal{F} := \left\{ f : [0,1]^d \to [0,1] : f \text{ is a } \mu \text{-measurable function} \right\};$$
(8)

the classifier $C_f(\boldsymbol{x}) := \mathbb{1}_{[1/2,1]}(f(\boldsymbol{x}))$ and label $y \in \{0,1\}$. We want to obtain the minimizer

$$C^* := \operatorname*{argmin}_{f \in \mathcal{F}} \mathcal{E}(C_f) \quad \text{with} \quad \mathcal{E} := \mathcal{E}_{\mathcal{L}^*} \quad \text{and} \quad \mathcal{L}^*(C_f(\boldsymbol{x}), y) := \mathbb{1}_{(2C_f(\boldsymbol{x}) - 1)(2y - 1) < 0}, \quad (9)$$

called the Bayes classifier which is the function that has the most accurate classification on average from a sample set. The classifier C^* could be approximated by the empirical risk minimization for the same loss function \mathcal{L}^* , that is

$$C^* \approx \operatorname*{argmin}_{f \in \mathcal{F}} \widehat{\mathcal{E}}_{\mathcal{L}^*, S}(C_f),$$

but finding the minimizer of $\widehat{\mathcal{E}}_{\mathcal{L}^*,S}$ is NP hard (see [4]). However, in [4, 22], it was proven that if the loss function is the hinge loss, denoted by ϕ and defined^{*} as

$$\phi: [0,1] \times \{0,1\} \to [0,\infty)$$

(x,y) $\mapsto \max\{0, 1 - (2y-1)(2x-1)\},$ (10)

then

$$C^* = \operatorname*{argmin}_{f \in \mathcal{F}} \mathcal{E}_{\phi}(C_f),$$

and this is one of the main reasons why from now on we continue to work with the hinge loss in the remainder of this paper. It is important to consider that the fast convergence rate of the excess \mathcal{L} -risk $\mathcal{E}_{\mathcal{L}}(f, C^*) := \mathcal{E}_{\mathcal{L}}(f) - \mathcal{E}_{\mathcal{L}}(C^*)$ does not always imply the fast convergence rate of the excess \mathcal{L}^* -risk $\mathcal{E}(f, C^*) := \mathcal{E}(f) - \mathcal{E}(C^*)$.

But for the hinge loss ϕ , some constant $C_{\phi} > 0$ (which depends on ϕ) and any $f \in \mathcal{F}$, the inequality

$$\mathcal{E}(f, C^*) \le C_{\phi} \mathcal{E}_{\phi}(f, C^*) \tag{11}$$

holds (see [16, inequality (2.3)]). This is the second reason why we choose to work with hinge loss, and inequality (11) will be used to prove our main results.

1.2.2 Covering entropy

We need to introduce the notion of covering entropy given in [18, Definition 3.9] from [21, Definitions 1 and 2].

Let $\mathcal{X} \neq \emptyset$ be a set and let dist : $\mathcal{X} \times \mathcal{X} \rightarrow [0, \infty]$ be a distance function. For a set $\emptyset \neq K \subset \mathcal{X}$ and $\epsilon > 0$, we call a set $G_{\epsilon} \subset \mathcal{X}$ an ϵ -net for $K \subset \mathcal{X}$ if

for all
$$x \in K$$
 there exists $y \in G_{\epsilon}$ satisfying $\operatorname{dist}(x, y) \leq \epsilon$. (12)

Then, we define

 $\mathcal{G}_{K,\text{dist}}(\epsilon) := \min\{|G| : G \subset \mathcal{X} \text{ is an } \epsilon \text{-net for } K\} \text{ and } V_{K,\text{dist}}(\epsilon) := \ln(\mathcal{G}_{K,\text{dist}}(\epsilon)),$

^{*}With a slight difference, since in this case ϕ is defined on $[0,1] \times \{0,1\}$ instead of $\mathbb{R} \times \{\pm 1\}$.

where $V_{K,\text{dist}}(\epsilon)$ is called the ϵ -covering entropy of K.

The next result provides an upper bound for the covering entropy of \mathcal{NN}_* and is a consequence of [14, Lemma 6.1] presented in [18, Remark 5.2].

Lemma 1. Fixing $\mathcal{X} := \mathcal{NN}_*(d, N, W, B)$, $K := [0, 1]^d$ and dist $:= \|\cdot\|_{\infty}$, we have

$$V_{[0,1]^{d},\|\cdot\|_{\infty}}(\delta) \le W \cdot (10 + \ln(1/\delta) + 5\ln(\lceil B \rceil) + 5\ln(\max\{d,W\}))$$

for $\delta \in (0, 1], d, N, W \in \mathbb{N}$ and B > 0.

1.3 Fourier-analytic Barron space

There are several slightly different interpretations of the term "Barron function" in the literature, see for example [8, Section 7]. We use this term to refer to the Fourier-analytic notion of Barron-type functions in [2, 8, 18]. Consequently, we adopt the definition of [8, Definition 2.1] with parameters $X := [0, 1]^d$ and $x_0 := \mathbf{0}$, i.e., as follows.

A function $f : [0,1]^d \to \mathbb{R}$ is said to be of Barron class with constant C > 0, if there are $c \in [-C, C]$ and a measurable function $F : \mathbb{R}^d \to \mathbb{C}$ satisfying

$$\int_{\mathbb{R}^d} |F(\boldsymbol{\xi})| \sup_{\boldsymbol{x} \in [0,1]^d} |\langle \boldsymbol{\xi}, \boldsymbol{x} \rangle| \, d\boldsymbol{\xi} \le C \quad \text{and} \quad f(\boldsymbol{x}) = c + \int_{\mathbb{R}^d} (e^{i\langle \boldsymbol{x}, \boldsymbol{\xi} \rangle} - 1) \cdot F(\boldsymbol{\xi}) d\boldsymbol{\xi} \tag{13}$$

for all $\boldsymbol{x} \in [0,1]^d$. The set of all these Barron functions is denoted as \mathcal{B}_C .

In [2] it was shown that for a certain class of functions with bounded variation, shallow NNs of N neurons achieve an approximation accuracy of order $N^{-1/2}$ in the L^2 norm. Subsequently, in [8, Proposition 2.2] and [12, Theorem 12], the same accuracy was demonstrated for approximating functions of the Barron class using shallow ReLU neural networks with N neurons under the supremum norm. This allows us to formulate [8, Definition 3.1].

For C > 0, we define the Barron approximation set \mathcal{BA}_C as the set of all functions $f:[0,1]^d \to \mathbb{R}$ such that for all $N \in \mathbb{N}$, there is a shallow ReLU neural network (shallow ReLU-NN) Φ with N neurons in the hidden layer such that

$$\|f - \Phi\|_{\infty} \le C\sqrt{d/N} \tag{14}$$

and all weights and biases of Φ are bounded in absolute value by

$$\sqrt{C}(5+\vartheta), \quad \text{where} \quad \vartheta := \sup_{\boldsymbol{\xi} \in \mathbb{R}^d \setminus \{0\}} \left(\frac{\|\boldsymbol{\xi}\|_{\ell^{\infty}}}{\sup_{\boldsymbol{x} \in [0,1]^d} |\langle \boldsymbol{\xi}, \boldsymbol{x} \rangle|} \right)$$

In addition, we know that $\sqrt{C}(5+\vartheta) \leq 7\sqrt{C}$, since

for
$$\widetilde{\boldsymbol{x}} = \frac{1}{2} \left(\frac{\boldsymbol{\xi}}{\|\boldsymbol{\xi}\|} + \mathbf{1} \right) \in [0, 1]^d, \quad \sup_{\boldsymbol{x} \in [0, 1]^d} |\langle \boldsymbol{\xi}, \boldsymbol{x} \rangle| \ge |\langle \boldsymbol{\xi}, \widetilde{\boldsymbol{x}} \rangle| = \frac{|\langle \boldsymbol{\xi}, \boldsymbol{\xi} \rangle|}{2\|\boldsymbol{\xi}\|} + \frac{|\langle \boldsymbol{\xi}, \mathbf{1} \rangle|}{2} \ge \frac{\|\boldsymbol{\xi}\|}{2}$$

and $\vartheta \leq \sup_{\boldsymbol{\xi} \in \mathbb{R}^d \setminus \{0\}} (2 \|\boldsymbol{\xi}\|_{\ell^{\infty}} / \|\boldsymbol{\xi}\|) \leq 2.$

The set $\mathcal{BA} = \bigcup_{C>0} \mathcal{BA}_C$ is called Barron approximation space. In fact, $\mathcal{B}_{C/\kappa_0} \subset \mathcal{BA}_C$ for every C > 0 and a constant $\kappa_0 > 0$ that is absolute (i.e. independent of all other quantities and objects) (see [8, Remark 3.2]).

1.3.1 Barron regular boundaries

We now define the Barron regular boundaries using [18, Definitions 2.2 and 2.3] which are compatible with [8, Definition 3.3] for \mathcal{BA}_C by [18, Remark 2.4]. Let $M, d \in \mathbb{N}$ with $d \geq 2$, let a set $\emptyset \neq \mathscr{C} \subset C([0, 1]^{d-1}; [0, 1])$ and $b \in \mathscr{C}$. We define the general horizon function associated with b as

$$h_b: [0,1]^d \to \{0,1\},$$

 $x = (x_1, \dots, x_d) \mapsto \mathbb{1}_{b(x_1,\dots, x_{d-1}) \le x_d},$

and the set of general horizon functions associated to \mathscr{C} as $H_{\mathscr{C}} := \{h_b : b \in \mathscr{C}\}$.

Moreover, we say that a compact set $\Omega \subset [0,1]^d$ has (\mathscr{C}, M) -regular decision boundary if there exist (closed, axis-aligned, non-degenerate) rectangles $Q_1, \ldots, Q_M \subset [0,1]^d$ such that $\Omega \subset \bigcup_{i=1}^M Q_i$, where the Q_i have disjoint interiors and such that either

$$\mathbb{1}_{\Omega} = g_i \circ P_i \quad \text{or} \quad \mathbb{1}_{\Omega} = 1 - g_i \circ P_i \quad \text{almost everywhere on} \quad Q_i, \tag{15}$$

for a general horizon function $g_i \in H_{\mathscr{C}}$ associated to \mathscr{C} and a *d*-dimensional permutation matrix P_i . Also, a family $\{Q_i\}_{i=1}^M$ of rectangles as above is called an associated cover of Ω . We write $\mathcal{R}_{\mathscr{C}}(d, M)$ for all sets with (\mathscr{C}, M) -regular decision boundary. Then, we define the set of all classifiers with (\mathscr{C}, M) -regular decision boundary as

$$\mathcal{C}\ell_{\mathscr{C}}(d,M) := \{\mathbb{1}_{\Omega} : \Omega \in \mathcal{R}_{\mathscr{C}}(d,M)\}.$$

Finally, Barron horizon functions and classifiers with Barron-regular decision boundary are the elements of $H_{\mathcal{BA}_C}$ and $\mathcal{C}\ell_{\mathcal{BA}_C}(d, M)$.

2 Main results

Our main results confirm the intuitive idea that using the margin condition with an appropriate margin exponent for sets with Barron-regular decision boundary can drastically improve the way NNs learn. To establish our results, we consider the following parameters:

- (P1) Let $d \in \mathbb{N}_{\geq 2}$; $M, N \in \mathbb{N}$; $C_1, C_2, C_3 > 0$; $\gamma > 0$, $\alpha \in (0, 1]$ and $\Omega \in \mathcal{R}_{\mathcal{BA}_{C_1}}(d, M)$, where d is the dimension of Ω ; M is the number of sets in the associated cover of Ω (see (15)); C_1 is the constant in the right hand side of (14); C_2 and γ are the margin condition parameters in (M); and finally, C_3 and α are the tube compatibility parameters in (3).
- (P2) With parameters (P1), let

$$\begin{split} \widehat{N}_n &:= \left\lceil (7Md)^{2/\gamma} (d-1) C_1^2 \max\{C_2, C_3\}^{2/\gamma} n^{2/(2+\gamma)} \right\rceil, \\ N_n &:= \left\lceil M(4(d+1) + \widehat{N}_n + 1) + d + 1 \right\rceil, \\ W_n &:= \left\lceil 41Md^2 \widehat{N}_n \right\rceil, \\ B_n &:= \left\lceil (1 + \sqrt{C_1}) \left(7 + N/C_1 + (N/C_1)^{\gamma/\alpha} \right) \right\rceil, \quad \text{for all} \quad n \in \mathbb{N}, \end{split}$$

where n = |S| and S is a training set as in (2).

Our first result shows that classifiers in $\mathcal{C}\ell_{\mathcal{BA}_{C_1}}(d, M)$ are well approximated by ReLU-NNs when the margin condition is assumed.

Theorem 2. Let the parameters be as defined in (P1). There exists a ReLU-NN Φ with three hidden layers such that, for all probability measures μ satisfying the margin condition (M), and the tube compatibility (3), it holds that

$$\mu\left(\left\{\boldsymbol{x}\in[0,1]^d:\mathbb{1}_{\Omega}(\boldsymbol{x})\neq\Phi(\boldsymbol{x})\right\}\right)\leq 3.5Md(d-1)^{\gamma/2}C_1^{\gamma}N^{-\gamma/2}\max\{C_2,C_3\}.$$

Moreover, $0 \leq \Phi(\mathbf{x}) \leq 1$ for all $\mathbf{x} \in [0,1]^d$ and the architecture of Φ is given by

$$\mathcal{A} = (d, M(2(d+1) + N), M(2d+2), M, 1).$$

Thus, Φ has at most M(4(d+1)+N+1)+d+1 neurons and at most $41Md^2N$ non-zero weights. The weights and biases of Φ are bounded in magnitude by

$$(1 + \sqrt{C_1}) (7 + N/C_1 + (N/C_1)^{\gamma/\alpha}).$$

To prove this result, we use the property defined in (15) for $\Omega \in \mathcal{R}_{\mathcal{BA}_{C_1}}(d, M)$, which allows us to identify Ω in each of the sets associated to its covering through Barron horizon functions. Then, in each set of the Ω covering we use the approximation (14) for shallow ReLU-NNs, and construct the three hidden layers using the ReLU function. Next, we sum the NNs created in each set of the covering and obtain the final NN for the whole Ω set. Lastly, we find the architecture of this final NN and bound in absolute value its weights and biases.

The next step is to measure the performance of this approximation using the 0-1 loss. Therefore, we present the following result.

Theorem 3. Let the parameters be as defined in (P1) and (P2). Then, for all probability measures μ satisfying the margin condition (M), and the tube compatibility (3), we have that

$$\mu\left(\mathbb{E}_{\boldsymbol{x}}\left[\mathbbm{1}_{[1/2,1]}\left(\widehat{f}_{\phi,S}(\boldsymbol{x})\right)\neq\mathbbm{1}_{\Omega}(\boldsymbol{x})\right]\gtrsim n^{-\gamma/(2+\gamma)}(1+\log n)\right)\lesssim n^{-1},\tag{16}$$

where

$$\widehat{f}_{\phi,S} := \operatorname*{argmin}_{f \in \mathcal{F}} \widehat{\mathcal{E}}_{\phi,S}(f) \in \mathcal{NN}_*(d, N_n, W_n, B_n)$$

 $\widehat{\mathcal{E}}_{\phi,S}(f)$ is the ϕ -empirical risk as in (7), \mathcal{F} is defined in (8) and \mathcal{NN}_* in (6).

In order to prove Theorem 3, we use [16, Theorem A.1] with a slight adaptation to our hypothesis and obtain Lemma 5 as an auxiliary result.

Remark 4. Theorem 3 implies

$$\mathbb{E}_{S,\boldsymbol{x}}\left[\mathbb{1}_{[1/2,1]}\left(\widehat{f}_{\phi,S}(\boldsymbol{x})\right)\neq\mathbb{1}_{\Omega}(\boldsymbol{x})\right]\lesssim n^{-\gamma/(2+\gamma)}(1+\log n).$$

Indeed, we denote for a moment $\Gamma_{S,\boldsymbol{x}} := \left\{ \mathbb{1}_{[1/2,1]} \left(\widehat{f}_{\phi,S}(\boldsymbol{x}) \right) \neq \mathbb{1}_{\Omega}(\boldsymbol{x}) \right\}$ and observe that (16) implies

$$\mathbb{E}_{S,oldsymbol{x}}\left[\Gamma_{S,oldsymbol{x}}
ight] = \mathbb{E}_{S}\left[\mathbb{E}_{oldsymbol{x}}\left[\Gamma_{S,oldsymbol{x}}
ight]
ight]$$

$$\begin{split} &\lesssim \mathbb{E}_{S} \left[\mathbb{E}_{\boldsymbol{x}} \left[\Gamma_{S,\boldsymbol{x}} \right] \mathbb{1}_{\mathbb{E}_{\boldsymbol{x}} \left[\Gamma_{S,\boldsymbol{x}} \right] \gtrsim n^{-\gamma/(2+\gamma)} (1+\log n)} \right] + n^{-\gamma/(2+\gamma)} (1+\log n) \\ &\leq \mu \left(\mathbb{E}_{\boldsymbol{x}} \left[\Gamma_{S,\boldsymbol{x}} \right] \gtrsim n^{-\gamma/(2+\gamma)} (1+\log n) \right) + n^{-\gamma/(2+\gamma)} (1+\log n) \\ &\lesssim n^{-1} + n^{-\gamma/(2+\gamma)} (1+\log n) \\ &\lesssim n^{-\gamma/(2+\gamma)} (1+\log n), \end{split}$$

here we use the fact that S was employed to define the probability measure μ and therefore $\mathbb{E}_S[X] = \int_{\Omega} X d\mu$ holds for every random variable X.

To close, we must emphasize that when γ is large we achieve in Theorem 2, the approximation of a high-dimensional discontinuous function at a rate comparable to that of a low-dimensional smooth function. Moreover, we get in Theorem 3 and Remark 4, a fast learning rate of $\approx n^{-1}(1 + \log n)$ if γ is large.

2.1 Numerical experiments

In this section, we present some numerical simulations that support the results of Theorems 2 and 3, in practical scenarios. We build classifiers using TensorFlow in Python with different training samples and architectures that fulfill the hypotheses of our results. For a clear explanation of our algorithm, we detail each step below.

2.1.1 Sample generation

We denote our samples as $S_{d,n}^{\gamma}$ where $d \in \{3, 50, 784\}$ is the dimension of $\Omega := \Omega^d \subset [0, 1]^d$, $n = |S_{d,n}^{\gamma}|$ is the number of examples, and γ is the margin exponent in (M). Subsequently, we define the characteristics of our samples and the distance dist $(\boldsymbol{x}, \partial \Omega^d)$ in order to apply the margin condition (M).

• For $d \in \{3, 50\}$, we construct a data set where we choose the hypersurface $\partial \Omega^d$ that separates the labels as the positive part of the hypersphere of dimension d and radius 1/2. Concretely, let

 $\mathcal{X}_r := \left\{ oldsymbol{x} \in [0,1]^d : oldsymbol{x} ext{ has positive entries and } \|oldsymbol{x}\| \leq r
ight\} ext{ and } \widetilde{\mathcal{X}}_r ext{ } := \left\{ oldsymbol{x}/r : oldsymbol{x} \in \mathcal{X}_r
ight\}.$

To avoid problems with the volume of hypespheres in high dimensions for r < 1, we start by taking r = 4 and generating 3×10^5 examples in \mathcal{X}_4 for training and 10^6 for testing, making sure that $|\mathcal{X}_4 \setminus \mathcal{X}_2| = |\mathcal{X}_2|$ (see Figure 1).



Figure 1: Examples test for d = 3.

However, our examples must be in $[0,1]^d$, so we define them such that $S_{d,n}^{\gamma} \subset \widetilde{\mathcal{X}}_4$ and the distance between any $\boldsymbol{x} \in S_{d,n}^{\gamma}$ and $\partial \Omega^d$ as dist $(\boldsymbol{x}, \partial \Omega^d) := |1/2 - ||\boldsymbol{x}|||$.

• For d = 784, we focus on an example based on real data. Therefore, we use the MNIST data set to have images of size 28×28 pixels, and we classify the images of numbers 0 and 1. Since MNIST does not have a large number of elements and we want several samples to observe the effect of the margin condition (M) and the number of elements, we use the SMOTE [9] technique (similar to the method used in [16] to generate artificial samples) to equalize the number of elements in each label and increase the sample of each label to 95% more than the larger original, obtaining in total 18576 examples for training and 12146 for test (see Figure 2).



Figure 2: Some examples after smote for d = 784.

In this sense, we define $S_{784,n}^{\gamma}$ such that its elements are vectors in $[0,1]^{784}$ constructed from matrices in $\mathbb{R}^{28\times28}$ flattened and with entries normalized to [0,1] representing images of 0 and 1. However, to apply the margin condition (M) we must have an idea of the distance between the examples and the decision boundary, so we use an initial NN[†] Φ_0 that assigns probabilities to each example and form a vector $\boldsymbol{w} = (w_i)_{i=1}^n$ with these predictions (and keeping only the well predicted examples), i.e. $w_i = \Phi_0(\boldsymbol{x})$ for some $\boldsymbol{x} \in S_{784,n}^{\gamma}$ and $i \in \{1, \ldots, n\}$, where $\mathbb{1}_{[1/2,1]}(\Phi_0(\boldsymbol{x}))$ corresponds to the original label of \boldsymbol{x} . Then, \boldsymbol{w} gives us an idea of the position of the examples and the possible decision boundary, using their probabilities in the interval [0,1]. Thus, we assume for a moment as the decision boundary, the set of $\boldsymbol{x} \in [0,1]^{784}$ with $\Phi_0(\boldsymbol{x}) \in [\ell, u]$ for some $\ell, u \in (0,1)$ such that $[\min_{i \in \{1,\ldots,n\}} \{w_i\}, \ell\} \cup (u, \max_{i \in \{1,\ldots,n\}} \{w_i\}]$ contains approximately 99.9% of the labels $\{w_i\}_{i=1}^n$ (see Figure 3).



Figure 3: Train sample probability applying Φ_0 and decision boundary for d = 784.

 $^{^{\}dagger}\Phi_{0}$ is a ReLU-NN with architecture (784, 256, 128, 64, 1), which was obtained using hinge loss and sigmoid function is applied to the output.

Next, we remove the 0.1% of elements with label in $[\ell, u]$ and rescale the values of \boldsymbol{w} with the mapping $t : [0, 1]^{784} \rightarrow [0, 1]$ that maps $\min_{i \in \{1, \dots, n\}} \{w_i\}$ to 0, ℓ, u to 1/2, and $\max_{i \in \{1, \dots, n\}} \{w_i\}$ to 1. Finally, after this process, we obtain 18337 elements for the training set (see Figure 4) and 12053 for the test set, with the same number of elements for each label. To simulate a margin condition, we require, for a given point $\boldsymbol{x} \in S^{\gamma}_{784,n}$, its distance to the decision boundary $\partial \Omega^{784}$. This is computationally very hard to identify. Therefore, we use the expression $|1/2 - t(\Phi_0(\boldsymbol{x}))|$ as a proxy for the distance, since

dist
$$(\boldsymbol{x}, \partial \Omega^{784}) \gtrsim |1/2 - t(\Phi_0(\boldsymbol{x}))|$$
 for all $i \in \{1, \dots, n\}.$ (17)

This proxy is reasonable because of the following considerations:

- $\partial \Omega^{784} = \partial \{ \boldsymbol{x} \in [0, 1]^d : \eta(\boldsymbol{x}) \approx t(\Phi_0(\boldsymbol{x})) \ge 1/2 \}.$
- $t(\Phi_0(\boldsymbol{x}))$ is a modified sigmoid function of the initial sigmoid $\Phi_0(\boldsymbol{x})$.
- t(Φ₀(**x**)) = 1/(1 + e^{-(⟨ŵ,x⟩+b)}) for some weight vector ŵ and bias b, after they have been modified by the use of the t function.
- $G := \{ \boldsymbol{x} \in [0,1]^d : \langle \hat{\boldsymbol{w}}, \boldsymbol{x} \rangle + b = 0 \}$ is the projection of the decision boundary in the characteristics space, therefore we assume

dist
$$(\boldsymbol{x}, \partial \Omega^{784}) \approx \text{dist} (\boldsymbol{x}, G) = |\langle \widehat{\boldsymbol{w}}, \boldsymbol{x} \rangle + b| / ||\widehat{\boldsymbol{w}}||.$$

• Approximating $1/(1+e^{-y})$ around y = 0, implies that $1/(1+e^{-y}) \approx 1/2 + y/4$. So, when $y = \langle \widehat{\boldsymbol{w}}, \boldsymbol{x} \rangle + b$, we obtain

$$|1/2 - t(\Phi_0(\boldsymbol{x}))| \approx |\langle \boldsymbol{\hat{w}}, \boldsymbol{x} \rangle + b|/4 \lesssim \operatorname{dist}(\boldsymbol{x}, \partial \Omega^{784})$$

In conclusion, to apply the margin condition (M) in this case, we use $|1/2-t(\Phi_0(\boldsymbol{x}))|$ instead of the exactly dist $(\boldsymbol{x}, \partial \Omega^{784})$, which is a stronger assumption.



Figure 4: Final train sample after SMOTE and location of the possible $\partial \Omega^{784}$.

To the basis sets defined in the previous items to describe the characteristics of the elements of $S_{d,n}^{\gamma}$, we apply the margin condition (M) with margin exponent

 $\gamma \in \Gamma := \{0.1, 0.644, 1.189, 1.733, 2.278, 2.822, 3.367, 3.911, 4.456, 5.0\},\$

defining a neighborhood centered at 0.5 of radius $c_d = 0.48$ when $d \in \{3, 50\}$ or $c_d \approx 0.5 - 2.8 \cdot 10^{-7}$ for d = 784, and assigning to each point a weight of 0 if it is outside this neighborhood or $1 - (\text{dist}(\boldsymbol{x}, \partial \Omega^d) / c_d)^{\gamma}$ if it is not. So, we randomly decide to remove from the sample with probability according to the weight. Then, for each γ we randomly select subsets of the samples, such that $n \in G_1$ with

 $G_1 := \{499, 514, 559, 643, 782, 1005, 1368, 1967, 2993, 4815, 8192, 14737, 28038, 56412, 120025\},\$

for train, and n = 400108 for test, when $d \in \{3, 50\}$; or

 $n \in G_2 := \{249, 321, 414, 533, 687, 885, 1140, 1468, 1890, 2435, 3135, 4038, 5200, 6696, 8624\}$

for test and n = 4837, when d = 784; and $|S_{d,n}^{\gamma_1}| = |S_{d,n}^{\gamma_2}|$ holds for all $\gamma_1, \gamma_2 \in \Gamma$ with $\gamma_1 \neq \gamma_2$ (see e.g. Figures 5 and 6).

 $\gamma = 2.278$ and n = 8192



Figure 5: Example of margin condition applied for d = 3.



Figure 6: Example of margin condition applied for d = 784. The second row contains the points of the first row but in ascending order.

Lastly, as the samples $S_{784,n}^{\gamma}$ have few elements and the decision boundary is not known beforehand, we repeat the process up to 56 iterations and obtain different[‡] samples to train and test, which will then be used to have an average result.

2.1.2 NN training

With the $S_{d,n}^{\gamma,\text{train}} \in \{S_{d,n}^{\gamma}\}_{n \in G_1 \cup G_2}$ samples previously generated as training sets, we train ReLU-NNs with architecture

$$(d, 3N, 2N, N, 1)$$
 where $N = \lceil n^{2/(\gamma+2)} \rceil$,

using the hinge loss ϕ and applying the sigmoid function to the output, we also use the optimizer Adam with a learning rate of 1e - 5, we apply early stop monitoring loss with patience equal to 1 for 50000 epochs, and we do not use validation sets as it is usually done, since our main generalization result does not describe this situation, but the errors minimizer of the training risk. Next, we compute the empirical 0 - 1 risk as in (7), using the samples $S_{d,n}^{\gamma,\text{test}} \in \{S_{d,n}^{\gamma}\}_{n \in \{4837,400108\}}$ previously defined as test sets, and obtain

$$\mathbb{E}_{S_{d,n}^{\gamma,\mathrm{train}},S_{d,n}^{\gamma,\mathrm{test}}}\left[\mathbbm{1}_{[1/2,1]}\circ\widehat{f}_{\phi,S_{d,n}^{\gamma,\mathrm{train}}}\neq\mathbbm{1}_{\Omega}\right]$$

as in the Remark 4. Our results can be seen in Figure 7, for each $d \in \{3, 50, 784\}$ and $\gamma \in \Gamma$, of n vs empirical 0 - 1 risk.

[‡]The number of examples of the base set for d = 784 is > 12053 and $n = |S_{784,n}^{\gamma}| < 8624$ for any $\gamma \in \Gamma$, also the selection of elements is done randomly, this ensures that the iterations have different elements.



Figure 7: Plot n vs 0-1 loss. The labels in the d = 784 plot are the same for the other plots. In the d = 784 case, a method of averaging over 56 iterations with an error shadow using the standard deviation was used, while in the $d \in \{3, 50\}$ cases, the empirical risk was only calculated in a single iteration.

• The $(d, \gamma) \in \{3, 50\} \times \{0.1\}$ cases could only be computed as far as seen in Figure 7 because the memory requirements of the data and architecture became excessive.

• For d = 784, we repeated the experiments obtaining different $S_{784,n}^{\gamma,\text{train}}$ and $S_{784,n}^{\gamma,\text{test}}$ in 56 iterations, then applied mean on all these results and plotted an error shadow based on the standard deviation of the results in these iterations.

In conclusion the simulations support our results (Theorem 3 and Remark 4). Furthermore, Figure 7 shows that the rate approaches n^{-1} for large margin and does not for small margin. The $\gamma = 5.0$ curve spans close to two orders of magnitude. The number of samples spans 1.5 orders of magnitude, so the error rate is close to or even slightly exceeding the rate n^{-1} . The case $\gamma = 0.1$, yields a reduction of the loss by roughly one order of magnitude, so the associated error rate is worse than n^{-1} . For the other examples, the x axis goes through 2.4 orders of magnitude. The curves with the highest margin also manage a bit more than 2 orders of magnitude. The ones with lower margin (green for example) only manage 1, so closer to the $n^{-1/2}$ rate.

Acknowledgements

J.G. and P.C.P. were supported by the Austrian Science Fund (FWF) Project P-37010.

References

- [1] A. R. BARRON, *Neural net approximation*, In Proc. 7th Yale Workshop on Adaptive and Learning Systems, 1 (1992), pp. 69–72.
- [2] A. R. BARRON, Universal approximation bounds for superpositions of a sigmoidal function, IEEE Transactions on Information Theory, 39 (1993), pp. 930–945.
- [3] A. R. BARRON, Approximation and estimation bounds for artificial neural networks, Machine learning, 14 (1994), pp. 115–133.
- [4] P. L. BARTLETT, M. I. JORDAN, AND J. D. MCAULIFFE, Convexity, classification, and risk bounds, Journal of the American Statistical Association, 101 (2006), pp. 138– 156.
- [5] C. M. BISHOP, *Pattern Recognition and Machine Learning*, Springer New York, 2016.
- [6] L. W. C. MA AND E. WEINAN, A priori estimates of the population risk for two-layer neural networks, Communications in Mathematical Sciences, 17 (2019), pp. 1407–1425.
- [7] E. J. CANDÈS AND D. L. DONOHO, New tight frames of curvelets and optimal representations of objects with piecewise c2 singularities, Communications on Pure and Applied Mathematics, 57 (2004), pp. 219–266.
- [8] A. CARAGEA, P. PETERSEN, AND F. VOIGTLAENDER, Neural network approximation and estimation of classifiers with classification boundary in a Barron class, The Annals of Applied Probability, 33 (2023), pp. 3039 – 3079.
- [9] N. CHAWLA, K. BOWYER, L. HALL, AND W. KEGELMEYER, Smote: Synthetic minority over-sampling technique, J. Artif. Intell. Res. (JAIR), 16 (2002), pp. 321– 357.
- [10] A. CHRISTMANN AND I. STEINWART, Support Vector Machines, Springer-Verlag New York, 2008.
- [11] L. DEVROYE, L. GYÖRFI, AND G. LUGOSI, A Probabilistic Theory of Pattern Recognition, Springer New York, 1996.
- [12] W. E, C. MA, S. WOJTOWYTSCH, AND L. WU., Towards a mathematical understanding of neural network-based machine learning: what we know and what we don't, arXiv:2009.10713, (2020).

- [13] W. E AND S. WOJTOWYTSCH, Representation formulas and pointwise properties for Barron functions, Calculus of Variations and Partial Differential Equations, 61 (2022), pp. 1–37.
- [14] P. GROHS AND F. VOIGTLAENDER, Proof of the theory-to-practice gap in deep learning via sampling complexity bounds for neural network approximation spaces, Foundations of Computational Mathematics, (2023).
- [15] T. HASTIE, R. TIBSHIRANI, AND J. FRIEDMAN, The Elements of Statistical Learning, Springer New York, 2009.
- [16] Y. KIM, I. OHN, AND D. KIM, Fast convergence rates of deep neural networks for classification, Neural Networks, 138 (2021), pp. 179–197.
- [17] P. PETERSEN AND F. VOIGTLAENDER, Optimal approximation of piecewise smooth functions using deep relu neural networks, Neural Networks, 108 (2018), pp. 296–330.
- [18] P. PETERSEN AND F. VOIGTLAENDER, Optimal learning of high-dimensional classification problems using deep neural networks, arXiv:2112.12555, (2021).
- [19] P. PETERSEN AND J. ZECH, Mathematical theory of deep learning, arXiv:2407.18384, (2024).
- [20] A. B. TSYBAKOV, Optimal aggregation of classifiers in statistical learning, The Annals of Statistics, 32 (2004), pp. 135 – 166.
- [21] Y. YANG AND A. BARRON, Information-theoretic determination of minimax rates of convergence, The Annals of Statistics, 27 (1999), pp. 1564 – 1599.
- [22] T. ZHANG, Statistical behavior and consistency of classification methods based on convex risk minimization, The Annals of Statistics, 32 (2004), pp. 56 – 85.

A Appendix

A.1 Proof of Theorem 2

Since $\Omega \in \mathcal{R}_{\mathcal{BA}_{C_1}}(d, M)$, there exists an associated cover of Ω , called $\{Q_i\}_{i=1}^M$, such that property (15) is satisfied for each $i \in \{1, \ldots, M\}$, with $P_i(\boldsymbol{x}) = (x_{i1}, \ldots, x_{id})$ where x_{i1}, \ldots, x_{id} are the entries of vector \boldsymbol{x} after being permuted via P_i , and $g_i(P_i(\boldsymbol{x})) =$ $\mathbb{1}_{b_i(x_{i1},\ldots,x_{i(d-1)}) \leq x_{id}}$ for some $b_i \in \mathcal{BA}_{C_1} \subset C([0,1]^{d-1};[0,1])$. Note that property (13) is fulfilled even for $f_i \in \mathcal{BA}_{C_1}$ defined as

$$f_i(\boldsymbol{x}^{(i)}) := f_i(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_d) = b_i(x_{i1}, \dots, x_{i(d-1)})$$
 and $x_i := x_{id}$,

given that it has been shown in [18, A.2] that in this case f_i can be seen as a composition of b_i with permutations of the vector $\boldsymbol{x}^{(i)}$. Therefore, fixing $m \in \{1, \ldots, M\}$, we have

$$\mathbb{1}_{Q_m \cap \Omega}(\boldsymbol{x}) = \mathbb{1}_{x_i \leq f_m(\boldsymbol{x}^{(i)})} \quad \text{or} \quad \mathbb{1}_{Q_m \cap \Omega}(\boldsymbol{x}) = \mathbb{1}_{x_i \geq f_m(\boldsymbol{x}^{(i)})} \quad \text{almost everywhere on} \quad Q_m.$$

Without loss of generality (otherwise the proof is analogous) we say

$$\mathbb{1}_{Q_m \cap \Omega}(\boldsymbol{x}) = \mathbb{1}_{x_i \le f_m(\boldsymbol{x}^{(i)})} \quad \text{for} \quad Q_m := \prod_{i=1}^d [a_i, b_i] \subseteq [0, 1]^d \quad \text{and} \quad \boldsymbol{x}^{(i)} \in Q_m \setminus \prod_{\substack{j=0\\j \ne i}}^d [a_j, b_j].$$

By inequality (14) and as $f_i \in \mathcal{BA}_{C_1}$, there exists a shallow ReLU-NN Φ_m such that

$$\|f_m - \Phi_m\|_{\infty} \le \delta := C_1 \sqrt{(d-1)/N} \tag{18}$$

and all weights and biases of Φ_m are bounded in absolute value by $7\sqrt{C_1}$. Let $G_m := \left\{ \boldsymbol{x} \in Q_m : \mathbb{1}_{Q_m \cap \Omega}(\boldsymbol{x}) \neq \mathbb{1}_{x_i \leq \Phi_m(\boldsymbol{x}^{(i)})} \right\}$ and observe that

$$G_{m} = \left\{ \boldsymbol{x} \in Q_{m} : \mathbb{1}_{x_{i} \leq f_{m}(\boldsymbol{x}^{(i)})} \neq \mathbb{1}_{x_{i} \leq \Phi_{m}(\boldsymbol{x}^{(i)})} \right\}$$

= $\left\{ \boldsymbol{x} \in Q_{m} : x_{i} - f_{m}(\boldsymbol{x}^{(i)}) \leq 0 \right\} \bigtriangleup \left\{ \boldsymbol{x} \in Q_{m} : x_{i} - f_{m}(\boldsymbol{x}^{(i)}) \leq \Phi_{m}(\boldsymbol{x}^{(i)}) - f_{m}(\boldsymbol{x}^{(i)}) \right\}$
 $\subseteq \left\{ \boldsymbol{x} \in Q_{m} : |x_{i} - f_{m}(\boldsymbol{x}^{(i)})| \leq |\Phi_{m}(\boldsymbol{x}^{(i)}) - f_{m}(\boldsymbol{x}^{(i)})| \leq \delta \right\},$

where \triangle denotes the symmetric difference of sets. If $\boldsymbol{x} = (x_1, \ldots, x_d) \in G_m$ and $\boldsymbol{z}^* =$ $(x_1,\ldots,x_{i-1},f_m(\boldsymbol{x}^{(i)}),x_{i+1},\ldots,x_d)$ is in the decision boundary $\partial\Omega$, we get

$$\operatorname{dist}(\boldsymbol{x},\partial\Omega) = \inf_{\boldsymbol{x}^* \in \partial\Omega} \|\boldsymbol{x} - \boldsymbol{x}^*\|_2 \le \|\boldsymbol{x} - \boldsymbol{z}^*\|_2 = |x_i - f_m(\boldsymbol{x}^{(i)})| \le \delta,$$
(19)

therefore $\boldsymbol{x} \in B^*_{\delta}$; this implies $G_m \subseteq B^*_{\delta}$. As the margin condition (M) is satisfied with $\gamma > 0$, we have

$$\mu(G_m) \le \mu(B^*_\delta) \le C_2 \delta^{\gamma}.$$
(20)

Now, we define the ReLU-NN

$$\widetilde{\Phi}_m(\boldsymbol{x}) := H_{\delta} \left(\Phi_m(\boldsymbol{x}^{(i)}) - x_i \right), \qquad \text{where} \quad H_{\delta}(x) := \begin{cases} 0 & \text{if } x \leq 0 \\ x/\delta & \text{if } 0 \leq x \leq \delta \\ 1 & \text{if } x \geq \delta \end{cases}$$
$$= \frac{1}{\delta} \left(\varrho(x) - \varrho(x - \delta) \right).$$

We also note that if

$$\boldsymbol{x} \in \widetilde{G}_m := \left\{ \boldsymbol{x} \in Q_m : \mathbb{1}_{x_i \le \Phi_m(\boldsymbol{x}^{(i)})} \neq \widetilde{\Phi}_m(\boldsymbol{x}) \right\} = \left\{ \boldsymbol{x} \in Q_m : 0 \le \Phi_m(\boldsymbol{x}^{(i)}) - x_i < \delta \right\},$$

then

$$|x_i - f_m(\boldsymbol{x}^{(i)})| - |f_m(\boldsymbol{x}^{(i)}) - \Phi_m(\boldsymbol{x}^{(i)})| \le |\Phi_m(\boldsymbol{x}^{(i)}) - f_m(\boldsymbol{x}^{(i)}) + f_m(\boldsymbol{x}^{(i)}) - x_i| < \delta.$$

Therefore, the above inequality and (18) imply that

$$|x_i - f_m(\boldsymbol{x}^{(i)})| \le |f_m(\boldsymbol{x}^{(i)}) - \Phi_m(\boldsymbol{x}^{(i)})| + \delta < 2\delta,$$

and with the argument of (19) we obtain $\boldsymbol{x} \in B_{2\delta}^*$; whereby $\widetilde{G}_m \subseteq B_{2\delta}^*$. So, by (M),

$$\mu(\widetilde{G}_m) \le \mu(B_{2\delta}^*) \le C_2(2\delta)^{\gamma}.$$
(21)

So far we constructed the sets G_m and \tilde{G}_m to approximate the indicator function $\mathbb{1}_{\Omega}$ with the ReLU-NN Φ_m , only on the set Q_m . In the following, we join these approximations and find another approximation for the same indicator function but on all Ω . To this end, we define

$$\widehat{\Phi}_m\left(\boldsymbol{x}, \widetilde{\Phi}_m(\boldsymbol{x})\right) := \begin{cases} 0 & \text{if } \exists i \in \{1, \dots, d\} : 0 < b_i - a_i < 2\widehat{\delta} \\ \varrho\left(\sum_{i=1}^d t_i(x_i) + \varrho(\widetilde{\Phi}_m(\boldsymbol{x})) - d\right) & \text{otherwise,} \end{cases}$$

with $\widehat{\delta} := \delta^{\gamma/\alpha}$,

$$t_{i}(u) := \begin{cases} 0 & u \in [0,1] \setminus [a_{i},b_{i}] \\ 1 & u \in [a_{i} + \hat{\delta},b_{i} - \hat{\delta}] \\ (u-a_{i})/\hat{\delta} & u \in [a_{i},a_{i} + \hat{\delta}] \\ (b_{i}-u)/\hat{\delta} & u \in [b_{i} - \hat{\delta},b_{i}]; \end{cases}$$
(22)

and

$$\widehat{G}_m := \left\{ \boldsymbol{x} \in [0,1]^d : \widehat{\Phi}_m\left(\boldsymbol{x}, \widetilde{\Phi}_m(\boldsymbol{x})\right) \neq \mathbb{1}_{Q_m} \cdot \widetilde{\Phi}_m(\boldsymbol{x}) \right\}$$

Note that when there exists $i \in \{1, \ldots, d\}$ such that $0 < b_i - a_i < 2\hat{\delta}$, we get $Q_m \subset T^{(i)}_{\frac{1}{2}(b_i - a_i) + a_i,\hat{\delta}}$, where $T^{(i)}_{\frac{1}{2}(b_i - a_i) + a_i,\hat{\delta}}$ is a tube as in (3), and therefore the tube compatibility of μ implies that

$$\mu(\widehat{G}_m) = \left(\left\{\boldsymbol{x} \in [0,1]^d : \mathbb{1}_{Q_m} \cdot \widetilde{\Phi}_m(\boldsymbol{x}) \neq 0\right\}\right) \le \mu(Q_m) \le \mu\left(T^{(i)}_{\frac{1}{2}(b_i - a_i) + a_i,\widehat{\delta}}\right) \le C_3\widehat{\delta}^{\alpha}.$$
(23)

If for all $i \in \{1, \ldots, d\}$, $b_i - a_i \ge 2\widehat{\delta}$ is satisfied, then

• If $\boldsymbol{x} \in [0,1]^d \setminus Q_m$, exists x_i such that $x_i \notin [a_i, b_i]$ and

$$\sum_{i=1}^{d} t_i(x_i) + \varrho(\widetilde{\Phi}_m(\boldsymbol{x})) - d \le \varrho(\widetilde{\Phi}_m(\boldsymbol{x})) - 1 \le 0,$$
(24)

this implies that $\widehat{\Phi}_m\left(\boldsymbol{x}, \widetilde{\Phi}_m(\boldsymbol{x})\right) = 0 = \mathbb{1}_{Q_m} \cdot \widetilde{\Phi}_m(\boldsymbol{x}).$

• If $\boldsymbol{x} \in \prod_{i=1}^{d} [a_i + \widehat{\delta}, b_i - \widehat{\delta}],$

$$\widehat{\Phi}_m\left(oldsymbol{x},\widetilde{\Phi}_m(oldsymbol{x})
ight) = arrho\left(arrho(\widetilde{\Phi}_m(oldsymbol{x}))
ight) = \mathbbm{1}_{Q_m}\cdot\widetilde{\Phi}_m(oldsymbol{x})$$

Then we conclude that $\widehat{G}_m \subset Q_m \setminus \prod_{i=1}^d [a_i + \widehat{\delta}, b_i - \widehat{\delta}]$ and analogously to (23) we get

$$\mu(\widehat{G}_m) \le \mu\left(Q_m \setminus \prod_{i=1}^d [a_i + \widehat{\delta}, b_i - \widehat{\delta}]\right) \le \mu\left(\bigcup_{i=1}^d \left[T^{(i)}_{a_i + \widehat{\delta}/2, \widehat{\delta}/2} \cup T^{(i)}_{b_i - \widehat{\delta}/2, \widehat{\delta}/2}\right]\right) \le 2dC_3(\widehat{\delta}/2)^{\alpha}.$$
(25)

Finally, we define

$$G_m' := \left\{ oldsymbol{x} \in [0,1]^d : \mathbb{1}_{Q_m \cap \Omega}(oldsymbol{x})
eq \widehat{\Phi}_m\left(oldsymbol{x}, \widetilde{\Phi}_m(oldsymbol{x})
ight)
ight\}$$

and obtain using (20), (21), (23) and (25), that

$$\mu(G'_{m}) \leq \mu(G_{m}) + \mu(\widetilde{G}_{m}) + \mu(\widehat{G}_{m})$$

$$\leq C_{2}\delta^{\gamma} + C_{2}(2\delta)^{\gamma} + 2dC_{3}(\widehat{\delta}/2)^{\alpha}$$

$$\leq (3C_{2} + 2dC_{3})\delta^{\gamma}$$

$$\leq (3C_{2} + 2dC_{3})\left(C_{1}\sqrt{(d-1)/N}\right)^{\gamma}$$

$$\leq 3.5d(d-1)^{\gamma/2}C_{1}^{\gamma}N^{-\gamma/2}\max\{C_{2},C_{3}\}.$$

We put, for $\boldsymbol{x} \in [0, 1]^d$

$$\Phi(\boldsymbol{x}) := \sum_{m=1}^{M} \widehat{\Phi}_m\left(\boldsymbol{x}, \widetilde{\Phi}_m(\boldsymbol{x})\right)$$
(26)

and since $\mathbb{1}_{\Omega} = \sum_{m=1}^{M} \mathbb{1}_{\Omega \cap Q_m}$ a.e., we conclude that

$$\mu\left(\left\{\boldsymbol{x}\in[0,1]^{d}:\mathbb{1}_{\Omega}(\boldsymbol{x})\neq\Phi(\boldsymbol{x})\right\}\right)\leq\sum_{m=1}^{M}\mu\left(G_{m}'\right)\leq3.5Md(d-1)^{\gamma/2}C_{1}^{\gamma}N^{-\gamma/2}\max\{C_{2},C_{3}\}.$$

Moreover, for some $m \in \{1, \ldots, M\}$, we know that $Q_m \cap Q_j = \emptyset$ for all $m \neq j$. If $\boldsymbol{x} \in Q_m$ is fixed[§], then for all $j \neq m$, $\widehat{\Phi}_j(\boldsymbol{x}, \widetilde{\Phi}_j(\boldsymbol{x})) = 0$ (see (24)) and

$$0 \le \Phi(\boldsymbol{x}) = \widehat{\Phi}_m\left(\boldsymbol{x}, \widetilde{\Phi}_m(\boldsymbol{x})\right) \le \varrho(\varrho(\widetilde{\Phi}_m(\boldsymbol{x}))) \le \widetilde{\Phi}_m(\boldsymbol{x}) \le 1.$$

This completes the first part of the theorem.

To conclude, we obtain information about the architecture, biases, and weights of Φ . Note that by (5) we can define

$$\Phi_m(\boldsymbol{x}^{(i)}) = b^{(1)} + \sum_{i=1}^N w_i^{(1)} \varrho\left(\left\langle \boldsymbol{w}_i^{(0)}, \boldsymbol{x}^{(i)} \right\rangle + b_i^{(0)}\right)$$
(27)

for some $w_i^{(1)}, b_i^{(0)}, b^{(1)} \in \mathbb{R}$ and $w_i^{(0)} \in \mathbb{R}^{d-1}$ for i = 1, ..., N. Also, $x_i = \varrho(x_i) - \varrho(-x_i)$ and

$$\widetilde{\Phi}_{m}(\boldsymbol{x}) = \frac{1}{\delta} \left(\varrho \left(\Phi_{m}(\boldsymbol{x}^{(i)}) - (\varrho(x_{i}) - \varrho(-x_{i})) \right) - \varrho \left(\Phi_{m}(\boldsymbol{x}^{(i)}) - (\varrho(x_{i}) - \varrho(-x_{i})) - \delta \right) \right).$$
(28)

Finally, $t_i(u) = \frac{t_i^1 - t_i^2 - t_i^3 + t_i^4}{\widehat{\delta}}(u)$ where

$$t_{i}^{1}(u) = \varrho(u - a_{i}), \quad t_{i}^{2}(u) = \varrho(u - a_{i} - \widehat{\delta}), \quad t_{i}^{3}(u) = \varrho(u - b_{i} + \widehat{\delta}), \quad t_{i}^{4}(u) = \varrho(u - b_{i}), \quad (29)$$

and

$$\widehat{\Phi}_{m}\left(\boldsymbol{x},\widetilde{\Phi}_{m}(\boldsymbol{x})\right) := \begin{cases} 0 & \text{if } \exists i: 0 < b_{i} - a_{i} < 2\widehat{\delta} \\ \varrho\left(\sum_{i=1}^{d} \left(\frac{t_{i}^{1} - t_{i}^{2} - t_{i}^{3} + t_{i}^{4}}{\widehat{\delta}}(x_{i})\right) + \varrho(\widetilde{\Phi}_{m}(\boldsymbol{x})) - d\right) & \text{otherwise.} \end{cases}$$

$$(30)$$

We count the number of neurons as follows.

[§]If for all $m \in \{1, \ldots, M\}$, $\boldsymbol{x} \notin Q_m$, then $\Phi(\boldsymbol{x}) = 0$ (see (22) and (24)).

- Input layer. We have d neurons by the numbers of coordinates in the vector $\boldsymbol{x} \in [0, 1]^d$.
- Hidden layer 1. In (27) we are using N neurons of Φ_m , for (28) we have to find before $\rho(\pm x_i)$ and $\rho(\pm(\rho(x_i) - \rho(-x_i)))$ so we need 2d and 2 more neurons. Then we get 2(d+1) + N for each $i = 1, \ldots, d$, but we was fixed m, thus in total we obtain M(2(d+1) + N).
- Hidden layer 2. For (28), we know that

$$\varrho\left(\Phi_m(\boldsymbol{x}^{(i)}) - (\varrho(x_i) - \varrho(-x_i))\right) \quad \text{and} \quad \varrho\left(\Phi_m(\boldsymbol{x}^{(i)}) - (\varrho(x_i) - \varrho(-x_i)) - \delta\right) \quad (31)$$

add up to two neurons, but in addition the decomposition (29) of t_i for the next layer is now done using 4 neurons for each i, so we have 4d more neurons. Then, in total we get 2M(2d+1).

- Hidden layer 3. For (30), we see that among the possibilities 0 or $\rho(\cdot)$ of $\widehat{\Phi}_m$, we use only one neuron, however $m \in \{1, \ldots, M\}$ is fixed, so in total we have M.
- Output layer. Here we only obtain $\Phi(\boldsymbol{x})$ and by its definition in (26) we use one neuron for the output.

In summary, the architecture of this NN is given by

$$(d, M(2(d+1)+N), 2M(d+1), M, 1)$$

and the number of all neurons is M(4(d+1) + N + 1) + d + 1. Also, an upper bound for the total weights can be obtained by summing the number of neurons of each layer when multiplied by its input dimension (which is given by the output dimension of the previous layer), plus the number of non-input neurons associated with the biases and Mto account for the weights of final output layer. Then

$$\begin{split} W(\Phi) &\leq dM(2(d+1)+N) + 2M(d+1)(2(d+1)+N) + 2M(d+1) \\ &+ M(4(d+1)+N+1) + M + 1 \\ &\leq 3dMN + 3MN + 6d^2M + 16dM + 12M + 1 \\ &\leq 41Md^2N \end{split}$$

By (18), we use $7\sqrt{C_1}$ to bound the number of weights and biases for Φ in absolute value. When there exists $i \in \{1, \ldots, d\} : 0 < b_i - a_i < 2\hat{\delta}$, then we have $\hat{\Phi}_m\left(\boldsymbol{x}, \tilde{\Phi}_m(\boldsymbol{x})\right) =$ 0 and we can choose all first layer weights of $\tilde{\Phi}_m$ and $\hat{\Phi}_m$ as zero. Otherwise, we can bound the weights and biases of $\tilde{\Phi}_m$ and $\hat{\Phi}_m$ with $7\sqrt{C_1}$. In both cases we can bound the weights and biases for the first layer using $1 + 7\sqrt{C_1}$. In the second layer by (31) we have the bound $1 + 7\sqrt{C_1}$ for each of the neurons, and by (29) we get $1 + \hat{\delta}^{-1}$, since $|t_i^j(u)|/\hat{\delta} \leq 1 + \hat{\delta}^{-1}$ for all $j \in \{1, 2, 3, 4\}$. Lastly, using (30) we know that for the third layer, the weights and biases are bounded by $\max\{\delta^{-1}, \hat{\delta}^{-1}, d\}$. Then, the weights and biases of Φ are bounded in magnitude by

$$\max\left\{1+7\sqrt{C_1}, 1+\delta^{-\gamma/\alpha}, \max\{\delta^{-1}, \delta^{-\gamma/\alpha}, d\}\right\}$$

$$\leq \max\left\{ (1+7\sqrt{C_1}), 1+\delta^{-\gamma/\alpha}, \delta^{-1} \right\}$$

$$\leq (1+\sqrt{C_1}) \left(7+\delta^{-\gamma/\alpha}+\delta^{-1}\right)$$

$$\leq (1+\sqrt{C_1}) \left(7+N/C_1+(N/C_1)^{\gamma/\alpha}\right),$$

since $\delta^{-1} = \left(C_1\sqrt{(d-1)/N}\right)^{-1} = (N/(d-1))^{1/2}/C_1 \leq N/C_1.$

A.2 Proof of Theorem 3

First, we show the following result which is a consequence of [16, Theorem A.1].

Lemma 5. Assume that C^* is the Bayes classifier as in (9), and:

(i) For a positive sequence $\{a_n\}_{n\in\mathbb{N}}$, there exists a sequence of function classes $\{\mathcal{F}_n\}_{n\in\mathbb{N}}$ such that

$$\mathcal{E}_{\phi}(f_n, C^*) \le a_n$$

for some $f_n \in \mathcal{F}_n \subset \mathcal{F}$, where \mathcal{F} is as in (8).

(ii) Let $\boldsymbol{x} \in [0,1]^d$ and $y \in \{0,1\}$. Then, for all $f_n \in \mathcal{F}_n$ and $n \in \mathbb{N}$,

$$\mathbb{E}_{\boldsymbol{x}}\left[\left(\phi(2f_n(\boldsymbol{x}) - 1, 2y - 1) - \phi(2C^*(\boldsymbol{x}) - 1, 2y - 1)\right)^2\right] \le c_2 \mathcal{E}_{\phi}(f_n, C^*)$$

for a constant $c_2 > 0$ depending only on η and ϕ , where η and ϕ are defined in (1) and (10).

(iii) There exist $\hat{c}_3 > 0$ and $\{\hat{\delta}_n\}_{n \in \mathbb{N}} \subset \mathbb{R}^+$ such that

$$V_{[0,1]^d,\|\cdot\|_n}(\hat{\delta}_n) \le \hat{c}_3 n \hat{\delta}_n, \quad for \ all \quad n \in \mathbb{N}.$$

Then with these assumptions and taking $\epsilon^2 := 2 \max\{a_n, 2^8 \hat{\delta}_n\}$, it holds for some universal constant c > 0, that

$$\mu\left(\mathcal{E}_{\phi}(\widehat{f}_{\phi,S},C^*) \ge \epsilon_n^2\right) \lesssim \exp\left(-cn\epsilon_n^2\right).$$
(32)

Proof. Note that in [16] the labels are in the set $\{\pm 1\}$, however, in this result we take them in $\{0, 1\}$, for this reason in (9) and (10) the mapping 2x - 1 from [0, 1] to [-1, 1] is used. Moreover, the functions $f \in \mathcal{F}$ here are understood to have the form 2f - 1 in [16]. Now, we see that conditions from (A1) to (A5) in [16] are satisfied as follows.

- By the definition of hinge loss we know that (A1) is true since ϕ is Lipschitz with constant $c_1 = 1$.
- Condition (A2) turns into (i) when the labels are changed from $\{\pm 1\}$ to $\{0, 1\}$.
- We know that $\sup_{f \in \mathcal{F}_n} \|f\|_{\infty} \leq 1$ with \mathcal{F}_n as in (i), but also $\sup_{f \in \mathcal{F}_n} \|2f 1\|_{\infty} \leq 1$. Then, in the condition (A3) we can fix $\{F_n\}_{n \in \mathbb{N}}$ where $F_n \gtrsim 1$, as $F_n := 1$, for all $n \in \mathbb{N}$.
- In (A4) if we set v = 1, as in the previous item we fixed $F_n = 1$, then (*ii*) becomes condition (A4).

• In this case, condition (A5) states the following: There exists a sequence $\{\delta_n\}_{n\in\mathbb{N}}$ such that

$$H_B := H_B(\delta_n, 2\mathcal{F}_n - 1, \|\cdot\|_2) \le c_3 n \left(\frac{\delta_n}{F_n}\right)^{2-\nu} = c_3 n \delta_n, \tag{33}$$

for some constant $c_3 > 0$, with $F_n = 1$ and v = 1 as in the two items above. Here, to avoid unnecessary details, H_B is called the δ_n -bracketing entropy and is bounded by the inequality [16, (A.1)], that is

$$H_B \le \log \mathcal{N}\left(\delta_n/2, 2\mathcal{F}_n - 1, \|\cdot\|_{\infty}\right) := \log \inf \mathcal{G}$$
(34)

where

$$\mathcal{G} := \left\{ N \in \mathbb{N} : \text{ exist } f_1, \dots, f_N \text{ such that } 2\mathcal{F}_n - 1 \subset \bigcup_{i=1}^N B_\infty(f_i, \delta_n) \right\}$$

and $B_{\infty}(f_i, \delta_n) := \{f \in 2\mathcal{F}_n - 1 : \|f - f_i\|_{\infty} \leq \delta_n\}$. Moreover, if we set $K := [-1, 1]^d$ and $G_{\delta_n} := \bigcup_{i=1}^N B_{\infty}(f_i, \delta_n)$ in (12), we know by (*iii*) that

$$\log \mathcal{N}\left(\delta_{n}/2, 2\mathcal{F}_{n}-1, \|\cdot\|_{\infty}\right) \le V_{[-1,1]^{d}, \|\cdot\|_{\infty}}(\delta_{n}/2) \le V_{[0,1]^{d}, \|\cdot\|_{\infty}}(\delta_{n}/4) \le \hat{c}_{3}n(\delta_{n}/4),$$
(35)

where $\hat{\delta}_n := \delta_n/4$. Then, taking $\hat{c}_3 := 4c_3$ we obtain from inequalities (34) and (35) that inequality (33) is satisfied, that is, condition (A5) is fulfilled.

In conclusion, condition (A1)-(A5) are met and we fix $c_3 := (2^{17} \max \{5c_2, 128\})^{-1}$ and $\epsilon^2 := 2 \max\{a_n, 2^8 \hat{\delta}_n\}$. Then, Theorem A.1 in [16] implies inequality (32).

We now continue with the proof of Theorem 3. By definition we can choose $C^* := \mathbb{1}_{\Omega}$, therefore $\mathcal{E}(C^*) = 0$,

$$\phi(2C^*(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1) = \max\{0, 1 - (2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1)^2\} = 0$$
(36)

and $\mathcal{E}_{\phi}(C^*) = 0$. So, $\mathcal{E}_{\phi}(f_n, C^*) = \mathcal{E}_{\phi}(f_n)$ and $\mathcal{E}(f_n, C^*) = \mathcal{E}(f_n)$. Now, we show that all hypotheses of Lemma 5 are satisfied. Let $\mathcal{F}_n := \mathcal{NN}_*(d, N_n, W_n, B_n)$, then

• By Theorem 2 we know that for all $n \in \mathbb{N}$, there exists $f_n \in \mathcal{F}_n$ such that

$$\mu\left(\left\{\boldsymbol{x}\in[0,1]^{d}:\mathbb{1}_{\Omega}(\boldsymbol{x})\neq f_{n}(\boldsymbol{x})\right\}\right) \leq 3.5Md(d-1)^{\gamma/2}C_{1}^{\gamma}\widehat{N}_{n}^{-\gamma/2}\max\{C_{2},C_{3}\} \\ \leq n^{-\gamma/(2+\gamma)}/2,$$

and this implies

$$\begin{split} \mathcal{E}_{\phi}(f_{n}) &= \mathbb{E}_{\boldsymbol{x}} \left[\phi(2f_{n}(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega} - 1) \right] \\ &= \int_{[0,1]^{d}} \max\{0, 1 - (2f_{n}(\boldsymbol{x}) - 1)(2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1)\} \, d\mu \\ &= \int_{\{\mathbb{1}_{\Omega}(\boldsymbol{x}) \neq f_{n}(\boldsymbol{x})\}} \max\{0, 1 - (2f_{n}(\boldsymbol{x}) - 1)(2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1)\} \, d\mu \\ &\leq \int_{[0,1]^{d}} 2\mathbb{1}_{\mathbb{1}_{\Omega}(\boldsymbol{x}) \neq f_{n}(\boldsymbol{x})} \, d\mu = 2\mu \left(\left\{ \boldsymbol{x} \in [0,1]^{d} : \mathbb{1}_{\Omega}(\boldsymbol{x}) \neq f_{n}(\boldsymbol{x}) \right\} \right) \leq n^{-\gamma/(2+\gamma)}, \end{split}$$

i.e, condition (i) is fulfilled with $a_n := n^{-\gamma/(2+\gamma)}$.

• Let $G = \{ \boldsymbol{x} \in [0,1]^d : \phi(2f_n(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1) \le 1 \}$. Using identity (36), we have

$$\mathbb{E}_{\boldsymbol{x}} \left[\left(\phi(2f_n(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1) - \phi(2C^*(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1) \right)^2 \right] \\
= \mathbb{E}_{\boldsymbol{x}} \left[\left(\phi(2f_n(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1) \right)^2 \right] \qquad (37)$$

$$= \int_{\mathbb{R}^d} \left(\phi(2f_n(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1) \right)^2 d\mu \\
\leq \int_G \phi(2f_n(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1) d\mu + \int_{G^c} 4d\mu \\
\leq \mathbb{E}_{\boldsymbol{x}} \left[\phi(2f_n(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1) \right] + 4\mu(G^c) \leq \mathcal{E}_{\phi}(f_n) + 4\mathbb{E}_{\boldsymbol{x}} \left[C_{f_n}(\boldsymbol{x}) \neq y \right] \\
\leq \mathcal{E}_{\phi}(f_n) + 4\mathcal{E}(f_n) \qquad (38)$$

since $|\phi(2f_n(\boldsymbol{x}) - 1, 2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1)| \le |1 - (2f_n(\boldsymbol{x}) - 1)(2\mathbb{1}_{\Omega}(\boldsymbol{x}) - 1)| \le 2$ and $x \in G^c$ implies

$$\phi(2f_n(\boldsymbol{x})-1,2\mathbb{1}_{\Omega}(\boldsymbol{x})-1)>1, \quad ext{that is} \quad (2f_n(\boldsymbol{x})-1)(2\mathbb{1}_{\Omega}(\boldsymbol{x})-1)<0,$$

i.e. $C_{f_n}(\boldsymbol{x}) \neq y$. Then, by inequalities (11) and (38),

$$\mathbb{E}_{\boldsymbol{x}}\left[\left(\phi(2f_n(\boldsymbol{x})-1,2\mathbb{1}_{\Omega}(\boldsymbol{x})-1)\right)^2\right] \leq \mathcal{E}_{\phi}(f_n) + 4\mathcal{E}(f_n) \leq (1+4C_{\phi})\mathcal{E}_{\phi}(f_n)$$

and condition (*ii*) with $c_2 := 1 + 4C_{\phi}$ is satisfied.

• Using Lemma 1 with $\delta = \hat{\delta}_n := a_n(1 + \log n)$ we obtain

$$V_{[0,1]^d,\|\cdot\|_{\infty}}(\hat{\delta}_n) \le W_n \cdot \left(10 + \log(1/\hat{\delta}_n) + 5\log(\lceil B_n \rceil) + 5\log(\max\{d, W_n\})\right)$$

$$\lesssim \widehat{N}_n(1 + \log n) \lesssim n \cdot n^{2/(2+\gamma)-1}(1 + \log n) = n\hat{\delta}_n, \quad \text{for all} \quad n \in \mathbb{N}.$$

So, there exists $\hat{c}_3 > 0$ such that

$$V_{[0,1]^d, \|\cdot\|_n}(\hat{\delta}_n) \le \hat{c}_3 n \hat{\delta}_n, \text{ for all } n \in \mathbb{N}$$

and condition (iii) is fulfilled.

Then, we use Lemma 5 with

$$\epsilon_n^2 = 2 \max\{a_n, 2^8 a_n (1 + \log n)\} = 2^9 n^{-\gamma/(2+\gamma)} (1 + \log n)$$

and we get

$$\mu\left(\mathcal{E}_{\phi}(\widehat{f}_{\phi,S}) \gtrsim n^{-\gamma/(2+\gamma)}(1+\log n)\right) \leq \mu\left(\mathcal{E}_{\phi}(\widehat{f}_{\phi,S}) \geq \epsilon_{n}^{2}\right)$$
$$\lesssim \exp\left(-n^{1-\gamma/(2+\gamma)}(1+\log n)\right) \lesssim n^{-1}. \tag{39}$$

Finally, by inequalities (11) and (39) we conclude (16).