

TRIVIALITY, ROTATIONAL SYMMETRY, AND CLASSIFICATION OF COMPLETE EXPANDING GRADIENT YAMABE SOLITONS

SHUN MAETA

ABSTRACT. In this paper, we rigorously analyze the scalar curvature of complete expanding gradient Yamabe solitons. We show that if the scalar curvature is less than the soliton constant by an arbitrarily small positive amount, then the soliton is trivial. Additionally, we show that if the scalar curvature is greater than the soliton constant by an arbitrarily small positive amount, then the soliton is rotationally symmetric. Furthermore, we completely classify nontrivial complete expanding gradient Yamabe solitons in both cases: when the scalar curvature is greater than the soliton constant and when it is less than the soliton constant.

1. INTRODUCTION

The Yamabe flow is a fundamental and highly effective tool for analyzing the structure of manifolds. The study of the Yamabe flow has developed rapidly in the last decades and has become a central area in geometry (cf. [2], [3], [6], [8], [11], [17], [19]). To analyze the Yamabe flow, and more generally, geometric flows, it is important to study singularity models. Yamabe solitons are special solutions to the Yamabe flow and are expected to serve as singularity models (cf. [8]). From this perspective, Yamabe solitons have also been studied extensively over the last few decades (cf. [4], [5], [9], [13], [14]).

An n -dimensional Riemannian manifold (M^n, g) is called a gradient Yamabe soliton if there exists a smooth function F on M and a constant $\lambda \in \mathbb{R}$, such that

$$\nabla \nabla F = (R - \lambda)g.$$

A gradient Yamabe soliton is represented as (M^n, g, F, λ) . If F is constant, then the gradient Yamabe soliton is called trivial. If $\lambda > 0$, $\lambda = 0$, or $\lambda < 0$, then the Yamabe soliton is called shrinking, steady, or expanding.

It is well known that there exist no nontrivial compact gradient Yamabe solitons (cf. [11]). One of the main problems in the study of Yamabe solitons is determining whether complete gradient Yamabe solitons are rotationally symmetric. In particular, the Yamabe soliton version of Perelman's conjecture, which states that "any nontrivial complete steady gradient Yamabe soliton with positive scalar curvature is rotationally symmetric," has stimulated significant interest in this problem. Partial answers have been obtained under the assumption of local conformal flatness (cf. [4], [9]). Catino, Mantegazza and Mazzieri also provided a partial answer to the conjecture under the assumption that $\text{Ric}(\nabla F, \nabla F) \geq 0$

2010 *Mathematics Subject Classification.* 53C21, 53C25, 53C20.

Key words and phrases. Expanding gradient Yamabe solitons; Yamabe flow; Scalar curvature.

The author is partially supported by the Grant-in-Aid for Scientific Research (C), No.23K03107, Japan Society for the Promotion of Science.

and $\text{Ric}(\nabla F, \nabla F) > 0$ holds at least at one point (cf. [5]). Here, Ric is the Ricci tensor of M . Finally, the author resolved the conjecture (cf. [13]). Furthermore, the author also proved the rotational symmetry of the complete shrinking gradient Yamabe solitons under $R > \lambda$, and showed that this assumption is optimal. Therefore, the remaining problem involves expanding solitons. However, the study of expanding solitons has been regarded as difficult so far. In fact, even in low-dimensional cases, expanding gradient Yamabe solitons exhibit richer geometry than steady or shrinking solitons (cf. [1], [14]), making it difficult to obtain rigidity results. To overcome this difficulty, we reduce the problem to the existence and nonexistence of solutions of a certain ODE, using the warped product structure provided by Tashiro's theorem (cf. [18], see also [12]).

Theorem 1.1 ([18], [12]). *Let (M^n, g, F, φ) be a complete Riemannian manifold which satisfies*

$$\varphi g = \nabla \nabla F$$

for smooth functions F and φ on M . Then (M^n, g, F, φ) is either

- (1) compact and rotationally symmetric, or
- (2) rotationally symmetric and equal to the warped product

$$([0, \infty), dr^2) \times_{|\nabla F|} (\mathbb{S}^{n-1}, \bar{g}_S),$$

where \bar{g}_S is the round metric on \mathbb{S}^{n-1} , or

- (3) the warped product

$$(\mathbb{R}, dr^2) \times_{|\nabla F|} (N^{n-1}, \bar{g}).$$

In Theorem 1.1, if $\varphi = R - \lambda$, then (M^n, g, F, λ) is a gradient Yamabe soliton. Tashiro's theorem is really powerful for studying complete gradient Yamabe solitons. In fact, by the proof of Theorem 1.1 (see [12]), it is easy to see that if $\text{Ric}(\nabla F, \nabla F) \geq 0$ and $\text{Ric}(\nabla F, \nabla F) > 0$ holds at least at one point, then any such complete Riemannian manifold (M, g, F, φ) with a non-constant function F is rotationally symmetric. Therefore, in the study of the rotational symmetry of complete expanding gradient Yamabe solitons, imposing conditions on the scalar curvature is of fundamental importance. To analyze the scalar curvature in cases (2) and (3) of Theorem 1.1, the following equations, derived from the structure of the warped product, are used.

$$|\nabla F|^2 R = \bar{R} - (n-1)(n-2)(R-\lambda)^2 - 2(n-1)g(\nabla F, \nabla R).$$

Here, we note that F depends only on r (cf. [18], see also [12]). Set $\rho = F'$. Then we have

$$(1.1) \quad 2(n-1)\rho\rho'' + (n-1)(n-2)\rho'^2 + \rho^2(\rho' + \lambda) = \bar{R}.$$

Hence, the scalar curvature \bar{R} of N is constant. Differentiating both sides of (1.1),

$$(1.2) \quad 2(n-1)^2\rho'\rho'' + 2(n-1)\rho\rho''' + 2\rho\rho'(\rho' + \lambda) + \rho^2\rho'' = 0.$$

2. EXAMPLES, TRIVIALITY AND ROTATIONAL SYMMETRY

To study rotational symmetry of complete expanding gradient Yamabe solitons, we first note that there exist nontrivial examples when $R = \lambda$ (cf. [13]).

Example. Let (N^{n-1}, \bar{g}) be an $(n-1)$ -dimensional complete Riemannian manifold with constant negative scalar curvature \bar{R} . Then, for any $\alpha \in \mathbb{R}$, $(M, g, F, \lambda) = (\mathbb{R} \times N^{n-1}, dr^2 + \frac{\bar{R}}{\lambda} \bar{g}, \sqrt{\frac{\bar{R}}{\lambda}} r + \alpha, \lambda)$ is an n -dimensional complete expanding gradient Yamabe soliton with $R = \lambda$.

In particular, $(M^3, g, F, \lambda) = (\mathbb{R} \times \mathbb{H}^2, dr^2 + \frac{\bar{R}}{\lambda} \bar{g}_H, \sqrt{\frac{\bar{R}}{\lambda}} r + \alpha, \lambda)$ is a 3-dimensional complete expanding gradient Yamabe soliton with $R = \lambda$, where $(\mathbb{H}^2, \bar{g}_H)$ is a hyperbolic surface.

Therefore, imposing conditions on the scalar curvature is crucial and fundamental for achieving rotational symmetry in complete expanding gradient Yamabe solitons. In particular, the problem reduces to the cases $R > \lambda$ or $R < \lambda$.

Interestingly, the following triviality result for expanding, steady, and shrinking solitons can be obtained for any $\varepsilon > 0$ provided that $R < \lambda - \varepsilon$.

Proposition 2.1. *If $R < \lambda - \varepsilon$ for some $\varepsilon > 0$, then there exists no nontrivial complete gradient Yamabe solitons.*

Proof. If M is compact, it is trivial. Hence, we consider the other cases. If $R < \lambda - \varepsilon$, then one has that the positive smooth function ρ satisfies $\rho' < -\varepsilon$. Therefore, we have a contradiction. □

On the other hand, if $R > \lambda + \varepsilon$, then the soliton is rotationally symmetric.

Proposition 2.2. *Let (M, g, F, λ) be a nontrivial complete expanding gradient Yamabe soliton. If $R > \lambda + \varepsilon$ for some $\varepsilon > 0$, then it is rotationally symmetric and equal to the warped product*

$$([0, \infty), dr^2) \times_{|\nabla F|} (\mathbb{S}^{n-1}, \bar{g}_S).$$

Proof. Since there exists no compact gradient Yamabe solitons, we only have to consider (3) of Theorem 1.1. Since $\rho' > \varepsilon$ and $\rho > 0$, we have a contradiction. □

As a corollary, we obtain rotational symmetry of complete expanding gradient Yamabe solitons with positive scalar curvature.

Corollary 2.3. *Any nontrivial complete expanding gradient Yamabe soliton with positive scalar curvature is rotationally symmetric.*

This generalizes Theorem 1.3 in [9], Corollary 3.3 in [5] and Corollary 1.6 in [4]. In particular, we remove the assumption of local conformal flatness.

The remaining cases are $R > \lambda$ and $R < \lambda$.

3. CASE $R < \lambda$

In this section, we consider the case $R < \lambda$, and completely classify complete expanding gradient Yamabe solitons with $R < \lambda$.

Proposition 3.1. *A complete nontrivial, expanding gradient Yamabe soliton (M^n, g, F, λ) with $R < \lambda$ is either*

(1) *the warped product*

$$(\mathbb{R}, dr^2) \times_{|\nabla F|} (N^{n-1}(0), \bar{g}),$$

where $(N^{n-1}(0), \bar{g})$ has flat scalar curvature, and the scalar curvature of M tends to λ at infinity, or

(2) *the warped product*

$$(\mathbb{R}, dr^2) \times_{|\nabla F|} (N^{n-1}(-c^2), \bar{g}),$$

where $(N^{n-1}(-c^2), \bar{g})$ has negative constant scalar curvature, and the scalar curvature of M tends to λ at infinity.

Furthermore, the potential function F satisfies $F''(r) < 0$, $F'''(r) > 0$ and $F'(r) \searrow \sqrt{\frac{\bar{R}}{\lambda}}$ as $r \nearrow +\infty$, where \bar{R} is the scalar curvature of N .

Proof. By the assumption and the soliton equation, $\rho(r) = F'(r)$ is monotone decreasing. Since ρ is positive, $\rho \searrow c (\geq 0)$, $\rho'(r) \nearrow 0$ and $\rho''(r) \searrow 0$ as $r \nearrow +\infty$. By the soliton equation again, one has $\lambda \lim_{r \nearrow +\infty} \rho^2(r) = \bar{R}$. Hence, $\bar{R} \leq 0$. Therefore, (2) of Theorem 1.1 cannot happen, and for (3) of Theorem 1.1, N has constant nonpositive scalar curvature.

Assume that there exists r_0 , such that $\rho''(r_0) = 0$. By (1.1) and the assumption, one has

$$\rho'''(r_0) = -\frac{1}{n-1} \rho'(r_0)(\rho'(r_0) + \lambda) < 0.$$

This indicates that there exists no open interval on which $\rho'' = 0$.

Assume that $\rho'' \leq 0$ on $(-\infty, r_0)$. In this case, we have $\rho'' \leq 0$ on \mathbb{R} . Since ρ is positive, we have that ρ is constant and $R = \lambda$, which is a contradiction. Hence, there must exist a subinterval $\Omega \subset (-\infty, r_0)$ on which $\rho'' > 0$ holds. If necessary, let us redefine r_0 to be the first point, such that $\rho''(r_0) = 0$, and we may assume that $\rho'' > 0$ on $(-\infty, r_0)$. In this case, r_0 is the maximum point of ρ' . Set $C = \rho'(r_0)$. Therefore, ρ satisfies $\rho'(r) \leq C (< 0)$ on \mathbb{R} . By Proposition 2.1, we have a contradiction. Therefore, we have $\rho'' > 0$ or $\rho'' < 0$ on \mathbb{R} . However, the second case cannot happen. Since $\rho > 0$, $\rho' < 0$, and $\rho'' < 0$, we have $\rho' \nearrow +\infty$. □

4. CASE $R > \lambda$

We finally consider the case $R > \lambda$. Interestingly, in this case, for (3) of Theorem 1.1, the scalar curvature of M must be negative:

Theorem 4.1. *Let (M^n, g, F, λ) be a complete nontrivial, expanding gradient Yamabe soliton with $R > \lambda$. Assume that F has no critical point, then it is the warped product*

$$(\mathbb{R}, dr^2) \times_{|\nabla F|} (N^{n-1}, \bar{g}),$$

where, N has constant scalar curvature \bar{R} . Furthermore, $\bar{R} \leq 0$ and $\lambda < R < 0$. Moreover, the potential function F satisfies $\lim_{r \rightarrow -\infty} F'(r) = \sqrt{\frac{\bar{R}}{\lambda}}$, $F''(r) > 0$ and $F'''(r) \geq 0$ on \mathbb{R} . However, there exists no open interval on which $F'''(r) = 0$.

As a corollary, one can classify expanding gradient Yamabe solitons with $R > \lambda$.

Corollary 4.2. *A complete nontrivial, expanding gradient Yamabe soliton (M^n, g, F, λ) with $R > \lambda$ is either*

- (1) *rotationally symmetric and equal to the warped product*

$$([0, +\infty), dr^2) \times_{|\nabla F|} (\mathbb{S}^{n-1}, \bar{g}_S),$$

where, \bar{g}_S is the round metric on \mathbb{S}^{n-1} , or

- (2) *the warped product*

$$(\mathbb{R}, dr^2) \times_{|\nabla F|} (N^{n-1}(0), \bar{g}),$$

which satisfies $\lambda < R < 0$, where $(N^{n-1}(0), \bar{g})$ has flat scalar curvature, or

- (3) *the warped product*

$$(\mathbb{R}, dr^2) \times_{|\nabla F|} (N^{n-1}(-c^2), \bar{g}),$$

which satisfies $\lambda < R < 0$, where $(N^{n-1}(-c^2), \bar{g})$ has negative constant scalar curvature.

Furthermore, in the cases (2) and (3), the potential function F satisfies $\lim_{r \rightarrow -\infty} F'(r) = \sqrt{\frac{\bar{R}}{\lambda}}$, $F''(r) > 0$ and $F'''(r) \geq 0$ on \mathbb{R} . However, there exists no open interval on which $F'''(r) = 0$.

Proof of Theorem 4.1. Assume that $\bar{R} > 0$. Since $\rho' > 0$, the positive function ρ is monotone increasing. Therefore, $\rho \searrow \alpha$ for some α , $\rho' \searrow 0$ and $\rho'' \nearrow 0$ as $r \searrow -\infty$. However, this contradicts (1.1). Therefore, one has $\bar{R} \leq 0$. Under the assumption the following claim holds.

Claim 1. If there exists a point $r_0 \in \mathbb{R}$, such that $\rho''(r_0) = 0$, then $\rho'''(r_0) > 0$. Furthermore, we also have $\sqrt{\frac{\bar{R}}{\lambda}} < \rho(r_0)$.

Proof of Claim 1.

Case 1. $\bar{R} = 0$. Assume that $\rho''(r_0) = 0$. By (1.1),

$$(n-1)(n-2)\rho'^2(r_0) + \rho^2(r_0)\rho'(r_0) = -\lambda\rho^2(r_0).$$

Hence, one has

$$\rho'(r_0) = \frac{-\rho^2(r_0) \pm \sqrt{\rho^4(r_0) - 4(n-1)(n-2)\lambda\rho^2(r_0)}}{2(n-1)(n-2)}.$$

Since $\rho' > 0$, it must be

$$(4.1) \quad \rho'(r_0) = \frac{-\rho^2(r_0) + \sqrt{\rho^4(r_0) - 4(n-1)(n-2)\lambda\rho^2(r_0)}}{2(n-1)(n-2)}.$$

Furthermore, by (1.2),

$$\rho'''(r_0) = -\frac{1}{n-1}\rho'(r_0)(\rho'(r_0) + \lambda).$$

Since $\rho' > 0$, we have 3 cases.

(1) Assume that $-\lambda < \rho'(r_0)$. Combining this inequality with (4.1), we have

$$-\lambda < \frac{-\rho^2(r_0) + \sqrt{\rho^4(r_0) - 4(n-1)(n-2)\lambda\rho^2(r_0)}}{2(n-1)(n-2)}.$$

A straightforward computation leads to a contradiction.

(2) Assume that $-\lambda = \rho'(r_0)$. A similar argument shows that $\lambda = 0$, which is a contradiction.

Hence, the inequality $-\lambda > \rho'(r_0)$ holds and one has $\rho'''(r_0) > 0$.

Case 2. $\bar{R} < 0$. Assume that $\rho''(r_0) = 0$. By (1.1),

$$(n-1)(n-2)\rho'^2(r_0) + \rho^2(r_0)\rho'(e_0) = \bar{R} - \lambda\rho^2(r_0).$$

Hence, one has

$$\rho^4(r_0) - 4\lambda(n-1)(n-2)\rho^2(r_0) + 4(n-1)(n-2)\bar{R} > 0,$$

and

$$\rho'(r_0) = \frac{-\rho^2(r_0) \pm \sqrt{\rho^4(r_0) + 4(n-1)(n-2)(\bar{R} - \lambda\rho^2(r_0))}}{2(n-1)(n-2)}.$$

Since $\rho' > 0$, one has

$$\sqrt{\frac{\bar{R}}{\lambda}} < \rho(r_0),$$

and

$$\rho'(r_0) = \frac{-\rho^2(r_0) + \sqrt{\rho^4(r_0) + 4(n-1)(n-2)(\bar{R} - \lambda\rho^2(r_0))}}{2(n-1)(n-2)}.$$

Furthermore, by (1.2),

$$\rho'''(r_0) = -\frac{1}{n-1}\rho'(r_0)(\rho'(r_0) + \lambda).$$

Since $\rho' > 0$, the similar argument shows that $\rho'''(r_0) > 0$. □

Assume that there exists $r_1 \in \mathbb{R}$, such that $\rho'(r_1) = -\lambda$. By (1.1), we have $\rho''(r_1) < 0$. Hence, there exists no point $r_{-1} \in (-\infty, r_1)$, such that $\rho''(r_{-1}) = 0$. In fact, if there exists such a point, then by the claim, $\rho'''(r_{-1}) > 0$ and the function ρ' is weakly monotone increasing on $(r_{-1}, +\infty)$. By the same reason, it cannot happen that $\rho'' > 0$ on $(-\infty, r_1)$. Therefore, we have $\rho'' < 0$ on $(-\infty, r_2)$ for some $r_2 (> r_1)$. However, since $\rho > 0$, $\rho' > 0$ and $\rho'' < 0$ on $(-\infty, r_2)$, we have a contradiction. Hence, such a point r_1 does not exist, and one has $\rho' > -\lambda$ or $\rho' < -\lambda$ on \mathbb{R} . By Proposition 2.2, the first case cannot happen. Therefore, $\lambda < R < 0$.

Claim 1 shows that for each critical point r_0 of ρ' , the function ρ' is weakly monotone increasing on $(r_0, +\infty)$. Assume that ρ' has a critical point r_0 , and furthermore, if there exists an interval on which ρ' is decreasing, then without loss of generality, we may assume that it is $(-\infty, r_0)$. However, by the same argument as before, we have a contradiction. Therefore, ρ' is weakly monotone increasing on \mathbb{R} . If there exists no critical point of ρ' , we have $\rho'' > 0$ or $\rho'' < 0$ on \mathbb{R} . However, $\rho'' < 0$ cannot happen. Hence, one has $\rho'' > 0$.

Therefore, $\rho'' \geq 0$ on \mathbb{R} and there exists no open interval on which $\rho'' = 0$. Since $\rho > 0$ and $\rho' > 0$, $\rho \searrow c$ for some $c \in \mathbb{R}$, $\rho' \searrow 0$ and $\rho'' \searrow 0$ as $r \searrow -\infty$. By (1.1), we have

$$\bar{R} = \lambda \lim_{r \rightarrow -\infty} \rho^2(r).$$

Therefore, $F'(r) > \sqrt{\frac{\bar{R}}{\lambda}}$ on \mathbb{R} . □

REFERENCES

1. J. Bernstein and T. Mettler, *Two-Dimensional Gradient Ricci Solitons Revisited*, Int. Math. Res Not., (2015), **2015**, 78–98.
2. S. Brendle, *Convergence of the Yamabe flow for arbitrary initial energy*, J. Differential Geom., (2005), **69**, 217–278.
3. S. Brendle, *Convergence of the Yamabe flow in dimension 6 and higher*, Invent. Math., (2007), **170**, 541–576.
4. H.-D. Cao, X. Sun and Y. Zhang, *On the structure of gradient Yamabe solitons*, Math. Res. Lett., (2012) **19**, 767–774.
5. G. Catino, C. Mantegazza and L. Mazzieri, *On the global structure of conformal gradient solitons with nonnegative Ricci tensor*, Commun. Contemp. Math., (2012) **14**, 12pp.
6. B. Chow, *The Yamabe flow on locally conformally flat manifolds with positive Ricci curvature*, Comm. Pure Appl. Math., (1992) **45**, 1003–1014.
7. B. Chow, S.-C. Chu, D. Glickenstein, C. Guenther, J. Isenberg, T. Ivey, D. Knopf, P. Lu, F. Luo and L. Ni, *The Ricci Flow: Techniques and Applications: Part I: Geometric Aspects*, Math. Surv. and Mono., Amer. Math. Soc., (2007), **135**.
8. B. Chow, P. Lu and L. Ni, *Hamilton's Ricci Flow*, Graduate Studies in Mathematics, **77**, Amer. Math. Soc., (2006).
9. P. Daskalopoulos and N. Sesum, *The classification of locally conformally flat Yamabe solitons*, Adv. Math., (2013) **240**, 346–369.
10. R. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom., (1982) **17**, 255–306.
11. R. Hamilton, *Lectures on geometric flows*, (1989), unpublished.
12. S. Maeta, *Classification of generalized Yamabe solitons*, arXiv:2107.05487[math DG].
13. S. Maeta, *Complete steady gradient Yamabe solitons with positive scalar curvature are rotationally symmetric*, arXiv:2309.09166[math DG].
14. S. Maeta, *Classification of low-dimensional complete gradient Yamabe solitons*, arXiv:2405.03921[math DG].
15. G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv math.DG/0211159, (2002).
16. P. Petersen, *Riemannian Geometry, Third edition*, Graduate Texts in Mathematics, (2016), **171**, Springer.
17. H. Schwetlick and M. Struwe, *Convergence of the Yamabe flow for large energies*, J. Reine Angew. Math., (2003), **562**, 59–100.
18. Y. Tashiro, *Complete Riemannian manifolds and some vector fields*, Trans. Amer. math. Soc., (1965), **117**, 251–275.
19. R. Ye, *Global existence and convergence of Yamabe flow* J. Differential Geom., (1994) **39**, 35–50.

DEPARTMENT OF MATHEMATICS, CHIBA UNIVERSITY, 1-33, YAYOICHO, INAGE, CHIBA, 263-8522, JAPAN.
 Email address: shun.maeta@faculty.gs.chiba-u.jp or shun.maeta@gmail.com