

# Criticality, splitting theorems under spectral Ricci bounds and the topology of stable minimal hypersurfaces

Giovanni Catino, Luciano Mari, Paolo Mastrolia, Alberto Roncoroni

*To the memory of Francesco "Franco" Mercuri and Celso Viana*

## Abstract

In this paper we prove general criticality criteria for operators  $\Delta + V$  on manifolds  $M$  with two ends, where  $V$  bounds the Ricci curvature of  $M$ , and a related spectral splitting theorem. We apply them to get new rigidity results for stable and  $\delta$ -stable minimal hypersurfaces in manifolds with non-negative bi-Ricci or sectional curvature, in ambient dimension up to 5 and 6, respectively. In the special case where the ambient space is  $\mathbb{R}^4$ , we prove that a  $1/3$ -stable minimal hypersurface must either have one end or be a catenoid, and consequently that a proper,  $\delta$ -stable minimal hypersurface with  $\delta > 1/3$  must be a hyperplane.

**MSC2020:** 53C24, 53C21, 53C42.

**Keywords:** Criticality, splitting, stable minimal hypersurfaces, spectral Ricci bounds, catenoid.

## 1 Introduction

The purpose of the present paper is to investigate the topological properties of complete, two-sided minimal hypersurfaces  $M^n \rightarrow N^{n+1}$  immersed into a non-negatively curved space. Hereafter, we always implicitly assume that  $M$  is connected. We will be interested in hypersurfaces of dimension  $n \geq 3$  which are stable or, more generally,  $\delta$ -stable, where  $\delta$ -stability means that the modified Jacobi operator

$$J_\delta \doteq \Delta + \delta \left( |A|^2 + \overline{\text{Ric}}(\nu, \nu) \right).$$

is non-negative<sup>1</sup>. Here,  $\overline{\text{Ric}}$  is the Ricci curvature of  $N$ ,  $\nu$  is a global unit normal vector field on  $M$  and  $A$  is the second fundamental form in direction  $\nu$ . "Classical" stability corresponds to  $\delta = 1$ , and one easily checks that  $\delta$ -stability implies  $\delta'$ -stability if  $\delta' < \delta$ . We shall focus on the range

$$\delta \in \left[ \frac{n-2}{n}, 1 \right].$$

---

<sup>1</sup>We follow the convention that  $\Delta = d^2/dx^2$  on  $\mathbb{R}$ , and that  $J_\delta \geq 0$  corresponds to the non-negativity of  $-(\phi, J_\delta \phi)_{L^2}$  for each test function  $\phi \in C_c^\infty(M)$ .

If  $N = \mathbb{R}^{n+1}$ , the recent solution to the stable Bernstein problem by Chodosh & Li [21, 22] ( $n = 3$ ), Catino, Mastrolia & Roncoroni [16] ( $n = 3$ ), Chodosh, Li, Minter & Stryker [24] ( $n = 4$ ) and Mazet [43] ( $n = 5$ ), and the results by Bellettini [6] for  $n = 6$ , provided new powerful tools to investigate the geometry of stable minimal hypersurfaces, with promising applications to treat the problem in more general ambient spaces. Considering the weaker notion of  $\delta$ -stability is natural in view of the following result:

**Theorem 1.1** ([3, 59, 31, 46]). *Let  $M^n \rightarrow \mathbb{R}^{n+1}$  be a complete, two-sided minimal hypersurface with finite total curvature, i.e.  $|A| \in L^n(M)$ . Then:*

- $M$  is  $\delta$ -stable with  $\delta > \frac{n-2}{n}$  if and only if  $M$  is a hyperplane;
- $M$  is  $\frac{n-2}{n}$ -stable if and only if  $M$  is either a hyperplane or a catenoid.

The theorem is due to Anderson [3] and Shen & Zhu [59] when  $\delta = 1$ , and to Fu & Li [31] in the present form. An alternative, self-contained proof is due to Pigola & Veronelli [46], which we also recommend for a detailed account of the literature. The  $\frac{n-2}{n}$ -stability of the higher dimensional catenoid and the finiteness of its total curvature were proved by Tam & Zhou in [60].

It is tempting to ask whether Theorem 1.1 holds in dimension  $n \leq 6$  (or in a smaller range) without assuming that  $M$  has finite total curvature. In [22], the authors obtained a first relevant step in this direction in dimension  $n = 3$  by proving that  $\delta$ -stability with  $\delta > 2/3$  implies flatness. The question was also addressed in the recent [33], a source of various interesting results in the full range  $\delta > \frac{n-2}{n}$ . As a direct consequence of [33, Theorem 1.7] and our Theorem 1.4 below, we get:

**Theorem 1.2.** *Let  $M^3 \rightarrow \mathbb{R}^4$  be a two-sided, properly immersed minimal hypersurface which is  $\delta$ -stable for some  $\delta > \frac{1}{3}$ . Then,  $M$  is a hyperplane.*

A bound on the number of ends of  $\delta$ -stable hypersurfaces is, in fact, the missing information we provide to get Theorem 1.2. In this regard, it is known by [10] that stable minimal hypersurfaces in  $\mathbb{R}^{n+1}$  must have only one end, a fact that plays a crucial role in the proofs in [22, 24, 43]. However, although the argument in [10] easily adapts to  $\delta$ -stability for any

$$\delta > \frac{n-1}{n},$$

justifying the bound  $\delta > 2/3$  in [22], it fails for  $\delta < \frac{n-1}{n}$ . To the best of our knowledge, in full generality no bound on the number of ends of  $M$  was established so far for  $\delta$  close to  $\frac{n-2}{n}$ , and neither the assertion that  $M$  has finitely many ends. We are only aware of the following result due to Cheng & Zhou, [20, Theorem 1.1]:

**Theorem 1.3** ([20]). *Let  $M^n \rightarrow \mathbb{R}^{n+1}$  be a  $\frac{n-2}{n}$ -stable, two-sided complete minimal hypersurface of dimension  $n \geq 3$ . If*

$$\begin{aligned} \lim_{R \rightarrow \infty} \left( \sup_{B_R(o)} |A| \right) / \log R = 0 & \quad \text{if } n = 3 \\ \text{or} & \\ \lim_{R \rightarrow \infty} \left( \sup_{B_R(o)} |A| \right) / R^{\frac{n-3}{2}} = 0 & \quad \text{if } n \geq 4, \end{aligned} \tag{1.1}$$

where  $B_R(o) \subset M$  is the intrinsic ball centered at  $o$  with radius  $R$ , then either  $M$  has only one end or it is a catenoid.

Observe that (1.1) holds, for instance, if  $|A|$  is bounded on  $M$ . Cheng and Zhou kindly suggested to the first and second author the question whether (1.1) is removable, which indeed was the starting point of the present investigation. We are able to positively answer their question in dimension  $n = 3$ :

**Theorem 1.4.** *Let  $M^3 \rightarrow \mathbb{R}^4$  be a complete, two-sided, 1/3-stable minimal hypersurface. Then, either  $M$  has only one end or it is a catenoid.*

Theorem 1.4 is a corollary of a more general result, Theorem 4.1 below, where we study the topology of 1/3-stable minimal hypersurfaces  $M^3 \rightarrow N^4$  into a 4-manifold with non-negative sectional curvature. The peculiarity of dimension  $n = 3$  will be clear in what follows, see in particular Theorem 3.3.

We focus on ambient spaces  $(N^{n+1}, \bar{g})$  with non-negative sectional curvature  $\overline{\text{Sec}}$  or, more generally, non-negative bi-Ricci curvature  $\overline{\text{BRic}}$ . Recall that, given an orthonormal set  $\{X, Y\} \subset T_p N$ , the bi-Ricci curvature is defined as

$$\overline{\text{BRic}}(X, Y) = \overline{\text{Ric}}(X, X) + \overline{\text{Ric}}(Y, Y) - \overline{\text{Sec}}(X, Y).$$

In particular, if  $n + 1 = 3$  then  $\overline{\text{BRic}} = \bar{R}/2$ , with  $\bar{R}$  the scalar curvature of  $M$ . Compared to the literature on minimal surfaces in non-negatively curved 3-manifolds, relatively few results on the topology of higher dimensional minimal hypersurfaces appeared so far. The list below collects the ones which are closer to the present paper. Let  $M^n \rightarrow N^{n+1}$  be a complete, non-compact, two-sided stable minimal immersion of dimension  $n \geq 3$ . Assume for simplicity that  $N$  is complete as well, although part of the theorems in the list do not need it.

- Li & Wang [37, Thm. 0.1]: if  $\overline{\text{Sec}} \geq 0$  and  $M$  is properly immersed, then either  $M$  has only one end or  $M$  is totally geodesic and  $M = \mathbb{R} \times P$  with the product metric, for some compact  $P$  with non-negative sectional curvature.
- Cheng [18, Thm. 3.2]: if  $\overline{\text{Sec}} \geq 0$  and  $\overline{\text{Ric}} > 0$ , then  $M$  is non-parabolic and it has only one end.
- Chodosh, Li & Stryker [23, Thm. 1.3 and Remark 1.4]: if  $n = 3$ ,  $0 \leq \overline{\text{Sec}} \leq \kappa$  for some  $\kappa \in \mathbb{R}^+$  and the scalar curvature of  $N$  satisfies  $\bar{R} \geq 1$ , then  $M$  is totally geodesic and  $\overline{\text{Ric}}(v, v) \equiv 0$ .
- Tanno [61, Thm. B]: if  $\overline{\text{BRic}} \geq 0$ , then the space of  $L^2$ -harmonic 1-forms on  $M$  is trivial. In particular, by Li-Tam's theory of harmonic functions in [34],  $M$  has only one non-parabolic end, and by [26] every cycle of codimension 1 must disconnect  $M$ . The result was obtained in [44] by assuming  $\overline{\text{Sec}} \geq 0$ . Related sharp vanishing theorems on spinnable minimal hypersurfaces in general ambient spaces were recently obtained by Bei & Pipoli [5].
- Shen & Zhu [58, Thm. 1 and Rem. 2]: if  $n \in \{3, 4\}$ ,  $\overline{\text{BRic}} \geq 0$  and  $M$  is topologically  $\mathbb{R} \times P$ , where  $P$  is compact and admits a metric of non-positive sectional curvature, then  $M$  is totally geodesic. Moreover,  $M$  is flat if  $\overline{\text{Sec}} \geq 0$ .
- Cheng [18, Thm. 2.1]: if  $n \in \{3, 4\}$  and  $\overline{\text{BRic}} > 0$ , then  $M$  must have only one end<sup>2</sup>.

**Remark 1.5.** For a detailed exposition of Li-Tam's theory of harmonic functions, we suggest [47].

<sup>2</sup>Cheng assumed  $M$  and  $N$  orientable, but her proof only uses that  $M$  is two-sided in  $N$ .

We are ready to state our main results in general target spaces. We first consider ambient manifolds with non-negative bi-Ricci curvature, and obtain the following splitting theorem that improves on [18, 58].

**Theorem 1.6.** *Let  $M^n \rightarrow (N^{n+1}, \bar{g})$  be a complete, non-compact, two-sided stable minimal hypersurface of dimension  $n \in \{3, 4\}$  in a manifold satisfying*

$$\overline{\text{BRic}} \geq 0.$$

*Then, one of the following mutually exclusive cases occurs:*

- (i)  *$M$  has only one end;*
- (ii)  *$M$  is totally geodesic and  $M = \mathbb{R} \times P$  with the product metric  $dt^2 + g_P$ , for some compact  $(n-1)$ -manifold  $(P, g_P)$  with non-negative Ricci curvature. Moreover,*

$$\overline{\text{Ric}}(v, v) \equiv 0, \quad \overline{\text{BRic}}(\partial_t, v) \equiv 0.$$

*Furthermore, if  $\overline{\text{Ric}} \geq 0$  then in (i) either  $M$  is non-parabolic, or  $M$  is parabolic, totally geodesic and  $\overline{\text{Ric}}(v, v) \equiv 0$ .*

**Remark 1.7.** Observe that  $\overline{\text{BRic}} \geq 0$  does not imply  $\overline{\text{Ric}} \geq 0$ , so the conclusion  $\overline{\text{Ric}}(v, v) \equiv 0$  in (ii) is not obvious. In dimension  $n = 2$ ,  $\overline{\text{BRic}} \geq 0$  becomes  $\bar{R} \geq 0$ , hence Theorem 1.6 can be seen as a higher dimensional analogue of the following well-known result by Fischer-Colbrie & Schoen, [29, Theorem 3]: a stable minimal surface  $M^2 \rightarrow N^3$  with more than one end in an ambient space with  $\bar{R} \geq 0$  must be conformally equivalent to a cylinder, flat and totally geodesic provided it has finite total curvature, i.e.

$$\int_M |K| < \infty.$$

In the same paper, the authors conjectured that the request of finite total curvature should be removable. This is the case, as shown by [44, 53, 7]. As a matter of fact, the argument in Theorem 1.6 works *verbatim* also in dimension  $n = 2$  (with  $P = \mathbb{S}^1$ ), thus providing a further different proof of Fischer-Colbrie and Schoen's claim.

We next consider ambient spaces with non-negative sectional curvature. Our goal is to improve on Li-Wang's remarkable splitting theorem [37, Thm. 0.1] quoted above by removing the condition that the immersion  $M \rightarrow N$  is proper. As a matter of fact, Li-Wang's result is the consequence of a more general statement where properness is not required a priori, see [37, Thm. 4.1]. However, in counting the number of ends the major issue therein is the possible presence of parabolic ends which are extrinsically bounded. In Theorem 1.6 we are able to overcome the problem in dimension  $n \leq 4$ . In the next theorem, we include one more dimension:

**Theorem 1.8.** *Let  $M^n \rightarrow (N^{n+1}, \bar{g})$  be a complete, non-compact, two-sided stable minimal hypersurface of dimension  $n \in \{3, 4, 5\}$  in a manifold satisfying*

$$\overline{\text{Sec}} \geq 0.$$

*Then, one of the following cases occurs:*

- (i)  *$M$  has only one end. Moreover, either  $M$  is non-parabolic, or  $M$  is parabolic, totally geodesic and  $\overline{\text{Ric}}(v, v) \equiv 0$ ;*

(ii)  $M$  is totally geodesic,  $\overline{\text{Ric}}(v, v) \equiv 0$  and  $M = \mathbb{R} \times P$  with the product metric, for some compact  $(n-1)$ -manifold  $P$  with non-negative sectional curvature.

Moreover, if  $n = 3$  then  $H^2(M, \mathbb{Z}) = 0$  or  $\mathbb{Z}$ .

**Remark 1.9.** Differently from [37], Theorems 1.6 and 1.8 do not need  $N$  to be complete.

**Remark 1.10.** The methods in [10, 37] to bound the number of ends of  $M$  ultimately rely on refined constructions of harmonic functions with controlled  $L^2$ -energy (more precisely, finite in [10] and slowly growing in [37]), and on the use of a geometric Poincaré formula due to Schoen & Yau, [55], for which stability is required. The formula still allows to conclude if stability is weakened to  $\delta$ -stability with  $\delta > \frac{n-1}{n}$ , but to reach lower  $\delta$  one needs further tools. On the other hand, the method we employ to prove Theorem 1.8 works *verbatim* for  $\delta$ -stable hypersurfaces with

$$\delta > \delta_n \doteq \frac{(n-1)^2}{4n},$$

and  $\delta_n \leq \frac{n-1}{n}$  if  $n \leq 5$ . Observe that  $\delta_3 = 1/3$ . The 3-dimensional case of Theorem 1.8 will be strengthened in Theorem 4.1 below.

**Remark 1.11.** Theorem 0.1 in [37] has a counterpart for properly immersed minimal hypersurfaces  $M \rightarrow N$  with finite index, which are shown to have finitely many ends, see [37, Thm. 4.1]. We think it would be very interesting to prove an analogous result in dimension  $n \leq 5$  by removing the properness assumption.

Our geometric applications to minimal hypersurfaces are consequences of two structure theorems for complete manifolds  $(M^n, g)$  for which the operator

$$\beta \Delta - \lambda_{\text{Ric}}$$

is non-negative in the spectral sense, where  $\lambda_{\text{Ric}} : M \rightarrow \mathbb{R}$  is the lowest eigenvalue of the Ricci tensor and  $\beta > 0$  is a suitable constant. In other words, we assume

$$\beta \int_M |\nabla \varphi|^2 + \int_M \lambda_{\text{Ric}} \varphi^2 \geq 0 \quad \forall \varphi \in \text{Lip}_c(M).$$

Note that the above can be rephrased as the existence of  $V \in C(M)$  so that

$$\text{Ric} \geq -\beta V g, \quad L \doteq \Delta + V \geq 0. \quad (1.2)$$

It is known that  $\lambda_{\text{Ric}}$  is a locally Lipschitz function (cf. [32]), so assuming  $V \in C_{\text{loc}}^{0,\alpha}(M)$  is not restrictive. Manifolds with property (1.2) were studied in depth by Li & Wang in [38], a continuation of their analysis of the topology of manifolds with positive spectral radius in [35, 36]. More recently, Carron [11], Carron & Rose [13] and Antonelli & Xu [2] also obtained relevant results. Three ranges of  $\beta$  play a significant role in the investigation in [38, 13, 2, 11]:

- (i)  $\beta \leq \frac{n-1}{n-2}$ : in this range, under some further assumptions on  $M$  one is able to control the number of non-parabolic ends, see [38, Theorem A and Corollary 4.2].
- (ii)  $\beta \leq \frac{4}{n-1}$ : this range also allows to control the number of parabolic ends of  $M$ , see [38, Theorems B,C].
- (iii)  $\beta < \frac{1}{n-2}$ : in this range,  $M$  looks close to a manifold with  $\text{Ric} \geq 0$ , as it will become apparent in Theorem 3.12 below. Analogies become tighter if one further assumes the *gaugeability* of  $L$ , i.e. the existence of a positive supersolution bounded below and above by positive constants, see [11, 13].

**Remark 1.12.** Note that  $\frac{n-1}{n-2} \geq \frac{4}{n-1}$ , with equality if and only if  $n = 3$ . This accounts for most of the extra information we obtain in dimension 3.

Our contributions are based on different ideas, and depend on a new point of view on the “conformal method” pioneered by Schoen & Yau [54], Fischer-Colbrie [28], Shen & Ye [56] and Shen & Zhu [58]. Loosely speaking, the method exploits a positive supersolution to  $Lu \leq 0$ , whose existence is granted by the second in (1.2), to conformally change the metric via  $\bar{g} = u^{2\beta}g$ . This allows to “push-up” the curvature contribution in the second variation formulas for  $\bar{g}$ -geodesics, and was mostly used to prove compactness results in the spirit of Bonnet-Myers’ one, see [53, 56, 57, 19, 27, 41, 17]. Very recently, the Bonnet-Myers’ type theorem in [56] found applications in [24] to prove the stable Bernstein theorem in  $\mathbb{R}^5$ , and in [15] to get rigidity of critical metrics. From a different perspective, the method was used in [28, 18, 16] to control the geometry of complete, non-compact minimal (hyper)-surfaces. Our investigation is inspired by these three last references, that we rework from a unified perspective.

First, we elaborate on [59, 18] to obtain a *criticality criterion* for the operator  $L$  on manifolds with more than one end. The criticality theory, developed by Murata [45] and Pinchover & Tintarev [49, 50, 51] (see also Zhao [64]), extends the standard dichotomy between parabolic and non-parabolic manifolds to operators of the type  $\Delta + V$ . The conclusions of our main Theorems 3.3 and 3.9 can be summarized as follows: either  $M$  has only one end, or  $L$  is critical and the first inequality in (1.2) is saturated at every point by at least one vector. More precisely:

- in Theorem 3.3, we obtain the above conclusion in the range  $\beta \in \left(0, \frac{n-1}{n-2}\right]$ . However, we need a further constraint in dimension  $n \geq 4$  (see Example 3.8), in accordance to statement (i) above: condition  $\beta \leq \frac{n-1}{n-2}$  hardly detects parabolic ends if  $n \geq 4$ . The constraint is not necessary if  $V_+$  is compactly supported, i.e. if  $\text{Ric} \geq 0$  outside of a compact set, see Theorem 3.5;
- in Theorem 3.9, we consider the ranges

$$\beta < \frac{4}{n-1} \quad \text{or}$$

$$\frac{4}{n-1} \leq \beta < \frac{n-1}{n-2} \quad \text{and } V_+ \text{ compactly supported.}$$

By exploiting a key property of the set of potentials  $V$  for which  $\Delta + V \geq 0$ , due to [50] (see Theorem 2.7), we are able to strengthen our conclusions to the following spectral splitting theorem: either

- $M$  has only one end, or
- $V \equiv 0$  and, consequently, by Cheeger-Gromoll’s theorem  $M$  splits as  $\mathbb{R} \times P$  for some compact manifold  $P$ .

One of its consequences is the obstruction result in Corollary 3.10, which can be seen as an extension of Cheng’s eigenvalue estimate.

In the range

$$\beta < \frac{1}{n-2},$$

we prove in Theorem 3.12 that for *any* choice of a positive (smooth) solution to  $Lu \leq 0$ , the metric

$$\bar{g} = u^{2\beta}g$$

is a *complete* metric with non-negative modified Bakry-Émery Ricci tensor

$$\overline{\text{Ric}}_f^N \doteq \overline{\text{Ric}} + \bar{\nabla}^2 f - \frac{1}{N-n} df \otimes df,$$

for  $f = (n-2)\beta \log u$  and a suitable  $N = N(n, \beta) > n$ . The result, which we obtain by adapting [16], establishes a higher-dimensional analogue of Fischer-Colbrie's [28, Theorem 1] and enables to exploit the extensive literature on manifolds with  $\overline{\text{Ric}}_f^N \geq 0$  and algebraic dimension  $N > n$ . These manifolds exhibit strong analogies to those with non-negative Ricci curvature: for instance, Corollaries 3.14 and 3.15 guarantee bounds on the Betti numbers of  $M$  and further topological consequences.

**Remark 1.13.** The criteria in Theorems 3.3, 3.5 and 3.12 also relate to an elegant rigidity result for surfaces satisfying

$$\Delta - \mu K \geq 0$$

for some  $\mu \geq 0$ , where  $K$  is the Gaussian curvature. The result, due to Castillon [14], was later extended by various authors, see in particular [7].

The paper is organized as follows: in Section 2, we recall the main results in criticality theory needed throughout the paper. In Section 3, we prove the abstract criteria in Theorems 3.3, 3.5, 3.9 and 3.12 and their corollaries. In Section 4, we deduce our geometric consequences.

**Note 1.14.** *When we were completing this paper, we learned that the authors of [1] had independently obtained results that partially overlap with ours, with different techniques. We agreed with them to post the results on arXiv independently at the same time.*

## 2 Criticality theory for $\Delta + V$

Let  $M$  be a Riemannian manifold. Hereafter, an exhaustion  $\{\Omega_j\}$  of  $M$  is a collection of relatively compact open sets with smooth boundary satisfying

$$\Omega_j \Subset \Omega_{j+1} \Subset M, \quad \bigcup_{j=1}^{\infty} \Omega_j = M.$$

Let  $V \in L_{\text{loc}}^{\infty}(M)$  and consider the operator

$$L_V \doteq \Delta + V.$$

Assume that  $L_V$  is non-negative in the spectral sense (we write  $L_V \geq 0$ ), namely, that the associated quadratic form

$$Q_V(\phi) \doteq \int_M \left[ |\nabla \phi|^2 - V \phi^2 \right]$$

is non-negative for each  $\phi \in \text{Lip}_c(M)$ , the set of compactly supported Lipschitz functions.

**Notation 2.1.** Hereafter, given  $V_1, V_2 \in L_{\text{loc}}^{\infty}(M)$  we will say that  $L_{V_1} \geq L_{V_2}$  whenever  $Q_{V_1}(\phi) \geq Q_{V_2}(\phi)$  for each  $\phi \in \text{Lip}_c(M)$ . This is equivalent to say that  $V_1 \leq V_2$  a.e. on  $M$ .

It is well-known, see for instance [47, Lemma 3.10], that  $L_V \geq 0$  is equivalent to the existence of a positive weak solution  $0 < u \in H_{\text{loc}}^1(M)$  to  $L_V u \leq 0$ , and also equivalent to the existence of a weak solution  $0 < u \in C^1(M)$  to  $L_V u = 0$ . The criticality theory discussed below, due to Murata [45] and Pinchover & Tintarev [49, 50], describes the geometry of the cone of positive solutions to  $L_V u \leq 0$ , see also the work of Zhao [64].

**Definition 2.2.** For  $V \in L_{\text{loc}}^\infty(M)$ , define  $L_V = \Delta + V$  and let  $\Omega \subseteq M$  be an open set.

- $L_V$  is **subcritical** in  $\Omega$  if there exists  $w \geq 0$ ,  $w \not\equiv 0$  in  $\Omega$  (called a **Hardy weight**) such that

$$\int_{\Omega} w |\phi|^2 \leq Q_V(\phi) \quad \forall \phi \in \text{Lip}_c(\Omega). \quad (2.1)$$

Otherwise, the operator  $L_V$  is said to be **critical**.

- $L_V$  has a **weighted spectral gap** in  $\Omega$  if there exists  $W \in C(\Omega)$ ,  $W > 0$  on  $\Omega$  such that

$$\int_{\Omega} W |\phi|^2 \leq Q_V(\phi) \quad \forall \phi \in \text{Lip}_c(\Omega). \quad (2.2)$$

- A sequence  $\{\phi_j\} \in L_c^\infty(\Omega) \cap H^1(\Omega)$  is said to be a **null sequence** if  $\phi_j \geq 0$  a.e. for each  $j$ ,  $Q_V(\phi_j) \rightarrow 0$  as  $j \rightarrow \infty$  and there exists a relatively compact open set  $B \Subset M$  and  $C > 1$  such that  $C^{-1} \leq \|\phi_j\|_{L^2(B)} \leq C$  for each  $j$ .
- A function  $0 \leq \eta \in H_{\text{loc}}^1(\Omega)$ ,  $\eta \geq 0$ ,  $\eta \not\equiv 0$  is a **ground state** for  $L_V$  on  $\Omega$  if it is the  $L_{\text{loc}}^2(\Omega)$  limit of a null sequence.

We have the following fundamental result, known as the ground state alternative, [45, 50]. The version stated below also includes some further implications, which can be found in [8, Theorem 4.1].

**Theorem 2.3.** *Let  $M$  be connected and non-compact, and consider an operator  $L_V \geq 0$  with  $V \in L_{\text{loc}}^\infty(M)$ . Then, either  $L_V$  has a weighted spectral gap or a ground state on  $M$ , and the two possibilities mutually exclude. Moreover, the following properties are equivalent:*

- (i)  $L_V$  is subcritical on  $M$ ;
- (ii)  $L_V$  has a weighted spectral gap on  $M$ ;
- (iii) There exist two positive solutions  $u_1, u_2 \in C(M) \cap H_{\text{loc}}^1(M)$  of  $L_V u \leq 0$  which are not proportional;
- (iv) for some (equivalently, each)  $o \in M$ , there exists a minimal positive distributional solution  $G$  to  $L_V G = -\delta_o$ ;
- (v) For some (equivalently, any)  $K \Subset M$  compact with non-empty interior, and for some (any)  $0 < \xi \in C(M) \cap H_{\text{loc}}^1(M)$  solving  $L_V \xi \leq 0$ , it holds

$$\text{cap}_V(K, \xi) \doteq \inf_{\phi \in \mathcal{D}(K, \xi)} Q_V(\phi) > 0,$$

where

$$\mathcal{D}(K, \xi) = \left\{ \phi \in \text{Lip}_c(M) : \phi \geq \xi \text{ in } K \right\}.$$



If  $L_V$  has a ground state  $\eta$ , then:

- $\eta > 0$  on  $M$  and solves  $L_V \eta = 0$  (in particular,  $\eta \in C_{\text{loc}}^{1,\mu}(M)$ ). Moreover, every other positive solution  $\xi$  to  $L_V \xi \leq 0$  is a multiple of  $\eta$ , hence a ground state;
- there exists a positive function  $W > 0$  such that for each  $\psi \in C_c^\infty(M)$  satisfying

$$\int \psi \eta \neq 0,$$

there exists a constant  $C = C(M, \psi, V)$  such that

$$C^{-1} \int_M W \phi^2 \leq Q_V(\phi) + C \left| \int_M \psi \phi \right|^2 \quad \forall \phi \in \text{Lip}_c(M). \quad (2.3)$$

*Proof (recap).* All the statements, apart from those regarding property (iv), are contained in [8, Theorem 4.1]. The equivalence (i)  $\Leftrightarrow$  (iv) is the content of [45, Theorem 2.4]. Inequality (2.3) was proved in [50, 51], see also [63] for a related result.  $\square$

**Remark 2.4.** Theorem 2.3 extends, to operators with a potential, the dichotomy between parabolic and non-parabolic manifolds, which are exactly the manifolds for which  $\Delta$  is critical and subcritical, respectively. If  $\Delta$  is subcritical, by [38] a Hardy weight can be constructed from any non-constant positive supersolution  $u$  of  $\Delta u \leq 0$  as

$$w = \frac{|\nabla u|^2}{4u^2}.$$

In particular, if  $u$  is the Green kernel of  $\Delta$  on  $\mathbb{R}^n$ ,  $n \geq 3$  and  $r$  is the distance to a fixed point, we recover the well-known Hardy weight

$$\frac{(n-2)^2}{4r^2}, \quad (2.4)$$

As shown in [12], the same Hardy weight also occurs for minimal submanifolds  $M^n \rightarrow N^p$  in a Cartan-Hadamard manifold, where now  $r$  is the restriction to  $M$  of the extrinsic distance from a fixed origin of  $N$ . More general Hardy weights, including sharpened ones for minimal submanifolds in hyperbolic space, can be found in [8, Section 5] and [38, Example 1.8], see also [9] for geometric applications.

**Remark 2.5.** As a direct application of the ground state alternative, if  $M$  satisfies a weighted Sobolev inequality

$$\left( \int_M \eta |\phi|^{\frac{vp}{v-p}} \right)^{\frac{v-p}{v}} \leq \mathcal{S} \int |\nabla \phi|^p \quad \forall \phi \in \text{Lip}_c(M)$$

for some  $0 < \eta \in C(M)$ ,  $\mathcal{S} > 0$  and  $v > p \geq 1$ , then  $\Delta$  is subcritical.

**Remark 2.6.** We stress that Theorem 2.3 also holds for homogeneous,  $p$ -Laplace operators

$$\text{div} (|\nabla u|^{p-2} \nabla u) + V |u|^{p-2} u,$$

see [51, 8]. Note that for  $p \neq 2$  one cannot avail of the Doob transform to reduce the result to the case  $V \equiv 0$  up to adding suitable weights to the Laplacian. See [11, Section 2.3] for more information on the Doob transform.

The proof of the following strict convexity property, which will be crucial for us in what follows, is contained in [51, Proposition 4.3] in Euclidean setting. The argument holds verbatim on manifolds, and we reproduce it here for the sake of completeness.

**Theorem 2.7.** *Let  $V_0, V_1 \in L_{\text{loc}}^\infty(M)$  and assume that  $L_{V_0} \geq 0$ ,  $L_{V_1} \geq 0$ . Then, setting  $V_t = (1-t)V_0 + tV_1$  for  $t \in [0, 1]$ , it holds  $L_{V_t} \geq 0$ . Moreover, if  $V_0$  does not coincide with  $V_1$  a.e., for each  $t \in (0, 1)$  the operator  $L_{V_t}$  is subcritical on  $M$ . In other words,*

$$\mathcal{K} \doteq \{V \in L_{\text{loc}}^\infty(M) : L_V \geq 0\}$$

is a convex set whose extremal points are those  $V$  for which  $L_V$  is critical.

*Proof.* Since

$$Q_{V_t}(\phi) = (1-t)Q_{V_0}(\phi) + tQ_{V_1}(\phi),$$

the assertion  $L_{V_t} \geq 0$  is immediate. Moreover, if any of  $L_{V_0}, L_{V_1}$  is subcritical, say  $L_{V_1}$ , and  $W_1$  is a Hardy weight,  $L_{V_t}$  is subcritical for  $t \in (0, 1)$  with Hardy weight  $tW_1$ . We are left to consider the case  $V_0 \not\equiv V_1$  and  $L_{V_0}, L_{V_1}$  critical. For  $j \in \{0, 1\}$  pick ground states  $\eta_j$  for  $L_{V_j}$ . Assume by contradiction that  $L_{V_t}$  is critical for some  $t \in (0, 1)$ , consider a null sequence  $\{\phi_k\}$  and its  $L_{\text{loc}}^2$ -limit  $\eta$ , a ground state. Since  $V_0 \not\equiv V_1$ ,  $\eta$  is neither proportional to  $\eta_0$  nor to  $\eta_1$ , thus there exist  $\psi_i \in C_c^\infty(M)$ ,  $i \in \{0, 1\}$  satisfying

$$\int_M \psi_i \eta_i \neq 0, \quad \int_M \psi_i \eta = 0 \quad \text{if } i \neq j. \quad (2.5)$$

By Theorem 2.3, there exist  $0 < W \in C(M)$  and a constant  $C = C(\psi_0, \psi_1, V_0, V_1, M)$  such that

$$C^{-1} \int_M W \phi^2 \leq Q_{V_t}(\phi) + C \left| \int_M \psi_i \phi \right|^2 \quad \forall \phi \in C_c^\infty(M).$$

By taking convex combination, we deduce

$$C^{-1} \int_M W \phi^2 \leq Q_{V_t}(\phi) + C(1-t) \left| \int_M \psi_1 \phi \right|^2 + Ct \left| \int_M \psi_2 \phi \right|^2 \quad \forall \phi \in C_c^\infty(M).$$

Hence, using  $\phi = \phi_k$ , from  $Q_{V_t}(\phi_k) \rightarrow 0$  and (2.5) we get

$$C^{-1} \int_M W \eta^2 \leq C(1-t) \left| \int_M \psi_1 \eta \right|^2 + Ct \left| \int_M \psi_2 \eta \right|^2 = 0, \quad (2.6)$$

contradiction.  $\square$

The notion of criticality can be localized on each end of  $M$  as follows: assume that  $L \doteq \Delta + V$  satisfies  $L \geq 0$  on  $M$ . Hereafter, a pair  $(K, \xi)$  is the data of a solution  $0 < \xi \in C_{\text{loc}}^{1,\alpha}(M)$  to  $L\xi = 0$  (equality here is important) and a compact set  $K$  with non-empty interior and smooth boundary. Let  $E$  be an end of  $M$  with respect to  $K$ , that is, a connected component of  $M \setminus K$  with non-compact closure. Fix an exhaustion  $\{\Omega_j\}$  of  $M$  with  $K \subset \Omega_1$ , set  $E_j = E \cap \Omega_j$  and consider the family of solutions  $u_j$  to

$$\begin{cases} Lu_j = 0 & \text{on } E_j, \\ u_j = \xi & \text{on } \partial E, \quad u_j = 0 & \text{on } E \cap \partial\Omega_j. \end{cases} \quad (2.7)$$

The maximum and comparison principles imply that  $0 < u_j \leq u_{j+1} \leq \xi$ , thus by elliptic estimates  $u_j \uparrow u_\xi$  in  $C_{\text{loc}}^1(\overline{E})$  for some  $u_\xi > 0$  solving

$$\begin{cases} Lu_\xi = 0 & \text{on } E, \\ u_\xi = \xi & \text{on } \partial E, \quad 0 < u_\xi \leq \xi \text{ on } E. \end{cases}$$

By comparison,  $u_\xi$  does not depend on the chosen exhaustion. We call  $u_\xi$  the minimal solution on  $E$  with respect to  $\xi$ . The strong maximum principle implies that either  $u_\xi \equiv \xi$  or  $u_\xi < \xi$  on  $E$ .

**Definition 2.8.** We say that  $L$  is  $\xi$ -critical on  $E$  if  $u_\xi \equiv \xi$  on  $E$ , and  $\xi$ -subcritical otherwise.

We have the following

**Theorem 2.9.** Let  $(M, g)$  be a Riemannian manifold,  $V \in L_{\text{loc}}^\infty(M)$  and assume that  $L \doteq \Delta + V \geq 0$ . Then, the following are equivalent:

- (i)  $L$  is critical on  $M$ ;
- (ii) for every pair  $(K, \xi)$ ,  $L$  is  $\xi$ -critical on each end  $E$  with respect to  $K$ ;
- (iii) for some pair  $(K, \xi)$ ,  $L$  is  $\xi$ -critical on each end  $E$  with respect to  $K$ ;

*Proof.* We introduce the following construction. Fix a pair  $(K, \xi)$ , and let  $E^{(1)}, \dots, E^{(k)}$  be the ends with respect to  $K$ . Up to attaching to  $K$  the bounded connected components of  $M \setminus K$ , we can assume that  $M \setminus K = E^{(1)} \cup \dots \cup E^{(k)}$ . For each  $i$ , let  $u^{(i)}$  be the minimal solution on  $E^{(i)}$  with respect to  $\xi$ . Let  $\{\Omega_j\}$  be a smooth exhaustion of  $M$ , and for each  $i$  let  $\{u_j^{(i)}\}$  be the sequence constructed in (2.7) on  $E^{(i)}$ . Define

$$\xi_j \doteq \begin{cases} \xi & \text{on } K, \\ u_j^{(i)} & \text{on } E^{(i)} \cap \Omega_j. \end{cases}$$

Then,  $0 < \xi_j \leq \xi$  on  $\Omega_j$ . Moreover, integrating by parts  $0 = \xi L \xi$  on  $K$  and  $0 = u_j^{(i)} L u_j^{(i)}$  on  $E^{(i)} \cap \Omega_j$  and subtracting the resulting identities we deduce

$$Q_V(\xi_j) = \sum_{i=1}^k \int_{\partial E^{(i)}} \xi \left[ \partial_\eta \xi - \partial_\eta u_j^{(i)} \right].$$

Moreover,

$$\xi_j \uparrow \hat{\xi} \doteq \begin{cases} \xi & \text{on } K, \\ u^{(i)} & \text{on } E^{(i)}, \end{cases}$$

and from  $u^{(i)} \leq \xi$  one easily deduces that  $L \hat{\xi} \leq 0$  in the weak sense on  $M$ . We are ready to prove our equivalence.

(i)  $\Rightarrow$  (ii). By contradiction, assume that for some pair  $(K, \xi)$  the operator  $L$  is subcritical on some end. Then, by construction  $\hat{\xi} \not\equiv \xi$ . Since  $\hat{\xi} = \xi$  on  $K$ ,  $\xi$  and  $\hat{\xi}$  are distinct supersolutions for  $L$  which are not proportional. By (iii) in Theorem 2.3 we deduce that  $L$  is subcritical on  $M$ , contradiction.

(ii)  $\Rightarrow$  (iii) is obvious.

(iii)  $\Rightarrow$  (i). Fix the pair  $(K, \xi)$  in (iii), and assume by contradiction that  $L$  is subcritical on  $M$ .

Then, the energy  $Q_V(\xi_j)$  cannot vanish as  $j \rightarrow \infty$  (otherwise,  $\{\xi_j\}$  would be a null sequence). Since

$$Q_V(\xi_j) \rightarrow \sum_{i=1}^k \int_{\partial E^{(i)}} \xi \left[ \partial_\eta \xi - \partial_\eta u^{(i)} \right],$$

we deduce that  $u^{(i)} \neq \xi$  for some  $i$ , so  $L$  is  $\xi$ -subcritical on  $E^{(i)}$ , contradiction.  $\square$

### 3 The conformal method, criticality and splitting criteria

Let us consider a conformal deformation

$$\bar{g} = e^{2\varphi} g, \quad \varphi \in C^\infty(M),$$

and denote with a bar superscript quantities in the metric  $\bar{g}$ . The Ricci curvatures of  $g$  and  $\bar{g}$  relate as follows:

$$\overline{\text{Ric}} = \text{Ric} - (n-2)(\nabla^2 \varphi - d\varphi \otimes d\varphi) - [\Delta \varphi + (n-2)|d\varphi|^2] g. \quad (3.1)$$

Let  $\gamma$  be a  $\bar{g}$ -geodesic, namely, a geodesic in the metric  $\bar{g}$ . Denoting by  $s$  and  $\bar{s}$ , respectively, the  $g$ -arclength and the  $\bar{g}$ -arclength of  $\gamma$ , we have  $d\bar{s} = e^\varphi ds$  and  $\partial_{\bar{s}} = e^{-\varphi} \partial_s$ , where  $\varphi$  is evaluated at  $\gamma(s)$ . Since the  $g$  and  $\bar{g}$ -unit tangent vectors to  $\gamma$  satisfy  $\gamma_{\bar{s}} = e^{-\varphi} \gamma_s$ , we have

$$0 = \bar{\nabla}_{\gamma_{\bar{s}}} \gamma_{\bar{s}} = e^{-2\varphi} \left( \nabla_{\gamma_s} \gamma_s + \gamma_s(\varphi) \gamma_s - \nabla \varphi \right)$$

Therefore,

$$\begin{aligned} (\varphi \circ \gamma)_{ss} &= \nabla^2 \varphi(\gamma_s, \gamma_s) + d\varphi(\nabla_{\gamma_s} \gamma_s) \\ &= \nabla^2 \varphi(\gamma_s, \gamma_s) + d\varphi \left( \nabla \varphi - \gamma_s(\varphi) \gamma_s \right) = \nabla^2 \varphi(\gamma_s, \gamma_s) + |d\varphi|^2 - d\varphi(\gamma_s)^2 \end{aligned} \quad (3.2)$$

Therefore, when restricted to  $\gamma$ , (3.1) implies the remarkable identity

$$\overline{\text{Ric}}(\gamma_s, \gamma_s) = \text{Ric}(\gamma_s, \gamma_s) - (n-2)(\varphi \circ \gamma)_{ss} - \Delta \varphi, \quad (3.3)$$

appearing implicitly in [56] and proved in the Appendix of [27]. Hereafter, the composition with  $\gamma$  will tacitly be assumed, thus we will write  $\varphi_s, \varphi_{ss}$  rather than  $(\varphi \circ \gamma)_s, (\varphi \circ \gamma)_{ss}$  and so forth.

We examine more closely the metric properties of the metric  $\bar{g} = e^{2\varphi} g$ . If  $f, \psi : M \rightarrow \mathbb{R}$  and smooth functions, note that

$$\begin{aligned} \bar{\nabla}^2 f &= \nabla^2 f - (df \otimes d\varphi + d\varphi \otimes df) + g(df, d\varphi)g, \\ \bar{\Delta} \psi &= e^{-2\varphi} \left( \Delta \psi + (n-2)g(d\varphi, d\psi) \right) \end{aligned}$$

Hence, from (3.1), choosing

$$f = (n-2)\varphi$$

the  $\infty$ -Bakry-Emery Ricci curvature

$$\overline{\text{Ric}}_f \doteq \overline{\text{Ric}} + \bar{\nabla}^2 f$$

satisfies:

$$\overline{\text{Ric}}_f = \text{Ric} - (n-2)d\varphi \otimes d\varphi - (\Delta\varphi)g. \quad (3.4)$$

Moreover, the associated weighted Laplacian

$$\bar{\Delta}_f \psi \doteq \bar{\Delta}\psi - \bar{g}(df, d\psi)$$

of a function  $\psi$  satisfies

$$\bar{\Delta}_f \psi = e^{-2\varphi} \Delta \psi. \quad (3.5)$$

Observe that, in dimension  $n = 2$ , formulas (3.4) and (3.5) become the well-known identities relating the Gaussian curvature and the Laplacian of conformally deformed metrics. The main computation, repeatedly used throughout the paper, is the following one essentially due to [56]. Hereafter, we write

$$C_0^k([a, b]) = \{\eta \in C^k([a, b]) : \eta(a) = \eta(b) = 0\}.$$

**Proposition 3.1.** *Assume that  $(M^n, g)$  is a Riemannian manifold, and let  $V \in C(M)$  satisfy*

$$L \doteq \Delta + V \geq 0.$$

Let  $0 < u \in C^2(M)$  solve  $Lu \leq 0$ , and for  $\beta > 0$  consider the metric

$$\bar{g} = u^{2\beta} g.$$

Assume that  $\gamma : [a, b] \rightarrow M$  is a curve parametrized by  $g$ -arclength  $s$ , and assume that  $\gamma$  is  $\bar{g}$ -length minimizing. Then, for each  $\eta \in C_0^1([a, b])$  it holds

$$\begin{aligned} & \beta \int_a^b \frac{-Lu}{u} \frac{\eta^2}{u^\beta} ds + \int_a^b \left( \text{Ric}(\gamma_s, \gamma_s) + \beta V \right) \frac{\eta^2}{u^\beta} ds + \beta \int_a^b |\nabla^\perp \log u|^2 \frac{\eta^2}{u^\beta} ds \\ & \leq (n-1) \int_a^b \frac{(\eta_s)^2}{\eta^2} \frac{\eta^2}{u^\beta} ds - \beta \int_a^b \frac{(u_s)^2}{u^2} \frac{\eta^2}{u^\beta} ds - \beta(n-2) \int_a^b \frac{u_s}{u} \left( \frac{\eta^2}{u^\beta} \right)_s ds, \end{aligned} \quad (3.6)$$

where  $\nabla^\perp$  is the component of the  $g$ -gradient  $\nabla$  orthogonal to  $\gamma_s$ .

*Proof.* Setting  $\varphi = \beta \log u$ , we deduce from (3.3) the identity

$$\overline{\text{Ric}}(\gamma_s, \gamma_s) = \text{Ric}(\gamma_s, \gamma_s) - (n-2)\beta(\log u)_{ss} - \beta \frac{Lu}{u} + \beta V + \beta |d \log u|^2.$$

Let  $\bar{s}$  denotes the  $\bar{g}$ -arclength. Since  $\gamma$  is  $\bar{g}$ -minimizing, the second variation formula in the metric  $\bar{g}$  yields

$$\begin{aligned} 0 & \leq \int_{\bar{s}(a)}^{\bar{s}(b)} \left[ (n-1)(\eta_{\bar{s}})^2 - \overline{\text{Ric}}(\gamma_{\bar{s}}, \gamma_{\bar{s}})\eta^2 \right] d\bar{s} \\ & = \int_a^b \left[ (n-1)(\eta_s)^2 - \overline{\text{Ric}}(\gamma_s, \gamma_s)\eta^2 \right] u^{-\beta} ds \\ & = (n-1) \int_a^b (\eta_s)^2 u^{-\beta} ds - \int_a^b \left( \text{Ric}(\gamma_s, \gamma_s) + \beta V \right) \eta^2 u^{-\beta} ds \\ & \quad - \beta \int_a^b \left( |d \log u|^2 - (n-2)(\log u)_{ss} \right) \eta^2 u^{-\beta} ds + \beta \int_a^b \frac{Lu}{u} \eta^2 u^{-\beta} ds. \end{aligned} \quad (3.7)$$

Using  $|d \log u|^2 = |\nabla^\perp \log u|^2 + (u_s)^2/u^2$  and integrating by parts the term with  $(\log u)_{ss}$  we get (3.6).  $\square$

The following proposition enables to construct a “shortest line”  $\gamma$  in a Riemannian manifold  $(M, h)$  with more than one end, so that the completeness of  $h$  is detected by the fact that  $\gamma$  is defined on the whole of  $\mathbb{R}$  once parametrized by  $h$ -arclength. The result improves on [18, Lemma 1.1].

**Proposition 3.2.** *Let  $(M, h)$  be a Riemannian manifold with more than one end. Then, there exist  $T, T^* \in (0, \infty]$  and a constant speed curve  $\gamma : (-T, T^*) \rightarrow M$  which satisfies:*

- (i)  $\gamma$  is length minimizing for every pair of its points,
- (ii)  $\gamma$  is a divergent curve, i.e.  $\gamma^{-1}(K)$  is compact for each  $K \subset M$  compact. Moreover, for  $\varepsilon$  small enough  $\gamma_{[T^*-\varepsilon, T^*)}$  and  $\gamma_{(-T, -T+\varepsilon]}$  belong to different ends of  $M$ .
- (iii)  $(M, h)$  is complete if and only if both the curves  $\gamma_{[0, T^*)}$  and  $\gamma_{(-T, 0]}$  have infinite length, that is,  $T = T^* = \infty$ .

*Proof.* To construct  $\gamma$ , fix a compact set  $K$  with smooth boundary such that  $M \setminus K$  has at least two ends. Up to including in  $K$  the components of  $M \setminus K$  with compact closure, we can assume that  $M \setminus K = \coprod_{i=1}^{\ell} E_i$ , with  $E_i$  an end. Pick an exhaustion  $\{\Omega_n\}$  of  $M$  with  $K \Subset \Omega_1$ . For each  $i, n$ , define  $E_i(n) = E_i \setminus \Omega_n$ . Define

$$\tau_n = \min \left\{ d(\partial E_i(n), \partial E_j(n)) : i, j \in \{1, \dots, \ell\}, i \neq j \right\}.$$

Since  $(\overline{\Omega_n}, h)$  is a compact manifold with boundary, there exists a segment (i.e. a unit speed, minimizing geodesic)  $\gamma_n$  in  $\overline{\Omega_n}$  realizing  $\tau_n$ , joining  $\partial E_{i_n}(n)$  to  $\partial E_{j_n}(n)$  for some indices  $i_n \neq j_n$ . By construction,  $\gamma_n$  intersects  $K$ . Thus, up to translation we can assume that  $\gamma : [-T_n, T_n^*] \rightarrow \overline{\Omega_n}$  for some positive  $T_n, T_n^*$ , and that

$$\gamma_n(0) \in K, \quad \gamma_n(-T_n) \in \partial E_{i_n}(n), \quad \gamma_n(T_n^*) \in \partial E_{j_n}(n).$$

By minimality  $\gamma_n((-T_n, T_n^*)) \subset \Omega_n$ , hence  $\gamma_n$  is a geodesic. Furthermore, by construction there exist indices  $i, j$  such that  $\gamma_n$  joins  $\partial E_i(n)$  and  $\partial E_j(n)$  for infinitely many  $n$ . Up to passing to a subsequence in  $n$  (still labelled the same), the family  $\{\gamma_n\}$  locally converges to a minimizing geodesic  $\gamma : (-T, T^*) \rightarrow M$  which crosses every  $\partial E_i(n)$  and  $\partial E_j(n)$ , say at instants  $t_n$  and  $t_n^*$ , respectively, and satisfies  $\gamma(0) = o \in K$ . In particular,  $\gamma$  satisfies (i). To prove (ii), it is enough to show that  $\gamma_{[0, T^*)}$  is divergent and eventually valued in  $E_j$ , the case of  $\gamma_{(-T, 0]}$  being analogous. By contradiction, assume that there exists a compact set  $K$  (which we can assume contains  $o$  in its interior) and a sequence  $\tau_n \rightarrow T^*$  so that  $\gamma(\tau_n) \in K$ . Pick  $n_0$  such that  $K \subset \Omega_{n_0}$ . Then, for each  $n \geq n_0$  we can estimate

$$|t_n^* - \tau_n| = d(\gamma(t_n^*), \gamma(\tau_n)) \geq d(\partial E_j(n), K) \geq d(\partial E_j(n_0), K).$$

From  $t_n, \tau_n \rightarrow T^*$  we infer that necessarily  $T^* = \infty$ . However, from

$$\tau_n = d(o, \gamma(\tau_n)) \leq \text{diam}(K)$$

and letting  $n \rightarrow \infty$  we get a contradiction. Thus,  $\gamma_{[0, T^*)}$  is divergent, and since  $\gamma(t_n) \in \partial E_j(n)$  we conclude that  $\gamma$  is eventually contained in  $E_j$ . This proves (ii).

To show (iii), assume that  $(M, h)$  is complete. Then, there exist  $R_n \rightarrow \infty$  such that  $B_{R_n}(o) \subset \Omega_n$ . Since  $\gamma$  is minimizing, by the definition of  $t_n, t_n^*$  it holds  $t_n \leq -R_n, t_n^* \geq R_n$ , hence  $T = T^* = \infty$ .

Viceversa, assume that  $T = T^* = \infty$ . In particular,  $T_n^* \rightarrow \infty$ . Let  $\sigma : [0, R) \rightarrow M$  be a

maximally extended unit speed geodesic issuing from  $o$ . We claim that  $R = \infty$ , from which the completeness follows. Assume by contradiction that  $R < \infty$ . Then, by ODE theory  $\sigma$  is a divergent curve, and we can fix  $k \in \{1, \dots, \ell\}$  so that  $\sigma([R_0, R]) \subset E_k$  for some  $R_0 > 0$ . Choose an index between  $i, j$  which is different from  $k$ , say  $i$ . For each  $n$ , let  $s_n$  be the first instant such that  $\sigma(s_n) \in \partial E_k(n)$ . We claim that

$$d(\gamma_n(0), \partial E_j(n)) \leq d(\gamma_n(0), \partial E_k(n)). \quad (3.8)$$

Otherwise, concatenating  $\gamma_n : (-T_n, 0] \rightarrow M$  to a segment  $\eta$  from  $\gamma_n(0)$  to  $\partial E_k(n)$  we would get a curve joining  $\partial E_i(n)$  to  $\partial E_k(n)$  shorter than  $\gamma_n$ , contradiction. From (3.8) we deduce

$$\begin{aligned} T_n^* &= d(\gamma_n(0), \partial E_j(n)) \leq d(\gamma_n(0), \partial E_k(n)) \\ &\leq d(\gamma_n(0), \gamma(0)) + d(\gamma(0), \partial E_k(n)) \\ &\leq d(\gamma_n(0), \gamma(0)) + s_n < d(\gamma_n(0), \gamma(0)) + R. \end{aligned}$$

Letting  $n \rightarrow \infty$  we get a contradiction.  $\square$

Here is our first main result:

**Theorem 3.3.** *Let  $(M, g)$  be a complete Riemannian manifold, and assume that there exists  $V \in C_{\text{loc}}^{0,\alpha}(M)$  such that*

$$\text{Ric} \geq -\beta V g \quad \text{on } M, \quad L \doteq \Delta + V \geq 0, \quad (3.9)$$

where

$$0 < \beta \leq \frac{n-1}{n-2}.$$

If either

- (i)  $n = 3$ , or
- (ii)  $n \geq 4$ , and there exists a compact set  $Z$  and a weak solution  $0 < w \in C(\overline{M \setminus Z})$  to  $Lw \leq 0$  satisfying

$$w(x) = o(r(x))^{\frac{n-1}{\beta(n-3)}} \quad \text{as } r(x) \rightarrow \infty, \quad (3.10)$$

where  $r(x)$  is the distance from a fixed origin, then the following holds: either

- $M$  has only one end, or
- $L$  is critical and, for each  $x \in M$ ,

$$\min_{X \in T_x^1 M} (\text{Ric}(X, X) + \beta V(x)) = 0. \quad (3.11)$$

*Proof.* Assume that  $M$  has at least two ends, and, by contradiction, that  $L$  is subcritical. Pick an exhaustion  $\{\Omega_j\}$  of  $M$  with  $Z \subset \Omega_1$  and so that  $M \setminus \Omega_1$  has at least two ends  $E_1$  and  $E_2$ . By (i)  $\iff$  (ii) in Theorem 2.3, there exists  $0 \leq W \in C_c^\infty(\Omega_3)$  which is positive on  $\overline{\Omega_2}$  and satisfies

$$\int_{\Omega} 2W |\eta|^2 dx \leq Q_V(\eta) \quad \forall \eta \in \text{Lip}_c(M).$$

Thus,  $L + W \geq 0$  and is subcritical. Picking a point  $o \in \Omega_1$ , by (iv) in Theorem 2.3 there exists a minimal positive solution  $G \in C_{\text{loc}}^{2,\alpha}(M \setminus \{o\})$  to  $LG + WG = -\delta_o$ . Fix also a global solution

$0 < v \in C_{\text{loc}}^{2,\alpha}(M)$  to  $Lv = 0$  on  $M$ . Since  $G$  is constructed as the limit of a sequence of kernels  $G_j$  on  $\Omega_j$  with Dirichlet boundary conditions, comparing  $G_j$  to  $w$  and  $v$  on  $\Omega_j \setminus \Omega_3$  (recall,  $LG_j = 0$  there) and letting  $j \rightarrow \infty$ , there exists a constant  $C > 0$  such that  $G \leq C \min\{w, v\}$  on  $M \setminus \Omega_3$ . Thus, up to increasing  $C$ ,

$$G \leq C \min\{w, v\} \quad \text{on } M \setminus \Omega_1. \quad (3.12)$$

The function  $z \doteq G/v$  therefore solves

$$\begin{cases} \operatorname{div}(v^2 \nabla z) \leq 0 & \text{on } M \setminus \{o\} \\ z(x) \rightarrow \infty & \text{as } x \rightarrow o \\ z \leq C & \text{on } M \setminus \Omega_1. \end{cases}$$

Consider a concave function  $\eta \in C^2([0, \infty))$  satisfying  $\eta(t) = t$  for  $t \in [0, 2C]$ ,  $\eta(t) = 3C$  for  $t \geq 4C$ . Then,  $\eta(z) \in C^2(M)$  solves

$$\operatorname{div}(v^2 \nabla \eta(z)) \leq 0 \quad \text{on } M,$$

and thus

$$u \doteq v\eta(z) \in C^2(M) \quad \text{solves} \quad \begin{cases} Lu \leq 0 & \text{on } M \\ u = G & \text{on } M \setminus \Omega_1. \end{cases}$$

Consider the conformal deformation

$$\bar{g} = u^{2\beta} g.$$

We construct a  $\bar{g}$ -line  $\gamma : (-T, T^*) \rightarrow M$  as in Proposition 3.2. By property (ii) of  $\gamma$  and since  $g$  is complete, it follows that once reparametrized by its  $g$ -arclength  $s$  the curve  $\gamma$  is defined on the whole of  $\mathbb{R}$ . Also, because  $M \setminus \Omega_1$  is disconnected and again by property (ii) we deduce that  $\gamma \cap \bar{\Omega}_1 \neq \emptyset$  and  $\gamma \cap (\Omega_2 \setminus \bar{\Omega}_1) \neq \emptyset$ . By Proposition 3.1, for each  $\eta \in C_c^1(\mathbb{R})$  it holds

$$\begin{aligned} & \beta \int_{-\infty}^{\infty} \frac{-Lu \eta^2}{u u^\beta} + \int_{-\infty}^{\infty} \left( \operatorname{Ric}(\gamma_s, \gamma_s) + \beta V \right) \frac{\eta^2}{u^\beta} \\ & \leq (n-1) \int_{-\infty}^{\infty} \frac{(\eta_s)^2 \eta^2}{\eta^2 u^\beta} - \beta \int_{-\infty}^{\infty} \frac{(u_s)^2 \eta^2}{u^2 u^\beta} - \beta(n-2) \int_{-\infty}^{\infty} \frac{u_s}{u} \left( \frac{\eta^2}{u^\beta} \right)_s, \end{aligned}$$

where we omitted to write  $ds$  for the ease of notation. From (3.9),

$$\beta \int_{-\infty}^{\infty} \frac{-Lu \eta^2}{u u^\beta} \leq (n-1) \int_{-\infty}^{\infty} \frac{(\eta_s)^2 \eta^2}{\eta^2 u^\beta} - \beta \int_{-\infty}^{\infty} \frac{(u_s)^2 \eta^2}{u^2 u^\beta} - \beta(n-2) \int_{-\infty}^{\infty} \frac{u_s}{u} \left( \frac{\eta^2}{u^\beta} \right)_s.$$

Write  $\eta = u^\theta \psi$  for some  $\theta \geq 0$  to be determined. Then,

$$\begin{aligned} \frac{\beta}{n-1} \int_{-\infty}^{\infty} \frac{-Lu u^{2\theta-\beta} \psi^2}{u} & \leq \int_{-\infty}^{\infty} \left[ \theta^2 - \frac{2\theta\beta(n-2)}{n-1} + \frac{\beta^2(n-2)}{n-1} - \frac{\beta}{n-1} \right] \frac{(u_s)^2}{u^2} u^{2\theta-\beta} \psi^2 \\ & \quad + \int_{-\infty}^{\infty} (\psi_s)^2 u^{2\theta-\beta} + 2 \left[ \theta - \frac{\beta(n-2)}{n-1} \right] \int_{-\infty}^{\infty} \frac{u_s \psi \psi_s}{u} u^{2\theta-\beta}. \end{aligned}$$

Choosing  $\theta = \frac{\beta(n-2)}{n-1}$ , we conclude

$$\frac{\beta}{n-1} \int_{-\infty}^{\infty} \frac{-Lu u^{\frac{\beta(n-3)}{n-1}} \psi^2}{u} \leq \int_{-\infty}^{\infty} (\psi_s)^2 u^{\frac{\beta(n-3)}{n-1}} \quad (3.13)$$



Let  $0 \leq \psi_R \leq 1$  be an even function satisfying  $\psi_R \equiv 1$  on  $[-R, R]$ ,  $\psi_R \in C_c^\infty((-2R, 2R))$  and  $|(\psi_R)_s| \leq 4/R$ . Then,

$$\frac{\beta}{n-1} \int_{-R}^R \frac{-Lu}{u} u^{\frac{\beta(n-3)}{n-1}} \leq \frac{16}{R^2} \left[ \int_R^{2R} u^{\frac{\beta(n-3)}{n-1}} + \int_{-2R}^{-R} u^{\frac{\beta(n-3)}{n-1}} \right]. \quad (3.14)$$

If  $n = 3$ , letting  $R \rightarrow \infty$  we deduce  $Lu \equiv 0$  along  $\gamma$ . Similarly, if  $n \geq 4$ , by (3.12) there exists  $C$  such that

$$u(\gamma(s)) \leq C(1 + w(\gamma(s))) \quad \forall s \in \mathbb{R}. \quad (3.15)$$

Assumption (3.10) (without loss of generality, we can assume that  $r$  is the distance from  $\gamma(0)$ ) and  $|\gamma_s| = 1$  imply

$$r(\gamma(s)) \leq |s|,$$

whence

$$u(\gamma(s)) = o(|s|^{\frac{n-1}{\beta(n-3)}}) \quad \text{as } |s| \rightarrow \infty.$$

Plugging into (3.14) and letting  $R \rightarrow \infty$  we again deduce  $Lu \equiv 0$  along  $\gamma$ . To conclude, we observe that

$$Lu = LG = -WG < 0 \quad \text{on } \gamma \cap (\Omega_2 \setminus \overline{\Omega_1}) \neq \emptyset,$$

leading to the desired contradiction. To prove (3.11), assume by contradiction that

$$\text{Ric} + \beta Vg > 0 \quad \text{at some } x_0 \in M,$$

and fix  $0 \leq w \in C^\infty(M)$  not identically zero satisfying

$$\text{Ric} + \beta(V - w)g \geq 0 \quad \text{on } M.$$

Since  $L - w \geq L \geq 0$ , and  $w \not\equiv 0$ ,  $L - w$  is subcritical. However, we can apply the first part of the proof with  $V - w$  replacing  $V$  to deduce that  $L - w$  is critical, contradiction.  $\square$

**Example 3.4.** We denote by  $C^n \rightarrow \mathbb{R}^{n+1}$ ,  $n \geq 3$ , the higher-dimensional catenoid. Following [60] (see also [25]),  $C^n$  depends on a real parameter  $\phi_0 > 0$  and is defined by the map

$$F : (-S, S) \times \mathbb{S}^{n-1} \rightarrow \mathbb{R}^{n+1}, \\ (s, \omega) \mapsto (\phi(s)\omega, s),$$

where

$$S = \int_{\phi_0}^{\infty} \frac{dt}{[(t/\phi_0)^{2(n-1)} - 1]^{1/2}} < \infty$$

and  $\phi$  solves

$$\begin{cases} \phi'' [1 + (\phi')^2]^{-\frac{3}{2}} - \frac{n-1}{\phi} [1 + (\phi')^2]^{-\frac{1}{2}} = 0 \\ \phi(0) = \phi_0 > 0 \\ \phi'(0) = 0. \end{cases}$$

By [60], the second fundamental  $|A|$  is positive on  $C^n$ , globally bounded (in fact,  $C^n$  has finite total curvature) and solves

$$\Delta |A|^{\frac{n-2}{n}} + \frac{n-2}{n} |A|^2 |A|^{\frac{n-2}{n}} = 0.$$

Therefore,  $C^n$  is  $\frac{n-2}{n}$ -stable:

$$L \doteq \Delta + V \geq 0, \quad \text{where } V \doteq \frac{n-2}{n}|A|^2.$$

By Gauss equations and the refined Kato inequality,

$$\text{Ric} \geq -\frac{n-1}{n}|A|^2 = \frac{n-1}{n-2}V,$$

Hence, we can apply Theorem 3.3 with  $w = |A|^{\frac{n-2}{n}}$  to deduce that  $L$  is critical on  $C^n$ .

We next examine the case where  $V_+$  has compact support, namely,  $\text{Ric} \geq 0$  outside of a compact set. In this case, we can drop the dimensional restriction and get:

**Theorem 3.5.** *Let  $(M, g)$  be a complete Riemannian manifold, and assume that there exists  $V \in C_{\text{loc}}^{0,\alpha}(M)$  with  $V_+$  compactly supported such that*

$$\text{Ric} \geq -\frac{n-1}{n-2}Vg \quad \text{on } M, \quad L \doteq \Delta + V \geq 0. \quad (3.16)$$

then, either

- $M$  has only one end, or
- $L$  is critical and, for each  $x \in M$ ,

$$\min_{X \in T_x^1 M} \left( \text{Ric}(X, X) + \frac{n-1}{n-2}V(x) \right) = 0.$$

*Proof.* Assuming, by contradiction, that  $M$  has at least two ends and that  $L$  is subcritical, we follow the proof of Theorem 3.3 and construct  $u, \gamma$ . For  $\tau \geq 1$ , let  $\xi : [0, \infty) \rightarrow \mathbb{R}$  be a concave, non-decreasing function satisfying

$$\xi(x) = x \quad \text{for } x \in [0, 1], \quad \xi(x) = 2 \quad \text{for } x \geq 4, \quad \xi'(x) \leq 1 \quad \forall x \in [0, \infty)$$

and define

$$\xi_\tau(x) = \tau\xi(x/\tau), \quad u_\tau = \xi_\tau(u)$$

Then,  $0 \leq \xi'_\tau \leq 1$ ,  $\xi_\tau - x\xi'_\tau \geq 0$ ,  $u_\tau \in L^\infty(M)$  and  $\xi'_\tau \uparrow 1$ ,  $u_\tau \uparrow u$  as  $\tau \rightarrow \infty$ . Moreover,

$$Lu_\tau \leq \xi'_\tau(u)Lu + V(u_\tau - u\xi'_\tau(u)) \leq \xi'_\tau(u)Lu + V_+(u_\tau - u\xi'_\tau(u))$$

By using  $\beta = \frac{n-1}{n-2}$  and  $u_\tau$  in place of  $u$ , and proceeding as in Theorem 3.3, we obtain the following analogue of (3.13):

$$\frac{1}{n-2} \int_{-\infty}^{\infty} \frac{-\xi'_\tau(u)Lu}{u_\tau} u_\tau^{\frac{n-3}{n-2}} \psi^2 - \frac{1}{n-2} \int_{-\infty}^{\infty} \frac{V_+(u_\tau - u\xi'_\tau(u))}{u_\tau} u_\tau^{\frac{n-3}{n-2}} \psi^2 \leq \int_{-\infty}^{\infty} (\psi_s)^2 u_\tau^{\frac{n-3}{n-2}} \quad (3.17)$$

Consider the family of cutoffs  $\psi_R$  as above, so  $\psi_R \uparrow 1$  pointwise as  $R \rightarrow \infty$ . Taking into account that the first integral in the left-hand side of (3.21) is non-negative and the second is restricted to  $\{s \in \mathbb{R} : V_+(\gamma(s)) \neq 0\}$ , a compact set since  $\gamma$  diverges, letting  $R \rightarrow \infty$  and applying monotone and Lebesgue convergence theorems we get

$$\int_{-\infty}^{\infty} \frac{-\xi'_\tau(u)Lu}{u_\tau} u_\tau^{\frac{n-3}{n-2}} - \int_{-\infty}^{\infty} \frac{V_+(u_\tau - u\xi'_\tau(u))}{u_\tau} u_\tau^{\frac{n-3}{n-2}} \leq 0.$$

In particular, the first integral is finite. Letting  $\tau \rightarrow \infty$  and applying Fatou's Lemma (to the first integral) and Lebesgue's Theorem (to the second), we conclude

$$\int_{-\infty}^{\infty} \frac{-Lu}{u} u^{\frac{n-3}{n-2}} \leq 0.$$

thus  $Lu \equiv 0$  along  $\gamma$ . The rest of the argument follows verbatim the one in Theorem 3.3.  $\square$

**Remark 3.6.** Theorem 3.5 should be compared to [38, Corollary 4.2]. There, the authors assume (3.16) with  $V \geq 0$  and the polynomial volume growth

$$|B_r| \leq Cr^{2(n-1)} \quad \text{for } r \geq 1,$$

a condition satisfied when  $V_+$  is compactly supported because of Bishop-Gromov volume comparison. Applying [38] we deduce that, in the setting of Theorem 3.5 with  $V \geq 0$ , either  $M$  has only one *non-parabolic* end, or  $M$  splits as Example 3.7 below.

To further comment on the assumptions in Theorem 3.3 and introduce our next result, denote by  $\lambda_1(M)$  the bottom of the Laplace spectrum on  $M$ .

**Example 3.7.** This example is taken from [38, Proposition 6.1]. Given  $n \geq 3$ , a manifold  $(P^{n-1}, h)$  and a function  $0 < \eta \in C^\infty(\mathbb{R})$ , consider the warped product

$$M = \mathbb{R} \times P, \quad g = dt^2 + \eta(t)^2 h$$

Under the assumptions

$$\eta'' > 0, \quad (n-2)(\log \eta)'' + \eta^{-2} \text{Ric}_P \geq 0,$$

the manifold  $M$  satisfies

$$\text{Ric} \geq -\frac{n-1}{n-2} V g \quad \text{with} \quad V = (n-2) \frac{\eta''}{\eta}.$$

The function

$$v = \int_0^t \frac{ds}{\eta(s)^{n-1}}$$

is harmonic and  $w = |\nabla v|^{\frac{n-2}{n-1}} = c \cdot \eta(t)^{2-n}$  is a positive solution to

$$Lw \doteq \Delta w + Vw = 0, \quad \text{so } L \geq 0.$$

Therefore, if  $\eta$  is bounded from below by a positive constant on  $\mathbb{R}$  and  $P$  is compact, by Theorem 3.3 the operator  $L$  is critical. An example is given by the choices

$$P \text{ compact with } \text{Ric}_P \geq -(n-2)h, \quad \eta(t) = \text{ch } t$$

considered in [35]. In this case, the resulting warped product metric  $dt^2 + (\text{ch } t)^2 h$  satisfies

$$\text{Ric} \geq -(n-1)g, \quad \Delta + (n-2) \geq 0 \quad \text{and is critical on } M.$$

In particular,  $\lambda_1(M) = n-2$ .

**Example 3.8.** The following example shows that a growth condition on  $w$  in Theorem 3.3 is necessary to prove criticality in dimension  $n \geq 4$ . Consider in Example 3.7 the choices

$$P \text{ compact with } \text{Ric}_P \geq 0, \quad \eta(t) = e^t,$$

see [35, Example 2.2]. The resulting warped product metric  $dt^2 + e^{2t}h$  satisfies

$$\text{Ric} \geq -(n-1)g, \quad L \doteq \Delta + (n-2) \geq 0.$$

In particular,  $\lambda_1(M) \geq n-2$ . However, the positive solution  $w = e^{(2-n)t}$  to  $Lw = 0$  singled out in Example 3.7 does not satisfy (3.10), thus in dimension  $n \geq 4$  Theorem 3.3 is not applicable to guarantee that  $L$  is critical. Indeed, we show that  $L$  is subcritical on  $M$ . To see this, for functions  $u$  of  $t$  only, equation  $\Delta u + (n-2)u$  becomes

$$u''(t) + (n-1)u'(t) + (n-2)u(t) = 0,$$

whose general solution is spanned by  $\{e^{-(n-2)t}, e^{-t}\}$  in dimension  $n \geq 4$  and by  $\{e^{-t}, te^{-t}\}$  in dimension 3. Thus, if  $n \geq 4$  the operator  $L$  admits the positive solutions  $e^{-(n-2)t}$  and  $e^{-t}$  which are not proportional, hence it is subcritical. On the other hand, if  $n = 3$  the only positive solution just depending on  $t$  is, up to scaling, the function  $e^{-t}$ . In fact, Theorem 3.3 guarantees that  $L$  is critical.

Let us examine this example further. The function

$$\hat{w} = e^{-\frac{n-1}{2}t} \quad \text{solves} \quad \Delta \hat{w} + \frac{(n-1)^2}{4} \hat{w} = 0;$$

hence,

$$\lambda_1(M) \geq \frac{(n-1)^2}{4} \tag{3.18}$$

(in fact, equality holds in (3.18) by Cheng's eigenvalue estimate). Equivalently, since  $\text{Ric} \geq -(n-1)$  this can be rewritten as follows:

$$\text{Ric} \geq -\frac{4}{n-1} \hat{V}, \quad \hat{L} \doteq \Delta + \hat{V} \geq 0 \quad \text{with } \hat{V} = \frac{(n-1)^2}{4}.$$

Note that  $\frac{(n-1)^2}{4} \geq (n-2)$ , with equality iff  $n = 3$ . We claim that  $\hat{L}$  is critical, and to see this we check each end separately. Let  $E^{(1)} = (-\infty, 0) \times P$  and  $E^{(2)} = (1, \infty) \times P$  be the ends with respect to  $K = [0, 1] \times P$ . Consider  $\hat{w}$  as a supersolution, and let  $u^{(i)}$  be the minimal solution on  $E^{(i)}$  with respect to  $\hat{w}$ . Then, by construction  $u^{(i)} = \lim_{j \rightarrow \infty} u_j^{(i)}$  where

$$\begin{cases} \hat{L}u_j^{(1)} = 0 & \text{on } [-j, 0] \times P, \\ u_j^{(1)} = 1 & \text{on } \{0\} \times P, \\ u_j^{(1)} = 0 & \text{on } \{-j\} \times P. \end{cases} \quad \begin{cases} \hat{L}u_j^{(2)} = 0 & \text{on } [1, j] \times P, \\ u_j^{(2)} = e^{-\frac{n-1}{2}} & \text{on } \{1\} \times P, \\ u_j^{(2)} = 0 & \text{on } \{j\} \times P. \end{cases}$$

By uniqueness of solutions,  $u_j^{(i)}$  only depends on  $t$  (since its average over  $P$  is still a solution). The general solution to the ODE

$$\hat{L}u = u''(t) + (n-1)u'(t) + \frac{(n-1)^2}{4}u(t) = 0$$

is  $e^{-\frac{n-1}{2}t}(a+bt)$  with  $a, b \in \mathbb{R}$ , whence

$$u_j^{(1)}(t) = \frac{j+t}{j} e^{-\frac{n-1}{2}t}, \quad u_j^{(2)}(t) = \frac{j-t}{j-1} e^{-\frac{n-1}{2}t}.$$

Letting  $j \rightarrow \infty$  we get  $u^{(1)} = \hat{w}$  and  $u^{(2)} = \hat{w}$ , therefore  $\hat{L}$  is  $\hat{w}$ -critical on each end. By Proposition 2.9,  $\hat{L}$  is critical on  $M$ .

Example 3.8 may suggest that if

$$\text{Ric} \geq -\frac{4}{n-1}Vg \quad \text{on } M, \quad L \doteq \Delta + V \geq 0,$$

then the dichotomy in the conclusion of Theorem 3.3 holds in any dimension  $n \geq 3$  without the growth condition (3.10). It is interesting, in this respect, to compare to [38, Theorem C] and to the detailed ODE analysis in Sections 6 and 8 therein. In the next result, we are able to prove the following stronger statement when  $\beta < \frac{4}{n-1}$ :

**Theorem 3.9.** *Let  $(M, g)$  be a complete Riemannian manifold of dimension  $n \geq 3$ , and assume that there exists  $V \in C_{\text{loc}}^{0,\alpha}(M)$  such that*

$$\text{Ric} \geq -\beta Vg \quad \text{on } M, \quad L \doteq \Delta + V \geq 0, \quad (3.19)$$

where either

$$0 < \beta < \frac{4}{n-1}, \quad \text{or}$$

$$\frac{4}{n-1} \leq \beta < \frac{n-1}{n-2} \quad \text{and } V_+ \text{ is compactly supported.}$$

Then, either

- (i)  $M$  has only one end, or
- (ii)  $V \equiv 0$  and  $M = \mathbb{R} \times P$  with the product metric, for some compact  $P$  with  $\text{Ric}_P \geq 0$ .

*Proof.* We first examine the case  $\beta \in \left(0, \frac{4}{n-1}\right)$ . Let us choose  $\sigma \in \left(\beta, \frac{4}{n-1}\right)$  and set

$$\hat{V} \doteq \frac{\beta}{\sigma}V.$$

Then, by Theorem 2.7

$$\text{Ric} \geq -\sigma\hat{V}g, \quad \hat{L} \doteq \Delta + \hat{V} \geq 0. \quad (3.20)$$

Moreover, since  $\Delta \geq 0$  and  $\beta/\sigma \in (0, 1)$ ,  $\hat{L}$  is subcritical unless  $\hat{V} \equiv 0$ , that is, unless  $V \equiv 0$ . Let  $0 < u \in C_{\text{loc}}^{2,\alpha}(M)$  solve  $\hat{L}u \leq 0$ , and consider the conformal deformation  $\bar{g} = u^{2\sigma}g$ . If we assume that  $M$  has at least two ends, using Proposition 3.2 we guarantee the existence of a  $\bar{g}$ -line  $\gamma : (T, T^*) \rightarrow M$  in  $(M, \bar{g})$ . By (ii) in Proposition 3.2 and since  $g$  is complete, it follows that once reparametrized by  $g$ -arclength  $s$  the curve  $\gamma$  is defined on the whole of  $\mathbb{R}$ . Proposition 3.1 implies

$$\begin{aligned} & \sigma \int_{-\infty}^{\infty} \frac{-\hat{L}u}{u} \frac{\eta^2}{u^\sigma} + \int_{-\infty}^{\infty} \left( \text{Ric}(\gamma_s, \gamma_s) + \sigma\hat{V} \right) \frac{\eta^2}{u^\sigma} \\ & \leq (n-1) \int_{-\infty}^{\infty} \frac{(\eta_s)^2}{\eta^2} \frac{\eta^2}{u^\sigma} - \sigma \int_{-\infty}^{\infty} \frac{(u_s)^2}{u^2} \frac{\eta^2}{u^\sigma} - \sigma(n-2) \int_{-\infty}^{\infty} \frac{u_s}{u} \left( \frac{\eta^2}{u^\sigma} \right)_s, \end{aligned}$$

holds for every  $\eta \in C_c^1(\mathbb{R})$ . Writing  $\eta = u^{\sigma/2}\psi$  and rearranging we get

$$\begin{aligned} & \sigma \int_{-\infty}^{\infty} \frac{-\hat{L}u}{u} \psi^2 + \int_{-\infty}^{\infty} \left( \text{Ric}(\gamma_s, \gamma_s) + \sigma \hat{V} \right) \psi^2 \\ & \leq \sigma \left[ \frac{(n-1)\sigma}{4} - 1 \right] \int_{-\infty}^{\infty} \frac{(u_s)^2}{u^2} \psi^2 - \sigma(n-3) \int_{-\infty}^{\infty} \frac{u_s \psi \psi_s}{u} + (n-1) \int_{-\infty}^{\infty} (\psi_s)^2. \end{aligned}$$

By (3.20) and since  $\sigma < \frac{4}{n-1}$ , applying Young's inequality to the second term of the right-hand side we deduce the inequality

$$\int_{-\infty}^{\infty} \frac{-\hat{L}u}{u} \psi^2 \leq C \int_{-\infty}^{\infty} (\psi_s)^2$$

for some constant  $C > 0$ . By choosing an even cut-off  $0 \leq \psi_R \leq 1$  such that  $\psi_R \equiv 1$  on  $[-R, R]$ ,  $\psi_R \in C_c^\infty((-2R, 2R))$  and  $|(\psi_R)_s| \leq 4/R$ , letting  $R \rightarrow \infty$  we deduce that  $\hat{L}u = 0$  along  $\gamma$ . As  $u$  can be any supersolution, we actually proved that any  $C_{\text{loc}}^{2,\alpha}$  supersolution  $u$  satisfy  $\hat{L}u = 0$  on some curve. If  $\hat{L}$  were subcritical, then by Theorem 2.3 there would exist  $0 < W \in C(M)$  such that  $\hat{L} + W \geq 0$ . By approximating  $W$  from below, we can assume that  $W \in C^\infty(M)$ . It would therefore exist a positive solution  $u \in C_{\text{loc}}^{2,\alpha}(M)$  to  $\hat{L}u = -Wu < 0$  on the entire  $M$ , contradiction. We conclude that  $\hat{L}$  is critical, thus  $V \equiv 0$ . The conclusion in (ii) thus follows from Cheeger-Gromoll's splitting theorem.

If  $\beta \in \left[ \frac{4}{n-1}, \frac{n-1}{n-2} \right)$  and  $V_+$  is compactly supported, fix  $\sigma \in \left( \beta, \frac{n-1}{n-2} \right)$  and  $\hat{V}, \hat{L}$  as above. We follow the proof of Theorem 3.5 with  $\sigma$  replacing  $\frac{n-1}{n-2}$  up to (3.21), which now becomes (see also (3.14))

$$\frac{\sigma}{n-1} \int_{-\infty}^{\infty} \frac{-\xi'_\tau(u) \hat{L}u}{u_\tau} u_\tau^{\frac{\sigma(n-3)}{n-1}} \psi^2 - \frac{\sigma}{n-1} \int_{-\infty}^{\infty} \frac{\hat{V}_+(u_\tau - u \xi'_\tau(u))}{u_\tau} u_\tau^{\frac{\sigma(n-3)}{n-1}} \psi^2 \leq \int_{-\infty}^{\infty} (\psi_s)^2 u_\tau^{\frac{\sigma(n-3)}{n-1}}. \quad (3.21)$$

Proceeding as in Theorem 3.5 we deduce that  $\hat{L}$  is critical. The conclusion follows as in the first part of the proof.  $\square$

As a corollary, we have the following obstruction theorem:

**Corollary 3.10.** *Let  $(M, g)$  be a compact Riemannian manifold of dimension  $n \geq 2$ , and assume that there exists  $0 \leq V \in C_{\text{loc}}^{0,\alpha}(M)$  such that*

$$\text{Ric} \geq -\beta V g \quad \text{on } M, \quad L \doteq \Delta + V \geq 0, \quad (3.22)$$

where

$$0 < \beta < \frac{4}{n-1}.$$

Then,  $V \equiv 0$ .

*Proof.* Consider the manifold  $\hat{M} = M \times \mathbb{R}$  with the product metric  $\hat{g}$  and the natural projection  $\pi : \hat{M} \rightarrow M$  onto the first factor. Since  $V \geq 0$ , the Ricci curvature of  $M$  satisfies  $\widehat{\text{Ric}} \geq -\beta(V \circ \pi)\hat{g}$ . Moreover, by lifting a positive solution to  $Lu \leq 0$  we have

$$\hat{\Delta} + (V \circ \pi) \geq 0.$$

By applying Theorem 3.9 to  $\hat{M}$  we conclude that  $V \circ \pi \equiv 0$ .  $\square$

**Remark 3.11.** Corollary 3.10 can be seen as an extension of Cheng's eigenvalue estimate: by taking  $V = \lambda_1(M) > 0$ , we deduce the non-existence of compact manifolds for which

$$\text{Ric} \geq -\beta \lambda_1(M)g \quad \text{with } 0 < \beta < \frac{4}{n-1}.$$

In other words, a compact manifold with  $\text{Ric} \geq -(n-1)$  must satisfy

$$\lambda_1(M) \leq (n-1) \frac{n-1}{4} = \frac{(n-1)^2}{4}.$$

In general, in case (i) of Theorem 3.9 the metric  $\bar{g} = u^{2\beta}g$  may not be complete. To see this, consider Euclidean space,  $V \equiv 0$  and  $u$  a smoothing of the Green kernel centered at the origin:

$$u(x) = \tau(|x|^{2-n})$$

for a concave function  $\tau : [0, \infty) \rightarrow \mathbb{R}$  satisfying  $\tau(t) = t$  for  $t \leq 1$  and  $\tau \equiv 2$  on  $[4, \infty)$ . Then,  $\Delta u \leq 0$  on  $\mathbb{R}^n$  but  $\bar{g}$  is incomplete for each

$$\beta \in \left( \frac{1}{n-2}, \frac{4}{n-1} \right).$$

The next theorem shows that for  $\beta < (n-2)^{-1}$  the fact does not occur.

**Theorem 3.12.** *Let  $(M, g)$  be a complete Riemannian manifold of dimension  $n \geq 3$ , and assume that there exists  $V \in C_{\text{loc}}^{0,\alpha}(M)$  such that*

$$\text{Ric} \geq -\beta Vg \quad \text{on } M, \quad L \doteq \Delta + V \geq 0 \quad (3.23)$$

with

$$0 < \beta < \frac{1}{n-2}.$$

Then, for each  $0 < u \in C^2(M)$  solving  $Lu \leq 0$ , the metric  $\bar{g} = u^{2\beta}g$  is complete and satisfies

$$\overline{\text{Ric}}_f^N \geq \text{Ric} + \beta Vg \geq 0,$$

where

$$f = (n-2)\beta \log u \quad \text{and} \quad N = n + \frac{\beta(n-2)}{1-\beta(n-2)} > n.$$

**Remark 3.13.** In the borderline case  $\beta = \frac{1}{n-2}$ , the same computations show that the  $\infty$ -Bakry-Emery Ricci tensor  $\overline{\text{Ric}}_f$  is non-negative on  $M$ . However, in this case we are not able to prove that  $\bar{g}$  is complete, nor we are aware of counterexamples to this fact.

*Proof.* By [41, Lemma 2.2], one can construct a "shortest ray"  $\gamma$  in  $(M, \bar{g})$  issuing from a fixed origin  $o$ , that is, a divergent curve minimizing  $\bar{g}$ -distance between any pair of its points, with the property that  $(M, \bar{g})$  is complete if and only if  $\gamma$  has infinite  $\bar{g}$ -length. Parametrizing  $\gamma$  by  $g$ -arclength  $s$ , since  $g$  is complete  $\gamma$  is defined for  $s \in [0, \infty)$ ; hence, to get completeness we only have to check that

$$\ell_{\bar{g}}(\gamma_{[0,\infty)}) = \int_0^\infty u^\beta ds = \infty.$$

Since  $\gamma$  is  $\bar{g}$ -minimizing, by equation (3.6) in Proposition 3.1 we deduce, using the hypothesis  $Lu \leq 0$  and  $\text{Ric} \geq -\beta Vg$ , that for every smooth function  $\eta$  with compact support in  $[0, \infty)$  and vanishing at 0, and for every  $\beta > 0$ ,

$$0 \leq (n-1) \int_0^\infty (\eta_s)^2 u^{-\beta} - \beta \int_0^\infty \frac{(u_s)^2}{u^2} \eta^2 u^{-\beta} - \beta(n-2) \int_0^\infty \frac{u_s}{u} (\eta^2 u^{-\beta})_s. \quad (3.24)$$

Writing  $\eta = u^\beta \psi$ , we have

$$\begin{aligned}\eta^2 u^{-\beta} &= u^\beta \psi^2, \\ \eta_s &= \beta \psi u^{\beta-1} u_s + u^\beta \psi_s, \\ (\eta_s)^2 u^{-\beta} &= \beta^2 \psi^2 u^{\beta-2} (u_s)^2 + u^\beta (\psi_s)^2 + 2\beta \psi \psi_s u^{\beta-1} u_s,\end{aligned}$$

and substituting in equation (3.24) we get

$$(n-1) \int_0^\infty u^\beta (\psi_s)^2 \geq -2\beta \int_0^\infty \psi \psi_s u^{\beta-1} u_s + \beta(1-\beta) \int_0^\infty \psi^2 u^{\beta-2} (u_s)^2. \quad (3.25)$$

Now we follow the same lines of the computations in [16, Lemma 2.3]. Define

$$I \doteq \int_0^\infty \psi \psi_s u^{\beta-1} u_s$$

and observe that

$$I = \frac{1}{\beta} \int_0^\infty \psi \psi_s (u^\beta)_s = -\frac{1}{\beta} \int_0^\infty u^\beta (\psi_s)^2 - \frac{1}{\beta} \int_0^\infty \psi \psi_{ss} u^\beta.$$

Moreover, for every  $t > 1$  and using Young's inequality with  $\varepsilon > 0$  to be chosen later, we have

$$\begin{aligned}2\beta I &= 2\beta t I + 2\beta(1-t)I \\ &= -2t \int_0^\infty u^\beta (\psi_s)^2 - 2t \int_0^\infty \psi \psi_{ss} u^\beta + 2\beta(1-t) \int_0^\infty \psi \psi_s u^{\beta-1} u_s \\ &\leq -2t \int_0^\infty u^\beta (\psi_s)^2 - 2t \int_0^\infty \psi \psi_{ss} u^\beta \\ &\quad + \beta(t-1)\varepsilon \int_0^\infty \psi^2 u^{\beta-2} (u_s)^2 + \frac{\beta(t-1)}{\varepsilon} \int_0^\infty u^\beta (\psi_s)^2.\end{aligned}$$

Recalling that  $\beta < (n-2)^{-1}$  and choosing

$$\varepsilon \doteq \frac{1-\beta}{t-1}$$

we obtain

$$\begin{aligned}2\beta I &\leq -2t \int_0^\infty \psi \psi_{ss} u^\beta + \beta(1-\beta) \int_0^\infty \psi^2 u^{\beta-2} (u_s)^2 \\ &\quad + \left[ \frac{\beta(t-1)^2}{1-\beta} - 2t \right] \int_0^\infty u^\beta (\psi_s)^2.\end{aligned}$$

From (3.25) we get

$$0 \leq \left[ \frac{\beta(t-1)^2}{1-\beta} - 2t + (n-1) \right] \int_0^\infty u^\beta (\psi_s)^2 - 2t \int_0^\infty \psi \psi_{ss} u^\beta \quad (3.26)$$

for every  $t > 1$  and every positive  $\beta < (n-2)^{-1}$ . Let

$$P(t, n, \beta) \doteq \frac{\beta(t-1)^2}{1-\beta} - 2t + (n-1) = \frac{1}{1-\beta} [\beta t^2 - 2t + (n-1) - \beta(n-2)];$$



it is not difficult to see that there exists  $\bar{t} = \bar{t}(\beta, n) > 1$  such that, for  $n \geq 3$ ,  $P(\bar{t}, n, \beta) < 0$ . Therefore, from (3.26) we deduce

$$0 \leq - \int_0^\infty u^\beta (\psi_s)^2 - C \int_0^\infty u^\beta \psi \psi_{ss}$$

for every  $\psi$  smooth with compact support in  $[0, \infty)$  and vanishing at 0, and for some positive constant  $C$  depending on  $\bar{t}$ ,  $n$  and  $\beta$ . Now we choose  $\psi = s\varphi$  with  $\varphi$  smooth with compact support in  $(0, +\infty)$ : thus

$$\psi_s = \varphi + s\varphi_s, \quad \psi_{ss} = 2\varphi_s + s\varphi_{ss},$$

and we get

$$\int_0^\infty u^\beta \varphi^2 \leq \int_0^\infty u^\beta (-2(C+1)s\varphi\varphi_s - Cs^2\varphi\varphi_{ss} - s^2(\varphi_s)^2).$$

Choose  $\varphi$  such that  $\varphi \equiv 1$  on  $[0, R]$ ,  $\varphi \equiv 0$  on  $[2R, +\infty)$  and with  $|\varphi_s|$  and  $|\varphi_{ss}|$  bounded by  $\tilde{C}/R$  and  $\tilde{C}/R^2$ , respectively, for  $R \leq s \leq 2R$  ( $\tilde{C}$  is a positive constant). Then

$$\int_0^R u^\beta \leq \int_0^\infty u^\beta \varphi^2 \leq C \int_R^\infty u^\beta$$

for some  $C > 0$  independent of  $R$ . We conclude that necessarily

$$\int_0^\infty u^\beta = \infty,$$

i.e.  $\bar{g} = u^{2\beta}g$  is complete.  $\square$

Two consequences of the previous theorem are the following topological properties. In the compact case, using the results in [52, 40, 62], we have the following

**Corollary 3.14.** *Let  $(M, g)$  be a compact Riemannian manifold of dimension  $n \geq 3$ , and assume that there exists  $V \in C_{\text{loc}}^{0,\alpha}(M)$  such that*

$$\text{Ric} \geq -\beta V g \quad \text{on } M, \quad L \doteq \Delta + V \geq 0, \quad (3.27)$$

where

$$0 < \beta \leq \frac{1}{n-2}.$$

Then:

- (i)  $b_1(M) \leq n$ . Equality holds iff  $M$  is a flat torus and  $V \equiv 0$ .
- (ii)  $\pi_1(M)$  has a free abelian subgroup of finite index of rank  $\leq n$ , with equality iff  $M$  is a flat torus and  $V \equiv 0$ .
- (iii)  $\pi_1(M)$  is finite if  $\text{Ric} + \beta V g > 0$  at one point.

*Proof.* Let  $0 < u \in C^2(M)$  solve  $Lu = 0$ . By Remark 3.13 the metric  $\bar{g} = u^{2\beta}g$  satisfies

$$\overline{\text{Ric}}_f \geq \text{Ric} + \beta V g \geq 0,$$

with  $f = (n-2)\beta \log u$ . Moreover, since  $M$  is compact  $\bar{g}$  is complete. By [62, Corollary 6.7]  $b_1(M) \leq n$  and  $\pi_1(M)$  has a free abelian subgroup of finite index of rank  $\leq n$ . Moreover, equality holds in any of the two inequalities if and only if  $f$  is constant (i.e.  $u$  is constant) and  $(M, \bar{g})$  is a flat torus, hence so is  $(M, g)$ . Moreover, in this case, from  $Lu = 0$ , we also conclude  $V \equiv 0$ . Property (iii) follows immediately from [62, Corollary 6.7].  $\square$

It is interesting to compare our theorem with those in [11, 13], where the authors assume similar but different spectral conditions to obtain topological conclusions. In the non-compact case, using [39, Theorem 1.6] we have the following:

**Corollary 3.15.** *Let  $(M, g)$  be a complete, non-compact, Riemannian manifold of dimension  $n \geq 3$ , and assume that there exists  $V \in C_{\text{loc}}^{0,\alpha}(M)$  such that*

$$\text{Ric} \geq -\beta V g \quad \text{on } M, \quad L \doteq \Delta + V \geq 0, \quad (3.28)$$

where

$$0 < \beta < \frac{1}{n-2}.$$

Then,  $H_{n-1}(M, \mathbb{Z}) = 0$  or  $\mathbb{Z}$ .

## 4 Applications to minimal hypersurfaces

Let  $M^n \rightarrow (N^{n+1}, \bar{g})$  be a complete, two-sided, minimally immersed hypersurface. Denote with a bar tensors associated to  $\bar{g}$ . Write

$$J_\delta \doteq \Delta + \delta \left( |A|^2 + \overline{\text{Ric}}(v, v) \right)$$

for the  $\delta$ -stability operator. Choose a Darboux frame  $\{e_i\}$  along  $M$ , and set  $e_0 = v$ . As the second fundamental form  $A$  in direction  $v$  is traceless, it satisfies the refined Kato inequality

$$\sum_{j=1}^n A_{1j}^2 \leq \frac{n-1}{n} |A|^2,$$

see [47, Lemma 1.5]. Moreover, equality holds at a point  $x$  where  $A \neq 0$  if and only if  $A(x)$  has only two eigenvalues  $\lambda, \mu$ , of multiplicities 1 and 2 respectively, and  $Ae_1 = \lambda e_1$ . By Gauss equations,

$$\text{Ric}_{11} = \sum_{\alpha \geq 2} \bar{R}_{1\alpha 1\alpha} - \sum_{j \geq 1} A_{1j}^2 \geq \sum_{\alpha \geq 2} \bar{R}_{1\alpha 1\alpha} - \frac{n-1}{n} |A|^2. \quad (4.1)$$

- Assume that  $\overline{\text{BRic}} \geq 0$ . Then,

$$\begin{aligned} \text{Ric}_{11} &\geq \sum_{\alpha \geq 2} \bar{R}_{1\alpha 1\alpha} - \frac{n-1}{n} |A|^2 = \overline{\text{BRic}}_{1v} - \overline{\text{Ric}}(v, v) - \frac{n-1}{n} |A|^2 \\ &\geq -\left( \overline{\text{Ric}}(v, v) + |A|^2 \right) + \frac{1}{n} |A|^2, \end{aligned} \quad (4.2)$$

- Assume that  $\overline{\text{Sec}} \geq 0$ . Then,

$$\text{Ric}_{11} \geq \sum_{\alpha \geq 2} \bar{R}_{1\alpha 1\alpha} - \frac{n-1}{n} |A|^2 \geq -\frac{n-1}{n} \left( \overline{\text{Ric}}(v, v) + |A|^2 \right). \quad (4.3)$$

We are ready to prove our results:

*Proof of Theorem 1.6.* Assume that  $M$  has at least two ends, for otherwise (i) holds. Defining  $V \doteq \overline{\text{Ric}}(v, v) + |A|^2$ , from (4.2) and the stability assumption we get

$$\text{Ric} \geq -Vg + \frac{1}{n} |A|^2 g, \quad \Delta + V \geq 0. \quad (4.4)$$

Since  $1 < \frac{4}{n-1}$  holds in dimension  $n \in \{3, 4\}$ , applying Theorem 3.9 we obtain that  $V \equiv 0$  and that  $M = \mathbb{R} \times P$  with the product metric  $ds^2 + g_P$ , for some compact  $(P, g_P)$  with non-negative Ricci curvature. From  $0 = \text{Ric}(\partial_s, \partial_s)$  and (4.4) we conclude  $|A| \equiv 0$ , thus  $\overline{\text{Ric}}(\nu, \nu) = V - |A|^2 \equiv 0$ . The conclusion  $\overline{\text{BRic}}(\partial_s, \nu) \equiv 0$  follows from (4.2). In case (i), if  $M$  is parabolic then one cannot deduce  $V \equiv 0$  (as the case of surfaces already suggests). However, if  $\overline{\text{Ric}} \geq 0$ , any positive solution to  $Lu = 0$  is superharmonic. Hence, the parabolicity of  $M$  forces  $u$  to be constant, so  $V \equiv 0$ .  $\square$

If  $\overline{\text{BRic}} \geq 0$  is strengthened to  $\overline{\text{Sec}} \geq 0$ , we gain one more dimension and further information in the previous result.

*Proof of Theorem 1.8.* The proof follows the same path as that of Theorem 1.6, we only point out the differences: setting  $V \doteq \text{Ric}(\nu, \nu) + |A|^2$ , Gauss equation (4.3) yields

$$\text{Ric} \geq -\frac{n-1}{n}V, \quad \Delta + V \geq 0.$$

Inequality  $\frac{n-1}{n} < \frac{4}{n-1}$  now holds up to  $n = 5$ , justifying the dimensional improvement. If  $M$  has more than one end, from  $V \equiv 0$  one deduces  $|A| \equiv 0$  and  $\overline{\text{Ric}}(\nu, \nu) \equiv 0$ . By Gauss equation,  $M$  has non-negative sectional curvature, hence so does  $(P, g_P)$ . If  $n = 3$ , the conclusion  $H^2(M, \mathbb{Z}) = 0$  or  $\mathbb{Z}$  follows by a direct application of Corollary 3.15, once observed that  $\frac{n-1}{n} < \frac{1}{n-2}$  holds in dimension 3.  $\square$

We are left to consider the case of  $\frac{n-2}{n}$ -stable hypersurfaces. Cheng & Zhou's strategy to obtain Theorem 1.3 is based on a clever refinement of [38, Theorem A], and exploits the Hardy inequality

$$\int_M \frac{(n-2)^2}{4r^2} \phi^2 \leq \int_M |\nabla \phi|^2 \quad \forall \phi \in \text{Lip}_c(M)$$

satisfied by minimal hypersurfaces in Euclidean space, see Remark 2.4. Crucial for the conclusion are both the completeness of the metric  $r^{-2}g$  on  $M$ , and a lower bound for the conformal factor  $r^{-2}$  in terms of the intrinsic distance on  $M$ . When trying to adapt the ideas to more general ambient spaces, this latter control is particularly difficult to achieve. In dimension  $n = 3$ , we can use Theorem 3.3 to overcome the problem and obtain information on  $1/3$ -stable hypersurfaces. The next theorem generalizes Theorem 1.4 in the Introduction.

**Theorem 4.1.** *Let  $M^3 \rightarrow (N^4, \bar{g})$  be a complete, non-compact, two-sided minimal hypersurface in a manifold satisfying  $\overline{\text{Sec}} \geq 0$ . Assume that  $M$  is  $\delta$ -stable with  $\delta \geq 1/3$ . Then, one of the following mutually exclusive cases occurs:*

- (i)  $M$  has only one end. Moreover, either  $M$  is non-parabolic, or  $M$  is parabolic, totally geodesic and  $\overline{\text{Ric}}(\nu, \nu) \equiv 0$ ;
- (ii)  $M$  is totally geodesic,  $\overline{\text{Ric}}(\nu, \nu) \equiv 0$  and  $M = \mathbb{R} \times P$  with the product metric, for some compact surface  $P$  with non-negative Gaussian curvature.
- (iii)  $\delta = 1/3$ ,  $\overline{\text{Ric}}(\nu, \nu) \equiv 0$ ,  $M$  has at least two ends and is non-parabolic, the set  $U \doteq \{|A| > 0\}$  is non-empty and  $A$  has only two eigenvalues  $\lambda, \mu$  on each connected component of  $U$ , of multiplicities 1 and 2 respectively. Moreover, if  $\{e_j\}$  is an orthonormal frame of eigenvectors with  $Ae_1 = \lambda e_1$  and  $Ae_\alpha = \mu e_\alpha$  for  $\alpha \geq 2$ , and setting  $e_0 = \nu$ , the components of the curvature tensor of  $N$  satisfy

$$\bar{R}_{1\alpha 1\alpha} = \bar{R}_{1010} = \bar{R}_{\alpha 0\alpha 0} = \bar{R}_{\alpha 0\alpha 1} = \bar{R}_{101\alpha} = \bar{R}_{\alpha\beta 0\beta} = \bar{R}_{\alpha 0\beta 0} = 0. \quad (4.5)$$

*Proof.* The restriction to dimension  $n = 3$  is only needed at one step of the proof, hence we prefer to keep writing  $n$ . The assumed  $\delta$ -stability and  $\overline{\text{Sec}} \geq 0$  therefore imply

$$J_{\frac{n-2}{n}} \doteq \Delta + \frac{n-2}{n} \left( |A|^2 + \overline{\text{Ric}}(v, v) \right) \geq 0.$$

Assume that  $M$  has at least two ends, for otherwise (i) holds. Define

$$L = \Delta + \frac{n-2}{n} |A|^2, \quad V \equiv \frac{n-2}{n} |A|^2.$$

By Gauss equations (4.1),

$$\text{Ric}_{11} = \sum_{\alpha \geq 2} \bar{R}_{1\alpha 1\alpha} - \sum_{j \geq 1} A_{1j}^2 \geq -\frac{n-1}{n} |A|^2 = -\frac{n-1}{n-2} V$$

On the other hand,

$$L \geq J_{\frac{n-2}{n}} \geq 0. \tag{4.6}$$

Therefore, applying Theorem 3.3 we deduce that  $L$  is critical and that, for each  $x \in M$ , there exists an orthonormal basis  $\{e_j\}$  whose first element  $e_1$  satisfies

$$\text{Ric}_{11} = \sum_{j \geq 1} A_{1j}^2 = -\frac{n-1}{n} |A|^2.$$

Namely,  $\bar{R}_{1\alpha 1\alpha} = 0$  for each  $\alpha \geq 2$ , and  $A$  satisfies equality in the refined Kato inequality. From (4.6) and the criticality of  $L$ ,

$$\overline{\text{Ric}}(v, v) \equiv 0 \quad \text{on } M.$$

If  $\delta > \frac{n-2}{n}$ , we deduce from

$$L \geq J_\delta \geq J_{\frac{n-2}{n}} \geq 0$$

and the criticality of  $L$  that  $J_\delta = J_{\frac{n-2}{n}}$ , which means  $|A| \equiv 0$ . From Gauss equations we deduce that  $M$  has non-negative sectional curvature. Since  $M$  has two ends, the splitting theorem implies  $M = \mathbb{R} \times P$  for some compact manifold  $P$  with non-negative sectional curvature.

It remains to consider the case  $\delta = \frac{n-2}{n}$  and  $U \doteq \{|A| > 0\} \neq \emptyset$ . Since  $A$  satisfies equality in the refined Kato inequality, we can choose an orthonormal frame  $\{e_i\}$  so that, at a given fixed point,

$$A_{1\alpha} = 0, \quad A_{11} = \lambda, \quad A_{\alpha\beta} = \mu \delta_{\alpha\beta} \quad \text{with } \lambda + (n-1)\mu = 0. \tag{4.7}$$

In each connected component of  $U$ , since  $\lambda \neq \mu$  the two eigenspace distributions have constant multiplicity and are therefore smooth. Thus, an orthonormal frame satisfying (4.7) can be chosen smoothly around any given point. Furthermore, from  $\bar{R}_{1\alpha 1\alpha} = 0$  and  $\overline{\text{Ric}}(v, v) \equiv 0$  we deduce

$$0 = \bar{R}_{1\alpha 1\alpha} = \bar{R}_{1010} = \bar{R}_{\alpha 0\alpha 0} \quad \forall \alpha \geq 2.$$

Setting

$$\begin{aligned} F(t, s) &= \bar{R}(e_1 + te_0, e_\alpha + se_0, e_1 + te_0, e_\alpha + se_0) \\ &= \bar{R}_{1\alpha 1\alpha} + 2t\bar{R}_{0\alpha 1\alpha} + 2s\bar{R}_{101\alpha} + t^2\bar{R}_{0\alpha 0\alpha} + 2ts\bar{R}_{0\alpha 10} + s^2\bar{R}_{1010} + o(\sqrt{t^2 + s^2}) \end{aligned}$$

from  $F(0, 0) = 0$  and  $F \geq 0$  we get

$$\begin{aligned} DF(0, 0) = 0 &\implies \bar{R}_{0\alpha 1\alpha} = 0 = \bar{R}_{10 1\alpha} \\ D^2F(0, 0) \geq 0 &\implies \bar{R}_{0\alpha 10} = 0. \end{aligned}$$

Applying an analogous argument switching the roles of  $e_1$  and  $e_0$ , we infer  $\bar{R}_{1\alpha 01} = 0$ . Next, considering

$$F(t) = \bar{R}(e_\beta, e_0 + te_\alpha, e_\beta, e_0 + te_\alpha) = \bar{R}_{\beta 0 \beta 0} + 2t\bar{R}_{\beta\alpha\beta 0} + t^2\bar{R}_{\beta\alpha\beta\alpha}.$$

From  $F(0) = 0$  and  $F \geq 0$  we deduce

$$F'(0) = 0 \implies \bar{R}_{\beta\alpha\beta 0} = 0.$$

Eventually, by using

$$F(t) = \bar{R}(e_\beta + te_\alpha, e_0, e_\beta + te_\alpha, e_0) = \bar{R}_{\beta 0 \beta 0} + 2t\bar{R}_{\beta 0 \alpha 0} + t^2\bar{R}_{\alpha 0 \alpha 0}$$

we deduce from  $F'(0) = 0$  that  $\bar{R}_{\beta 0 \alpha 0} = 0$ , concluding the proof.  $\square$

**Remark 4.2.** Theorem 4.1 holds in dimension  $n \geq 4$ , with  $(n-2)/n$  replacing  $1/3$  and principal eigenspaces of dimensions 1 and  $(n-1)$ , whenever  $M$  supports a positive solution to  $J_{\frac{n-2}{n}} w \leq 0$  satisfying

$$w(x) = o\left(r(x)^{\frac{n-2}{n-3}}\right) \quad \text{as } r(x) \rightarrow \infty.$$

However, to present we do not have a manageable geometric condition to guarantee the existence of such  $w$ .

It is unclear to us whether to expect the vanishing of the mixed components  $\bar{R}_{01\alpha\beta}$  or not. If this were the case, by the Codazzi-Mainardi equations  $A$  would be a Codazzi tensor with only two eigenvalues. One may therefore obtain a local (likely, global) splitting of  $M$  as a warped product. This is the case of ambient Euclidean space:

*Proof of Theorem 1.4.* Denote by  $f : M \rightarrow \mathbb{R}^4$  the minimal immersion. If  $M$  has at least two ends, by Theorem 4.1 and since case (ii) does not occur in Euclidean space we deduce that  $M$  is non-parabolic, that  $U \doteq \{|A| > 0\}$  is non-empty and that  $A$  has only two eigenvalues on any connected, relatively compact subset  $U' \subset U$ . By a result due to Do Carmo and Dajczer [25, Theorem 4.4],  $f(U')$  is a piece of a catenoid  $C$ , and by continuity so is  $f(\bar{U}')$ . If by contradiction  $U' \not\equiv M$ , then  $A$  should vanish on  $\partial U'$ . However, the second fundamental form of  $C$  is positive on the compact set  $\bar{f(U')} \supset f(\bar{U}')$ , contradiction. Therefore,  $U' \equiv M$  and  $f(M) \subset C$ . Since  $M$  is complete and  $f : M \rightarrow C$  is a local isometry, by Ambrose theorem  $f$  is surjective and a Riemannian covering, hence a diffeomorphism since the 3-dimensional catenoid is simply connected.  $\square$

*Proof of Theorem 1.2.* Up to passing to the universal covering, we can assume that  $M$  is simply connected. By Corollary 1.4, either  $M$  has only one end or  $M$  is the catenoid. However, the catenoid is not  $\delta$ -stable for  $\delta > 1/3$ , as can be deduced by the criticality of  $\Delta + 1/3|A|^2$  in Example 3.4. The conclusion follows from [33, Theorem 1.7].  $\square$

## Acknowledgements

The authors are members of GNSAGA (Gruppo Nazionale per le Strutture Algebriche, Geometriche e loro Applicazioni) and are partially supported by the PRIN project no. 20225J97H5 (Italy) “Differential-geometric aspects of manifolds via Global Analysis”.

**Conflict of Interest.** The authors have no conflict of interest.

## References

- [1] G. Antonelli, M. Pozzetta, K. Xu, *A sharp spectral splitting theorem*. Preprint. [7](#)
- [2] G. Antonelli, K. Xu, *New spectral Bishop-Gromov and Bonnet-Myers theorems and applications to isoperimetry*. Available at arXiv:2405.08918. [5](#)
- [3] M. Anderson, *The compactification of a minimal submanifold by its Gauss map*. Available at <http://www.math.sunysb.edu/~anderson/papers.html> [2](#)
- [4] M. Anderson, *On area-minimizing hypersurfaces in manifolds of nonnegative curvature*. Indiana Univ. Math. J. 32 (1983), 745–760.
- [5] F. Bei, G. Pipoli,  *$L^2$ -harmonic forms and spinors on stable minimal hypersurfaces*. Available at arXiv:2407.18836. [3](#)
- [6] C. Bellettini, *Extensions of Schoen–Simon–Yau and Schoen–Simon theorems via iteration à la De Giorgi*. Available at arXiv:2310.01340. [2](#)
- [7] P. Bérard, P. Castillon, *Inverse spectral positivity for surfaces*. Rev. Mat. Iberoam. 30 (2014), no.4, 1237–1264. [4](#), [7](#)
- [8] B. Bianchini, L. Mari, M. Rigoli, *Yamabe type equations with a sign-changing nonlinearity, and the prescribed curvature problem*. J. Differential Equations 260 (2016), 7416–7497. [8](#), [9](#)
- [9] B. Bianchini, L. Mari, M. Rigoli, *On some aspects of oscillation theory and geometry*. Mem. Amer. Math. Soc. 225 (2013), no. 1056, vi+195 pp. [9](#)
- [10] H.-D. Cao, Y. Shen, S. Zhu, *The structure of stable minimal hypersurfaces in  $\mathbb{R}^{n+1}$* . Math. Res. Lett. 4 (1997), no. 5, 637–644. [2](#), [5](#)
- [11] G. Carron, *Geometric inequalities for manifolds with Ricci curvature in the Kato class*. Ann. Inst. Fourier 69 (2019), no. 7, 3095–3167. [5](#), [9](#), [26](#)
- [12] G. Carron, *Inégalités de Hardy sur les variétés riemanniennes non-compactes*. J. Math. Pures Appl. (9) 76 (1997), no. 10, 883–891 [9](#)
- [13] G. Carron, C. Rose, *Geometric and spectral estimates based on spectral Ricci curvature assumptions*. J. Reine Angew. Math. 772 (2021), 121–145. [5](#), [26](#)
- [14] P. Castillon, *An inverse spectral problem on surfaces*. Comment. Math. Helv. 81 (2006), no. 2, 271–286. [7](#)
- [15] G. Catino, P. Mastrolia, D. Monticelli, *Uniqueness of critical metrics for a quadratic curvature functional*. Available at arXiv:2303.08025. [6](#)

- [16] G. Catino, P. Mastrolia, A. Roncoroni, *Two rigidity results for stable minimal hypersurfaces*. *Geom. Funct. Anal.* 34 (2024), no. 1, 1–18. [2](#), [6](#), [7](#), [24](#)
- [17] G. Catino, A. Roncoroni, *A closure result for globally hyperbolic spacetimes*. *Proc. Amer. Math. Soc.* 152 (2024), 5339–5354. [6](#)
- [18] X. Cheng, *One end theorem and application to stable minimal hypersurfaces*. *Arch. Math.* 90 (2008), 461–470. [3](#), [4](#), [6](#), [14](#)
- [19] X. Cheng, *On constant mean curvature hypersurfaces with finite index*. *Arch. Math. (Basel)* 86 (2006), no. 4, 365–374. [6](#)
- [20] X. Cheng, D. Zhou, *Manifolds with weighted Poincaré inequality and uniqueness of minimal hypersurfaces*. *Comm. Anal. Geom.* 17 (2009), no. 1, 139–154. [2](#)
- [21] O. Chodosh, C. Li, *Stable minimal hypersurfaces in  $\mathbb{R}^4$* . *Acta Math.* 233 (2024), no. 1, 1–31. [2](#)
- [22] O. Chodosh, C. Li, *Stable anisotropic minimal hypersurfaces in  $\mathbb{R}^4$* . *Forum Math. Pi* 11 (2023), Paper No. e3, 22 pp. [2](#)
- [23] O. Chodosh, C. Li, D. Stryker, *Complete stable minimal hypersurfaces in positively curved 4-manifolds*. *J. Eur. Math. Soc. (Online first)*, DOI 10.4171/JEMS/1499. [3](#)
- [24] O. Chodosh, C. Li, P. Minter, D. Stryker, *Stable minimal hypersurfaces in  $\mathbb{R}^5$* . Available at arXiv:2401.01492. [2](#), [6](#)
- [25] M.P. do Carmo, M. Dajczer, *Rotation Hypersurfaces in Spaces of Constant Curvature*. *Trans. Amer. Math. Soc.* 277 (1983), no. 2, 685–709. [17](#), [29](#)
- [26] J. Dodziuk,  *$L^2$  harmonic forms on complete manifolds*. *Ann. of Math. Stud.*, 102 (1982), 291–302. [3](#)
- [27] M.F. Elbert, B. Nelli, H. Rosenberg, *Stable constant mean curvature hypersurfaces*. *Proc. Amer. Math. Soc.* 135 (2007), no. 10, 3359–3366. [6](#), [12](#)
- [28] D. Fischer-Colbrie, *On complete minimal surfaces with finite Morse index in three-manifolds*. *Invent. Math.* 82 (1985), 121–132. [6](#), [7](#)
- [29] D. Fischer-Colbrie, R. Schoen, *The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature*. *Comm. Pure Appl. Math.* 33 (1980), no.2, 199–211. [4](#)
- [30] F. Fang, X.-D. Li, Z. Zhang, *Two generalizations of Cheeger–Gromoll splitting theorem via Bakry–Emery Ricci curvature*. *Ann. Inst. Fourier* 59 (2009), no. 2, 563–573.
- [31] H. Fu, Z. Li, *The structure of complete manifolds with weighted Poincaré’ inequalities and minimal hypersurfaces*. *International J. Math.* 21 (2010), 1–8. [2](#)
- [32] A.J. Hoffman, H.W. Wielandt, *The variation of the spectrum of a normal matrix*. *Duke Math. J.* 20 (1953), 37–39. [5](#)
- [33] H. Hong, H. Li, G. Wang, *On  $\delta$ -stable minimal hypersurfaces in  $\mathbb{R}^{n+1}$* . Available at arXiv:2407.03222. [2](#), [29](#)

- [34] P. Li, L.-F. Tam, *Harmonic functions and the structure of complete manifolds*. J. Differential Geom. 35 (1992), no. 2, 359–383. [3](#)
- [35] P. Li, J. Wang, *Complete manifolds with positive spectrum. I*. J. Differential Geom. 58 (2001), no. 3, 501–534. [5](#), [19](#), [20](#)
- [36] P. Li, J. Wang, *Complete manifolds with positive spectrum. II*. J. Differential Geom. 62 (2002), no. 1, 143–162. [5](#)
- [37] P. Li, J. Wang, *Stable minimal hypersurfaces in a nonnegatively curved manifold*. J. Reine Angew. Math. 566 (2004), 215–230. [3](#), [4](#), [5](#)
- [38] P. Li, J. Wang, *Weighted Poincaré inequality and rigidity of complete manifolds*. Ann. Sci. École Norm. Sup. 39 (4) (2006), no.6, 921–982. [5](#), [9](#), [19](#), [21](#), [27](#)
- [39] A. Lim, *The splitting theorem and topology of noncompact spaces with nonnegative  $N$ -Bakry Émery Ricci curvature*. Proc. Amer. Math. Soc. 149 (2021), no. 8, 3515–3529. [26](#)
- [40] J. Lott, *Some geometric properties of the Bakry–Emery–Ricci tensor*. Comment. Math. Helv. 78 (2003) 865–883. [25](#)
- [41] M. Magliaro, L. Mari, F. Roing, A. Savas-Halilaj, *Sharp pinching theorems for complete submanifolds in the sphere*. J. Reine Angew. Math. 814 (2024), 117–134. [6](#), [23](#)
- [42] L. Mari, M. Rigoli, A.G. Setti, *On the  $1/H$ -flow by  $p$ -Laplace approximation: new estimates via fake distances under Ricci lower bounds*. Amer. J. Math. 144 (2022), no. 3, 779–849.
- [43] L. Mazet, *Stable minimal hypersurfaces in  $\mathbb{R}^6$* . Available at arXiv:2405.14676. [2](#)
- [44] R. Miyaoka,  *$L^2$  harmonic 1-forms on a complete stable minimal hypersurface*. In Geometry and global analysis (Sendai, 1993), 289–293. Tohoku Univ., Sendai, 1993. [3](#), [4](#)
- [45] M. Murata, *Structure of positive solutions to  $(-\Delta + V)u = 0$  in  $\mathbb{R}^n$* . Duke Math. J. 53 (1986), no. 4, 869–943. [6](#), [8](#), [9](#)
- [46] S. Pigola, G. Veronelli, *Remarks on  $L^p$ -vanishing results in geometric analysis*. Internat. J. Math. 23 (2012), no. 1, 1250008, 18 pp. [2](#)
- [47] S. Pigola, M. Rigoli, A.G. Setti, *Vanishing and finiteness results in geometric analysis*. Progr. Math. 266, Birkhäuser Verlag, Basel, 2008, xiv+282 pp. [3](#), [8](#), [26](#)
- [48] Y. Pinchover, *On positive solutions of second-order elliptic equations, stability results, and classification*. Duke Math. J. 57, (1988), 955–980. [6](#), [8](#)
- [49] Y. Pinchover, *On criticality and ground states of second order elliptic equations, II*. J. Diff. Equ. 87 (1990), 353–364. [6](#), [8](#)
- [50] Y. Pinchover, K. Tintarev, *A ground state alternative for singular Schrödinger operators*. J. Funct. Anal. 230 (2006), no. 1, 65–77. [6](#), [8](#), [9](#)
- [51] Y. Pinchover, C. Tintarev, *Ground state alternative for  $p$ -Laplacian with potential term*. Calc. Var. Partial Differential Equations 28 (2007), no.2, 179–201. [6](#), [9](#), [10](#)



- [52] Z. Qian, *Estimates for weighted volumes and applications*. Quart. J. Math. Oxford Ser. (2) 48 (1997), no. 190, 235–242. 25
- [53] R. Schoen, S.-T. Yau, *Complete three dimensional manifolds with positive Ricci curvature and scalar curvature*. In Seminar on differential geometry, 209–228. Ann. Math. Stud. 102, Princeton Univ. Press, Princeton, NJ, 1982. 4, 6
- [54] R. Schoen, S.-T. Yau, *The existence of a black hole due to condensation of matter*. Comm. Math. Phys. 90 (1983), 575–579. 6
- [55] R. Schoen, S.-T. Yau, *Harmonic maps and the topology of stable hypersurfaces and manifolds with nonnegative Ricci curvature*. Comment. Math. Helvetici 51 (1976), 333–341. 5
- [56] Y. Shen, R. Ye, *On stable minimal surfaces in manifolds of positive bi-Ricci curvatures*. Duke Math. J. 85 (1996), no. 1, 109–116. 6, 12, 13
- [57] Y. Shen, R. Ye, *On the geometry and topology of manifolds of positive bi-Ricci curvature*. Available at arXiv:9708014 (1997). 6
- [58] Y. Shen, S. Zhu, *Rigidity of stable minimal hypersurfaces*. Math. Ann. 309 (1997), no. 1, 107–116. 3, 4, 6
- [59] Y.-B. Shen, X.-H. Zhu, *On stable complete minimal hypersurfaces in  $R^{m+1}$* . Amer. J. Math. 120 (1998), 103–116. 2, 6
- [60] L.-F. Tam, D. Zhou, *Stability properties for the higher dimensional catenoid in  $\mathbb{R}^{m+1}$* . Proc. Amer. Math. Soc. 137 (2009), 3451–3461. 2, 17
- [61] S. Tanno,  *$L^2$  harmonic forms and stability of minimal hypersurfaces*. J. Math. Soc. Japan. 48 (1996), 761–768 3
- [62] G. Wei, W. Wylie, *Comparison geometry for the Bakry-Emery Ricci tensor*. J. Differential Geom. 83 (2009), no. 2, 377–405. 25
- [63] T. Weidl, *Remarks on virtual bound states for semi-bounded operators*, Comm. Partial Differential Equations 24 (1999), 25–60. 9
- [64] Z. Zhao, *Subcriticality, positivity and gaugeability of the Schrödinger operator*. Bull. Am. Math. Soc. 23 (1990), no. 2, p. 513–517. 6, 8

G. Catino, DIPARTIMENTO DI MATEMATICA, POLITECNICO DI MILANO, PIAZZA LEONARDO DA VINCI 32, 20133, MILANO, ITALY.

*E-mail address*, G. Catino: giovanni.catino@polimit.it

L. Mari, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA CESARE SALDINI 50, 20133 MILANO, ITALY

*E-mail address*, L. Mari: luciano.mari@unimi.it

P. Mastrolia, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA CESARE SALDINI 50, 20133 MILANO, ITALY

*E-mail address*, P. Mastrolia: paolo.mastrolia@unimi.it

A. Roncoroni, DIPARTIMENTO DI MATEMATICA, POLITECNICO DI MILANO, PIAZZA LEONARDO DA VINCI 32, 20133, MILANO, ITALY.

*E-mail address*, A. Roncoroni: alberto.roncoroni@polimit.it