POLYNOMIALLY OSCILLATORY MULTIPLIERS ON GELFAND-SHILOV SPACES

ALEXANDRE ARIAS JUNIOR AND PATRIK WAHLBERG

ABSTRACT. We study continuity of the multiplier operator e^{iq} acting on Gelfand–Shilov spaces, where q is a polynomial on \mathbf{R}^d of degree at least two with real coefficients. In the parameter quadrant for the spaces we identify a wedge that depends on the polynomial degree for which the operator is continuous. We also show that in a large part of the complement region the operator is not continuous in dimension one. The results give information on well-posedness for linear evolution equations that generalize the Schrödinger equation for the free particle.

1. INTRODUCTION

Let q be a polynomial on \mathbf{R}^d with real coefficients. We consider in this paper the multiplication operator $T = T_g$ with $g(x) = e^{iq(x)}$ defined by

$$(Tf)(x) = e^{iq(x)}f(x), \quad x \in \mathbf{R}^d.$$

$$(1.1)$$

This operator is unitary on $L^2(\mathbf{R}^d)$ and obviously acts continuously on the Schwartz space $\mathscr{S}(\mathbf{R}^d)$ of smooth functions whose derivatives decay rapidly. The Schwartz space can equivalently be defined as the space of functions f that decay superpolynomially at infinity, plus the same condition on the Fourier transform \hat{f} .

Our goal is to sort out for which parameters of Gelfand–Shilov spaces the operator T is continuous. The Gelfand–Shilov spaces are scales of spaces smaller than the Schwartz space. In fact a Gelfand–Shilov space has two parameters $\theta, s > 0$ and can be defined by the exponential decay conditions

$$\sup_{x \in \mathbf{R}^d} e^{a|x|^{\frac{1}{\theta}}} |f(x)| < \infty, \quad \sup_{\xi \in \mathbf{R}^d} e^{a|\xi|^{\frac{1}{s}}} |\widehat{f}(\xi)| < \infty, \tag{1.2}$$

for some a > 0. The parameter θ thus controls the decay rate of f, and the parameter s controls the decay rate of \hat{f} , that is the smoothness of f.

We use two types of Gelfand–Shilov spaces: The Roumieu spaces $S^s_{\theta}(\mathbf{R}^d)$ for which (1.2) holds for some a > 0, and the Beurling spaces $\Sigma^s_{\theta}(\mathbf{R}^d)$ for which (1.2) holds for all a > 0.

Our first result concerns sufficient conditions for continuity on Gelfand–Shilov spaces.

Theorem 1.1. Define T by (1.1) where q is a polynomial on \mathbb{R}^d with real coefficients and degree $m \ge 2$, and let $s, \theta > 0$.

Key words and phrases. Gelfand–Shilov spaces, oscillatory multiplier operators, linear evolution equations, well-posedness.

- (i) If $s \ge (m-1)\theta \ge 1$ then T is continuous on $\mathcal{S}^s_{\theta}(\mathbf{R}^d)$.
- (ii) If $s \ge (m-1)\theta \ge 1$ and $(\theta, s) \ne \left(\frac{1}{m-1}, 1\right)$ then T is continuous on $\Sigma^s_{\theta}(\mathbf{R}^d)$.

An immediate consequence is the corresponding claim for ultradistributions $(S^s_{\theta})'(\mathbf{R}^d)$ and $(\Sigma^s_{\theta})'(\mathbf{R}^d)$, see Corollary 4.1.

Secondly we prove negative results for d = 1 in a parameter region which is close to complementary to $s \ge (m-1)\theta$. The first result generalizes [1, Proposition 2].

Theorem 1.2. Define T by (1.1) where q is a polynomial on **R** with real coefficients and degree $m \ge 2$, and let $s, \theta > 0$. If

$$1 \leq s < \theta m - \max(\theta, 1),$$

and $\theta \ge 1$ if m = 3, then $T\mathcal{S}^s_{\theta}(\mathbf{R}) \not\subseteq \mathcal{S}^s_{\theta}(\mathbf{R})$.

Theorem 1.3. Define T by (1.1) where q is a polynomial on **R** with real coefficients and degree $m \ge 2$, and let $s, \theta > 0$. If

$$1 < s < \theta m - \max(\theta, 1),$$

and $\theta > 1$ if m = 3, then $T\Sigma^s_{\theta}(\mathbf{R}) \not\subseteq \Sigma^s_{\theta}(\mathbf{R})$.

Note that the statements in Theorems 1.2 and 1.3 are stronger than the lack of continuity on the spaces $S^s_{\theta}(\mathbf{R})$ and $\Sigma^s_{\theta}(\mathbf{R})$ respectively. We illustrate Theorems 1.1, 1.2 and 1.3 in Figure 1.

Our results can be applied to well-posedness of the initial value Cauchy problem for linear evolution equations of the form

$$\begin{cases} \partial_t u(t,x) + ip(D_x)u(t,x) = 0, \quad x \in \mathbf{R}^d, \quad t \in \mathbf{R}, \\ u(0,\cdot) = u_0 \end{cases}$$
(1.3)

where $p : \mathbf{R}^d \to \mathbf{R}$ is a polynomial with real coefficients of degree $m \ge 2$. This generalizes the Schrödinger equation for the free particle where m = 2and $p(\xi) = |\xi|^2$. The solution operator (propagator) to (1.3) is

$$u(t,x) = e^{-itp(D_x)}u_0 = (2\pi)^{-\frac{d}{2}} \int_{\mathbf{R}^d} e^{i\langle x,\xi\rangle - itp(\xi)} \widehat{u}_0(\xi) \mathrm{d}\xi$$

for $u_0 \in \mathscr{S}(\mathbf{R}^d)$. This means that the propagator $e^{-itp(D_x)} = \mathscr{F}^{-1}T\mathscr{F}$ is the conjugation by the Fourier transform of the multiplier operator $(Tf)(x) = e^{-itp(x)}f(x)$ of the form (1.1). Since the Fourier transform maps the Gelfand–Shilov spaces into themselves with an interchange of indices as $\mathscr{F} : \mathscr{S}^s_{\theta}(\mathbf{R}^d) \to \mathscr{S}^s_s(\mathbf{R}^d)$ and $\mathscr{F} : \Sigma^s_{\theta}(\mathbf{R}^d) \to \Sigma^s_s(\mathbf{R}^d)$ we obtain the following consequence of Theorems 1.1, 1.2 and 1.3.

Corollary 1.4. Let $p : \mathbf{R}^d \to \mathbf{R}$ be a polynomial with real coefficients of degree $m \ge 2$, consider the solution operator $e^{-itp(D_x)}$ to the Cauchy problem (1.3), let $s, \theta > 0$ and let $t \in \mathbf{R}$.

- (i) If $\theta \ge (m-1)s \ge 1$ then $e^{-itp(D_x)}$ is continuous on $\mathcal{S}^s_{\theta}(\mathbf{R}^d)$.
- (ii) If $\theta \ge (m-1)s \ge 1$ and $(s,\theta) \ne \left(\frac{1}{m-1},1\right)$ then $e^{-itp(D_x)}$ is continuous on $\Sigma^s_{\theta}(\mathbf{R}^d)$.
- (iii) If d = 1, $1 \leq \theta < sm \max(s, 1)$, $s \geq 1$ if m = 3, and $t \neq 0$ then $e^{-itp(D_x)} \mathcal{S}^s_{\theta}(\mathbf{R}) \notin \mathcal{S}^s_{\theta}(\mathbf{R})$.




FIGURE 1. The (θ, s) -parameter quadrant with behavior of the operator T acting on Gelfand–Shilov spaces indicated when $m \ge 4$ and d = 1.

(iv) If d = 1, $1 < \theta < sm - \max(s, 1)$, s > 1 if m = 3, and $t \neq 0$ then $e^{-itp(D_x)}\Sigma^s_{\theta}(\mathbf{R}) \notin \Sigma^s_{\theta}(\mathbf{R})$.

The proof of Theorem 1.1 uses Debrouwere and Neyt's recent results [4] concerning characterization of smooth functions that acts continuously by multiplication on Gelfand–Shilov spaces, plus a result from [10]. The proofs of Theorems 1.2 and 1.3 are based on ideas from the proof of [1, Proposition 2] which concerns the multiplier function e^{-itx^2} , together with an investigation of the polynomials p_k that appear upon differentiation as $\partial^k e^{iq(x)} = p_k(x) e^{iq(x)}$ for $k \in \mathbf{N}$, where q is a monomial.

The paper is organized as follows. In Section 2 we specify notation, conventions and background material. Then in Section 3 we work out the structure and estimates for the derivatives of exponential monomials in one variable of the form $e^{\lambda x^m/m}$ for $x \in \mathbf{R}$, $m \in \mathbf{N}$, $m \ge 2$ and $\lambda \in \mathbb{C} \setminus \{0\}$. The results are used as tools for the negative results Theorems 1.2 and 1.3. Finally Section 4 is devoted to the proof of Theorem 1.1 and Section 5 to the proofs of Theorems 1.2 and 1.3.

2. Preliminaries

2.1. Notation. The floor function is denoted $\lfloor x \rfloor$ for $x \in \mathbf{R}$. Multiindices $\alpha \in \mathbf{N}^d$ are measured with the 1-norm $|\alpha| = \sum_{j=1}^d \alpha_j$, whereas vectors $x \in \mathbf{R}^d$ are measured with the 2-norm $|x| = \left(\sum_{j=1}^d x_j^2\right)^{\frac{1}{2}}$. We use the bracket $\langle x \rangle = \left(1 + |x|^2\right)^{\frac{1}{2}}$ for $x \in \mathbf{R}^d$. Partial differential operators on \mathbf{R}^d are denoted ∂^{α} for $\alpha \in \mathbf{N}^d$, and $D^{\alpha} = i^{-|\alpha|}\partial^{\alpha}$. The normalization of the

Fourier transform is

$$\mathscr{F}f(\xi) = \widehat{f}(\xi) = (2\pi)^{-\frac{d}{2}} \int_{\mathbf{R}^d} f(x) e^{-i\langle x,\xi\rangle} \,\mathrm{d}x, \qquad \xi \in \mathbf{R}^d,$$

for $f \in \mathscr{S}(\mathbf{R}^d)$ (the Schwartz space), where $\langle \cdot, \cdot \rangle$ denotes the scalar product on \mathbf{R}^d . The conjugate linear action of a (ultra-)distribution u on a test function ϕ is written (u, ϕ) , consistent with the L^2 inner product $(\cdot, \cdot) =$ $(\cdot, \cdot)_{L^2}$ which is conjugate linear in the second argument.

2.2. Gelfand–Shilov spaces. The Schwartz space $\mathscr{S}(\mathbf{R}^d)$ is the subspace of the smooth functions for which the seminorms

$$f \mapsto \sup_{x \in \mathbf{R}^d} |x^{\alpha} D^{\beta} f(x)| := C_{\alpha\beta}$$
(2.1)

are finite for all $\alpha, \beta \in \mathbf{N}^d$.

In this paper we work with Gelfand–Shilov spaces and their dual ultradistribution spaces [5]. For Gelfand–Shilov spaces you impose certain restrictions of the constants $C_{\alpha\beta}$ in (2.1) which leads to spaces that are smaller than $\mathscr{S}(\mathbf{R}^d)$.

Let $\theta, s, h > 0$. The space denoted $\mathcal{S}^s_{\theta,h}(\mathbf{R}^d)$ is the set of all $f \in C^{\infty}(\mathbf{R}^d)$ such that

$$\|f\|_{\mathcal{S}^{s}_{\theta,h}} \equiv \sup \frac{|x^{\alpha}D^{\beta}f(x)|}{h^{|\alpha+\beta|}\alpha!^{\theta}\beta!^{s}}$$
(2.2)

is finite, where the supremum is taken over all $\alpha, \beta \in \mathbf{N}^d$ and $x \in \mathbf{R}^d$. The function space $\mathcal{S}^s_{\theta,h}$ is a Banach space which increases with h, s and θ , and $\mathcal{S}^s_{\theta,h} \subseteq \mathscr{S}$. The topological dual $(\mathcal{S}^s_{\theta,h})'(\mathbf{R}^d)$ is a Banach space which contains the tempered distributions: $\mathscr{S}'(\mathbf{R}^d) \subseteq (\mathcal{S}^s_{\theta,h})'(\mathbf{R}^d)$.

The Beurling type *Gelfand–Shilov space* $\Sigma^s_{\theta}(\mathbf{R}^d)$ is the projective limit of $\mathcal{S}^s_{\theta,h}(\mathbf{R}^d)$ with respect to h [5]. This means

$$\Sigma^{s}_{\theta}(\mathbf{R}^{d}) = \bigcap_{h>0} \mathcal{S}^{s}_{\theta,h}(\mathbf{R}^{d})$$
(2.3)

and the Fréchet space topology of $\Sigma^s_{\theta}(\mathbf{R}^d)$ is defined by the seminorms $\|\cdot\|_{\mathcal{S}^s_{\theta,h}}$ for h > 0.

We have $\Sigma_{\theta}^{s}(\mathbf{R}^{d}) \neq \{0\}$ if and only if $s + \theta > 1$ [8]. The topological dual of $\Sigma_{\theta}^{s}(\mathbf{R}^{d})$ is the space of (Beurling type) *Gelfand–Shilov ultradistributions* [5, Section I.4.3]

$$(\Sigma^s_{\theta})'(\mathbf{R}^d) = \bigcup_{h>0} (\mathcal{S}^s_{\theta,h})'(\mathbf{R}^d).$$
(2.3)'

The Roumieu type Gelfand–Shilov space is the union

$$\mathcal{S}^s_{\theta}(\mathbf{R}^d) = \bigcup_{h>0} \mathcal{S}^s_{\theta,h}(\mathbf{R}^d)$$

equipped with the inductive limit topology [9], that is the strongest topology such that each inclusion $S^s_{\theta,h}(\mathbf{R}^d) \subseteq S^s_{\theta}(\mathbf{R}^d)$ is continuous. We have $S^s_{\theta}(\mathbf{R}^d) \neq \{0\}$ if and only if $s + \theta \geq 1$ [5]. The corresponding (Roumieu type) Gelfand–Shilov ultradistribution space is

$$(\mathcal{S}^s_{\theta})'(\mathbf{R}^d) = \bigcap_{\substack{h>0\\4}} (\mathcal{S}^s_{\theta,h})'(\mathbf{R}^d).$$

For every $s, \theta > 0$ such that $s + \theta > 1$, and for any $\varepsilon > 0$ we have

$$\Sigma^{s}_{\theta}(\mathbf{R}^{d}) \subseteq \mathcal{S}^{s}_{\theta}(\mathbf{R}^{d}) \subseteq \Sigma^{s+\varepsilon}_{\theta+\varepsilon}(\mathbf{R}^{d}).$$
(2.4)

The dual spaces $(\Sigma_{\theta}^{s})'(\mathbf{R}^{d})$ and $(\mathcal{S}_{\theta}^{s})'(\mathbf{R}^{d})$ may be equipped with several topologies. In this paper we use the weak* topologies on $(\Sigma_{\theta}^{s})'(\mathbf{R}^{d})$ and $(\mathcal{S}_{\theta}^{s})'(\mathbf{R}^{d})$ respectively.

The Gelfand–Shilov (ultradistribution) spaces enjoy invariance properties, with respect to translation, dilation, tensorization, coordinate transformation and (partial) Fourier transformation. The Fourier transform extends uniquely to homeomorphisms on $\mathscr{S}'(\mathbf{R}^d)$, from $(\mathcal{S}^s_{\theta})'(\mathbf{R}^d)$ to $(\mathcal{S}^{\theta}_s)'(\mathbf{R}^d)$, and from $(\Sigma^s_{\theta})'(\mathbf{R}^d)$ to $(\Sigma^{\theta}_s)'(\mathbf{R}^d)$, and restricts to homeomorphisms on $\mathscr{S}(\mathbf{R}^d)$, from $\mathcal{S}^s_{\theta}(\mathbf{R}^d)$ to $\mathcal{S}^{\theta}_s(\mathbf{R}^d)$, and from $\Sigma^s_{\theta}(\mathbf{R}^d)$ to $\Sigma^{\theta}_s(\mathbf{R}^d)$, and to a unitary operator on $L^2(\mathbf{R}^d)$.

Chung, Chung and Kim characterized in [3] the Roumieu space $S^s_{\theta}(\mathbf{R}^d)$ as the space of functions $f \in C^{\infty}(\mathbf{R}^d)$ that satisfy (1.2) for some a > 0. It also follows that the Beurling space $\Sigma^s_{\theta}(\mathbf{R}^d)$ can be characterized as the space of functions $f \in C^{\infty}(\mathbf{R}^d)$ that satisfy (1.2) for all a > 0.

We need the following result which says that we may use an alternative family of seminorms instead of the seminorms (2.2) indexed by h > 0 for the spaces $\Sigma^s_{\theta}(\mathbf{R}^d)$ and $\mathcal{S}^s_{\theta}(\mathbf{R}^d)$. This is the family of seminorms

$$\|f\|_a \equiv \sup_{\beta \in \mathbf{N}^d, \ x \in \mathbf{R}^d} e^{a|x|^{\frac{1}{\theta}}} \beta!^{-s} a^{|\beta|} |D^{\beta} f(x)|$$
(2.5)

indexed by a > 0. The result may be considered quite well known but we write down a proof for the benefit of the reader.

Lemma 2.1. Suppose $\theta, s > 0$ and $\theta + s > 1$. For any a > 0 there exists C, h > 0 such that

$$||f||_a \leqslant C ||f||_{\mathcal{S}^s_{\theta,h}}, \quad f \in \Sigma^s_{\theta}(\mathbf{R}^d).$$
(2.6)

For any h > 0 there exists C, a > 0 such that

$$\|f\|_{\mathcal{S}^{s}_{\theta,h}} \leqslant C \|f\|_{a}, \quad f \in \Sigma^{s}_{\theta}(\mathbf{R}^{d}).$$

$$(2.7)$$

Proof. Let $f \in \Sigma^s_{\theta}(\mathbf{R}^d)$. From (2.2) we have for any h > 0

$$|x^{\alpha}D^{\beta}f(x)| \leq ||f||_{\mathcal{S}^{s}_{\theta,h}} \alpha!^{\theta}\beta!^{s}h^{|\alpha+\beta|}, \quad \alpha,\beta \in \mathbf{N}^{d}, \quad x \in \mathbf{R}^{d}.$$

This gives for any $n \in \mathbf{N}$ and any $\beta \in \mathbf{N}^d$

$$|x|^{n}|D^{\beta}f(x)| \leq d^{\frac{n}{2}} \max_{|\alpha|=n} |x^{\alpha}D^{\beta}f(x)| \leq d^{\frac{n}{2}} ||f||_{\mathcal{S}^{s}_{\theta,h}} n!^{\theta}\beta!^{s}h^{n+|\beta|}, \quad x \in \mathbf{R}^{d}.$$

Thus for a > 0 we have

$$\begin{split} \exp\left(\frac{a}{\theta}|x|^{\frac{1}{\theta}}\right)|D^{\beta}f(x)|^{\frac{1}{\theta}} &= \sum_{n=0}^{\infty} \frac{|x|^{\frac{n}{\theta}}|D^{\beta}f(x)|^{\frac{1}{\theta}}\left(d^{\frac{1}{2}}h\right)^{-\frac{n}{\theta}}}{n!} \left(\frac{a\left(d^{\frac{1}{2}}h\right)^{\frac{1}{\theta}}}{\theta}\right)^{n} \\ &\leqslant \|f\|_{\mathcal{S}^{s}_{\theta,h}}^{\frac{1}{\theta}}\beta!^{\frac{s}{\theta}}h^{\frac{|\beta|}{\theta}}\sum_{n=0}^{\infty}2^{-n}, \quad x \in \mathbf{R}^{d}, \quad \beta \in \mathbf{N}^{d}, \end{split}$$

provided $2a\left(d^{\frac{1}{2}}h\right)^{\frac{1}{\theta}} \leq \theta$. Hence

$$e^{a|x|^{\frac{1}{\theta}}}|D^{\beta}f(x)| \leq 2^{\theta} \|f\|_{\mathcal{S}^{s}_{\theta,h}}\beta!^{s}h^{|\beta|}$$

which gives

$$\|f\|_a \leqslant 2^{\theta} \|f\|_{\mathcal{S}^s_{\theta,h}}$$

provided $h \leq \min\left(a^{-1}, d^{-\frac{1}{2}}\left(\theta 2^{-1}a^{-1}\right)^{\theta}\right)$. We have shown (2.6).

In order to show (2.7) we let $f \in \Sigma^s_{\theta}(\mathbf{R}^d)$. From (2.6) we know that $||f||_a < \infty$ for any a > 0. Hence we have for any a > 0, $\beta \in \mathbf{N}^d$ and $x \in \mathbf{R}^d$

$$\sum_{n=0}^{\infty} \frac{|x|^{\frac{n}{\theta}} |D^{\beta} f(x)|^{\frac{1}{\theta}}}{n!} \left(\frac{a}{\theta}\right)^{n} = e^{\frac{a}{\theta}|x|^{\frac{1}{\theta}}} |D^{\beta} f(x)|^{\frac{1}{\theta}} \leqslant \|f\|_{a}^{\frac{1}{\theta}} \beta!^{\frac{s}{\theta}} a^{-\frac{|\beta|}{\theta}},$$

which gives

,

$$|x|^{n}|D^{\beta}f(x)| \leq ||f||_{a}n!^{\theta}\beta!^{s}a^{-|\beta|} \left(\frac{\theta}{a}\right)^{\theta n}, \quad n \in \mathbf{N}, \quad \beta \in \mathbf{N}^{d}, \quad x \in \mathbf{R}^{d},$$

and thus, using [7, Eq. (0.3.3)],

$$|x^{\alpha}D^{\beta}f(x)| \leq ||f||_{a}\alpha!^{\theta}\beta!^{s}a^{-|\beta|}\left(\frac{d\theta}{a}\right)^{\theta|\alpha|}, \quad \alpha, \beta \in \mathbf{N}^{d}, \quad x \in \mathbf{R}^{d}.$$

From this it follows that

$$\|f\|_{\mathcal{S}^s_{\theta,h}} \leq \|f\|_a, \quad f \in \Sigma^s_{\theta}(\mathbf{R}^d),$$

for any h > 0 provided $a \ge \max(h^{-1}, d\theta h^{-\frac{1}{\theta}})$. This proves (2.7).

2.3. Faà di Bruno's formula. Of the several available versions of Faà di Bruno's formula we will use the following two. If $f, g \in C^{\infty}(\mathbf{R})$ then we have for any $k \in \mathbf{N}$

$$\frac{\mathrm{d}^{k}}{\mathrm{d}x^{k}} (f(g(x)))$$

$$= \sum_{m_{1}+2m_{2}+\dots+km_{k}=k} \frac{k!}{m_{1}!m_{2}!\dots m_{k}!} f^{(m_{1}+\dots+m_{k})}(g(x)) \prod_{j=1}^{k} \left(\frac{g^{(j)}(x)}{j!}\right)^{m_{j}}.$$
(2.8)

The second version of Faà di Bruno's formula concerns $f : \mathbf{R} \to \mathbf{R}$ and $g : \mathbf{R}^d \to \mathbf{R}$ with $f \in C^{\infty}(\mathbf{R})$ and $g \in C^{\infty}(\mathbf{R}^d)$, and reads

$$\partial^{\alpha} \big(f(g(x)) \big) = \sum_{j=1}^{|\alpha|} \frac{f^{(j)}(g(x))}{j!} \sum_{\substack{\alpha_1 + \dots + \alpha_j = \alpha \\ |\alpha_\ell| \ge 1, \ 1 \le \ell \le j}} \frac{\alpha!}{\alpha_1! \cdots \alpha_j!} \prod_{\ell=1}^j \partial^{\alpha_\ell} g(x)$$
(2.9)

for any $\alpha \in \mathbf{N}^d \setminus \{0\}$ [6, Eq. (2.3)]. For an even more general version of Faà di Bruno's formula we refer the reader to [2, Proposition 4.3].

3. Derivatives of exponential monomials

Let $m \in \mathbf{N}$, $m \ge 2$, $\lambda \in \mathbf{C} \setminus \{0\}$, and consider the function

$$g_{\lambda}(x) = e^{\lambda x^m/m}, \quad x \in \mathbf{R}.$$
 (3.1)

This function can be considered a generalized Gaussian. Clearly we have for $k \in \mathbf{N}$

$$\partial^k g_\lambda(x) = p_{\lambda,k}(x) \, g_\lambda(x) \tag{3.2}$$

where $p_{\lambda,k}$ is a polynomial of degree deg $p_{\lambda,k} = k(m-1)$. In fact we have $p_{\lambda,0}(x) = 1, p_{\lambda,1}(x) = \lambda x^{m-1}$, and the recursive relation

$$p_{\lambda,k+1}(x) = \lambda x^{m-1} p_{\lambda,k}(x) + p'_{\lambda,k}(x)$$
(3.3)

for $k \in \mathbf{N}$. When $m = 2 = -\lambda$ then $(-1)^k p_{\lambda,k}$ are the Hermite polynomials.

Lemma 3.1. Let $m \in \mathbf{N}$, $m \ge 2$, $\lambda \in \mathbf{C} \setminus \{0\}$, and let g_{λ} be defined by (3.1) and $p_{\lambda,k}$ by (3.2) for $k \in \mathbf{N}$. Then the polynomials $p_{\lambda,k}$ have the form

$$p_{\lambda,k}(x) = \sum_{n=0}^{\lfloor k\frac{m-1}{m} \rfloor} \lambda^{k-n} x^{(m-1)k-nm} C_{k,n}$$
(3.4)

for $k \in \mathbf{N} \setminus \{0\}$, where $C_{k,n} \in \mathbf{N} \setminus \{0\}$ for $0 \leq n \leq \lfloor k \frac{m-1}{m} \rfloor$, and $C_{k,0} = 1$ for all $k \in \mathbf{N} \setminus \{0\}$.

Proof. First we note that for $k \ge 1$ we have

$$\frac{k-1}{2} \leqslant \left\lfloor \frac{k}{2} \right\rfloor \leqslant \left\lfloor k \frac{m-1}{m} \right\rfloor \leqslant k-1.$$
(3.5)

If k = 1 then the sum (3.4) contains only one term of the stated form as confirmed above, with $C_{1,0} = 1$. In an induction argument we suppose that (3.4) holds true for a fixed $k \ge 1$. By (3.5) we have

$$\left\lfloor (k+1)\frac{m-1}{m} \right\rfloor \ge \left\lfloor \frac{k+1}{2} \right\rfloor \ge 1$$

so the sum (3.4) with k replaced by k + 1 does contain the term with index n = 1. We obtain from (3.3) and the induction hypothesis (3.4) with $C_{k,0} = 1$

$$p_{\lambda,k+1}(x) = \lambda x^{m-1} p_{\lambda,k}(x) + p'_{\lambda,k}(x)$$

$$= \lambda^{k+1} x^{(m-1)(k+1)} + \sum_{1 \le n \le \lfloor k \frac{m-1}{m} \rfloor} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n}$$

$$+ \lambda^k x^{(m-1)k-1} (m-1)k$$

$$+ \sum_{1 \le n < k \frac{m-1}{m}} \lambda^{k-n} x^{(m-1)k-nm-1} C_{k,n} \Big((m-1)k - nm \Big)$$

$$= \lambda^{k+1} x^{(m-1)(k+1)} + \sum_{1 \le n \le \lfloor k \frac{m-1}{m} \rfloor} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n}$$

$$+ \lambda^k x^{(m-1)(k+1)-m} (m-1)k$$

$$+ \sum_{2 \le n < k \frac{m-1}{m} + 1} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n-1} \Big((m-1)k - (n-1)m \Big).$$
(3.6)

Note that the term $\lambda^k x^{(m-1)(k+1)-m}(m-1)k$ fits into (3.4) with k replaced by k+1 and index n=1. In the last sum the indices n satisfy $2m \leq mn \leq mn$ (m-1)k + m - 1 which gives $2 \le n \le (k+1)\frac{m-1}{m}$. Hence

$$2 \leqslant n \leqslant \left\lfloor (k+1)\frac{m-1}{m} \right\rfloor.$$

We may conclude that all terms in (3.6) can be absorbed into the formula (3.4) with k replaced by k + 1 for certain coefficients $C_{k+1,n} \in \mathbf{N} \setminus \{0\}$ for $0 \leq n \leq \lfloor (k+1)\frac{m-1}{m} \rfloor$. In fact the coefficients $C_{k+1,n}$ are linear combinations of $\{C_{k,n}\}_{0 \le n \le \lfloor k \frac{m-1}{m} \rfloor}$ with positive integer coefficients. It also follows that $C_{k+1,0} = 1$. This proves the induction step which guarantees that (3.4) holds for any $k \in \mathbf{N} \setminus \{0\}$, and $C_{k,0} = 1$ for all $k \in \mathbf{N} \setminus \{0\}$.

In order to understand more about the polynomials $p_{\lambda,k}$ we would like to gain knowledge about the coefficients $C_{k,n}$. We know from Lemma 3.1 that $C_{k,0} = 1$ for all $k \in \mathbf{N} \setminus \{0\}$.

We need a simple lemma.

Lemma 3.2. Suppose $m \in \mathbf{N}$, $m \ge 2$ and $k \in \mathbf{N} \setminus \{0\}$. Then

$$k \in m\mathbf{N} \implies \left\lfloor (k+1)\frac{m-1}{m} \right\rfloor = \left\lfloor k\frac{m-1}{m} \right\rfloor$$
(3.7)

and

$$k \notin m\mathbf{N} \implies \left\lfloor (k+1)\frac{m-1}{m} \right\rfloor = \left\lfloor k\frac{m-1}{m} \right\rfloor + 1.$$
 (3.8)

Proof. If $k \in m\mathbf{N}$ then there exists $p \in \mathbf{N} \setminus \{0\}$ such that k = mp which gives

$$k\frac{m-1}{m} = p(m-1) = \left\lfloor k\frac{m-1}{m} \right\rfloor$$

and $(k+1)\frac{m-1}{m} = p(m-1) + 1 - \frac{1}{m}$. Since $\frac{1}{2} \le 1 - \frac{1}{m} < 1$ we get
 $\left\lfloor (k+1)\frac{m-1}{m} \right\rfloor = p(m-1) = \left\lfloor k\frac{m-1}{m} \right\rfloor$

which proves (3.7).

If instead $k \notin m\mathbf{N}$ then there exist $p, q \in \mathbf{N}$ with $1 \leq q \leq m-1$ such that k = mp + q. Then $(q + 1)(m - 1) \ge qm$ which yields

$$(k+1)\frac{m-1}{m} = p(m-1) + (q+1)\frac{m-1}{m} \ge p(m-1) + q$$
$$> p(m-1) + q\left(1 - \frac{1}{m}\right) = k\frac{m-1}{m}.$$

The implication (3.8) follows.

The following result gives a recursion formula for the coefficients $C_{k,n}$. First note that

$$C_{2,1} = m - 1. (3.9)$$

Lemma 3.3. If $m \ge 2$ and $k \ge 2$ then

$$C_{k+1,n} = C_{k,n} + C_{k,n-1} \left((m-1)k - m(n-1) \right), \quad 1 \le n \le \left\lfloor k \frac{m-1}{m} \right\rfloor.$$
(3.10)

If $k \ge 2$ and $k \notin m\mathbf{N}$ then we have also

$$C_{k+1,n} = C_{k,n-1} \left((m-1)k - m(n-1) \right), \quad n = \left\lfloor (k+1)\frac{m-1}{m} \right\rfloor. \quad (3.11)$$

Proof. We use (3.3) and (3.4). If $k \in m\mathbf{N}$ then the last term in (3.4) is constant. Thus (3.3) gives

$$p_{\lambda,k+1}(x) = \sum_{n=0}^{\lfloor k\frac{m-1}{m} \rfloor} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n} + \sum_{n=0}^{\lfloor k\frac{m-1}{m} \rfloor -1} \lambda^{k-n} x^{(m-1)k-nm-1} C_{k,n} \Big((m-1)k - nm \Big) = \sum_{n=0}^{\lfloor k\frac{m-1}{m} \rfloor} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n} + \sum_{n=1}^{\lfloor k\frac{m-1}{m} \rfloor} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n-1} \Big((m-1)k - m(n-1) \Big).$$

In view of Lemma 3.2 and (3.7) this proves (3.10) when $k \in m\mathbf{N}$.

If $k \notin m\mathbf{N}$ then $\lfloor k\frac{m-1}{m} \rfloor < k\frac{m-1}{m}$. In fact if we assume $k\frac{m-1}{m} = p \in \mathbf{N}$ then we get the contradiction $k = m(k-p) \in m\mathbf{N}$. Thus the last term in (3.4) contains a positive power of x. Thus again using (3.3) we get

$$p_{\lambda,k+1}(x) = \sum_{n=0}^{\lfloor k\frac{m-1}{m} \rfloor} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n} + \sum_{n=0}^{\lfloor k\frac{m-1}{m} \rfloor} \lambda^{k-n} x^{(m-1)k-nm-1} C_{k,n} \Big((m-1)k - nm \Big) = \sum_{n=0}^{\lfloor k\frac{m-1}{m} \rfloor} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n} + \sum_{n=1}^{\lfloor k\frac{m-1}{m} \rfloor+1} \lambda^{k+1-n} x^{(m-1)(k+1)-nm} C_{k,n-1} \Big((m-1)k - m(n-1) \Big).$$

Again Lemma 3.2 and (3.8) prove (3.10) when $k \notin m\mathbf{N}$, and it also shows (3.11) when $k \notin m\mathbf{N}$.

Remark 3.4. Note that (3.9), written as $C_{2,1} = m - 1 = C_{1,1} + C_{1,0}(m-1)$ using $C_{1,0} = 1$ and forcing $C_{1,1} = 0$, fits into the formula (3.10) for k = n = 1(without the upper bound for n). Indeed $C_{1,1}$ is not well defined due to $\lfloor \frac{m-1}{m} \rfloor = 0$.

In general it is challenging to find explicit expressions for the coefficients $C_{k,n}$ when $n \neq 0$, but n = 1 is an exception.

Lemma 3.5. If $k \ge 2$ then

$$C_{k,1} = \frac{1}{2}(m-1)k(k-1).$$
(3.12)

Proof. We observe that $C_{2,1} = m - 1$ may be written as (3.12) for k = 2. If $k \ge 3$ then (3.5) implies $\lfloor (k-1)\frac{m-1}{m} \rfloor \ge 1$. From $C_{k-1,0} = 1$, Lemma 3.3 and (3.10) we get for $k \ge 3$ and n = 1

$$C_{k,1} = C_{k-1,1} + (m-1)(k-1).$$

If we assume that (3.12) holds with k replaced by k-1 then we get

$$C_{k,1} = C_{k-1,1} + (m-1)(k-1)$$

= $\frac{1}{2}(m-1)(k-1)(k-2) + (m-1)(k-1)$
= $(m-1)(k-1)\left(\frac{1}{2}(k-2) + 1\right)$
= $\frac{1}{2}(m-1)k(k-1).$

By induction this proves (3.12) for all $k \ge 2$.

In the next proposition we need the following lemma.

Lemma 3.6. If $m \ge 2$ and $\theta \ge \frac{2}{m}$ then

$$f(x) = (1+x)^{m\theta} - \left(1 - \frac{1}{m}\right) x^{m\theta - 1} - 1 \ge 0$$
 (3.13)

for all $0 \leq x \leq 1$.

Proof. We have f(0) = 0 and $f(1) = 2^{m\theta} - 2 + \frac{1}{m} > 0$. If 0 < x < 1 then $f'(x) = m\theta(1+x)^{m\theta-1} - (m\theta-1)\left(1 - \frac{1}{m}\right)x^{m\theta-2}$

$$= m\theta(1+x)^{m\theta-2}\left(1+x-\left(1-\frac{1}{m}\right)\left(1-\frac{1}{m\theta}\right)\left(\frac{x}{1+x}\right)^{m\theta-2}\right).$$

It follows that f'(x) > 0 for all 0 < x < 1. The function f is thus strictly increasing in [0, 1] which proves the claim (3.13) for $0 \le x \le 1$.

We may now state and prove our main result on the coefficients $C_{k,n}$, which concerns a recursive bound.

Proposition 3.7. Suppose $\lambda \in \mathbb{C} \setminus \{0\}$, $m \in \mathbb{N}$, $m \ge 2$, $\theta \ge \frac{2}{m}$, and consider the polynomials $p_{\lambda,k}$ defined by (3.2) having the form (3.4) involving the coefficients $\{C_{k,n}\} \subseteq \mathbb{N} \setminus \{0\}$. Then we have the bound

$$C_{k,n+1} \leq C_{k,n}mk^{m\theta}, \quad k \geq 2, \quad 0 \leq n \leq \left\lfloor k\frac{m-1}{m} \right\rfloor - 1.$$
 (3.14)

Proof. If k = 2 then by (3.5) we have $\left\lfloor k \frac{m-1}{m} \right\rfloor = 1$. We have $C_{2,1} = m - 1 \leq C_{2,0}m2^{m\theta} = m2^{m\theta}$ so (3.14) is true for k = 2. In an induction argument we suppose that (3.14) holds for a fixed $k \geq 2$.

First Lemma 3.5 yields

$$\frac{C_{k+1,1}}{m(k+1)^{m\theta}} = \frac{(m-1)k(k+1)}{2m(k+1)^{m\theta}} = \frac{1}{2}\left(1-\frac{1}{m}\right)\frac{k}{(k+1)^{m\theta-1}} \leqslant 1$$

due to the assumption $m\theta \ge 2$. Thus (3.14) holds when k is replaced by k+1 and n=0.

Next let $1 \leq n \leq \left\lfloor (k+1)\frac{m-1}{m} \right\rfloor - 1$. If $k \in m\mathbf{N}$ then by Lemma 3.2 and (3.7) we have $\left\lfloor (k+1)\frac{m-1}{m} \right\rfloor = \left\lfloor k\frac{m-1}{m} \right\rfloor$. Lemma 3.3 and (3.10) gives $C_{k+1,n} \geq C_{k,n}$ and therefore by the induction hypothesis

$$\frac{C_{k+1,n+1}}{C_{k+1,n}} = \frac{C_{k,n+1} + C_{k,n} \left((m-1)k - mn\right)}{C_{k+1,n}} \\
\leqslant \frac{C_{k,n+1}}{C_{k,n}} + (m-1)k \\
\leqslant mk^{m\theta} + (m-1)k \\
= m(k+1)^{m\theta} \left(\frac{k}{k+1}\right)^{m\theta} \left(1 + \left(1 - \frac{1}{m}\right)k^{1-m\theta}\right) \\
\leqslant m(k+1)^{m\theta}$$

in the final inequality using Lemma 3.6, in the form

$$1 + \left(1 - \frac{1}{m}\right)k^{1 - m\theta} \le \left(\frac{k + 1}{k}\right)^{m\theta}$$

for all $k \ge 1$. We have now shown the induction step which shows that (3.14) holds when k is replaced by k + 1 and $0 \le n \le \left\lfloor (k+1)\frac{m-1}{m} \right\rfloor - 1$, provided $k \in m\mathbf{N}$.

We also need to consider $k \notin m\mathbf{N}$ for which Lemma 3.2 and (3.8) yields $\left\lfloor (k+1)\frac{m-1}{m} \right\rfloor = \left\lfloor k\frac{m-1}{m} \right\rfloor + 1$. Thus we assume $1 \leqslant n \leqslant \left\lfloor k\frac{m-1}{m} \right\rfloor$. If $1 \leqslant n \leqslant \left\lfloor k\frac{m-1}{m} \right\rfloor - 1$ then Lemma 3.3 and (3.10) gives $C_{k+1,n} \geqslant C_{k,n}$ and thus by the induction hypothesis

$$\frac{C_{k+1,n+1}}{C_{k+1,n}} = \frac{C_{k,n+1} + C_{k,n} \left((m-1)k - mn \right)}{C_{k+1,n}} \\
\leqslant \frac{C_{k,n+1}}{C_{k,n}} + (m-1)k \\
\leqslant mk^{m\theta} + (m-1)k \\
= m(k+1)^{m\theta} \left(\frac{k}{k+1}\right)^{m\theta} \left(1 + \left(1 - \frac{1}{m}\right)k^{1-m\theta} \\
\leqslant m(k+1)^{m\theta}\right)$$

as before.

Finally if $n = \lfloor k \frac{m-1}{m} \rfloor$ then again by Lemma 3.3 and (3.10) we have $C_{k+1,n} \ge C_{k,n}$. From (3.11) we obtain

$$\frac{C_{k+1,n+1}}{C_{k+1,n}} \leq \frac{C_{k+1,n+1}}{C_{k,n}} = (m-1)k - mn \leq m(k+1)^{m\theta}$$

again due to the assumption $m\theta \ge 2$. We have shown the induction step in all cases.

As a consequence of Proposition 3.7 we have

$$C_{k,2} \leqslant m^2 k^{2m\theta} \tag{3.15}$$

for $k \ge 4$ and $\theta \ge \frac{2}{m}$. In fact (3.5) implies $\left\lfloor k \frac{m-1}{m} \right\rfloor - 1 \ge 1$ if $k \ge 4$ so Proposition 3.7 applies to n = 0 and n = 1.

In the next result we improve the estimate (3.15).

Proposition 3.8. Let $m \in \mathbf{N}$, $m \ge 2$, $\lambda \in \mathbf{C} \setminus \{0\}$ and consider the polynomials $p_{\lambda,k}$ defined by (3.1) and (3.2) having the form (3.4) involving the coefficients $\{C_{k,n}\} \subseteq \mathbf{N} \setminus \{0\}$. If $k \ge 4$ then

$$C_{k,2} \leq \frac{1}{2}m^2k^4.$$
 (3.16)

Proof. From Faà di Bruno's formula (2.9) we obtain if $k \ge 1$

$$p_{\lambda,k}(x) = \frac{\partial^k g_{\lambda}(x)}{g_{\lambda}(x)} = \sum_{j=1}^k \frac{\lambda^j}{m^j j!} \sum_{\substack{k_1 + \dots + k_j = k \\ 1 \le k_\ell \le m}} \frac{k!}{k_1! \cdots k_j!} \prod_{\ell=1}^j \partial^{k_\ell}_x x^m$$

$$= \sum_{j=1}^k \lambda^j x^{mj-k} \frac{k!}{m^j j!} \sum_{\substack{k_1 + \dots + k_j = k \\ 1 \le k_\ell \le m}} \prod_{\substack{\ell=1 \\ k_\ell \le m}}^j \frac{m!}{(m-k_\ell)! k_\ell!}$$

$$= \sum_{n=0}^{k-1} \lambda^{k-n} x^{(m-1)k-nm} \frac{k!}{m^{k-n}(k-n)!} \sum_{\substack{k_1 + \dots + k_{k-n} = k \\ 1 \le k_\ell \le m}} \prod_{\substack{\ell=1 \\ k_\ell \le m}}^{k-n} \frac{m!}{(m-k_\ell)! k_\ell!}.$$
(3.17)

The smallest power of x is (m-1)k - (k-1)m = m - k when n = k - 1. This power seems to be negative if k > m which would be absurd. But negative powers are in fact excluded by (3.17): If k - n = 1 then the sum over $k_1 + \cdots + k_{k-n} = k$ reduces to $k_1 = k$ and $1 \le k \le m$ is postulated. Thus the smallest power of x is non-negative in (3.17).

Comparing (3.17) with (3.4) we observe that the upper limits for the summation index n seems to be possibly different. In fact by (3.5) we know that $\lfloor k \frac{m-1}{m} \rfloor \leq k-1$. Suppose $n = k-1 > \lfloor k \frac{m-1}{m} \rfloor$. Then

$$k-2 \geqslant \left\lfloor k \frac{m-1}{m} \right\rfloor > k \frac{m-1}{m} - 1 = k-1 - \frac{k}{m}$$

which implies k > m. This contradicts the conditions in the sum over $1 \leq k_1 = k \leq m$ above, which is interpreted as zero then. So in fact (3.4) and (3.17) are identical.

The assumption $k \ge 4$ and (3.5) implies that n = 2 has a nonzero coefficient in (3.4). From the identity of (3.4) and (3.17) it follows in particular that

$$C_{k,2} = \frac{k!}{m^{k-2}(k-2)!} \sum_{\substack{k_1 + \dots + k_{k-2} = k \\ 1 \le k_\ell \le m}} \prod_{\ell=1}^{k-2} \frac{m!}{(m-k_\ell)!k_\ell!}.$$

Since $k_{\ell} \ge 1$ for all $1 \le \ell \le k - 2$ and $k_1 + \dots + k_{k-2} = k$, we have

$$k_{\ell} \leq 3, \quad \ell = 1, 2, \dots, k - 2.$$

Hence

$$C_{k,2} = \frac{k!}{m^{k-2}(k-2)!} \sum_{\substack{k_1 + \dots + k_{k-2} = k \\ 1 \le k_\ell \le 3}} \prod_{\ell=1}^{k-2} \frac{m!}{(m-k_\ell)!k_\ell!}.$$
 (3.18)

For k_1, \ldots, k_{k-2} satisfying

$$k_1 + \dots + k_{k-2} = k$$
 and $k_\ell \ge 1$

we must have $k_{\ell_1} = 3$ for some $1 \leq \ell_1 \leq k-2$, and $k_\ell = 1$ for $\ell \neq \ell_1$, or $k_{\ell_1} = k_{\ell_2} = 2$ for some $1 \leq \ell_1 \neq \ell_2 \leq k-2$, and $k_\ell = 1$ for $\ell \notin \{\ell_1, \ell_2\}$. In the first case we have

$$\prod_{\ell=1}^{k-2} \frac{m!}{(m-k_{\ell})!k_{\ell}!} = \frac{m(m-1)(m-2)}{6}m^{k-3} \le m^k \tag{3.19}$$

and in the second case we have

$$\prod_{\ell=1}^{k-2} \frac{m!}{(m-k_{\ell})!k_{\ell}!} = \frac{m^2(m-1)^2}{4} m^{k-4} \le m^k.$$
(3.20)

Combinatorics give

$$\sum_{\substack{k_1+\dots+k_{k-2}=k\\1\leqslant k_{\ell}\leqslant 3}}\leqslant k-2+\binom{k-2}{2}=\frac{1}{2}(k-1)(k-2).$$
(3.21)

Finally we insert (3.19), (3.20) and (3.21) into (3.18) which gives

$$C_{k,2} \leq \frac{k!}{m^{k-2}(k-2)!} m^k \frac{1}{2} (k-1)(k-2) = \frac{1}{2} m^2 k(k-1)^2 (k-2) \leq \frac{1}{2} m^2 k^4.$$

In the next result we look at the particular case of (3.1) where $\lambda = \pm im$, that is $g_{\lambda}(x) = e^{\pm ix^m}$, and the corresponding polynomials $p_{\lambda,k}$ defined by (3.2).

Proposition 3.9. Let $m \in \mathbf{N}$, $m \ge 2$, $\lambda = \pm im$ and consider the polynomials $p_{\lambda,k}$ defined by (3.1) and (3.2) having the form (3.4) involving the coefficients $\{C_{k,n}\} \subseteq \mathbf{N} \setminus \{0\}$. If $\theta \ge \frac{2}{m}$ then there exists a sequence $\{k_j\}_{j=1}^{+\infty} \subseteq \mathbf{N}$ such that

$$\left| p_{\lambda,k_j}(k_j^{\theta}) \right| \ge \frac{1}{2} m^{k_j} k_j^{\theta k_j(m-1)}, \quad j \in \mathbf{N} \setminus \{0\}.$$
(3.22)

Proof. Let $j \in \mathbf{N} \setminus \{0\}$. Due to $1 < \frac{m}{m-1} \leq 2$, the interval $[4j\frac{m}{m-1}, (4j + 1)\frac{m}{m-1}]$ must contain positive integers. Denote by $k_j \in \mathbf{N} \setminus \{0\}$ the smallest of them. Hence we have $4j \leq k_j \frac{m-1}{m} \leq 4j+1$, and therefore

$$4j \leqslant \left\lfloor k_j \frac{m-1}{m} \right\rfloor \leqslant 4j+1 \tag{3.23}$$

as well as

$$\left\lfloor \frac{1}{2} \left\lfloor k_j \frac{m-1}{m} \right\rfloor \right\rfloor = 2j. \tag{3.24}$$

The bound $k_j \ge 4j \frac{m}{m-1} > 4j$ implies that $k_j \ge 4$ for all $j \in \mathbf{N} \setminus \{0\}$.

For notational simplicity we write $k = k_j$. Lemma 3.1 and (3.4) gives

$$p_{\lambda,k}(x) = (\pm 1)^k i^k m^k x^{(m-1)k} \sum_{n=0}^{\lfloor k\frac{m-1}{m} \rfloor} (\pm 1)^n i^{-n} m^{-n} x^{-nm} C_{k,n}$$

which in turn gives

$$\begin{aligned} \left| p_{\lambda,k}(k^{\theta}) \right| &= m^{k} k^{\theta(m-1)k} \left| \sum_{n=0}^{\lfloor k \frac{m-1}{m} \rfloor} (\pm 1)^{n} i^{-n} m^{-n} k^{-\theta nm} C_{k,n} \right| \\ &\geqslant m^{k} k^{\theta(m-1)k} \left| \sum_{\substack{n=0\\n \text{ even}}}^{\lfloor k \frac{m-1}{m} \rfloor} i^{-n} m^{-n} k^{-\theta nm} C_{k,n} \right| \\ &= m^{k} k^{\theta(m-1)k} \left| \sum_{\substack{n=0\\n=0}}^{\lfloor \frac{1}{2} \lfloor k \frac{m-1}{m} \rfloor \rfloor} (-1)^{n} m^{-2n} k^{-2\theta nm} C_{k,2n} \right|. \end{aligned}$$
(3.25)

Using (3.24), and noting that $m^{-4j}k^{-4\theta jm}C_{k,4j} \ge 0$, we have

$$\begin{split} & \sum_{n=0}^{\lfloor \frac{1}{2} \lfloor k \frac{m-1}{m} \rfloor \rfloor} (-1)^n m^{-2n} k^{-2\theta n m} C_{k,2n} \geqslant \sum_{n=0}^{2j-1} (-1)^n m^{-2n} k^{-2\theta n m} C_{k,2n} \\ &= 1 - \frac{C_{k,2}}{m^2 k^{2\theta m}} \\ &\quad + \frac{C_{k,4}}{m^4 k^{4\theta m}} - \frac{C_{k,6}}{m^6 k^{6\theta m}} + \dots + \frac{C_{k,2(2j-2)}}{m^{2(2j-2)} k^{2(2j-2)\theta m}} - \frac{C_{k,2(2j-1)}}{m^{2(2j-1)} k^{2(2j-1)\theta m}}. \end{split}$$
By Proposition 3.7 we have if $0 \leq 2n \leq \lfloor k \frac{m-1}{m} \rfloor - 2,$

 $C_{k,2(n+1)} \leqslant C_{k,2n} m^2 k^{2\theta m}$

which gives

$$\frac{C_{k,2n}}{m^{2n}k^{2n\theta m}} - \frac{C_{k,2(n+1)}}{m^{2(n+1)}k^{2(n+1)\theta m}} \ge 0$$

for all $n \in \mathbb{N} \setminus \{0\}$ such that $0 \leq 2n \leq \lfloor k \frac{m-1}{m} \rfloor - 2$. By (3.23) this includes all $n \in \mathbb{N} \setminus \{0\}$ such that $2 \leq n \leq 2j - 2$. We may conclude that

$$\sum_{n=0}^{\lfloor \frac{1}{2} \lfloor k \frac{m-1}{m} \rfloor \rfloor} (-1)^n m^{-2n} k^{-2\theta nm} C_{k,2n} \ge 1 - \frac{C_{k,2}}{m^2 k^{2\theta m}}.$$
 (3.27)

Finally the assumption $\theta \ge \frac{2}{m}$ and Proposition 3.8 yield

$$\frac{C_{k,2}}{m^2 k^{2\theta m}} \leqslant \frac{m^2 k^4}{2m^2 k^{2\theta m}} = \frac{1}{2} k^{2(2-\theta m)} \leqslant \frac{1}{2}$$

for all k. Combining this with (3.25) and (3.27), we obtain (3.22).

4. Proof Theorem 1.1

In [4, Theorem 5.7] the authors identify the subspace of $g \in C^{\infty}(\mathbf{R}^d)$ such that the multiplier operator $T_g f = fg$ is continuous on $\Sigma^s_{\theta}(\mathbf{R}^d)$ and on $\mathcal{S}^s_{\theta}(\mathbf{R}^d)$, respectively. They prove a characterization of the multiplier space formulated in terms of estimates of the form

$$|\partial^{\alpha}g(x)| \leq Ch^{|\alpha|}|\alpha|!^{s}e^{\lambda^{-\frac{1}{\theta}}|x|^{\frac{1}{\theta}}}, \quad \alpha \in \mathbf{N}^{d}, \quad x \in \mathbf{R}^{d}.$$
(4.1)

In fact $T_g : S^s_{\theta}(\mathbf{R}^d) \to S^s_{\theta}(\mathbf{R}^d)$ is continuous if and only if (4.1) holds for all $\lambda > 0$, some $C = C(\lambda) > 0$, and some $h = h(\lambda) > 0$. Moreover $T_g : \Sigma^s_{\theta}(\mathbf{R}^d) \to \Sigma^s_{\theta}(\mathbf{R}^d)$ is continuous if and only if (4.1) holds for all h > 0, some C = C(h) > 0, and some $\lambda = \lambda(h) > 0$.

Proof of Theorem 1.1.

Theorem 1.1 concerns multiplier functions $g(x) = e^{iq(x)}$ where q is a polynomial of degree $m \ge 2$, that is

$$q(x) = \sum_{|\alpha| \le m} c_{\alpha} x^{\alpha}, \quad c_{\alpha} \in \mathbf{R}, \quad x \in \mathbf{R}^{d},$$
(4.2)

and $c_{\alpha} \neq 0$ for some $\alpha \in \mathbf{N}^d$ with $|\alpha| = m$.

If $\gamma \in \mathbf{N}^d$ and $|\gamma| \leq m$ then

$$\partial^{\gamma} q(x) = \sum_{\substack{|\alpha|=m\\\alpha \geqslant \gamma}} c_{\alpha} \frac{\alpha!}{(\alpha - \gamma)!} x^{\alpha - \gamma} + \text{L.O.T.}$$
(4.3)

Hence

$$|\partial^{\gamma} q(x)| \leq C_{q,m} \max(1, |x|^{m-|\gamma|}), \tag{4.4}$$

where $C_{q,m} > 0$ is a constant depending only on m and on the coefficients of q.

From Faà di Bruno's formula (2.9) one gets for $\alpha \in \mathbf{N}^d \setminus \{0\}$

$$\partial^{\alpha}g(x) = \sum_{j=1}^{|\alpha|} i^{j} \frac{g(x)}{j!} \sum_{\substack{\alpha_{1}+\dots+\alpha_{j}=\alpha\\1\leqslant|\alpha_{\ell}|\leqslant m, \ 1\leqslant\ell\leqslant j}} \frac{\alpha!}{\alpha_{1}!\cdots\alpha_{j}!} \prod_{\ell=1}^{j} \partial^{\alpha_{\ell}}q(x).$$
(4.5)

Now (4.4) entails if $|x| \ge 1$

$$\begin{aligned} |\partial^{\alpha}g(x)| &\leq \sum_{j=1}^{|\alpha|} \frac{1}{j!} \sum_{\substack{\alpha_{1}+\dots+\alpha_{j}=\alpha\\1\leq|\alpha_{\ell}|\leq m,\ 1\leq\ell\leq j}} \frac{\alpha!}{\alpha_{1}!\cdots\alpha_{j}!} \prod_{\ell=1}^{j} C_{q,m} |x|^{m-|\alpha_{\ell}|} \\ &\leq \alpha! \sum_{j=1}^{|\alpha|} \frac{C_{q,m}^{j}}{j!} |x|^{jm-|\alpha|} \sum_{\substack{\alpha_{1}+\dots+\alpha_{j}=\alpha\\1\leq|\alpha_{\ell}|\leq m,\ 1\leq\ell\leq j}} \frac{1}{\alpha_{1}!\cdots\alpha_{j}!}. \end{aligned}$$

$$\tag{4.6}$$

On the one hand we get from [7, Eq. (0.3.15)]

$$\sum_{\substack{\alpha_1 + \dots + \alpha_j = \alpha \\ 1 \le |\alpha_\ell| \le m, \ 1 \le \ell \le j}} \frac{1}{\alpha_1! \cdots \alpha_j!} \le \binom{m+d}{m}^j \le 2^{(m+d)j}.$$
(4.7)

On the other hand if $|x| \ge 1$ we have for any $\lambda > 0$, using $j \le |\alpha| \le jm$,

$$|x|^{jm-|\alpha|} = \left(\frac{\left(\theta^{-1}\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}\right)^{jm-|\alpha|}}{(jm-|\alpha|)!}\right)^{\theta} (jm-|\alpha|)!^{\theta} \left(\theta^{\theta}\lambda\right)^{jm-|\alpha|}$$

$$\leq e^{\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}} (j(m-1))!^{\theta} \left(\theta^{\theta}\lambda\right)^{jm-|\alpha|}$$

$$(4.8)$$

and, using [7, Eq. (0.3.12)],

$$(j(m-1))!^{\theta} \leq (j(m-1))^{j(m-1)\theta} = (m-1)^{(m-1)\theta j} j^{j(m-1)\theta}$$

$$\leq (e(m-1))^{(m-1)\theta j} j!^{(m-1)\theta}.$$
(4.9)

If $|x| \ge 1$ then insertion of (4.7), (4.8) and (4.9) into (4.6) gives, using the assumption $(m-1)\theta \ge 1$,

$$\begin{aligned} |\partial^{\alpha}g(x)| &\leq \alpha ! e^{\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}} \sum_{j=1}^{|\alpha|} \left(C_{q,m} \left(e(m-1) \right)^{(m-1)\theta} 2^{m+d} \right)^{j} \left(\theta^{\theta} \lambda \right)^{jm-|\alpha|} j!^{(m-1)\theta-1} \\ &\leq C^{|\alpha|} |\alpha|!^{(m-1)\theta} e^{\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}} \sum_{j=1}^{|\alpha|} \lambda^{jm-|\alpha|} \end{aligned}$$

$$(4.10)$$

for some C > 0.

If $|x| \le 1$ then (4.4), (4.5), (4.7), (4.8) with |x| = 1, and (4.9) give for any $\lambda > 0$

$$\begin{aligned} |\partial^{\alpha}g(x)| &\leq \alpha! \sum_{j=1}^{|\alpha|} \left(C_{q,m} 2^{m+d} \right)^{j} j!^{-1} \\ &\leq \alpha! \sum_{j=1}^{|\alpha|} \left(C_{q,m} 2^{m+d} \right)^{j} j!^{-1} e^{\lambda^{-\frac{1}{\theta}}} (j(m-1))!^{\theta} \left(\theta^{\theta} \lambda \right)^{jm-|\alpha|} \\ &\leq \alpha! e^{\lambda^{-\frac{1}{\theta}}} \sum_{j=1}^{|\alpha|} \left(C_{q,m} \left(e(m-1) \right)^{(m-1)\theta} 2^{m+d} \right)^{j} \left(\theta^{\theta} \lambda \right)^{jm-|\alpha|} j!^{(m-1)\theta-1} \\ &\leq C^{|\alpha|} |\alpha|!^{(m-1)\theta} e^{\lambda^{-\frac{1}{\theta}}} \sum_{j=1}^{|\alpha|} \lambda^{jm-|\alpha|} \end{aligned}$$

$$(4.11)$$

for some C > 0.

The estimate (4.10) for $|x| \ge 1$ and the estimate (4.11) for $|x| \le 1$ can be combined into the estimate for $x \in \mathbf{R}^d$

$$\left|\partial^{\alpha}g(x)\right| \leqslant e^{\lambda^{-\frac{1}{\theta}}}C^{|\alpha|}|\alpha|!^{(m-1)\theta}e^{\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}}\sum_{j=1}^{|\alpha|}\lambda^{jm-|\alpha|}$$
(4.12)

for some C > 0.

Proof of claim (i).

The assumption $s \ge (m-1)\theta$ and (4.12) imply if $\lambda \ge 1$

$$\begin{aligned} |\partial^{\alpha}g(x)| &\leq e^{\lambda^{-\frac{1}{\theta}}} C^{|\alpha|} |\alpha|!^{s} e^{\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}} \sum_{j=1}^{|\alpha|} \lambda^{|\alpha|(m-1)} \\ &\leq e^{\lambda^{-\frac{1}{\theta}}} \left(2C\lambda^{m-1}\right)^{|\alpha|} |\alpha|!^{s} e^{\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}} \end{aligned}$$
(4.13)

and if $0 < \lambda < 1$

$$\begin{aligned} |\partial^{\alpha}g(x)| &\leq e^{\lambda^{-\frac{1}{\theta}}} C^{|\alpha|} |\alpha|!^{s} e^{\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}} \sum_{j=1}^{|\alpha|} \lambda^{-|\alpha|} \\ &\leq e^{\lambda^{-\frac{1}{\theta}}} \left(2C\lambda^{-1}\right)^{|\alpha|} |\alpha|!^{s} e^{\left(\lambda^{-1}|x|\right)^{\frac{1}{\theta}}} \end{aligned}$$
(4.14)

Combining (4.13) and (4.14) we may by the criterion in [4, Theorem 5.7]conclude that T is continuous on $\mathcal{S}^s_{\theta}(\mathbf{R}^d)$. Claim (i) has been proved.

Proof of claim (ii).

The assumptions imply that either $s \ge \theta(m-1) > 1$ or $s > \theta(m-1) \ge 1$. First we suppose that $s \ge \theta(m-1) > 1$. Then [10, Theorem 7.1] shows that T is continuous on $\Sigma^s_{\theta}(\mathbf{R}^d)$.

It remains to consider $s > \theta(m-1) \ge 1$. Then $\varepsilon := s - \theta(m-1) > 0$. From (4.12) with $\lambda = 1$ we obtain for any h > 0

$$\begin{aligned} |\partial^{\alpha}g(x)| &\leq e \, C^{|\alpha|} |\alpha|!^{s-\varepsilon} e^{|x|^{\frac{1}{\theta}}} |\alpha| \\ &\leq e \, (2Ch)^{|\alpha|} \, |\alpha|!^s e^{|x|^{\frac{1}{\theta}}} \left(\frac{h^{-\frac{|\alpha|}{\varepsilon}}}{|\alpha|!}\right)^{\varepsilon} \\ &\leq e^{1+\varepsilon h^{-\frac{1}{\varepsilon}}} \, (2Ch)^{|\alpha|} \, |\alpha|!^s e^{|x|^{\frac{1}{\theta}}}. \end{aligned}$$

Again by the criterion in [4, Theorem 5.7] we may conclude that T is continuous on $\Sigma^s_{\theta}(\mathbf{R}^d)$. Claim (ii) has been proved.

Corollary 4.1. Define T by (1.1) where q is a polynomial on \mathbf{R}^d with real coefficients and degree $m \ge 2$, and let $s, \theta > 0$.

- (i) If $s \ge (m-1)\theta \ge 1$ then T is continuous on $(\mathcal{S}^s_{\theta})'(\mathbf{R}^d)$.
- (ii) If $s \ge (m-1)\theta \ge 1$ and $(\theta, s) \ne \left(\frac{1}{m-1}, 1\right)$ then T is continuous on $(\Sigma^s_{\rho})'(\mathbf{R}^d).$

5. Proofs of Theorems 1.2 and 1.3

Lemma 5.1. Let $f, g \in C^{\infty}(\mathbf{R})$. If there exists $A, B, a, \theta, \nu > 0$, and $s \ge 1$ such that

$$|D^k f(x)| \leq B^k k! |f(x)| \langle x \rangle^{k \max\left(\frac{1}{\nu} - 1, 0\right)}, \quad k \in \mathbf{N},$$
(5.1)

and

$$|D^{k}(g(x)f(x))| \leq AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}}, \quad k \in \mathbf{N},$$
(5.2)

then

$$|(D^{k}g)(x)f(x)| \leq A(2B)^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}}\langle x \rangle^{k\max\left(\frac{1}{\nu}-1,0\right)}, \quad k \in \mathbf{N}.$$
(5.3)

Proof. The assumption (5.2) for k = 0 implies (5.3) for k = 0. In an induction proof we let $k \in \mathbb{N} \setminus \{0\}$ and assume that (5.3) holds for orders smaller than k. We use

$$(D^{k}g)(x)f(x) = D^{k}(g(x)f(x)) - \sum_{n=0}^{k-1} \binom{k}{n} D^{n}g(x)D^{k-n}f(x)$$

(5.1), (5.2), the induction hypothesis, and $s \ge 1$. If $\nu \ge 1$ then max $\left(\frac{1}{\nu} - 1, 0\right) =$ 0 which gives

$$\begin{split} |(D^{k}g)(x)f(x)| &\leq |D^{k}(g(x)f(x))| + \sum_{n=0}^{k-1} \binom{k}{n} |D^{n}g(x)| |D^{k-n}f(x)| \\ &\leq AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} + \sum_{n=0}^{k-1} \binom{k}{n} |D^{n}g(x)| |f(x)| B^{k-n}(k-n)! \\ &\leq AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} + \sum_{n=0}^{k-1} \binom{k}{n} A(2B)^{n}n!^{s}e^{-a|x|^{\frac{1}{\theta}}} B^{k-n}(k-n)! \\ &= AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} \left(1 + \sum_{n=0}^{k-1} \binom{n!}{k!}\right)^{s-1} 2^{n} \right) \\ &\leq AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} \left(1 + \sum_{n=0}^{k-1} 2^{n} \right) \\ &= A(2B)^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}}. \end{split}$$

This proves the induction step, and hence (5.3), provided $\nu \ge 1$. It remains to consider $0 < \nu < 1$. Then $\max\left(\frac{1}{\nu} - 1, 0\right) = \frac{1}{\nu} - 1$ and we have

$$\begin{split} |(D^{k}g)(x)f(x)| \\ &\leqslant |D^{k}(g(x)f(x))| + \sum_{n=0}^{k-1} \binom{k}{n} |D^{n}g(x)| |D^{k-n}f(x)| \\ &\leqslant AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} + \sum_{n=0}^{k-1} \binom{k}{n} |D^{n}g(x)| |f(x)| \langle x \rangle^{(k-n)(\frac{1}{\nu}-1)} B^{k-n}(k-n)! \\ &\leqslant AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} + \sum_{n=0}^{k-1} \binom{k}{n} A(2B)^{n}n!^{s}e^{-a|x|^{\frac{1}{\theta}}} \langle x \rangle^{k(\frac{1}{\nu}-1)} B^{k-n}(k-n)! \\ &= AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} \left(1 + \langle x \rangle^{k(\frac{1}{\nu}-1)} \sum_{n=0}^{k-1} \binom{n!}{k!} \sum_{n=0}^{s-1} 2^{n}\right) \\ &\leqslant AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} \left(1 + \langle x \rangle^{k(\frac{1}{\nu}-1)} \sum_{n=0}^{k-1} 2^{n}\right) \\ &= AB^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} \left(1 + \langle x \rangle^{k(\frac{1}{\nu}-1)} (2^{k}-1)\right) \\ &\leqslant A(2B)^{k}k!^{s}e^{-a|x|^{\frac{1}{\theta}}} \langle x \rangle^{k(\frac{1}{\nu}-1)} \end{split}$$

which proves the induction step, and hence (5.3), when $0 < \nu < 1$.

Consider the function $\langle x \rangle^t = (1+x^2)^{\frac{t}{2}}$ for $x \in \mathbf{R}$ with $t \in \mathbf{R}$. We have the following estimate for its derivatives.

Lemma 5.2. If $t \in \mathbf{R}$ then there exists $C_t \ge 1$ such that for all $k \in \mathbf{N}$

$$\left| D^{k} \langle x \rangle^{t} \right| \leqslant C_{t} 2^{3k} k! \langle x \rangle^{t-k}, \quad x \in \mathbf{R}.$$
(5.4)

Proof. First we assume $|t| \leq 1$ and show

$$\left| D^{k} \langle x \rangle^{t} \right| \leq 2^{2k+5} k! \langle x \rangle^{t-k}, \quad x \in \mathbf{R},$$
(5.5)

for all $k \in \mathbf{N}$.

The function $\langle x \rangle^t = (1+x^2)^{\frac{t}{2}}$ is a composition of $f(x) = x^{\frac{t}{2}}$ with g(x) = $1 + x^2$. The estimate (5.5) is trivial when k = 0, and can be verified for k = 1 and k = 2 so we may assume that $k \ge 3$.

For $n \in \mathbf{N}$ we have

$$f^{(n)}(x) = \frac{t}{2} \left(\frac{t}{2} - 1\right) \cdots \left(\frac{t}{2} - (n-1)\right) x^{\frac{t}{2} - n}$$
(5.6)

and if $m_j = 0$ for $3 \leq j \leq k$ then

$$\prod_{j=1}^{k} \left(\frac{g^{(j)}(x)}{j!}\right)^{m_j} = \left(g'(x)\right)^{m_1} \left(\frac{g''(x)}{2}\right)^{m_2} = 2^{m_1} x^{m_1}.$$
 (5.7)

If instead $m_j > 0$ for some $3 \leq j \leq k$ then the product is zero.

Inserting (5.6) and (5.7) into Faà di Bruno's formula (2.8) yields by means of [7, Eq. (0.3.5)]

$$\begin{aligned} \frac{|D^{k}\langle x\rangle^{t}|}{k!\langle x\rangle^{t-k}} \\ &\leqslant \sum_{m_{1}+2m_{2}=k} \frac{1}{m_{1}!m_{2}!} \left| \frac{t}{2} \left(\frac{t}{2} - 1 \right) \cdots \left(\frac{t}{2} - (m_{1} + m_{2} - 1) \right) \right| \langle x \rangle^{t-2(m_{1}+m_{2})+k-t} 2^{m_{1}} x^{m_{1}} \\ &\leqslant \frac{|t|}{2} \sum_{m_{1}+2m_{2}=k} \frac{1}{m_{1}!m_{2}!} \left(\frac{|t|}{2} + 1 \right) \cdots \left(\frac{|t|}{2} + m_{1} + m_{2} + 1 \right) \langle x \rangle^{-(m_{1}+2m_{2})+k} 2^{m_{1}} \\ &\leqslant \sum_{m_{1}+2m_{2}=k} \frac{2^{m_{1}-1}(m_{1} + m_{2} + 2)!}{m_{1}!m_{2}!} \leqslant \sum_{m_{1}+2m_{2}=k} 2^{2m_{1}+m_{2}+1}(m_{2} + 2)(m_{2} + 1) \\ &\leqslant \sum_{m_{1}+2m_{2}=k} 2^{2m_{1}+3m_{2}+4} = \sum_{m_{2}=0}^{\lfloor \frac{k}{2} \rfloor} 2^{2(k-2m_{2})+3m_{2}+4} = 2^{2k+4} \sum_{m_{2}=0}^{\lfloor \frac{k}{2} \rfloor} 2^{-m_{2}} \leqslant 2^{2k+5} \end{aligned}$$

which proves (5.5) provided $|t| \leq 1$.

Next we consider |t| > 1 and put $m = \lfloor |t| \rfloor$. Writing

$$\langle x \rangle^t = \langle x \rangle^{\pm |t|} = \langle x \rangle^{\pm (|t|-m)} \prod_{j=1}^m \langle x \rangle^{\pm 1}$$

we use Leibniz' rule

$$D^{k}\langle x \rangle^{t} = \sum_{k_{0}+k_{1}+\dots+k_{m}=k} \frac{k!}{k_{0}!k_{1}!\cdots k_{m}!} D^{k_{0}}\langle x \rangle^{\pm (|t|-m)} \prod_{j=1}^{m} D^{k_{j}}\langle x \rangle^{\pm 1}.$$
19

Since $0 \leq |t| - m < 1$ we may use (5.5) which yields, using [7, Eq. (0.3.16)],

$$\begin{split} \left| D^k \langle x \rangle^t \right| &\leq \sum_{k_0 + k_1 + \dots + k_m = k} \frac{k!}{k_0! k_1! \cdots k_m!} \left| D^{k_0} \langle x \rangle^{\pm (|t| - m)} \right| \prod_{j=1}^m \left| D^{k_j} \langle x \rangle^{\pm 1} \right| \\ &\leq \sum_{k_0 + k_1 + \dots + k_m = k} \frac{k!}{k_0! k_1! \cdots k_m!} 2^{2k_0 + 5} k_0! \langle x \rangle^{\pm (|t| - m) - k_0} \prod_{j=1}^m 2^{2k_j + 5} k_j! \langle x \rangle^{\pm 1 - k_j} \\ &\leq 2^{2k + 5(m+1)} k! \langle x \rangle^{t-k} \sum_{k_0 + k_1 + \dots + k_m = k} \\ &= 2^{2k + 5(m+1)} k! \langle x \rangle^{t-k} \binom{k+m}{m} \\ &\leq 2^{3k + 6m + 5} k! \langle x \rangle^{t-k} \leqslant 2^{3k + 6|t| + 5} k! \langle x \rangle^{t-k}. \end{split}$$

Combined with (5.5) for $|t| \leq 1$ we have now shown (5.4) for all $t \in \mathbf{R}$. \Box

Let $\theta > 0$ and consider the function

$$f(x) = e^{-\langle x \rangle^{\frac{1}{\theta}}}, \quad x \in \mathbf{R}.$$
 (5.8)

Lemma 5.3. If f is defined by (5.8) with $\theta > 0$ then there exists $C_{\theta} \ge 1$ such that for all $k \in \mathbf{N} \setminus \{0\}$

$$|f^{(k)}(x)| \leq C_{\theta}^{k} k! f(x) \sum_{j=1}^{k} \frac{1}{j!} \langle x \rangle^{j\left(\frac{1}{\theta}-1\right)}.$$

Proof. Faà di Bruno's formula (2.9), Lemma 5.2 and [7, Eq. (0.3.16)] give for some $C_{\theta} \ge 1$ if $k \ge 1$

$$\begin{split} \frac{|f^{(k)}(x)|}{k!f(x)} &= \left| \sum_{j=1}^{k} \frac{(-1)^{j}}{j!} \sum_{\substack{k_{1}+\dots+k_{j}=k\\k_{\ell}\geqslant 1,\ 1\leqslant \ell\leqslant j}} \frac{1}{k_{1}!\cdots k_{j}!} \prod_{\ell=1}^{j} D^{k_{\ell}}\langle x \rangle^{\frac{1}{\theta}} \right| \\ &\leqslant \sum_{j=1}^{k} \frac{1}{j!} \sum_{\substack{k_{1}+\dots+k_{j}=k\\k_{\ell}\geqslant 1,\ 1\leqslant \ell\leqslant j}} \prod_{\ell=1}^{j} C_{\theta} 2^{3k_{\ell}} \langle x \rangle^{\frac{1}{\theta}-k_{\ell}} \\ &\leqslant C_{\theta}^{k} 2^{3k} \sum_{j=1}^{k} \frac{1}{j!} \langle x \rangle^{\frac{j}{\theta}-k} \sum_{\substack{k_{1}+\dots+k_{j}=k\\k_{\ell}\geqslant 1,\ 1\leqslant \ell\leqslant j}} \\ &\leqslant C_{\theta}^{k} 2^{3k} \sum_{j=1}^{k} \frac{1}{j!} \langle x \rangle^{\frac{j}{\theta}-k} \binom{k+j-1}{j-1} \\ &\leqslant C_{\theta}^{k} 2^{3k} \sum_{j=1}^{k} \frac{1}{j!} \langle x \rangle^{\frac{j}{\theta}-k} 2^{k+j-1} \\ &\leqslant C_{\theta}^{k} 2^{5k} \sum_{j=1}^{k} \frac{1}{j!} \langle x \rangle^{j(\frac{1}{\theta}-1)}. \end{split}$$

Corollary 5.4. If f is defined by (5.8) with $\theta > 0$ then there exists $C_{\theta} \ge 1$ such that for all $k \in \mathbf{N}$

$$|f^{(k)}(x)| \leq C_{\theta}^{k} k! f(x) \langle x \rangle^{k \max\left(\frac{1}{\theta} - 1, 0\right)}.$$

Proof. The estimate is trivial when k = 0. Lemma 5.3 gives for some $C_{\theta} \ge 1$ and $k \ge 1$ with $\tau = \max\left(\frac{1}{\theta} - 1, 0\right)$

$$|f^{(k)}(x)| \leq C_{\theta}^{k} k! f(x) \sum_{j=1}^{k} \frac{1}{j!} \langle x \rangle^{j\tau}$$
$$\leq (2C_{\theta})^{k} k! f(x) \langle x \rangle^{k\tau}.$$

Proposition 5.5. If $\theta > 0$ and the function f is defined by (5.8) then $f \in S^1_{\theta}(\mathbf{R})$.

Proof. If $\theta \ge 1$ then Corollary 5.4 gives

$$|f^{(k)}(x)| \leq C_{\theta}^{k} k! e^{-\langle x \rangle^{\frac{1}{\theta}}}$$

for some $C_{\theta} \ge 1$, and thus it follows from Lemma 2.1 that $f \in \mathcal{S}_{\theta}^{1}(\mathbf{R})$. Let $0 < \theta < 1$. We have for $x \in \mathbf{R}$ and $j \in \mathbf{N}$

$$\langle x \rangle^j = \left(\frac{\langle x \rangle^{\frac{j}{\theta}}}{j!}
ight)^{\theta} j!^{\theta} \leqslant e^{\theta \langle x \rangle^{\frac{1}{\theta}}} j!^{\theta}$$

which combined with Lemma 5.3 gives for some $C_{\theta} \ge 1$ and $k \ge 1$

$$|f^{(k)}(x)| \leq C_{\theta}^{k} k! f(x) \sum_{j=1}^{k} \frac{1}{j!} \langle x \rangle^{j\frac{1-\theta}{\theta}}$$
$$\leq C_{\theta}^{k} k! f(x) e^{(1-\theta)\langle x \rangle^{\frac{1}{\theta}}} \sum_{j=1}^{k} j!^{-1+1-\theta}$$
$$\leq (2C_{\theta})^{k} k! e^{-\theta\langle x \rangle^{\frac{1}{\theta}}}.$$

Lemma 2.1 again shows that $f \in \mathcal{S}^1_{\theta}(\mathbf{R})$.

We assume that q is a polynomial on **R** with real coefficients of degree $m \ge 2$, that is

$$q(x) = \sum_{k=0}^{m} c_k x^k, \quad c_k \in \mathbf{R}, \quad c_m \neq 0, \quad x \in \mathbf{R}.$$

In order to show Theorems 1.2 and 1.3, we may by a rescaling argument assume that $c_m = \pm 1$, that is

$$q(x) = \pm x^m + q_{m-1}(x) \tag{5.9}$$

where q_{m-1} is a polynomial with real coefficients of degree deg $q_{m-1} \leq m-1$.

Lemma 5.6. Let $m \in \mathbf{N}$, $m \ge 2$, and let $\theta > \frac{2}{m}$. Let the polynomial q be defined by (5.9) and let $f \in C^{\infty}(\mathbf{R})$.

(i) Suppose $\theta \ge 1$ if m = 3. If

$$e^{\pm ix^m} f \notin \bigcup_{1 \le s < \theta m - \max(\theta, 1)} \mathcal{S}^s_{\theta}(\mathbf{R})$$
(5.10)

then

$$e^{iq(x)}f \notin \bigcup_{1 \leq s < \theta m - \max(\theta, 1)} S^s_{\theta}(\mathbf{R}).$$
 (5.11)

(ii) Suppose $\theta > 1$ if m = 3. If $\frac{2}{m} < \nu < \theta$ and

$$e^{\pm ix^m} f \notin \bigcup_{1 < s < \nu m - \max(\nu, 1)} \Sigma^s_{\theta}(\mathbf{R})$$
(5.12)

then

$$e^{iq(x)}f \notin \bigcup_{1 < s < \nu m - \max(\nu, 1)} \Sigma^{s}_{\theta}(\mathbf{R}).$$
(5.13)

Proof. We start by proving statement (i). First we discuss the case m = 2in which $q_{m-1} = q_1$ is a polynomial of degree one. By assumption $\theta > 1$. Suppose $e^{\pm ix^2} f \notin S^s_{\theta}(\mathbf{R})$ for some $1 \leq s < \theta$. Then $e^{iq(x)} f = e^{iq_1(x)} e^{\pm ix^2} f \notin S^s_{\theta}(\mathbf{R})$ follows from the invariance of $S^s_{\theta}(\mathbf{R})$ with respect to modulation. This shows a strengthened form of statement (i).

It remains to consider the case $m \ge 3$, in which the assumptions imply that $(m-2)\theta \ge 1$. Assumption (5.10) means that $e^{\pm ix^m}f \notin S^s_{\theta}(\mathbf{R})$ for all

$$1 \leq s < \theta m - \max(\theta, 1).$$

In a contradictory argument we suppose that (5.11) is not true, that is $e^{iq} f \in S^s_{\theta}(\mathbf{R})$ for some $1 \leq s < \theta m - \max(\theta, 1)$.

If $(m-2)\theta \leq s < \theta m - \max(\theta, 1)$ then Theorem 1.1 (i) implies that $e^{-iq_{m-1}}$ is continuous on $\mathcal{S}^s_{\theta}(\mathbf{R})$, which gives $e^{\pm ix^m}f = e^{-iq_{m-1}}e^{iq}f \in \mathcal{S}^s_{\theta}(\mathbf{R})$. This contradicts the assumption (5.10).

If instead $1 \leq s < (m-2)\theta$ then $e^{iq}f \in \mathcal{S}^s_{\theta}(\mathbf{R}) \subseteq \mathcal{S}^{(m-2)\theta}_{\theta}(\mathbf{R})$. Theorem 1.1 (i) again implies that $e^{-iq_{m-1}}$ is continuous on $\mathcal{S}^{(m-2)\theta}_{\theta}(\mathbf{R})$, which gives $e^{\pm ix^m}f = e^{-iq_{m-1}}e^{iq}f \in \mathcal{S}^{(m-2)\theta}_{\theta}(\mathbf{R})$. This again contradicts the assumption (5.10). We have shown statement (i) for all $m \geq 2$.

It remains to prove of statement (ii). If m = 2 then $q_{m-1} = q_1$ is a polynomial of degree one and $\theta > 1$. Suppose $e^{\pm ix^2}f \notin \Sigma_{\theta}^s(\mathbf{R})$ for some $1 < s < \nu$. Then $e^{iq(x)}f = e^{iq_1(x)}e^{\pm ix^2}f \notin \Sigma_{\theta}^s(\mathbf{R})$ follows from the invariance of $\Sigma_{\theta}^s(\mathbf{R})$ with respect to modulation. This shows a strengthened form of statement (ii).

Let $m \ge 3$. The assumptions imply $(m-2)\theta > 1$. Assumption (5.12) means that $e^{\pm ix^m} f \notin \Sigma^s_{\theta}(\mathbf{R})$ for all

$$1 < s < \nu m - \max(\nu, 1).$$

Suppose that (5.13) is not true, that is $e^{iq}f \in \Sigma^s_{\theta}(\mathbf{R})$ for some $1 < s < \nu m - \max(\nu, 1)$.

If $(m-2)\theta \leq s < \nu m - \max(\nu, 1)$ then Theorem 1.1 (ii) implies that $e^{-iq_{m-1}}$ is continuous on $\Sigma^s_{\theta}(\mathbf{R})$, which gives $e^{\pm ix^m}f = e^{-iq_{m-1}}e^{iq}f \in \Sigma^s_{\theta}(\mathbf{R})$. This contradicts the assumption (5.12).

If instead $1 < s < (m-2)\theta$ then $e^{iq}f \in \Sigma^s_{\theta}(\mathbf{R}) \subseteq \Sigma^{(m-2)\theta}_{\theta}(\mathbf{R})$. Theorem 1.1 (ii) again implies that $e^{-iq_{m-1}}$ is continuous on $\Sigma^{(m-2)\theta}_{\theta}(\mathbf{R})$, which gives $e^{\pm ix^m}f = e^{-iq_{m-1}}e^{iq}f \in \Sigma_{\theta}^{(m-2)\theta}(\mathbf{R})$. This again contradicts the assumption (5.12). We have shown statement (ii) for all $m \ge 2$.

Proof of Theorem 1.2.

The assumptions for Lemma 5.6 (i) are satisfied. If m = 2 then the assumptions for Theorem 1.2 are $1 \leq s < \theta$. The proof of [1, Proposition 2] shows that there exists $f \in \mathcal{S}^1_{\theta}(\mathbf{R})$ such that $e^{\pm ix^2} f \notin \mathcal{S}^s_{\theta}(\mathbf{R})$ for all $1 \leq s < \theta$. Lemma 5.6 (i) then implies that $e^{iq(x)}f \notin \mathcal{S}^s_{\theta}(\mathbf{R})$ for all $1 \leq s < \theta$ which proves Theorem 1.2 when m = 2.

Suppose next that $m \ge 3$. The assumptions imply $(m-2)\theta \ge 1$. First we show that there exists $f \in \mathcal{S}^1_{\theta}(\mathbf{R})$ such that $e^{\pm ix^m} f \notin \mathcal{S}^s_{\theta}(\mathbf{R})$ for all s > 0such that $1 \leq s < \theta m - \max(\theta, 1)$.

In fact for the function (5.8) we have $f \in \mathcal{S}^1_{\theta}(\mathbf{R})$ by Proposition 5.5. In order to show $e^{\pm ix^m} f \notin \mathcal{S}^s_{\theta}(\mathbf{R})$ for all s > 0 such that $1 \leq s < \theta m - \max(\theta, 1)$ we argue by contradiction. Let $1 \leq s < \theta m - \max(\theta, 1)$ and suppose that $e^{\pm ix^m} f \in \mathcal{S}^s_{\mathcal{A}}(\mathbf{R})$. By Lemma 2.1 the estimate (5.2) with $g(x) = e^{\pm ix^m}$ holds for some A, B, a > 0. By Corollary 5.4 and Lemma 5.1 with $\nu = \theta$ we may conclude that (5.3) holds. Thus (5.3) for $x = k^{\theta}$ gives with $\tau =$ $\max(\frac{1}{\theta} - 1, 0)$

$$\begin{aligned} |(D^{k}g)(k^{\theta})f(k^{\theta})| &\leq A(2B)^{k}k!^{s}e^{-ak}\langle k^{\theta}\rangle^{k\tau} \\ &\leq A(2B)^{k}k!^{s}e^{-ak}2^{\frac{1}{2}k\tau}k^{k(1-\min(\theta,1))} \\ &\leq A(2^{1+\tau}B)^{k}k^{k(s+1-\min(\theta,1))}e^{-ak}, \quad k \in \mathbf{N} \setminus \{0\}. \end{aligned}$$

$$(5.14)$$

On the other hand we may apply Proposition 3.9 since the assumptions imply $\theta > \frac{2}{m}$. Hence

$$\left| (D^{k}g)(k_{j}^{\theta})f(k_{j}^{\theta}) \right| = \left| p_{\pm im,k_{j}}(k_{j}^{\theta}) \right| e^{-\langle k_{j}^{\theta} \rangle^{\frac{1}{\theta}}} \ge \frac{1}{2} m^{k_{j}} k_{j}^{\theta k_{j}(m-1)} e^{-2\frac{1}{2\theta}k_{j}}, \quad j \in \mathbf{N} \setminus \{0\}$$

$$(5.15)$$

for some sequence $\{k_j\}_{j=1}^{+\infty} \subseteq \mathbf{N}$, using $k_j^{\theta} \ge 1$. Combining (5.15) and (5.14) yields

$$\frac{1}{2}m^{k_j}k_j^{k_j\theta(m-1)}e^{-2\frac{1}{2\theta}k_j} \leqslant A(2^{1+\tau}B)^{k_j}k_j^{k_j(s+1-\min(\theta,1))}e^{-ak_j}$$

for $j \in \mathbf{N} \setminus \{0\}$ which contradicts the assumption $s < \theta m - \max(\theta, 1)$. The assumption that $e^{\pm ix^m} f \in \mathcal{S}^s_{\theta}(\mathbf{R})$ hence must be wrong. Thus

$$e^{\pm ix^m} f \notin \bigcup_{1 \leqslant s < \theta m - \max(\theta, 1)} \mathcal{S}^s_{\theta}(\mathbf{R}),$$

and Lemma 5.6 (i) yields

$$Tf \notin \bigcup_{1\leqslant s < \theta m - \max(\theta, 1)} \mathcal{S}^s_{\theta}(\mathbf{R}).$$

Since $f \in \mathcal{S}^1_{\theta}(\mathbf{R}) \subseteq \mathcal{S}^s_{\theta}(\mathbf{R})$ when $s \ge 1$ we have in particular $T\mathcal{S}^s_{\theta}(\mathbf{R}) \not\subseteq \mathcal{S}^s_{\theta}(\mathbf{R})$ for any s > 0 such that $1 \leq s < \theta m - \max(\theta, 1)$.

Proof of Theorem 1.3.

Pick $\nu < \theta$ such that $\nu > \frac{2}{m}$, $1 < \nu < \theta$ if $\theta > 1$, and $1 < s < \nu m \max(\nu, 1)$. Set

$$f(x) = e^{-\langle x \rangle^{\frac{1}{\nu}}}, \quad x \in \mathbf{R}.$$

By Proposition 5.5 we have $f \in S^1_{\nu}(\mathbf{R})$. Let $g(x) = g_{\pm im}(x) = e^{\pm ix^m}$ and suppose that $T_g f \in \Sigma^{\sigma}_{\theta}(\mathbf{R})$ for some $\sigma > 0$ such that $1 < \sigma < \nu m - \max(\nu, 1)$. Then by Lemma 2.1 the estimate

$$|D^k(g(x)f(x))| \leqslant AB^k k!^{\sigma} e^{-a|x|^{\frac{1}{\theta}}}, \quad k \in \mathbf{N},$$

holds for all B, a > 0, and some A > 0 depending on B and a. By Corollary 5.4 and Lemma 5.1 we may conclude that

$$|(D^k g)(x)f(x)| \leq A(2B)^k k!^{\sigma} e^{-a|x|^{\frac{1}{\theta}}} \langle x \rangle^{k \max\left(\frac{1}{\nu} - 1, 0\right)}, \quad k \in \mathbf{N},$$

holds for some A, B, a > 0. This gives for $x = k^{\nu}$ with $\tau = \max\left(\frac{1}{\nu} - 1, 0\right)$

$$\begin{aligned} |(D^{k}g)(k^{\nu})f(k^{\nu})| &\leq A(2B)^{k}k!^{\sigma}e^{-ak\frac{\overline{\theta}}{\theta}}\langle k^{\nu}\rangle^{k\tau} \\ &\leq A(2^{1+\tau}B)^{k}k^{k(\sigma+1-\min(\nu,1))}e^{-ak\frac{\overline{\theta}}{\theta}}, \quad k \in \mathbf{N} \backslash \{0\}. \end{aligned}$$
(5.16)

Due to $\nu > \frac{2}{m}$ the assumptions for Proposition 3.9 with θ replaced by ν are satisfied. We obtain from Proposition 3.9

$$|(D^{k}g)(k_{j}^{\nu})f(k_{j}^{\nu})| = \left|p_{\pm im,k_{j}}(k_{j}^{\nu})\right| e^{-\langle k_{j}^{\nu} \rangle^{\frac{1}{\nu}}} \ge \frac{1}{2}m^{k_{j}}k_{j}^{\nu k_{j}(m-1)}e^{-2\frac{1}{2\nu}k_{j}}, \quad j \in \mathbf{N} \setminus \{0\}$$
(5.17)

for some sequence $\{k_j\}_{j=1}^{+\infty} \subseteq \mathbf{N}$, using $k_j^{\nu} \ge 1$.

Combining (5.17) and (5.16) yields finally

$$\frac{1}{2}m^{k_j}k_j^{k_j\nu(m-1)}e^{-2\frac{1}{2\nu}k_j} \leqslant A(2^{1+\tau}B)^{k_j}k_j^{k_j(\sigma+1-\min(\nu,1))}e^{-ak_j^{\frac{\nu}{\theta}}}$$

for $j \in \mathbf{N} \setminus \{0\}$ which contradicts the assumption $\sigma < \nu m - \max(\nu, 1)$. The assumption that $T_g f \in \Sigma_{\theta}^{\sigma}(\mathbf{R})$ hence must be wrong. Thus

$$T_g f \notin \bigcup_{1 < \sigma < \nu m - \max(\nu, 1)} \Sigma_{\theta}^{\sigma}(\mathbf{R}).$$

The assumptions for Lemma 5.6 (ii) are satisfied so we obtain

$$Tf \notin \bigcup_{1 < \sigma < \nu m - \max(\nu, 1)} \Sigma_{\theta}^{\sigma}(\mathbf{R}).$$

Since $f \in \mathcal{S}^1_{\nu}(\mathbf{R}) \subseteq \Sigma^s_{\theta}(\mathbf{R})$ and $1 < s < \nu m - \max(\nu, 1)$ we get $T\Sigma^s_{\theta}(\mathbf{R}) \notin$ $\Sigma^s_{\theta}(\mathbf{R}).$

Remark 5.7. When $g(x) = g_{\pm im}(x) = e^{\pm ix^m}$ it is possible to prove the lack of continuity of the operator $T = T_g$ in Theorems 1.2 and 1.3 on $\mathcal{S}^s_{\theta}(\mathbf{R})$ and $\Sigma^s_{\theta}(\mathbf{R})$ respectively, by other means. Here we may weaken the assumptions into $0 < s < (m-1)\theta$ and $\theta \ge \frac{2}{m}$. In fact we may use the criterion (4.1) and [4, Theorem 5.7].

Consider the assumptions of Theorem 1.2 relaxed into $0 < s < (m-1)\theta$ and $\theta \ge \frac{2}{m}$. By Proposition 3.9 we have

$$\left| (D^k g)(k_j^{\theta}) \right| = \left| p_{\pm im, k_j}(k_j^{\theta}) \right| \ge \frac{1}{2} m^{k_j} k_j^{\theta k_j(m-1)} \quad j \in \mathbf{N} \setminus \{0\},$$
(5.18)

for some sequence $\{k_j\}_{j=1}^{+\infty} \subseteq \mathbf{N}$.

Suppose that (4.1) holds for all $\lambda > 0$, some $C = C(\lambda) > 0$, and some $h = h(\lambda) > 0$. Then we obtain from (5.18)

$$\frac{1}{2}m^{k_j}k_j^{\theta k_j(m-1)} \leqslant Ch^{k_j}k_j!^s e^{\lambda^{-\frac{1}{\theta}}k_j} \leqslant Ch^{k_j}k_j^{sk_j}e^{\lambda^{-\frac{1}{\theta}}k_j}, \quad j \in \mathbf{N} \setminus \{0\}.$$
(5.19)

If $s < (m-1)\theta$ this is a contradiction. By [4, Theorem 5.7] it follows that T_g is not continuous on $\mathcal{S}^s_{\theta}(\mathbf{R})$.

Consider finally the assumptions of Theorem 1.3 relaxed into $0 < s < (m-1)\theta$ and $\theta \ge \frac{2}{m}$. Suppose that (4.1) holds for all h > 0, some C = C(h) > 0, and some $\lambda = \lambda(h) > 0$. Again the estimate (5.19) gives a contradiction if $s < (m-1)\theta$. It follows by [4, Theorem 5.7] that T_g is not continuous on $\Sigma^s_{\theta}(\mathbf{R})$.

References

- A. Arias Junior, Some remarks on the Cauchy problem for Schrödinger type equations in Gelfand-Shilov spaces, arXiv:2309.08194 [math.AP] (2023).
- [2] E. Bierstone and P. D. Milman, Resolution of singularities in Denjoy-Carleman classes, Selecta Math. (N. S.) 10 (1) (2004), 1–28.
- [3] J. Chung, S. Y. Chung and D. Kim, Characterization of the Gelfand-Shilov spaces via Fourier transforms, Proc. AMS 124 (7), 2101–2108, 1996.
- [4] A. Debrouwere and L. Neyt, Weighted (PLB)-spaces of ultradifferentiable functions and multiplier spaces, Monatsh. Math. 198 (1) (2022), 31–60.
- [5] I. M. Gel'fand and G. E. Shilov, *Generalized Functions*, Vol. II, Academic Press, New York London, 1968.
- [6] T. Gramchev and P. Popivanov, Partial differential equations. Approximate solutions in scales of functional spaces, Math. Res. 108, Wiley-VCH Verlag Berlin GmbH, Berlin, 2000.
- [7] F. Nicola and L. Rodino, Global Pseudo-Differential Calculus on Euclidean Spaces, Birkhäuser, Basel, 2010.
- [8] A. Petersson, Fourier characterizations and non-triviality of Gelfand-Shilov spaces, with applications to Toeplitz operators, J. Fourier Anal. Appl. **29** (3) 29 (2023).
- [9] H. H. Schaefer and M. P. Wolff, *Topological Vector Spaces*, Springer-Verlag, New York, 1999.
- [10] P. Wahlberg, Propagation of anisotropic Gelfand-Shilov wave front sets, J. Pseudo-Differ. Oper. Appl. 14 (1) 7 (2023).

DEPARTMENT OF COMPUTER SCIENCE AND MATHEMATICS (DCM - FFCLRP), UNI-VERSITY OF SÃO PAULO (USP), RIBEIRÃO PRETO, SP, 14040-901, BRAZIL

Email address: alexandre.ariasjunior[AT]usp.br

DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY

Email address: patrik.wahlberg[AT]polito.it