

ON THE EQUIVALENCE OF L^p -PARABOLICITY, L^q -LIOUVILLE PROPERTY ON WEIGHTED GRAPHS

LU HAO AND YUHUA SUN

ABSTRACT. We study the relationship between the L^p -parabolicity, the L^q -Liouville property for positive superharmonic functions, and the existence of nonharmonic positive solutions to the system

$$\begin{cases} -\Delta u \geq 0, \\ \Delta(|\Delta u|^{p-2}\Delta u) \geq 0, \end{cases}$$

on weighted graphs, where $1 \leq p < \infty$ and (p, q) are Hölder conjugate exponent pair. Moreover, some new technique is developed to establish the estimate of green function under volume doubling and Poincaré inequality conditions, and the sharp volume growth conditions for the L^p -parabolicity can be derived on some graphs.

1. INTRODUCTION

Throughout the paper, let $G = (V, E)$ be an infinite, connected, locally finite graph, where V denotes the vertex set, and E denotes the edge set. If there exists an edge connecting x and y , that is, $(x, y) \in E$, we denote it by $x \sim y$. The edge (x, x) is called a loop. A graph G is called simple if it has no loops.

Let $\mu : V \times V \rightarrow [0, \infty)$ be an edge weight, and denote it by $\mu_{xy} := \mu(x, y)$. Note $\mu_{xy} > 0$ if and only if $x \sim y$. Moreover, $\mu_{xy} = \mu_{yx}$. The weight of a vertex x is defined as

$$\mu(x) = \sum_{y \sim x} \mu_{xy}.$$

Such graph (V, E, μ) is called a weighted graph, and usually simplified as (G, μ) or (V, μ) .

A random walk $\{X_n\}$ on a locally finite weighted graph (V, μ) is a Markov chain with the following transition probability

$$P(x, y) = \begin{cases} \frac{\mu_{xy}}{\mu(x)}, & \text{if } x \sim y, \\ 0, & \text{otherwise,} \end{cases} \quad (1.1)$$

Such Markov chain is also denoted by the pair (G, P) . Since $\mu(x)P(x, y) = \mu(y)P(y, x)$, then the above Markov Chain is called reversible. Conversely a reversible Markov Chain on V can determine a weighted graph (V, E, μ) by letting

$$\mu_{xy} := P(x, y)\mu(x),$$

and edge set $E = \{x \sim y | \mu_{xy} > 0\}$. Henceforth, a locally finite weighted graph (V, μ) is also referred to the related reversible Markov Chain (G, P) and the corresponding random walk $\{X_n\}$.

2020 *Mathematics Subject Classification.* Primary 31B35, 31E05; Secondary 35J91.

Key words and phrases. weighted graph, L^p parabolicity, L^q Liouville property, green function, volume growth.

Lu Hao was funded by the Deutsche Forschungsgemeinschaft(DFG, German Research Foundation)-Project-ID317210226-SFB 1283. Yuhua Sun was funded by the National Natural Science Foundation of China (Grant No.12371206).

For our convenience, let us denote the n -step transition function as

$$P_n(x, y) := \mathbb{P}_x[X_n = y] = \mathbb{P}[X_0 = x, X_n = y],$$

Hence, $P_0(x, y) = \delta_x(y)$, and $P_1(x, y) = P(x, y)$.

Let $\ell(U)$ be the collection of all real functions on $U \subset V$, $\ell_0(U)$ be the subset of $\ell(U)$ with compact support, and $\ell^+(U)$ be the set of nonnegative functions. Moreover, for $1 \leq p < \infty$, we define $L^p(U) = \{f \in \ell(U) : \sum_{x \in V} |f(x)|^p \mu(x) < \infty\}$. Additionally, we define $L_+^p(U) = L^p(U) \cap \ell^+(U)$.

Let us define the Laplace operator $\Delta : \ell(V) \rightarrow \ell(V)$ on weighted graph (V, μ) by

$$\begin{aligned} \Delta u(x) &= \frac{1}{\mu(x)} \sum_{y \sim x} \mu_{xy} (u(y) - u(x)) \\ &= \sum_{y \in V} P(x, y) (u(y) - u(x)) \\ &= (P - I)u(x), \end{aligned}$$

where the Markov operator P is defined by

$$Pu(x) = \sum_{y \in V} P(x, y) u(y).$$

Let

$$p_n(x, y) = \frac{P_n(x, y)}{\mu(y)},$$

It follows by reversibility that

$$p_n(x, y) = p_n(y, x).$$

The Green function of Δ on (V, μ) can be defined by

$$g(x, y) = \sum_{n=0}^{\infty} p_n(x, y), \tag{1.2}$$

and hence $g(x, y) = g(y, x)$.

For $u \in \ell(V)$, we define Green operator G by

$$Gu(x) = \sum_{y \in V} g(x, y) u(y) \mu(y).$$

Fix $A \subset V$ and $\nu \in \ell^+(V)$, define

$$\nu(A) = \sum_{x \in A} \nu(x).$$

Now fix $1 \leq p < \infty$, let us define the L^p -capacity of a finite set $K \subset V$ by

$$C_p(K) = \sup\{\nu(K)^p : \nu = f\mu, f \in \ell^+(K), \|Gf\|_{L^q(V)} \leq 1\}, \tag{1.3}$$

where the norm $\|Gf\|_{L^q(V)}$ is defined by

$$\|Gf\|_{L^q(V)} = \left(\sum_{x \in V} |Gf(x)|^q \mu(x) \right)^{\frac{1}{q}},$$

and q is the Hölder conjugate number of p , namely, $\frac{1}{p} + \frac{1}{q} = 1$.

Definition 1.1. A graph (V, μ) is called L^p -parabolic if $C_p(K) = 0$ for every finite subset $K \subset V$.

Definition 1.2. We say a graph (V, μ) admits L^q -Liouville property if any positive L^q -superharmonic function on (V, μ) is constant.

A random walk is called recurrent if it returns to the starting vertex infinitely many times. If every random walk on graph is recurrent, then the underlying graph is called parabolic. There are various equivalent characterizations of parabolicity in terms of different fields, for instance, any positive superharmonic function being constant or equivalently; and the capacity of any finite set being zero, see [2],[10],[19]. These characterizations provide critical insights into the connections between stochastic processes, graph theory and potential theory. If the graph is not parabolic, the random walk (or, the graph) is called transient.

A random walk is called recurrent if it returns to the starting vertex infinitely many times. If every random walk on graph is recurrent, then the underlying graph is called parabolic. There are various equivalent characterizations of parabolicity in terms of different fields, for instance, any positive superharmonic function being constant or equivalently; and the capacity of any finite set being zero, see [2],[10],[19]. These characterizations provide critical insights into the connections between stochastic processes, graph theory and potential theory.

For $1 < p < \infty$, we call a manifold (resp. graph) is p -parabolic if for any compact (resp. finite) K , where the p -capacity of K is defined on manifold M (resp. graph (V, μ)) by

$$\text{Cap}_p(K) = \inf\left\{ \int_M |\nabla u|^p : u \in W_0^{1,p}(M) \cap C_0^0(M), u \geq 1 \text{ on } K \right\},$$

respectively,

$$\text{Cap}_p(K) = \inf\left\{ \sum_{x,y \in V} \mu_{xy} |u(y) - u(x)|^p : u \in \ell_0(V), u \geq 1 \text{ on } K \right\}.$$

The p -parabolicity has been well studied in both manifolds and graphs, see [12], [18], [5], [20], [17], and [15]. In particular, Holopainen and Saloff-Coste, working on manifolds [12] and graphs [15], respectively, proved that p -parabolicity is equivalent to the Liouville property for p -superharmonic functions. This Liouville property states that any nonnegative solution of $-\Delta_p u \geq 0$ must be constant. Moreover, parabolicity is equivalent to 2-parabolicity and it is also worth noting that, on a graph G , the p -Laplacian is defined as

$$-\Delta_p u(x) = \sum_{y \sim x} \frac{\mu_{xy}}{\mu(x)} |u(y) - u(x)|^{p-2} (u(y) - u(x)), \quad \text{for } u \in \ell(V).$$

Furthermore, in both manifold and graph settings, they obtained a volume growth criterion for p -parabolicity, see [5] and [15]. We present a simple corollary here: if on a connected, complete manifold M (resp. a connected graph $G = (V, \mu)$) satisfies, for some $o \in M$ (resp. $o \in V$), the volume growth condition

$$V(o, r) \lesssim r^p (\log r)^{p-1},$$

then M (resp. G) is p -parabolic.

A manifold M is called biparabolic if any nonnegative solution of the system

$$\begin{cases} -\Delta u \geq 0, \\ \Delta^2 u \geq 0. \end{cases} \quad (1.4)$$

on M is harmonic, that is, $\Delta u = 0$. In [7] Faraji and Grigor'yan studied the biparabolicity of Riemannian manifolds, and obtained a nearly optimal criterion condition, namely, if

the manifold M is geodesically complete and satisfies

$$V(x_0, r) \lesssim \frac{r^4}{\log r}, \quad (1.5)$$

then the manifold M is bipolarabolic. Here $V(x_0, r)$ stands for the Riemannian volume of geodesic ball centered at x_0 with radius r .

Very recently, Grigor'yan, Pessoa and Setti [8] studied the equivalence between the L^p -parabolicity and the L^q -Liouville property on Riemannian manifolds, where p and q are Hölder conjugate exponents. Moreover, they obtained that bipolarabolicity of manifold is equivalent to L^2 -parabolicity on manifold.

Motivated by these results, our object is to study the relationship on graph (V, μ) between L^p -parabolicity, L^q -Liouville property, and the existence of nonnegative nonharmonic solution to the following system

$$\begin{cases} -\Delta u \geq 0, \\ \Delta(|\Delta u|^{p-2}\Delta u) \geq 0, \end{cases} \quad (1.6)$$

where $\frac{1}{p} + \frac{1}{q} = 1$. As a particular case $p = 2$, (1.6) is simplified to (1.4). When $1 \leq p < 2$, the second inequality of (1.6) is understood as $-\Delta(|\Delta u|^{p-1}) \geq 0$. We emphasize that the equivalence of existence of nonnegative nonharmonic solutions to the system (1.6) with L^p -parabolicity and L^q -Liouville property is only obtained when $p = q = 2$ on manifold case in [8].

Definition 1.3. A graph (V, μ) is called bipolarabolic if any nonnegative solution of system (1.4) on (V, μ) is harmonic.

Our main result is announced as follows.

Theorem 1.4. For $1 \leq p < \infty$, the following three conditions are equivalent.

- (I). (V, μ) is L^p -parabolic.
- (II). (V, μ) admits L^q -Liouville property.
- (III). Any nonnegative solution of system (1.6) on (V, μ) is harmonic.

Remark 1.5. We have four motivational comments:

- (1) We emphasize there the equivalence of (III) and the other conditions was not obtained in manifold case for general $1 \leq p < \infty$, and the equivalence on manifold case is only obtained for $p = 2$. However, we can deal with general case of p on weighted graphs.
- (2) For $p = 1$, L^1 -parabolicity is equivalent to parabolicity, and thus equivalent to that (V, μ) admits L^∞ -Liouville property. By noting that the minimum of a superharmonic function and a constant is again a superharmonic function, condition (III) is equivalent to the parabolicity of graph.
- (3) For $p = 2$, Theorem 1.4 implies that L^2 -parabolicity is equivalent to bipolarabolicity on (V, μ) .

For any two vertices x and y , let us define $d(x, y)$ to be the minimal number of edges among all possible paths connecting x and y on graph (V, μ) . Then $d(\cdot, \cdot)$ is a distance function on $V \times V$, and called the graph distance. Fix some vertex $o \in V$, and for $r > 0$, denote

$$B(o, r) := \{x \in V \mid d(o, x) \leq r\},$$

and

$$V(o, r) = \mu(B(o, r)).$$

We are ready to give some sufficient condition for L^p -parabolicity in terms of volume growth. First, we give a nearly optimal volume condition.

Theorem 1.6. For $1 < p \leq 2$, assume there exists a positive constant $b > 0$ such that $\mu(x) \geq b$ for any $x \in V$. If for some $o \in V$, we have

$$V(o, r) \lesssim \frac{r^{2p}}{\log r}, \quad \text{for all large enough } r. \quad (1.7)$$

Then (V, μ) is L^p -parabolic.

To obtain a sharp volume growth condition, some other geometric conditions on graph are also needed.

Definition 1.7. We say that the weighted graph (V, μ) satisfies volume doubling condition $\text{VD}(C_1)$, if there exists some positive constant C_1 such that for all $x \in V$ and all $r > 0$, the following holds

$$V(x, 2r) \leq C_1 V(x, r). \quad (\text{VD})$$

Definition 1.8. We say a weighted graph (V, μ) admits the Poincaré inequality $\text{PI}(C_2)$, if there exists a positive constant C_2 such that for all $x_0 \in V$, all $r > 0$, and all $f \in \ell(V)$, there holds

$$\sum_{x \in B(x_0, r)} |f(x) - f_B|^2 \mu(x) \leq C_2 r^2 \sum_{x, y \in B(x_0, 2r)} \mu_{xy} (f(y) - f(x))^2, \quad (\text{PI})$$

where

$$f_B = \frac{1}{V(x_0, r)} \sum_{x \in B(x_0, r)} f(x) \mu(x).$$

Definition 1.9. We say that the weighted graph (V, μ) satisfies $P_0(\alpha)$ -condition

$$\frac{\mu_{xy}}{\mu(x)} \geq \alpha \quad \text{when } y \sim x. \quad (P_0)$$

Under the conditions of (VD), (PI) and (P_0) , we introduce a new operation technique on (V, μ) which is different from the one in the existing literature (see [3, 6]), and we can drop the loop assumption in Delmotte's heat kernel estimate in [6]. Using this improvement, we can derive the following Li-Yau type Green function estimate:

Theorem 1.10. Assume (V, μ) satisfies (VD), (PI) and (P_0) , then there exist constants $C, C' > 0$ such that

$$C \sum_{n=d(x,y)}^{\infty} \frac{n}{V(x, n)} \leq g(x, y) \leq C' \sum_{n=d(x,y)}^{\infty} \frac{n}{V(x, n)}. \quad (1.8)$$

Theorem 1.11. Let $1 < p < \infty$, assume that conditions (VD), (PI), and (P_0) are satisfied on (V, μ) . If there exists some $o \in V$ such that

$$V(o, r) \lesssim r^{2p} (\log r)^{p-1}, \quad \text{for all large enough } r, \quad (1.9)$$

then (V, μ) is L^p -parabolic.

Notations. In the above and below, let f, g be functions from \mathbb{R}_+ to \mathbb{R}_+ . $f \lesssim g$ means that the quotient of f and g is bounded from the above for all $r > 0$ and $f \sim g$ means both $f \lesssim g$ and $g \lesssim f$ hold.

2. PRELIMINARY

Fix a subset $U \subset V$. We introduce the transition probability for the process $\{X_n\}$ killed on exiting from U (see [2, section 1.5])

$$P_n^U(x, y) = \mathbb{P}[X_0 = x, X_n = y, n < \tau_U],$$

where $\tau_U = \min\{n \geq 0 : X_n \notin U\}$ is the first exit time from U .

Then the heat kernel is given by

$$p_n^U(x, y) = \frac{1}{\mu(y)} P_n^U(x, y).$$

Clearly,

$$\begin{cases} P_n^V(x, y) = P_n(x, y), \\ p_n^U(x, y) = p_n^U(y, x), \\ p_n^U(x, y) = 0, \quad \text{if } x \in U^c \text{ or } y \in U^c. \end{cases}$$

Introduce the following operators:

$$\begin{cases} I_U f(x) = \mathbf{1}_U(x) f(x), \\ P^U f(x) = \sum_{y \in U} P_1^U(x, y) f(y), \\ \Delta_U f(x) = (P^U - I_U) f(x), \\ P_n^U f(x) = \sum_{y \in U} P_n^U(x, y) f(y), \end{cases}$$

where $\mathbf{1}_U(x) = 1$ when $x \in U$ and $\mathbf{1}_U(x) = 0$ otherwise. From the above definitions, we can see that $P^U f(x) = P_1^U f(x)$.

The Green's function of Δ_U on U is defined as

$$g^U(x, y) = \sum_{n=0}^{\infty} p_n^U(x, y),$$

Consequently, the following properties hold:

$$g^V(x, y) = g(x, y) \quad \text{and} \quad g^U(x, y) = g^U(y, x).$$

Furthermore, it is known that

$$-\Delta g^U(x, x_0) = \frac{1}{\mu(x_0)} \mathbf{1}_{\{x_0\}}(x) \quad \text{for } x, x_0 \in U.$$

The following result is known for existing literature, for example, see [2, Theorem 1.31].

Proposition 2.1. $g^U(x, y) < \infty$ for any $x, y \in U$ provided that either of the following case holds

- (i). (V, μ) is transient. (ii). $U \neq V$.

For $u \in \ell(V)$, define the Green operator G by

$$G^U u(x) = \sum_{y \in U} g^U(x, y) u(y) \mu(y),$$

where $G^U u(x) = 0$ for any $x \in U^c$.

Now, for $1 < q < \infty$, the L^q -Green function $g_q(x, y)$ is defined as

$$g_q(x, y) = \sum_{z \in V} g(x, z) g(z, y)^{q-1} \mu(z). \quad (2.1)$$

It follows that if $g(x, y) \equiv \infty$, then $g_q(x, y) \equiv \infty$.

From the Markov property of random walk, we can deduce the following properties (for details, see [2, Section 1.5 and 1.6]).

Proposition 2.2. For $n \in \mathbb{N}_+$, we have

- (i) $P_{n+1}^U(x, y) = \sum_{z \in U} P_1^U(x, z) P_n^U(z, y)$.
- (ii) $P_0^U f(x) = \mathbf{1}_U f(x)$, and $P_n^U f(x) = (P^U)^n f(x)$.
- (iii) For $u \in \ell(V)$, $G^U u(x) = \sum_{n=0}^{\infty} P_n^U u(x) = \mathbb{E}[\sum_{n=0}^{\tau_U-1} u(X_n) | X_0 = x]$, where $\mathbb{E}(f)$ is the expectation of random variable f .

Using these properties, we obtain the following lemmas.

Lemma 2.3. If $g_q(x_0, y_0) < \infty$ holds for some $x_0, y_0 \in V$, then for any $x, y \in V$, $g_q(x, y) < \infty$.

Proof. Since (V, μ) is connected, then for any x, y , there exist some nonnegative integer i and j such that

$$P_i(x_0, x) > 0, \quad P_j(y_0, y) > 0.$$

Recalling

$$P_{n+m}(x, y) = \sum_{z \in V} P_n(x, z) P_m(z, y),$$

and taking the sum over m

$$\sum_{m=0}^{\infty} P_{n+m}(x, y) = \sum_{m=0}^{\infty} \sum_{z \in V} P_n(x, z) P_m(z, y) = \sum_{z \in V} P_n(x, z) \sum_{m=0}^{\infty} P_m(z, y),$$

and by noting that

$$g(x, y) \mu(y) = \sum_{n=0}^{\infty} P_n(x, y),$$

we have for all $x, y, z \in V$, and all $n \in \mathbb{N}$

$$g(x, y) \geq P_n(x, z) g(z, y).$$

Specially, by using that

$$\begin{cases} g(x_0, z) \geq P_i(x_0, x) g(x, z), \\ g(y_0, z) \geq P_j(y_0, y) g(z, y), \end{cases}$$

we obtain

$$\begin{aligned} g_q(x_0, y_0) &= \sum_{z \in V} g(x_0, z) g(z, y_0)^{q-1} \mu(z) \\ &\geq P_i(x_0, x) P_j(y_0, y)^{q-1} \sum_{z \in V} g(x, z) g(z, y)^{q-1} \mu(z) \\ &= P_i(x, x_0) P_j(y_0, y)^{q-1} g_q(x, y). \end{aligned}$$

Since $P_i(x, x_0), P_j(y, y_0) > 0$, thus we can complete the proof. \square

Lemma 2.4. Let (V, μ) be transient or $U \neq V$. If $G^U f(x) \in \ell(U)$, then

$$(-\Delta_U) G^U f(x) = f(x), \quad \text{for } x \in U. \quad (2.2)$$

Proof. Using local finiteness of graph, we obtain

$$\begin{aligned} P^U G^U f &= P^U \sum_{n=0}^{\infty} P_n^U f \\ &= \sum_{n=0}^{\infty} P_{n+1}^U f \\ &= G^U f - I_U f, \end{aligned}$$

whence $(-\Delta_U)G^U f(x) = (I_U - P^U)G^U f(x) = I_U f(x)$, the proof is complete. \square

3. L^p -CAPACITY

Throughout this section, since we need to use local Green function G^U to define capacity, hence we emphasize here that (V, μ) is transient or $U \neq V$, see Proposition 2.1.

Definition 3.1. Fix any finite subset $K \subset U$ and $1 \leq p < \infty$, let L^p -capacity of (K, U) be defined by

$$C_p(K, U) = \sup\{\nu(K)^p : \nu = f\mu, f \in \ell^+(K), \|G^U f\|_{L^q(U)} \leq 1\}, \quad (3.1)$$

where q is the Hölder conjugate exponent of p . When $U = V$, we denote by $C_p(K) := C_p(K, V)$ for simplicity, see (1.3).

Remark 3.2. From the definition of $C_p(K, U)$, it is easy to verify that

$$C_p(K, U) = \sup\{\nu(K)^p : \nu = f\mu, f \in \ell^+(K), \|G^U f\|_{L^q(U)} = 1\}, \quad (3.2)$$

Now we claim that

$$\min_{\nu \in \ell^+(K)} \frac{\|G^U f\|_{L^q(U)}}{\nu(K)} = \frac{1}{C_p(K, U)^{\frac{1}{p}}}. \quad (3.3)$$

Noting that (V, μ) is transient or $U \neq V$, we always have $C_p(K, U) > 0$. First, for any small enough $\epsilon > 0$, there exists $f \in \ell^+(K)$ such that $\|G^U f\|_{L^q(U)} \leq 1$, and $\nu = f\mu$ such that

$$\nu(K) \geq (C_p(K, U) - \epsilon)^{\frac{1}{p}},$$

which gives that

$$\frac{\|G^U f\|_{L^q(U)}}{\nu(K)} \leq \frac{1}{(C_p(K, U) - \epsilon)^{\frac{1}{p}}},$$

It follows that

$$\min_{\nu \in \ell^+(K)} \frac{\|G^U f\|_{L^q(U)}}{\nu(K)} \leq \frac{1}{(C_p(K, U) - \epsilon)^{\frac{1}{p}}},$$

Letting $\epsilon \rightarrow 0$, we obtain that

$$\min_{\nu \in \ell^+(K)} \frac{\|G^U f\|_{L^q(U)}}{\nu(K)} \leq \frac{1}{C_p(K, U)^{\frac{1}{p}}},$$

On the other hand, for any $\nu = f\mu$ such that $\|G^U f\|_{L^q(U)} = 1, f \in \ell^+(V)$, we have

$$\frac{1}{C_p(K, U)^{\frac{1}{p}}} \leq \frac{1}{\nu(K)} = \frac{\|G^U f\|_{L^q(U)}}{\nu(K)}, \quad (3.4)$$

Since ν is arbitrary, we obtain that

$$\frac{1}{C_p(K, U)^{\frac{1}{p}}} \leq \min_{\nu \in \ell^+(K)} \frac{\|G^U f\|_{L^q(U)}}{\nu(K)}, \quad (3.5)$$

Hence, we obtain that (3.3).

Remark 3.3. Indeed, there exist many other capacities in the existing literature, for example, the harmonic capacity $\text{Cap}(K, U)$

$$\text{Cap}(K, U) = \inf \left\{ \sum_{x, y \in U} (f(x) - f(y))^2 \mu_{xy} : f \in \ell_0(U), f \geq 1 \text{ on } K \right\}.$$

If $U = V$, denote $\text{Cap}(K) := \text{Cap}(K, V)$. It is known that for any finite subset $K \subset U$ with U finite or $U = V$, we have

$$\text{Cap}(K, U) = C_1(K, U).$$

see [2, Proposition 7.9].

We introduce two different equivalent characterization of $C_p(K, U)$ for $1 < p < \infty$, which are more convenient for us. For $1 < p < \infty$, and let K be a finite set of U , and define the following two capacities

$$\hat{C}_p(K, U) = \inf \left\{ \|f\|_{L^p(U)}^p : f \in L_+^p(U), G^U f \geq 1 \text{ on } K \right\}$$

and

$$\bar{C}_p(K, U) = \inf \left\{ \|\Delta f\|_{L^p(U)}^p : f \in \ell_0(U), f \geq 1 \text{ on } K \right\}$$

Theorem 3.4. For $1 < p < \infty$, and let $K \subset U$ be a finite subset of U , then

$$C_p(K, U) = \hat{C}_p(K, U) = \bar{C}_p(K, U). \quad (3.6)$$

Remark 3.5. For $p = 1$, the second equality still holds, and the proof for this case remains valid. However, the first equality does not hold.

Proof. We show the first equality of (3.6) by using the similar argument of [1, Theorem 2.5.1] as in \mathbb{R}^n . For our convenience, we give a simple argument here. Define a bilinear functional $\mathfrak{E}(\cdot, \cdot)$ by

$$\mathfrak{E}(\nu, u) = \sum_{x \in U} G^U u(x) \nu(x), \quad \text{for } (\nu, u) \in X \times Y,$$

where

$$\begin{aligned} X &= \{ \nu : \nu \in \ell^+(U), \nu(K) = 1, \nu(x) = 0 \text{ when } x \in V \setminus K \}, \\ Y &= \{ u : u \in L_+^p(U), \|u\|_{L^p(U)} \leq 1 \}. \end{aligned}$$

Note that

$$\sum_{x \in U} G^U u(x) \nu(x) = \sum_{x, y \in U} g^U(x, y) u(y) \mu(y) \nu(x) = \sum_{y \in U} G^U f(y) u(y) \mu(y),$$

where $f(x) = \frac{\nu(x)}{\mu(x)}$. Thus,

$$\sup_{u \in Y} \mathfrak{E}(\nu, u) = \sup_{u \in Y} \sum_{y \in U} G^U f(y) u(y) \mu(y) = \|G^U f\|_{L^q(U)}.$$

It follows that

$$\min_{\nu \in X} \sup_{u \in Y} \mathfrak{E}(\nu, u) = \min_{\nu \in \ell^+(K)} \frac{\|G^U f\|_{L^q(U)}}{\nu(K)} = \frac{1}{C_p(K, U)^{\frac{1}{p}}}.$$

Similarly

$$\min_{\nu \in X} \mathfrak{E}(\nu, f) = \min_{\nu \in X} \sum_{x \in U} G^U f(x) \nu(x) = \min_{x \in K} G^U f(x),$$

we derive

$$\sup_{f \in Y} \min_{\nu \in X} \mathcal{E}(\nu, f) = \sup_{f \in L_+^p(U)} \frac{\min_{x \in K} G^U f(x)}{\|f\|_{L^p(U)}} = \frac{1}{\hat{C}_p(K, U)^{\frac{1}{p}}}.$$

Since X and Y are convex, X is a close subset of $\mathbb{R}^{|K|}$, and the function $\mathcal{E}(\nu, f)$ is continuous in ν for fixed f . Here $|K|$ stands for the number of vertices in K . By Mini-Max Theorem of [1, Theorem 2.4.1], we obtain

$$C_p(K, U) = \hat{C}_p(K, U),$$

which shows the first equality of (3.6).

Now let us prove the second equality of (3.6). Let $\{U_n\}$ be an increasing exhaustion sequence of U , for any $f \in L_+^p(U)$ with $G^U f \geq 1$ on K , define

$$u_{n,\epsilon} := (1 + \epsilon)G^{U_n} f,$$

where $(n, \epsilon) \in \mathbb{N} \times \mathbb{R}_+$.

Noting that for any $x, y \in U$, $g^{U_n}(x, y)$ monotone increasingly converges to $g^U(x, y)$, and hence we have

$$G^{U_n} f(x) \uparrow G^U f(x), \quad \text{for all } x \in K,$$

which implies that $u_{n,\epsilon} \geq 1$ on K holds for large enough n .

By Lemma 2.4, we have

$$\|\Delta u_{n,\epsilon}(x)\|_{L^p(U_n)}^p = (1 + \epsilon)^p \sum_{x \in U_n} |\Delta G^{U_n} f(x)|^p \mu(x) = (1 + \epsilon)^p \sum_{x \in U_n} |f(x)|^p \mu(x).$$

Therefore

$$\bar{C}_p(K, U) \leq (1 + \epsilon)^p \|f\|_{L^p(U_n)}^p. \quad (3.7)$$

By letting $n \rightarrow \infty$, we obtain

$$\bar{C}_p(K, U) \leq (1 + \epsilon)^p \hat{C}_p(K, U).$$

By the arbitrariness of ϵ , we derive that

$$\bar{C}_p(K, U) \leq \hat{C}_p(K, U). \quad (3.8)$$

On the other hand, for any $f \in \ell_0(U)$ satisfying $f \geq 1$ on K , we have

$$G^U(-\Delta_U)f = \sum_{n=0}^{\infty} P_n^U(I_U - P^U)f = \sum_{n=0}^{\infty} (P_n^U f - P_{n+1}^U f) \quad (3.9)$$

Noting that (V, μ) is transient or satisfies $U \neq V$, and $f \in \ell_0(U)$, we obtain

$$-\Delta f(x) = -\Delta_U f(x) \quad \text{and} \quad \lim_{n \rightarrow \infty} P_n^U f(x) = 0, \quad \forall x \in U$$

Hence, (3.9) implies that for all $x \in U$,

$$\begin{aligned} G^U(-\Delta)f(x) &= G^U(-\Delta_U)f(x) \\ &= \lim_{l \rightarrow \infty} \sum_{n=0}^l (P_n^U f(x) - P_{n+1}^U f(x)) \\ &= \lim_{l \rightarrow \infty} (f(x) - P_{l+1}^U f(x)) \\ &= f(x). \end{aligned}$$

Lastly, letting $u = |\Delta f|$, we obtain that

$$u \in \ell_0(V) \subset L_+^p(U), \quad G^U u \geq G^U(-\Delta)f = f \geq 1 \quad \text{on } K,$$

and

$$\|u\|_{L^p(U)}^p = \sum_U u^p \mu = \sum_U |\Delta f|^p \mu = \|\Delta f\|_{L^p(U)}^p,$$

which implies

$$\hat{C}_p(K, U) \leq \bar{C}_p(K, U). \quad (3.10)$$

Combining (3.8) and (3.10), we derive $\hat{C}_p(K, U) = \bar{C}_p(K, U)$. Hence, we complete the proof. \square

Proposition 3.6. Let $1 < p < \infty$, and $K_1 \subset K_2 \subset U_1 \subset U_2$, where K_1 and K_2 are finite sets, then

$$C_p(K_1, U_1) \leq C_p(K_2, U_1), \quad C_p(K_1, U_2) \leq C_p(K_1, U_1), \quad (3.11)$$

and

$$C_p(K_1 \cup K_2, U) \leq C_p(K_1, U) + C_p(K_2, U). \quad (3.12)$$

Proof. (3.11) can be derived by the definition of $C_p(K, U)$. (3.12) can be derived by the definition of $\hat{C}_p(K, U)$ and Theorem 3.4. Indeed, Fix $\epsilon > 0$, we can choose $f_i \in L_+^p(U)$ ($i=1, 2$) such that $G^U f_i \geq 1$ on K_i and $\|f_i\|_{L^p(U)}^p \leq C_p(K_i, U) + \frac{\epsilon}{2}$. Let us define $f(x) = \max\{f_1, f_2\}$, obviously, $G^U f \geq 1$ on $K_1 \cup K_2$, and

$$\hat{C}_p(K_1 \cup K_2, U) \leq \sum_U f^p \mu \leq \sum_U f_1^p \mu + \sum_U f_2^p \mu \leq C_p(K_1, U) + C_p(K_2, U) + \epsilon.$$

Letting $\epsilon \rightarrow 0$, we obtain

$$\hat{C}_p(K_1 \cup K_2, U) \leq C_p(K_1, U) + C_p(K_2, U),$$

Thus, we complete the proof. \square

Proposition 3.7. Let $\{U_n\}$ be an increasing exhaustion of U and $K \subset U$ be a finite set. Then, for any $1 < p < \infty$,

$$\lim_{n \rightarrow \infty} C_p(K, U_n) = C_p(K, U).$$

Proof. By (3.11), it suffices to show

$$\lim_{n \rightarrow \infty} C_p(K, U_n) \leq C_p(K, U).$$

Let $f \in L_+^p(U)$ and satisfy that $G^U f \geq 1$ on K . Define

$$f_{n,\epsilon} = (1 + \epsilon)f \mathbf{1}_{U_n},$$

where $(n, \epsilon) \in \mathbb{N} \times \mathbb{R}_+$.

Since

$$\lim_{n \rightarrow \infty} G^{U_n} f_{n,\epsilon}(x) = (1 + \epsilon)G^U f(x) \geq (1 + \epsilon), \quad \text{for any } x \in K,$$

It follows that $G^{U_n} f_{n,\epsilon} \geq 1$ on K holds for all large enough n .

Noting that $f_{n,\epsilon} \in L_+^p(U_n)$, for large enough n , we have

$$\hat{C}_p(K, U_n) \leq \|f_{n,\epsilon}\|_{L^p(U_n)}^p \leq (1 + \epsilon) \|f\|_{L^p(U)}^p, \quad (3.13)$$

By the arbitrariness of f , we obtain from (3.13) that

$$\lim_{n \rightarrow \infty} \hat{C}_p(K, U_n) \leq (1 + \epsilon) \hat{C}_p(K, U).$$

Letting $\epsilon \rightarrow 0$ in the above and by Theorem 3.4, we complete the proof. \square

4. PROOF OF THEOREM 1.4

For the case $p = 1$, notice that (1.6) reduces to $-\Delta u \geq 0$, Theorem 1.4 can be derived by the well-established equivalent conditions of parabolicity as follows.

Theorem 4.1. [19, Theorems 1.16 and 2.12] Let (V, μ) be an infinite, connected, locally finite graph. The following statements are equivalent.

- (1) (V, μ) is parabolic.
- (2) Any nonnegative superharmonic function is constant.
- (3) For some (or, all) $x, y \in V$, $g(x, y) = \infty$.
- (4) For some (or, every) $x \in V$, $\text{Cap}(\{x\}) = 0$. Here $\text{Cap}(\{x\}) := \text{Cap}(\{x\}, V)$

Noting that the three conditions in Theorem 1.4 are always valid when (V, μ) is parabolic, so without loss of generality we always assume (V, μ) is non-parabolic, which means there exists a finite nonnegative green function corresponding to Δ on graph.

The next theorem is devoted to the general case $p > 1$.

Theorem 4.2. For $1 < p < \infty$, let (V, μ) be an infinite, connected, locally finite graph. Then the following conditions are equivalent.

- (a) (V, μ) is L^p -parabolic.
- (b) (V, μ) admits L^q -Liouville property.
- (c) Any nonnegative solution to (1.6) is harmonic.
- (d) For some (or, all) $x, y \in V$, $g_q(x, y) = \infty$.

Here p, q are Hölder conjugate exponents, and $g_q(x, y)$ is defined as in (2.1).

Proof. We complete the proof by using contradiction argument.

(a) \Rightarrow (d). Assume that (d) is not valid, we know from Lemma 2.3, there exists some x_0 such that

$$g_q(x_0, x_0) < \infty.$$

For any finite set $K \subset V$ with $x_0 \in K$, define

$$h(x) = g(x, x_0).$$

Since $-\Delta h(x) = \frac{\delta_{x_0}(x)}{\mu(x_0)}$, for any function $v \in \ell_0(V)$ such that $v(x_0) \geq 1$, we have

$$\begin{aligned} 1 &\leq \sum_{x \in V} (-\Delta h(x))v(x)\mu(x) = \sum_{x \in V} h(x)(-\Delta v(x))\mu(x) \\ &\leq \left(\sum_{x \in V} h(x)^q \mu(x) \right)^{\frac{1}{q}} \left(\sum_{x \in V} |\Delta v(x)|^p \mu(x) \right)^{\frac{1}{p}}. \end{aligned}$$

Hence

$$\sum_{x \in V} |\Delta v(x)|^p \mu(x) \geq \left(\sum_{x \in V} h(x)^q \mu(x) \right)^{-\frac{p}{q}} = [g_q(x_0, x_0)]^{-\frac{p}{q}} \mu(x_0)^{-p},$$

By the definition of $C_p(\{x_0\})$ and Theorem 3.4, we obtain that

$$C_p(\{x_0\}) = \bar{C}_p(\{x_0\}) > 0,$$

which yields that (V, μ) is not L^p -parabolic, and this contradicts with (a). Thus, (d) is valid.

(d) \Rightarrow (a). Assume that (V, μ) is not L^p -parabolic, then there exists some finite set A such that $C_p(A) > 0$. By Proposition 3.6, there exists some $x_0 \in A$ such that $C_p(\{x_0\}) > 0$.

Given a finite set $U \subset V$ with $x_0 \in U$, let us set

$$g_q^U(x, x_0) := G([g^U(x, x_0)]^{q-1}) = \sum_{z \in V} g(x, z) [g^U(z, x_0)]^{q-1} \mu(z),$$

and

$$f(x) := \frac{[g^U(x, x_0)]^{q-1}}{g_q^U(x_0, x_0)}.$$

It follows that $g_q^U(x, x_0) < \infty$, $f \in \ell_0(V)$, and $Gf(x_0) = 1$. Moreover, note that

$$\begin{aligned} 0 < C_p(\{x_0\}) &= \hat{C}_p(\{x_0\}) \leq \|f(x)\|_{L^p(V)}^p \\ &= \frac{\sum_{x \in V} [g^U(x, x_0)]^q \mu(x)}{[g_q^U(x_0, x_0)]^p} \end{aligned}$$

Since $\sum_{x \in V} [g^U(x, x_0)]^q \mu(x) \leq g_q^U(x_0, x_0)$, then

$$g_q^U(x_0, x_0) \leq \left(\frac{1}{C_p(\{x_0\})} \right)^{\frac{1}{p-1}}.$$

Similarly, letting $\{U_n\}$ be an exhaustion sequence of V contain $\{x_0\}$, we arrive

$$g_q^{U_n}(x_0, x_0) \leq \left(\frac{1}{C_p(\{x_0\})} \right)^{\frac{1}{p-1}}.$$

Letting $n \rightarrow \infty$ and using Monotone Convergence theorem, we obtain

$$g_q(x_0, x_0) \leq \left(\frac{1}{C_p(\{x_0\})} \right)^{\frac{1}{p-1}}.$$

which contradicts with (d) by Lemma 2.3, hence (a) holds.

(b) \Rightarrow (d). Assume that (d) is not valid, then there exists some x_0 such that $g_q(x_0, x_0) < \infty$. Noting that

$$\|g(x, x_0)\|_{L^q(V)}^q = g_q(x_0, x_0) < \infty,$$

and $g(x, x_0)$ is a non-trivial positive superharmonic function, which contradicts with (b). Thus, we obtain (d).

(d) \Rightarrow (b). Assume that (b) fails, then there exists $f \in L^q(V)$ which is a non-trivial positive superharmonic function, hence there exists some $x_0 \in V$ such that $-\Delta f(x_0) > 0$. Let us choose $\lambda > 0$ such that $\lambda f(x_0) \geq g(x_0, x_0)$.

Let $\{U_n\}$ be an exhaustion sequence of V with $x_0 \in U_n$. By Maximum principle (cf. [10, Lemma 1.39]), we have

$$g^{U_n}(x, x_0) \leq \lambda f(x), \quad \text{for all } x \in U_n.$$

It follows by letting $n \rightarrow \infty$ that

$$g(x, x_0) \leq \lambda f(x).$$

Noting $\sum_{x \in V} f(x)^q \mu(x) < \infty$, we obtain

$$g_q(x_0, x_0) = \sum_{x \in V} g(x, x_0)^q \mu(x) < \infty,$$

which contradicts with (d). Thus, it shows that (b) is true.

(c) \Rightarrow (d). Assume (d) fails, then fix $x_0 \in V$, we know $g_q(x, x_0) < \infty$, by Lemma 2.4, a direct calculation shows that $g_q(x, x_0)$ is a positive solution to (1.6), but $g_q(x, x_0)$ is not harmonic, which contradicts with (c).

(d) \Rightarrow (c). Assume (c) fails, then there exists a nonnegative nonharmonic function h which is a solution to (1.6). Set

$$h_1 := -\Delta h \geq 0,$$

then

$$-\Delta h_1^{p-1} = \Delta(|\Delta h|^{p-2} \Delta h) \geq 0.$$

Since h is not harmonic and h_1^{p-1} is superharmonic, by Maximum principle, h_1 is strictly positive. Then for fixed x_0 , by Maximum principle again, we obtain that there exists some constant $C > 0$ such that

$$h_1(x)^{p-1} \geq Cg(x, x_0).$$

Noting that for any positive integer l ,

$$\sum_{n=0}^l P_n(-\Delta)h = \sum_{n=0}^l P_n(I - P)h = h - P_{l+1}h \leq h,$$

we obtain

$$\sum_{y \in V} g(x, y)h_1(y)\mu(y) = Gh_1(x) = \sum_{n=0}^{\infty} P_n(-\Delta)h(x) \leq h(x).$$

Letting $C' = C^{\frac{1}{p-1}}$, we have

$$\begin{aligned} h(x_0) &\geq \sum_{y \in V} g(x_0, y)h_1(y)\mu(y) \\ &\geq C' \sum_{y \in V} g(x_0, y)g(y, x_0)^{\frac{1}{p-1}}\mu(y) \\ &= C' \sum_{y \in V} g(y, x_0)^q \mu(y) \\ &= g_q(x_0, x_0). \end{aligned}$$

Combining with Lemma 2.3, we have $g_q(x, y) < \infty$ for any $x, y \in V$, which contradicts with (d). Thus, we complete the proof. \square

Corollary 4.3. For every $1 \leq s < t < \infty$, if (V, μ) is L^s -parabolic, then (V, μ) is L^t -parabolic.

Proof. Let s' and t' denote the Hölder conjugate exponent corresponding to s and t respectively.

For $s = 1$ and $t > 1$, the parabolicity of (V, μ) is equivalent to that any nonnegative superharmonic function on (V, μ) is constant. Hence (V, μ) also admits $L^{t'}$ -Liouville property. Then by Theorem 4.2, (V, μ) is L^t -parabolic.

Now for $1 < s < t < \infty$, without loss of generality, let us assume that (V, μ) is not parabolic. Then for fixed $x_0 \in V$, we have

$$g(x, x_0) = h(x, x_0)g(x_0, x_0) \leq g(x_0, x_0), \quad \text{for all } x \in V,$$

where

$$h(x, y) := \mathbb{P}[X_0 = x, \exists n \in \mathbb{N}, \text{ s.t. } X_n = y].$$

Hence

$$g_{s'}(x_0, x_0) = \sum_{x \in V} g(x, x_0)^{s'} \mu(x) \leq g(x_0, x_0)^{t-s'} g_{t'}(x_0, x_0),$$

which finishes the proof by Theorem 4.2. \square

5. VOLUME GROWTH CONDITION

The volume growth condition for L^1 -parabolicity (or equivalently, parabolicity) has been well studied in the existing literature, see [10, 19, 15, 16]. We introduce a sufficient volume condition for parabolicity, which is a direct consequence from the Nash-Williams' test: if for some $o \in V$, the following

$$\sum_{n=1}^{\infty} \frac{n}{\mu(B(o, n))} = \infty, \quad (5.1)$$

holds, then (V, μ) is parabolic, see [10, Theorem 6.13].

If (V, μ) is L^1 -parabolic, by Corollary 4.3, we derive that (V, μ) is also L^p -parabolic for $p \geq 1$. Thus, without loss of generality, throughout this section, we always assume that (V, μ) is nonparabolic.

Now let us move to the general case $p > 1$. First, we consider $1 < p \leq 2$.

Theorem 5.1. For $1 < p \leq 2$, assume that $\mu_0 := \inf_{x \in V} \mu(x) > 0$. If there exists $o \in V$ such that

$$V(o, r) \lesssim \frac{r^{2p}}{\log r}, \quad \text{for all large enough } r, \quad (5.2)$$

then (V, μ) is L^p -parabolic.

Proof. Letting $q = \frac{p}{p-1}$, and by using Fubini's theorem and Jensen's inequality, we have

$$\begin{aligned} g_q(o, o) &= \sum_{x \in V} g(o, x) g(x, o)^{q-1} \mu(x) \\ &= \sum_{x \in V} \sum_{n=0}^{\infty} p_n(o, x) g(x, o)^{q-1} \mu(x) \\ &\geq \sum_{n=0}^{\infty} \left(\sum_{x \in V} p_n(o, x) g(x, o) \mu(x) \right)^{q-1} \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{\infty} \sum_{x \in V} p_n(o, x) p_m(x, o) \mu(x) \right)^{q-1} \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=n}^{\infty} p_m(o, o) \right)^{q-1}. \end{aligned} \quad (5.3)$$

where we have used that $\sum_{x \in V} p_n(o, x) \mu(x) = \sum_{x \in V} P_n(o, x) = 1$.

Since $\mu_0 := \inf_{x \in V} \mu(x) > 0$, and by the diagonal heat kernel lower estimate of [13], we obtain there exists some n_0 such that

$$p_n(o, o) \geq \frac{1/4}{V(o, \sqrt{cn \log n})}, \quad \text{for all } n \geq n_0.$$

Substituting the above into (5.3), and combining with (5.2), we obtain

$$\begin{aligned} g_q(o, o) &\gtrsim \sum_{n=n_0}^{\infty} \left(\sum_{m=n}^{\infty} m^{-p} (\log m)^{1-p} \right)^{q-1}, \\ &\sim \sum_{n=n_0}^{\infty} (n \log n)^{-1} = \infty. \end{aligned}$$

Thus by Theorem 4.2, we complete the proof. Here \asymp means that the ratio of are both bounded from above and below. \square

Remark 5.2. The volume condition (5.2) in Theorem 5.1 is not sharp. In fact, if we have diagonal lower bound of heat kernel

$$p_n(o, o) \gtrsim \frac{1}{V(o, \sqrt{n})},$$

then (5.2) can be improved to

$$V(o, r) \lesssim r^{2p}(\log r)^{p-1}.$$

In the last part of section 5, we try to prove a Li-Yau type estimate of Green function via volume growth, namely, (5.14) in Theorem 5.6. Then by using the estimate of Green function, we can obtain a sharp volume condition for L^p -capacity on a class of graphs with more strict geometric property, see Theorem 5.7.

We say that (V, μ) satisfies (Δ) if it admits (P_0) and every vertex has a loop, namely

$$\begin{cases} y \sim x \Rightarrow \frac{\mu_{xy}}{\mu(x)} \geq \alpha, \\ x \sim x, \quad \text{for all } x \in V, \end{cases} \quad (\Delta)$$

Now Let us introduce the following equivalent Gaussian estimate of heat kernel which was obtained by Delmotte in [6].

Theorem 5.3. Let (V, μ) satisfy (Δ) . The following two conditions are equivalent:

- (A) (V, μ) admits conditions (VD) and (PI).
- (B) There exist $c_l, C_l, c_r, C_r > 0$, such that

$$\frac{c_l}{V(x, \sqrt{n})} e^{-\frac{C_l d(x, y)^2}{n}} \leq p_n(x, y) \leq \frac{c_r}{V(x, \sqrt{n})} e^{-\frac{C_r d(x, y)^2}{n}}, \quad (5.4)$$

holds for all $x, y \in V$ and all $n \geq d(x, y)$.

Let us emphasize that in Theorem 5.3, the condition (Δ) implies that $\mu_{xx} > 0$ for all $x \in V$. Thus Theorem 5.3 can not be directly applied to even for the simple random walk in \mathbb{Z}^d , since $\mu_{xx} = 0$ in \mathbb{Z}^d . By overcoming such inconvenience, there is a traditional technique to deal with this problem by establishing a new heat kernel $p'(x, y) = p_2(x, y)$, see [3, 6]. But it may be difficult to prove that this transformation preserves (PI).

Inspired by this technique, we find another way to drop the loop condition and establish the estimate of Green functions, see Theorem 5.6.

For any $x, y \in V$, let us define a new edge weight by

$$\hat{\mu}_{xy} = \frac{1}{2}\mu_{xy} + \frac{1}{2} \sum_{z \in V} \frac{\mu_{xz}\mu_{zy}}{\mu(z)}. \quad (5.5)$$

Under this new edge weight, we let $x \sim y$ if and only if $\hat{\mu}(xy) > 0$.

According to the definition of $\hat{\mu}$, we have

1. If $x \sim y$ on (V, μ) , then $x \sim y$ on $(V, \hat{\mu})$.
2. If $x \not\sim y$, but $x \sim z, z \sim y$ on (V, μ) , then $x \sim y$ on $(V, \hat{\mu})$.

We now give the relationship between Green functions corresponding to different weights μ and $\hat{\mu}$.

Lemma 5.4. Let $g(x, y)$ and $\hat{g}(x, y)$ be the corresponding Green functions on (V, μ) and $(V, \hat{\mu})$ respectively. Then

$$\frac{1}{2}g(x, y) \leq \hat{g}(x, y) \leq g(x, y). \quad (5.6)$$

Proof. From (5.5), we have

$$\hat{\mu}(x) = \mu(x), \quad (5.7)$$

and

$$\hat{P}(x, y) = \frac{1}{2}P(x, y) + \frac{1}{2}P_2(x, y). \quad (5.8)$$

Combining with (5.8), we have

$$\begin{aligned} \hat{P}_2(x, y) &= \frac{1}{2} \sum_{z \in V} P(x, z) \hat{P}(z, y) + \frac{1}{2} \sum_{z \in V} P_2(x, z) \hat{P}(z, y) \\ &= \frac{1}{2} \left(\frac{1}{2} \sum_{z \in V} P(x, z) P(z, y) + \frac{1}{2} \sum_{z \in V} P(x, z) P_2(z, y) \right) \\ &\quad + \frac{1}{2} \left(\frac{1}{2} \sum_{z \in V} P_2(x, z) P(z, y) + \frac{1}{2} \sum_{z \in V} P_2(x, z) P_2(z, y) \right) \\ &= \frac{1}{4} (P_2(x, y) + 2P_3(x, y) + P_4(x, y)). \end{aligned}$$

We claim that for all $n \geq 0$

$$\hat{P}_n(x, y) = \frac{1}{2^n} \sum_{m=0}^n \binom{n}{m} P_{n+m}(x, y), \quad (5.9)$$

where the binomial coefficient $\binom{n}{m} = \frac{n!}{m!(n-m)!}$.

Assume that (5.9) holds for $n \leq k$, we will show that (5.9) is also valid for $n = k + 1$. Since

$$\begin{aligned} \hat{P}_{k+1}(x, y) &= \sum_{z \in V} \hat{P}(x, z) \hat{P}_k(z, y) \\ &= \sum_{z \in V} \left(\frac{1}{2}P(x, z) + \frac{1}{2}P_2(x, z) \right) \left(\frac{1}{2^k} \sum_{m=0}^k \binom{k}{m} P_{k+m}(z, y) \right) \\ &= \frac{1}{2^{k+1}} \sum_{m=0}^k \binom{k}{m} \sum_{z \in V} P(x, z) P_{k+m}(z, y) \\ &\quad + \frac{1}{2^{k+1}} \sum_{m=0}^k \binom{k}{m} \sum_{z \in V} P_2(x, z) P_{k+m}(z, y) \\ &= \frac{1}{2^{k+1}} \sum_{m=0}^k \binom{k}{m} P_{k+1+m}(x, y) + \frac{1}{2^{k+1}} \sum_{m=0}^k \binom{k}{m} P_{k+2+m}(x, y) \\ &= \frac{1}{2^{k+1}} \sum_{m=0}^k \binom{k}{m} P_{k+1+m}(x, y) + \frac{1}{2^{k+1}} \sum_{m=1}^{k+1} \binom{k}{m-1} P_{k+1+m}(x, y), \end{aligned}$$

Combining with Pascal's formula

$$\binom{k}{m} + \binom{k}{m-1} = \binom{k+1}{m},$$

we obtain

$$\hat{P}_{k+1}(x, y) = \frac{1}{2^{k+1}} P_{k+1}(x, y) + \frac{1}{2^{k+1}} \sum_{m=1}^k \left(\binom{k}{m} + \binom{k}{m-1} \right) P_{k+1+m}(x, y)$$

$$\begin{aligned}
& + \frac{1}{2^{k+1}} P_{2k+2}(x, y) \\
& = \frac{1}{2^{k+1}} \sum_{m=0}^{k+1} \binom{k+1}{m} P_{k+1+m}(x, y).
\end{aligned} \tag{5.10}$$

Thus the claim (5.9) is valid.

For any $i, k \in \mathbb{N}$, define

$$\begin{cases} a_k = 2^{2k} c_{2k} = \sum_{m=0}^k 2^m \binom{2k-m}{m}, \\ \bar{a}_k = 2^{2k+1} c_{2k+1} = \sum_{m=0}^k 2^m \binom{2k+1-m}{m}, \end{cases}$$

and

$$\begin{cases} b_{k+2} = a_{k+2} - a_{k+1}, \\ \bar{b}_{k+2} = \bar{a}_{k+2} - \bar{a}_{k+1}, \end{cases}$$

and

$$c_i = \sum_{(m,n) \in A_i} \frac{1}{2^n} \binom{n}{m},$$

where

$$A_i = \{(m, n) | m + n = i, 0 \leq m \leq n\}.$$

When $k \geq 2$, using Pascal's formula, we obtain

$$\begin{aligned}
b_k & = \sum_{m=0}^k 2^m \binom{2k-m}{m} - \sum_{m=0}^{k-1} 2^m \binom{2k-2-m}{m} \\
& = \sum_{m=0}^{k-1} 2^m \binom{2k-2-m}{m} + \sum_{m=1}^{k-1} 2^{m+1} \binom{2k-2-m}{m-1} \\
& \quad + \sum_{m=2}^k 2^m \binom{2k-2-m}{m-2} - \sum_{m=0}^{k-1} 2^m \binom{2k-2-m}{m} \\
& = \sum_{m=1}^{k-1} 2^{m+1} \binom{2k-2-m}{m-1} + \sum_{m=2}^k 2^m \binom{2k-2-m}{m-2} \\
& = 4 \left(\sum_{m=0}^{k-2} 2^m \binom{2k-3-m}{m} + \sum_{m=0}^{k-2} 2^m \binom{2k-4-m}{m} \right) \\
& = 4 \left(\sum_{m=0}^{k-2} 2^m \binom{2k-3-m}{m} + \sum_{m=1}^{k-1} 2^m \binom{2k-3-m}{m-1} \right. \\
& \quad \left. - \sum_{m=0}^{k-2} 2^m \binom{2k-4-m}{m} \right) \\
& = 4 \left(\sum_{m=0}^{k-1} 2^m \binom{2k-2-m}{m} - \sum_{m=0}^{k-2} 2^m \binom{2k-4-m}{m} \right) \\
& = 4b_{k-1}.
\end{aligned}$$

Noting $b_2 = a_2 - a_1 = 8$, thus $b_k = 2^{2k-1}$. Hence, for $k \geq 2$,

$$a_k = a_1 + \sum_{m=2}^k 2^{2m-1} = 3 + \sum_{m=2}^k 2^{2m-1}.$$

By the similar arguments, we derive that

$$\bar{b}_k = 2^{2k},$$

and

$$\bar{a}_k = 5 + \sum_{m=2}^k 2^{2m}.$$

For $k \geq 2$, it follows that

$$c_{2k} = \frac{1}{2^{2k}} \left(3 + \sum_{m=2}^k 2^{2m-1} \right),$$

and

$$c_{2k+1} = \frac{1}{2^{2k+1}} \left(5 + \sum_{m=2}^k 2^{2m} \right).$$

Noting $c_0 = c_1 = 1$, $c_2 = \frac{3}{4}$ and $c_3 = \frac{5}{8}$, and by induction method, we obtain for any $i \in \mathbb{N}$,

$$\frac{1}{2} \leq c_i \leq 1. \quad (5.11)$$

Summing up $\hat{P}_n(x, y)$ for n from 0 to l , we obtain

$$\begin{aligned} \sum_{n=0}^l \hat{P}_n(x, y) &= \sum_{n=0}^l \frac{1}{2^n} \left(\sum_{m=0}^n \binom{n}{m} P_{n+m}(x, y) \right) \\ &= \sum_{i=0}^{2l} \left(\sum_{(m,n) \in B_{i,l}} \frac{1}{2^n} \binom{n}{m} \right) P_i(x, y) \\ &= \sum_{i=0}^l \left(\sum_{(m,n) \in A_i} \frac{1}{2^n} \binom{n}{m} \right) P_i(x, y) \\ &\quad + \sum_{i=l+1}^{2l} \left(\sum_{(m,n) \in B_{i,l}} \frac{1}{2^n} \binom{n}{m} \right) P_i(x, y), \end{aligned} \quad (5.12)$$

where

$$B_{i,l} = \{(m, n) | m + n = i, 0 \leq m \leq n \leq l\},$$

thus, $B_{i,l} \subset A_i$, and in particular, $B_{i,l} = A_i$ for $i \leq l$.

Combining (5.12) with the definition of c_i , we obtain

$$\sum_{n=0}^l c_n P_n(x, y) \leq \sum_{n=0}^l \hat{P}_n(x, y) \leq \sum_{n=0}^{2l} c_n P_n(x, y). \quad (5.13)$$

It follows from (5.11) that

$$\frac{1}{2} \sum_{n=0}^l P_n(x, y) \leq \sum_{n=0}^l \hat{P}_n(x, y) \leq \sum_{n=0}^{2l} P_n(x, y).$$

By letting $l \rightarrow \infty$, and combining with (5.7), we obtain (5.6). \square

Assume condition (P_0) (resp. (VD), (PI)) is satisfied on (V, μ) , then the same condition is also satisfied on $(V, \hat{\mu})$, see the following proposition.

Proposition 5.5. The statements are as follows.

- (1). The condition (P_0) on (V, μ) implies (Δ) on $(V, \hat{\mu})$.
- (2). The volum doubling property (VD) on (V, μ) implies that on $(V, \hat{\mu})$.
- (3). The Poincaré inequality (PI) on (V, μ) implies that on $(V, \hat{\mu})$.

Proof. (1). For any $x \in V$, let us note that

$$\{y : y \sim x \text{ on } (V, \hat{\mu})\} = S(x, 1) \cup S(x, 2) \cup \{x\},$$

where $S(x, n) = \{y : d(x, y) = n\}$ for $n = 1, 2$.

Then from the condition (P_0) on (V, μ) , we have

$$\frac{\hat{\mu}_{xy}}{\hat{\mu}(x)} \geq \frac{1}{2} \frac{\mu_{xy}}{\mu(x)} \geq \frac{1}{2} \alpha, \quad \text{for } y \in S(x, 1),$$

and

$$\frac{\hat{\mu}_{xy}}{\hat{\mu}(x)} \geq \frac{1}{2} \sum_{z \in V} \frac{\mu_{xz} \mu_{zy}}{\mu(x) \mu(z)} \geq \frac{1}{2} \alpha^2, \quad \text{for } y \in S(x, 2) \cup \{x\}.$$

Hence $(V, \hat{\mu})$ satisfies the condition (Δ) .

(2). Let us note that a path of length n on $(V, \hat{\mu})$ corresponds a path on (V, μ) whose length is less than $2n$, on the other hand, a path of length $2n$ or $2n-1$ on (V, μ) corresponds a path of length n on $(V, \hat{\mu})$. Hence for all $x \in V$ and $n \in \mathbb{N}$, we have

$$\hat{B}(x, n) = B(x, 2n),$$

which completes the proof of (2).

(3). By the Poincaré inequality (PI) on (V, μ) , we obtain that for all $r > 0$, $x_0 \in V$ and $f \in \ell(V)$,

$$\begin{aligned} \sum_{x \in \hat{B}(x_0, r)} |f(x) - f_{\hat{B}}|^2 \hat{\mu}(x) &\leq \sum_{x \in \hat{B}(x_0, [r])} |f(x) - f_{\hat{B}}|^2 \hat{\mu}(x) \\ &\leq \sum_{x \in B(x_0, 2[r])} |f(x) - f_B|^2 \mu(x) \\ &\leq 4C_2 [r]^2 \sum_{x, y \in B(x_0, 4[r])} \mu_{xy} (f(y) - f(x))^2 \\ &\leq 8C_2 [r]^2 \sum_{x, y \in \hat{B}(x_0, 2[r])} \hat{\mu}_{xy} (f(y) - f(x))^2 \\ &\leq 8C_2 r^2 \sum_{x, y \in \hat{B}(x_0, 2r)} \hat{\mu}_{xy} (f(y) - f(x))^2, \end{aligned}$$

where

$$f_B = \frac{1}{V(x_0, 2[r])} \sum_{x \in B(x_0, 2[r])} f(x) \mu(x) = \frac{1}{\hat{V}(x_0, [r])} \sum_{x \in \hat{B}(x_0, [r])} f(x) \hat{\mu}(x) = f_{\hat{B}}.$$

where $[r]$ is greatest integer function (or floor function). Thus, we complete the proof. \square

Now we are ready to prove the following estimate of Green function.

Theorem 5.6. Assume conditions (VD), (PI) and (P_0) are satisfied on (V, μ) , then there exist constants $C, C' > 0$ such that

$$C \sum_{n=d(x,y)}^{\infty} \frac{n}{V(x,n)} \leq g(x,y) \leq C' \sum_{n=d(x,y)}^{\infty} \frac{n}{V(x,n)}. \quad (5.14)$$

Proof. Let us denote $d := \hat{d}(x, y)$. and note that $\hat{p}_n(x, y) = 0$ when $n < d$. Since conditions (VD), (PI) and (P_0) are satisfied on (V, μ) , by Proposition 5.5, we know conditions (VD), (PI) and (Δ) hold on $(V, \hat{\mu})$. Then combining with Theorem 5.3, we obtain

$$\sum_{n=d}^{\infty} \frac{c_l}{\hat{V}(x, \sqrt{n})} e^{-\frac{c_l d^2}{n}} \leq \hat{g}(x, y) \leq \sum_{n=d}^{\infty} \frac{c_r}{\hat{V}(x, \sqrt{n})} e^{-\frac{c_r d^2}{n}} \quad (5.15)$$

Indeed, we have

$$\begin{aligned} \sum_{n=d^2}^{\infty} \frac{c_l}{\hat{V}(x, \sqrt{n})} e^{-\frac{c_l d^2}{n}} &\geq \frac{c_l}{e^{C_l}} \sum_{i=0}^{\infty} \sum_{n=(d+i)^2+1}^{(d+i+1)^2} \frac{1}{\hat{V}(x, \sqrt{n})} \\ &\geq \frac{c_l}{e^{C_l}} \sum_{i=0}^{\infty} \frac{2(d+i)+1}{\hat{V}(x, d+i+1)} \\ &\geq \frac{c_l}{e^{C_l}} \sum_{i=0}^{\infty} \frac{d+i+1}{\hat{V}(x, d+i+1)} \\ &= \frac{c_l}{e^{C_l}} \sum_{n=d+1}^{\infty} \frac{n}{\hat{V}(x, n)}. \end{aligned} \quad (5.16)$$

Obviously, (VD) implies that

$$\frac{d}{\hat{V}(x, d)} \leq \frac{C_1}{2} \frac{2d}{\hat{V}(x, 2d)}.$$

Combining the above with (5.15) and (5.16), letting $C = \frac{2c_l}{e^{C_l}(2+C_1)}$, and by Lemma 5.4, we derive the lower bound of (5.14) by

$$\begin{aligned} g(x, y) &\geq \hat{g}(x, y) \geq \frac{2c_l}{e^{C_l}} \sum_{n=d+1}^{\infty} \frac{n}{\hat{V}(x, n)} \\ &\geq \frac{2c_l}{e^{C_l}(2+C_1)} \sum_{n=d+1}^{\infty} \frac{n}{\hat{V}(x, n)} + \frac{c_l C_1}{e^{C_l}(2+C_1)} \frac{2d}{\hat{V}(x, 2d)} \\ &\geq \frac{2c_l}{e^{C_l}(2+C_1)} \sum_{n=d}^{\infty} \frac{n}{\hat{V}(x, n)} \\ &\geq \frac{2c_l C_1}{e^{C_l}(2+C_1)} \sum_{n=d}^{\infty} \frac{n}{V(x, n)}. \end{aligned}$$

Similarly, for the upper bound of (5.14), we have

$$\begin{aligned} \sum_{n=d^2}^{\infty} \hat{p}_n(x, y) &\leq \sum_{n=d^2}^{\infty} \frac{c_r}{\hat{V}(x, \sqrt{n})} e^{-\frac{c_r d^2}{n}} \leq c_r \sum_{i=0}^{\infty} \sum_{n=(d+i)^2}^{(d+i)^2-1} \frac{1}{\hat{V}(x, \sqrt{n})} \\ &\leq c_r \sum_{i=0}^{\infty} \frac{2(d+i)+1}{\hat{V}(x, d+i)} \\ &\leq 2c_r \sum_{n=d+1}^{\infty} \frac{n}{\hat{V}(x, n)}. \end{aligned}$$

Then it suffices to prove

$$\sum_{n=d}^{d^2} \hat{p}_n(x, y) \leq (C' - 2c_r) \sum_{n=d}^{\infty} \frac{n}{\hat{V}(x, n)}. \quad (5.17)$$

Indeed by (5.6) and (VD), we have

$$g(x, y) \leq 2\hat{g}(x, y) \leq C' \sum_{n=d}^{\infty} \frac{n}{\hat{V}(x, n)} \leq C' \sum_{n=d}^{\infty} \frac{n}{V(x, n)}.$$

To prove (5.17), for any $d \geq 1$, let us define

$$t = \min\{m : 2^m \geq d, m \in \mathbb{N}\},$$

and for $0 \leq l \leq t$, define

$$s_l = \lfloor \frac{d^2}{2^l} \rfloor.$$

Combining with (5.4), we obtain

$$\begin{aligned} \sum_{n=d}^{d^2} \hat{p}_n(x, y) &\leq \sum_{n=d}^{d^2} \frac{c_r}{\hat{V}(x, \sqrt{n})} e^{-\frac{c_r d^2}{n}} \\ &\leq c_r \sum_{l=0}^t \sum_{n=s_{l+1}}^{s_{l+1}+1} \frac{1}{\hat{V}(x, \sqrt{n})} e^{-\frac{c_r d^2}{n}} \\ &\leq c_r \sum_{l=0}^t \left(\frac{d^2}{2^{l+1}} + 1 \right) e^{-\frac{c_r d^2}{s_l}} \frac{1}{\hat{V}(x, \sqrt{s_{l+1}})} \\ &\leq c_r \sum_{l=0}^t \left(\frac{d^2}{2^{l+1}} + 1 \right) e^{-C_r 2^l} \frac{1}{\hat{V}(x, \sqrt{\frac{d^2}{2^{l+1}}})}. \end{aligned} \quad (5.18)$$

And using (VD), we have

$$\frac{1}{\hat{V}(x, \sqrt{n})} \leq C_1^l \frac{1}{\hat{V}(x, 2^l \sqrt{n})}. \quad (5.19)$$

Hence

$$\begin{aligned}
\sum_{n=d}^{d^2} \hat{p}_n(x, y) &\leq c_r \sum_{l=0}^t 2^l d^2 e^{-C_r 2^l} C_1^{l+1} \frac{1}{\hat{V}(x, \sqrt{2^{l+1} d^2})} \\
&\leq c_r \sum_{l=0}^t e^{-C_r 2^l} C_1^{l+1} \sum_{n=2^l d^2+1}^{2^{l+1} d^2} \frac{1}{\hat{V}(x, \sqrt{n})} \\
&\leq c_r \sum_{l=0}^t e^{-C_r (2^l - k(l+1))} \sum_{n=2^l d^2+1}^{2^{l+1} d^2} \frac{1}{\hat{V}(x, \sqrt{n})},
\end{aligned} \tag{5.20}$$

where $k := \min\{m : e^{C_r m} \geq C_1\}$.

Noting that the function $2^x - k(x+1)$ is bounded from below by

$$c := \frac{k}{\log 2} - k \log_2 \left(\frac{k}{\log 2} \right) - k,$$

we obtain

$$\begin{aligned}
\sum_{n=d}^{d^2} \hat{p}_n(x, y) &\leq c_r e^{-C_r c} \sum_{n=d^2}^{\infty} \frac{1}{\hat{V}(x, \sqrt{n})} \\
&\leq c_r e^{-C_r c} \sum_{i=0}^{\infty} \sum_{n=(d+i)^2}^{(d+i+1)^2-1} \frac{1}{\hat{V}(x, \sqrt{n})} \\
&\leq c_r e^{-C_r c} \sum_{i=0}^{\infty} \frac{2(d+i)+1}{\hat{V}(x, d+i)} \\
&\leq 4c_r e^{-C_r c} \sum_{n=d}^{\infty} \frac{n}{\hat{V}(x, n)}.
\end{aligned}$$

By setting $C' = 4c_r e^{-C_r c} + 2c_r$, the upper bound of (5.14) follows. Hence, we complete the proof. \square

Theorem 5.7. Let $1 < p < \infty$. Assume that conditions (VD), (PI) and (P_0) are satisfied on (V, μ) , then (V, μ) is L^p -parabolic if and only if for some $o \in V$,

$$\sum_{n=0}^{\infty} \left(\sum_{m=n}^{\infty} \frac{m}{V(o, m)} \right)^q V(S(o, n)) = \infty. \tag{5.21}$$

where $S(o, n) = \{x \in V \mid d(x, o) = n\}$.

Proof. Noting that

$$g_q(o, o) = \sum_{z \in V} g(z, 0)^q \mu(z), \tag{5.22}$$

and combining with Theorem 5.6, we obtain

$$\begin{aligned}
C \sum_{n=0}^{\infty} \left(\sum_{m=n}^{\infty} \frac{m}{V(o, m)} \right)^q V(S(o, n)) &\leq g_q(o, o) \\
&\leq C' \sum_{n=0}^{\infty} \left(\sum_{m=n}^{\infty} \frac{m}{V(o, m)} \right)^q V(S(o, n)),
\end{aligned}$$

By Theorem 4.2, we obtain that (5.21) is equivalent to L^p -parabolicity. Thus, the proof is complete. \square

Theorem 5.8. Let $1 < p < \infty$. Assume that conditions (VD), (PI) and (P_0) are satisfied on (V, μ) . If for some $o \in V$,

$$\sum_{n=0}^{\infty} n \left(\sum_{m=n}^{\infty} \frac{m}{V(o, m)} \right)^{\frac{1}{p-1}} = \infty, \quad (5.23)$$

then (V, μ) is L^p -parabolic.

Proof. Noting that (V, μ) is non-parabolic, hence for fixed $y \in V$, we obtain

$$\sum_{m=0}^{\infty} \frac{m}{V(o, m)} \leq Cg(o, y) < \infty,$$

from Theorem 5.6.

Set

$$a_n = \sum_{m=n}^{\infty} \frac{m}{V(o, m)}, \quad \text{for } n \geq 0.$$

It is clear that $\{a_n\}$ is a decreasing sequence and by (VD).

Then for any positive integer l , we have

$$\begin{aligned} \sum_{n=0}^l a_n^q V(S(o, n)) &\geq \sum_{n=1}^l a_n^q (V(o, n) - V(o, n-1)) + a_0^q V(o, 0) \\ &= \sum_{n=0}^l a_n^q V(o, n) - \sum_{n=1}^l a_n^q V(o, n-1) \\ &\geq \sum_{n=0}^l (a_n^q - a_{n+1}^q) V(o, n) \\ &\geq q \sum_{n=0}^l a_{n+1}^{q-1} (a_n - a_{n+1}) V(o, n) \\ &= Cq \sum_{n=0}^l n a_{n+1}^{q-1}, \end{aligned}$$

where we have used the mean value Theorem.

By letting $l \rightarrow \infty$, we get

$$\sum_{n=0}^{\infty} \left(\sum_{m=n}^{\infty} \frac{m}{V(o, m)} \right)^q V(S(o, n)) \geq Cq \sum_{n=0}^{\infty} n \left(\sum_{m=n}^{\infty} \frac{m}{V(o, m)} \right)^{\frac{1}{p-1}}.$$

Finally, we complete the proof by Theorem 5.7. \square

Corollary 5.9. Let $1 < p < \infty$. Assume that conditions (VD), (PI) and (P_0) are satisfied on (V, μ) . If there exists some $o \in V$ such that for large enough r

$$V(o, r) \leq Cr^{2p}(\log r)^{p-1},$$

then (V, μ) is L^p -parabolic.

6. EXAMPLES

Let us now introduce a class of graphs known as Cayley graphs. Assume that G is a group and $S \subset G$ is a subset, which satisfies that if $s \in S$, then $s^{-1} \in S$. Such subset S is called symmetric.

The group G and subset S determines a graph (V, E) as follows: the set V of vertices coincides with G , and the set of edges E is defined by $x \sim y$ if and only if $x^{-1}y \in S$. The edge weight is defined by

$$\mu_{xy} = \begin{cases} \frac{1}{|S|} & \text{when } x^{-1}y \in S, \\ 0 & \text{otherwise,} \end{cases}$$

which implies $u(x) = 1$ for all $x \in G$. It is clear that if the neutral element $e \in S$, every vertex in the graph (V, E) contains a loop, otherwise, the graph contains no loops.

Moreover, since $\deg(x) = |S|$ for any $x \in V$, hence $V(x, n) = V(e, n)$ for any $x \in G$.

Definition 6.1. Let (V, μ) be an infinite graph, if there exist constant a $D > 0$ and an element $o \in V$ such that

$$V(o, r) \asymp r^D, \quad \text{for all } r > 0. \quad (6.1)$$

Then we call (V, μ) has polynomial growth.

Proposition 6.2. Let (G, μ) and (G, μ') be an infinite Cayley graph generated by finite set S and S' respectively. If (G, μ) has polynomial growth, then (G, μ') also has polynomial growth.

Proof. Assume $S = \{s_1, s_2, \dots, s_k\}$ and $S' = \{s'_1, s'_2, \dots, s'_l\}$, Let $B(e, r)$ and $B'(e, r)$ be the ball centered at e with radius r in the corresponding graph (G, S) and (G, S') respectively. Then any $z \in B'(e, n)$ can be represented in the form $z = s'_{l_1} s'_{l_2} \cdots s'_{l_t}$ where $s'_{l_i} \in S'$ and $l_t = d'(e, z) \leq n$.

For $1 \leq i \leq l$, setting $a_i = d(e, s'_i)$ and $a = \max\{a_1, \dots, a_l\}$, we have

$$d(e, z) \leq ad'(e, z),$$

which implies that

$$V'(e, r) \leq V(e, ar) \lesssim r^D.$$

By the same argument, we have $r^D \lesssim V'(e, r)$, which concludes our claim. \square

Definition 6.3. We say a finitely generated group G has polynomial growth, if its corresponding Cayley graph (G, μ) with some generating set S has polynomial growth.

Proposition 6.4. Let (G, μ) be the Cayley graph of a finitely generated group G . If (G, μ) satisfies the volume doubling property (VD), then it also satisfies the Poincaré inequality (PI).

This proposition can be found in [4] without proof, for completeness, we also provide a proof here.

Proof. For any $x_0 \in G$, any positive integer n and any $f \in \ell(G)$, letting

$$f_B = \frac{1}{V(x_0, n)} \sum_{x \in B(x_0, n)} f(x),$$

and applying Jensen's inequality, we obtain

$$\sum_{x \in B(x_0, n)} |f(x) - f_B|^2 \leq \frac{1}{V(x_0, n)} \sum_{x \in B(x_0, n)} \sum_{y \in B(x_0, n)} |f(x) - f(y)|^2$$

$$= \frac{1}{V(x_0, n)} \sum_{z \in B(e, 2n)} \sum_{x \in A_z} |f(x) - f(xz)|^2 \quad (6.2)$$

where e is the identity element of G and

$$A_z = \{x \in G : x \in B(x_0, n) \text{ and } xz \in B(x_0, n)\}.$$

Since any $z \in B(e, 2n)$ can be represented in the form $z = s_1 s_2 \cdots s_k$ where $s_i \in S$ and $k \leq 2n$, we have

$$\begin{aligned} \sum_{x \in A_z} |f(x) - f(xz)|^2 &= \sum_{x \in A_z} |f(x) - f(xs_1) + f(xs_1) - f(xs_1 s_2) \\ &\quad + \cdots + f(xs_1 \cdots s_{k-1}) - f(xs_1 \cdots s_k)|^2 \\ &\leq \sum_{x \in A_z} \sum_{0 \leq i, j \leq k} |f(xs_1 \cdots s_{i-1}) - f(xs_1 \cdots s_i)| \\ &\quad \times |f(xs_1 \cdots s_{j-1}) - f(xs_1 \cdots s_j)| \\ &\leq 2n \sum_{x \in A_z} \sum_{0 \leq i \leq k} |f(xs_1 \cdots s_{i-1}) - f(xs_1 \cdots s_i)|^2. \end{aligned} \quad (6.3)$$

For any $x \in A_z$ and any positive integers $0 \leq i \leq k$, noting that $x \in B(x_0, n)$ and $xz \in B(x_0, n)$, we have

$$xs_1 \cdots s_i \in B(x_0, 2n).$$

It follows that

$$\sum_{x \in A_z} |f(xs_1 \cdots s_{i-1}) - f(xs_1 \cdots s_i)|^2 \leq |S| \sum_{x, y \in B(x_0, 2n)} \mu_{xy} |f(x) - f(y)|^2. \quad (6.4)$$

Substituting this back into (6.3), we obtain

$$\sum_{x \in A_z} |f(x) - f(xz)|^2 \leq 4n^2 |S| \sum_{x, y \in B(x_0, 2n)} \mu_{xy} |f(x) - f(y)|^2. \quad (6.5)$$

Finally, combining (6.5) with (6.2), we conclude that

$$\sum_{x \in B(x_0, n)} |f(x) - f_B|^2 \leq 4 \frac{V(e, 2n)}{V(x_0, n)} |S| n^2 \sum_{x, y \in B(x_0, 2n)} \mu_{xy} |f(x) - f(y)|^2.$$

Since $V(x, n) = V(e, n)$ for any $x \in G$ and (G, μ) satisfies (VD), the above implies that

$$\sum_{x \in B(x_0, n)} |f(x) - f_B|^2 \leq C n^2 \sum_{x, y \in B(x_0, 2n)} \mu_{xy} |f(x) - f(y)|^2, \quad (6.6)$$

which yields the Poincaré inequality (PI). \square

Corollary 6.5. If a finitely generated group G has polynomial growth (6.1), then its Cayley graph (G, μ) satisfies the volume doubling property (VD) and the Poincaré inequality (PI). Moreover, (G, μ) is L^p -parabolic for $p \geq \frac{D}{2}$.

Remark 6.6. Let us emphasize that the finitely generated groups with polynomial growth form a large and well-studied class of groups. In fact, from Gromov's famous work [11], a finitely generated group G has polynomial growth if and only if it is virtually nilpotent, which means it contains a nilpotent subgroup of finite index.

Proof. The volume doubling condition (VD) follows directly from the polynomial growth condition (6.1). Then, applying Proposition 6.4 and Corollary 5.9, the proof is complete. \square

Example 6.7. \mathbb{Z}^d is L^p -parabolic for $p \geq \frac{d}{2}$, and is not L^p -parabolic for $p < \frac{d}{2}$. This is because that $S(e, n) \asymp n^{d-1}$ and $V(e, n) \asymp n^d$, we derive this example using Theorem 5.7.

Example 6.8. The discrete Heisenberg group

$$\left\{ \left(\begin{array}{ccc} 1 & b & c \\ 0 & 1 & a \\ 0 & 0 & 1 \end{array} \right) \mid a, b, c \in \mathbb{Z} \right\},$$

is L^p -parabolic for $p \geq 2$. This follows from the well-known fact that $V(e, n) \asymp n^4$ on discrete Heisenberg group and using Corollary 6.5.

Acknowledgements. The authors would like to thank Prof. Grigor'yan from University of Bielefeld for many valuable suggestion and discussion.

REFERENCES

- [1] D. R. Adams, L. I. Hedberg, Function Spaces and Potential Theory. Grundlehren der mathematischen Wissenschaften **314** (1996). 2nd edition, Springer-Verlag Berlin Heidelberg 1999, p.366.
- [2] M. T. Barlow, Random walks and heat kernels on graphs, LMS Lecture Note Series, 2017.
- [3] T. Coulhon, A. Grigoryan, Random walks on graphs with regular volume growth, Geom. Funct. Anal. **8**(1998), 656-701.
- [4] T. Coulhon, L. Saloff-Coste, Isopérimétrie pour les groupes et les variétés, Rev. Mat. Iberoam. **9**(1993), 293-314.
- [5] T. Coulhon, I. Holopainen, L. Saloff-Coste, Harnack inequality and hyperbolicity for subelliptic p-Laplacians with applications to Picard type theorems, Geom. Funct. Anal. **11**(2001), 1139-1191.
- [6] T. Delmotte, Parabolic Harnack inequality and estimates of Markov chains on graphs, Revista Matemática Iberoamericana **15**(1999), 181-232.
- [7] S. Faraji, A. Grigor'yan, On biparabolicity of Riemannian manifolds, Rev. Mat. Iberoam. **35**(2019), 2025-2034.
- [8] A. Grigor'yan, L. F. Pessoa, A.G. Setti, L^p -parabolicity of Riemannian manifolds, (2024), preprint.
- [9] A. Grigor'yan, On the existence of positive fundamental solution of the Laplace equation on Riemannian manifolds, Matem. Sb. 128 (1985) 354-363. English transl. Math. USSR Sb. **56** (1987), 349-358.
- [10] A. Grigor'yan, Introduction to analysis on graphs, AMS University Lecture Series, 2018.
- [11] M. Gromov, Groups of polynomial growth and expanding maps, Publ. Math. IHES **53** (1981), 53-73.
- [12] I. Holopainen, Nonlinear potential theory and quasiregular mappings on Riemannian manifolds, Ann. Acad. Sc. Fenn. Ser. A I Math. Diss. **74**(1990), 1-45.
- [13] F. Lust-Piquard, Lower bounds on $\|K^n\|_{1 \rightarrow \infty}$ for some contractions K of $L^2(\mu)$, with some applications to Markov operators, Math. Ann. **303** (1995), 699-712.
- [14] W. Woess, Random walks on infinite graphs and groups, Cambridge U. Process, Cambridge, 2000.
- [15] L. Saloff-Coste, Inequalities for p-superharmonic functions on networks. Rend. Sem. Mat. Fis. Milano, **65**(1995), 139-158.
- [16] L. Saloff-Coste, Some inequalities for superharmonic functions on graphs, Potential Anal. **6**(1997), no.2, 163-181.
- [17] P. Soardi, Parabolic networks and polynomial growth, Colloq. Math. **LX/LXI** (1990), 65-70.
- [18] M. Troyanov, Parabolicity of manifolds, Siberian Adv. Math. **9** (1999), 125-150.
- [19] W. Woess, Random walks on infinite graphs and groups, Cambridge U. Process, Cambridge, 2000.
- [20] M. Yamasaki, Parabolic and hyperbolic infinite networks, Hiroshima Math. J. **7**(1977), 135-146.

UNIVERSITÄT BIELEFELD, FAKULTÄT FÜR MATHEMATIK, POSTFACH 100131, D-33501, BIELEFELD, GERMANY

Email address: lhao@math.uni-bielefeld.de

SCHOOL OF MATHEMATICAL SCIENCES AND LPMC, NANKAI UNIVERSITY, 300071 TIANJIN, P. R. CHINA

Email address: sunyuhua@nankai.edu.cn