ON SOME SOBOLEV AND PÓLYA-SEZGÖ TYPE INEQUALITIES WITH WEIGHTS AND APPLICATIONS

TRUNG HIEU GIANG^{1,2}, NGUYEN MINH TRI², AND DANG ANH TUAN³

ABSTRACT. We are motivated by studying a boundary-value problem for a class of semilinear degenerate elliptic equations

$$\begin{cases} -\Delta_x u - |x|^{2\alpha} \frac{\partial^2 u}{\partial y^2} = f(x, y, u), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$
(P)

where $x = (x_1, x_2) \in \mathbb{R}^2$, Ω is a bounded smooth domain in \mathbb{R}^3 , $(0, 0, 0) \in \Omega$, and $\alpha > 0$.

In this paper, we will study this problem by establishing embedding theorems for weighted Sobolev spaces. To this end, we need a new Pólya-Szegö type inequality, which can be obtained by studying an isoperimetric problem for the corresponding weighted area. Our results then extend the existing ones in [1, 3] to the three-dimensional context.

1. INTRODUCTION

In [1], Nga, Tri, and Tuan established a Pólya–Szegö type inequality for a weighted gradient of a function on \mathbb{R}^2 with respect to a weighted area. Then, they applied this inequality to prove embedding theorems for weighted Sobolev spaces. Their results are motivated by studying the following equation

$$\begin{cases} -\frac{\partial^2 u}{\partial x^2} - |x|^{2\alpha} \frac{\partial^2 u}{\partial y^2} = f(x, y, u), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$
(1.1)

where Ω is a bounded smooth domain in \mathbb{R}^2 , $(0,0) \in \Omega$, and $\alpha > 0$. The existence results for this equation are then given by Luyen et al. (see [3]) based on the aforementioned weighted Sobolev inequalities.

In this paper, we aim to extend the results in [1, 3] to the three-dimensional context. More specifically, we will establish the existence and nonexistence results for the equations (P). The nonexistence result will be derived from a Pohozaev-type identity, while the existence result will be studied via the associated weighted Sobolev inequalities.

The organization of this paper is as follows. In Section 2 we first state and prove our isoperimetric inequality with weights and then use it to prove our new Pólya-Szegö type inequality. This result will be utilized to establish some Sobolev inequalities with weights in Section 3. Here, we would like to emphasize that our method, base on [1], are only able to give lower estimates for the best constants

²⁰²⁰ Mathematics Subject Classification. Primary: 28A20, 46E30, 46E35; Secondary: 51M16, 35J70.

Key words and phrases. Isoperimetric Inequality, Best Sobolev Constant, Pólya–Szegö Inequality, Rearrangement, Degenerate Elliptic Equations.

appearing in the weighted Sobolev inequalities, and calculating these constants still remains an open question. Finally, in Section 4 we will apply the above results and the Pohozaev-type identity to study the existence and nonexistence results for the equations (\mathbf{P}) .

2. A Pólya-Szegö type inequality

In this section, we will state and prove a new Pólya-Szegö type inequality mentioned in Section 1. To this end, we first introduce some notations.

Throughout this paper, let *alpha* be a fixed given positive real number and $n(\alpha)$ be the smallest positive integer such that $n(\alpha) \ge \alpha + 1$. We denote

$$\mathbb{R}^3_{s_j} := \left\{ (x_1, x_2, y) = (r \cos \theta, r \sin \theta, y) \in \mathbb{R}^3, \ r > 0, \ \theta \in \left(\frac{(j-1)\pi}{n(\alpha)}, \frac{j\pi}{n(\alpha)}\right) \right\},\$$

for each $j = 1, ..., 2n(\alpha)$.

Let $E \subset \mathbb{R}^3$ be a bounded open set with Lipschitz boundary ∂E . The notation denotes $\nu = (\nu_1, \nu_2, \nu_3)$ the outward unit normal to ∂E . We also denote

$$\partial_{s_j} E := \partial E \cap \mathbb{R}^3_{s_j},$$

for each $j = 1, ..., 2n(\alpha)$, and

$$|x| := \sqrt{x_1^2 + x_2^2}$$
, for each $(x_1, x_2) \in \mathbb{R}^2$.

We have the following definitions.

Definition 1. The $(2, \alpha)$ -volume of E is defined by

$$|E|_{2,\alpha} = \int\limits_E |x|^{2\alpha} dx_1 dx_2 dy.$$

Definition 2. The $(2, \alpha)$ -area of E is defined by

$$P_{2,\alpha}(E) = \int_{\partial E} |x|^{\alpha} \sqrt{\nu_1^2 + \nu_2^2 + |x|^{2\alpha} \nu_3^2} d\mathcal{H}^2,$$

where \mathcal{H}^2 denotes the two-dimensional Hausdorff measure of a surface in \mathbb{R}^3 .

Definition 3. For each $j = 1, ..., 2n(\alpha)$, the $(2, \alpha, j)$ -area of E is defined by

$$P_{2,\alpha,j}(E) = \int_{\partial_{s_j} E} |x|^{\alpha} \sqrt{\nu_1^2 + \nu_2^2 + |x|^{2\alpha} \nu_3^2} d\mathcal{H}^2.$$

Notice that

$$P_{2,\alpha}(E_{\lambda}) = \int_{\partial E_{\lambda}} |X|^{\alpha} \sqrt{\mu_{1}^{2} + \mu_{2}^{2} + |X|^{2\alpha} \mu_{3}^{2}} d\mathcal{H}^{2} = \lambda^{2\alpha+2} P_{2,\alpha}(E),$$

$$P_{2,\alpha,j}(E_{\lambda}) = \int_{\partial_{s_{j}} E_{\lambda}} |X|^{\alpha} \sqrt{\mu_{1}^{2} + \mu_{2}^{2} + |X|^{2\alpha} \mu_{3}^{2}} d\mathcal{H}^{2} = \lambda^{2\alpha+2} P_{2,\alpha,j}(E),$$

and

$$|E_{\lambda}|_{2,\alpha} = \int\limits_{E_{\lambda}} |X|^{2\alpha} dX_1 dX_2 dY = \lambda^{3\alpha+3} |E|_{2,\alpha}$$

where $E_{\lambda} = \{(\lambda x_1, \lambda x_2, \lambda^{\alpha+1}y) : (x_1, x_2, y) \in E\}, \mu = (\mu_1, \mu_2, \mu_3)$ is the outward unit normal to ∂E_{λ} , the fractions

$$\frac{P_{2,\alpha}(E)^{3/2}}{|E|_{2,\alpha}}$$
 and $\frac{P_{2,\alpha,j}(E)^{3/2}}{|E|_{2,\alpha}}$

for each $j = 1, ..., n(\alpha)$, are invariant under the scaling $(x_1, x_2, y) \mapsto (\lambda x_1, \lambda x_2, \lambda^{\alpha+1} y)$. Our weighted isoperimetric inequality is stated as follows.

Theorem 1 (The weighted isoperimetric inequality). Let $E \subset \mathbb{R}^3$ be a nonempty bounded open set with Lipschitz boundary ∂E such that $P_{2,\alpha}(E)$ is finite. Then the following inequality holds for every $j = 1, ..., 2n(\alpha)$

$$\frac{P_{2,\alpha,j}(B_{s_j})^{3/2}}{|B_{s_j}|_{2,\alpha}} \le \frac{P_{2,\alpha}(E)^{3/2}}{|E|_{2,\alpha}},$$

$$where \ B_{s_j} := \left\{ (x_1, x_2, y) \in \mathbb{R}^3_{s_j}, \ \frac{|x|^{2\alpha+2}}{(\alpha+1)^2} + y^2 < 1 \right\}.$$
(2.1)

Proof. For readers' convenience, we will divide the proof into two steps. **Step 1:** Assuming that there exists an integer j such that $1 \leq j \leq 2n(\alpha)$ and $E \subset \mathbb{R}^3_{s_j}$ with Lipschitz boundary. We will show that

$$\frac{P_{2,\alpha,j}(B_{s_j})^{3/2}}{|B_{s_j}|_{2,\alpha}} \le \frac{P_{2,\alpha,j}(E)^{3/2}}{|E|_{2,\alpha}}.$$
(2.2)

Without loss of generality, we can assume that j = 1. We define $\Phi_1 : \mathbb{R}_+ \times \left(0, \frac{\pi}{n(\alpha)}\right) \times \mathbb{R} \to \mathbb{R}^3_{s_1}$ by $\Phi_1(r, \theta, y) = (r \cos \theta, r \sin \theta, y).$

It is easy to see that
$$\Phi_1$$
 is a homeomorphism. Next, we define $\Phi_2 : \mathbb{R}_+ \times \left(0, \frac{\pi}{n(\alpha)}\right) \times \mathbb{R} \to \mathbb{R}^3_{\tilde{s}_1}$ by

$$\Phi_2(r,\theta,\eta) = \left(\frac{r^{\alpha+1}\cos(\alpha+1)\theta}{\alpha+1}, \frac{r^{\alpha+1}\sin(\alpha+1)\theta}{\alpha+1}, \eta\right),\,$$

where

$$\mathbb{R}^3_{\widetilde{s}_1} := \left\{ (\xi_1, \xi_2, \eta) = (\rho \cos \varphi, \rho \sin \varphi, y) \in \mathbb{R}^3, \ \rho > 0, \ \varphi \in \left(0, \frac{(\alpha + 1)\pi}{n(\alpha)}\right) \right\}.$$

It is also easy to verify that Φ_2 is a homeomorphism. Hence

$$\Psi = \Phi_1^{-1} \circ \Phi_2$$

is a homeomorphism.

Next, let $\widetilde{E} = \Psi(E)$ and $\widetilde{B}_{s_1} = \Psi(B_{s_1})$. We deduce that

$$\widetilde{B}_{s_1} = \left\{ (\xi_1, \xi_2, \eta) \in \mathbb{R}^3_{\widetilde{s}_1}, \ \xi_1^2 + \xi_2^2 + \eta^2 < 1 \right\}.$$

By some straightforward calculations, we obtain

$$|E|_{2,\alpha} = \int_{\widetilde{E}} d\xi_1 d\xi_2 d\eta = |\widetilde{E}|, \quad |B_{s_1}|_{2,\alpha} = \int_{\widetilde{B}_{s_1}} d\xi_1 d\xi_2 d\eta = |\widetilde{B}_{s_1}|,$$

$$P_{2,\alpha,1}(E) = \int_{\partial_{\tilde{s}_1}\tilde{E}} d\mathcal{H}^2 =: P_{s_1}(\tilde{E}), \quad P_{2,\alpha,1}(B_{s_1}) = \int_{\partial_{\tilde{s}_1}\tilde{B}_{s_1}} d\mathcal{H}^2 =: P_{s_1}(\tilde{B}_{s_1}).$$

where

 $\partial_{\widetilde{s}_1}\widetilde{E} = \partial \widetilde{E} \cap \mathbb{R}^3_{\widetilde{s}_1}, \quad \partial_{\widetilde{s}_1}\widetilde{B}_{s_1} = \partial \widetilde{B}_{s_1} \cap \mathbb{R}^3_{\widetilde{s}_1}.$

Now, by applying [4, Theorem 1.1], we deduce that

$$\frac{P_{s_1}(\tilde{E})^{3/2}}{|\tilde{E}|} \ge \frac{P_{s_1}(\tilde{B}_{s_1})^{3/2}}{|\tilde{B}_{s_1}|}$$

and thus (2.2) follows.

Step 2: For the open, bounded set $E \subset \mathbb{R}^3$ with Lipschitz boundary, we set

$$E = \bigcup_{i=1}^{2n(\alpha)} E_{s_i}$$

where $E_{s_i} := E \cap \overline{\mathbb{R}}_{s_i}^3$. It is easy to see that

$$\bigcup_{i=1}^{2n(\alpha)} \partial_{s_i} E \subset \partial E.$$
(2.3)

Notice that the sets E_{s_i} are respectively open, bounded subsets in $\overline{\mathbb{R}}_{s_i}^3$ with Lipschitz boundaries. Hence we have the inequality (2.2) for each E_{s_i} . Since 3/2 > 1, we have that

$$P_{2,\alpha}(E)^{3/2} \ge \sum_{j=1}^{2n(\alpha)} P_{2,\alpha,j}(E_{s_j})^{3/2}$$

By combining this and the inequality (2.2) for every E_{s_i} , our proof is complete. \Box

Next, we have the following definition.

Definition 4. Let $u \in C_0^{\infty}(\mathbb{R}^3; \mathbb{R}_+)$. The rearrangement $u^* : \overline{\mathbb{R}}_{s_1}^3 \to \mathbb{R}_+$ is defined by

$$u^*(x_1, x_2, y) = \phi((|x|^{2\alpha+2} + (\alpha+1)^2 y^2)^{1/2}) = \phi(r)$$

where $r := (|x|^{2\alpha+2} + (\alpha+1)^2 y^2))^{1/2}$, $\phi : \mathbb{R}_+ \to \mathbb{R}_+$ and

$$|\{u^*>t\}|_{2,\alpha}=|\{u>t\}|_{2,\alpha}\,,\ for\ all\ t>0.$$

Remark 1. Put $M = \max_{\mathbb{R}^3} u$. It is not difficult to see that the map

$$\lambda: t \mapsto |\{u > t\}|_{2,\alpha}$$

is nonincreasing and right-continuous. Therefore the map

$$t \mapsto |\{t < u \le M\} \cap \{\nabla u = 0\}|_{2,\alpha}$$

is nonincreasing and the function ϕ is nonincreasing, right-continuous. Moreover, the set

$$\{t \in \mathbb{R} : \exists s \in \mathbb{R}, \phi(s) = t, \phi'(s) = 0\}$$

has Lebesgue measure 0 in \mathbb{R} . As in [15] the map $h: [0, M] \to [0, \infty)$ defined by

$$h(t) = |\{t < u^* \le M\} \cap \{\nabla u^* = 0\}|_{2,\alpha}$$

is nonincreasing. Moreover h'(t) = 0 a.e. on [0, M].

We now obtain the following new Pólya–Szegö type inequality.

Theorem 2 (The Pólya-Szegö type inequality). Let $u \in C_0^{\infty}(\mathbb{R}^3; \mathbb{R}_+)$. Then

$$\int_{\mathbb{R}^3_{s_1}} |\nabla_G u^*|^2 \, dx_1 dx_2 dy \le \int_{\mathbb{R}^3} |\nabla_G u|^2 \, dx_1 dx_2 dy,$$

where $|\nabla_G u| = (u_{x_1}^2 + u_{x_2}^2 + |x|^{2\alpha} u_y^2)^{1/2}$.

We need the following lemma to prove Theorem 2.

Lemma 1. Let $u \in C_0^{\infty}(\mathbb{R}^3; \mathbb{R}_+)$. Assume that $u \neq 0$. Denote $M = \max_{\mathbb{R}^3} u$. Then

$$\int_{\{u^*=t\}} \frac{|x|^{2\alpha}}{|\nabla u^*|} d\mathcal{H}^2 \ge \int_{\{u=t\}} \frac{|x|^{2\alpha}}{|\nabla u|} d\mathcal{H}^2 \text{ for a.e. } t \in [0, M].$$
(2.4)

Proof. The proof is similar to the one of [1, Lemma 1] and for this reason, it is omitted here.

Proof of Theorem 2. We follow the process of the proof of [1, Theorem 2]. The case $u \equiv 0$ is simple. We may now assume $u \not\equiv 0$. Let $M = \max_{\mathbb{R}^3} u$. Notice that $u \in C_0^{\infty}(\mathbb{R}^3; \mathbb{R}_+)$. Then, by Sard's Theorem, the set

$$\left\{t \in [0,M] : \exists (x_1, x_2, y) \in \mathbb{R}^3 \text{ s.t } u(x_1, x_2, y) = t, \ \nabla u(x_1, x_2, y) = 0\right\}$$

has Lebesgue measure 0 in \mathbb{R} . By the definition of u^* , the set

$$\left\{t \in [0,M] : \exists (x_1, x_2, y) \in \overline{\mathbb{R}}_{s_1}^3 \text{ s.t } u^*(x_1, x_2, y) = t, \ \nabla u^*(x_1, x_2, y) = 0\right\}$$

has Lebesgue measure 0 in \mathbb{R} . Using the co-area formula, we deduce that

$$\int_{\mathbb{R}^3} |\nabla_G u|^2 dx_1 dx_2 dy = \int_0^M dt \int_{u^{-1}(t)} |\nabla_G u| d\mu_G,$$

and

$$\int_{\mathbb{R}^{3}_{s_{1}}} |\nabla_{G}u^{*}|^{2} dx_{1} dx_{2} dy = \int_{0}^{M} dt \int_{u^{*-1}(t)} |\nabla_{G}u^{*}| d\mu_{G}^{*},$$

where $d\mu_G = \frac{|\nabla_G u|}{|\nabla u|} d\mathcal{H}^2$ and $d\mu_G^* = \frac{|\nabla_G u^*|}{|\nabla u^*|} d\mathcal{H}^2$. To prove Theorem 2, we only need to show that for $t \in [0, M]$ such that t is not a critical value of u and u^* , the following inequality holds

$$\int_{u^{*-1}(t)} |\nabla_G u^*| d\mu_G^* \le \int_{u^{-1}(t)} |\nabla_G u| d\mu_G.$$
(2.5)

Notice that

$$d\mu_G = \frac{|\nabla_G u|}{|\nabla u|} d\mathcal{H}^2 = \sqrt{\nu_1^2 + \nu_2^2 + |x|^{2\alpha} \nu_3^2} d\mathcal{H}^2.$$

From this and the Hölder inequality, we obtain

$$(P_{2,\alpha}(\{u > t\}))^{2} = \left(\int_{u^{-1}\{t\}} |x|^{\alpha} d\mu_{G}\right)^{2}$$
$$\leq \left(\int_{u^{-1}\{t\}} |\nabla_{G}u| d\mu_{G}\right) \left(\int_{u^{-1}\{t\}} \frac{|x|^{2\alpha}}{|\nabla_{G}u|} d\mu_{G}\right).$$
(2.6)

By definition of u^* , we have

$$\nabla_G u^*(x_1, x_2, y)| = (\alpha + 1)|x|^{\alpha}|g'(r)|.$$

Since $d\mu_G^* = \frac{|\nabla_G u^*|}{|\nabla u^*|} d\mathcal{H}^2$ and r is constant on $u^{*-1}{t} \cap \mathbb{R}^3_{s_1}$, we deduce that

$$(P_{2,\alpha,1}(\{u^* > t\}))^2 = \left(\int_{u^{*-1}\{t\}} |x|^{\alpha} d\mu_G^*\right)^2$$
$$= \left(\int_{u^{*-1}\{t\}} |\nabla_G u^*| d\mu_G^*\right) \left(\int_{u^{*-1}\{t\}} \frac{|x|^{2\alpha}}{|\nabla_G u^*|} d\mu_G^*\right).$$
(2.7)

Recall that

$$|\{u>t\}|_{2,\alpha}=|\{u^*>t\}|_{2,\alpha}.$$

Thus, Theorem 1 infers that

$$P_{2,\alpha,1}(\{u^* > t\}) \le P_{2,\alpha}(\{u > t\}).$$
(2.8)

The inequality (2.5) then follows from Lemma 1, (2.6), (2.7), (2.8) and the fact that $\frac{d\mu_G}{|\nabla_G u|} = \frac{d\mathcal{H}^2}{|\nabla u|}$. Our proof is complete.

3. Sobolev-type inequalities with weights

Definition 5. Let q > 1. We define $W_0^{1,2\alpha,q}(\mathbb{R}^3)$ as the completion of $C_0^{\infty}(\mathbb{R}^3)$ with respect to the norm

$$\|u\|_{W_0^{1,2\alpha,q}} = \left(\int_{\mathbb{R}^3} |\nabla_G u|^2 dx_1 dx_2 dy\right)^{1/2} + \left(\int_{\mathbb{R}^3} |x|^{2\alpha} |u|^q dx_1 dx_2 dy\right)^{1/q},$$

where $|x| = \sqrt{x_1^2 + x_2^2}$.

For $u \in W_0^{1,2\alpha,q}(\mathbb{R}^3) \setminus \{0\}$, we consider the ratio

$$C_{2\alpha,q}(u) = \frac{\left(\int_{\mathbb{R}^3} |\nabla_G u|^2 dx_1 dx_2 dy\right)^{1/2}}{\left(\int_{\mathbb{R}^3} |x|^{2\alpha} |u|^q dx_1 dx_2 dy\right)^{1/q}}.$$
(3.1)

By rescaling $X_1 = \lambda x_1, X_2 = \lambda x_2$, and $Y = \lambda^{\alpha+1}y$, we have $U(x, y) = u(\lambda x, \lambda^2 y)$ and

$$\int_{\mathbb{R}^3} |\nabla_G U|^2 dx_1 dx_2 dy = \lambda^{-(\alpha+1)} \int_{\mathbb{R}^3} |\nabla_G u|^2 dX_1 dX_2 dY,$$

and

$$\int_{\mathbb{R}^3} |x|^{2\alpha} |U|^q dx_1 dx_2 dy = \lambda^{-(3\alpha+3)} \int_{\mathbb{R}^3} |X|^2 |u|^q dX_1 dX_2 dY,$$

so that

$$C_{2\alpha,q}(U) = \lambda^{\frac{3\alpha+3}{q} - \frac{\alpha+1}{2}} C_{2\alpha,q}(u).$$

Hence, to have

$$\inf_{W_0^{1,2\alpha,q}(\mathbb{R}^3)\setminus\{0\}} C_{2\alpha,q}(u) > 0$$

we need q = 6 to be the critical exponent. For this case, we obtain the best Sobolev inequality as follows.

Theorem 3. The following inequality holds

$$\left(\int_{\mathbb{R}^3} |x|^{2\alpha} |u|^6 dx_1 dx_2 dy\right)^{1/6} \le C_{2\alpha,6}^{-1} \left(\int_{\mathbb{R}^3} |\nabla_G u|^2 dx_1 dx_2 dy\right)^{1/2}, \qquad (3.2)$$

The best constant $C_{2\alpha,6}$ satisfies

$$C_{2\alpha,6} \ge \left(\frac{2\pi}{n(\alpha)}\right)^{\frac{-1}{3}} (\alpha+1)^{\frac{-1}{3}} D_{2,6,3}$$

where $D_{2,6,3}$ is a constant which will be given in Lemma 2.

To prove Theorem 3, we first need to recall here the results of [19, Lemma 2].

Lemma 2. Let m, p, q be real numbers such that

$$1$$

Let $\phi : \mathbb{R}_+ \to \mathbb{R}_+$ be a Lipschitz function and such that

$$\int_{0}^{\infty} r^{m-1} |\phi'(r)|^p dr < \infty, \ \phi(r) \to 0 \ when \ r \to \infty.$$
(3.3)

Then

$$\frac{\left(\int_0^\infty r^{m-1} |\phi'(r)|^p\right)^{1/p}}{\left(\int_0^\infty r^{m-1} |\phi(r)|^q\right)^{1/q}} \ge D_{p,q,m}$$

with the best constant given by

$$D_{p,q,m} = m^{\frac{1}{p}} \left(\frac{p-1}{m-1}\right)^{-\frac{1}{p'}} \left[\frac{1}{p'} B\left(\frac{m}{p}, \frac{m}{p'}\right)\right]^{\frac{1}{m}},$$

where p' = p/(p-1). The equality sign in (3.3) holds with

$$\phi(r) = (a + br^{p'})^{1 - m/p},$$

where a, b are positive constant.

We also need some calculations. Using the polar coordinates

$$x_1 = (r\sin\varphi)^{1/(\alpha+1)}\sin\theta, \ x_2 = (r\sin\varphi)^{1/(\alpha+1)}\cos\theta, \ y = \frac{r\cos\varphi}{\alpha+1},$$

we have $dx_1 dx_2 dy = \frac{r^{2/(\alpha+1)}}{(\alpha+1)^2} (\sin \varphi)^{(1-\alpha)/(\alpha+1)} dr d\theta d\varphi$. Next, consider the case $u(x_1, x_2, y) = \phi(r)$, where $r = (|x|^{2\alpha+2} + (\alpha+1)^2 y^2)^{1/2}$. We have

$$\int_{\mathbb{R}^3_{s_1}} |x|^{2\alpha} |u|^6 dx_1 dx_2 dy = \frac{2\pi}{n(\alpha)(\alpha+1)^2} \int_0^\infty r^2 |\phi(r)|^6 dr,$$
(3.4)

and

$$\int_{\mathbb{R}^{3}_{s_{1}}} |\nabla_{G}u|^{2} dx_{1} dx_{2} dy = \frac{2\pi}{n(\alpha)} \int_{0}^{\infty} r^{2} |\phi'(r)|^{2} dr.$$
(3.5)

Now we are able to give the proof of Theorem 3.

Proof of Theorem 3. Since $C_0^{\infty}(\mathbb{R}^3)$ is dense in $W_0^{1,2\alpha,6}(\mathbb{R}^3)$, we can assume that $u \in C_0^{\infty}(\mathbb{R}^3)$. Notice that

$$|\nabla_G|u|| \le |\nabla_G u|$$

thus we can assume u is a nonnegative function. It is easy to see that we can approximate a nonnegative function $w \in W_0^{1,2\alpha,6}(\mathbb{R}^3)$ by a sequence of nonnegative functions $\{w_j\}_{j=1}^{\infty}$, we only need to give the proof for $u \in C_0^{\infty}(\mathbb{R}^3)$ and $u \ge 0$.

Let u^* be respectively the rearrangement of u. The property of rearrangement infers that

$$\int_{\mathbb{R}^3_{s_1}} |x|^{2\alpha} |u^*|^6 dx_1 dx_2 dy = \int_{\mathbb{R}^3} |x|^{2\alpha} |u|^6 dx_1 dx_2 dy.$$
(3.6)

Note that $u \in C_0^{\infty}(\mathbb{R}^3)$, $u \ge 0$. It follows from Theorem 2 that

$$\int_{\mathbb{R}^{3}_{s_{1}}} |\nabla_{G} u^{*}|^{2} dx_{1} dx_{2} dy \leq \int_{\mathbb{R}^{3}} |\nabla_{G} u|^{2} dx_{1} dx_{2} dy.$$
(3.7)

On the other hand, we have

$$u^*(x_1, x_2, y) = \phi(r),$$

where $r = (|x|^{2\alpha+2} + (\alpha+1)^2 y^2)^{1/2}$. Note that the function ϕ satisfies Lemma 2. Then, from (3.4), (3.5) and Lemma 2, we deduce that

$$L_{2\alpha,6} \left(\int_{\mathbb{R}^3_{s_1}} |x|^{2\alpha} |u^*|^6 dx_1 dx_2 dy \right)^{1/6} \le \left(\int_{\mathbb{R}^3_{s_1}} |\nabla_G u^*|^2 dx_1 dx_2 dy \right)^{1/2}, \qquad (3.8)$$

where

$$L_{2\alpha,6} = \left(\frac{2\pi}{n(\alpha)}\right)^{\frac{-1}{3}} (\alpha+1)^{\frac{-1}{3}} D_{2,6,3}$$
(3.9)

with $D_{2,6,3}$ is given in Lemma 2. From (3.6) - (3.9), our Sobolev-type inequality (3.2) follows. Our proof is complete.

Remark 2. Our proof only gives an lower bound for the best constant $C_{2\alpha,6}$. The exact value of $C_{2\alpha,6}$ still remains an open question.

4. Applications to a class of degenerate elliptic equations

In this section, we are mainly concerned with the existence and nonexistence of nontrivial solutions to the semilinear subelliptic Dirichlet problem

$$\begin{cases} -\Delta_x u - |x|^{2\alpha} \frac{\partial^2 u}{\partial y^2} = f(x, y, u), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$
(4.1)

where $x = (x_1, x_2) \in \mathbb{R}^2$, Ω is a bounded smooth domain in \mathbb{R}^3 , $(0, 0, 0) \in \Omega$, and $\alpha > 0$. Notice that our methods are similar to the ones that have been used in [3], and thus many details will be omitted.

4.1. Nonexistence result. In this subsection, we will derive the nonexistence of nontrivial solutions to the problem (4.1) for $f(x_1, x_2, y, \xi) = |x|^{2\alpha} |\xi|^{p-1} \xi$, $p \ge 1$. To this end, let

$$F(x_1, x_2, y, \xi) := \int_0^{\xi} f(x_1, x_2, y, \tau) d\tau = \frac{|x|^{2\alpha}}{p+1} |\xi|^{p+1}$$

We denote by $\mathcal{S}_2(\overline{\Omega})$ the linear space of functions $C_0^1(\overline{\Omega})$ such that

$$\frac{\partial^2 u}{\partial x_1^2}$$
, $\frac{\partial^2 u}{\partial x_2^2}$, and $|x|^{2\alpha} \frac{\partial^2 u}{\partial y^2}$ (in distribution sense)

are continuous in Ω and can be continuously extended to $\overline{\Omega}$.

A function $u(x_1, x_2, y) \in S_2(\overline{\Omega})$ is said to be a solution to the problem (4.1) if

$$\begin{cases} -\Delta_x u - |x|^{2\alpha} \frac{\partial^2 u}{\partial y^2} = |x|^{2\alpha} |u|^{p-1} u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega. \end{cases}$$

Lemma 3. Let $u(x_1, x_2, y) \in S_2(\overline{\Omega})$ be a solution of (4.1). Denote by $\nu = (\nu_1, \nu_2, \nu_3)$ the unit outward normal on $\partial\Omega$. Then we have

$$\left(\frac{3\alpha+3}{p+1} - \frac{\alpha+1}{2}\right) \int_{\Omega} |x|^{2\alpha} |u|^{p+1} dx_1 dx_2 dy$$

=
$$\int_{\partial \Omega} [x_1\nu_1 + x_2\nu_2 + (1+\alpha)y\nu_3](\nu_1^2 + \nu_2^2 + \nu_3^2|x|^{2\alpha}) \left(\frac{\partial u}{\partial \nu}\right)^2 ds. \quad (4.2)$$

Proof. The proof is similar to the one of the lemma in [24], and for this reason, it is omitted here. \Box

Definition 6. A domain Ω is called G_{α} – star-shaped with respect to the origin if $(0,0,0) \in \Omega$ and $x_1\nu_1 + x_2\nu_2 + (1+\alpha)y\nu_3 \ge 0$ at every point of $\partial\Omega$.

The following result is obtained directly from Lemma 3.

Theorem 4. Let Ω be G_{α} – star-shaped with respect to the origin and p > 5. Then the problem (4.1) has no nontrivial solution $u \in S_2(\overline{\Omega})$. 4.2. Existence result. In this subsection, we will give the existence result for the problem (4.1). We have the following definitions.

Definition 7. Let Ω be a bounded domain in \mathbb{R}^3 . We denote by $L^p_{|x|^{2\alpha}}(\Omega)$ the set of all measurable functions $u: \Omega \to \mathbb{R}$ such that

$$\int_{\Omega} |x|^{2\alpha} |u|^p dx_1 dx_2 dy < \infty.$$

We define the norm in $L^p_{|x|^{2\alpha}}(\Omega)$ as follows

$$\|u\|_{L^p_{|x|^{2\alpha}}(\Omega)} := \left(\int_{\Omega} |x|^{2\alpha} |u|^p dx_1 dx_2 dy\right)^{1/p},$$

for every $u \in L^p_{|x|^{2\alpha}}(\Omega)$.

Definition 8. Let Ω be a bounded domain in \mathbb{R}^3 . We denote by $S_1^2(\Omega)$ the set of all functions $u \in L^2(\Omega)$ such that

$$\frac{\partial u}{\partial x_1}, \ \frac{\partial u}{\partial x_2}, \ and \ |x|^\alpha \frac{\partial u}{\partial y} \in L^2(\Omega).$$

We define the norm in $\mathcal{S}_1^2(\Omega)$ as follows

$$\|u\|_{\mathcal{S}^{2}_{1}(\Omega)} := \left(\int_{\Omega} |u|^{2} dx_{1} dx_{2} dy\right)^{1/2} + \left(\int_{\Omega} |\nabla_{G} u|^{2} dx_{1} dx_{2} dy\right)^{1/2}$$

where $\nabla_G u := \left(\frac{\partial u}{\partial x_1}, \frac{\partial u}{\partial x_2}, |x|^{\alpha} \frac{\partial u}{\partial y}\right)$. We can also define the scalar product in $\mathcal{S}_1^2(\Omega)$ by

$$(u,v)_{\mathcal{S}^{2}_{1}(\Omega)} := (u,v)_{L^{2}(\Omega)} + (\nabla_{G}u, \nabla_{G}v)_{L^{2}(\Omega)}.$$

Definition 9. The space $S_{1,0}^2(\Omega)$ is defined as the closure of $C_0^1(\Omega)$ in the space $S_1^2(\Omega)$.

Remark 3. By a similar argument as in the proof of [24, Theorem 6], we also have the two norms $||u||_{S^2_1(\Omega)}$ and

$$\|u\|_{\mathcal{S}^{2}_{1,0}(\Omega)} := \left(\int_{\Omega} |\nabla_{G} u|^{2} dx_{1} dx_{2} dy\right)^{1/2}$$

are equivalent in $\mathcal{S}^2_{1,0}(\Omega)$.

We need the following embedding theorem.

Theorem 5. Let Ω be a bounded domain in \mathbb{R}^3 with smooth boundary $\partial\Omega$ such that $(0,0,0) \in \Omega$. Then the embedding

 $\mathcal{S}^2_{1,0}(\Omega) \hookrightarrow L^q_{|x|^{2\alpha}}(\Omega), \text{ where } 1 \leq q \leq 6,$

is continuous, i.e. there exists a constant $C_q > 0$ such that

 $\|u\|_{L^q_{|x|^{2\alpha}}(\Omega)} \le C_q \|u\|_{\mathcal{S}^2_1(\Omega)}, \ \forall u \in \mathcal{S}^2_{1,0}(\Omega).$

Moreover, the embedding

$$\mathcal{S}^2_{1,0}(\Omega) \hookrightarrow L^q_{|x|^{2\alpha}}(\Omega), \text{ where } 1 \leq q < 6,$$

is compact.

Proof. The proof of this lemma is similar to the one of [3, Theorem 3.2]. We omit the details. \Box

Now we are able to study the existence result for the problem (4.1). To this end, we first have the following definition of weak solutions to the problem (4.1).

Definition 10. A function $u \in S^2_{1,0}(\Omega)$ is called a weak solution of the problem (4.1) if the following identity

$$\int_{\Omega} \nabla_G u \cdot \nabla_G \varphi dx_1 dx_2 dy - \int_{\Omega} f(x, y, u) \varphi dx_1 dx_2 dy = 0$$

holds for every $\varphi \in \mathcal{S}^2_{1,0}(\Omega)$.

The following lemma is obtained from the Hölder inequality and Theorem 5.

Lemma 4. Assume that $f: \Omega \times \mathbb{R} \to \mathbb{R}$ is a Carathéodory function such that there exist $q \in (2,6), f_1(x,y) \in L^{p_1}_{|x|^{2\alpha}}(\Omega, \mathbb{R}_+), f_2(x,y) \in L^{p_2}_{|x|^{2\alpha}}(\Omega, \mathbb{R}_+),$ where $p_2 > 1$, $qp_2/(p_2-1) \leq 6$, and $p_1 > \frac{6p_2}{p_2(q-1)+6}$ such that

$$|f(x_1, x_2, y, \xi)| \le |x|^{2\alpha} (f_1(x_1, x_2, y) + f_2(x_1, x_2, y)|\xi|^{q-1})$$

for a.e. $(x_1, x_2, y) \in \Omega, \forall \xi \in \mathbb{R}$. Then $\Psi_1(u) \in C^1(\mathcal{S}^2_{1,0}(\Omega), \mathbb{R})$ and

$$\Psi_1'(u)(v) = \int_{\Omega} f(x_1, x_2, y, u) v dx_1 dx_2 dy$$

for all $v \in \mathcal{S}^2_{1,0}(\Omega)$, where

$$\Psi_1(u) = \int_{\Omega} F(x_1, x_2, y, u) dx_1 dx_2 dy,$$

and $F(x_1, x_2, y, \xi) = \int_0^{\xi} f(x_1, x_2, y, \tau) d\tau$.

Next, we assume that $f: \Omega \times \mathbb{R} \to \mathbb{R}$ is a Carathéodory function such that

(A1) f satisfies the assumption in Lemma 4 and the additional condition

$$p_1 > \frac{3}{2};$$
 (4.3)

(A2) there exist $C \in [0, +\infty)$ and $\psi \in L^1_{|x|^{2\alpha}}(\Omega)$ such that

$$|f(x_1, x_2, y, \xi)| \le |x|^{2\alpha} \psi(x_1, x_2, y)$$

for a.e. (x_1, x_2, y) in Ω and for every $|\xi| \leq C$;

(A3) there exists a non-positive function φ such that

$$\int_{\Omega} |\varphi(x_1, x_2, y)| dx_1 dx_2 dy < \infty$$

and

$$\varphi(x_1, x_2, y) \le \frac{f(x_1, x_2, y, \xi)}{\xi}$$

for a.e. $(x_1, x_2, y) \in \Omega$ and for every $\xi \in \mathbb{R}_+$;

(A4) $f(x_1, x_2, y, 0) = 0$ for a.e. (x_1, x_2, y) in Ω and the following limit holds uniformly for a.e. (x_1, x_2, y) in Ω

$$\lim_{\xi \to 0} \frac{f(x_1, x_2, y, \xi)}{|x|^{2\alpha} \xi} = 0 \text{ and } \lim_{|\xi| \to +\infty} \frac{f(x_1, x_2, y, \xi)}{|x|^{2\alpha} \xi} = +\infty;$$

(A5) $\frac{f(x_1, x_2, y, \xi)}{\xi}$ is increasing in $\xi \ge C$ and decreasing in $\xi \le -C$ for a.e. (x_1, x_2, y) in Ω .

In the following we define the energy functional $\Phi : \mathcal{S}^2_{1,0}(\Omega) \to \mathbb{R}$ associated with the problem (4.1) by letting

$$\Phi(u) := \frac{1}{2} \int_{\Omega} |\nabla_G u|^2 dx_1 dx_2 dy - \int_{\Omega} F(x, y, u) dx_1 dx_2 dy$$
(4.4)

for all $u \in S_{1,0}^2(\Omega)$, where F is defined as in Lemma 4. It follows from Lemma 4 and the condition (A1) that Φ is well-defined on $S_{1,0}^2(\Omega)$ and $\Phi \in C^1(S_{1,0}^2(\Omega), \mathbb{R})$ with

$$\langle \Phi'(u), v \rangle = \int_{\Omega} \nabla_G u \cdot \nabla_G v dx_1 dx_2 dy - \int_{\Omega} f(x, y, u) v dx_1 dx_2 dy,$$

for all $v \in S^2_{1,0}(\Omega)$. Therefore, the weak solutions of the problem (4.1) are critical points of the functional Φ .

We need to use the Mountain Pass Lemma to study the existence of critical points of the functional Φ . First, let us recall the notion of the $(C)_c$ condition, which is an important notion in the statement of this lemma.

Definition 11. Let X be a real Banach space with its dual space X^* and let $\Psi \in C^1(X, \mathbb{R})$. For $c \in \mathbb{R}$ we say that Ψ satisfies the $(C)_c$ condition if for any sequence $\{u_n\}_{n=1}^{\infty} \subset X$ with

$$\Psi(u_n) \to c \text{ and } (1 + ||u_n||_{\mathbf{X}}) ||\Psi'(u_n)||_{\mathbf{X}^*} \to 0,$$

there exists a subsequence $\{u_{n_k}\}_{k=1}^{\infty}$ that converges strongly in X.

We will need the following version of the Mountain Pass Lemma (see also [5, 6]).

Lemma 5. Let X be a real Banach space and let $\Psi \in C^1(X, \mathbb{R})$ satisfy the $(C)_c$ condition for any $c \in \mathbb{R}$, $\Psi(0) = 0$ and

(i) There exist constants $\rho, \alpha > 0$ such that $\Psi(u) \ge \alpha, \forall u \in \mathbf{X}, ||u||_{\mathbf{X}} = \rho$;

(ii) There exists an $u_1 \in \mathbf{X}$, $||u_1||_{\mathbf{X}} \ge \rho$ such that $\Psi(u_1) \le 0$.

Then $\beta := \inf_{\lambda \in \Lambda} \max_{0 \le t \le 1} \Psi(\lambda(t)) \ge \alpha$ is a critical value of Ψ , where

$$\Lambda := \{ \lambda \in C([0;1], \mathbf{X}) : \lambda(0) = 0, \lambda(1) = u_1 \}.$$

We have the following existence theorem.

Theorem 6. Suppose that f satisfies (A1)-(A5). Then the boundary value problem (4.1) has a nontrivial weak solution.

Proof. The proof follows the same arguments presented in the one of [3, Theorem 4.3] and for this reason, the details will be omitted here. The idea is that based on the fact that f satisfies the conditions (A1)-(A5) and Theorem 5, one can verify that the functional Φ satisfies the conditions of Lemma 5. Thus our proof is complete.

Acknowledgement

The second author of this paper is a member of a VAST project.

12

WEIGHTED INEQUALITIES

References

- NGA, N.Q., TRI, N.M., TUAN, D.A. Pólya–Szegö type inequality and imbedding theorems for weighted Sobolev spaces. Anal. Math. Phys. 14(2), 1-12 (2024).
- [2] LUYEN, D.T., TRI, N.M. Infinitely many solutions for a class of perturbed degenerate elliptic equations involving the Grushin operator. *Complex Var. Elliptic Equ.* 65(12), 2135–2150 (2020)
- [3] LUYEN, D.T., TRI, N.M., TUAN, D.A. Nontrivial solutions to the Dirichlet problems for semilinear degenerate elliptic equations. *Math. Notes.* 116(5), 1051-1063 (2024)
- [4] LIONS, P.L., PACELLA, F. Isoperimetric inequalities for convex cones. Proc. Amer. Math. Soc. 109 (1990), no. 2, 477–485
- [5] CERAMI, G. An existence criterion for the critical points on unbounded manifolds. (Italian) Istit. Lombardo Accad. Sci. Lett. Rend. A 112 (1978), no. 2, 332–336.
- [6] CERAMI, G. On the existence of eigenvalues for a nonlinear boundary value problem. (Italian) ANN. MAT. PURA APPL. (4) 124 (1980), 161–179
- [7] ALVINO, A., BROCK, F., CHIACCHIO, F., MERCALDO, A., POSTERARO, M.R. Some isoperimetric inequalities with respect to monomial weights. em ESAIM: control. Optim. Calc. Var. 27(Supplement), 1–29 (2021)
- [8] AUBIN, T. Problèmes isopérimétriques et espaces de Sobolev. J. Differ. Geom. 11(4), 573–598 (1976)
- BECKNER, W. On the Grushin operator and hyperbolic symmetry. Proc. AMS. 129(4), 1233–1246 (2000)
- [10] BROCK, F., CHIACCHIO, F., MERCALDO, A. Weighted isoperimetric inequalities in cones and applications. Nonlinear Anal. 75(2012), no.15, 5737–5755.
- BROCK, F., CHIACCHIO, F., MERCALDO, A. A weighted isoperimetric inequality in an orthant. *Potential Anal.* 41(1), 171–186 (2014)
- [12] CAPOGNA, L., DANIELLI, D., GAROFALO, N. An embedding theorem and the Harnack inequality for nonlinear subelliptic equations. *Comm. Partial Differ. Equ.* 18(9–10), 1765–1794 (1993)
- [13] CHEN, H., CHEN, H.G., LI, J.N. Estimates of Dirichlet eigenvalues for degenerate Δ_{μ} -Laplace operator. Calc. Var. Partial Differ. Equ. 59(4), 27 (2020)
- [14] CABRE, X., ROS-OTON, X. Sobolev and isoperimetric inequalities with monomial weights. J. Differ. Equ. 225(11), 4312–4336 (2013)
- [15] CIANCHI, A., FUSCO, N. Functions of bounded variation and rearrangements. Arch. Ration. Mech. Anal. 165, 1–40 (2002)
- [16] DOU, J., SUN, L., WANG, L., ZHU, M. Divergent operator with degeneracy and related sharp inequalities. J. Funct. Anal. 282(2), 109–294 (2022)
- [17] MONTI, R., MORBIDELLI, D. Isoperimetric inequality in the Grushin plane. J. Geom. Anal. 14(2), 355–368 (2004)
- [18] PÓLYA, G., SZEGÖ, G. Isoperimetric inequalities in mathematical physics, annals of mathematics studies. Princeton University Press, Princeton (1951)
- [19] TALENTI, G. Best constant in Sobolev inequality. Ann. Mat. Pura Appl. 110, 353–372 (1976)
- [20] MADERNA, C., SALSA, S. Sharp estimates for solutions to a certain type of singular elliptic boundary value problems in two dimensions. *Appl. Anal.* 12(4), 307–321 (1981)
- [21] KOGOJ, A.E., LANCONELLI, E. On semilinear $\Delta_\lambda\text{-Laplace}$ equation. Nonlinear Anal. 75, 4637–4649 (2012)
- [22] TRI, N.M. On Grushin's equation. Math. Notes. 63(1), 84–93 (1998)
- [23] TRI, N.M. Recent progress in the theory of semilinear equations involving degenerate elliptic differential operators. p. 376. Publishing House for Science and Technology, Vietnam (2014)
- [24] TRI, N.M. Critical Sobolev exponent for degenerate elliptic operators. Acta Math. Vietnam. 23 (1998), 83–94.
- [25] LIU, S. On superlinear problems without the Ambrosetti and Rabinowitz condition. Nonlinear Anal. 73 (2010), no. 3, 788–795.

Trung Hieu Giang 1,2

 1 Department of Mathematics, City University of Hong Kong, 83 Tat Chee Avenue,

Kowloon, Hong Kong;
 ² Institute of Mathematics, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam.

Email address: thgiang2-c@my.cityu.edu.hk

Nguyen Minh Tri²

 2 Institute of Mathematics, Vietnam Academy of Science and Technology, 18 Hoang QUOC VIET, CAU GIAY, HANOI, VIETNAM.

Email address: triminh@math.ac.vn

Dang Anh Tuan 3

 3 University of Sciences, Vietnam National University, 334 Nguyen Trai, Thanh Xuan, HANOI, VIETNAM.

Email address: datuan1105@gmail.com