Spatiotemporal pattern formations in a two-layer coupled reaction-diffusion Lengyel-Epstein system

Qidong Wu, Fengqi Yi^{*}

School of Mathematical Sciences, Dalian University of Technology, Dalian, Liaoning Province, 116024, China.

Abstract

Spatiotemporal pattern formations in two-layer coupled reaction-diffusion Lengyel-Epstein system with distributed delayed couplings are investigated. Firstly, for the original decoupled system, it is proved that when the intra-reactor diffusion rate ϵ of the inhibitor is sufficiently small and the intra-reactor diffusion rate d of the inhibitor is large enough, then the subsystem can exhibit non-constant positive steady state $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ with large amplitude, and that as the parameter τ varies, the stability of $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ changes, leading to the emergence of periodic solutions via Hopf bifurcation. Secondly, for the two-layer coupled system, the stability of the symmetric steady state $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is studied by treating k_1, k_2 (the inter-reactor diffusion rates) and α (the delay parameter) as the main parameters. In case of non-delayed couplings, the first quadrant of the (k_1, k_2) parameter space can be divided into two regions: one is stable region, the other one is unstable region, and the two regions have the common boundary, which is the primary Turing bifurcation curve. In case of delayed couplings, it is shown that the first quadrant of the (k_1, k_2) parameter space can be re-divided into three regions: the first one is unstable region, the second one is stable region, while the third one is the potential "bifurcation" region, where Hopf bifurcation may occur for suitable α . Our analysis is mainly based on the singular perturbation techniques and the implicit function theorem, and the results show some different phenomena from those of the original decoupled system in one reactor.

Keywords: Coupled Lengyel-Epstein system; Distributed-delay coupling; Non-constant layered solution; Turing bifurcation and Hopf bifurcation; Singular limit eigenvalue problem. **MSC**: 35B32, 35P20, 35Q92, 92E20.

1 Introduction

One of the crucial issues in developmental biology is to concern the spatial pattern formation in the early embryo. In a remarkable work [31], Turing predicted that a system of reacting and

^{*}Corresponding Author (F. Yi). Email: yi@dlut.edu.cn

[†]The authors were partially by National Key R & D Program of China (2023YFA1009200), the National Natural Science Foundation of China (11971088, 12371160, 12271144) and the Fundamental Research Funds for the Central Universities (DUT18RC(3)070, DUT23LAB304).

diffusing chemicals could generate spatial patterns in chemical concentrations from initial near homogeneity. He suggested that this process might lead to a variety of morphogenetic phenomena. The possibility of such diffusion-driven instability, now usually termed Turing instability, has been widely investigated in the chemical and biological literatures.

In 1990, Kepper and the collaborators [3] discovered the formation of the stationary patterns in the chlorite–iodide–malonic acid-starch (CIMA) reaction in an open unstirred gel reactor, which is the first unambiguous experimental evidence of the existence of the symmetry breaking reactiondiffusion structures predicted by Turing in 1952. In this CIMA chemical reaction, there are five reactants, which makes the mathematical description very complicated. However, numerical calculations showed that during oscillations in the CIMA reaction, only the concentrations of I^- (the activator iodide) and ClO_2^- (the inhibitor chlorite) vary significantly, and they react to produce the immobile complexing reactant malonic acid (MA).

The partial differential equations reflecting the evolutionary behaviors of the concentrations $u := [I^-], v := [\text{ClO}_2^-]$ and w := [MA] are given by (see [11, 12, 13])

(1.1)
$$\begin{cases} \frac{\partial u}{\partial t} = d_1 \Delta u + a - u - \frac{4uv}{1 + u^2} - k_+ u + k_- w, \\ \frac{\partial v}{\partial t} = d_2 \Delta v + b \left(u - \frac{uv}{1 + u^2} \right), \\ \frac{\partial w}{\partial t} = k_+ u - k_- w, \end{cases}$$

where $d_1 > 0$ and $d_2 > 0$ are the diffusion rates of the activator I^- and the inhibitor ClO_2^- , respectively. Since malonic acid is assumed to be immobile, there is no diffusion in the third equation of (1.1). The constants a, b, k_+ and k_- are all positive. Moreover, the homogeneous Neumann boundary conditions are imposed on the boundary.

If the formation and the dissociation of the complex are rapid, then w = ku for some k > 0 (see [12] for details). Thus, by setting $\sigma = 1 + k$, we can reduce system (1.1) to the following so-called reaction-diffusion Lengyel-Epstein system ([12])

(1.2)
$$\begin{cases} \frac{\partial u}{\partial t} = \frac{1}{\sigma} \left(d_1 \Delta u + a - u - \frac{4uv}{1 + u^2} \right), \\ \frac{\partial v}{\partial t} = d_2 \Delta v + b \left(u - \frac{uv}{1 + u^2} \right). \end{cases}$$

Setting $t' = \sigma t$ and $x' = x/\sqrt{d_1}$, one can reduce system (1.2) to

(1.3)
$$\begin{cases} \frac{\partial u}{\partial t} = \Delta u + a - u - \frac{4uv}{1 + u^2}, \\ \frac{\partial v}{\partial t} = \sigma c \Delta v + \sigma b \left(u - \frac{uv}{1 + u^2} \right) \end{cases}$$

where we drop the ' from t' and x' for convenience, and c is the effective diffusion ratio d_2/d_1 .

The dynamics of the reaction-diffusion Lengyel-Epstein system mainly concentrate on the dynamics of the rescaled system (1.3). For example, in [22], Ni and Tang studied Turing instability of the unique constant steady state of system (1.3) and further considered the existence/nonexistence of the non-constant steady states by using topological degree theory. In [8], Jang, Ni and Tang studied the global bifurcation structure of the set of the non-constant steady states for system (1.3). In [9], Jin, Shi, Wei and Yi studied the interactions between Hopf bifurcations and steady state bifurcations of system (1.3). In [35], Yi studied Turing instability of the spatially homogeneous periodic solutions of system (1.3) by using the regular perturbation methods. In [39], Zhao, Yu and Wang introduced discrete time delay into the system and studied Turing bifurcation, Hopf bifurcation, Turing-Turing bifurcation, Turing-Hopf bifurcation and Bogdanov-Takens bifurcation. These results have the common features: the existence of the non-constant steady states has been established, however, the stability of these solutions was unsettled. As one of the purposes of this paper, we shall give an affirmative answer for the stability issue.

On the other hand, since bilayer membranes or multilayer tissues are often found in biological, physical and chemical systems, spatiotemporal pattern formation in coupled system that consists of two or more coupled chemical reacting subsystems is of great significance ([10, 12, 14, 24]). Generally speaking, the width of the membrane cannot be negligible compared with the size of the system, it is then necessary to include the time delay in the coupling terms (see [1, 2, 7, 23, 32, 36, 37]).

Motivated by [7, 12, 14, 32, 33], we make the following assumptions:

- (H1). There are two identical well-stirred reactors (layers), where the chemical reactions and initial or feed concentrations of the reactants are exactly the same. They are connected by a semipermeable membrane through which the chemicals can diffuse (termed "interreactor" diffusion) by Fick's first law (see [12, 32]): the mass flux J_X of a species (denoted by X) from reactor 1 to reactor 2 is directly proportional to the concentration difference $\Delta X = X_2 - X_1$, that is, $J_X = -\rho_X \Delta X$, where ρ_X is the constant of proportionality. The amount of X leaving reactor 1 through the membrane of area A in time Δt is $J_X A \Delta t$. Then, the rate of change of concentration X in reactor 1 (resp., reactor 2) of volume V is $k_X(X_2 - X_1)$ (resp., $-k_X(X_2 - X_1)$), where $k_X := \rho_X A/V$ is the "inter-reactor effective diffusion constant".
- (H2). In absence of coupling, in each of the reactors, there also have "intra-reactor" diffusions, which are the random diffusions in the spatial domain Ω of one reactor and are described by the Laplacian terms in each equation. The dynamics of each of the two reactors (designated 1 and 2) is described by the following reaction-diffusion equations in the absence of coupling:

$$\text{Reactor } i \ (i = 1, 2) : \begin{cases} \frac{\partial u_i}{\partial t} = \frac{1}{\sigma} \left(d_1 \Delta u_i + a - u_i - \frac{4u_i v_i}{1 + u_i^2} \right), & (x, t) \in \Omega \times (0, +\infty), \\ \frac{\partial v_i}{\partial t} = d_2 \Delta v_i + b \left(u_i - \frac{u_i v_i}{1 + u_i^2} \right), & (x, t) \in \Omega \times (0, +\infty), \end{cases}$$

subject to homogeneous Neumann boundary conditions.

(H3). There has the distributed time delay occurring when the species cross the membrane (see [7, 15, 26, 38] for some practical backgrounds). For our technical reasons, we assume that distributed time delay only occurs in the couplings of activators. Although it might be more interesting if both the couplings of the activators and the inhibitors involve distributed delays (see [7]), mathematical analysis tends to be much more difficult.

Based on (H1)-(H3), if we couple system (1.2) with an identical system, then the resulting four-component coupled system can be described by

$$(1.4) \quad \begin{cases} \frac{\partial u_i}{\partial t} = \frac{1}{\sigma} \Big(d_1 \Delta u_i + a - u_i - \frac{4u_i v_i}{1 + u_i^2} \Big) + k_1 \Big(\int_{-\infty}^t F(t - s) u_j(x, s) ds - u_i \Big), & \text{in } \Omega_{\infty}, \\ \frac{\partial v_i}{\partial t} = d_2 \Delta v_i + b \Big(u_i - \frac{u_i v_i}{1 + u_i^2} \Big) + k_2 (v_j - v_i), & \text{in } \Omega_{\infty}, \\ \frac{\partial u_i}{\partial u_i} = \partial v_i \end{cases}$$

$$\left(\frac{\partial u_i}{\partial n} = \frac{\partial v_i}{\partial n} = 0, \quad \text{on } \partial\Omega_{\infty},\right.$$

where $i, j \in \{1, 2\}$ and $i \neq j$; $\Omega_{\infty} := \Omega \times (0, \infty)$ and $\partial \Omega_{\infty} := \partial \Omega \times (0, \infty)$, in which $\Omega \subset \mathbf{R}^N$ $(N \geq 1)$ is the open bounded domain with a sufficient smooth boundary; n is the outer unit normal derivative across the boundary $\partial \Omega$; u_i and v_i (i = 1, 2) stand for the concentrations of the activators and the inhibitors in the reactor i, respectively; k_1 and k_2 are the inter-reactor diffusion rates of the activators and inhibitors, respectively; the kernel F(s) describes the distribution of delay effect. For our interests, we specify F(t) to be the weak kernel $\alpha e^{-\alpha t}$ with $\alpha > 0$ and $t \geq 0$. Clearly, $\int_{-\infty}^t F(t-s)ds = 1$ (see [25, 30]).

Motivated by [19, 33], by making the following change of variables

$$\epsilon = \sqrt{d_1/\sigma}, \ \tau = b\sqrt{\sigma/d_1}, \ d = d_2/b, \ \hat{t} = bt, \ \hat{k}_1 = k_1\sqrt{\sigma/d_1},$$

we can reduce system (1.4) to the following system

$$(1.5) \begin{cases} \tau \frac{\partial u_i}{\partial t} = \epsilon \Delta u_i + \frac{1}{\sigma \epsilon} \left(a - u_i - \frac{4u_i v_i}{1 + u_i^2} \right) + k_1 \left(\int_{-\infty}^t F(t - s) u_j(x, s) ds - u_i \right), & \text{in } \Omega_{\infty}, \\ \frac{\partial v_i}{\partial t} = d\Delta v_i + u_i - \frac{u_i v_i}{1 + u_i^2} + k_2 (v_j - v_i), & \text{in } \Omega_{\infty}, \end{cases}$$

$$\Delta \frac{\partial u_i}{\partial n} = \frac{\partial v_i}{\partial n} = 0, \qquad \qquad \text{on } \partial \Omega_{\infty}$$

where $i, j \in \{1, 2\}$ and $i \neq j$; we still use t (resp., k_1) to stand for \hat{t} (resp., k_1) if there is no confusion. In system (1.5), the parameter τ has the significant chemical meanings. Indeed, in the absence of coupling, if $\tau = O(1)$, u species reacts faster than v species does, and u species diffuses slower than v species does; if $\tau = O(1/\epsilon)$, u species and v species react to the same degree, but uspecies diffuses much slower than v species does; if $\tau = O(\epsilon)$, u species reacts faster than v species does, u species and v species react to the same degree (see [33] for more details).

When there is no couplings in system (1.5) (say, $k_1 = k_2 = 0$), we can obtain the following subsystem (indeed two subsystems, but they are exactly identical)

(1.6)
$$\begin{cases} \tau \frac{\partial u}{\partial t} = \epsilon \Delta u + \frac{1}{\sigma \epsilon} \left(a - u - \frac{4uv}{1 + u^2} \right), & \text{in } \Omega_{\infty}, \\ \frac{\partial v}{\partial t} = d\Delta v + u - \frac{uv}{1 + u^2}, & \text{in } \Omega_{\infty}, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & \text{on } \partial \Omega_{\infty}. \end{cases}$$

Clearly, if (U(x), V(x)) is the steady state of subsystem (1.6), then by $\int_{-\infty}^{t} F(t-s)ds = 1$, (U(x), V(x), U(x), V(x)) is the steady state of system (1.5), which is called the symmetric steady state (or reactor-based "spatially homogeneous" steady state).

In the remaining part of the paper, we shall always concentrate on the dynamics of both the coupled system (1.5) and its decoupled subsystem (1.6). For the convenience of our discussions, throughout the paper, we always assume that $\Omega = (0, \ell)$, with $\ell > 0$.

Firstly, for the two-component decoupled subsystem (1.6), it is found that when ϵ is sufficiently small and d is large enough, it can exhibit the non-constant positive steady state $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ with large amplitudes (see Theorem 2.1). Moreover, it is proved that as the parameter τ varies, the stability of $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ changes, leading to Hopf bifurcation. More precisely, there exists a $\tau_c^{\epsilon} \in (0, +\infty)$, such that $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is stable when $\tau \in (\tau_c^{\epsilon}, +\infty)$, unstable when $\tau \in (0, \tau_c^{\epsilon})$, while Hopf bifurcation occurs at $\tau = \tau_c^{\epsilon}$ (see Theorem 2.2).

Secondly, for the four-component coupled system (1.5), we are mainly interested in examining how k_1, k_2 , and α can affect the stability of the symmetric steady state $(\tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon), \tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon))$. To that end, we first assume that $\tau \in (\tau_c^{\epsilon}, +\infty)$ so that $(\tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon), \tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon))$ is stable when there are no couplings between the two reactors. Need to note that $T = 1/\alpha$ is the average delay. Thus, if $\alpha = +\infty$, then there is no time delay. If $0 < \alpha < +\infty$, then there involves time delay. Moreover, the larger (resp., smaller) α is, the smaller (resp., larger) the average time delay T is. We have the following results:

- The non-delayed coupling (α = +∞). It is proved that, roughly speaking, the first quadrant of the (k₁, k₂) parameter space can be divided into two distinct regions: one is stable region Γ^ε₂ ∪ Γ^ε₃ and the other one is the "Turing unstable" region Γ^ε₁, where Γ^ε₁ and Γ^ε₂ have the common boundary C^ε, which is indeed the primary steady state bifurcation curve (see Figure 2). More precisely, for sufficiently small ε > 0, there exists a constant ρ^{*}₀ > 0 such that if k₁ > ρ^{*}₀/2, then (ũ(x; ε), ῦ(x; ε), ũ(x; ε), ῦ(x; ε)) is always stable regardless of k₂ > 0, and that if k₁ < ρ^{*}₀/2, then there exists a k^ε₂ > 0, such that (ũ(x; ε), ῦ(x; ε), ũ(x; ε), ῦ(x; ε)) is stable when k₂ < k^ε₂, while Turing unstable when k₂ > k^ε₂. At k₂ = k^ε₂, near the symmetric solution (ũ(x; ε), ῦ(x; ε), ũ(x; ε), ῦ(x; ε)), the Turing-type steady state bifurcation occurs (see Theorem 3.3). Meanwhile, Hopf bifurcations around (ũ(x; ε), ῦ(x; ε), ũ(x; ε), ῦ(x; ε)) can never be possible. Moreover, it seems that the smaller ρ^{*}₀ is, the larger the "stable" region is, where ρ^{*}₀ depends on the diffusion rate d and the spatial interval Ω (see (A.1) in Appendix). Need to note that the special case of k₁ = 0 and k₂ > 0 has been studied by Takaishi, Mimura and Nishiura [33], and our results indeed extend the results of [33] to the general case of k₁ > 0 and k₂ > 0.
- 2. The delayed coupling $(0 < \alpha < +\infty)$. We choose α to be the main bifurcation parameter, where $T = 1/\alpha$ is the average delay. It is shown that the first quadrant of (k_1, k_2) parameter space can be re-divided into four distinct regions (see Fig. 5): Γ_1 is the unstable region (regardless of α) and Γ_{3-2} is the stable region (regardless of α), indicating that in regions Γ_1 and Γ_{3-2} , the coupled system is robust to time delay (see Theorem 4.5); $\Gamma_2 \cup \Gamma_{3-1}$ is the potential "bifurcation" region, where Hopf bifurcation may occur for some suitable α . The regions Γ_1 and Γ_2 have the common boundary C, where the bifurcation of higher co-dimension (with respect to α) might occur. More precisely, in the potential "bifurcation" region $\Gamma_2 \cup$

 Γ_{3-1} , under the conditions of Theorem 4.12, there exists a critical value $\alpha_H^{\epsilon} > 0$ depending on ϵ , such that the symmetric solution $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is stable when $\alpha \in (\alpha_H^{\epsilon}, \infty)$, while unstable when α is slightly less than α_H^{ϵ} ; at $\alpha = \alpha_H^{\epsilon}$, system (1.5) undergoes a Hopf bifurcation near the symmetric solution.

3. Similarities and differences (between $\alpha = +\infty$ and $0 < \alpha < +\infty$). As $\epsilon \to 0^+$, $\Gamma_1^{\epsilon} \to \Gamma_1$, $\Gamma_2^{\epsilon} \to \Gamma_2$ and $\mathcal{C}^{\epsilon} \to \mathcal{C}$. For sufficiently small $\epsilon > 0$, $\Gamma_3^{\epsilon} \equiv \Gamma_{3-1} \cup \Gamma_{3-2}$. When $\epsilon > 0$ is sufficiently small, there exists at least a subregion $\Gamma_1^{\epsilon,u}$ of Γ_1^{ϵ} such that in this subregion $(\widetilde{u}(x;\epsilon), \widetilde{v}(x;\epsilon), \widetilde{u}(x;\epsilon), \widetilde{v}(x;\epsilon))$ is always unstable in both $\alpha = +\infty$ and $0 < \alpha < +\infty$, showing that in such case time delay can never change the stability of the steady state; there also exists at least a subregion $\Gamma_2^{\epsilon,u}$ of $\Gamma_2^{\epsilon} \cup \Gamma_{3-2}$ such that in this subregion $(\widetilde{u}(x;\epsilon), \widetilde{v}(x;\epsilon), \widetilde{u}(x;\epsilon), \widetilde{v}(x;\epsilon))$ is stable when $\alpha = +\infty$, but unstable for some $\alpha \in (0, +\infty)$, implying that in such case, time delay does change the stability. In the subregion Γ_{3-2} of Γ_3^{ϵ} , $(\widetilde{u}(x;\epsilon), \widetilde{v}(x;\epsilon), \widetilde{u}(x;\epsilon), \widetilde{v}(x;\epsilon))$ is always stable in both $\alpha = +\infty$ and $0 < \alpha < +\infty$, showing again that in such case time delay can never change the stability of the steady state.

Finally, our main novelties are listed in the following way: for system (1.6), it is firstly reported that it can exhibit Hopf bifurcation branching from non-constant steady state. For system (1.5), it is the first to report Turing instability of its non-constant steady state. A different scenery is that in our model there are two different kinds of diffusions: intra-reactor diffusions (ϵ and d) and inter-reactor diffusions (k_1 and k_2). Our Turing instability results are different from previous works ([12, 9, 16, 17, 18, 22, 5, 25, 27, 28, 29, 30, 34]). Moreover, our analysis for system (1.5) follows the lines of [21], but we consider the problem in a quite different setting (4-component coupled system with distributed delays). We show that the basic framework of [21] works for the coupled system (1.5), but the generalization is non-trivial as analysis for the eigenvalue problems is difficult.

The present paper is organized as follows. In §2, for system (1.6), we study the existence and stability of positive non-constant steady state $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$. In particular, we show the existence of Hopf bifurcation branching from the non-constant steady state solution. In §3 and §4, we study the stability and instability of $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ with respect to system (1.5) in the case of $\alpha = +\infty$ and $\alpha \in (0, +\infty)$, respectively. In §5, we end up our investigations by drawing some conclusions. In Appendix, we list some well-known results on the spectral properties of the elliptic operator.

2 Dynamics of the decoupled subsystem (1.6): one reactor

In this section, we consider the dynamics of subsystem (1.6). For convenience, we copy (1.6) here:

(2.1)
$$\begin{cases} \tau \frac{\partial u}{\partial t} = \epsilon u_{xx} + \frac{1}{\epsilon \sigma} \left(a - u - \frac{4uv}{1 + u^2} \right), & x \in (0, \ell), \ t > 0, \\ \frac{\partial v}{\partial t} = dv_{xx} + u - \frac{uv}{1 + u^2}, & x \in (0, \ell), \ t > 0, \\ u_x(0, t) = u_x(\ell, t) = v_x(0, t) = v_x(\ell, t) = 0, & t > 0. \end{cases}$$

Clearly, system (2.1) has a unique constant steady state solution $(u_*, v_*) := \left(\frac{a}{5}, 1 + \frac{a^2}{25}\right)$. The linearized operator of system (2.1) evaluated at (u_*, v_*) is given by $L := [L_{ij}]_{2\times 2}$, where

$$L_{11} := \frac{3a^2 - 125}{\sigma\epsilon\tau(a^2 + 25)}, \ L_{12} := -\frac{20a}{\sigma\epsilon\tau(a^2 + 25)}, \ L_{21} := \frac{2a^2}{a^2 + 25}, \ L_{22} := -\frac{5a}{a^2 + 25}.$$

Assume that $a > \frac{5}{3}\sqrt{15}$ so that system (2.1) is an activator-inhibitor system in the sense that $L_{11} > 0, L_{12} < 0, L_{21} > 0$ and $L_{22} < 0$.

The steady state problem of (2.1) is governed by the following elliptic equations

(2.2)
$$\begin{cases} \epsilon^2 u_{xx} + \frac{1}{\sigma} \left(a - u - \frac{4uv}{1 + u^2} \right) = 0, & x \in (0, \ell), \\ dv_{xx} + u - \frac{uv}{1 + u^2} = 0, & x \in (0, \ell), \\ u'(0) = u'(\ell) = v'(0) = v'(\ell) = 0. \end{cases}$$

We denote $C^{2}_{\epsilon}\left([0,\ell]\right)$ by the subspace of $C^{2}\left([0,\ell]\right)$ with norm

$$\|u\|_{C^2_{\epsilon}([0,\ell])} := \sum_{k=0}^2 \max_{x \in [0,\ell]} \left| \left(\epsilon \frac{d}{dx} \right)^k u\left(x\right) \right|.$$

Then, we have the following results on the existence of positive singularly perturbed steady state solutions of system (2.1) for small $\epsilon > 0$.

Theorem 2.1. Suppose that $a > \frac{5}{3}\sqrt{15}$ holds. Then, the following conclusions hold true:

- 1. There exist positive constants d_0 and ϵ_0 , such that for any $0 < \epsilon < \epsilon_0$ and $d > d_0$, system (2.1) has an ϵ -family of positive non-constant steady state solutions $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon)) \in C^2_{\epsilon}([0,\ell]) \times C^2([0,\ell]).$
- 2. Moreover, for any $\kappa > 0$ and $I_{\kappa} := (x^* \kappa, x^* + \kappa)$,
 - (2.3) $\lim_{\epsilon \to 0^+} \widetilde{u}(x;\epsilon) = U^*(x) \text{ uniformly on } (0,\ell) \setminus I_{\kappa}, \lim_{\epsilon \to 0^+} \widetilde{v}(x;\epsilon) = V^*(x) \text{ uniformly on } (0,\ell),$

where $x^* \in (0, \ell)$ is the layer position, and $(U^*(x), V^*(x))$ is the discontinuous solution of the reduced system ($\epsilon = 0$) of (2.1), which is defined in (2.8).

Proof. Our proof falls into the general framework of [21]. To prove the desired results, we need to verify (A0)-(A5) stated in [21] (pp. 484-485).

Step 1. To show that there is an intersection of

$$f(u,v) := \frac{1}{\sigma} \left(a - u - \frac{4uv}{1 + u^2} \right) = 0$$
, and $g(u,v) := u - \frac{uv}{1 + u^2} = 0$.

This is obvious since the unique intersection point is (u_*, v_*) .



Figure 1: Nullclines of u and v: f(u, v) = 0 and g(u, v) = 0.

Step 2. To check that: f(u, v) = 0 is sigmoidal and it consists of three continuous curves $u = h_{-}(v)$, $u = h_{0}(v)$ and $u = h_{+}(v)$ which are defined in the intervals I_{-} , I_{0} , I_{+} , respectively, such that $h_{-}(v) < h_{0}(v) < h_{+}(v)$ holds for $v \in (\underline{v}, \overline{v})$ and $h_{+}(v)$ (resp., $h_{-}(v)$) coincides with $h_{0}(v)$ at only one point $v = \overline{v}$ (resp., $v = \underline{v}$), where $\underline{v} := \min I_{-}, \overline{v} := \max I_{+}$.

Indeed, by Lemmas 4.1 and 4.2 of [8], if $a > \frac{5}{3}\sqrt{15}$, then f(u, v) = 0 is sigmoidal and it consists of three continuous curves $u = h_{-}(v)$, $u = h_{0}(v)$, $u = h_{+}(v)$ (see Fig. 1). Moreover, by (4.9) and (4.10) of [8], we have

$$\max\left\{1,\frac{8}{a}\right\} < \underline{u} := h_{-}(\underline{v}) < \frac{a}{5}, \ \frac{2a}{5} < \overline{u} := h_{+}(\overline{v}) < \frac{a}{2},$$
$$\max\left\{2, 1 + \frac{64}{a^{2}}\right\} < \underline{v} < 1 + \frac{a^{2}}{25} < \overline{v} < \frac{a^{2}}{12} - \frac{1}{4}.$$

Step 3. To prove that M(v) = 0 has a unique root \hat{v} in $(\underline{v}, \overline{v})$ and that $M'(\hat{v}) < 0$, where

$$M(v) := \int_{h_{-}(v)}^{h_{+}(v)} f(s,v) ds = \frac{1}{\sigma} \int_{h_{-}(v)}^{h_{+}(v)} \left(a - s - \frac{4sv}{1 + s^{2}}\right) ds.$$

Indeed, for any u, v > 0, $f_v(u, v) < 0$. Thus, in the first quadrant, f(u, v) < 0 in the region above f(u, v) = 0; while f(u, v) > 0 in the region below f(u, v) = 0. Thereby, we have $M(\underline{v}) > 0$ and $M(\overline{v}) < 0$. Moreover, by $f(h_+(v), v) = f(h_-(v), v) = 0$, we have

$$M'(v) = -\frac{4}{\sigma} \int_{h_{-}(v)}^{h_{+}(v)} \frac{s}{1+s^{2}} ds < 0,$$

implying that M(v) = 0 has a unique root \hat{v} in $(\underline{v}, \overline{v})$ and $M'(\hat{v}) < 0$.

Step 4. To verify that for any $(u, v) \in \mathscr{R}_+ \cup \mathscr{R}_-$, $f_u(u, v) < 0$, where

$$\mathscr{R}_+ := \big\{ (u,v) | u = h_+(v) \text{ for } \widehat{v} \le v < \overline{v} \big\}, \ \mathscr{R}_- := \big\{ (u,v) | u = h_-(v) \text{ for } \underline{v} \le v < \widehat{v} \big\}.$$

In fact, for any u, v > 0, $f_v(u, v) < 0$. Then, in the left (resp., right) side of \mathscr{R}_{\pm} , we have f(u, v) > 0 (resp., < 0). Thus, $f_u(u, v)|_{\mathscr{R}_+} < 0$.

Step 5. To show that

$$(2.4) g(u,v)\big|_{(u,v)\in\mathscr{R}_-} < 0 < g(u,v)\big|_{(u,v)\in\mathscr{R}_+}, \ \det\left[\frac{\partial(f,g)}{\partial(u,v)}\right]\Big|_{(u,v)\in\mathscr{R}_\pm} > 0.$$

Indeed, for any u, v > 0, $g_v(u, v) < 0$. Thus, one can verify that $g(u, v)|_{\mathscr{R}_-} < 0 < g(u, v)|_{\mathscr{R}_+}$, since the intersection point (u_*, v_*) lies on the middle branch of f(u, v) = 0. Moreover,

$$\frac{\partial(f,g)}{\partial(u,v)}\Big|_{(u,v)\in\mathscr{R}_{\pm}} = \frac{5u}{\sigma(1+u^2)^3}\Big|_{(u,v)\in\mathscr{R}_{\pm}} > 0.$$

So far, we have verified (A0)-(A5) in [21]. Then, by [21], we can conclude the existence of the positive non-constant steady state solution $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ for $0 < \epsilon < \epsilon_0$ and $d > d_0$ with $\epsilon_0 > 0$ and $d_0 > 0$.

In case of $\epsilon = 0$, the elliptic system (2.2) is reduced to

(2.5)
$$\begin{cases} a - u - \frac{4uv}{1 + u^2} = 0, & x \in (0, \ell), \\ dv_{xx} + u - \frac{uv}{1 + u^2} = 0, & x \in (0, \ell), \\ u'(0) = u'(\ell) = v'(0) = v'(\ell) = 0. \end{cases}$$

Solving u from the first equation of (2.5), we have $u = h_i(v)$ for i = 0, +, -. Substituting $u = h_i(v)$ into the second equation of (2.5), we have (see [19])

$$\begin{cases} dv_{xx} + G_i(v) = 0, & x \in (0, \ell), \\ v'(0) = v'(\ell) = 0, \end{cases}$$

where, for i = 0, +, -,

$$G_i(v) := h_i(v) - \frac{h_i(v)v}{1 + h_i(v)^2}$$

Choosing $\hat{v} \in I_- \cup I_+$, given by Step 3, as the separating point, we can combine $G_{\pm}(v)$ by

$$G(v; \widehat{v}) = \begin{cases} G_-(v), & v \in I_- \cup \{v < \widehat{v}\}, \\ G_+(v), & v \in I_+ \cup \{v > \widehat{v}\}, \end{cases}$$

which is discontinuous at \hat{v} of the first kind.

As in [19, 20, 21], there exists a $x^* \in I$, such that (2.5) can be decomposed by the following way:

(2.6)
$$\begin{cases} dV_{xx} + h_{-}(V) - \frac{h_{-}(V)V}{1 + h_{-}(V)^{2}} = 0, \ x \in (0, x^{*}), \\ V'(0) = 0, \ V(x^{*}) = \widehat{v}, \end{cases}$$

and

(2.7)
$$\begin{cases} dV_{xx} + h_+(V) - \frac{h_+(V)V}{1 + h_+(V)^2} = 0, \ x \in (x^*, \ell), \\ V(x^*) = \widehat{v}, \ V'(\ell) = 0, \end{cases}$$

with C^1 matching condition: $\frac{dV^-}{dx}(x^*) = \frac{dV^+}{dx}(x^*)$, where V^- and V^+ are the solutions of (2.6) and (2.7), respectively.

Then, $(U^*(x), V^*(x))$ is the weak solution of system (2.5), where

(2.8)
$$V^*(x) = \begin{cases} V^-, \ x \in (0, x^*), \\ V^+, \ x \in (x^*, \ell). \end{cases}, \ U^*(x) = \begin{cases} h_-(V^-), \ x \in (0, x^*), \\ h_+(V^+), \ x \in (x^*, \ell). \end{cases}$$

Finally, the asymptotic behaviors in terms of (2.3) can be obtained by Appendix 1 in [20] and Theorem 1.1 in [21]. For more details, we refer readers to [19, 20, 21].

Next, we concentrate on the stability/instability of $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ for $\epsilon \in (0, \epsilon_0)$. We have the following results.

Theorem 2.2. Suppose that $a > \frac{5}{3}\sqrt{15}$ holds so that $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ exists for any $\epsilon \in (0, \epsilon_0)$. Then, there exists a $\tau_c^{\epsilon} > 0$, such that $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is locally asymptotically stable with respect to system (2.1) when $\tau > \tau_c^{\epsilon}$, while unstable when $\tau < \tau_c^{\epsilon}$. In particular, at $\tau = \tau_c^{\epsilon}$, Hopf bifurcation occurs around $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$.

Proof. Our results are really the direct consequences of Theorem 3.1 in [21] since all the conditions (A0)-(A5) in [21] are satisfied. For the convenience of our later discussions in next sections, we list the sketch of the proof below. The linearized system of (2.1) evaluated at $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is given by

(2.9)
$$\begin{cases} \epsilon \tau \frac{\partial u}{\partial t} = \epsilon^2 u_{xx} + f_u^{\epsilon} u + f_v^{\epsilon} v, & x \in (0, \ell), \ t > 0, \\ \frac{\partial v}{\partial t} = dv_{xx} + g_u^{\epsilon} u + g_v^{\epsilon} v, & x \in (0, \ell), \ t > 0, \\ u_x(x, t) = v_x(x, t) = 0, & x = 0, \ell, \ t > 0, \end{cases}$$

where

(2.10)
$$f_{u}^{\epsilon} := -\frac{1}{\sigma} \left(1 + \frac{4(1 - \widetilde{u}^{2}(x;\epsilon))\widetilde{v}(x;\epsilon)}{(1 + \widetilde{u}^{2}(x;\epsilon))^{2}} \right),$$
$$f_{v}^{\epsilon} := -\frac{4\widetilde{u}(x;\epsilon)}{\sigma(1 + \widetilde{u}^{2}(x;\epsilon))}, \ g_{u}^{\epsilon} := 1 - \frac{(1 - \widetilde{u}^{2}(x;\epsilon))\widetilde{v}(x;\epsilon)}{(1 + \widetilde{u}^{2}(x;\epsilon))^{2}}, \ g_{v}^{\epsilon} := -\frac{\widetilde{u}(x;\epsilon)}{1 + \widetilde{u}^{2}(x;\epsilon)}.$$

The eigenvalue problem associated with (2.9) is given by

(2.11)
$$\begin{cases} \begin{pmatrix} \epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} & f_v^{\epsilon} \\ g_u^{\epsilon} & d\frac{d^2}{dx^2} + g_v^{\epsilon} \end{pmatrix} \begin{pmatrix} \phi \\ \psi \end{pmatrix} = \lambda \begin{pmatrix} \epsilon \tau & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \phi \\ \psi \end{pmatrix}, \ x \in (0, \ell), \\ \phi'(0) = \phi'(\ell) = \psi'(0) = \psi'(\ell) = 0, \end{cases}$$

where $(\phi, \psi)^T \in (H^2(0, \ell) \cap H^1_N(0, \ell)) \times (H^2(0, \ell) \cap H^1_N(0, \ell))$, and

$$H^1_N(0,\ell) = \left\{ \phi \in H^1(0,\ell) | \phi'(0) = \phi'(\ell) = 0 \right\}.$$

By Theorem 3.1 in [21], it follows that there exists a unique $\tau_c^{\epsilon} > 0$ such that for any $\tau \in (\tau_c^{\epsilon}, \infty)$, all the eigenvalues of the eigenvalue problem (2.11) have strictly negative real parts, while for any $\tau \in (0, \tau_c^{\epsilon})$, the eigenvalue problem (2.11) has at least one eigenvalue with positive real part. In particular, at $\tau = \tau_c^{\epsilon}$, the eigenvalue problem (2.11) has a pair of simple purely imaginary eigenvalues and the transversality condition holds. So far, based on these observations, we can complete the proof.

3 Dynamics of the coupled system (1.5): the case of $\alpha \to +\infty$

In this section, we shall consider the following non-delayed system

$$(3.1) \qquad \begin{cases} \tau \frac{\partial u_i}{\partial t} = \epsilon \frac{\partial^2 u_i}{\partial x^2} + \frac{1}{\sigma \epsilon} \left(a - u_i - \frac{4u_i v_i}{1 + u_i^2} \right) + k_1 (u_j - u_i), & x \in (0, \ell), \ t > 0, \\ \frac{\partial v_i}{\partial t} = d \frac{\partial^2 v_i}{\partial x^2} + u_i - \frac{u_i v_i}{1 + u_i^2} + k_2 (v_j - v_i), & x \in (0, \ell), \ t > 0, \\ \frac{\partial u_i}{\partial x} = \frac{\partial v_i}{\partial x} = 0, & x = 0, \ell, \ t > 0, \end{cases}$$

where $i, j \in \{1, 2\}$ and $i \neq j$; $\epsilon \in (0, \epsilon_0)$, in which $\epsilon_0 > 0$ is stated in Theorems 2.1 and 2.2. Clearly, $(u_1, v_1, u_2, v_2) = (\tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon), \tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon))$ is the symmetric steady state solution of system (3.1), where $\tilde{u}(x; \epsilon)$ and $\tilde{v}(x; \epsilon)$ are stated in Theorem 2.1.

We would like to mention that the non-delayed system (3.1) is indeed the limiting system of system (1.5) when $\alpha \to +\infty$. In fact, we can extend the domain of the weak kernel F(t) to $(-\infty, +\infty)$ by defining

$$\widetilde{F}(t) := \begin{cases} F(t) = \alpha e^{-\alpha t}, & t \ge 0, \\ 0, & t < 0. \end{cases}$$

Then, we have

$$\int_{-\infty}^{+\infty} \widetilde{F}(s)ds = \int_{0}^{+\infty} F(s)ds = 1, \lim_{\alpha \to +\infty} \widetilde{F}(t) = \delta(t) = \begin{cases} +\infty, & \text{if } t = 0\\ 0, & \text{if } t \neq 0 \end{cases}$$

Therefore, as $\alpha \to +\infty$, for i = 1, 2, we have

$$\int_{-\infty}^{t} F(t-s)u_i(x,s)ds = \int_{-\infty}^{\infty} \widetilde{F}(t-s)u_i(x,s)ds \to \int_{-\infty}^{\infty} \delta(s-t)u_i(x,s)ds = u_i(x,t),$$

where we use the fact that $\delta(t-s) = \delta(s-t)$. Thus, as $\alpha \to +\infty$, the limiting system of system (1.5) is indeed the non-delayed system (3.1).

The linearized system of (3.1) evaluated at $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is given by

$$(3.2) \qquad \begin{cases} \tau \frac{\partial \phi_i}{\partial t} = \epsilon \frac{\partial^2 \phi_i}{\partial x^2} + \frac{1}{\epsilon} \left(f_u^{\epsilon} \phi_i + f_v^{\epsilon} \psi_i \right) + k_1 (\phi_j - \phi_i), & x \in (0, \ell), \ t > 0, \\ \frac{\partial \psi_i}{\partial t} = d \frac{\partial^2 \psi_i}{\partial x^2} + g_u^{\epsilon} \phi_i + g_v^{\epsilon} \psi_i + k_2 (\psi_j - \psi_i), & x \in (0, \ell), \ t > 0, \\ \frac{\partial \phi_i}{\partial x} = \frac{\partial \psi_i}{\partial x} = 0, & x = 0, \ell, \ t > 0, \end{cases}$$

where $i, j \in \{1, 2\}$ and $i \neq j$, and $f_u^{\epsilon}, f_v^{\epsilon}, g_u^{\epsilon}, g_v^{\epsilon}$ are precisely defined in (2.10).

For i = 1, 2, rewrite $\phi_i(x, t)$ and $\psi_i(x, t)$ in the following form (separation of variables)

$$\phi_1(x,t) := w_1(x)e^{\lambda t}, \ \phi_2(x,t) := w_2(x)e^{\lambda t}, \ \psi_1(x,t) := z_1(x)e^{\lambda t}, \ \psi_2(x,t) := z_2(x)e^{\lambda t},$$

where λ is to be determined later.

Following [12, 33], by introducing

(3.3)
$$w_s(x) := \frac{1}{2}(w_1(x) + w_2(x)), \ w_a(x) := \frac{1}{2}(w_1(x) - w_2(x)),$$
$$z_s(x) := \frac{1}{2}(z_1(x) + z_2(x)), \quad z_a(x) := \frac{1}{2}(z_1(x) - z_2(x)),$$

we can reduce the linearized system (3.2) to the following two eigenvalue problems

(3.4)
$$\begin{cases} \epsilon^2 w_s''(x) + f_u^{\epsilon} w_s(x) + f_v^{\epsilon} z_s(x) = \epsilon \tau \lambda w_s(x), & x \in (0, \ell), \\ dz_s''(x) + g_u^{\epsilon} w_s(x) + g_v^{\epsilon} z_s(x) = \lambda z_s(x), & x \in (0, \ell), \\ w_s'(0) = w_s'(\ell) = z_s'(0) = z_s'(\ell) = 0, \end{cases}$$

and

(3.5)
$$\begin{cases} \epsilon^2 w_a''(x) + (f_u^{\epsilon} - 2\epsilon k_1) w_a(x) + f_v^{\epsilon} z_a(x) = \epsilon \tau \lambda w_a(x), & x \in (0, \ell), \\ d z_a''(x) + g_u^{\epsilon} w_a(x) + (g_v^{\epsilon} - 2k_2) z_a(x) = \lambda z_a(x), & x \in (0, \ell), \\ w_a'(0) = w_a'(\ell) = z_a'(\ell) = z_a'(\ell) = 0. \end{cases}$$

Clearly, the eigenvalue problem (3.4) is exactly (2.11) in Section 2, and the distributions of its eigenvalues are clear (see the proof of Theorem 2.2 in this paper, or Theorem 3.1 in [21]). It remains to consider the eigenvalue problem (3.5).

Following [20, 21], by rewriting

$$\lambda =: \lambda_R + i\lambda_I, \ w_a =: w_R + iw_I, \ z_a =: z_R + iz_I,$$

we can reduce the eigenvalue problem (3.5) to the following equations

$$(3.6) \qquad \begin{cases} (\epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} - 2\epsilon k_1)w_R + f_v^{\epsilon} z_R = \epsilon \tau (\lambda_R w_R - \lambda_I w_I), \\ (\epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} - 2\epsilon k_1)w_I + f_v^{\epsilon} z_I = \epsilon \tau (\lambda_R w_I + \lambda_I w_R), \\ (d \frac{d^2}{dx^2} + g_v^{\epsilon} - 2k_2)z_R + g_u^{\epsilon} w_R = \lambda_R z_R - \lambda_I z_I, \\ (d \frac{d^2}{dx^2} + g_v^{\epsilon} - 2k_2)z_I + g_u^{\epsilon} w_I = \lambda_I z_R + \lambda_R z_I, \end{cases}$$

subject to homogeneous Neumann boundary conditions.

Define

$$\mathcal{L}^{\epsilon} := \epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} - 2\epsilon k_1.$$

Choose μ^* satisfying

(3.7)
$$0 < \mu^* < \inf_{x \in (0,\ell)} \frac{f_u^* g_v^* - f_v^* g_u^*}{-f_u^*}$$

where

$$f_u^* := f_u(U^*, V^*), \ f_v^* := f_v(U^*, V^*), \ g_u^* := g_u(U^*, V^*), \ g_v^* := g_v(U^*, V^*),$$

in which $(U^*, V^*) := (U^*(x), V^*(x))$ is stated in Theorem 2.1, and $(f_u^* g_v^* - f_v^* g_u^*)/(-f_u^*) > 0$ (see the proof of Theorem 2.1).

For sufficiently small $\epsilon > 0$, by the similar argument of Lemma 2.1 in [20] and Remark 2.2 in [21], $(\mathcal{L}^{\epsilon} - \epsilon \tau \lambda)^{-1}$ is well-defined for Re $\lambda > -\mu^*$. In the following, we always assume that Re $\lambda > -\mu^*$. In case of Re $\lambda \leq -\mu^*$, all the eigenvalues of (3.2) have negative real parts, which do not affect the stability of $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$.

Then, by solving the first two equations of (3.6) with respect to w_R and w_I , we have

$$w_{R} = \left[I + (\epsilon\tau\lambda_{I})^{2} (\mathcal{L}^{\epsilon} - \epsilon\tau\lambda_{R})^{-2}\right]^{-1} \left[\epsilon\tau\lambda_{I} (\mathcal{L}^{\epsilon} - \epsilon\tau\lambda_{R})^{-2} (f_{v}^{\epsilon}z_{I}) - (\mathcal{L}^{\epsilon} - \epsilon\tau\lambda_{R})^{-1} (f_{v}^{\epsilon}z_{R})\right],$$

$$w_{I} = -\left[I + (\epsilon\tau\lambda_{I})^{2} (\mathcal{L}^{\epsilon} - \epsilon\tau\lambda_{R})^{-2}\right]^{-1} \left[(\mathcal{L}^{\epsilon} - \epsilon\tau\lambda_{R})^{-1} (f_{v}^{\epsilon}z_{I}) + \epsilon\tau\lambda_{I} (\mathcal{L}^{\epsilon} - \epsilon\tau\lambda_{R})^{-2} (f_{v}^{\epsilon}z_{R})\right].$$

where I is the identity operator.

By Lemma A.1 (Appendix), the eigenvalue problem

$$\begin{cases} \epsilon^2 \frac{d^2 \phi}{dx^2} + f_u^{\epsilon} \phi = \mu \phi, \ x \in (0, \ell), \\ \phi'(0) = \phi'(\ell) = 0, \end{cases}$$

has complete orthonormal pairs (in $L^2(0, \ell)$ sense) of eigenvalues and eigenfunctions $\{(\mu_n^{\epsilon}, \phi_n^{\epsilon}(x))\}_{n=0}^{\infty}$.

By eigenfunction expansions, we have

(3.8)
$$w_R = \sum_{n \ge 0} w_R^n \phi_n^{\epsilon} = w_R^0 \phi_0^{\epsilon} + w_R^{\dagger}, \ w_I = \sum_{n \ge 0} w_I^n \phi_n^{\epsilon} = w_I^0 \phi_0^{\epsilon} + w_I^{\dagger},$$

where $w_R^{\dagger} := \sum_{n \ge 1} w_R^n \phi_n^{\epsilon}, \, w_I^{\dagger} := \sum_{n \ge 1} w_I^n \phi_n^{\epsilon}$, in which

$$w_R^n = \frac{-\mu_n^{\epsilon} + 2\epsilon k_1 + \epsilon\tau\lambda_R}{(\mu_n^{\epsilon} - 2\epsilon k_1 - \epsilon\tau\lambda_R)^2 + (\epsilon\tau\lambda_I)^2} \langle f_v^{\varepsilon} z_R, \phi_n^{\varepsilon} \rangle + \frac{\epsilon\tau\lambda_I}{(\mu_n^{\epsilon} - 2\epsilon k_1 - \epsilon\tau\lambda_R)^2 + (\epsilon\tau\lambda_I)^2} \langle f_v^{\varepsilon} z_I, \phi_n^{\varepsilon} \rangle,$$

$$w_I^n = \frac{-\mu_n^{\epsilon} + 2\epsilon k_1 + \epsilon\tau\lambda_R}{(\mu_n^{\epsilon} - 2\epsilon k_1 - \epsilon\tau\lambda_R)^2 + (\epsilon\tau\lambda_I)^2} \langle f_v^{\varepsilon} z_I, \phi_n^{\varepsilon} \rangle - \frac{\epsilon\tau\lambda_I}{(\mu_n^{\epsilon} - 2\epsilon k_1 - \epsilon\tau\lambda_R)^2 + (\epsilon\tau\lambda_I)^2} \langle f_v^{\varepsilon} z_R, \phi_n^{\varepsilon} \rangle.$$

Hereafter, $\langle\cdot,\cdot\rangle$ denotes the inner product in $L^2(0,\ell)$ sense.

By (3.8), we have

$$w_R^{\dagger} = \left[I + (\epsilon\tau\lambda_I)^2 (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{-2}\right]^{\dagger} \left[\epsilon\tau\lambda_I (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{2\dagger} (f_v^{\epsilon}z_I) - (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{\dagger} (f_v^{\epsilon}z_R)\right],$$

$$w_I^{\dagger} = -\left[I + (\epsilon\tau\lambda_I)^2 (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{-2}\right]^{\dagger} \left[(\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{\dagger} (f_v^{\epsilon}z_I) + \epsilon\tau\lambda_I (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{2\dagger} (f_v^{\epsilon}z_R)\right],$$

where

$$(\mathcal{L}_{1}^{\epsilon} - \epsilon \tau \lambda_{R})^{\dagger} = \sum_{n \ge 1} \frac{\langle \cdot, \phi_{n}^{\epsilon} \rangle}{\mu_{n}^{\epsilon} - 2\epsilon k_{1} - \epsilon \tau \lambda_{R}} \phi_{n}^{\epsilon}, \quad (\mathcal{L}_{1}^{\epsilon} - \epsilon \tau \lambda_{R})^{2\dagger} = \sum_{n \ge 1} \frac{\langle \cdot, \phi_{n}^{\epsilon} \rangle}{(\mu_{n}^{\epsilon} - 2\epsilon k_{1} - \epsilon \tau \lambda_{R})^{2}} \phi_{n}^{\epsilon};$$

and

$$\left[I + (\epsilon \tau \lambda_I)^2 (\mathcal{L}_1^{\epsilon} - \epsilon \tau \lambda_R)^{-2}\right]^{\dagger} = \sum_{n \ge 1} \frac{(\mu_n^{\epsilon} - 2\epsilon k_1 - \epsilon \tau \lambda_R)^2}{(\mu_n^{\epsilon} - 2\epsilon k_1 - \epsilon \tau \lambda_R)^2 + (\epsilon \tau \lambda_I)^2} \langle \cdot, \phi_n^{\epsilon} \rangle \phi_n^{\epsilon}.$$

Substituting (3.8) into the last two equations of (3.6), we have

(3.9)
$$\begin{pmatrix} T_{\lambda_R}^{\epsilon} & -\lambda_I + \epsilon \tau \lambda_I S^{\epsilon} \\ \lambda_I + \epsilon \tau \lambda_I S^{\epsilon} & T_{\lambda_R}^{\epsilon} \end{pmatrix} \begin{pmatrix} z_R \\ z_I \end{pmatrix} = -\frac{g_u^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\varepsilon}} M^{\varepsilon} \begin{pmatrix} \langle z_R, \frac{f_v^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\varepsilon}} \rangle \\ \langle z_I, \frac{f_v^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\varepsilon}} \rangle \end{pmatrix},$$

where

$$(3.10) \quad \begin{cases} T_{\lambda_R}^{\epsilon} := -d\frac{d^2}{dx^2} - g_v^{\epsilon} + 2k_2 + \lambda_R + g_u^{\epsilon} \left[I + (\epsilon\tau\lambda_I)^2 (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{-2}\right]^{\dagger} (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{\dagger} f_v^{\epsilon}, \\ S^{\epsilon} := -g_u^{\epsilon} \left[I + (\epsilon\tau\lambda_I)^2 (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{-2}\right]^{\dagger} (\mathcal{L}_1^{\epsilon} - \epsilon\tau\lambda_R)^{2\dagger} f_v^{\epsilon}, \\ M^{\epsilon} := \begin{pmatrix} \frac{\rho(\epsilon) - 2k_1 - \tau\lambda_R}{(\rho(\epsilon) - 2k_1 - \tau\lambda_R)^2 + (\tau\lambda_I)^2} & -\frac{\tau\lambda_I}{(\rho(\epsilon) - 2k_1 - \tau\lambda_R)^2 + (\tau\lambda_I)^2} \\ \frac{\tau\lambda_I}{(\rho(\epsilon) - 2k_1 - \tau\lambda_R)^2 + (\tau\lambda_I)^2} & \frac{\rho(\epsilon) - 2k_1 - \tau\lambda_R}{(\rho(\epsilon) - 2k_1 - \tau\lambda_R)^2 + (\tau\lambda_I)^2} \end{pmatrix}. \end{cases}$$

By Lemma A.1 (Appendix), we have

$$\lim_{\epsilon \to 0^+} \frac{f_v^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}} = -c_1^* \delta(x - x^*), \ \lim_{\epsilon \to 0^+} \frac{g_u^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}} = c_2^* \delta(x - x^*), \ \text{in } H^{-1} \text{ sense},$$

where

$$(3.11) \quad c_1^* := -\kappa^* M'(\widehat{v}) > 0, \ c_2^* := \kappa^* \left[h_+(\widehat{v}) - h_-(\widehat{v}) + \left(\frac{h_-(\widehat{v})}{1 + (h_-(\widehat{v}))^2} - \frac{h_+(\widehat{v})}{1 + (h_+(\widehat{v}))^2} \right) \widehat{v} \right] > 0.$$

Set $\lim_{\epsilon \to 0^+} \lambda =: \lambda^* =: \lambda^*_R + i\lambda^*_I$ and $\lim_{\epsilon \to 0^+} z =: z^* =: z^*_R + iz^*_I$. Then, as $\epsilon \to 0^+$, (3.9) can be reduced to

(3.12)
$$\begin{pmatrix} T_{\lambda_R^*} & -\lambda_I^* \\ \lambda_I^* & T_{\lambda_R^*} \end{pmatrix} \begin{pmatrix} z_R^* \\ z_I^* \end{pmatrix} = c_1^* c_2^* \delta(x - x^*) M^* \begin{pmatrix} \langle z_R^*, \delta(x - x^*) \rangle \\ \langle z_I^*, \delta(x - x^*) \rangle \end{pmatrix}, \text{ in } H^{-1} \text{ sense,}$$

where

$$\begin{pmatrix} T_{\lambda_R^*} := -d\frac{d^2}{dx^2} - \frac{f_u^* g_v^* - f_v^* g_u^*}{f_u^*} + \lambda_R^* + 2k_2, \\ M^* := \begin{pmatrix} \frac{\rho_0^* - 2k_1 - \tau \lambda_R^*}{(\rho_0^* - 2k_1 - \tau \lambda_R^*)^2 + (\tau \lambda_I^*)^2} & -\frac{\tau \lambda_I^*}{(\rho_0^* - 2k_1 - \tau \lambda_R^*)^2 + (\tau \lambda_I^*)^2} \\ \frac{\tau \lambda_I^*}{(\rho_0^* - 2k_1 - \tau \lambda_R^*)^2 + (\tau \lambda_I^*)^2} & \frac{\rho_0^* - 2k_1 - \tau \lambda_R^*}{(\rho_0^* - 2k_1 - \tau \lambda_R^*)^2 + (\tau \lambda_I^*)^2} \end{pmatrix}.$$

If for any given $h \in H^{-1}(I)$, z is uniquely determined by

$$\langle T^{\epsilon}_{\lambda_R} z, \psi \rangle = \langle h, \psi \rangle \quad (\forall \psi \in H^1_N(I)),$$

then we define the generalized inverse operator of $T_{\lambda_R}^{\epsilon}$ as mapping $(T_{\lambda_R}^{\epsilon})^{-1} : H^{-1}(I) \to H_N^1(I)$ such that $(T_{\lambda_R}^{\epsilon})^{-1}h = z$. The other generalized operators in this paper are defined similarly.

By [20, 21, 33], $T_{\lambda_R^*}$ has a generalized inverse operator $T_{\lambda_R^*}^{-1}$ when $\lambda_R^* + 2k_2 > -\mu^*$. In the following, we always treat the case when $\lambda_R^* + 2k_2 > -\mu^*$. In case of $\lambda_R^* + 2k_2 \leq -\mu^*$, all the eigenvalues of (3.12) have negative real parts, which do not affect the stability of the solution in the sense of $\epsilon \to 0^+$.

Applying $T_{\lambda_R^*}^{-1}$ to system (3.12) yields

$$\begin{pmatrix} I & -\lambda_I^* T_{\lambda_R^*}^{-1} \\ \lambda_I^* T_{\lambda_R^*}^{-1} & I \end{pmatrix} \begin{pmatrix} z_R^* \\ z_I^* \end{pmatrix} = c_1^* c_2^* T_{\lambda_R^*}^{-1} \delta(x - x^*) M^* \begin{pmatrix} \langle z_R^*, \delta(x - x^*) \rangle \\ \langle z_I^*, \delta(x - x^*) \rangle \end{pmatrix}, \text{ in } H^{-1} \text{ sense},$$

or equivalently

$$(3.13) \quad \begin{pmatrix} z_R^* \\ z_I^* \end{pmatrix} = c_1^* c_2^* G \begin{pmatrix} I & \lambda_I^* T_{\lambda_R^*}^{-1} \\ -\lambda_I^* T_{\lambda_R^*}^{-1} & I \end{pmatrix} T_{\lambda_R^*}^{-1} \delta(x - x^*) M^* \begin{pmatrix} \langle z_R^*, \delta(x - x^*) \rangle \\ \langle z_I^*, \delta(x - x^*) \rangle \end{pmatrix}, \text{ in } H^{-1} \text{ sense,}$$

where $G := [I + (\lambda_I^*)^2 (T_{\lambda_R^*}^{-1})^2]^{-1}$.

Taking the inner product with $\delta(x - x^*)$ in (3.13), we have

(3.14)
$$\mathcal{N}\begin{pmatrix} \langle z_R^*, \delta(x-x^*) \rangle \\ \langle z_I^*, \delta(x-x^*) \rangle \end{pmatrix} = 0,$$

where

$$\mathcal{N} := \left\langle c_1^* c_2^* G \begin{pmatrix} I & \lambda_I^* T_{\lambda_R^*}^{-1} \\ -\lambda_I^* T_{\lambda_R^*}^{-1} & I \end{pmatrix} T_{\lambda_R^*}^{-1} \delta(x - x^*) M^*, \ \delta(x - x^*) \right\rangle - I.$$

Let $\{(\gamma_n, \psi_n)\}_{n=0}^{\infty}$ be the complete orthonormal pairs of eigenvalues and eigenfunctions of the operator $-d\frac{d^2}{dx^2} - (f_u^*g_v^* - f_v^*g_u^*)/f_u^*$ subject to the homogeneous Neumann boundary condition in $L^2(0, \ell)$ sense. Moreover, $\gamma_n > 0$ holds for any integer $n \ge 0$.

Then, by using lengthy but straightforward eigenfunction expansions, we can reduce ${\mathcal N}$ to

(3.15)
$$\mathcal{N} = \begin{pmatrix} \frac{(\rho_0^* - 2k_1 - \tau\lambda_R^*)X + \tau(\lambda_I^*)^2Y}{(\rho_0^* - 2k_1 - \tau\lambda_R^*)^2 + (\tau\lambda_I^*)^2} - 1 & -\lambda_I^* \frac{\tau X - (\rho_0^* - 2k_1 - \tau\lambda_R^*)Y}{(\rho_0^* - 2k_1 - \tau\lambda_R^*)^2 + (\tau\lambda_I^*)^2} \\ \lambda_I^* \frac{\tau X - (\rho_0^* - 2k_1 - \tau\lambda_R^*)Y}{(\rho_0^* - 2k_1 - \tau\lambda_R^*)^2 + (\tau\lambda_I^*)^2} & \frac{(\rho_0^* - 2k_1 - \tau\lambda_R^*)X + \tau(\lambda_I^*)^2}{(\rho_0^* - 2k_1 - \tau\lambda_R^*)^2 + (\tau\lambda_I^*)^2} - 1 \end{pmatrix},$$

where

$$(3.16)$$

$$X = X(\lambda_R^*, (\lambda_I^*)^2, k_2) := \left\langle c_1^* c_2^* G(T_{\lambda_R^*})^{-1} \delta(x - x^*), \delta(x - x^*) \right\rangle = \sum_{n \ge 0} \frac{(\gamma_n + \lambda_R^* + 2k_2) c_1^* c_2^* \psi_n^2(x^*)}{(\gamma_n + \lambda_R^* + 2k_2)^2 + (\lambda_I^*)^2}$$

$$Y = Y(\lambda_R^*, (\lambda_I^*)^2, k_2) := \left\langle c_1^* c_2^* G(T_{\lambda_R^*})^{-2} \delta(x - x^*), \delta(x - x^*) \right\rangle = \sum_{n \ge 0} \frac{c_1^* c_2^* \psi_n^2(x^*)}{(\gamma_n + \lambda_R^* + 2k_2)^2 + (\lambda_I^*)^2}.$$

To expect the existence of nonzero solution of (3.14), one requires $det(\mathcal{N}) = 0$, or equivalently

(3.17)
$$\begin{cases} \frac{\rho_0^* - 2k_1 - \tau \lambda_R^*}{(\rho_0^* - 2k_1 - \tau \lambda_R^*)^2 + (\tau \lambda_I^*)^2} X + \frac{\tau (\lambda_I^*)^2}{(\rho_0^* - 2k_1 - \tau \lambda_R^*)^2 + (\tau \lambda_I^*)^2} Y = 1, \\ \frac{\tau \lambda_I^*}{(\rho_0^* - 2k_1 - \tau \lambda_R^*)^2 + (\tau \lambda_I^*)^2} X - \frac{\lambda_I^* (\rho_0^* - 2k_1 - \tau \lambda_R^*)}{(\rho_0^* - 2k_1 - \tau \lambda_R^*)^2 + (\tau \lambda_I^*)^2} Y = 0, \end{cases}$$

which is called the real singular limit eigenvalue problem (SLEP) of (3.5).

Proposition 3.1. Let $X(\lambda_R^*, (\lambda_I^*)^2, k_2)$ and $Y(\lambda_R^*, (\lambda_I^*)^2, k_2)$ be defined in (3.16). Then,

$$\begin{split} X(0,0,0) &= \sum_{n\geq 0} \frac{c_1^* c_2^*}{\gamma_n} \psi_n^2(x^*) > \rho_0^*, \ Y(0,0,0) = \sum_{n\geq 0} \frac{c_1^* c_2^*}{\gamma_n^2} \psi_n^2(x^*) > 0, \\ \lim_{k_2 \to +\infty} X(\lambda_R^*, (\lambda_I^*)^2, k_2) &= \lim_{k_2 \to +\infty} Y(\lambda_R^*, (\lambda_I^*)^2, k_2) = 0, \\ \lim_{(\lambda_I^*)^2 \to +\infty} X(\lambda_R^*, (\lambda_I^*)^2, k_2) &= \lim_{(\lambda_I^*)^2 \to +\infty} Y(\lambda_R^*, (\lambda_I^*)^2, k_2) = 0, \\ \frac{\partial X}{\partial (\lambda_I^*)^2} < 0, \ \frac{\partial Y}{\partial (\lambda_I^*)^2} < 0, \ \frac{1}{2} \frac{\partial X}{\partial k_2} \Big|_{\lambda_I^* = 0} = \frac{\partial X}{\partial \lambda_R^*} \Big|_{\lambda_I^* = 0} < 0, \quad \frac{1}{2} \frac{\partial Y}{\partial k_2} = \frac{\partial Y}{\partial \lambda_R^*} < 0. \end{split}$$

In particular, it holds true that

$$\frac{\partial X}{\partial \lambda_R^*}(0,0,k_2) = \frac{1}{2} \frac{\partial X}{\partial k_2}(0,0,k_2) = -Y(0,0,k_2) < 0.$$

Proof. Firstly, $X(0,0,0) > \rho_0^*$ is proved by [21]. Secondly, the rest of the results can be proved by using the eigenfunction expansions of (3.16), and we shall omit the proofs.

Note that to expect zero eigenvalue of (3.17), we need to require that $2k_1 \neq \rho_0^*$. The following results are useful for our later stability analysis for $\epsilon \in (0, \epsilon_0)$.

Proposition 3.2. Let c_1^*, c_2^*, ρ_0^* be defined (3.11) and (A.1), respectively, and let $\{(\gamma_n, \psi_n)\}_{n=0}^{\infty}$ be the complete orthonormal pairs of eigenvalues and eigenfunctions of the operator $-d\frac{d^2}{dx^2} - (f_u^*g_v^* - f_v^*g_u^*)/f_u^*$ subject to the homogeneous Neumann boundary condition in the $L^2(0, \ell)$ sense. Suppose that $\tau \in (\tau^*, +\infty)$ holds, where

(3.18)
$$\tau^* := \sum_{n \ge 0} \frac{c_1^* c_2^*}{\gamma_n^2} \psi_n^2(x^*) = Y(0, 0, 0).$$

Then, the following conclusions hold true:

1. For any $k_1 \in (0, \rho_0^*/2)$, there exists a unique curve $\Gamma = \{(k_1, k_2) | k_2 = \xi(k_1), 0 < k_1 < \rho_0^*/2\}$ in the first quadrant of (k_1, k_2) plane, such that for any $(k_1, k_2) \in \Gamma$, the limiting $(\epsilon \rightarrow 0^+)$ eigenvalue problem (3.12) has zero as its eigenvalue, where $k_2 = \xi(k_1)$ is a positive, continuous and increasing function of k_1 satisfying

(3.19)
$$\lim_{k_1 \to 0^+} \xi(k_1) > 0, \ \lim_{k_1 \to (\rho_0^*/2)^-} \xi(k_1) = \infty.$$

Moreover, for any fixed $k_1 \in (0, \rho_0^*/2)$, all the eigenvalues of the limiting $(\epsilon \to 0^+)$ eigenvalue problem (3.12) have negative real parts for $k_2 \in (0, \xi(k_1))$, while (3.12) has a positive eigenvalue for $k_2 \in (\xi(k_1), \infty)$.

2. For any $k_1 \in (\rho_0^*/2, \infty)$, all the eigenvalues of the limiting $(\epsilon \to 0^+)$ eigenvalue problem (3.12) have negative real parts regardless of $k_2 > 0$.

Proof. To begin with, we note that when $\lambda_I^* \neq 0$, (3.17) can be simplified into

(3.20)
$$\tau \lambda_R^* = \rho_0^* - 2k_1 - X(\lambda_R^*, (\lambda_I^*)^2, k_2), \quad Y(\lambda_R^*, (\lambda_I^*)^2, k_2) = \tau.$$

While if $\lambda_I^* = 0$, then from the first equation of (3.17), we have

(3.21)
$$\tau \lambda_R^* = \rho_0^* - 2k_1 - X(\lambda_R^*, 0, k_2).$$

We firstly claim that the limiting eigenvalue problem (3.12) has no complex eigenvalues $\lambda_R^* \pm i \lambda_I^*$ satisfying $\lambda_R^* \ge 0$ and $\lambda_I^* \ne 0$. Suppose not. Then, (3.20) holds. In particular, $Y(\lambda_R^*, (\lambda_I^*)^2, k_2) = \tau$. It follows from Proposition 3.1 and (3.16) that

(3.22)
$$\tau > \tau^* = \sum_{n \ge 0} \frac{c_1^* c_2^*}{\gamma_n^2} \psi_n^2(x^*) = Y(0, 0, 0) > Y(\lambda_R^*, (\lambda_I^*)^2, k_2),$$

which contradicts with $Y(\lambda_R^*, (\lambda_I^*)^2, k_2) = \tau$. Thus, we complete the proof of the claim.

1. Suppose that the limiting eigenvalue problem (3.12) has zero as its eigenvalue, say $\lambda_R^* = \lambda_I^* = 0$. By $\lambda_R^* = \lambda_I^* = 0$, (3.21) is equivalent to

(3.23)
$$\mathcal{H}(k_1, k_2) := \rho_0^* - 2k_1 - X(0, 0, k_2) = 0.$$

For any fixed $k_1 \in (0, \rho_0^*/2)$, in view of Proposition 3.1, we have

(3.24)
$$\lim_{k_2 \to 0^+} \mathcal{H}(k_1, k_2) = \rho_0^* - 2k_1 - X(0, 0, 0) < 0, \quad \lim_{k_2 \to +\infty} \mathcal{H}(k_1, k_2) = \rho_0^* - 2k_1 > 0.$$

Moreover, by Proposition 3.1, we have $\frac{\partial \mathcal{H}}{\partial k_2}(k_1, k_2) = -\frac{\partial X}{\partial k_2}(0, 0, k_2) > 0$. Then, by (3.24), for any fixed $k_1 \in (0, \rho_0^*/2)$, there exists a unique $k_2 = \hat{k}_2$ such that $\mathcal{H}(k_1, \hat{k}_2) = 0$. Clearly, \hat{k}_2 depends on k_1 continuously. Define $\xi(k_1) := \hat{k}_2$. Then, we have

$$\mathcal{H}(k_1,\xi(k_1)) = 0.$$

Differentiating (3.25) with respect to k_1 , we have $\frac{\partial \mathcal{H}}{\partial k_1} + \frac{\partial \mathcal{H}}{\partial k_2} \frac{d\xi(k_1)}{dk_1} = 0$. Since $\frac{\partial \mathcal{H}}{\partial k_1} = -2 < 0$, we have

(3.26)
$$\frac{d\xi(k_1)}{dk_1} = -2/\frac{\partial X}{\partial k_2}(0,0,k_2) > 0.$$

Then, $\xi(k_1)$ is an increasing function of k_1 . So far, we have proved that if $\lambda_R^* = \lambda_I^* = 0$, then $(k_1, k_2) \in \Gamma = \{(k_1, k_2) | k_2 = \xi(k_1), 0 < k_1 < \rho_0^*/2\}$. On the other hand, it is clear that

for any $(k_1, k_2) \in \Gamma$, (3.12) has zero as its eigenvalue. Moreover, we claim that (3.19) holds. Suppose otherwise that the first term of (3.19) is false. Then, we must have $\xi(k_1) \to 0^+$ as $k_1 \to 0^+$ since $\xi(k_1) > 0$. Then, as $k_1 \to 0^+$, $0 = \mathcal{H}(k_1, \xi(k_1)) \to \mathcal{H}(0, 0) = \rho_0^* - X(0, 0, 0) < 0$, which is a contradiction. Suppose otherwise that the second term of (3.19) is false. Then, as $k_1 \to (\rho_0^*/2)^-$, $0 = \mathcal{H}(k_1, \xi(k_1)) \to \mathcal{H}(\rho_0^*/2, \xi(\rho_0^*/2)) = -X(0, 0, \xi(\rho_0^*/2)) < 0$ since $\xi(\rho_0^*/2) < +\infty$ and $X(0, 0, \xi(\rho_0^*/2)) > 0$. Again, it reaches a contradiction.

Next, we shall show that the transversality condition holds. To that end, we need to resort to the SLEP system of (3.5) in its complex form, instead of the real SLEP equations (3.17). In the following, we always assume that $k_1 \in (0, \rho_0^*/2)$.

Recall that

$$\lim_{\epsilon \to 0^+} \lambda =: \lambda^* =: \lambda^*_R + i\lambda^*_I, \ \lim_{\epsilon \to 0^+} z =: z^* =: z^*_R + iz^*_I.$$

Define

$$T_{\lambda^*} := -d\frac{d^2}{dx^2} - \frac{f_u^* g_v^* - f_v^* g_u^*}{f_u^*} + 2k_2 + \lambda^*.$$

Then, by Remark 2.3 in [21], for any λ^* satisfying $\lambda_R^* + 2k_2 > -\mu^*$, T_{λ^*} has a generalized inverse operator $(T_{\lambda^*})^{-1}$.

Multiplying the second equation of (3.12) by *i* and adding the first equation of (3.12), we have

$$T_{\lambda^*} z^* = \frac{\langle z^*, \delta(x - x^*) \rangle}{\rho_0^* - 2k_1 - \tau \lambda^*} c_1^* c_2^* \delta(x - x^*),$$

or equivalently

(3.27)
$$z^* = \frac{\langle z^*, \delta(x - x^*) \rangle}{\rho_0^* - 2k_1 - \tau \lambda^*} (T_{\lambda^*})^{-1} c_1^* c_2^* \delta(x - x^*).$$

Then, taking the inner product with $\delta(x - x^*)$ on the two sides of (3.27), we have

(3.28)
$$\left(1 - \frac{\left\langle (T_{\lambda}^*)^{-1} c_1^* c_2^* \delta(x - x^*), \delta(x - x^*) \right\rangle}{\rho_0^* - 2k_1 - \tau \lambda^*} \right) \left\langle z^*, \delta(x - x^*) \right\rangle = 0.$$

where the eigenfunction expansion shows that

$$\langle (T_{\lambda}^*)^{-1} c_1^* c_2^* \delta(x - x^*), \delta(x - x^*) \rangle = \sum_{n=0}^{\infty} \frac{c_1^* c_2^*}{\gamma_n + 2k_2 + \lambda^*} \psi_n^2(x^*).$$

Then, (3.28) is equivalent to

(3.29)
$$\frac{\mathcal{F}^*\left(\lambda^*, k_1, k_2\right)}{\rho_0^* - 2k_1 - \tau\lambda^*} \cdot \left\langle z^*, \delta(x - x^*) \right\rangle = 0,$$

where

(3.30)
$$\mathcal{F}^*(\lambda^*, k_1, k_2) := \rho_0^* - 2k_1 - \tau \lambda^* - \sum_{n=0}^{\infty} \frac{c_1^* c_2^*}{\gamma_n + 2k_2 + \lambda^*} \psi_n^2(x^*).$$

Thus, to expect nonzero solutions of (3.29), we need $\mathcal{F}^*(\lambda^*, k_1, k_2) = 0$, which is called the complex SLEP system of (3.5) and is equivalent to

(3.31) Re
$$\mathcal{F}^* := \rho_0^* - 2k_1 - \tau \lambda_R^* - X(\lambda_R^*, (\lambda_I^*)^2, k_2) = 0$$
, Im $\mathcal{F}^* := \lambda_I^*(Y(\lambda_R^*, (\lambda_I^*)^2, k_2) - \tau) = 0$,

where Re and Im stand for the real part and imaginary part, respectively.

It follows from (3.30) and (3.23) that

$$\mathcal{F}^{*}(0,k_{1},\xi(k_{1})) = \rho_{0}^{*} - 2k_{1} - X(0,0,\xi(k_{1})) = \mathcal{H}(k_{1},\xi(k_{1})) = 0,$$
(3.32)
$$\frac{\partial \mathcal{F}^{*}}{\partial \lambda^{*}}(0,k_{1},\xi(k_{1})) = -\tau + \sum_{n=0}^{\infty} \frac{c_{1}^{*}c_{2}^{*}}{(\gamma_{n}+2k_{2})^{2}}\psi_{n}^{2}(x^{*}) = -\tau + Y(0,0,k_{2}) < 0.$$

Thus, by the implicit function theorem, for any $(k_1, k_2) \in \mathcal{O}$, the sufficiently small neighborhood of Γ , there exists a unique $\lambda^* = \lambda^*(k_1, k_2)$ such that

$$\mathcal{F}^*(\lambda^*(k_1,k_2),k_1,k_2) = 0, \ \lambda^*(k_1,\xi(k_1)) = 0.$$

We claim that $\lambda^* = \lambda^*(k_1, k_2)$ is real. Suppose not. Then, $\lambda_I^* \neq 0$. Thus, from the second equation of (3.31), we have $Y(\lambda_R^*, (\lambda_I^*)^2, k_2) - \tau = 0$. However, this is impossible since in the small neighborhood of Γ , we have $Y(\lambda_R^*, (\lambda_I^*)^2, k_2) - \tau < 0$ in that $Y(0, 0, k_2) - \tau < 0$.

Then, we have $\lambda^* = \lambda_R^*$ and $\lambda_I^* = 0$. Therefore, by (3.30) and $\lambda^* = \lambda_R^*$, we have

(3.33)
$$\mathcal{F}^*(\lambda^*(k_1,k_2),k_1,k_2) = \rho_0^* - 2k_1 - \tau \lambda^*(k_1,k_2) - X(\lambda^*(k_1,k_2),0,k_2) = 0.$$

For any fixed $k_1 \in (0, \rho_0^*/2)$, differentiating (3.33) and setting $k_2 = \xi(k_1)$, one obtains

(3.34)
$$\frac{\partial \lambda^*}{\partial k_2}(k_1,\xi(k_1)) = -\frac{\frac{\partial X}{\partial k_2}(0,0,k_2)}{\tau + \frac{\partial X}{\partial \lambda_R^*}(0,0,k_2)} = -\frac{\frac{\partial X}{\partial k_2}(0,0,k_2)}{\tau - Y(0,0,k_2)} > 0.$$

Thus, for any fixed $k_1 \in (0, \rho_0^*/2)$, whenever k_2 is sufficiently close to $\xi(k_1)$, $\lambda^* = \lambda^*(k_1, k_2)$ exists and is real. Clearly, $\lambda^*(k_1, k_2)$ persists for any larger k_2 (say, $k_2 > \xi(k_1)$) and is always positive, since (3.12) has no complex eigenvalues $\lambda^* = \lambda_R^* \pm i\lambda_I^*$ satisfying $\lambda_R^* \ge 0$ and $\lambda_I^* \ne 0$. On the other hand, $\lambda^*(k_1, k_2)$ persists for any smaller k_2 (say, $k_2 < \xi(k_1)$) and it will always have a negative real part. Suppose not. Then, there exists a critical value in $(0, \xi(k_1))$, denoted by \tilde{k}_2^* , of k_2 such that at $k_2 = \tilde{k}_2^*$, the real part of $\lambda^*(k_1, k_2)$ is zero, however, this is impossible since for one hand, (3.12) has no complex eigenvalues $\lambda^* = \lambda_R^* \pm i\lambda_I^*$ satisfying $\lambda_R^* \ge 0$ and $\lambda_I^* \ne 0$, thus eigenvalue with zero real part can never be purely imaginary number; on the other hand, this eigenvalue with zero real part can never be zero since only when $(k_1, k_2) \in \Gamma$, can the eigenvalue be zero.

Thus, we can conclude that for any fixed $k_1 \in (0, \rho_0^*/2)$, all the eigenvalues of (3.12) have negative real parts for $k_2 \in (0, \xi(k_1))$, while (3.12) has a positive eigenvalue for $k_2 \in (\xi(k_1), \infty)$. This proves 1.

2. For any fixed $k_1 \in (\rho_0^*/2, \infty)$, regardless of $k_2 > 0$, we have $\mathcal{H}(k_1, k_2) < 0$. Thus, the eigenvalue problem (3.12) has no zero eigenvalues. Note that Γ is the boundary of the stable



Figure 2: Stability diagram of $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ with respect to (3.1) at any fixed $\epsilon \in (0, \epsilon_0)$, where $\Gamma_1^{\epsilon} = (0, \rho_0^*/2) \times (k_2^{\epsilon}, \infty)$, $\Gamma_2^{\epsilon} = (0, \rho_0^*/2) \times (0, k_2^{\epsilon})$, and $\Gamma_3^{\epsilon} = (\rho_0^*/2, \infty) \times (0, \infty)$. In Γ_1^{ϵ} , $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is Turing unstable with respect to (3.1) driven by suitable k_1 and k_2 . In $\Gamma_2^{\epsilon} \cup \Gamma_3^{\epsilon}$, $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{v}(x;\epsilon))$ is stable with respect to (3.1).

region and unstable region. Since $\{(k_1, k_2) : k_1 \in (\rho_0^*/2, \infty), k_2 \in (0, +\infty)\}$ is in the right side of Γ , and the stable region is in the left side of Γ , it follows that all the eigenvalues of (3.12) have negative real parts. This thus proves 2.

Finally, we have the following results describing the stability of $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ with respect to (3.1) for small $\epsilon > 0$.

Theorem 3.3. Suppose that $\tau \in (\tau^*, +\infty)$ holds. Let τ^* and ρ_0^* be precisely defined in (3.22) and (A.1), respectively. Then, for any fixed $\epsilon \in (0, \epsilon_0)$, the following conclusions hold true:

- 1. If $k_1 > \rho_0^*/2$, then the symmetric solution $(\widetilde{u}(x;\epsilon), \widetilde{v}(x;\epsilon), \widetilde{u}(x;\epsilon), \widetilde{v}(x;\epsilon))$ is stable with respect to system (3.1) for any $k_2 > 0$ (see Fig. 2).
- If k₁ < ρ₀^{*}/2, then there is a critical value k₂^ϵ > 0 such that the symmetric solution is stable with respect to system (3.1) for k₂ < k₂^ϵ, while unstable if k₂ > k₂^ϵ. At k₂ = k₂^ϵ, the Turing-type steady state bifurcation occurs near the solution (ũ(x; ϵ), ῦ(x; ϵ), ũ(x; ϵ), ῦ(x; ϵ)) (see Fig. 2). In particular, k₂^ϵ is an increasing function of k₁, denoted by k₂^ϵ = ξ^ϵ(k₁), satisfying lim_{ϵ→0⁺} ξ^ϵ(k₁) = ξ(k₁) ∈ (0,∞).

Proof. We shall derive the complex characteristic equation of (3.9) firstly.

Solving the first equation of (3.5) after omitting subscription "a" yields

(3.35)
$$w = -(\mathcal{L}_1^{\epsilon} - \epsilon \tau \lambda)^{-1} (f_v^{\epsilon} z) = -\frac{\langle f_v^{\epsilon} z, \phi_0^{\epsilon} \rangle}{\mu_0^{\epsilon} - 2\epsilon k_1 - \epsilon \tau \lambda} \phi_0^{\epsilon} - (\mathcal{L}_1^{\epsilon} - \epsilon \tau \lambda)^{\dagger} (f_v^{\epsilon} z).$$

We define

$$T_{\lambda}^{\epsilon} := -d\frac{d^2}{dx^2} - g_v^{\epsilon} + 2k_2 + g_u^{\epsilon} (\mathcal{L}_1^{\epsilon} - \epsilon \tau \lambda)^{\dagger} f_v^{\epsilon} + \lambda.$$

Substituting (3.35) into the second equation of (3.5), we have

(3.36)
$$T_{\lambda}^{\epsilon}z = -\frac{\langle f_{v}^{\epsilon}z, \phi_{0}^{\epsilon} \rangle}{\mu_{0}^{\epsilon} - 2\epsilon k_{1} - \epsilon\tau\lambda} g_{u}^{\epsilon} \phi_{0}^{\epsilon} = -\frac{\left\langle z, \frac{f_{v}^{\epsilon}\phi_{0}^{\epsilon}}{\sqrt{\epsilon}} \right\rangle}{\rho(\epsilon) - 2k_{1} - \tau\lambda} \frac{g_{u}^{\epsilon}\phi_{0}^{\epsilon}}{\sqrt{\epsilon}}.$$

By using the methods in Lemma 4.1 of [21], one can check that, for any λ satisfying Re $\lambda + 2k_2 > -\mu^*$, T^{ϵ}_{λ} has an inverse operator $(T^{\epsilon}_{\lambda})^{-1}$.

Applying $(T_{\lambda}^{\epsilon})^{-1}$ into (3.36) and taking the inner product with $\frac{f_v^{\epsilon}\phi_0^{\epsilon}}{\sqrt{\epsilon}}$ leads to

(3.37)
$$\left(1 + \frac{\left\langle (T_{\lambda}^{\epsilon})^{-1} \frac{g_{u}^{\epsilon} \phi_{0}^{\epsilon}}{\sqrt{\epsilon}}, \frac{f_{v}^{\epsilon} \phi_{0}^{\epsilon}}{\sqrt{\epsilon}} \right\rangle}{\rho(\epsilon) - 2k_{1} - \tau\lambda} \left\langle z, \frac{f_{v}^{\epsilon} \phi_{0}^{\epsilon}}{\sqrt{\epsilon}} \right\rangle =: \frac{\mathcal{F}(\lambda, k_{1}, k_{2}, \epsilon)}{\rho(\epsilon) - 2k_{1} - \tau\lambda} \left\langle z, \frac{f_{v}^{\epsilon} \phi_{0}^{\epsilon}}{\sqrt{\epsilon}} \right\rangle = 0$$

To expect the nonzero $\langle z, \frac{f_{\psi}^* \phi_0^*}{\sqrt{\epsilon}} \rangle$ and zero eigenvalue in (3.37), we need to require $k_1 \neq \rho_0^*/2$ (for $\epsilon \in (0, \epsilon_0)$) and

(3.38)
$$\mathcal{F}(\lambda, k_1, k_2, \epsilon) := \rho(\epsilon) - 2k_1 - \tau\lambda + \left\langle (T_{\lambda}^{\epsilon})^{-1} \frac{g_u^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}}, \frac{f_v^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}} \right\rangle = 0.$$

As $\epsilon \to 0^+$, we have

(3.39)
$$\mathcal{F}(\lambda, k_1, k_2, \epsilon) \to \mathcal{F}^*(\lambda, k_1, k_2),$$

where $\mathcal{F}^*(\lambda, k_1, k_2)$ is given by (3.30).

1. Suppose that $k_1 > \rho_0^*/2$, then by Proposition 3.2, all the eigenvalues of the limiting eigenvalue problem (3.12) have negative real parts. Thus, for sufficiently small ϵ , say for $0 < \epsilon < \epsilon_1$, all the eigenvalues of the eigenvalue problem (3.5) have negative real parts owing to the continuity of eigenvalues with respect to ϵ . Setting $\hat{\epsilon}_0 = \min \{\epsilon_0, \epsilon_1\}$, and dropping the hat of $\hat{\epsilon}_0$. This proves 1 for any fixed $\epsilon \in (0, \epsilon_0)$.

2. Suppose that $2k_1 < \rho_0^*$. By (3.32) and (3.39), we have

$$\mathcal{F}(0,k_1,\xi(k_1),0) = \mathcal{F}^*(0,k_1,\xi(k_1)) = 0, \quad \frac{\partial \mathcal{F}}{\partial \lambda}(0,k_1,\xi(k_1),0) = \frac{\partial \mathcal{F}^*}{\partial \lambda}(0,k_1,\xi(k_1)) < 0.$$

Then, by the implicit function theorem, for sufficiently small ϵ , say $\epsilon \in (0, \epsilon_2)$ for some $\epsilon_2 > 0$, there exists a $\lambda = \lambda (k_1, k_2, \epsilon)$, such that

(3.40)
$$\mathcal{F}\left(\lambda\left(k_{1},k_{2},\epsilon\right),k_{1},k_{2},\epsilon\right)=0,$$

and that as $\epsilon \to 0^+$,

(3.41)
$$\lambda(k_1, k_2, \epsilon) \to \lambda^*(k_1, k_2).$$

In particular,

$$(3.42) \qquad 0 = \operatorname{Im} \mathcal{F}\left(\lambda(k_1, k_2, \epsilon), k_1, k_2, \epsilon\right) = -\tau \operatorname{Im} \lambda(k_1, k_2, \epsilon) - \operatorname{Im}\left\langle (T_{\lambda}^{\epsilon})^{-1} \frac{g_u^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}}, \frac{f_v^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}} \right\rangle.$$

We claim that $\lambda(k_1, k_2, \epsilon)$ is real. Suppose not. Then, Im $\lambda(k_1, k_2, \epsilon) \neq 0$. Then, by (3.42), we have

(3.43)
$$0 = -\tau - \frac{\operatorname{Im}\left\langle (T_{\lambda}^{\epsilon})^{-1} g_{u}^{\epsilon} \phi_{0}^{\epsilon} / \sqrt{\epsilon}, f_{v}^{\epsilon} \phi_{0}^{\epsilon} / \sqrt{\epsilon} \right\rangle}{\operatorname{Im}\lambda\left(k_{1}, k_{2}, \epsilon\right)}$$

Then, as $\epsilon \to 0^+$, by (3.43), (3.39) and (3.31), we have $0 = -\tau + Y(\lambda^*, 0, k_2)$. However, since $\tau > \tau^*$, we have $-\tau + Y(\lambda^*, 0, k_2) < 0$, which is a contradiction.

From (3.34) and (3.41), we can obtain

(3.44)
$$\lambda(k_1, \xi(k_1), 0) = 0, \quad \frac{\partial \lambda(k_1, \xi(k_1), 0)}{\partial k_2} = \frac{\partial \lambda^*(k_1, \xi(k_1))}{\partial k_2} > 0.$$

Then, by the implicit function theorem, we can obtain the existence of $k_2 = \hat{\xi}(k_1, \epsilon)$ for small ϵ , satisfying

(3.45)
$$\lambda\left(k_1,\widehat{\xi}\left(k_1,\epsilon\right),\epsilon\right) = 0, \ \widehat{\xi}\left(k_1,\epsilon\right) \to \xi(k_1) \ (\text{as } \epsilon \to 0^+).$$

Thus, $\frac{\partial \hat{\xi}(k_1,\epsilon)}{\partial k_1} > 0$ for $\epsilon \in (0,\epsilon_3)$ due to $\frac{\partial \xi(k_1)}{\partial k_1} > 0$ (given by (3.26)).

It follows from (3.44) and (3.45) that the real eigenvalue $\lambda(k_1, k_2, \epsilon)$ changes from negative to positive as k_2 increases around $\hat{\xi}(k_1, \epsilon)$. Define

$$\widehat{\xi}^{\epsilon}(k_1) := \widehat{\xi}(k_1, \epsilon).$$

Setting $\hat{\epsilon}_0 = \min \{\epsilon_0, \epsilon_2, \epsilon_3\}$, and dropping the hat of $\hat{\epsilon}_0$. Then for any fixed $\epsilon \in (0, \epsilon_0)$, we can conclude

$$\frac{\partial \widehat{\xi}^{\epsilon}(k_{1})}{\partial k_{1}} = \frac{\partial \widehat{\xi}(k_{1},\epsilon)}{\partial k_{1}} > 0$$

Thus, $\hat{\xi}^{\epsilon}(k_1)$ is increasing in k_1 for any fixed $\epsilon \in (0, \epsilon_0)$, such that the symmetric solution $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is stable with respect to system (3.1) when $k_2 < k_2^{\epsilon} := \hat{\xi}^{\epsilon}(k_1)$, while unstable when $k_2 > k_2^{\epsilon}$.

Finally, by applying the similar argument in [21, 33], we can derive the simplicity of the critical real eigenvalue $\lambda(k_1, k_2, \epsilon)$, which ends up the proof of 2 for any fixed $\epsilon \in (0, \epsilon_0)$.

4 Dynamics of the coupled system (1.5): case of $0 < \alpha < \infty$

In this section, we shall consider system (1.5) in case of $0 < \alpha < \infty$. For i = 1, 2, we define

$$U_i(x,t) = \int_{-\infty}^t \alpha e^{-\alpha(t-s)} u_i(x,s) ds.$$

Then, system (1.5) can be reduced to

$$(4.1) \qquad \begin{cases} \tau \frac{\partial u_i}{\partial t} = \epsilon \frac{\partial^2 u_i}{\partial x^2} + \frac{1}{\sigma \epsilon} \left(a - u_i - \frac{4u_i v_i}{1 + u_i^2} \right) + k_1 (U_j - u_i), & x \in (0, \ell), \ t > 0, \\ \frac{\partial v_i}{\partial t} = d \frac{\partial^2 v_i}{\partial x^2} + u_i - \frac{u_i v_i}{1 + u_i^2} + k_2 (v_j - v_i), & x \in (0, \ell), \ t > 0, \\ \frac{\partial U_j}{\partial t} = \alpha u_j - \alpha U_j, & x \in (0, \ell), \ t > 0, \\ \frac{\partial u_i}{\partial x} = \frac{\partial v_i}{\partial x} = 0, & x = 0, \ell, \ t > 0, \end{cases}$$

where $i, j \in \{1, 2\}$ and $i \neq j$. Clearly, the fact that $(\tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon), \tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon))$ is the steady state solution of system (1.5) is equivalent to the fact that

$$(u_1, v_1, u_2, v_2, U_1, U_2) = (\widetilde{u}(x; \epsilon), \widetilde{v}(x; \epsilon), \widetilde{u}(x; \epsilon), \widetilde{v}(x; \epsilon), \widetilde{u}(x; \epsilon), \widetilde{u}(x; \epsilon))$$

is the steady state solution of system (4.1).

The eigenvalue problem associated with the linearized system of (4.1) around $(\tilde{u}, \tilde{v}, \tilde{u}, \tilde{v}, \tilde{u}, \tilde{u}, \tilde{u})$ is determined by

(4.2)
$$\begin{cases} \epsilon^2 w_1''(x) + f_u^{\epsilon} w_1 + f_v^{\epsilon} z_1 + \epsilon k_1 (W_2 - w_1) = \epsilon \tau \lambda w_1, \\ dz_1''(x) + g_u^{\epsilon} w_1 + g_v^{\epsilon} z_1 + k_2 (z_2 - z_1) = \lambda z_1, \\ \epsilon^2 w_2''(x) + f_u^{\epsilon} w_2 + f_v^{\epsilon} z_2 + \epsilon k_1 (W_1 - w_2) = \epsilon \tau \lambda w_2, \\ dz_2''(x) + g_u^{\epsilon} w_2 + g_v^{\epsilon} z_2 + k_2 (z_1 - z_2) = \lambda z_2, \\ \alpha w_1 - \alpha W_1 = \lambda W_1, \alpha w_2 - \alpha W_2 = \lambda W_2, \end{cases}$$

subject to homogeneous Neumann boundary conditions for w_1 , z_1 , w_2 and z_2 .

By the last two equations of (4.2), for i = 1, 2, we have

(4.3)
$$W_i = \theta_{\alpha} w_i := \frac{\alpha}{\alpha + \lambda} w_i.$$

Substituting (4.3) into (4.2), we have

(4.4)
$$\begin{cases} \epsilon^2 w_1''(x) + f_u^{\epsilon} w_1 + f_v^{\epsilon} z_1 + \epsilon k_1 (\theta_{\alpha} w_2 - w_1) = \epsilon \tau \lambda w_1, \\ dz_1''(x) + g_u^{\epsilon} w_1 + g_v^{\epsilon} z_1 + k_2 (z_2 - z_1) = \lambda z_1, \\ \epsilon^2 w_2''(x) + f_u^{\epsilon} w_2 + f_v^{\epsilon} z_2 + \epsilon k_1 (\theta_{\alpha} w_1 - w_2) = \epsilon \tau \lambda w_2, \\ dz_2''(x) + g_u^{\epsilon} w_2 + g_v^{\epsilon} z_2 + k_2 (z_1 - z_2) = \lambda z_2, \end{cases}$$

subject to homogeneous Neumann boundary conditions.

By virtue of transformation (3.3), one can reduce (4.4) to the following two eigenvalue problems

(4.5)
$$\begin{cases} \epsilon^2 w_s''(x) + (f_u^{\epsilon} + \epsilon k_1(\theta_{\alpha} - 1))w_s + f_v^{\epsilon} z_s = \epsilon \tau \lambda w_s, & x \in (0, \ell), \\ dz_s''(x) + g_u^{\epsilon} w_s + g_v^{\epsilon} z_s = \lambda z_s, & x \in (0, \ell), \\ w_s'(0) = w_s'(\ell) = z_s'(0) = z_s'(\ell) = 0, \end{cases}$$

and

(4.6)
$$\begin{cases} \epsilon^2 w_a''(x) + (f_u^{\epsilon} - \epsilon k_1(\theta_{\alpha} + 1))w_a + f_v^{\epsilon} z_a = \epsilon \tau \lambda w_a, & x \in (0, \ell), \\ dz_a''(x) + g_u^{\epsilon} w_a + (g_v^{\epsilon} - 2k_2)z_a = \lambda z_a, & x \in (0, \ell), \\ w_a'(0) = w_a'(\ell) = z_a'(0) = z_a'(\ell) = 0. \end{cases}$$

Hereafter, the subscriptions of w_a , z_s , w_a and w_a are ignored if there is no confusion.

4.1 Analysis of the eigenvalue problem (4.5)

Rewriting θ_{α} as $\theta_{\alpha} =: \theta_{\alpha,R} + i\theta_{\alpha,I}$ and $\lambda =: \lambda_R + i\lambda_I$, we have

$$\theta_{\alpha,R} := \frac{\alpha(\alpha + \lambda_R)}{(\alpha + \lambda_R)^2 + \lambda_I^2}, \quad \theta_{\alpha,I} := -\frac{\alpha\lambda_I}{(\alpha + \lambda_R)^2 + \lambda_I^2}.$$

Then, combining with $w =: w_R + iw_I$, $z =: z_R + iz_I$, by (4.5), we have

$$(4.7) \qquad \begin{cases} \left(\epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} + \epsilon k_1(\theta_{\alpha,R} - 1)\right) w_R - \epsilon k_1 \theta_{\alpha,I} w_I + f_v^{\epsilon} z_R = \epsilon \tau (\lambda_R w_R - \lambda_I w_I), \\ \left(\epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} + \epsilon k_1(\theta_{\alpha,R} - 1)\right) w_I + \epsilon k_1 \theta_{\alpha,I} w_R + f_v^{\epsilon} z_I = \epsilon \tau (\lambda_R w_I + \lambda_I w_R), \\ d \frac{d^2}{dx^2} z_R + g_u^{\epsilon} w_R + g_v^{\epsilon} z_R = \lambda_R z_R - \lambda_I z_I, \\ d \frac{d^2}{dx^2} z_I + g_u^{\epsilon} w_I + g_v^{\epsilon} z_I = \lambda_R z_I + \lambda_I z_R. \end{cases}$$

Define

$$\mathcal{L}_{\alpha}^{\epsilon} := \epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} + \epsilon k_1 (\theta_{\alpha,R} - 1) - \epsilon \tau \lambda_R.$$

Clearly, $\mathcal{L}^{\epsilon}_{\alpha}$ has complete and orthonormal pairs of eigenvalues and eigenfunctions

$$\{(\mu_n^{\epsilon} + \epsilon k_1(\theta_{\alpha,R} - 1) - \epsilon \tau \lambda_R, \phi_n^{\epsilon})\}_{n=0}^{\infty}$$

with respect to homogeneous Neumann boundary condition in $L^2(0,\ell)$ sense.

Solving the first two equations of (4.7) with respect to w_R and w_I , we have

$$w_{R} = \left[I + (\epsilon \tau \lambda_{I} - \epsilon k_{1} \theta_{\alpha,I})^{2} (\mathcal{L}_{\alpha}^{\epsilon})^{-2}\right]^{-1} \left[(\epsilon \tau \lambda_{I} - \epsilon k_{1} \theta_{\alpha,I}) (\mathcal{L}_{\alpha}^{\epsilon})^{-2} (f_{v}^{\epsilon} z_{I}) - (\mathcal{L}_{\alpha}^{\epsilon})^{-1} (f_{v}^{\epsilon} z_{R}) \right],$$

$$w_{I} = - \left[I + (\epsilon \tau \lambda_{I} - \epsilon k_{1} \theta_{\alpha,I})^{2} (\mathcal{L}_{\alpha}^{\epsilon})^{-2}\right]^{-1} \left[(\mathcal{L}_{\alpha}^{\epsilon})^{-1} (f_{v}^{\epsilon} z_{I}) + (\epsilon \tau \lambda_{I} - \epsilon k_{1} \theta_{\alpha,I}) (\mathcal{L}_{\alpha}^{\epsilon})^{-2} (f_{v}^{\epsilon} z_{R}) \right],$$

where the invertibility of $\mathcal{L}^{\epsilon}_{\alpha}$ for small $\epsilon > 0$ and $\operatorname{Re} \lambda > -\mu^*$ can be shown similarly by Lemma 2.1 of [20] and Remark 2.2 of [21].

By using the eigenfunction expansions, we have

(4.8)
$$w_R = \sum_{n \ge 0} w_R^n \phi_n^{\epsilon} = w_R^0 \phi_0^{\epsilon} + w_R^{\dagger}, \quad w_I = \sum_{n \ge 0} w_I^n \phi_n^{\epsilon} = w_I^0 \phi_0^{\epsilon} + w_I^{\dagger},$$

where
$$w_R^{\dagger} = \sum_{n \ge 1} w_R^n \phi_n^{\epsilon}$$
, $w_I^{\dagger} = \sum_{n \ge 1} w_I^n \phi_n^{\epsilon}$, and for $n \ge 0$, it holds true that
 $w_R^n = \frac{\left(-\mu_n^{\epsilon} + \epsilon k_1 (1 - \theta_{\alpha,R}) + \epsilon \tau \lambda_R\right) \langle f_v^{\epsilon} z_R, \phi_n^{\epsilon} \rangle + \left(\epsilon \tau \lambda_I - \epsilon k_1 \theta_{\alpha,I}\right) \langle f_v^{\epsilon} z_I, \phi_n^{\epsilon} \rangle}{(\mu_n^{\epsilon} + \epsilon k_1 (\theta_{\alpha,R} - 1) - \epsilon \tau \lambda_R)^2 + (\epsilon \tau \lambda_I - \epsilon k_1 \theta_{\alpha,I})^2},$
 $w_I^n = \frac{\left(-\mu_n^{\epsilon} + \epsilon k_1 (1 - \theta_{\alpha,R}) + \epsilon \tau \lambda_R\right) \langle f_v^{\epsilon} z_I, \phi_n^{\epsilon} \rangle - (\epsilon \tau \lambda_I - \epsilon k_1 \theta_{\alpha,I}) \langle f_v^{\epsilon} z_R, \phi_n^{\epsilon} \rangle}{(\mu_n^{\epsilon} + \epsilon k_1 (\theta_{\alpha,R} - 1) - \epsilon \tau \lambda_R)^2 + (\epsilon \tau \lambda_I - \epsilon k_1 \theta_{\alpha,I}) \langle f_v^{\epsilon} z_R, \phi_n^{\epsilon} \rangle}{(\mu_n^{\epsilon} + \epsilon k_1 (\theta_{\alpha,R} - 1) - \epsilon \tau \lambda_R)^2 + (\epsilon \tau \lambda_I - \epsilon k_1 \theta_{\alpha,I})^2}.$

Similar to (3.8), we have

$$w_{R}^{\dagger} = \left[I + (\epsilon \tau \lambda_{I} - \epsilon k_{1} \theta_{\alpha,I})^{2} (\mathcal{L}_{\alpha}^{\epsilon})^{-2}\right]^{\dagger} \left[(\epsilon \tau \lambda_{I} - \epsilon k_{1} \theta_{\alpha,I}) (\mathcal{L}_{\alpha}^{\epsilon})^{2\dagger} (f_{v}^{\epsilon} z_{I}) - (\mathcal{L}_{\alpha}^{\epsilon})^{\dagger} (f_{v}^{\epsilon} z_{R}) \right],$$

$$w_{I}^{\dagger} = -\left[I + (\epsilon \tau \lambda_{I} - \epsilon k_{1} \theta_{\alpha,I})^{2} (\mathcal{L}_{\alpha}^{\epsilon})^{-2}\right]^{\dagger} \left[(\mathcal{L}_{\alpha}^{\epsilon})^{\dagger} (f_{v}^{\epsilon} z_{I}) + (\epsilon \tau \lambda_{I} - \epsilon k_{1} \theta_{\alpha,I}) (\mathcal{L}_{\alpha}^{\epsilon})^{2\dagger} (f_{v}^{\epsilon} z_{R}) \right],$$

where

$$(\mathcal{L}_{\alpha}^{\epsilon})^{\dagger} = \sum_{n \ge 1} \frac{\langle \cdot, \phi_{n}^{\epsilon} \rangle}{\mu_{n}^{\epsilon} - \epsilon \tau \lambda_{R} + \epsilon k_{1}(\theta_{\alpha,R} - 1)} \phi_{n}^{\epsilon}, \quad (\mathcal{L}_{\alpha}^{\epsilon})^{2\dagger} = \sum_{n \ge 1} \frac{\langle \cdot, \phi_{n}^{\epsilon} \rangle}{(\mu_{n}^{\epsilon} - \epsilon \tau \lambda_{R} + \epsilon k_{1}(\theta_{\alpha,R} - 1))^{2}} \phi_{n}^{\epsilon},$$

and

$$\left[I + (\epsilon \tau \lambda_I - \epsilon k_1 \theta_{\alpha,I})^2 (\mathcal{L}_{\alpha}^{\epsilon})^{-2}\right]^{\dagger} = \sum_{n \ge 1} \frac{(\mu_n^{\epsilon} - \epsilon \tau \lambda_R + \epsilon k_1 (\theta_{\alpha,R} - 1))^2 \langle \cdot, \phi_n^{\epsilon} \rangle}{(\mu_n^{\epsilon} - \epsilon \tau \lambda_R + \epsilon k_1 (\theta_{\alpha,R} - 1))^2 + (\epsilon \tau \lambda_I - \epsilon k_1 \theta_{\alpha,I})^2} \phi_n^{\epsilon}$$

Substituting (4.8) into the last two equations of (4.7) yields

(4.9)
$$\begin{pmatrix} \widehat{T}_{\lambda_R}^{\varepsilon} & -\lambda_I + \widehat{S}^{\epsilon} \\ \lambda_I - \widehat{S}^{\epsilon} & \widehat{T}_{\lambda_R}^{\varepsilon} \end{pmatrix} \begin{pmatrix} z_R \\ z_I \end{pmatrix} = -\frac{g_u^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}} \widehat{M}^{\varepsilon} \begin{pmatrix} \left\langle z_R, \frac{f_v^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}} \right\rangle \\ \left\langle z_I, \frac{f_v^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}} \right\rangle \end{pmatrix},$$

where

$$\begin{cases} \widehat{T}_{\lambda_{R}}^{\epsilon} := -d\frac{d^{2}}{dx^{2}} - g_{v}^{\epsilon} + g_{u}^{\epsilon} \left\{ I + (\epsilon\tau\lambda_{I} - \epsilon k_{1}\theta_{\alpha,I})^{2} (\mathcal{L}_{\alpha}^{\epsilon})^{-2} \right\}^{\dagger} (\mathcal{L}_{\alpha}^{\epsilon})^{\dagger} f_{v}^{\epsilon} + \lambda_{R}, \\ \widehat{S}^{\epsilon} := -(\epsilon\tau\lambda_{I} - \epsilon k_{1}\theta_{\alpha,I}) g_{u}^{\epsilon} \left\{ I + (\epsilon\tau\lambda_{I} - \epsilon k_{1}\theta_{\alpha,I})^{2} (\mathcal{L}_{\alpha}^{\epsilon})^{-2} \right\}^{\dagger} (\mathcal{L}_{\alpha}^{\epsilon})^{2\dagger} f_{v}^{\epsilon}, \\ \widehat{M}^{\epsilon} := \begin{pmatrix} \frac{\rho(\epsilon) - k_{1}(1 - \theta_{\alpha,R}) - \tau\lambda_{R}}{(\rho(\epsilon) - k_{1}(1 - \theta_{\alpha,R}) - \tau\lambda_{R})^{2} + (\tau\lambda_{I} - k_{1}\theta_{\alpha,I})^{2}} & \frac{k_{1}\theta_{\alpha,I} - \tau\lambda_{I}}{(\rho(\epsilon) - k_{1}(1 - \theta_{\alpha,R}) - \tau\lambda_{R})^{2} + (\tau\lambda_{I} - k_{1}\theta_{\alpha,I})^{2}} \\ \frac{\rho(\epsilon) - k_{1}(1 - \theta_{\alpha,R}) - \tau\lambda_{R}}{(\rho(\epsilon) - k_{1}(1 - \theta_{\alpha,R}) - \tau\lambda_{R})^{2} + (\tau\lambda_{I} - k_{1}\theta_{\alpha,I})^{2}} & \frac{\rho(\epsilon) - k_{1}(1 - \theta_{\alpha,R}) - \tau\lambda_{R}}{(\rho(\epsilon) - k_{1}(1 - \theta_{\alpha,R}) - \tau\lambda_{R})^{2} + (\tau\lambda_{I} - k_{1}\theta_{\alpha,I})^{2}} \end{pmatrix}.$$

Analogous to (3.12), we can obtain the following singular limit system of (4.9)

(4.10)
$$\begin{pmatrix} \widehat{T}_{\lambda_R^*} & -\lambda_I^* \\ \lambda_I^* & \widehat{T}_{\lambda_R^*} \end{pmatrix} \begin{pmatrix} z_R^* \\ z_I^* \end{pmatrix} = c_1^* c_2^* \delta(x - x^*) \widehat{M}^* \begin{pmatrix} \langle z_R^*, \delta(x - x^*) \rangle \\ \langle z_I^*, \delta(x - x^*) \rangle \end{pmatrix}, \text{ in } H^{-1} \text{ sense,}$$

where

$$\begin{split} \widehat{T}_{\lambda_{R}^{*}} &:= -d\frac{d^{2}}{dx^{2}} - \frac{f_{u}^{*}g_{v}^{*} - f_{v}^{*}g_{u}^{*}}{f_{u}^{*}} + \lambda_{R}^{*}, \\ \widehat{M}^{*} &:= \begin{pmatrix} \frac{\rho_{0}^{*} - k_{1}(1 - \theta_{\alpha,R}^{*}) - \tau\lambda_{R}^{*}}{\left(\rho_{0}^{*} - k_{1}(1 - \theta_{\alpha,R}^{*}) - \tau\lambda_{R}^{*}\right)^{2} + \left(\tau\lambda_{I}^{*} - k_{1}\theta_{\alpha,I}^{*}\right)^{2}}{\tau\lambda_{I}^{*} - k_{1}\theta_{\alpha,I}^{*}} & \frac{k_{1}\theta_{\alpha,I}^{*} - \tau\lambda_{I}^{*}}{\left(\rho_{0}^{*} - k_{1}(1 - \theta_{\alpha,R}^{*}) - \tau\lambda_{R}^{*}\right)^{2} + \left(\tau\lambda_{I}^{*} - k_{1}\theta_{\alpha,I}^{*}\right)^{2}} & \frac{\rho_{0}^{*} - k_{1}(1 - \theta_{\alpha,R}^{*}) - \tau\lambda_{R}^{*}}{\left(\rho_{0}^{*} - k_{1}(1 - \theta_{\alpha,R}^{*}) - \tau\lambda_{R}^{*}\right)^{2} + \left(\tau\lambda_{I}^{*} - k_{1}\theta_{\alpha,I}^{*}\right)^{2}} & \frac{\rho_{0}^{*} - k_{1}(1 - \theta_{\alpha,R}^{*}) - \tau\lambda_{R}^{*}}{\left(\rho_{0}^{*} - k_{1}(1 - \theta_{\alpha,R}^{*}) - \tau\lambda_{R}^{*}\right)^{2} + \left(\tau\lambda_{I}^{*} - k_{1}\theta_{\alpha,I}^{*}\right)^{2}} \end{pmatrix}, \end{split}$$

in which ρ_0^* is defined by (A.1) in Appendix, and

$$\theta_{\alpha,R}^* := \frac{\alpha(\alpha + \lambda_R^*)}{(\alpha + \lambda_R^*)^2 + (\lambda_I^*)^2}, \quad \theta_{\alpha,I}^* := -\frac{\alpha\lambda_I^*}{(\alpha + \lambda_R^*)^2 + (\lambda_I^*)^2}.$$

By Lemma 2.4 of [21], the generalized inverse operator $(\hat{T}_{\lambda_R^*})^{-1}$ exists for $\lambda_R^* > -\mu^*$, where μ^* is given by (3.7). Similar to the derivation of (3.13) in §2, we have

$$(4.11) \qquad \begin{pmatrix} z_R^* \\ z_I^* \end{pmatrix} = c_1^* c_2^* \widehat{G} \begin{pmatrix} I & \lambda_I^* (\widehat{T}_{\lambda_R^*})^{-1} \\ -\lambda_I^* (\widehat{T}_{\lambda_R^*})^{-1} & I \end{pmatrix} (\widehat{T}_{\lambda_R^*})^{-1} \delta(x - x^*) \widehat{M}^* \begin{pmatrix} \langle z_R^*, \delta(x - x^*) \rangle \\ \langle z_I^*, \delta(x - x^*) \rangle \end{pmatrix},$$

where $\widehat{G} := \{I + (\lambda_I^*)^2 (\widehat{T}_{\lambda_R^*})^{-2}\}^{-1}.$

Taking the inner product with $\delta(x - x^*)$ in (4.11), we have

$$\mathcal{N}_{\alpha}\left(\begin{pmatrix} \langle z_{R}^{*}, \delta(x-x^{*}) \rangle \\ \langle z_{I}^{*}, \delta(x-x^{*}) \rangle \end{pmatrix} = 0,$$

where

(4.12)
$$\mathcal{N}_{\alpha} := \left\langle c_1^* c_2^* \widehat{G} \begin{pmatrix} I & \lambda_I^* (\widehat{T}_{\lambda_R^*})^{-1} \\ -\lambda_I^* (\widehat{T}_{\lambda_R^*})^{-1} & I \end{pmatrix} (\widehat{T}_{\lambda_R^*})^{-1} \delta(x - x^*) \widehat{M}^*, \delta(x - x^*) \right\rangle - I.$$

Similarly, we can check that \mathcal{N}_{α} is equivalent to

$$\begin{pmatrix} \frac{\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)\widehat{X}+\lambda_{I}^{*}\left(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*}\right)\widehat{Y}}{\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)\widehat{X}+\left(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*}\right)^{2}}-1 & -\frac{\left(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*}\right)\widehat{X}-\lambda_{I}^{*}\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)\widehat{Y}}{\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)\widehat{X}-\lambda_{I}^{*}\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)\widehat{Y}} & \frac{\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)\widehat{X}+\lambda_{I}^{*}\left(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*}\right)\widehat{Y}}{\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)\widehat{X}+\left(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*}\right)\widehat{Y}}-1 \end{pmatrix},$$

where $\widehat{X} = \widehat{X}(\lambda_R^*, (\lambda_I^*)^2)$ and $\widehat{Y} = \widehat{Y}(\lambda_R^*, (\lambda_I^*)^2)$ are respectively given by

(4.13)

$$\widehat{X} := \left\langle c_1^* c_2^* \widehat{G}(\widehat{T}_{\lambda_R^*})^{-1} \delta(x - x^*), \delta(x - x^*) \right\rangle = \sum_{n \ge 0} \frac{(\gamma_n + \lambda_R^*) c_1^* c_2^*}{(\gamma_n + \lambda_R^*)^2 + (\lambda_I^*)^2} \psi_n^2(x^*), \\
\widehat{Y} := \left\langle c_1^* c_2^* \widehat{G}(\widehat{T}_{\lambda_R^*})^{-2} \delta(x - x^*), \delta(x - x^*) \right\rangle = \sum_{n \ge 0} \frac{c_1^* c_2^*}{(\gamma_n + \lambda_R^*)^2 + (\lambda_I^*)^2} \psi_n^2(x^*).$$

To expect the existence of nonzero solutions of (4.12), one requires $det(\mathcal{N}_{\alpha}) = 0$, or equivalently

$$(\text{SLEP-1}) \qquad \begin{cases} \frac{\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)\widehat{X}+\lambda_{I}^{*}\left(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*}\right)\widehat{Y}}{\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)^{2}+\left(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*}\right)^{2}}=1,\\ \frac{(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*})\widehat{X}-\lambda_{I}^{*}(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*})\widehat{Y}}{\left(\rho_{0}^{*}-k_{1}(1-\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)^{2}+\left(\tau\lambda_{I}^{*}-k_{1}\theta_{\alpha,I}^{*}\right)^{2}}=0, \end{cases}$$

which is called the real singular limit eigenvalue problem (SLEP) of (4.5).

Similar to Proposition 3.1, we have

Proposition 4.1. Let $\widehat{X}(\lambda_R^*, (\lambda_I^*)^2)$ and $\widehat{Y}(\lambda_R^*, (\lambda_I^*)^2)$ be defined in (4.13). Then it holds true that

$$\begin{split} \widehat{X}(0,0) &= \sum_{n\geq 0} \frac{c_1^* c_2^*}{\gamma_n} \psi_n^2(x^*) > \rho_0^*, \quad \widehat{Y}(0,0) = \sum_{n\geq 0} \frac{c_1^* c_2^*}{\gamma_n^2} \psi_n^2(x^*) > 0, \quad \frac{\partial \widehat{Y}}{\partial \lambda_R^*} < 0, \\ \frac{\partial \widehat{X}}{\partial (\lambda_I^*)^2} < 0, \quad \lim_{(\lambda_I^*)^2 \to \infty} \widehat{X}(\lambda_R^*, (\lambda_I^*)^2) = \lim_{(\lambda_I^*)^2 \to \infty} \widehat{Y}(\lambda_R^*, (\lambda_I^*)^2) = 0. \end{split}$$

It turns out that the distributed-delay coupling cannot affect the stability of symmetric solution through (SLEP-1).

Lemma 4.2. Suppose that $\tau \in (\tau^*, +\infty)$ holds. Then, for any $k_1 > 0$, $k_2 > 0$ and $\alpha > 0$, all the eigenvalues of (SLEP-1) lie on the left side of imaginary axis.

Proof. It suffices to show that for any $k_1 > 0$, $k_2 > 0$ and $\alpha > 0$, both zero and pure imaginary numbers are impossible to become eigenvalues of (SLEP-1).

If $\lambda^* = 0$ is a eigenvalue for some (k_1, k_2, α) , then (SLEP-1) is reduced to

$$X(0,0) - \rho_0^* = 0,$$

which is a contradiction with Proposition 4.1.

If $\lambda^* = i\lambda_I^* \neq 0$ is a pure imaginary eigenvalue for some (k_1, k_2, α) , then (SLEP-1) is simplified into

$$\widehat{X}(0, (\lambda_I^*)^2) = \rho_0^* - \frac{k_1(\lambda_I^*)^2}{\alpha^2 + (\lambda_I^*)^2}, \ \widehat{Y}(0, (\lambda_I^*)^2) = \tau + \frac{k_1\alpha}{\alpha^2 + (\lambda_I^*)^2},$$

where the second equation does not hold true owing to

$$\tau + \frac{k_1 \alpha}{\alpha^2 + (\lambda_I^*)^2} > \tau > \tau^* = \widehat{Y}(0,0) > \widehat{Y}(0,(\lambda_I^*)^2),$$

which completes the proof.

4.2 Analysis of the eigenvalue problem (4.6)

Repeating the similar procedure in $\S4.1$, we can derive the following real SLEP equations of the eigenvalue problem (4.6)

$$(\text{SLEP-2}) \qquad \left\{ \begin{aligned} \frac{\left(\rho_{0}^{*}-k_{1}(1+\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)X+\lambda_{I}^{*}\left(\tau\lambda_{I}^{*}+k_{1}\theta_{\alpha,I}^{*}\right)Y}{\left(\rho_{0}^{*}-k_{1}(1+\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)^{2}+\left(\tau\lambda_{I}^{*}+k_{1}\theta_{\alpha,I}^{*}\right)^{2}}=1,\\ \frac{(\tau\lambda_{I}^{*}+k_{1}\theta_{\alpha,I}^{*})X-\lambda_{I}^{*}(\rho_{0}^{*}-k_{1}(1+\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*})Y}{\left(\rho_{0}^{*}-k_{1}(1+\theta_{\alpha,R}^{*})-\tau\lambda_{R}^{*}\right)^{2}+\left(\tau\lambda_{I}^{*}+k_{1}\theta_{\alpha,I}^{*}\right)^{2}}=0, \end{aligned}$$

where $X = X(\lambda_R^*, (\lambda_I^*)^2, k_2)$ and $Y = Y(\lambda_R^*, (\lambda_I^*)^2, k_2)$ are given by (3.16).



Figure 3: Graph of $X(0, (\lambda_I^*)^2, k_2), P((\lambda_I^*)^2), Y(0, (\lambda_I^*)^2, k_2) \text{ and } Q((\lambda_I^*)^2).$

To show the stability of $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ with respect to (4.6), we need to investigate the existence of zero or pure imaginary eigenvalues of (SLEP-2) with respect to $\alpha \in (0, +\infty)$ under some fixed $(k_1, k_2) \in \mathbb{R}^2_+ \setminus \Gamma$, where Γ is given by Proposition 3.2.

Suppose that $\lambda_R^* = \lambda_I^* = 0$. Then, we can reduce (SLEP-2) to

(4.14)
$$X(0,0,k_2) - \rho_0^* + 2k_1 = c_1^* c_2^* \sum_{n \ge 0} \frac{1}{\gamma_n + 2k_2} \psi_n^2(x^*) - \rho_0^* + 2k_1 = 0.$$

Clearly, (4.14) is independent of α . In §3, we have shown that, if $\alpha = +\infty$, then for any $(k_1, k_2) \in \mathbb{R}^2_+ \setminus \Gamma$, the eigenvalue problem has no zero eigenvalues. Thus, we can conclude that the results can be extended to the case of $\alpha \in (0, +\infty)$ and $(k_1, k_2) \in \mathbb{R}^2_+ \setminus \Gamma$.

Suppose that $\lambda_R^* = 0$ and $\lambda_I^* \neq 0$. Then, (SLEP-2) can be reduced to

(4.15a)
$$\begin{cases} X(0, (\lambda_I^*)^2, k_2) = \rho_0^* - k_1 \left(1 + \frac{\alpha^2}{\alpha^2 + (\lambda_I^*)^2} \right) =: P((\lambda_I^*)^2), \\ k_1 = \frac{\alpha^2}{\alpha^2 + (\lambda_I^*)^2} = \frac{\alpha^2}{\alpha^2 + (\lambda_$$

(4.15b)
$$\left(Y(0, (\lambda_I^*)^2, k_2) = \tau - \frac{k_1 \alpha}{\alpha^2 + (\lambda_I^*)^2} =: Q((\lambda_I^*)^2),\right)$$

where (see (3.16))

$$X(0, (\lambda_I^*)^2, k_2) = \sum_{n \ge 0} \frac{(\gamma_n + 2k_2)c_1^* c_2^* \psi_n^2(x^*)}{(\gamma_n + 2k_2)^2 + (\lambda_I^*)^2}, \ Y(0, (\lambda_I^*)^2, k_2) = \sum_{n \ge 0} \frac{c_1^* c_2^* \psi_n^2(x^*)}{(\gamma_n + 2k_2)^2 + (\lambda_I^*)^2}.$$

We have the following results on the properties of $P((\lambda_I^*)^2)$ and $Q((\lambda_I^*)^2)$ (see Fig. 3).

Proposition 4.3. Let $P((\lambda_I^*)^2)$ and $Q((\lambda_I^*)^2)$ be defined by (4.15). Then, the following conclusions hold true:

$$P(0) = \rho_0^* - 2k_1, \lim_{(\lambda_I^*)^2 \to +\infty} P((\lambda_I^*)^2) = \rho_0^* - k_1, \ Q(0) = \tau - \frac{k_1}{\alpha}, \lim_{(\lambda_I^*)^2 \to +\infty} Q((\lambda_I^*)^2) = \tau,$$

and

$$\frac{\partial P}{\partial (\lambda_I^*)^2} = \frac{k_1 \alpha^2}{(\alpha^2 + (\lambda_I^*)^2)^2} > 0, \quad \frac{\partial Q}{\partial (\lambda_I^*)^2} = \frac{k_1 \alpha}{(\alpha^2 + (\lambda_I^*)^2)^2} > 0$$

Proof. The proof is trivial and we shall omit it.

Then, by Propositions 3.1 and 4.3, we have the following results (see Fig. 3).

Lemma 4.4. Suppose that $\tau \in (\tau^*, +\infty)$ holds. Define

(4.16)
$$\Gamma_{1} := \{ (k_{1}, k_{2}) | k_{1} \in (0, \rho_{0}^{*}/2), X(0, 0, k_{2}) < \rho_{0}^{*} - 2k_{1} \},$$

$$\Gamma_{2} := \{ (k_{1}, k_{2}) | k_{1} \in (0, \rho_{0}^{*}/2), X(0, 0, k_{2}) > \rho_{0}^{*} - 2k_{1} \},$$

$$\Gamma_{3-1} := (\rho_{0}^{*}/2, \rho_{0}^{*}) \times (0, \infty), \ \Gamma_{3-2} := [\rho_{0}^{*}, \infty) \times (0, \infty).$$

If one of the following conditions holds:

- (1). $P(0) = \rho_0^* 2k_1 > X(0, 0, k_2)$, and $\alpha \in (0, +\infty)$ (see region Γ_1);
- (2). $P(\infty) = \rho_0^* k_1 < 0$, and $\alpha \in (0, +\infty)$ (see region Γ_{3-2});
- (3). $Q(0) = \tau \frac{k_1}{\alpha} \ge Y(0, 0, k_2)$, or equivalently $\alpha \in [\alpha_0, +\infty)$, where

(4.17)
$$\alpha_0 := \frac{k_1}{\tau - Y(0, 0, k_2)} > 0,$$

then the eigenvalue problem (SLEP-2) has no purely imaginary eigenvalues.

Then, by Lemma 4.4, (4.16) and the continuity of eigenvalues with respect to ϵ up to $\epsilon \to 0^+$, one can immediately obtain the following results.

Theorem 4.5. Suppose that $\tau \in (\tau^*, +\infty)$ holds. Then, there exists a $\epsilon_0 > 0$, such that for any fixed $\epsilon \in (0, \epsilon_0)$, regardless of $\alpha \in (0, +\infty)$, $(\tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon), \tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon))$ is always unstable when $(k_1, k_2) \in \Gamma_1$, while always stable when $(k_1, k_2) \in \Gamma_{3-2}$.

In view of (3) in Lemma 4.4 and Theorem 4.5, to expect zero eigenvalue or purely imaginary eigenvalues, it is sufficient to consider the case when $(k_1, k_2) \in \Gamma_2 \cup \Gamma_{3-1}$ and $\alpha \in (0, \alpha_0)$.

It is clear that when $(k_1, k_2) \in \Gamma_2 \cup \Gamma_{3-1}$, (4.15a) always has a unique positive root, denoted by $\lambda_{I,1}^*(\alpha)$, which is defined in $(0, +\infty)$. On the other hand, when $(k_1, k_2) \in \Gamma_2 \cup \Gamma_{3-1}$ and $\alpha \in (0, \alpha_0)$, (4.15b) has a unique root (non-zero), denoted by $\lambda_{I,2}^*(\alpha)$, which is defined in $(0, \alpha_0)$.

We have the following lemma on the properties of $\lambda_{I,1}^*(\alpha)$ in $(0, +\infty)$ (see Fig. 4).

Lemma 4.6. Suppose that $\tau \in (\tau^*, +\infty)$, $(k_1, k_2) \in \Gamma_2 \cup \Gamma_{3-1}$ and $\alpha \in (0, +\infty)$. Let $\lambda_{I,1}^*(\alpha)$ be the solution of (4.15a). Then, $\lambda_{I,1}^*(\alpha)$ is increasing with respect to $\alpha \in (0, +\infty)$. Moreover, the following conclusions hold true:

1. Suppose that $k_1 \ge 2\rho_0^*/3$ holds. Then, for any $\alpha \in (0, +\infty)$, $\lambda_{I,1}^*(\alpha) > \alpha$.

2. Suppose that $k_1 \in (0, \rho_0^*/2) \cup (\rho_0^*/2, 2\rho_0^*/3)$ holds. Then, there exists a $\hat{k}_2^* > 0$ such that if $k_2 \ge \hat{k}_2^*$, then for any $\alpha \in (0, +\infty)$, $\lambda_{I,1}^*(\alpha) < \alpha$; while if $k_2 < \hat{k}_2^*$, then there is a unique $\alpha_1 \in (0, +\infty)$, such that

(4.18)
$$\lambda_{I,1}^*(\alpha) \begin{cases} > \alpha, & \text{if } \alpha \in (0, \alpha_1), \\ = \alpha, & \text{if } \alpha = \alpha_1, \\ < \alpha, & \text{if } \alpha \in (\alpha_1, +\infty). \end{cases}$$

In particular, $\hat{k}_2^* < \xi(k_1)$, where $\xi(k_1)$ is given by (3.25).

Proof. For $(k_1, k_2) \in \Gamma_2 \cup \Gamma_{3-1}$ and $\alpha \in (0, +\infty)$, we define

(4.19)
$$\widehat{F}_1(\eta, \alpha, k_1, k_2) := X(0, \eta \alpha^2, k_2) - \rho_0^* + k_1 \left(1 + \frac{1}{1+\eta}\right), \ \eta_1^*(\alpha) = \left(\frac{\lambda_{I,1}^*(\alpha)}{\alpha}\right)^2,$$

where $\eta > 0$.

Thus, by (4.15a), we have $\widehat{F}_1(\eta_1^*(\alpha), \alpha, k_1, k_2) = 0$. Differentiating it with respect to α , we have

$$\frac{\partial \lambda_{I,1}^*(\alpha)}{\partial \alpha} = -\frac{k_1 \alpha \lambda_{I,1}^*(\alpha)}{(\alpha^* + (\lambda_{I,1}^*(\alpha))^2) \frac{\partial X(0,(\lambda_I^*)^2,k_2)}{\partial (\lambda_I^*)^2} - k_1 \alpha} > 0,$$

where $\frac{\partial X(0,(\lambda_I^*)^2,k_2)}{\partial (\lambda_I^*)^2} < 0$ is used. Then, $\lambda_{I,1}^*(\alpha)$ is increasing with respect to $\alpha \in (0,+\infty)$.

By Proposition 3.1, we have

$$\frac{\partial F_1}{\partial \eta} = \alpha^2 \frac{\partial X}{\partial (\lambda_I^*)^2} (0, \eta \alpha^2, k_2) - \frac{k_1}{(1+\eta)^2} < 0,$$

which implies that

(4.20)
$$\widehat{F}_{1}(\eta, \alpha, k_{1}, k_{2}) \begin{cases} > 0, \text{ if } \eta < \eta_{1}^{*}(\alpha), \\ = 0, \text{ if } \eta = \eta_{1}^{*}(\alpha), \\ < 0, \text{ if } \eta > \eta_{1}^{*}(\alpha). \end{cases}$$

One can check that

(4.21)
$$\widehat{F}_1(1,\alpha,k_1,k_2) = X(0,\alpha^2,k_2) + \frac{3k_1}{2} - \rho_0^*,$$

which is decreasing with respect to $\alpha \in (0, +\infty)$ due to Proposition 3.1.

Suppose that $k_1 \ge 2\rho_0^*/3$ holds. Since $X(0, \alpha^2, k_2)$ is positive, we have $\widehat{F}_1(1, \alpha, k_1, k_2) > 0$. Then, by (4.20), $1 < \eta_1^*(\alpha)$ or equivalently $\alpha < \lambda_{I,1}^*(\alpha)$.

Suppose that $0 < k_1 < 2\rho_0^*/3$ and $k_1 \neq \rho_0^*/2$ hold. For one hand, we have

$$\lim_{\alpha \to +\infty} \widehat{F}_1(1, \alpha, k_1, k_2) = \frac{3k_1}{2} - \rho_0^* < 0.$$

On the other hand, we have

(4.22)
$$\lim_{\alpha \to 0^+} \widehat{F}_1(1, \alpha, k_1, k_2) = X(0, 0, k_2) + \frac{3k_1}{2} - \rho_0^* \begin{cases} > 0, & \text{if } k_2 \in (0, \widehat{k}_2^*), \\ = 0, & \text{if } k_2 = \widehat{k}_2^*, \\ < 0, & \text{if } k_2 \in (\widehat{k}_2^*, \infty), \end{cases}$$

where $\hat{k}_2^* = \hat{k}_2^*(k_1)$ is the unique solution of

(4.23)
$$X(0,0,\hat{k}_2^*) = \rho_0^* - \frac{3k_1}{2}.$$

The existence and uniqueness of \hat{k}_2^* is really induced by the monotonicity of $X(0, 0, k_2)$ on k_2 (see Proposition 3.1) and the facts that

$$X(0,0,0) > \rho_0^* > \rho_0^* - \frac{3k_1}{2}, \ X(0,0,\infty) = 0 < \rho_0^* - \frac{3k_1}{2}.$$

Moreover, by comparing (3.25) with (4.23), we conclude that $\hat{k}_2^* < \xi(k_1)$ for $k_1 \in (0, \rho_0^*/2)$.

In summary, we have

(1). In case of $k_2 \in (0, \widehat{k}_2^*)$, we have $\lim_{\alpha \to 0^+} \widehat{F}_1(1, \alpha, k_1, k_2) > 0$, which indicates that there exists a unique solution $\alpha_1 = \alpha_1(k_1, k_2) \in (0, \infty)$ such that

(4.24)
$$\widehat{F}_{1}(1,\alpha,k_{1},k_{2}) \begin{cases} >0, & \text{if } \alpha \in (0,\alpha_{1}), \\ =0, & \text{if } \alpha = \alpha_{1}, \\ <0, & \text{if } \alpha \in (\alpha_{1},+\infty). \end{cases}$$

since $\frac{\partial \widehat{F}_1(1,\alpha,k_1,k_2)}{\partial \alpha} < 0$. Then, for any $\alpha \in (0,\alpha_1)$, by (4.24) and (4.20), we have $1 < \eta_1^*(\alpha)$ or equivalently $\alpha < \lambda_{I,1}^*(\alpha)$. For any $\alpha \in (\alpha_1, +\infty)$, we have $1 > \eta_1^*(\alpha)$ or equivalently $\alpha > \lambda_{I,1}^*(\alpha)$. In particular, at $\alpha = \alpha_1$, we have $\alpha = \lambda_{I,1}^*(\alpha)$.

Moreover, one can check that $\alpha_1(k_1, k_2)$ is decreasing in k_2 , and

(4.25)
$$\lim_{k_2 \to (\widehat{k}_2^*)^-} \alpha_1(k_1, k_2) = 0.$$

(2). In case of $k_2 \in [\widehat{k}_2^*, \infty)$, we have $\lim_{\alpha \to 0^+} \widehat{F}_1(1, \alpha, k_1, k_2) \leq 0$. Thus, for any $\alpha \in (0, +\infty)$, $\widehat{F}_1(1, \alpha, k_1, k_2) < 0$. Then, by (4.24) and (4.20), we have $1 > \eta_1^*(\alpha)$ or equivalently $\alpha > \lambda_{I,1}^*(\alpha)$.

We have the following lemma on the properties of $\lambda_{I,2}^*(\alpha)$ (see Fig. 4).

Lemma 4.7. Suppose that $\tau \in (\tau^*, +\infty)$, $(k_1, k_2) \in \Gamma_2 \cup \Gamma_{3-1}$, and $\alpha \in (0, \alpha_0)$, where α_0 is defined in (4.17). Let $\lambda_{I,2}^*(\alpha)$ be the solution of (4.15b). Then,

(4.26)
$$\lim_{\alpha \to 0^+} \lambda_{I,2}^*(\alpha) = \lim_{\alpha \to \alpha_0^-} \lambda_{I,2}^*(\alpha) = 0,$$



Figure 4: The rough figures of $\lambda_{I,1}^*(\alpha)$ (red) and $\lambda_{I,2}^*(\alpha)$ (blue). (a). $k_1 \ge 2\rho_0^*/3$; (b). $k_1 \in (0, \rho_0^*/2) \cup (\rho_0^*/2, 2\rho_0^*/3)$ and $k_2 < \hat{k}_2^*$; (c). $k_1 \in (0, \rho_0^*/2), k_2 \in [\hat{k}_2^*, \xi(k_1))$, or $k_1 \in (\rho_0^*/2, 2\rho_0^*/3), k_2 \in [\hat{k}_2^*, \infty)$.

and there exists a unique $\alpha_2 \in (0, \alpha_0)$, such that $\lambda_{I,2}^*(\alpha)$ is increasing in $(0, \alpha_2)$, while decreasing in (α_2, α_0) , and at $\alpha = \alpha_2$, $\lambda_{I,2}^*(\alpha)$ attains its local maximal value. Moreover,

(4.27)
$$\lambda_{I,2}^*(\alpha) \begin{cases} > \alpha, & \text{if } \alpha \in (0, \alpha_2), \\ = \alpha, & \text{if } \alpha = \alpha_2, \\ < \alpha, & \text{if } \alpha \in (\alpha_2, \alpha_0). \end{cases}$$

Proof. 1. By (4.17), one can directly check that $\lim_{\alpha \to \alpha_0^-} \lambda_{I,2}^*(\alpha) = 0.$

We then show that $\lambda_{I,2}^*(\alpha) \to 0$ as $\alpha \to 0^+$. Suppose not. Then, $\lambda_{I,2}^*(\alpha) \to c$ for some $c \in (0,\infty]$. By (4.15b), we have

(4.28)
$$F_2((\lambda_{I,2}^*(\alpha))^2, \alpha, k_1, k_2) := Y(0, (\lambda_{I,2}^*(\alpha))^2, k_2) - \tau + \frac{k_1 \alpha}{\alpha^2 + (\lambda_{I,2}^*(\alpha))^2} = 0.$$

Then, from (4.28) and $\lambda_{I,2}^*(\alpha) \to c$, we have $Y(0,c,k_2) - \tau = 0$. However, this is impossible since $\tau > \tau^* = Y(0,0,0) > Y(0,c,k_2)$.

2. For $\alpha \in (0, \alpha_0)$, we define

(4.29)
$$\eta_2^*(\alpha) = \left(\frac{\lambda_{I,2}^*(\alpha)}{\alpha}\right)^2.$$

For $(k_1, k_2) \in \Gamma_2 \cup \Gamma_{3-1}$ and $\alpha \in (0, \alpha_0)$, we define

(4.30)
$$\widehat{F}_{2}(\eta, \alpha, k_{1}, k_{2}) := Y(0, \eta \alpha^{2}, k_{2}) - \tau + \frac{k_{1}}{\alpha + \eta \alpha},$$

where $\eta > 0$.

Then, by (4.28), we have

(4.31)
$$F_2(\eta_2^*(\alpha), \alpha, k_1, k_2) = 0.$$

By Proposition 3.1, we have

(4.32)
$$\frac{\partial \widehat{F}_2}{\partial \eta} = \alpha^2 \frac{\partial Y}{\partial (\lambda_I^*)^2} (0, \eta \alpha^2, k_2) - \frac{k_1}{(1+\eta)^2 \alpha} < 0.$$

Then, by (4.31) and (4.32), we have

(4.33)
$$\widehat{F}_{2}(\eta, \alpha, k_{1}, k_{2}) \begin{cases} > 0, \text{ if } \eta < \eta_{2}^{*}(\alpha), \\ = 0, \text{ if } \eta = \eta_{2}^{*}(\alpha), \\ < 0, \text{ if } \eta > \eta_{2}^{*}(\alpha). \end{cases}$$

On the other hand, we can check that

(4.34)
$$\widehat{F}_2(1,\alpha,k_1,k_2) = Y(0,\alpha^2,k_2) + \frac{k_1}{2\alpha} - \tau.$$

By Proposition 3.1, $\widehat{F}_2(1, \alpha, k_1, k_2)$ is decreasing in α . Then, by

$$\lim_{\alpha \to 0^+} \widehat{F}_2(1, \alpha, k_1, k_2) = +\infty,$$

$$\lim_{\alpha \to \alpha_0^-} \widehat{F}_2(1, \alpha, k_1, k_2) = Y(0, \alpha_0^2, k_2) - \frac{Y(0, 0, k_2)}{2} - \tau < \frac{Y(0, 0, k_2) - \tau}{2} < 0.$$

there exists a unique $\alpha_2 = \alpha_2(k_1, k_2) \in (0, \alpha_0)$ satisfying

(4.35)
$$\widehat{F}_2(1,\alpha,k_1,k_2) \begin{cases} >0, \text{ if } \alpha < \alpha_2, \\ =0, \text{ if } \alpha = \alpha_2, \\ <0, \text{ if } \alpha > \alpha_2. \end{cases}$$

Therefore, combining (4.35) with (4.33) and (4.29), we can obtain (4.27).

Moreover, by differentiating $\widehat{F}_2(1, \alpha_2, k_1, k_2) = 0$ with respect to k_2 (resp., k_1), we can conclude that $\alpha_2 = \alpha_2(k_1, k_2)$ is decreasing (resp., increasing) in k_2 (resp., k_1). In particular, it follows from (4.34) that

(4.36)
$$\lim_{k_2 \to +\infty} \alpha_2(k_1, k_2) = \frac{k_1}{2\tau}.$$

Finally, we show the monotone properties of $\lambda_{I,2}^*(\alpha)$ with respect to α . Differentiating (4.28) with respect to α yields

$$\frac{d\lambda_{I,2}^*(\alpha)}{d\alpha} = \frac{k_1(\alpha + \lambda_{I,2}^*(\alpha))(\alpha - \lambda_{I,2}^*(\alpha))}{2\lambda_{I,2}^*(\alpha)\left[\alpha^2 + (\lambda_{I,2}^*(\alpha))^2\right]^2 \frac{\partial Y(0,(\lambda_I^*)^2,k_2)}{\partial(\lambda_I^*)^2}} \begin{cases} > 0, \text{ if } \alpha < \alpha_2 \\ = 0, \text{ if } \alpha = \alpha_2 \\ < 0, \text{ if } \alpha > \alpha_2 \end{cases}$$

where $\frac{\partial Y(0,(\lambda_I^*)^2,k_2)}{\partial (\lambda_I^*)^2} < 0$ is used.

r	-	
L		

We have the following two lemmas.

Lemma 4.8. Suppose that $\tau \in (\tau^*, +\infty)$ and

(4.37)
$$k_1 \in (0, \rho_0^*/2) \cup (\rho_0^*/2, 2\rho_0^*/3), \quad k_2 \in (k_2^* - \kappa_1, \widehat{k}_2^*),$$

where $\kappa_1 > 0$ is sufficiently small. Then, there exist 2m (counting multiplicity, $m \geq 1$ is an integer) points $\alpha_j^* \in (0, \alpha_0), \ 1 \leq j \leq 2m$, satisfying

$$0 < \alpha_1^* \le \alpha_2^* \le \dots \le \alpha_{2m-1}^* < \alpha_2 < \alpha_{2m}^* < \alpha_0,$$

such that at $\alpha = \alpha_j^*$, $1 \leq j \leq 2m$, $\lambda_{I,1}^*(\alpha) = \lambda_{I,2}^*(\alpha)$. In particular, $0 < \lambda_{I,1}^*(\alpha_{2m}^*) = \lambda_{I,2}^*(\alpha_{2m}^*) < 1$ α_{2m}^* .

Proof. Since $\alpha_2(k_1, k_2)$ is decreasing in k_2 , by (4.36) and (4.25), we have

$$\alpha_2(k_1, \hat{k}_2^*) > \lim_{k_2 \to +\infty} \alpha_2(k_1, k_2) = \frac{k_1}{2\tau} > 0 = \alpha_1(k_1, \hat{k}_2^*),$$

which indicates that $\alpha_2 = \alpha_2(k_1, k_2) > \alpha_1 = \alpha_1(k_1, k_2)$ for $k_2 \in (\widehat{k}_2^* - \kappa_1, \widehat{k}_2^*)$ with some sufficiently small $\kappa_1 > 0$.

Under (4.37), we can obtain the existence of 2m (counting multiplicity, $m \ge 1$ is an integer) points $\alpha_j^* \in (0, \alpha_0), 1 \leq j \leq 2m$, such that at $\alpha = \alpha_j^*, 1 \leq j \leq 2m, \lambda_{I,1}^*(\alpha) = \lambda_{I,2}^*(\alpha)$. One can see Fig. 4b. Moreover, $\lambda_{I,1}^*(\alpha) - \lambda_{I,2}^*(\alpha)$ is increasing in $\alpha \in (\alpha_2, \alpha_0)$ and satisfies

$$\lambda_{I,1}^*(\alpha_2) - \lambda_{I,2}^*(\alpha_2) < 0, \quad \lambda_{I,1}^*(\alpha_0) - \lambda_{I,2}^*(\alpha_0) > 0.$$

Then, there exists a unique α such that $\lambda_{I,1}^*(\alpha) - \lambda_{I,2}^*(\alpha) = 0$ in (α_2, α_0) . This proves that $\alpha_{2m}^* \in (\alpha_2, \alpha_0)$ but $\alpha_{2m-1}^* \notin (\alpha_2, \alpha_0)$. In particular, $\lambda_{I,1}^*(\alpha_{2m}^*) < \alpha_{2m}^*$.

Lemma 4.9. Suppose that $\tau \in (\tau^*, +\infty)$ and that one of the following conditions holds true:

 $k_1 \in (0, \rho_0^*/2), \ k_2 \in [\widehat{k}_2^*, \xi(k_1)),$ (4.38)

(4.39)
$$k_1 \in (\rho_0^*/2, 2\rho_0^*/3), \ k_2 \in [\widehat{k}_2^*, \infty),$$

 $\kappa_1 \in (\rho_0/2, 2\rho_0/3), \ \kappa_2 \in [\kappa_2, \infty),$ $k_1 \in [2\rho_0^*/3, \rho_0^*), \ k_2 \text{ is sufficiently large.}$ (4.40)

Then, there exists a unique $\alpha^* \in (0, \alpha_0)$, such that at $\alpha = \alpha^*$, the eigenvalue problem (4.6) has a pair of purely imaginary eigenvalues $\pm i\lambda^*$. Moreover,

- 1. if either (4.38) or (4.39) holds, then $\alpha_2 < \lambda^* < \alpha^* < \alpha_0$.
- 2. if either (4.39) or (4.40) holds, then in the limit of $k_2 \to +\infty$, we have

(4.41)
$$\alpha^* = \frac{\rho_0^* - k_1}{\tau}, \ \lambda^* = \sqrt{\frac{(2k_1 - \rho_0^*)(\rho_0^* - k_1)}{\tau^2}}.$$

Proof. Suppose that (4.38) holds. Then, by Lemma 4.7, for any $\alpha \in (0, \alpha_2)$, we have $\lambda_{I,2}^*(\alpha) > \alpha$. By Lemma 4.6, for any $\alpha \in (0, \alpha_2)$, we have $\lambda_{I,1}^*(\alpha) < \alpha$. Thus, in $(0, \alpha_2)$, $\lambda_{I,1}^*(\alpha) < \alpha < \lambda_{I,2}^*(\alpha)$. By Lemma 4.6, $\lambda_{I,1}^*(\alpha)$ is increasing in (α_2, α_0) and $\lim_{\alpha \to \alpha_0^-} \lambda_{I,1}^*(\alpha) > 0$. By Lemma 4.7, $\lambda_{I,2}^*(\alpha)$ is decreasing in (α_2, α_0) and $\lim_{\alpha \to \alpha_0^-} \lambda_{I,2}^*(\alpha) = 0$. Moreover, $\lim_{\alpha \to \alpha_2^+} \lambda_{I,1}^*(\alpha) < \lim_{\alpha \to \alpha_2^+} \lambda_{I,2}^*(\alpha)$. Then, in (α_2, α_0) , there exists a unique α^* , such that at $\alpha = \alpha^*$, we have $\lambda_{I,1}^*(\alpha^*) = \lambda_{I,2}^*(\alpha^*)$.

Similarly, we can show the existence and uniqueness of α^* in the case of (4.39). And it is clear to show that under either (4.38) or (4.39), it holds true that $\alpha_2 < \lambda_{I,1}^*(\alpha^*) = \lambda_{I,2}^*(\alpha^*) < \alpha^* < \alpha_0$.

Suppose that either (4.39) or (4.40) holds. Then,

$$\lim_{k_2 \to \infty} \lambda_{I,1}^*(\alpha) = \alpha \sqrt{\frac{2k_1 - \rho_0^*}{\rho_0^* - k_1}}, \ \lim_{k_2 \to \infty} \lambda_{I,2}^*(\alpha) = \sqrt{\frac{k_1 \alpha}{\tau} - \alpha^2}.$$

Solving $\lim_{k_2 \to \infty} \lambda_{I,1}^*(\alpha) = \lim_{k_2 \to \infty} \lambda_{I,2}^*(\alpha)$, we can obtain (4.41). This completes the proof. \Box

Define

(4.42)
$$\alpha_H^* := \begin{cases} \alpha_{2m}^*, & \text{in case of Lemma 4.8,} \\ \alpha^*, & \text{in case of Lemma 4.9,} \end{cases} \lambda_{I,H}^* = \lambda_{I,1}^*(\alpha_H^*) = \lambda_{I,2}^*(\alpha_H^*) > 0.$$

We have the following results.

Lemma 4.10. Let α_H^* and λ_H^* be defined in (4.42). Then, there exists a sufficiently small $\kappa_2 > 0$, such that for any $\alpha \in (\alpha_H^* - \kappa_2, \alpha_H^* + \kappa_2)$, the eigenvalue problem (4.6) has the simple complex conjugate eigenvalues $\lambda^*(\alpha) := \lambda_R^*(\alpha) \pm i\lambda_I^*(\alpha)$ such that $\lambda_R^*(\alpha_H^*) = 0$, $\lambda_I^*(\alpha_H^*) = \lambda_{I,H}^*$. Moreover, if in addition $d > d^*$ for some $d^* \ge d_0$, where d_0 is defined in Theorem 2.1, then $\frac{d}{d\alpha}\lambda_R^*(\alpha_H^*) < 0$.

Proof. Similar to (3.30), we can obtain the following complex SLEP system of (4.6)

(4.43)
$$\mathcal{G}^{*}(\lambda^{*},\alpha) := \rho_{0}^{*} - \tau \lambda^{*} - k_{1} \left(\frac{\alpha}{\alpha + \lambda^{*}} + 1\right) - \sum_{n=0}^{\infty} \frac{c_{1}^{*} c_{2}^{*}}{\gamma_{n} + \lambda^{*} + 2k_{2}} \psi_{n}^{2}(x^{*}) = 0,$$

where $\lambda^* = \lambda_R^* + i\lambda_I^*$.

We only consider the existence of the desired eigenvalue $\lambda_R^*(\alpha) + i\lambda_I^*(\alpha)$, while the existence of the eigenvalue $\lambda_R^*(\alpha) - i\lambda_I^*(\alpha)$ can be proved similarly.

Differentiating (4.43) with respect to λ^* at $(i\lambda_{I,H}^*, \alpha_H^*)$ yields

(4.44)
$$\frac{\partial \mathcal{G}^*}{\partial \lambda^*} (i\lambda_{I,H}^*, \alpha_H^*) = -\tau + \frac{k_1 \alpha_H^*}{(\alpha_H^* + i\lambda_{I,H}^*)^2} + \sum_{n=0}^{\infty} \frac{c_1^* c_2^*}{(\gamma_n + i\lambda_{I,H}^* + 2k_2)^2} \psi_n^2(x^*).$$

One can directly check that

$$(4.45) \operatorname{Im}\left\{\frac{\partial \mathcal{G}^{*}}{\partial \lambda^{*}}(i\lambda_{I,H}^{*},\alpha_{H}^{*})\right\} = -\frac{2k_{1}(\alpha_{H}^{*})^{2}\lambda_{I,H}^{*}}{\left((\alpha_{H}^{*})^{2} + (\lambda_{I,H}^{*})^{2}\right)^{2}} - \sum_{n=0}^{\infty} \frac{2c_{1}^{*}c_{2}^{*}\lambda_{I,H}^{*}(\gamma_{n}+2k_{2})}{\left((\gamma_{n}+2k_{2})^{2} + (\lambda_{I,H}^{*})^{2}\right)^{2}}\psi_{n}^{2}(x^{*}) < 0.$$

Then, $\frac{\partial \mathcal{G}^*}{\partial \lambda^*}(i\lambda_{I,H}^*, \alpha_H^*) \neq 0$. Therefore, by the implicit function theorem, there exists a sufficiently small $\kappa_2 > 0$, such that for any $\alpha \in (\alpha_H^* - \kappa_2, \alpha_H^* + \kappa_2)$, the eigenvalue problem (4.6) has the complex eigenvalue $\lambda^*(\alpha) = \lambda_R^*(\alpha) + i\lambda_I^*(\alpha)$ such that $\lambda_R^*(\alpha_H^*) = 0$, $\lambda_I^*(\alpha_H^*) = \lambda_{I,H}^*$.

Next, we show that the transversality condition holds. That is, $\frac{d}{d\alpha}\lambda_R^*(\alpha_H^*) < 0$.

Substituting $\lambda^*(\alpha)$ into (4.43), differentiating (4.43) with respect to α , and setting ($\lambda^*(\alpha), \alpha$) = $(i\lambda_H^*, \alpha_H^*)$, we can obtain

(4.46)
$$\frac{\partial \mathcal{G}^*}{\partial \lambda^*} (i\lambda_H^*, \alpha_H^*) \frac{d\lambda^*(\alpha_H^*)}{d\alpha} + \frac{\partial \mathcal{G}^*}{\partial \alpha} (i\lambda_H^*, \alpha_H^*) = 0,$$

where $\frac{\partial \mathcal{G}^*}{\partial \lambda^*}(i\lambda_H^*, \alpha_H^*)$ is given by (4.44), and $\frac{\partial \mathcal{G}^*}{\partial \alpha}(i\lambda_H^*, \alpha_H^*) = -\frac{k_1 i \lambda_H^*}{(i \lambda_H^* + \alpha_H^*)^2}$.

Separating the real and imaginary parts of (4.46), we have

(4.47)
$$\frac{d}{d\alpha}\lambda_R^*(\alpha_H^*) = -\frac{\operatorname{Re}\left\{\frac{\partial\mathcal{G}^*}{\partial\lambda^*}\right\}\operatorname{Re}\left\{\frac{\partial\mathcal{G}^*}{\partial\alpha}\right\} + \operatorname{Im}\left\{\frac{\partial\mathcal{G}^*}{\partial\lambda^*}\right\}\operatorname{Im}\left\{\frac{\partial\mathcal{G}^*}{\partial\alpha}\right\}}{\left(\operatorname{Re}\left\{\frac{\partial\mathcal{G}^*}{\partial\lambda^*}\right\}\right)^2 + \left(\operatorname{Im}\left\{\frac{\partial\mathcal{G}^*}{\partial\lambda^*}\right\}\right)^2}$$

where $\frac{\partial \mathcal{G}^*}{\partial \lambda^*}$ and $\frac{\partial \mathcal{G}^*}{\partial \alpha}$ are evaluated at $(i\lambda_{I,H}^*, \alpha_H^*)$. In particular, $\operatorname{Im}\left\{\frac{\partial \mathcal{G}^*}{\partial \lambda^*}\right\}$ is given by (4.45), and

$$\operatorname{Re}\left\{\frac{\partial\mathcal{G}^{*}}{\partial\lambda^{*}}\right\} = -\tau + \frac{\alpha_{H}^{*}k_{1}\left((\alpha_{H}^{*})^{2} - (\lambda_{I,H}^{*})^{2}\right)}{\left((\alpha_{H}^{*})^{2} + (\lambda_{I,H}^{*})^{2}\right)^{2}} + c_{1}^{*}c_{2}^{*}\sum_{n=0}^{\infty} \frac{(\gamma_{n} + 2k_{2})^{2} - (\lambda_{I,H}^{*})^{2}}{\left((\gamma_{n} + 2k_{2})^{2} + (\lambda_{I,H}^{*})^{2}\right)^{2}}\psi_{n}^{2}(x^{*}),$$

$$\operatorname{Re}\left\{\frac{\partial\mathcal{G}^{*}}{\partial\alpha}\right\} = \frac{-2\alpha_{H}^{*}k_{1}(\lambda_{I,H}^{*})^{2}}{\left((\alpha_{H}^{*})^{2} + (\lambda_{I,H}^{*})^{2}\right)^{2}}, \operatorname{Im}\left\{\frac{\partial\mathcal{G}^{*}}{\partial\alpha}\right\} = \frac{\lambda_{I,H}^{*}k_{1}\left((\lambda_{I,H}^{*})^{2} - (\alpha_{H}^{*})^{2}\right)}{\left((\alpha_{H}^{*})^{2} + (\lambda_{I,H}^{*})^{2}\right)^{2}}.$$

One can check that

(4.48)
$$\frac{d}{d\alpha}\lambda_R^*(\alpha_H^*) = -\frac{I_1 + I_2}{\left(\operatorname{Re}\left\{\frac{\partial \mathcal{G}^*}{\partial \lambda^*}(i\lambda_{I,H}^*, \alpha_H^*)\right\}\right)^2 + \left(\operatorname{Im}\left\{\frac{\partial \mathcal{G}^*}{\partial \lambda^*}(i\lambda_{I,H}^*, \alpha_H^*)\right\}\right)^2}$$

where

(4.49)

$$I_{1} := \left(\tau - c_{1}^{*}c_{2}^{*}\sum_{n=0}^{\infty} \frac{(\gamma_{n} + 2k_{2})^{2} - (\lambda_{I,H}^{*})^{2}}{((\gamma_{n} + 2k_{2})^{2} + (\lambda_{I,H}^{*})^{2})^{2}}\psi_{n}^{2}(x^{*})\right) \frac{2\alpha_{H}^{*}k_{1}(\lambda_{I,H}^{*})^{2}}{((\alpha_{H}^{*})^{2} + (\lambda_{I,H}^{*})^{2})^{2}} I_{2} := \frac{2k_{1}(\lambda_{I,H}^{*})^{2}((\alpha_{H}^{*})^{2} - (\lambda_{I,H}^{*})^{2})}{((\alpha_{H}^{*})^{2} + (\lambda_{I,H}^{*})^{2})^{2}} \sum_{n=0}^{\infty} \frac{c_{1}^{*}c_{2}^{*}(\gamma_{n} + 2k_{2})}{((\gamma_{n} + 2k_{2})^{2} + (\lambda_{I,H}^{*})^{2})^{2}}\psi_{n}^{2}(x^{*}).$$

Since $\lambda_{I,H}^* < \alpha_H^*$, we have $I_2 > 0$. It remains to show that $I_1 > 0$.

Case 1: For any integer $n \ge 0$, if $k_2 = +\infty$, then I_1 is reduced to

(4.50)
$$I_1 = \frac{2\tau \alpha_H^* k_1 (\lambda_{I,H}^*)^2}{((\alpha_H^*)^2 + (\lambda_{I,H}^*)^2)^2} > 0.$$

Case 2: For any integer $n \ge 0$ and $k_2 \in (0, +\infty)$.

It is clear from (4.17), Lemmas 4.8 and 4.9 that

(4.51)
$$\lambda_{I,H}^* < \alpha_H^* < \alpha_0 = \alpha_0(k_2) := \frac{k_1}{\tau - Y(0, 0, k_2)}$$

Define

$$h(k_2) := \frac{\alpha_0(k_2)}{\sqrt{3}} - 2k_2.$$

Then, one can check that $h'(k_2) < 0$ since $\alpha'_0(k_2) < 0$. Note that

$$\lim_{k_2 \to 0^+} h(k_2) = \frac{k_1}{\sqrt{3}(\tau - Y(0, 0, 0))} = \frac{k_1}{\sqrt{3}(\tau - \tau^*)} > 0.$$

Recall that $\gamma_0 > 0$ is the principle eigenvalue of the operator $-d\frac{d^2}{dx^2} - (f_u^*g_v^* - f_v^*g_u^*)/f_u^*$ subject to the homogeneous Neumann boundary condition in the $L^2(0, \ell)$ sense. In particular, γ_0 is increasing in d. On the other hand, one can check that τ^* is decreasing in d (see (3.18)), and so is $\lim_{k_2 \to 0^+} h(k_2)$. Then, there exists a $d_1 > 0$, such that when $d > d_1$, $\gamma_0 > \lim_{k_2 \to 0^+} h(k_2)$.

Choose $d^* = \max\{d_0, d_1\}$. Then, for any $d > d^*$, we have $\gamma_0 > h(k_2)$. Thus, for any $n \ge 0$, by $\alpha_0 = \alpha_0(k_2) > \lambda_{I,H}^*$, we have

$$\gamma_n \ge \gamma_0 > h(k_2) = \frac{\alpha_0(k_2)}{\sqrt{3}} - 2k_2 > \frac{\lambda_{I,H}^*}{\sqrt{3}} - 2k_2,$$

which implies that $\lambda_{I,H}^* < \sqrt{3}(\gamma_n + 2k_2)$ or equivalently $(\lambda_{I,H}^*)^2 \in (0, 3(\gamma_n + 2k_2)^2)$.

One can check that, if $(\lambda_I^*)^2 \in (0, 3(\gamma_n + 2k_2)^2)$, then

$$\frac{d}{d(\lambda_I^*)^2} \left\{ \frac{(\gamma_n + 2k_2)^2 - (\lambda_I^*)^2}{\left((\gamma_n + 2k_2)^2 + (\lambda_I^*)^2\right)^2} \right\} = \frac{\left((\lambda_I^*)^2 - 3(\gamma_n + 2k_2)^2\right)\left((\lambda_I^*)^2 + (\gamma_n + 2k_2)^2\right)}{\left((\gamma_n + 2k_2)^2 + (\lambda_I^*)^2\right)^4} < 0,$$

which yields

$$\frac{(\gamma_n + 2k_2)^2 - (\lambda_I^*)^2}{\left((\gamma_n + 2k_2)^2 + (\lambda_I^*)^2\right)^2} \le \frac{(\gamma_n + 2k_2)^2 - (\lambda_I^*)^2}{\left((\gamma_n + 2k_2)^2 + (\lambda_I^*)^2\right)^2}\Big|_{(\lambda_I^*)^2 = 0} = \frac{1}{(\gamma_n + 2k_2)^2}.$$

Then, by $\lambda_{I,H}^* < \sqrt{3}(\gamma_n + 2k_2)$, we can obtain

$$c_1^* c_2^* \sum_{n=0}^{\infty} \frac{(\gamma_n + 2k_2)^2 - (\lambda_{I,H}^*)^2}{\left((\gamma_n + 2k_2)^2 + (\lambda_{I,H}^*)^2\right)^2} \psi_n^2(x^*) \le c_1^* c_2^* \sum_{n=0}^{\infty} \frac{1}{(\gamma_n + 2k_2)^2} \psi_n^2(x^*) = Y(0,0,k_2),$$

which leads to

$$\tau - c_1^* c_2^* \sum_{n=0}^{\infty} \frac{(\gamma_n + 2k_2)^2 - (\lambda_{I,H}^*)^2}{\left((\gamma_n + 2k_2)^2 + (\lambda_{I,H}^*)^2\right)^2} \psi_n^2(x^*) \ge \tau - Y(0,0,k_2) > \tau - \tau^* > 0.$$

Then, $I_1 > 0$.

So far, we have proved that under either $k_2 = +\infty$ or $k_2 \in (0, +\infty)$, $I_1 > 0$ so that $\frac{d}{d\alpha} \lambda_R^*(\alpha_H^*) < 0$. This completes the proof.



Figure 5: Stability diagram of the symmetric solution $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ with respect to (1.5) at any fixed $\epsilon \in (0, \epsilon_0)$, where Γ_1 , Γ_2 , Γ_{3-1} and Γ_{3-2} are explicitly given by (4.16). The symmetric solution is always unstable in Γ_1 , while stable in Γ_{3-2} regardless of $\alpha > 0$. In the dotted subregion of $\Gamma_2 \cup \Gamma_{3-1}$, i.e. $(k_1, k_2) \in ((0, \rho_0^*/2) \times (\hat{k}_2^*, \xi(k_1))) \cup ((\rho_0^*/2, 2\rho_0^*/3) \times (\hat{k}_2^*, \infty))$, system (1.5) undergoes Hopf bifurcation near the symmetric solution at $\alpha = \alpha_H^{\epsilon}$, which can also be triggered if k_2 is slightly less than \hat{k}_2^* ; or k_2 is sufficiently large with $k_1 \in [2\rho_0^*/3, \rho_0^*)$.

As the consequence of Lemma 4.10, we have the following results (in the sense of $\epsilon \to 0^+$).

Lemma 4.11. Suppose that all the conditions of Lemma 4.10 hold. Then, the following conditions hold true:

- 1. When $\alpha > \alpha_H^*$, all the eigenvalues of the limiting eigenvalue problem (SLEP-2) have negative real parts.
- 2. At $\alpha = \alpha_H^*$, the limiting eigenvalue problem (SLEP-2) has a pair of simple purely imaginary eigenvalues $\pm i\lambda_H^*$, with all the remaining eigenvalues having strictly negative real parts.
- 3. When $\alpha \in (\alpha_H^* \kappa_3, \alpha_H^*)$ for some sufficiently small $\kappa_3 > 0$, the eigenvalue problem (SLEP-2) has one eigenvalue with strictly positive real part. In particular, if (4.39) holds, then for any $\alpha \in (0, \alpha_H^*)$, the limiting eigenvalue problem (SLEP-2) has one eigenvalue with strictly positive real part.

Finally, we shall state the main results on Hopf bifurcation of (1.5) around the symmetric solution $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ for any fixed $\epsilon \in (0, \epsilon_0)$, see Fig. 5.

Theorem 4.12. Suppose that $\tau \in (\tau^*, +\infty)$ and $d > d^*$, where d^* is defined in Lemma 4.10. Let (k_1, k_2) be fixed such that one of (4.37)-(4.40) is satisfied. Then for any fixed $\epsilon \in (0, \epsilon_0)$, there exists a critical value $\alpha_H^{\epsilon} > 0$, such that the solution $(\tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon), \tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon))$ is stable with respect to system (1.5) when $\alpha > \alpha_H^{\epsilon}$, while unstable when α is slightly less than α_H^{ϵ} . In particular, system (1.5) undergoes a Hopf bifurcation at $\alpha = \alpha_H^{\epsilon}$ for $\epsilon \in (0, \epsilon_0)$. That is to say, a family of periodic solutions of (1.5) occurs around $(\tilde{u}(x; \epsilon), \tilde{v}(x; \epsilon), \tilde{v}(x; \epsilon))$.

Proof. Similar to (3.38), for (4.6), we can derive the following complex characteristic equation

(4.52)
$$\mathcal{G}(\lambda,\alpha,\epsilon) := \rho(\epsilon) - k_1 \left(1 + \frac{\alpha}{\lambda + \alpha}\right) - \tau \lambda - \left\langle (T_{\lambda,\alpha}^{\epsilon})^{-1} \frac{g_u^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}}, \frac{-f_v^{\epsilon} \phi_0^{\epsilon}}{\sqrt{\epsilon}} \right\rangle = 0,$$

where $(T_{\lambda,\alpha}^{\epsilon})^{-1}$, the inverse operator of

$$T_{\lambda,\alpha}^{\epsilon} := -d\frac{d^2}{dx^2} - g_v^{\epsilon} + 2k_2 + g_u^{\epsilon} \left(\epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} - \epsilon k_1 \left(1 + \frac{\alpha}{\lambda + \alpha}\right) - \epsilon \tau \lambda\right)^{\dagger} f_v^{\epsilon} + \lambda,$$

exists for $\text{Re}\lambda + 2k_2 > -\mu^*$ by Lemma 4.1 of [21], where μ^* could be chosen smaller if necessary.

One can observe that $\mathcal{G}(\lambda, \alpha, \epsilon) \to \mathcal{G}^*(\lambda, \alpha)$ as $\epsilon \to 0^+$, where $\mathcal{G}^*(\lambda, \alpha)$ is given by (4.43), then it follows that

$$\mathcal{G}(\pm i\lambda_{I,H}^*, \alpha_H^*, 0) = \mathcal{G}^*(\pm i\lambda_{I,H}^*, \alpha_H^*) = 0, \ \frac{\partial \mathcal{G}}{\partial \lambda}(\pm i\lambda_{I,H}^*, \alpha_H^*, 0) = \frac{\partial \mathcal{G}^*}{\partial \lambda}(\pm i\lambda_{I,H}^*, \alpha_H^*) \neq 0,$$

where Lemma 4.10 is used.

By the implicit function theorem, for (α, ϵ) in a small neighborhood \mathcal{O} of $(\alpha_H^*, 0)$, there exists a unique $\lambda^{\epsilon}(\alpha) := \lambda_R(\alpha, \epsilon) + i\lambda_I(\alpha, \epsilon)$, such that $\mathcal{G}(\lambda^{\epsilon}(\alpha), \alpha, \epsilon) = 0$ and in particular,

$$\lim_{\epsilon \to 0^+} \lambda^{\epsilon}(\alpha) = \lambda^*(\alpha), \ \lim_{\epsilon \to 0^+} \lambda^{\epsilon}(\alpha_H^*) = \lambda^*(\alpha_H^*) = \pm i \lambda_{I,H}^*$$

Since

$$\lambda_R(\alpha_H^*, 0) = \lambda_R^*(\alpha_H^*) = 0, \ \lambda_I(\alpha_H^*, 0) = \lambda_{I,H}^* \neq 0, \ \frac{d\lambda_R}{d\alpha}(\alpha_H^*, 0) = \frac{d\lambda_R^*}{d\alpha}(\alpha_H^*) < 0,$$

by the implicit function theorem, for small $\epsilon > 0$, there exists a unique continuous differentiable function $\alpha_H^{\epsilon} = \alpha_H(\epsilon)$ such that

$$\lambda_R(\alpha_H^{\epsilon},\epsilon) = 0, \ \lambda_{I,H}^{\epsilon} := \lambda_I(\alpha_H^{\epsilon},\epsilon) \neq 0, \ \frac{d\lambda_R}{d\alpha}(\alpha_H^{\epsilon},\epsilon) < 0.$$

Thus, at $\alpha = \alpha_H^{\epsilon}$, the eigenvalue problem (4.52) has a pair of simple purely imaginary eigenvalues and the transversality condition holds. Hence, by the Hopf bifurcation theorem [4, 6], the system undergoes Hopf bifurcation at $\alpha = \alpha_H^{\epsilon}$ for small $\epsilon > 0$.

Therefore, when $\alpha > \alpha_H^{\epsilon}$, all the eigenvalues of (4.4) have negative real parts. On the contrary, if $\alpha \in (\alpha_H^{\epsilon} - \kappa_4, \alpha_H^{\epsilon})$ for sufficiently small $\kappa_4 > 0$, the eigenvalue problem (4.4) has a pair of complex eigenvalues with positive real part.

Let $(0, \hat{\epsilon}_0)$ be the new definition region of ϵ and drop the hat of $\hat{\epsilon}_0$. Then the simplicity of the pair of critical complex eigenvalues of (4.6) for $\epsilon \in (0, \epsilon_0)$ can be obtained by the similar argument in [21], which completes the proof.

5 Concluding remarks

In this paper, we mainly consider the dynamics of a kind of two-layer coupled Lengyel-Epstein model characterizing the spatiotemporal pattern formations in the CIMA chemical reactions. We make the following concluding remarks:

- 1. In the classical activator-inhibitor system (decoupled), if the diffusion rate ϵ of the activator u is small but the diffusion rate d of the inhibitor v is large, then we can expect the occurrence of Turing patterns (see for instance [8, 9]). In this paper, for the two-layered coupled system, we can obtain the same phenomena, that is, if the inter-reactor diffusion rate k_1 of the activator u is not that large but the inter-reactor diffusion rate k_2 of the inhibitor v is large enough, then we can also find Turing patterns (see Theorem 3.3).
- 2. A slightly large k_1 tends to offset the impacts of k_2 and α in the sense that once k_1 is larger than ρ_0^* (see Theorem 4.5), the symmetric steady state solution is always stable regardless of k_2 and α .
- 3. It is well-known that delay can induce the instability of the steady state solutions and trigger Hopf bifurcations. However, in the existing literatures, most of the works focus on Hopf bifurcations branching from the constant steady state solutions. It is rare to consider the Hopf bifurcation from non-constant steady state solutions as delay varies. For our first attempt, we study in this paper how the distributed-delayed coupling can affect the stability of the non-constant steady state solution and bring out the existence of Hopf bifurcation. Another interesting question is what happens if we replace distributed-delayed coupling with the discrete-delayed coupling. This is still an open question, which will be our next attempt.
- 4. In the present paper, we mainly concentrate on the stability/Turing instability of the symmetric steady state solution $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon), \tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ affected by k_1, k_2 and α . However, we are more than willing to know how k_1 and k_2 affect the stability of the symmetric periodic solution $(\tilde{u}_p(t, x; \epsilon), \tilde{v}_p(t, x; \epsilon), \tilde{v}_p(t, x; \epsilon), \tilde{v}_p(t, x; \epsilon))$. We would like to mention that in [35], the Turing instability of the spatially homogeneous periodic solutions is considered. We conjecture that the perturbation technique used in [35] seems to work well for our model when studying the Turing instability of the symmetric periodic solution.
- 5. In system with only one kind of diffusion, it is meaningless to define Turing instability of the spatially non-homogeneous solutions (steady state solutions and periodic solutions). However, when system has more than one kind of diffusion, Turing instability of the above solutions can be well-defined.

A Appendix: The spectral properties of
$$\epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon}$$
 for small ϵ

The following results are due to Nishiura and Fujii [20] (see also Lemmas 2.2 and 2.3 of [21]):

Lemma A.1 (Spectral properties ([20, 21])). Assume that $0 < \epsilon < \epsilon_0$, so that $(\tilde{u}(x;\epsilon), \tilde{v}(x;\epsilon))$ is the singularly perturbed steady state solution of system (2.1). Then, the eigenvalue problem

$$\begin{cases} \epsilon^2 \frac{d^2 \phi}{dx^2} + f_u^{\epsilon} \phi = \mu \phi, \ x \in (0, \ell), \\ \phi'(0) = \phi'(\ell) = 0, \end{cases}$$

has complete and orthonormal pairs of eigenvalues and eigenfunctions $\{(\mu_n^{\epsilon}, \phi_n^{\epsilon}(x))\}_{n=0}^{\infty}$ in $L^2(0, \ell)$ sense. Moreover,

1. There exists a $\mu_* < 0$ independent of ϵ , such that $\mu_0^{\epsilon} > 0 > \mu_* > \mu_1^{\epsilon} > \cdots$. In particular, $\mu_0^{\epsilon} = \rho(\epsilon)\epsilon$, where $\rho(\epsilon)$ is a positive continuous function up to $\epsilon \to 0^+$ and

(A.1)
$$\rho_0^* = \lim_{\epsilon \to 0^+} \rho(\epsilon) = \frac{(\kappa^*)^2}{d} M'(\hat{v}) \int_0^{x^*} g(U^*, V^*) dx,$$

where $x^* \in (0, \ell)$ is the layer position of the reduced solution $(U^*(x), V^*(x))$, \hat{v} is the zero of $M(\hat{v})$, and κ^* is a positive constant.

2. Let F(u, v) be a smooth function of u and v. Then, as $\epsilon \to 0^+$,

$$(\epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} - \epsilon \tau \lambda)^{\dagger} (F(\widetilde{u}(x;\epsilon),\widetilde{v}(x;\epsilon))h) \to \frac{F(U^*(x),V^*(x))h}{f_u(U^*(x),V^*(x))} \text{ in strongly } L^2 \text{-sense}$$

for any function $h \in L^2(0,\ell) \cap L^{\infty}(0,\ell)$, $\tau \in \mathbb{R}$, and $\lambda \in \mathbb{C}$, where for any $\hat{h} \in L^2(0,\ell) \cap L^{\infty}(0,\ell)$,

$$(\epsilon^2 \frac{d^2}{dx^2} + f_u^{\epsilon} - \epsilon \tau \lambda)^{\dagger} \hat{h} := \sum_{n \ge 1} \frac{\langle \hat{h}, \phi_n^{\epsilon} \rangle}{\mu_0^{\epsilon} - \epsilon \tau \lambda} \phi_n^{\epsilon}.$$

3. Let $\delta(x - x^*)$ be the Dirac's delta function at $x = x^*$ and κ^* be stated in (A.1). Then, in the H^{-1} sense, it holds true that

$$\lim_{\epsilon \to 0^+} \frac{f_v^\epsilon \phi_0^\epsilon}{\sqrt{\epsilon}} = \kappa^* M'(\widehat{v}) \delta(x - x^*),$$

$$\lim_{\epsilon \to 0^+} \frac{g_u^\epsilon \phi_0^\epsilon}{\sqrt{\epsilon}} = \kappa^* \Big[h_+(\widehat{v}) - h_-(\widehat{v}) + \Big(\frac{h_-(\widehat{v})}{1 + (h_-(\widehat{v}))^2} - \frac{h_+(\widehat{v})}{1 + (h_+(\widehat{v}))^2}\Big) \widehat{v} \Big] \delta(x - x^*),$$

where $M'(\widehat{v}) < 0, \ h_+(\widehat{v}) - h_-(\widehat{v}) + \left(\frac{h_-(\widehat{v})}{1 + (h_-(\widehat{v}))^2} - \frac{h_+(\widehat{v})}{1 + (h_+(\widehat{v}))^2}\right)\widehat{v} > 0.$

References

- [1] N. Awal, I. Epstein, T. Kaper, T. Vo, Strong symmetry breaking in coupled, identical Lengyel-Epstein oscillator via
- [2] K. Bar-Eli, Coupling of chemical oscillators, J. Phys. Chem. 88 (16) (1984) 3616-3622.
- [3] V. Castets, E. Dulos, J. Boissonade, P. De Kepper, Experimental evidence of a sustained standing Turing-type noned

- [4] M. Crandall, P. Rabinowitz, The Hopf Bifurcation Theorem in Infinite Dimensions, Arch. Ration. Mech. An. 67 (1) (
- [5] L. Deoliveira, Instability of homogeneous periodic solutions of parabolic-delay equations, J. Differ. Equ. 109 (1) (1994)
- [6] B. Hassard, N. Kazarinoff, Y. Wan, Theory and Application of Hopf Bifurcation, Cambridge University Press, Cambridge, 1981.
- [7] L. Ji, Q. Li, Turing pattern formation in coupled reaction-diffusion system with distributed delays, J. Chem. Phys. 1
- [8] J. Jang, W. Ni, M. Tang, Global bifurcation and structure of Turing patterns in 1-D Lengyel-Epstein model, J. Dyna
- [9] J. Jin, J. Shi, J. Wei, F. Yi, Bifurcations of patterned solutions in a diffusive Lengyel–Epstein system of CIMA chem
- [10] M. Kawato, R. Suzuki, Two coupled neural oscillators as model of the circadian pacemaker, J. Theor. Biol. 86 (3) (19)
- [11] I. Lengyel, I. Epstein, Modeling of Turing structures in the chlorite-iodide-malonic acid-starch reaction system, Scien
- [12] I. Lengyel, I. Epstein, Diffusion-induced instability in chemically reacting systems: steady state multiplicity, oscillation
- [13] I. Lengyel, I. Epstein, A chemical approach to designing Turing patterns in reaction-diffusion systems, Proc. Natl. Ac
- [14] J. Li, H. Wang, Q. Ouyang, Square Turing patterns in reaction-diffusion systems with coupled layers, Chaos 24 (2) (2)
- [15] P. Liu, F. Yi, Q. Guo, J. Yang, W. Wu, Analysis on global exponential robust stability of reaction-diffusion neural ne
- [16] K. Maginu, Stability of spatially homogeneous periodic solutions of reaction-diffusion equations, J. Differ. Equ. 31 (1)
- [17] Y. Morita, Instability of spatially homogeneous periodic solutions to delay-diffusion equations, Lecture Notes in Num[18] Y. Morita, Destabilization of periodic solutions arising in delay-diffusion systems in several space dimensions, Japan.
- [19] M. Mimura, M. Tabata, Y. Hosono, Multiple solutions of two-point boundary value problems of Neumann type with
- [20] Y. Nishiura, H. Fujii, Stability of singularly perturbed solutions to systems of reaction-diffusion equations, SIAM J. I
- [21] Y. Nishiura, M. Mimura, Layer oscillations in reaction-diffusion systems, SIAM J. Appl. Math. 49 (2) (1989) 481-514
- [22] W. Ni, M. Tang, Turing patterns in the Lengyel-Epstein system for the CIMA reactions, Trans. Amer. Math. Soc. 35
- [23] I. Prigogine, R. Lefever, Symmetry breaking instabilities in dissipative systems. II, J. Chem. Phys. 48 (1968) 1695-17
- [24] K. Pal, S. Paul, D. Ray, Spatiotemporal antiresonance in coupled reaction-diffusion systems, Phys. Rev. E 101 (2020)
- [25] S. Roy Choudhury, C. Fosser, Turing bifurcation in nonlinear competition models with delay, Q. Appl. Math. 54 (1)
- [26] S. Ruan, R. Filfil, Dynamics of a two-neuron system with discrete and distributed delays, Physica D 191 (2004) 323-3
- [27] G. She, F. Yi, Dynamics and bifurcations in a non-degenerate homogeneous diffusive SIR rabies model, SIAM J. App
- [28] G. She, F. Yi, Stability and bifurcation analysis in a reaction-diffusion SIRS model with the general saturated incident
- [29] Q. Song, F. Yi, Spatiotemporal patterns and bifurcations of a delayed diffusive predator-prey system with fear effects
- [30] Q. Song, F. Yi, Turing instability of the periodic solutions for the general reaction-diffusion system involving nonlocal delays, submitted.
- [31] A. Turing, The chemical basis of morphogenesis, Philos. Trans. R. Soc. Lond. B 237 (1952) 37–72.
- [32] J. Tyson, S. Kauffman, Control of Mitosis by a continuous biochemical oscillation: synchronization; spatially inhomo
- [33] T. Takaishi, M. Mimura, Y. Nishiura, Pattern formation in coupled reaction-diffusion systems, Japan J. Indust. Appl
- [34] M. Wang, F. Yi, On the dynamics of the diffusive Field-Noyes model for the Belousov-Zhabotinskii reaction, J. Differ
- [35] F. Yi, Turing instability of the periodic solutions for reaction-diffusion systems with cross-diffusion and the patch me
- [36] L. Yang, M. Dolnik, A. Zhabontinsky, I. Epstein, Spatial resonances and superposition patterns in a reaction-diffusio
- [37] L. Yang, I. Epstein, Oscillatory Turing patterns in reaction-diffusion systems with two coupled layers, Phys. Rev. Let
- [38] X. Zhang, C. Li, H. Li, J. Xu, Synchronization of neural networks involving distributed-delay coupling: A distributed
- [39] S. Zhao, P. Yu, H. Wang, Spatiotemporal patterns in a Lengyel-Epstein model near a Turing-Hopf singular point, SL