# **BERNIS ESTIMATES FOR HIGHER-DIMENSIONAL DOUBLY-DEGENERATE NON-NEWTONIAN THIN-FILM EQUATIONS**

#### CHRISTINA LIENSTROMBERG AND KATERINA NIK

ABSTRACT. For the doubly-degenerate parabolic non-Newtonian thin-film equation

 $u_t + \text{div}\left(u^n | \nabla \Delta u|^{p-2} \nabla \Delta u\right) = 0,$ 

we derive (local versions) of Bernis estimates of the form

$$
\int_{\Omega} u^{n-2p} |\nabla u|^{3p} \, \mathrm{d} x + \int_{\Omega} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} \, \mathrm{d} x \le c(n, p, d) \int_{\Omega} u^{n} |\nabla \Delta u|^{p} \, \mathrm{d} x,
$$

for functions  $u \in W_p^2(\Omega)$  with Neumann boundary condition, where  $2 \le p < \frac{19}{3}$  and *n* lies in a certain range. Here, Ω *⊂* ℝ*<sup>𝑑</sup>* is a smooth convex domain with *𝑑 <* 3*𝑝*. A particularly important consequence is the estimate

$$
\int_{\Omega} |\nabla \Delta(u^{\frac{n+p}{p}})|^p \, \mathrm{d}x \le c(n,p,d) \int_{\Omega} u^n |\nabla \Delta u|^p \, \mathrm{d}x.
$$

The methods used in this article follow the approach of [\[Grü01](#page-22-0)] for the Newtonian case, while addressing the specific challenges posed by the nonlinear higher-order term  $|\nabla \Delta u|^{p-2} \nabla \Delta u$  and the additional degeneracy. The derived estimates are key to establishing further qualitative results, such as the existence of weak solutions, finite propagation of support, and the appearance of a waiting-time phenomenon.

### MSC (2020): 76A05, 76A20, 35A23, 46B70,35Q35, 35K35, 35K65

KEYWORDS: NON-NEWTONIAN FLUIDS, POWER-LAW FLUIDS, FOURTH-ORDER DEGENERATE PARABOLIC EQUATIONS, THIN-FILM EQUATIONS, BERNIS ESTIMATES

#### <span id="page-0-0"></span>1. INTRODUCTION AND MAIN RESULTS

1.1. **Background.** In this article, we derive so-called Bernis estimates for fourth-order doubly-degenerate parabolic problems of the form

$$
\begin{cases}\n u_t + \operatorname{div}\left(u^n|\nabla\Delta u|^{p-2}\nabla\Delta u\right) = 0 & \text{on } (0, \infty) \times \Omega \subset \mathbb{R}^{d+1}, \\
 \nabla u \cdot v = \nabla\Delta u \cdot v = 0 & \text{in } (0, \infty) \times \partial\Omega, \\
 u(0, \cdot) = u_0(\cdot) & \text{in } \Omega,\n\end{cases}
$$
\n(1.1)

as they arise in the modeling of non-Newtonian thin-film flows with a strain rate-dependent power-law rheology.

Problem [\(1.1\)](#page-0-0) describes the evolution of the height  $u = u(t, x) \ge 0$  of a thin layer of a viscous, non-Newtonian, and incompressible fluid over an impermeable flat bottom, as sketched in Figure [1.](#page-1-0)

The partial differential equation  $(1.1)<sub>1</sub>$  is a fourth-order quasilinear equation that is doubly-degenerate parabolic, meaning that parabolicity gets lost when either the film height *u* or the third-order term  $\nabla \Delta u$  vanishes.

The term  $u^n$  represents the mobility of the flow, with the exponent  $n > 0$  determined by the choice of the slip condition at the lower boundary (i.e., the interface between the solid bottom and the fluid film). For a no-slip condition, we have  $n = p + 1$ , while for a Navier-slip condition, we have  $n = p$ . Other values of  $n \in (p-2, p+1)$ have been proposed to model weaker or stronger slippage; see, for example, [\[Gre78](#page-22-1)]. The term  $|\nabla \Delta u|^{p-2}\nabla \Delta u$ has a dual physical interpretation: the higher-order term ∇Δ*𝑢* accounts for surface tension effects (we assume that the dynamics of the flow is driven solely by surface tension, thus neglecting gravitational effects), while its appearance in the nonlinear form  $|s|^{p-2}$ *s* is due to the non-Newtonian power-law rheology of the fluid.

The evolution equation  $(1.1)_1$  is supplemented with Neumann-type boundary conditions  $\nabla u \cdot v = 0$  and  $\nabla \Delta u \cdot$  $ν = 0$ , which represent a zero-contact angle condition and a no-flux condition at the lateral boundary (0, ∞)×*∂*Ω, respectively. Finally,  $u_0$  denotes the initial film height.

<span id="page-1-3"></span>

<span id="page-1-0"></span>FIGURE 1. Sketch of a fluid film on an impermeable solid bottom.

The natural energy inequality for non-negative solutions to  $(1.1)$  is given by

$$
\frac{1}{2} \int_{\Omega} |\nabla u(t)|^2 dx + \int_0^t \int_{\Omega} u^n |\nabla \Delta u|^p dx ds \le \frac{1}{2} \int_{\Omega} |\nabla u_0|^2 dx, \quad t \ge 0.
$$
\n(1.2)

However, even in the one-dimensional case or for the classical Newtonian thin-film equation  $(p = 2)$ , this estimate on the dissipation  $\int_{\Omega} u^n |\nabla \Delta u|^p dx$  is not sufficient for many purposes. Further estimates on the integrals  $\int_{\Omega} |\nabla (u^{\frac{n+p}{3p}})$  $\int_{\Omega}^{3p} dx, \int_{\Omega} |\Delta(u^{\frac{2(n+p)}{3p}})|$ |  $\frac{3p}{2}$  dx, and  $\int_{\Omega} |\nabla \Delta (u^{\frac{n+p}{p}})|^2$  $\vert \rho \vert$  dx are needed to obtain qualitative results such as finite propagation of support or the occurrence of a waiting-time phenomenon. See Section [1.3](#page-3-0) for more details on related results.

1.2. **Aim of the article and main results.** The main goal of this article is to derive inequalities of the form

$$
\int_{\Omega}\varphi^{3p}u^{n-2p}|\nabla u|^{3p}\, {\rm d} x+\int_{\Omega}\varphi^{3p}u^{n-\frac{p}{2}}|\Delta u|^{\frac{3p}{2}}\, {\rm d} x\leq c(n,p,d)\left(\int_{\Omega}\varphi^{3p}u^{n}|\nabla\Delta u|^{p}\, {\rm d} x+\int_{\Omega}|\nabla\varphi|^{3p}u^{n+p}\, {\rm d} x\right)
$$

for a given non-negative test function  $\varphi$ , and for positive functions  $u \in W_p^2(\Omega)$ ,  $2 \le p < \frac{19}{3}$ , satisfying the Neumann boundary condition  $\nabla u \cdot v = 0$  on  $\partial \Omega$  and having finite dissipation  $\int_{\Omega} u^n |\nabla \Delta u|^p dx < \infty$ . These are established in Theorem [1.1](#page-1-1) below.

From these inequalities, we can deduce further estimates of the form

$$
\int_{\Omega} \left| \nabla \left( u^{\frac{n+p}{3p}} \right) \right|^{3p} dx + \int_{\Omega} \left| \Delta \left( u^{\frac{2(n+p)}{3p}} \right) \right|^{3p} dx + \int_{\Omega} \left| \nabla \Delta \left( u^{\frac{n+p}{p}} \right) \right|^{p} dx \le c(n,p,d) \int_{\Omega} u^n |\nabla \Delta u|^p dx
$$

for *u* as above. This result is proven in Theorem [1.3](#page-2-0) below.

These inequalities, established in Theorems [1.1](#page-1-1) and [1.3,](#page-2-0) provide a basis for applying appropriate interpolation results, which are in turn crucial for proving qualitative properties of solutions to [\(1.1\)](#page-0-0), such as finite speed of propagation or waiting-time phenomena.

Our results generalize the findings of [\[Grü01](#page-22-0)], where the Newtonian analogue of [\(1.1\)](#page-0-0) is addressed, to non-Newtonian thin-film equations of the form [\(1.1\)](#page-0-0). In the one-dimensional non-Newtonian case, similar Bernis inequalities were shown in [\[AG04](#page-22-2)] in the parameter range  $p \geq \frac{4}{3}$  $\frac{4}{3}$  and  $\frac{p-1}{2} < n < 2p - 1$ .

<span id="page-1-1"></span>In the following, we present the main results of this article, Theorems [1.1](#page-1-1) and [1.3,](#page-2-0) in detail.

**Theorem 1.1.** Let  $\Omega \subset \mathbb{R}^d$ ,  $d < 3p$ , be a bounded, convex domain with smooth boundary  $\partial \Omega$ . Let  $2 \le p < \frac{19}{3}$ *and*

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}
$$

*Assume*  $u \in W_p^2(\Omega)$  *is positive, satisfies*  $\nabla u \cdot v = 0$  *on*  $\partial \Omega$ *, and* 

$$
\int_{\Omega} u^n |\nabla \Delta u|^p \, dx < \infty.
$$

*Then there exist positive constants*  $c_i = c_i(n, p, d) > 0$ ,  $i = 1, 2, 3$ , such that the estimates

<span id="page-1-2"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \le c_1 \left( \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)
$$
\n(1.3)

<span id="page-2-1"></span>*and*

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, \mathrm{d}x + \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 \, \mathrm{d}x \n+ \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \mathrm{II}(\nabla u, \nabla u) \, \mathrm{d}H^{d-1} \leq c_2 \left( \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p \, \mathrm{d}x + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} \, \mathrm{d}x \right)
$$
\n(1.4)

<span id="page-2-2"></span>*and*

$$
\int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} dx \le c_3 \left( \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)
$$
\n(1.5)

*hold true for all non-negative test functions*  $\varphi \in W^1_{\infty}(\Omega)$  *satisfying*  $\int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx < \infty$ . Here, II denotes the *(positive semi-definite) second fundamental form of*  $\partial Ω$ *.* 

- **Remark 1.2.** (i) For  $p = 2$  and  $\varphi \equiv 1$ , the inequalities in [\(1.3\)](#page-1-2) and [\(1.4\)](#page-2-1) recover exactly the results in [\[Grü01](#page-22-0), Theorem 1.1]. The localized version for  $p = 2$  with a non-negative localization function  $\varphi$  is mentioned in [\[Grü05](#page-22-3)], where a proof is sketched for the range  $2 - \sqrt{\frac{8}{8+d}} < n < 3$ .
	- (ii) For  $d = 1$ , estimates [\(1.3\)](#page-1-2) and [\(1.5\)](#page-2-2) are derived in [\[AG04](#page-22-2), Proposition 4.1] for  $p \geq \frac{4}{3}$  $\frac{4}{3}$  and  $\frac{p-1}{2} < n < 2p-1$ . The restriction  $p \ge 2$  in our case (instead of  $p \ge \frac{4}{3}$ )  $\frac{4}{3}$ , as in *d* = 1) is due to the term  $|\nabla u|^{3p-6}$  in the integral

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, \mathrm{d} x.
$$

In the case  $d = 1$ , this integral simplifies to

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 \, \mathrm{d} x.
$$

Note, however, that we can estimate  $|D^2u\nabla u|^2 \leq |D^2u|^2|\nabla u|^2$ , where  $|D^2u| = \sqrt{\sum_{i,j=1}^d (\partial_{ij}u)^2}$  denotes the Frobenius norm of the Hessian matrix  $D^2u = (\partial_{ij}u)_{i,j=1,\dots,d}$ .

- (iii) The restriction  $p < \frac{19}{3}$  is used only once, in [\(2.8\)](#page-8-0) of the proof of Proposition [2.3,](#page-6-0) to ensure the nonnegativity of the integral  $\mathcal{I}_4$ . This integral does not appear for  $d = 1$ , as in this case  $D^2u = \Delta u$ .
- (iv) For smooth functions, Theorem [1.1](#page-1-1) is valid for any dimension  $d \ge 1$ . The critical dimension  $d < 3p$  in Theorem [1.1](#page-1-1) increases monotonically with *p*. It can be computed from the conditions  $W_p^3(\Omega) \hookrightarrow W_q^2(\Omega)$ for  $1 \le q < \frac{pd}{d-p}$  and  $W_q^2(\Omega) \hookrightarrow C(\overline{\Omega})$  for  $q > \frac{d}{2}$ , as evident from the proof of Theorem [1.1.](#page-1-1)

(v) For 
$$
p \ge 2
$$
, we have

$$
2p - 2 \frac{2 + \sqrt{(3p-2)^2 + d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2 + d(3p-4)}} < 2p-2.
$$

For  $p = 2$ , this reduces to  $2 - \sqrt{\frac{8}{8+d}} < 2$ , which is consistent with [\[Grü01;](#page-22-0) [Grü05](#page-22-3)]. For  $d = 1$ , we have

$$
2p - 2 \frac{2 + \sqrt{(3p - 2)^2 + d(3p - 4)}}{-3p + 6 + \sqrt{(3p - 2)^2 + d(3p - 4)}} = \frac{4p - 2 - 2\sqrt{p(p - 1)}}{3},
$$

which is greater than  $\frac{p-1}{2}$  for all  $p \ge 1$ . This shows that the bound for *n* derived in this article is stricter than the one provided in [\[AG04](#page-22-2)].

<span id="page-2-0"></span>**Theorem 1.3.** *Let*  $\Omega \subset \mathbb{R}^d$ ,  $d < 3p$ , be a bounded, convex domain with smooth boundary  $\partial \Omega$ . *Let*  $2 \le p < \frac{19}{3}$ *and*

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}
$$

*Assume*  $u \in W_p^2(\Omega)$  *is positive, satisfies*  $\nabla u \cdot v = 0$  *on*  $\partial \Omega$ *, and* 

$$
\int_{\Omega} u^n |\nabla \Delta u|^p \, dx < \infty.
$$

*Then, we have*

<span id="page-3-4"></span>
$$
u^{\frac{n+p}{3p}} \in W^1_{3p}(\Omega), \quad u^{\frac{2(n+p)}{3p}} \in W^2_{\frac{3p}{2}}(\Omega), \quad \text{and} \quad u^{\frac{n+p}{p}} \in W^3_p(\Omega).
$$
 (1.6)

*Moreover, there exists a constant*  $c_4 = c_4(n, p, d) > 0$  such that

<span id="page-3-1"></span>
$$
\int_{\Omega} \left| \nabla \left( u^{\frac{n+p}{3p}} \right) \right|^{3p} dx + \int_{\Omega} \left| \Delta \left( u^{\frac{2(n+p)}{3p}} \right) \right|^{3p} dx + \int_{\Omega} \left| \nabla \Delta \left( u^{\frac{n+p}{p}} \right) \right|^{p} dx \leq c_4 \int_{\Omega} u^n |\nabla \Delta u|^p dx. \tag{1.7}
$$

**Remark 1.4.** For the first two terms in [\(1.7\)](#page-3-1), we also obtain localized estimates of the form

$$
\int_{\Omega} \varphi^{3p} \left| \nabla \left( u^{\frac{n+p}{3p}} \right) \right|^{3p} dx + \int_{\Omega} \varphi^{3p} \left| \Delta \left( u^{\frac{2(n+p)}{3p}} \right) \right|^{3p} dx \le c(n,p,d) \left( \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} dx \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)
$$

for all non-negative test functions  $\varphi \in W^1_{\infty}(\Omega)$  satisfying  $\int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx < \infty$ , see Corollary [3.1.](#page-20-0) However, deriving a localized version for the third-order term in [\(1.7\)](#page-3-1) turns out to be significantly more challenging. This issue is discussed in Remark [3.2](#page-21-0) in detail.

We emphasize again that the strategy used to prove inequalities [\(1.3\)](#page-1-2) and [\(1.4\)](#page-2-1) in Theorem [1.1](#page-1-1) is strongly inspired by the approach in [\[Grü01\]](#page-22-0), while incorporating the necessary generalizations. In the Newtonian case  $p = 2$ , these two inequalities alone are sufficient to derive the weighted analogue of Theorem [1.3.](#page-2-0) Inequality [\(1.5\)](#page-2-2) in Theorem [1.1](#page-1-1) can be established similarly to the one-dimensional case in [\[AG04](#page-22-2)]. Nonetheless, the results presented here are novel and play a significant role in studying weak solutions to [\(1.1\)](#page-0-0), as well as in proving qualitative properties such as finite speed of propagation and the occurrence of waiting-time phenomena. Further details on potential applications of Theorems [1.1](#page-1-1) and [1.3](#page-2-0) are given in Section [1.3,](#page-3-0) where we summarize the state of the art for the one-dimensional non-Newtonian case and for the Newtonian case in dimensions  $1 \leq d < 6$ .

<span id="page-3-0"></span>1.3. **Related results.** In the one-dimensional Newtonian case, the corresponding thin-film equation is given by

<span id="page-3-2"></span>
$$
u_t + \left(u^n u_{xxx}\right)_x = 0.\tag{1.8}
$$

This equation was first studied in the seminal paper [\[BF90](#page-22-4)]. The property of finite speed of propagation for [\(1.8\)](#page-3-2) was shown in [\[Ber96a](#page-22-5); [Ber96b](#page-22-6)] for  $0 < n < 2$  and  $2 \le n < 3$ , respectively, with upper bounds on the propagation rate for  $0 < n < 2$  derived in [\[Ber96a](#page-22-5)]. The ideas used in the proofs differ depending on the mobility exponent *n*. For  $0 \lt n \lt 2$ , the proof relies heavily on the so-called  $\alpha$ -entropy estimates, while for  $2 \le n \lt 3$ , it is based on localized versions of the energy inequality  $(1.2)$  in  $d = 1$ . In [\[HS98\]](#page-22-7), the authors showed a finite speed of propagation result in terms of the  $L_1$ -norm. The occurrence of a waiting-time phenomenon was proved in [\[DGG01\]](#page-22-8), even for higher space dimensions.

The one-dimensional non-Newtonian analogue of [\(1.1\)](#page-0-0) is given by

$$
u_t + (u^n | u_{xxx} |^{p-2} u_{xxx})_x = 0,
$$

and Bernis estimates in the style of Theorems [1.1](#page-1-1) and [1.3](#page-2-0) were derived in [\[AG04](#page-22-2)], where the authors used these estimates in particular to guarantee the zero-contact angle condition for the constructed global non-negative solutions. Earlier, estimates of the form [\(1.3\)](#page-1-2) and [\(1.5\)](#page-2-2) were obtained in [\[Ber96c](#page-22-9)], where the parameter range  $p \geq \frac{4}{5}$  $\frac{4}{3}$  and  $n \in (\frac{p-1}{2}, \frac{p+1}{2})$  $\frac{+1}{2} - \frac{1}{3}$  $\frac{1}{3p}$  was considered. Let us note that *α*-entropy estimates are not available in the non-Newtonian setting.

The higher-dimensional Newtonian thin-film equation reads

<span id="page-3-3"></span>
$$
u_t + \operatorname{div}(u^n \nabla \Delta u) = 0,\tag{1.9}
$$

and Bernis estimates for this equation were derived in [\[Grü01](#page-22-0)]. An entropy estimate for [\(1.9\)](#page-3-3) was shown in [\[DGG98\]](#page-22-10), and based on this, further qualitative results for the parameter range  $0 < n < 2$  – including finite speed of propagation, waiting-time phenomena, and the existence of solutions to [\(1.9\)](#page-3-3) emanating from measure-valued initial data – were obtained in [\[Ber+98](#page-22-11)], [\[DG99\]](#page-22-12), and [\[DGG01](#page-22-8)].

To the best of the authors' knowledge, this is the first paper to address Bernis estimates for the higherdimensional non-Newtonian thin-film equation.

1.4. **Notation.** Let  $\Omega \subset \mathbb{R}^d$  be a domain. The standard  $L_q(\Omega)$ -norm for  $1 \leq q \leq \infty$  is denoted by  $\|\cdot\|_{L_q(\Omega)}$ . The Sobolev space  $W_q^m(\Omega)$ , where  $m \in \mathbb{N}$ , consists of functions whose generalized derivatives up to order  $m$  belong to  $L_q(\Omega)$ . The corresponding norm is denoted by  $\|\cdot\|_{W_q^m(\Omega)}$ .

*<del><i>v*</del> *i* is the *v*<sup>*m*</sup> in *v*<sub>*a*</sub> (ε*x*) *i* is *m<sub><i>na*</sub></sup> (ε*x*) *i* is denoted by  $∇$ *u*, and the Hessian *For* a sufficiently differentiable function *u* :  $Ω → ℝ$ , the gradient of *u* is denoted by  $∇$ *u*, and t matrix of *u* is denoted by  $D^2u$ . The Laplacian  $\Delta u$  is defined as the trace of the Hessian matrix  $D^2u$ .

The exterior unit normal vector to the boundary *∂*Ω is denoted by *ν*. Depending on the context, |⋅| represents the absolute value for scalars in ℝ, the Euclidean norm for vectors in ℝ<sup>d</sup>, or the Frobenius norm for matrices in  $\mathbb{R}^{d \times d}$ .

This article is organized into three sections: in Section [2,](#page-4-0) we prove Theorem [1.1,](#page-1-1) and in Section [3,](#page-18-0) we prove Theorem [1.3.](#page-2-0)

### 2. PROOF OF THEOREM [1.1](#page-1-1)

<span id="page-4-0"></span>This main section of the paper is devoted to the proof of Theorem [1.1,](#page-1-1) and we follow, more or less, the approach of Grün [\[Grü01\]](#page-22-0), where Bernis estimates for the higher-dimensional thin-film equation were established in the Newtonian setting.

The proof proceeds in three main steps. First, we derive a technical auxiliary integral identity, presented in Lemma [2.2.](#page-5-0) Next, we establish the desired estimates  $(1.3)$  and  $(1.4)$  for two specific ranges of *n*. For the range 2*p*−2 < *n* < 2*p*−1, the estimates are proved in Proposition [2.3.](#page-6-0) For the range

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}
$$

the estimates are obtained in Proposition [2.5.](#page-12-0) In both cases, these results are shown for smooth functions. The inequality [\(1.5\)](#page-2-2) is proved directly in Proposition [2.7](#page-16-0) for the full range

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}
$$

This result is also established for smooth functions. Finally, an approximation argument is used to extend these results to  $u \in W_p^2(\Omega)$ . Since

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}} < 2p-2 \quad \text{and} \quad 2p-2 \frac{6p-6+\sqrt{(3p-2)^2+d(3p-4)}}{9p-10+\sqrt{(3p-2)^2+d(3p-4)}} < 2p-1,
$$

we conclude Theorem [1.1](#page-1-1) for the entire range

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}
$$

Lemma [2.2,](#page-5-0) as mentioned earlier, provides the auxiliary integral identity. To prove this lemma, we use the following relation between the normal derivative of  $|\nabla u|^2$  and the second fundamental form of the boundary of Ω.

<span id="page-4-1"></span>**Lemma 2.1.** *Let*  $\Omega \subset \mathbb{R}^d$  *be a bounded smooth domain, and let* II *denote the second fundamental form of*  $\partial \Omega$ *. For any function*  $u \in C^2(\overline{\Omega})$  *with*  $\nabla u \cdot v = 0$  *on*  $\partial \Omega$ *, we have*  $\nabla |\nabla u|^2 \cdot v = -2\Pi(\nabla u, \nabla u)$ *. In particular, if*  $\Omega$  *is*  $convex, then \nabla |\nabla u|^2 \cdot v \leq 0 \text{ on } \partial \Omega.$ 

This control of the normal derivative of  $|\nabla u|^2$  on  $\partial \Omega$  is widely used in the literature; see, for example, [\[CH78](#page-22-13)], [\[SZ98](#page-22-14)], and the references cited in [\[Grü01\]](#page-22-0). For details, we refer to these works.

<span id="page-5-0"></span>**Lemma 2.2.** Let  $\Omega \subset \mathbb{R}^d$  be a bounded smooth domain, and let  $p \geq 2$ . For any positive function  $u \in C^\infty(\overline{\Omega})$ *satisfying*  $\nabla u \cdot v = 0$  *on*  $\partial \Omega$ *, and for any*  $\varphi \in W^1_{\infty}(\Omega)$ *, the following identity holds:* 

<span id="page-5-1"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 dx
$$
\n
$$
= -\frac{3p}{3p-4} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot D^2 u \nabla u dx
$$
\n
$$
- \frac{n-2p+2}{3p-4} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} \nabla u \cdot D^2 u \nabla u dx
$$
\n
$$
- \frac{1}{3p-4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 dx - \frac{1}{3p-4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla u \cdot \nabla \Delta u dx
$$
\n
$$
- \frac{1}{3p-4} \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) d\mathcal{H}^{d-1}.
$$
\n(2.1)

*Proof.* We first use the identities

$$
D^2u\nabla u = \frac{1}{2}\nabla|\nabla u|^2 \quad \text{and} \quad \nabla|\nabla u|^2 \cdot \nabla|\nabla u|^2 = \text{div}\left(|\nabla u|^2 \nabla|\nabla u|^2\right) - |\nabla u|^2 \Delta|\nabla u|^2
$$

to obtain

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, dx
$$
\n
$$
= \frac{1}{4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} \nabla |\nabla u|^2 \cdot \nabla |\nabla u|^2 \, dx
$$
\n
$$
= \frac{1}{4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} \operatorname{div} (|\nabla u|^2 \nabla |\nabla u|^2) \, dx - \frac{1}{4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Delta |\nabla u|^2 \, dx.
$$

Applying integration by parts on the first integral on the right-hand side and using that  $\nabla |\nabla u|^2 \cdot v = -2\Pi(\nabla u, \nabla u)$ on  $\partial Ω$ , see Lemma [2.1,](#page-4-1) then yields

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, dx
$$
\n
$$
= -\frac{3p}{4} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot \nabla |\nabla u|^2 \, dx - \frac{n-2p+2}{4} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} \nabla u \cdot \nabla |\nabla u|^2 \, dx
$$
\n
$$
- \frac{3p-6}{4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} D^2 u \nabla u \cdot \nabla |\nabla u|^2 \, dx - \frac{1}{4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Delta |\nabla u|^2 \, dx
$$
\n
$$
- \frac{1}{2} \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) \, d\mathcal{H}^{d-1}.
$$

With the identities  $\nabla |\nabla u|^2 = 2D^2 u \nabla u$  and  $\Delta |\nabla u|^2 = 2(|D^2 u|^2 + \nabla u \cdot \nabla \Delta u)$ , this can be rewritten as

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 dx
$$
\n
$$
= -\frac{3p}{2} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot D^2 u \nabla u dx - \frac{n-2p+2}{2} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} D^2 u \nabla u \cdot \nabla u dx
$$
\n
$$
- \frac{3p-6}{2} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 dx - \frac{1}{2} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 dx
$$
\n
$$
- \frac{1}{2} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla u \cdot \nabla \Delta u dx - \frac{1}{2} \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi (\nabla u, \nabla u) d\mathcal{H}^{d-1}.
$$

Since the third integrand on the right-hand side equals the integrand on the left-hand side and since  $p \ge 2 > \frac{4}{3}$ 3 ensures that  $\frac{1}{3p-4} > 0$ , the proof is complete.

We remark that in Lemma [2.2,](#page-5-0) we need  $p \geq 2 > \frac{4}{3}$  $\frac{4}{3}$  in the final step of the proof to ensure that  $\frac{2}{3p-4} > 0$ . In this case, we have  $\frac{n-2p+2}{3p-4} > 0$  if and only if  $n > 2p-2$ . Next, we establish Theorem [1.1](#page-1-1) for fixed  $2 \le p < \frac{19}{3}$  in the case  $2p-2 < n < 2p-1$ . Note that this corresponds to the bounds  $2 < n < 3$  in the Newtonian case  $p = 2$ , as detailed in [\[Grü01,](#page-22-0) Lemma 2.3].

<span id="page-6-0"></span>**Proposition 2.3.** *Let*  $\Omega \subset \mathbb{R}^d$  *be a bounded, convex and smooth domain, and let*  $2 \leq p < \frac{19}{3}$  *be fixed. Assume fhat*  $u \in C^{\infty}(\bar{\Omega})$  *is positive and satisfies*  $\nabla u \cdot v = 0$  *on*  $\partial \Omega$ *, and let*  $\varphi \in W_{\infty}^{1}(\Omega)$  *be non-negative. If* 

$$
2p-2 < n < 2p-1
$$

*then there exist constants*  $C_i = C_i(n, p) > 0$ ,  $i = 1, 2$ , such that the following estimates hold:

<span id="page-6-1"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \le C_1 \left[ \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right]
$$
\n(2.2)

<span id="page-6-2"></span>*and*

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, dx + \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 \, dx \n+ \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 \, dx + \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) \, d\mathcal{H}^{d-1} \n\leq C_2 \left[ \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p \, dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} \, dx \right].
$$
\n(2.3)

Note that Proposition [2.3](#page-6-0) works for any dimension  $d \geq 1$ .

**Remark 2.4.** The constants  $C_1$  and  $C_2$  in Proposition [2.3](#page-6-0) are given by

$$
C_1 = \max \left\{ \frac{3p}{2} \left| \frac{2(3p-1)}{(n-2p+1)(n-2p+2)} \right|^p, \right\}
$$
  

$$
p^{\frac{3p}{2}} \left( \frac{9p^2}{2(n-2p+1)^2} + \frac{9p^2(3p-2)^2}{\epsilon_2(n-2p+1)^2(n-2p+2)^2} + \frac{9p^2}{\epsilon_3(n-2p+1)^2(n-2p+2)^2} \right)^{\frac{3p}{2}}
$$

and

$$
C_2 = \frac{5}{3p} + C_1 \left[ \frac{p-1}{p} \left( \frac{3p-1}{C_{\min}} \right)^{\frac{p}{p-1}} + \frac{3p-2}{3p} \left( \frac{C_{\delta_0} + C_{\delta_1} + C_{\delta_2}}{C_{\min}} \right)^{\frac{3p}{3p-2}} + \frac{1}{C_{\min}} \right],
$$

where

$$
C_{\delta_0} = \frac{9p^2(n-2p+2)^2}{4} > 0,
$$

and  $C_{\min}$ ,  $C_{\delta_1}$ , and  $C_{\delta_2}$  are positive constants depending only on the fixed parameter *p*, but not on *n*. Hence, both  $C_1$  and  $C_2$  blow up as *n* goes to 2*p*−2. Moreover, we need *n* > 2*p*−2 in the proof to guarantee that the involved integrals have the 'right' sign. However, it turns out that the lower bound  $2p-2$  can be slightly improved (see Proposition [2.5](#page-12-0) below).

*Proof.* We first prove [\(2.2\)](#page-6-1). To this end, we use the identity  $\nabla u \cdot \nabla u = \text{div}(u \nabla u) - u \Delta u$  and integration by parts, together with the boundary condition for  $u$ , to rewrite the left-hand side of  $(2.2)$  as

$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx = \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p-2} \nabla u \cdot \nabla u dx
$$
\n
$$
= \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p-2} \operatorname{div}(u \nabla u) dx - \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-2} \Delta u dx
$$
\n
$$
= -3p \int_{\Omega} \varphi^{3p-1} u^{n-2p+1} |\nabla u|^{3p-2} \nabla \varphi \cdot \nabla u dx - (n-2p) \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx
$$
\n
$$
- (3p-2) \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} D^2 u \nabla u \cdot \nabla u dx - \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-2} \Delta u dx.
$$

Since the second integrand on the right-hand side equals the integrand on the left-hand side and since  $n < 2p - 1$ by assumption, we find that

<span id="page-7-0"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx = -\frac{3p}{n-2p+1} \int_{\Omega} \varphi^{3p-1} u^{n-2p+1} |\nabla u|^{3p-2} \nabla \varphi \cdot \nabla u dx \n- \frac{3p-2}{n-2p+1} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} D^2 u \nabla u \cdot \nabla u dx \n- \frac{1}{n-2p+1} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-2} \Delta u dx.
$$
\n(2.4)

In the last term of [\(2.4\)](#page-7-0) we use again the identity  $\nabla u \cdot \nabla u = \text{div}(u\nabla u) - u\Delta u$  and integrate by parts to obtain

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-2} \Delta u \, dx
$$
\n
$$
= \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} \operatorname{div}(u \nabla u) \Delta u \, dx - \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^{2} \, dx
$$
\n
$$
= -3p \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \Delta u \nabla \varphi \cdot \nabla u \, dx - \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^{2} \, dx
$$
\n
$$
-(n-2p+1) \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-2} \Delta u \, dx - (3p-4) \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} \Delta u D^{2} u \nabla u \cdot \nabla u \, dx
$$
\n
$$
- \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \Delta u \cdot \nabla u \, dx.
$$

Noting that the integrand on the left-hand side equals the third integrand on the right-hand side and recalling that  $n > 2p - 2$  by assumption, we deduce

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-2} \Delta u \, dx
$$
\n
$$
= -\frac{3p}{n-2p+2} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \Delta u \nabla \varphi \cdot \nabla u \, dx - \frac{1}{n-2p+2} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^{2} \, dx
$$
\n
$$
- \frac{3p-4}{n-2p+2} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} \Delta u D^{2} u \nabla u \cdot \nabla u \, dx
$$
\n
$$
- \frac{1}{n-2p+2} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \Delta u \cdot \nabla u \, dx.
$$

Inserting this into [\(2.4\)](#page-7-0), we obtain

<span id="page-7-1"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx = -\frac{3p}{n-2p+1} \int_{\Omega} \varphi^{3p-1} u^{n-2p+1} |\nabla u|^{3p-2} \nabla \varphi \cdot \nabla u dx \n- \frac{3p-2}{n-2p+1} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} D^2 u \nabla u \cdot \nabla u dx \n+ \frac{3p}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \Delta u \nabla \varphi \cdot \nabla u dx \n+ \frac{1}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 dx \n+ \frac{3p-4}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} \Delta u D^2 u \nabla u \cdot \nabla u dx \n+ \frac{1}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \Delta u \cdot \nabla u dx.
$$
\n(2.5)

Next, we use Lemma [2.2](#page-5-0) to rewrite the second term on the right-hand side of [\(2.5\)](#page-7-1) as follows

<span id="page-8-1"></span>
$$
-\frac{3p-2}{n-2p+1} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} D^2 u \nabla u \cdot \nabla u \, dx
$$
  
= 
$$
\frac{(3p-2)(3p-4)}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, dx
$$
  
+ 
$$
\frac{3p(3p-2)}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot D^2 u \nabla u \, dx
$$
  
+ 
$$
\frac{3p-2}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 \, dx
$$
  
+ 
$$
\frac{3p-2}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla u \cdot \nabla \Delta u \, dx
$$
  
+ 
$$
\frac{3p-2}{(n-2p+1)(n-2p+2)} \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) \, d\mathcal{H}^{d-1}.
$$
 (2.6)

Applying now the Cauchy–Schwarz inequality and Young's inequality with weight  $\varepsilon_0 = \frac{1}{15}$  on the fifth term on the right-hand side of [\(2.5\)](#page-7-1) we obtain

<span id="page-8-2"></span>
$$
-\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} \Delta u D^2 u \nabla u \cdot \nabla u \, dx
$$
  
\n
$$
\leq \left| \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} \Delta u D^2 u \nabla u \cdot \nabla u \, dx \right|
$$
  
\n
$$
\leq \frac{1}{15} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 \, dx + \frac{15}{4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, dx.
$$
\n(2.7)

Inserting  $(2.6)$  back into  $(2.5)$  and using  $(2.7)$  yields

<span id="page-8-0"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx
$$
\n
$$
\leq \frac{(3p-4)(12p-23)}{4(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 dx
$$
\n
$$
+ \frac{3p-2}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 dx
$$
\n
$$
+ \frac{3p-1}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla u \cdot \nabla \Delta u dx
$$
\n
$$
+ \frac{19-3p}{15(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 dx
$$
\n
$$
+ \frac{3p-2}{(n-2p+1)(n-2p+2)} \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) d\mathcal{H}^{d-1}
$$
\n
$$
- \frac{3p}{n-2p+1} \int_{\Omega} \varphi^{3p-1} u^{n-2p+1} |\nabla u|^{3p-2} \nabla \varphi \cdot \nabla u dx
$$
\n
$$
+ \frac{3p(3p-2)}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot D^2 u \nabla u dx
$$
\n
$$
+ \frac{3p}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot \nabla u dx
$$
\n
$$
=: \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3 + \mathcal{I}_4 + \mathcal{I}_5 + \mathcal{I}_6
$$

Due to the choices of  $p \ge 2$  and  $2p - 2 < n < 2p - 1$  the pre factors of  $\mathcal{I}_1, \mathcal{I}_2$  and  $\mathcal{I}_5$  are negative, while the integrals are non-negative. The non-negativity of  $II(\nabla u, \nabla u)$  follows from the convexity of the domain, see Lemma [2.1.](#page-4-1) Moreover, the choices of  $p < \frac{19}{3}$  and  $2p - 2 < n < 2p - 1$  yield  $\mathcal{I}_4 \le 0$ . We directly use  $\mathcal{I}_2, \mathcal{I}_5 \le 0$ in [\(2.8\)](#page-8-0). To control the last three integrals  $\tilde{\mathcal{I}}_6$ ,  $\mathcal{I}_7$  and  $\mathcal{I}_8$ , we use the Cauchy–Schwarz inequality and Young's inequality such that  $\mathcal{I}_6$  can be partly absorbed by the integral on the left-hand side of [\(2.8\)](#page-8-0) and the integrals  $\mathcal{I}_7$  and  $\mathcal{I}_8$  are absorbed by  $\mathcal{I}_1$ , respectively  $\mathcal{I}_4$ . Indeed, applying the Cauchy-Schwarz inequality and Young's inequality with weight  $\varepsilon_1 = \frac{1}{2}$  $\frac{1}{2}$  to  $\mathcal{I}_6$ , we have

<span id="page-9-5"></span>
$$
-\frac{3p}{n-2p+1} \int_{\Omega} \varphi^{3p-1} u^{n-2p+1} |\nabla u|^{3p-2} \nabla \varphi \cdot \nabla u \, dx
$$
  

$$
\leq \frac{1}{2} \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} \, dx + C_{\varepsilon_1} \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} \, dx
$$
 (2.9)

with  $C_{\epsilon_1} = C(\epsilon_1, n, p) := \frac{9p^2}{2(n-2p+1)^2} > 0$ . Inserting this inequality in [\(2.8\)](#page-8-0), we obtain

<span id="page-9-2"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \le 2(\mathcal{I}_1 + \mathcal{I}_3 + \mathcal{I}_4 + \mathcal{I}_7 + \mathcal{I}_8) + 2C_{\varepsilon_1} \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} dx.
$$
 (2.10)

We can further estimate  $\mathcal{I}_7$  by

<span id="page-9-0"></span>
$$
\left| \frac{6p(3p-2)}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot D^2 u \nabla u \, dx \right|
$$
\n
$$
\leq \varepsilon_2 \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, dx + C_{\varepsilon_2} \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} \, dx,
$$
\n(2.11)

as well as  $\mathcal{I}_8$  by

<span id="page-9-1"></span>
$$
\left| \frac{6p}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \Delta u \nabla \varphi \cdot \nabla u \, dx \right|
$$
  
\n
$$
\leq \varepsilon_3 \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 \, dx + C_{\varepsilon_3} \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} \, dx,
$$
\n(2.12)

where  $\varepsilon_2, \varepsilon_3 > 0$  can be chosen arbitrarily and where  $C_{\varepsilon_2}, C_{\varepsilon_3} > 0$  are given by

$$
C_{\varepsilon_2} = C(\varepsilon_2, n, p) := \frac{9p^2(3p - 2)^2}{\varepsilon_2(n - 2p + 1)^2(n - 2p + 2)^2}
$$

and

$$
C_{\varepsilon_3} = C(\varepsilon_3, n, p) := \frac{9p^2}{\varepsilon_3(n-2p+1)^2(n-2p+2)^2}.
$$

We choose  $\varepsilon_2 = \varepsilon_2(n, p)$  and  $\varepsilon_3 = \varepsilon_3(n, p)$  small enough so that

<span id="page-9-3"></span>
$$
0 < \varepsilon_2 < \left| \frac{(3p-4)(12p-23)}{2(n-2p+1)(n-2p+2)} \right| \quad \text{and} \quad 0 < \varepsilon_3 < \left| \frac{2(19-3p)}{15(n-2p+1)(n-2p+2)} \right|.
$$

Using  $(2.11)$  and  $(2.12)$  in  $(2.10)$ , we can absorb the terms in  $(2.10)$  to get the estimate

$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \le \frac{2(3p-1)}{(n-2p+1)(n-2p+2)} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla u \cdot \nabla \Delta u dx + \left(2C_{\varepsilon_1} + C_{\varepsilon_2} + C_{\varepsilon_3}\right) \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} dx.
$$
\n(2.13)

Applying Hölder's inequality to both integrals on the right-hand side of [\(2.13\)](#page-9-3), we obtain

<span id="page-9-4"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx
$$
\n
$$
\leq \left| \frac{2(3p-1)}{(n-2p+1)(n-2p+2)} \right| \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{p-1}{p}} \left( \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} dx \right)^{\frac{1}{p}}
$$
\n
$$
+ (2C_{\epsilon_1} + C_{\epsilon_2} + C_{\epsilon_3}) \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{3p-2}{3p}} \left( \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)^{\frac{2}{3p}}.
$$
\n(2.14)

We then use Young's inequality on the first line on the right-hand side of  $(2.14)$  to deduce

$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx
$$
\n
$$
\leq \frac{p-1}{p} \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx + \frac{1}{p} \left| \frac{2(3p-1)}{(n-2p+1)(n-2p+2)} \right|^p \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx
$$
\n
$$
+ (2C_{\epsilon_1} + C_{\epsilon_2} + C_{\epsilon_3}) \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{3p-2}{3p}} \left( \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)^{\frac{2}{3p}},
$$

which immediately gives

$$
\begin{aligned} \int_{\Omega}\varphi^{3p}u^{n-2p}|\nabla u|^{3p}\,\mathrm{d} x &\leq \left|\frac{2(3p-1)}{(n-2p+1)(n-2p+2)}\right|^p\int_{\Omega}\varphi^{3p}u^n|\nabla\Delta u|^p\,\mathrm{d} x\\&\qquad\qquad\qquad\left.+p\big(2C_{\varepsilon_1}+C_{\varepsilon_2}+C_{\varepsilon_3}\big)\left(\int_{\Omega}\varphi^{3p}u^{n-2p}|\nabla u|^{3p}\,\mathrm{d} x\right)^{\frac{3p-2}{3p}}\left(\int_{\Omega}|\nabla\varphi|^{3p}u^{n+p}\,\mathrm{d} x\right)^{\frac{2}{3p}}. \end{aligned}
$$

Applying again Young's inequality, but now to the second line on the right-hand side of the above inequality, leads to

$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \le \frac{3p}{2} \left| \frac{2(3p-1)}{(n-2p+1)(n-2p+2)} \right|^p \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx + p^{\frac{3p}{2}} \left( 2C_{\varepsilon_1} + C_{\varepsilon_2} + C_{\varepsilon_3} \right)^{\frac{3p}{2}} \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx,
$$

and from this we conclude estimate [\(2.2\)](#page-6-1) with the constant  $C_1 > 0$  given by

$$
C_1 = \max \left\{ \frac{3p}{2} \left| \frac{2(3p-1)}{(n-2p+1)(n-2p+2)} \right|^p, \right\}
$$
  

$$
p^{\frac{3p}{2}} \left( \frac{9p^2}{(n-2p+1)^2} + \frac{9p^2(3p-2)^2}{\epsilon_2(n-2p+1)^2(n-2p+2)^2} + \frac{9p^2}{\epsilon_3(n-2p+1)^2(n-2p+2)^2} \right)^{\frac{3p}{2}} \right\}.
$$

It remains to prove [\(2.3\)](#page-6-2). Noting that the left-hand side in [\(2.8\)](#page-8-0) is non-negative and multiplying (2.8) by  $(n 2p+1(n-2p+2) < 0$  gives

<span id="page-10-1"></span><span id="page-10-0"></span>
$$
(n-2p+1)(n-2p+2)\sum_{i=1}^{8} \mathcal{I}_i \le 0
$$

and by rearranging this inequality, we see that

$$
\frac{(3p-4)(12p-23)}{4} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 dx + (3p-2) \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 dx \n+ \frac{19-3p}{15} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 dx + (3p-2) \int_{\partial\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) dH^{d-1} \n\leq -(3p-1) \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla u \cdot \nabla \Delta u dx \n+ 3p(n-2p+2) \int_{\Omega} \varphi^{3p-1} u^{n-2p+1} |\nabla u|^{3p-2} \nabla \varphi \cdot \nabla u dx \n- 3p(3p-2) \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot D^2 u \nabla u dx \n- 3p \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \Delta u \nabla \varphi \cdot \nabla u dx
$$
\n(2.15)\n  
\n=:  $J_1 + J_2 + J_3 + J_4$ .

Now, we estimate the terms  $\mathcal{J}_2$ ,  $\mathcal{J}_3$ , and  $\mathcal{J}_4$  by similar arguments as those used for [\(2.9\)](#page-9-5), [\(2.11\)](#page-9-0) and [\(2.12\)](#page-9-1). Indeed, to estimate  $\mathcal{J}_2$ , we use Young's inequality with  $\delta_0 = 1$  and [\(2.2\)](#page-6-1). We obtain

$$
\mathcal{J}_2 \leq C_{\delta_0} \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} \, \mathrm{d}x + C_1 \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p \, \mathrm{d}x + C_1 \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} \, \mathrm{d}x \tag{2.16}
$$

with  $C_{\delta_0} = C(\delta_0, n, p) := \frac{9p^2(n-2p+2)^2}{4}$  $\frac{(2p+2)^2}{4} > 0$ . For  $\mathcal{J}_3$  and  $\mathcal{J}_4$ , we obtain the estimates

$$
\mathcal{J}_3 \le \delta_1 \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, \mathrm{d}x + C_{\delta_1} \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} \, \mathrm{d}x \tag{2.17}
$$

<span id="page-11-0"></span>and

$$
\mathcal{J}_4 \le \delta_2 \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 \, \mathrm{d}x + C_{\delta_2} \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} \, \mathrm{d}x,\tag{2.18}
$$

where  $\delta_1$ ,  $\delta_2$  > 0 can be chosen arbitrarily and with

$$
C_{\delta_1} = C(\delta_1, p) := \frac{9p^2(3p - 2)^2}{4\delta_1} > 0 \quad \text{and} \quad C_{\delta_2} = C(\delta_2, p) := \frac{9p^2}{4\delta_2} > 0.
$$

Inserting  $(2.16)$ – $(2.18)$  into  $(2.15)$  and choosing

$$
0 < \delta_1 < \frac{(3p-4)(12p-23)}{8} \quad \text{and} \quad 0 < \delta_2 < \frac{19-3p}{30}
$$

yields

$$
\frac{(3p-4)(12p-23)}{8} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 dx + (3p-2) \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 dx \n+ \frac{19-3p}{30} \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 dx + (3p-2) \int_{\partial\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) dH^{d-1} \n\leq -(3p-1) \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla u \cdot \nabla \Delta u dx \n+ (C_{\delta_0} + C_{\delta_1} + C_{\delta_2}) \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} dx \n+ C_1 \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx + C_1 \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx.
$$
\n(2.19)

Setting

$$
C_{\min} := \min\left\{\frac{(3p-4)(12p-23)}{8}, \frac{19-3p}{30}, 3p-2\right\} > 0
$$

and, applying Hölder's and Young's inequalities, further yields

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 dx + \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 dx \n+ \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |\Delta u|^2 dx + \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) dH^{d-1} \n\leq \frac{(3p-1)}{C_{\min}} \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{p-1}{p}} \left( \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^p dx \right)^{\frac{1}{p}} \n+ \frac{(C_{\delta_0} + C_{\delta_1} + C_{\delta_2})}{C_{\min}} \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{3p-2}{3p}} \left( \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)^{\frac{2}{3p}} \n+ \frac{C_1}{C_{\min}} \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^p dx + \frac{C_1}{C_{\min}} \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \n\leq \frac{p-1}{p} \left( \frac{3p-1}{C_{\min}} \right)^{\frac{p}{p-1}} \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx + \frac{1}{p} \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^p dx \n+ \frac{3p-2}{3p} \left( \frac{C_{\delta_0} + C_{\delta_1} + C_{\delta_2}}{C_{\min}} \right)^{\frac{3p}{3p-2}} \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx + \frac{2}{3p} \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \n+ \frac{C_1}{C_{\min}}
$$

Using [\(2.2\)](#page-6-1), we finally obtain the estimate [\(2.3\)](#page-6-2) with the constant  $C_2 > 0$  given by

$$
C_2 := \frac{5}{3p} + C_1 \left[ \frac{p-1}{p} \left( \frac{3p-1}{C_{\min}} \right)^{\frac{p}{p-1}} + \frac{3p-2}{3p} \left( \frac{C_{\delta_0} + C_{\delta_1} + C_{\delta_2}}{C_{\min}} \right)^{\frac{3p}{3p-2}} + \frac{1}{C_{\min}} \right].
$$

<span id="page-12-0"></span>With the following result we extend the lower bound for *n*, obtained in Proposition [2.3.](#page-6-0)

**Proposition 2.5.** *Let*  $\Omega \subset \mathbb{R}^d$  *be a bounded, convex and smooth domain, and let*  $2 \leq p < \frac{19}{3}$  *be fixed. Assume fhat*  $u \in C^{\infty}(\bar{\Omega})$  *is positive and satisfies*  $\nabla u \cdot v = 0$  *on*  $\partial \Omega$ *, and let*  $\varphi \in W_{\infty}^{1}(\Omega)$  *be non-negative. If* 

$$
2p-2\frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}
$$

*then there exist positive constants*  $C_i = C_i(n, p, d) > 0$ ,  $i = 3, 4$ , *such that the following estimates hold:* 

<span id="page-12-1"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \le C_3 \left[ \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right]
$$
\n(2.20)

*and*

$$
\int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 \, \mathrm{d}x + \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 \, \mathrm{d}x \n+ \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \mathrm{II}(\nabla u, \nabla u) \, \mathrm{d}H^{d-1} \leq C_4 \left[ \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p \, \mathrm{d}x + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} \, \mathrm{d}x \right].
$$
\n(2.21)

**Remark 2.6.** (i) Observe that Proposition [2.5](#page-12-0) is valid for any dimension  $d \ge 1$ . Moreover, as

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}<2p-2
$$

for all  $p \ge 2$ , Proposition [2.5](#page-12-0) extends the statement of Proposition [2.3](#page-6-0) to values of *n* that are slightly smaller than  $2p-2$ . However, in Proposition [2.5](#page-12-0) the constants  $C_3, C_4 > 0$  depend not only on *n* and *p*, but also on the dimension *d*.

(ii) Note that for  $p = 2$ , the upper bound

$$
2p - 2 \frac{6p - 6 + \sqrt{(3p - 2)^2 + d(3p - 4)}}{9p - 10 + \sqrt{(3p - 2)^2 + d(3p - 4)}}
$$
  
equals  $2 + \frac{2}{4 + \sqrt{\frac{8 + d}{8}}}$ , i.e. it coincides with the one in [Gri01].

*Proof.* We start by introducing the notation

$$
\mathcal{K}_1 := \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx,
$$
  
\n
$$
\mathcal{K}_2 := \int_{\Omega} \varphi^{3p-2} |\nabla \varphi|^2 u^{n-2p+2} |\nabla u|^{3p-2} dx,
$$
  
\n
$$
\mathcal{K}_3 := \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-6} |D^2 u \nabla u|^2 dx,
$$
  
\n
$$
\mathcal{K}_4 := \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} |D^2 u|^2 dx,
$$
  
\n
$$
\mathcal{B} := \int_{\partial \Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \Pi(\nabla u, \nabla u) d\mathcal{H}^{d-1}.
$$

Using this, we can rewrite [\(2.1\)](#page-5-1) from Lemma [2.2](#page-5-0) as

<span id="page-13-0"></span>
$$
(3p-4)\mathcal{K}_3 + \mathcal{K}_4 + \mathcal{B} = -3p \int_{\Omega} \varphi^{3p-1} u^{n-2p+2} |\nabla u|^{3p-4} \nabla \varphi \cdot D^2 u \nabla u \, dx
$$

$$
- (n-2p+2) \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} \nabla u \cdot D^2 u \nabla u \, dx
$$

$$
- \int_{\Omega} \varphi^{3p} u^{n-2p+2} |\nabla u|^{3p-4} \nabla u \cdot \nabla \Delta u \, dx.
$$
 (2.22)

We want to estimate the right-hand side of [\(2.22\)](#page-13-0) in terms of  $\mathcal{K}_3$  and  $\mathcal{K}_4$ . For that purpose, we apply the Cauchy–Schwarz and Hölder inequalities, obtaining that

<span id="page-13-3"></span>
$$
(3p-4)\mathcal{K}_3 + \mathcal{K}_4 + \mathcal{B} \le 3p\sqrt{\mathcal{K}_2}\sqrt{\mathcal{K}_3} + |n-2p+2|\sqrt{\mathcal{K}_1}\sqrt{\mathcal{K}_3} + \mathcal{K}_1^{\frac{p-1}{p}} \left( \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p \, dx \right)^{\frac{1}{p}}.
$$
\n
$$
(2.23)
$$

**Step 1. Estimates for**  $\mathcal{K}_1$  **and**  $\mathcal{K}_2$  **in terms of**  $\mathcal{K}_3$  **and**  $\mathcal{K}_4$ **. First, we observe that, by Hölder's inequality,**  $\mathcal{K}_2$ can be estimated in terms of  $\mathcal{K}_1$  by

<span id="page-13-1"></span>
$$
\mathcal{K}_2 \leq \mathcal{K}_1^{\frac{3p-2}{3p}} \left( \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} \, \mathrm{d}x \right)^{\frac{2}{3p}}.
$$
\n
$$
(2.24)
$$

For the integral  $\mathcal{K}_1$ , we obtain from equation [\(2.4\)](#page-7-0), by using the Cauchy–Schwarz inequality and the estimate

$$
\frac{1}{d}|\Delta u|^2 \le \sum_{i=1}^d (\partial_{ii} u)^2 \le \sum_{i,j=1}^d (\partial_{ij} u)^2 = |D^2 u|^2 \quad \text{in } \Omega,
$$

that

$$
\mathcal{K}_1 = -\frac{3p}{n-2p+1} \int_{\Omega} \varphi^{3p-1} u^{n-2p+1} |\nabla u|^{3p-2} \nabla \varphi \cdot \nabla u \, dx \n- \frac{3p-2}{n-2p+1} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-4} D^2 u \nabla u \cdot \nabla u \, dx \n- \frac{1}{n-2p+1} \int_{\Omega} \varphi^{3p} u^{n-2p+1} |\nabla u|^{3p-2} \Delta u \, dx \n\leq \frac{3p}{|n-2p+1|} \sqrt{\mathcal{K}_1} \sqrt{\mathcal{K}_2} + \frac{3p-2}{|n-2p+1|} \sqrt{\mathcal{K}_1} \sqrt{\mathcal{K}_3} + \frac{\sqrt{d}}{|n-2p+1|} \sqrt{\mathcal{K}_1} \sqrt{\mathcal{K}_4},
$$

and dividing both sides by  $\sqrt{\mathcal{K}_1}$  gives

<span id="page-13-4"></span>
$$
\sqrt{\mathcal{K}_1} \le \frac{1}{|n-2p+1|} \left( 3p\sqrt{\mathcal{K}_2} + (3p-2)\sqrt{\mathcal{K}_3} + \sqrt{d}\sqrt{\mathcal{K}_4} \right).
$$
\n(2.25)

Hence, by using Young's inequality, we see that

<span id="page-13-2"></span>
$$
\mathcal{K}_1 \le \frac{1}{|n-2p+1|^2} \left( 3p\sqrt{\mathcal{K}_2} + (3p-2)\sqrt{\mathcal{K}_3} + \sqrt{d}\sqrt{\mathcal{K}_4} \right)^2
$$
  

$$
\le \frac{18p^2}{|n-2p+1|^2} \mathcal{K}_2 + \frac{2}{|n-2p+1|^2} \left( (3p-2)\sqrt{\mathcal{K}_3} + \sqrt{d}\sqrt{\mathcal{K}_4} \right)^2.
$$
 (2.26)

Now, inserting the inequality [\(2.24\)](#page-13-1) for  $\mathcal{K}_2$  into [\(2.26\)](#page-13-2) and applying Young's inequality with weight  $\frac{1}{2}$ , we deduce the estimate

$$
\mathcal{K}_1 \le \frac{1}{2}\mathcal{K}_1 + \frac{2}{3p} \left(\frac{3p}{6p-4}\right)^{\frac{2-3p}{2}} \left(\frac{18p^2}{|n-2p+1|^2}\right)^{\frac{3p}{2}} \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx + \frac{2}{|n-2p+1|^2} \left((3p-2)\sqrt{\mathcal{K}_3} + \sqrt{d}\sqrt{\mathcal{K}_4}\right)^2.
$$

This directly implies the following estimate for  $\mathcal{K}_1$ :

<span id="page-13-5"></span>
$$
\mathcal{K}_1 \le \frac{4}{|n-2p+1|^2} \left( (3p-2)\sqrt{\mathcal{K}_3} + \sqrt{d}\sqrt{\mathcal{K}_4} \right)^2 + C \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx,
$$
\n(2.27)

where the constant  $C$  is given by

$$
C = C(n,p) := \frac{4}{3p} \left( \frac{3p}{6p-4} \right)^{\frac{2-3p}{2}} \left( \frac{18p^2}{|n-2p+1|^2} \right)^{\frac{3p}{2}} > 0.
$$

**Step 2. Further estimate inequality** [\(2.23\)](#page-13-3). Applying in (2.23) the estimate [\(2.25\)](#page-13-4) for  $\sqrt{K_1}$  on the second term and Young's inequality with weight  $\tilde{\varepsilon}_1$  on the third term, we find that

$$
(3p-4)\mathcal{K}_3 + \mathcal{K}_4 + \mathcal{B} \le 3p\sqrt{\mathcal{K}_2}\sqrt{\mathcal{K}_3} + \frac{|n-2p+2|}{|n-2p+1|} \left(3p\sqrt{\mathcal{K}_2}\sqrt{\mathcal{K}_3} + (3p-2)\mathcal{K}_3 + \sqrt{d}\sqrt{\mathcal{K}_3}\sqrt{\mathcal{K}_4}\right)
$$
  
+  $\tilde{\epsilon}_1 \mathcal{K}_1 + C_{\tilde{\epsilon}_1} \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx$   
=  $3p \left(1 + \frac{|n-2p+2|}{|n-2p+1|}\right) \sqrt{\mathcal{K}_2}\sqrt{\mathcal{K}_3} + \frac{|n-2p+2|}{|n-2p+1|} \left((3p-2)\mathcal{K}_3 + \sqrt{d}\sqrt{\mathcal{K}_3}\sqrt{\mathcal{K}_4}\right)$   
+  $\tilde{\epsilon}_1 \mathcal{K}_1 + C_{\tilde{\epsilon}_1} \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx$ 

with

$$
\tilde{\varepsilon}_1 := \varepsilon_1 \frac{(n-2p+1)^2}{4} > 0 \quad \text{and} \quad C_{\tilde{\varepsilon}_1} := \frac{1}{p} \left( \frac{p}{p-1} \tilde{\varepsilon}_1 \right)^{1-p} > 0
$$

for some arbitrary constant  $\epsilon_1 > 0$ . By Young's inequality with weight  $\frac{\epsilon_2}{2}$  for the second term on the right-hand side of the above inequality and [\(2.27\)](#page-13-5), we further obtain

<span id="page-14-0"></span>
$$
(3p-4)\mathcal{K}_3 + \mathcal{K}_4 + \mathcal{B} \le 3p \left( 1 + \frac{|n-2p+2|}{|n-2p+1|} \right) \sqrt{\mathcal{K}_2} \sqrt{\mathcal{K}_3}
$$
  
+ 
$$
\frac{|n-2p+2|}{|n-2p+1|} \left[ \left( (3p-2) + \frac{\sqrt{d}}{2\epsilon_2} \right) \mathcal{K}_3 + \frac{\epsilon_2 \sqrt{d}}{2} \mathcal{K}_4 \right]
$$
  
+ 
$$
\epsilon_1 \left( (3p-2) \sqrt{\mathcal{K}_3} + \sqrt{d} \sqrt{\mathcal{K}_4} \right)^2
$$
  
+ 
$$
\tilde{\epsilon}_1 C \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx
$$
  
+ 
$$
C_{\tilde{\epsilon}_1} \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx
$$
 (2.28)

for some arbitrary  $\varepsilon_2 > 0$ .

Now we estimate the first term on the right-hand side of [\(2.28\)](#page-14-0). Recalling the estimate [\(2.24\)](#page-13-1) for  $\mathcal{K}_2$  and applying twice Young's inequality with weights  $\frac{\epsilon_3}{2}$  and  $\epsilon_4$ , we have

<span id="page-14-1"></span>
$$
3p\left(1+\frac{|n-2p+2|}{|n-2p+1|}\right)\sqrt{\mathcal{K}_{2}}\sqrt{\mathcal{K}_{3}}
$$
\n
$$
\leq 3p\left(1+\frac{|n-2p+2|}{|n-2p+1|}\right)\mathcal{K}_{1}^{\frac{3p-2}{6p}}\left(\int_{\Omega}|\nabla\varphi|^{3p}u^{n+p}\,dx\right)^{\frac{1}{3p}}\sqrt{\mathcal{K}_{3}}
$$
\n
$$
\leq \frac{\varepsilon_{3}}{2}\mathcal{K}_{3}+\frac{9p^{2}}{2\varepsilon_{3}}\left(1+\frac{|n-2p+2|}{|n-2p+1|}\right)^{2}\mathcal{K}_{1}^{\frac{3p-2}{3p}}\left(\int_{\Omega}|\nabla\varphi|^{3p}u^{n+p}\,dx\right)^{\frac{2}{3p}}
$$
\n
$$
\leq \frac{\varepsilon_{3}}{2}\mathcal{K}_{3}+\varepsilon_{4}\mathcal{K}_{1}+\left(\frac{3p}{3p-2}\varepsilon_{4}\right)^{\frac{2-3p}{2}}\frac{2}{3p}\left[\frac{9p^{2}}{2\varepsilon_{3}}\left(1+\frac{|n-2p+2|}{|n-2p+1|}\right)^{2}\right]^{\frac{3p}{2}}\int_{\Omega}|\nabla\varphi|^{3p}u^{n+p}\,dx
$$
\n(2.29)

with some  $\varepsilon_3$ ,  $\varepsilon_4$  > 0. Inserting [\(2.29\)](#page-14-1) and [\(2.27\)](#page-13-5) back into [\(2.28\)](#page-14-0) and rearranging, we have

<span id="page-15-0"></span>
$$
(3p-4)\mathcal{K}_3 + \mathcal{K}_4 + \mathcal{B}
$$
  
\n
$$
\leq \frac{\varepsilon_3}{2}\mathcal{K}_3 + \left(\frac{4\varepsilon_4}{|n-2p+1|^2} + \varepsilon_1\right) \left((3p-2)\sqrt{\mathcal{K}_3} + \sqrt{d}\sqrt{\mathcal{K}_4}\right)^2
$$
  
\n
$$
+ \frac{|n-2p+2|}{|n-2p+1|} \left[ \left((3p-2) + \frac{\sqrt{d}}{2\varepsilon_2}\right) \mathcal{K}_3 + \frac{\varepsilon_2\sqrt{d}}{2}\mathcal{K}_4 \right]
$$
  
\n
$$
+ \left[ (\varepsilon_4 + \tilde{\varepsilon}_1)C + \left(\frac{3p}{3p-2}\varepsilon_4\right)^{\frac{2-3p}{2}} \frac{2}{3p} \left[ \frac{9p^2}{2\varepsilon_3} \left(1 + \frac{|n-2p+2|}{|n-2p+1|}\right)^2 \right]^{\frac{3p}{2}} \right] \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx
$$
  
\n
$$
+ C_{\tilde{\varepsilon}_1} \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx.
$$
 (2.30)

**Step 3. Finish the proof.** We now want to absorb the terms on the right-hand side of [\(2.30\)](#page-15-0) into its left-hand side. Therefore, we choose  $\varepsilon_2$  as the unique positive solution of

$$
(3p-2) + \frac{\sqrt{d}}{2\varepsilon_2} = (3p-4)\frac{\varepsilon_2\sqrt{d}}{2},
$$

since this absorbs the second line on the right-hand side of [\(2.30\)](#page-15-0) in an optimal way and also leads us to the desired bound for *n*. Solving this equation for  $\varepsilon_2 > 0$  gives

$$
\varepsilon_2 = \frac{3p - 2 + \sqrt{(3p - 2)^2 + d(3p - 4)}}{\sqrt{d(3p - 4)}}.
$$

<span id="page-15-1"></span>If

$$
\frac{|n-2p+2|}{|n-2p+1|} \varepsilon_2 \sqrt{d} < 2,\tag{2.31}
$$

then we may absorb the second line on the right-hand side into the left-hand side. Inserting the equation for  $\epsilon_2$ into  $(2.31)$  gives us a bound for *n* in terms of *p* and *d*. Indeed, the inequality

$$
\frac{|n-2p+2|}{|n-2p+1|}\frac{3p-2+\sqrt{(3p-2)^2+d(3p-4)}}{3p-4}<2
$$

is satisfied for

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}
$$

Next, using Young's inequality in the second term on the right-hand side of [\(2.30\)](#page-15-0) and applying [\(2.31\)](#page-15-1), we find that [\(2.30\)](#page-15-0) can be further estimated as follows:

<span id="page-15-2"></span>
$$
(3p-4)\mathcal{K}_3 + \mathcal{K}_4 + \mathcal{B}
$$
\n
$$
\leq 2\left[\frac{\varepsilon_3}{2} + \left(\frac{4\varepsilon_4}{|n-2p+1|^2} + \varepsilon_1\right)(3p-2)\left((3p-2) + \sqrt{d}\right)\right]\mathcal{K}_3
$$
\n
$$
+ 2\left(\frac{4\varepsilon_4}{|n-2p+1|^2} + \varepsilon_1\right)\sqrt{d}\left((3p-2) + \sqrt{d}\right)\mathcal{K}_4
$$
\n
$$
+ 2\left[\left(\varepsilon_4 + \tilde{\varepsilon}_1\right)C + \left(\frac{3p}{3p-2}\varepsilon_4\right)^{\frac{2-3p}{2}}\frac{2}{3p}\left[\frac{9p^2}{2\varepsilon_3}\left(1 + \frac{|n-2p+2|}{|n-2p+1|}\right)^2\right]^{\frac{3p}{2}}\right]\int_{\Omega}|\nabla\varphi|^{3p}u^{n+p}\,dx
$$
\n
$$
+ 2C_{\tilde{\varepsilon}_1}\int_{\Omega}\varphi^{3p}u^n|\nabla\Delta u|^p\,dx.
$$
\n(2.32)

Now we can choose  $\epsilon_1$ ,  $\epsilon_3$  and  $\epsilon_4$  such that we can absorb the first two lines on the right-hand side of [\(2.32\)](#page-15-2) into its left-hand side. Hence, we find a constant  $\tilde{C} = \tilde{C}(n, p, d) > 0$  such that

$$
\mathcal{K}_3 + \mathcal{K}_4 + \mathcal{B} \le \tilde{C} \left[ \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} \, \mathrm{d}x + \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p \, \mathrm{d}x \right].
$$

Combining this estimate with [\(2.27\)](#page-13-5) concludes the proof.  $\Box$ 

The final Bernis-type inequality, presented in the following proposition, provides an estimate for an integral involving the product of the function *u* and its Laplacian. The proof does not require a case-by-case analysis based on the mobility exponent *n* but can be directly established for the entire range of *n*. The key ingredients of the proof are the inequalities [\(2.2\)](#page-6-1) and [\(2.20\)](#page-12-1), which control the integral  $\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx$ . Additionally, we note that the corresponding result for the one-dimensional case is presented in [\[AG04\]](#page-22-2).

<span id="page-16-0"></span>**Proposition 2.7.** *Let*  $\Omega \subset \mathbb{R}^d$  *be a bounded, convex and smooth domain, and let*  $2 \leq p < \frac{19}{3}$  *be fixed. Assume fhat*  $u \in C^{\infty}(\bar{\Omega})$  *is positive and satisfies*  $\nabla u \cdot v = 0$  *on*  $\partial \Omega$ *, and let*  $\varphi \in W_{\infty}^{1}(\Omega)$  *be non-negative. If* 

$$
2p-2 \frac{2+\sqrt{(3p-2)^2+d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2+d(3p-4)}}
$$

*then there exists a positive constant*  $C_5 = C_5(n, p, d) > 0$  *such that the following estimate holds:* 

$$
\int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} \, \mathrm{d}x \leq \, C_5 \left( \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p \, \mathrm{d}x + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} \, \mathrm{d}x \right). \tag{2.33}
$$

*Proof.* Using integration by parts, we can rewrite

$$
\int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} dx = -3p \int_{\Omega} \varphi^{3p-1} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p-4}{2}} \Delta u \nabla \varphi \cdot \nabla u dx \n- \left(n - \frac{p}{2}\right) \int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}-1} |\nabla u|^2 |\Delta u|^{\frac{3p-4}{2}} \Delta u dx \n- \left(\frac{3p-2}{2}\right) \int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p-4}{2}} \nabla u \cdot \nabla \Delta u dx.
$$
\n(2.34)

Applying now Hölder's and Young's inequalities on each of the integrals on the right-hand side yields the estimates

<span id="page-16-1"></span>
$$
\int_{\Omega} \varphi^{3p-1} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p-4}{2}} \Delta u \nabla \varphi \cdot \nabla u \, dx
$$
\n
$$
\leq \left( \int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} dx \right)^{\frac{3p-2}{3p}} \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{1}{3p}} \left( \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)^{\frac{1}{3p}}
$$
\n
$$
\leq \epsilon_1 \int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} dx + C_{\epsilon_1} \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx + \tilde{C}_{\epsilon_1} \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx,
$$
\n(2.35)

for the first integral,

$$
\int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}-1} |\nabla u|^{2} |\Delta u|^{\frac{3p-4}{2}} \Delta u \, dx \leq \left( \int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} dx \right)^{\frac{3p-2}{3p}} \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{2}{3p}} \leq \varepsilon_{2} \int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} dx + C_{\varepsilon_{2}} \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \tag{2.36}
$$

for the second integral, and

<span id="page-16-2"></span>
$$
\int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p-4}{2}} \nabla u \cdot \nabla \Delta u \, dx
$$
\n
$$
\leq \left( \int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} \, dx \right)^{\frac{3p-4}{3p}} \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} \, dx \right)^{\frac{1}{3p}} \left( \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} \, dx \right)^{\frac{1}{p}} \n\leq \varepsilon_{3} \int_{\Omega} \varphi^{3p} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} \, dx + C_{\varepsilon_{3}} \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} \, dx + \tilde{C}_{\varepsilon_{3}} \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} \, dx \n\tag{2.37}
$$

for the third one, where we chose  $0 < \varepsilon_1 + \varepsilon_2 + \varepsilon_3 < 1$ . By the inequalities [\(2.2\)](#page-6-1), respectively [\(2.20\)](#page-12-1) we can estimate the integral  $\int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx$  in [\(2.35\)](#page-16-1)–[\(2.37\)](#page-16-2) in terms of  $\int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} dx$  and  $\int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx$ . This yields the desired inequality.  $\Box$ 

To complete the proof of Theorem [1.1,](#page-1-1) it remains to perform the approximation argument to establish the result in the desired functional setting, i.e., for functions  $u \in W_p^2(\Omega)$  with  $2 \le p < \frac{19}{3}$ . Note that all results in this section are valid for arbitrary dimensions  $d$ . The restriction  $d < 3p$  arises only during the approximation argument, where Sobolev embeddings are employed.

*Proof of Theorem [1.1](#page-1-1).* The proof follows the idea of the proof of [\[Grü01,](#page-22-0) Theorem 1.1], using elliptic regularity theory in  $L_p(\Omega)$  instead of  $L_2(\Omega)$ . Let  $u \in W_p^2(\Omega)$ , with  $2 \le p < \frac{19}{3}$ , be a positive function satisfying  $\nabla u \cdot v = 0$ on *𝜕*Ω,

<span id="page-17-0"></span>
$$
\int_{\Omega} u^n |\nabla \Delta u|^p \, dx < \infty. \tag{2.38}
$$

We may assume that there is  $\delta > 0$  such that  $u \ge \delta > 0$  a.e. in  $\Omega$ . Using [\(2.38\)](#page-17-0), this implies  $\int_{\Omega} |\nabla \Delta u|^p dx < \infty$ . To prove the claim, we construct a sequence of functions  $(u_k)_{k \in \mathbb{N}} \subset C^\infty(\bar{\Omega})$  with the following properties:

- $(1)$   $u_k \geq \frac{\delta}{2}$  $\frac{\delta}{2}$  in  $\bar{\Omega}$  for  $k \in \mathbb{N}$  sufficiently large;
- (2)  $\nabla u_k \cdot \vec{v} = 0$  on  $\partial \Omega$ ;
- (3)  $u_k \longrightarrow u$  strongly in  $W_p^2(\Omega)$ , as  $k \longrightarrow \infty$ ;
- (4)  $\nabla \Delta u_k \longrightarrow \nabla \Delta u$  strongly in  $L_p(\Omega)$ , as  $k \to \infty$ .

Once this sequence  $u_k$  is constructed, we know it satisfies the inequalities of Propositions [2.3,](#page-6-0) [2.5](#page-12-0) and [2.7](#page-16-0) for the whole range

$$
2p - 2 \frac{2 + \sqrt{(3p-2)^2 + d(3p-4)}}{-3p+6+\sqrt{(3p-2)^2 + d(3p-4)}} < n < 2p-1.
$$

We can then take the limit on both sides of the corresponding inequalities. For the right-hand side, we apply Vitali's convergence theorem, while for the left-hand side, we use Fatou's lemma. This yields the corresponding estimates  $(1.3)$ ,  $(1.4)$  and  $(1.5)$  for the limit function *.* 

We now construct the sequence  $u_k$ . Since  $\nabla \Delta u \in L_p(\Omega)$ , we have  $\Delta u \in W^1_p(\Omega)$ . Thus, there exists a sequence  $(f_k)_{k \in \mathbb{N}}$  ⊂  $C^\infty(\overline{\Omega})$  satisfying

$$
f_k \longrightarrow -\Delta u
$$
 strongly in  $W_p^1(\Omega)$  and  $\int_{\Omega} f_k dx = 0$ ,

see [\[GT01\]](#page-22-15). Here, the zero-mean condition can be obtained since  $\int_{\Omega} \Delta u \, dx = \int_{\partial \Omega} \nabla u \cdot v = 0$  by the Neumann boundary condition on *u*. For each  $k \in \mathbb{N}$ , let  $u_k$  be the solution to the Neumann problem

$$
\begin{cases}\n-\Delta u_k = f_k, & x \in \Omega \\
\nabla u_k \cdot v = 0, & x \in \partial \Omega \\
\int_{\Omega} u_k \, dx = \int_{\Omega} u \, dx.\n\end{cases}
$$

By elliptic regularity theory [\[Gri11,](#page-22-16) Theorem 2.5.1.1, Remark 2.5.1.2], we find that  $u_k \in C^\infty(\bar{\Omega})$  and

 $u_k \longrightarrow u$  strongly in  $W_p^3(\Omega)$ .

In particular,  $u_k$  satisfies (2), (3), and (4). Moreover, by the Sobolev embedding theorem, we have

$$
W_p^3(\Omega) \hookrightarrow W_q^2(\Omega) \quad \text{for } 1 \le q < \frac{dp}{d-p} \quad \text{and} \quad W_q^2(\Omega) \hookrightarrow C(\bar{\Omega}) \quad \text{for } q > \frac{d}{2}.
$$

Consequently, we find that

$$
\begin{cases} u_k \longrightarrow u & \text{strongly in } W_q^2(\Omega) \quad \text{for } 1 \le q < \frac{dp}{d-p} \\ u_k \longrightarrow u & \text{strongly in } C(\bar{\Omega}) \quad \text{for } d < 3p. \end{cases}
$$

The strong convergence  $u_k \to u$  in  $C(\overline{\Omega})$  and  $u \ge \delta > 0$  imply (1) for  $k \in \mathbb{N}$  large enough. This concludes the construction of the sequence  $u_k$  satisfying properties (1)–(4), thereby completing the proof of the theorem.  $\Box$ 

*.*

## 3. PROOF OF THEOREM [1.3](#page-2-0)

<span id="page-18-0"></span>We now prove Theorem [1.3](#page-2-0) which is a consequence of the inequalities [\(1.3\)](#page-1-2) and [\(1.5\)](#page-2-2) of Theorem [1.1.](#page-1-1)

*Proof of Theorem [1.3](#page-2-0)*. The first parts of [\(1.6\)](#page-3-4), respectively [\(1.7\)](#page-3-1), follow from [\(1.3\)](#page-1-2), while parts two and three follow from [\(1.3\)](#page-1-2) and [\(1.5\)](#page-2-2). We mention again that we actually prove weighted versions of the first-order and second-order estimate in [\(1.7\)](#page-3-1).

**Parts one of** [\(1.6\)](#page-3-4), [\(1.7\)](#page-3-1): We first show that  $u^{\frac{n+p}{3p}}$  lies in  $W^1_{3p}(\Omega)$  and that the inequality

$$
\int_{\Omega} \varphi^{3p} \left| \nabla \left( u^{\frac{n+p}{3p}} \right) \right|^{3p} dx \le C \left( \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)
$$
\n(3.1)

holds for some constant  $C = C(n, p, d) > 0$  and for all non-negative test functions  $\varphi \in W^1_{\infty}(\Omega)$  satisfying  $\int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx <$  $\infty$ . Since  $u \in W_p^2(\Omega)$  and  $W_p^2(\Omega) \hookrightarrow L_{n+p}(\Omega)$ , we have  $u^{\frac{n+p}{3p}} \in L_{3p}(\Omega)$ . In addition, we compute that

$$
\nabla \left( u^{\frac{n+p}{3p}} \right) = \frac{n+p}{3p} u^{\frac{n-2p}{3p}} \nabla u,
$$
\n(3.2)

and together with [\(1.3\)](#page-1-2) from Theorem [1.1,](#page-1-1) we obtain that  $u^{\frac{n+p}{3p}} \in W^1_{3p}(\Omega)$  with the inequality

$$
\int_{\Omega} \varphi^{3p} \left| \nabla \left( u^{\frac{n+p}{3p}} \right) \right|^{3p} dx = \left( \frac{n+p}{3p} \right)^{3p} \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx
$$
  

$$
\leq \left( \frac{n+p}{3p} \right)^{3p} c_1 \left( \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right) < \infty.
$$

Setting  $\varphi \equiv 1$  yields the first estimate.

**Parts two of** [\(1.6\)](#page-3-4), [\(1.7\)](#page-3-1): Next, we show that  $u^{\frac{2(n+p)}{3p}}$  belongs to  $W^2_{\frac{3p}{2}}$  $(\Omega)$  and that there exists a positive constant  $\tilde{C} = \tilde{C}(n, p, d)$  such that

<span id="page-18-2"></span>
$$
\int_{\Omega} \varphi^{3p} \left| \Delta \left( u^{\frac{2(n+p)}{3p}} \right) \right|^{3p} dx \leq \tilde{C} \left( \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)
$$
\n(3.3)

holds for all non-negative test functions  $\varphi \in W^1_{\infty}(\Omega)$  satisfying  $\int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx < \infty$ . Since  $u \in W^2_{p}(\Omega) \hookrightarrow$  $L_{n+p}(\Omega)$ , we obtain as before that  $u^{\frac{2(n+p)}{3p}} \in L_{\frac{3p}{2}}(\Omega)$ . For  $\nabla(u^{\frac{2(n+p)}{3p}})$ , we compute

<span id="page-18-3"></span>
$$
\nabla \left( u^{\frac{2(n+p)}{3p}} \right) = \frac{2(n+p)}{3p} u^{\frac{2n-p}{3p}} \nabla u \tag{3.4}
$$

and use Hölder's inequality together with [\(1.3\)](#page-1-2) (with  $\varphi \equiv 1$ ) to find that

$$
\int_{\Omega} \left| \nabla \left( u^{\frac{2(n+p)}{3p}} \right) \right|^{\frac{3p}{2}} dx = \left( \frac{2(n+p)}{3p} \right)^{\frac{3p}{2}} \int_{\Omega} u^{\frac{n+p}{2}} u^{\frac{n}{2} - p} |\nabla u|^{\frac{3p}{2}} dx
$$
  

$$
\leq \left( \frac{2(n+p)}{3p} \right)^{\frac{3p}{2}} c_1^{\frac{1}{2}} \left( \int_{\Omega} u^{n+p} dx \right)^{\frac{1}{2}} \left( \int_{\Omega} u^n |\nabla \Delta u|^p dx \right)^{\frac{1}{2}} < \infty.
$$

Hence,  $\nabla(u^{\frac{2(n+p)}{3p}}) \in L_{\frac{3p}{2}}(\Omega;\mathbb{R}^d)$ . Now, we compute

$$
\Delta(u^{\frac{2(n+p)}{3p}}) = \frac{2(n+p)}{3p} \left( \frac{2n-p}{3p} u^{\frac{2(n-2p)}{3p}} |\nabla u|^2 + u^{\frac{2n-p}{3p}} \Delta u \right),
$$

and by Jensen's inequality we further have that

<span id="page-18-1"></span>
$$
\left|\Delta\left(u^{\frac{2(n+p)}{3p}}\right)\right|^{\frac{3p}{2}} \le c\left(u^{n-2p}|\nabla u|^{3p} + u^{n-\frac{p}{2}}|\Delta u|^{\frac{3p}{2}}\right) \tag{3.5}
$$

for some constant  $c = c(n, p) > 0$ . Using [\(1.3\)](#page-1-2) and [\(1.5\)](#page-2-2), we infer from [\(3.5\)](#page-18-1) that  $\Delta(u^{\frac{2(n+p)}{3p}})$  belongs to  $L_{\frac{3p}{2p}}(\Omega)$ and that [\(3.3\)](#page-18-2) holds. We deduce the second estimate in [\(1.7\)](#page-3-1) by setting  $\varphi \equiv 1$  in (3.3). It remains to check that  $D^2(u^{\frac{2(n+p)}{3p}}) \in L_{\frac{3p}{2}}(\Omega;\mathbb{R}^{d \times d})$ . It follows from [\(3.4\)](#page-18-3) and the Neumann boundary condition for u that  $\nabla(u^{\frac{2(n+p)}{3p}}) \cdot v =$ 0 on  $\partial \Omega$ . Elliptic *L<sub>p</sub>*-theory (see for instance [\[Gri11](#page-22-16), Thm. 2.4.2.7 & 2.5.5.1]) then proves the claim.

**Parts three of** [\(1.6\)](#page-3-4), [\(1.7\)](#page-3-1): Finally, we show that  $u^{\frac{n+p}{p}}$  belongs to  $W_p^3(\Omega)$  and that there exists a constant  $\hat{C}$  =  $\hat{C}(n, p, d) > 0$  such that

<span id="page-19-4"></span>
$$
\int_{\Omega} \left| \nabla \Delta \left( u^{\frac{n+p}{p}} \right) \right|^{p} dx \leq \hat{C} \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx.
$$
\n(3.6)

Because  $u \in W_p^2(\Omega) \hookrightarrow L_{n+p}(\Omega)$ , we have  $u^{\frac{n+p}{p}} \in L_p(\Omega)$ . For the gradient of  $u^{\frac{n+p}{p}}$ , we calculate

<span id="page-19-0"></span>
$$
\nabla \left( u^{\frac{n+p}{p}} \right) = \frac{n+p}{p} u^{\frac{n}{p}} \nabla u,
$$
\n(3.7)

and using Hölder's inequality along with [\(1.3\)](#page-1-2) for  $\varphi \equiv 1$  we deduce that

$$
\int_{\Omega} \left| \nabla \left( u^{\frac{n+p}{p}} \right) \right|^{p} dx = \left( \frac{n+p}{p} \right)^{p} \int_{\Omega} u^{n} |\nabla u|^{p} dx
$$
\n
$$
\leq \left( \frac{n+p}{p} \right)^{p} \left( \int_{\Omega} u^{n+p} dx \right)^{\frac{2}{3}} \left( \int_{\Omega} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{1}{3}}
$$
\n
$$
\leq \left( \frac{n+p}{p} \right)^{p} c_{1}^{\frac{1}{3}} \left( \int_{\Omega} u^{n+p} dx \right)^{\frac{2}{3}} \left( \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx \right)^{\frac{1}{3}} < \infty.
$$

Thus,  $\nabla (u^{\frac{n+p}{p}}) \in L_p(\Omega;\mathbb{R}^d)$ . For the Laplacian of  $u^{\frac{n+p}{p}}$ , we compute

$$
\Delta\left(u^{\frac{n+p}{p}}\right) = \frac{n+p}{p}u^{\frac{n}{p}}\Delta u + \frac{n(n+p)}{p^2}u^{\frac{n-p}{p}}|\nabla u|^2.
$$

By applying Jensen's and Hölder's inequalities, along with [\(1.3\)](#page-1-2) and [\(1.5\)](#page-2-2), again for  $\varphi \equiv 1$ , we obtain

$$
\int_{\Omega} \left| \Delta \left( u^{\frac{n+p}{p}} \right) \right|^{p} dx \leq c \left( \int_{\Omega} u^{n} |\Delta u|^{p} dx + \int_{\Omega} u^{n-p} |\nabla u|^{2p} dx \right)
$$
  

$$
\leq c \left( \int_{\Omega} u^{n+p} dx \right)^{\frac{1}{3}} \left[ \left( \int_{\Omega} u^{n-\frac{p}{2}} |\Delta u|^{\frac{3p}{2}} dx \right)^{\frac{2}{3}} + \left( \int_{\Omega} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{2}{3}} \right]
$$
  

$$
\leq c \left( c_1^{\frac{2}{3}} + c_3^{\frac{2}{3}} \right) \left( \int_{\Omega} u^{n+p} dx \right)^{\frac{1}{3}} \left( \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx \right)^{\frac{2}{3}} < \infty
$$

for some constant  $c = c(n, p) > 0$ . Hence,  $\Delta(u^{\frac{n+p}{p}}) \in L_p(\Omega)$ . To verify that  $D^2(u^{\frac{n+p}{p}})$  belongs to  $L_p(\Omega; \mathbb{R}^{d \times d})$ , observe that from [\(3.7\)](#page-19-0) and the Neumann boundary condition for *u*, we have  $\nabla (u^{\frac{n+p}{p}}) \cdot v = 0$  on  $\partial \Omega$ . The claim then follows by applying elliptic  $L_p$ -theory (see, e.g., [\[Gri11](#page-22-16), Thm. 2.4.2.7]). Next, it follows from

$$
\nabla\Delta\left(u^{\frac{(n+p)}{p}}\right) = \frac{(n+p)n(n-p)}{p^3} u^{\frac{n-2p}{p}} |\nabla u|^2 \nabla u + \frac{2(n+p)n}{p^2} u^{\frac{n-p}{p}} D^2 u \nabla u + \frac{(n+p)n}{p^2} u^{\frac{n-p}{p}} \Delta u \nabla u + \frac{n+p}{p} u^{\frac{n}{p}} \nabla\Delta u
$$

and Jensen's inequality that there is a constant  $\tilde{c} = \tilde{c}(n, p) > 0$  such that

<span id="page-19-3"></span>
$$
|\nabla\Delta(u^{\frac{n+p}{p}})|^p \le \tilde{c}\left(u^{n-2p}|\nabla u|^{3p} + u^{n-p}|D^2u\nabla u|^p + u^{n-p}|\Delta u\nabla u|^p + u^n|\nabla\Delta u|^p\right).
$$
\n(3.8)

The main task is to estimate the second-order term  $u^{n-p} | D^2 u \nabla u |^p$ . To this end, we compute for  $s \in \mathbb{R} \setminus \{0\}$  the following two identities:

$$
D^{2}u\nabla u = \frac{1}{s}u^{1-s}D^{2}u^{s}\nabla u - (s-1)u^{-1}|\nabla u|^{2}\nabla u \text{ and } \Delta u\nabla u = \frac{1}{s}u^{1-s}\Delta u^{s}\nabla u - (s-1)u^{-1}|\nabla u|^{2}\nabla u.
$$

Multiplying these identities by  $u^{\frac{n-p}{p}}$ , we rewrite them as

<span id="page-19-1"></span>
$$
u^{\frac{n-p}{p}} D^2 u \nabla u = \frac{1}{s} u^{\frac{n}{p}-s} D^2 u^s \nabla u - (s-1) u^{\frac{n-2p}{p}} |\nabla u|^2 \nabla u \tag{3.9}
$$

<span id="page-19-2"></span>and

$$
u^{\frac{n-p}{p}} \Delta u \nabla u = \frac{1}{s} u^{\frac{n}{p} - s} \Delta u^s \nabla u - (s - 1) u^{\frac{n-2p}{p}} |\nabla u|^2 \nabla u.
$$
 (3.10)

For the term  $u^{\frac{n-2}{p}} |\nabla u|^2 \nabla u$ , appearing in [\(3.9\)](#page-19-1) and [\(3.10\)](#page-19-2), a straightforward calculation gives the identity

$$
u^{\frac{n-2p}{p}}|\nabla u|^2 \nabla u = \left(\frac{3p}{n+p}\right)^3 |\nabla (u^{\frac{n+p}{3p}})|^2 \nabla (u^{\frac{n+p}{3p}}).
$$

$$
u^{n-p} |D^2 u \nabla u|^p \leq \hat{c} \left( u^{n-sp} |D^2 u^s \nabla u|^p + |\nabla \left( u^{\frac{n+p}{3p}} \right)|^{3p} \right)
$$

and

$$
u^{n-p} |\Delta u \nabla u|^p \leq \hat{c} \left( u^{n-sp} |\Delta u^s \nabla u|^p + |\nabla \left( u^{\frac{n+p}{3p}} \right)|^{3p} \right)
$$

with a constant  $\hat{c} > 0$ , depending only on *n*, *p* and *s*. Setting  $s = \frac{2(n+p)}{n}$  $\frac{(3+1)^{n+p}}{3p}$ , it follows from [\(3.8\)](#page-19-3), by applying the above inequalities and Hölder's inequality, along with [\(1.3\)](#page-1-2) and the estimates from the first and second part of  $(1.7)$  with  $\varphi \equiv 1$ , that

<span id="page-20-2"></span>
$$
\int_{\Omega} \left| \nabla \Delta (u^{\frac{n+p}{p}}) \right|^{p} dx
$$
\n
$$
\leq \tilde{c} \int_{\Omega} (u^{n-2p} |\nabla u|^{3p} + u^{n-p} |D^{2}u \nabla u|^{p} + u^{n-p} |\Delta u \nabla u|^{p} + u^{n} |\nabla \Delta u|^{p}) dx
$$
\n
$$
\leq c_{*} \left( \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx + \int_{\Omega} |\nabla (u^{\frac{n+p}{3p}})|^{3p} dx + \int_{\Omega} u^{\frac{n-2p}{3}} |\Delta (u^{\frac{2(n+p)}{3p}}) \nabla u|^{p} dx \right.
$$
\n
$$
+ \int_{\Omega} u^{\frac{n-2p}{3}} |D^{2} (u^{\frac{2(n+p)}{3p}}) \nabla u|^{p} dx \right)
$$
\n
$$
\leq c_{*} \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx + c_{*} \left( \int_{\Omega} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{1}{3}} \left( \int_{\Omega} |\Delta (u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx \right)^{\frac{2}{3}}
$$
\n
$$
+ c_{*} \left( \int_{\Omega} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{1}{3}} \left( \int_{\Omega} |D^{2} (u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx \right)^{\frac{2}{3}}
$$
\n
$$
\leq c_{*} \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx + c_{*} \left( \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx \right)^{\frac{1}{3}} \left( \int_{\Omega} |D^{2} (u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx \right)^{\frac{2}{3}},
$$
\n
$$
\geq 0 \text{ is a generic constant that may change from line to line but depends only on } n \text{, and } d \text{. Due}
$$

where  $c_* > 0$  is a generic constant that may change from line to line, but depends only on *n*, *p*, and *d*. Due to the fact that  $D^2(u^{\frac{2(n+p)}{3p}}) \in L_{\frac{3p}{2}}(\Omega;\mathbb{R}^{d \times d})$ , we deduce that  $\nabla \Delta(u^{\frac{n+p}{p}}) \in L_p(\Omega;\mathbb{R}^d)$ . Therefore, applying again elliptic  $L_p$ -theory (see [\[Gri11](#page-22-16), Thm. 2.5.5.1]), we conclude that  $u^{\frac{n+p}{p}} \in W_p^3(\Omega)$ . Furthermore, using [\[Gri11,](#page-22-16) Thm. 2.3.3.6], we obtain the estimate

<span id="page-20-1"></span>
$$
\|D^{2}(u^{\frac{2(n+p)}{3p}})\|_{L_{\frac{3p}{2}}(\Omega)} \leq c^{*}\left(\|\Delta(u^{\frac{2(n+p)}{3p}})\|_{L_{\frac{3p}{2}}(\Omega)} + \|u^{\frac{2(n+p)}{3p}}\|_{L_{\frac{3p}{2}}(\Omega)}\right)
$$
  

$$
\leq c^{*}\left(\|\Delta(u^{\frac{2(n+p)}{3p}})\|_{L_{\frac{3p}{2}}(\Omega)} + \|\nabla(u^{\frac{2(n+p)}{3p}})\|_{L_{\frac{3p}{2}}(\Omega)}\right)
$$
(3.12)

for some generic constant  $c^*$  > 0 depending on *n*, *p*, and  $\Omega$ . To establish inequality [\(3.6\)](#page-19-4) we insert the elliptic estimate [\(3.12\)](#page-20-1) into [\(3.11\)](#page-20-2) and apply Young's inequality as well as the first two statements of the theorem. This leads to

$$
\int_{\Omega} \left| \nabla \Delta \left( u^{\frac{n+p}{p}} \right) \right|^{p} dx \leq c_{*} \left( \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx + \int_{\Omega} |\Delta \left( u^{\frac{2(n+p)}{3p}} \right)|^{\frac{3p}{2}} dx + \int_{\Omega} |\nabla \left( u^{\frac{2(n+p)}{3p}} \right)|^{\frac{3p}{2}} dx \right)
$$
  

$$
\leq c_{*} \int_{\Omega} u^{n} |\nabla \Delta u|^{p} dx.
$$

Thus, the proof is complete.  $\Box$ 

<span id="page-20-0"></span>We can improve the estimates in Theorem [1.3](#page-2-0) to localized versions as follows.

**Corollary 3.1.** *Under the assumptions of Theorem [1.3](#page-2-0) we have the additional weighted estimates*

$$
\int_{\Omega} \varphi^{3p} \left| \nabla \left( u^{\frac{n+p}{3p}} \right) \right|^{3p} dx + \int_{\Omega} \varphi^{3p} \left| \Delta \left( u^{\frac{2(n+p)}{3p}} \right) \right|^{3p} dx \leq c_5 \left( \int_{\Omega} \varphi^{3p} u^{n} |\nabla \Delta u|^{p} dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)
$$

*for all non-negative test functions*  $\varphi \in W^1_\infty(\Omega)$  *satisfying*  $\int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx < \infty$ *, and where*  $c_5$  *is a positive constant depending only on n, p, and d.* 

<span id="page-21-0"></span>The proof of these two estimates is already contained in the proof of Theorem [1.3.](#page-2-0)

**Remark 3.2.** In Corollary [3.1,](#page-20-0) we address the estimates for the first- and second-order terms. However, for the third-order term, we can only establish the following result. Assuming, in addition to the assumptions of Theorem [1.3,](#page-2-0) that an elliptic estimate of the form

<span id="page-21-1"></span>
$$
\int_{\Omega} \psi |D^2 v|^s dx \leq c_s \left( \int_{\Omega} \psi |\Delta v|^s dx + \int_{\Omega} |D^2 v|^{s-1} |\nabla \psi| |\nabla v| dx + \text{l.o.t.} \right),\tag{3.13}
$$

for  $1 < s < \infty$  and suitable test functions  $\psi \geq 0$ , holds true for solutions v to the Neumann problem

$$
\begin{cases}\n-\Delta v + v = f & \text{in } \Omega \\
\nabla v \cdot v = 0 & \text{on } \partial \Omega,\n\end{cases}
$$

then we also get the weighted estimate

<span id="page-21-2"></span>
$$
\int_{\Omega} \varphi^{3p} \left| \nabla \Delta \left( u^{\frac{n+p}{p}} \right) \right|^p dx \le c_6 \left( \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right) \tag{3.14}
$$

for suitable non-negative test functions  $\varphi$  satisfying  $\int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx < \infty$  and for some constant  $c_6 = c_6(n, p, d) >$ 0. The authors are not aware whether an estimate of the form [\(3.13\)](#page-21-1) holds true for the Neumann problem. For the Dirichlet case, weighted elliptic estimates are available (cf. [\[CD18](#page-22-17); [DST08](#page-22-18); [DST10\]](#page-22-19)), provided the weight belongs to some Muckenhoupt class. In this case, the second integral on the right-hand side of [\(3.13\)](#page-21-1) becomes superfluous. However, since the test functions  $\varphi$  in this paper might vanish on sets of positive measure, they do not belong to the Muckenhoupt class. As a result, [\(3.13\)](#page-21-1), if it holds at all, would require the additional integral involving lower-order terms. It is also possible that the test function might need  $W_{\infty}^2$ -regularity for [\(3.13\)](#page-21-1) to be valid.

*Sketch of proof of* [\(3.14\)](#page-21-2) *under the assumption* [\(3.13\)](#page-21-1)*.* If an estimate of the form [\(3.13\)](#page-21-1) is satisfied, where we ignore the lower-order terms in the calculation, we can choose

$$
\psi = \varphi^{3p}
$$
,  $v = u^{\frac{2(n+p)}{3p}}$  and  $s = \frac{3p}{2}$ 

and apply Hölder's and Young's inequalities to obtain

$$
\int_{\Omega} \varphi^{3p} |D^{2}(u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx
$$
\n
$$
\leq c_{p} \int_{\Omega} \varphi^{3p} |\Delta(u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx + 2c_{p}(n+p) \int_{\Omega} \varphi^{3p-1} u^{\frac{2n-p}{3p}} |D^{2}(u^{\frac{2(n+p)}{3p}})|^{\frac{3p-2}{2}} |\nabla \varphi||\nabla u| dx
$$
\n
$$
\leq c_{p} \int_{\Omega} \varphi^{3p} |\Delta(u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx
$$
\n
$$
+ 2c_{p}(n+p) \left( \int_{\Omega} \varphi^{3p} |D^{2}(u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx \right)^{\frac{3p-2}{3p}} \left( \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)^{\frac{1}{3p}} \left( \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)^{\frac{1}{3p}}
$$
\n
$$
\leq c_{p} \int_{\Omega} \varphi^{3p} |\Delta(u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx + \varepsilon \int_{\Omega} \varphi^{3p} |D^{2}(u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx
$$
\n
$$
+ C_{\varepsilon} \left( \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx + \int_{\Omega} \varphi^{3p} u^{n-2p} |\nabla u|^{3p} dx \right)
$$

for some  $0 < \varepsilon < 1$ , where  $C_{\varepsilon} > 0$  depends only on *n*, *p*, and  $\varepsilon$ . Absorbing the second term on the right-hand side in the left-hand side, estimating the first term by  $(1.7)$  and the fourth term by  $(1.3)$ , we obtain the weighted estimate

<span id="page-21-3"></span>
$$
\int_{\Omega} \varphi^{3p} |D^2(u^{\frac{2(n+p)}{3p}})|^{\frac{3p}{2}} dx \le c_7 \left( \int_{\Omega} \varphi^{3p} u^n |\nabla \Delta u|^p dx + \int_{\Omega} |\nabla \varphi|^{3p} u^{n+p} dx \right)
$$
\n(3.15)

with a constant  $c_7 > 0$  depending on *n*, *p*, and *d*. Following the steps for [\(3.11\)](#page-20-2) and using [\(3.15\)](#page-21-3) we finally obtain  $(3.14).$  $(3.14).$ 

### REFERENCES 23

#### ACKNOWLEDGEMENTS

The authors would like to thank Jonas Jansen for fruitful discussions.

### FUNDING

KN was supported by ENW Vidi grant VI.Vidi.223.019 entitled 'Codimension two free boundary problems' of the Dutch Research Council NWO and by the APART-MINT Fellowship of the Austrian Academy of Sciences (ÖAW-APART-MINT Grant No. 12114)

## **REFERENCES**

- <span id="page-22-11"></span><span id="page-22-2"></span>[AG04] L. Ansini and L. Giacomelli. "Doubly Nonlinear Thin-Film Equations in One Space Dimension". In: *Archive for Rational Mechanics and Analysis* 173.1 (2004), pp. 89–131.
- <span id="page-22-5"></span>[Ber+98] M. Bertsch, R. Dal Passo, H. Garcke, and G. Grün. "The thin viscous flow equation in higher space dimensions". In: *Adv. Differential Equations* 3.3 (1998), pp. 417–440.
- <span id="page-22-6"></span>[Ber96a] F. Bernis. "Finite speed of propagation and continuity of the interface for thin viscous flows". In: *Adv. Differential Equations* 1.3 (1996), pp. 337–368.
- <span id="page-22-9"></span>[Ber96b] F. Bernis. "Finite speed of propagation for thin viscous flows when  $2 \le n < 3$ ". In: *C. R. Acad. Sci. Paris Sér. I Math.* 322.12 (1996), pp. 1169–1174.
- <span id="page-22-4"></span>[Ber96c] F. Bernis. "Integral inequalities with applications to nonlinear degenerate parabolic equations". In: *Nonlinear problems in applied mathematics*. SIAM, Philadelphia, PA, 1996, pp. 57–65.
- <span id="page-22-17"></span>[BF90] F. Bernis and A. Friedman. "Higher order nonlinear degenerate parabolic equations". In: *J. Differential Equations* 83.1 (1990), pp. 179–206.
- <span id="page-22-13"></span>[CD18] M. E. Cejas and R. G. Durán. "Weighted a priori estimates for elliptic equations". In: *Studia Math.* 243.1 (2018), pp. 13–24.
- [CH78] R. G. Casten and C. J. Holland. "Instability results for reaction diffusion equations with Neumann boundary conditions". In: *J. Differential Equations* 27.2 (1978), pp. 266–273.
- <span id="page-22-12"></span><span id="page-22-8"></span>[DG99] R. Dal Passo and H. Garcke. "Solutions of a fourth order degenerate parabolic equation with weak initial trace". In: *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* 28.1 (1999), pp. 153–181.
- [DGG01] R. Dal Passo, L. Giacomelli, and G. Grün. "A waiting time phenomenon for thin film equations". In: *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* 30.2 (2001), pp. 437–463.
- <span id="page-22-10"></span>[DGG98] R. Dal Passo, H. Garcke, and G. Grün. "On a fourth-order degenerate parabolic equation: global entropy estimates, existence, and qualitative behavior of solutions". In: *SIAM J. Math. Anal.* 29.2 (1998), pp. 321–342.
- <span id="page-22-19"></span><span id="page-22-18"></span>[DST08] R. G. Durán, M. Sanmartino, and M. Toschi. "Weighted a priori estimates for the Poisson equation". In: *Indiana Univ. Math. J.* 57.7 (2008), pp. 3463–3478.
- [DST10] R. G. Durán, M. Sanmartino, and M. Toschi. "Weighted a priori estimates for solution of (−Δ)*𝑚𝑢*= *𝑓* with homogeneous Dirichlet conditions". In: *Anal. Theory Appl.* 26.4 (2010), pp. 339–349.
- <span id="page-22-16"></span><span id="page-22-1"></span>[Gre78] H. P. Greenspan. "On the motion of a small viscous droplet that wets a surface". In: *Journal of Fluid Mechanics* 84.1 (1978), pp. 125–143.
- <span id="page-22-0"></span>[Gri11] P. Grisvard. *Elliptic Problems in Nonsmooth Domains*. Society for Industrial and Applied Mathematics, 2011.
- [Grü01] G. Grün. "On Bernis' Interpolation Inequalities in Multiple Space Dimensions". In: *Zeitschrift für Analysis und ihre Anwendungen* 20.4 (2001), pp. 987–998.
- <span id="page-22-3"></span>[Grü05] G. Grün. "Droplet Spreading Under Weak Slippage—Existence for the Cauchy Problem". In: *Communications in Partial Differential Equations* 29.11-12 (2005), pp. 1697–1744.
- <span id="page-22-15"></span>[GT01] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Reprint of the 1998 edition. Springer-Verlag, Berlin, 2001, pp. xiv+517.
- <span id="page-22-7"></span>[HS98] J. Hulshof and A. E. Shishkov. "The thin film equation with  $2 \le n < 3$ : finite speed of propagation in terms of the  $L^1$ -norm". In: *Adv. Differential Equations* 3.5 (1998), pp. 625–642.
- <span id="page-22-14"></span>[SZ98] P. Sternberg and K. Zumbrun. "Connectivity of phase boundaries in strictly convex domains". In: *Arch. Rational Mech. Anal.* 141.4 (1998), pp. 375–400.

### 24 REFERENCES

*Christina Lienstromberg:* INSTITUTE OF ANALYSIS, DYNAMICS AND MODELING, UNIVERSITY OF STUTTGART, PFAFFEN-WALDRING 57, 70569 STUTTGART, GERMANY

*Email address*: christina.lienstromberg@mathematik.uni-stuttgart.de

*Katerina Nik:* KING ABDULLAH UNIVERSITY OF SCIENCE AND TECHNOLOGY (KAUST), CEMSE DIVISION, THUWAL 23955- 6900, SAUDI ARABIA

*Email address*: katerina.nik@kaust.edu.sa