# Asymptotic Properties of the Maximum Likelihood Estimator for Markov-switching Observation-driven Models

Frederik Krabbe<sup>(a)</sup>\*

(a) Aarhus University, Denmark

#### Abstract

A Markov-switching observation-driven model is a stochastic process  $((S_t, Y_t))_{t \in \mathbb{Z}}$  where (i)  $(S_t)_{t \in \mathbb{Z}}$  is an unobserved Markov process taking values in a finite set and (ii)  $(Y_t)_{t \in \mathbb{Z}}$  is an observed process such that the conditional distribution of  $Y_t$  given all past Y's and the current and all past S's depends only on all past S's and  $S_t$ . In this paper, we prove the consistency and asymptotic normality of the maximum likelihood estimator for such model. As a special case hereof, we give conditions under which the maximum likelihood estimator for the widely applied Markov-switching generalised autoregressive conditional heteroscedasticity model introduced by Haas et al. (2004b) is consistent and asymptotic normal.

### 1 Introduction

A state space model is a stochastic process  $((X_t, Y_t))_{t \in \mathbb{Z}}$  where (i)  $(X_t)_{t \in \mathbb{Z}}$  is an unobserved Markov process taking values in X and (ii)  $(Y_t)_{t \in \mathbb{Z}}$  is an observed process taking values in Y such that the conditional distribution of  $Y_t$  given  $\mathbf{Y}_{-\infty}^{t-1}$  where  $\mathbf{Y}_i^j := (Y_i, ..., Y_j)$  and  $\mathbf{X}_{-\infty}^t$  where  $\mathbf{X}_i^j := (X_i, ..., X_j)$  depends only on  $X_t$ . State space models and hidden Markov models, which are special cases of state space models in which  $X_t = S_t$  where  $(S_t)_{t \in \mathbb{Z}}$  is a Markov chain on a finite set have been applied in numerous areas. Applications of the former are found in economics (Stock and Watson (1989), Harvey and Chung (2000), and Bräuning and Koopman (2020)), finance (Harvey and Shephard (1996), Jacquier et al. (2004), Yu (2005), and Catania (2022)), and climatology (Bennedsen et al. (2023)) whereas applications of the latter are found in finance (Rydén et al. (1998)), biology (Churchill (1989)), ecology (Langrock et al. (2012)), meteorology (Zucchini and Guttorp (1991) and Bulla et al. (2012)), and quantum mechanics (Gammelmark et al. (2014)).

Statistical inference for state space models is therefore of significant practical importance. Leroux (1992) and Bickel et al. (1998) proved consistency and asymptotic normality of the maximum likelihood (ML) estimator for hidden Markov models, respectively. Later, Jensen and Petersen (1999) proved (local) consistency and asymptotic normality for state space

<sup>\*</sup>I would like to thank Leopoldo Catania, Morten Ørregaard Nielsen, Christian Francq, Jean-Michel Zakoïan, and Christian Gourieroux for useful discussions. This research is supported by the Danish National Research Foundation (DNRF Chair grant number DNRF154).

models where X is compact, and Douc et al. (2011) proved (global) consistency for general state space models.

Several extensions of state space models have been proposed especially in economics and finance. The arguably most famous extension is the autoregressive state space model of order  $p \in \mathbb{N}$  in which the conditional distribution of  $Y_t$  given  $\mathbf{Y}_{-\infty}^{t-1}$  and  $\mathbf{X}_{-\infty}^t$  depends on both  $\mathbf{Y}_{t-p}^{t-1}$  and  $X_t$ . One such example is the Markov-switching autoregressive model introduced by Hamilton (1989) to model economic growth in which X is finite. Another one in which X is also finite is the Markov-switching autoregressive conditional heteroscedasticity (ARCH) model introduced by Cai (1994) and Hamilton and Susmel (1994) to model financial returns which is a generalisation of the ARCH model introduced by Engle (1982). See, for instance, Hamilton (2010) and Ang and Timmermann (2012) for more examples in economics and finance, respectively.

Statistical inference for autoregressive state space models has also attracted much attention in the literature. Consistency of the ML estimator for autoregressive state space models where X is finite was proved by Francq and Roussignol (1998) and Krishnamurthy and Ryden (1998). This was later generalised by Douc et al. (2004) who proved consistency and asymptotic normality for autoregressive state space models where X is compact and not necessarily finite. More recently, Kasahara and Shimotsu (2019) relax some of the assumptions in Douc et al. (2004).

Another extension which has attracted much attention in the literature in more recent years is the observation-driven state space model in which the conditional distribution of  $Y_t$  given  $\mathbf{Y}_{-\infty}^{t-1}$  and  $\mathbf{X}_{-\infty}^t$  now depends on both  $\mathbf{Y}_{-\infty}^{t-1}$  and  $X_t$ . Two such examples are the Markov-switching generalised ARCH (GARCH) model introduced by Haas et al. (2004b) in which X is finite, which is a generalisation of the GARCH model introduced by Bollerslev (1986), and the score-driven state space model introduced by Monache et al. (2021) in which X is not finite. Other applications of such models are found in finance (Haas et al. (2004a), Broda et al. (2013), Ardia et al. (2018), and Bernardi and Catania (2019)) and meteorology (Harvey and Palumbo (2023)) in which X is finite as well as in economics (Monache et al. (2016) and Angelini and Gorgi (2018)) and finance (Buccheri et al. (2021) and Buccheri and Corsi (2021)) in which X is not finite. In contrast to state space models and autoregressive state space models, statistical inference for observation-driven state space models is to a large extent undiscovered land to the best of our knowledge. Only recently, Kandji and Misko (2024) gave conditions under which the ML estimator for the Markov-switching GARCH model by Haas et al. (2004b) is consistent.

In this paper, we prove consistency (Theorem 2) and asymptotic normality (Theorem 3) of the ML estimator for an observation-driven state space model where X is finite which we call a Markov-switching observation-driven model. The fact that the filter forgets its initialisation asymptotically (Lemma 2) is crucial in the proofs as in Douc et al. (2004) (Corollary 1) and Kasahara and Shimotsu (2019) (Lemma 1). However, in contrast to Douc et al. (2004) and Kasahara and Shimotsu (2019) who use theory for Markov chains, we prove this using theory for stochastic difference equations which is most commonly used for standard observation-

<sup>&</sup>lt;sup>1</sup>Note that there exist two types of MS-GARCH models namely the one by Gray (1996) in which the conditional distribution of  $Y_t$  given  $\mathbf{Y}_{-\infty}^{t-1}$  and  $\mathbf{X}_{-\infty}^t$  depends on both  $\mathbf{Y}_{-\infty}^{t-1}$  and  $\mathbf{X}_{-\infty}^t$  so ML estimation is infeasible and the one by Haas et al. (2004b) just discussed in which ML estimation is feasible. See Francq and Zakoian (2019) for more details.

driven models, see, for instance, Straumann and Mikosch (2006), Blasques et al. (2018), and Blasques et al. (2022). As a special case of the general theory, we give conditions under which the ML estimator for the widely applied Markov-switching GARCH model by Haas et al. (2004b) is both consistent (Theorem 5) and asymptotic normal (Theorem 7) thus extending the work by Kandji and Misko (2024).

The rest of the paper is organised as follows. Section 2 introduces the Markov-switching observation-driven model, and Section 3 gives some examples of Markov-switching observation-driven models. In Section 4, stationarity and ergodicity of the model is considered. Consistency and asymptotic normality of the ML estimator is then considered in Section 5. Section 6 gives conditions under which the ML estimator for the Markov-switching GARCH model by Haas et al. (2004b) is consistent and asymptotic normal. An application to real data is presented in Section ??. Section 7 concludes. All proofs are collected in the appendix.

First, a word on notation:  $||\mathbf{x}||_p := (\sum_{i=1}^n |x_i|^p)^{1/p}, \mathbf{x} \in \mathbb{R}^n$  denotes the *p*-norm of  $\mathbf{x}$  and  $||\mathbf{X}||_{p,p} := (\sum_{i=1}^n \sum_{j=1}^m |x_{ij}|^p)^{1/p}, \mathbf{X} \in \mathbb{R}^{n \times m}$  denotes the *p*-norm of  $\mathbf{X}$ . Moreover, for a function  $\mathbf{x} \mapsto \mathbf{f}(\mathbf{x}) : \mathbb{R}^n \to \mathbb{R}^k, \ \nabla_{\mathbf{x}'}\mathbf{f}(\mathbf{x}) := \left(\frac{\partial \mathbf{f}(\mathbf{x})}{\partial x_1}, ..., \frac{\partial \mathbf{f}(\mathbf{x})}{\partial x_n}\right)$  denotes the first order derivative of  $\mathbf{f}$  with respect to  $\mathbf{x}$ ,  $\nabla_{\mathbf{x}}\mathbf{f}(\mathbf{x}) := (\nabla_{\mathbf{x}'}\mathbf{f}(\mathbf{x}))'$ , and if k = 1, then  $\nabla_{\mathbf{x}\mathbf{x}}f(\mathbf{x}) := \nabla_{\mathbf{x}'}\nabla_{\mathbf{x}}f(\mathbf{x})$  denotes the second order derivative of  $\mathbf{f}$  with respect to  $\mathbf{x}$ .

### 2 The Markov-switching Observation-driven Model

A Markov-switching observation-driven model is a stochastic process  $((S_t, Y_t))_{t \in \mathbb{Z}}$  where (i)  $(S_t)_{t \in \mathbb{Z}}$  is an unobserved first-order homogenous Markov chain taking values in  $\{1, ..., J\}$  with transition probabilities

$$p_{ij} := \mathbb{P}(S_{t+1} = j \mid S_t = i), \quad i, j \in \{1, ..., J\},$$

and transition probability matrix

$$\mathbf{P} := \begin{bmatrix} p_{11} & \cdots & p_{1J} \\ \vdots & \ddots & \vdots \\ p_{J1} & \cdots & p_{JJ} \end{bmatrix},$$

and (ii)  $(Y_t)_{t\in\mathbb{Z}}$  is an observed stochastic process taking values in  $\mathcal{Y}\subseteq\mathbb{R}$  such that the conditional distribution of  $Y_t$  given  $\mathbf{Y}_{-\infty}^{t-1}$  and  $\mathbf{S}_{-\infty}^t$  depends only on  $\mathbf{Y}_{-\infty}^{t-1}$  and  $S_t$  as follows

$$Y_t \mid (\mathbf{Y}_{-\infty}^{t-1}, S_t) \sim \mathcal{D}_{S_t}(X_{S_t,t}, \boldsymbol{v}_{S_t}),$$

where  $X_{j,t}, j \in \{1, ..., J\}$  is a time-varying parameter taking values in a complete set  $\mathcal{X}_j \subseteq \mathbb{R}$  given by

$$X_{j,t+1} = \phi_j(Y_t, ..., Y_{t-p+1}, X_{j,t}, ..., X_{j,t-q+1}; \boldsymbol{v}_j),$$

and  $v_j, j \in \{1, ..., J\}$  is a vector of constant parameters taking values in a set  $\Upsilon_j \subseteq \mathbb{R}^{d_j}$ . The Markov-switching observation-driven model is called a mixture observation-driven model if  $(S_t)_{t \in \mathbb{Z}}$  is an independent and identically distributed (i.i.d.) chain, that is, if

$$p_{1j} = \cdots = p_{Jj}$$

for all  $j \in \{1, ..., J\}$ .

The unobserved Markov chain  $(S_t)_{t\in\mathbb{Z}}$  is estimated via the conditional distribution

$$\pi_{j,t|s} := \mathbb{P}(S_t = j \mid \mathbf{Y}_{-\infty}^s), \quad j \in \{1, ..., J\},$$

which is called the filtering distribution when t = s, the smoothing distribution when t < s, and the predictive distribution when t > s. The one-step-ahead prediction is given by

$$\pi_{j,t+1|t} = \sum_{i=1}^{J} p_{ij} \pi_{i,t|t},$$

and the filter is given by

$$\pi_{j,t|t} = \frac{\pi_{j,t|t-1} f_j(Y_t; X_{j,t}, v_j)}{f(Y_t)},$$

where  $f(y), y \in \mathcal{Y}$  is the conditional probability density function (pdf) of  $Y_t$  given  $\mathbf{Y}_{-\infty}^{t-1}$  given by

$$f(y) = \sum_{k=1}^{J} \pi_{k,t|t-1} f_k(y; X_{k,t}, \boldsymbol{v}_k),$$

and  $f_j(y; X_{j,t}, v_j), y \in \mathcal{Y}$  is the conditional pdf of  $Y_t$  given  $\mathbf{Y}_{-\infty}^{t-1}$  and  $S_t = j$ , see Hamilton (1994) for details.<sup>2</sup> Define  $\boldsymbol{\pi}_{t|s} := (\pi_{1,t|s}, ..., \pi_{J,t|s})'$ . Then, the one-step-ahead prediction is given by

$$\pi_{t+1|t} = \mathbf{P}' \pi_{t|t},$$

and the filter is given by

$$\boldsymbol{\pi}_{t|t} = \mathbf{F}_t(\boldsymbol{\pi}_{t|t-1})\boldsymbol{\pi}_{t|t-1},$$

where  $\mathbf{F}_{t}(\boldsymbol{\pi}_{t|t-1})$  is a diagonal matrix with generic element

$$[\mathbf{F}_{t}(\boldsymbol{\pi}_{t|t-1})]_{ii} = \frac{f_{i}(Y_{t}; X_{i,t}, \boldsymbol{v}_{i})}{\sum_{k=1}^{J} \pi_{k|t|t-1} f_{k}(Y_{t}; X_{k|t}, \boldsymbol{v}_{k})}, \quad i \in \{1, ..., J\}.$$

Moreover, the smoother is given by

$$\pi_{j,t|T} = \pi_{j,t|t} \sum_{i=1}^{J} p_{ji} \frac{\pi_{i,t+1|T}}{\pi_{i,t+1|t}}, \quad t < T,$$

see Hamilton (1994) for details once again. We also refer to Hamilton (1994) for details on prediction of the observed stochastic process  $(Y_t)_{t\in\mathbb{Z}}$ .

Finally, note that the Markov-switching observation-driven model reduces to the Markov-switching autoregressive model of order  $p \in \mathbb{N}$  if  $\phi_j(Y_t, ..., Y_{t-p+1}, X_{j,t}, ..., X_{j,t-q+1}; \boldsymbol{v}_j) = \phi_j(Y_t, ..., Y_{t-p+1}; \boldsymbol{v}_j)$  for all  $j \in \{1, ..., J\}$  and to the hidden Markov model if  $X_{j,t} \equiv X_j$  for all  $j \in \{1, ..., J\}$ .

In the following, we restrict our attention to Markov-switching observation-driven models where p = 1 and q = 1. All results can straightforward be generalised by using the same techniques.

$$\pi_{j,t+h|t} = \sum_{i=1}^{J} p_{ij}^{(h)} \pi_{i,t|t},$$

where  $p_{ij}^{(h)} := \mathbb{P}(S_{t+h} = j \mid S_t = i)$  is the (i, j)'th element of  $\mathbf{P}^h := \prod_{i=1}^h \mathbf{P}$ .

 $<sup>^{2}</sup>$ More generally, the h-step-ahead prediction is given by

### 3 Examples of Markov-switching Observation-driven Models

In this section, we give some examples of Markov-switching observation-driven models.

**Example 1.** An example of a Markov-switching observation-driven model is

$$Y_t = X_{S_t,t} + \sigma_{S_t} \varepsilon_t, \tag{1}$$

where  $(\varepsilon_t)_{t\in\mathbb{Z}}$  is a sequence of independent standard normal distributed random variables independent of  $(S_t)_{t\in\mathbb{Z}}$  and, for each  $j \in \{1, ..., J\}$ ,

$$X_{i,t+1} = \omega_i + \alpha_i Y_t + \beta_i X_{i,t},$$

where  $\omega_j \in \mathbb{R}$ ,  $\alpha_j \in \mathbb{R}$ ,  $\beta_j \in \mathbb{R}$ , and  $\sigma_j^2 > 0$ . Here,  $\mathcal{Y} = \mathbb{R}$ ,  $\mathcal{D}_{S_t}$  is the normal distribution with mean  $X_{S_t,t}$  and variance  $\sigma_{S_t}^2$ ,  $\mathcal{X}_j = \mathbb{R}$ ,  $\phi_j(y,x_j;\boldsymbol{v}_j) = [\boldsymbol{v}_j]_1 + [\boldsymbol{v}_j]_2 y + [\boldsymbol{v}_j]_3 x_j$ ,  $\boldsymbol{v}_j = (\omega_j,\alpha_j,\beta_j,\sigma_j^2)'$ , and  $\Upsilon_j = \mathbb{R} \times \mathbb{R} \times \mathbb{R} \times (0,\infty)$ .

A related model is the Markov-switching autoregressive model of order  $p \in \mathbb{N}$  by Hamilton (1989). This model is given by

$$Y_t = a_{S_t} + \sum_{i=1}^{p} b_{S_t}^{(i)} Y_{t-i} + \sigma_{S_t} \varepsilon_t,$$

where  $(\varepsilon_t)_{t\in\mathbb{Z}}$  is a sequence of independent standard normal distributed random variables independent of  $(S_t)_{t\in\mathbb{Z}}$  as above.

It can be shown that if  $(Y_t)_{t\in\mathbb{Z}}$  is stationary and ergodic with  $\mathbb{E}[\log^+|Y_t|] < \infty$  where  $\log^+ x = \max(\log x, 0)$  for all x > 0 and  $|\beta_j| < 1$  for all  $j \in \{1, ..., J\}$ , then

$$X_{j,t} = \frac{\omega_j}{1 - \beta_j} + \sum_{i=1}^{\infty} \alpha_j \beta_j^{i-1} Y_{t-i} \quad a.s.$$

for all  $j \in \{1, ..., J\}$ . The Markov-switching observation-driven model in Equation (1) can thus be thought of as a Markov-switching autoregressive model of order infinity given by

$$Y_t = a_{S_t} + \sum_{i=1}^{\infty} b_{S_t}^{(i)} Y_{t-i} + \sigma_{S_t} \varepsilon_t,$$

where

$$a_{S_t} = \frac{\omega_{S_t}}{1 - \beta_{S_t}}$$
 and  $b_{S_t}^{(i)} = \alpha_{S_t} \beta_{S_t}^{i-1}$ .

**Example 2.** The Markov-switching GARCH model by Haas et al. (2004b) is given by

$$Y_t = \sqrt{X_{S_t,t}} \varepsilon_t,$$

where  $(\varepsilon_t)_{t\in\mathbb{Z}}$  is a sequence of independent standard normal distributed random variables independent of  $(S_t)_{t\in\mathbb{Z}}$  and, for each  $j\in\{1,...,J\}$ ,

$$X_{j,t+1} = \omega_j + \alpha_j Y_t^2 + \beta_j X_{j,t},$$

where  $\omega_j \geq 0$ ,  $\alpha_j \geq 0$ , and  $\beta_j \geq 0$ . This is also an example of a Markov-switching observationdriven model where  $\mathcal{Y} = \mathbb{R}$ ,  $\mathcal{D}_{S_t}$  is the normal distribution with mean zero and variance  $X_{S_t,t}$ ,  $\mathcal{X}_{j} = [0, \infty), \ \phi_{j}(y, x_{j}; \mathbf{v}_{j}) = [\mathbf{v}_{j}]_{1} + [\mathbf{v}_{j}]_{2}y^{2} + [\mathbf{v}_{j}]_{3}x_{j}, \ \mathbf{v}_{j} = (\omega_{j}, \alpha_{j}, \beta_{j})', \ and \ \Upsilon_{j} = [0, \infty) \times [0, \infty).$ 

Note that the Markov-switching GARCH model reduces to the mixture GARCH model by Haas et al. (2004a) if  $(S_t)_{t\in\mathbb{Z}}$  is an i.i.d. chain.

**Example 3.** Let  $y \mapsto F(y; x, \tilde{v})$  be a cumulative distribution function (cdf) with support  $\mathcal{N} \subseteq [0, \infty)$  indexed by the mean x and a vector of parameters  $\tilde{v}$  such that, for all  $u \in (0, 1)$ ,

$$x \le x^* \quad \Rightarrow \quad F^-(u; x, \tilde{\boldsymbol{v}}) \le F^-(u; x^*, \tilde{\boldsymbol{v}}),$$

where  $F^{-}(u; x, \tilde{\boldsymbol{v}}) = \inf\{y \in \mathcal{N} : F(y; x, \tilde{\boldsymbol{v}}) \ge u\}.$ 

The (present-regime dependent) Markov-switching positive linear conditional mean model by Aknouche and Francq (2022) is given by

$$Y_t = F_{S_t}^-(U_t; X_{S_t,t}, \tilde{\boldsymbol{v}}_{S_t}),$$

where  $(U_t)_{t\in\mathbb{Z}}$  is a sequence of independent uniform distributed random variables on [0, 1] independent of  $(S_t)_{t\in\mathbb{Z}}$  and, for each  $j \in \{1, ..., J\}$ ,

$$X_{i,t+1} = \omega_i + \alpha_i Y_t + \beta_i X_{i,t},$$

where  $\omega_j \geq 0$ ,  $\alpha_j \geq 0$ ,  $\beta_j \geq 0$ , and  $\tilde{\boldsymbol{v}}_j \in \tilde{\boldsymbol{\Upsilon}}_j$ . This is another example of a Markov-switching observation-driven model where  $\mathcal{Y} = \mathcal{N}$ ,  $F_{S_t}$  is the cdf of  $\mathcal{D}_{S_t}$ ,  $\mathcal{X}_j = [0, \infty)$ ,  $\phi_j(y, x_j; \boldsymbol{v}_j) = [\boldsymbol{v}_j]_1 + [\boldsymbol{v}_j]_2 y + [\boldsymbol{v}_j]_3 x_j$ ,  $\boldsymbol{v}_j = (\omega_j, \alpha_j, \beta_j, \tilde{\boldsymbol{v}}_j)'$ , and  $\boldsymbol{\Upsilon}_j = [0, \infty) \times [0, \infty) \times \tilde{\boldsymbol{\Upsilon}}_j$ .

## 4 Probabilistic Properties of the Model

First, we study the probabilistic properties of the Markov-switching observation-driven model. We restrict our attention to the Markov-switching observation-driven models that can be written as

$$Y_t = \mathbf{1}_{S_t} \mathbf{g}(\boldsymbol{\varepsilon}_t; \mathbf{X}_t, \boldsymbol{v}), \tag{2}$$

with  $\mathbf{1}_{S_t} := (1_{\{S_t = 1\}}, ..., 1_{\{S_t = J\}})$  where  $(S_t)_{t \in \mathbb{Z}}$  is stationary, irreducible, and aperiodic (thus ergodic), and  $\mathbf{g}(\boldsymbol{\varepsilon}_t; \mathbf{X}_t, \boldsymbol{v}) := (g_1(\varepsilon_{1,t}; X_{1,t}, \boldsymbol{v}_1), ..., g_J(\varepsilon_{J,t}; X_{J,t}, \boldsymbol{v}_J))'$  where  $\boldsymbol{\varepsilon}_t := (\varepsilon_{1,t}, ..., \varepsilon_{J,t})'$ ,  $(\varepsilon_{j,t})_{t \in \mathbb{Z}}$  is a sequence of i.i.d. random variables taking values in  $\mathcal{E}_j$  with distribution  $\mathcal{D}_j^{\varepsilon}(\boldsymbol{v}_j)$  for all  $j \in \{1, ..., J\}$ , and  $(\varepsilon_{i,t})_{t \in \mathbb{Z}}$  and  $(\varepsilon_{j,t})_{t \in \mathbb{Z}}$  are independent for all  $i, j \in \{1, ..., J\}$  such that  $i \neq j$ . Moreover,  $\mathbf{X}_t := (X_{1,t}, ..., X_{J,t})'$  is given by

$$\mathbf{X}_{t+1} = \boldsymbol{\phi}(S_t, \boldsymbol{\varepsilon}_t, \mathbf{X}_t; \boldsymbol{v})$$

with

$$[\phi(S_t, \varepsilon_t, \mathbf{X}_t; \boldsymbol{v})]_j = \phi_j(\mathbf{1}_{S_t} \mathbf{g}(\varepsilon_t; \mathbf{X}_t, \boldsymbol{v}), X_{j,t}; \boldsymbol{v}_j), \quad j \in \{1, ..., J\},$$

where  $\mathbf{x} \mapsto \phi(S_t, \boldsymbol{\varepsilon}_t, \mathbf{x}; \boldsymbol{v})$  is stationary, ergodic, and Lipschitz, and  $\boldsymbol{v} := (\boldsymbol{v}_1, ..., \boldsymbol{v}_J)'$ . Finally,  $(\varepsilon_{j,t})_{t \in \mathbb{Z}}$  and  $(S_t)_{t \in \mathbb{Z}}$  are independent for all  $j \in \{1, ..., J\}$ .

<sup>&</sup>lt;sup>3</sup>All examples in Section 3 can be written like this because if  $\varepsilon_{i,t} \stackrel{d}{=} \varepsilon_{j,t}$  for all  $i, j \in \{1, ..., J\}$ , then let  $\varepsilon_{j,t} = \varepsilon_t$  for all  $j \in \{1, ..., J\}$  where  $(\varepsilon_t)_{t \in \mathbb{Z}}$  is a sequence of i.i.d. random variables taking values in  $\mathcal{E}$  with distribution  $\mathcal{D}^{\varepsilon}(\boldsymbol{v})$ .

### 4.1 Stationarity and Ergodicity

Theorem 1, which follows from an application of Theorem 3.1 in Bougerol (1993) (see also Theorem 2.8 in Straumann and Mikosch (2006)), gives conditions under which the model is stationary and ergodic. Let

$$\Lambda(\boldsymbol{\phi}_t) := \sup_{\substack{\mathbf{x}, \mathbf{y} \in \mathcal{X}_1 \times \cdots \times \mathcal{X}_J \\ \mathbf{x} \neq \mathbf{y}}} \frac{||\boldsymbol{\phi}_t(\mathbf{x}) - \boldsymbol{\phi}_t(\mathbf{y})||_2}{||\mathbf{x} - \mathbf{y}||_2},$$

where  $\phi_t(\mathbf{x}) := \phi(S_t, \varepsilon_t, \mathbf{x}; \boldsymbol{v}).$ 

Theorem 1. Assume that

- (i) there exists an  $\mathbf{x} \in \mathcal{X}_1 \times \cdots \times \mathcal{X}_J$  such that  $\mathbb{E}[\log^+ ||\phi_t(\mathbf{x}) \mathbf{x}||_2] < \infty$ ,
- (ii)  $\mathbb{E}[\log^+ \Lambda(\phi_t)] < \infty$ , and
- (iii) there exists an  $r \in \mathbb{N}$  such that

$$\mathbb{E}\left[\log\Lambda\left(\boldsymbol{\phi}_{t}^{(r)}\right)\right]<0,$$

where 
$$\phi_t^{(r)}(\mathbf{x}) := \phi_t \circ \cdots \circ \phi_{t-r+1}(\mathbf{x})$$
.

Then,  $(Y_t)_{t\in\mathbb{Z}}$  is stationary and ergodic.

### 5 Asymptotic Properties of the Maximum Likelihood Estimator

We now study the asymptotic properties of the maximum likelihood estimator for the Markov-switching observation-driven model discussed in the previous section.

First, the parameter vector is given by

$$\boldsymbol{\theta} := (p_{ij}, i = 1, ..., J, j = 1, ..., J - 1, \boldsymbol{v}_j, j = 1, ..., J)'$$

and takes values in

$$oldsymbol{\Theta} \subset \left\{oldsymbol{ heta}: p_{ij} > 0, i = 1, ..., J, j = 1, ..., J - 1, \sum_{j=1}^{J} p_{ij} = 1, i = 1, ..., J, oldsymbol{v}_j \in oldsymbol{\Upsilon}_j, j = 1, ..., J
ight\} \subset \mathbb{R}^d,$$

where  $d := J(J-1) + \sum_{j=1}^{J} d_j$ . Assume that a realisation,  $(y_t)_{t=1}^T$ , of the Markov-switching observation-driven model,  $(Y_t)_{t \in \mathbb{Z}}$ , satisfying Equation (2) for  $\boldsymbol{\theta} = \boldsymbol{\theta}_0$  is observed. Then, the average log-likelihood function is given by

$$\hat{L}_T(\boldsymbol{\theta}) = \frac{1}{T} \sum_{t=1}^{T} \log \hat{f}(y_t; \boldsymbol{\theta}),$$

where

$$\hat{f}(y_t; \boldsymbol{\theta}) = \sum_{j=1}^{J} \hat{\pi}_{j,t|t-1}(\boldsymbol{\theta}) f_j(y_t; \hat{X}_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j).$$

In the average log-likelihood function, for each  $j \in \{1, ..., J\}$ ,  $X_{j,t}(v_j)$  is given by

$$\hat{X}_{j,t+1}(\boldsymbol{v}_j) = \phi_j(y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)$$

for some initialisation  $\hat{X}_{i,1}(v_i) \in \mathcal{X}_i$ , and  $\hat{\pi}_{t|t-1}(\theta)$  is given by

$$\hat{\boldsymbol{\pi}}_{t+1|t}(\boldsymbol{\theta}) = \mathbf{P}'\hat{\boldsymbol{\pi}}_{t|t}(\boldsymbol{\theta})$$

with

$$\hat{oldsymbol{\pi}}_{t|t}(oldsymbol{ heta}) = \hat{f F}_t(\hat{oldsymbol{\pi}}_{t|t-1}(oldsymbol{ heta});oldsymbol{ heta})\hat{oldsymbol{\pi}}_{t|t-1}(oldsymbol{ heta})$$

for some initialisation  $\hat{\pi}_{0|0}(\theta) \in \mathcal{S}$  with  $\mathcal{S} := \{\mathbf{x} \in \mathbb{R}^J : x_j \geq 0, j = 1, ..., J, \sum_{j=1}^J x_j = 1\}$  where

$$[\hat{\mathbf{F}}_t(\hat{\boldsymbol{\pi}}_{t|t-1}(\boldsymbol{\theta});\boldsymbol{\theta})]_{ii} = \frac{f_i(y_t; \hat{X}_{i,t}(\boldsymbol{v}_i), \boldsymbol{v}_i)}{\sum_{k=1}^J \hat{\pi}_{k,t|t-1}(\boldsymbol{\theta}) f_k(y_t; \hat{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k)}, \quad i \in \{1, ..., J\}.$$

The maximum likelihood estimator is then given by

$$\hat{\boldsymbol{\theta}}_T = \operatorname*{arg\,max} \hat{L}_T(\boldsymbol{\theta}).$$

#### 5.1 Consistency

We assume the following.

**Assumption 1.** The conditions in Theorem 1 hold for  $\theta = \theta_0$ .

Assumption 2.  $\Theta$  is compact.

**Assumption 3.** For each  $j \in \{1, ..., J\}$ ,

- (i)  $(x_i, \mathbf{v}_i) \mapsto f_i(y; x_i, \mathbf{v}_i)$  is continuous for all  $y \in \mathcal{Y}$  and
- (ii)  $x_i \mapsto f_i(y; x_i, \mathbf{v}_i)$  is differentiable for all  $y \in \mathcal{Y}$  and  $\mathbf{v}_i \in \Upsilon_i$ .

**Assumption 4.** For each  $j \in \{1, ..., J\}$ ,

- (i)  $x_i \mapsto \phi_i(y, x_i; v_i)$  is Lipschitz for all  $y \in \mathcal{Y}$  and  $v_i \in \Upsilon_i$ ,
- (ii)  $(x_i, \mathbf{v}_i) \mapsto \phi_i(y, x_i; \mathbf{v}_i)$  is continuous for all  $y \in \mathcal{Y}$ , and
- (iii)  $x_j \mapsto \phi_j(y, x_j; \mathbf{v}_j)$  is differentiable for all  $y \in \mathcal{Y}$  and  $\mathbf{v}_j \in \Upsilon_j$ .

First, the behaviour of the initialised sequences must be studied in order to ensure an appropriate form of convergence of the log-likelihood function because the initialised sequences are non-stationary by construction.

The following lemma, which follows from an application of Theorem 3.1 in Bougerol (1993), gives conditions under which, for each  $j \in \{1, ..., J\}$ , the non-stationary sequence  $(\hat{X}_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{N}}$ converges uniformly exponentially fast almost surely (e.a.s.) to a unique stationary and ergodic sequence  $(X_{i,t}(\boldsymbol{v}_i))_{t\in\mathbb{Z}}$ . For each  $j\in\{1,...,J\}$ , let

$$\Lambda_{j,t}(oldsymbol{v}_j) := \sup_{x_j \in \mathcal{X}_j} \left| 
abla_{x_j} \phi_j(Y_t, x_j; oldsymbol{v}_j) \right|.$$

 $<sup>\</sup>Lambda_{j,t}(\boldsymbol{v}_j) := \sup_{x_j \in \mathcal{X}_j} \left| \nabla_{x_j} \phi_j(Y_t, x_j; \boldsymbol{v}_j) \right|.$ <sup>4</sup>A sequence  $(\mathbf{Z}_t)_{t \in \mathbb{Z}}$  of random matrices is said to converge to zero e.a.s. if there exists a  $\gamma > 1$  such that  $\gamma^t ||\mathbf{Z}_t||_{p,p} \stackrel{a.s.}{\to} 0 \quad \text{as} \quad t \to \infty.$ 

**Lemma 1.** Assume that Assumption 1, 2, and 4 hold. Moreover, assume that, for each  $j \in \{1,...,J\}$ ,

- (i) there exists an  $x_j \in \mathcal{X}_j$  such that  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\phi_j(Y_t, x_j; \boldsymbol{v}_j) x_j|] < \infty$ ,
- (ii)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_i \in \boldsymbol{\Upsilon}_i} \Lambda_{j,t}(\boldsymbol{v}_j)] < \infty$ , and
- (iii)  $\mathbb{E}[\log \sup_{\boldsymbol{v}_i \in \boldsymbol{\Upsilon}_i} \Lambda_{j,t}(\boldsymbol{v}_j)] < 0.$

Then, for each  $j \in \{1,...,J\}$ , the sequence  $(X_{j,t}(v_j))_{t \in \mathbb{Z}}$  given by

$$X_{j,t+1}(\boldsymbol{\upsilon}_j) = \phi_j(Y_t, X_{j,t}(\boldsymbol{\upsilon}_j); \boldsymbol{\upsilon}_j)$$

is stationary and ergodic for all  $v_j \in \Upsilon_j$  and

$$\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left| \hat{X}_{j,t}(\boldsymbol{v}_j) - X_{j,t}(\boldsymbol{v}_j) \right| \overset{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\hat{X}_{j,1}(\boldsymbol{v}_j) \in \mathcal{X}_j$ .

Theorem 3.1 in Bougerol (1993) cannot be used for  $(\hat{\pi}_{t|t-1}(\boldsymbol{\theta}))_{t\in\mathbb{N}}$  because  $(\hat{\pi}_{t|t-1}(\boldsymbol{\theta}))_{t\in\mathbb{N}}$  depends on  $(\hat{X}_{1,t}(\boldsymbol{v}_1))_{t\in\mathbb{N}},...,(\hat{X}_{J,t}(\boldsymbol{v}_J))_{t\in\mathbb{N}}$  which are non-stationary. The next lemma, which gives conditions under which the non-stationary sequence  $(\hat{\pi}_{t|t}(\boldsymbol{\theta}))_{t\in\mathbb{N}}$  converges uniformly e.a.s. to a unique stationary and ergodic sequence  $(\pi_{t|t}(\boldsymbol{\theta}))_{t\in\mathbb{Z}}$ , follows instead by using similar arguments as in the proof of Theorem 2.10 in Straumann and Mikosch (2006).

**Lemma 2.** Assume that Assumption 1-4 and the conditions in Lemma 1 hold. Moreover, assume that for each  $j \in \{1, ..., J\}$ , there exists an  $m_j > 0$  such that

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \nabla_{x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right|^{m_j} \right] < \infty.$$

Then, the sequence  $(\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}))_{t\in\mathbb{Z}}$  given by

$$oldsymbol{\pi}_{t|t}(oldsymbol{ heta}) = \mathbf{F}_t(\mathbf{P}'oldsymbol{\pi}_{t-1|t-1}(oldsymbol{ heta});oldsymbol{ heta})\mathbf{P}'oldsymbol{\pi}_{t-1|t-1}(oldsymbol{ heta})$$

is stationary and ergodic for all  $\theta \in \Theta$  and

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \hat{\boldsymbol{\pi}}_{t|t}(\boldsymbol{\theta}) - \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}) \right| \right|_{2} \stackrel{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\hat{\boldsymbol{\pi}}_{0|0}(\boldsymbol{\theta}) \in \mathcal{S}$ .

The next corollary is a direct consequence hereof.

Corollary 1. Under the assumptions in Lemma 2, the sequence  $(\pi_{t|t-1}(\theta))_{t\in\mathbb{Z}}$  given by

$$oldsymbol{\pi}_{t+1|t}(oldsymbol{ heta}) = \mathbf{P}' oldsymbol{\pi}_{t|t}(oldsymbol{ heta})$$

is stationary and ergodic for all  $\theta \in \Theta$  and

$$\sup_{\boldsymbol{\theta} \in \Theta} \left| \left| \hat{\boldsymbol{\pi}}_{t|t-1}(\boldsymbol{\theta}) - \boldsymbol{\pi}_{t|t-1}(\boldsymbol{\theta}) \right| \right|_2 \stackrel{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\hat{\boldsymbol{\pi}}_{0|0}(\boldsymbol{\theta}) \in \mathcal{S}$ .

**Remark 1.** Note that  $\pi_{t|t-1}(\boldsymbol{\theta}) \in \mathcal{S}_{\boldsymbol{\theta}}$  for all  $t \in \mathbb{Z}$  where  $\mathcal{S}_{\boldsymbol{\theta}} := \{\mathbf{x} \in \mathbb{R}^J : x_j \geq \min_{i \in \{1,...,J\}} p_{ij}, j = 1,...,J, \sum_{j=1}^J x_j = 1\}.$ 

The result in Lemma 2 is not surprising: the Markov chain itself forgets its initialisation asymptotically (in case it is initialised) as  $p_{ij} > 0$  for all  $i, j \in \{1, ..., J\}$ , so it is not surprising that the filter also forgets its initialisation asymptotically provided that the time-varying parameters do the same.

Moreover, we assume the following.

**Assumption 5.** For each  $j \in \{1, ..., J\}$ ,

$$\mathbb{E}\left[\sup_{oldsymbol{v}_j\inoldsymbol{\Upsilon}_j}|\log f_j(Y_t;X_{j,t}(oldsymbol{v}_j),oldsymbol{v}_j)|
ight]<\infty.$$

Finally, we assume the following as in Francq and Roussignol (1998) where  $f^{(m)}(\mathbf{y}; \boldsymbol{\theta}), \mathbf{y} \in \mathcal{Y}^m$  denotes the conditional pdf of  $\mathbf{Y}_{t-m+1}^t$  given  $\mathbf{Y}_{-\infty}^{t-m}$ .

**Assumption 6.** There exists an  $m \in \mathbb{N}$  such that

$$f^{(m)}(Y_t,...,Y_{t-m+1};\boldsymbol{\theta}) = f^{(m)}(Y_t,...,Y_{t-m+1};\boldsymbol{\theta}_0)$$
 a.s.

implies that

$$\theta = \theta_0$$
.

Theorem 2 gives conditions under which the ML estimator is consistent.

**Theorem 2.** Assume that Assumption 1-6 and the conditions in Lemma 1 and 2 hold. Then,

$$\hat{\boldsymbol{\theta}}_T \stackrel{a.s.}{\rightarrow} \boldsymbol{\theta}_0 \quad as \quad T \rightarrow \infty.$$

#### 5.2 Asymptotic Normality

For a function  $\boldsymbol{v} \mapsto f(X(\boldsymbol{v}), \boldsymbol{v}) : \boldsymbol{\Upsilon} \to \mathbb{R}$  where  $\boldsymbol{v} \mapsto X(\boldsymbol{v}) : \boldsymbol{\Upsilon} \to \mathbb{R}$  is another function, let  $\bar{\nabla}_x f(X(\boldsymbol{v}), \boldsymbol{v})$  and  $\bar{\nabla}_{\boldsymbol{v}} f(X(\boldsymbol{v}), \boldsymbol{v})$  be given by

$$\bar{\nabla}_x f(X(\boldsymbol{v}), \boldsymbol{v}) := \nabla_{\bar{x}} f(\bar{x}, \bar{\boldsymbol{v}})|_{\bar{x} = X(\boldsymbol{v}), \bar{\boldsymbol{v}} = \boldsymbol{v}} \quad \text{and} \quad \bar{\nabla}_{\boldsymbol{v}} f(X(\boldsymbol{v}), \boldsymbol{v}) := \nabla_{\bar{\boldsymbol{v}}} f(\bar{x}, \bar{\boldsymbol{v}})|_{\bar{x} = X(\boldsymbol{v}), \bar{\boldsymbol{v}} = \boldsymbol{v}},$$

let  $\nabla_{\boldsymbol{v}x} f(X(\boldsymbol{v}), \boldsymbol{v})$  be given by

$$\bar{\nabla}_{\boldsymbol{v}x} f(X(\boldsymbol{v}), \boldsymbol{v}) := \nabla_{\bar{\boldsymbol{v}}\bar{x}} f(\bar{x}, \bar{\boldsymbol{v}})|_{\bar{x}=X(\boldsymbol{v}), \bar{\boldsymbol{v}}=\boldsymbol{v}},$$

and let  $\bar{\nabla}_{xx} f(X(\boldsymbol{v}), \boldsymbol{v})$  and  $\bar{\nabla}_{\boldsymbol{v}\boldsymbol{v}} f(X(\boldsymbol{v}), \boldsymbol{v})$  be given by

$$\bar{\nabla}_{xx} f(X(\boldsymbol{v}), \boldsymbol{v}) := \nabla_{\bar{x}\bar{x}} f(\bar{x}, \bar{\boldsymbol{v}})|_{\bar{x} = X(\boldsymbol{v}), \bar{\boldsymbol{v}} = \boldsymbol{v}} \quad \text{and} \quad \bar{\nabla}_{\boldsymbol{v}\boldsymbol{v}} f(X(\boldsymbol{v}), \boldsymbol{v}) := \nabla_{\bar{\boldsymbol{v}}\bar{\boldsymbol{v}}} f(\bar{x}, \bar{\boldsymbol{v}})|_{\bar{x} = X(\boldsymbol{v}), \bar{\boldsymbol{v}} = \boldsymbol{v}}.$$

In addition to Assumption 1-6, we assume the following.

Assumption 7.  $\theta_0 \in \text{int}(\Theta)$ .

**Assumption 8.** For each  $j \in \{1, ..., J\}$ ,

- (i)  $(x_j, \mathbf{v}_j) \mapsto f_j(y; x_j, \mathbf{v}_j)$  is twice continuously differentiable for all  $y \in \mathcal{Y}$  and
- (ii)  $x_j \mapsto \nabla_{(x_j, \boldsymbol{v}_j)(x_j, \boldsymbol{v}_j)} f_j(y; x_j, \boldsymbol{v}_j)$  is differentiable for all  $y \in \mathcal{Y}$  and  $\boldsymbol{v}_j \in \Upsilon_j$ .

**Assumption 9.** For each  $j \in \{1, ..., J\}$ ,

(i)  $(x_j, \mathbf{v}_j) \mapsto \phi_j(y, x_j; \mathbf{v}_j)$  is twice continuously differentiable for all  $y \in \mathcal{Y}$ .

In the same vein as above, the behaviour of the initialised derivative sequences must first be studied in order to ensure an appropriate form of convergence of the score function and the observed information.

The next two lemmata give conditions under which, for each  $j \in \{1, ..., J\}$ , the non-stationary sequences  $(\nabla_{\boldsymbol{v}_j} \hat{X}_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{N}}$  and  $(\nabla_{\boldsymbol{v}_j} \hat{X}_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{N}}$  converge uniformly e.a.s. to the unique stationary and ergodic sequences  $(\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{Z}}$  and  $(\nabla_{\boldsymbol{v}_j} v_j X_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{Z}}$ , respectively.

**Lemma 3.** Assume that Assumption 1, 2, 4, 9, and the conditions in Lemma 1 hold. Moreover, assume that, for each  $j \in \{1, ..., J\}$ ,

(i) 
$$\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_i \in \Upsilon_i} ||\bar{\nabla}_{\boldsymbol{v}_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)||_2] < \infty.$$

Then, for each  $j \in \{1, ..., J\}$ ,  $(\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{v}_j \in \Upsilon_j$ . Finally, assume that, for each  $j \in \{1, ..., J\}$ ,

- (i) there exists a  $k_j > 1$  such that  $\mathbb{E}[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j)||_2^{2k_j}] < \infty$ ,
- (ii)  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\bar{\nabla}_{\boldsymbol{v}_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) \bar{\nabla}_{\boldsymbol{v}_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)||_2 \stackrel{e.a.s.}{\longrightarrow} 0 \text{ as } t \to \infty, \text{ and } t \to \infty,$

(iii) 
$$\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\bar{\nabla}_{x_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) - \bar{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)| \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty.$$

Then, for each  $j \in \{1, ..., J\}$ ,

$$\sup_{\boldsymbol{v}_i \in \boldsymbol{\Upsilon}_i} \left| \left| \nabla_{\boldsymbol{v}_j} \hat{X}_{j,t}(\boldsymbol{v}_j) - \nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j) \right| \right|_2 \stackrel{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\nabla_{\boldsymbol{v}_i} \hat{X}_{i,1}(\boldsymbol{v}_i) \in \mathbb{R}^{d_i}$ .

**Lemma 4.** Assume that Assumption 1, 2, 4, 9, and the conditions in Lemma 1 and 3 hold. Moreover, assume that, for each  $j \in \{1, ..., J\}$ ,

- (i)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_i \in \boldsymbol{\Upsilon}_i} |\bar{\nabla}_{x_i x_i} \phi_i(Y_t, X_{j,t}(\boldsymbol{v}_i); \boldsymbol{v}_i)|] < \infty,$
- (ii)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_i \in \boldsymbol{\Upsilon}_i} ||\bar{\nabla}_{\boldsymbol{v}_i x_i} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)||_2] < \infty$ , and
- (iii)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \Upsilon_i} ||\bar{\nabla}_{\boldsymbol{v}_j \boldsymbol{v}_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)||_{2,2}] < \infty.$

Then, for each  $j \in \{1,...,J\}$ ,  $(\nabla_{\boldsymbol{v}_j\boldsymbol{v}_j}X_{j,t}(\boldsymbol{v}_j))_{t\in\mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j$ . Finally, assume that, for each  $j \in \{1,...,J\}$ ,

(i)  $\mathbb{E}[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j)||_{2,2}^{k_j}] < \infty$  where  $k_j$  is given in Lemma 3,

(ii) 
$$\sup_{\boldsymbol{v}_j \in \Upsilon_i} |\bar{\nabla}_{x_j x_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) - \bar{\nabla}_{x_j x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)| \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty,$$

(iii) 
$$\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_i} ||\bar{\nabla}_{\boldsymbol{v}_j x_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) - \bar{\nabla}_{\boldsymbol{v}_j x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)||_2 \overset{e.a.s.}{\to} 0 \text{ as } t \to \infty, \text{ and}$$

$$(iv) \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} || \bar{\nabla}_{\boldsymbol{v}_j \boldsymbol{v}_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) - \bar{\nabla}_{\boldsymbol{v}_j \boldsymbol{v}_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) ||_{2,2} \overset{e.a.s.}{\to} 0 \ as \ t \to \infty.$$

Then, for each  $j \in \{1, ..., J\}$ ,

$$\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left| \left| \nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} \hat{X}_{j,t}(\boldsymbol{v}_j) - \nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j) \right| \right|_{2,2} \stackrel{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} \hat{X}_{j,1}(\boldsymbol{v}_j) \in \mathbb{R}^{d_j \times d_j}$ .

Moreover, we assume the following where  $k_i$  is given in Lemma 3.

**Assumption 10.** For each  $j \in \{1, ..., J\}$ ,

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}\left|\bar{\nabla}_{x_j}\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)\right|^{\frac{2k_j}{k_j-1}}\right]<\infty\quad and\quad \mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}\left||\bar{\nabla}_{\boldsymbol{v}_j}\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)\right||_2^2\right]<\infty.$$

Moreover, for each  $j \in \{1, ..., J\}$ ,

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}||\bar{\nabla}_{\boldsymbol{v}_jx_j}\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)||_2^{\frac{k_j}{k_j-1}}\right]<\infty.$$

Finally, for each  $j \in \{1, ..., J\}$ ,

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}\left|\bar{\nabla}_{x_jx_j}\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)\right|^{\frac{k_j}{k_j-1}}\right]<\infty\quad and\quad \mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}\left||\bar{\nabla}_{\boldsymbol{v}_j\boldsymbol{v}_j}\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)\right|\right|_{2,2}\right]<\infty.$$

In the next two and only the next two lemmata, let, for a function  $\mathbf{x} \mapsto \mathbf{f}(\mathbf{x}) : \mathbb{R}^n \to \mathbb{R}^k$ ,  $\nabla_{\mathbf{x}} \mathbf{f}(\mathbf{x})$  be given by

$$[\nabla_{\mathbf{x}} \mathbf{f}(\mathbf{x})]_{(j-1)n+i} = \nabla_{x_i} f_j(\mathbf{x}), \quad (i,j) \in \{1,...,n\} \times \{1,...,k\}$$

and  $\nabla_{\mathbf{x}\mathbf{x}}\mathbf{f}(\mathbf{x})$  be given by

$$[\nabla_{\mathbf{x}\mathbf{x}}\mathbf{f}(\mathbf{x})]_{(i-1)n^2+(i-1)n+l} = \nabla_{x_ix_l}f_i(\mathbf{x}), \quad (i,j,l) \in \{1,...,n\} \times \{1,...,k\} \times \{1,...,n\}.$$

With this notation, the following two lemmata give conditions under which the non-stationary sequences  $(\nabla_{\theta}\hat{\pi}_{t|t}(\theta))_{t\in\mathbb{N}}$  and  $(\nabla_{\theta\theta}\hat{\pi}_{t|t}(\theta))_{t\in\mathbb{N}}$  converge uniformly e.a.s. to the unique stationary and ergodic sequences  $(\nabla_{\theta}\pi_{t|t}(\theta))_{t\in\mathbb{Z}}$  and  $(\nabla_{\theta\theta}\pi_{t|t}(\theta))_{t\in\mathbb{Z}}$ , respectively.

**Lemma 5.** Assume that Assumption 1-4, 8-10, and the conditions in Lemma 1-3 hold. Moreover, assume that for each  $j \in \{1, ..., J\}$ , there exists an  $m_j > 0$  such that

(i) 
$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \left| \bar{\nabla}_{\boldsymbol{v}_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right| \right|_2^{m_j} \right] < \infty$$
,

(ii) 
$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \bar{\nabla}_{x_j x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right|^{m_j} \right] < \infty$$
, and

(iii) 
$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \left| \bar{\nabla}_{\boldsymbol{v}_j x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right| \right|_2^{m_j} \right] < \infty.$$

Then,  $(\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  with  $\mathbb{E}[\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_2^2] < \infty$  and

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \nabla_{\boldsymbol{\theta}} \hat{\boldsymbol{\pi}}_{t|t}(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}) \right| \right|_{2} \stackrel{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\nabla_{\boldsymbol{\theta}} \hat{\boldsymbol{\pi}}_{0|0}(\boldsymbol{\theta}) \in \mathbb{R}^{(J-1)d}$ .

Corollary 2. Under the assumptions in Lemma 5,  $(\nabla_{\theta} \pi_{t|t-1}(\theta))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\theta \in \Theta$  with  $\mathbb{E}[\sup_{\theta \in \Theta} ||\nabla_{\theta} \pi_{t|t-1}(\theta)||_2^2] < \infty$  and

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \nabla_{\boldsymbol{\theta}} \hat{\boldsymbol{\pi}}_{t|t-1}(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t-1}(\boldsymbol{\theta}) \right| \right|_{2} \stackrel{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\nabla_{\boldsymbol{\theta}} \hat{\pi}_{0|0}(\boldsymbol{\theta}) \in \mathbb{R}^{(J-1)d}$ .

**Lemma 6.** Assume that Assumption 1-4, 8-10, and the conditions in Lemma 1-5 hold. Moreover, assume that for each  $j \in \{1, ..., J\}$ , there exists an  $m_j > 0$  such that

(i) 
$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}\sup_{x_j\in\mathcal{X}_j}\left|\left|\bar{\nabla}_{\boldsymbol{v}_j\boldsymbol{v}_j}\log f_j(Y_t;x_j,\boldsymbol{v}_j)\right|\right|_{2,2}^{m_j}\right]<\infty,$$

(ii) 
$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \bar{\nabla}_{x_j x_j x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right|^{m_j} \right] < \infty,$$

(iii) 
$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \left| \bar{\nabla}_{\boldsymbol{v}_j x_j x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right| \right|_2^{m_j} \right] < \infty$$
, and

(iv) 
$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}\sup_{x_j\in\mathcal{X}_j}\left|\left|\bar{\nabla}_{\boldsymbol{v}_j\boldsymbol{v}_jx_j}\log f_j(Y_t;x_j,\boldsymbol{v}_j)\right|\right|_{2,2}^{m_j}\right]<\infty.$$

Then,  $(\nabla_{\theta\theta}\pi_{t|t}(\theta))_{t\in\mathbb{Z}}$  is stationary and ergodic for all  $\theta \in \Theta$  with  $\mathbb{E}[\sup_{\theta \in \Theta} ||\nabla_{\theta\theta}\pi_{t|t}(\theta)||_2] < \infty$  and

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \hat{\boldsymbol{\pi}}_{t|t}(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}) \right| \right|_2 \stackrel{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{\pi}_{0|0}(\boldsymbol{\theta}) \in \mathbb{R}^{(J-1)d^2}$ .

Corollary 3. Under the assumptions in Lemma 6,  $(\nabla_{\theta\theta}\pi_{t|t-1}(\theta))_{t\in\mathbb{Z}}$  is stationary and ergodic for all  $\theta \in \Theta$  with  $\mathbb{E}[\sup_{\theta \in \Theta} ||\nabla_{\theta\theta}\pi_{t|t-1}(\theta)||_2] < \infty$  and

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \hat{\boldsymbol{\pi}}_{t|t-1}(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \boldsymbol{\pi}_{t|t-1}(\boldsymbol{\theta}) \right| \right|_2 \stackrel{e.a.s.}{\to} 0 \quad as \quad t \to \infty$$

for any initialisation  $\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{\boldsymbol{\pi}}_{0|0}(\boldsymbol{\theta}) \in \mathbb{R}^{(J-1)d^2}$ .

Theorem 3 gives conditions under which the ML estimator is also asymptotic normal. In the theorem,

$$\mathbf{I}(\boldsymbol{\theta}) := -\mathbb{E} \left[ \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \log f(Y_t; \boldsymbol{\theta}) \right].$$

is the Fisher information.

**Theorem 3.** Assume that Assumption 1-10 and the conditions in Lemma 1-6 hold and that  $I(\theta_0)$  is invertible. Then,

$$\sqrt{T}(\hat{\boldsymbol{\theta}}_T - \boldsymbol{\theta}_0) \stackrel{d}{\to} \mathcal{N}(\mathbf{0}, \mathbf{I}(\boldsymbol{\theta}_0)^{-1}) \quad as \quad T \to \infty.$$

### 6 The Markov-switching GARCH Model

In this section, we prove the consistency and asymptotic normality of the ML estimator for the Markov-switching GARCH model by Haas et al. (2004b).<sup>5</sup>

Recall that the Markov-switching GARCH model by Haas et al. (2004b) is given by

$$Y_t = \sqrt{X_{S_t,t}} \varepsilon_t,$$

where  $(S_t)_{t\in\mathbb{Z}}$  is a stationary, irreducible, and aperiodic (thus ergodic) Markov chain with transition probabilities  $p_{ij,0} \in (0,1), i,j \in \{1,...,J\}$ , for each  $j \in \{1,...,J\}$ ,

$$X_{j,t+1} = \omega_{j,0} + \alpha_{j,0} Y_t^2 + \beta_{j,0} X_{j,t},$$

where  $\omega_{j,0} > 0$ ,  $\alpha_{j,0} \geq 0$ , and  $\beta_{j,0} \geq 0$ , and  $(\varepsilon_t)_{t \in \mathbb{Z}}$  is a sequence of independent normal distributed random variables with zero mean and unit variance independent of  $(S_t)_{t \in \mathbb{Z}}$ .

Liu (2006) gave conditions under which the model is stationary and ergodic. In the following,  $\mathbf{M}_0$  is a  $J^2 \times J^2$  matrix given by

$$[\mathbf{M}_0]_{ij} = p_{ji,0} (\boldsymbol{\alpha}_0 \mathbf{e}'_i + \boldsymbol{\beta}_0), \quad i, j \in \{1, ..., J\},$$

where  $\alpha_0 = (\alpha_{1,0}, ..., \alpha_{J,0})'$ ,  $\beta_0 = \operatorname{diag}(\beta_{1,0}, ..., \beta_{J,0})$ , and  $\mathbf{e}_i$  is the *i*'th unit vector in  $\mathbb{R}^J$ .

**Theorem 4** (Liu (2006)).  $(Y_t)_{t\in\mathbb{Z}}$  is stationary and ergodic with  $\mathbb{E}[Y_t^2] < \infty$  if and only if  $\rho(\mathbf{M}_0) < 1$  where  $\rho(\mathbf{M}_0)$  is the spectral radius of  $\mathbf{M}_0$ .

We now give conditions under which the ML estimator is consistent and asymptotic normal. The ML estimator is consistent under the following assumptions.

Assumption 11.  $\theta_0 \in \Theta$ .

Assumption 12.  $\rho(\mathbf{M}_0) < 1$ .

Assumption 13.  $\Theta$  is compact.

**Assumption 14.** For all  $\theta \in \Theta$ ,  $\beta_j < 1$  for all  $j \in \{1,...,J\}$ .

**Assumption 15.** There exists an  $m \in \mathbb{N}$  such that  $f^{(m)}(Y_t, ..., Y_{t-m+1}; \boldsymbol{\theta}) = f^{(m)}(Y_t, ..., Y_{t-m+1}; \boldsymbol{\theta}_0)$  a.s. implies that  $\boldsymbol{\theta} = \boldsymbol{\theta}_0$ .

Theorem 5. If Assumption 11-15 hold, then

$$\hat{\boldsymbol{\theta}}_T \overset{a.s.}{\to} \boldsymbol{\theta}_0 \quad as \quad T \to \infty.$$

*Proof.* In the following,  $\underline{\theta} = \inf_{\theta \in \Theta} \theta$  and  $\overline{\theta} = \sup_{\theta \in \Theta} \theta$ . Note that  $(Y_t)_{t \in \mathbb{Z}}$  is stationary and ergodic by Theorem 4. We thus only need to verify Assumption 2-6 and the conditions in Lemma 1 and 2.

<sup>&</sup>lt;sup>5</sup>Recall that the Markov-switching GARCH model reduces to the mixture GARCH model by Haas et al. (2004a) if  $(S_t)_{t\in\mathbb{Z}}$  is an i.i.d. chain.

Let  $j \in \{1, ..., J\}$  be given. First, Assumption 2 is true by assumption and Assumption 3 and 4 are trivially satisfied.

We now verify the conditions in Lemma 1 and 2. First, by Lemma 2.2 in Straumann and Mikosch (2006),

$$\mathbb{E}\left[\log^{+}\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\phi_{j}(Y_{t},x_{j};\boldsymbol{v}_{j})-x_{j}\right|\right]=\mathbb{E}\left[\log^{+}\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\omega_{j}+\alpha_{j}Y_{t}^{2}+\beta_{j}x_{j}-x_{j}\right|\right]\leq C_{j}+2\mathbb{E}\left[\log^{+}\left|Y_{t}\right|\right]$$

for all  $x_j \in \mathcal{X}_j$  where  $C_j = 6 \log 2 + \log^+ \overline{\omega}_j + \log^+ \overline{\alpha}_j + \log^+ \overline{\beta}_j + 2 \log^+ x_j < \infty$ , so Condition (i) in Lemma 1 is satisfied since  $\mathbb{E}[Y_t^2] < \infty$  implies that  $\mathbb{E}[\log^+ |Y_t|] < \infty$  by Lemma 2.2 in Straumann and Mikosch (2006) once again. Moreover,

$$\mathbb{E}\left[\log\sup_{oldsymbol{v}_j\inoldsymbol{\Upsilon}_j}\Lambda_{j,t}(oldsymbol{v}_j)
ight]=\mathbb{E}\left[\log\sup_{oldsymbol{v}_j\inoldsymbol{\Upsilon}_j}eta_j
ight]=\log\overline{eta}_j,$$

so Condition (ii) and (iii) in Lemma 1 are also satisfied. Finally,

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \nabla_{x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right| \right] = \mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| -\frac{1}{2} \frac{1}{x_j} + \frac{1}{2} \frac{Y_t^2}{x_j^2} \right| \right] \leq \frac{1}{2\underline{x}_j} + \frac{1}{2\underline{x}_j^2} \mathbb{E}\left[Y_t^2\right],$$

where  $\underline{x}_j = \frac{\underline{\omega}_j}{1-\underline{\beta}_j} > 0$ , so the condition in Lemma 2 is also satisfied since  $\mathbb{E}[Y_t^2] < \infty$ . Moreover, note that

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})\right|\right] = \mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|-\frac{1}{2}\log 2\pi - \frac{1}{2}\log X_{j,t}(\boldsymbol{v}_{j}) - \frac{1}{2}\frac{Y_{t}^{2}}{X_{j,t}(\boldsymbol{v}_{j})}\right|\right] \\ \leq \frac{1}{2}\log 2\pi + \frac{1}{2}\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\log X_{j,t}(\boldsymbol{v}_{j})\right|\right] + \frac{1}{2\underline{x}_{j}}\mathbb{E}\left[Y_{t}^{2}\right],$$

so Assumption 5 is also satisfied since  $\mathbb{E}\left[Y_t^2\right] < \infty$ . Indeed,  $\mathbb{E}\left[Y_t^2\right] < \infty$  implies that  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left|\log X_{j,t}(\boldsymbol{v}_j)\right|\right] < \infty$  which we now show. Note that  $\log x = \log^+ x - \log^- x$  for all x > 0 where  $\log^+ x = \max(\log x, 0)$  and  $\log^- x = -\min(\log x, 0)$ . Thus,

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}\left|\log X_{j,t}(\boldsymbol{v}_j)\right|\right]\leq \mathbb{E}\left[\log^+\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}X_{j,t}(\boldsymbol{v}_j)\right]+\log^-\underline{x}_j.$$

Now, by Lemma 1,

$$X_{j,t}(\boldsymbol{v}_j) = \frac{\omega_j}{1 - \beta_j} + \alpha_j \sum_{i=0}^{\infty} \beta_j^i Y_{t-1-i}^2 \quad a.s.$$

for all  $v_j \in \Upsilon_j$ . Thus,

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\Upsilon_j}X_{j,t}(\boldsymbol{v}_j)\right]\leq \frac{\overline{\omega}_j}{1-\overline{\beta}_j}+\frac{\overline{\alpha}_j}{1-\overline{\beta}_j}\mathbb{E}\left[Y_{t-1}^2\right].$$

Therefore,  $\mathbb{E}\left[Y_t^2\right] < \infty$  implies that  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left|\log X_{j,t}(\boldsymbol{v}_j)\right|\right] < \infty$  since  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} X_{j,t}(\boldsymbol{v}_j)\right] < \infty$  implies that  $\mathbb{E}\left[\log^+ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} X_{j,t}(\boldsymbol{v}_j)\right] < \infty$  by Lemma 2.2 in Straumann and Mikosch (2006). Finally, Assumption 6 is true by assumption.

To give conditions under which the ML estimator is also asymptotic normal, we need the following result where, for all  $s \in \mathbb{N}$ ,  $\Sigma_0^{(\otimes s)}$  is a  $J^{s+1} \times J^{s+1}$  matrix given by

$$\left[\boldsymbol{\Sigma}_{0}^{(\otimes s)}\right]_{ij} = p_{ji,0} \mathbb{E}\left[\mathbf{A}_{it,0}^{(\otimes s)}\right], \quad i, j \in \{1, ..., J\}$$

with

$$\mathbf{A}_{it,0} = \varepsilon_t^2 \boldsymbol{\alpha}_0 \mathbf{e}_i' + \boldsymbol{\beta}_0,$$

where  $\otimes$  denotes the Kronecker product.

**Theorem 6** (Liu (2006)). If 
$$\rho(\mathbf{M}_0) < 1$$
 and  $\rho(\Sigma_0^{(\otimes s)}) < 1$ , then  $\mathbb{E}[Y_t^{2s}] < \infty$ .

In addition to the assumptions above, the ML estimator is asymptotic normal under the following assumptions.

Assumption 16.  $\theta_0 \in int(\Theta)$ .

Assumption 17. 
$$\rho\left(\Sigma_0^{(\otimes 4)}\right) < 1$$
.

**Theorem 7.** If Assumption 11-17 hold and  $I(\theta_0)$  is invertible, then

$$\sqrt{T}(\hat{\boldsymbol{\theta}}_T - \boldsymbol{\theta}_0) \stackrel{d}{\to} \mathcal{N}(\mathbf{0}, \mathbf{I}(\boldsymbol{\theta}_0)^{-1}) \quad as \quad T \to \infty.$$

*Proof.* We need to verify Assumption 7-10 and the conditions in Lemma 3-6.

Let, as in the proof of Theorem 5,  $j \in \{1, ..., J\}$  be given. First, Assumption 7 is true by assumption and Assumption 8 and 9 are trivially satisfied.

We now verify the conditions in Lemma 3. First, by Lemma 2.2 in Straumann and Mikosch (2006),

$$\mathbb{E}\left[\log^{+}\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\left|\bar{\nabla}_{\boldsymbol{v}_{j}}\phi_{j}(Y_{t},X_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\right|\right|_{2}\right] \leq \mathbb{E}\left[\log^{+}\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left(1+Y_{t}^{2}+X_{j,t}(\boldsymbol{v}_{j})\right)\right] \\
\leq 4\log 2 + 2\mathbb{E}\left[\log^{+}|Y_{t}|\right] + \mathbb{E}\left[\log^{+}\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}X_{j,t}(\boldsymbol{v}_{j})\right],$$

so Condition (i) is satisfied since  $\mathbb{E}\left[Y_t^8\right] < \infty$  implies that  $\mathbb{E}\left[\log^+|Y_t|\right] < \infty$  by Lemma 2.2 in Straumann and Mikosch (2006) once again and that  $\mathbb{E}\left[\log^+\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}X_{j,t}(\boldsymbol{v}_j)\right] < \infty$ , see the proof of Theorem 5. We now show that  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}\left|\left|\nabla_{\boldsymbol{v}_j}X_{j,t}(\boldsymbol{v}_j)\right|\right|_2^4\right] < \infty$ . First,

$$\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j) = \sum_{i=0}^{\infty} \beta_j^i \mathbf{v}_{j,t-1-i}(\boldsymbol{v}_j) \quad a.s.$$

for all  $v_j \in \Upsilon_j$  where  $\mathbf{v}_{j,t}(v_j) = (1, Y_t^2, X_{j,t}(v_j))'$ , so, by Minkowskis inequality,

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\left|\nabla_{\boldsymbol{v}_{j}}X_{j,t}(\boldsymbol{v}_{j})\right|\right|_{2}^{4}\right]^{\frac{1}{4}} \leq \mathbb{E}\left[\left(\sum_{i=0}^{\infty}\overline{\beta}_{j}^{i}\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\left|\mathbf{v}_{j,t-1-i}(\boldsymbol{v}_{j})\right|\right|_{2}\right)^{4}\right]^{\frac{1}{4}}$$

$$\leq \sum_{i=0}^{\infty}\overline{\beta}_{j}^{i}\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\left|\mathbf{v}_{j,t-1}(\boldsymbol{v}_{j})\right|\right|_{2}^{4}\right]^{\frac{1}{4}}$$

$$\leq \frac{1}{1-\overline{\beta}_j} \left( 4 + 16\mathbb{E}\left[Y_{t-1}^8\right] + 16\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} X_{j,t-1}^4(\boldsymbol{v}_j)\right] \right)^{\frac{1}{4}},$$

where the last inequality follows from the inequality  $|x+y|^p \le 2^p |x|^p + 2^p |y|^p$  for all  $x, y \in \mathbb{R}$  and  $p \in (0, \infty)$ . Moreover, by Lemma 1,

$$X_{j,t}(\boldsymbol{v}_j) = \frac{\omega_j}{1 - \beta_j} + \alpha_j \sum_{i=0}^{\infty} \beta_j^i Y_{t-1-i}^2 \quad a.s.$$

for all  $v_j \in \Upsilon_j$ , so

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}X_{j,t}^{4}(\boldsymbol{v}_{j})\right]^{\frac{1}{4}} \leq \mathbb{E}\left[\left(\frac{\overline{\omega}_{j}}{1-\overline{\beta}_{j}}+\overline{\alpha}_{j}\sum_{i=0}^{\infty}\overline{\beta}_{j}^{i}Y_{t-1-i}^{2}\right)^{4}\right]^{\frac{1}{4}}$$

$$\leq \frac{\overline{\omega}_{j}}{1-\overline{\beta}_{j}}+\overline{\alpha}_{j}\sum_{i=0}^{\infty}\overline{\beta}_{j}^{i}\mathbb{E}\left[Y_{t-1}^{8}\right]^{\frac{1}{4}}$$

$$=\frac{\overline{\omega}_{j}}{1-\overline{\beta}_{j}}+\frac{\overline{\alpha}_{j}}{1-\overline{\beta}_{j}}\mathbb{E}\left[Y_{t-1}^{8}\right]^{\frac{1}{4}}$$

by Minkowskis inequality once again. Therefore,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left|\left|\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j)\right|\right|_2^4\right] < \infty$  since  $\mathbb{E}\left[Y_t^8\right] < \infty$ . Finally,

$$\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left| \left| \bar{\nabla}_{\boldsymbol{v}_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) - \bar{\nabla}_{\boldsymbol{v}_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) \right| \right|_2 = \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left| \hat{X}_{j,t}(\boldsymbol{v}_j) - X_{j,t}(\boldsymbol{v}_j) \right|,$$

and

$$\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left| \bar{\nabla}_{x_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) - \bar{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) \right| = 0,$$

so Condition (ii) and (iii) are also satisfied since  $\sup_{\boldsymbol{v}_j \in \Upsilon_j} \left| \hat{X}_{j,t}(\boldsymbol{v}_j) - X_{j,t}(\boldsymbol{v}_j) \right| \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$ . The conditions in Lemma 4 can be verified similarly.

Moving on to Assumption 10,

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\bar{\nabla}_{x_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})\right|^{4}\right] = \mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|-\frac{1}{2}\frac{1}{X_{j,t}(\boldsymbol{v}_{j})} + \frac{1}{2}\frac{Y_{t}^{2}}{X_{j,t}^{2}(\boldsymbol{v}_{j})}\right|^{4}\right]$$

$$\leq \frac{1}{\underline{x}_{j}^{4}} + \frac{1}{\underline{x}_{j}^{8}}\mathbb{E}\left[Y_{t}^{8}\right]$$

and

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\bar{\nabla}_{x_{j}x_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})\right|^{2}\right] = \mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\left|\frac{1}{2}\frac{1}{X_{j,t}^{2}(\boldsymbol{v}_{j})} - \frac{Y_{t}^{2}}{X_{j,t}^{3}(\boldsymbol{v}_{j})}\right|^{2}\right]$$

$$\leq \frac{2}{\underline{x}_{j}^{4}} + \frac{4}{\underline{x}_{j}^{6}}\mathbb{E}\left[Y_{t}^{4}\right],$$

so Assumption 10 is also satisfied since  $\mathbb{E}\left[Y_t^8\right] < \infty$ .

Finally, we verify the condition in Lemma 5. Note that

$$\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \bar{\nabla}_{x_j x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right| \right] = \mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \frac{1}{2} \frac{1}{x_j^2} - \frac{Y_t^2}{x_j^3} \right| \right]$$

$$\leq \frac{1}{2\underline{x}_{j}^{2}} + \frac{1}{\underline{x}_{j}^{3}} \mathbb{E}\left[Y_{t}^{2}\right],$$

so that condition is also satisfied since  $\mathbb{E}\left[Y_t^8\right]<\infty$ . The condition in Lemma 6 can be verified similarly.

Note that Assumption 17 is a strong assumption that might be relaxed. We leave this for future research.

### 7 Conclusion

State space models, autoregressive state space models, and observation-driven state space models have been applied in numerous areas so statistical inference for these models is of significant practical importance. While it has attracted much attention in the literature for both state space models and autoregressive state space models, it is to a large extent undiscovered land for observation-driven state space models.

In this paper, we proved consistency and asymptotic normality of the ML estimator for an observation-driven state space model where X is finite. As a special case of the general theory, we gave conditions under which the ML estimator for the widely applied Markov-switching GARCH model by Haas et al. (2004b) is both consistent and asymptotic normal thus extending the work by Kandji and Misko (2024).

An interesting extension of the paper could be to extend the theory to observation-driven state space models where X is not necessarily finite to cover more of the examples in the introduction. We leave this for future research.

### References

- A. Aknouche and C. Francq. Stationarity and ergodicity of markov switching positive conditional mean models. *Journal of Time Series Analysis*, 43(3):436–459, 2022.
- A. Ang and A. Timmermann. Regime changes and financial markets. *Annu. Rev. Financ. Econ.*, 4(1):313–337, 2012.
- G. Angelini and P. Gorgi. Dsge models with observation-driven time-varying volatility. *Economics Letters*, 171:169–171, 2018.
- D. Ardia, K. Bluteau, K. Boudt, and L. Catania. Forecasting risk with markov-switching garch models: A large-scale performance study. *International Journal of Forecasting*, 34(4):733–747, 2018.
- M. Bennedsen, E. Hillebrand, and S. J. Koopman. A multivariate dynamic statistical model of the global carbon budget 1959–2020. *Journal of the Royal Statistical Society Series A:* Statistics in Society, 186(1):20–42, 2023.
- M. Bernardi and L. Catania. Switching generalized autoregressive score copula models with application to systemic risk. *Journal of Applied Econometrics*, 34(1):43–65, 2019.
- P. J. Bickel, Y. Ritov, and T. Ryden. Asymptotic normality of the maximum-likelihood estimator for general hidden markov models. *The Annals of Statistics*, 26(4):1614–1635, 1998.
- P. Billingsley. The lindeberg-levy theorem for martingales. *Proceedings of the American Mathematical Society*, 12(5):788–792, 1961.
- F. Blasques, P. Gorgi, S. J. Koopman, and O. Wintenberger. Feasible invertibility conditions and maximum likelihood estimation for observation-driven models. 2018.
- F. Blasques, J. van Brummelen, S. J. Koopman, and A. Lucas. Maximum likelihood estimation for score-driven models. *Journal of Econometrics*, 227(2):325–346, 2022.
- T. Bollerslev. Generalized autoregressive conditional heteroskedasticity. *Journal of econometrics*, 31(3):307–327, 1986.
- P. Bougerol. Kalman filtering with random coefficients and contractions. SIAM Journal on Control and Optimization, 31(4):942–959, 1993.
- F. Bräuning and S. J. Koopman. The dynamic factor network model with an application to international trade. *Journal of Econometrics*, 216(2):494–515, 2020.
- S. A. Broda, M. Haas, J. Krause, M. S. Paolella, and S. C. Steude. Stable mixture garch models. *Journal of Econometrics*, 172(2):292–306, 2013.
- G. Buccheri and F. Corsi. Hark the shark: Realized volatility modeling with measurement errors and nonlinear dependencies. *Journal of Financial Econometrics*, 19(4):614–649, 2021.

- G. Buccheri, G. Bormetti, F. Corsi, and F. Lillo. A score-driven conditional correlation model for noisy and asynchronous data: An application to high-frequency covariance dynamics. *Journal of Business & Economic Statistics*, 39(4):920–936, 2021.
- J. Bulla, F. Lagona, A. Maruotti, and M. Picone. A multivariate hidden markov model for the identification of sea regimes from incomplete skewed and circular time series. *Journal of Agricultural, Biological, and Environmental Statistics*, 17:544–567, 2012.
- J. Cai. A markov model of switching-regime arch. *Journal of Business & Economic Statistics*, 12(3):309–316, 1994.
- L. Catania. A stochastic volatility model with a general leverage specification. *Journal of Business & Economic Statistics*, 40(2):678–689, 2022.
- G. A. Churchill. Stochastic models for heterogeneous dna sequences. *Bulletin of mathematical biology*, 51(1):79–94, 1989.
- R. Douc, E. Moulines, and T. Rydén. Asymptotic properties of the maximum likelihood estimator in autoregressive models with markov regime. 2004.
- R. Douc, É. Moulines, J. Olsson, and R. van Handel. Consistency of the maximum likelihood estimator for general hidden markov models. *Annals of Statistics*, 39(1):474–513, 2011.
- R. F. Engle. Autoregressive conditional heteroscedasticity with estimates of the variance of united kingdom inflation. *Econometrica: Journal of the econometric society*, pages 987–1007, 1982.
- C. Francq and M. Roussignol. Ergodicity of autoregressive processes with markov-switching and consistency of the maximum-likelihood estimator. *Statistics: A Journal of Theoretical and Applied Statistics*, 32(2):151–173, 1998.
- C. Francq and J.-M. Zakoian. *GARCH models: structure, statistical inference and financial applications*. John Wiley & Sons, 2019.
- S. Gammelmark, K. Mølmer, W. Alt, T. Kampschulte, and D. Meschede. Hidden markov model of atomic quantum jump dynamics in an optically probed cavity. *Physical Review A*, 89(4): 043839, 2014.
- S. F. Gray. Modeling the conditional distribution of interest rates as a regime-switching process. Journal of Financial Economics, 42(1):27–62, 1996.
- M. Haas, S. Mittnik, and M. S. Paolella. Mixed normal conditional heteroskedasticity. *Journal of financial Econometrics*, 2(2):211–250, 2004a.
- M. Haas, S. Mittnik, and M. S. Paolella. A new approach to markov-switching garch models. *Journal of financial Econometrics*, 2(4):493–530, 2004b.
- J. D. Hamilton. A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica: Journal of the econometric society*, pages 357–384, 1989.

- J. D. Hamilton. *Time series analysis*. Princeton university press, 1994.
- J. D. Hamilton. Regime switching models. In *Macroeconometrics and time series analysis*, pages 202–209. Springer, 2010.
- J. D. Hamilton and R. Susmel. Autoregressive conditional heteroskedasticity and changes in regime. *Journal of econometrics*, 64(1-2):307–333, 1994.
- A. Harvey and C.-H. Chung. Estimating the underlying change in unemployment in the uk. Journal of the Royal Statistical Society: Series A (Statistics in Society), 163(3):303–309, 2000.
- A. Harvey and D. Palumbo. Regime switching models for circular and linear time series. *Journal of Time Series Analysis*, 44(4):374–392, 2023.
- A. C. Harvey and N. Shephard. Estimation of an asymmetric stochastic volatility model for asset returns. *Journal of Business & Economic Statistics*, 14(4):429–434, 1996.
- E. Jacquier, N. G. Polson, and P. E. Rossi. Bayesian analysis of stochastic volatility models with fat-tails and correlated errors. *Journal of Econometrics*, 122(1):185–212, 2004.
- J. L. Jensen and N. V. Petersen. Asymptotic normality of the maximum likelihood estimator in state space models. *The Annals of Statistics*, 27(2):514–535, 1999.
- B. M. Kandji and A. Misko. Markov-switching normal-mixture garch. 2024.
- H. Kasahara and K. Shimotsu. Asymptotic properties of the maximum likelihood estimator in regime switching econometric models. *Journal of econometrics*, 208(2):442–467, 2019.
- V. Krishnamurthy and T. Ryden. Consistent estimation of linear and non-linear autoregressive models with markov regime. *Journal of time series analysis*, 19(3):291–307, 1998.
- R. Langrock, R. King, J. Matthiopoulos, L. Thomas, D. Fortin, and J. M. Morales. Flexible and practical modeling of animal telemetry data: hidden markov models and extensions. *Ecology*, 93(11):2336–2342, 2012.
- B. G. Leroux. Maximum-likelihood estimation for hidden markov models. *Stochastic processes* and their applications, 40(1):127–143, 1992.
- J.-C. Liu. Stationarity of a markov-switching garch model. *Journal of Financial Econometrics*, 4(4):573–593, 2006.
- D. D. Monache, I. Petrella, and F. Venditti. Common faith or parting ways? a time varying parameters factor analysis of euro-area inflation. In *Dynamic factor models*, volume 35, pages 539–565. Emerald Group Publishing Limited, 2016.
- D. D. Monache, I. Petrella, and F. Venditti. Price dividend ratio and long-run stock returns: A score-driven state space model. *Journal of Business & Economic Statistics*, 39(4):1054–1065, 2021.

- B. M. Pötscher and I. Prucha. *Dynamic nonlinear econometric models: Asymptotic theory*. Springer Science & Business Media, 1997.
- R. R. Rao. Relations between weak and uniform convergence of measures with applications. *The Annals of Mathematical Statistics*, pages 659–680, 1962.
- T. Rydén, T. Teräsvirta, and S. Åsbrink. Stylized facts of daily return series and the hidden markov model. *Journal of applied econometrics*, 13(3):217–244, 1998.
- J. H. Stock and M. W. Watson. New indexes of coincident and leading economic indicators. NBER macroeconomics annual, 4:351–394, 1989.
- D. Straumann and T. Mikosch. Quasi-maximum-likelihood estimation in conditionally heteroscedastic time series: A stochastic recurrence equations approach. 2006.
- J. Yu. On leverage in a stochastic volatility model. *Journal of Econometrics*, 127(2):165–178, 2005.
- W. Zucchini and P. Guttorp. A hidden markov model for space-time precipitation. Water Resources Research, 27(8):1917–1923, 1991.

### Appendix

In the following,  $C \in \mathbb{R}$  is an arbitrary constant that can change from line to line.

### A Main Proofs

#### A.1 Proof of Lemma 1

For each  $j \in \{1, ..., J\}$ ,  $(X_{j,t}(v_j))_{t \in \mathbb{Z}}$  is a stochastic process taking values in  $(\mathcal{X}_j, |\cdot|)$  given by

$$X_{j,t+1}(\boldsymbol{v}_j) = \phi_{j,t}(X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j),$$

where  $(\phi_{j,t}(\cdot; v_j))_{t \in \mathbb{Z}}$  is a sequence of stationary and ergodic Lipschitz functions given by

$$\phi_{j,t}(x_j; \boldsymbol{v}_j) = \phi_j(Y_t, x_j; \boldsymbol{v}_j)$$

with Lipschitz coefficient

$$\Lambda(\phi_{j,t}; \boldsymbol{v}_j) = \sup_{\substack{x_j, y_j \in \mathcal{X}_j \\ x_j \neq y_j}} \frac{|\phi_{j,t}(x_j; \boldsymbol{v}_j) - \phi_{j,t}(y_j; \boldsymbol{v}_j)|}{|x_j - y_j|} \leq \sup_{x_j \in \mathcal{X}_j} |\nabla_{x_j} \phi_{j,t}(x_j; \boldsymbol{v}_j)| = \Lambda_{j,t}(\boldsymbol{v}_j).$$

The conclusion follows from Theorem 3.1 in Bougerol (1993), see also Lemma C.1, since, for each  $j \in \{1, ..., J\}$ ,

- (i) there exists an  $x_j \in \mathcal{X}_j$  such that  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_i} |\phi_{j,t}(x_j; \boldsymbol{v}_j) x_j|] < \infty$ ,
- (ii)  $\mathbb{E}[\log^{+}\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}\Lambda_{j,t}(\boldsymbol{v}_{j})]<\infty$ , and
- (iii)  $\mathbb{E}[\log \sup_{\boldsymbol{v}_i \in \boldsymbol{\Upsilon}_i} \Lambda_{j,t}(\boldsymbol{v}_j)] < 0$

by assumption.

#### A.2 Proof of Lemma 2

The proof consists of two parts. The first part considers the stochastic process  $(\pi_{t|t}(\boldsymbol{\theta}))_{t\in\mathbb{Z}}$ , and the second part considers the stochastic process  $(\hat{\pi}_{t|t}(\boldsymbol{\theta}))_{t\in\mathbb{N}}$ .

First,  $(\pi_{t|t}(\boldsymbol{\theta}))_{t\in\mathbb{Z}}$  is a stochastic process taking values in  $(\mathcal{S}, ||\cdot||_2)$  given by

$$\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}) = \boldsymbol{\phi}_t(\boldsymbol{\pi}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}),$$

where  $(\phi_t(\cdot;\boldsymbol{\theta}))_{t\in\mathbb{Z}}$  is a sequence of stationary and ergodic Lipschitz functions given by

$$\phi_t(\mathbf{s}; \boldsymbol{\theta}) = \mathbf{F}_t(\mathbf{P}'\mathbf{s}; \boldsymbol{\theta})\mathbf{P}'\mathbf{s}$$

with Lipschitz coefficient

$$\Lambda(\phi_t; \boldsymbol{\theta}) = \sup_{\substack{\mathbf{x}, \mathbf{y} \in \mathcal{S} \\ \mathbf{x} \neq \mathbf{y}}} \frac{||\phi_t(\mathbf{x}; \boldsymbol{\theta}) - \phi_t(\mathbf{y}; \boldsymbol{\theta})||_2}{||\mathbf{x} - \mathbf{y}||_2}.$$

The fact that  $(\pi_{t|t}(\boldsymbol{\theta}))_{t\in\mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  follows from Theorem 3.1 in Bougerol (1993) if

- (i) there exists an  $\mathbf{s} \in \mathcal{S}$  such that  $\mathbb{E}[\log^+ \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\boldsymbol{\phi}_t(\mathbf{s}; \boldsymbol{\theta}) \mathbf{s}||_2] < \infty$ ,
- (ii)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \Lambda(\boldsymbol{\phi}_t; \boldsymbol{\theta})] < \infty$ , and
- (iii) there exists an  $r \in \mathbb{N}$  such that  $\mathbb{E}[\log \sup_{\theta \in \Theta} \Lambda(\phi_t^{(r)}; \theta)] < 0$ .

Condition (i) is trivial.

Note that

$$\mathbb{P}_{\boldsymbol{\theta}}(S_t = j \mid \mathbf{Y}_{-\infty}^u) = \sum_{i=1}^J \mathbb{P}_{\boldsymbol{\theta}}(S_t = j \mid S_{t-1} = i, \mathbf{Y}_{-\infty}^u) \mathbb{P}_{\boldsymbol{\theta}}(S_{t-1} = i \mid \mathbf{Y}_{-\infty}^u)$$

for all  $t, u \in \mathbb{Z}$  such that  $t \leq u$  where

$$\mathbb{P}_{\theta}(S_{t} = j \mid S_{t-1} = i, \mathbf{Y}_{-\infty}^{u}) = \frac{\mathbb{P}_{\theta}(\mathbf{Y}_{t}^{u} \in d\mathbf{y}_{t}^{u}, S_{t} = j \mid S_{t-1} = i, \mathbf{Y}_{-\infty}^{t-1})}{\sum_{k=1}^{J} \mathbb{P}_{\theta}(\mathbf{Y}_{t}^{u} \in d\mathbf{y}_{t}^{u}, S_{t} = k \mid S_{t-1} = i, \mathbf{Y}_{-\infty}^{t-1})} \\
= \frac{\mathbb{P}_{\theta}(\mathbf{Y}_{t}^{u} \in d\mathbf{y}_{t}^{u} \mid \mathbf{Y}_{-\infty}^{t-1}, S_{t} = j, S_{t-1} = i) \mathbb{P}_{\theta}(S_{t} = j \mid S_{t-1} = i, \mathbf{Y}_{-\infty}^{t-1})}{\sum_{k=1}^{J} \mathbb{P}_{\theta}(\mathbf{Y}_{t}^{u} \in d\mathbf{y}_{t}^{u} \mid \mathbf{Y}_{-\infty}^{t-1}, S_{t} = k, S_{t-1} = i) \mathbb{P}_{\theta}(S_{t} = k \mid S_{t-1} = i, \mathbf{Y}_{-\infty}^{t-1})} \\
= \frac{\mathbb{P}_{\theta}(\mathbf{Y}_{t}^{u} \in d\mathbf{y}_{t}^{u} \mid \mathbf{Y}_{-\infty}^{t-1}, S_{t} = j) \mathbb{P}_{\theta}(S_{t} = j \mid S_{t-1} = i)}{\sum_{k=1}^{J} \mathbb{P}_{\theta}(\mathbf{Y}_{t}^{u} \in d\mathbf{y}_{t}^{u} \mid \mathbf{Y}_{-\infty}^{t-1}, S_{t} = k) \mathbb{P}_{\theta}(S_{t} = k \mid S_{t-1} = i)}.$$

Hence,  $\pi_{j,t|u}(\boldsymbol{\theta}) := \mathbb{P}_{\boldsymbol{\theta}}(S_t = j \mid \mathbf{Y}^u_{-\infty}), j \in \{1,...,J\}$  is given by

$$\pi_{j,t|u}(\boldsymbol{\theta}) = \sum_{i=1}^{J} \frac{\mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{t}^{u} \in \mathrm{d}\mathbf{y}_{t}^{u} \mid \mathbf{Y}_{-\infty}^{t-1}, S_{t} = j) p_{ij}}{\sum_{k=1}^{J} \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{t}^{u} \in \mathrm{d}\mathbf{y}_{t}^{u} \mid \mathbf{Y}_{-\infty}^{t-1}, S_{t} = k) p_{ik}} \pi_{i,t-1|u}(\boldsymbol{\theta})$$

for all  $t, u \in \mathbb{Z}$  such that  $t \leq u$ , and  $\pi_{t|u}(\theta) := (\pi_{1,t|u}(\theta), ..., \pi_{J,t|u}(\theta))'$  is given by

$$oldsymbol{\pi}_{t|u}(oldsymbol{ heta}) = \mathbf{M}_{t|u}'(oldsymbol{ heta}) oldsymbol{\pi}_{t-1|u}(oldsymbol{ heta})$$

for all  $t, u \in \mathbb{Z}$  such that  $t \leq u$  where  $\mathbf{M}_{t|u}(\boldsymbol{\theta})$  is a stochastic matrix, that is, a matrix with non-negative elements where each row sum to one, with generic element

$$[\mathbf{M}_{t|u}(\boldsymbol{\theta})]_{ij} = \frac{\mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_t^u \in d\mathbf{y}_t^u \mid \mathbf{Y}_{-\infty}^{t-1}, S_t = j)p_{ij}}{\sum_{t=1}^{J} \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_t^u \in d\mathbf{y}_t^u \mid \mathbf{Y}_{-\infty}^{t-1}, S_t = k)p_{ik}}, \quad i, j \in \{1, ..., J\}.$$

Thus, we have that

$$\pi_{t|t}(\boldsymbol{\theta}) = \mathbf{M}'_{t|t}(\boldsymbol{\theta}) \cdots \mathbf{M}'_{t-r+1|t}(\boldsymbol{\theta}) \pi_{t-r|t}(\boldsymbol{\theta})$$
 (3)

for all  $r \in \mathbb{N}$  where

$$[\boldsymbol{\pi}_{t-r|t}(\boldsymbol{\theta})]_{j} = \frac{\mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{t-r+1}^{t} \in \mathrm{d}\mathbf{y}_{t-r+1}^{t} \mid \mathbf{Y}_{-\infty}^{t-r}, S_{t-r} = j)[\boldsymbol{\pi}_{t-r|t-r}(\boldsymbol{\theta})]_{j}}{\sum_{k=1}^{J} \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{t-r+1}^{t} \in \mathrm{d}\mathbf{y}_{t-r+1}^{t} \mid \mathbf{Y}_{-\infty}^{t-r}, S_{t-r} = k)[\boldsymbol{\pi}_{t-r|t-r}(\boldsymbol{\theta})]_{k}}, \quad j \in \{1, ..., J\}.$$

This observation leads to Lemma A.2 proved using the following lemma.

**Lemma A.1.** If there exists an  $\varepsilon \in (0,1)$  such that

$$p_{ij} \geq \varepsilon$$

for all  $i, j \in \{1, ..., J\}$ , then, for all  $\tau, t \in \mathbb{Z}$  such that  $\tau \leq t$ , there exists a  $\mathbf{v}_{\tau|t}(\boldsymbol{\theta}) \in \mathcal{S}$  such that

$$[\mathbf{M}_{\tau|t}(\boldsymbol{\theta})]_{ij} \geq \varepsilon [\mathbf{v}_{\tau|t}(\boldsymbol{\theta})]_{j}$$

for all  $i, j \in \{1, ..., J\}$ .

*Proof.* Let  $\mathbf{v}_{\tau|t}(\boldsymbol{\theta})$  be a stochastic vector, that is, a vector with non-negative elements that sum to one, with generic element

$$[\mathbf{v}_{\tau|t}(\boldsymbol{\theta})]_j = \frac{\mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{\tau}^t \in \mathrm{d}\mathbf{y}_{\tau}^t \mid \mathbf{Y}_{-\infty}^{\tau-1}, S_{\tau} = j)}{\sum_{k=1}^J \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{\tau}^t \in \mathrm{d}\mathbf{y}_{\tau}^t \mid \mathbf{Y}_{-\infty}^{\tau-1}, S_{\tau} = k)}, \quad j \in \{1, ..., J\}.$$

Then,

$$[\mathbf{M}_{\tau|t}(\boldsymbol{\theta})]_{ij} = \frac{\mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{\tau}^{t} \in d\mathbf{y}_{\tau}^{t} \mid \mathbf{Y}_{-\infty}^{\tau-1}, S_{\tau} = j)p_{ij}}{\sum_{k=1}^{J} \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{\tau}^{t} \in d\mathbf{y}_{\tau}^{t} \mid \mathbf{Y}_{-\infty}^{\tau-1}, S_{\tau} = k)p_{ik}}$$

$$\geq \varepsilon \frac{\mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{\tau}^{t} \in d\mathbf{y}_{\tau}^{t} \mid \mathbf{Y}_{-\infty}^{\tau-1}, S_{\tau} = j)}{\sum_{k=1}^{J} \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{\tau}^{t} \in d\mathbf{y}_{\tau}^{t} \mid \mathbf{Y}_{-\infty}^{\tau-1}, S_{\tau} = k)} = \varepsilon [\mathbf{v}_{\tau|t}(\boldsymbol{\theta})]_{j}$$

for all  $i, j \in \{1, ..., J\}$ .

**Lemma A.2.** Assume that there exists an  $\varepsilon \in (0,1)$  such that

$$p_{ij} \geq \varepsilon$$

for all  $i, j \in \{1, ..., J\}$ . Then, there exists an  $\alpha \in (0, 1)$  such that

$$||\phi_t^{(r)}(\mathbf{x}; \boldsymbol{\theta}) - \phi_t^{(r)}(\mathbf{y}; \boldsymbol{\theta})||_2 \le \alpha^r C||\mathbf{x} - \mathbf{y}||_2$$

for all  $r \in \mathbb{N}$  and  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$ .

*Proof.* Let  $r \in \mathbb{N}$  be given. First,  $||\mathbf{z}||_2 \leq ||\mathbf{z}||_1$  for all  $\mathbf{z} \in \mathbb{R}^J$  implies that

$$||\phi_t^{(r)}(\mathbf{x};\boldsymbol{\theta}) - \phi_t^{(r)}(\mathbf{y};\boldsymbol{\theta})||_2 \le ||\phi_t^{(r)}(\mathbf{x};\boldsymbol{\theta}) - \phi_t^{(r)}(\mathbf{y};\boldsymbol{\theta})||_1$$
(4)

for all  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$ . To ease the notation, let  $p_{j,t} := \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{t-r+1}^t \in d\mathbf{y}_{t-r+1}^t \mid \mathbf{Y}_{-\infty}^{t-r}, S_{t-r} = j)$ . Equation (3) shows that

$$oldsymbol{\phi}_t^{(r)}(\mathbf{z};oldsymbol{ heta}) = \mathbf{M}_{t|t}'(oldsymbol{ heta}) \cdots \mathbf{M}_{t-r+1|t}'(oldsymbol{ heta}) ilde{\mathbf{z}}$$

for all  $\mathbf{z} \in \mathcal{S}$  where

$$\tilde{z}_j = \frac{p_{j,t} z_j}{\sum_{k=1}^J p_{k,t} z_k}.$$

An application of Lemma D.1 together with Lemma A.1 thus shows that there exists an  $\alpha \in (0, 1)$  such that

$$||\phi_t^{(r)}(\mathbf{x}; \boldsymbol{\theta}) - \phi_t^{(r)}(\mathbf{y}; \boldsymbol{\theta})||_1 \le \alpha^r ||\tilde{\mathbf{x}} - \tilde{\mathbf{y}}||_1$$

for all  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$  where

$$||\tilde{\mathbf{x}} - \tilde{\mathbf{y}}||_1 = \sum_{j=1}^{J} \left| \frac{p_{j,t} x_j}{\sum_{k=1}^{J} p_{k,t} x_k} - \frac{p_{j,t} y_j}{\sum_{k=1}^{J} p_{k,t} y_k} \right|.$$

Note that

$$\left| \frac{p_{j,t}x_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} - \frac{p_{j,t}y_{j}}{\sum_{k=1}^{J} p_{k,t}y_{k}} \right| = \left| \frac{p_{j,t}x_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} - \frac{p_{j,t}y_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} + \frac{p_{j,t}y_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} - \frac{p_{j,t}y_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} \right|$$

$$\leq \left| \frac{p_{j,t}x_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} - \frac{p_{j,t}y_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} \right| + \left| \frac{p_{j,t}y_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} - \frac{p_{j,t}y_{j}}{\sum_{k=1}^{J} p_{k,t}x_{k}} \right|$$

$$= \frac{p_{j,t}}{\sum_{k=1}^{J} p_{k,t} x_{k}} |x_{j} - y_{j}| + \frac{p_{j,t} y_{j}}{\sum_{k=1}^{J} p_{k,t} y_{k}} \frac{1}{\sum_{k=1}^{J} p_{k,t} x_{k}} \left| \sum_{k=1}^{J} p_{k,t} (x_{k} - y_{k}) \right|$$

$$\leq \frac{p_{j,t}}{\sum_{k=1}^{J} p_{k,t} x_{k}} |x_{j} - y_{j}| + \sum_{k=1}^{J} \frac{p_{k,t}}{\sum_{k=1}^{J} p_{k,t} x_{k}} |x_{k} - y_{k}|$$

$$\leq \frac{1}{\varepsilon} |x_{j} - y_{j}| + \frac{1}{\varepsilon} \sum_{k=1}^{J} |x_{k} - y_{k}|$$

since

$$p_{i,t} = \sum_{j=1}^{J} \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{Y}_{t-r+1}^{t} \in d\mathbf{y}_{t-r+1}^{t} \mid \mathbf{Y}_{-\infty}^{t-r}, S_{t-r+1} = j) p_{ij}.$$

Hence,

$$||\phi_t^{(r)}(\mathbf{x}; \boldsymbol{\theta}) - \phi_t^{(r)}(\mathbf{y}; \boldsymbol{\theta})||_1 \le \alpha^r \frac{J+1}{\varepsilon} ||\mathbf{x} - \mathbf{y}||_1$$
 (5)

for all  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$ . Finally,  $||\mathbf{z}||_1 \leq \sqrt{J}||\mathbf{z}||_2$  for all  $\mathbf{z} \in \mathbb{R}^J$  implies that

$$\alpha^r \frac{J+1}{\varepsilon} ||\mathbf{x} - \mathbf{y}||_1 \le \alpha^r \frac{J+1}{\varepsilon} \sqrt{J} ||\mathbf{x} - \mathbf{y}||_2$$
 (6)

for all  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$ . Altogether, Equation (4)-(6) show that

$$||\phi_t^{(r)}(\mathbf{x}; \boldsymbol{\theta}) - \phi_t^{(r)}(\mathbf{y}; \boldsymbol{\theta})||_2 \le \alpha^r \frac{J+1}{\varepsilon} \sqrt{J} ||\mathbf{x} - \mathbf{y}||_2$$

for all  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$  which concludes the proof.

Condition (ii) and (iii) follow straightforwardly from Lemma A.2.

Moreover,  $(\hat{\pi}_{t|t}(\boldsymbol{\theta}))_{t\in\mathbb{N}}$  is a stochastic process taking values in  $(\mathcal{S}, ||\cdot||_2)$  given by

$$\hat{\boldsymbol{\pi}}_{t|t}(\boldsymbol{\theta}) = \hat{\boldsymbol{\phi}}_t(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}),$$

where  $(\hat{\phi}_t(\cdot; \boldsymbol{\theta}))_{t \in \mathbb{N}}$  is a sequence of non-stationary Lipschitz functions given by

$$\hat{\phi}_t(\mathbf{s}; \boldsymbol{\theta}) = \hat{\mathbf{F}}_t(\mathbf{P}'\mathbf{s}; \boldsymbol{\theta})\mathbf{P}'\mathbf{s}.$$

The fact that also  $\sup_{\theta \in \Theta} ||\hat{\pi}_{t|t}(\theta) - \pi_{t|t}(\theta)||_2 \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  follows if

- (i)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_2] < \infty$ ,
- (ii) there exists an  $\mathbf{s} \in \mathcal{S}$  such that  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\hat{\phi}_t(\mathbf{s}; \boldsymbol{\theta}) \phi_t(\mathbf{s}; \boldsymbol{\theta})||_2 \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$ , and
- (iii)  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \Lambda(\hat{\boldsymbol{\phi}}_t \boldsymbol{\phi}_t; \boldsymbol{\theta}) \stackrel{e.a.s.}{\rightarrow} 0 \text{ as } t \to \infty$

by using similar arguments as in the proof of Theorem 2.10 in Straumann and Mikosch (2006). As above, Condition (i) is trivial.

We have that

$$\begin{aligned} & |[\hat{\phi}_{t}(\mathbf{s};\boldsymbol{\theta}) - \phi_{t}(\mathbf{s};\boldsymbol{\theta})]_{j}| \\ & = \left| \frac{\sum_{i=1}^{J} f_{j}(Y_{t}; \hat{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) p_{ij} s_{i}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}} - \frac{\sum_{i=1}^{J} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) p_{ij} s_{i}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; X_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}} \right| \end{aligned}$$

for all  $j \in \{1, ..., J\}$ . Let  $m_{1,t}^{\theta} : \mathcal{X}_1 \times \cdots \times \mathcal{X}_J \to \mathbb{R}$  be given by

$$m_{1,t}^{\theta}(\mathbf{x}) = \frac{\sum_{i=1}^{J} f_j(Y_t; x_j, \mathbf{v}_j) p_{ij} s_i}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_k(Y_t; x_k, \mathbf{v}_k) p_{lk} s_l}.$$

An application of the mean value theorem and the Cauchy–Schwarz inequality shows that there exists an  $\bar{\mathbf{X}}_t(\boldsymbol{v}) \in \{c\hat{\mathbf{X}}_t(\boldsymbol{v}) + (1-c)\mathbf{X}_t(\boldsymbol{v}) : c \in [0,1]\}$  such that

$$|m_{1,t}^{\theta}(\hat{\mathbf{X}}_{t}(v)) - m_{1,t}^{\theta}(\mathbf{X}_{t}(v))| \le ||\nabla_{\mathbf{x}} m_{1,t}^{\theta}(\bar{\mathbf{X}}_{t}(v))||_{2}||\hat{\mathbf{X}}_{t}(v) - \mathbf{X}_{t}(v)||_{2},$$

where, if  $m \neq j$ ,

$$\begin{split} &|[\nabla_{\mathbf{x}} m_{1,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_{t}(\boldsymbol{v}))]_{m}|\\ &\leq \frac{(\sum_{i=1}^{J} f_{j}(Y_{t}; \bar{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})p_{ij}s_{i})(\sum_{l=1}^{J} |\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})|p_{lm}s_{l})}{(\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})p_{lk}s_{l})^{2}} \\ &\leq \frac{f_{j}(Y_{t}; \bar{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})|}{(\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})p_{lk}s_{l})^{2}} \\ &\leq C\frac{f_{j}(Y_{t}; \bar{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})|}{(\sum_{k=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}))^{2}} \\ &\leq C|\nabla_{x_{m}} \log f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})| \end{split}$$

and, if m = j,

$$\begin{split} &|[\nabla_{\mathbf{x}} m_{1,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_{t}(\boldsymbol{\upsilon}))]_{m}|\\ &\leq \frac{\sum_{i=1}^{J} |\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})| p_{im} s_{i}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}) p_{lk} s_{l}}\\ &+ \frac{(\sum_{i=1}^{J} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m}) p_{im} s_{i})(\sum_{l=1}^{J} |\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})| p_{lm} s_{l})}{(\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}) p_{lk} s_{l})^{2}}\\ &\leq \frac{|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}) p_{lk} s_{l}}\\ &+ \frac{f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|}{(\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}) p_{lk} s_{l})^{2}}\\ &\leq C \frac{|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|}{\sum_{k=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k})}\\ &+ C \frac{f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|}{(\sum_{k=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}))^{2}}\\ &\leq C |\nabla_{x_{m}} \log f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})| \\ &\leq C |\nabla_{x_{m}} \log f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})| \end{aligned}$$

so

$$||\nabla_{\mathbf{x}} m_{1,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_t(\boldsymbol{v}))||_2 \leq C \sum_{m=1}^{J} |\nabla_{x_m} \log f_m(Y_t; \bar{X}_{m,t}(\boldsymbol{v}_m), \boldsymbol{v}_m)|.$$

Hence, we have that

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |[\hat{\boldsymbol{\phi}}_t(\mathbf{s}; \boldsymbol{\theta}) - \boldsymbol{\phi}_t(\mathbf{s}; \boldsymbol{\theta})]_j|$$

$$\leq C \sum_{m,n=1}^{J} \sup_{\boldsymbol{v}_m \in \boldsymbol{\Upsilon}_m} \sup_{x_m \in \mathcal{X}_m} |\nabla_{x_m} \log f_m(Y_t; x_m, \boldsymbol{v}_m)| \sup_{\boldsymbol{v}_n \in \boldsymbol{\Upsilon}_n} |\hat{X}_{n,t}(\boldsymbol{v}_n) - X_{n,t}(\boldsymbol{v}_n)|$$

for all  $j \in \{1,...,J\}$ . Condition (ii) then follows from Lemma 2.1 and 2.2 in Straumann and Mikosch (2006) since  $(Y_t)_{t\in\mathbb{Z}}$  is stationary by Assumption 1, for each  $j \in \{1,...,J\}$ , there exists an  $m_j > 0$  such that  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left|\nabla_{x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j)\right|^{m_j}\right] < \infty$  by assumption, and, for each  $j \in \{1,...,J\}$ ,  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left|\hat{X}_{j,t}(\boldsymbol{v}_j) - X_{j,t}(\boldsymbol{v}_j)\right| \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  by Lemma 1.

Moreover, we have that

$$\sup_{\substack{\mathbf{x}, \mathbf{y} \in \mathcal{S} \\ \mathbf{x} \neq \mathbf{y}}} \frac{|[(\hat{\phi}_t(\mathbf{x}; \boldsymbol{\theta}) - \phi_t(\mathbf{x}; \boldsymbol{\theta})) - (\hat{\phi}_t(\mathbf{y}; \boldsymbol{\theta}) - \phi_t(\mathbf{y}; \boldsymbol{\theta}))]_j|}{||\mathbf{x} - \mathbf{y}||_2} \le C \sup_{\mathbf{s} \in \mathcal{S}} ||\nabla_{\mathbf{s}}[\hat{\phi}_t(\mathbf{s}; \boldsymbol{\theta})]_j - \nabla_{\mathbf{s}}[\phi_t(\mathbf{s}; \boldsymbol{\theta})]_j||_2$$

for all  $j \in \{1, ..., J\}$  where

$$\begin{split} & | [\nabla_{\mathbf{s}} [\hat{\phi}_{t}(\mathbf{s}; \boldsymbol{\theta})]_{j} - \nabla_{\mathbf{s}} [\phi_{t}(\mathbf{s}; \boldsymbol{\theta})]_{j}]_{r} | \\ & \leq \left| \frac{f_{j}(Y_{t}; \hat{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) p_{rj}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}} - \frac{f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) p_{rj}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}} \right| \\ & + C \left| \frac{\sum_{i=1}^{J} f_{j}(Y_{t}; \hat{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) p_{ij} s_{i}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}} - \frac{\sum_{k=1}^{J} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) p_{ij} s_{i}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}} \right| \\ & + C \left| \frac{\sum_{k=1}^{J} f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{rk}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}} \right| \\ & + C \left| \frac{\sum_{k=1}^{J} f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; X_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}} \right| \end{aligned}$$

for all  $r \in \{1, ..., J\}$  since  $\hat{a}\hat{b} - ab = (\hat{a} - a)(\hat{b} - b) + (\hat{a} - a)b + a(\hat{b} - b)$ . First, let  $m_{2,t}^{\theta}$ :  $\mathcal{X}_1 \times \cdots \times \mathcal{X}_J \to \mathbb{R}$  be given by

$$m_{2,t}^{\theta}(\mathbf{x}) = \frac{f_j(Y_t; x_j, \mathbf{v}_j) p_{rj}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_k(Y_t; x_k, \mathbf{v}_k) p_{lk} s_l}.$$

Then, there exists an  $\bar{\mathbf{X}}_t(\boldsymbol{v}) \in \{c\hat{\mathbf{X}}_t(\boldsymbol{v}) + (1-c)\mathbf{X}_t(\boldsymbol{v}) : c \in [0,1]\}$  (not necessarily equal to the one above) such that

$$|m_{2,t}^{\boldsymbol{\theta}}(\hat{\mathbf{X}}_t(\boldsymbol{v})) - m_{2,t}^{\boldsymbol{\theta}}(\mathbf{X}_t(\boldsymbol{v}))| \leq ||\nabla_{\mathbf{x}} m_{2,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_t(\boldsymbol{v}))||_2 ||\hat{\mathbf{X}}_t(\boldsymbol{v}) - \mathbf{X}_t(\boldsymbol{v})||_2,$$

where, if  $m \neq j$ ,

$$\begin{split} &|[\nabla_{\mathbf{x}} m_{2,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_{t}(\boldsymbol{v}))]_{m}|\\ &\leq \frac{f_{j}(Y_{t}; \bar{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})p_{rj}(\sum_{l=1}^{J}|\nabla_{X_{m}}f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})|p_{lm}s_{l})}{(\sum_{k=1}^{J}\sum_{l=1}^{J}f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})p_{lk}s_{l})^{2}} \\ &\leq \frac{f_{j}(Y_{t}; \bar{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})|\nabla_{X_{m}}f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})|}{(\sum_{k=1}^{J}\sum_{l=1}^{J}f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})p_{lk}s_{l})^{2}} \end{split}$$

$$||\nabla_{\mathbf{s}}^R f(\mathbf{s})||_2 \le C||\nabla_{\mathbf{s}}^U f(\mathbf{s})||_2$$

where  $[\nabla_{\mathbf{s}}^{U} f(\mathbf{s})]_{j} := \nabla_{s_{j}} f(\mathbf{s}), j \in \{1, ..., J\}.$ 

<sup>&</sup>lt;sup>6</sup>Note that for a function  $f: \mathcal{S} \to \mathbb{R}$  where  $s_J = 1 - \sum_{j=1}^{J-1} s_j$ ,  $[\nabla_{\mathbf{s}}^R f(\mathbf{s})]_j := \nabla_{s_J} f(\mathbf{s}) - \nabla_{s_J} f(\mathbf{s})$ ,  $j \in \{1, ..., J-1\}$  satisfies

$$\leq C \frac{f_j(Y_t; \bar{X}_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j) |\nabla_{x_m} f_m(Y_t; \bar{X}_{m,t}(\boldsymbol{v}_m), \boldsymbol{v}_m)|}{(\sum_{k=1}^J f_k(Y_t; \bar{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k))^2}$$
  
$$\leq C |\nabla_{x_m} \log f_m(Y_t; \bar{X}_{m,t}(\boldsymbol{v}_m), \boldsymbol{v}_m)|$$

and, if m = j,

$$\begin{split} &|[\nabla_{\mathbf{x}} m_{2,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_{t}(\boldsymbol{\upsilon}))]_{m}| \\ &\leq \frac{|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})| p_{rm}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}) p_{lk} s_{l}} \\ &+ \frac{f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m}) p_{rm}(\sum_{l=1}^{J} |\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})| p_{lm} s_{l})}{(\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}) p_{lk} s_{l})^{2}} \\ &\leq \frac{|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}) p_{lk} s_{l}} \\ &+ \frac{f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|}{(\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}) p_{lk} s_{l})^{2}} \\ &\leq C \frac{|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|}{\sum_{k=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k})} \\ &+ C \frac{f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})|}{(\sum_{k=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{\upsilon}_{k}), \boldsymbol{\upsilon}_{k}))^{2}} \\ &\leq C |\nabla_{x_{m}} \log f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{\upsilon}_{m}), \boldsymbol{\upsilon}_{m})| \end{aligned}$$

so

$$||\nabla_{\mathbf{x}} m_{2,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_t(\boldsymbol{v}))||_2 \leq C \sum_{m=1}^J |\nabla_{x_m} \log f_m(Y_t; \bar{X}_{m,t}(\boldsymbol{v}_m), \boldsymbol{v}_m)|.$$

Second, let  $m_{3,t}^{\theta}: \mathcal{X}_1 \times \cdots \times \mathcal{X}_J \to \mathbb{R}$  be given by

$$m_{3,t}^{\theta}(\mathbf{x}) = \frac{\sum_{k=1}^{J} f_k(Y_t; x_k, \mathbf{v}_k) p_{rk}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_k(Y_t; x_k, \mathbf{v}_k) p_{lk} s_l}.$$

Again, there exists an  $\bar{\mathbf{X}}_t(v) \in \{c\hat{\mathbf{X}}_t(v) + (1-c)\mathbf{X}_t(v) : c \in [0,1]\}$  (not necessarily equal to the ones above) such that

$$|m_{3,t}^{\boldsymbol{\theta}}(\hat{\mathbf{X}}_t(\boldsymbol{v})) - m_{3,t}^{\boldsymbol{\theta}}(\mathbf{X}_t(\boldsymbol{v}))| \leq ||\nabla_{\mathbf{X}}m_{3,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_t(\boldsymbol{v}))||_2 ||\hat{\mathbf{X}}_t(\boldsymbol{v}) - \mathbf{X}_t(\boldsymbol{v})||_2,$$

where

$$\begin{split} &|[\nabla_{\mathbf{x}} m_{3,t}^{\theta}(\bar{\mathbf{X}}_{t}(\boldsymbol{v}))]_{m}|\\ &\leq \frac{|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})| p_{rm}}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}}\\ &+ \frac{(\sum_{k=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{rk}) (\sum_{l=1}^{J} |\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})| p_{lm} s_{l})}{(\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l})^{2}}\\ &\leq \frac{|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})|}{\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l}}\\ &+ \frac{(\sum_{k=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}))|\nabla_{x_{m}} f_{m}(Y_{t}; \bar{X}_{m,t}(\boldsymbol{v}_{m}), \boldsymbol{v}_{m})|}{(\sum_{k=1}^{J} \sum_{l=1}^{J} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) p_{lk} s_{l})^{2}} \end{split}$$

$$\leq C \frac{|\nabla_{x_m} f_m(Y_t; \bar{X}_{m,t}(\boldsymbol{v}_m), \boldsymbol{v}_m)|}{\sum_{k=1}^J f_k(Y_t; \bar{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k)}$$

$$+ C \frac{(\sum_{k=1}^J f_k(Y_t; \bar{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k))|\nabla_{x_m} f_m(Y_t; \bar{X}_{m,t}(\boldsymbol{v}_m), \boldsymbol{v}_m)|}{(\sum_{k=1}^J f_k(Y_t; \bar{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k))^2}$$

$$\leq C |\nabla_{x_m} \log f_m(Y_t; \bar{X}_{m,t}(\boldsymbol{v}_m), \boldsymbol{v}_m)|$$

so

$$||\nabla_{\mathbf{x}} m_{3,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_t(\boldsymbol{v}))||_2 \leq C \sum_{m=1}^J |\nabla_{x_m} \log f_m(Y_t; \bar{X}_{m,t}(\boldsymbol{v}_m), \boldsymbol{v}_m)|.$$

Hence, we have that

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \sup_{\mathbf{s} \in \mathcal{S}} |[\nabla_{\mathbf{s}} [\hat{\boldsymbol{\phi}}_t(\mathbf{s}; \boldsymbol{\theta})]_j - \nabla_{\mathbf{s}} [\boldsymbol{\phi}_t(\mathbf{s}; \boldsymbol{\theta})]_j]_r|$$

$$\leq C \sum_{m,n=1}^J \sup_{\boldsymbol{v}_m \in \boldsymbol{\Upsilon}_m} \sup_{x_m \in \mathcal{X}_m} |\nabla_{x_m} \log f_m(Y_t; x_m, \boldsymbol{v}_m)| \sup_{\boldsymbol{v}_n \in \boldsymbol{\Upsilon}_n} |\hat{X}_{n,t}(\boldsymbol{v}_n) - X_{n,t}(\boldsymbol{v}_n)|$$

for all  $j \in \{1, ..., J\}$  and  $r \in \{1, ..., J\}$ . Condition (iii) then follows by using the same arguments as above.

#### A.3 Proof of Theorem 2

First, recall that  $\hat{L}_T(\boldsymbol{\theta})$  is given by  $\hat{L}_T(\boldsymbol{\theta}) = \frac{1}{T} \sum_{t=1}^T \log \hat{f}(Y_t; \boldsymbol{\theta})$  where  $\hat{f}(Y_t; \boldsymbol{\theta}) = \sum_{j=1}^J \hat{\pi}_{j,t|t-1}(\boldsymbol{\theta}) f_j(Y_t; \hat{X}_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)$ . Moreover, let  $L_T(\boldsymbol{\theta})$  be given by  $L_T(\boldsymbol{\theta}) = \frac{1}{T} \sum_{t=1}^T \log f(Y_t; \boldsymbol{\theta})$  where  $f(Y_t; \boldsymbol{\theta}) = \sum_{j=1}^J \pi_{j,t|t-1}(\boldsymbol{\theta}) f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)$  and let  $L(\boldsymbol{\theta})$  be given by  $L(\boldsymbol{\theta}) = \mathbb{E}[\log f(Y_t; \boldsymbol{\theta})]$ .

The conclusion follows from Lemma 3.1 and 4.1 in Pötscher and Prucha (1997) if

- (i)  $\Theta$  is compact,
- (ii)  $\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}|\hat{L}_T(\boldsymbol{\theta})-L(\boldsymbol{\theta})| \stackrel{a.s.}{\to} 0 \text{ as } T\to\infty,$
- (iii)  $\boldsymbol{\theta} \mapsto L(\boldsymbol{\theta})$  is continuous, and
- (iv)  $L(\boldsymbol{\theta}) \leq L(\boldsymbol{\theta}_0)$  for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  with equality if and only if  $\boldsymbol{\theta} = \boldsymbol{\theta}_0$ .

Condition (i) follows from Assumption 2. Condition (ii) follows from Lemma A.3 and A.4 since

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\hat{L}_T(\boldsymbol{\theta}) - L(\boldsymbol{\theta})| \le \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\hat{L}_T(\boldsymbol{\theta}) - L_T(\boldsymbol{\theta})| + \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |L_T(\boldsymbol{\theta}) - L(\boldsymbol{\theta})|,$$

and Condition (iii) is a by-product of the law of large numbers used in the proof of Lemma A.4.

**Lemma A.3.** Under the assumptions in Theorem 2,

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\hat{L}_T(\boldsymbol{\theta}) - L_T(\boldsymbol{\theta})| \stackrel{a.s.}{\to} 0 \quad as \quad T \to \infty.$$

*Proof.* We have that

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\hat{L}_T(\boldsymbol{\theta}) - L_T(\boldsymbol{\theta})| \le \frac{1}{T} \sum_{t=1}^T \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\log \hat{f}(Y_t; \boldsymbol{\theta}) - \log f(Y_t; \boldsymbol{\theta})|.$$

The conclusion thus follows from Lemma 2.1 in Straumann and Mikosch (2006) if  $\sup_{\theta \in \Theta} |\log \hat{f}(Y_t; \theta) - \log f(Y_t; \theta)| \stackrel{e.a.s.}{\longrightarrow} 0$  as  $t \to \infty$ .

We have that

$$|\log \hat{f}(Y_t; \boldsymbol{\theta}) - \log f(Y_t; \boldsymbol{\theta})| \le |\log \hat{f}(Y_t; \boldsymbol{\theta}) - \log \tilde{f}(Y_t; \boldsymbol{\theta})| + |\log \tilde{f}(Y_t; \boldsymbol{\theta}) - \log f(Y_t; \boldsymbol{\theta})|,$$

where  $\tilde{f}(Y_t; \boldsymbol{\theta}) = \sum_{j=1}^J \pi_{j,t|t-1}(\boldsymbol{\theta}) f_j(Y_t; \hat{X}_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)$ . First, let  $m_{1,t}^{\boldsymbol{\theta}} : (0, \infty)^J \to \mathbb{R}$  be given by

$$m_{1,t}^{\boldsymbol{\theta}}(\mathbf{s}) = \log \sum_{j=1}^{J} s_j f_j(Y_t; \hat{X}_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j).$$

An application of the mean value theorem and the Cauchy-Schwarz inequality shows that there exists a  $\bar{\pi}_{t|t-1}(\theta) \in \{c\hat{\pi}_{t|t-1}(\theta) + (1-c)\pi_{t|t-1}(\theta) : c \in [0,1]\}$  such that

$$|m_{1,t}^{\theta}(\hat{\pi}_{t|t-1}(\theta)) - m_{1,t}^{\theta}(\pi_{t|t-1}(\theta))| \leq ||\nabla_{\mathbf{s}} m_{1,t}^{\theta}(\bar{\pi}_{t|t-1}(\theta))||_2 ||\hat{\pi}_{t|t-1}(\theta) - \pi_{t|t-1}(\theta)||_2,$$

where

$$\begin{split} |[\nabla_{\mathbf{s}} m_{1,t}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\pi}}_{t|t-1}(\boldsymbol{\theta}))]_{k}| &= \frac{f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})}{\sum_{j=1}^{J} \bar{\pi}_{j,t|t-1}(\boldsymbol{\theta}) f_{j}(Y_{t}; \hat{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})} \\ &= \frac{1}{\bar{\pi}_{k,t|t-1}(\boldsymbol{\theta})} \frac{\bar{\pi}_{k,t|t-1}(\boldsymbol{\theta}) f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})}{\sum_{j=1}^{J} \bar{\pi}_{j,t|t-1}(\boldsymbol{\theta}) f_{j}(Y_{t}; \hat{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})} \\ &\leq C \end{split}$$

by Remark 1 so

$$||\nabla_{\mathbf{s}} m_{1,t}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\pi}}_{t|t-1}(\boldsymbol{\theta}))||_2 \leq C.$$

Moreover, let  $m_{2,t}^{\theta}: \mathcal{X}_1 \times \cdots \times \mathcal{X}_J \to \mathbb{R}$  be given by

$$m_{2,t}^{\boldsymbol{\theta}}(\mathbf{x}) = \log \sum_{j=1}^{J} \pi_{j,t|t-1}(\boldsymbol{\theta}) f_j(Y_t; x_j, \boldsymbol{v}_j).$$

Another application of the mean value theorem and the Cauchy-Schwarz inequality shows that there exists an  $\bar{\mathbf{X}}_t(\boldsymbol{v}) \in \{c\hat{\mathbf{X}}_t(\boldsymbol{v}) + (1-c)\mathbf{X}_t(\boldsymbol{v}) : c \in [0,1]\}$  such that

$$|m_{2,t}^{\boldsymbol{\theta}}(\hat{\mathbf{X}}_t(\boldsymbol{v})) - m_{2,t}^{\boldsymbol{\theta}}(\mathbf{X}_t(\boldsymbol{v}))| \leq ||\nabla_{\mathbf{x}} m_{2,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_t(\boldsymbol{v}))||_2 ||\hat{\mathbf{X}}_t(\boldsymbol{v}) - \mathbf{X}_t(\boldsymbol{v})||_2,$$

where

$$\begin{split} |[\nabla_{\mathbf{x}} m_{2,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_{t}(\boldsymbol{v}))]_{k}| &= \frac{\pi_{k,t|t-1}(\boldsymbol{\theta})|\nabla_{x_{k}} f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})|}{\sum_{j=1}^{J} \pi_{j,t|t-1}(\boldsymbol{\theta}) f_{j}(Y_{t}; \bar{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})} \\ &= \frac{\pi_{k,t|t-1}(\boldsymbol{\theta}) f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})}{\sum_{j=1}^{J} \pi_{j,t|t-1}(\boldsymbol{\theta}) f_{j}(Y_{t}; \bar{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})} |\nabla_{x_{k}} \log f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})| \\ &\leq |\nabla_{x_{k}} \log f_{k}(Y_{t}; \bar{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k})| \end{split}$$

so

$$||\nabla_{\mathbf{x}} m_{2,t}^{\boldsymbol{\theta}}(\bar{\mathbf{X}}_t(\boldsymbol{v}))||_2 \leq \sum_{k=1}^{J} |\nabla_{x_k} \log f_k(Y_t; \bar{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k)|$$
.

Hence, we have that

$$|\log \hat{f}_{t|t-1}(Y_t; \boldsymbol{\theta}) - \log f_{t|t-1}(Y_t; \boldsymbol{\theta})| \leq C||\hat{\boldsymbol{\pi}}_{t|t-1}(\boldsymbol{\theta}) - \boldsymbol{\pi}_{t|t-1}(\boldsymbol{\theta})||_2$$

$$+ \sum_{k,l=1}^{J} |\nabla_{x_k} \log f_k(Y_t; \bar{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k)||\hat{X}_{l,t}(\boldsymbol{v}_l) - X_{l,t}(\boldsymbol{v}_l)|.$$

It thus follows from Lemma 2.1 and 2.2 in Straumann and Mikosch (2006) that  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\log \hat{f}(Y_t; \boldsymbol{\theta}) - \log f(Y_t; \boldsymbol{\theta})| \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  since  $(Y_t)_{t \in \mathbb{Z}}$  is stationary by Assumption 1, for each  $j \in \{1, ..., J\}$ , there exists an  $m_j > 0$  such that  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} |\nabla_{x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j)|^{m_j}\right] < \infty$  by assumption, for each  $j \in \{1, ..., J\}$ ,  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\hat{X}_{j,t}(\boldsymbol{v}_j) - X_{j,t}(\boldsymbol{v}_j)| \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  by Lemma 1, and  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\hat{\boldsymbol{\pi}}_{t|t-1}(\boldsymbol{\theta}) - \boldsymbol{\pi}_{t|t-1}(\boldsymbol{\theta})||_2 \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  by Corollary 1.

**Lemma A.4.** Under the assumptions in Theorem 2,

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |L_T(\boldsymbol{\theta}) - L(\boldsymbol{\theta})| \stackrel{a.s.}{\to} 0 \quad as \quad T \to \infty.$$

*Proof.* The conclusion follows from the uniform law of large numbers by Rao (1962) if  $(\log f(Y_t; \boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  and  $\mathbb{E}\left[\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\log f(Y_t; \boldsymbol{\theta})|\right] < \infty$ .

First,  $(\log f(Y_t; \boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  since  $(Y_t)_{t \in \mathbb{Z}}$  is stationary and ergodic by Assumption 1, for each  $j \in \{1, ..., J\}$ ,  $(X_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j$  by Lemma 1, and  $(\boldsymbol{\pi}_{t|t-1}(\boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  by Corollary 1.

Note that

$$|\log f(Y_t; \boldsymbol{\theta})| \leq \sum_{j=1}^{J} |\log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)|$$

since

$$\begin{split} \log f(Y_t; \boldsymbol{\theta}) &\geq \log \min_{j \in \{1, ..., J\}} f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j) \\ &= \min_{j \in \{1, ..., J\}} \log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j) \\ &\geq \min_{j \in \{1, ..., J\}} -|\log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)| \geq -\sum_{j=1}^J |\log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)| \end{split}$$

and

$$\begin{split} \log f(Y_t; \boldsymbol{\theta}) &\leq \log \max_{j \in \{1, \dots, J\}} f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j) \\ &= \max_{j \in \{1, \dots, J\}} \log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j) \\ &\leq \max_{j \in \{1, \dots, J\}} |\log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)| \leq \sum_{j=1}^J |\log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)|. \end{split}$$

Hence,  $\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}|\log f(Y_t;\boldsymbol{\theta})|\right]<\infty$  since, for each  $j\in\{1,...,J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}|\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)|\right]<\infty$  by Assumption 5.

Finally, Condition (iv) follows from Lemma A.5.

**Lemma A.5.** Under the assumptions in Theorem 2,

$$L(\boldsymbol{\theta}) \leq L(\boldsymbol{\theta}_0)$$

for all  $\theta \in \Theta$  with equality if and only if  $\theta = \theta_0$ .

*Proof.* We have that

$$L(\boldsymbol{\theta}) - L(\boldsymbol{\theta}_0) = \mathbb{E}\left[\log \frac{f(Y_t; \boldsymbol{\theta})}{f(Y_t; \boldsymbol{\theta}_0)}\right].$$

Observe that

$$\mathbb{E}\left[\log \frac{f^{(m)}(Y_t, ..., Y_{t-m+1}; \boldsymbol{\theta})}{f^{(m)}(Y_t, ..., Y_{t-m+1}; \boldsymbol{\theta})}\right] = \mathbb{E}\left[\log \prod_{i=1}^m \frac{f(Y_{t-i+1}; \boldsymbol{\theta})}{f(Y_{t-i+1}; \boldsymbol{\theta}_0)}\right]$$
$$= \sum_{i=1}^m \mathbb{E}\left[\log \frac{f(Y_{t-i+1}; \boldsymbol{\theta})}{f(Y_{t-i+1}; \boldsymbol{\theta}_0)}\right] = m\mathbb{E}\left[\log \frac{f(Y_t; \boldsymbol{\theta})}{f(Y_t; \boldsymbol{\theta}_0)}\right],$$

SO

$$\mathbb{E}\left[\log\frac{f(Y_t;\boldsymbol{\theta})}{f(Y_t;\boldsymbol{\theta}_0)}\right] = \frac{1}{m}\mathbb{E}\left[\log\frac{f^{(m)}(Y_t,...,Y_{t-m+1};\boldsymbol{\theta})}{f^{(m)}(Y_t,...,Y_{t-m+1};\boldsymbol{\theta}_0)}\right].$$

Note that  $\log x \le x - 1$  for all x > 0 with equality if and only if x = 1. Thus,

$$\mathbb{E}\left[\log \frac{f^{(m)}(Y_t, ..., Y_{t-m+1}; \boldsymbol{\theta})}{f^{(m)}(Y_t, ..., Y_{t-m+1}; \boldsymbol{\theta}_0)}\right] \le 0$$

for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  with equality if and only if  $f^{(m)}(Y_t,...,Y_{t-m+1};\boldsymbol{\theta}) = f^{(m)}(Y_t,...,Y_{t-m+1};\boldsymbol{\theta}_0)$  a.s. Hence, we have that

$$L(\boldsymbol{\theta}) - L(\boldsymbol{\theta}_0) \le 0$$

for all  $\theta \in \Theta$  with equality if and only if  $\theta = \theta_0$  by Assumption 6.

#### A.4 Proof of Theorem 3

The conclusion follows from Lemma 8.1 in Pötscher and Prucha (1997) if

- (i)  $\boldsymbol{\theta}_0 \in \operatorname{int}(\boldsymbol{\Theta})$ ,
- (ii)  $\hat{\boldsymbol{\theta}}_T \stackrel{\mathbb{P}}{\to} \boldsymbol{\theta}_0 \text{ as } T \to \infty,$
- (iii)  $\theta \mapsto \hat{L}_T(\theta)$  is twice continuously differentiable a.s.,
- (iv)  $\sqrt{T}\nabla_{\boldsymbol{\theta}}\hat{L}_T(\boldsymbol{\theta}_0) \stackrel{d}{\to} \mathcal{N}(\mathbf{0}, \mathbf{I}(\boldsymbol{\theta}_0))$  as  $T \to \infty$ ,
- (v)  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{L}_T(\boldsymbol{\theta}) (-\mathbf{I}(\boldsymbol{\theta}))||_{2,2} \stackrel{\mathbb{P}}{\to} 0 \text{ as } T \to \infty, \text{ and}$
- (vi)  $\mathbf{I}(\boldsymbol{\theta}_0)$  is invertible.

First, Condition (i) follows from Assumption 7, Condition (ii) from Theorem 2, and Condition (iii) follows from Assumption 8 and 9. Moreover, Condition (iv) follows from Lemma A.6, A.7, and A.8.

**Lemma A.6.** Under the assumptions in Theorem 3,

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\sqrt{T} \nabla_{\boldsymbol{\theta}} \hat{L}_T(\boldsymbol{\theta}) - \sqrt{T} \nabla_{\boldsymbol{\theta}} L_T(\boldsymbol{\theta})||_2 \stackrel{a.s.}{\to} 0 \quad as \quad T \to \infty.$$

*Proof.* We have that

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\sqrt{T} \nabla_{\boldsymbol{\theta}} \hat{L}_T(\boldsymbol{\theta}) - \sqrt{T} \nabla_{\boldsymbol{\theta}} L_T(\boldsymbol{\theta})||_2 \le \frac{1}{\sqrt{T}} \sum_{t=1}^T \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_t}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}} f_t}{f_t} \right| \right|_2,$$

where  $\hat{f}_t$  denotes  $\hat{f}(Y_t; \boldsymbol{\theta})$  and  $f_t$  denotes  $f(Y_t; \boldsymbol{\theta})$  and that

$$\begin{split} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{t}}{f_{t}} \right| \right|_{2} &\leq \sum_{j=1}^{J} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} f_{j,t}}{f_{t}} \right| \right|_{2} \\ &+ \sum_{j=1}^{J} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\pi_{j,t|t-1} \nabla_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right| \right|_{2}, \end{split}$$

where  $\hat{\pi}_{j,t|t-1}$  denotes  $\hat{\pi}_{j,t|t-1}(\boldsymbol{\theta})$ ,  $\hat{f}_{j,t}$  denotes  $f_j(Y_t; \hat{X}_{j,t}, \boldsymbol{v}_j)$ ,  $\hat{X}_{j,t}$  denotes  $\hat{X}_{j,t}(\boldsymbol{v}_j)$ ,  $\pi_{j,t|t-1}$  denotes  $\pi_{j,t|t-1}(\boldsymbol{\theta})$ ,  $f_{j,t}$  denotes  $f_j(Y_t; X_{j,t}, \boldsymbol{v}_j)$ , and  $X_{j,t}$  denotes  $X_{j,t}(\boldsymbol{v}_j)$ .

As in the proof of Lemma A.3, the conclusion thus follows from Lemma 2.1 in Straumann and Mikosch (2006) if  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_t}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}} f_t}{f_t} \right| \right|_2 \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$ . This follows if, for each  $j \in \{1, ..., J\}$ ,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} f_{j,t}}{f_t} \right| \right|_2 \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  and  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\pi_{j,t|t-1} \nabla_{\boldsymbol{\theta}} f_{j,t}}{f_t} \right| \right|_2 \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$ . First.

$$\begin{aligned} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} f_{j,t}}{f_{t}} \right| \right|_{2} &\leq \left| \left| \nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} - \nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \right| \right|_{2} \left| \frac{\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{f_{j,t}}{f_{t}} \right| \\ &+ C \left| \left| \nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} - \nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \right| \right|_{2} \\ &+ \left| \left| \nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \right| \right|_{2} \left| \frac{\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{f_{j,t}}{f_{t}} \right|. \end{aligned}$$

We have that

$$\left| \frac{\hat{f}_{j,t}}{\hat{f}_t} - \frac{f_{j,t}}{f_t} \right| \le \left| \frac{\hat{f}_{j,t}}{\hat{f}_t} - \frac{\hat{f}_{j,t}}{\tilde{f}_t} \right| + \left| \frac{\hat{f}_{j,t}}{\tilde{f}_t} - \frac{f_{j,t}}{f_t} \right|,$$

where  $\tilde{f}_t$  denotes  $\tilde{f}(Y_t; \boldsymbol{\theta}) = \sum_{j=1}^J \pi_{j,t|t-1}(\boldsymbol{\theta}) f_j(Y_t; \hat{X}_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)$ . By using the same arguments as in the proof of Lemma A.3, we have that

$$\left| \frac{\hat{f}_{j,t}}{\hat{f}_t} - \frac{f_{j,t}}{f_t} \right| \le C \left| \left| \hat{\boldsymbol{\pi}}_{t|t-1} - \boldsymbol{\pi}_{t|t-1} \right| \right|_2 + C \sum_{k,l=1}^{J} \left| \bar{\nabla}_{x_k} \log \bar{f}_{k,t} \right| |\hat{X}_{l,t} - X_{l,t}|,$$

where  $\bar{f}_{j,t}$  denotes  $f_j(Y_t; \bar{X}_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)$ . It thus follows from Lemma 2.1 and 2.2 in Straumann and Mikosch (2006) that, for each  $j \in \{1, ..., J\}$ ,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} f_{j,t}}{f_t} \right\|_2^{e.a.s.} 0$  as  $t \to \infty$  since, in addition to the arguments used in the proof of Lemma A.3, for each  $j \in \{1, ..., J\}$ ,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \|\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} - \nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1}\|_2 \overset{e.a.s.}{\to} 0$  as

 $t \to \infty$  where  $(\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1})_{t \in \mathbb{Z}}$  is stationary for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  and  $\mathbb{E}\left[\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1}||_2^2\right] < \infty$  by Corollary 2.

Moreover,

$$\left\| \frac{\hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\pi_{j,t|t-1} \nabla_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right\|_{2} \leq \left| \hat{\pi}_{j,t|t-1} - \pi_{j,t|t-1} \right| \left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right\|_{2} + \left| \hat{\pi}_{j,t|t-1} - \pi_{j,t|t-1} \right| \left\| \frac{\nabla_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right\|_{2} + \left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right\|_{2},$$

where

$$\left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right\|_{2} \leq \left\| \frac{\bar{\nabla}_{x_{j}} \hat{f}_{j,t} \nabla_{\boldsymbol{\theta}} \hat{X}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}} f_{j,t} \nabla_{\boldsymbol{\theta}} X_{j,t}}{f_{t}} \right\|_{2} + \left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right\|_{2}.$$

First, we have that

$$\left\| \frac{\bar{\nabla}_{x_{j}}\hat{f}_{j,t}\nabla_{\boldsymbol{\theta}}\hat{X}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}}f_{j,t}\nabla_{\boldsymbol{\theta}}X_{j,t}}{f_{t}} \right\|_{2} \leq \left| \frac{\bar{\nabla}_{x_{j}}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}}f_{j,t}}{f_{t}} \right| \left| \left| \nabla_{\boldsymbol{\theta}}\hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}}X_{j,t} \right| \right|_{2} + \left| \frac{\bar{\nabla}_{x_{j}}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}}f_{j,t}}{f_{t}} \right| \left| \left| \nabla_{\boldsymbol{\theta}}X_{j,t} \right| \right|_{2} + \left| \frac{\bar{\nabla}_{x_{j}}f_{j,t}}{f_{t}} \right| \left| \left| \nabla_{\boldsymbol{\theta}}\hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}}X_{j,t} \right| \right|_{2},$$

where

$$\left| \frac{\bar{\nabla}_{x_j} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{x_j} f_{j,t}}{f_t} \right| \leq \left| \frac{\bar{\nabla}_{x_j} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{x_j} \hat{f}_{j,t}}{\tilde{f}_t} \right| + \left| \frac{\bar{\nabla}_{x_j} \hat{f}_{j,t}}{\tilde{f}_t} - \frac{\bar{\nabla}_{x_j} f_{j,t}}{f_t} \right|.$$

By using the same arguments as in the proof of Lemma A.3 again, we have that

$$\left| \frac{\bar{\nabla}_{x_{j}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}} f_{j,t}}{f_{t}} \right| \leq C \left| \bar{\nabla}_{x_{j}} \log \hat{f}_{j,t} \right| \left| \left| \hat{\pi}_{t|t-1} - \pi_{t|t-1} \right| \right|_{2} 
+ C \sum_{l=1}^{J} \left| \bar{\nabla}_{x_{j}x_{j}} \log \bar{f}_{j,t} \right| \left| \hat{X}_{l,t} - X_{l,t} \right| 
+ C \sum_{k,l=1}^{J} \left| \bar{\nabla}_{x_{j}} \log \bar{f}_{j,t} \right| \left| \bar{\nabla}_{x_{k}} \log \bar{f}_{k,t} \right| \left| \hat{X}_{l,t} - X_{l,t} \right|.$$

Second, we have that

$$\left| \left| \frac{\bar{\nabla}_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}} f_{j,t}}{f_t} \right| \right|_2 \le \left| \left| \frac{\bar{\nabla}_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\tilde{f}_t} \right| \right|_2 + \left| \left| \frac{\bar{\nabla}_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\tilde{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}} f_{j,t}}{f_t} \right| \right|_2.$$

By using the same arguments as in the proof of Lemma A.3 once again, we have that

$$\left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right\|_{2} \leq C \sum_{h=1}^{d_{j}} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}} \log \hat{f}_{j,t} \right| \left| \left| \hat{\boldsymbol{\pi}}_{t|t-1} - \boldsymbol{\pi}_{t|t-1} \right| \right|_{2}$$

$$+ C \sum_{h=1}^{d_{j}} \sum_{l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}x_{j}} \log \bar{f}_{j,t} \right| |\hat{X}_{l,t} - X_{l,t}|$$

$$+ C \sum_{h=1}^{d_{j}} \sum_{k,l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}} \log \bar{f}_{j,t} \right| |\bar{\nabla}_{x_{k}} \log \bar{f}_{k,t}| |\hat{X}_{l,t} - X_{l,t}|.$$

It thus follows from Lemma 2.1 and 2.2 in Straumann and Mikosch (2006) that, for each  $j \in \{1,...,J\}$ , also  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\pi_{j,t|t-1} \nabla_{\boldsymbol{\theta}} f_{j,t}}{f_t} \right\|_{2}^{e.a.s.} 0$  as  $t \to \infty$  since, in addition to the arguments used in the proof of Lemma A.3, for each  $j \in \{1,...,J\}$ ,  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j} \hat{X}_{j,t} - \nabla_{\boldsymbol{v}_j} X_{j,t}||_{2}^{e.a.s.} 0$  as  $t \to \infty$  where  $(\nabla_{\boldsymbol{v}_j} X_{j,t})_{t \in \mathbb{Z}}$  is stationary for all  $\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j$  and  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j} X_{j,t}||_{2}^{2k_j}\right] < \infty$  by Lemma 3. Moreover, for each  $j \in \{1,...,J\}$ , there exists an  $m_j > 0$  such that  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} ||\bar{\nabla}_{\boldsymbol{v}_j} \log f_{j,t}||_{2}^{m_j}\right] < \infty$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} |\bar{\nabla}_{\boldsymbol{v}_j x_j} \log f_{j,t}||_{2}^{m_j}\right] < \infty$ , and  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} ||\bar{\nabla}_{\boldsymbol{v}_j x_j} \log f_{j,t}||_{2}^{m_j}\right] < \infty$  by assumption.

**Lemma A.7.** Under the assumptions in Theorem 3,

$$\sqrt{T}\nabla_{\boldsymbol{\theta}}L_T(\boldsymbol{\theta}_0) \stackrel{d}{\to} \mathcal{N}\left(\mathbf{0}, \mathbb{E}\left[\nabla_{\boldsymbol{\theta}}\log f(Y_t; \boldsymbol{\theta}_0)\nabla_{\boldsymbol{\theta}'}\log f(Y_t; \boldsymbol{\theta}_0)\right]\right) \quad as \quad T \to \infty.$$

*Proof.* The conclusion follows from the central limit theorem by Billingsley (1961) together with the Cramér–Wold theorem if  $(\nabla_{\boldsymbol{\theta}} \log f(Y_t; \boldsymbol{\theta}_0))_{t \in \mathbb{Z}}$  is a stationary and ergodic martingale difference sequence and  $\mathbb{E}\left[||\nabla_{\boldsymbol{\theta}} \log f(Y_t; \boldsymbol{\theta}_0)||_2^2\right] < \infty$ .

First,  $(\nabla_{\boldsymbol{\theta}} \log f(Y_t; \boldsymbol{\theta}_0))_{t \in \mathbb{Z}}$  is a stationary and ergodic martingale difference sequence since, in addition to the arguments used in the proof of Lemma A.4, for each  $j \in \{1, ..., J\}$ ,  $(\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j$  by Lemma 3, for each  $j \in \{1, ..., J\}$ ,  $(\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1}(\boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  by Corollary 2, and

$$\mathbb{E}\left[\nabla_{\boldsymbol{\theta}} \log f(Y_t; \boldsymbol{\theta}_0) \mid \mathbf{Y}_{-\infty}^{t-1}\right] = \int_{\mathcal{Y}} \nabla_{\boldsymbol{\theta}} f(y; \boldsymbol{\theta})|_{\boldsymbol{\theta} = \boldsymbol{\theta}_0} \, \mathrm{d}y = \left. \nabla_{\boldsymbol{\theta}} \int_{\mathcal{Y}} f(y; \boldsymbol{\theta}) \, \mathrm{d}y \right|_{\boldsymbol{\theta} = \boldsymbol{\theta}_0} = \mathbf{0}.$$

We have that

$$||\nabla_{\boldsymbol{\theta}} \log f(Y_t; \boldsymbol{\theta}_0)||_2^2 = \frac{1}{f^2(Y_t; \boldsymbol{\theta}_0)} ||\nabla_{\boldsymbol{\theta}} f(Y_t; \boldsymbol{\theta}_0)||_2^2$$

where

$$\begin{split} &||\nabla_{\boldsymbol{\theta}}f(Y_{t};\boldsymbol{\theta}_{0})||_{2}^{2} \\ &\leq \sum_{i,j=1}^{J} ||\nabla_{\boldsymbol{\theta}}\pi_{i,t|t-1}(\boldsymbol{\theta}_{0})||_{2}f_{i}(Y_{t};X_{i,t}(\boldsymbol{\upsilon}_{i,0}),\boldsymbol{\upsilon}_{i,0})||\nabla_{\boldsymbol{\theta}}\pi_{j,t|t-1}(\boldsymbol{\theta}_{0})||_{2}f_{j}(Y_{t};X_{j,t}(\boldsymbol{\upsilon}_{j,0}),\boldsymbol{\upsilon}_{j,0}) \\ &+ 2\sum_{i,j=1}^{J} ||\nabla_{\boldsymbol{\theta}}\pi_{i,t|t-1}(\boldsymbol{\theta}_{0})||_{2}f_{i}(Y_{t};X_{i,t}(\boldsymbol{\upsilon}_{i,0}),\boldsymbol{\upsilon}_{i,0})\pi_{j,t|t-1}(\boldsymbol{\theta}_{0})||\nabla_{\boldsymbol{\theta}}f_{j}(Y_{t};X_{j,t}(\boldsymbol{\upsilon}_{j,0}),\boldsymbol{\upsilon}_{j,0})||_{2} \\ &+ \sum_{i,j=1}^{J} \pi_{i,t|t-1}(\boldsymbol{\theta}_{0})||\nabla_{\boldsymbol{\theta}}f_{i}(Y_{t};X_{i,t}(\boldsymbol{\upsilon}_{i,0}),\boldsymbol{\upsilon}_{i,0})||_{2}\pi_{j,t|t-1}(\boldsymbol{\theta}_{0})||\nabla_{\boldsymbol{\theta}}f_{j}(Y_{t};X_{j,t}(\boldsymbol{\upsilon}_{j,0}),\boldsymbol{\upsilon}_{j,0})||_{2}, \end{split}$$

so

$$\mathbb{E}\left[||\nabla_{\boldsymbol{\theta}}\log f(Y_t;\boldsymbol{\theta}_0)||_2^2\right]$$

$$\leq C \left( \sum_{j=1}^{J} \left( \mathbb{E} \left[ ||\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1}(\boldsymbol{\theta}_{0})||_{2}^{2} \right] \right)^{1/2} \right)^{2}$$

$$+ C \sum_{i,j=1}^{J} \left( \mathbb{E} \left[ ||\nabla_{\boldsymbol{\theta}} \pi_{i,t|t-1}(\boldsymbol{\theta}_{0})||_{2}^{2} \right] \right)^{1/2} \left( \mathbb{E} \left[ ||\nabla_{\boldsymbol{\theta}} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})||_{2}^{2} \right] \right)^{1/2}$$

$$+ \left( \sum_{j=1}^{J} \left( \mathbb{E} \left[ ||\nabla_{\boldsymbol{\theta}} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})||_{2}^{2} \right] \right)^{1/2} \right)^{2}.$$

Moreover, we have that

$$||\nabla_{\boldsymbol{\theta}} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})||_{2}^{2} = \frac{1}{f_{j}^{2}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})} ||\nabla_{\boldsymbol{\theta}} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})||_{2}^{2}$$

where

$$\begin{split} &||\nabla_{\boldsymbol{\theta}} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})||_{2}^{2} \\ &\leq \left|\bar{\nabla}_{x_{j}} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})\right|^{2} ||\nabla_{\boldsymbol{\theta}} X_{j,t}(\boldsymbol{v}_{j,0})||_{2}^{2} \\ &+ 2\left|\bar{\nabla}_{x_{j}} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})\right| ||\nabla_{\boldsymbol{\theta}} X_{j,t}(\boldsymbol{v}_{j,0})||_{2} ||\bar{\nabla}_{\boldsymbol{\theta}} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})||_{2} \\ &+ ||\bar{\nabla}_{\boldsymbol{\theta}} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})||_{2}^{2} \end{split}$$

SO

$$\mathbb{E}\left[\left|\left|\nabla_{\boldsymbol{\theta}} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})\right|\right|_{2}^{2}\right] \\
\leq \left(\mathbb{E}\left[\left|\bar{\nabla}_{x_{j}} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})\right|^{2k_{j}/(k_{j}-1)}\right]\right)^{(k_{j}-1)/k_{j}} \left(\mathbb{E}\left[\left|\left|\nabla_{\boldsymbol{\theta}} X_{j,t}(\boldsymbol{v}_{j,0})\right|\right|_{2}^{2k_{j}}\right]\right)^{1/k_{j}} \\
+ 2\left(\mathbb{E}\left[\left|\bar{\nabla}_{x_{j}} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})\right|^{2k_{j}/(k_{j}-1)}\right]\right)^{(k_{j}-1)/2k_{j}} \left(\mathbb{E}\left[\left|\left|\nabla_{\boldsymbol{\theta}} X_{j,t}(\boldsymbol{v}_{j,0})\right|\right|_{2}^{2k_{j}}\right]\right)^{1/2k_{j}} \\
\cdot \left(\mathbb{E}\left[\left|\left|\bar{\nabla}_{\boldsymbol{\theta}} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})\right|\right|_{2}^{2}\right]\right)^{1/2} \\
+ \mathbb{E}\left[\left|\left|\bar{\nabla}_{\boldsymbol{\theta}} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j,0}), \boldsymbol{v}_{j,0})\right|\right|_{2}^{2}\right].$$

Hence, 
$$\mathbb{E}\left[||\nabla_{\boldsymbol{\theta}}\log f(Y_t; \boldsymbol{\theta}_0)||_2^2\right] < \infty$$
 since, for each  $j \in \{1, ..., J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j)||_2^{2k_j}\right] < \infty$  by Lemma 3, for each  $j \in \{1, ..., J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1}(\boldsymbol{\theta})||_2^2\right] < \infty$  by Corollary 2, and, for each  $j \in \{1, ..., J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\bar{\nabla}_{x_j}\log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)|^{2k_j/(k_j-1)}\right] < \infty$  and  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\bar{\nabla}_{\boldsymbol{v}_j}\log f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)||_2^2\right] < \infty$  by Assumption 10.

**Lemma A.8.** Under the assumptions in Theorem 3,

$$\mathbf{I}(\boldsymbol{\theta}_0) = \mathbb{E}\left[\nabla_{\boldsymbol{\theta}} \log f(Y_t; \boldsymbol{\theta}_0) \nabla_{\boldsymbol{\theta}'} \log f(Y_t; \boldsymbol{\theta}_0)\right].$$

*Proof.* We have that

$$\mathbb{E}\left[\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\log f(Y_t;\boldsymbol{\theta}_0)\mid \mathbf{Y}_{-\infty}^{t-1}\right] = -\mathbb{E}\left[\frac{1}{f^2(Y_t;\boldsymbol{\theta}_0)}\nabla_{\boldsymbol{\theta}}f(Y_t;\boldsymbol{\theta}_0)\nabla_{\boldsymbol{\theta}'}f(Y_t;\boldsymbol{\theta}_0)\mid \mathbf{Y}_{-\infty}^{t-1}\right] + \int_{\mathcal{Y}}\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}f(y;\boldsymbol{\theta})|_{\boldsymbol{\theta}=\boldsymbol{\theta}_0}\,\mathrm{d}y \\ = -\mathbb{E}\left[\nabla_{\boldsymbol{\theta}}\log f(Y_t;\boldsymbol{\theta}_0)\nabla_{\boldsymbol{\theta}'}\log f(Y_t;\boldsymbol{\theta}_0)\mid \mathbf{Y}_{-\infty}^{t-1}\right]$$

$$+ \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \int_{\mathcal{Y}} f(y; \boldsymbol{\theta}) dy \bigg|_{\boldsymbol{\theta} = \boldsymbol{\theta}_{0}}$$

$$= -\mathbb{E} \left[ \nabla_{\boldsymbol{\theta}} \log f(Y_{t}; \boldsymbol{\theta}_{0}) \nabla_{\boldsymbol{\theta}'} \log f(Y_{t}; \boldsymbol{\theta}_{0}) \mid \mathbf{Y}_{-\infty}^{t-1} \right]$$

and thus that

$$\mathbb{E}\left[\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\log f(Y_t;\boldsymbol{\theta}_0)\right] = -\mathbb{E}\left[\nabla_{\boldsymbol{\theta}}\log f(Y_t;\boldsymbol{\theta}_0)\nabla_{\boldsymbol{\theta}'}\log f(Y_t;\boldsymbol{\theta}_0)\right].$$

Condition (v) follows from Lemma A.9 and A.10 since

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{L}_T(\boldsymbol{\theta}) - (-\mathbf{I}(\boldsymbol{\theta}))||_{2,2} \leq \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{L}_T(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} L_T(\boldsymbol{\theta})||_{2,2} + \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} L_T(\boldsymbol{\theta}) - (-\mathbf{I}(\boldsymbol{\theta}))||_{2,2}.$$

**Lemma A.9.** Under the assumptions in Theorem 3,

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{L}_T(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} L_T(\boldsymbol{\theta})||_{2,2} \stackrel{a.s.}{\to} 0 \quad as \quad T \to \infty.$$

*Proof.* We have that

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{L}_{T}(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} L_{T}(\boldsymbol{\theta})||_{2,2} \leq \frac{1}{T} \sum_{t=1}^{T} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{t}}{\hat{f}_{t}} \frac{\nabla_{\boldsymbol{\theta'}} \hat{f}_{t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{t}}{f_{t}} \frac{\nabla_{\boldsymbol{\theta'}} f_{t}}{f_{t}} \right\|_{2,2} + \frac{1}{T} \sum_{t=1}^{T} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{f}_{t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} f_{t}}{f_{t}} \right\|_{2,2},$$

where  $\hat{f}_t$  denotes  $\hat{f}(Y_t; \boldsymbol{\theta})$  and  $f_t$  denotes  $f(Y_t; \boldsymbol{\theta})$ , that

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{t}}{\hat{f}_{t}} \frac{\nabla_{\boldsymbol{\theta}'} \hat{f}_{t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{t}}{f_{t}} \frac{\nabla_{\boldsymbol{\theta}'} f_{t}}{f_{t}} \right\|_{2,2} \leq \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{t}}{f_{t}} \right\|_{2}^{2} + 2 \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta}} f_{t}}{\hat{f}_{t}} \right\|_{2} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{t}}{\hat{f}_{t}} \right\|_{2}^{2},$$

and that

$$\begin{split} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{f}_t}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} f_t}{f_t} \right| \right|_{2,2} &\leq \sum_{j=1}^{J} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \pi_{j,t|t-1} f_{j,t}}{f_t} \right| \right|_{2,2} \\ &+ 2 \sum_{j=1}^{J} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta'}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \nabla_{\boldsymbol{\theta'}} f_{j,t}}{f_t} \right| \right|_{2,2} \\ &+ \sum_{j=1}^{J} \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\pi_{j,t|t-1} \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} f_{j,t}}{f_t} \right| \right|_{2,2}, \end{split}$$

where  $\hat{\pi}_{j,t|t-1}$  denotes  $\hat{\pi}_{j,t|t-1}(\boldsymbol{\theta})$ ,  $\hat{f}_{j,t}$  denotes  $f_j(Y_t; \hat{X}_{j,t}, \boldsymbol{v}_j)$ ,  $\hat{X}_{j,t}$  denotes  $\hat{X}_{j,t}(\boldsymbol{v}_j)$ ,  $\pi_{j,t|t-1}$  denotes  $\pi_{j,t|t-1}(\boldsymbol{\theta})$ ,  $f_{j,t}$  denotes  $f_j(Y_t; X_{j,t}, \boldsymbol{v}_j)$ , and  $X_{j,t}$  denotes  $X_{j,t}(\boldsymbol{v}_j)$ .

The conclusion thus follows from Lemma 2.1 in Straumann and Mikosch (2006) if  $\sup_{\theta \in \Theta} \left| \left| \frac{\nabla_{\theta} \hat{f}_t}{\hat{f}_t} \frac{\nabla_{\theta'} \hat{f}_t}{\hat{f}_t} - \frac{\nabla_{\theta} f_t}{f_t} \frac{\nabla_{\theta'} f_t}{f_t} \right| \right|_{2,2} \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  and  $\sup_{\theta \in \Theta} \left| \left| \frac{\nabla_{\theta\theta} \hat{f}_t}{\hat{f}_t} - \frac{\nabla_{\theta\theta} f_t}{f_t} \right| \right|_{2,2} \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$ . The former follows from the arguments used in the proof of Lemma A.6 and

the latter follows if, for each  $j \in \{1, ..., J\}$ ,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \pi_{j,t|t-1} f_{j,t}}{f_t} \right\|_{2,2} \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$ ,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta}'} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \nabla_{\boldsymbol{\theta}'} f_{j,t}}{f_t} \right\|_{2,2} \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$ , and  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\pi_{j,t|t-1} \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} f_{j,t}}{f_t} \right\|_{2,2} \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$ .

First.

$$\left\| \frac{\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{\pi}_{j,t|t-1}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\pi_{j,t|t-1}f_{j,t}}{f_{t}} \right\|_{2,2} \leq \left| \left| \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{\pi}_{j,t|t-1} - \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\pi_{j,t|t-1} \right| \right|_{2,2} \left| \frac{\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{f_{j,t}}{f_{t}} \right| + C \left| \left| \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{\pi}_{j,t|t-1} - \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\pi_{j,t|t-1} \right| \right|_{2,2} + \left| \left| \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\pi_{j,t|t-1} \right| \right|_{2,2} \left| \frac{\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{f_{j,t}}{f_{t}} \right|.$$

It thus follows from Lemma 2.1 and 2.2 in Straumann and Mikosch (2006) that, for each  $j \in \{1,...,J\}$ ,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \pi_{j,t|t-1} f_{j,t}}{f_t} \right\|_{2,2} \stackrel{e.a.s.}{\longrightarrow} 0$  as  $t \to \infty$  since, in addition to the arguments used in the proofs of Lemma A.3 and A.6, for each  $j \in \{1,...,J\}$ ,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \|\nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} - \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \pi_{j,t|t-1}\|_{2,2} \stackrel{e.a.s.}{\longrightarrow} 0$  as  $t \to \infty$  where  $(\nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \pi_{j,t|t-1})_{t \in \mathbb{Z}}$  is stationary for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  and  $\mathbb{E}\left[\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \|\nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \pi_{j,t|t-1}\|_{2,2}\right] < \infty$  by Corollary 3.

Moreover,

$$\begin{aligned} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta}'} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \nabla_{\boldsymbol{\theta}'} f_{j,t}}{f_{t}} \right| \right|_{2,2} &\leq \left| \left| \nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} - \nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \right| \right|_{2} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right| \right|_{2} \\ &+ \left| \left| \nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} - \nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \right| \right|_{2} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{f_{t}} \right| \right|_{2} \\ &+ \left| \left| \nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \right| \right|_{2} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\boldsymbol{\theta}} f_{j,t}}{f_{t}} \right| \right|_{2} \end{aligned}$$

It thus follows from the arguments used in the proof of Lemma A.6 that, for each  $j \in \{1, ..., J\}$ , also  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\nabla_{\boldsymbol{\theta}} \hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta'}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\nabla_{\boldsymbol{\theta}} \pi_{j,t|t-1} \nabla_{\boldsymbol{\theta'}} f_{j,t}}{f_t} \right| \right| \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty.$ 

Finally,

$$\left\| \frac{\hat{\pi}_{j,t|t-1} \nabla_{\theta\theta} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\pi_{j,t|t-1} \nabla_{\theta\theta} f_{j,t}}{f_{t}} \right\|_{2,2} \leq \left| \hat{\pi}_{j,t|t-1} - \pi_{j,t|t-1} \right| \left\| \frac{\nabla_{\theta\theta} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\theta\theta} f_{j,t}}{f_{t}} \right\|_{2,2} + \left| \hat{\pi}_{j,t|t-1} - \pi_{j,t|t-1} \right| \left\| \frac{\nabla_{\theta\theta} f_{j,t}}{f_{t}} \right\|_{2,2} + \left\| \frac{\nabla_{\theta\theta} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\theta\theta} f_{j,t}}{f_{t}} \right\|_{2,2},$$

where

$$\left\| \frac{\nabla_{\theta\theta} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\nabla_{\theta\theta} f_{j,t}}{f_{t}} \right\|_{2,2} \leq \left\| \frac{\bar{\nabla}_{x_{j}x_{j}} \hat{f}_{j,t} \nabla_{\theta} \hat{X}_{j,t} \nabla_{\theta'} \hat{X}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}x_{j}} f_{j,t} \nabla_{\theta} X_{j,t} \nabla_{\theta'} X_{j,t}}{f_{t}} \right\|_{2,2} + \left\| \frac{\bar{\nabla}_{x_{j}} \hat{f}_{j,t} \nabla_{\theta\theta} \hat{X}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}} f_{j,t} \nabla_{\theta\theta} X_{j,t}}{f_{t}} \right\|_{2,2}$$

$$+ 2 \left\| \frac{\bar{\nabla}_{\theta x_{j}} \hat{f}_{j,t} \nabla_{\theta'} \hat{X}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\theta x_{j}} f_{j,t} \nabla_{\theta'} X_{j,t}}{f_{t}} \right\|_{2,2} \\ + \left\| \frac{\bar{\nabla}_{\theta \theta} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\theta \theta} f_{j,t}}{f_{t}} \right\|_{2,2}.$$

First, we have that

$$\begin{split} & \left| \left| \frac{\bar{\nabla}_{x_{j}x_{j}}\hat{f}_{j,t}\nabla_{\boldsymbol{\theta}}\hat{X}_{j,t}\nabla_{\boldsymbol{\theta}'}\hat{X}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}x_{j}}f_{j,t}\nabla_{\boldsymbol{\theta}}X_{j,t}\nabla_{\boldsymbol{\theta}'}X_{j,t}}{f_{t}} \right| \right|_{2,2} \\ & \leq \left| \frac{\bar{\nabla}_{x_{j}x_{j}}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}x_{j}}f_{j,t}}{f_{t}} \right| \left| \left| \nabla_{\boldsymbol{\theta}}\hat{X}_{j,t}\nabla_{\boldsymbol{\theta}'}\hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}}X_{j,t}\nabla_{\boldsymbol{\theta}'}X_{j,t} \right| \right|_{2,2} \\ & + \left| \frac{\bar{\nabla}_{x_{j}x_{j}}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}x_{j}}f_{j,t}}{f_{t}} \right| \left| \left| \nabla_{\boldsymbol{\theta}}X_{j,t} \right|^{2} \\ & + \left| \frac{\bar{\nabla}_{x_{j}x_{j}}f_{j,t}}{f_{t}} \right| \left| \left| \nabla_{\boldsymbol{\theta}}\hat{X}_{j,t}\nabla_{\boldsymbol{\theta}'}\hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}}X_{j,t}\nabla_{\boldsymbol{\theta}'}X_{j,t} \right| \right|_{2,2}, \end{split}$$

where

$$\left|\frac{\bar{\nabla}_{x_j x_j} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{x_j x_j} f_{j,t}}{f_t}\right| \leq \left|\frac{\bar{\nabla}_{x_j x_j} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{x_j x_j} \hat{f}_{j,t}}{\tilde{f}_t}\right| + \left|\frac{\bar{\nabla}_{x_j x_j} \hat{f}_{j,t}}{\tilde{f}_t} - \frac{\bar{\nabla}_{x_j x_j} f_{j,t}}{f_t}\right|$$

where  $\tilde{f}_t$  denotes  $\tilde{f}(Y_t; \boldsymbol{\theta}) = \sum_{j=1}^J \pi_{j,t|t-1}(\boldsymbol{\theta}) f_j(Y_t; \hat{X}_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)$  and

$$\left\| \left| \nabla_{\boldsymbol{\theta}} \hat{X}_{j,t} \nabla_{\boldsymbol{\theta}'} \hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}} X_{j,t} \nabla_{\boldsymbol{\theta}'} X_{j,t} \right| \right\|_{2,2} \leq \left\| \left| \nabla_{\boldsymbol{\theta}} \hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}} X_{j,t} \right| \right\|_{2}^{2} + 2 \left\| \left| \nabla_{\boldsymbol{\theta}} \hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}} X_{j,t} \right| \right\|_{2}.$$

By using the same arguments as in the proof of Lemma A.3, we have that

$$\begin{split} \left| \frac{\bar{\nabla}_{x_{j}x_{j}}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}x_{j}}f_{j,t}}{f_{t}} \right| &\leq C \left| \bar{\nabla}_{x_{j}x_{j}} \log \hat{f}_{j,t} \right| \left| \left| \hat{\pi}_{t|t-1} - \pi_{t|t-1} \right| \right|_{2} \\ &+ C \left| \bar{\nabla}_{x_{j}} \log \hat{f}_{j,t} \right|^{2} \left| \left| \hat{\pi}_{t|t-1} - \pi_{t|t-1} \right| \right|_{2} \\ &+ C \sum_{l=1}^{J} \left| \bar{\nabla}_{x_{j}x_{j}x_{j}} \log \bar{f}_{j,t} \right| \left| \hat{X}_{l,t} - X_{l,t} \right| \\ &+ C \sum_{k,l=1}^{J} \left| \bar{\nabla}_{x_{j}x_{j}} \log \bar{f}_{j,t} \right| \left| \bar{\nabla}_{x_{k}} \log \bar{f}_{k,t} \right| \left| \hat{X}_{l,t} - X_{l,t} \right| \\ &+ C \sum_{k,l=1}^{J} \left| \bar{\nabla}_{x_{j}} \log \bar{f}_{j,t} \right|^{2} \left| \bar{\nabla}_{x_{k}} \log \bar{f}_{k,t} \right| \left| \hat{X}_{l,t} - X_{l,t} \right|, \end{split}$$

where  $\bar{f}_{j,t}$  denotes  $f_j(Y_t; \bar{X}_{j,t}(v_j), v_j)$ . Second, we have that

$$\left\| \frac{\bar{\nabla}_{x_{j}} \hat{f}_{j,t} \nabla_{\theta \theta} \hat{X}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}} f_{j,t} \nabla_{\theta \theta} X_{j,t}}{f_{t}} \right\|_{2,2} \leq \left| \frac{\bar{\nabla}_{x_{j}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}} f_{j,t}}{f_{t}} \right| \left\| \nabla_{\theta \theta} \hat{X}_{j,t} - \nabla_{\theta \theta} X_{j,t} \right\|_{2,2}$$

$$+ \left| \frac{\bar{\nabla}_{x_{j}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{x_{j}} f_{j,t}}{f_{t}} \right| \left\| \nabla_{\theta \theta} \hat{X}_{j,t} \right\|_{2,2}$$

$$+ \left| \frac{\bar{\nabla}_{x_{j}} f_{j,t}}{f_{t}} \right| \left\| \nabla_{\theta \theta} \hat{X}_{j,t} - \nabla_{\theta \theta} X_{j,t} \right\|_{2,2},$$

where

$$\left|\frac{\bar{\nabla}_{x_j}\hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{x_j}f_{j,t}}{f_t}\right| \leq \left|\frac{\bar{\nabla}_{x_j}\hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{x_j}\hat{f}_{j,t}}{\tilde{f}_t}\right| + \left|\frac{\bar{\nabla}_{x_j}\hat{f}_{j,t}}{\tilde{f}_t} - \frac{\bar{\nabla}_{x_j}f_{j,t}}{f_t}\right|$$

as in the proof of Lemma A.6. Third, we have that

$$\left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}x_{j}}\hat{f}_{j,t}\nabla_{\boldsymbol{\theta}'}\hat{X}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\boldsymbol{\theta}x_{j}}f_{j,t}\nabla_{\boldsymbol{\theta}'}X_{j,t}}{f_{t}} \right\|_{2,2} \leq \left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}x_{j}}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\boldsymbol{\theta}x_{j}}f_{j,t}}{f_{t}} \right\|_{2} \left\| |\nabla_{\boldsymbol{\theta}}\hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}}X_{j,t}| \right\|_{2} + \left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}x_{j}}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\boldsymbol{\theta}x_{j}}f_{j,t}}{f_{t}} \right\|_{2} \left\| |\nabla_{\boldsymbol{\theta}}X_{j,t}| \right\|_{2} + \left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}x_{j}}f_{j,t}}{f_{t}} \right\|_{2} \left\| |\nabla_{\boldsymbol{\theta}}\hat{X}_{j,t} - \nabla_{\boldsymbol{\theta}}X_{j,t}| \right\|_{2},$$

where

$$\left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}x_j} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}x_j} f_{j,t}}{f_t} \right\|_2 \le \left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}x_j} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}x_j} \hat{f}_{j,t}}{\tilde{f}_t} \right\|_2 + \left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}x_j} \hat{f}_{j,t}}{\tilde{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}x_j} f_{j,t}}{f_t} \right\|_2.$$

By using the same arguments as in the proof of Lemma A.3 again, we have that

$$\begin{split} \left\| \frac{\bar{\nabla}_{\theta x_{j}} \hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\theta x_{j}} f_{j,t}}{f_{t}} \right\|_{2} &\leq C \sum_{h=1}^{d_{j}} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h} x_{j}} \log \hat{f}_{j,t} \right| \left| |\hat{\boldsymbol{\pi}}_{t|t-1} - \boldsymbol{\pi}_{t|t-1}| \right|_{2} \\ &+ C \sum_{h=1}^{d_{j}} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}} \log \hat{f}_{j,t} \right| \left| \bar{\nabla}_{x_{j}} \log \hat{f}_{j,t} \right| \left| |\hat{\boldsymbol{\pi}}_{t|t-1} - \boldsymbol{\pi}_{t|t-1}| \right|_{2} \\ &+ C \sum_{h=1}^{d_{j}} \sum_{l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h} x_{j} x_{j}} \log \bar{f}_{j,t} \right| \left| \hat{\boldsymbol{X}}_{l,t} - \boldsymbol{X}_{l,t} \right| \\ &+ C \sum_{h=1}^{d_{j}} \sum_{k,l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h} x_{j}} \log \bar{f}_{j,t} \right| \left| \bar{\nabla}_{x_{k}} \log \bar{f}_{k,t} \right| \left| \hat{\boldsymbol{X}}_{l,t} - \boldsymbol{X}_{l,t} \right| \\ &+ C \sum_{h=1}^{d_{j}} \sum_{l=1}^{J} \left| \bar{\nabla}_{x_{j} x_{j}} \log \bar{f}_{j,t} \right| \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}} \log \bar{f}_{j,t} \right| \left| \bar{\boldsymbol{X}}_{l,t} - \boldsymbol{X}_{l,t} \right| \\ &+ C \sum_{h=1}^{d_{j}} \sum_{k,l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}} \log \bar{f}_{j,t} \right| \left| \bar{\nabla}_{x_{j}} \log \bar{f}_{j,t} \right| \left| \bar{\nabla}_{x_{k}} \log \bar{f}_{k,t} \right| \left| \hat{\boldsymbol{X}}_{l,t} - \boldsymbol{X}_{l,t} \right| . \end{split}$$

Finally, we have that

$$\left| \left| \frac{\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}f_{j,t}}{f_t} \right| \right|_{2,2} \leq \left| \left| \frac{\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{f}_{j,t}}{\hat{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{f}_{j,t}}{\tilde{f}_t} \right| \right|_{2,2} + \left| \left| \frac{\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{f}_{j,t}}{\tilde{f}_t} - \frac{\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}f_{j,t}}{f_t} \right| \right|_{2,2}.$$

By using the same arguments as in the proof of Lemma A.3 once again, we have that

$$\left\| \frac{\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{f}_{j,t}}{\hat{f}_{t}} - \frac{\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}f_{j,t}}{f_{t}} \right\|_{2,2} \leq C \sum_{h,i=1}^{d_{j}} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}[\boldsymbol{v}_{j}]_{i}} \log \hat{f}_{j,t} \right| \left| \left| \hat{\boldsymbol{\pi}}_{t|t-1} - \boldsymbol{\pi}_{t|t-1} \right| \right|_{2}$$

$$+ C \sum_{h,i=1}^{d_{j}} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}} \log \hat{f}_{j,t} \right| \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{i}} \log \hat{f}_{j,t} \right| \left| \left| \hat{\boldsymbol{\pi}}_{t|t-1} - \boldsymbol{\pi}_{t|t-1} \right| \right|_{2}$$

$$+ C \sum_{h,i=1}^{d_{j}} \sum_{l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}[\boldsymbol{v}_{j}]_{i}x_{j}} \log \bar{f}_{j,t} \right| |\hat{X}_{l,t} - X_{l,t}|$$

$$+ C \sum_{h,i=1}^{d_{j}} \sum_{k,l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}[\boldsymbol{v}_{j}]_{i}} \log \bar{f}_{j,t} \right| |\bar{\nabla}_{x_{k}} \log \bar{f}_{k,t}| |\hat{X}_{l,t} - X_{l,t}|$$

$$+ C \sum_{h,i=1}^{d_{j}} \sum_{l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}x_{j}} \log \bar{f}_{j,t} \right| |\bar{\nabla}_{[\boldsymbol{v}_{j}]_{i}} \log \bar{f}_{j,t}| |\hat{X}_{l,t} - X_{l,t}|$$

$$+ C \sum_{h,i=1}^{d_{j}} \sum_{l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{i}x_{j}} \log \bar{f}_{j,t} \right| |\bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}} \log \bar{f}_{j,t}| |\hat{X}_{l,t} - X_{l,t}|$$

$$+ C \sum_{h,i=1}^{d_{j}} \sum_{k,l=1}^{J} \left| \bar{\nabla}_{[\boldsymbol{v}_{j}]_{h}} \log \bar{f}_{j,t} \right| |\bar{\nabla}_{[\boldsymbol{v}_{j}]_{i}} \log \bar{f}_{j,t}| |\bar{\nabla}_{x_{k}} \log \bar{f}_{k,t}| |\hat{X}_{l,t} - X_{l,t}|.$$

It thus follows from Lemma 2.1 and 2.2 in Straumann and Mikosch (2006) that, for each  $j \in \{1, ..., J\}$ , also  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \left| \frac{\hat{\pi}_{j,t|t-1} \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} \hat{f}_{j,t}}{\hat{f}_t} - \frac{\pi_{j,t|t-1} \nabla_{\boldsymbol{\theta} \boldsymbol{\theta}} f_{j,t}}{f_t} \right| \right|_{2,2} \stackrel{e.a.s.}{\longrightarrow} 0$  as  $t \to \infty$  since, in addition to the arguments used in the proofs of Lemma A.3 and A.6, for each  $j \in \{1, ..., J\}$ ,  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} \hat{X}_{j,t} - \nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t}||_{2,2} \stackrel{e.a.s.}{\longrightarrow} 0$  as  $t \to \infty$  where  $(\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t})_{t \in \mathbb{Z}}$  is stationary for all  $\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j$  and  $\mathbb{E} \left[ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t}||_{2,2}^{k_j} \right] < \infty$  by Lemma 4. Moreover, for each  $j \in \{1, ..., J\}$ , there exists an  $m_j > 0$  such that  $\mathbb{E} \left[ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} ||\bar{\nabla}_{\boldsymbol{v}_j \boldsymbol{v}_j} \log f_{j,t}||_{2,2}^{m_j} \right] < \infty$ ,  $\mathbb{E} \left[ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} ||\bar{\nabla}_{\boldsymbol{v}_j x_j x_j} \log f_{j,t}||_{2}^{m_j} \right] < \infty$ ,  $\mathbb{E} \left[ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} ||\bar{\nabla}_{\boldsymbol{v}_j x_j x_j} \log f_{j,t}||_{2}^{m_j} \right] < \infty$  by assumption.  $\square$ 

**Lemma A.10.** Under the assumptions in Theorem 3,

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} L_T(\boldsymbol{\theta}) - (-\mathbf{I}(\boldsymbol{\theta}))||_{2,2} \stackrel{a.s.}{\to} 0 \quad as \quad T \to \infty.$$

*Proof.* As in the proof of Lemma A.4, the conclusion follows from the uniform law of large numbers by Rao (1962) if  $(\nabla_{\theta\theta} \log f(Y_t; \theta))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\theta \in \Theta$  and  $\mathbb{E} \left[ \sup_{\theta \in \Theta} ||\nabla_{\theta\theta} \log f(Y_t; \theta)||_{2,2} \right] < \infty$ .

First,  $(\nabla_{\theta\theta} \log f(Y_t; \theta))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\theta \in \Theta$  since, in addition to the arguments used in the proofs of Lemma A.4 and A.7, for each  $j \in \{1, ..., J\}$ ,  $(\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{v}_j \in \Upsilon_j$  by Lemma 4, and, for each  $j \in \{1, ..., J\}$ ,  $(\nabla_{\theta\theta} \pi_{j,t|t-1}(\theta))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\theta \in \Theta$  by Corollary 3.

We have that

$$||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\log f(Y_t;\boldsymbol{\theta})||_{2,2} \leq \frac{1}{f^2(Y_t;\boldsymbol{\theta})}||\nabla_{\boldsymbol{\theta}}f(Y_t;\boldsymbol{\theta})||_2^2 + \frac{1}{f(Y_t;\boldsymbol{\theta})}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}f(Y_t;\boldsymbol{\theta})||_{2,2}$$

where

$$||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} f(Y_t; \boldsymbol{\theta})||_{2,2}$$

$$\leq \sum_{j=1}^{J} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \pi_{j,t|t-1}(\boldsymbol{\theta})||_{2,2} f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)$$

$$+2\sum_{j=1}^{J}||\nabla_{\boldsymbol{\theta}}\pi_{j,t|t-1}(\boldsymbol{\theta})||_{2}||\nabla_{\boldsymbol{\theta}}f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2}$$

$$+\sum_{j=1}^{J}\pi_{j,t|t-1}(\boldsymbol{\theta})||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2,2},$$

so

$$\begin{split} & \mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\Theta}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\log f(Y_{t};\boldsymbol{\theta})||_{2,2}\right] \\ & \leq \mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\Theta}||\nabla_{\boldsymbol{\theta}}\log f(Y_{t};\boldsymbol{\theta})||_{2}^{2}\right] \\ & + C\sum_{j=1}^{J}\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\Theta}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\pi_{j,t|t-1}(\boldsymbol{\theta})||_{2,2}\right] \\ & + C\sum_{j=1}^{J}\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\Theta}||\nabla_{\boldsymbol{\theta}}\pi_{j,t|t-1}(\boldsymbol{\theta})||_{2}^{2}\right]\right)^{1/2}\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\Theta}||\nabla_{\boldsymbol{\theta}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2}^{2}\right]\right)^{1/2} \\ & + \sum_{j=1}^{J}\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\Theta}||\nabla_{\boldsymbol{\theta}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2}^{2}\right] \\ & + \sum_{j=1}^{J}\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\Theta}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2,2}\right]. \end{split}$$

Moreover, we have that

$$||\nabla_{\theta\theta} \log f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})||_{2,2} \leq \frac{1}{f_{j}^{2}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})} ||\nabla_{\theta} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})||_{2}^{2} + \frac{1}{f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})} ||\nabla_{\theta\theta} f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j})||_{2,2}$$

where

$$\begin{split} ||\nabla_{\theta\theta}f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2,2} &\leq |\bar{\nabla}_{x_{j}x_{j}}f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})|||\nabla_{\theta}X_{j,t}(\boldsymbol{v}_{j})||_{2}^{2} \\ &+ |\bar{\nabla}_{x_{j}}f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})|||\nabla_{\theta\theta}X_{j,t}(\boldsymbol{v}_{j})||_{2,2} \\ &+ 2||\bar{\nabla}_{\theta x_{j}}f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2}||\nabla_{\theta}X_{j,t}(\boldsymbol{v}_{j})||_{2} \\ &+ ||\bar{\nabla}_{\theta\theta}f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2,2}, \end{split}$$

so

$$\begin{split} & \mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{\upsilon}_{j}),\boldsymbol{\upsilon}_{j})||_{2,2}\right] \\ & \leq \mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{\upsilon}_{j}),\boldsymbol{\upsilon}_{j})||_{2}^{2}\right] \\ & + \left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}|\bar{\nabla}_{x_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{\upsilon}_{j}),\boldsymbol{\upsilon}_{j})|^{2k_{j}/(k_{j}-1)}\right]\right)^{(k_{j}-1)/k_{j}}\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}X_{j,t}(\boldsymbol{\upsilon}_{j})||_{2}^{2k_{j}}\right]\right)^{1/k_{j}} \\ & + \left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}|\bar{\nabla}_{x_{j}x_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{\upsilon}_{j}),\boldsymbol{\upsilon}_{j})|^{k_{j}/(k_{j}-1)}\right]\right)^{(k_{j}-1)/k_{j}}\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}X_{j,t}(\boldsymbol{\upsilon}_{j})||_{2}^{2k_{j}}\right]\right)^{1/k_{j}} \\ & + \left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}|\bar{\nabla}_{x_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{\upsilon}_{j}),\boldsymbol{\upsilon}_{j})|^{k_{j}/(k_{j}-1)}\right]\right)^{(k_{j}-1)/k_{j}}\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}X_{j,t}(\boldsymbol{\upsilon}_{j})||_{2,2}^{2k_{j}}\right]\right)^{1/k_{j}} \end{split}$$

$$+2\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}|\bar{\nabla}_{x_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})|^{2k_{j}/(k_{j}-1)}\right]\right)^{(k_{j}-1)/2k_{j}}$$

$$\cdot\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\bar{\nabla}_{\boldsymbol{\theta}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2}^{2}\right]\right)^{1/2}\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}X_{j,t}(\boldsymbol{v}_{j})||_{2}^{2k_{j}}\right]\right)^{1/2k_{j}}$$

$$+2\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\bar{\nabla}_{\boldsymbol{\theta}x_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2}^{k_{j}/(k_{j}-1)}\right]\right)^{(k_{j}-1)/k_{j}}\left(\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}X_{j,t}(\boldsymbol{v}_{j})||_{2}^{k_{j}}\right]\right)^{1/k_{j}}$$

$$+\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\bar{\nabla}_{\boldsymbol{\theta}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2}^{2}\right]$$

$$+\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\bar{\nabla}_{\boldsymbol{\theta}\boldsymbol{\theta}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2,2}\right].$$

Hence,  $\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\log f(Y_t;\boldsymbol{\theta})||_{2,2}\right]<\infty$  since, in addition to the arguments used in the proof of Lemma A.7, for each  $j\in\{1,...,J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}||\nabla_{\boldsymbol{v}_j\boldsymbol{v}_j}X_{j,t}(\boldsymbol{v}_j)||_{2,2}^{k_j}\right]<\infty$  by Lemma 4, for each  $j\in\{1,...,J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\pi_{j,t|t-1}(\boldsymbol{\theta})||_{2,2}\right]<\infty$  by Corollary 3, and, for each  $j\in\{1,...,J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}||\bar{\nabla}_{x_jx_j}\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)||_{k_j/(k_j-1)}^{k_j/(k_j-1)}\right]<\infty$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}||\bar{\nabla}_{\boldsymbol{v}_jx_j}\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)||_{2,2}^{k_j/(k_j-1)}\right]<\infty$ , and  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j\in\boldsymbol{\Upsilon}_j}||\bar{\nabla}_{\boldsymbol{v}_jv_j}\log f_j(Y_t;X_{j,t}(\boldsymbol{v}_j),\boldsymbol{v}_j)||_{2,2}\right]<\infty$  by Assumption 10.

Finally, Condition (vi) is true by assumption.

# B Other Proofs

#### B.1 Proof of Lemma 3

For each  $j \in \{1, ..., J\}$ ,  $(\nabla_{v_j} X_{j,t}(v_j))_{t \in \mathbb{Z}}$  is a stochastic process taking values in  $(\mathbb{R}^{d_j}, ||\cdot||_2)$  given by

$$\nabla_{\boldsymbol{v}_j} X_{j,t+1}(\boldsymbol{v}_j) = \phi_{j,t}^x (\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j),$$

where  $(\phi_{j,t}^x(\cdot; v_j))_{t \in \mathbb{Z}}$  is a sequence of stationary and ergodic Lipschitz functions given by

$$\phi_{i,t}^{x}(\mathbf{x}_{j}; \boldsymbol{v}_{j}) = \bar{\nabla}_{x_{i}}\phi_{j}(Y_{t}, X_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j})\mathbf{x}_{j} + \bar{\nabla}_{\boldsymbol{v}_{j}}\phi_{j}(Y_{t}, X_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j})$$

with Lipschitz coefficient

$$\Lambda(\phi_{j,t}^x; \boldsymbol{v}_j) = |\bar{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)|.$$

The first conclusion follows from Lemma C.2 since, for each  $j \in \{1, ..., J\}$ ,

- (i)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\bar{\nabla}_{\boldsymbol{v}_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)||_2] < \infty,$
- (ii)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \Upsilon_j} |\bar{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)|] < \infty$ , and
- (iii)  $\mathbb{E}[\log \sup_{\boldsymbol{v}_i \in \boldsymbol{\Upsilon}_i} |\bar{\nabla}_{x_i} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)|] < 0$

by assumption.<sup>7</sup>

<sup>&</sup>lt;sup>7</sup>Condition (ii) and (iii) follow from Condition (ii) and (iii) in Lemma 1.

Moreover, for each  $j \in \{1, ..., J\}$ ,  $(\nabla_{\boldsymbol{v}_j} \hat{X}_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{N}}$  is a stochastic process taking values in  $(\mathbb{R}^{d_j}, ||\cdot||_2)$  given by

$$\nabla_{\boldsymbol{v}_j} \hat{X}_{j,t+1}(\boldsymbol{v}_j) = \hat{\phi}_{j,t}^x (\nabla_{\boldsymbol{v}_j} \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j),$$

where  $(\hat{\phi}_{j,t}^x(\cdot; \boldsymbol{v}_j))_{t\in\mathbb{N}}$  is a sequence of non-stationary Lipschitz functions given by

$$\hat{\phi}_{j,t}^{x}(\mathbf{x}_{j};\boldsymbol{v}_{j}) = \bar{\nabla}_{x_{j}}\phi_{j}(Y_{t},\hat{X}_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\mathbf{x}_{j} + \bar{\nabla}_{\boldsymbol{v}_{j}}\phi_{j}(Y_{t},\hat{X}_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j}).$$

The second conclusion follows from Lemma C.2 as well since, for each  $j \in \{1, ..., J\}$ ,

- (i)  $\mathbb{E}[\sup_{\boldsymbol{v}_i \in \boldsymbol{\Upsilon}_i} ||\nabla_{\boldsymbol{v}_i} X_{i,t}(\boldsymbol{v}_i)||_2^{2k_j}] < \infty$ ,
- (ii)  $\sup_{\boldsymbol{v}_{j} \in \boldsymbol{\Upsilon}_{j}} ||\bar{\nabla}_{\boldsymbol{v}_{j}} \phi_{j}(Y_{t}, \hat{X}_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j}) \bar{\nabla}_{\boldsymbol{v}_{j}} \phi_{j}(Y_{t}, X_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j})||_{2} \overset{e.a.s.}{\to} 0 \text{ as } t \to \infty, \text{ and } t \to \infty$
- (iii)  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\bar{\nabla}_{x_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) \bar{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)| \overset{e.a.s.}{\to} 0 \text{ as } t \to \infty$

by assumption.

### B.2 Proof of Lemma 4

For each  $j \in \{1, ..., J\}$ ,  $(\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{Z}}$  is a stochastic process taking values in  $(\mathbb{R}^{d_j \times d_j}, ||\cdot||_{2,2})$  given by

$$\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t+1}(\boldsymbol{v}_j) = \phi_{j,t}^{xx} (\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j),$$

where  $(\phi_{i,t}^{xx}(\cdot; v_j))_{t \in \mathbb{Z}}$  is a sequence of stationary and ergodic Lipschitz functions given by

$$\phi_{j,t}^{xx}(\mathbf{X}_{j};\boldsymbol{v}_{j}) = \bar{\nabla}_{x_{j}}\phi_{j}(Y_{t},X_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\mathbf{X}_{j} + \bar{\nabla}_{x_{j}x_{j}}\phi_{j}(Y_{t},X_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}}X_{j,t}(\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}'}X_{j,t}(\boldsymbol{v}_{j}) + \nabla_{\boldsymbol{v}_{j}}X_{j,t}(\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}'}X_{j,t}(\boldsymbol{v}_{j}) + \bar{\nabla}_{\boldsymbol{v}_{j}x_{j}}\phi_{j}(Y_{t},X_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}'}X_{j,t}(\boldsymbol{v}_{j}) + \bar{\nabla}_{\boldsymbol{v}_{j}x_{j}}\phi_{j}(Y_{t},X_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}'}X_{j,t}(\boldsymbol{v}_{j}) + \bar{\nabla}_{\boldsymbol{v}_{j}x_{j}}\phi_{j}(Y_{t},X_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}'}X_{j,t}(\boldsymbol{v}_{j})$$

with Lipschitz coefficient

$$\Lambda(\phi_{j,t}^{xx}; \boldsymbol{v}_j) = |\bar{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)|$$

as above.

The first conclusion follows from Lemma C.2 since, for each  $j \in \{1, ..., J\}$ ,

- (i)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\bar{\nabla}_{x_j x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)|] < \infty, \mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\bar{\nabla}_{\boldsymbol{v}_j x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)||_2] < \infty$  $\infty \text{ and } \mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\bar{\nabla}_{\boldsymbol{v}_j \boldsymbol{v}_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)||_{2,2}] < \infty,$
- (ii)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\bar{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)|] < \infty$ , and
- (iii)  $\mathbb{E}[\log \sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\bar{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)|] < 0$

by assumption.

Moreover, for each  $j \in \{1, ..., J\}$ ,  $(\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} \hat{X}_{j,t}(\boldsymbol{v}_j))_{t \in \mathbb{N}}$  is a stochastic process taking values in  $(\mathbb{R}^{d_j \times d_j}, ||\cdot||_{2,2})$  given by

$$\nabla_{\boldsymbol{v}_{j}\boldsymbol{v}_{j}}\hat{X}_{j,t+1}(\boldsymbol{v}_{j}) = \hat{\phi}_{j,t}^{xx}(\nabla_{\boldsymbol{v}_{j}\boldsymbol{v}_{j}}\hat{X}_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j}),$$

where  $(\hat{\phi}_{i,t}^{xx}(\cdot; \boldsymbol{v}_j))_{t\in\mathbb{N}}$  is a sequence of non-stationary Lipschitz functions given by

$$\begin{split} \hat{\phi}_{j,t}^{xx}(\mathbf{X}_{j};\boldsymbol{v}_{j}) &= \bar{\nabla}_{x_{j}}\phi_{j}(Y_{t},\hat{X}_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\mathbf{X}_{j} + \bar{\nabla}_{x_{j}x_{j}}\phi_{j}(Y_{t},\hat{X}_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}}\hat{X}_{j,t}(\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}'}\hat{X}_{j,t}(\boldsymbol{v}_{j}) \\ &+ \nabla_{\boldsymbol{v}_{j}}\hat{X}_{j,t}(\boldsymbol{v}_{j})\bar{\nabla}_{x_{j}\boldsymbol{v}_{j}}\phi_{j}(Y_{t},\hat{X}_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j}) + \bar{\nabla}_{\boldsymbol{v}_{j}x_{j}}\phi_{j}(Y_{t},\hat{X}_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j})\nabla_{\boldsymbol{v}_{j}'}\hat{X}_{j,t}(\boldsymbol{v}_{j}) \\ &+ \bar{\nabla}_{\boldsymbol{v}_{j}\boldsymbol{v}_{j}}\phi_{j}(Y_{t},\hat{X}_{j,t}(\boldsymbol{v}_{j});\boldsymbol{v}_{j}). \end{split}$$

The second conclusion follows from Lemma C.2 as well since, for each  $j \in \{1, ..., J\}$ ,

- (i)  $\mathbb{E}[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_i} ||\nabla_{\boldsymbol{v}_j \boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j)||_{2,2}^{k_j}] < \infty$ ,
- (ii)  $\sup_{\boldsymbol{v}_{j} \in \boldsymbol{\Upsilon}_{j}} |\bar{\nabla}_{x_{j}x_{j}} \phi_{j}(Y_{t}, \hat{X}_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j}) \bar{\nabla}_{x_{j}x_{j}} \phi_{j}(Y_{t}, X_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j})| \stackrel{e.a.s.}{\longrightarrow} 0 \text{ as } t \to \infty,$   $\sup_{\boldsymbol{v}_{j} \in \boldsymbol{\Upsilon}_{j}} ||\bar{\nabla}_{\boldsymbol{v}_{j}x_{j}} \phi_{j}(Y_{t}, \hat{X}_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j}) - \bar{\nabla}_{\boldsymbol{v}_{j}x_{j}} \phi_{j}(Y_{t}, X_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j})||_{2} \stackrel{e.a.s.}{\longrightarrow} 0 \text{ as } t \to \infty \text{ and }$  $\sup_{\boldsymbol{v}_{j} \in \boldsymbol{\Upsilon}_{j}} ||\bar{\nabla}_{\boldsymbol{v}_{j}\boldsymbol{v}_{j}} \phi_{j}(Y_{t}, \hat{X}_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j}) - \bar{\nabla}_{\boldsymbol{v}_{j}\boldsymbol{v}_{j}} \phi_{j}(Y_{t}, X_{j,t}(\boldsymbol{v}_{j}); \boldsymbol{v}_{j})||_{2,2} \stackrel{e.a.s.}{\longrightarrow} 0 \text{ as } t \to \infty, \text{ and }$
- (iii)  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\overline{\nabla}_{x_j} \phi_j(Y_t, \hat{X}_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j) \overline{\nabla}_{x_j} \phi_j(Y_t, X_{j,t}(\boldsymbol{v}_j); \boldsymbol{v}_j)| \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty$ by assumption.<sup>8</sup>

#### B.3 Proof of Lemma 5

First,  $(\nabla_{\boldsymbol{\theta}} \pi_{t|t}(\boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is a stochastic process taking values in  $(\mathbb{R}^{(J-1)d}, ||\cdot||_2)$  given by

$$\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}) = \boldsymbol{\phi}_t^{\boldsymbol{\pi}}(\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}),$$

where  $(\phi_t^{\pi}(\cdot; \boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is a sequence of stationary and ergodic Lipschitz functions given by

$$\begin{split} \left[ \boldsymbol{\phi}_{t}^{\boldsymbol{\pi}}(\mathbf{x}; \boldsymbol{\theta}) \right]_{(n-1)d+m} &= \sum_{j=1}^{J-1} \left( \bar{\nabla}_{\left[\boldsymbol{\pi}\right]_{j}} \left[ \boldsymbol{\phi}_{t}(\boldsymbol{\pi}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{\left[\boldsymbol{\pi}\right]_{J}} \left[ \boldsymbol{\phi}_{t}(\boldsymbol{\pi}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} \right) [\mathbf{x}]_{(j-1)d+m} \\ &+ \bar{\nabla}_{\left[\boldsymbol{\theta}\right]_{m}} \left[ \boldsymbol{\phi}_{t}(\boldsymbol{\pi}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n}, \quad (n, m) \in \{1, ..., J-1\} \times \{1, ..., d\}, \end{split}$$

see Appendix E for details, with Lipschitz coefficient

$$\Lambda(\phi_t^{\boldsymbol{\pi}};\boldsymbol{\theta}) = \sup_{\substack{\mathbf{x},\mathbf{y} \in \mathbb{R}^{(J-1)d} \\ \mathbf{x} \neq \mathbf{y}}} \frac{||\phi_t^{\boldsymbol{\pi}}(\mathbf{x};\boldsymbol{\theta}) - \phi_t^{\boldsymbol{\pi}}(\mathbf{y};\boldsymbol{\theta})||_2}{||\mathbf{x} - \mathbf{y}||_2}.$$

The fact that  $(\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is stationary and ergodic for all  $\boldsymbol{\theta} \in \boldsymbol{\Theta}$  with  $\mathbb{E}[\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_2^2] < \infty$  follows from Lemma C.2 if

- (i)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\phi_t^{\boldsymbol{\pi}}(\boldsymbol{0}; \boldsymbol{\theta})||_2] < \infty$ ,
- (ii)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \Lambda(\boldsymbol{\phi}_t^{\boldsymbol{\pi}}; \boldsymbol{\theta})] < \infty$ , and
- (iii) there exists an  $r \in \mathbb{N}$  such that  $\mathbb{E}[\log \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \Lambda((\boldsymbol{\phi}_t^{\boldsymbol{\pi}})^{(r)}; \boldsymbol{\theta})] < 0$ ;

the former is a direct consequence of Lemma C.2, and the latter is a bi-product of Lemma C.2, see later.

<sup>&</sup>lt;sup>8</sup>Condition (iii) follows from Condition (iii) in Lemma 3.

Condition (i) follows if, for each  $(n,m) \in \{1,...,J-1\} \times \{1,...,d\}$ ,  $\mathbb{E}[\log^+ \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\overline{\nabla}_{[\boldsymbol{\theta}]_m} [\boldsymbol{\phi}_t(\boldsymbol{\pi}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta})]_n|] < \infty$ . This is true if, for each  $j \in \{1,...,J-1\}$ ,

$$\mathbb{E}\left[\log^{+}\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}\left|\frac{\pi_{a,t-1|t-1}(\boldsymbol{\theta})f_{b}(Y_{t};X_{b,t}(\boldsymbol{v}_{b}),\boldsymbol{v}_{b})}{\sum_{k=1}^{J}\sum_{l=1}^{J}p_{lk}\pi_{l,t-1|t-1}(\boldsymbol{\theta})f_{k}(Y_{t};X_{k,t}(\boldsymbol{v}_{k}),\boldsymbol{v}_{k})}\right|\right]<\infty$$

and

$$\mathbb{E}\left[\log^{+}\sup_{\boldsymbol{\theta}\in\Theta}\left|\frac{\left(\sum_{i=1}^{J}p_{ij}\pi_{i,t-1|t-1}(\boldsymbol{\theta})f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})\right)\pi_{a,t-1|t-1}(\boldsymbol{\theta})f_{b}(Y_{t};X_{b,t}(\boldsymbol{v}_{b}),\boldsymbol{v}_{b})}{\left(\sum_{k=1}^{J}\sum_{l=1}^{J}p_{lk}\pi_{l,t-1|t-1}(\boldsymbol{\theta})f_{k}(Y_{t};X_{k,t}(\boldsymbol{v}_{k}),\boldsymbol{v}_{k})\right)^{2}}\right|\right]<\infty$$

for all  $a, b \in \{1, ..., J\}$  and

$$\mathbb{E}\left[\log^{+}\sup_{\boldsymbol{\theta}\in\Theta}\left|\frac{\sum_{i=1}^{J}p_{ij}\pi_{i,t-1|t-1}(\boldsymbol{\theta})\nabla_{[\boldsymbol{\upsilon}_{a}]_{b}}f_{a}(Y_{t};X_{a,t}(\boldsymbol{\upsilon}_{a}),\boldsymbol{\upsilon}_{a})}{\sum_{k=1}^{J}\sum_{l=1}^{J}p_{lk}\pi_{l,t-1|t-1}(\boldsymbol{\theta})f_{k}(Y_{t};X_{k,t}(\boldsymbol{\upsilon}_{k}),\boldsymbol{\upsilon}_{k})}\right|\right]<\infty$$

and

$$\mathbb{E}\left[\log^{+}\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}\left|\frac{\left(\sum_{i=1}^{J}p_{ij}\pi_{i,t-1|t-1}(\boldsymbol{\theta})f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})\right)\left(\sum_{l=1}^{J}p_{la}\pi_{l,t-1|t-1}(\boldsymbol{\theta})\nabla_{[\boldsymbol{v}_{a}]_{b}}f_{a}(Y_{t};X_{a,t}(\boldsymbol{v}_{a}),\boldsymbol{v}_{a})\right)}{\left(\sum_{k=1}^{J}\sum_{l=1}^{J}p_{lk}\pi_{l,t-1|t-1}(\boldsymbol{\theta})f_{k}(Y_{t};X_{k,t}(\boldsymbol{v}_{k}),\boldsymbol{v}_{k})\right)^{2}}\right|\right]<\infty$$

for all  $a \in \{1, ..., J\}$  and  $b \in \{1, ..., d_a\}$ , see Appendix E for details. Condition (i) thus follows since, for each  $j \in \{1, ..., J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left| \bar{\nabla}_{x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j) \right|^{m_j} \right] < \infty$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \left|\left|\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j)\right|\right|_2^{2k_j}\right] < \infty$ , and  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{x_j \in \mathcal{X}_j} \left|\left|\bar{\nabla}_{\boldsymbol{v}_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j)\right|\right|_2^{m_j} \right] < \infty$  by assumption.

Let  $r \in \mathbb{N}$  be given. For each  $(n, m) \in \{1, ..., J - 1\} \times \{1, ..., d\}$ , we have that

$$\begin{split} \left[ (\boldsymbol{\phi}_{t}^{\boldsymbol{\pi}})^{(r)}(\mathbf{x};\boldsymbol{\theta}) \right]_{(n-1)d+m} &= \sum_{j=1}^{J-1} \left( \bar{\nabla}_{\left[\boldsymbol{\pi}\right]_{j}} \left[ \boldsymbol{\phi}_{t}^{(r)}(\boldsymbol{\pi}_{t-r|t-r}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{\left[\boldsymbol{\pi}\right]_{J}} \left[ \boldsymbol{\phi}_{t}^{(r)}(\boldsymbol{\pi}_{t-r|t-r}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} \right) [\mathbf{x}]_{(j-1)d+m} \\ &+ \bar{\nabla}_{\left[\boldsymbol{\theta}\right]_{m}} \left[ \boldsymbol{\phi}_{t}^{(r)}(\boldsymbol{\pi}_{t-r|t-r}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} \end{split}$$

for all  $\mathbf{x} \in \mathbb{R}^{(J-1)d}$ . Note that

$$||\phi_t^{(r)}(\mathbf{x}; \boldsymbol{\theta}) - \phi_t^{(r)}(\mathbf{y}; \boldsymbol{\theta})||_2 \le \Lambda(\phi_t^{(r)}; \boldsymbol{\theta})||\mathbf{x} - \mathbf{y}||_2$$

for all  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$  so, for each  $j, n \in \{1, ..., J-1\}$ ,

$$\left| \bar{\nabla}_{\left[\boldsymbol{\pi}\right]_{j}} \left[ \boldsymbol{\phi}_{t}^{(r)}(\boldsymbol{\pi}_{t-r|t-r}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{\left[\boldsymbol{\pi}\right]_{J}} \left[ \boldsymbol{\phi}_{t}^{(r)}(\boldsymbol{\pi}_{t-r|t-r}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} \right| \leq \Lambda(\boldsymbol{\phi}_{t}^{(r)};\boldsymbol{\theta}),$$

see Straumann and Mikosch (2006) for a similar argument. Hence, for each  $(n, m) \in \{1, ..., J - 1\} \times \{1, ..., d\}$ , we have that

$$\left| \left[ (\boldsymbol{\phi}_t^{\boldsymbol{\pi}})^{(r)}(\mathbf{x}; \boldsymbol{\theta}) \right]_{(n-1)d+m} - \left[ (\boldsymbol{\phi}_t^{\boldsymbol{\pi}})^{(r)}(\mathbf{y}; \boldsymbol{\theta}) \right]_{(n-1)d+m} \right| \leq \Lambda(\boldsymbol{\phi}_t^{(r)}; \boldsymbol{\theta}) \sum_{j=1}^J \left| [\mathbf{x}]_{(j-1)d+m} - [\mathbf{y}]_{(j-1)d+m} \right|$$

for all  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^{(J-1)d}$ . Thus,

$$\Lambda((\boldsymbol{\phi}_t^{\boldsymbol{\pi}})^{(r)};\boldsymbol{\theta}) \leq C\Lambda(\boldsymbol{\phi}_t^{(r)};\boldsymbol{\theta})$$

for all  $r \in \mathbb{N}$ . Condition (ii) and (iii) thus follow by using the same arguments as in the proof of Lemma 2.

To show that  $\mathbb{E}[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_2^2]<\infty$ , we have that

$$\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}) = \sum_{k=0}^{\infty} ((\boldsymbol{\phi}_t^{\boldsymbol{\pi}})^{(k)}(\mathbf{V}_{t-k}(\boldsymbol{\theta}); \boldsymbol{\theta}) - (\boldsymbol{\phi}_t^{\boldsymbol{\pi}})^{(k)}(\mathbf{0}; \boldsymbol{\theta})) \quad a.s.,$$

see Lemma C.2, where, by convention,  $(\phi_t^{\boldsymbol{\pi}})^{(0)}(\mathbf{v};\boldsymbol{\theta}) = \mathbf{v}$  and

$$[\mathbf{V}_{t-k}(\boldsymbol{\theta})]_{(n-1)d+m} = \bar{\nabla}_{[\boldsymbol{\theta}]_m} \left[ \phi_{t-k}(\boldsymbol{\pi}_{t-k-1|t-k-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_n, \quad (n,m) \in \{1,...,J-1\} \times \{1,...,d\}.$$

Thus,

$$||\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_{2}^{2} \leq \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} ||(\boldsymbol{\phi}_{t}^{\boldsymbol{\pi}})^{(k)}(\mathbf{V}_{t-k}(\boldsymbol{\theta});\boldsymbol{\theta}) - (\boldsymbol{\phi}_{t}^{\boldsymbol{\pi}})^{(k)}(\mathbf{0};\boldsymbol{\theta})||_{2}||(\boldsymbol{\phi}_{t}^{\boldsymbol{\pi}})^{(l)}(\mathbf{V}_{t-l}(\boldsymbol{\theta});\boldsymbol{\theta}) - (\boldsymbol{\phi}_{t}^{\boldsymbol{\pi}})^{(l)}(\mathbf{0};\boldsymbol{\theta})||_{2} \quad a.s.$$

Note that there exists an  $\alpha \in (0,1)$  such that

$$\sup_{\mathbf{x},\mathbf{y}\in\mathbb{R}^{(J-1)d}}\frac{||(\boldsymbol{\phi}_t^{\boldsymbol{\pi}})^{(k)}(\mathbf{x};\boldsymbol{\theta})-(\boldsymbol{\phi}_t^{\boldsymbol{\pi}})^{(k)}(\mathbf{y};\boldsymbol{\theta})||_2}{||\mathbf{x}-\mathbf{y}||_2}\leq C\alpha^k$$

for all  $k \in \mathbb{N}_0$ . Thus,

$$||\nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_{2}^{2} \leq C \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \alpha^{k} ||\mathbf{V}_{t-k}(\boldsymbol{\theta})||_{2} \alpha^{l} ||\mathbf{V}_{t-l}(\boldsymbol{\theta})||_{2} \quad a.s.$$

Hence, we have that

$$\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_{2}^{2}\right]\leq C\mathbb{E}\left[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\mathbf{V}_{t}(\boldsymbol{\theta})||_{2}^{2}\right],$$

so  $\mathbb{E}[\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_{2}^{2}] < \infty$  since, for each  $j \in \{1,...,J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}|\bar{\nabla}_{x_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})|^{2k_{j}/(k_{j}-1)}\right] < \infty$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}||\nabla_{\boldsymbol{v}_{j}}X_{j,t}(\boldsymbol{v}_{j})||_{2}^{2k_{j}}\right] < \infty$ , and  $\mathbb{E}\left[\sup_{\boldsymbol{v}_{j}\in\boldsymbol{\Upsilon}_{j}}||\bar{\nabla}_{\boldsymbol{v}_{j}}\log f_{j}(Y_{t};X_{j,t}(\boldsymbol{v}_{j}),\boldsymbol{v}_{j})||_{2}^{2}\right] < \infty$  by assumption.

Moreover,  $(\nabla_{\boldsymbol{\theta}} \hat{\pi}_{t|t}(\boldsymbol{\theta}))_{t \in \mathbb{N}}$  is a stochastic process taking values in  $(\mathbb{R}^{(J-1)d}, ||\cdot||_2)$  given by

$$\nabla_{\boldsymbol{\theta}} \hat{\boldsymbol{\pi}}_{t|t}(\boldsymbol{\theta}) = \hat{\boldsymbol{\phi}}_t^{\boldsymbol{\pi}}(\nabla_{\boldsymbol{\theta}} \hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}),$$

where  $(\hat{\phi}_t^{\pi}(\cdot; \boldsymbol{\theta}))_{t \in \mathbb{N}}$  is a sequence of non-stationary Lipschitz functions given by

$$\left[\hat{\boldsymbol{\phi}}_{t}^{\boldsymbol{\pi}}(\mathbf{x};\boldsymbol{\theta})\right]_{(n-1)d+m} = \sum_{j=1}^{J-1} \left(\bar{\nabla}_{[\boldsymbol{\pi}]_{j}} \left[\hat{\boldsymbol{\phi}}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta})\right]_{n} - \bar{\nabla}_{[\boldsymbol{\pi}]_{J}} \left[\hat{\boldsymbol{\phi}}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta})\right]_{n}\right) [\mathbf{x}]_{(j-1)d+m} + \bar{\nabla}_{[\boldsymbol{\theta}]_{m}} \left[\hat{\boldsymbol{\phi}}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta})\right]_{n}, \quad (n,m) \in \{1,...,J-1\} \times \{1,...,d\}.$$

The fact that also  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\nabla_{\boldsymbol{\theta}} \hat{\boldsymbol{\pi}}_{t|t}(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta}} \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_2 \overset{e.a.s.}{\to} 0$  as  $t \to \infty$  follows also from Lemma C.2 if

- (i)  $\mathbb{E}[\log^{+}\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}}\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_{2}]<\infty$ ,
- (ii)  $\sup_{\boldsymbol{\theta}\in\Theta}||\hat{\phi}_t^{\boldsymbol{\pi}}(\mathbf{0};\boldsymbol{\theta}) \phi_t^{\boldsymbol{\pi}}(\mathbf{0};\boldsymbol{\theta})||_2 \stackrel{e.a.s.}{\to} 0 \text{ as } t\to\infty, \text{ and}$
- (iii)  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \Lambda(\hat{\phi}_t^{\boldsymbol{\pi}} \phi_t^{\boldsymbol{\pi}}; \boldsymbol{\theta}) \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty.$

Condition (i) follows since  $\mathbb{E}[\sup_{\theta \in \Theta} ||\nabla_{\theta} \pi_{t|t}(\theta)||_2^2] < \infty$ .

Condition (ii) follows if, for each  $(n,m) \in \{1,...,J-1\} \times \{1,...,d\}$ ,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} |\bar{\nabla}_{[\boldsymbol{\theta}]_m} \left[ \hat{\phi}_t(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_n - \bar{\nabla}_{[\boldsymbol{\theta}]_m} \left[ \phi_t(\boldsymbol{\pi}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_n | \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty.$  This is true if, for each  $j \in \{1,...,J-1\}$ ,

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \frac{\hat{\pi}_{a,t-1|t-1}(\boldsymbol{\theta}) f_b(Y_t; \hat{X}_{b,t}(\boldsymbol{v}_b), \boldsymbol{v}_b)}{\sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \hat{\pi}_{l,t-1|t-1}(\boldsymbol{\theta}) f_k(Y_t; \hat{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k)} - \frac{\pi_{a,t-1|t-1}(\boldsymbol{\theta}) f_b(Y_t; X_{b,t}(\boldsymbol{v}_b), \boldsymbol{v}_b)}{\sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \pi_{l,t-1|t-1}(\boldsymbol{\theta}) f_k(Y_t; X_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k)} \right| \stackrel{e.a.s.}{\to} 0 \quad \text{as} \quad t \to \infty$$

and

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \frac{\left( \sum_{i=1}^{J} p_{ij} \hat{\pi}_{i,t-1|t-1}(\boldsymbol{\theta}) f_{j}(Y_{t}; \hat{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) \right) \hat{\pi}_{a,t-1|t-1}(\boldsymbol{\theta}) f_{b}(Y_{t}; \hat{X}_{b,t}(\boldsymbol{v}_{b}), \boldsymbol{v}_{b})}{\left( \sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \hat{\pi}_{l,t-1|t-1}(\boldsymbol{\theta}) f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) \right)^{2}} - \frac{\left( \sum_{i=1}^{J} p_{ij} \pi_{i,t-1|t-1}(\boldsymbol{\theta}) f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) \right) \pi_{a,t-1|t-1}(\boldsymbol{\theta}) f_{b}(Y_{t}; X_{b,t}(\boldsymbol{v}_{b}), \boldsymbol{v}_{b})}{\left( \sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \pi_{l,t-1|t-1}(\boldsymbol{\theta}) f_{k}(Y_{t}; X_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) \right)^{2}} \right| \stackrel{e.a.s.}{\to} 0 \quad \text{as} \quad t \to \infty$$

for all  $a, b \in \{1, ..., J\}$  and

$$\sup_{\boldsymbol{\theta} \in \Theta} \left| \frac{\sum_{i=1}^{J} p_{ij} \hat{\pi}_{i,t-1|t-1}(\boldsymbol{\theta}) \nabla_{[\boldsymbol{v}_a]_b} f_a(Y_t; \hat{X}_{a,t}(\boldsymbol{v}_a), \boldsymbol{v}_a)}{\sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \hat{\pi}_{l,t-1|t-1}(\boldsymbol{\theta}) f_k(Y_t; \hat{X}_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k)} - \frac{\sum_{i=1}^{J} p_{ij} \pi_{i,t-1|t-1}(\boldsymbol{\theta}) \nabla_{[\boldsymbol{v}_a]_b} f_a(Y_t; X_{a,t}(\boldsymbol{v}_a), \boldsymbol{v}_a)}{\sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \pi_{l,t-1|t-1}(\boldsymbol{\theta}) f_k(Y_t; X_{k,t}(\boldsymbol{v}_k), \boldsymbol{v}_k)} \right| \stackrel{e.a.s.}{\to} 0 \quad \text{as} \quad t \to \infty$$

and

$$\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left| \frac{\left( \sum_{i=1}^{J} p_{ij} \hat{\pi}_{i,t-1|t-1}(\boldsymbol{\theta}) f_{j}(Y_{t}; \hat{X}_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) \right) \left( \sum_{l=1}^{J} p_{la} \hat{\pi}_{l,t-1|t-1}(\boldsymbol{\theta}) \nabla_{[\boldsymbol{v}_{a}]_{b}} f_{a}(Y_{t}; \hat{X}_{a,t}(\boldsymbol{v}_{a}), \boldsymbol{v}_{a}) \right)}{\left( \sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \hat{\pi}_{l,t-1|t-1}(\boldsymbol{\theta}) f_{k}(Y_{t}; \hat{X}_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) \right)^{2}} - \frac{\left( \sum_{i=1}^{J} p_{ij} \pi_{i,t-1|t-1}(\boldsymbol{\theta}) f_{j}(Y_{t}; X_{j,t}(\boldsymbol{v}_{j}), \boldsymbol{v}_{j}) \right) \left( \sum_{l=1}^{J} p_{la} \pi_{l,t-1|t-1}(\boldsymbol{\theta}) \nabla_{[\boldsymbol{v}_{a}]_{b}} f_{a}(Y_{t}; X_{a,t}(\boldsymbol{v}_{a}), \boldsymbol{v}_{a}) \right)}{\left( \sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \pi_{l,t-1|t-1}(\boldsymbol{\theta}) f_{k}(Y_{t}; X_{k,t}(\boldsymbol{v}_{k}), \boldsymbol{v}_{k}) \right)^{2}} \right| \stackrel{e.a.s.}{\longrightarrow} 0 \quad \text{as} \quad t \to \infty$$

for all  $a \in \{1, ..., J\}$  and  $b \in \{1, ..., d_a\}$ , see Appendix E for details once again. Condition (ii) thus follows by using the same arguments as in the proof of Lemma 2 since  $(Y_t)_{t \in \mathbb{Z}}$  is stationary by Assumption 1. Moreover, for each  $j \in \{1, ..., J\}$ ,  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} |\hat{X}_{j,t}(\boldsymbol{v}_j) - X_{j,t}(\boldsymbol{v}_j)| \overset{e.a.s.}{\to} 0$  as  $t \to \infty$  by Lemma 1,  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\hat{\boldsymbol{\pi}}_{t|t}(\boldsymbol{\theta}) - \boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_2 \overset{e.a.s.}{\to} 0$  as  $t \to \infty$  by Lemma 2, and, for each  $j \in \{1, ..., J\}$ ,  $\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j} \hat{X}_{j,t}(\boldsymbol{v}_j) - \nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j)||_2 \overset{e.a.s.}{\to} 0$  as  $t \to \infty$  by Lemma 3. Finally, for each  $j \in \{1, ..., J\}$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{\boldsymbol{x}_j \in \mathcal{X}_j} |\bar{\nabla}_{x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j)|^{m_j}\right] < \infty$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} ||\nabla_{\boldsymbol{v}_j} X_{j,t}(\boldsymbol{v}_j)||_2^{2k_j}\right] < \infty$ ,  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{\boldsymbol{x}_j \in \mathcal{X}_j} ||\bar{\nabla}_{\boldsymbol{v}_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j)||_2^{m_j}\right] < \infty$ , and  $\mathbb{E}\left[\sup_{\boldsymbol{v}_j \in \boldsymbol{\Upsilon}_j} \sup_{\boldsymbol{x}_j \in \mathcal{X}_j} ||\bar{\nabla}_{\boldsymbol{v}_j x_j} \log f_j(Y_t; x_j, \boldsymbol{v}_j)||_2^{m_j}\right] < \infty$  by assumption. By using the same arguments as above,

$$\Lambda(\hat{\phi}_t^{\pi} - \phi_t^{\pi}; \boldsymbol{\theta}) \le C\Lambda(\hat{\phi}_t - \phi_t; \boldsymbol{\theta}).$$

Condition (iii) thus follows by using the same arguments as in the proof of Lemma 2.

#### B.4 Proof of Lemma 6

First,  $(\nabla_{\theta\theta}\pi_{t|t}(\theta))_{t\in\mathbb{Z}}$  is a stochastic process taking values in  $(\mathbb{R}^{(J-1)d^2}, ||\cdot||_2)$  given by

$$\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta}) = \phi_t^{\boldsymbol{\pi}\boldsymbol{\pi}}(\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\boldsymbol{\pi}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}),$$

where  $(\phi_t^{\pi\pi}(\cdot; \boldsymbol{\theta}))_{t \in \mathbb{Z}}$  is a sequence of stationary and ergodic Lipschitz functions given by

$$\begin{split} &[\phi_{t}^{\pi\pi}(\mathbf{x};\boldsymbol{\theta})]_{(n-1)d^{2}+(m_{1}-1)d+m_{2}} \\ &= \sum_{j=1}^{J-1} \left( \bar{\nabla}_{[\pi]_{j}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\pi]_{J}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} \right) [\mathbf{x}]_{(j-1)d^{2}+(m_{1}-1)d+m_{2}} \\ &+ \sum_{j=1}^{J-1} \left( \bar{\nabla}_{[\pi]_{j}[\boldsymbol{\theta}]_{m_{2}}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\pi]_{J}[\boldsymbol{\theta}]_{m_{2}}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} \right) \nabla_{[\boldsymbol{\theta}]_{m_{1}}} \left[ \pi_{t-1|t-1}(\boldsymbol{\theta}) \right]_{j} \\ &+ \sum_{j=1}^{J-1} \sum_{i=1}^{J-1} \left[ \left( \bar{\nabla}_{[\pi]_{j}[\pi]_{i}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\pi]_{J}[\pi]_{J}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} \right) \\ &- \left( \bar{\nabla}_{[\pi]_{J}[\pi]_{i}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\pi]_{J}[\pi]_{J}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} \right) \right] \\ &\cdot \nabla_{[\boldsymbol{\theta}]_{m_{1}}} \left[ \pi_{t-1|t-1}(\boldsymbol{\theta}) \right]_{j} \nabla_{[\boldsymbol{\theta}]_{m_{2}}} \left[ \pi_{t-1|t-1}(\boldsymbol{\theta}) \right]_{i} \\ &+ \sum_{i=1}^{J-1} \left( \bar{\nabla}_{[\boldsymbol{\theta}]_{m_{1}}[\pi]_{i}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\boldsymbol{\theta}]_{m_{1}}[\pi]_{J}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n} \right) \nabla_{[\boldsymbol{\theta}]_{m_{2}}} \left[ \pi_{t-1|t-1}(\boldsymbol{\theta}) \right]_{i} \\ &+ \bar{\nabla}_{[\boldsymbol{\theta}]_{m_{1}}[\boldsymbol{\theta}]_{m_{2}}} \left[ \phi_{t}(\pi_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_{n}, \quad (n,m_{1},m_{2}) \in \{1,...,J-1\} \times \{1,...,d\} \times \{1,...,d\}, \end{split}$$

with Lipschitz coefficient

$$\Lambda(\phi_t^{\pi\pi}; \boldsymbol{\theta}) = \sup_{\substack{\mathbf{x}, \mathbf{y} \in \mathbb{R}^{(J-1)d^2} \\ \mathbf{x} \neq \mathbf{y}}} \frac{||\phi_t^{\pi\pi}(\mathbf{x}; \boldsymbol{\theta}) - \phi_t^{\pi\pi}(\mathbf{y}; \boldsymbol{\theta})||_2}{||\mathbf{x} - \mathbf{y}||_2}.$$

The fact that  $(\nabla_{\theta\theta}\pi_{t|t}(\theta))_{t\in\mathbb{Z}}$  is stationary and ergodic for all  $\theta\in\Theta$  with  $\mathbb{E}[\sup_{\theta\in\Theta}||\nabla_{\theta\theta}\pi_{t|t}(\theta)||_2]<\infty$  follows from Lemma C.2 if

- (i)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\phi_t^{\boldsymbol{\pi}\boldsymbol{\pi}}(\boldsymbol{0}; \boldsymbol{\theta})||_2] < \infty$ ,
- (ii)  $\mathbb{E}[\log^+ \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \Lambda(\boldsymbol{\phi}_t^{\boldsymbol{\pi}\boldsymbol{\pi}}; \boldsymbol{\theta})] < \infty$ , and
- (iii) there exists an  $r \in \mathbb{N}$  such that  $\mathbb{E}[\log \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \Lambda((\boldsymbol{\phi}_t^{\boldsymbol{\pi}\boldsymbol{\pi}})^{(r)}; \boldsymbol{\theta})] < 0$ .

The proof follows the same lines as the proof of Lemma 5 so we omit the details. Moreover,  $(\nabla_{\theta\theta}\hat{\pi}_{t|t}(\theta))_{t\in\mathbb{N}}$  is a stochastic process taking values in  $(\mathbb{R}^{(J-1)d^2}, ||\cdot||_2)$  given by

$$\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{\pi}_{t|t}(\boldsymbol{\theta}) = \hat{\phi}_t^{\boldsymbol{\pi}\boldsymbol{\pi}}(\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\hat{\pi}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}),$$

where  $(\hat{\phi}_t^{\pi\pi}(\cdot;\boldsymbol{\theta}))_{t\in\mathbb{N}}$  is a sequence of non-stationary Lipschitz functions given by

$$\begin{split} & \left[ \hat{\phi}_t^{\boldsymbol{\pi}\boldsymbol{\pi}}(\mathbf{x};\boldsymbol{\theta}) \right]_{(n-1)d^2 + (m_1 - 1)d + m_2} \\ &= \sum_{j=1}^{J-1} \left( \bar{\nabla}_{[\boldsymbol{\pi}]_j} \left[ \hat{\phi}_t(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_n - \bar{\nabla}_{[\boldsymbol{\pi}]_J} \left[ \hat{\phi}_t(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta});\boldsymbol{\theta}) \right]_n \right) [\mathbf{x}]_{(j-1)d^2 + (m_1 - 1)d + m_2} \end{split}$$

$$\begin{split} &+\sum_{j=1}^{J-1} \left( \bar{\nabla}_{[\boldsymbol{\pi}]_{j}[\boldsymbol{\theta}]_{m_{2}}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\boldsymbol{\pi}]_{J}[\boldsymbol{\theta}]_{m_{2}}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} \right) \nabla_{[\boldsymbol{\theta}]_{m_{1}}} \left[ \hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}) \right]_{j} \\ &+\sum_{j=1}^{J-1} \sum_{i=1}^{J-1} \left[ \left( \bar{\nabla}_{[\boldsymbol{\pi}]_{j}[\boldsymbol{\pi}]_{i}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\boldsymbol{\pi}]_{j}[\boldsymbol{\pi}]_{J}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} \right) \\ &- \left( \bar{\nabla}_{[\boldsymbol{\pi}]_{J}[\boldsymbol{\pi}]_{i}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\boldsymbol{\pi}]_{J}[\boldsymbol{\pi}]_{J}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} \right) \right] \\ &\cdot \nabla_{[\boldsymbol{\theta}]_{m_{1}}} \left[ \hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}) \right]_{j} \nabla_{[\boldsymbol{\theta}]_{m_{2}}} \left[ \hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}) \right]_{i} \\ &+ \sum_{i=1}^{J-1} \left( \bar{\nabla}_{[\boldsymbol{\theta}]_{m_{1}}[\boldsymbol{\pi}]_{i}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} - \bar{\nabla}_{[\boldsymbol{\theta}]_{m_{1}}[\boldsymbol{\pi}]_{J}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n} \right) \nabla_{[\boldsymbol{\theta}]_{m_{2}}} \left[ \hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}) \right]_{i} \\ &+ \bar{\nabla}_{[\boldsymbol{\theta}]_{m_{1}}[\boldsymbol{\theta}]_{m_{2}}} \left[ \hat{\phi}_{t}(\hat{\boldsymbol{\pi}}_{t-1|t-1}(\boldsymbol{\theta}); \boldsymbol{\theta}) \right]_{n}, \quad (n, m_{1}, m_{2}) \in \{1, ..., J-1\} \times \{1, ..., d\} \times \{1, ..., d\}, \end{split}$$

The fact that also  $\sup_{\boldsymbol{\theta} \in \Theta} ||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \hat{\pi}_{t|t}(\boldsymbol{\theta}) - \nabla_{\boldsymbol{\theta}\boldsymbol{\theta}} \pi_{t|t}(\boldsymbol{\theta})||_2 \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  follows also from Lemma C.2 if

- (i)  $\mathbb{E}[\log^{+}\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}}||\nabla_{\boldsymbol{\theta}\boldsymbol{\theta}}\boldsymbol{\pi}_{t|t}(\boldsymbol{\theta})||_{2}]<\infty$ ,
- (ii)  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} ||\hat{\phi}_t^{\boldsymbol{\pi}\boldsymbol{\pi}}(\mathbf{0};\boldsymbol{\theta}) \phi_t^{\boldsymbol{\pi}\boldsymbol{\pi}}(\mathbf{0};\boldsymbol{\theta})||_2 \overset{e.a.s.}{\to} 0 \text{ as } t \to \infty, \text{ and}$
- (iii)  $\sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \Lambda(\hat{\phi}_t^{\boldsymbol{\pi}\boldsymbol{\pi}} \phi_t^{\boldsymbol{\pi}\boldsymbol{\pi}}; \boldsymbol{\theta}) \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty.$

We omit the details once again for the same reason as above.

#### C Technical Lemmata

Let  $c \in \mathbb{C}$  where  $\mathbb{C} \subset \mathbb{R}^d$  is a compact set be given. Let  $(Y_t(c))_{t \in \mathbb{Z}}$  be a stochastic process taking values in  $(Y, ||\cdot||_{2,2})$  where  $Y \subseteq \mathbb{R}^{d_1' \times d_2'}$  is a complete set and  $||y||_{2,2} = (\sum_{i=1}^{d_1'} \sum_{j=1}^{d_2'} |y_{ij}|^2)^{1/2}$  given by

$$Y_t(c) = \phi_t(Y_{t-1}(c); c),$$

where  $(\phi_t(\cdot;c))_{t\in\mathbb{Z}}$  is a sequence of stationary and ergodic Lipschitz functions with Lipschitz coefficient

$$\Lambda(\phi_t; c) = \sup_{\substack{x, y \in Y \\ x \neq y}} \frac{||\phi_t(x; c) - \phi_t(y; c)||_{2,2}}{||x - y||_{2,2}}.$$

Assume that  $(y,c) \mapsto \phi_t(y;c)$  is continuous. Then,  $(Y_t)_{t \in \mathbb{Z}}$  is a stochastic process taking values in  $(\mathcal{C}(C,Y),||\cdot||_C)$  where  $\mathcal{C}(C,Y)$  is the set of continuous functions from C to Y and  $||Y||_C = \sup_{c \in C} ||Y(c)||_{2,2}$  given by

$$Y_t = \Phi_t(Y_{t-1})$$

with

$$\Phi_t(Y) = \phi_t(Y(\cdot); \cdot),$$

where  $(\Phi_t(\cdot))_{t\in\mathbb{Z}}$  is a sequence of stationary and ergodic Lipschitz maps with Lipschitz coefficient

$$\Lambda(\Phi_t) = \sup_{\substack{X,Y \in \mathcal{C}(\mathbf{C},\mathbf{Y})\\X \neq Y}} \frac{||\Phi_t(X) - \Phi_t(Y)||_{\mathbf{C}}}{||X - Y||_{\mathbf{C}}}.$$

First, we have the following result.

#### Lemma C.1. Assume that

- (i) there exists a  $y \in Y$  such that  $\mathbb{E}[\log^+ \sup_{c \in C} ||\phi_t(y;c) y||_{2,2}] < \infty$ ,
- (ii)  $\mathbb{E}[\log^+ \sup_{c \in \mathcal{C}} \Lambda(\phi_t; c)] < \infty$ , and
- (iii) there exists an  $r \in \mathbb{N}$  such that  $\mathbb{E}[\log \sup_{c \in \mathcal{C}} \Lambda(\phi_t^{(r)}; c)] < 0$ .

Then, the stochastic process  $(Y_t)_{t\in\mathbb{Z}}$  is stationary and ergodic and

$$Y_t = \lim_{m \to \infty} \Phi_t \circ \cdots \circ \Phi_{t-m}(Y)$$
 a.s.

for all  $Y \in \mathcal{C}(C, Y)$ .

If  $(X_t)_{t\in\mathbb{N}}$  is another stochastic process given by

$$X_t = \Phi_t(X_{t-1})$$

with

$$\Phi_t(X) = \phi_t(X(\cdot); \cdot),$$

then

$$||X_t - Y_t||_{\mathcal{C}} \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty.$$

Proof. The conclusions follow from Theorem 3.1 in Bougerol (1993) if

- (I) there exists a  $Y \in \mathcal{C}(C, Y)$  such that  $\mathbb{E}[\log^+ ||\Phi_t(Y) Y||_C] < \infty$ ,
- (II)  $\mathbb{E}[\log^+ \Lambda(\Phi_t)] < \infty$ , and
- (III) there exists an  $r \in \mathbb{N}$  such that  $\mathbb{E}[\log \Lambda(\Phi_t^{(r)})] < 0$ .

Let  $Y \in \mathcal{C}(C, Y)$  be given by Y(c) = y. Condition (I) then follows since  $\mathbb{E}[\log^+ ||\Phi_t(Y) - Y||_C] = \mathbb{E}[\log^+ \sup_{c \in C} ||\phi_t(y; c) - y||_{2,2}] < \infty$  by Condition (i). We have that

$$\begin{split} &\Lambda(\Phi_t) = \sup_{\substack{X,Y \in \mathcal{C}(\mathbf{C},\mathbf{Y})\\X \neq Y}} \frac{||\Phi_t(X) - \Phi_t(Y)||_{\mathbf{C}}}{||X - Y||_{\mathbf{C}}} \\ &= \sup_{\substack{X,Y \in \mathcal{C}(\mathbf{C},\mathbf{Y})\\X \neq Y}} \sup_{c \in \mathbf{C}} \frac{||\phi_t(X(c);c) - \phi_t(Y(c);c)||_{2,2}}{||X(c) - Y(c)||_{2,2}} \frac{||X(c) - Y(c)||_{2,2}}{\sup_{c \in \mathbf{C}} ||X(c) - Y(c)||_{2,2}} \\ &\leq \sup_{c \in \mathbf{C}} \sup_{x,y \in \mathbf{Y}} \frac{||\phi_t(x;c) - \phi_t(y;c)||_{2,2}}{||x - y||_{2,2}} = \sup_{c \in \mathbf{C}} \Lambda(\phi_t;c). \end{split}$$

Condition (II) and (III) then follow since  $\mathbb{E}[\log^+ \Lambda(\Phi_t)] \leq \mathbb{E}[\log^+ \sup_{c \in \mathcal{C}} \Lambda(\phi_t; c)] < \infty$  and there exists an  $r \in \mathbb{N}$  such that  $\mathbb{E}[\log \Lambda(\Phi_t^{(r)})] \leq \mathbb{E}[\log \sup_{c \in \mathcal{C}} \Lambda(\phi_t^{(r)}; c)] < 0$  by Condition (ii) and (iii), respectively.

Moreover, we have the following result when  $Y = \mathbb{R}^{d_1' \times d_2'}$ .

#### Lemma C.2. Assume that

(i)  $\mathbb{E}[\log^+ \sup_{c \in C} ||\phi_t(0; c)||_{2,2}] < \infty$ ,

- (ii)  $\mathbb{E}[\log^+ \sup_{c \in \mathcal{C}} \Lambda(\phi_t; c)] < \infty$ , and
- (iii) there exists an  $r \in \mathbb{N}$  such that  $\mathbb{E}[\log \sup_{c \in C} \Lambda(\phi_t^{(r)}; c)] < 0$ .

Then, the stochastic process  $(Y_t)_{t\in\mathbb{Z}}$  is stationary and ergodic and

$$Y_t = \lim_{m \to \infty} \Phi_t \circ \cdots \circ \Phi_{t-m}(Y)$$
 a.s.

for all  $Y \in \mathcal{C}(C, \mathbb{R}^{d'_1 \times d'_2})$ .

Assume that  $\mathbb{E}[\log^+ ||Y_t||_{\mathcal{C}}] < \infty$ . Let  $(X_t)_{t \in \mathbb{N}}$  be another stochastic process given by

$$X_t = \hat{\Phi}_t(X_{t-1})$$

with

$$\hat{\Phi}_t(X) = \hat{\phi}_t(X(\cdot); \cdot),$$

where  $(\hat{\Phi}_t(\cdot))_{t\in\mathbb{N}}$  is a sequence of Lipschitz maps. Assume that

(iv) 
$$\sup_{c \in \mathbb{C}} ||\hat{\phi}_t(0;c) - \phi_t(0;c)||_{2,2} \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty \text{ and}$$

$$(v) \sup_{c \in C} \Lambda(\hat{\phi}_t - \phi_t; c) \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty.$$

Then,

$$||X_t - Y_t||_C \stackrel{e.a.s.}{\to} 0$$
 as  $t \to \infty$ .

*Proof.* The first conclusion follows from Lemma C.1. The second one follows from Theorem 2.10 in Straumann and Mikosch (2006) if

(IV) 
$$||\hat{\Phi}_t(0) - \Phi_t(0)||_{\mathcal{C}} \stackrel{e.a.s.}{\to} 0$$
 as  $t \to \infty$  and

(V) 
$$\Lambda(\hat{\Phi}_t - \Phi_t) \stackrel{e.a.s.}{\to} 0 \text{ as } t \to \infty.$$

Condition (IV) follows since  $||\hat{\Phi}_t(0) - \Phi_t(0)||_{\mathcal{C}} = \sup_{c \in \mathcal{C}} ||\hat{\phi}_t(0;c) - \phi_t(0;c)||_{2,2} \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  by Condition (iv). As in the proof of Lemma C.1, we have that

$$\begin{split} &\Lambda(\hat{\Phi}_t - \Phi_t) = \sup_{X,Y \in \mathcal{C}(\mathbf{C}, \mathbb{R}^{d_1' \times d_2'})} \frac{||(\hat{\Phi}_t(X) - \Phi_t(X)) - (\hat{\Phi}_t(Y) - \Phi_t(Y))||_{\mathbf{C}}}{||X - Y||_{\mathbf{C}}} \\ &= \sup_{X,Y \in \mathcal{C}(\mathbf{C}, \mathbb{R}^{d_1' \times d_2'})} \sup_{c \in \mathbf{C}} \frac{||(\hat{\phi}_t(X(c); c) - \phi_t(X(c); c)) - (\hat{\phi}_t(Y(c); c) - \phi_t(Y(c); c))||_{2,2}}{||X(c) - Y(c)||_{2,2}} \frac{||X(c) - Y(c)||_{2,2}}{\sup_{c \in \mathbf{C}} ||X(c) - Y(c)||_{2,2}} \\ &\leq \sup_{c \in \mathbf{C}} \sup_{x,y \in \mathbb{R}^{d_1' \times d_2'}} \frac{||(\hat{\phi}_t(x; c) - \phi_t(x; c)) - (\hat{\phi}_t(y; c) - \phi_t(y; c))||_{2,2}}{||x - y||_{2,2}} = \sup_{c \in \mathbf{C}} \Lambda(\hat{\phi}_t - \phi_t; c). \end{split}$$

Condition (V) then follows since  $\Lambda(\hat{\Phi}_t - \Phi_t) \leq \sup_{c \in \mathcal{C}} \Lambda(\hat{\phi}_t - \phi_t; c) \stackrel{e.a.s.}{\to} 0$  as  $t \to \infty$  by Condition (v).

# D Other Lemmata

We have the following result.

**Lemma D.1.** Let **P** be a  $J \times J$ -dimensional stochastic matrix with generic element  $p_{ij}$ . Assume that there exist an  $\varepsilon \in (0,1)$  and a  $\rho \in \mathcal{S}$  such that

$$p_{ij} \ge \varepsilon \rho_j$$

for all  $i, j \in \{1, ..., J\}$ . Then, there exists an  $\alpha \in (0, 1)$  such that

$$||\mathbf{P}'\mathbf{x} - \mathbf{P}'\mathbf{y}||_1 \le \alpha ||\mathbf{x} - \mathbf{y}||_1$$

for all  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$ .

*Proof.* Let  $\tilde{\mathbf{P}}$  be a  $J \times J$ -dimensional matrix with generic element  $\tilde{p}_{ij} = (1 - \varepsilon)^{-1}(p_{ij} - \varepsilon \rho_j)$ . Note that  $\tilde{\mathbf{P}}$  is a stochastic matrix. Thus,

$$||\mathbf{P}'\mathbf{x} - \mathbf{P}'\mathbf{y}||_{1} = \sum_{j=1}^{J} \left| \sum_{i=1}^{J} p_{ij}(x_{i} - y_{i}) \right|$$

$$= (1 - \varepsilon) \sum_{j=1}^{J} \left| \sum_{i=1}^{J} \tilde{p}_{ij}(x_{i} - y_{i}) \right|$$

$$\leq (1 - \varepsilon) \sum_{j=1}^{J} \sum_{i=1}^{J} \tilde{p}_{ij}|x_{i} - y_{i}|$$

$$= (1 - \varepsilon) \sum_{i=1}^{J} |x_{i} - y_{i}| = (1 - \varepsilon)||\mathbf{x} - \mathbf{y}||_{1}$$

for all  $\mathbf{x}, \mathbf{y} \in \mathcal{S}$ .

# E Supplementary Material

In the following,  $\pi_{j,t|t}$  denotes  $\pi_{j,t|t}(\boldsymbol{\theta})$  and  $f_{j,t}$  denotes  $f_j(Y_t; X_{j,t}(\boldsymbol{v}_j), \boldsymbol{v}_j)$ . For each  $j \in \{1, ..., J-1\}$ ,

$$\begin{split} \nabla_{p_{ab}}\pi_{j,t|t} &= 1_{\{j=b\}} \frac{\pi_{a,t-1|t-1}f_{j,t}}{\sum_{k=1}^{J}\sum_{l=1}^{J}p_{lk}\pi_{l,t-1|t-1}f_{k,t}} \\ &+ \frac{\sum_{i=1}^{J-1}(p_{ij}-p_{Jj})\nabla_{p_{ab}}\pi_{i,t-1|t-1}f_{j,t}}{\sum_{k=1}^{J}\sum_{l=1}^{J}p_{lk}\pi_{l,t-1|t-1}f_{k,t}} \\ &- \frac{\left(\sum_{i=1}^{J}p_{ij}\pi_{i,t-1|t-1}f_{j,t}\right)\pi_{a,t-1|t-1}(f_{b,t}-f_{J,t})}{\left(\sum_{k=1}^{J}\sum_{l=1}^{J}p_{lk}\pi_{l,t-1|t-1}f_{k,t}\right)^{2}} \\ &- \frac{\left(\sum_{i=1}^{J}p_{ij}\pi_{i,t-1|t-1}f_{j,t}\right)\left(\sum_{k=1}^{J-1}\sum_{l=1}^{J-1}(p_{lk}-p_{Jk})\nabla_{p_{ab}}\pi_{l,t-1|t-1}(f_{k,t}-f_{J,t})\right)}{\left(\sum_{k=1}^{J}\sum_{l=1}^{J}p_{lk}\pi_{l,t-1|t-1}f_{k,t}\right)^{2}} \end{split}$$

for all  $a \in \{1,...,J\}$  and  $b \in \{1,...,J-1\}$  and

$$\begin{split} \nabla_{[\boldsymbol{v}_a]_b} \pi_{j,t|t} &= \frac{\sum_{i=1}^{J-1} (p_{ij} - p_{Jj}) \nabla_{[\boldsymbol{v}_a]_b} \pi_{i,t-1|t-1} f_{j,t}}{\sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \pi_{l,t-1|t-1} f_{k,t}} \\ &+ 1_{\{j=a\}} 1_{\{a \in \{1, \dots, J-1\}\}} \frac{\sum_{i=1}^{J} p_{ij} \pi_{i,t-1|t-1} \nabla_{[\boldsymbol{v}_a]_b} f_{j,t}}{\sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \pi_{l,t-1|t-1} f_{k,t}} \\ &- \frac{\left(\sum_{i=1}^{J} p_{ij} \pi_{i,t-1|t-1} f_{j,t}\right) \left(\sum_{k=1}^{J-1} \sum_{l=1}^{J-1} (p_{lk} - p_{Jk}) \nabla_{[\boldsymbol{v}_a]_b} \pi_{l,t-1|t-1} (f_{k,t} - f_{J,t})\right)}{\left(\sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \pi_{l,t-1|t-1} f_{k,t}\right)^2} \\ &- \frac{\left(\sum_{i=1}^{J} p_{ij} \pi_{i,t-1|t-1} f_{j,t}\right) \left(\sum_{l=1}^{J} p_{la} \pi_{l,t-1|t-1} \nabla_{[\boldsymbol{v}_a]_b} f_{a,t}\right)}{\left(\sum_{k=1}^{J} \sum_{l=1}^{J} p_{lk} \pi_{l,t-1|t-1} f_{k,t}\right)^2} \end{split}$$

for all  $a \in \{1, ..., J\}$  and  $b \in \{1, ..., d_a\}$ .