# On the Missing Factor in Some Concentration Inequalities for Martingales

Arun Kumar Kuchibhotla

Department of Statistics & Data Science, Carnegie Mellon University

#### Abstract

In this note, we improve some concentration inequalities for martingales with bounded increments. These results recover the missing factor in Freedman-style inequalities and are near optimal. We also provide minor refinements of concentration inequalities for functions of independent random variables. These proofs use techniques from the works of Bentkus and Pinelis.

#### **1** Introduction

Concentration inequalities such as Hoeffding's inequality are central to modern (e.g., high-dimensional) statistical theory and inference. For sums of independent random variables that are mean zero and bounded, the classical Hoeffding, Bernstein, and Bennett inequalities have been improved in a series of works by Bentkus and Pinelis; see, for example, Bentkus (2002, 2004); Pinelis (2006a, 2014a). These improvements recover the so-called "missing factor" in classical inequalities (Talagrand, 1995). Moreover, these inequalities are nearly optimal.

The goal of this article is to bring about similar improvements in martingale inequalities with bounded increments. In particular, we improve on Freedman's inequality and its refinements derived in Fan et al. (2012, 2015). To briefly mention the result, note that the traditional concentration inequalities are proved using  $\mathbf{1}\{u \ge 0\} \le \exp(\lambda u)$  for any  $\lambda \ge 0$ . The improvements proposed in the works of Bentkus and Pinelis start with the inequality

$$\mathbf{1}\{u \ge 0\} \leqslant \left(1 + \frac{\lambda u}{\alpha}\right)_{+}^{\alpha} \leqslant \exp(\lambda u) \quad \text{for all} \quad \lambda \ge 0, u \in \mathbb{R}$$

Hence, for any  $x \in \mathbb{R}$  and any set of random variables  $X_1, \ldots, X_n$ ,

$$\mathbb{P}\left(\sum_{i=1}^{n} X_i \ge x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^{n} X_i - x\right)\right)_+^{\alpha}\right] \le \inf_{\lambda \ge 0} \mathbb{E}\left[\exp\left(\lambda\left(\sum_{i=1}^{n} X_i - x\right)\right)\right].$$

The right hand side is the classical Cramer-Chernoff bound on the tail probability. The intermediate bound is the improvement suggested in Bentkus (2002), and we refer to these bounds as the Bentkus bound. There are two important differences between these bounds: (1) unlike the Cramer-Chernoff bound, the Bentkus bound is finite as long as the random variables have a finite  $\alpha$ -th moment; and (2) unlike the Cramer-Chernoff bound, Bentkus bound cannot be written as a product if  $X_i$ 's are mutually independent. This second aspect makes it difficult to work with the Bentkus bound in many cases.

The results available for independent random variables are as follows.

1. If  $X_i \in [a_i, b_i], 1 \leq i \leq n$  with  $a_i \leq 0 \leq b_i$  and  $\mathbb{E}[X_i] \leq 0$ , then for any  $x \geq 0$ ,

$$\mathbb{P}\left(\sum_{i=1}^{n} X_i \ge x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \lambda \left(\sum_{i=1}^{n} G_{1,i} - x\right)\right)_+\right] \le \inf_{\lambda \ge 0} \mathbb{E}\left[\exp\left(\lambda \left(\sum_{i=1}^{n} G_{1,i} - x\right)\right)\right],$$

where  $G_{1,i} \in \{a_i, b_i\}, 1 \leq i \leq n$  are independent mean zero random variables. More formally,  $G_{1,i}, 1 \leq i \leq n$  are two-point random variables such that

$$\mathbb{P}(G_{1,i} = a_i) = \frac{b_i}{b_i - a_i}$$
 and  $\mathbb{P}(G_{1,i} = b_i) = \frac{-a_i}{b_i - a_i}$ . (1)

2. If  $X_i \leq b_i, 1 \leq i \leq n$  with  $\mathbb{E}[X_i] \leq 0$ ,  $\operatorname{Var}(X_i) \leq \sigma_i^2$ , then for any  $x \geq 0$ ,

$$\mathbb{P}\left(\sum_{i=1}^{n} X_i \ge x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^{n} G_{2,i} - x\right)\right)_{+}^{2}\right] \le \inf_{\lambda \ge 0} \mathbb{E}\left[\exp\left(\lambda\left(\sum_{i=1}^{n} G_{2,i} - x\right)\right)\right],$$

where  $G_{2,i} \in \{-\sigma_i^2/b_i, b_i\}, 1 \leq i \leq n$  are independent mean zero random variables. More formally,  $G_{2,i}, 1 \leq i \leq n$  are two-point random variables such that

$$\mathbb{P}(G_{2,i} = -\sigma_i^2/b_i) = \frac{b_i^2}{\sigma_i^2 + b_i^2}$$
 and  $\mathbb{P}(G_{2,i} = b_i) = \frac{\sigma_i^2}{\sigma_i^2 + b_i^2}$ .

Note that in both cases, we have lower bounds in terms of  $G_{1,i}$ 's and  $G_{2,i}$ 's, i.e.,

$$\sup_{X_i \in [a_i, b_i], \mathbb{E}[X_i] = 0} \mathbb{P}\left(\sum_{i=1}^n X_i \ge x\right) \ge \mathbb{P}\left(\sum_{i=1}^n G_{1,i} \ge x\right),$$

and

$$\sup_{X_i \leqslant b_i, \mathbb{E}[X_i]=0, \operatorname{Var}(X_i) \leqslant \sigma_i^2} \mathbb{P}\left(\sum_{i=1}^n X_i \geqslant x\right) \geqslant \mathbb{P}\left(\sum_{i=1}^n G_{2,i} \geqslant x\right).$$

The near-optimality of Bentkus bounds is that for  $\alpha \in \{1, 2\}$ , and x in the support of  $\sum_{i=1}^{n} G_{\alpha,i}$ , we have

$$\inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{\alpha} \left(\sum_{i=1}^{n} G_{\alpha,i} - x\right)\right)_{+}^{\alpha}\right] \leq \frac{e^{\alpha}}{\alpha} \mathbb{P}\left(\sum_{i=1}^{n} G_{\alpha,i} \ge x\right).$$
(2)

Therefore, the Bentkus bounds presented above cannot be improved by more than a factor of  $e^{\alpha}/\alpha$ . It should be noted that the left-hand side of (2) is at most one for any  $x \ge 0$  (e.g. by taking  $\lambda = 0$ ), while the right hand side can be larger than 1. Moreover, we note that the optimal Cramer-Chernoff bounds presented above are known in Bennett (1962) and Hoeffding (1963). For readers unfamiliar with the Bentkus bounds, Bentkus (2002) provides an accessible introduction to this literature. As noted in Hoeffding (1963) and Bentkus (2004), the optimal Cramer-Chernoff bound is better than the classical Bernstein, Bennett, and Prokhorov inequalities. The Bentkus bound is also better than Talagrand's inequality (Talagrand, 1995).

One more point to note in comparison with classical concentration inequalities is that the Bentkus bound is not just a function of the variance of the sum, but is a nontrivial function of individual variances. Using some domination inequalities for sums of Bernoulli random variables (Bentkus, 2004, Lemma 4.5), one can obtain (less precise) Bentkus bounds that depend only on  $\sum_{i=1}^{n} (b_i - a_i)^2$  and  $\sum_{i=1}^{n} \sigma_i^2$ , respectively. For example, Pinelis (2006b) proves

$$\sup_{X_i \in [a_i, b_i], \mathbb{E}[X_i] \le 0} \mathbb{P}\left(\sum_{i=1}^n X_i \ge x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{5}(vZ - x)\right)_+^5\right], \quad \text{where} \quad v^2 = \sum_{i=1}^n (b_i - a_i)^2,$$

and Bentkus (2004) proves

$$\sup_{X_i \leq b_i, \mathbb{E}[X_i] \leq 0, \operatorname{Var}(X_i) \leq \sigma_i^2} \mathbb{P}\left(\sum_{i=1}^n X_i \geq x\right) \leq \inf_{\lambda \geq 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^n G'_i - x\right)\right)_+\right]$$

where  $G'_i \in \{-v^2/b, b\}$  with  $b = \max_{1 \le i \le n} b_i$ , and  $v^2 = \sum_{i=1}^n \sigma_i^2$ . These inequalities have been extended to super-martingales in the works cited above, but with the restrictive assumption that  $\sigma_i^2$  represents an almost sure upper bound on the conditional variance of  $X_i$  given  $\mathcal{F}_{i-1}$ . This makes it difficult to obtain concentration inequalities for functions of independent random variables via the Doob martingale. The goal of this note is to obtain results without assuming that the conditional variances are almost surely bounded.

**On the Proof Techniques.** Most of the martingale analogues of the classical concentration inequalities (for independent data) are obtained using the Cramer-Chernoff technique. Some of the inequalities obtained time uniformly (i.e., simultaneously valid for all sample sizes) have also been obtained using the Cramer-Chernoff technique. The main idea in these proofs is the construction of an exponential supermartingale and an application of Ville's or Doob's maximal inequality (Howard et al., 2021). The construction of this exponential supermartingale is based on bounds on the moment generating function. Unfortunately, there is no such simple supermartingale in relation to the Bentkus style bounds, and it is not obvious if there is a general way to extend/improve all the existing Cramer-Chernoff results to the Bentkus style bounds. In fact, it is not clear whether such analogues hold true, but we know that if such an analogue exists, then it is a significant improvement over the Cramer-Chernoff ones.

Our proofs are closely related to the Lindeberg swapping technique commonly used in proving central limit theorems (Lalley, 2014). In the literature on concentration inequalities, the closest result is Lemma 5.2 of Pinelis (1998). Our results arise as direct corollaries of the proof of this lemma. This note exists only because these results have not appeared in the literature in this form before and might be of interest to the broader community.

#### 2 Improving Freedman-style Inequality

Suppose  $(X_i, \mathcal{F}_i)_{i \ge 0}$  with  $X_0 = 0$  forms a (super)martingale difference sequence bounded above by 1, i.e.,  $X_i$  is  $\mathcal{F}_i$ -measurable, and  $\mathbb{E}[X_i|\mathcal{F}_{i-1}] \le 0$ . Let  $\sigma_i^2 := \mathbb{E}[X_i^2|\mathcal{F}_{i-1}]$ . Consider the event

$$\mathcal{E}_k := \left\{ \sum_{i=1}^k X_i \ge x \quad \text{and} \quad \sum_{i=1}^k \sigma_i^2 \le v^2 \right\}.$$

For any  $x \ge 0$  and  $0 < v < \infty$ , Freedman (1975) proved inequalities for  $\mathbb{P}(\bigcup_{k\ge 1}\mathcal{E}_k)$  that can be considered as extensions of the classical Bernstein inequality for independent random variables. Fan et al. (2015) improved Freedman inequalities by proving

$$\mathbb{P}\left(\bigcup_{k=1}^{n} \mathcal{E}_{k}\right) \leqslant \inf_{\lambda \ge 0} \mathbb{E}\left[\exp\left(\lambda\left(\sum_{i=1}^{n} G_{i} - x\right)\right)\right],\tag{3}$$

where  $G_i \in \{-v^2/n, 1\}, 1 \leq i \leq n$  are mutually independent random variables with mean zero, i.e.,

$$\mathbb{P}(G_i = -v^2/n) = \frac{1}{1 + v^2/n} \quad \text{and} \quad \mathbb{P}(G_i = 1) = \frac{v^2/n}{1 + v^2/n}.$$
(4)

(Although Theorem 2.1 of Fan et al. (2015) is not presented in this form, it is equivalent to the form above; see Eq. (41) of Fan et al. (2015).) Moreover, Fan et al. (2015) show that this bound improves many results as described in Remark 2.1 and Corollary 2.2 there.

The following is the main result of this section.

**Theorem 2.1.** Suppose  $(X_i, \mathcal{F}_i), i \ge 0$  forms a supermartingale difference sequence with  $X_0 = 0$ and  $\mathbb{P}(X_i \le 1) = 1^1$  Then

$$\mathbb{P}\left(\bigcup_{k=1}^{n} \mathcal{E}_{k}\right) \leqslant \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^{n} G_{i} - x\right)\right)_{+}^{2}\right] \leqslant \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}(\widetilde{\Pi}_{v^{2}} - x)\right)_{+}^{2}\right], \quad (5)$$

where  $G_i$ 's are independent and identically distributed random variables with distribution specified in (4), and  $\widetilde{\Pi}_{v^2} \sim \text{Poisson}(v^2) - v^2$ . In particular, we obtain

$$\mathbb{P}\left(\sum_{i=1}^{n} X_{i} \geqslant x \quad and \quad \sum_{i=1}^{n} \sigma_{i}^{2} \leqslant v^{2} \quad for \ some \quad n \geqslant 1\right) \leqslant \inf_{\lambda \geqslant 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}(\widetilde{\Pi}_{v^{2}} - x)\right)_{+}^{2}\right] \qquad (6)$$
$$\leqslant \frac{e^{2}}{2} \mathbb{P}^{\circ}(\widetilde{\Pi}_{v^{2}} \geqslant x),$$

where  $x \mapsto \mathbb{P}^{\circ}(\widetilde{\Pi}_{v^2} \ge x)$  is the least log-concave majorant of  $x \mapsto \mathbb{P}(\widetilde{\Pi}_{v^2} \ge x)$ .

To our knowledge, Theorem 2.1 is new. The second inequality of (5) follows from Proposition 2.8 of Pinelis (2014a). Inequality (6) follows from the second inequality of (5) and Lemma 4.2 of Bentkus (2004). The first inequality of (5) is already near optimal for independent data because it is exactly the Bentkus bound for sums of IID mean zero random variables that are bounded above. Moreover, from the works cited in Section 1, the first inequality of (5) is an improvement of Theorem 2.1 of Fan et al. (2015), and therefore an improvement of the Freedman and de la Pena inequalities. In particular, Theorem 2.1 recovers the missing factor similar to how Bentkus' inequality for independent data recovers the missing factor in Bernstein's inequality.

The proof of Theorem 2.1 is provided in Section S.1. Traditional proofs of Freedman's inequality and its improvements are based on the exponential function, and the fact that the exponential of a sum is the product of exponentials significantly helps the analysis. Our proof is a repeated application of Lemma 4.4(i) and Lemma 4.5 of Bentkus (2004).

## 3 Improving Azuma-Hoeffding Inequality

In Section 3, we considered the case of supremartingale differences that are bounded above by a non-random constant. In this section, we obtain an inequality that allows for a random upper bound and yields a generalization of the Azuma-Hoeffding inequality. Suppose  $(X_i, \mathcal{F}_i)_{i\geq 0}$  is a supermartingale difference sequence such that for some  $\mathcal{F}_{i-1}$ -measurable random variables  $B_i$ , we have  $\mathbb{P}(X_i \leq B_i | \mathcal{F}_{i-1}) = 1$  and  $\sigma_i^2 = \mathbb{E}[X_i^2 | \mathcal{F}_{i-1}]$ . Define

$$s_i := \frac{1}{2} \left( B_i + \frac{\sigma_i^2}{B_i} \right)$$
 and  $V_k^2 = \sum_{i=1}^k s_i^2$ .

<sup>&</sup>lt;sup>1</sup>We assume the bound to be 1, without loss of generality.

Consider the event

$$\mathcal{E}_k = \left\{ \sum_{i=1}^k X_i \ge x \text{ and } V_k^2 \le v^2 \right\}.$$

With this notation, we have the following result.

**Theorem 3.1.** For any  $n \ge 1$ ,  $x \ge 0$ , and  $0 < v < \infty$ , with  $Z \sim N(0, 1)$ ,

$$\mathbb{P}\left(\bigcup_{k=1}^{n} \mathcal{E}_{k}\right) \leq \inf_{\lambda \geq 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{5}(vZ - x)\right)_{+}^{5}\right] \leq 5!(e/5)^{5}\mathbb{P}(vZ \geq x).$$

In particular,

$$\mathbb{P}\left(\sum_{i=1}^{n} X_i \ge x \quad and \quad \sum_{i=1}^{n} s_i^2 \le v^2 \text{ for some } n \ge 1\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{5}(vZ - x)\right)_+^5\right]$$

Theorem 3.1 improves inequality (2.22) of Fan et al. (2015) and fills the gap in the literature; see the discussion following Eq. (2.22) of Fan et al. (2015). The proof of Theorem 3.1 follows a structure similar to that of Theorem 2.1 using Lemma 5.1.3 of Pinelis (2006b).

Note that if  $\mathbb{P}(A_i \leq X_i \leq B_i | \mathcal{F}_{i-1}) = 1$  for some non-negative  $\mathcal{F}_{i-1}$ -measurable random variables  $A_i, B_i$ , then  $\sigma_i^2 \leq -A_i B_i$  and hence  $s_i \leq (B_i - A_i)/2$ . This fact combined with Theorem 3.1 implies the following corollary.

**Corollary 3.2.** Suppose  $(X_i, \mathcal{F}_i)_{i \ge 0}$  is a supermartingale difference sequence such that  $\mathbb{P}(A_i \le X_i \le B_i | \mathcal{F}_{i-1}) = 1$  almost surely. Then

$$\mathbb{P}\left(\sum_{i=1}^{n} X_i \ge x \quad and \quad \sum_{i=1}^{n} (B_i - A_i)^2 \le 4v^2 \text{ for some } n \ge 1\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{5}(vZ - x)\right)_+^5\right].$$

Corollary 3.2 improves the classical Azuma-Hoeffding and McDiarmid inequalities. It also improves Theorems 2.5 and 2.6 of van de Geer (2002).

## 4 Unbounded Martingale Increments and Functions of Independent Random Variables

The results in previous sections are concerned with bounded supermartingale differences, and in many practical settings, one often encounters differences that are unbounded but bounded with high probability, for example, via certain tail assumption on the differences. This leads to generalizations of McDiarmid's inequality with unbounded influences. There is a thriving literature on this problem Li and Liu (2023, 2024); Marchina (2018); Maurer and Pontil (2021); Kutin (2002); Maurer (2019); Kontorovich (2014). As with the classical concentration inequalities, most of these are based on the moment generating function and Cramer-Chernoff technique. In the following, we provide some results of the Bentkus type for these problems. Some results in this direction are provided in Marchina (2018).

The following results are based on the framework and closely related to Theorem 2.1 of Marchina (2018), which itself is based on the results of Bentkus (2008, 2010). To present the main result, we need some notation and preliminary results. For any two random variables U and V, we say U is stochastically dominated by V of order  $\alpha \ge 0$  (denoted by  $\le_{\alpha}$ ) if

$$\mathbb{E}[(U-t)_{+}^{\alpha}] \leq \mathbb{E}[(V-t)_{+}^{\alpha}] \quad \text{for all} \quad t \in \mathbb{R}.$$

In particular,  $U \leq_0 V$  is equivalent to  $\mathbb{P}(U > t) \leq \mathbb{P}(V > t)$  for all  $t \in \mathbb{R}$ . Note that  $U \leq_{\alpha} V$  is not equivalent to  $-V \leq_{\alpha} -U$  in general, unless V is absolutely continuous with respect to the Lebesgue measure.

**Lemma 4.1.** Suppose  $X \leq_1 W$  and  $-X \leq_1 -T$  for some random variables T and W satisfying  $T \leq_0 W$ . Then  $\mathbb{E}[T] \leq \mathbb{E}[X] \leq \mathbb{E}[W]$ . Moreover, for any  $q \in [0,1]$ , if  $a_q := \inf\{x : \mathbb{P}(T \leq x) \geq 1-q\}$  and  $b_q := \inf\{x : \mathbb{P}(W \leq x) \geq 1-q\}$ , then the random variable  $\xi_q$  defined by

$$\mathbb{P}(\xi_q \leqslant x) = \begin{cases} \mathbb{P}(T \leqslant x), & \text{if } x < a_q, \\ 1 - q, & \text{if } a_q \leqslant x < b_q, \\ \mathbb{P}(W \leqslant x), & \text{if } x \ge b_q, \end{cases}$$

satisfies the following:

- 1.  $\xi_0 \stackrel{d}{=} T, \, \xi_1 \stackrel{d}{=} W, \, \mathbb{E}[T] \leq \mathbb{E}[\xi_q] \leq \mathbb{E}[W] \text{ for all } q \in [0,1].$
- 2.  $q \mapsto \mathbb{E}[\xi_q]$  is a differentiable non-decreasing function.
- 3. If  $q_0$  is the largest number such that  $\mathbb{E}[\xi_{q_0}] = \mathbb{E}[X]$ , then  $X \leq_1 \xi_{q_0}$ , i.e.,  $\mathbb{E}[(X t)_+] \leq \mathbb{E}[(\xi_{q_0} t)_+]$  for all  $t \in \mathbb{R}$ .

This lemma is an expanded version of Lemma 4.3 of Marchina (2018) and follows from Theorem 1 of Bentkus (2010). In what follows, we are concerned with mean zero random variables and, therefore, we propose the notation  $\xi_{T,W}$  for  $\xi_{q_0}$  in Lemma 4.1 with  $q_0$  chosen so that  $\mathbb{E}[\xi_{q_0}] = 0$ . The reader should note that  $\xi_q$  depends on T and W only through their distributions.

The following result is one of the main results in this section. We remark/stress here that this is a known result and essentially follows by combining Theorem 1 of Bentkus (2010) and the basic proof techniques of Bentkus (2008, 2004). Given the importance of such a result for applications in statistics and machine learning, this result needs to be spelled out explicitly.

**Theorem 4.2.** Suppose  $(X_i, \mathcal{F}_i)_{i \ge 0}$  is a real-valued martingale difference sequence and suppose there exist random variables  $T_i \leq_0 W_i$  such that  $X_i \leq_1 W_i$  and  $-X_i \leq -T_i$  (conditional on  $\mathcal{F}_{i-1}$ ), *i.e.*, with probability 1,

$$\mathbb{E}[(X_i - t)_+ | \mathcal{F}_{i-1}] \leq \mathbb{E}[(T_i - t)_+] \quad and \quad \mathbb{E}[(t - X_i)_+ | \mathcal{F}_{i-1}] \leq \mathbb{E}[(t - W_i)_+] \quad for \ all \quad t \in \mathbb{R}$$

Then, for mutually independent random variables  $\xi_{T_i,W_i}, 1 \leq i \leq n$ , we have

$$\mathbb{E}\left[\left(\sum_{i=1}^{n} X_{i} - t\right)_{+}\right] \leqslant \mathbb{E}\left[\left(\sum_{i=1}^{n} \xi_{T_{i},W_{i}} - t\right)_{+}\right] \quad for \ all \quad t \in \mathbb{R}.$$
(7)

In addition, if  $T_i, W_i$  are absolutely continuous with respect to the Lebesgue measure, then

$$\mathbb{E}\left[\left(t-\sum_{i=1}^{n}X_{i}\right)_{+}\right] \leq \mathbb{E}\left[\left(t+\sum_{i=1}^{n}\xi_{-W_{i},-T_{i}}\right)_{+}\right] \quad for \ all \quad t \in \mathbb{R}.$$
(8)

Consequently, for any  $x \ge 0$ ,

$$\mathbb{P}\left(\sum_{i=1}^{n} X_{i} \ge x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \lambda \left(\sum_{i=1}^{n} \xi_{T_{i},W_{i}} - x\right)\right)_{+}\right],$$

$$\mathbb{P}\left(\sum_{i=1}^{n} X_{i} \le -x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \lambda \left(\sum_{i=1}^{n} \xi_{-W_{i},-T_{i}} - x\right)\right)_{+}\right],$$
(9)

It should be stressed here that  $\xi_{T_i,W_i}$ ,  $1 \leq i \leq n$  are independent and  $T_i, W_i$  are independent of  $\mathcal{F}_n$ . The proof of Theorem 4.2 is omitted because it easily follows from the proofs in Bentkus (2008). Moreover, inequalities (7) and (8) can be combined to yield an improved tail bound in (9) by noting that if  $\mathbb{E}[(R_1-t)_+] \leq \mathbb{E}[(R_2-t)_+]$  for all  $t \in \mathbb{R}$  and  $\mathbb{E}[R_1] = \mathbb{E}[R_2]$ , then  $\mathbb{E}[(t-R_1)_+] \leq \mathbb{E}[(t-R_2)_+]$  for all  $t \in \mathbb{R}$  and  $\mathbb{E}[R_1] = \mathbb{E}[R_2]$ , then  $\mathbb{E}[(t-R_1)_+] \leq \mathbb{E}[(t-R_2)_+]$  for all  $t \in \mathbb{R}$ ; see Bentkus (2008, Proposition 3). This result yields the following important corollary for functions of independent random variables.

**Corollary 4.3.** Suppose  $X_1, \ldots, X_n$  are independent random variables taking values in some space  $\mathcal{X}$  and  $Z = F(X_1, \ldots, X_n)$  for some function  $F : \mathcal{X}^n \to \mathbb{R}$ . For  $1 \leq i \leq n$ , let

$$Z_i = F(X_1, \dots, X_{i-1}, X'_i, X_{i+1}, \dots, X_n), \quad and \quad \Delta_i := Z - Z_i$$

where  $X'_i$  is an independent copy of  $X_i$  (independent of  $X_1, \ldots, X_n$ ). Suppose that we have random variables  $T_i \leq_0 W_i$  such that

$$\mathbb{E}[(\Delta_i - t)_+ | \mathcal{F}_{i-1}] \leq \mathbb{E}[(W_i - t)_+] \quad and \quad \mathbb{E}[(t - \Delta_i)_+ | \mathcal{F}_{i-1}] \leq \mathbb{E}[(t - T_i)_+].$$
(10)

Then

$$\mathbb{E}[(Z - \mathbb{E}[Z] - t)_+] \leq \mathbb{E}\left[\left(\sum_{i=1}^n \xi_{T_i, W_i} - t\right)_+\right] \quad \text{for all} \quad t \in \mathbb{R}$$

Consequently, for all  $x \in \mathbb{R}$ ,

$$\mathbb{P}\left(Z - \mathbb{E}[Z] \ge x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \lambda \left(\sum_{i=1}^{n} \xi_{T_i, W_i} - x\right)\right)_+\right].$$

Corollary 4.3 follows from Theorem 4.2 by taking  $\mathcal{F}_i = \sigma(\{X_1, \ldots, X_i\})$  and  $X_i = \mathbb{E}[\Delta_i | \mathcal{F}_i]$ which is mean zero and forms a martingale difference sequence with respect to  $\mathcal{F}_i$ . More formally, the Doob martingale decomposition yields

$$Z - \mathbb{E}[Z] = \sum_{i=1}^{n} (\mathbb{E}[Z|\mathcal{F}_i] - \mathbb{E}[Z|\mathcal{F}_{i-1}]) = \sum_{i=1}^{n} \mathbb{E}[\Delta_i|\mathcal{F}_i],$$

where  $\mathcal{F}_0$  is the trivial  $\sigma$ -field. This shows that condition (10) can be relaxed by replacing  $\Delta_i$  with  $\mathbb{E}[\Delta_i|\mathcal{F}_i]$ . Given the use of the (first-order) influence in traditional concentration inequalities for functions of independent random variables, we feel that the condition (10) is more natural and intuitive. In Theorem 4.2 and Corollary 4.3, if  $T_i$  and  $W_i$  are taken to be degenerate random variables, then they yield refinements of the Azuma-Hoeffding and McDiarmid inequalities, respectively. This holds because for the degenerate  $T_i, W_i, \xi_{T_i,W_i}$  is the two-point random variable presented in (1). In the special case where  $-T_i \stackrel{d}{=} W_i \stackrel{d}{=} B_i$  for some constant  $B_i > 0$ , we get  $\xi_{T_i,W_i} \stackrel{d}{=} B_i \varepsilon_i$  for a Rademacher random variable  $\varepsilon_i$  independent of  $B_i$  (i.e.,  $\mathbb{P}(\varepsilon_i = -1) = \mathbb{P}(\varepsilon_i = 1) = 1/2$ ). Corollary 4.3 should be compared with Theorems 3 & 4 of Maurer and Pontil (2021), Lemma 3 & Theorem 4 of Li and Liu (2024), and Theorems 3.4 & 3.5 of Li and Liu (2023). Unlike the results of these papers, our result is derived without making any specific assumptions about the tail behavior of  $T_i, W_i$ . For example, one can allow  $T_i, W_i$  to have tails that decay as slowly as quadratically. On the other hand, we require  $\Delta_i$  to be stochastically bounded conditional on  $\mathcal{F}_{i-1}$  for each *i* individually, while in contrast, Maurer and Pontil (2021) only require the average (over *i*) conditional sub-Gaussianity/sub-exponentially constant to be bounded.

In the following, we discuss a few applications of Corollary 4.3 to show its generality.

Sums of Random Variables. Suppose  $X_1, \ldots, X_n$  are independent random variables taking values in a Banach space  $(\mathbb{B}, \|\cdot\|)$  for some norm  $\|\cdot\|$ . Take

$$Z = \left\| \sum_{i=1}^{n} X_i \right\|.$$

Note that we do not require  $X_i$ 's to be centered. By the triangle inequality, we have

$$|\Delta_i| \leq ||X_i - X_i'|| \quad \text{for all} \quad 1 \leq i \leq n.$$

Hence, we can take  $-T_i = W_i = ||X_i - X'_i||$ . Corollary 4.3 implies that for all  $x \in \mathbb{R}$ 

$$\mathbb{P}\left(\left\|\sum_{i=1}^{n} X_{i}\right\| - \mathbb{E}\left\|\sum_{i=1}^{n} X_{i}\right\| \ge x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \lambda \left(\sum_{i=1}^{n} \varepsilon_{i} \|X_{i} - X_{i}'\| - x\right)\right)_{+}\right] \quad \text{for all} \quad x \in \mathbb{R}.$$
(11)

If one has a tail assumption such as  $\mathbb{P}(||X_i - X'_i|| \ge t) \le \overline{F}_i(t)$  for some non-increasing function  $\overline{F}(\cdot)$ , then the right hand side can be further bounded by

$$\inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \lambda \left(\sum_{i=1}^{n} \varepsilon_i W_i - x\right)\right)_+\right],\$$

where  $W_i, 1 \leq i \leq n$  are independent random variables whose survival functions are  $\overline{F}_i(t)$ . For instance, if  $||X_i - X'_i||$  is  $\sigma_i$ -subGaussian, i.e.,  $\mathbb{P}(||X_i - X'_i|| \geq t) \leq \exp(-t^2/(2\sigma_i^2))$  for all  $t \geq 0$  and some  $\sigma_i > 0$ , then we can take  $W_i$ 's to be Weibull random variables with shape parameter 2.

Inequality (11) should be compared with Proposition 7 of Maurer and Pontil (2021) and Theorem 35 of Li and Liu (2024).

**Lipschitz Functions.** Suppose  $X_1, \ldots, X_n$  are independent random variables taking values in a Banach space  $(\mathbb{B}, \|\cdot\|)$  for some norm  $\|\cdot\|$ . Take  $Z = F(X_1, \ldots, X_n)$  for some function F satisfying

$$|F(x_1, \dots, x_n) - F(y_1, \dots, y_n)| \leq \sum_{i=1}^n L_i ||x_i - y_i||$$
 for all  $x_i, y_i$ .

Then  $|\Delta_i| \leq L_i ||X_i - X'_i||$  for all  $1 \leq i \leq n$  and hence, for all  $x \in \mathbb{R}$ ,

$$\mathbb{P}\left(\left\|\sum_{i=1}^{n} X_{i}\right\| - \mathbb{E}\left\|\sum_{i=1}^{n} X_{i}\right\| \ge x\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \lambda \left(\sum_{i=1}^{n} L_{i}\varepsilon_{i}\|X_{i} - X_{i}'\| - x\right)\right)_{+}\right].$$

This inequality should be compared with Theorem 5 of Li and Liu (2024).

### 5 Comments and Conjectures for Unbounded Martingale Increments

Theorem 4.2 provides tail bounds for martingales with unbounded martingale increments, but requires stochastic dominance of the increments conditional on their past. This can be a restrictive assumption, and additionally, the bounds do not account for variances. One way to tackle both of these issues is to consider winsorized random variables, similar to Corollary 2.4 of Pinelis (2006a).

For example, if  $(X_i, \mathcal{F}_i)_{i \ge 0}$  is a real-valued martingale difference sequence, then for any y > 0,  $(\min\{X_i, y\}, \mathcal{F}_i)_{i \ge 0}$  is a super-martingale difference sequence and hence, setting  $\sigma_i^2 = \mathbb{E}[X_i^2|\mathcal{F}_{i-1}]$ , Theorem 2.1 implies

$$\mathbb{P}\left(\sum_{i=1}^{k} X_{i} \ge x \quad \text{and} \quad \sum_{i=1}^{k} \sigma_{i}^{2} \le v^{2} \text{ for some } 1 \le k \le n\right) \\
\le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}(y\widetilde{\Pi}_{v^{2}/y} - x)\right)_{+}^{2}\right] + \mathbb{P}\left(\max_{1 \le k \le n} X_{i} > y\right), \quad \text{for any} \quad y \ge 0.$$
(12)

We can replace  $\sigma_i^2$  with  $\sigma_i^2(y) = \mathbb{E}[(\min\{X_i, y\})^2 | \mathcal{F}_{i-1}]$ . From Theorem 2.1, it is clear that one can replace the centered Poisson,  $\widetilde{\Pi}_{v^2/y}$ , in (12) with a sum of *n* independent identically distributed twopoint random variables. The main result of Hahn and Klass (1997) suggests that if  $X_i$ 's are not too heterogeneously distributed, then for the optimal choice of *y*, inequality (12) may be unimprovable except for a constant multiplier. Finally, inequality (12) can be considered as an improvement of Corollary 2.2 of Fan et al. (2012).

Allowing the winsorization to be different for the martingale increments, one can get the following result. Following Theorem 2.3 of Pinelis (2014b), for any random variable U, define

$$Q_{\alpha}(U;\delta) = \inf_{t \in \mathbb{R}} t + \frac{\left(\mathbb{E}\left[(U-t)^{\alpha}_{+}\right]\right)^{1/\alpha}}{\delta^{1/\alpha}}, \quad \text{for} \quad \alpha \in (0,\infty), \delta \in (0,1).$$

Note that  $Q_{\alpha}(rU;\delta) = rQ_{\alpha}(U;\delta)$  for any  $r \ge 0$ .

**Corollary 5.1.** Suppose  $(X_i, \mathcal{F}_i)_{i \ge 0}$  is a martingale difference sequence such that, for non-increasing, non-negative function  $t \mapsto g(t)$ ,

$$\mathbb{E}\left[\left(\frac{X_i}{\overline{\sigma}_i} - t\right)_+ \middle| \mathcal{F}_{i-1}\right] \leq g(t), \quad \text{for all} \quad t \ge 0,$$
(13)

where  $\overline{\sigma}_i$  is an  $\mathcal{F}_{i-1}$ -measurable random variable. Additionally, set  $\sigma_i^2 := \mathbb{E}[X_i^2|\mathcal{F}_{i-1}]$ . Then for any  $\delta \in (0,1), y \ge 0$ , and v > 0, with  $Z \sim N(0,1)$ ,

$$\mathbb{P}\left(\bigcup_{k=1}^{n}\left\{\sum_{i=1}^{k}X_{i} \geqslant vQ_{5}\left(Z;\frac{\delta}{2}\right) + \frac{4v^{2}g(y)}{y\delta} \quad and \quad \sum_{i=1}^{k}\left(\frac{\overline{\sigma}_{i}y}{2} + \frac{\sigma_{i}^{2}}{2\overline{\sigma}_{i}y}\right) \leqslant v^{2}\right\}\right) \leqslant \delta.$$
(14)

Moreover,  $Q_5(vZ; \delta/2) \le v \min \left\{ \sqrt{2\log(2/\delta)}, \, \Phi^{-1}(1 - \delta/11.4) \right\}.$ 

A proof of Corollary 5.1 can be found in Section S.4. Assumption (13) is weaker than a tail assumption of the type  $\mathbb{P}(X_i/\overline{\sigma}_i > t | \mathcal{F}_{i-1}) \leq \overline{F}(t)$  for some survival function. Also, note that assumption (13) is equivalent to

$$\mathbb{E}\left[\left(\left(X_{i}/\overline{\sigma}_{i}\right)_{+}-t\right)_{+}|\mathcal{F}_{i-1}\right] \leq g(t) \quad \text{for all} \quad t \geq 0.$$
(15)

See, for example, Theorem 2.1 of Marsiglietti and Melbourne (2022) for cases where (15) might hold without the tail being dominated. Corollary 5.1 can be thought of as an analogue of Fuk-Nagaev type inequalities (Nagaev, 1979). Also, see Rio (2017). If  $\mathbb{P}(X_i/\overline{\sigma}_i > u|\mathcal{F}_{i-1}) \leq u^{-q}$  for some  $q \geq 2$ and for all  $u \geq 0$ , then

$$\mathbb{E}[(X_i/\overline{\sigma}_i - t)_+ | \mathcal{F}_{i-1}] \leq \int_t^\infty \mathbb{P}(X_i/\overline{\sigma}_i > u | \mathcal{F}_{i-1}) du = \int_t^\infty u^{-q} du = \frac{t^{-q+1}}{q-1}$$

In this case, inequality (14) becomes

$$\mathbb{P}\left(\bigcup_{k=1}^{n} \left\{\sum_{i=1}^{k} X_{i} \ge Q_{5}(vZ;\delta/2) + \frac{4v^{2}}{(q-1)\delta y^{q}} \quad \text{and} \quad \sum_{i=1}^{k} \left(\frac{\overline{\sigma}_{i}y}{2} + \frac{\sigma_{i}^{2}}{2\overline{\sigma}_{i}y}\right) \le v^{2}\right\}\right) \le \delta.$$
(16)

This result resembles the conclusion of Corollary 4.2 of Rio (2017) but with the leading term replaced by a Gaussian quantile rather than the crude upper bound typically obtained by the Cramer-Chernoff technique. However, it should be stressed that in our inequality y is left as a free parameter. Unfortunately, this makes the bound weaker because changing y with  $\delta$  implies that the Gaussian leading term will no longer be a Gaussian term.

In both inequalities (12) and (14), improvements exist in the case where  $\sigma_i^2$  are non-stochastic or bounded almost surely. For example, the results of Bentkus (2010) provide extensions of Theorem 2.1 with almost sure boundedness replaced by stochastic boundedness but requiring nonstochastic bound on the conditional variances. We could not combine the proof techniques in this note with the results of Bentkus (2010), because we need a stochastic dominance for the "worst-case" random variables that appear in these bounds. In detail, in the context of improved Hoeffding and Bernstein inequalities, the worst-case random variables are two-point random variables (1) and we know that sums of independent non-identically distributed Bernoulli's are stochastically dominated by sums of independent identically distributed Bernoulli with parameter being some aggregate of individual probabilities. The same holds with Gaussian and Poisson random variables, which lead to Theorem 3.1.

Ideally, when using winsorization, one should control  $\sum_{i=1}^{n} \min\{X_i, y\}$  or  $\sum_{i=1}^{n} \min\{X_i, \overline{\sigma}_i y\}$  by using the fact that in addition to the conditional variance bounded by  $\sigma_i^2$ , these random variables also satisfy a tail assumption (governed by (13)) before hitting y as their upper bound. As stated currently, the inequalities (12) and (14) do not make use of this information, as was done in Rio (2017) (for example). Using such tail information along with variance and bound, one could employ Theorem 3 of Bentkus (2010) for any martingale increment, but due to the lack of readily available stochastic domination of such sums, our bounds are weaker. For example, in the context of moment generating function (MGF) based proof in Rio (2017), one gets three regimes in the MGF for winsorized sums; see Proposition 3.5 and Eq. (3.14) there. These regimes show that the Gaussian leading term is unaffected by the winsorization (unlike ours), and the Poisson tail when balanced with winsorization yields the Fuk-Nagaev inequality with optimal constants. It might be possible to use Lemmas 4.1-4.8 of Pinelis (2014a) or the results of Bentkus and Juškevičius (2008) to obtain improvements of the Fuk-Nagaev inequalities, but our attempts to date have been unsuccessful. In summary, it would be interesting to explore Bentkus style bounds analogues to those in Fan et al. (2015).

In another direction, there are some results where truncation is introduced in the event of conditional variances. For example, Dzhaparidze and Van Zanten (2001) and Fan et al. (2017) proved the following result for a supermartingale difference sequence  $(X_i, \mathcal{F}_i)_{i \ge 0}$ :

$$\mathbb{P}\left(\bigcup_{k=1}^{n}\left\{\sum_{i=1}^{k}X_{i} \ge x \text{ and } \sum_{i=1}^{k}\mathbb{E}[X_{i}^{2}\mathbf{1}\{X_{i} \le y\}] + \sum_{i=1}^{k}X_{i}^{2}\mathbf{1}\{X_{i} > y\} \le v^{2}\right\}\right) \\ \leqslant \inf_{\lambda \ge 0} \mathbb{E}\left[\exp(-\lambda x + \lambda y\widetilde{\Pi}_{v^{2}/y})\right].$$

Although Theorem 2.1 of Fan et al. (2017) is not stated in this form, it is equivalent; see Eq. (33) there. It stands to reason that a stronger inequality of the Bentkus type might also hold. In

particular, we *conjecture* that the following holds

$$\mathbb{P}\left(\bigcup_{k=1}^{n}\left\{\sum_{i=1}^{k}X_{i} \ge x \quad \text{and} \quad \sum_{i=1}^{k}\mathbb{E}[X_{i}^{2}\mathbf{1}\{X_{i} \le y\}] + \sum_{i=1}^{k}X_{i}^{2}\mathbf{1}\{X_{i} > y\} \le v^{2}\right\}\right) \\ \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}(y\widetilde{\Pi}_{v^{2}/y} - x)\right)_{+}^{2}\right].$$

It is not clear how to generalize the proof in Fan et al. (2017) to the positive part polynomials.

#### References

- Bennett, G. (1962). Probability inequalities for the sum of independent random variables. *Journal* of the American Statistical Association, 57(297):33–45.
- Bentkus, V. (2002). A remark on Bernstein, Prokhorov, Bennett, Hoeffding, and Talagrand inequalities. *Lithuanian Mathematical Journal*, 42(3):262–269.
- Bentkus, V. (2004). On Hoeffding's inequalities. The Annals of Probability, 32(2):1650–1673.
- Bentkus, V. (2008). An extension of the Hoeffding inequality to unbounded random variables. Lithuanian Mathematical Journal, 48(2):137–157.
- Bentkus, V. (2010). Bounds for the stop loss premium for unbounded risks under the variance constraints. *Preprint. Available at https://www.math.uni-bielefeld. de/sfb701/preprints/view/423.*
- Bentkus, V. and Juškevičius, T. (2008). Bounds for tail probabilities of martingales using skewness and kurtosis. *Lithuanian Mathematical Journal*, 48(1):30–37.
- Dzhaparidze, K. and Van Zanten, J. (2001). On bernstein-type inequalities for martingales. Stochastic processes and their applications, 93(1):109–117.
- Fan, X., Grama, I., and Liu, Q. (2012). Hoeffding's inequality for supermartingales. Stochastic Processes and their Applications, 122(10):3545–3559.
- Fan, X., Grama, I., and Liu, Q. (2015). Exponential inequalities for martingales with applications. *Electron. J. Probab.*, 20(1):1–22.
- Fan, X., Grama, I., and Liu, Q. (2017). Martingale inequalities of type dzhaparidze and van zanten. Statistics, 51(6):1200–1213.
- Freedman, D. A. (1975). On tail probabilities for martingales. the Annals of Probability, pages 100–118.
- Hahn, M. G. and Klass, M. J. (1997). Approximation of partial sums of arbitrary iid random variables and the precision of the usual exponential upper bound. *The Annals of Probability*, 25(3):1451–1470.
- Hoeffding, W. (1963). Probability inequalities for sums of bounded random variables. Journal of the American Statistical Association, 58(301):13–30.
- Howard, S. R., Ramdas, A., McAuliffe, J., and Sekhon, J. (2021). Time-uniform, nonparametric, nonasymptotic confidence sequences. *The Annals of Statistics*, 49(2).

- Kontorovich, A. (2014). Concentration in unbounded metric spaces and algorithmic stability. In *International conference on machine learning*, pages 28–36. PMLR.
- Kutin, S. (2002). Extensions to mcdiarmid's inequality when differences are bounded with high probability. Dept. Comput. Sci., Univ. Chicago, Chicago, IL, USA, Tech. Rep. TR-2002-04.
- Lalley, S. P. (2014). The martingale central limit theorem. Lecture notes, University of Chicago Department of Statistics.
- Li, S. and Liu, Y. (2023). Distribution-dependent mcdiarmid-type inequalities for functions of unbounded interaction. In *International Conference on Machine Learning*, pages 19789–19810. PMLR.
- Li, S. and Liu, Y. (2024). Concentration and moment inequalities for general functions of independent random variables with heavy tails. *Journal of Machine Learning Research*, 25(268):1–33.
- Marchina, A. (2018). Concentration inequalities for separately convex functions. *Bernoulli*, 24(4A):2906–2933.
- Marsiglietti, A. and Melbourne, J. (2022). Moments, concentration, and entropy of log-concave distributions. arXiv preprint arXiv:2205.08293.
- Maurer, A. (2019). A bernstein-type inequality for functions of bounded interaction. *Bernoulli*, 25(2):1451–1471.
- Maurer, A. and Pontil, M. (2021). Concentration inequalities under sub-gaussian and subexponential conditions. Advances in Neural Information Processing Systems, 34:7588–7597.
- Nagaev, S. V. (1979). Large deviations of sums of independent random variables. The Annals of Probability, pages 745–789.
- Pinelis, I. (1998). Optimal tail comparison based on comparison of moments. In *High dimensional probability*, pages 297–314. Springer.
- Pinelis, I. (2006a). Binomial upper bounds on generalized moments and tail probabilities of (super) martingales with differences bounded from above. In *High dimensional probability*, pages 33–52. Institute of Mathematical Statistics.
- Pinelis, I. (2006b). On normal domination of (super)martingales. *Electron. J. Probab.*, 11:no. 39, 1049–1070.
- Pinelis, I. (2014a). On the Bennett-Hoeffding inequality. In Annales de l'IHP Probabilités et statistiques, volume 50, pages 15–27.
- Pinelis, I. (2014b). An optimal three-way stable and monotonic spectrum of bounds on quantiles: A spectrum of coherent measures of financial risk and economic inequality. *Risks*, 2(3):349–392.
- Rio, E. (2017). About the constants in the Fuk-Nagaev inequalities. *Electronic Communications* in Probability, 22.
- Talagrand, M. (1995). The missing factor in hoeffding's inequalities. In Annales de l'IHP Probabilités et statistiques, volume 31, pages 689–702.
- van de Geer, S. A. (2002). On hoeffding's inequality for dependent random variables. In *Empirical* process techniques for dependent data, pages 161–169. Springer.

## Supplement to "On the Missing Factor in Some Concentration Inequalities for Martingales"

#### Abstract

This supplement contains the proofs of all the main results in the paper and some supporting lemmas.

### S.1 Proof of Theorem 2.1

The proof proceeds as follows. We first provide the bound for  $\mathbb{P}(\mathcal{E}_n)$  with  $\mathcal{E}_n$  defined based on any supermartingale difference sequence satisfying the assumptions. Then we prove the bound for  $\mathbb{P}(\mathcal{E}_k \text{ for some } 1 \leq k \leq n)$  by creating a new supermartingale difference sequence  $(\widetilde{X}_i, \mathcal{F}_i)_{i\geq 0}$  for which  $\cup_{k=1}^n \mathcal{E}_k = \widetilde{\mathcal{E}}_n$ , with  $\widetilde{\mathcal{E}}_n$  defined using  $\widetilde{X}_i, 1 \leq i \leq n$ . The proof of the inequality with  $n = \infty$ follows from Proposition 4.8 of Pinelis (2014a).

Clearly,

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^n X_i - x\right)\right)_+^2 \mathbf{1}\{\mathcal{E}_n\}\right]$$
$$\leq \mathbb{E}\left[\mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^n X_i - x\right)\right)_+^2 \mathbf{1}\left\{\sum_{i=1}^n \sigma_i^2 \leq v^2\right\} \middle| \mathcal{F}_{n-1}\right]\right]$$

Observe that

$$\mathbb{E}[X_n|\mathcal{F}_{n-1}] \leq 0, \quad \operatorname{Var}(X_n|\mathcal{F}_{n-1}) = v^2 - \sum_{i=1}^{n-1} \sigma_i^2, \quad X_n \leq 1 \text{ almost surely.}$$

This implies that

$$\mathbb{E}\left[\left(1+\frac{\lambda}{2}\left(\sum_{i=1}^{n}X_{i}-x\right)\right)_{+}^{2}\mathbf{1}\{\mathcal{E}_{n}\}\middle|\mathcal{F}_{n-1}\right] \leq \mathbb{E}\left[\left(1+\frac{\lambda}{2}\sum_{i=1}^{n-1}X_{i}+\frac{\lambda}{2}(G_{n}-x)\right)_{+}^{2}\mathbf{1}\left\{\sum_{i=1}^{n}\sigma_{i}^{2}\leq v^{2}\right\}\middle|\mathcal{F}_{n-1}\right],$$

where  $G_n \in \{-(v^2 - \sum_{i=1}^{n-1} \sigma_i^2), 1\}$  with mean zero and variance bounded by  $v^2 - \sum_{i=1}^{n-1} \sigma_i^2$ . Because  $\sum_{i=1}^{n-1} \sigma_i^2$  is  $\mathcal{F}_{n-2}$  measurable, we can think of  $G_n$  (conditional on  $\mathcal{F}_{n-2}$ ) to be a centered and scaled Bernoulli random variable. Let that (uncentered and unscaled) Bernoulli be  $B_n$ , independent of  $\mathcal{F}_n$ . Then there exists  $\mathcal{F}_{n-2}$  measurable constants  $a_n, b_n$  such that  $G_n = a_n B_n + b_n$  and  $\mathbb{E}[G_n | \mathcal{F}_{n-2}] = 0$ .

Therefore,

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\left(1 + \frac{\lambda}{2} \left(\sum_{i=1}^{n-1} X_i + G_n - x\right)\right)_+^2\right].$$

Now consider the conditional expectation given  $\mathcal{F}_{n-2} \cup \{B_n\}$ . We know

$$\mathbb{E}[X_{n-1}|\mathcal{F}_{n-2}\cup\{B_n\}] \leq 0, \quad \operatorname{Var}(X_{n-1}|\mathcal{F}_{n-2}\cup\{B_n\}) = \sigma_{n-1}^2, \quad X_{n-1} \leq 1 \quad \text{almost surely.}$$

This implies

$$\mathbb{E}[(1+(\lambda/2)(X_{n-1}-u))_+^2|\mathcal{F}_{n-2}] \leq \mathbb{E}[(1+(\lambda/2)(G_{n-1}-u))_+^2|\mathcal{F}_{n-2}], \text{ for all } \lambda \geq 0, u \in \mathbb{R}.$$

Here  $G_{n-1} \in \{-\sigma_{n-1}^2, 1\}$  with mean zero (conditional on  $\mathcal{F}_{n-2}$ ) and variance  $\sigma_{n-1}^2$ . As before, we can write  $G_{n-1} = a_{n-1}B_{n-1} + b_{n-1}$  for some  $\mathcal{F}_{n-2}$  measurable random variables  $a_{n-1}, b_{n-1}$  and Bernoulli random variable  $B_{n-1}$  independent of  $\mathcal{F}_n$  so that  $\mathbb{E}[G_{n-1}|\mathcal{F}_{n-2}] = 0$ . Therefore,

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^{n-2} X_i + G_{n-1} + G_n - x\right)\right)_+^2\right],$$

Note that conditional on  $\mathcal{F}_{n-2}$ ,  $G_n$  and  $G_{n-1}$  are two mean zero independent random variables and  $\operatorname{Var}(G_n + G_{n-1}|\mathcal{F}_{n-2}) = \sigma_{n-1}^2 + v^2 - \sum_{i=1}^{n-1} \sigma_i^2 = v^2 - \sum_{i=1}^{n-2} \sigma_i^2$ . Hence, by Lemma 4.5 of Bentkus (2004), we can bound the right with

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\left(1 + \frac{\lambda}{2} \left(\sum_{i=1}^{n-2} X_i + \tilde{G}_{n-1} + \tilde{G}_n - x\right)\right)_+^2\right],$$

where  $\tilde{G}_{n-1}$ ,  $\tilde{G}_n$  are IID conditional on  $\mathcal{F}_{n-2}$  and have mean zero, bounded above by 1, and equal variances of  $(v^2 - \sum_{i=1}^{n-2} \sigma_i^2)/2$ . Note that this variance is  $\mathcal{F}_{n-3}$ -measurable, which implies that we can take  $\tilde{G}_{n-1}$ ,  $\tilde{G}_n$  to be IID conditional on  $\mathcal{F}_{n-3}$ .

Now consider this expectation conditional on  $\mathcal{F}_{n-3} \cup \{\tilde{B}_{n-1}, \tilde{B}_n\}$ . We have

 $\mathbb{E}[X_{n-2}|\mathcal{F}_{n-3}] \leq 0, \operatorname{Var}(X_{n-2}|\mathcal{F}_{n-3}) \leq \sigma_{n-2}^2, \text{ and } X_{n-3} \leq 1 \text{ almost surely.}$ 

Hence, we can replace  $X_{n-2}$  with  $G_{n-2}$ . Observe that conditional on  $\mathcal{F}_{n-3}$ ,  $G_{n-2}$ ,  $\tilde{G}_{n-1}$ ,  $\tilde{G}_n$  are independent random variables taking only two values, and  $\operatorname{Var}(G_{n-2} + \tilde{G}_{n-1} + \tilde{G}_n | \mathcal{F}_{n-3}) = v^2 - \sum_{i=1}^{n-3} \sigma_i^2$ . Applying Lemma 4.5 of Bentkus (2004) again and iterating this procedure, we can bound the probability by

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^n \tilde{G}_i - x\right)\right)_+^2\right],$$

where  $\tilde{G}_i, 1 \leq i \leq n$  are IID binary random variables taking values in  $\{-v^2/n, 1\}$  with mean zero and variance  $v^2/n$ .

To bound the probability of  $\bigcup_{k=1}^{n} \mathcal{E}_k$ , define  $T \in \{1, 2, ..., n\}$  be the minimum k such that  $\sum_{i=1}^{k} X_i \ge x$  and  $\sum_{i=1}^{k} \sigma_i^2 \le v^2$ . If  $\bigcup_{k=1}^{n} \mathcal{E}_k$  does not occur, set T = n. This is a stopping time because  $\{T \le i\}$  can be determined by looking at  $\mathcal{F}_i$ , or equivalently,  $\{T \le i\} \in \mathcal{F}_i$ . Then

$$\mathbf{1}\{\mathcal{E}_n\} \leq \mathbf{1}\left\{\sum_{i=1}^n \widetilde{X}_i \ge x, \quad \text{and} \quad \sum_{i=1}^n \widetilde{\sigma}_i^2 \le v^2\right\},\$$

where  $\widetilde{X}_i = X_i \mathbf{1}\{T \ge i\}$  and  $\widetilde{\sigma}_i^2 = \mathbb{E}[\widetilde{X}_i^2 | \mathcal{F}_{i-1}]$ . Observe that  $\sum_{i=1}^n \widetilde{X}_i = \sum_{i=1}^T X_i$ . Also, note that  $(\widetilde{X}_k, \mathcal{F}_k)_{k\ge 0}$  is a supermartingale difference sequence if  $(X_k, \mathcal{F}_k)$  is a supermartingale difference sequence. This follows because

$$\mathbb{E}\left[\sum_{i=1}^{k+1} \tilde{X}_i \middle| \mathcal{F}_k\right] = \sum_{i=1}^{k+1} \mathbb{E}[X_i \mathbf{1}\{T \ge i\} \middle| \mathcal{F}_k] = \sum_{i=1}^k X_i \mathbf{1}\{T \ge i\} + \mathbb{E}[X_{k+1} \mathbf{1}\{T \ge k+1\} \middle| \mathcal{F}_k],$$

where the last equality follows because  $X_i \in \mathcal{F}_k$  for any  $k \ge i$  and  $\mathbf{1}\{T \ge i\} = 1 - \mathbf{1}\{T \le i-1\} \in \mathcal{F}_{i-1} \subseteq \mathcal{F}_k$  for any  $k \ge i$ . Finally, because  $\mathbf{1}\{T \ge k+1\} \in \mathcal{F}_k$ , we get  $\mathbb{E}[X_{k+1}\mathbf{1}\{T \ge k\}|\mathcal{F}_k] = 1$ 

 $1{T \ge k}\mathbb{E}[X_{k+1}|\mathcal{F}_k] \le 0$ . This shows that if  $(X_k, \mathcal{F}_k)_{k\ge 0}$  is a supermartingale, then  $(\widetilde{X}_k, \mathcal{F}_k)_{k\ge 0}$  is also a supermartingale. Hence, applying the previous result, we get

$$\mathbb{P}\left(\bigcup_{k=1}^{n} \mathcal{E}_{k}\right) \leqslant \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}\left(\sum_{i=1}^{n} G_{i} - x\right)\right)_{+}^{2}\right],$$

where  $G_i, 1 \leq i \leq n$  are IID mean zero random variables such that  $\mathbb{P}(G_i \in \{-v^2/n, 1\}) = 1$ .

Finally, from Proposition 2.8 of Pinelis (2014a), it follows that for any  $\lambda \ge 0$ ,

$$\mathbb{E}\left[\left(1+\frac{\lambda}{2}\left(\sum_{i=1}^{n}\widetilde{G}_{i}-x\right)\right)_{+}^{2}\right] \leq \mathbb{E}\left[\left(1+\frac{\lambda}{2}(\widetilde{\Pi}_{v^{2}}-x)\right)_{+}^{2}\right],$$

where  $\widetilde{\Pi}_{v^2}$  is a centered Poisson $(v^2)$  random variable. Given that the right hand side is independent of n, we get

$$\mathbb{P}\left(\sum_{i=1}^{k} X_i \ge x \text{ and } \sum_{i=1}^{k} \sigma_i^2 \le v^2 \text{ for some } k \ge 1\right) \le \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{2}(\widetilde{\Pi}_{v^2} - x)\right)_+^2\right] \le \frac{e^2}{2} \mathbb{P}^{\circ}(\widetilde{\Pi}_{v^2} \ge x),$$

where  $x \mapsto \mathbb{P}^{\circ}(\widetilde{\Pi}_{v^2} \ge x)$  is the log-concave majorant of  $x \mapsto \mathbb{P}(\widetilde{\Pi}_{v^2} \ge x)$ .

### S.2 Proof of Theorem 3.1

We have, as before,

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\mathbb{E}\left[\left(1 + \frac{\lambda}{5}\left(\sum_{i=1}^n X_i - x\right)\right)_+^5 \middle| \mathcal{F}_{n-1}\right] \mathbf{1}\left\{s_n^2 \leq v^2 - V_{n-1}^2\right\}\right].$$

Conditional on  $\mathcal{F}_{n-1}$ ,  $X_n$  has a non-positive conditional mean, bounded by  $B_n$ , and conditional variance bounded by  $\sigma_n^2$  so that  $(1/2)(B_n + \sigma_n^2/B_n) \leq (v^2 - V_{n-1}^2)^{1/2}$ . Applying Lemma 5.1.3 of Pinelis (2006b), we conclude

$$\mathbb{E}\left[\left(1+\frac{\lambda}{5}\left(\sum_{i=1}^{n}X_{i}-x\right)\right)_{+}^{5}\middle|\mathcal{F}_{n-1}\right]\mathbf{1}\left\{s_{n}^{2}\leqslant v^{2}-V_{n-1}^{2}\right\}\right]$$
$$\leq \mathbb{E}\left[\left(1+\frac{\lambda}{5}\left(\sum_{i=1}^{n-1}X_{i}+G_{n}-x\right)\right)_{+}^{5}\middle|\mathcal{F}_{n-1}\right]\mathbf{1}\left\{V_{n-1}\leqslant v^{2}\right\},$$

where  $G_n|\mathcal{F}_{n-1} \sim N(0, v^2 - V_{n-1}^2)$ . Observe that  $v^2 - V_{n-1}^2$  is  $\mathcal{F}_{n-2}$ -measurable, which implies that  $G_n|\mathcal{F}_{n-2} \sim N(0, v^2 - V_{n-1}^2)$ . This implies

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\left(1 + \frac{\lambda}{5} \left(\sum_{i=1}^{n-1} X_i + G_n - x\right)\right)_+^5 \mathbf{1}\{V_{n-1}^2 \leq v^2\}\right]$$
$$\leq \mathbb{E}\left[\mathbb{E}\left[\left(1 + \frac{\lambda}{5} \left(\sum_{i=1}^{n-2} X_i + G_n + X_{n-1} - x\right)\right)_+^5 \mathbf{1}\{V_{n-1}^2 \leq v^2\} \middle| \mathcal{F}_{n-2}, G_n\right]\right].$$

Observe now that conditional on  $\mathcal{F}_{n-2}, G_n, X_{n-1}$  has a non-positive mean, variance  $\sigma_{n-1}^2$ , bounded by  $B_{n-1}$  so that  $(1/2)(B_{n-1} + \sigma_{n-1}^2/B_{n-1}) \leq s_{n-1} \in \mathcal{F}_{n-2}$ . Therefore, by Lemma 5.1.3 of Pinelis (2006b), we conclude that on the event  $V_{n-1}^2 \leq v^2$ ,

$$\mathbb{E}\left[\left(1+\frac{\lambda}{5}\left(\sum_{i=1}^{n-2}X_i+G_n+X_{n-1}-x\right)\right)_+^5 \middle| \mathcal{F}_{n-2},G_n\right]$$
  
$$\leqslant \mathbb{E}\left[\left(1+\frac{\lambda}{5}\left(\sum_{i=1}^{n-2}X_i+G_n+G_{n-1}\right)\right)_+^5 \middle| \mathcal{F}_{n-2},G_n\right],$$

where  $G_{n-1}|\mathcal{F}_{n-2}, G_n \sim N(0, s_{n-1}^2)$ . This implies that  $G_{n-1} + G_n|\mathcal{F}_{n-2} \sim N(0, v^2 - V_{n-1}^2 + s_{n-1}^2) = N(0, v^2 - V_{n-2}^2)$ . Combining, we get

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\left(1 + \frac{\lambda}{5} \left(\sum_{i=1}^{n-2} X_i + (v^2 - V_{n-2}^2)^{1/2} Z - x\right)\right)_+^5 \mathbf{1}\{V_{n-2}^2 \leq v^2\}\right],\$$

where  $Z \sim N(0,1)$  independent of  $\mathcal{F}_n$ . Repeating the argument iteratively, we conclude that

$$\mathbb{P}(\mathcal{E}_n) \leq \mathbb{E}\left[\left(1 + \frac{\lambda}{5}(vZ - x)\right)_+^5\right].$$

Hence, taking the infimum over  $\lambda \ge 0$ , we get

$$\mathbb{P}(\mathcal{E}_n) \leq \inf_{\lambda \geq 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{5}(vZ - x)\right)_+^5\right] \leq 5!(e/5)^5 \mathbb{P}\left(vZ \geq x\right).$$

Now consider the bigger event

$$\widetilde{\mathcal{E}}_n = \left\{ \sum_{i=1}^k X_i \ge x \text{ and } V_k^2 \le v^2 \text{ for some } 1 \le k \le n \right\}.$$

As before, define T as the smallest  $k \in \{1, 2, ..., n\}$  such that  $\sum_{i=1}^{k} X_i \ge x$  and  $V_k^2 \le v^2$ . If no such k exists, then set T = n. This is a stopping time, and set  $\tilde{X}_i = X_i \mathbf{1}\{T \ge i\}$ . Then it is as before easy to verify that  $(\tilde{X}_i, \mathcal{F}_i), 1 \le i \le n$  is a supermartingale if  $(X_i, \mathcal{F}_i)$  is a supermartingale. Moreover,  $\tilde{\sigma}_i^2 = \operatorname{Var}(\tilde{X}_i | \mathcal{F}_{i-1}) = \sigma_i^2 \mathbf{1}\{T \ge i\}$  and set  $\tilde{s}_i = (1/2)(B_i + \tilde{\sigma}_i^2/B_i)$ . Hence,

$$\mathbf{1}\{\widetilde{\mathcal{E}}_n\} \leq \mathbf{1}\left\{\sum_{i=1}^n \widetilde{X}_i \geqslant x \text{ and } \sum_{i=1}^n \widetilde{s}_i^2 \leq v^2\right\}.$$
$$\mathbb{P}(\widetilde{\mathcal{E}}_n) \leq \inf_{\lambda \ge 0} \mathbb{E}\left[\left(1 + \frac{\lambda}{5}(vZ - x)\right)_+^5\right].$$

and

### S.3 Proof of Lemma 4.1

**Lemma S.3.1.** Suppose X and Y are two real-valued random variables satisfying

$$\mathbb{E}[(X-t)_+] \leq \mathbb{E}[(Y-t)_+] \quad for \ all \quad t \in \mathbb{R}.$$
(E.1)

Then  $\mathbb{E}[X] \leq \mathbb{E}[Y]$ . In other words,  $X \leq_1 Y$  implies  $\mathbb{E}[X] \leq \mathbb{E}[Y]$ .

*Proof.* The fact that  $X \leq_1 Y$  implies  $\mathbb{E}[X] \leq \mathbb{E}[Y]$  follows from Proposition 4 of Bentkus (2008, Lithuanian Mathematical Journal, 48(3)). The following facts will be important:  $x = x_+ - x_= x_+ - (-x)_+$ . The proof is short and is as follows: Observe that

$$\mathbb{E}[X-t] = \mathbb{E}[(X-t)_{+}] - \mathbb{E}[(t-X)_{+}] \le \mathbb{E}[(Y-t)_{+}] = \mathbb{E}[Y-t] + \mathbb{E}[(t-Y)_{+}].$$

Therefore,

 $\mathbb{E}[X] \leq \mathbb{E}[Y] + \mathbb{E}[(t - Y)_+] \quad \text{for all} \quad t \in \mathbb{R}.$ 

Taking the limiting as  $t \to -\infty$  implies the result.

Proof of Lemma 4.1. The inequality  $\mathbb{E}[T] \leq \mathbb{E}[X] \leq \mathbb{E}[W]$  follows from Lemma S.3.1. Because  $T \leq_0 W$  (i.e.,  $\mathbb{P}(T > x) \leq \mathbb{P}(W > x)$  for all x), we get  $a_q \leq b_q$  for all  $q \in [0,1]$ . If q = 0, then  $a_q$  is the right end of the support of T and hence,  $\mathbb{P}(\xi_q \leq x) = \mathbb{P}(T \leq x)$  for all  $x < a_q$  and  $\mathbb{P}(\xi_q \leq a_q) = 1$ . This implies  $\xi_0 \stackrel{d}{=} T$ . Similarly,  $\xi_1 \stackrel{d}{=} W$ .

Setting 
$$F_{\xi_q}(x) = \mathbb{P}(\xi_q \leq x), F_T(x) = \mathbb{P}(T \leq x)$$
, and  $F_W(x) = \mathbb{P}(W \leq x)$ , we get

$$\mathbb{E}[\xi_q] = \int_0^1 F_{\xi_q}^{-1}(\delta) d\delta = \int_0^{1-q} F_T^{-1}(\delta) d\delta + \int_{1-q}^1 F_W^{-1}(\delta) d\delta = \mathbb{E}[\eta] + \int_{1-q}^1 (F_W^{-1}(\delta) - F_T^{-1}(\delta)) d\delta.$$

Because  $T \leq_0 W$ , we get differentiability and non-decreasing property of  $q \mapsto \mathbb{E}[\xi_q]$ .

To show the final property  $X \leq_1 \xi_{q_0}$ , we consider three cases: (i)  $t \leq a_{q_0}$ , (ii)  $a_{q_0} < t < b_{q_0}$ , (iii)  $t \geq b_{q_0}$ . The first and third cases are easy to prove from  $-X \leq_1 -T$  and  $X \leq_1 W$ . For example, if  $t \leq a_{q_0}$ ,

$$\mathbb{E}[(X-t)_{+}] = \mathbb{E}[(X-t)] + \mathbb{E}[(t-X)_{+}] \leq \mathbb{E}[(\xi_{q_{0}}-t)] + \mathbb{E}[(t-T)_{+}].$$

For  $t \leq a_{q_0}$ ,

$$\mathbb{E}[(t-T)_{+}] = \int_{0}^{\infty} \mathbb{P}(t-T>s) ds = \int_{-\infty}^{t} \mathbb{P}(T$$

where the condition  $t \leq a_{q_0}$  is used in the penultimate equality. Therefore,  $\mathbb{E}[(X-t)_+] \leq \mathbb{E}[(\xi_{q_0}-t)_+]$  for  $t \leq a_{q_0}$ .

If  $a_{q_0} < t < b_{q_0}$ , then note that

$$\mathbb{E}[(\xi_{q_0}-t)_+] = \int_t^{b_{q_0}} \mathbb{P}(\xi_{q_0} > s)ds + \int_{b_{q_0}}^{\infty} \mathbb{P}(\xi_{q_0} > s)ds = q_0(b_{q_0}-t) + \int_{b_{q_0}}^{\infty} \mathbb{P}(W > s)ds = q_0(b_{q_0}-t) + \mathbb{E}[(W-t)_+]$$

On the other hand, we can write

$$t = \frac{b_{q_0} - t}{b_{q_0} - a_{q_0}} a_{q_0} + \frac{t - a_{q_0}}{b_{q_0} - a_{q_0}} b_{q_0},$$

which implies

$$\mathbb{E}[(X-t)_{+}] \leq \frac{b_{q_{0}}-t}{b_{q_{0}}-a_{q_{0}}} \mathbb{E}[(X-a_{q_{0}})_{+}] + \frac{t-a_{q_{0}}}{b_{q_{0}}-a_{q_{0}}} \mathbb{E}[(X-b_{q_{0}})_{+}].$$

From case (i) & (iii), this can be upper bounded as

$$\mathbb{E}[(X-t)_{+}] \leq \frac{b_{q_{0}}-t}{b_{q_{0}}-a_{q_{0}}} \mathbb{E}[(\xi_{q_{0}}-a_{q_{0}})_{+}] + \frac{t-a_{q_{0}}}{b_{q_{0}}-a_{q_{0}}} \mathbb{E}[(W-b_{q_{0}})_{+}] = \mathbb{E}[(\xi_{q_{0}}-t)_{+}],$$

where the last equality follows from the expression for  $\mathbb{E}[(\xi_{q_0} - t)_+]$ .

### S.4 Proof of Corollary 5.1

For any  $\delta \in (0, 1)$ , set

$$x_1 = \inf_{t \in \mathbb{R}} t + \frac{(\mathbb{E}[(vZ - t)_+])^{1/5}}{(\delta/2)^5}, \quad \text{and} \quad x_2 = \frac{4v^2g(y)}{y\delta}.$$
 (E.2)

From the proof of Theorem 2.1, it suffices to prove that  $\mathbb{P}(\mathcal{E}_n) \leq \delta$ , where

$$\mathcal{E}_n = \left\{ \sum_{i=1}^n X_i \ge x_1 + x_2 \quad \text{and} \quad \sum_{i=1}^n \left( \frac{\overline{\sigma}_i y}{2} + \frac{\sigma_i^2}{2\overline{\sigma}_i y} \right) \le v^2 \right\}.$$

Observe that  $\mathcal{E}_n \subseteq \mathcal{E}_n^{(1)} \cup \mathcal{E}_n^{(2)}$  where

$$\mathcal{E}_n^{(1)} := \left\{ \sum_{i=1}^n \min\{X_i, \overline{\sigma}_i y\} \ge x_1 \text{ and } \sum_{i=1}^n \left( \frac{\overline{\sigma}_i y}{2} + \frac{\sigma_i^2}{2\overline{\sigma}_i y} \right) \le v^2 \right\},\$$
$$\mathcal{E}_n^{(2)} := \left\{ \sum_{i=1}^n (X_i - \overline{\sigma}_i y)_+ \ge x_2 \text{ and } \sum_{i=1}^n \overline{\sigma}_i \le \frac{2v^2}{y} \right\}.$$

Because  $(\min\{X_i, \overline{\sigma}_i y\}, \mathcal{F}_i)_{i \ge 0}$  is a supermartingale difference sequence, Theorem 3.1 implies that

$$\mathbb{P}(\mathcal{E}_{n}^{(1)}) \leq \inf_{\lambda \geq 0} \mathbb{E}\left[ \left( 1 + \frac{\lambda}{5} (vZ - x_{1}) \right)_{+}^{5} \right] \leq \min\left\{ e^{-x_{1}^{2}/(2v^{2})}, \frac{5!e^{5}}{5^{5}} (1 - \Phi(x_{1}/v)) \right\}.$$
(E.3)

From Eq. (2.2), Theorem 2.3 of Pinelis (2014b), and our choice of  $x_1$  in (E.2), we get  $\mathbb{P}(\mathcal{E}_n^{(1)}) \leq \delta/2$ . The second inequality of (E.3) implies that

$$Q_5(vZ;\delta/2) \le \min\left\{v\sqrt{2\log(2/\delta)}, v\Phi^{-1}(1-\delta/11.4)\right\}.$$

To bound the probability of  $\mathcal{E}_n^{(2)}$ , set  $\mathcal{V}_k := \mathbf{1}\{\sum_{i=1}^k \overline{\sigma}_i \leq 2v^2/y\}$  for  $1 \leq k \leq n$ . Note that, by assumption,  $\overline{\sigma}_i \in \mathcal{F}_{i-1}, \mathcal{V}_k \in \mathcal{F}_{k-1}$ . This fact will be used in the following without specific reference. Note that, by Markov inequality,

$$\mathbb{P}(\mathcal{E}_{n}^{(2)}) \leq \frac{1}{x_{2}} \mathbb{E}\left[\sum_{i=1}^{n} (X_{i} - \overline{\sigma}_{i}y)_{+} \mathbf{1}\{\mathcal{V}_{n}\}\right]$$

$$\leq \frac{1}{x_{2}} \mathbb{E}\left[\sum_{i=1}^{n-1} (X_{i} - \overline{\sigma}_{i}y)_{+} \mathbf{1}\{\mathcal{V}_{n-1}\} + \left(\frac{2v^{2}}{y} - \sum_{i=1}^{n-1} \overline{\sigma}_{i}\right) \mathbf{1}\{\mathcal{V}_{n-1}\} \mathbb{E}[(X_{n}/\overline{\sigma}_{n} - y)_{+}|\mathcal{F}_{n-1}]\right]$$

$$\leq \frac{1}{x_{2}} \mathbb{E}\left[\sum_{i=1}^{n-1} (X_{i} - \overline{\sigma}_{i}y)_{+} \mathbf{1}\{\mathcal{V}_{n-1}\} + \left(\frac{2v^{2}}{y} - \sum_{i=1}^{n-1} \overline{\sigma}_{i}\right) g(y) \mathbf{1}\{\mathcal{V}_{n-1}\}\right].$$

This implies, again by (13),

$$\mathbb{P}(\mathcal{E}_{n}^{(2)}) \leq \frac{1}{x_{2}} \mathbb{E}\left[\sum_{i=1}^{n-2} (X_{i} - \overline{\sigma}_{i}y)_{+} \mathbf{1}\{\mathcal{V}_{n-2}\} + \left(\frac{2v^{2}}{y} - \sum_{i=1}^{n-1} \overline{\sigma}_{i}\right) \mathbf{1}\{\mathcal{V}_{n-1}\}g(y)\right] \\ + \mathbb{E}\left[\overline{\sigma}_{n-1}\mathbf{1}\{\mathcal{V}_{n-1}\}\mathbb{E}\left[(X_{n-1}/\overline{\sigma}_{n-1} - y)_{+}|\mathcal{F}_{n-2}\right]\right] \\ \leq \frac{1}{x_{2}} \mathbb{E}\left[\sum_{i=1}^{n-2} (X_{i} - \overline{\sigma}_{i}y)_{+}\mathbf{1}\{\mathcal{V}_{n-2}\} + \left(\frac{2v^{2}}{y} - \sum_{i=1}^{n-2} \overline{\sigma}_{i}\right)\mathbf{1}\{\mathcal{V}_{n-2}\}g(y)\right].$$

Repeating this argument further, we get

$$\mathbb{P}(\mathcal{E}_n^{(2)}) \leqslant \frac{2v^2 g(y)}{yx_2}.$$

Our choice of  $x_2$  in (E.2) implies that  $\mathbb{P}(\mathcal{E}_n^{(2)}) \leq \delta/2$ . This completes the proof of the inequality  $\mathbb{P}(\mathcal{E}_n) \leq \delta$  and proves the result.