On the supremum of the sets of copulas with given

curvilinear section

Yao Ouyang^{*1}, Yonghui Sun¹, and Hua-Peng Zhang^{†2}

¹Faculty of Science, Huzhou University, Huzhou, Zhejiang 313000, China

> ²School of Science, Nanjing University of Posts and Telecommunications, Nanjing 210023, China

> > December 31, 2024

Abstract

By employing the total variation of certain functions, we give an explicit formula for the supremum of the set of all copulas with a given curvilinear section. This supremum being a copula is characterized and the relationship between this supremum and that of the set of all quasi-copulas with the same curvilinear section is considered.

Keywords: Copula; Quasi-copula; Curvilinear section; Total variation

[∗] oyy@zjhu.edu.cn

[†]Corresponding author. huapengzhang@163.com

1 Introduction

In the realm of statistical modeling and data analysis, understanding and modeling the dependencies among variables is crucial for accurate predictions and informed decision-making. Traditional methods often rely on assumptions about the joint distribution of variables, which can be limiting and restrictive, especially when dealing with complex, real-world data. Copula-based method [\[4,](#page-23-0) [5,](#page-23-1) [14\]](#page-24-0) allows for the separate modeling of marginal distributions and the dependency structure among variables, providing a more nuanced and realistic representation of the underlying data generating process. Thus it offers a flexible and versatile alternative to these traditional methods.

Copulas, which were first introduced by Sklar in [\[13\]](#page-24-1), find uses in various domains such as finance and insurance. In finance, for example, copulas have been instrumental in modeling the joint behavior of asset returns, risk management, and portfolio optimization [\[1\]](#page-23-2). In insurance [\[3\]](#page-23-3), they have been used to assess the risk of simultaneous occurrences of extreme events.

Over the years, the investigation of the diagonal section of (quasi-)copulas has attracted much attention [\[2,](#page-23-4) [9,](#page-24-2) [10\]](#page-24-3), since the diagonal section provides some information about the tail dependence of a bivariate random vector and the diagonal δ_C itself is the distribution function of the random variable $\max\{X, Y\}$ provided that the copula of (X, Y) is C. For a given diagonal δ, the set of all copulas with this diagonal section δ is nonempty. The infimum of this set is known, leaving the supremum of this set unknown for a long time. It was in [\[7\]](#page-23-5) that Kokol Bukovšek, Mojškerc and Stopar gave the precise expression of this supremum.

The study of the diagonal section of copulas has recently been extended

to that of diagonal curves. For each diagonal curvilinear Γ_{ϕ} (we will give its definition in the next section), the set of all (quasi-)copulas with given diagonal curvilinear section Γ_{ϕ} is nonempty. The infimum of the set of all quasi-copulas with given diagonal curvilinear section is a copula (called Bertino copula with the curvilinear section Γ_{ϕ} in [\[6\]](#page-23-6)), thus it is also the infimum of the set of all copulas with given diagonal curvilinear section. The supremum of the set of all quasi-copulas with a given diagonal curvilinear section Γ_{ϕ} is known as $A_{\Gamma_{\phi}}$ and is not a copula in general. The authors characterized $A_{\Gamma_{\phi}}$ as a copula in [\[11\]](#page-24-4), while a partial result can be found in [\[15\]](#page-24-5). However, the supremum of the set of all copulas with a given diagonal curvilinear section Γ_{ϕ} is yet unknown.

The central task of this paper is to give the explicit formula of the set of all copulas with a given diagonal curvilinear section Γ_{ϕ} . After recalling some necessary knowledge that will be used in this paper, we give this formula in Section 3. It is expected that it is only a quasi-copula in general, so we characterize when it is a copula in Section 4. Section 5 is devoted to discuss the relationship between two infima, namely, the infimum of the set of all quasi-copulas with given diagonal curvilinear section Γ_{ϕ} and that of the set of all copulas with the same diagonal curvilinear section. We end this paper with some concluding remarks.

2 Preliminaries

In this section, we recall some basic knowledge that will be used throughout the paper.

In the whole paper, we will not use the term "nondecreasing", instead we use the term "increasing" and "strictly increasing".

Definition 2.1. [\[8\]](#page-23-7) A (bivariate) *quasi-copula* is a function $Q : [0,1]^2 \rightarrow$

- $[0, 1]$ satisfying the following conditions:
- (Q1) $Q(x, 0) = Q(0, x) = 0$ and $Q(x, 1) = Q(1, x) = x \ (\forall x \in [0, 1]);$
- $(Q2)$ Q is increasing in each variable;
- (Q3) Q is 1-Lipschitz, *i.e.*, for all $x_1, x_2, y_1, y_2 \in [0, 1]$, it holds that

$$
|Q(x_1, y_1) - Q(x_2, y_2)| \le |x_1 - x_2| + |y_1 - y_2|.
$$

Definition 2.2. [\[8\]](#page-23-7) A (bivariate) copula is a function $C : [0,1]^2 \rightarrow [0,1]$ satisfying:

 $(C1)$ the boundary condition, *i.e.*,

$$
C(x, 0) = C(0, x) = 0
$$
 and $C(x, 1) = C(1, x) = x \ (\forall x \in [0, 1]);$

(C2) the 2-increasing property, *i.e.*, $V_C(R) \geq 0$ for any rectangle $R =$ $[x_1, x_2] \times [y_1, y_2]$, where $V_C(R) = C(x_2, y_2) - C(x_2, y_1) - C(x_1, y_2) + C(x_1, y_1)$ is called the C-volume of R.

It is not difficult to see that every copula is also a quasi-copula, but not vice versa. Those quasi-copulas that are not copulas are called proper quasi-copulas. An important copula is the independence copula Π given by $\Pi(x, y) = xy$. Other typical copulas are W, M , given by

$$
W(x, y) = \max\{x + y - 1, 0\}, M(x, y) = \min\{x, y\}.
$$

They are known as the Fréchet-Hoeffding bounds of the set of all quasicopulas, *i.e.*, for any quasi-copula Q , it holds that $W \le Q \le M$. In the presence of additional information on the quasi-copula, however, the Fréchet-Hoeffding can be narrowed, see [\[8\]](#page-23-7) for the case when the copula is known at a single point, [\[12\]](#page-24-6) for the case when the value of the copula on a given set is restricted and see [\[9\]](#page-24-2) for the case when the diagonal section is given.

Definition 2.3. [\[9\]](#page-24-2) A *diagonal* is a function $\delta : [0,1] \rightarrow [0,1]$ with the properties:

(D1)
$$
\delta(1) = 1
$$
 and $\delta(t) \le t$ for all $t \in [0, 1]$;
(D2) $0 \le \delta(t_2) - \delta(t_1) \le 2(t_2 - t_1)$ for all $t_1, t_2 \in [0, 1]$ with $t_1 \le t_2$.

The diagonal section $\delta_Q:[0,1]\to[0,1]$ of any quasi-copula Q defined by $\delta_Q(t) = Q(t, t)$ is a diagonal. Denote \mathbb{C}_{δ} (resp. \mathbb{Q}_{δ}) the set of all copulas (resp. quasi-copulas) with given diagonal section δ . Since for any diagonal δ there exists at least one copula with diagonal section δ, the set \mathbb{C}_{δ} as well as \mathbb{Q}_{δ} is nonempty. It is known [\[9\]](#page-24-2) that the copula B_{δ} given by

$$
B_{\delta}(x, y) = \begin{cases} y - \min_{t \in [y, x]} \hat{\delta}(t) & \text{if } y \leq x \\ x - \min_{t \in [x, y]} \hat{\delta}(t) & \text{if } y > x, \end{cases}
$$

is the infimum of \mathbb{C}_{δ} and \mathbb{Q}_{δ} , *i.e.*, inf $\mathbb{C}_{\delta} = \inf \mathbb{Q}_{\delta} = B_{\delta}$, where $\hat{\delta}(t) = t - \delta(t)$. The supremum of \mathbb{Q}_{δ} is the quasi-copula A_{δ} given by

$$
A_{\delta}(x, y) = \begin{cases} \min \left\{ y, x - \max_{t \in [y, x]} \hat{\delta}(t) \right\} & \text{if } y \leq x \\ \min \left\{ x, y - \max_{t \in [x, y]} \hat{\delta}(t) \right\} & \text{if } y > x. \end{cases}
$$

For the characterization of A_{δ} being a copula, we refer to [\[10\]](#page-24-3). Unlike the case of infimum, the supremum of \mathbb{Q}_{δ} and that of \mathbb{C}_{δ} are not equal, *i.e.*, $A_{\delta} \neq \sup \mathbb{C}_{\delta}$. The explicit formula of $\sup \mathbb{C}_{\delta} \triangleq \overline{C}_{\delta}$ has only recently been given by Kokol Bukovšek, Mojškerc and Stopar [\[7\]](#page-23-5):

$$
\bar{C}_{\delta} = \begin{cases}\n\min \left\{ y, x - \frac{1}{2} \left(\hat{\delta}(x) + \hat{\delta}(y) + \mathbb{V}_y^x(\hat{\delta}) \right) \right\} & \text{if } y \leq x \\
\min \left\{ x, y - \frac{1}{2} \left(\hat{\delta}(x) + \hat{\delta}(y) + \mathbb{V}_x^y(\hat{\delta}) \right) \right\} & \text{if } y > x.\n\end{cases}
$$

The study of the diagonal section of copulas has been extended to that of diagonal curves. Let $\phi : [0,1] \rightarrow [0,1]$ be an *automorphism*, *i.e.*, ϕ is continuous and strictly increasing with $\phi(0) = 0$ and $\phi(1) = 1$. A diagonal curvilinear related to ϕ is a function $\Gamma_{\phi} \colon [0,1] \to [0,1]$ with the properties: (i) $\max\{0, t + \phi(t) - 1\} \leq \Gamma_{\phi}(t) \leq \min\{t, \phi(t)\}$ for all $t \in [0, 1]$;

(ii) $0 \leq \Gamma_{\phi}(t_2) - \Gamma_{\phi}(t_1) \leq t_2 - t_1 + \phi(t_2) - \phi(t_1)$ for all $t_1, t_2 \in [0, 1]$ with $t_1 < t_2.$

Let Φ be the set of all automorphism of [0, 1] and for any $\phi \in \Phi$, denote by Δ_{ϕ} the set of all functions Γ_{ϕ} with properties (i) and (ii), then the curvilinear section $g_{Q,\phi} : [0,1] \to [0,1]$ of a quasi-copula Q defined by $g_{Q,\phi}(t) = Q(t, \phi(t))$ belongs to Δ_{ϕ} . Obviously, id $\in \Phi$, where id is the identity map on [0, 1], *i.e.*, $id(x) = x$ for all $x \in [0, 1]$ and Δ_{id} is nothing else but the set of all diagonals.

Let $\phi \in \Phi$. For each $\Gamma_{\phi} \in \Delta_{\phi}$, there exists at least one copula with curvilinear section Γ_{ϕ} . Indeed, the function given by

$$
B_{\Gamma_{\phi}}(x, y) = \begin{cases} y - \min_{t \in [\phi^{-1}(y), x]} \tilde{\Gamma}_{\phi}(t) & \text{if } y \le \phi(x) \\ x - \min_{t \in [x, \phi^{-1}(y)]} \hat{\Gamma}_{\phi}(t) & \text{if } y > \phi(x) \end{cases}
$$

is such a copula(called *Bertino copula with the curvilinear section* Γ_{ϕ} in [\[6\]](#page-23-6)), where $\tilde{\Gamma}_{\phi}(t) = \phi(t) - \Gamma_{\phi}(t)$ and $\hat{\Gamma}_{\phi}(t) = t - \Gamma_{\phi}(t)$. It is also the smallest (quasi-)copula with the curvilinear section Γ_{ϕ} . Hence it is the infima of $\mathbb{C}_{\Gamma_{\phi}}$ and $\mathbb{Q}_{\Gamma_{\phi}}$, where $\mathbb{C}_{\Gamma_{\phi}}$ (resp. $\mathbb{Q}_{\Gamma_{\phi}}$) is the set of all copulas (resp. quasicopulas) with the curvilinear section. The supremum of $\mathbb{Q}_{\Gamma_{\phi}}$ is $A_{\Gamma_{\phi}}$ given by

$$
A_{\Gamma_{\phi}}(x,y) = \begin{cases} \min \left\{ y, x - \max_{t \in [\phi^{-1}(y),x]} \hat{\Gamma}_{\phi}(t) \right\} & \text{if } y \le \phi(x) \\ \min \left\{ x, y - \max_{t \in [x,\phi^{-1}(y)]} \tilde{\Gamma}_{\phi}(t) \right\} & \text{if } y > \phi(x). \end{cases}
$$

For the characterization of $A_{\Gamma_{\phi}}$ being a copula, we refer to [\[11\]](#page-24-4)(see [\[15\]](#page-24-5) for a partial characterization). In general, $A_{\Gamma_{\phi}}$ is not the supremum of $\mathbb{C}_{\Gamma_{\phi}}$ and

the supremum of $\mathbb{C}_{\Gamma_{\phi}}$ is unknown. In the next section, by employing the total variation of certain functions, we give an explicit formula of $\sup \mathbb{C}_{\Gamma_{\phi}}$.

Let f be a real-valued function defined on $[a, b]$. The *total variation* $\mathbb{V}_a^b(f)$ of f over $[a, b]$ is

$$
\mathbb{V}_a^b(f) = \sup \left\{ \sum_{k=1}^n |f(x_k) - f(x_{k-1})| \mid a = x_0 \le x_1 \le \dots \le x_n = b \right\}.
$$

Note 2.4. Clearly, if $f(x_{i-1}) \leq f(x_i) \leq f(x_{i+1})$ (or $f(x_{i-1}) \geq f(x_i)$ ≥ $f(x_{i+1})$) then we can use $|f(x_{i+1})-f(x_{i-1})|$ to substitute $|f(x_i)-f(x_{i-1})|+$ $|f(x_{i+1})-f(x_i)|$ with the summation $\sum_{k=1}^n |f(x_k)-f(x_{k-1})|$ unchanged. So, when we calculate the variation, we can assume that $f(x_{2k}) \ge f(x_{2k+1}), f(x_{2k+2}) \ge$ $f(x_{2k+1}), k \geq 0$ (we can let $x_1 = x_0$ if needed).

We use the convention $\mathbb{V}_b^a(f) = -\mathbb{V}_a^b(f)$, then $\mathbb{V}_a^a(f) = 0$ and $\mathbb{V}_a^c(f) = 0$ $\mathbb{V}_a^b(f) + \mathbb{V}_b^c(f)$ for any $a, b, c \in \mathbb{R}$. It is well known that $\mathbb{V}_a^b(f) = |f(b) - f(a)|$ if f is monotone and for any f, g we have $\mathbb{V}_a^b(f+g) \leq \mathbb{V}_a^b(f) + \mathbb{V}_a^b(g)$.

3 Precise representation of the supremum

The following proposition provides a new upper bound of $\mathbb{C}_{\Gamma_{\phi}}$.

Proposition 3.1. Let $\phi \in \Phi$ and $\Gamma_{\phi} \in \Delta_{\phi}$. Then for any $C \in \mathbb{C}_{\Gamma_{\phi}}$ we have $C(x, y) \leq F(x, y)$, where $F: [0, 1]^2 \rightarrow [0, 1]$ is given by

$$
F(x,y) = \begin{cases} \min\left\{y, x - \frac{1}{2}\Big(\mathbb{V}_{\phi^{-1}(y)}^x(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x) + \hat{\Gamma}_{\phi}(\phi^{-1}(y))\Big) \right\}, & \text{if } y \le \phi(x) \\ \min\left\{x, y - \frac{1}{2}\Big(\mathbb{V}_x^{\phi^{-1}(y)}(\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x) + \tilde{\Gamma}_{\phi}(\phi^{-1}(y))\Big) \right\}, & \text{if } y \ge \phi(x). \end{cases}
$$
\n(1)

Proof. Note that when $y = \phi(x)$ we have

$$
x - \frac{1}{2} \left(\mathbb{V}_{\phi^{-1}(y)}^x(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x) + \hat{\Gamma}_{\phi}(\phi^{-1}(y)) \right) = \Gamma_{\phi}(x)
$$

and

$$
y - \frac{1}{2} \Big(\mathbb{V}_x^{\phi^{-1}(y)} (\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x) + \tilde{\Gamma}_{\phi}(\phi^{-1}(y)) \Big) = \Gamma_{\phi}(x).
$$

Thus F is well-defined.

Suppose now that $y \ge \phi(x)$. For any $\epsilon > 0$, by Note [2.4](#page-6-0) there exists a sequence of points $\phi(x) = x_0 \le x_1 \le \cdots \le x_{2m} = y$ such that $\tilde{\Gamma}_{\phi}(x_{2k}) >$ $\tilde{\Gamma}_{\phi}(x_{2k+1}), \tilde{\Gamma}_{\phi}(x_{2k+2}) > \tilde{\Gamma}_{\phi}(x_{2k+1})$ for all $k \geq 0$ and

$$
\mathbb{V}_x^{\phi^{-1}(y)}(\tilde{\Gamma}_{\phi}) < \sum_{i=1}^n |\tilde{\Gamma}_{\phi}(\phi^{-1}(r_i)) - \tilde{\Gamma}_{\phi}(\phi^{-1}(r_{i-1}))| + \epsilon.
$$

For convenience, let $x_{2k} \triangleq s_k, k = 0, 1, \cdots, m$ and $x_{2k-1} \triangleq t_k, k = 1, 2, \cdots, m$. Then $\tilde{\Gamma}_{\phi}(\phi^{-1}(s_{i-1})) \leq \tilde{\Gamma}_{\phi}(\phi^{-1}(t_i)), \ \tilde{\Gamma}_{\phi}(\phi^{-1}(s_i)) \leq \tilde{\Gamma}_{\phi}(\phi^{-1}(t_i))$ for all $1 \leq$ $i\leq m$ and

$$
\sum_{i=1}^{n} |\tilde{\Gamma}_{\phi}(\phi^{-1}(x_i)) - \tilde{\Gamma}_{\phi}(\phi^{-1}(x_{i-1}))| = \sum_{i=1}^{m} (\tilde{\Gamma}_{\phi}(\phi^{-1}(t_i)) - \tilde{\Gamma}_{\phi}(\phi^{-1}(s_{i-1})) + \sum_{i=1}^{m} (\tilde{\Gamma}_{\phi}(\phi^{-1}(t_i)) - \tilde{\Gamma}_{\phi}(\phi^{-1}(s_i))).
$$

As

$$
0 \leq V_C([x,\phi^{-1}(t_1)] \times [0,t_1]) + V_C([x,1] \times [t_m,y]) + \sum_{k=1}^{m-1} V_C([x,\phi^{-1}(t_{k+1})] \times [t_k,t_{k+1}])
$$

\n
$$
= \Gamma_{\phi}(\phi^{-1}(t_1)) - C(x,t_1) + y + C(x,t_m) - C(x,y) - t_m
$$

\n
$$
+ \sum_{k=1}^{m-1} \Big(C(x,t_k) + \Gamma_{\phi}(\phi^{-1}(t_{k+1})) - C(x,t_{k+1}) - C(\phi^{-1}(t_{k+1}),t_k) \Big),
$$

we have that

$$
C(x,y) \le y - t_m + \sum_{k=1}^m \Gamma_{\phi}(\phi^{-1}(t_k)) - \sum_{k=1}^{m-1} C(\phi^{-1}(t_{k+1}), t_k).
$$
 (2)

Moreover, by

$$
V_C([\phi^{-1}(s_k), \phi^{-1}(t_{k+1})] \times [0, s_k]) + V_C([\phi^{-1}(t_{k+1}), 1] \times [t_k, s_k]) \ge 0
$$

we have for each k that

$$
C(\phi^{-1}(t_{k+1}), t_k) \ge t_k - \tilde{\Gamma}_{\phi}(\phi^{-1}(s_k)).
$$
\n(3)

Substituting [\(3\)](#page-8-0) into [\(2\)](#page-7-0) we have that

$$
C(x,y) \leq y - t_m + \sum_{k=1}^{m} \Gamma_{\phi}(\phi^{-1}(t_k)) - \sum_{k=1}^{m-1} (t_k - \tilde{\Gamma}_{\phi}(\phi^{-1}(s_k)))
$$

\n
$$
= y + \sum_{k=1}^{m-1} \tilde{\Gamma}_{\phi}(\phi^{-1}(s_k)) - \sum_{k=1}^{m} \tilde{\Gamma}_{\phi}(\phi^{-1}(t_k))
$$

\n
$$
= y - \frac{1}{2} \left(\sum_{k=1}^{m} (\tilde{\Gamma}_{\phi}(\phi^{-1}(t_k)) - \tilde{\Gamma}_{\phi}(\phi^{-1}(s_{k-1})) \right)
$$

\n
$$
+ \sum_{k=1}^{m} (\tilde{\Gamma}_{\phi}(\phi^{-1}(t_k)) - \tilde{\Gamma}_{\phi}(\phi^{-1}(s_k)) + \tilde{\Gamma}_{\phi}(\phi^{-1}(s_0)) + \tilde{\Gamma}_{\phi}(s_m))
$$

\n
$$
\leq y - \frac{1}{2} (\mathbb{V}_x^{\phi^{-1}(y)}(\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x) + \tilde{\Gamma}_{\phi}(\phi^{-1}(y)) + \frac{\epsilon}{2}.
$$

By the arbitrariness of ϵ , we have

$$
C(x,y) \le y - \frac{1}{2} \Big(\mathbb{V}_x^{\phi^{-1}(y)}(\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x) + \tilde{\Gamma}_{\phi}(\phi^{-1}(y)) \Big).
$$

Since $C(x, y) \leq x$ is obvious, we get that $C(x, y) \leq F(x, y)$ for the case $y \ge \phi(x)$ as desired. In a similar way we can show that $C(x, y) \le F(x, y)$ for the case $y \leq \phi(x)$. \Box

For convenience, we define $f_1, f_2 \colon [0,1]^2 \to [0,1]$ as

$$
f_1(x,y) = x - \frac{1}{2} \Big(\mathbb{V}_{\phi^{-1}(y)}^x(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x) + \hat{\Gamma}_{\phi}(\phi^{-1}(y)) \Big) f_2(x,y) = y - \frac{1}{2} \Big(\mathbb{V}_x^{\phi^{-1}(y)}(\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x) + \tilde{\Gamma}_{\phi}(\phi^{-1}(y)) \Big).
$$

Note 3.2. Let $y \leq \phi(x)$ be given and $t^* \in [\phi^{-1}(y), x]$ be such that $\hat{\Gamma}_{\phi}(t^*) =$ $\max_{t \in [\phi^{-1}(y),x]} \{\hat{\Gamma}_{\phi}(t)\}.$ Then $\mathbb{V}_{\phi^{-1}(y)}^{x}(\hat{\Gamma}_{\phi}) \geq (\hat{\Gamma}_{\phi}(t^{*}) - \hat{\Gamma}_{\phi}(x)) + (\hat{\Gamma}_{\phi}(t^{*}) - \hat{\Gamma}_{\phi}(\phi^{-1}(y)))$. As a consequence, we have that

$$
f_1(x,y)\leq x-\widehat{\Gamma}_\phi(t^*)=x-\max_{t\in[\phi^{-1}(y),x]}\{t-\Gamma_\phi(t)\}.
$$

In a similar way we have $f_2(x, y) \leq y - \max_{t \in [x, \phi^{-1}(y)]} {\{\phi(t) - \Gamma_{\phi}(t)\}}$ for any $(x, y) \in [0, 1] \times [\phi(x), 1]$. Thus we conclude that $F \leq A_{\Gamma_{\phi}}, i.e.,$ we have obtained a better upper bound of $\mathbb{C}_{\Gamma_{\phi}}$ than $A_{\Gamma_{\phi}}$.

The central task of this paper is to prove that F is nothing else but the supremum of $\mathbb{C}_{\Gamma_{\phi}}$. To do this, let us start by proving that both f_1 and f_2 are increasing in each argument and 1-Lipschitz.

Proposition 3.3. Both f_1 and f_2 are increasing in each argument.

Proof. We only prove that f_1 is increasing since the proof of f_2 is similar. Let $x_2 > x_1$ be given. Then, for any y we have

$$
f_1(x_2, y) - f_1(x_1, y) = x_2 - x_1 - \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x_2) - \hat{\Gamma}_{\phi}(x_1) \Big).
$$

It follows from

$$
\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) \leq \mathbb{V}_{x_1}^{x_2}(t) + \mathbb{V}_{x_1}^{x_2}(\Gamma_{\phi}) = (x_2 - x_1) + (\Gamma_{\phi}(x_2) - \Gamma_{\phi}(x_1)).
$$

that $f_1(x_2, y) - f_1(x_1, y) \geq 0$, *i.e.*, f_1 is increasing in the first argument.

Let $y_2 > y_1$. Then, for any x we have

$$
f(x,y_2) - f(x,y_1) = \frac{1}{2} \Big(\mathbb{V}_{\phi^{-1}(y_1)}^{\phi^{-1}(y_2)} (\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(\phi^{-1}(y_2)) + \hat{\Gamma}_{\phi}(\phi^{-1}(y_1)) \Big) \ge 0,
$$

i.e., f_1 is increasing in the second argument. \Box

Before proving that both f_1 and f_2 are 1-Lipschitz, we need two lemmas which play a crucial role in our discussion.

Lemma 3.4. Let $\phi \in \Phi$ and $\Gamma_{\phi} \in \Delta_{\phi}$. Then for any $t_2 > t_1$ we have

(i)
$$
|\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)| \leq 2(\phi(t_2) - \phi(t_1)) + (\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1));
$$

\n(ii) $|\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1)| \leq 2(t_2 - t_1) + (\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1));$
\n(iii) $|\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)| + |\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1)| \leq (t_2 - t_1) + (\phi(t_2) - \phi(t_1)).$

Proof. (i) We distinguish two possible cases.

(a) $\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1) \ge 0.$

In this case, we have

$$
\begin{array}{rcl}\n|\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)| & = & \hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1) \\
& \leq & 2(\phi(t_2) - \phi(t_1)) + \hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1).\n\end{array}
$$

(b) $\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1) < 0.$

In this case, since
$$
\Gamma_{\phi}(t_2) - \Gamma_{\phi}(t_1) \leq (t_2 - t_1) + (\phi(t_2) - \phi(t_1))
$$
, we have
\n
$$
\phi(t_1) - \phi(t_2) \leq \hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1) < 0.
$$
 Thus we have
\n
$$
|\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)| \leq \phi(t_2) - \phi(t_1)
$$
\n
$$
\leq \phi(t_2) - \phi(t_1) + ((\phi(t_2) - \phi(t_1)) + (t_2 - t_1) - (\Gamma_{\phi}(t_2) - \Gamma_{\phi}(t_1)))
$$
\n
$$
= 2(\phi(t_2) - \phi(t_1)) + \hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1).
$$

(ii) The proof is similar to that of (i).

(iii) If both $\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1) \ge 0$ and $\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1) \ge 0$ then

$$
|\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)| + |\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1)|
$$

= $(t_2 - t_1) + (\phi(t_2) - \phi(t_1)) - 2(\Gamma_{\phi}(t_2) - \Gamma_{\phi}(t_1))$
 $\leq (t_2 - t_1) + (\phi(t_2) - \phi(t_1)).$

If both $\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1) \leq 0$ and $\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1) \leq 0$ then $|\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)|$ $\hat{\Gamma}_{\phi}(t_1) \leq \phi(t_2) - \phi(t_1)$ and $|\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1)| \leq t_2 - t_1$. So the inequality holds.

If
$$
\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1) > 0
$$
 and $\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1) < 0$ then
\n
$$
|\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)| + |\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1)| = (t_2 - t_1) - (\phi(t_2) - \phi(t_1))
$$
\n
$$
\leq (t_2 - t_1) + (\phi(t_2) - \phi(t_1)).
$$

The case $\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1) < 0$ and $\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1) > 0$ is similar. \Box

Lemma 3.5. Let $\phi \in \Phi$ and $\Gamma_{\phi} \in \Delta_{\phi}$. Then for any $t_2 > t_1$ we have

(i)
$$
\mathbb{V}_{t_1}^{t_2}(\hat{\Gamma}_{\phi}) \leq 2(\phi(t_2) - \phi(t_1)) + (\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1));
$$

\n(ii) $\mathbb{V}_{t_1}^{t_2}(\tilde{\Gamma}_{\phi}) \leq 2(t_2 - t_1) + (\tilde{\Gamma}_{\phi}(t_2) - \tilde{\Gamma}_{\phi}(t_1));$
\n(iii) $\mathbb{V}_{t_1}^{t_2}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{t_1}^{t_2}(\tilde{\Gamma}_{\phi}) \leq (t_2 - t_1) + (\phi(t_2) - \phi(t_1)).$

Proof. (i) For any $\epsilon > 0$ there exists a sequence $\{x_i\}_{i=0}^n$ with $t_1 = x_0 <$ $x_1 < \cdots < x_n = t_2$ such that

$$
\mathbb{V}_{t_1}^{t_2}(\hat{\Gamma}_{\phi}) < \sum_{i=1}^n |\hat{\Gamma}_{\phi}(x_i) - \hat{\Gamma}_{\phi}(x_{i-1})| + \epsilon.
$$

By (i) of Lemma [3.4,](#page-9-0) we have that

$$
\sum_{i=1}^{n} |\hat{\Gamma}_{\phi}(x_i) - \hat{\Gamma}_{\phi}(x_{i-1})| \leq \sum_{i=1}^{n} \left(2(\phi(x_i) - \phi(x_{i-1})) + (\hat{\Gamma}_{\phi}(x_i) - \hat{\Gamma}_{\phi}(x_{i-1})) \right)
$$

= 2(\phi(t_2) - \phi(t_1)) + (\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)).

Thus $\mathbb{V}_{t_1}^{t_2}(\hat{\Gamma}_{\phi}) < 2(\phi(t_2) - \phi(t_1)) + (\hat{\Gamma}_{\phi}(t_2) - \hat{\Gamma}_{\phi}(t_1)) + \epsilon$. The conclusion follows from the arbitrariness of $\epsilon.$

(ii) The proof is similar to that of (i).

(iii) For any $\epsilon > 0$ there exists a sequence $\{x_i\}_{i=0}^n$ with $t_1 = x_0 < x_1 <$ $\cdots < x_n = t_2$ such that

$$
\mathbb{V}_{t_1}^{t_2}(\hat{\Gamma}_{\phi}) < \sum_{i=1}^n |\hat{\Gamma}_{\phi}(x_i) - \hat{\Gamma}_{\phi}(x_{i-1})| + \epsilon
$$

and

$$
\mathbb{V}_{t_1}^{t_2}(\tilde{\Gamma}_{\phi}) < \sum_{i=1}^n |\tilde{\Gamma}_{\phi}(x_i) - \tilde{\Gamma}_{\phi}(x_{i-1})| + \epsilon
$$

By (iii) of Lemma [3.4,](#page-9-0) we have that

$$
\sum_{i=1}^{n} |\hat{\Gamma}_{\phi}(x_i) - \hat{\Gamma}_{\phi}(x_{i-1})| + \sum_{i=1}^{n} |\tilde{\Gamma}_{\phi}(x_i) - \tilde{\Gamma}_{\phi}(x_{i-1})|
$$
\n
$$
= \sum_{i=1}^{n} (|\hat{\Gamma}_{\phi}(x_i) - \hat{\Gamma}_{\phi}(x_{i-1})| + |\tilde{\Gamma}_{\phi}(x_i) - \tilde{\Gamma}_{\phi}(x_{i-1})|)
$$
\n
$$
\leq \sum_{i=1}^{n} ((x_i - x_{i-1}) + (\phi(x_i) - \phi(x_{i-1})) = (t_2 - t_1) + (\phi(t_2) - \phi(t_1)).
$$

Thus $\mathbb{V}_{t_1}^{t_2}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{t_1}^{t_2}(\tilde{\Gamma}_{\phi}) < (t_2 - t_1) + (\phi(t_2) - \phi(t_1)) + 2\epsilon$. The conclusion follows from the arbitrariness of ϵ . \Box

We are in the position to prove the Lipschitz property of f_1 and f_2 .

Proposition 3.6. Both f_1 and f_2 are 1-Lipschitz.

Proof. Let $x_1 < x_2$. Then for any y we have

$$
f_1(x_2, y) - f_1(x_1, y) = (x_2 - x_1) - \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x_2) - \hat{\Gamma}_{\phi}(x_1) \Big) \le x_2 - x_1.
$$

Since f_1 is increasing in the first argument, we have that f_1 is 1-Lipschitz w.r.t. the first argument.

Let $y_1 < y_2$. For any x, it follows from (i) of Lemma [3.5](#page-11-0) that

$$
f_1(x, y_2) - f_1(x, y_1) = \frac{1}{2} \Big(\mathbb{V}_{\phi^{-1}(y_1)}^{\phi^{-1}(y_2)} (\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(\phi^{-1}(y_2)) + \hat{\Gamma}_{\phi}(\phi^{-1}(y_1)) \Big) \leq y_2 - y_1.
$$

Since f_1 is increasing in the second argument, we have that f_1 is 1-Lipschitz w.r.t. the second argument.

Thus f_1 is 1-Lipschitz. In a similar way we can show that f_2 is 1-Lipschitz. \square

To improve that $F = \bar{C}_{\Gamma_{\phi}}$, it suffices to prove that for any upper bound Q of \mathbb{C}_{Γ_ϕ} we have $Q \geq F.$ Before doing this, let

$$
C_1(x, y) = \min\{x, y, f_1(x, y)\}\tag{4}
$$

and

$$
C_2(x, y) = \min\{x, y, f_2(x, y)\}.
$$

We have the following result.

Proposition 3.7. Let $\phi \in \Phi$ and $\Gamma_{\phi} \in \Delta_{\phi}$. Then $C_1 \in \mathbb{C}_{\Gamma_{\phi}}$ and $C_2 \in \mathbb{C}_{\Gamma_{\phi}}$.

Proof. We only prove that $C_1 \in \mathbb{C}_{\Gamma_{\phi}}$ since the proof of $C_2 \in \mathbb{C}_{\Gamma_{\phi}}$ is similar. Since $C_1(x, \phi(x)) = \Gamma_{\phi}(x)$ is obvious, it is enough to prove that C_1 is a copula.

Boundary conditions: For any x , we have

$$
\mathbb{V}_{\phi^{-1}(1)}^x(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x) + \hat{\Gamma}_{\phi}(\phi^{-1}(1)) = \hat{\Gamma}_{\phi}(x) - \mathbb{V}_x^1(\hat{\Gamma}_{\phi}) \le 0,
$$

Thus $f_1(x, 1) \geq x$ and which implies $C_1(x, 1) = x$. On the other hand, by (i) of Lemma [3.5,](#page-11-0) we have

$$
\mathbb{V}_{\phi^{-1}(x)}^1(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(1) + \hat{\Gamma}_{\phi}(\phi^{-1}(x) \le 2(\phi(1) - x) + 2\hat{\Gamma}_{\phi}(1) = 2(1 - x).
$$

Thus $f_1(1, x) \geq x$ and which implies $C_1(1, x) = x$. Since f_1 is increasing in each argument, C_1 is also increasing in each argument. Thus we have $C_1(0, x) = C_1(x, 0) = 0.$

2-increase: Note that any rectangle $R = [a, b] \times [c, d]$ can be partitioned into at most three non-overlapping rectangles R_i and $V_{C_1}(R)$ is the sum of all $V_{C_1}(R_i)$, where each R_i belongs to one of the three types of rectangles: Type 1: $R \subset \{(x, y) \mid y \leq \phi(x)\}, i.e., R$ is below the curve $\{(x, \phi(x)) \mid x \in$ $[0, 1]$;

Type 2: $R \subset \{(x, y) \mid y \ge \phi(x)\}\$, *i.e.*, R is above the curve $\{(x, \phi(x)) \mid x \in \mathbb{R}\}$ $[0, 1]$;

Type 3: $R = [x_1, x_2] \times [\phi(x_1), \phi(x_2)]$, *i.e.*, R has two vertices on the curve $\{(x, \phi(x)) \mid x \in [0,1]\}.$

We only need to prove that $V_{C_1}(R) \geq 0$ for each of the three types of rectangles $R = [x_1, x_2] \times [y_1, y_2]$.

Type 1: $R = [x_1, x_2] \times [y_1, y_2] \subset \{(x, y) | y \le \phi(x)\}.$

In this case we have $C_1(x, y) = \min\{y, f_1(x, y)\}\)$ for any $(x, y) \in R$.

If $C_1(x_1, y_1) = y_1$ then by the increasingness of C_1 we know that $V_{C_1}(R) =$ $C_1(x_2, y_2) - C_1(x_1, y_2) \geq 0$. So, without of generality, we suppose that $C_1(x_1, y_1) = f_1(x_1, y_1)$. There are two subcases to be considered.

(a)
$$
C_1(x_2, y_2) = y_2
$$
.

In this subcase we have

$$
V_{C_1}(R) \ge f_1(x_1, y_1) + y_2 - y_1 - f_1(x_1, y_2) \ge 0,
$$

where the last inequality is due to the 1-Lipschitz of f_1 .

(b) $C_1(x_2, y_2) = f_1(x_2, y_2)$.

In this subcase we have

$$
V_{C_1}(R) \ge f_1(x_1, y_1) + f_1(x_2, y_2) - f_1(x_1, y_2) - f_1(x_2, y_1) = V_{f_1}(R) = 0.
$$

Type 2: $R = [x_1, x_2] \times [y_1, y_2] \subset \{(x, y) \mid y \ge \phi(x)\}.$

If $C_1(x_1, y_1) = x_1$ or $C_1(x_1, y_1) = y_1$ then by the increasingness of C_1 implies that $V_{C_1}(R) \geq 0$. So, without of generality, we suppose that $C_1(x_1, y_1) = f_1(x_1, y_1)$. There are three subcases to be considered.

(a) $C_1(x_2, y_2) = x_2$.

In this subcase we have

$$
V_{C_1}(R) \ge f_1(x_1, y_1) + x_2 - x_1 - f_1(x_2, y_1) \ge 0.
$$

(b) $C_1(x_2, y_2) = y_2$.

In this subcase we have

$$
V_{C_1}(R) \ge f_1(x_1, y_1) + y_2 - y_1 - f_1(x_1, y_2) \ge 0.
$$

(c) $C_1(x_2, y_2) = f_1(x_2, y_2)$.

In this subcase we have

$$
V_{C_1}(R) \ge f_1(x_1, y_1) + f_1(x_2, y_2) - f_1(x_1, y_2) - f_1(x_2, y_1) = V_{f_1}(R) = 0.
$$

Type 3: $y_1 = \phi(x_1)$ and $y_2 = \phi(x_2)$.

In this case $C_1(x_1, y_1) = f_1(x_1, y_1) = \Gamma_{\phi}(x_1)$ and $C_1(x_2, y_2) = f_1(x_2, y_2) =$ $\Gamma_{\phi}(x_2)$. Thus

$$
V_{C_1}(R) \geq \Gamma_{\phi}(x_1) + \Gamma_{\phi}(x_2) - f_1(x_1, y_2) - f(x_2, y_1)
$$

= $\Gamma_{\phi}(x_1) + \Gamma_{\phi}(x_2) - x_1 + \frac{1}{2} \Big(\mathbb{V}_{x_2}^{x_1}(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x_1) + \hat{\Gamma}_{\phi}(x_2) \Big)$
 $-x_2 + \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x_1) + \hat{\Gamma}_{\phi}(x_2) \Big)$
= $\Gamma_{\phi}(x_1) + \Gamma_{\phi}(x_2) - \Gamma_{\phi}(x_1) - \Gamma_{\phi}(x_2) = 0.$

Thus C_1 is 2-increasing and hence $C_1 \in \mathbb{C}_{\Gamma_{\phi}}$. \Box

Note 3.8. From the viewpoint of [\[6\]](#page-23-6), F is the ϕ -splicing of two copulas C_1 and C_2 , *i.e.*,

$$
F(x,y) = (C_2 \boxtimes_{\phi} C_1)(x,y) = \begin{cases} C_2(x,y), & \text{if } y \le \phi(x) \\ C_1(x,y), & \text{if } y \ge \phi(x), \end{cases}
$$

thus F is a quasi-copula.

Our main theorem states that $F = \bar{C}_{\Gamma_{\phi}}$.

Theorem 3.9. The function F given by [\(1\)](#page-6-1) is the supremum of $\mathbb{C}_{\Gamma_{\phi}}$, i.e., $F = \bar{C}_{\Gamma_{\phi}}$.

Proof. Let Q be an arbitrary upper bound of $\overline{\mathbb{C}}_{\Gamma_{\phi}}$. For any (x, y) with $y \leq \phi(x)$ we have

$$
Q(x, y) \ge C_1(x, y) = F(x, y).
$$

For any (x, y) with $y \ge \phi(x)$ we have

$$
Q(x, y) \ge C_2(x, y) = F(x, y).
$$

So we have $Q \geq F$ which implies that $F = \bar{C}_{\Gamma_{\phi}}$. \Box

4 Characterization for \bar{C}_{Γ_ϕ} being a copula

In last section we have proved that for each $\Gamma_{\phi} \in \Delta_{\phi}$, the supremum of $\mathbb{C}_{\Gamma_{\phi}}$ is given by

$$
\bar{C}_{\Gamma_{\phi}} = \begin{cases} \min \left\{ y, x - \frac{1}{2} \Big(\mathbb{V}_{\phi^{-1}(y)}^x (\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x) + \hat{\Gamma}_{\phi}(\phi^{-1}(y)) \Big) \right\}, & \text{if } y \le \phi(x) \\ \min \left\{ x, y - \frac{1}{2} \Big(\mathbb{V}_{x}^{\phi^{-1}(y)} (\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x) + \tilde{\Gamma}_{\phi}(\phi^{-1}(y)) \Big) \right\}, & \text{if } y \ge \phi(x). \end{cases}
$$

This section is devoted to explore the conditions such that $\bar{C}_{\Gamma_{\phi}}$ is a copula.

Since C_1 and C_2 are copulas, to ensure $\bar{C}_{\Gamma_{\phi}}$ being a copula, it suffices to guarantee that $V_{\bar{C}_{\Gamma_{\phi}}}(R) \ge 0$ for each rectangle $R = [x_1, x_2] \times [\phi(x_1), \phi(x_2)].$ For such a rectangle R we have

$$
V_{\bar{C}_{\Gamma_{\phi}}}(R) = \Gamma_{\phi}(x_1) - \min \left\{ x_1, \phi(x_2) - \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x_1) + \tilde{\Gamma}_{\phi}(x_2) \Big) \right\} - \min \left\{ \phi(x_1), x_2 - \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x_1) + \hat{\Gamma}_{\phi}(x_2) \Big) \right\} + \Gamma_{\phi}(x_2).
$$

If

$$
\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x_1) - \hat{\Gamma}_{\phi}(x_2) \ge 0 \tag{5}
$$

then

$$
V_{\bar{C}_{\Gamma_{\phi}}}(R) \geq \Gamma_{\phi}(x_1) - x_1 - x_2 + \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x_1) + \hat{\Gamma}_{\phi}(x_2) \Big) + \Gamma_{\phi}(x_2)
$$

=
$$
\frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x_1) - \hat{\Gamma}_{\phi}(x_2) \Big) \geq 0.
$$

Similarly, if

$$
\mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) - \tilde{\Gamma}_{\phi}(x_1) - \tilde{\Gamma}_{\phi}(x_2) \ge 0 \tag{6}
$$

then

$$
V_{\bar{C}_{\Gamma_{\phi}}}(R) \geq \Gamma_{\phi}(x_1) - \phi(x_2) + \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x_1) + \tilde{\Gamma}_{\phi}(x_2) \Big) - \phi(x_1) + \Gamma_{\phi}(x_2)
$$

=
$$
\frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) - \tilde{\Gamma}_{\phi}(x_1) - \tilde{\Gamma}_{\phi}(x_2) \Big) \geq 0.
$$

Note that by (iii) of Lemma [3.5](#page-11-0) $\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) - (x_2 - x_1) - (\phi(x_2) \phi(x_1)$) \leq 0. So, if

$$
\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) - (x_2 - x_1) - (\phi(x_2) - \phi(x_1)) = 0 \tag{7}
$$

then

$$
V_{\bar{C}_{\Gamma_{\phi}}}(R) \geq \Gamma_{\phi}(x_1) - \phi(x_2) + \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) + \tilde{\Gamma}_{\phi}(x_1) + \tilde{\Gamma}_{\phi}(x_2) \Big) -x_2 + \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x_1) + \hat{\Gamma}_{\phi}(x_2) \Big) + \Gamma_{\phi}(x_2) = \frac{1}{2} \Big(\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) - (x_2 - x_1) - (\phi(x_2) - \phi(x_1)) \Big) \geq 0.
$$

Finally, if

$$
\Gamma_{\phi}(x_1) + \Gamma_{\phi}(x_2) - x_1 - \phi(x_1) \ge 0 \tag{8}
$$

then we also have $V_{\bar{C}_{\Gamma_{\phi}}}(R) \geq 0$.

Let us stress that if $\bar{C}_{\Gamma_{\phi}}$ is a copula then [\(8\)](#page-17-0) cannot occur in isolation. Before doing this, we need the following result.

Lemma 4.1. Let $x_1 < x_2$.

(i) If $\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x_1) - \hat{\Gamma}_{\phi}(x_2) \geq 0$ then for any $x \geq x_2$ we have $\mathbb{V}_{x_1}^x(\hat{\Gamma}_\phi) - \hat{\Gamma}_\phi(x_1) - \hat{\Gamma}_\phi(x) \geq 0;$ (ii) If $\mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) - \tilde{\Gamma}_{\phi}(x_1) - \tilde{\Gamma}_{\phi}(x_2) \geq 0$ then for any $x \geq x_2$ we have $\mathbb{V}_{x_1}^x(\tilde{\Gamma}_\phi) - \tilde{\Gamma}_\phi(x_1) - \tilde{\Gamma}_\phi(x) \geq 0$

Proof. (i) For any $x \geq x_2$ we have

$$
\begin{array}{rcl}\n\mathbb{V}_{x_1}^x(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x_1) - \hat{\Gamma}_{\phi}(x) & = & \mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x_1) - \hat{\Gamma}_{\phi}(x_2) \\
& & + (\mathbb{V}_{x_2}^x(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x) + \hat{\Gamma}_{\phi}(x_2)) \ge 0.\n\end{array}
$$

(ii) The proof is similar to that of (i). \Box

Now, we suppose that [\(8\)](#page-17-0) holds but [\(5\)](#page-16-0)-[\(7\)](#page-17-1) fail. Then we have $\Gamma_{\phi}(x_1)$ < $\min\{x_1, \phi(x_1)\}$ (if not, then either [\(5\)](#page-16-0) or [\(6\)](#page-17-2) holds), as a consequence $\Gamma_{\phi}(t)$ + $\Gamma_\phi(x_1) - x_1 - \phi(x_1) < 0$ whenever t is close enough to x_1 . Let

$$
t^* = \sup\{t \mid t \ge x_1 \text{ and } \Gamma_{\phi}(t) + \Gamma_{\phi}(x_1) - x_1 - \phi(x_1) < 0\}.
$$

Then $t^* \in]x_1, x_2]$ and $\Gamma_{\phi}(t^*) + \Gamma_{\phi}(x_1) - x_1 - \phi(x_1) = 0$. By Lemma [4.1](#page-18-0) we have $\mathbb{V}_{x_1}^{t^*}$ $x_1^{t^*}(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x_1) - \hat{\Gamma}_{\phi}(t^*)$ < 0 and $\mathbb{V}_{x_1}^{t^*}$ $\begin{aligned} t_{x_1}^*(\tilde{\Gamma}_\phi)-\tilde{\Gamma}_\phi(x_1)-\tilde{\Gamma}_\phi(t^*)\,<\,0. \end{aligned}$ Thus,

$$
\mathbb{V}_{x_1}^{t^*}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{x_1}^{t^*}(\tilde{\Gamma}_{\phi}) - (t^* - x_1) - (\phi(t^*) - \phi(x_1))
$$
\n
$$
= \left(\mathbb{V}_{x_1}^{t^*}(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x_1) - \hat{\Gamma}_{\phi}(t^*) \right) + \left(\mathbb{V}_{x_1}^{t^*}(\tilde{\Gamma}_{\phi}) - \tilde{\Gamma}_{\phi}(x_1) - \tilde{\Gamma}_{\phi}(t^*) \right)
$$
\n
$$
- 2\left(\Gamma_{\phi}(t^*) + \Gamma_{\phi}(x_1) - x_1 - \phi(x_1) \right) < 0.
$$

By the continuity of Γ_{ϕ} and ϕ we can choose $t_0 \in]x_1, t^*[$ such that $(5)-(7)$ $(5)-(7)$ fail for $[x_1, t_0]$. But, by the definition of t^* , [\(8\)](#page-17-0) also fails for $[x_1, t_0]$, *i.e.*,

$$
\Gamma_{\phi}(t_0) + \Gamma_{\phi}(x_1) - x_1 - \phi(x_1) < 0.
$$

Thus $V_{\bar{C}_{\Gamma_{\phi}}}([x_1, t_0] \times [\phi(x_1), \phi(t_0)]) < 0$ and $\bar{C}_{\Gamma_{\phi}}$ is not a copula.

Thus we have the following result.

Proposition 4.2. Let $\phi \in \Phi$ and $\Gamma_{\phi} \in \Delta_{\phi}$. $\bar{C}_{\Gamma_{\phi}}$ is a copula if and only if, for any rectangle $R = [x_1, x_2] \times [\phi(x_1), \phi(x_2))]$, one of [\(5\)](#page-16-0)-[\(7\)](#page-17-1) holds.

Example 4.3. Let $\phi \in \Phi$ and

$$
\Gamma_{\phi}(t) = \begin{cases}\n\min\{a_i, \phi(a_i)\} & \text{if } t \in [a_i, u_i^*] \\
\phi(t) + t - \max\{b_i, \phi(b_i)\} & \text{if } t \in [u_i^*, b_i] \\
m_{\phi}(t) & \text{otherwise,} \n\end{cases}
$$

where $u_i^* \in]a_i, b_i[$ is such that $\phi(u_i^* \in]a_i, a_i]$ $\{e_i^*\} + u_i^* = m_\phi(a_i) + \max\{b_i, \phi(b_i)\}.$ We will show that one of [\(5\)](#page-16-0)-[\(7\)](#page-17-1) holds for any rectangle of the form $R = [x_1, x_2] \times$ $[\phi(x_1), \phi(x_2)]$ and thus $\bar{C}_{\Gamma_{\phi}}$ is a copula:

(i) If there is $t \in [\phi(x_1), \phi(x_2)]$ such that $\Gamma_{\phi}(t) = t$, *i.e.*, $\hat{\Gamma}_{\phi}(t) = 0$, then

$$
\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) - \hat{\Gamma}_{\phi}(x_1) - \hat{\Gamma}_{\phi}(x_2) \ge -2\hat{\Gamma}_{\phi}(t) = 0.
$$

That is, [\(5\)](#page-16-0) holds. Similarly, if $\Gamma_{\phi}(t) = \phi(t)$ then [\(6\)](#page-17-2) holds.

(ii) We now suppose that $[x_1, x_2] \subset]a_i, b_i[$ for some i. If $x_1, x_2 \leq u_i^*$ i ^{*} then both $\hat{\Gamma}_{\phi}(t)$ and $\tilde{\Gamma}_{\phi}(t)$ are increasing on $[x_1, x_2]$. Then by a simple calculation we have

$$
\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) = x_2 - x_1 + \phi(x_2) - \phi(x_1),
$$

i.e., [\(7\)](#page-17-1) holds. Similarly, if $x_1, x_2 \ge u_i^*$ then (7) also holds since $\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi})$ + $\mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) = x_2 - x_1 + \phi(x_2) - \phi(x_1)$. For the last case that $x_1 < u_i^* < x_2$ we have

$$
\begin{array}{rcl}\n\mathbb{V}_{x_1}^{x_2}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{x_1}^{x_2}(\tilde{\Gamma}_{\phi}) & = & \left(\mathbb{V}_{x_1}^{u_i^*}(\hat{\Gamma}_{\phi}) + \mathbb{V}_{x_1}^{u_i^*}(\hat{\Gamma}_{\phi}) \right) + \left(\mathbb{V}_{u_i^*}^{x_2}(\tilde{\Gamma}_{\phi}) + \mathbb{V}_{u_i^*}^{x_2}(\tilde{\Gamma}_{\phi}) \right) \\
& = & \left(u_i^* - x_1 + \phi(u_i^*) - \phi(x_1) \right) + \left(x_2 - u_i^* + \phi(x_2) - \phi(u_i^*) \right) \\
& = & \left(x_2 - x_1 + \phi(x_2) - \phi(x_1) \right),\n\end{array}
$$

again [\(7\)](#page-17-1) holds.

It should be pointed out that the quasi-copula $A_{\Gamma_{\phi}}$ may not be a copula since the ϕ may not be compatible with $\{ |a_i, b_i| \}_{i \in I}$, *i.e.*, $\min(a_i, \phi(a_i)) \ge$ $\max\{b_{i-1}, \phi(b_{i-1})\}$ may fail for some i. Clearly, if ϕ is not compatible with $\{|a_i,b_i|\}_{i\in I}$ then $A_{\Gamma_{\phi}} > \bar{C}_{\Gamma_{\phi}}$.

Now we consider the case that $\phi = id$, *i.e.*, $\phi(t) = t, \forall t \in [0, 1]$. In this case, both [\(5\)](#page-16-0) and [\(6\)](#page-17-2) become to

$$
\mathbb{V}_{x_1}^{x_2}(\hat{\delta}) \ge \hat{\delta}(x_1) + \hat{\delta}(x_2)
$$
\n(9)

and [\(7\)](#page-17-1) becomes to

$$
\mathbb{V}_{x_1}^{x_2}(\hat{\delta}) = x_2 - x_1. \tag{10}
$$

If $\hat{\delta}(x_1) = 0$, i.e., $\delta(x_1) = x_1$ then [\(9\)](#page-20-0) holds. Otherwise, we can choose $x_2 > x_1$ such that $x_2 - x_1 < \hat{\delta}(x_1)$. Then, by the 1-Lipschitz of $\hat{\delta}$, we have $\mathbb{V}_{x_1}^{x_2}(\hat{\delta}) \le x_2 - x_1 < \hat{\delta}(x_1) + \hat{\delta}(x_2)$, *i.e.*, [\(9\)](#page-20-0) fails. To ensure \bar{C}_{δ} to be a copula, [\(10\)](#page-20-1) must hold. But [\(10\)](#page-20-1) holds if and only if $|\hat{\delta}'| = 1$ almost everywhere on $[x_1, x_2]$. Thus Proposition [4.2](#page-19-0) retrieves to Theorem 3.8 of [\[7\]](#page-23-5).

Corollary 4.4. [\[7\]](#page-23-5) Let δ be a given diagonal. Then \overline{C}_{δ} is a copula if and only if $\delta'(t) \in \{0,2\}$ almost everywhere on the set $\{x \mid \delta(x) < x\}.$

5 Relationship between two suprema

In this section, we consider the relationship between $\bar{C}_{\Gamma_{\phi}}$ and $A_{\Gamma_{\phi}}$. The following characterizes when they are equal to each other.

Proposition 5.1. Let $\phi \in \Phi$ and $\Gamma_{\phi} \in \Delta_{\phi}$. Then $\bar{C}_{\Gamma_{\phi}} = A_{\Gamma_{\phi}}$ if and only if the following two conditions are satisfied:

(i) $y-x \ge \max{\{\tilde{\Gamma}_{\phi}(x), \tilde{\Gamma}_{\phi}(\phi^{-1}(y))\}}$ whenever x, y are such that $y > \phi(x)$ and $\min_{t\in[x,\phi^{-1}(y)]}\hat{\Gamma}_{\phi}(t) < \min\{\hat{\Gamma}_{\phi}(x),\hat{\Gamma}_{\phi}(\phi^{-1}(y))\};$ (ii) $x-y \ge \max\{\hat{\Gamma}_{\phi}(\phi^{-1}(y)), \hat{\Gamma}_{\phi}(x)\}\$ whenever x, y are such that $y < \phi(x)$ and $\min_{t \in [\phi^{-1}(y),x]} \hat{\Gamma}_{\phi}(t) < \min \{ \hat{\Gamma}_{\phi}(x), \hat{\Gamma}_{\phi}(\phi^{-1}(y)) \}$.

Proof. Necessity. (i) Suppose that $t_0, t^* \in [x, \phi^{-1}(y)]$ such that $\tilde{\Gamma}_{\phi}(t_0) =$ $\min_{t\in[x,\phi^{-1}(y)]}\tilde{\Gamma}_{\phi}(t)$ and $\tilde{\Gamma}_{\phi}(t^*) = \max_{t\in[x,\phi^{-1}(y)]}\tilde{\Gamma}_{\phi}(t)$, then $t_0 \in]x,\phi^{-1}(y)[$ and $t_0 \neq$ t ∗ . There are two cases to be considered.

(a) $t_0 > t^*$.

In this case, we have $A_{\Gamma_{\phi}}(t^*, y) = \min\{t^*, y - \tilde{\Gamma}_{\phi}(t^*)\}$. Since

$$
\mathbb{V}_{t^*}^{\phi^{-1}(y)}(\tilde{\Gamma}_{\phi}) \ge \tilde{\Gamma}_{\phi}(\phi^{-1}(y)) + \tilde{\Gamma}_{\phi}(t^*) - 2\tilde{\Gamma}_{\phi}(t_0) > \tilde{\Gamma}_{\phi}(t^*) - \tilde{\Gamma}_{\phi}(\phi^{-1}(y)),
$$

which implies that $f_2(t^*, y) < y - \tilde{\Gamma}_{\phi}(t^*)$. To ensure $\bar{C}_{\Gamma_{\phi}}(t^*, y) =$ $\min\{t^*, f_2(x, y)\} = A_{\Gamma_{\phi}}(t^*, y)$, the only possibility is $t^* < y - \tilde{\Gamma}_{\phi}(t^*),$ $i.e., y - x \ge y - t^* > \tilde{\Gamma}_{\phi}(t^*) \ge \max{\{\tilde{\Gamma}_{\phi}(x), \tilde{\Gamma}_{\phi}(\phi^{-1}(y))\}}.$

(b) $t_0 < t^*$.

In this case, we have $A_{\Gamma_{\phi}}(x, \phi(t^*)) = \min\{x, \phi(t^*) - \tilde{\Gamma}_{\phi}(t^*)\}.$ Since

$$
\mathbb{V}_x^{t^*}(\tilde{\Gamma}_{\phi}) \ge \tilde{\Gamma}_{\phi}(t^*) + \tilde{\Gamma}_{\phi}(x) - 2\tilde{\Gamma}_{\phi}(t_0) > \tilde{\Gamma}_{\phi}(t^*) - \tilde{\Gamma}_{\phi}(x),
$$

which implies that $f_2(x, \phi(t^*)) < \phi(t^*) - \tilde{\Gamma}_{\phi}(t^*)$. To ensure $\bar{C}_{\Gamma_{\phi}}(x, \phi(t^*))$ = $\min\{x, f_2(x, \phi(t^*))\} = A_{\Gamma_\phi}(x, \phi(t^*))$, the only possibility is $x < \phi(t^*)$ – $\tilde{\Gamma}_{\phi}(t^*) \leq y - \tilde{\Gamma}_{\phi}(t^*), \ i.e., y - x > \tilde{\Gamma}_{\phi}(t^*) \geq \max\{\tilde{\Gamma}_{\phi}(x), \tilde{\Gamma}_{\phi}(\phi^{-1}(y))\}.$

(ii) The proof is similar to that of (i).

Sufficiency. Let $x, y \in [0, 1]$. If $y = \phi(x)$ then $\overline{C}_{\Gamma_{\phi}}(x, y) = A_{\Gamma_{\phi}}(x, y)$ $\Gamma_{\phi}(x)$. For $y \neq \phi(x)$ there are two cases to be considered, namely $y < \phi(x)$

and $y > \phi(x)$. We only consider the former since the latter is similar. We distinguish two possible cases to prove that $\bar{C}_{\Gamma_{\phi}}(x,y) = A_{\Gamma_{\phi}}(x,y)$.

(a) For any $[y_1, x_1] \subset [\phi^{-1}(y), x]$ we have $\min{\{\hat{\Gamma}_{\phi}(y_1), \hat{\Gamma}_{\phi}(x_1)\}} = \min_{t \in [y_1, x_1]} \hat{\Gamma}_{\phi}(t)$. In this case, there is $t_0 \in [\phi^{-1}(y), x]$ such that $\hat{\Gamma}_{\phi}(t)$ is increasing in $[\phi^{-1}(y), t_0]$ and decreasing in $[t_0, x]$. Thus

$$
\mathbb{V}_{\phi^{-1}(y)}^x \hat{\Gamma}_{\phi}(t) = 2\hat{\Gamma}_{\phi}(t_0) - \hat{\Gamma}_{\phi}(\phi^{-1}(y)) - \hat{\Gamma}_{\phi}(x),
$$

which implies that $f_1(x,y) = x - \hat{\Gamma}_{\phi}(t_0)$ and

$$
\bar{C}_{\Gamma_{\phi}}(x,y) = \min\{y, x - \hat{\Gamma}_{\phi}(t_0)\} = \min\{y, x - \max_{t \in [\phi^{-1}(y), x]} \hat{\Gamma}_{\phi}(t)\} = A_{\Gamma_{\phi}}(x,y).
$$

(b) There is $[y_1, x_1] \subset [\phi^{-1}(y), x]$ such that $\min{\{\hat{\Gamma}_{\phi}(y_1), \hat{\Gamma}_{\phi}(x_1)\}} > \min_{t \in [y_1, x_1]} \hat{\Gamma}_{\phi}(t)$.

In this case there is $t_1 \in]y_1, x_1[$ such that $\hat{\Gamma}_{\phi}(t_1) = \min_{t \in [y_1, x_1]} \hat{\Gamma}_{\phi}(t)$. Let $y_0 = \inf\{s \mid \hat{\Gamma}_{\phi}(t) = \hat{\Gamma}_{\phi}(t_1), \forall t \in [s, t_1]\}\$ and $x_0 = \sup\{s \mid \hat{\Gamma}_{\phi}(t) =$ $\hat{\Gamma}_{\phi}(t_1), \forall t \in [t_1, s]$. Then, by the continuity of $\hat{\Gamma}_{\phi}, \hat{\Gamma}_{\phi}(t) \equiv \hat{\Gamma}_{\phi}(t_1), t \in$ [y_0, x_0]. By the definition of x_0, y_0 , for any sufficiently small $\epsilon > 0$ there are $y_{\epsilon} \in]y_0 - \epsilon, y_0[$ and $x_{\epsilon} \in]x_0, x_0 + \epsilon[$ such that $\hat{\Gamma}_{\phi}(y_{\epsilon}) > \hat{\Gamma}_{\phi}(t_1)$ and $\hat{\Gamma}_{\phi}(x_{\epsilon}) > \hat{\Gamma}_{\phi}(t_1), i.e.,$

$$
\min_{t\in[\phi^{-1}(\phi(y_{\epsilon})),x_{\epsilon}]} \hat{\Gamma}_{\phi}(t) = \hat{\Gamma}_{\phi}(t_1) < \min\left\{\hat{\Gamma}_{\phi}(\phi^{-1}(\phi(y_{\epsilon})),\hat{\Gamma}_{\phi}(x_{\epsilon})\right\}.
$$

Thus, by (ii) we have

$$
x_{\epsilon} - y_{\epsilon} \ge \max{\{\hat{\Gamma}_{\phi}(\phi^{-1}(\phi(y_{\epsilon})), \hat{\Gamma}_{\phi}(x_{\epsilon})\}} > \hat{\Gamma}_{\phi}(t_1),
$$

which implies that $x_0 - y_0 > \hat{\Gamma}_{\phi}(t_1) - 2\epsilon$. Letting $\epsilon \to 0$ we have $x_0 - y_0 \geq \hat{\Gamma}_{\phi}(t_1)$. Now

$$
\bar{C}_{\Gamma_{\phi}}(x_0, \phi(y_0)) = \min \left\{ \phi(y_0), x_0 - \frac{1}{2} \Big(\mathbb{V}_{y_0}^{x_0}(\hat{\Gamma}_{\phi}) + \hat{\Gamma}_{\phi}(x_0) + \hat{\Gamma}_{\phi}(y_0) \Big) \right\}
$$

= $\min \{ \phi(y_0), x_0 - \hat{\Gamma}_{\phi}(t_1) \} = \phi(y_0).$

Since $\bar{C}_{\Gamma_{\phi}}$ is a quasi-copula, it is increasing and 1-Lipschitz. Thus,

$$
\overline{C}_{\Gamma_{\phi}}(x,y) \ge \overline{C}_{\Gamma_{\phi}}(x_0,y) \ge \overline{C}_{\Gamma_{\phi}}(x_0,\phi(y_0)) - (\phi(y_0) - y) = y.
$$

Thus $\bar{C}_{\Gamma_{\phi}}(x,y) = A_{\Gamma_{\phi}}(x,y)$ also holds.

$$
\Box
$$

References

- [1] U. Cherubini, E. Luciano, W. Vecchiato, Copula Methods in Finance, John Wiley & Sons, Chichester, 2004.
- [2] F. Durante, P. Jaworski, Absolutely continuous copulas with given diagonal sections, Commun. Stat., Theory Methods 37 (2008) 2924-2942.
- [3] Y. Fang, L. Madsen, Modified Gaussian pseudo-copula: Applications in insurance and finance, Insur. Math. Econ. 53 (2013) 292-301.
- [4] K. Goda, J. Ren, Assessment of seismic loss dependence using copula, Risk Analysis 30 (2020) 1076-1091.
- [5] H. Joe, Dependence Modeling with Copulas, CRC Press, Boca Raton, doi:10.1201/b17116, 2015.
- [6] T. Jwaid, H. De Meyer, A. Haj Ismail, B. De Baets, Curved splicing of copulas, Inform. Sci. 556 (2021) 95-110.
- [7] D. Kokol Bukovšek, B. Mojškerc, N. Stopar, Exact upper bound for copulas with a given diagonal section, Fuzzy Sets Syst. 480 (2024) 108865.
- [8] R.B. Nelsen, An Introduction to Copulas, Springer Science & Business Media, New York, 2006.
- [9] R.B. Nelsen, J.J. Quesada-Molina, J.A. Rodríguez-Lallena, M. Úbeda-Flores, Best-possible bounds on sets of bivariate distribution functions, J. Multivar. Anal. 90 (2004) 348-358.
- [10] R.B. Nelsen, J.J. Quesada-Molina, J.A. Rodríguez-Lallena, M. Úbeda-Flores, On the construction of copulas and quasi-copulas with given diagonal sections, Insur. Math. Econ. 42 (2008) 473-483.
- [11] Y. Ouyang, Y. Sun, H. Zhang, On an upper bound of the set of copulas with a given curvilinear section, Fuzzy Sets Syst. 500 (2025) 109199.
- [12] S. T. Rachev, L. Rüschendorf, Solution of some transportation problems with relaxed or additional constraints, SIAM J. Control Optim. 32 (1994) 673-689.
- [13] A. Sklar, Fonctions de répartition à n dimensions et leurs marges, Publ. Inst. Stat. Univ. Paris 8 (1959) 229-231.
- [14] Y. Tang, H.J. Wang, Y. Sun, A.S. Hering, Copula-based semiparametric models for spatiotemporal data, Biometrics, 75 (2019) 1156-1167.
- [15] W. Zou, L. Sun, J. Xie, Best-possible bounds on the sets of copulas and quasi-copulas with given curvilinear sections, Fuzzy Sets Syst. 441 (2022) 335-365.