Structural controllability and management of cascading regime shifts

Juan C. Rocha¹, Anne-Sophie Crépin²

¹Stockholm Resilience Centre, Stockholm University

²Beijer Institute of Ecological Economics, The Royal Swedish Academy of Sciences

Abstract

Abrupt transitions in ecosystems can be interconnected, raising challenges for science and management in identifying sufficient interventions to prevent them or recover from undesirable shifts. Here we use principles of network controllability to explore how difficult it is to manage coupled regime shifts. We find that coupled regime shifts are easier to manage when they share drivers, but can become harder to manage if new feedbacks are formed when coupled. Simulation experiments showed that both network structure and coupling strength matter in our ability to manage interconnected systems. This theoretical observation calls for an empirical assessment of cascading regime shifts in ecosystems and warns about our limited ability to control cascading effects.

Introduction

Regime shifts are large, abrupt and persistent changes in the function and structure of systems^{1,2}. They have been documented in a variety of social-ecological systems^{3,4}, as well as climate⁵, finance⁶, or health⁷, to name a few examples. Changes in structure and function of ecosystems can diminish the benefits people get from nature such as food production, employment opportunities, climate regulation, or water purification⁸. For example, the North Atlantic cod collapse represented millions of economic loss and compromised more than 35000 jobs in 400 coastal communities⁹. The weakening or late arrival of the Indian summer monsoon can compromise food production in one of the most densely populated regions of the world. How management strategies avoid regime shifts or recover natural ecosystems to desirable regimes are key research questions and an active area of research.

But one theoretical fact that complicates their study and management is that regime shifts can be interconnected¹⁰. The occurrence of one can impact the likelihood of another through sequential tipping, like a domino effect^{10,11}; or through two-way interactions where new feedbacks can amplify or dampen the probability of tipping $over^{10}$. Conceptual work have looked at these connections in the context of committed risk 5,12 , and recent modeling work has studied at interactions between climate tipping elements to understand additional challenges of prediction¹³, or the role of minimal network configurations¹⁴. Besides climate, cascading effects have also been studied in the context of engineered systems 15,16 , social and economic systems¹⁹, or diseases²⁰.

The possibility of cascading regime shifts raises a few additional challenges for management. First, long distance coupling or teleconnections²¹ between regime shifts implies that optimal local or regional managerial practices might not be enough to reach managerial objectives^{10,22}. This not only illustrates the governance issue of with whom to coordinate for successful management of local resources; it also implies a tragedy of the commons with potential power asymmetries. Classical examples include transnational resource systems where pollution or exploitation of a resource on one side of the border impacts its quality on the other side, or downstream^{23,24}. Second, and most importantly, connected regime shifts pose the challenge of identifying sufficient and appropriate points of intervention for managerial actions. In an interconnected world, how can we identify leverage points to manage regime shifts? Are there regime shifts – or their interactions - that are harder to manage?

Here we answer these questions in the light of controllability of complex systems. A system is controllable if one can drive it from any initial state to any desired final state in finite time^{25–27}. Hence, controllability quantifies our ability to steer dynamical systems²⁷. While control theory has been fairly well developed in the realm of linear systems, many open problems remain for nonlinear ones^{26,28}. Strict global controllability of nonlinear systems remain elusive, but weaker notions of controllability can be described $26,29,30$. For example, open loop control has been used to steer

Figure 1: **Structural controllability of individual regime shifts** The number of nodes of the control set per regime shift (A), and their proportion with respect to the full causal network (B). Top controlling variables ranked by the number of regime shifts where they are part of the control set.

chaotic systems to desired trajectories in the phase space. Here we focus on feedback vertex sets as a proxy of structural controllability^{26,29} of regime shifts and their interactions. We use network topology to gain insights on the controllability of regime shift and the management challenges of their cascading effects. We further test our intuitions with minimal models of interconnected resources and pollution recipient systems.

Methods

We study structural controllability first in a static setting, and then using minimal simulation experiments to approximate the dynamics of interconnected systems. For the static setting we used causal diagrams derived from the regime shifts database, an open online repository of regime shift reviews⁴. The database offers causal graphs that summarize scientific hypotheses of how regime shifts work. Each directed graph (N=30) can be interpreted as a network where a link between two variables exists if a scientific paper has suggested a causal link to the occurrence of the regime shift⁴. We use these networks to study structural controllability (Fig [S1\)](#page-1-0).

Feedback control: To steer a nonlinear system to a desired attractor, one needs to find the feedback vertex set of the system. When the system is described as a network, its state variables are represented by nodes (or vertices), and the dependence between variables as directed links (or arcs). A feedback in this context is a collection of nodes and links that form a circular pathway. Feedbacks are often responsible for the nonlinear dynamics of the system. If a network has feedbacks, it is called a directed cyclic graph (DCG), and if it lacks them a directed acyclic graph (DAG). The feedback vertex set is the set of nodes that if deleted would render a DAG. The feedback arc set is the set of links that if eliminated would render a DAG. Finding the feedback vertex set or the arc set are equivalent problems³¹, yet finding the minimum set is a nondeterministic polynomial-time hard (NPhard) problem. Once the system is reduced to a DAG, linear control methods can be applied, such as finding unmatched nodes or the Kalman rank condition for linear systems^{26,27}. Hence, the set of nodes that need to be controlled to gain leverage on the behaviour of a nonlinear system are the feedback vertex set plus the unmatched nodes of the remaining linear system 26,29 .

Structural control on causal networks We compute the proportion of nodes that needs to be controlled for all causal networks from the regime shifts database (Fig [S1\)](#page-1-0), as well as their pair-wise combinations. The pair-wise combinations have been used to study plausible mechanisms that can couple different types of regime shifts known as cascading effects¹⁰. We then compare how structural controllability changes if ecological regime shifts are interconnected.

Dynamic control on minimal models To further test our intuitions, we compared the results from structural controllability with modeling experiments in simplified resource and pollutant models. We chose these models because they are easier to parametrise than the qualitative networks offered by the database, yet

Figure 2: **Structural controllability of coupled regime shifts** The proportion of nodes that need to be controlled to manage individual regime shifts (A) compared against the number of nodes (B) and ratio of nodes (C) that one needs to control to manage coupled regime shifts. Top controlling nodes are ranked by the number of pair-wise compbinations where they appear on the control set (D). For a sample of 30 regime shifts, there are 900 pair-wise combinations or 870 combinations of non-identical regime shifts. Thus, for example, fertilizers use and sewage appear in more than 175 of these combinations.

they encapsulate well understood and common dynamics underlying many real world regime shifts. The pollution system is defined by:

$$
\frac{dx_i}{dt} = u_i - s_i x_i + v_i \frac{x_i^{\alpha_i}}{z_i^{\alpha_i} + x_i^{\alpha_i}} - \sum_{j \neq i} (\delta_{ij} x_i - \delta_{ji} x_j) A_{ij}
$$

where x_i is the level of pollutants of patch i (e.g. a lake), u_i denotes the level of pollutants from human activities, and s_i the internal loss rate (e.g. sedimentation). The term $v_i \frac{x_i^{\alpha_i}}{z_i^{\alpha_i} + x_i^{\alpha_i}}$ is a pollutant release nonlinear function where v_i is the maximum level of internal pollutants, α_i indicates the sharpness of the function, and z_i is a threshold level at which the system flips from low to high nutrient regimes. Connectivity across patches is represented with the term $\sum_{j\neq i} (\delta_{ij}x_i - \delta_{ji}x_j)A_{ij}$ where δ_{ij} is the diffusion coefficient from patch $i \rightarrow j$, given that δ_{ij} and δ_{ji} are not necessarily the same; while A_{ij} is the adjacency matrix. Similarly, the resource system is defined by:

$$
\frac{dy_i}{dt} = r_i y_i \left(1 - \frac{y_i}{k_i} \right) - c_i \frac{y_i^{\beta_i}}{q_i^{\beta_i} + y_i^{\beta_i}} + \sum_{j \neq i} (\delta_{ij} y_j - \delta_{ji} y_i) A_{ij}
$$

The term $c_i \frac{y_i^{\beta_i}}{q_i^{\beta_i} + y_i^{\beta_i}}$ is a Holling type III predation function where c_i represent refuges from predators in patch i, q_i is the half saturation biomass for predators (e.g. a threshold for high to low abundance), and β_i is the curvature parameter that defines the abruptness of the shift³². The last term indicates mobility of the resource between patches at a diffusion rate $\delta_{ij} \neq \delta_{ji}$, and an adjacency matrix A_{ij} that takes 1 if patches *i* and *j* are connected, or zero otherwise. For both systems the diffusion coefficient $(\delta_{i,j})$ is normalized by the out degree, or the number of outgoing connections from each system. This is to ensure that the amount of pollutants or resources flowing from one system to another never exceeds the amount available at run time.

where y_i is the state of the population *i*, r_i is the $(\delta_{i,j} : \{low = 0.01 : 0.05, high = 0.25 : 0.5\})$. Thus, intrinsic growth rate, and k_i its carrying capacity. we have 96 different experimental conditions for which Here we concentrate our efforts in exploring numerical simulations of medium size interconnected systems $(N < 10²)$, with potentially multiple feedbacks. Analytical solutions for the low dimensional diffusion problem of our pollution and resource systems can be found in^{33} . For each of the dynamical systems (pollution, resource) we generated networks varying its size $(N:\{25, 50, 75, 100\})$; the network generative process which could be random (Erdös-Rényi model), β preferential attachment (Barabási-Albert model), or small-world (Watts-Strogatz model); the network density (*d* : {0*.*05*,* 0*.*1*,* 0*.*3*,* 0*.*5}); and the coupling strength

we generated 100 replicates resulting in 9600 simulation experiments for each dynamical system. Each model was run for 200 time steps and integration steps of 0.001 *dt*.

For each experiment we identified the feedback vertex set and the minimum controlling set of the network. Then we manipulated the driver at run time to decrease nutrients in the pollution system (*u* decreased from 5 to -5 allowing for management that can extract nutrients from the system), or decrease harvest in the resource system (*c* decreased from 10 to 0) on the controlling set only. We then assess if management of the control set successfully recovers the networked systems, the proportion of nodes recovered, or time to recovery when comparing strong versus weak coupling (δ) . The computer code used for our simulations as well as the curated networks from the regime shifts database are available at: <https://github.com/juanrocha/imperio>

Results

Individual regime shifts cannot be controlled or managed by only targeting variables that have been coded as drivers in the regime shifts database. Drivers there are defined as variables outside feedbacks that influence the dynamics of the system, but that are not in turn influenced by the system itself⁴; which inherits the logic first proposed by the Millennium Ecosystem Assessment³⁴. As recent advances in network controllability show26,29, to steer the system towards a desired state, an ideal manager needs to intervene not only on drivers but also on some of the variables involved in feedback dynamics. In all the individual regime shifts analyzed, an ideal manager needs to account for the minimum control set of the DAG and the feedback vertex set (Fig [1\)](#page-1-0). By number of driving nodes, regime shifts such as forest to savanna or sea grass transitions require higher efforts; but by proportion of nodes: thermokarst lakes, primary productivity in the Arctic ocean, tundra to forest, or marine food webs are among the regime shifts where $>60\%$ of nodes need some intervention. The top controlling variables are, as expected, related to climate change, biodiversity loss, or food production (Fig [1C](#page-1-0)). We observe that trophic groups such as zooplankton, meso predators, or top predators are often part of the feedback vertex set, while fertilizers use, sewage or soil erosion are always part of the minimum control set of the DAG.

Coupled regime shifts are harder to manage because the ideal manager needs to influence more variables. However, the number and proportion of nodes to be controlled varies depending on the coupling (Fig [2\)](#page-2-0). On average, coupled regime shifts are harder to manage than individual counterparts (Fig [2\)](#page-2-0). For example, preventing the West Antarctica Ice Sheet collapse (WAIS) implies being able to intervene on a minimum of 40% of the variables describing processes underlying the regime shift. But when combined with other regime shifts, the percentage increases to 49% on average, and a maximum of 64%. Similarly, managing marine food webs regime shifts requires intervening on $~66\%$ of the variables describing the individual shift; but when coupled it varies from as low as 53% with kelp transitions to as high as 72% with Arctic benthos borealization. This is because sharing common variables on the minimum driving set reduces the proportion of nodes to control, while new feedbacks formed when coupling the network, increase the feedback vertex set. The most common variables to manage coupled regime shifts are fertiliser use, sewage, fishing and soil erosion. Climate related variables are important but down the rank in the list, such as droughts, temperature, and precipitation. Many variables that have not been previously reported in similar studies about key drivers of regime shifts are reported here because they are part of the feedback vertex set, such as management of apex predators, zooplankton, or fish populations. This however aligns with management strategies that attempt to manipulate feedback strength to recover ecosystems from regime shifts, for example fish population manipulations to recover from eutrophication³⁵.

These results suggest that network structure can have implications for management. If two regime shifts are solely coupled by sharing drivers, then controlling the common drivers will help the ideal manager to tackle both regime shift problems. But if the coupling of regime shifts generates new feedbacks, the minimum number of processes that need to be controlled increases, making it more difficult to manage the coupled system than the independent ones. To test that intuition, we developed simple toy models of resource and pollutant systems where we varied network structure. The models were not as complex as the causal graphs previously studied, but they still capture the essential dynamics. Both typeS of systems in isolation can be controlled by one parameter, the addition or removal of pollutants *uⁱ* , and the availability of refuges from predators c_i respectively. Thus when coupled, any gains or losses on the ability to manage the regime shifts is due to network structure alone (the coupling matrix $A_{i,j}$).

In the pollution system, the higher the network density the higher the proportion of nodes recovered in our simulations (Fig $3A$). This effect is accentuated when the coupling strength δ is high. However, the propor-

 \overline{A} Proportion of nodes recovered 0.75 0.50 0.2 0.00 $_{0.3}$ 0.05 0.1 0.5 0.05 $\overline{0.1}$ $0.\overline{3}$ 0.5 vork density $\sf B$ Proportion of nodes recovered
that belongs to control set 0.05 01 0.3 0.5 0.5 o'i 0.3 Network density \mathbf{C} $20₀$ Mean recovery time 150 $10[°]$ $\overline{\mathbf{a}}$ Propo rtion of nodes recovered Туре .
DAG DCG ilgh

Figure 3: **Pollution system simulations** The proportion of nodes recovered by network density for different types of network architecture and coupling strength (A). The proportion of nodes recovered that belongs to the control set (B), and the mean recovery time by proportion of nodes recovered (C). Boxplots and scaterplots summarize $N = 9600$ networks (100) replicates * 96 experimental settings).

tion of nodes recovered that belong to the control set is higher when coupling strength is low (Fig [3B](#page-4-0)) for all network types tested. This means that coupling strength increases the probability that a larger proportion of the system is recovered by managerial actions on the control set, but the control set in turn becomes vulnerable through incoming connections that impede complete recovery. We also observe that mean recovery time is on average faster with high coupling strength, but in networks whose generative process is preferential attachment (scale-free like degree distributions) and strong presence of dyadic acyclic graphs (DAG), recovery can sometimes be slower under high coupling (Fig [3C](#page-4-0)). Regardless of network architecture, the lower bound for proportion of node recovery is as low as 60% for preferential attachment networks, around 30% for random and small-world networks under high coupling strength, and almost none for low

Figure 4: **Resource system simulations** The proportion of nodes recovered by network density for different types of network architecture and coupling strength (A). The proportion of nodes recovered that belongs to the control set (B), and the mean recovery time by proportion of nodes recovered (C). Boxplots and scaterplots summarize $N = 9600$ networks (100) replicates * 96 experimental settings).

coupling.

We find contrasting results for the resource simulations. First, the proportion of nodes recovered declines on average with increasing network density, although for the case of preferential attachment, all nodes were recovered regardless of network density in the low coupling strength scenario (Fig [4A](#page-4-1)). Second, the proportion of nodes recovered that belong to the control set increases with network density for the cases of random and small-world networks, but there are no differences between the high and low coupling strength (Fig [4B](#page-4-1)). In the case of preferential attachment, the proportion of recovered nodes that belong to the control set is higher in networks under the low coupling strength. High coupling strength reduces the chances of recovery for the control set. While most simulated systems under low coupling strength recovered by time step ~130, recovery time varied much more in networks

with high coupling strength (Fig $4C$).

Discussion

Our simulation results confirm that network structure matters when trying to manage coupled regime shifts. The type of network architecture (random, preferential attachment, small-world) changes the likelihood of recovering the full system, as well as the coupling strength between systems. Surprisingly, even under the same network configurations (N=9600) the type of dynamics occurring in the networks (pollutant versus resource equations) substantially influenced the qualitative patterns identified for the full ensemble (Figs [3,](#page-4-0) [4\)](#page-4-1).

In many of our experimental settings the system only recovered partially despite having full information of the control set at the start of the simulation. One plausible explanation is that the system required longer time to recover. A more likely explanation however is that our network setting allowed for changes in the amount and direction of flow of nutrients and pollutants. In both systems the diffusion term depended on the difference in concentration between patches $(\sum_{j\neq i} (\delta_{ij}y_j - \delta_{ji}y_i)A_{i,j})$. While the adjacency matrix $A_{i,j}$ and the diffusion coefficients $\delta i, j$ were constant through the simulation, the direction of flow could change if the difference $\delta_{ij}y_j - \delta_{ji}y_i$ was positive or negative. In practice, this means that the feedback structure and thus the control set can change over time. Another plausible explanation is that the adjacency matrix (A_{ij}) is non-normal. Networks are non-normal if $A\ddot{A}^T \neq A^T A$, which is the case in our directed graphs, although matrix asymmetry is not sufficient for strong non-normality³⁶. Strong non-normality emerges when the network is directed (asymmetrical), has low reciprocity, lacks feedback, and has hierarchical organization³⁶. While our networks have feedbacks, we cannot discard the possibility that the lack of recovery is due to non-normal induced features on the dynamics. The non-normality of networks increases their return time to equilibrium, thus reducing their resilience³⁷.

Knowing the control set at one moment in time is insufficient to gain full control, and thus recovery, of the connected system. Deriving the control set at the beginning of the simulation resulted in insufficient efforts to successfully stir the system towards full recovery (Figs [3,](#page-4-0) [4\)](#page-4-1). In practice, this is likely to be the case if one is managing a meta-population where organisms are free to move from patch to another and likely to go to places where they find better resources, which change over time. It is less likely to occur in settings where the flow of resources or pollutants is bounded by a physical parameter. For example, water between lakes flows according to gravity, so nutrients are unlikely to be transported against it. While our assumption can be unrealistic for certain cases, it opens the case when the control set changes over time. In most real settings we do not even know the real empirical structure of the network –how regime shifts are connected¹⁰. Our results suggest that even if we knew the structure at a certain point in time, successfully steering the networked system requires monitoring programs that enable learning the structure, direction and coupling strength of the system in real time.

This dynamical feature is unlikely to affect our structural controllability results derived from causal networks (Fig [2\)](#page-2-0), because the direction of causality is unlikely to change. However, in a dynamical setting the feedback strength or the coupling of different variables could change, akin to δi , *j* in our simulations. Thus to apply efficient managerial practices that avoid tipping cascades, one could gain further insights on the priority of variables for management, or sequence of actions required. Our results showed that while climate is important, other coupling mechanisms can be more relevant in the sense that they appear more in pair-wise combinations (Fig [2\)](#page-2-0). Our results also confirm that dealing with climate related variables alone won't be enough to prevent tipping cascades¹⁰, after all, these variables are a small share of the control set.

Our modelling framework opens potential avenues for future research. Thus far our assumption is that an ideal manager can intervene in all subsystems of the controlling set simultaneously. Relaxing thus assumption would question whether it is possible to recover the system with asynchronous control. We have not yet considered the cost of interventions in the system, nor time bounds. With limited budget or time to achieve the controllability goals, what would be an optimal strategy? or a sequence of strategies? Another interesting avenue for future research is a case with multiple managers who have to coordinate to avoid the tipping cascade, adding a game theoretic layer to the problem. We believe that relaxing any of our simplifying assumptions to make the problem more realistic would inevitably make it harder to control.

Our results show that interconnected systems put certain subsystems and their managers in a privileged position to steer the system, for good and bad. Given these power asymmetries, is coordination and cooperation likely to emerge to avoid tipping cascades? The question is imperative since both the climate and biodiversity crises underlie a similar networked structure to the framework here proposed, where a few

actors have disproportional agency to influence the overall dynamics of the system. Moreover, our results show that when systems are interconnected, there is no guarantee that one can recover the full system. This implies that current climate narratives that rely on the potential recovery from overshoot scenarios³⁸ should be taken with skepticism, because the probability of recovery seen in simple one-dimensional models does not necessarily hold in higher dimensions. Here we show that the ability to control is reduced. In fact, if one is to believe that tipping points can be interconnected^{5,10,11}, each connections becomes a codimension (a driver) of other tipping points, so the problem is never one-dimensional.

Conclusion

By applying principles of structural controllability, we show here that connected regime shifts are more difficult to manage and recover than independent systems. Sharing common drivers can perhaps simplify the problem, but the emergence of feedbacks increases the number of variables that need interventions. Interconnected systems come with additional dynamic challenges for network control and management: being able to observe and measure how the control set changes over time. In more realistic settings, the network structure imposes additional challenges for the emergence of cooperation, power asymmetries, the cost, timing, and sequence of managerial actions. These questions remain open areas of research. Yet, tackling the climate or biodiversity crises requires learning such networks from empirical data. Our theoretical exercise shows that management practices that can work on one dimensional systems such as overshooting scenarios for climate may not necessarily succeed in high dimensional systems. We appeal for the precautionary principle when dealing with interconnected crises.

Acknowledgements

We acknowledge support from VR grant 2022-04122, Formas grants 2015-00731, 2020-00454 and 2019-02316, the latter through the support of the Belmont Forum.

References

1. Folke, C. *et al.* Regime shifts, resilience, and biodiversity in ecosystem management. *Annu Rev Ecol Evol S* **35,** 557–581 (2004).

- 2. Scheffer, M., Carpenter, S., Foley, J., Folke, C. & Walker, B. Catastrophic shifts in ecosystems. *Nature* **413,** 591–596 (2001).
- 3. Scheffer, M. *Critical Transitions in Nature and Society*. (Princeton University Press, 2009).
- 4. Biggs, R., Peterson, G. & Rocha, J. The Regime Shifts Database: a framework for analyzing regime shifts in social-ecological systems. *Ecology and Society* **23,** art9 (2018).
- 5. Lenton, T. M. *et al.* Climate tipping points too risky to bet against. *Nature* **575,** 592–595 (2019).
- 6. Guttal, V., Raghavendra, S., Goel, N. & Hoarau, Q. Lack of Critical Slowing Down Suggests that Financial Meltdowns Are Not Critical Transitions, yet Rising Variability Could Signal Systemic Risk. *PLoS ONE* **11,** e0144198 (2016).
- 7. Leemput, I. A. van de *et al.* Critical slowing down as early warning for the onset and termination of depression. *P Natl Acad Sci Usa* **111,** 87–92 (2014).
- 8. Carpenter, S. R. *et al.* Science for managing ecosystem services: Beyond the Millennium Ecosystem Assessment. *P Natl Acad Sci Usa* **106,** 1305–1312 (2009).
- 9. Gien, L. T. Land and sea connection: The east coast fishery closure, unemployment and health. *Canadian Journal of Public Health* **91,** 121–124 $(2000).$
- 10. Rocha, J. C., Peterson, G., Bodin, O. & Levin, S. Cascading regime shifts within and across scales. *Science* **362,** 1379–1383 (2018).
- 11. Wunderling, N., Donges, J. F., System, J. K. E. & Zubov, K. Interacting tipping elements increase risk of climate domino effects under global warming. *esd.copernicus.org*
- 12. Steffen, W. *et al.* Trajectories of the Earth System in the Anthropocene. **115,** 8252–8259 (2018).
- 13. Dekker, S. C. *et al.* Biogeophysical feedbacks trigger shifts in the modelled vegetation-atmosphere system at multiple scales. *BIOGEOSCIENCES* **7,** 1237–1245 (2010).
- 14. Wunderling, N. *et al.* How motifs condition critical thresholds for tipping cascades in complex networks: Linking micro- to macro-scales. *Chaos* **30,** 043129 (2020).
- 15. Brummitt, C. D., D'Souza, R. M. & Leicht, E. A. Suppressing cascades of load in interdependent networks. *P Natl Acad Sci Usa* **109,** E680–9 (2012).
- 16. D'Souza, R. M. Curtailing cascading failures. *Science* **358,** 860–861 (2017).
- 17. Lee, K. *et al.* Impact of the topology of global macroeconomic network on the spreading of economic crises. *PLoS ONE* **6,** e18443 (2011).
- 18. Brummitt, C. D., Barnett, G. & D'Souza, R. M. Coupled catastrophes: sudden shifts cascade and hop among interdependent systems. *J. R. Soc. Interface* **12,** 20150712–20150712 (2015).
- 19. Brummitt, C. D., Huremović, K., Pin, P., Bonds, M. H. & Vega-Redondo, F. Contagious disruptions and complexity traps in economic development. *Nat. hum. behav.* **77,** 1–672 (2017).
- 20. Watts, D. J., Muhamad, R., Medina, D. C. & Dodds, P. S. Multiscale, resurgent epidemics in a hierarchical metapopulation model. **102,** 11157– 11162 (2005).
- 21. Liu, J. *et al.* Systems integration for global sustainability. *Science* **347,** 1258832–1258832 (2015).
- 22. Rocha, J. C., Peterson, G. D. & Biggs, R. Regime Shifts in the Anthropocene: Drivers, Risks, and Resilience. *PLoS ONE* **10,** e0134639 (2015).
- 23. Young, O. R. *The institutional dimensions of environmental change: Fit, interplay, and scale*. (MIT press, 2002).
- 24. Keys, P. W., Wang-Erlandsson, L. & Gordon, L. J. Megacity precipitationsheds reveal tele-connected water security challenges. *PLoS One* **13,** e0194311 (2018).
- 25. Kalman, R. E. Mathematical Description of Linear Dynamical Systems. *Journal of the Society for Industrial and Applied Mathematics Series A Control* **1,** 152–192 (1963).
- 26. Liu, Y.-Y. & Barabasi, A.-L. [Control Principles of](https://arxiv.org/abs/1508.05384v2) [Complex Networks.](https://arxiv.org/abs/1508.05384v2) 247 (2015).
- 27. Liu, Y.-Y., Slotine, J.-J. & Barabasi, A.-L. Controllability of complex networks. *Nature* **473,** 167–173 (2011).
- 28. Cowan, N. J., Chastain, E. J., Vilhena, D. A., Freudenberg, J. S. & Bergstrom, C. T. Nodal Dynamics, Not Degree Distributions, Determine the Structural Controllability of Complex Networks. *PLoS ONE* **7,** e38398 (2012).
- 29. Zañudo, J. G. T., Yang, G. & Albert, R. Structurebased control of complex networks with nonlinear dynamics. **335,** 201617387 (2017).
- 30. Mochizuki, A., Fiedler, B., Kurosawa, G. & Saito, D. Dynamics and control at feedback vertex sets. II: A faithful monitor to determine the diversity of molecular activities in regulatory networks. *J. Theor. Biol.* **335,** 130–146 (2013).
- 31. Zhou, H.-J. A spin glass approach to the directed feedback vertex set problem. *Journal of Statistical Mechanics: Theory and Experiment* **2016,** 073303 (2016).
- 32. Holling, C. S. Some Characteristics of Simple Types of Predation and Parasitism. *Can Entomol* **91,** 385–398 (2012).
- 33. Crépin, A.-S. & Rocha, J. C. Cascading regime shifts in pollution recipients and resource systems. (2021).
- 34. Ecosystem Assessment, M. *Ecosystems and human well-being: Synthesis*. 26 (Island Press, 2005).
- 35. Smith, V. H. & Schindler, D. W. Eutrophication science: Where do we go from here? *Trends in ecology & evolution* **24,** 201–207 (2009).
- 36. Asllani, M., Lambiotte, R. & Carletti, T. [Structure](https://doi.org/10.1126/sciadv.aau9403) [and dynamical behavior of non-normal networks.](https://doi.org/10.1126/sciadv.aau9403) *Science Advances* **4,** (2018).
- 37. Asllani, M. & Carletti, T. [Topological resilience in](https://doi.org/10.1103/physreve.97.042302) [non-normal networked systems.](https://doi.org/10.1103/physreve.97.042302) *Physical Review E* **97,** (2018).
- 38. Ritchie, P. D. L., Clarke, J. J., Cox, P. M. & Huntingford, C. [Overshooting tipping point thresholds](https://doi.org/10.1038/s41586-021-03263-2) [in a changing climate.](https://doi.org/10.1038/s41586-021-03263-2) *Nature* **592,** 517–523 (2021).

Supplementary Material

Figure S1: **Causal networks** The regime shifts database (www.regimeshifts.org) reports causal hypothesis of how regime shifts work synthesized in causal loop diagrams, where variables are connected by arrows if a causal mechanisms is reported in scientific literature. Variables in red are drivers (outside feedbacks), in grey are variables that belong to feedback mechanisms, links in blue are positive and orange negative. We used the version of the database reported in ref 10.