# ANALYZING SCALOGRAM RIDGES IN THE PRESENCE OF NOISE

GI-REN LIU, YUAN-CHUNG SHEU, AND HAU-TIENG WU

ABSTRACT. While ridges in the scalogram, determined by the squared modulus of analytic wavelet transform (AWT), have been widely applied in time series analysis and time-frequency analysis, their behavior in noisy environments remains underexplored. We fill in this gap in this paper. We define ridges as paths formed by local maximizers of the scalogram along the scale axis, and analyze their properties when the signal is contaminated by stationary Gaussian noise. In addition to establishing several key properties of the AWT for random processes, we investigate the probabilistic characteristics of the resulting random ridge points in the scalogram. Specifically, we establish the uniqueness property of the ridge point at individual time instances and prove the upper hemicontinuity of the set of ridge points. Furthermore, we derive bounds on the probability that the deviation between the ridge points of noisy and clean signals exceeds a specified threshold. These bounds are expressed in terms of constants related to the signal-to-noise ratio and apply when the signal satisfies the adaptive harmonic model and is corrupted by a stationary Gaussian process. To achieve these results, we derive maximal inequalities for the complex modulus of nonstationary Gaussian processes, leveraging classical tools such as the Borell-TIS inequality and Dudley's theorem, which might be of independent interest.

#### 1. INTRODUCTION

Ridge analysis plays a vital role in connecting time-frequency analysis [8, 15] and signal processing [26], particularly when the notion of "instantaneous frequency" (IF) is of interest. Ridges in the time-frequency representation (TFR) of nonstationary time series provide critical information about IF, and the extracted ridge is utilized as the estimate of IF, serving as a cornerstone for various data analysis applications [15, 26]. Although the concept of ridges is widely utilized in practice, and numerous algorithms have been developed to extract them [4, 5, 6, 19, 21, 22, 28, 30, 38, 41], the properties of ridges, particularly in the presence of noise, remain underexplored. This paper aims to establish a theoretical foundation to address this gap. Before proceeding, it is worth noting that the term "ridge" has a long history of usage in statistics [16], image analysis [13], etc. However, due to differences in dataset characteristics and data analysis mission, this usage is not equivalent to ridges in the TFR. In the following, we focus exclusively on ridges within TFRs.

Traditionally, a ridge in a TFR is often understood in an "ad hoc" manner as a sequence of local maxima along the time-indexed frequency or scale axis of the TFR. This interpretation likely stems from the visualization of a time series as a series of mountain-shaped bumps within the time-frequency or time-scale domain, with the associated ridge forming a continuous curve. This continuity property is explicitly incorporated into the definition of ridges for adapted harmonic signals [23] and plays a critical role in the development of ridge extraction algorithms [4, 5, 6, 19, 21, 22, 28, 30, 38, 41]. However, in noisy environments, numerical experiments reveal that the TFRs of noise-affected signals exhibit more complex patterns compared to their noise-free counterparts. See Figure 1 for a comparison. In the top portion of Figure 1, the input is a clean signal,  $f(t) = A(t) \cos(2\pi\phi(t))$ , where A represents a smooth, positive and slowly varying amplitude modulation and  $\phi$  represents a smooth, monotonically increasing phase function with a slowly varying derivative  $\phi'(t)$ ; that is, f(t) is locally close to a harmonic function. In the bottom portion of Figure 1, the input is a noisy signal, which is a realization of  $Y(t) = f(t) + \Phi(t)$ , where  $\Phi(t)$  is a Gaussian random process with mean zero and finite variance. The TFRs clearly exhibit multiple local maxima along each timeindexed frequency or scale axis, with different behaviors caused by noise. Another notable phenomenon is that the maxima on each cross-section of the TFR for a noise-affected adapted harmonic signal may exhibit abrupt jumps at certain times; for example, around the 23rd second (see Figures 2 and 4 for more details). These observations highlight the need for a more rigorous definition of ridges under noisy conditions and a detailed investigation of the statistical behavior of ridges.

In this work, we consider the noisy signal as  $Y = f + \Phi$ , where f fulfills the adaptive harmonic model; that is, it comprises multiple oscillatory components with time-varying amplitude modulation and IF (details below) [9] and  $\Phi$  is a Gaussian random process exhibiting either long- or short-range dependence. We focus on the TFR obtained using the continuous wavelet transform with an analytic mother wavelet, referred to as the analytic wavelet transform (AWT). For the clean signal f, the AWT produces the TFR of f, which is a complex-valued function  $\{W_f(t,s) \in \mathbb{C} \mid t \in \mathbb{R}, s > 0\}$  defined in the time-scale domain, indexed by time and scale variables. The squared modulus of this function, known as the scalogram and denoted by  $\{S_f(t,s) \in [0,\infty) \mid t \in \mathbb{R}, s > 0\}$ , captures the spectral energy distribution of f as it varies over time. In the time-scale domain, the graph of the map

$$s_f(t) = \arg \max_{s \in B(t)} S_f(t,s), \ t \in \mathbb{R},$$

where  $B(t) \subset (0, \infty)$  represents either the entire scale axis  $(0, \infty)$  or a bounded scale region, specified by the user, is referred to as the wavelet ridge [26] or the wavelet skeleton [2, pp. 14-18]. We may directly extend this concept to the random process Y by defining

$$s_Y(t) = \arg \max_{s \in B(t)} S_Y(t,s), \ t \in \mathbb{R},$$

and refer to  $s_Y$  as the ridge function of Y. Note  $s_f(t)$  is generally a set-valued function, and  $s_Y(t)$  is in general a random process due to the randomness of  $\Phi$ , and its structure is more complicated. It is important to note that focusing on the adaptive harmonic model is less restrictive than it might initially appear. To our knowledge, analyzing a generic signal under the notion of IF, and hence the ridge, remains a challenging task. However, when the analysis centers on IF, researchers typically employ the adaptive harmonic model to facilitate the estimation of IFs by extracting energy-concentrated curves from the scalograms (or spectrograms) of noisy signals [4, 5, 19, 37, 38].

Our contributions in this paper are multifaceted. First, we conduct a detailed analysis of the AWT and its inverse transform applied to random processes, covering its regularity properties and a novel observation regarding the covariance structure of the scalogram of  $\Phi$ . To our knowledge, these aspects are underexplored in the literature, yet they are crucial for statistical inference in signal processing using AWT. Second, we propose a definition of ridges for noisy input signals that aligns with practical needs. Under this framework, we establish the uniqueness and upper hemicontinuity properties of ridges. Specifically, under mild conditions, we prove that the set-valued random process  $t \mapsto s_Y(t)$  is single-valued with probability one on any countable and dense subset of  $\mathbb{R}$ , and upper hemicontinuous on  $\mathbb{R}$  almost surely. This means that for any time t, as t' approaches t, the maxima of  $S_Y(t', \cdot)$  converge arbitrarily close to some of the maxima of  $S_Y(t, \cdot)$ . Based on these properties, we further discuss the local  $C^1$ -smoothness of  $s_Y(t)$ . Third, we analyze how  $s_Y(t)$  deviates from  $s_f(t)$  under the adaptive harmonic model. Since the scalogram is derived by composing the modulus and a family of convolution operators, explicitly characterizing the local maximum in  $S_Y(t, \cdot)$  for a fixed time t is challenging. This makes it difficult to analyze  $|s_Y(t) - s_f(t)|$ , particularly from the perspectives of time-frequency analysis and probability. This problem has received limited attention in the literature, with the exception of [36], where the authors considered a frequency-modulated signal contaminated with additive Gaussian white noise and provided approximations for the mean and variance of the difference between the peak locations of  $S_f(t, \cdot)$  and  $S_Y(t, \cdot)$ . We extend this result by conducting a systematic ridge analysis, which is based on establishing the maximal inequalities for the scalogram of  $\Phi$  and its derivatives. This result extends classical maximal inequalities, such as the Borell-TIS inequality and Dudley's theorem, to the complex-valued setting within the time-frequency analysis framework. This extension is of independent interest because the classical Gaussian concentration inequalities [20], Dudley's theorem, and the Borell-TIS inequality [39, 40] are formulated for real-valued Gaussian elements.

The rest of the paper is organized as follows. In Section 2, we summarize the necessary material for AWT and present the definitions of the adapted harmonic model and the noise we consider. Section 3 is dedicated to AWT for Gaussian random processes, where we present both old and new results. In Section 4, we introduce a definition of the ridge that takes into account practical needs when the data is noisy. In Section 5, we state our main results about the probability of the occurrence of ridge deviation. The proofs of our main results and some technical lemmas are provided in Section 6. The appendix includes a table of symbols frequently used throughout the manuscript for the reader's convenience.

#### 2. Preliminary

Consider the following *adaptive harmonic model (AHM)*. Take a real-valued signal f:

(1) 
$$f(t) = \sum_{m=1}^{M} A_m(t) \cos\left(2\pi\phi_m(t)\right), \ t \in \mathbb{R},$$

where  $A_m(t) \in C^1(\mathbb{R})$  are positive functions representing the amplitude modulation and  $\phi_m(t) \in C^2(\mathbb{R})$  are strictly monotonically increasing functions representing the phase so that  $\phi'_m(t) > 0$  represents the IF. We impose two conditions on the



FIGURE 1. The top section illustrates a frequency-modulated signal f along with its 3-dimensional and image representations in the time-scale domain. The bottom section presents the same for its noise-affected counterpart Y, with a signal-to-noise ratio of -6.72 dB. Here, Y is the sum of f and a sample path of the Gaussian process  $\Phi$  satisfying Assumption 1.

AHM model. The first one is the *slowly varying* condition:  $|A'_m(t)| \leq \epsilon \phi'_m(t)$  and  $|\phi''_m(t)| \leq \epsilon \phi'_m(t)$ , where  $\epsilon \geq 0$  is a small constant. The second one is the *spectral separation* condition:

(2) 
$$\phi'_1(t) < \phi'_2(t) < \dots < \phi'_M(t)$$

so that  $\phi'_{m+1}(t) - \phi'_m(t) \geq \delta(\phi'_{m+1}(t) + \phi'_m(t))$  for some  $\delta \in (0,1)$  and for all  $t \in \mathbb{R}$ . When M = 1, we denote  $A(t) = A_1(t)$  and  $\phi(t) = \phi_1(t)$ . We call  $A_m(t) \cos(2\pi\phi_m(t))$  the *m*-th *intrinsic mode type (IMT)* function. While it is possible to consider more complicated models, we focus on this model in this paper.

In the presence of noise, the noise-affected AHM may be expressed as

(3) 
$$Y(t) = f(t) + \Phi(t), \ t \in \mathbb{R},$$

where  $\Phi$  is a random process satisfying the following assumption.

**Assumption 1.** The noise  $\Phi$  is a mean-square continuous and stationary Gaussian process. Additionally, we assume that the covariance function of  $\Phi$  has a spectral density function  $p : \mathbb{R} \to (0, \infty)$  such that

$$\operatorname{Cov}(\Phi(t), \Phi(0)) = \int_{\mathbb{R}} e^{it\lambda} p(\lambda) d\lambda$$

Under Assumption 1, the process  $\Phi$  can be represented by

(4) 
$$\Phi(t) = \mathbb{E}[\Phi(0)] + \int_{\mathbb{R}} e^{it\lambda} \sqrt{p(\lambda)} Z(d\lambda),$$

where Z is an orthogonally scattered Gaussian random measure on  $\mathbb{R}$  satisfying

(5) 
$$Z(\Delta_1) = \overline{Z(-\Delta_1)}, \quad \mathbb{E}[Z(\Delta_1)] = 0, \text{ and } \quad \mathbb{E}\left[Z(\Delta_1)\overline{Z(\Delta_2)}\right] = \operatorname{Leb}(\Delta_1 \cap \Delta_2)$$

for any  $\Delta_1, \Delta_2 \in \mathcal{B}(\mathbb{R})$ , where Leb is the Lebesgue measure on  $\mathbb{R}$  and  $\mathcal{B}(\mathbb{R})$  is the Borel  $\sigma$ -algebra on  $\mathbb{R}$  [25].

The analytic wavelet transform (AWT) of an  $L^2$  function f, denoted by  $\{W_f(t,s) \mid t \in \mathbb{R}, s > 0\}$ , is defined as follows [8]:

(6) 
$$W_f(t,s) = \int_{\mathbb{R}} f(\tau) \frac{1}{s} \psi^* \left(\frac{\tau - t}{s}\right) d\tau$$

where  $t \in \mathbb{R}$  represents time, s > 0 represents scale, the asterisk represents the complex conjugate, and  $\psi \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$  is the chosen analytic mother wavelet. Recall that the mother wavelet  $\psi$  is called analytic if  $\hat{\psi}(\omega) = 0$  for  $\omega \leq 0$ , where  $\hat{\psi}$  is the Fourier transform of  $\psi$  defined by

$$\widehat{\psi}(\omega) = \int_{\mathbb{R}} e^{-it\omega} \psi(t) dt, \ \omega \in \mathbb{R}.$$

The squared magnitude of the complex-valued function  $W_f$ ,

(7) 
$$S_f(t,s) := |W_f(t,s)|^2,$$

is called the *scalogram*. See Figure 1 for an example.

Similar to (6) and (7), we denote the AWTs of  $\Phi$  and Y by  $W_{\Phi}$  and  $W_Y$ , respectively, and the scalograms of  $\Phi$  and Y by  $S_{\Phi}$  and  $S_Y$ , respectively. By the linearity of AWT, we know  $W_Y = W_f + W_{\Phi}$ . By (4) and the stochastic Fubini Theorem [31],

(8) 
$$W_{\Phi}(t,s) = \int_0^\infty e^{it\lambda} \overline{\widehat{\psi}(s\lambda)} \sqrt{p(\lambda)} Z(d\lambda).$$

The AWT of both  $\Phi$  and Y are complex-valued and nonstationary Gaussian random processes on the scale axis for each  $t \in \mathbb{R}$ .

#### 3. AWT FOR GAUSSIAN RANDOM PROCESS-SOME OLD AND NEW RESULTS

The first result focuses on the covariance structure of  $W_{\Phi}$ , which serves as a foundational step in analyzing the ridges of the TFR determined by AWT.

**Theorem 1.** Suppose that  $\Phi$  is a stationary Gaussian process satisfying Assumption 1.

(a) For any  $t \in \mathbb{R}$ ,  $d \in \mathbb{N}$ , and  $s_1, s_2, \ldots, s_d > 0$ , the complex Gaussian random vector

$$\mathbf{W} := [W_{\Phi}(t,s_1) \ W_{\Phi}(t,s_2) \ \cdots \ W_{\Phi}(t,s_d)]^{\top}$$

is circularly symmetric, where  $\top$  represents the transpose. That is,  $e^{i\theta}\mathbf{W}$  has the same probability distribution as  $\mathbf{W}$  for any  $\theta \in \mathbb{R}$ . (b) For any  $t \in \mathbb{R}$ ,  $s_1, s_2 > 0$  with  $s_1 \neq s_2$ ,

 $\frac{|W_{\Phi}(t,s_1) - W_{\Phi}(t,s_2)|^2}{d_{W_{\Phi}}^2(s_1,s_2)}$ 

follows an exponential distribution with a mean of one, where

(9) 
$$d_{W_{\Phi}}(s_1, s_2) = \left\{ \mathbb{E} \left[ |W_{\Phi}(0, s_1) - W_{\Phi}(0, s_2)|^2 \right] \right\}^{1/2}.$$

(c) For any  $t \in \mathbb{R}$  and  $s_1, s_2 > 0$ ,

$$\operatorname{Cov}\left(S_{\Phi}(t,s_{1}), S_{\Phi}(t,s_{2})\right) = 4\left\{\mathbb{E}\left[W_{\Phi}^{R}(t,s_{1})W_{\Phi}^{R}(t,s_{2})\right]\right\}^{2} + 4\left\{\mathbb{E}\left[W_{\Phi}^{R}(t,s_{1})W_{\Phi}^{I}(t,s_{2})\right]\right\}^{2},$$

where  $W_{\Phi}^{R}$  and  $W_{\Phi}^{I}$  are the real and imaginary parts of  $W_{\Phi}$ , respectively. Especially, for any s > 0,

Var 
$$(S_{\Phi}(t,s)) = 4 \left\{ \mathbb{E} \left[ |W_{\Phi}^{R}(t,s)|^{2} \right] \right\}^{2} = \left\{ \mathbb{E} \left[ S_{\Phi}(t,s) \right] \right\}^{2}$$

It is worth noting that when the noise is modeled as a stationary Gaussian process, the scalogram  $S_{\Phi}$  at a given time is a nonstationary random process along the scale axis. Theorem 1(c) shows that for any  $t \in \mathbb{R}$ , the nonstationary process  $S_{\Phi}(t, \cdot)$  is non-negatively correlated. To our knowledge, this property is not found in existing literature, so we provide its proof in Section 6.1.

The goal of this work is to analyze the location of the maximum of  $|W_Y(t, \cdot)|^2$ along the scale axis and to examine its deviation from the location of the maximum of  $|W_f(t, \cdot)|^2$  along the same axis. For any fixed  $t \in \mathbb{R}$ , finding the maximizers of  $|W_Y(t, \cdot)|^2$  requires computing the partial derivatives of both  $W_f$  and  $W_{\Phi}$  with respect to the scale variable since  $W_Y = W_f + W_{\Phi}$ . Furthermore, it is essential to ensure the finiteness of the expectation of the maximum of  $|W_{\Phi}|$  and  $|\partial W_{\Phi}/\partial s|$ (see Theorem 4 and Lemma 4 below), so we make the following assumption.

**Assumption 2.** The Fourier transform  $\widehat{\psi}$  of the analytic mother wavelet  $\psi$  is three times differentiable on  $(0, \infty)$ , and the following conditions hold:

$$\begin{split} (\mathbf{D}_{1}^{0}-\mathbf{D}_{2}^{0}) &: \sup_{\lambda > 1} |\lambda^{p} \widehat{\psi}(\lambda)| < \infty, \ p = 1, 2; \\ (\mathbf{D}_{0}^{1}-\mathbf{D}_{2}^{1}) &: \sup_{\lambda > 0} |\lambda^{p} D \widehat{\psi}(\lambda)| < \infty, \ p = 0, 1, 2; \\ (\mathbf{D}_{2}^{2}-\mathbf{D}_{3}^{2}) &: \sup_{\lambda > 0} |\lambda^{p} D^{2} \widehat{\psi}(\lambda)| < \infty, \ p = 2, 3; \\ (\mathbf{D}_{3}^{3}) &: \sup_{\lambda > 0} |\lambda^{3} D^{3} \widehat{\psi}(\lambda)| < \infty, \end{split}$$

where  $D^k$  represents the kth derivative operator. Without loss of generality, we also assume that  $|\hat{\psi}|$  is a unimodal function.

**Example 1.** The set of wavelets that satisfies Assumption 2 includes some of the generalized Morse wavelets discussed in [23, (14)] and the analytic wavelets derived in [17, (6)]. For instance, the generalized Morse wavelets, whose Fourier transform takes the form

$$\widehat{\psi}(\lambda) = a_{\beta_1,\beta_2} \lambda^{\beta_1} e^{-\lambda^{\beta_2}} \mathbf{1}_{[0,\infty)}(\lambda), \ \beta_1 \ge 1, \ \beta_2 > 0,$$

where  $a_{\beta_1,\beta_2}$  is a normalizing constant. Another example is the Klauder wavelet, whose Fourier transform is given by

$$\widehat{\psi}(\lambda) = \lambda^{\alpha} e^{-\gamma\lambda} e^{i\beta \log \lambda} \mathbb{1}_{[0,\infty)}(\lambda),$$

where  $\alpha \geq 1$ ,  $\beta \in \mathbb{R}$ , and  $\gamma \in \mathbb{C}$  with  $\operatorname{Re}(\gamma) > 0$ .

**Proposition 1.** Under Assumptions 1 and 2, the sample paths of  $W_{\Phi}(\cdot, \cdot)$  are two times continuously differentiable on  $\mathbb{R} \times (0, \infty)$  almost surely.

The proof of Proposition 1 is provided in Section 6.2. It shows that if the mother wavelet  $\psi$  has higher regularity, the sample paths of  $W_Y$  will also exhibit higherorder smoothness. A surface plot of  $|W_Y|$  is shown in the bottom portion of Figure 1, where the signal-to-noise ratio is -6.72 dB.

**Assumption 3.** For the behavior of the spectral density function p of the noise  $\Phi$  at infinity, we assume that (a) there exist constants  $\gamma \in (0,2)$  and  $C_1 > 0$  such that

$$p(\lambda) \leq C_1 |\lambda|^{-(1+\gamma)}$$
 for any  $\lambda \neq 0$ .

For the behavior of p near the origin, we consider two scenarios.

(b1) Long-range dependence scenario:

(10) 
$$p(\lambda) = L(|\lambda|^{-1})|\lambda|^{H-1},$$

where H is a long-memory parameter belonging to (0,1) and  $L: (0,\infty) \rightarrow (0,\infty)$  is a locally bounded function which is slowly varying at infinity, meaning  $L(cr)/L(r) \rightarrow 1$  as  $r \rightarrow \infty$  for any c > 0.

(b2) Short-range dependence scenario: p is a bounded function continuous at the origin.

**Example 2.** Let us consider the stationary random process  $\Phi$  with the generalized Linnik covariance function [24], defined as

$$C_{\Phi}(t) = (1 + |t|^{\gamma})^{-H/\gamma}, \ \gamma \in (0, 2], \ H > 0.$$

According to [24, Propositions 3.1 and 3.4],  $p(\lambda) = \mathcal{O}(|\lambda|^{-(1+\gamma)})$  when  $|\lambda| \to \infty$ . For the case H > 1, because  $C_{\Phi} \in L^1(\mathbb{R})$ , the spectral density function p is continuous everywhere. For the case 0 < H < 1, by [24, Corollaries 3.3 and 3.10],

$$p(\lambda) \sim |\lambda|^{H-1} \Gamma(\frac{1-H}{2}) \left[ 2^H \pi^{\frac{1}{2}} \Gamma\left(\frac{H}{2}\right) \right]^{-1}$$

as  $|\lambda| \to 0$ .

**Corollary 1.** Suppose that Assumption 1 and conditions  $(D_1^0)$ ,  $(D_0^1)$ , and  $(D_1^1)$  in Assumption 2 hold.

(a) If Assumption 3(a) is satisfied, then

(11) 
$$\mathbb{P}\left(\lim_{s\to 0+} W_{\Phi}(t,s) = 0 \text{ for all } t \in \mathbb{R}\right) = 1.$$

(b) If either Assumption 3(b1) or Assumption 3(b2) is satisfied, then

(12) 
$$\mathbb{P}\left(\lim_{s \to \infty} W_{\Phi}(t,s) = 0 \text{ for all } t \in \mathbb{R}\right) = 1.$$

We note that the definition of the AWT does not include the scale s = 0. Corollary 1 shows that, under Assumption 3, the continuity of  $W_{\Phi}$  can be extended to the boundary s = 0 by defining  $W_{\Phi}(\cdot, 0) = 0$ . The proof of Corollary 1 is provided in Section 6.3.

Finally, we conclude this section by presenting the inverse AWT and the holomorphic properties of the AWT of Gaussian processes. While these properties are not utilized in the ridge analysis, they hold independent interest. The results are derived by adapting findings from [8] and [17, Theorem 2], which were originally established for the deterministic case.

**Proposition 2.** Suppose that  $\Phi$  is a stationary Gaussian process satisfying Assumption 1.

(a) If the analytic mother wavelet  $\psi$  and the spectral density function p of  $\Phi$  satisfy

(13) 
$$\int_0^\infty \left[\int_0^\infty \left|\widehat{\psi}(s\lambda)\sqrt{p(\lambda)}s^{-1}\right|^2 d\lambda\right]^{1/2} ds < \infty,$$

then, almost surely, we have the inverse AWT:

$$\Phi(t) = \operatorname{Re}\left[C_{\psi}^{-1} \int_{0}^{\infty} W_{\Phi}(t,s)s^{-1}ds\right], \ t \in \mathbb{R},$$

where  $C_{\psi} = 2^{-1} \int_0^{\infty} \overline{\widehat{\psi}(\lambda)} \lambda^{-1} d\lambda$ . (b) If the Fourier transform of the mother wavelet  $\psi$  has the form

(14) 
$$\widehat{\psi}(\lambda) = \lambda^{\alpha} e^{-\gamma \lambda} e^{i\beta \log \lambda} \mathbb{1}_{[0,\infty)}(\lambda)$$

where  $\alpha \geq 1$ ,  $\beta \in \mathbb{R}$ , and  $\gamma \in \mathbb{C}$  with  $\operatorname{Re}(\gamma) > 0$ , then, almost surely, the function  $h_{\Phi}: \mathbb{C} \mapsto \mathbb{C}, defined by$ 

$$h_{\Phi} : \{ z \in \mathbb{C} : \operatorname{Im}(z) > 0 \} \mapsto \mathbb{C}$$
$$t + is \mapsto g(s) W_{\Phi} \left( t - \frac{\operatorname{Im}(\gamma)}{\operatorname{Re}(\gamma)} s, \frac{s}{\operatorname{Re}(\gamma)} \right),$$

is holomorphic, where

$$g(s) = s^{-\alpha} e^{i\beta \log s}.$$

It is important to note that condition (13) is necessary for applying the stochastic Fubini theorem [31]. This condition holds for wavelets in Example 1 and covariance models in Example 2. In the second part of Proposition 2, we consider a specific type of wavelets, which is not essential for the main results of this paper. However, the holomorphic property derived from this class may simplify the proofs of some results below. The proof of Proposition 2 is provided in Section 6.4.

# 4. Definition of ridges

We start the definition of ridges by recalling existing results when the input signal is noise-free. With AHM and the slowly varying condition specified in (1), we have the following approximation for the AWT of f fulfilling the AHM [9, Estimate 3.5]:

$$|W_{f}(t,s)| = \left| \int_{\mathbb{R}} \frac{1}{s} \psi^{*} \left( \frac{\tau - t}{s} \right) \sum_{m=1}^{M} A_{m}(\tau) \cos \left( 2\pi \phi_{m}(\tau) \right) d\tau \right|$$
$$= \left| \sum_{m=1}^{M} A_{m}(t) \int_{\mathbb{R}} \frac{1}{s} \psi^{*} \left( \frac{\tau - t}{s} \right) \cos \left( 2\pi (\phi_{m}(t) + \phi'_{m}(t)(\tau - t)) \right) d\tau + \epsilon E(t,s) \right|$$
$$(15) = \left| \frac{1}{2} \sum_{m=1}^{M} A_{m}(t) e^{i2\pi \phi_{m}(t)} \widehat{\psi} \left( 2\pi s \phi'_{m}(t) \right) + \epsilon E(t,s) \right|,$$

where

$$0 \le E(t,s) \le s \left[ \mathcal{I}_1 \sum_{m=1}^M |\phi'_m(t)| + \frac{1}{2} \mathcal{I}_2 s \sum_{m=1}^M [M''_m + |A_m(t)||\phi'_m(t)|] + \frac{1}{6} \mathcal{I}_3 s^2 \sum_{m=1}^M M''_k |A_m(t)| \right] + \frac{1}{6} \mathcal{I}_3 s^2 \sum_{m=1}^M M''_k |A_m(t)| = \frac{1}{$$

and  $M_m'' := \sup_{t\in\mathbb{R}} |\phi_m''(t)| < \infty$ . In other words, when  $\epsilon > 0$  is sufficiently small, we have

(16) 
$$|W_f(t,s)| \approx \frac{1}{2} \left| \sum_{m=1}^M A_m(t) e^{i2\pi\phi_m(t)} \widehat{\psi}(2\pi s \phi'_m(t)) \right|.$$

To analyze the ridges in the scalogram of f, we make the following observations. First, if the amplitude of the *m*-th IMT function in f, where  $m \in \{1, \ldots, M\}$ , is very small, its contribution to the scalogram may be buried by spectral leakage caused by the time-varying nature of the frequencies and amplitudes. This can make detecting the *m*-th IF,  $\phi'_m$ , from the scalogram challenging. Conversely, if one component has a very large amplitude, its spectral leakage may dominate the scalogram, potentially masking the contributions of other components. Second, if the spectral energy of the mother wavelet  $\psi$  is spread over a broad frequency range rather than being tightly concentrated, the components in (16) may overlap significantly, making it harder to isolate individual components or accurately infer the IFs. To build intuition about ridge definitions for noise-free AMH, we focus on the scenario where the impact of spectral leakage in the scalogram of the noise-free AHM is negligible. This holds, for example, when the support of  $\psi$  is concentrated around [1-B, 1+B], where  $0 < B < \delta/(1+\delta)$  and  $\delta$  is the constant mentioned in the spectral separation condition (2). Under this assumption, (16) can be further approximated as follows:

(17) 
$$|W_f(t,s)| \approx \frac{1}{2} \sum_{m=1}^{M} A_m(t) \left| \hat{\psi} \left( 2\pi s \phi'_m(t) \right) \right|.$$

Denote

$$\omega_{\psi} = \frac{1}{2\pi} \arg \max_{\omega > 0} |\widehat{\psi}(\omega)|,$$

which is a singleton under Assumption 2. By (17), for any  $t \in \mathbb{R}$ , the local maxima of  $|W_f(t, \cdot)|$  approximately occur at

$$\frac{\omega_{\psi}}{\phi_1'(t)}, \frac{\omega_{\psi}}{\phi_2'(t)}, \dots, \frac{\omega_{\psi}}{\phi_M'(t)},$$

which are inversely related to the IFs  $\{\phi'_1(t), \phi'_2(t), \ldots, \phi'_M(t)\}$ . The observation above leads to the following definition of ridges [23, Definitions 2.1 and 2.2] (see also [27, Section B]).

**Definition 1.** Suppose f satisfies AHM (1). A ridge point of  $S_f(t, s)$  is a time/scale pair (t, s) satisfying the two conditions:

$$(R1) \ \frac{\partial}{\partial s} S_f(t,s) = 0 \quad and \quad (R2) \ \frac{\partial^2}{\partial s^2} S_f(t,s) < 0;$$

that is  $S_f(t,s)$  is a local maximum along the scale axis. A ridge curve in  $S_f(t,s)$  is the graph of a function of t such that the conditions (R1) and (R2) are satisfied along the curve, and (R3) the function is continuously differentiable.

Note that for a signal fulfilling the AHM, there might be multiple ridge curves, and a ridge curve is a set of ridge points fulfilling the regularity condition.

To utilize this definition, consider

(18) 
$$s_f(t) = \arg\max_{s>0} S_f(t,s).$$

When f satisfies the AHM (1) with M = 1,  $s_f$  is a single-valued function. By (17),

$$s_f(t) \approx \arg \max_{s>0} \left| \widehat{\psi} \left( 2\pi s \phi'(t) \right) \right| = \frac{\omega_{\psi}}{\phi'(t)};$$

that is, the ridge captures the IF information. See Figure 1 for an example of the graph of  $s_f$  and its relationship with the IF  $\phi'(t)$ .

When f satisfies the AHM (1) with  $M \ge 2$ , (18) might be problematic. First, due to the time-varying amplitude,  $s_f(t)$  might alternately capture the IF of different IMT functions. For example, consider the case where M = 2,  $\phi'_m(t) = \xi_m > 0$ are constants, and  $\xi_2 > 2\xi_1$ . If  $A_1(t) > 2A_2(t)$  for t < -1 and  $A_2(t) > 2A_1(t)$  for t > 1, then  $s_f(t) \approx \omega_{\psi}/\phi'_1(t)$  when t < -1 and  $s_f(t) \approx \omega_{\psi}/\phi'_2(t)$  when t > 1. On the other hand, note that when the spectral leakage from other IMT functions of f is smaller than  $A_m(t)|\widehat{\psi}(2\pi s \phi'_m(t))|$ ,  $s_{f,m}(t)$  is close to  $\phi'_m(t)$  and we have M local maxima. In this case, to utilize the definition of ridges, consider

(19) 
$$\frac{\omega_{\psi}}{\phi'_m(t)} \in B_m(t) := \left[\underline{b}_m(t), \overline{b}_m(t)\right] \text{ for } 1 \le m \le M,$$

where  $\underline{b}_1, \overline{b}_1, \underline{b}_2, \overline{b}_2, \dots, \underline{b}_M, \overline{b}_M$  are continuous functions defined on  $\mathbb{R}$  satisfying

$$0 < \underline{b}_M(t) < \overline{b}_M(t) \le \dots \le \underline{b}_2(t) < \overline{b}_2(t) \le \underline{b}_1(t) < \overline{b}_1(t) < \infty.$$

Then, define

$$s_{f,m}(t) = \arg \max_{s \in B_m(t)} S_f(t,s).$$

The functions  $s_{f,m}(t)$  become the ridge curves by definition if they are continuously differentiable. For the spectrogram of a function fulfilling the AHM, there are analogous definition of ridge curves and their differentiability has been established [7]. A similar proof can be carried out to the scalogram and we skip the details here. Hereafter, we use the graphs of  $s_f$  and  $\{s_{f,m}\}_{m=1,\ldots,M}$  to represent the ridge curves of the noise-free function satisfying the AHM and call them the *ridge curves* to simplify the nomination. **Remark 1.** It is worth noting that, in practice, information about the IFs of f, and hence  $B_m(t)$ , may sometimes be available through prior knowledge. However, more often than not, this information is unavailable and must be estimated. For an example of an algorithm addressing this estimation challenge, see [21]. Nonetheless, this estimation problem lies beyond the scope of this paper. From a broader signal processing perspective, once this definition is established, the next step is to design an appropriate algorithm to extract the ridge curves  $\{s_{f,m}\}_{m=1,...,M}$  from the scalogram. See [10] and [5, 4, 30, 19, 28, 41, 6, 22, 21, 38] for various proposed algorithms in the literature. However, the design and analysis of such algorithms lie beyond the scope of this study.

4.1. **Definition of ridges for noise-affected AHM.** We proceed to the main focus of this paper: the ridges when the signal is noisy. Motivated by Definition 1 and real data experience, we consider the following definition.

**Definition 2.** Denote  $S_Y(t,s) = |W_Y(t,s)|^2$ . For the noise-affected AHM (3) with  $M \ge 1$ , and analogous to (18), we define

(20) 
$$s_Y(t) = \arg\max_{s>0} S_Y(t,s)$$

for any  $t \in \mathbb{R}$ , and refer the graph of  $s_Y$  as a ridge curve of Y, noting that  $s_Y$  may be a multivalued function at some time points.

For the noise-affected AHM (3) with  $M \ge 1$ , we also define

(21) 
$$s_{Y,m}(t) = \arg \max_{s \in B_m(t)} S_Y(t,s)$$

for any  $t \in \mathbb{R}$  and  $m \in \{1, 2, ..., M\}$ , and refer to the graph of  $s_{Y,m}$  as the mth ridge curve of Y.

Note that compared with the ridge curve defined in Definition 1 when the signal is noise-free, in the above definition we do not specify the regularity of the ridge curve. Instead, we need to explore its regularity. To this end, observe that unlike  $|W_f(t, \cdot)|$  shown in Figure 1,  $|W_Y(t, \cdot)|$  has multiple local maxima even in the case where M = 1, as illustrated in Figure 2. We further show the scatter plot of the local maximizers of  $|W_Y(t, \cdot)|$  for any  $t \in \mathbb{R}$  in Figure 4. We find many paths formed by the local maximizers of  $|W_Y(t, \cdot)|$  for each  $t \in \mathbb{R}$ , and they are piecewise continuous. Moreover, we find that near the "true" IF, the associated local maximum that we expect to be the ridge occasionally jumps; for example, around the 5th, 10th, and 23rd seconds in the top panel of Figure 4. This motivates us to analyze the graphs of  $s_Y$  and  $\{s_{Y,m}\}_{m=1}^M$ , as well as the difference between  $s_f$  (and  $\{s_{f,m}\}_{m=1}^M$ ) and  $s_Y$  (and  $\{s_{Y,m}\}_{m=1}^M$ ).

#### 5. Main Results

In this section, we present our main results. The first part is about the singleton property of the set-valued functions  $s_Y$  (or  $s_{Y,1}, \ldots, s_{Y,M}$  when M > 1), as well as the hemicontinuity and local  $C^1$ -smoothness properties of their graphs. The second part is the deviation of  $s_Y$  (or  $s_{Y,1}, \ldots, s_{Y,M}$  when M > 1) from  $s_f$  (or  $s_{f,1}, \ldots, s_{f,M}$  when M > 1).



FIGURE 2. Example of the complex modulus of the AWT for a 30second noise-contaminated, frequency-modulated signal. The first row displays the frequency-modulated signal f, while the second row shows the noise, a sample from the stationary Gaussian process  $\Phi$ , identical to that in Figure 1. The third row illustrates the complex modulus of the AWT, i.e., the spectral amplitude, of the noise-contaminated signal Y, where  $Y = f + \Phi$ . In the last three columns, the black curves represent the spectral amplitude of Y at specific times, corresponding to cross-sections of the image in the third row. For comparison, the spectral amplitudes of f are shown in gray.

## 5.1. Local properties of ridge curves.

**Theorem 2.** Suppose  $f \in L^2$ ,  $\Phi$  a stationary Gaussian process satisfying Assumptions 1 and 3, and  $\psi$  a mother wavelet satisfying Assumption 2. Then, we have

 $\mathbb{P}(s_Y(t) \text{ is a singleton at any } t \in \mathbb{Q}) = 1,$ 

where  $\mathbb{Q}$  denotes the set of rational numbers. Moreover, when f satisfies the AHM defined in (1) with  $M \geq 1$ , we have

 $\mathbb{P}(s_{Y,m}(t) \text{ is a singleton for any } t \in \mathbb{Q} \text{ and } m \in \{1, \ldots, M\}) = 1.$ 

We note that Theorem 2 holds regardless of whether  $f \equiv 0$  or not, and in general, the singleton property of  $s_Y$  does not require the AHM assumption. The proof of Theorem 2 is provided in Section 6.5. In addition to investigating the singleton property of the sets  $s_Y(t)$ , or  $s_{Y,1}(t), \ldots, s_{Y,M}(t)$  for f satisfying the AHM, at  $\mathbb{Q}$ , we are also interested in the behavior of these sets for the remaining  $t \in \mathbb{R} \setminus \mathbb{Q}$ .

**Theorem 3.** Suppose  $f \in L^2$ ,  $\Phi$  a stationary Gaussian process satisfying Assumptions 1 and 3, and  $\psi$  a mother wavelet satisfying Assumption 2. With probability one, the set-valued function  $s_Y$  is upper hemicontinuous on  $\mathbb{R}$ . That is, almost surely, for any  $t \in \mathbb{R}$  and any open set U with  $s_Y(t) \subset U$ , there exists  $\delta > 0$  such that  $s_Y(t') \subset U$  for any  $t' \in (t - \delta, t + \delta)$ . Moreover, if f is an adapted harmonic signal as defined in (1) with  $M \geq 1$ , then almost surely, the set-valued functions  $s_{Y,1}, \ldots, s_{Y,M}$ , defined in (21), are also upper hemicontinuous on  $\mathbb{R}$ .

The upper hemicontinuity of  $s_Y$  implies that as t' approaches t, the maximizers of  $S_Y(t', \cdot)$  get arbitrarily close to some of the maximizers of  $S_Y(t, \cdot)$ . The proof of Theorem 3 is provided in Section 6.6, which relies on Berge's maximum theorem [18, Theorem 3.4] (see also [1, p. 570] and Lemma 5 in the appendix).

Theorems 2 and 3 imply that, with probability one, the restriction of  $s_Y$  on the dense set  $\mathbb{Q}$ , denoted by  $s_Y|_{\mathbb{Q}}$ , is continuous. By [14, Lemma (4.3.16)] (see also Lemma 6 in the appendix),  $s_Y|_{\mathbb{Q}}$  is extendable to a continuous function defined on a  $G_{\delta}$ -set containing  $\mathbb{Q}$ . Here, a  $G_{\delta}$ -set refers to a subset of  $\mathbb{R}$  that is a countable intersection of open sets.

Next, we provide more insights into this  $G_{\delta}$ -set.

**Corollary 2.** Assume that Assumptions 1-3 hold. For any  $t_0 \in \mathbb{Q}$ , if

(22) 
$$\mathbb{P}\left(\frac{\partial^2 S_Y}{\partial s^2}\left(t_0, s_Y(t_0)\right) = 0\right) = 0,$$

then, almost surely, there exists a unique continuously differentiable function r defined on  $(t_0 - \delta, t_0 + \delta)$  for some  $\delta > 0$  such that

- r(t) = s<sub>Y</sub>(t) for any t ∈ (t<sub>0</sub> − δ, t<sub>0</sub> + δ) ∩ Q, and
  r(t) ∈ s<sub>Y</sub>(t) for any t ∈ (t<sub>0</sub> − δ, t<sub>0</sub> + δ) ∩ Q<sup>c</sup>.

The proof of Corollary 2 is provided in Section 6.7. Corollary 2 provides additional insight into the nature of the  $G_{\delta}$ -set, on which  $s_Y$  is continuous, under the assumption (22). This assumption is supported by the histogram of  $\frac{\partial^2 S_Y}{\partial s^2}(t, s_Y(t))$ shown in Figure 3. Corollary 2 also explains why the ridge curves observed in the numerical experiments consist of piecewise continuous curves. See Figure 4 for an illustration. Note that while it is visually possible to identify if the ridge points form a continuous path, the local maxima plot further highlights and emphasizes any discontinuities.

Furthermore, if the signal f fulfills the AHM, we have the following property of the paths  $\{(t, s_{Y,m}(t)) \mid t \in \mathbb{R}\}_{m=1,\dots,M}$ , which is parallel to Corollary 2.



FIGURE 3. Histogram of the random variable  $\frac{\partial^2 S_Y}{\partial s^2}(t, s_Y(t))$  for the special case  $Y(t) = f(t) + \Phi(t)$ , where  $\Phi$  is a stationary Gaussian process. The time variable is fixed throughout the experiment, and the histogram is based on  $10^5$  realizations of  $\Phi$ .



FIGURE 4. Example of ridge points (local maxima) and a ridge curve  $s_Y$ , where  $Y = f + \Phi$ , and  $s_f$ . The signal f and the realization of  $\Phi$  are identical to those in Figure 1. The zoomed-in blue box in the bottom-right region highlights the discontinuity of the ridge in the presence of noise.

**Corollary 3.** Assume that Assumptions 1-3 hold. For any  $t_0 \in \mathbb{Q}$  and  $m \in \{1, \ldots, M\}$ , if  $s_{Y,m}(t_0)$  lies in the interior of  $B_m(t_0)$  and

$$\mathbb{P}\left(\frac{\partial^2 S_Y}{\partial s^2}\left(t_0, s_{Y,m}(t_0)\right) = 0\right) = 0,$$

then, almost surely, there exists a unique continuously differentiable function  $r_m$  defined on  $(t_0 - \delta, t_0 + \delta)$  for some  $\delta > 0$  such that

- $r_m(t) \in B_m(t)$  for any  $t \in (t_0 \delta, t_0 + \delta)$ ,
- $r_m(t) = s_{Y,m}(t)$  for any  $t \in (t_0 \delta, t_0 + \delta) \cap \mathbb{Q}$ , and
- $r_m(t) \in s_{Y,m}(t)$  for any  $t \in (t_0 \delta, t_0 + \delta) \cap \mathbb{Q}^c$ .

Based on our literature review, the analysis of the uniqueness of maxima for a random field determined by AWT along the scale axis, as well as the behavior of the locations of these maxima along another axis, has rarely been discussed.

5.2. Deviation of ridge points caused by noise. We continue to quantify the probability that  $|s_Y(t) - s_f(t)|$  and  $|s_{Y,m}(t) - s_{f,m}(t)|$  exceed a given threshold, where  $m = 1, \ldots, M$ .

**Theorem 4.** Suppose that the stationary Gaussian process  $\Phi$  satisfies Assumptions 1 and 3, and the mother wavelet  $\psi$  satisfies Assumption 2. Fix  $t \in \mathbb{R}$ . For any interval I containing  $s_f(t)$ , denote

(23) 
$$\Delta_I = |W_f(t, s_f(t))| - \max_{s \in I^c} |W_f(t, s)|,$$
$$\mu_I = \mathbb{E}\left[|W_{\Phi}(t, s_f(t))|\right] + \mathbb{E}\left[\max_{s \in I^c} |W_{\Phi}(t, s)|\right],$$

and

(24) 
$$\sigma_I = \sqrt{\mathbb{E}\left[|W_{\Phi}(t, s_f(t))|^2\right]} + \max_{s \in I^c} \sqrt{\mathbb{E}\left[|W_{\Phi}(t, s)|^2\right]}.$$

If  $\triangle_I > \mu_I$ , then

(25) 
$$\mathbb{P}\left(s_Y(t) \in I\right) \ge 1 - \exp\left(-\frac{(\triangle_I - \mu_I)^2}{\sigma_I^2}\right)$$

where  $s_Y$  is defined in (20).

Theorem 4 shows that if the peak of  $|W_f(t, \cdot)|$  is distinct (i.e.  $\Delta_I$  is "large", or  $\epsilon E(t, \cdot)$  in (15) is sufficiently small) and much higher than the peaks of  $\{|W_{\Phi}(t,s)| | s \in I^c\}$  (i.e.  $\Delta_I > \mu_I$ ), then the deviation of the ridge point  $s_f(t)$  caused by noise is controlled with high probability. It is worth noting that deviations in the ridge point of f may also occur if the spectral component of f at the scale  $s_f(t)$  is dominated by the spectral component of the noise at the same scale. Hence, the moments of the wavelet coefficients of the noise  $\Phi$  at the scale  $s_f(t)$ , including  $\mathbb{E}[|W_{\Phi}(t, s_f(t))|^2]$ , are contained in  $\mu_I$  and  $\sigma_I$ . Most components in (23) and (24) can be calculated as follows:

$$\mathbb{E}\left[|W_{\Phi}(t,s_f(t))|\right] = \frac{1}{2}\pi^{\frac{1}{2}} \left[\int_0^\infty |\widehat{\psi}\left(s_f(t)\lambda\right)|^2 p(\lambda) d\lambda\right]^{\frac{1}{2}}$$

and

$$\sigma_{I} = \sqrt{\int_{0}^{\infty} |\widehat{\psi}(s_{f}(t)\lambda)|^{2} p(\lambda) d\lambda} + \max_{s \in I^{c}} \sqrt{\int_{0}^{\infty} |\widehat{\psi}(s\lambda)|^{2} p(\lambda) d\lambda}.$$

The proof of Theorem 4 is presented in Section 6.8. To prove Theorem 4 and derive an upper bound for the second component

$$\mathbb{E}\left[\max_{s\in I^c} |W_{\Phi}(t,s)|\right]$$

in (23), we need to extend the classical Gaussian concentration inequality for Lipschitz functions of real-valued Gaussian elements [20], Dudley's theorem, and the Borell-TIS inequality [39, 40], to account for complex-valued Gaussian elements arising from the AWT of  $\Phi$ . To keep the focus on the discussion of ridge deviation, these extensions are provided in the next subsection.

**Corollary 4.** Suppose that f satisfies the AHM, the stationary Gaussian process  $\Phi$  satisfies Assumptions 1 and 3, and the mother wavelet  $\psi$  satisfies Assumption 2. Fix  $t \in \mathbb{R}$ . For any  $m \in \{1, \ldots, M\}$  and any interval  $I_m \subset B_m(t)$  containing  $s_{f,m}(t)$ , denote

(26)  

$$\Delta_{I_m} = |W_f(t, s_{f,m}(t))| - \max_{s \in B_m(t) \setminus I_m} |W_f(t, s)|,$$

$$\mu_{I_m} = \mathbb{E}\left[|W_{\Phi}(t, s_{f,m}(t))|\right] + \mathbb{E}\left[\max_{s \in B_m(t) \setminus I_m} |W_{\Phi}(t, s)|\right],$$

and

$$\sigma_{I_m} = \sqrt{\mathbb{E}\left[|W_{\Phi}(t, s_{f,m}(t))|^2\right]} + \max_{s \in B_m(t) \setminus I_m} \sqrt{\mathbb{E}\left[|W_{\Phi}(t, s)|^2\right]}.$$



FIGURE 5. Illustration of the relationship between the scale region  $B_m(t)$  and the interval  $[\underline{i}_m(t), \overline{i}_m(t)]$  considered in Theorem 5.

If  $\triangle_{I_m} > \mu_{I_m}$ , then

$$\mathbb{P}\left(s_{Y,m}(t) \in I_m\right) \ge 1 - \exp\left(-\frac{(\triangle_{I_m} - \mu_{I_m})^2}{\sigma_{I_m}^2}\right)$$

where  $s_{Y,m}$  is defined in (21).

For every  $m \in \{1, 2, ..., M\}$  and  $t \in \mathbb{R}$ , let  $\underline{i}_m(t)$  and  $\overline{i}_m(t)$  be the inflection points of  $S_f(t, \cdot)$  closest to  $s_{f,m}(t)$  with  $\underline{i}_m(t) < s_{f,m}(t) < \overline{i}_m(t)$ , as depicted in Figure 5. As shown in Figures 2 and 4, local maxima of  $S_Y(t, \cdot)$  are likely to occur within neighborhoods of the local maxima of  $S_f(t, \cdot)$ . Hence, for the function  $s_{Y,m}$  defined in (21) with  $B_m(t) \subset (\underline{i}_m(t), \overline{i}_m(t))$ , we are also interested in the probability that  $|s_{Y,m}(t) - s_{f,m}(t)|$  exceeds a threshold conditioned on the event  $s_{Y,m}(t) \in B_m^\circ(t)$ , where  $B_m^\circ(t)$  is the interior of  $B_m(t)$ .

Before presenting our final main result, we introduce the following notation: For every  $m \in \{1, 2, ..., M\}$  and  $t \in \mathbb{R}$ , let

$$\mu_{1,m} = \mathbb{E}\left[\max_{s \in B_m(t)} \left| \frac{\partial W_f}{\partial s}(t,s) W_{\Phi}(t,s) \right| \right], \qquad \sigma_{1,m}^2 = \max_{s \in B_m(t)} \left| \frac{\partial W_f}{\partial s}(t,s) \right|^2 \mathbb{E}\left[ |W_{\Phi}(t,s)|^2 \right],$$
$$\mu_{2,m} = \mathbb{E}\left[\max_{s \in B_m(t)} \left| W_f(t,s) \frac{\partial W_{\Phi}}{\partial s}(t,s) \right| \right], \qquad \sigma_{2,m}^2 = \max_{s \in B_m(t)} |W_f(t,s)|^2 \mathbb{E}\left[ \left| \frac{\partial W_{\Phi}}{\partial s}(t,s) \right|^2 \right],$$
(27)

$$\mu_{3,m} = \mathbb{E}\left[\max_{s\in B_m(t)} |W_{\Phi}(t,s)|\right], \qquad \sigma_{3,m}^2 = \max_{s\in B_m(t)} \mathbb{E}\left[|W_{\Phi}(t,s)|^2\right],$$
$$\mu_{4,m} = \mathbb{E}\left[\max_{s\in B_m(t)} \left|\frac{\partial W_{\Phi}}{\partial s}(t,s)\right|\right], \qquad \sigma_{4,m}^2 = \max_{s\in B_m(t)} \mathbb{E}\left[\left|\frac{\partial W_{\Phi}}{\partial s}(t,s)\right|^2\right],$$

where

$$\mathbb{E}\left[\left|W_{\Phi}(t,s)\right|^{2}\right] = \int_{0}^{\infty} |\widehat{\psi}(s\lambda)|^{2} p(\lambda) d\lambda,$$
$$\mathbb{E}\left[\left|\frac{\partial W_{\Phi}}{\partial s}(t,s)\right|^{2}\right] = \int_{0}^{\infty} |\lambda[D\widehat{\psi}](s\lambda)|^{2} p(\lambda) d\lambda,$$

and  $D\widehat{\psi}$  represents the derivative of  $\widehat{\psi}$ .

**Theorem 5.** Suppose that f satisfies the AHM, the stationary Gaussian process  $\Phi$  satisfies Assumptions 1 and 3, and the mother wavelet  $\psi$  satisfies Assumption 2.

For every  $m \in \{1, 2, ..., M\}$ ,  $t \in \mathbb{R}$ , and  $B_m(t) \subset (\underline{i}_m(t), \overline{i}_m(t))$ , where the interval  $B_m(t)$  is defined in (19), let

(28) 
$$L_m = \min_{s \in B_m(t)} \left| \frac{\partial^2 S_f}{\partial s^2}(t, s) \right|$$

For any  $\varepsilon > 0$  with

(29) 
$$\min_{s \in B_m(t)} |s - s_{f,m}(t)| > \varepsilon > \frac{6}{L_m} \max\left\{\mu_{1,m}, \mu_{2,m}, \mu_{3,m}^2, \mu_{4,m}^2\right\},$$

we have

$$\mathbb{P}\left(|s_{Y,m}(t) - s_{f,m}(t)| > \varepsilon \mid s_{Y,m}(t) \in B_m^{\circ}(t)\right)$$

$$\leq \left[1 - \exp\left(-\frac{\left(\triangle_{B_m^{\circ}(t)} - \mu_{B_m^{\circ}(t)}\right)^2}{\sigma_{B_m^{\circ}(t)}^2}\right)\right]^{-1}$$

$$(30) \times \left\{\sum_{k=1}^2 \exp\left[-\frac{L_m^2}{\sigma_{k,m}^2} \left(\frac{\varepsilon}{6} - \frac{\mu_{k,m}}{L_m}\right)^2\right] + \sum_{k=3}^4 \exp\left[-\frac{L_m}{\sigma_{k,m}^2} \left(\sqrt{\frac{\varepsilon}{6}} - \frac{\mu_{k,m}}{\sqrt{L_m}}\right)^2\right]\right\},$$

where  $\triangle_{B_m^{\circ}(t)}$ ,  $\mu_{B_m^{\circ}(t)}$ , and  $\sigma_{B_m^{\circ}(t)}$  are defined in (26) with  $I = B_m^{\circ}(t)$  and  $B_m(t) \setminus I = \partial B_m(t)$ , i.e., the endpoints of the interval  $B_m(t)$ .

Intuitively, when the noise level is relatively low compared to the strength of the clean signal, the ridge point  $s_{Y,m}(t)$  of the noisy signal Y is unlikely to deviate too far from the ridge point  $s_{f,m}(t)$  of the clean signal f. Theorem 5 formalizes this intuition by providing a bound on the probability that the deviation  $|s_{Y,m}(t)-s_{f,m}(t)|$  exceeds a given threshold  $\varepsilon$ . The bound is expressed in terms of the threshold  $\varepsilon$  and the following ratios:

$$\frac{\mu_{1,m}}{L_m}, \frac{\mu_{2,m}}{L_m}, \frac{\mu_{3,m}^2}{L_m}, \frac{\mu_{4,m}^2}{L_m}, \frac{\sigma_{m,1}^2}{L_m^2}, \frac{\sigma_{m,2}^2}{L_m^2}, \frac{\sigma_{m,3}^2}{L_m}, \frac{\sigma_{m,4}^2}{L_m},$$

which are small when the noise strength is low relative to the clean signal. The connection between these ratios and the signal-to-noise ratio is discussed in the appendix. The proof of Theorem 5 is presented in Section 6.9.

5.3. Dudley's theorem and the Borell-TIS inequality for AWT. To derive Theorems 4 and 5, a key challenge is estimating the tail probabilities of the following random variables arising from the AWT of the Gaussian process  $\Phi$ :

• The sum of  $|W_{\Phi}(t, s_f(t))|$  and the supremum of  $|W_{\Phi}(t, s)|$  over  $s \in I^c$ , where I is an interval containing  $s_f(t)$  and  $I^c$  is the complement of I, i.e.,

(31) 
$$|W_{\Phi}(t, s_f(t))| + \max_{s \in I^c} |W_{\Phi}(t, s)|$$

• The supremum of  $|W_{\Phi}(t,s)|$  and  $\left|\frac{\partial W_{\Phi}}{\partial s}(t,s)\right|$  over  $s \in I$ , i.e.,

(32) 
$$\max_{s \in I} |W_{\Phi}(t,s)| \text{ and } \max_{s \in I} \left| \frac{\partial W_{\Phi}}{\partial s}(t,s) \right|.$$

• The maximum of the products

(33) 
$$\left| \frac{\partial W_f}{\partial s}(t,s) \right| |W_{\Phi}(t,s)| \text{ and } |W_f(t,s)| \left| \frac{\partial W_{\Phi}}{\partial s}(t,s) \right|$$

over  $s \in I$ .

The tail probabilities of the random variables above, with the replacement of the interval I and its complement  $I^c$  by a subinterval  $I_m$  of  $B_m(t)$  and the relative complement  $B_m(t) \setminus I_m$ , respectively, also play an important role in the proof of Theorem 5.

Our estimation method is inspired by the Borell-TIS inequality [39, 40], which provides a tail probability estimate for the supremum of a real-valued non-stationary Gaussian process over an index set. In the proof of the Borell-TIS inequality, the Gaussian concentration inequality for Lipschitz functions of real-valued Gaussian random vectors [20] plays an important role. Lemma 1 below generalizes the concentration inequality in [20] to accommodate complex-valued and circularly symmetric Gaussian random vectors, whose proof is provided in Section 6.10.

**Lemma 1.** Let **J** be a circularly symmetric Gaussian random vector in  $\mathbb{C}^{d\times 1}$  with  $\mathbb{E}[\mathbf{J}] = \mathbf{0}$ , where  $d \in \mathbb{N}$ . For any Lipschitz continuous function  $g : \mathbb{C}^{d\times 1} \to \mathbb{R}$  and any  $u > \mathbb{E}[g(\mathbf{J})]$ ,

$$\mathbb{P}\left(g(\mathbf{J}) > u\right) \le \exp\left(-\frac{(u - \mathbb{E}[g(\mathbf{J})])^2}{2\|g(\mathbf{A} \bullet)\|_{\mathrm{Lip}}^2}\right),\$$

where **A** is a matrix in  $\mathbb{C}^{d \times d}$  satisfying  $\mathbf{AA}^* = 2^{-1}\mathbb{E}\left[\mathbf{JJ}^*\right]$  and

$$\|g(\mathbf{A}\bullet)\|_{\mathrm{Lip}} = \sup_{\substack{\mathbf{x},\mathbf{y}\in\mathbb{C}^d\\\mathbf{x}\neq\mathbf{y}}} \frac{|g(\mathbf{A}\mathbf{x}) - g(\mathbf{A}\mathbf{y})|}{|\mathbf{x}-\mathbf{y}|}.$$

The Lipschitz continuous function g in Lemma 1 can be chosen as follows:

$$g(\mathbf{x}) = |x_1| + \max_{2 \le \ell \le d} |x_\ell|$$

for case (31), and

$$g(\mathbf{x}) = \|\mathbf{x}\|_{\ell^{\infty}} := \max_{1 \le \ell \le d} |x_{\ell}|,$$

for case (32), where  $\mathbf{x} = (x_1, x_2, \dots, x_d) \in \mathbb{C}^d$ .

Note that in cases (31) and (32), the supremum is taken over the continuous index sets I and  $I^c$ , respectively. In order to apply the dominated convergence theorem to extend the result in Lemma 1 to the continuous index set case, we need to ensure the convergence of the expected value of the supremum. Details of this extension are provided in the proof of Theorem 4. In [11, 39], Dudley's theorem provides an upper bound for the expected maximum of a centered real-valued Gaussian field over a compact set. Because the AWT of the Gaussian process  $\Phi$  is a complexvalued random process and the index set  $I^c$  for case (31) is noncompact, we present a complex-valued version of Dudley's theorem for the expectation of the supremum of the process  $|W_{\Phi}(t, \cdot)|$  over the noncompact set  $[0, \infty)$  in Lemma 3. The statement and proof of Lemma 3 rely on estimates of  $\mathbb{E}[|W_{\Phi}(0, s)|^2]$  for large scale s and the distance function  $d_{W_{\Phi}}$  as follows.

**Lemma 2.** Suppose that  $\psi$  is an analytic wavelet belonging to  $L^1 \cap L^2$ , and its Fourier transform  $\widehat{\psi}$  is Lipschitz continuous on  $[0, \infty)$ . For the random process  $\Phi$ , if Assumption 1 and Assumptions 3(a) and 3(b1) hold, then for any constant  $H^- \in (0, H)$ , where H is the long-memory parameter of  $\Phi$  defined in (10), there exists a threshold T > 0 such that for any  $s \geq T$ ,

(34) 
$$\mathbb{E}[|W_{\Phi}(0,s)|^2] \le s^{-H^-}.$$

Under the same assumptions, for any  $s_1, s_2 > 0$  satisfying  $|s_1 - s_2| \leq 1$ , the distance  $d_{W_{\Phi}}(s_1, s_2)$ , defined in (9), is  $\gamma/2$ -Hölder continuous:

(35) 
$$d_{W_{\Phi}}(s_1, s_2) \le C_2 |s_1 - s_2|^{\gamma/2},$$

where  $\gamma$  is given in Assumption 3(a),

(36) 
$$C_2 = 1 \vee \left[ C_1 \left( \frac{1}{\gamma} + \frac{2}{2 - \gamma} \right) (2 \| \widehat{\psi} \|_{\infty})^{2 - \gamma} \| \widehat{\psi} \|_{\mathrm{Lip}}^{\gamma} + 2 \| \widehat{\psi} \|_{\infty}^2 \mathrm{Var} \left( \Phi(0) \right) \right]^{1/2}$$

and  $\|\psi\|_{\text{Lip}}$  represents the Lipschitz constant of  $\psi$ .

If the spectral density function of  $\Phi$  satisfies Assumption 3(b2) instead of Assumption 3(b1), the results in (34) and (35) still hold with the substitution  $H^- = 1$ .

**Lemma 3** (Dudley's theorem for AWT). Suppose that  $\psi$  is an analytic wavelet belonging to  $L^1 \cap L^2$ , and its Fourier transform  $\widehat{\psi}$  is Lipschitz continuous on  $[0, \infty)$ . For the random process  $\Phi$ , if Assumptions 1 and 3 hold, then

$$\mathbb{E}\left[\sup_{s>0}|W_{\Phi}(0,s)|\right] \le 4\left[\sqrt{\ln \mathcal{D}}\frac{\mathcal{D}^3}{(\mathcal{D}-1)^2}\sqrt{\frac{1}{\gamma}+\frac{1}{H^-}} + \left(\sqrt{\frac{1}{\gamma}\ln C_2} + \sqrt{\ln T}\right)\frac{\mathcal{D}^2}{\mathcal{D}-1}\right]$$

where  $H^-$ ,  $C_2$ , and T are constants mentioned in Lemma 2 and

(38) 
$$\mathcal{D} = 2 \vee \left\{ \sup_{s>0} \mathbb{E} \left[ |W_{\Phi}(0,s)|^2 \right] \right\}^{\frac{1}{2}}.$$

In [11, 39], the expectation of the maximum of a centered real-valued Gaussian field over a compact set is bounded above by the Dudley entropy integral. Lemma 3 extends this bound, providing an exact upper bound for the expectation of the maximum of the modulus of a circularly symmetric and nonstationary complex Gaussian process over  $(0, \infty)$ . The upper bound given in (37) increases as the long-memory parameter H and the decay rate parameter  $\gamma$  of the spectral density function of  $\Phi$  decrease to zero. This relationship echoes Slepian's inequality and the Sudakov-Fernique inequality [40, Chapter 7], which state that, for two realvalued Gaussian processes with the same mean, the process with higher correlation exhibits a greater expected maximum. The proof of Lemma 3 is provided in Section 6.12. Finally, to estimate the tail probabilities of the random variables in (33), we introduce the complex-valued version of the Borell-TIS inequality as follows.

**Lemma 4** (Borell-TIS inequality for AWT). Suppose that the stationary Gaussian process  $\Phi$  satisfies Assumptions 1 and 3, and the mother wavelet  $\psi$  satisfies Assumption 2. For any  $a, b \in (0, \infty)$  and any bounded and continuous function  $w : [a, b] \to [0, \infty)$ , define either

$$\begin{cases} M_{[a,b]} = \max_{s \in [a,b]} w(s) |W_{\Phi}(0,s)|, \\ \sigma_{[a,b]} = \max_{s \in [a,b]} w(s) \left(\int_{0}^{\infty} |\widehat{\psi}(s\lambda)|^{2} p(\lambda) d\lambda\right)^{\frac{1}{2}}, \end{cases}$$

or

$$M_{[a,b]} = \max_{s \in [a,b]} w(s) \left| \frac{\partial W_{\Phi}}{\partial s}(0,s) \right|,$$
  
$$\sigma_{[a,b]} = \max_{s \in [a,b]} w(s) \left( \int_0^\infty |\lambda[D\widehat{\psi}](s\lambda)|^2 p(\lambda) d\lambda \right)^{\frac{1}{2}}$$

In both cases, we have  $\mathbb{E}\left[M_{[a,b]}\right] < \infty$ , and for any  $u \ge \mathbb{E}\left[M_{[a,b]}\right]$ ,

$$\mathbb{P}\left(M_{[a,b]} \ge u\right) \le \exp\left(-\frac{\left(u - \mathbb{E}\left[M_{[a,b]}\right]\right)^2}{\sigma_{[a,b]}^2}\right).$$

The proof of Lemma 4 is provided in Section 6.13. Last but not least, our work bridges and extends classical maximum theorems and maximum inequalities to address a key question in signal processing: how noise perturbs the ridge curves of adapted harmonic signals that encode the IF information.

# 6. Proofs

6.1. **Proof of Theorem 1.** (a) To prove that **W** is circularly symmetric, it suffices to show that any element in the pseudo-covariance matrix of **W** is zero. Denote  $W_{\Phi}^{R} = \operatorname{Re}(W_{\Phi})$  and  $W_{\Phi}^{I} = \operatorname{Im}(W_{\Phi})$ . For any  $k, k' \in \{1, \ldots, d\}$ ,

$$\mathbb{E}\left[W_{\Phi}(t,s_{k})W_{\Phi}(t,s_{k'})\right] = \mathbb{E}\left[W_{\Phi}^{R}(t,s_{k})W_{\Phi}^{R}(t,s_{k'})\right] - \mathbb{E}\left[W_{\Phi}^{I}(t,s_{k})W_{\Phi}^{I}(t,s_{k'})\right]$$

$$(39) \qquad \qquad + i\mathbb{E}\left[W_{\Phi}^{R}(t,s_{k})W_{\Phi}^{I}(t,s_{k'})\right] + i\mathbb{E}\left[W_{\Phi}^{I}(t,s_{k})W_{\Phi}^{R}(t,s_{k'})\right].$$

Denote  $\psi_R = \operatorname{Re}(\psi)$  and  $\psi_I = \operatorname{Im}(\psi)$ . Analogous to the representation in (8),

(40) 
$$W_{\Phi}^{R}(t,s) = \int_{\mathbb{R}} e^{it\lambda} \overline{\widehat{\psi}_{R}(s\lambda)} \sqrt{p(\lambda)} Z(d\lambda)$$

and

(41) 
$$W^{I}_{\Phi}(t,s) = \int_{\mathbb{R}} e^{it\lambda} \overline{\widehat{\psi_{I}}(s\lambda)} \sqrt{p(\lambda)} Z(d\lambda).$$

Because  $\psi$  is an analytic wavelet, the Fourier transforms of  $\psi_R$  and  $\psi_I$  are related by the following equation:

(42) 
$$\widehat{\psi}_{I}(\zeta) = -i \operatorname{sgn}(\zeta) \widehat{\psi}_{R}(\zeta), \ \zeta \in \mathbb{R},$$

where sgn is the signum function. By the orthogonal property (5) of the Gaussian random measure Z and (42),

(43)  

$$\mathbb{E}\left[W_{\Phi}^{I}(t,s_{k})W_{\Phi}^{I}(t,s_{k'})\right] = \int_{-\infty}^{\infty} \overline{\widehat{\psi}_{I}(s_{k}\lambda)}\widehat{\psi}_{I}(s_{k'}\lambda)p(\lambda)d\lambda$$

$$= \int_{-\infty}^{\infty} \overline{\widehat{\psi}_{R}(s_{k}\lambda)}\widehat{\psi}_{R}(s_{k'}\lambda)p(\lambda)d\lambda$$

$$= \mathbb{E}\left[W_{\Phi}^{R}(t,s_{k})W_{\Phi}^{R}(t,s_{k'})\right]$$

and

(44)  

$$\mathbb{E}\left[W_{\Phi}^{I}(t,s_{k})W_{\Phi}^{R}(t,s_{k'})\right] = \int_{-\infty}^{\infty} \overline{\widehat{\psi_{I}}(s_{k}\lambda)}\widehat{\psi_{R}}(s_{k'}\lambda)p(\lambda)d\lambda$$

$$= -\int_{-\infty}^{\infty} \overline{\widehat{\psi_{R}}(s_{k}\lambda)}\widehat{\psi_{I}}(s_{k'}\lambda)p(\lambda)p(\lambda)d\lambda$$

$$= -\mathbb{E}\left[W_{\Phi}^{R}(t,s_{k})W_{\Phi}^{I}(t,s_{k'})\right].$$

By substituting (43) and (44) into (39), we obtain  $\mathbb{E}[W_{\Phi}(t, s_k)W_{\Phi}(t, s_{k'})] = 0$ . (b) For any  $t \in \mathbb{R}$ ,  $s_1, s_2 > 0$  with  $s_1 \neq s_2$ , by (40) and (41),

$$W_{\Phi}^{R}(t,s_{1}) - W_{\Phi}^{R}(t,s_{2}) = \int_{\mathbb{R}} e^{it\lambda} \left[ \overline{\widehat{\psi}_{R}(s_{1}\lambda)} - \overline{\widehat{\psi}_{R}(s_{2}\lambda)} \right] \sqrt{p(\lambda)} Z(d\lambda)$$

and

$$W_{\Phi}^{I}(t,s_{1}) - W_{\Phi}^{I}(t,s_{2}) = \int_{\mathbb{R}} e^{it\lambda} \left[ \overline{\widehat{\psi}_{I}(s_{1}\lambda)} - \overline{\widehat{\psi}_{I}(s_{2}\lambda)} \right] \sqrt{p(\lambda)} Z(d\lambda).$$

By the orthogonal property (5) of the Gaussian random measure Z,

(45) 
$$\mathbb{E}\left[\left(W_{\Phi}^{R}(t,s_{1}) - W_{\Phi}^{R}(t,s_{2})\right)\left(W_{\Phi}^{I}(t,s_{1}) - W_{\Phi}^{I}(t,s_{2})\right)\right] \\ = \int_{\mathbb{R}}\left[\widehat{\psi}_{R}(s_{1}\lambda) - \widehat{\psi}_{R}(s_{2}\lambda)\right]\left[\overline{\widehat{\psi}_{I}(s_{1}\lambda)} - \overline{\widehat{\psi}_{I}(s_{2}\lambda)}\right]p(\lambda)d\lambda.$$

Applying (42) again, and noting that the spectral density function p is even, we can rewrite (45) as follows

$$\mathbb{E}\left[\left(W_{\Phi}^{R}(t,s_{1})-W_{\Phi}^{R}(t,s_{2})\right)\left(W_{\Phi}^{I}(t,s_{1})-W_{\Phi}^{I}(t,s_{2})\right)\right]$$
$$=i\int_{\mathbb{R}}\left|\widehat{\psi_{R}}(s_{1}\lambda)-\widehat{\psi_{R}}(s_{2}\lambda)\right|^{2}\operatorname{sgn}(\lambda)p(\lambda)d\lambda=0,$$

which shows that the real and imaginary parts of  $W_{\Phi}(t, s_1) - W_{\Phi}(t, s_2)$  are independent. Following the same procedure, we can obtain that

(46) 
$$\mathbb{E}\left[\left|W_{\Phi}^{R}(t,s_{1}) - W_{\Phi}^{R}(t,s_{2})\right|^{2}\right] = \mathbb{E}\left[\left|W_{\Phi}^{I}(t,s_{1}) - W_{\Phi}^{I}(t,s_{2})\right|^{2}\right] = \frac{1}{2}d_{W_{\Phi}}^{2}(s_{1},s_{2}).$$

The equality (46) implies that

$$\begin{aligned} & \frac{|W_{\Phi}(t,s_1) - W_{\Phi}(t,s_2)|^2}{d_{W_{\Phi}}^2(s_1,s_2)} \\ &= & \frac{1}{2} \left( \frac{|W_{\Phi}^R(t,s_1) - W_{\Phi}^R(t,s_2)|^2}{\mathbb{E} \left[|W_{\Phi}^R(t,s_1) - W_{\Phi}^R(t,s_2)|^2\right]} + \frac{|W_{\Phi}^I(t,s_1) - W_{\Phi}^I(t,s_2)|^2}{\mathbb{E} \left[|W_{\Phi}^I(t,s_1) - W_{\Phi}^I(t,s_2)|^2\right]} \right). \end{aligned}$$

Theorem 1(b) follows from the fact that the distribution of half the sum of two independent chi-square random variables, each with one degree of freedom, is an exponential distribution with a mean of one.

(c) By (40), (41), and Itô's formula,

$$\left(W_{\Phi}^{R}(t,s)\right)^{2} = \mathbb{E}\left[\left(W_{\Phi}^{R}(t,s)\right)^{2}\right] + \int_{\mathbb{R}^{2}}^{\prime} e^{it(\lambda_{1}+\lambda_{2})} \overline{\widehat{\psi}_{R}(s\lambda_{1})\widehat{\psi}_{R}(s\lambda_{2})} \sqrt{p(\lambda_{1})p(\lambda_{2})} Z(d\lambda_{1}) Z(d\lambda_{2}),$$

where  $\int'$  means that the integral excludes the diagonal hyperplanes  $\lambda_1 = \pm \lambda_2$  [25]. Similarly,

$$\left(W_{\Phi}^{I}(t,s)\right)^{2} = \mathbb{E}\left[\left(W_{\Phi}^{I}(t,s)\right)^{2}\right] + \int_{\mathbb{R}^{2}}^{\prime} e^{it(\lambda_{1}+\lambda_{2})} \overline{\widehat{\psi_{I}}(s\lambda_{1})\widehat{\psi_{I}}(s\lambda_{2})} \sqrt{p(\lambda_{1})p(\lambda_{2})} Z(d\lambda_{1}) Z(d\lambda_{2}).$$

Hence, for any s > 0,

$$S_{\Phi}(t,s) - \mathbb{E}\left[S_{\Phi}(t,s)\right]$$
$$= \int_{\mathbb{R}^2}' e^{it(\lambda_1 + \lambda_2)} \left[ \overline{\widehat{\psi_R}(s\lambda_1)\widehat{\psi_R}(s\lambda_2)} + \overline{\widehat{\psi_I}(s\lambda_1)\widehat{\psi_I}(s\lambda_2)} \right] \sqrt{p(\lambda_1)p(\lambda_2)} Z(d\lambda_1) Z(d\lambda_2).$$

By the product formula for the second Wiener chaos [25, 29],

$$Cov \left(S_{\Phi}(t, s_{1}), S_{\Phi}(t, s_{2})\right) = 2 \int_{\mathbb{R}^{2}} \left[ \overline{\widehat{\psi_{R}}(s_{1}\lambda_{1})\widehat{\psi_{R}}(s_{1}\lambda_{2})} + \overline{\widehat{\psi_{I}}(s_{1}\lambda_{1})\widehat{\psi_{I}}(s_{1}\lambda_{2})} \right]$$

$$(47) \qquad \times \left[ \widehat{\psi_{R}}(s_{2}\lambda_{1})\widehat{\psi_{R}}(s_{2}\lambda_{2}) + \widehat{\psi_{I}}(s_{2}\lambda_{1})\widehat{\psi_{I}}(s_{2}\lambda_{2}) \right] p(\lambda_{1})p(\lambda_{2})d\lambda_{1}d\lambda_{2}$$

$$= \sum_{i=1}^{2} \sum_{j=1}^{2} \varphi_{i,j}(s_{1}, s_{2}),$$

where

(48) 
$$\varphi_{i,j}(s_1, s_2) = 2 \left( \int_{\mathbb{R}} \widehat{\psi_{\alpha(i)}(s_1\lambda)} \widehat{\psi_{\alpha(j)}(s_2\lambda)} p(\lambda) d\lambda \right)^2.$$

 $\alpha(1) = R$ , and  $\alpha(2) = I$ . Because  $\psi$  is an analytic wavelet, the relation (42) implies that  $\varphi_{1,1}(s_1, s_2) = \varphi_{2,2}(s_1, s_2)$  and  $\varphi_{1,2}(s_1, s_2) = \varphi_{2,1}(s_1, s_2)$ .

On the other hand, from (40) and (41), we have

(49) 
$$\mathbb{E}\left[W_{\Phi}^{R}(t,s_{1})W_{\Phi}^{R}(t,s_{2})\right] = \int_{\mathbb{R}}\overline{\widehat{\psi}_{R}(s_{1}\lambda)}\widehat{\psi}_{R}(s_{2}\lambda)p(\lambda)d\lambda$$

and

(50) 
$$\mathbb{E}\left[W_{\Phi}^{R}(t,s_{1})W_{\Phi}^{I}(t,s_{2})\right] = \int_{\mathbb{R}} \overline{\widehat{\psi}_{R}(s_{1}\lambda)}\widehat{\psi}_{I}(s_{2}\lambda)p(\lambda)d\lambda.$$

By (48), (50), and (49),

(51) 
$$\varphi_{i,j}(s_1, s_2) = \begin{cases} 2\left(\mathbb{E}\left[W_{\Phi}^R(t, s_1)W_{\Phi}^R(t, s_2)\right]\right)^2 & \text{if } i = j, \\ 2\left(\mathbb{E}\left[W_{\Phi}^R(t, s_1)W_{\Phi}^I(t, s_2)\right]\right)^2 & \text{if } i \neq j. \end{cases}$$

The proof of the first equality in Theorem 1(c) is concluded by substituting (51) into (48).

When  $s_1 = s_2 = s > 0$ , (44) shows that

(52) 
$$\mathbb{E}\left[W_{\Phi}^{R}(t,s)W_{\Phi}^{I}(t,s)\right] = 0.$$

The second equality in Theorem 1(c) implies that

$$\operatorname{Var}\left(S_{\Phi}(t,s)\right) = 4\left(\mathbb{E}\left[|W_{\Phi}^{R}(t,s)|^{2}\right]\right)^{2}$$

The proof of the second equality in Theorem 1(c) is concluded by noting that

$$\mathbb{E}[S_{\Phi}(t,s)] = \mathbb{E}\left[|W_{\Phi}^{R}(t,s)|^{2}\right] + \mathbb{E}\left[|W_{\Phi}^{I}(t,s)|^{2}\right] \stackrel{(43)}{=} 2\mathbb{E}\left[|W_{\Phi}^{R}(t,s)|^{2}\right].$$

6.2. **Proof of Proposition 1.** The proof consists of three steps. First, we show that the sample paths of  $W_{\Phi}$  are almost surely continuous on  $\mathbb{R} \times (0, \infty)$ . Next, we prove that the sample paths of  $W_{\Phi}$  are continuously differentiable on  $\mathbb{R} \times (0, \infty)$  almost surely. Finally, the second-order differentiability of the sample paths of  $W_{\Phi}$  follows by the same arguments.

Step 1. To prove that the sample paths of the Gaussian field  $W_{\Phi}$  are almost surely continuous on  $\mathbb{R} \times (0, \infty)$ , according to [12, p.73 and p.93], it suffices to show the existence of a constant q > 0 such that for any  $\delta > 0$ , there exists a constant  $C_{\delta}$  satisfying

(53) 
$$\left\{ \mathbb{E}\left[ \left| W_{\Phi}(t',s') - W_{\Phi}(t,s) \right|^2 \right] \right\}^{1/2} \le C_{\delta} \left( |t - t'|^2 + |s - s'|^2 \right)^{q/2}$$

for any  $t, t' \in \mathbb{R}$  and  $s, s' > \delta$ . We prove (53) as follows. By the triangle inequality and (8),

$$\left\{ \mathbb{E} \left[ \left| W_{\Phi}(t',s') - W_{\Phi}(t,s) \right|^{2} \right] \right\}^{1/2}$$
  
 
$$\leq \left\{ \mathbb{E} \left[ \left| W_{\Phi}(t',s') - W_{\Phi}(t',s) \right|^{2} \right] \right\}^{1/2} + \left\{ \mathbb{E} \left[ \left| W_{\Phi}(t',s) - W_{\Phi}(t,s) \right|^{2} \right] \right\}^{1/2}$$
  
(54) 
$$= \sqrt{\mathcal{E}_{1}} + \sqrt{\mathcal{E}_{2}},$$

where

$$\mathcal{E}_1 = \int_{\mathbb{R}} |\widehat{\psi}(s'\lambda) - \widehat{\psi}(s\lambda)|^2 p(\lambda) d\lambda$$

and

$$\mathcal{E}_2 = \int_{\mathbb{R}} |e^{i(t'-t)\lambda} - 1|^2 |\widehat{\psi}(s\lambda)|^2 p(\lambda) d\lambda.$$

*Estimate of*  $\mathcal{E}_1$ : By the mean value theorem, there exists a constant  $c_{s,s',\lambda}$  lying between  $s\lambda$  and  $s'\lambda$  such that

$$\mathcal{E}_1 = \int_0^\infty |D\widehat{\psi}(c_{s,s',\lambda})(s'\lambda - s\lambda)|^2 p(\lambda) d\lambda.$$

Because  $s, s' > \delta$ , we have  $c_{s,s',\lambda} > \delta\lambda$  and

(55) 
$$\mathcal{E}_{1} \leq \delta^{-2} |s'-s|^{2} \int_{0}^{\infty} |D\widehat{\psi}(c_{s,s',\lambda})c_{s,s',\lambda}|^{2} p(\lambda) d\lambda$$
$$\leq \delta^{-2} |s'-s|^{2} \left( \sup_{z>0} |zD\widehat{\psi}(z)|^{2} \right) \mathbb{E} \left[ \Phi^{2}(0) \right],$$

where the supremum is finite under Assumptions  $2(D_1^1)$ .

Estimate of  $\mathcal{E}_2$ : Given that  $|e^{ix} - 1| \le |x|$  for any  $x \in \mathbb{R}$ , and  $s > \delta$ ,

(56) 
$$\mathcal{E}_2 \leq |t'-t|^2 \int_{\mathbb{R}} |\widehat{\psi}(s\lambda)|^2 \lambda^2 p(\lambda) d\lambda \leq \delta^{-2} |t'-t|^2 \left( \sup_{z>0} |z\widehat{\psi}(z)|^2 \right) \mathbb{E} \left[ |\Phi(0)|^2 \right],$$

where the supremum is finite under Assumption  $2(D_1^0)$ .

Substituting (55) and (56) into (54) shows that (53) holds with q = 1 and

$$C_{\delta} = \delta^{-1} \left\{ \left( \sup_{z>0} |z\widehat{\psi}(z)|^2 \right) + \left( \sup_{z>0} |zD\widehat{\psi}(z)|^2 \right) \right\}^{1/2} \left\{ \mathbb{E} \left[ \Phi^2(0) \right] \right\}^{1/2}.$$

Step 2. Motivated from (8), we consider two random fields

(57) 
$$\partial_t W_{\Phi}(t,s) = i \int_0^\infty \lambda e^{it\lambda} \overline{\widehat{\psi}(s\lambda)} \sqrt{p(\lambda)} Z(d\lambda),$$

(58) 
$$\partial_s W_{\Phi}(t,s) = \int_0^\infty \lambda e^{it\lambda} \overline{[D\widehat{\psi}](s\lambda)} \sqrt{p(\lambda)} Z(d\lambda),$$

where  $D\hat{\psi}$  represents the derivative of  $\hat{\psi}$ . Under Assumption 2, conditions  $(D_1^0)$  and  $(D_1^1)$  ensure that

$$\mathbb{E}|\partial_t W_{\Phi}(t,s)|^2 < \infty \text{ and } \mathbb{E}|\partial_s W_{\Phi}(t,s)|^2 < \infty.$$

To prove that sample paths of  $W_{\Phi}$  are continuously differentiable almost surely, according to [33], it suffices to show that (a) the difference quotient of  $W_{\Phi}$  along the time direction converges to  $\partial_t W_{\Phi}$  in quadratic mean, (b) the difference quotient of  $W_{\Phi}$  along the scale direction converges to  $\partial_s W_{\Phi}$  in quadratic mean, and (c) both  $\partial_t W_{\Phi}$  and  $\partial_s W_{\Phi}$  are continuous in quadratic mean and almost surely have continuous realizations. We verify these conditions as follows.

Proof of (a): By (8) and (57), for any  $h \in \mathbb{R} \setminus \{0\}$ ,

$$\frac{W_{\Phi}(t+h,s) - W_{\Phi}(t,s)}{h} - \partial_t W_{\Phi}(t,s)$$
$$= \int_0^\infty e^{it\lambda} \left(\frac{e^{ih\lambda} - 1}{h} - i\lambda\right) \overline{\widehat{\psi}(s\lambda)} \sqrt{p(\lambda)} Z(d\lambda).$$

By the isometry property of the Wiener integral,

(59) 
$$\mathbb{E}\left[\left|\frac{W_{\Phi}(t+h,s) - W_{\Phi}(t,s)}{h} - \partial_{t}W_{\Phi}(t,s)\right|^{2}\right]$$
$$= \int_{0}^{\infty} \left|\frac{e^{ih\lambda} - 1}{h} - i\lambda\right|^{2} |\widehat{\psi}(s\lambda)|^{2} p(\lambda) d\lambda.$$

Note that for any  $h \in \mathbb{R} \setminus \{0\}$  and  $\lambda \in \mathbb{R}$ ,

(60) 
$$\left|\frac{e^{ih\lambda}-1}{h}-i\lambda\right| \le 2|\lambda|.$$

Assumption  $2(D_1^0)$ , together with (60), enables us to apply the dominated convergence theorem to (59) and obtain

$$\lim_{h \to 0} \mathbb{E}\left[ \left| \frac{W_{\Phi}(t+h,s) - W_{\Phi}(t,s)}{h} - \partial_t W_{\Phi}(t,s) \right|^2 \right] = 0,$$

which shows that the difference quotient of  $W_{\Phi}$  along the time direction converges to  $\partial_t W_{\Phi}$  in quadratic mean.

Proof of (b): By (8) and (58), for any  $h \in (-s/2, s/2) \setminus \{0\}$ ,

$$\frac{W_{\Phi}(t,s+h) - W_{\Phi}(t,s)}{h} - \partial_s W_{\Phi}(t,s) \\= \int_0^\infty e^{it\lambda} \left[ \frac{h}{\widehat{\psi}((s+h)\lambda)} - \overline{\widehat{\psi}(s\lambda)}}{h} - \lambda \overline{[D\widehat{\psi}](s\lambda)} \right] \sqrt{p(\lambda)} Z(d\lambda).$$

By the isometry property of the Wiener integral,

(61) 
$$\mathbb{E}\left[\left|\frac{W_{\Phi}(t,s+h) - W_{\Phi}(t,s)}{h} - \partial_s W_{\Phi}(t,s)\right|^2\right] = \int_0^\infty |K_h(s,\lambda)|^2 p(\lambda) d\lambda,$$

where

$$K_h(s,\lambda) = \frac{\widehat{\psi}((s+h)\lambda) - \widehat{\psi}(s\lambda)}{h} - \lambda [D\widehat{\psi}](s\lambda).$$

For any  $\lambda > 0$  and  $h \in (-s/2, s/2) \setminus \{0\}$ ,

$$\begin{split} \left| \frac{\widehat{\psi}((s+h)\lambda) - \widehat{\psi}(s\lambda)}{h} \right| &= \left| \lambda \int_0^1 [D\widehat{\psi}] \left( (s+h)\lambda u + s\lambda(1-u) \right) du \right| \\ &\leq \left( \sup_{z>0} \left| zD\widehat{\psi}(z) \right| \right) \lambda \int_0^1 \left| \frac{1}{(s+h)\lambda u + s\lambda(1-u)} \right| du \\ &= \left( \sup_{z>0} \left| zD\widehat{\psi}(z) \right| \right) \frac{1}{|h|} \ln \left| 1 + \frac{h}{s} \right| \\ &\leq \left( \sup_{z>0} \left| zD\widehat{\psi}(z) \right| \right) \left( \frac{2\ln 2}{s} \right), \end{split}$$

where the supremum is finite under Assumption  $2(D_1^1)$ . Hence, for any s > 0,

(62) 
$$\sup_{h \in (-s/2, s/2) \setminus \{0\}} \sup_{\lambda > 0} |K_h(s, \lambda)| < \left( \sup_{z > 0} \left| z D \widehat{\psi}(z) \right| \right) \left( \frac{1 + 2 \ln 2}{s} \right) < \infty.$$

The differentiability of  $\widehat{\psi}$  implies that  $K_h(s, \lambda) \to 0$  as  $h \to 0$ . The uniform bound (62) enables us to apply the dominated convergence theorem to (61), yielding

$$\lim_{h \to 0} \mathbb{E}\left[ \left| \frac{W_{\Phi}(t,s+h) - W_{\Phi}(t,s)}{h} - \partial_s W_{\Phi}(t,s) \right|^2 \right] = 0.$$

This result shows that the difference quotient of  $W_{\Phi}$  along the scale direction converges to  $\partial_s W_{\Phi}$  in quadratic mean.

Proof of (c): To prove that  $\partial_t W_{\Phi}$  and  $\partial_s W_{\Phi}$  not only are continuous in the quadratic mean but also almost surely have modification that is sample continuous, according to [12, p.73 and p.93] (see also [32, Corollary 4.6]), it suffices to show that for any  $\delta > 0$ , there exists a constant  $C_{\delta}$  such that

$$\left\{ \mathbb{E}\left[ \left| \partial_t W_{\Phi}(t', s') - \partial_t W_{\Phi}(t, s) \right|^2 \right] \right\}^{1/2} \le C_{\delta} \left( |t - t'|^2 + |s - s'|^2 \right)^{1/2}$$

and

$$\left\{ \mathbb{E}\left[ \left| \partial_s W_{\Phi}(t',s') - \partial_s W_{\Phi}(t,s) \right|^2 \right] \right\}^{1/2} \le C_{\delta} \left( |t - t'|^2 + |s - s'|^2 \right)^{1/2}$$

for any  $t, t' \in \mathbb{R}$  and  $s, s' > \delta$ . This step requires additional conditions, including conditions  $(D_2^0)$ ,  $(D_2^1)$  and  $(D_2^2)$  in Assumption 2. Since the proof is similar to the argument used to prove the continuity of the sample paths of  $W_{\Phi}$  on  $\mathbb{R} \times (0, \infty)$ , it is omitted here. Based on the verification process (a)-(c), we conclude that the sample paths of  $W_{\Phi}$  are continuously differentiable almost surely.

Step 3. The twice continuous differentiability of  $W_{\Phi}$  follows by applying the same arguments, i.e., the verification process (a)-(c), to the derivatives  $\partial_t W_{\Phi}$  and  $\partial_s W_{\Phi}$ . This part of proof relies on conditions (D<sub>3</sub><sup>2</sup>) and (D<sub>3</sub><sup>3</sup>) in Assumption 2.

6.3. **Proof of Corollary 1.** Proof of (11): Based on the continuity of the sample paths of  $W_{\Phi}$  on  $\mathbb{R} \times (0, \infty)$ , it suffices to show that the sample paths of the Gaussian field  $W_{\Phi}$  are almost surely continuous at the boundary  $\mathbb{R} \times \{0\}$  if we define  $W_{\Phi}(t,0) = 0$  for any  $t \in \mathbb{R}$ . According to [12, p.73 and p.93], it suffices to show the existence of a constant q > 0 such that there exists a constant C satisfying

(63) 
$$\mathbb{E}\left[\left|W_{\Phi}(t,s)\right|^{2}\right] = \int_{0}^{\infty} |\widehat{\psi}(s\lambda)|^{2} p(\lambda) d\lambda \leq Cs^{q}$$

for any  $t \in \mathbb{R}$  and sufficiently small s > 0. Under Assumption 2(D<sub>0</sub><sup>1</sup>), we have  $|\widehat{\psi}(s\lambda)| \leq ||D\widehat{\psi}||_{\infty} |s\lambda|$ . Hence,

(64) 
$$\int_0^{s^{-2/(2+\gamma)}} |\widehat{\psi}(s\lambda)|^2 p(\lambda) d\lambda \le \|D\widehat{\psi}\|_{\infty}^2 \mathbb{E}[\Phi^2(0)] s^{\frac{2\gamma}{2+\gamma}}.$$

On the other hand, by Assumption 3(a),

(65) 
$$\int_{s^{-2/(2+\gamma)}}^{\infty} |\widehat{\psi}(s\lambda)|^2 p(\lambda) d\lambda \le \|\widehat{\psi}\|_{\infty}^2 C_1 \gamma^{-1} s^{\frac{2\gamma}{2+\gamma}}.$$

By combining (64) and (65), we obtain that

$$\mathbb{E}\left[\left|W_{\Phi}(t,s)\right|^{2}\right] \leq \left[\left\|D\widehat{\psi}\right\|_{\infty}^{2} \mathbb{E}[\Phi^{2}(0)] + \left\|\widehat{\psi}\right\|_{\infty}^{2} C_{1} \gamma^{-1}\right] s^{\frac{2\gamma}{2+\gamma}}.$$

Hence, (63) holds with  $q = 2\gamma/(2+\gamma)$ .

*Proof of (12):* We first introduce a Gaussian random field  $\varphi : \mathbb{R} \times [0, \infty) \mapsto \mathbb{C}$  as follows

$$\varphi(t,\tau) = \begin{cases} W_{\Phi}(t,1/\tau) & \text{ for } \tau > 0, \\ 0 & \text{ for } \tau = 0. \end{cases}$$

Similarly to the proof of (11), proving (12) is equivalent to showing that the function  $\varphi$  is continuous at the boundary  $\mathbb{R} \times \{0\}$ . For any  $t \in \mathbb{R}$  and  $\tau > 0$ ,

(66) 
$$\mathbb{E}\left[\left|\varphi(t,\tau)\right|^{2}\right] = \int_{0}^{\infty} |\widehat{\psi}(\frac{\lambda}{\tau})|^{2} p(\lambda) d\lambda = \tau \int_{0}^{\infty} |\widehat{\psi}(u)|^{2} p(\tau u) du.$$

Under Assumption 3(b1), (66) can be rewritten as

$$\mathbb{E}\left[\left|\varphi(t,\tau)\right|^{2}\right] = \tau \int_{0}^{\infty} |\widehat{\psi}(u)|^{2} \frac{L(|\tau u|^{-1})}{|\tau u|^{1-H}} du.$$

By [3, Theorem 1.5.3],

$$\lim_{\tau \to 0+} \left[ L(\tau^{-1})\tau^H \right]^{-1} \mathbb{E} \left[ |\varphi(t,\tau)|^2 \right] = \lim_{\tau \to 0+} \int_0^\infty \frac{|\widehat{\psi}(u)|^2}{|u|^{1-H}} \frac{L(|\tau u|^{-1})}{L(\tau^{-1})} du = \int_0^\infty \frac{|\widehat{\psi}(u)|^2}{|u|^{1-H}} du$$

Hence,  $\mathbb{E}\left[|\varphi(t,\tau)|^2\right] \lesssim L(\tau^{-1})\tau^H$  as  $\tau$  is sufficiently small. On the other hand, under Assumption 3(b2), it can similarly be shown that  $\mathbb{E}\left[|\varphi(t,\tau)|^2\right] \lesssim \tau$  as  $\tau$  is sufficiently small. Therefore, the sample paths of  $\varphi$  are continuous at the boundary  $\mathbb{R} \times \{0\}$  almost surely.

6.4. Proof of Proposition 2. For the squared integrable function f, [9] shows that the inverse wavelet transform

(67) 
$$f(t) = \operatorname{Re}\left[C_{\psi}^{-1} \int_{0}^{\infty} W_{f}(t,s)s^{-1}ds\right],$$

holds. In the following, we show that the relation in (67) remains valid when f is replaced by  $\Phi$ . By the spectral representation of  $W_{\Phi}$  in (8)

$$\int_0^\infty W_{\Phi}(t,s)s^{-1}ds = \int_0^\infty \left[\int_0^\infty e^{it\lambda}\overline{\widehat{\psi}(s\lambda)}\sqrt{p(\lambda)}Z(d\lambda)\right]s^{-1}ds.$$

The condition (13) allows us to use the stochastic Fubini theorem [31], yielding

(68) 
$$\int_{0}^{\infty} W_{\Phi}(t,s)s^{-1}ds = \int_{0}^{\infty} e^{it\lambda} \left[ \int_{0}^{\infty} \overline{\widehat{\psi}(s\lambda)}s^{-1}ds \right] \sqrt{p(\lambda)}Z(d\lambda)$$
$$= 2C_{\psi} \int_{0}^{\infty} e^{it\lambda} \sqrt{p(\lambda)}Z(d\lambda).$$

Because  $\overline{Z([a,b])} = Z([-b,-a])$  for any b > a > 0,

(69) 
$$\operatorname{Re}\left[2\int_{0}^{\infty}e^{it\lambda}\sqrt{p(\lambda)}Z(d\lambda)\right] = \Phi(t).$$

By (68) and (69), we obtain that

$$\Phi(t) = \operatorname{Re}\left[C_{\psi}^{-1} \int_{0}^{\infty} W_{\Phi}(t,s)s^{-1}ds\right].$$

For the proof of the second part of Proposition 2, because the Fourier transform of  $\psi$ , given in (14), satisfies Assumption 2, Proposition 1 implies that the real and imaginary parts of  $h_{\Phi}$  are continuously differentiable with respect to t and s. Furthermore, the Cauchy-Riemann condition holds:

(70) 
$$\frac{\partial h_{\Phi}}{\partial t} = -i\frac{\partial h_{\Phi}}{\partial s}.$$

The verification of (70) follows the same steps as in [17, Theorem 2]. Thus, the function  $h_{\Phi}$  is holomorphic.

6.5. **Proof of Theorem 2.** (a) Because the time variable is fixed in the statement of Theorem 2, we omit it in the following derivation. That is, we denote  $W_Y(s) := W_Y(t,s)$  and  $S_Y(s) := S_Y(t,s)$  for a given time point t. Next, we define the following matrices

$$\mathbf{W}_{f}(s) = \begin{bmatrix} W_{f}^{R}(s) \ W_{f}^{I}(s) \end{bmatrix}^{\top}, \mathbf{W}_{\Phi}(s) = \begin{bmatrix} W_{\Phi}^{R}(s) \ W_{\Phi}^{I}(s) \end{bmatrix}^{\top}, \mathbf{W}_{Y}(s) = \begin{bmatrix} W_{Y}^{R}(s) \ W_{Y}^{I}(s) \end{bmatrix}^{\top}, \\ \text{where } W_{f}^{R}(s) = \text{Re}(W_{f}(s)), \ W_{f}^{I}(s) = \text{Im}(W_{f}(s)), \text{ and similarly for other terms.} \\ \text{Clearly, } \mathbf{W}_{Y}(s) = \mathbf{W}_{f}(s) + \mathbf{W}_{\Phi}(s) \text{ and } S_{Y}(s) = \mathbf{W}_{Y}^{\top}(s)\mathbf{W}_{Y}(s). \text{ To prove the uniqueness of the global maximizer of } S_{Y}, \text{ it suffices to show that for any different } s_{1}, s_{2} > 0, \text{ there exist neighborhoods } N(s_{1}) \text{ and } N(s_{2}) \text{ of } s_{1} \text{ and } s_{2} \text{ such that } \end{cases}$$

(71) 
$$\mathbb{P}\left(\sup_{s\in N(s_1)}S_Y(s) = \sup_{s\in N(s_2)}S_Y(s)\right) = 0$$

because

$$\mathbb{P}\left(S_Y(s_1) = \sup_{s>0} S_Y(s) = S_Y(s_2) \text{ for some different } s_1, s_2 > 0\right)$$
$$\leq \mathbb{P}\left(\sup_{s \in N(s_1)} S_Y(s) = \sup_{s \in N(s_2)} S_Y(s) \text{ for some different } s_1, s_2 \in \mathbb{Q}_{>0}\right)$$
$$(72) \qquad \leq \sum_{\substack{s_1, s_2 \in \mathbb{Q}_{>0} \\ s_1 \neq s_2}} \mathbb{P}\left(\sup_{s \in N(s_1)} S_Y(s) = \sup_{s \in N(s_2)} S_Y(s)\right),$$

where  $\mathbb{Q}_{>0}$  represents the set of positive rational numbers, and the first inequality follows from the property that, almost surely, the sample path of  $S_Y$  is continuous, as demonstrated in Proposition 1.

Denote the covariance matrix function of the process  $\mathbf{W}_Y$  by  $\Gamma$ . Because  $\mathbb{E}[\mathbf{W}_Y(s)] = \mathbf{W}_f(s)$  for any s > 0,

$$\Gamma(s_1, s_2) = \mathbb{E}\left[\mathbf{W}_{\Phi}(s_1)\mathbf{W}_{\Phi}(s_2)^{\top}\right] = \left[\begin{array}{cc} \mathbb{E}\left[W_{\Phi}^R(s_1)W_{\Phi}^R(s_2)\right] & \mathbb{E}\left[W_{\Phi}^R(s_1)W_{\Phi}^I(s_2)\right] \\ \mathbb{E}\left[W_{\Phi}^I(s_1)W_{\Phi}^R(s_2)\right] & \mathbb{E}\left[W_{\Phi}^I(s_1)W_{\Phi}^I(s_2)\right] \end{array}\right].$$

By (43) and (52), for  $k \in \{1, 2\}$ ,

$$\Gamma(s_k, s_k) = \mathbb{E}\left[|W_{\Phi}^R(s_k)|^2\right] \mathbf{I}_{2 \times 2} = 2^{-1} \sqrt{\operatorname{Var}\left(S_{\Phi}(s_k)\right)} \mathbf{I}_{2 \times 2},$$

where  $\mathbf{I}_{2\times 2}$  is the 2 × 2 identity matrix. By the condition that the spectral density p is positive almost everywhere in Assumption 1, we have  $\operatorname{Var}(S_{\Phi}(s_1)) > 0$  and  $\operatorname{Var}(S_{\Phi}(s_2)) > 0$ . Without loss of generality, we assume that

(73) 
$$\operatorname{Var}\left(S_{\Phi}(s_1)\right) > \operatorname{Var}\left(S_{\Phi}(s_2)\right) > 0.$$

Denote

$$\frac{1}{\Gamma(s_1, s_1)} = \left\{ \mathbb{E}\left[ |W_{\Phi}^R(s_1)|^2 \right] \right\}^{-1}$$

and consider the following decomposition. For any s > 0,

$$\mathbf{W}_Y(s) = \mathbf{W}_{\perp}(s) + \frac{\Gamma(s, s_1)}{\Gamma(s_1, s_1)} \mathbf{W}_{\Phi}(s_1),$$

where

$$\mathbf{W}_{\perp}(s) = \mathbf{W}_{f}(s) + \mathbf{W}_{\Phi}(s) - \frac{\Gamma(s, s_{1})}{\Gamma(s_{1}, s_{1})} \mathbf{W}_{\Phi}(s_{1}).$$

Let  $\delta_1, \delta_2 > 0$  that will be determined later. For  $k \in \{1, 2\}$  and any  $\mathbf{z} = [z_1 \ z_2]^\top \in \mathbb{R}^2$ , we define

$$M_k(\mathbf{z}) = \sup_{s \in (s_k - \delta_k, s_k + \delta_k)} Q(\mathbf{z}; s),$$

where  $Q(\cdot; s)$  is a quadratic function of  $\mathbf{z}$  defined as follows

$$Q(\mathbf{z};s) = \left(\mathbf{W}_{\perp}(s) + \frac{\Gamma(s,s_1)}{\Gamma(s_1,s_1)}\mathbf{z}\right)^{\top} \left(\mathbf{W}_{\perp}(s) + \frac{\Gamma(s,s_1)}{\Gamma(s_1,s_1)}\mathbf{z}\right).$$

By noting that

$$M_k\left(\mathbf{W}_{\Phi}(s_1)\right) = \sup_{s \in (s_k - \delta_k, s_k + \delta_k)} Q(\mathbf{W}_{\Phi}(s_1); s) = \sup_{s \in (s_k - \delta_k, s_k + \delta_k)} S_Y(s),$$

the left-hand side of (71) with  $N(s_1) = (s_1 - \delta_1, s_1 + \delta_1)$  and  $N(s_2) = (s_2 - \delta_2, s_2 + \delta_2)$  can be rewritten as

$$\mathbb{P}\left(\sup_{s\in(s_1-\delta_1,s_1+\delta_1)}S_Y(s)=\sup_{s\in(s_2-\delta_2,s_2+\delta_2)}S_Y(s)\right)=\mathbb{E}\left[\mathbb{P}\left(M_1(\mathbf{W}_{\Phi}(s_1))=M_2(\mathbf{W}_{\Phi}(s_1))\mid\mathbf{W}_{\perp}\right)\right].$$

Thus, to prove (71), it suffices to show that conditioned on  $\mathbf{W}_{\perp}$ ,

(74) 
$$\mathbb{P}\left(M_1(\mathbf{W}_{\Phi}(s_1)) = M_2(\mathbf{W}_{\Phi}(s_1)) \mid \mathbf{W}_{\perp}\right) = 0.$$

Our strategy is to show that the Lebesgue measure of the set

$$\mathcal{I} := \{ \mathbf{z} \in \mathbb{R}^2 \mid M_1(\mathbf{z}) = M_2(\mathbf{z}) \}$$

is zero.

By a direct expansion,

$$Q(\mathbf{z};s) = \mathbf{W}_{\perp}^{\top}(s)\mathbf{W}_{\perp}(s) + 2\mathbf{W}_{\perp}^{\top}(s)\frac{\Gamma(s,s_1)}{\Gamma(s_1,s_1)}\mathbf{z} + \mathbf{z}^{\top}\frac{\Gamma(s,s_1)}{\Gamma(s_1,s_1)}\frac{\Gamma(s,s_1)}{\Gamma(s_1,s_1)}\mathbf{z}.$$

By (43) and (44),

$$\frac{\Gamma(s,s_1)^{\top}}{\Gamma(s_1,s_1)} \frac{\Gamma(s,s_1)}{\Gamma(s_1,s_1)} = \frac{\left(\mathbb{E}\left[W_{\Phi}^R(s)W_{\Phi}^R(s_1)\right]\right)^2 + \left(\mathbb{E}\left[W_{\Phi}^I(s)W_{\Phi}^R(s_1)\right]\right)^2}{\left(\mathbb{E}\left[W_{\Phi}^R(s_1)W_{\Phi}^R(s_1)\right]\right)^2} \mathbf{I}_{2\times 2}$$
$$= \frac{\operatorname{Cov}(S_{\Phi}(s), S_{\Phi}(s_1))}{\operatorname{Var}\left(S_{\Phi}(s_1)\right)} \mathbf{I}_{2\times 2},$$

where the second equality stems from Theorem 1(c). Therefore, for any s > 0,

$$Q(\mathbf{z};s) = \mathbf{W}_{\perp}^{\top}(s)\mathbf{W}_{\perp}(s) + 2\mathbf{W}_{\perp}^{\top}(s)\frac{\Gamma(s,s_1)}{\Gamma(s_1,s_1)}\mathbf{z} + \frac{\operatorname{Cov}(S_{\Phi}(s), S_{\Phi}(s_1))}{\operatorname{Var}(S_{\Phi}(s_1))}\|\mathbf{z}\|^2$$

where  $\|\mathbf{z}\|^2 = \mathbf{z}^\top \mathbf{z}$ . By the result Cov  $(S_{\Phi}(s_1), S_{\Phi}(s_2)) \ge 0$  given in Theorem 1(c), the Cauchy-Schwarz inequality, the condition in Assumption 1 that the spectral density p is positive almost everywhere, and (73), we have

(75) 
$$0 \le \operatorname{Cov}(S_{\Phi}(s_2), S_{\Phi}(s_1)) < \operatorname{Var}(S_{\Phi}(s_1)) = \operatorname{Cov}(S_{\Phi}(s_1), S_{\Phi}(s_1)).$$

The continuity of the function  $s \mapsto \operatorname{Cov}(S_{\Phi}(s), S_{\Phi}(s_1))$  and (75) ensure that there exist constants  $c_1 > c_2 > 0$  and sufficiently small  $\delta_1, \delta_2 > 0$  such that the quadratic coefficients of  $\{Q(\cdot; s) \mid s \in (s_1 - \delta_1, s_1 + \delta_1)\}$  and  $\{Q(\cdot; s) \mid s \in (s_2 - \delta_2, s_2 + \delta_2)\}$  can be separated as follows

(76) 
$$\sup_{s \in (s_1 - \delta_1, s_1 + \delta_1)} \frac{\operatorname{Cov}(S_{\Phi}(s, s_1))}{\operatorname{Var}(S_{\Phi}(s_1))} > c_1 > c_2 > \sup_{s \in (s_2 - \delta_2, s_2 + \delta_2)} \frac{\operatorname{Cov}(S_{\Phi}(s, s_1))}{\operatorname{Var}(S_{\Phi}(s_1))}.$$

Furthermore, for any fixed s > 0, the graph of  $Q(\cdot, s)$  can be categorized as follows:

- For s satisfying  $\operatorname{Cov}(S_{\Phi}(s), S_{\Phi}(s_1)) > 0$ , the surface  $\{(\mathbf{z}, Q(\mathbf{z}, s)) \mid \mathbf{z} \in \mathbb{R}^2\}$  is an upward-opening paraboloid;
- For s satisfying  $\operatorname{Cov}(S_{\Phi}(s), S_{\Phi}(s_1)) = 0$ , by Theorem 1(c), we have  $\mathbb{E}\left[W_{\Phi}^R(s)W_{\Phi}^R(s_1)\right] = 0$  and  $\mathbb{E}\left[W_{\Phi}^R(s)W_{\Phi}^I(s_1)\right] = 0$ , which imply that  $\Gamma(s, s_1) = \mathbf{0}_{2\times 2}$  and  $Q(\mathbf{z}; s) = \mathbf{W}_{\perp}^T(s)\mathbf{W}_{\perp}(s)$  for any  $\mathbf{z} \in \mathbb{R}^2$ . This indicates that  $\{(\mathbf{z}, Q(\mathbf{z}, s)) \mid \mathbf{z} \in \mathbb{R}^2\}$  is a horizontal plane.

By the fact that the supremum of a family of convex functions is still convex, both  $M_1$  and  $M_2$  are convex functions defined on  $\mathbb{R}^2$ . According to Alexandrov's theorem [34, Theorem 2.1],  $M_1$  and  $M_2$  have the second derivatives almost everywhere. More precisely, there exists a set  $D \subset \mathbb{R}^2$ , whose complement has Lebesgue measure zero, such that for  $k \in \{1, 2\}$  and every point  $\mathbf{z} \in D$ , there is a quadratic expansion in the form

(77) 
$$M_k(\mathbf{z} + \mathbf{w}) = M_k(\mathbf{z}) + \mathbf{v}_k(\mathbf{z})^\top \mathbf{w} + \frac{1}{2} \mathbf{w}^\top \mathbf{H}_k(\mathbf{z}) \mathbf{w} + o(\|\mathbf{w}\|^2), \ \mathbf{w} \in \mathbb{R}^2,$$

where  $\mathbf{v}_k(\mathbf{z})$  is the gradient, and  $\mathbf{H}_k(\mathbf{z})$  is the Hessian matrix of  $M_k$  at  $\mathbf{z}$ . The function  $o(\|\mathbf{w}\|^2)$  satisfies  $o(\|\mathbf{w}\|^2) \|\mathbf{w}\|^{-2} \to 0$  as  $\|\mathbf{w}\| \to 0$ . By (76) (see also [35, Example 13.10]),

(78) 
$$\frac{1}{2}\mathbf{w}^{\top}\mathbf{H}_{1}(\mathbf{z})\mathbf{w} \geq c_{1}\|\mathbf{w}\|^{2} > c_{2}\|\mathbf{w}\|^{2} \geq \frac{1}{2}\mathbf{w}^{\top}\mathbf{H}_{2}(\mathbf{z})\mathbf{w},$$

which implies that

(79) 
$$\operatorname{Leb}\left(\mathcal{I}\right) = 0.$$

where Leb represents the Lebesgue measure on  $\mathbb{R}^2$ . For a further explanation of (79), see Remark 2 at the end of this section. From the covariance computation, the normal random vector  $\mathbf{W}_{\Phi}(s_1)$  is independent of the Gaussian random process  $\mathbf{W}_{\perp}$ . Thus, conditioning on  $\mathbf{W}_{\perp}$ , the probability density of  $\mathbf{W}_{\Phi}(s_1)$  remains absolutely continuous with respect to the Lebesgue measure on  $\mathbb{R}^2$ . Therefore, the observation (79) implies that

$$\mathbb{P}\left(M_1\left(\mathbf{W}_{\Phi}(s_1)\right) = M_2\left(\mathbf{W}_{\Phi}(s_1)\right) \mid \mathbf{W}_{\perp}\right) = \mathbb{P}\left(\mathbf{W}_{\Phi}(s_1) \in \mathcal{I} \mid \mathbf{W}_{\perp}\right) = 0,$$

which establishes (74).

(b) Regarding the second part of Theorem 2, similar to (72), for any  $t \in \mathbb{R}$  and  $m \in \{1, \ldots, M\}$ , we have

 $\mathbb{P}(s_{Y,m}(t) \text{ contains at least two scale points})$ 

$$= \mathbb{P}\left(S_Y(t,s_1) = \sup_{s \in B_m(t)} S_Y(t,s) = S_Y(t,s_2) \text{ for some different } s_1, s_2 \in B_m(t)\right)$$
  
$$\leq \mathbb{P}\left(\sup_{s \in N(s_1) \cap B_m(t)} S_Y(t,s) = \sup_{s \in N(s_2) \cap B_m(t)} S_Y(t,s) \text{ for some different } s_1, s_2 \in \mathbb{Q} \cap B_m(t)\right)$$
  
$$\leq \sum_{\substack{s_1, s_2 \in \mathbb{Q} \cap B_m(t) \\ s_1 \neq s_2}} \mathbb{P}\left(\sup_{s \in N(s_1) \cap B_m(t)} S_Y(t,s) = \sup_{s \in N(s_2) \cap B_m(t)} S_Y(t,s)\right).$$

Thus, to prove the second part of Theorem 2, it suffices to show that, for any different  $s_1, s_2 \in \mathbb{Q} \cap B_m(t)$ , there exist neighborhoods  $N(s_1)$  and  $N(s_2)$  of  $s_1$  and  $s_2$ , respectively, such that

(80) 
$$\mathbb{P}\left(\sup_{s\in N(s_1)\cap B_m(t)}S_Y(t,s) = \sup_{s\in N(s_2)\cap B_m(t)}S_Y(t,s)\right) = 0.$$

The proof of (80) is similar to that of (71). Therefore, we omit the details here.  $\Box$ 

**Remark 2.** Suppose that (79) does not hold. Then,  $\text{Leb}(\mathcal{I} \cap D) > 0$ . By the Lebesgue density theorem, there exists  $\mathbf{z} \in \mathcal{I} \cap D$  such that for any  $\varepsilon > 0$ ,

$$\operatorname{Leb}\left(\mathcal{I}\cap D\cap B_{\varepsilon}(\mathbf{z})\right)>0,$$

where  $B_{\varepsilon}(\mathbf{z})$  is the disk centered at  $\mathbf{z}$  and with radius  $\varepsilon$ . For any  $\varepsilon > 0$  and  $\mathbf{w} \in \mathbb{R}^2$ satisfying  $\mathbf{z} + \mathbf{w} \in \mathcal{I} \cap B_{\varepsilon}(\mathbf{z})$ , because  $M_1(\mathbf{z} + \mathbf{w}) = M_2(\mathbf{z} + \mathbf{w})$ , (77) implies that

(81) 
$$(\mathbf{v}_1(\mathbf{z}) - \mathbf{v}_2(\mathbf{z}))^\top \mathbf{w} + \frac{1}{2} \mathbf{w}^\top (\mathbf{H}_1(\mathbf{z}) - \mathbf{H}_2(\mathbf{z})) \mathbf{w} = o(\|\mathbf{w}\|^2).$$

For the left-hand side of (81), from (78),

$$(\mathbf{v}_1(\mathbf{z}) - \mathbf{v}_2(\mathbf{z}))^\top \mathbf{w} + \frac{1}{2} \mathbf{w}^\top (\mathbf{H}_1(\mathbf{z}) - \mathbf{H}_2(\mathbf{z})) \mathbf{w}$$
  
 
$$\geq (\mathbf{v}_1(\mathbf{z}) - \mathbf{v}_2(\mathbf{z}))^\top \mathbf{w} + (c_1 - c_2) \|\mathbf{w}\|^2,$$

which contradicts the right-hand side of (81).

30

6.6. **Proof of Theorem 3.** Let  $\Omega$  be the space of random elements. For any  $\omega \in \Omega$ , we denote the associated sample path of the random process  $\Phi$  by  $\{\Phi(t;\omega) \mid t \in \mathbb{R}\}$ . Besides, to emphasize the dependence of the scalogram of Y on the sample path of  $\Phi$ , we denote the scalogram by  $S_Y(\cdot, \cdot; \omega)$  for the noisy signal  $Y(t;\omega) = f(t) + \Phi(t;\omega)$ . Proposition 1 shows that there exists a subset  $\widetilde{\Omega}$  of  $\Omega$  with  $\mathbb{P}(\widetilde{\Omega}) = 1$  such that for any  $\omega \in \widetilde{\Omega}$ ,  $S_Y(\cdot, \cdot; \omega)$  is two times continuously differentiable on  $\mathbb{R} \times (0, \infty)$ .

In the following, we focus on the case where  $\omega \in \tilde{\Omega}$ . In order to show that the set-valued function  $t \in \mathbb{R} \mapsto s_Y(t; \omega) \subseteq [0, \infty)$  is upper hemicontinuous, we classify time points into two groups:

$$\mathcal{M} := \{ t \in \mathbb{R} \mid S_Y(t, s; \omega) > 0 \text{ for a certain } s > 0 \}$$

and

$$\mathcal{N} := \left\{ t \in \mathbb{R} \mid S_Y(t, s; \omega) = 0 \text{ for all } s > 0 \right\}.$$

For any  $t_0 \in \mathcal{N}$ , it is clear that  $s_Y(t_0; \omega) = [0, \infty)$ . Thus,  $s_Y(t'; \omega) \subseteq s_Y(t_0; \omega)$  for any time point t'. This directly implies that  $s_Y(\cdot; \omega)$  is upper hemicontinuous at every time point in  $\mathcal{N}$ . It remains to prove that  $s_Y(\cdot; \omega)$  is also upper hemicontinuous at every  $t_0 \in \mathcal{M}$ . The proof is structured as follows.

Step 1. We begin by showing that there exist an interval [a, b], whose interior contains the time point  $t_0$ , and a threshold  $S(\omega) < \infty$  such that

(82) 
$$s_Y(t;\omega) = \arg\max_{s>0} S_Y(t,s;\omega) \subseteq [0,\mathcal{S}(\omega)]$$

for all  $t \in [a, b]$ . Because  $S_Y(t_0, s_0; \omega) > 0$  for some  $s_0 > 0$  and  $S_Y(\cdot, \cdot; \omega)$  is continuous on  $\mathbb{R} \times (0, \infty)$ , there exist  $\delta > 0$  and an interval [a, b] with  $a < t_0 < b$  such that

(83) 
$$S_Y(\cdot, \cdot; \omega) > \frac{S_Y(t_0, s_0; \omega)}{2} \text{ on } [a, b] \times [s_0 - \delta, s_0 + \delta].$$

We now fix the interval [a, b] and proceed to prove the finiteness of the threshold  $\mathcal{S}(\omega)$ . Assume, for the sake of contradiction, that no such threshold exists. This would imply that for any  $n \in \mathbb{N}$ , there exist  $t_n \in [a, b]$  and  $s_n \geq n$  such that  $s_n \in s_Y(t_n; \omega)$ . That is,

(84) 
$$S_Y(t_n, s_n; \omega) = \sup_{s>0} S_Y(t_n, s; \omega).$$

To establish the impossibility of this scenario, we define a function  $\mathcal{R}$  on  $[a, b] \times [0, 1]$  as follows

(85) 
$$\mathcal{R}(t, u; \omega) = \begin{cases} S_Y(t, 1/u; \omega) & \text{if } u \in (0, 1], \\ 0 & \text{if } u = 0. \end{cases}$$

Because  $S_Y$  is continuous on  $\mathbb{R} \times (0, \infty)$  and  $\mathbb{P}(\lim_{s \to \infty} S_Y(t, s) = 0) = 1$  for any  $t \in \mathbb{R}$ , as shown in Corollary 1, the function  $\mathcal{R}$  is continuous on  $[a, b] \times [0, 1]$ . Because  $\{t_n\}_{n \in \mathbb{N}} \subset [a, b]$ , there exists a convergent subsequence of  $\{t_n\}_{n \in \mathbb{N}}$  which converges to some time point  $t^* \in [a, b]$ . For simplicity of notation, assume that  $t_n \to t^*$  as  $n \to \infty$ . Because the pointwise supremum of any collection of continuous functions is lower semi-continuous, we have

$$\sup_{s>0} S_Y(t^*, s; \omega) \leq \liminf_{n \to \infty} \sup_{s>0} S_Y(t_n, s; \omega)$$

$$\stackrel{(84)}{=} \liminf_{n \to \infty} S_Y(t_n, s_n; \omega)$$

$$\stackrel{(85)}{=} \liminf_{n \to \infty} \mathcal{R}(t_n, 1/s_n; \omega)$$

$$= \mathcal{R}(t^*, 0; \omega) = 0.$$

The derivation above implies that  $S_Y(t^*, s; \omega) = 0$  for any s > 0, which contradicts (83). Hence, (82) holds.

Step 2. By (82), for any  $t \in [a, b]$ ,

$$s_Y(t;\omega) = \arg\max_{s>0} S_Y(t,s;\omega) = \arg\max_{s\in[0,\mathcal{S}(\omega)]} S_Y(t,s;\omega).$$

The continuity of  $S_Y(t, s; \omega)$  and the compactness of  $[0, \mathcal{S}(\omega)]$  allow us to apply Berge's maximum theorem [18, Theorem 3.4] (see also [1, p. 570] or Lemma 5 in the appendix), implying that the set-valued function  $s_Y(\cdot; \omega)$  is upper hemicontinuous at every time point  $t_0 \in \mathcal{M}$ .

6.7. **Proof of Corollary 2.** Using the notation introduced at the beginning of Section 6.6, Theorems 2 and 3 show that there exists a subset  $\widetilde{\Omega}$  of  $\Omega$  with  $\mathbb{P}(\widetilde{\Omega}) = 1$  such that for any  $\omega \in \widetilde{\Omega}$ ,

- $S_Y(\cdot, \cdot; \omega)$  is two times continuously differentiable on  $\mathbb{R} \times (0, \infty)$ ,
- $s_Y(t;\omega)$  is a singleton for every  $t \in \mathbb{Q}$ , and
- $t \mapsto s_Y(t; \omega)$  is upper hemicontinuous on  $\mathbb{R}$ .

In the following, we focus on the case where  $\omega \in \tilde{\Omega}$  and consider an arbitrary point  $t_0 \in \mathbb{Q}$ . The proof of Corollary 2 is organized as follows.

Step 1. According to the definition of upper hemicontinuity in Theorem 3, for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

(86) 
$$s_Y(t';\omega) \subset (s_Y(t_0;\omega) - \varepsilon, s_Y(t_0;\omega) + \varepsilon)$$

for all  $t' \in (t_0 - \delta, t_0 + \delta)$ . The inclusion relation (86) implies the continuity of the restriction of  $s_Y$  on  $\mathbb{Q}$ .

Step 2: We aim to demonstrate the existence of a continuously differentiable function r defined on  $(t_0 - \delta, t_0 + \delta)$  for some  $\delta > 0$  such that  $r(t_0) = s_Y(t_0; \omega)$  and

(87) 
$$\frac{\partial S_Y}{\partial s}(t, r(t)) = 0$$

for any  $t \in (t_0 - \delta, t_0 + \delta)$ . Because the function  $s \mapsto S_Y(t, s; \omega)$  reaches its global maximum at only  $s_Y(t_0; \omega)$ ,

(88) 
$$\frac{\partial S_Y}{\partial s}(t_0, s_Y(t_0); \omega) = 0.$$

On the other hand, if the assumption (22) holds, then

(89) 
$$\frac{\partial^2 S_Y}{\partial s^2}(t_0, s_Y(t_0); \omega) \neq 0$$

for almost every  $\omega \in \widetilde{\Omega}$ . By (88) and (89), the implicit function theorem guarantees the existence of the function r and the positive constant  $\delta$ .

Step 3: We want to show that the function r determined by the implicit function theorem coincides with  $s_Y(\cdot; \omega)$  on rational numbers in a neighborhood of  $t_0$ . That is, there exists a constant  $\delta' > 0$  with  $\delta' \leq \delta$  such that

(90) 
$$r(t) = s_Y(t;\omega) \text{ for any } t \in (t_0 - \delta', t_0 + \delta') \cap \mathbb{Q}.$$

Suppose not. Then, (a) there exists a sequence  $\{q_n\}_{n\in\mathbb{N}}\subset (t_0-\delta',t_0+\delta')\cap\mathbb{Q}\setminus\{t_0\}$ such that  $q_n \to t_0$  as  $n \to \infty$ , (b)  $r(q_n) \neq s_Y(q_n;\omega)$  for any  $n \in \mathbb{N}$ . For the condition (b), note that if (90) does not hold for any  $\delta'$ , then there exists  $q_1 \in$  $(t_0 - \delta', t_0 + \delta') \cap \mathbb{Q} \setminus \{t_0\}$  such that  $r(q_1) \neq s_Y(q_1;\omega)$ . We can iteratively choose  $q_n \in (t_0 - \delta', t_0 + \delta') \cap \mathbb{Q} \setminus \{t_0\}$  from the intervals  $(q_{n-1}, t_0)$  or  $(t_0, q_{n-1})$  for n > 1such that  $r(q_n) \neq s_Y(q_n;\omega)$ . Because

- the continuity of r implies that  $r(q_n) \to r(t_0)$  as  $n \to \infty$ ,
- the continuity of  $s_Y(\cdot; \omega)$  on  $\mathbb{Q}$  implies that  $s_Y(q_n; \omega) \to s_Y(t_0; \omega)$  as  $n \to \infty$ , and
- $s_Y(t_0;\omega) = r(t_0),$

we obtain that  $r(q_n) - s_Y(q_n; \omega) \to 0$  and

(91)  

$$\left[\frac{\partial S_Y}{\partial s}(q_n, r(q_n); \omega) - \frac{\partial S_Y}{\partial s}(q_n, s_Y(q_n); \omega)\right] [r(q_n) - s_Y(q_n)]^{-1} \to \frac{\partial^2 S_Y}{\partial s^2}(t_0, s_Y(t_0); \omega)$$

as  $n \to \infty$ . Here, we used the two times continuous differentiability of  $S_Y(\cdot, \cdot; \omega)$  to obtain the limit in (91). However, (87) and (88) imply that the left-hand side of (91) is equal to zero. This leads to

$$\frac{\partial^2 S_Y}{\partial s^2}(t_0, s_Y(t_0); \omega) = 0,$$

which contradicts (89). Therefore, (90) holds.

Step 4: We finally demonstrate that

(92) 
$$r(t) \in s_Y(t;\omega) \text{ for } t \in (t_0 - \delta', t_0 + \delta') \cap \mathbb{Q}^c.$$

Indeed, for any  $t \in (t_0 - \delta', t_0 + \delta') \cap \mathbb{Q}^c$ , there exists a sequence  $\{q_n\}_{n \in \mathbb{N}} \subset (t_0 - \delta', t_0 + \delta') \cap \mathbb{Q}$  such that  $q_n \to t$  as  $n \to \infty$ . For any s > 0,

(93)  

$$S_{Y}(t, r(t); \omega) = \lim_{n \to \infty} S_{Y}(q_{n}, r(q_{n}); \omega)$$

$$\geq \lim_{n \to \infty} S_{Y}(q_{n}, s; \omega)$$

$$= S_{Y}(t, s; \omega),$$

where the equalities follow from the continuity of r on the interval  $(t_0 - \delta', t_0 + \delta')$ and the continuity of  $S_Y(\cdot, \cdot; \omega)$ . The observation (93) implies that

$$S_Y(t, r(t); \omega) \ge \sup_{s>0} S_Y(t, s; \omega).$$

Hence, (92) holds.

6.8. **Proof of Theorem 4.** By the definition of  $s_Y(t)$  in (20), for any interval I containing  $s_f(t)$ , we have

$$\mathbb{P}(s_{Y}(t) \in I) = \mathbb{P}\left(\max_{s \in I} |W_{Y}(t,s)| > \max_{s \in I^{c}} |W_{Y}(t,s)|\right)$$
  

$$\geq \mathbb{P}\left(|W_{f}(t,s_{f}(t))| - |W_{\Phi}(t,s_{f}(t))| > \max_{s \in I^{c}} |W_{f}(t,s)| + \max_{s \in I^{c}} |W_{\Phi}(t,s)|\right)$$
  

$$= \mathbb{P}\left(|W_{f}(t,s_{f}(t))| - \max_{s \in I^{c}} |W_{f}(t,s)| > |W_{\Phi}(t,s_{f}(t))| + \max_{s \in I^{c}} |W_{\Phi}(t,s)|\right)$$
  

$$= \mathbb{P}\left(\bigtriangleup_{I} > |W_{\Phi}(t,s_{f}(t))| + \max_{s \in I^{c}} |W_{\Phi}(t,s)|\right),$$

where  $\Delta_I = |W_f(t, s_f(t))| - \max_{s \in I^c} |W_f(t, s)|$ . Hence, to prove (25), it suffices to show that

(94) 
$$\mathbb{P}\left(|W_{\Phi}(t,s_f(t))| + \max_{s \in I^c} |W_{\Phi}(t,s)| \ge \Delta_I\right) \le \exp\left(-\frac{(\Delta_I - \mu_I)^2}{\sigma_I^2}\right),$$

where  $\mu_I$  and  $\sigma_I$  are defined in (23) and (24). The proof of (94) consists of two steps.

Step 1: Establish a discrete version of (94). For any  $d \in \mathbb{N}$  and  $\mathbf{x} = [x_1 \ x_2 \ \cdots \ x_d]^\top \in \mathbb{C}^d$ , we define

$$g(\mathbf{x}) = |x_1| + \max_{2 \le \ell \le d} |x_\ell|,$$

which is Lipschitz continuous. For any  $s_1, s_2, \ldots, s_d > 0$ , let

$$\mathbf{J} = [W_{\Phi}(t, s_1) \ W_{\Phi}(t, s_2) \ \cdots \ W_{\Phi}(t, s_d)]^{\top} \in \mathbb{C}^d.$$

Because the wavelet  $\psi$  is assumed to be analytic, **J** is a circularly symmetric complex Gaussian random vector, as shown in Theorem 1(a). By Lemma 1, for any  $u > \mathbb{E}[g(\mathbf{J})]$ ,

(95)

$$\mathbb{P}\left(|W_{\Phi}(t,s_1)| + \max_{2 \le \ell \le d} |W_{\Phi}(t,s_\ell)| > u\right) = \mathbb{P}\left(g(\mathbf{J}) > u\right) \le \exp\left(-\frac{(u - \mathbb{E}[g(\mathbf{J})])^2}{2\|g(\mathbf{A} \bullet)\|_{\operatorname{Lip}}^2}\right),$$

where  $\mathbf{A} \in \mathbb{C}^{d \times d}$  satisfying

(96) 
$$\mathbf{A}\mathbf{A}^* = 2^{-1}\mathbb{E}\left[\mathbf{J}\mathbf{J}^*\right].$$

About  $||g(\mathbf{A}\bullet)||_{\text{Lip}}$ , for any  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^d$  with  $\mathbf{z} = \mathbf{x} - \mathbf{y}$ ,

(97)  

$$|g(\mathbf{A}\mathbf{x}) - g(\mathbf{A}\mathbf{y})| \leq |e_{1}^{\top}\mathbf{A}\mathbf{z}| + \max_{2\leq\ell\leq d} |e_{\ell}^{\top}\mathbf{A}\mathbf{z}|$$

$$\leq |e_{1}^{\top}\mathbf{A}||\mathbf{z}| + \max_{2\leq\ell\leq d} |e_{\ell}^{\top}\mathbf{A}||\mathbf{z}|$$

$$= \left(\sqrt{e_{1}^{\top}\mathbf{A}\mathbf{A}^{*}e_{1}} + \max_{2\leq\ell\leq d}\sqrt{e_{\ell}^{\top}\mathbf{A}\mathbf{A}^{*}e_{\ell}}\right)|\mathbf{z}|,$$

where  $e_{\ell}$  is the column vector in  $\mathbb{R}^d$  with one in position  $\ell$  and zeros elsewhere. From (96), for  $\ell \in \{1, 2, \ldots, d\}$ ,

(98) 
$$e_{\ell}^{\top} \mathbf{A} \mathbf{A}^* e_{\ell} = 2^{-1} \mathbb{E} \left[ |W_{\Phi}(t, s_{\ell})|^2 \right].$$

Combining (97) and (98) yields

(99) 
$$||g(\mathbf{A}\bullet)||_{\text{Lip}} \le 2^{-\frac{1}{2}} \left\{ \sqrt{\mathbb{E}\left[|W_{\Phi}(t,s_1)|^2\right]} + \max_{2 \le \ell \le d} \sqrt{\mathbb{E}\left[|W_{\Phi}(t,s_\ell)|^2\right]} \right\}$$

Step 2: Extend the discrete version to the continuous case. By setting  $s_1 = s_f(t)$ and defining  $T_d = \{s_2, \ldots, s_d\}$  as a subset of  $I^c$  such that  $T_d \subset T_{d+1}$  and  $T_d$ increases to a dense subset of  $I^c$ , the inequality (95) becomes

$$\mathbb{P}\left(|W_{\Phi}(t,s_f(t))| + \max_{s \in T_d} |W_{\Phi}(t,s)| > u\right) \le \exp\left(-\frac{(u - \mathbb{E}[g(\mathbf{J})])^2}{2\|g(\mathbf{A} \bullet)\|_{\operatorname{Lip}}^2}\right)$$

As d increases to infinity, the continuity of the sample paths of  $W_{\Phi}$  given in Proposition 1, implies that

$$|W_{\Phi}(t, s_f(t))| + \max_{s \in T_d} |W_{\Phi}(t, s)| \uparrow |W_{\Phi}(t, s_f(t))| + \max_{s \in I^c} |W_{\Phi}(t, s)|$$

almost surely, the upper bound of  $||g(\mathbf{A}\bullet)||_{\text{Lip}}$  in (99) increases to

$$2^{-\frac{1}{2}} \left\{ \sqrt{\mathbb{E}\left[ |W_{\Phi}(t, s_f(t))|^2 \right]} + \max_{s \in I^c} \sqrt{\mathbb{E}\left[ |W_{\Phi}(t, s)|^2 \right]} \right\}$$

by the continuity of  $s \mapsto \mathbb{E}\left[|W_{\Phi}(t,s)|^2\right]$ , and

(100) 
$$\mathbb{E}[g(\mathbf{J})] \uparrow \mathbb{E}\left[|W_{\Phi}(t, s_f(t))|\right] + \mathbb{E}\left[\max_{s \in I^c} |W_{\Phi}(t, s)|\right],$$

where the finiteness of the expectation in (100) is guaranteed by Lemma 3.

6.9. **Proof of Theorem 5.** Step 1. For any  $m \in \{1, 2, ..., M\}$ , we condition on the event  $s_{Y,m}(t) \in B_m^{\circ}(t) \subset (\underline{i}_m(t), \overline{i}_m(t))$ . Under this condition, we derive the following formula:

$$s_{Y,m}(t) - s_{f,m}(t) = -\left[\frac{\partial^2 S_f}{\partial s^2}(t, c(t))\right]^{-1} \frac{\partial}{\partial s} \left[W_f(t, s)\overline{W_{\Phi}(t, s)} + \overline{W_f(t, s)}W_{\Phi}(t, s)\right]\Big|_{s=s_{Y,m}(t)}$$

$$(101) \qquad -\left[\frac{\partial^2 S_f}{\partial s^2}(t, c(t))\right]^{-1} \frac{\partial S_{\Phi}}{\partial s}(t, s_{Y,m}(t)),$$

where c(t) lies between  $s_{f,m}(t)$  and  $s_{Y,m}(t)$ .

Indeed, from the following relationship among the scalograms of the clean signal f, the noisy signal Y, and the noise  $\Phi$ :

$$S_Y(t,s) = |W_f(t,s) + W_{\Phi}(t,s)|^2$$
$$= S_f(t,s) + W_f(t,s)\overline{W_{\Phi}(t,s)} + \overline{W_f(t,s)}W_{\Phi}(t,s) + S_{\Phi}(t,s),$$

we have

$$(102) \\ \frac{\partial S_Y}{\partial s}(t,s) = \frac{\partial S_f}{\partial s}(t,s) + \frac{\partial}{\partial s} \left[ W_f(t,s)\overline{W_\Phi(t,s)} + \overline{W_f(t,s)}W_\Phi(t,s) \right] + \frac{\partial S_\Phi}{\partial s}(t,s)$$

Because Theorem 2 shows that  $s_{Y,m}(t)$  is a singleton almost surely, the condition  $s_{Y,m}(t) \in B_m^{\circ}(t)$  implies that

$$\frac{\partial S_Y}{\partial s}(t, s_{Y,m}(t)) = 0.$$

,s).

By substituting  $s = s_{Y,m}(t)$  into (102),

$$(103)$$

$$0 = \frac{\partial S_f}{\partial s}(t, s_{Y,m}(t)) + \frac{\partial}{\partial s} \left[ W_f(t, s) \overline{W_{\Phi}(t, s)} + \overline{W_f(t, s)} W_{\Phi}(t, s) \right] + \frac{\partial S_{\Phi}}{\partial s}(t, s) \Big|_{s=s_{Y,m}(t)}$$

From the condition (R1) in Definition 1, we have

$$\frac{\partial S_f}{\partial s}(t, s_{f,m}(t)) = 0.$$

By applying the mean value theorem to the first term on the right-hand side of (102), there exists a random variable c(t) between  $s_{f,m}(t)$  and  $s_{Y,m}(t)$  such that

$$\frac{\partial^2 S_f}{\partial s^2}(t,c(t)) \left(s_{Y,m}(t) - s_{f,m}(t)\right)$$

$$(104) = -\frac{\partial}{\partial s} \left[ W_f(t,s) \overline{W_{\Phi}(t,s)} + \overline{W_f(t,s)} W_{\Phi}(t,s) \right] \Big|_{s=s_{Y,m}(t)} - \frac{\partial S_{\Phi}}{\partial s}(t,s_{Y,m}(t)).$$

Since the condition  $B_m^{\circ}(t) \subset (\underline{\mathfrak{i}}_m(t), \overline{\mathfrak{i}}_m(t))$  implies that

$$\frac{\partial^2 S_f}{\partial s^2}(t, c(t)) < 0,$$

we obtain (101) from (104).

Step 2. We derive an upper bound for  $|s_{Y,m}(t) - s_{f,m}(t)|$ . Let  $L_m$  be the constant defined in (28). On the event  $\{s_Y(t) \in B_m^\circ(t)\},\$ 

$$\left. \frac{\partial^2 S_f}{\partial s^2}(t, c(t)) \right| \ge L_m.$$

Hence, (101) implies that

(105)

$$\begin{split} &L_{m}|s_{Y,m}(t) - s_{f,m}(t)| \\ \leq \left| \frac{\partial}{\partial s} \left[ W_{f}(t,s) \overline{W_{\Phi}(t,s)} + \overline{W_{f}(t,s)} W_{\Phi}(t,s) \right] \right|_{s=s_{Y,m}(t)} \right| + \left| \frac{\partial S_{\Phi}}{\partial s}(t,s_{Y,m}(t)) \right| \\ \leq \left| \frac{\partial W_{f}}{\partial s}(t,s_{Y,m}(t)) \right| \left| W_{\Phi}(t,s_{Y,m}(t)) \right| + \left| W_{f}(t,s_{Y,m}(t)) \right| \left| \frac{\partial W_{\Phi}}{\partial s}(t,s_{Y,m}(t)) \right| \\ + \left| \frac{\partial W_{f}}{\partial s}(t,s_{Y,m}) \right| \left| W_{\Phi}(t,s_{Y,m}(t)) \right| + \left| W_{f}(t,s_{Y,m}(t)) \right| \left| \frac{\partial W_{\Phi}}{\partial s}(t,s_{Y,m}(t)) \right| + \left| \frac{\partial S_{\Phi}}{\partial s}(t,s_{Y,m}(t)) \right| \\ = 2 \left| \frac{\partial W_{f}}{\partial s}(t,s_{Y,m}(t)) \right| \left| W_{\Phi}(t,s_{Y,m}(t)) \right| + 2 \left| W_{f}(t,s_{Y,m}(t)) \right| \left| \frac{\partial W_{\Phi}}{\partial s}(t,s_{Y,m}(t)) \right| + \left| \frac{\partial S_{\Phi}}{\partial s}(t,s_{Y,m}(t)) \right| \\ \end{split}$$

Under the condition  $B_m(t) \subset (\underline{i}_m(t), \overline{i}_m(t)), L_m > 0$ . By (105), for any  $\varepsilon > 0$ ,

$$\mathbb{P}\left(|s_{Y,m}(t) - s_{f,m}(t)| > \varepsilon \text{ and } s_{Y,m}(t) \in B_{m}^{\circ}(t)\right)$$

$$\leq \mathbb{P}\left(\left|\frac{\partial W_{f}}{\partial s}(t, s_{Y,m}(t))W_{\Phi}(t, s_{Y,m}(t))\right| > \frac{\varepsilon L_{m}}{6} \text{ and } s_{Y,m} \in B_{m}^{\circ}(t)\right)$$

$$+ \mathbb{P}\left(\left|W_{f}(t, s_{Y,m})\frac{\partial W_{\Phi}}{\partial s}(t, s_{Y,m})\right| > \frac{\varepsilon L_{m}}{6} \text{ and } s_{Y,m}(t) \in B_{m}^{\circ}(t)\right)$$

$$\left(106\right) \qquad + \mathbb{P}\left(\left|\frac{\partial S_{\Phi}}{\partial s}(t, s_{Y,m})\right| > \frac{\varepsilon L_{m}}{3} \text{ and } s_{Y,m}(t) \in B_{m}^{\circ}(t)\right).$$

36

For the first term on the right-hand side of (106), by the first part of Lemma 4, we have

$$\left\| \mathbb{P}\left( \left| \frac{\partial W_f}{\partial s}(t, s_{Y,m}(t)) W_{\Phi}(t, s_{Y,m}(t)) \right| > \frac{\varepsilon L_m}{6} \text{ and } s_{Y,m}(t) \in B_m^{\circ}(t) \right) \\ \leq \mathbb{P}\left( \max_{s \in B_m(t)} \left| \frac{\partial W_f}{\partial s}(t, s) \right| |W_{\Phi}(t, s)| > \frac{\varepsilon L_m}{6} \right) \\ (107) \qquad \leq \exp\left[ -\frac{1}{\sigma_{m,1}^2} \left( \frac{\varepsilon L_m}{6} - \mu_{m,1} \right)^2 \right]$$

for any  $\varepsilon > 6\mu_{m,1}L_m^{-1}$ , where  $\mu_{m,1}$  and  $\sigma_{m,1}^2$  are defined in (27). For the second term on the right-hand side of (106), by the second part of Lemma 4, we have

$$\left\| \left( \left| W_f(t, s_{Y,m}(t)) \frac{\partial W_{\Phi}}{\partial s}(t, s_{Y,m}(t)) \right| > \frac{\varepsilon L_m}{6} \text{ and } s_{Y,m}(t) \in B_m^{\circ}(t) \right) \\ \leq \left\| \left( \max_{s \in B_m(t)} |W_f(t, s)| \left| \frac{\partial W_{\Phi}}{\partial s}(t, s) \right| > \frac{\varepsilon L_m}{6} \right) \\ (108) \qquad \leq \exp\left[ -\frac{1}{\sigma_{m,2}^2} \left( \frac{\varepsilon L_m}{6} - \mu_{m,2} \right)^2 \right]$$

for any  $\varepsilon > 6\mu_2 L_m^{-1}$ , where  $\mu_{m,2}$  and  $\sigma_{m,2}^2$  are defined in (27). For the third term in (106), because

$$\begin{aligned} \frac{\partial S_{\Phi}}{\partial s}(t,s) & = \left| \frac{\partial}{\partial s} \left( W_{\Phi}(t,s) \overline{W_{\Phi}(t,s)} \right) \right| \\ & = \left| \left( \frac{\partial}{\partial s} W_{\Phi}(t,s) \right) \overline{W_{\Phi}(t,s)} + W_{\Phi}(t,s) \left( \frac{\partial}{\partial s} \overline{W_{\Phi}(t,s)} \right) \right| \\ & \leq 2 \left| W_{\Phi}(t,s) \right| \left| \frac{\partial W_{\Phi}}{\partial s}(t,s) \right|, \end{aligned}$$

we have

$$\mathbb{P}\left(\left|\frac{\partial S_{\Phi}}{\partial s}(t, s_{Y,m}(t))\right| > \frac{\varepsilon L_m}{3} \text{ and } s_{Y,m}(t) \in B_m^{\circ}(t)\right)$$
$$\leq \mathbb{P}\left(\left|W_{\Phi}(t, s_{Y,m}(t))\right| > \sqrt{\frac{\varepsilon L_m}{6}} \text{ and } s_{Y,m}(t) \in B_m^{\circ}(t)\right)$$
$$+ \mathbb{P}\left(\left|\frac{\partial W_{\Phi}}{\partial s}(t, s_{Y,m}(t))\right| > \sqrt{\frac{\varepsilon L_m}{6}} \text{ and } s_{Y,m}(t) \in B_m^{\circ}(t)\right).$$

By the first part of Lemma 4,

$$\mathbb{P}\left(|W_{\Phi}(t, s_{Y,m}(t))| > \sqrt{\frac{\varepsilon L_m}{6}} \text{ and } s_{Y,m}(t) \in B_m^{\circ}(t)\right)$$

$$(109) \leq \mathbb{P}\left(\max_{s \in B_m(t)} |W_{\Phi}(t, s)| > \sqrt{\frac{\varepsilon L_m}{6}}\right) \leq \exp\left[-\frac{1}{\sigma_{m,3}^2} \left(\sqrt{\frac{\varepsilon L_m}{6}} - \mu_{m,3}\right)^2\right]$$

if  $\sqrt{\varepsilon L_m} > \mu_{m,3}/\sqrt{6}$ , where  $\mu_{m,3}$  and  $\sigma_{m,3}^2$  are defined in (27). Similarly, by the second part of Lemma 4,

$$\mathbb{P}\left(\left|\frac{\partial W_{\Phi}}{\partial s}(t, s_{Y,m}(t))\right| > \sqrt{\frac{\varepsilon L_m}{6}} \text{ and } s_{Y,m} \in B_m^{\circ}(t)\right)$$

$$(110) \leq \mathbb{P}\left(\max_{s \in B_m(t)} \left|\frac{\partial W_{\Phi}}{\partial s}(t, s)\right| > \sqrt{\frac{\varepsilon L_m}{6}}\right) \leq \exp\left[-\frac{1}{\sigma_{m,4}^2} \left(\sqrt{\frac{\varepsilon L_m}{6}} - \mu_{m,4}\right)^2\right]$$

if  $\sqrt{\varepsilon L_m} > \mu_{m,4}/\sqrt{6}$ , where  $\mu_4$  and  $\sigma_4^2$  are defined in (27). The proof of Theorem 5 is completed by summing (107), (108), (109), and (110).

6.10. **Proof of Lemma 1.** Because the complex Gaussian random vector **J** is assumed to be circularly symmetric, there exists a matrix  $\mathbf{A} \in \mathbb{C}^{d \times d}$  such that

(111) 
$$\mathbf{J} \stackrel{\text{Law}}{=} \mathbf{AN}$$

where  $\mathbf{N} = [N_1 + iN_{d+1} \ N_2 + iN_{d+2} \ \cdots \ N_d + iN_{2d}]^\top \in \mathbb{C}^d$  and  $\{N_1, N_2, \ldots, N_{2d}\}$  is a set of independent standard normal random variables. The matrix  $\mathbf{A}$  in (111) satisfies

$$\mathbb{E}\left[\mathbf{J}\mathbf{J}^*\right] = \mathbf{A}\mathbb{E}\left[\mathbf{N}\mathbf{N}^*\right]\mathbf{A}^* = 2\mathbf{A}\mathbf{A}^*.$$

By the following vector augmentation:

$$\mathbf{N}_{2d} = \begin{bmatrix} \operatorname{Re}\left(\mathbf{N}\right) \\ \operatorname{Im}\left(\mathbf{N}\right) \end{bmatrix} \in \mathbb{R}^{2d}, \ \mathbf{J}_{2d} = \begin{bmatrix} \operatorname{Re}\left(\mathbf{J}\right) \\ \operatorname{Im}\left(\mathbf{J}\right) \end{bmatrix} \in \mathbb{R}^{2d} \text{ and } \mathbf{A}_{2d} = \begin{bmatrix} \operatorname{Re}\left(\mathbf{A}\right) & -\operatorname{Im}\left(\mathbf{A}\right) \\ \operatorname{Im}\left(\mathbf{A}\right) & \operatorname{Re}\left(\mathbf{A}\right) \end{bmatrix} \in \mathbb{R}^{2d \times 2d}$$

we have  $\mathbf{J}_{2d} \stackrel{\text{Law}}{=} \mathbf{A}_{2d} \mathbf{N}_{2d}$  from (111), and for any  $u \in \mathbb{R}$ ,

(112) 
$$\mathbb{P}(g(\mathbf{J}) > u) = \mathbb{P}(g_{2d}(\mathbf{J}_{2d}) > u) = \mathbb{P}(g_{2d}(\mathbf{A}_{2d}\mathbf{N}_{2d}) > u)$$

where  $g_{2d}: \mathbb{R}^{2d} \mapsto \mathbb{R}$  is defined by

$$g_{2d}\left(\begin{bmatrix}\operatorname{Re}(\mathbf{w})\\\operatorname{Im}(\mathbf{w})\end{bmatrix}\right) = g(\mathbf{w}), \ \mathbf{w} \in \mathbb{C}^d.$$

By the Gaussian concentration inequality [20, Corollary 8.5], for any  $u > \mathbb{E}[g_{2d}(\mathbf{A}_{2d}\mathbf{N}_{2d})]$ ,

(113) 
$$\mathbb{P}\left(g_{2d}(\mathbf{A}_{2d}\mathbf{N}_{2d}) > u\right) \le \exp\left(-\frac{\left(u - \mathbb{E}\left[g_{2d}(\mathbf{A}_{2d}\mathbf{N}_{2d})\right]\right)^2}{2\|g_{2d}(\mathbf{A}_{2d}\bullet)\|_{\mathrm{Lip}}^2}\right),$$

where

$$\|g_{2d}(\mathbf{A}_{2d}\bullet)\|_{\mathrm{Lip}} = \sup_{\substack{\mathbf{x},\mathbf{y}\in\mathbb{R}^{2d}\\\mathbf{x}\neq\mathbf{y}}} \frac{|g_{2d}(\mathbf{A}_{2d}\mathbf{x}) - g_{2d}(\mathbf{A}_{2d}\mathbf{y})|}{|\mathbf{x}-\mathbf{y}|} = \|g(\mathbf{A}\bullet)\|_{\mathrm{Lip}}.$$

The proof of Lemma 1 is completed by substituting (113) into (112) and observing that  $\mathbb{E}[g_{2d}(\mathbf{A}_{2d}\mathbf{N}_{2d})] = \mathbb{E}[g(\mathbf{J})].$ 

6.11. **Proof of Lemma 2.** (a) By the spectral representation of  $W_{\Phi}(t, s)$  in (8) and the orthogonal property (5),

(114) 
$$\mathbb{E}[|W_{\Phi}(0,s)|^2] = \int_0^\infty |\widehat{\psi}(s\lambda)|^2 p(\lambda) d\lambda = s^{-1} \int_0^\infty |\widehat{\psi}(\lambda)|^2 p\left(s^{-1}\lambda\right) d\lambda.$$

Under Assumption 3(b1),  $p(\lambda) = L(|\lambda|^{-1})|\lambda|^{H-1}$ , where L is slowly varying at infinity. According to [3, Theorem 1.5.3], for any slowly varying function L and any  $\delta > 0$ ,  $L(s)s^{-\delta} \to 0$  as  $s \to \infty$ . Thus, (114) implies that

(115)  

$$\lim_{s \to \infty} L(s)^{-1} s^H \mathbb{E}[|W_{\Phi}(0,s)|^2] \\
= \lim_{s \to \infty} \int_0^\infty |\widehat{\psi}(\lambda)|^2 |\lambda|^{H-1} \frac{L(s/|\lambda|)}{L(s)} d\lambda \\
= \int_0^\infty |\widehat{\psi}(\lambda)|^2 |\lambda|^{H-1} d\lambda.$$

The inequality (34) follows from (115).

(b) By (8),

$$W_{\Phi}(0,s_1) - W_{\Phi}(0,s_2) = \int_0^\infty \left[ \ \overline{\widehat{\psi}(s_1\lambda)} - \overline{\widehat{\psi}(s_2\lambda)} \ \right] \sqrt{p(\lambda)} Z(d\lambda).$$

By the orthogonal property (5),

(116)  
$$d_{W_{\Phi}}(s_1, s_2) = \left\{ \mathbb{E} \left[ |W_{\Phi}(0, s_1) - W_{\Phi}(0, s_2)|^2 \right] \right\}^{\frac{1}{2}}$$
$$= \left( \int_0^\infty |\widehat{\psi}(s_1\lambda) - \widehat{\psi}(s_2\lambda)|^2 p(\lambda) d\lambda \right)^{\frac{1}{2}}$$

Because  $\hat{\psi}$  is assumed to be Lipschitz continuous on  $(0,\infty)$  with Lipschitz constant  $\|\hat{\psi}\|_{\text{Lip}}$ , (116) can be estimated as follows

(117) 
$$d_{W_{\Phi}}^{2}(s_{1},s_{2}) \leq \int_{0}^{\infty} \left( \min\left\{2\|\widehat{\psi}\|_{\infty}, \|\widehat{\psi}\|_{\operatorname{Lip}}|s_{1}-s_{2}|\lambda\right\} \right)^{2} p(\lambda) d\lambda$$
$$= 4\|\widehat{\psi}\|_{\infty}^{2} \int_{\Lambda}^{\infty} p(\lambda) d\lambda + \|\widehat{\psi}\|_{\operatorname{Lip}}^{2}|s_{1}-s_{2}|^{2} \int_{0}^{\Lambda} \lambda^{2} p(\lambda) d\lambda,$$

where  $\Lambda = 2 \|\widehat{\psi}\|_{\infty} \|\widehat{\psi}\|_{\text{Lip}}^{-1} |s_1 - s_2|^{-1}$ . Under Assumption 3(a),

(118) 
$$\int_{\Lambda}^{\infty} p(\lambda) d\lambda \le C_1 \int_{\Lambda}^{\infty} \lambda^{-(1+\gamma)} d\lambda = \frac{C_1}{\gamma} \left( \frac{2\|\widehat{\psi}\|_{\infty}}{\|\widehat{\psi}\|_{\mathrm{Lip}} |s_1 - s_2|} \right)^{-\gamma}$$

Let  $\kappa = 2 \|\widehat{\psi}\|_{\infty} \|\widehat{\psi}\|_{\text{Lip}}^{-1}$ . When  $|s_1 - s_2| \leq 1$ , we have  $\Lambda \geq \kappa$  and

$$\int_{0}^{\Lambda} \lambda^{2} p(\lambda) d\lambda \leq \int_{0}^{\kappa} \lambda^{2} p(\lambda) d\lambda + C_{1} \int_{\kappa}^{\Lambda} \lambda^{1-\gamma} d\lambda$$

$$(119) \qquad \leq 2 \frac{\|\widehat{\psi}\|_{\infty}^{2}}{\|\widehat{\psi}\|_{\text{Lip}}^{2}} \text{Var}\left(\Phi(0)\right) + \frac{2^{2-\gamma} C_{1}}{2-\gamma} \left(\frac{\|\widehat{\psi}\|_{\infty}}{\|\widehat{\psi}\|_{\text{Lip}}}\right)^{2-\gamma} \left(|s_{1}-s_{2}|^{\gamma-2}-1\right).$$

By substituting (118) and (119) into (117),

$$\begin{aligned} d_{W_{\Phi}}^{2}(s_{1},s_{2}) &\leq C_{1}\left(\frac{1}{\gamma} + \frac{1}{2-\gamma}\right)(2\|\widehat{\psi}\|_{\infty})^{2-\gamma}\|\widehat{\psi}\|_{\mathrm{Lip}}^{\gamma}|s_{1} - s_{2}|^{\gamma} \\ &+ \left(2\|\widehat{\psi}\|_{\infty}^{2}\mathrm{Var}\left(\Phi(0)\right) - \frac{C_{1}}{2-\gamma}(2\|\widehat{\psi}\|_{\infty})^{2-\gamma}\|\widehat{\psi}\|_{\mathrm{Lip}}^{\gamma}\right)|s_{1} - s_{2}|^{2} \\ &\leq \left[C_{1}\left(\frac{1}{\gamma} + \frac{2}{2-\gamma}\right)(2\|\widehat{\psi}\|_{\infty})^{2-\gamma}\|\widehat{\psi}\|_{\mathrm{Lip}}^{\gamma} + 2\|\widehat{\psi}\|_{\infty}^{2}\mathrm{Var}\left(\Phi(0)\right)\right]|s_{1} - s_{2}|^{\gamma} \end{aligned}$$
for  $|s_{1} - s_{2}| \leq 1$ 

for  $|s_1 - s_2| \le 1$ .

6.12. Proof of Lemma 3. The proof idea can be traced back to the chaining argument in [39]. We firstly create a sequence of sets, denoted as  $\{\pi_j\}_{j\in\mathbb{N}\cup\{0\}}$ , of finite points in  $[0, \infty)$  such that for each  $j \in \mathbb{N} \cup \{0\}$ ,

(120) 
$$\sup_{s>0} \inf_{s'\in\pi_j} d_{W_{\Phi}}(s,s') \le \mathcal{D}^{1-j},$$

where  $\mathcal{D}$  is a constant greater than two, as defined in (38). Because  $\lim_{s \to 0^+} W_{\Phi}(\cdot, s) =$ 0 almost surely, as shown in Corollary 1, we define  $W_{\Phi}(\cdot, 0) = 0$ . Hence, for any  $s \ge 0,$ 

$$d_{W_{\Phi}}(0,s) = \left\{ \mathbb{E}\left[ |W_{\Phi}(0,s)|^2 \right] \right\}^{\frac{1}{2}} \le \mathcal{D},$$

where the inequality follows from the definition of  $\mathcal{D}$  in (38). Hence, we can choose  $\pi_0 = \{0\}$ . For every  $j \in \mathbb{N}$ , denote

(121) 
$$S_j = \max\{(2^{-1}\mathcal{D}^{1-j})^{-2/H^-}, T\}$$

By Lemma 2(a), for any  $s \ge S_j$ ,

(122)

$$d_{W_{\Phi}}(s, S_j) \leq \left\{ \mathbb{E}\left[ |W_{\Phi}(0, s)|^2 \right] \right\}^{\frac{1}{2}} + \left\{ \mathbb{E}\left[ |W_{\Phi}(0, S_j)|^2 \right] \right\}^{\frac{1}{2}} \leq s^{-H^-/2} + S_j^{-H^-/2} \leq \mathcal{D}^{1-j}$$

Consider a set of partition points on  $[0, S_i)$  with lag size  $2(C_2 \mathcal{D}^{j-1})^{-2/\gamma}$  as follows

(123) 
$$\pi_{j,k} = 2k(C_2\mathcal{D}^{j-1})^{-\frac{2}{\gamma}}, \ k = 0, 1, 2, \dots, \lfloor S_j 2^{-1} (C_2\mathcal{D}^{j-1})^{\frac{2}{\gamma}} \rfloor$$

where  $C_2$  is a constant defined in (36). For any  $s \in [0, S_j)$ , one of the following situations occurs:  $|s - S_j| \leq (C_2 \mathcal{D}^{j-1})^{-2/\gamma} \leq 1$ , or there exists an integer  $k \in \{0, 1, 2, \dots, \lfloor S_j 2^{-1} (C_2 \mathcal{D}^{j-1})^{2/\gamma} \rfloor\}$  such that  $|s - \pi_{j,k}| \leq (C_2 \mathcal{D}^{j-1})^{-2/\gamma} \leq 1$ . If the first situation occurs, by Lemma 2(b),

(124) 
$$d_{W_{\Phi}}(s, S_j) \le C_2 |s - S_j|^{\gamma/2} \le \mathcal{D}^{1-j}.$$

If the second situation occurs, by Lemma 2(b) again, we have

(125) 
$$d_{W_{\Phi}}(s,\pi_{j,k}) \le |s-\pi_{j,k}|^{\gamma/2} \le \mathcal{D}^{1-j}.$$

Hence, the set

$$\pi_j = \{\pi_{j,0}, \pi_{j,1}, \ldots\} \cup \{S_j\}$$

fulfills the condition (120). Denote the cardinality of  $\pi_i$  by  $|\pi_i|$ . By (121) and (123), for  $j \in \mathbb{N}$ ,

(126) 
$$|\pi_j| \le C_2^{\frac{2}{\gamma}} \left[ 2^{\frac{2}{H^-}} \mathcal{D}^{\left(\frac{2}{\gamma} + \frac{2}{H^-}\right)(j-1)} \lor T \mathcal{D}^{\frac{2}{\gamma}(j-1)} \right].$$

For any s > 0 and  $j \in \mathbb{N} \cup \{0\}$ , we choose a proxy for s from the set  $\pi_j$  as follows

$$\pi_j(s) = \arg\min_{s' \in \pi_j} |s - s'|.$$

According to (122), (124), and (125),

$$d_{W_{\Phi}}(s,\pi_j(s)) \le \mathcal{D}^{1-j}$$

Because  $\pi_0 = \{0\}$  and  $W_{\Phi}(0,0) = 0$  by definition, we can represent  $|W_{\Phi}(0,s)|$  via a telescoping sum as follows

(127) 
$$|W_{\Phi}(0,s)| = \sum_{j=0}^{\infty} |W_{\Phi}(0,\pi_{j+1}(s))| - |W_{\Phi}(0,\pi_j(s))|.$$

Let  $\{a_j\}_{j=0}^{\infty} \subset [0,\infty)$  satisfying  $A := \sum_{j=0}^{\infty} a_j < \infty$ , where  $a_j$  will be chosen later. By (127), for any u > 0,

$$\mathbb{P}\left(\sup_{s>0} |W_{\Phi}(0,s)| \ge u\right) = \mathbb{P}\left(\sup_{s>0} \sum_{j=0}^{\infty} |W_{\Phi}(0,\pi_{j+1}(s))| - |W_{\Phi}(0,\pi_{j}(s))| \ge uA^{-1}\sum_{j=0}^{\infty} a_{j}\right) \\
(128) \qquad \qquad \leq \sum_{j=0}^{\infty} \mathbb{P}\left(\sup_{s>0} |W_{\Phi}(0,\pi_{j+1}(s))| - |W_{\Phi}(0,\pi_{j}(s))| \ge uA^{-1}a_{j}\right).$$

For any s > 0 and  $j \in \mathbb{N} \cup \{0\}$ ,

$$\mathbb{P}\left(|W_{\Phi}(0,\pi_{j+1}(s))| - |W_{\Phi}(0,\pi_{j}(s))| \ge uA^{-1}a_{j}\right)$$
  
$$\leq \mathbb{P}\left(|W_{\Phi}(0,\pi_{j+1}(s)) - W_{\Phi}(0,\pi_{j}(s))| \ge uA^{-1}a_{j}\right)$$
  
(129) 
$$= \mathbb{P}\left(\frac{|W_{\Phi}(0,\pi_{j+1}(s)) - W_{\Phi}(0,\pi_{j}(s))|^{2}}{d^{2}_{W_{\Phi}}(\pi_{j+1}(s),\pi_{j}(s))} \ge \frac{u^{2}A^{-2}a_{j}^{2}}{d^{2}_{W_{\Phi}}(\pi_{j+1}(s),\pi_{j}(s))}\right),$$

where

$$d_{W_{\Phi}}^{2}(\pi_{j+1}(s),\pi_{j}(s)) = \mathbb{E}\left[ |W_{\Phi}(0,\pi_{j+1}(s)) - W_{\Phi}(0,\pi_{j}(s))|^{2} \right].$$

By Theorem 1(b), the random variable

$$\frac{|W_{\Phi}(0,\pi_{j+1}(s)) - W_{\Phi}(0,\pi_{j}(s))|^2}{d^2_{W_{\Phi}}(\pi_{j+1}(s),\pi_{j}(s))}$$

follows an exponential distribution with a mean of one. Continuing from (129), for any s > 0 and  $j \in \mathbb{N} \cup \{0\}$ ,

(130) 
$$\mathbb{P}\left(|W_{\Phi}(0,\pi_{j+1}(s))| - |W_{\Phi}(0,\pi_{j}(s))| \ge uA^{-1}a_{j}\right) \le \exp\left(-\frac{u^{2}A^{-2}a_{j}^{2}}{d_{W_{\Phi}}^{2}(\pi_{j+1}(s),\pi_{j}(s))}\right) \le \exp\left(-u^{2}A^{-2}a_{j}^{2}\mathcal{D}^{2(j-1)}\right)$$

where the last inequality follows from that  $d_{W_{\Phi}}(\pi_{j+1}(s), \pi_j(s)) \leq \mathcal{D}^{1-j}$ . Because the number of pairs  $\{(\pi_j(s), \pi_{j+1}(s)) \mid s \geq 0\}$  is less than  $|\pi_j||\pi_{j+1}|$ , (128) and (130) imply that

$$\mathbb{P}\left(\sup_{s>0} |W_{\Phi}(0,s)| \ge u\right) \le \sum_{j=0}^{\infty} |\pi_j| |\pi_{j+1}| \exp\left(-u^2 A^{-2} a_j^2 \mathcal{D}^{2(j-1)}\right)$$

for any  $u \geq 0$ , and

$$\mathbb{E}\left[\sup_{s>0}|W_{\Phi}(0,s)|\right] = \int_0^\infty \mathbb{P}\left(\sup_{s>0}|W_{\Phi}(0,s)| \ge u\right) du$$
(131)
$$\leq A + \int_A^\infty \sum_{j=0}^\infty |\pi_j| |\pi_{j+1}| \exp\left(-u^2 A^{-2} a_j^2 \mathcal{D}^{2(j-1)}\right) du$$

By choosing

$$a_j = \mathcal{D}^{1-j} \sqrt{\ln\left(2^{j+1} |\pi_j| |\pi_{j+1}|\right)},$$

the integrand function in (131) can be estimated as follows (132)

$$\sum_{j=0}^{\infty} |\pi_j| |\pi_{j+1} |\exp\left(-u^2 A^{-2} a_j^2 \mathcal{D}^{2(j-1)}\right) = \sum_{j=0}^{\infty} |\pi_j| |\pi_{j+1}| \left(2^{j+1} |\pi_j| |\pi_{j+1}|\right)^{-u^2 A^{-2}} \le 2^{-u^2 A^{-2}}$$

for any  $u \ge A$ . By substituting (132) into (131),

$$\mathbb{E}\left[\sup_{s>0} |W_{\Phi}(0,s)|\right] \le A\left(1 + \int_{1}^{\infty} 2^{-v^2} dv\right) \le \frac{4}{3}A.$$

Because

(133) 
$$A = \sum_{j=0}^{\infty} \mathcal{D}^{1-j} \sqrt{\ln\left(2^{j+1}|\pi_j||\pi_{j+1}|\right)},$$

by substituting (126) into (133) and performing elementary calculation,

$$A \le \sqrt{3\ln \mathcal{D}} \frac{\mathcal{D}^3}{(\mathcal{D}-1)^2} \sqrt{\frac{1}{\gamma} + \frac{1}{H^-}} + \left(\sqrt{\frac{4}{\gamma}\ln C_2} + \sqrt{2\ln T}\right) \frac{\mathcal{D}^2}{\mathcal{D}-1}.$$

6.13. **Proof of Lemma 4.** Since the proofs of the first and second parts of Lemma 4 are similar, with the first being simpler, we provide only the proof of the second part below. Consider a complex-valued Gaussian process as follows

$$V(s) = w(s) \frac{\partial W_{\Phi}}{\partial s}(0,s), \ s \in [a,b]$$

As shown in Proposition 1,  $W_{\Phi}$  is two times continuously differentiable, which implies that V is continuous on [a, b] almost surely, and

(134) 
$$\mathbb{P}\left(\|V\|_{[a,b]} < \infty\right) = 1, \text{ where } \|V\|_{[a,b]} := \max_{s \in [a,b]} |V(s)|.$$

Using (8) and denoting the derivative of  $\widehat{\psi}$  by  $D\widehat{\psi}$ , we have

(135)  

$$V(s) = w(s) \int_{0}^{\infty} \overline{[D\widehat{\psi}](s\lambda)} \lambda \sqrt{p(\lambda)} Z(d\lambda)$$

$$= w(s) \int_{-\infty}^{0} \overline{[D\widehat{\psi}](-s\lambda)} \lambda \sqrt{p(\lambda)} \overline{Z(d\lambda)},$$

which implies that for any  $s_1, s_2 \in [a, b]$ ,

(136) 
$$\mathbb{E}[V(s_1)V(s_2)] = 0.$$

Let  $\{T_d\}_{d\in\mathbb{N}}$  be a sequence of subsets of [a, b] such that  $T_d$  has the cardinality d,  $T_d \subset T_{d+1}$ , and  $T_d$  increases to a dense subset of [a, b]. By (136),  $\mathbf{V}_d := \{V(s) \mid d\}$ 

 $s \in T_d$  is a circularly symmetric complex Gaussian random vector. By Lemma 1, for any u > 0,

(137) 
$$\mathbb{P}\left(\pm\left(\|\mathbf{V}_d\|_{\ell^{\infty}} - \mathbb{E}\left[\|\mathbf{V}_d\|_{\ell^{\infty}}\right]\right) > u\right) \le \exp\left(-\frac{u^2}{2\|\|\mathbf{A}_d \bullet\|_{\ell^{\infty}}\|_{\mathrm{Lip}}^2}\right),$$

where  $\mathbf{A}_d$  is a matrix in  $\mathbb{C}^{d \times d}$  satisfying

(138) 
$$\mathbf{A}_d \mathbf{A}_d^* = 2^{-1} \mathbb{E} \left[ \mathbf{V}_d \mathbf{V}_d^* \right].$$

For any  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^{d \times 1}$  with  $\mathbf{x} \neq \mathbf{y}$ ,

(139)  

$$\begin{aligned} \|\mathbf{A}_{d}\mathbf{x}\|_{\ell^{\infty}} - \|\mathbf{A}_{d}\mathbf{y}\|_{\ell^{\infty}} &\leq \max_{1 \leq k \leq d} \left| e_{k}^{\top} \mathbf{A}_{d} \left( \mathbf{x} - \mathbf{y} \right) \right| \\ &\leq \max_{1 \leq k \leq d} \left| e_{k}^{\top} \mathbf{A}_{d} \right| \left| \mathbf{x} - \mathbf{y} \right| \\ &= \left( \max_{1 \leq k \leq d} e_{k}^{\top} \mathbf{A}_{d} \mathbf{A}_{d}^{*} e_{k} \right) \left| \mathbf{x} - \mathbf{y} \right| \end{aligned}$$

where  $e_k$  is the column vector in  $\mathbb{R}^d$  with one in position k and zeros elsewhere. If we denote the elements of  $T_d$  by  $s_{d,1}, s_{d,2}, \ldots, s_{d,d}$ , then (135) and (138) imply that

(140) 
$$e_k^{\top} \mathbf{A}_d \mathbf{A}_d^* e_k = 2^{-1} w^2(s_{d,k}) \int_0^\infty |[D\widehat{\psi}](s_{d,k}\lambda)|^2 \lambda^2 p(\lambda) d\lambda.$$

By combining (137), (139), and (140), we obtain that for any u > 0,

(141) 
$$\mathbb{P}\left(\pm\left(\|\mathbf{V}_d\|_{\ell^{\infty}} - \mathbb{E}\left[\|\mathbf{V}_d\|_{\ell^{\infty}}\right]\right) > u\right) \le \exp\left(-\frac{u^2}{\sigma_{T_d}^2}\right),$$

where

$$\sigma_{T_d}^2 = \max_{1 \le \ell \le d} w^2(s_{d,\ell}) \int_0^\infty |[D\widehat{\psi}](s_{d,\ell}\lambda)|^2 \lambda^2 p(\lambda) d\lambda.$$

Because  $T_d$  increases to a dense subset of [a, b] as  $d \to \infty$ , the continuity of V on [a, b] implies that

(142) 
$$\|\mathbf{V}_d\|_{\ell^{\infty}} \uparrow \|V\|_{[a,b]} < \infty$$

almost surely, and under Assumption 2,

(143) 
$$\sigma_{T_d}^2 \uparrow \sigma_{[a,b]}^2 = \max_{s \in [a,b]} w^2(s) \int_0^\infty |[D\widehat{\psi}](s\lambda)|^2 \lambda^2 p(\lambda) d\lambda < \infty.$$

On the other hand, when d increases,

(144) 
$$\mathbb{E}\left[\|\mathbf{V}_d\|_{\ell^{\infty}}\right] \uparrow \mathbb{E}\left[\|V\|_{[a,b]}\right].$$

If the limit  $\mathbb{E}[||V||_{[a,b]}]$  is finite, then for any u > 0, (142) and (144) imply that

$$\mathbb{P}\left(\pm\left(\|V\|_{[a,b]} - \mathbb{E}\left[\|V\|_{[a,b]}\right]\right) > u\right) = \lim_{d \to \infty} \mathbb{P}\left(\pm\left(\|\mathbf{V}_d\|_{\ell^{\infty}} - \mathbb{E}\left[\|\mathbf{V}_d\|_{\ell^{\infty}}\right]\right) > u\right)$$
$$\overset{(141)}{\leq} \lim_{d \to \infty} \exp\left(-\frac{u^2}{\sigma_{T_d}^2}\right) \stackrel{(143)}{=} \exp\left(-\frac{u^2}{\sigma_{[a,b]}^2}\right)$$

Therefore, to complete the proof, it suffices to show that  $\mathbb{E}[\|V\|_{[a,b]}] < \infty$ .

We proceed by contradiction. Thus, assume that  $\mathbb{E}[||V||_{[a,b]}] = \infty$ , and choose  $u_0 > 0$  such that

$$\exp\left(-\frac{u_0^2}{\sigma_{[a,b]}^2}\right) < \frac{1}{3} \text{ and } \mathbb{P}\left(\|V\|_{[a,b]} < u_0\right) \ge \frac{1}{2},$$

which is attainable due to (134). Now choose  $d \ge 1$  such that  $\mathbb{E}[\|\mathbf{V}_d\|_{\ell^{\infty}}] > 2u_0$ , which is attainable due to (144) and  $\mathbb{E}[\|V\|_{[a,b]}] = \infty$ . By (141),

$$\begin{aligned} \frac{1}{3} > \exp\left(-\frac{u_0^2}{\sigma_{[a,b]}^2}\right) &\geq \exp\left(-\frac{u_0^2}{\sigma_{T_d}^2}\right) \\ \geq \mathbb{P}\left(-\left(\|\mathbf{V}_d\|_{\ell^{\infty}} - \mathbb{E}[\|\mathbf{V}_d\|_{\ell^{\infty}}]\right) > u_0\right) \\ &= \mathbb{P}\left(\|\mathbf{V}_d\|_{\ell^{\infty}} < \mathbb{E}[\|\mathbf{V}_d\|_{\ell^{\infty}}] - u_0\right) \\ \geq \mathbb{P}\left(\|\mathbf{V}_d\|_{\ell^{\infty}} < u_0\right) \\ &\geq \mathbb{P}\left(\|\mathbf{V}\|_{[a,b]} < u_0\right) \geq \frac{1}{2}, \end{aligned}$$

which is contradictory.

#### References

- Charalambos D Aliprantis and Kim C Border. Infinite Dimensional Analysis: A Hitchhiker's Guide. Springer Science & Business Media, 2006.
- [2] Jean-Pierre Antoine, Romain Murenzi, Pierre Vandergheynst, and Syed Twareque Ali. Twodimensional wavelets and their relatives. Cambridge University Press, 2008.
- [3] Nicholas H Bingham, Charles M Goldie, Jozef L Teugels, and JL Teugels. *Regular variation*. Number 27. Cambridge University Press, 1989.
- [4] René A Carmona, Wen L Hwang, and Bruno Torrésani. Multiridge detection and timefrequency reconstruction. *IEEE Trans. Signal Process.*, 47(2):480–492, 1999.
- [5] René A Carmona, Wen-Liang Hwang, and Bruno Torrésani. Characterization of signals by the ridges of their wavelet transforms. *IEEE Trans. Signal Process.*, 45(10):2586–2590, 1997.
- [6] Marcelo A. Colominas, Sylvain Meignen, and Duong-Hung Pham. Fully adaptive ridge detection based on STFT phase information. *IEEE Signal Process. Lett.*, 27:620–624, 2020.
- [7] Marcelo A Colominas and Hau-Tieng Wu. An iterative warping and clustering algorithm to estimate multiple wave-shape functions from a nonstationary oscillatory signal. *IEEE Trans.* Signal Process., 71:701–712, 2023.
- [8] Ingrid Daubechies. Ten lectures on wavelets. SIAM, 1992.
- [9] Ingrid Daubechies, Jianfeng Lu, and Hau-Tieng Wu. Synchrosqueezed wavelet transforms: An empirical mode decomposition-like tool. Appl. Comput. Harmon. Anal., 30(2):243–261, 2011.
- [10] Nathalie Delprat, Bernard Escudié, Philippe Guillemain, Richard Kronland-Martinet, Philippe Tchamitchian, and Bruno Torresani. Asymptotic wavelet and Gabor analysis: Extraction of instantaneous frequencies. *IEEE Trans. Inform. Theory*, 38(2):644–664, 1992.
- [11] Richard M Dudley. The sizes of compact subsets of Hilbert space and continuity of Gaussian processes. J. Funct. Anal., 1(3):290–330, 1967.
- [12] RM Dudley. Sample functions of the Gaussian process. Ann. Probab., 1(5):66–103, 1973.
- [13] David Eberly. Ridges in image and data analysis, volume 7. Springer Science & Business Media, 2012.
- [14] Ryszard Engelking. General topology, volume 6. Heldermann Verlag, Berlin, 2nd edition, 1989.
- [15] Patrick Flandrin. Time-frequency/time-scale analysis. Academic Press, 1998.
- [16] Peter Hall, Wei Qian, and DM Titterington. Ridge finding from noisy data. J. Comput. Graph. Statist., 1(3):197–211, 1992.
- [17] Nicki Holighaus, Gunther Koliander, Zdenek Prusa, and Luis Daniel Abreu. Characterization of analytic wavelet transforms and a new phaseless reconstruction algorithm. *IEEE Trans. Signal Process.*, 67(15):3894–3908, 2019.

- [18] Shouchuan Hu and Nikolaos S Papageorgiou. Handbook of multivalued analysis: Volume I: Theory, Kluwer, 1997.
- [19] Dmytro Iatsenko, Peter VE McClintock, and Aneta Stefanovska. Extraction of instantaneous frequencies from ridges in time-frequency representations of signals. *Signal Process.*, 125:290– 303, 2016.
- [20] Franziska Kühn and René L Schilling. Maximal inequalities and some applications. Probab. Surv., 20:382–485, 2023.
- [21] Nils Laurent and Sylvain Meignen. A novel ridge detector for nonstationary multicomponent signals: Development and application to robust mode retrieval. *IEEE Trans. Signal Process.*, 69:3325–3336, 2021.
- [22] Quentin Legros and Dominique Fourer. A novel pseudo-bayesian approach for robust multiridge detection and mode retrieval. In Proc. 29th Eur. Signal Process. Conf., pages 1925–1929, 2021.
- [23] Jonathan M Lilly and Sofia C Olhede. On the analytic wavelet transform. *IEEE Trans. Inform. Theory*, 56(8):4135–4156, 2010.
- [24] S. C. Lim and L Teo. Analytic and asymptotic properties of multivariate generalized Linnik's probability densities. J. Fourier Anal. Appl., 16:715–747, 2010.
- [25] Péter Major. Muliple Wiener-Itô integrals, volume 849. Springer, 2nd edition, 2014.
- [26] Stéphane Mallat. A wavelet tour of signal processing. Elsevier, 1999.
- [27] Sylvain Meignen and Marcelo Colominas. A new ridge detector localizing strong interference in multicomponent signals in the time-frequency plane. *IEEE Trans. Signal Process.*, 71:3413– 3425, 2023.
- [28] Sylvain Meignen, Duong-Hung Pham, and Stephen McLaughlin. On demodulation, ridge detection, and synchrosqueezing for multicomponent signals. *IEEE Trans. Signal Process.*, 65(8):2093–2103, 2017.
- [29] I. Nourdin and G. Peccati. Normal approximations with Malliavin calculus: from Stein's method to universality. Number 192. Cambridge University Press, 2012.
- [30] N. Ozkurt and F.A. Savaci. Determination of wavelet ridges of nonstationary signals by singular value decomposition. *IEEE Trans. Circuits Syst. II Express Briefs*, 52(8):480–485, 2005.
- [31] Vladas Pipiras and Murad S Taqqu. Regularization and integral representations of Hermite processes. Statist. Probab. Lett., 80(23-24):2014–2023, 2010.
- [32] Jürgen Potthoff. Sample properties of random fields I: Separability and measurability. Commun. Stoch. Anal., 3(1):9, 2009.
- [33] Jürgen Potthoff. Sample properties of random fields III: Differentiability. Commun. Stoch. Anal., 4(3):3, 2010.
- [34] R. Tyrrell Rockafellar. Second-order convex analysis. J. Nonlinear Convex Anal., 1(1):1–16, 2000.
- [35] R Tyrrell Rockafellar and Roger J-B Wets. Variational analysis, volume 317. Springer Science & Business Media, 2009.
- [36] Ervin Sejdic, Igor Djurovic, and LJubisa Stankovic. Quantitative performance analysis of scalogram as instantaneous frequency estimator. *IEEE Trans. Signal Process.*, 56(8):3837– 3845, 2008.
- [37] Martin K Stiles, David Clifton, Neil R Grubb, James N Watson, and Paul S Addison. Wavelet-based analysis of heart-rate-dependent ECG features. Ann. Noninvasive Electrocardiol., 9(4):316–322, 2004.
- [38] Yan-Wei Su, Gi-Ren Liu, Yuan-Chung Sheu, and Hau-Tieng Wu. Ridge detection for nonstationary multicomponent signals with time-varying wave-shape functions and its applications. *IEEE Trans. Signal Process.*, 2024.
- [39] Michel Talagrand. Upper and lower bounds for stochastic processes, volume 60. Springer, 2014.
- [40] Roman Vershynin. High-dimensional probability: An introduction with applications in data science, volume 47. Cambridge University Press, 2018.
- [41] Xiangxiang Zhu, Zhuosheng Zhang, Jinghuai Gao, and Wenting Li. Two robust approaches to multicomponent signal reconstruction from STFT ridges. *Mech. Syst. Signal Process.*, 115:720–735, 2019.

# Appendix

Symbol	Description
f	Adapted harmonic signal with $M$ components (see (1)).
$\phi_1',\ldots,\phi_M'$	IFs of f with $\phi'_1(t) < \cdots < \phi'_M(t)$ for any $t \in \mathbb{R}$ .
$A_1,\ldots,A_M$	Amplitude functions of harmonic components of $f$ .
$\Phi$	Stationary Gaussian process (see Assumption 1).
p	Spectral density function of $\Phi$ .
Y	Noisy signal modeled as $Y = f + \Phi$ .
Z	Orthogonally scattered Gaussian random measure on $\mathbb{R}$ .
$\psi,\widehat{\psi}$	Analytic mother wavelet and its Fourier transform.
$\psi_R,  \psi_I$	Real and imaginary parts of the wavelet $\psi$ .
$D\widehat{\psi}, D^2\widehat{\psi}$	First and second derivatives of $\widehat{\psi}$ .
$W_f, W_{\Phi}, W_Y$	Continuous wavelet transform of $f$ , $\Phi$ , and $Y$ .
$W^R_{\Phi}, W^I_{\Phi}$	Real and imaginary parts of $W_{\Phi}$ .
$S_f, S_{\Phi}, S_Y$	Scalograms of $f$ , $\Phi$ , and $Y$ , i.e., the squared modulus of $W_f$ ,
	$W_{\Phi}$ , and $W_Y$ .
$s_f(t)$	Argmax of the scalogram of $f$ at time $t$ , i.e., $\arg \max_{s>0} S_f(t,s)$ .
$s_{f,m}(t)$	Local maxima of f's scalogram at time t, ordered as $s_{f,1} >$
	$\cdots > s_{f,M}.$
$s_Y(t)$	Set of scales maximizing $S_Y(t, \cdot)$ , i.e., $s_Y(t) = \arg \max_{s>0} S_Y(t, s)$ .
$\omega_\psi$	Argmax of $ \widehat{\psi} $ .
$B_m(t)$	IF information of $f: \omega_{\psi}(\phi'_m(t))^{-1} \in B_m(t)$ (see (19)).
$s_{Y,m}(t)$	Scales in $B_m(t)$ maximizing $S_Y(t, \cdot)$ , i.e., $s_{Y,m}(t) =$
	$\arg\max_{s\in B_m(t)}S_Y(t,s).$

TABLE 1. List of frequently used symbols.

**Lemma 5** (Berge's maximum theorem). Let  $\mathbb{X}$  and  $\mathbb{Y}$  be Hausdorff topological spaces. If  $u : \mathbb{X} \times \mathbb{Y} \to \mathbb{R}$  is a continuous function and  $F : \mathbb{X} \to \{K \subseteq \mathbb{Y} : K \text{ is a nonempty compact set}\}$  is a continuous multifunction, then

$$\arg \sup_{y \in F(x)} u(x,y) := \left\{ z \in F(x) : \ u(x,z) = \sup_{y \in F(x)} u(x,y) \right\}$$

is an upper hemicontinuous multifunction.

Interpretation of Theorem 5 by the single component case M = 1: In the following, we omit the component-specific index m. That is, the symbols  $A_m$ ,  $L_m$ ,  $\mu_{m,1},...,\mu_{m,4}$ , and  $\sigma_{m,1},...,\sigma_{m,4}$  are replaced by A, L,  $\mu_1,...,\mu_4$ , and  $\sigma_1,...,\sigma_4$ , respectively, to reflect the single-component scenario.

• L: A larger value of the second derivative of  $|W_f(t, \cdot)|^2$  with respect to the scale variable at  $s_f(t)$  indicates that  $|W_f(t, \cdot)|$  exhibits a sharper peak at  $s_f(t)$ . This can be associated with a rapid change in the spectral content of the signal f. Hence, when L is larger, the ridge point of f is less likely

to be disturbed. If the magnitude function A varies slowly enough to allow the approximation

(145) 
$$S_f(t,s) \approx A^2(t) S_{f/A}(t,s),$$

then L is approximately proportional to  $A^2(t)$ . In this case, a larger A similarly reduces the possibility of ridge point disturbance.

•  $\mu_1, \mu_2, \mu_3, \mu_4$ : Under the approximation (145),

$$\mu_1 \approx A(t) \sqrt{\operatorname{Var}(\Phi)} \mathbb{E} \left[ \max_{s \in I} \left| \frac{\partial W_{f/A}}{\partial s}(t,s) W_{\widetilde{\Phi}}(t,s) \right| \right],$$

where  $\tilde{\Phi} = \Phi/\sqrt{\operatorname{Var}(\Phi)}$ . Similarly,  $\mu_2$  is approximately proportional to  $A(t)\sqrt{\operatorname{Var}(\Phi)}$ . Both  $\mu_3$  and  $\mu_4$  are approximately proportional to  $\sqrt{\operatorname{Var}(\Phi)}$ . This implies that

(146) 
$$\frac{6}{L} \max\left\{\mu_1, \mu_2, \mu_3^2, \mu_4^2\right\} \approx \max\left\{\mathcal{O}\left(\frac{\sqrt{\operatorname{Var}(\Phi)}}{A}\right), \mathcal{O}\left(\frac{\operatorname{Var}(\Phi)}{A^2}\right)\right\}$$

Therefore, the available region for  $\varepsilon$  in (29) becomes larger when the ratio  $A(t)/\sqrt{\operatorname{Var}(\Phi)}$  increases.

• For any fixed  $\varepsilon$  satisfying (29), (146) shows that the exponents

$$\frac{\varepsilon}{6} - \frac{\mu_1}{L}, \ \frac{\varepsilon}{6} - \frac{\mu_2}{L}, \ \sqrt{\frac{\varepsilon}{6}} - \frac{\mu_3}{\sqrt{L}}, \ \sqrt{\frac{\varepsilon}{6}} - \frac{\mu_4}{\sqrt{L}}$$

become larger when the ratio  $A/\sqrt{\operatorname{Var}(\Phi)}$  increases. Thus, (30) implies that the probability of  $s_Y(t)$  deviating from  $s_f(t)$  by more than  $\varepsilon$  decreases exponentially fast when the ratio  $A/\sqrt{\operatorname{Var}(\Phi)}$  increases.

• Exponents  $L^2/\sigma_1^2$ ,  $L^2/\sigma_2^2$ ,  $L/\sigma_3^2$ , and  $L/\sigma_4^2$ : Since both  $\sigma_1^2$  and  $\sigma_2^2$  are approximately proportional to  $A^2 \operatorname{Var}(\Phi)$ , and both  $\sigma_3^2$  and  $\sigma_4^2$  are approximately proportional to  $\operatorname{Var}(\Phi)$ , we have

$$L^2/\sigma_k^2 = \mathcal{O}\left(\frac{A^2}{\operatorname{Var}(\Phi)}\right) \text{ for } k = 1, 2,$$

and

$$L/\sigma_k^2 = \mathcal{O}\left(\frac{A^2}{\operatorname{Var}(\Phi)}\right) \text{ for } k = 3, 4.$$

This also implies that, for any fixed  $\varepsilon$  satisfying (29), the probability that  $s_Y(t)$  deviates from  $s_f(t)$  by more than  $\varepsilon$  decreases exponentially fast when the ratio  $A/\sqrt{\operatorname{Var}(\Phi)}$  increases.

**Lemma 6.** [14, Lemma (4.3.16)] If  $\mathbb{X}$  is a topological space,  $\mathbb{Y}$  a complete metric space and  $r : \mathbb{D} \to \mathbb{Y}$  a continuous mapping defined on a dense subset  $\mathbb{D}$  of the space  $\mathbb{X}$ , then the mapping r is extendable to a continuous mapping defined on the set consisting of all points of  $\mathbb{X}$  at which the oscillation of r is equal to zero. Here, we say that the oscillation of the mapping r at a point  $x \in \mathbb{X}$  is equal to zero if for every  $\varepsilon > 0$  there exists a neighborhood U of the point x such that

$$\sup_{x',x''\in\mathbb{D}\cap U}r(x')-r(x'')<\varepsilon.$$

The set of all points at which the oscillation of r is equal to zero is a  $G_{\delta}$ -set containing the dense set  $\mathbb{D}$ .

DEPARTMENT OF MATHEMATICS, NATIONAL CHENG-KUNG UNIVERSITY, TAINAN, TAIWAN AND NATIONAL CENTER FOR THEORETICAL SCIENCES, NATIONAL TAIWAN UNIVERSITY, TAIPEI, TAIWAN Email address: girenliu@ncku.edu.tw

Department of Applied Mathematics, National Yang Ming Chiao Tung University, Hsinchu, Taiwan

 $Email \ address: \ {\tt sheu@math.nctu.edu.tw}$ 

Courant Institute of Mathematical Sciences, New York University, New York, NY, 10012 $\rm USA$ 

Email address: hauwu@cims.nyu.edu