SELF-INTERSECTION NUMBER OF NEGATIVE CURVES ON FERMAT SURFACES

ZHENJIAN WANG

ABSTRACT. We give an explicit formula for the self-intersection number of negative curves on Fermat surfaces. The formula offers us hints to either prove or disprove the Bounded Negativity Conjecture for the Fermat surfaces.

CONTENTS

1.	Introduction	1
2.	Curves on surfaces of Sebastiani-Thom type	5
3.	Strategy of Proof	6
4.	Self-intersection number for curves on Fermat surfaces	9
5.	Discussions and Problems	12
6.	Puiseux series, resultants, and intersection multiplicities	13
References		

1. INTRODUCTION

Let \mathbb{P}^3 be the three-dimensional complex projective space, with homogeneous coordinate ring $S = \mathbb{C}[x, y, z, w]$. The Fermat surface of degree d, denoted as X_d , is the smooth hypersurface in \mathbb{P}^3 defined by the equation:

$$x^d + y^d + z^d + w^d = 0.$$

The main objective in this paper is to establish the following theorem.

Theorem 1.1 (Theorem 4.5). Let $X_d \subset \mathbb{P}^3$ be the Fermat surface defined by $x^d + y^d + z^d + w^d = 0$ for $d \geq 5$, and C a reduced and irreducible curve on X_d with $C^2 < 2 - d < 0$. Then

(1) There exists an irreducible homogeneous polynomial g(x, y, z) which vanishes on C.

Denote by $e = \deg g$. Define two curves E and D in \mathbb{P}^2 by

E: g = 0 and $D: x^d + y^d + z^d = 0.$

²⁰¹⁰ Mathematics Subject Classification. Primary 14C17, Secondary 14H20.

Key words and phrases. Bounded Negativity Conjecture, Fermat surface, intersection multiplicity, Puiseux series, resultant.

(2) There exists $k \geq 2$ and k|d such that

$$C^{2} = \frac{d}{k}e^{2} - \frac{d(d-1)}{k}e - \frac{d}{k}\sum_{p\in D\cap E}\mu_{p}(E) + \sum_{[x,y,z]\in D\cap E}\mu_{[x,y,z,0]}(C) + \left(\frac{d}{k} - 1\right)\#(D\cap E),$$

where $\mu_p(E)$ denotes the Milnor number of E at p. In particular, when d is prime, then k = d and

$$C^{2} = e^{2} - (d-1)e - \sum_{p \in D \cap E} \mu_{p}(E) + \sum_{[x,y,z] \in D \cap E} \mu_{[x,y,z,0]}(C).$$

This result is closely connected with an old folklore conjecture among experts in algebraic surface theory, commonly known as the *Bounded Negativity Conjecture*. We formulate this conjecture as follows.

Conjecture 1.2 (Bounded Negativity Conjecture). For any given smooth complex projective surface X, there exists a number $B_X \ge 0$ which depends only on X such that for any reduced curve C on X, the self-intersection number C^2 is bounded below by $-B_X$, i.e., $C^2 \ge -B_X$.

The origins of this conjecture are unclear, but it can at least be traced back to the Italian mathematician Federigo Enriques (1871-1946), and it has been an open problem for one hundred years. For a more comprehensive overview, we refer to [3, p.1878].

An alternative formulation of the Bounded Negativity Conjecture is as follows.

Conjecture 1.3 (Bounded Negativity Conjecture for integral curves). For any given smooth complex projective surface X, there exists a number $b_X \ge 0$ which depends only on X such that for any reduced and irreducible curve C on X, the self-intersection number C^2 is bounded below by $-b_X$, i.e., $C^2 \ge -b_X$.

Indeed, as proven in [3, Prop.5.1, p.1891], we can take the number B_X in Conjecture 1.2 as $B_X = (\rho(X) - 1) b_X$, where $\rho(X)$ is the Picard number of X. We will refer to b_X as the *negativity bound* for X.

Our result in Theorem 1.1 implies that under some conditions, the Bounded Negativity Conjecture is true for the Fermat surface X_d .

Corollary 1.4 (Corollary 4.6). The Bounded Negativity Conjecture holds for the Fermat surface X_d if one of the following conditions is satisfied: (a)

$$\liminf_{e \to \infty} \inf_{E} \left(e^2 - (d-1)e - \sum_{p \in D \cap E} \mu_p(E) \right) \ge -\lambda$$

for some constant $\lambda > 0$ which may depend on d, where E ranges over all irreducible projective plane curves of degree e and D is the projective plane curve defined by $x^d + y^d + z^d = 0$; or (b)

$$\limsup_{e\to\infty}\sup_E \sup_E \frac{\sum\limits_{p\in D\cap E} \mu_p(E)}{e^2} < 1,$$

where E ranges over all irreducible projective plane curves of degree e and D is the projective plane curve defined by $x^d + y^d + z^d = 0$.

On the other hand, the formula in Theorem 1.1 also gives us hints to construct counterexamples for the Bounded Negativity Conjecture on Fermat surfaces; see Section 5.

In the sequel, we will only investigate the self-intersection number of reduced and irreducible curves. For convenience, we denote C such a curve on a given smooth projective surface.

It is evident that the Bounded Negativity Conjecture holds for X with $-K_X \ge 0$, as can be seen by applying the adjunction formula:

$$K_X \cdot C + C^2 = 2p_a(C) - 2 \Longrightarrow C^2 \ge -2.$$

In particular, for $d \leq 4$, the bounded negativity of X_d follows from this argument, since the canonical divisor of X_d is $K_{X_d} = (d-4)H$ where H is the hyperplane section. However, for $d \geq 5$, the canonical divisor K_{X_d} is ample and X_d is of general type. To the author's knowledge, there are currently no effective methods available to deal with the Bounded Negativity of such surfaces, except for the weak Bounded Negativity Conjecture which will be quoted below.

Another natural approach to establishing the Bounded Negativity for a surface is to show that there are only finitely many negative curves on the surface. However, it is proved in [3, Thm.4.1] that for any m > 0, there exists a smooth projective surface which admits infinitely many smooth irreducible curves of self-intersection number -m. For the Fermat surface X_d , we do not know whether the number of negative curves on X_d is finite or not.

A third potential method to prove the Bounded Negativity Conjecture is to establish the boundedness of the geometric genera of reduced and irreducible curves. In fact, the weak Bounded Negativity Conjecture, proven in [10], states that on a given smooth projective surface X, the inequality

$$C^2 \ge -b_{X,g(C)}$$

holds for any reduced and irreducible curve C, where $b_{X,g(C)}$ is a constant depending only on X and the geometric genus g(C) of C. Consequently, the Bounded Negativity Conjecture would follow if we could establish a bound for the genera g(C). Nevertheless, such a bound does not exist for almost all surfaces. For the Fermat surface X_d , consider a generic curve C in the linear series |kH|, then C is a smooth irreducible curve on X_d with geometric genus

$$g(C) = \frac{1}{2}(K_{X_d} \cdot C + C^2) + 1 = \frac{1}{2}\left(d(d-4)k + dk^2\right) + 1,$$

thus the genus g(C) goes to infinity as k tends to infinity. Therefore, the Bounded Negativity Conjecture for X_d cannot be proved by bounding the genera of reduced and irreducible curves.

Other strategies to prove the Bounded Negativity Conjecture include investigating the non-isomorphic surjective endomorphisms or establishing dominance by other surfaces that have bounded negativity [3, Prop.2.1], exploring the birational invariance of bounded negativity and the *H*-constant [2], using the equivalence between bounded negativity and the boundedness of the denominators in the Zariski decomposition [4], discussing the weighted Bounded Negativity Conjecture [11, 8], and studying the connections between the Bounded Negativity Conjecture and other unsolved problems, such as SHGH conjecture or the bounded cohomology conjecture [1, 7]. However, despite providing invaluable insights in understanding the Bounded Negativity Conjecture, these methods either deal with only very special cases, e.g., rational surfaces or K3 surfaces, or just offer general frameworks that reduce the conjecture to other open questions. They rely on general formulas, such as the adjunction formula and Riemann-Roch formula, to establish bounds for the intersection number C^2 of a reduced and irreducible curve C, without using specific information about C, such as its equation.

Can we estimate C^2 from the equation of the curve C? Can we give a concrete example of a surface with the Bounded Negativity by doing explicit computation through the defining equations of the surface?

To explore these questions, we consider smooth surfaces in \mathbb{P}^3 , and give the following conjecture.

Conjecture 1.5. For any $d \ge 1$, there exists a constant c_d that depends only on d such that all smooth surfaces of degree d in \mathbb{P}^3 have negativity bound c_d .

Since all the degree d surfaces in \mathbb{P}^3 are contained in a connected family, they can be deformed into one another. We further conjecture that the bounded negativity property is invariant under deformations.

Conjecture 1.6. Suppose X and Y are two smooth complex projective surfaces that are in a single family of complex projective surfaces, then the Bounded Negativity Conjecture is true for X if and only if it is true for Y.

Assuming the conjecture above, to establish the Bounded Negativity Conjecture for surfaces in \mathbb{P}^3 , it suffices to prove its validity for a specific surface. Our Theorem 1.1 is proved for this purpose.

The formulas in Theorem 1.1 show how the Bounded Negativity Conjecture is connected with the singularity theory of curves. The key term in the expression of the self-intersection number C^2 is the sum $\sum_{p \in D \cap E} \mu_p(E)$. It is notable that this sum

is taken over the intersection points of D and E, not all the singular points of E.

To conclude the introduction, we give an outline of the proof of Theorem 1.1. Given a negative curve C on X_d , we represent the Fermat surface X_d as a branched Galois cover of \mathbb{P}^2 :

$$\rho: X_d \to \mathbb{P}^2, \quad [x, y, z, w] \mapsto [x, y, z]$$

Let $E = \rho(C)$ and $\mathscr{C} = \rho^* E$. Then E is a reduced and irreducible curve in \mathbb{P}^2 ; denote by g its defining equation. It is clear that g can be seen as a section of the line bundle $\mathscr{O}_{X_d}((\deg g)H)$ which vanishes on C. Then we prove that \mathscr{C} is a reduced curve on X_d ; see Proposition 4.4. Moreover, \mathscr{C} is of the following form

$$\mathscr{C} = C_1 + \dots + C_k,$$

with $C_i^2 = C$ for all i = 1, ..., k. This result holds because C_i 's lie in the orbit of C under the Galois group action for ρ . It follows that

$$kC^2 = d(\deg g)^2 - \sum_{i \neq j} C_i \cdot C_j.$$

Hence, to compute C^2 , we need to calculate the sum on the right hand side. Finally, the intersection numbers can be calculated in a local way at any intersection point. Locally, the curve \mathscr{C} can be represented as the curve in $(\mathbb{C}^3, 0)$ defined by

$$h(u, v) = 0, \ l(u, v) + \omega^d = 0.$$

We will first solve v as Puiseux series of u from the first equation, and then use $l(u, v(u)) + \omega^d = 0$ to investigate the intersection multiplicities between different irreducible components of \mathscr{C} . This is the most technical part in this paper and will be completed in Proposition 4.4 and Section 6.

The author would like to thank the organizers of International Symposium on Singularities and Applications, Dec. 9–13, 2024 in Sanya, China. The stimulating atmosphere of the conference inspired him to conceive the idea of performing computation using Puiseux series. The author was partially supported by the grant YSBR-001 and is currently supported by NSFC No. 12301100.

2. Curves on surfaces of Sebastiani-Thom type

Assume that $X \subset \mathbb{P}^3$ is a smooth surface defined by the following equation:

$$X: f(x, y, z) + w^d = 0,$$

where f is a form of degree d. Such a surface is said to be of *Sebastiani-Thom type*. The Fermat surface is a typical example. Let C be a reduced and irreducible curve on X.

In this section, we show that C is "almost" defined by a homogeneous polynomial on X in the following result.

Proposition 2.1. Let $X : f(x, y, z) + w^d = 0$ be a smooth surface in \mathbb{P}^3 and C an reduced and irreducible curve on X. Then there exists an irreducible form g(x, y, z) satisfying the following properties:

(i) The divisor on X cut out by g can be written as

$$(g) = m(C_1 + \dots + C_k),$$

where $C_1 = C, C_2, \ldots, C_k$ are distinct reduced and irreducible curves and $m \ge 1$ is an integer.

(ii) The self-intersection numbers C_i^2 , i = 1, ..., k are all equal to C^2 .

Proof. Consider the following projection morphism

$$\rho: X \to \mathbb{P}^2, \ [x, y, z, w] \mapsto [x, y, z].$$

It is a branched Galois covering map with Galois group $\mathbb{Z}_d = \mathbb{Z}/d\mathbb{Z}$, and the Galois action is given by

$$a \cdot [x, y, z, w] = [x, y, z, e^{2\pi\sqrt{-1}a/d}w], \ \forall a \in \mathbb{Z}_d.$$

For the curve $C \subset X$, we consider its image under the morphism ρ , denoted by $E = \rho(C)$. Then E is a reduced and irreducible curve in \mathbb{P}^2 .

Let g(x, y, z) be an irreducible defining equation for E. Then the inverse image $\rho^{-1}(E)$ is exactly the curve defined by g = 0 on X (here we see g as a homogeneous polynomial in x, y, z, w restricted to X). Write the divisor (g) cut out by g on X as

$$(g) = m_1 C_1 + \dots + m_k C_k, \ C_1 = C,$$

where C_1, \ldots, C_k are reduced and irreducible curves on X and $m_1, \ldots, m_k \ge 1$.

Since the action \mathbb{Z}_d for the morphism ρ is a Galois action, we see that $\rho^{-1}(E)$ is exactly the union of the curves in the \mathbb{Z}_d -orbit of C. It follows that $C_i, i > 1$ are all isomorphic to C by automorphisms of X. Hence $C_i^2 = C^2$ for all $i = 1, \ldots, k$; and (i) follows.

Moreover, it is obvious that g is a \mathbb{Z}_d -invariant polynomial in (x, y, z, w). For each pair $i \neq j$, there exists an element α in the Galois group \mathbb{Z}_d such that $\alpha \cdot C_i = C_j$, and then

$$m_i = \operatorname{ord}_{C_i}(g) = \operatorname{ord}_{\alpha \cdot C_i}(\alpha \cdot g) = \operatorname{ord}_{C_i}(g) = m_j$$

Hence (ii) is true.

3. Strategy of Proof

Let $X_d: x^d + y^d + z^d + w^d = 0$ be the Fermat surface in \mathbb{P}^3 and $C \subset X_d$ a reduced and irreducible curve. We shall investigate the lower bound of the self-intersection number C^2 .

By Proposition 2.1, there exists an irreducible polynomial g(x, y, z) which cuts out a divisor on X of the following form

$$(g) = m(C_1 + \dots + C_k), \ C_1 = C$$

and $C_i^2 = C^2$ for i = 1, ..., k.

To investigate the Bounded Negativity Conjecture for X_d , we can safely assume that k > 1; otherwise, $m^2C^2 = d(\deg g)^2$ and thus $C^2 \ge 0$. Therefore, we assume $k \ge 2$ in the sequel.

For convenience, denote by D the curve defined by $x^d + y^d + z^d = 0$ and E the curve defined by g = 0 in \mathbb{P}^2 . Namely,

$$D: x^{d} + y^{d} + z^{d} = 0$$
 and $E: g(x, y, z) = 0.$

Denote $e = \deg g = \deg E$.

Denote \mathscr{C} the scheme on X_d defined by g = 0:

$$\mathscr{C}: g(x, y, z) = 0.$$

Then C_1, \ldots, C_k are the irreducible components of \mathscr{C} .

3.1. Strategy. From the equation

$$(g) = m(C_1 + \dots + C_k),$$

we get

$$de^{2} = m^{2}(C_{1} + \dots + C_{k})^{2}$$

= $m^{2} \sum_{i=1}^{k} C_{i}^{2} + m^{2} \sum_{i \neq j} C_{i} \cdot C_{j}$
= $m^{2}kC^{2} + m^{2} \sum_{i \neq j} C_{i} \cdot C_{j},$

where $C_i \cdot C_j$ denotes the intersection number on X_d .

To study the Bounded Negativity Conjecture on X_d , we need to obtain a lower bound for C^2 . From the formula above, we are led to give an upper bound for the sum

$$\sum_{i \neq j} C_i \cdot C_j.$$

The result will be a function of e where d is viewed as a constant, and so is the difference

$$de^2 - m^2 \sum_{i \neq j} C_i \cdot C_j.$$

If this function is bounded below for $e \ge 1$, then the Bounded Negativity Conjecture for X_d will follow.

The main part of this paper is to calculate the intersection numbers using local computations. The results are summarized in Proposition 4.4, and the lengthy proof will be presented in Section 6.

Fix an arbitrary intersection point $P = [x_0, y_0, z_0, w_0]$ of some C_i and C_j with $i \neq j$, we will give an explicit formula for the sum

$$\sum_{i \neq j} I_P(C_i, C_j),$$

where $I_P(C_i, C_j)$ denotes the intersection multiplicity (that will be explored in detail in Section 6).

Since we consider all curves C on X_d , we should view g as an abstract polynomial. From $P \in X_d$, we have that $x_0^d + y_0^d + z_0^d + w_0^d = 0$, hence not all x_0, y_0, z_0 are zero. By the symmetry of the coordinates x, y, z in the equation $X_d : x^d + y^d + z^d + w^d = 0$, we may without loss of generality assume that $x_0 \neq 0$. To decide where the intersection point $P = [x_0, y_0, z_0, w_0]$ may situate on X_d , we propose the following Lemma which will be proved at the end of this section.

Lemma 3.2. Following the notations above, for any point Q lying on at least two irreducible components of C, we have

$$w(Q) = 0,$$

where w is the fourth homogeneous coordinate for \mathbb{P}^3 . Moreover, $Q \in C_i$ for all $i = 1, \ldots, k$, i.e., Q lies on every irreducible component of \mathscr{C} .

In particular, for the point $P = [x_0, y_0, z_0, w_0]$ above, we have $w_0 = 0$ and $P \in C_i$ for all $i \leq k$.

Let $p = \rho(P) = [x_0, y_0, z_0]$. Then $p \in E \cap D$.

Recall that $w_0 = 0$ and we assumed $x_0 \neq 0$. To calculate the intersection multiplicities at P, we introduce affine coordinates

$$u = \frac{y}{x} - \frac{y_0}{x_0}, \qquad v = \frac{z}{x} - \frac{z_0}{x_0}, \qquad \omega = \frac{w}{x}.$$

Then the curve $\mathscr{C}: g = 0$ on X_d is defined in the affine piece $\mathbb{C}^3 = \{x \neq 0\} \subset \mathbb{P}^3$ by

$$h(u, v) = 0, \quad l(u, v) + \omega^d = 0,$$

where

$$h(u, v) = g\left(1, u + \frac{y_0}{x_0}, v + \frac{z_0}{x_0}\right)$$

and

$$l(u, v) = 1 + \left(u + \frac{y_0}{x_0}\right)^d + \left(v + \frac{z_0}{x_0}\right)^d$$

Moreover, since $x_0^d + y_0^d + z_0^d = 0$ and $x_0 \neq 0$, at least one of y_0, z_0 does not vanish. We may without loss of generality assume that $z_0 \neq 0$. Then

$$\frac{\partial \left(l(u,v) + \omega^d \right)}{\partial v} \bigg|_{(0,0,0)} = d \left(\frac{z_0}{x_0} \right)^{d-1} \neq 0$$

Therefore, we can locally express v as an analytic function of (u, ω) by applying the implicit function theorem to the equation $l(u, v) + \omega^d = 0$. Then the local equation for \mathscr{C} is $h(u, v(u, \omega)) = 0$. By analyzing this local equation, we can investigate the local branches and compute the local intersection multiplicities. This is the usual way to analyze the curve \mathscr{C} .

Our method is, however, first to solve v from the equation h(u, v) = 0 by using the Puiseux series method (which will be recalled in Section 6), say v = s(u). Then, using the equation $l(u, s(u)) + \omega^d = 0$, we discuss the properties of the curve \mathscr{C} and calculate the intersection multiplicities. To this end, we need that h is v-general (the related notions will be discussed in Section 6).

At this point, we specially consider the exceptional case where h is not v-general, or equivalently, $h(0, v) \equiv 0$; the v-general case will be investigated in Section 6.

Recall that h is transformed from an irreducible polynomial g, thus h is an irreducible polynomial in u, v. If h is not v-general, then u is a factor of h; therefore, by irreducibility, h is a constant multiple of u. Back to g, we see that g is a linear function y - ax (up to multiplicative factor). Therefore, the curve \mathscr{C} is defined as

$$\mathscr{C}: y = ax, \ (1 + a^d)x^d + z^d + w^d = 0.$$

If $a^d + 1 \neq 0$, then \mathscr{C} is itself irreducible and hence it must be equal to C. Thus, $C^2 = d$ in this case.

If $a^d + 1 = 0$, then \mathscr{C} splits into d lines. Let

$$L_i: y = ax, \ z = e^{(1+2i)\pi\sqrt{-1/d}}w,$$

for $i = 1, \ldots, d$. Then

$$(g) = L_1 + \dots + L_d.$$

It is straightforward that $L_i \cdot L_j = 1$ for $i \neq j$. Hence, by the above discussions, we have

$$d = dC^2 + d(d-1),$$

and so $C^2 = 2 - d$.

3.3. **Proof of Lemma 3.2.** Recall that we have the degree *d* branched Galois cover $\rho : X_d \to \mathbb{P}^2$ defined by $[x, y, z, w] \mapsto [x, y, z]$. The ramification locus of ρ is the closed subset of X_d defined by the equation w = 0.

If Q is a point on at least two components of \mathscr{C} , then $\#(\rho^{-1}(\rho(Q))) < d$, where #A means the number of points in a set A. It follows that Q lies on the ramification locus of ρ . Thus, w(Q) = 0 as desired.

Furthermore, denote $q = \rho(Q)$, then $\rho^{-1}(q) = \{Q\}$. For all $i \leq k$, the restriction morphism $\rho|_{C_i} : C_i \to E$ is a finite morphism and thus $\rho^{-1}(q) \cap C_i \neq \emptyset$. But $\rho^{-1}(q)$ consists of only one point Q, it follows that $Q \in C_i$. We are done.

4. Self-intersection number for curves on Fermat surfaces

Following the notations above, we will give an explicit formula for the self-intersection number C^2 .

4.1. Position of intersection points. From Lemma 3.2, we see that any intersection point Q of two irreducible components of \mathscr{C} satisfies w(Q) = 0. It follows that $q = \rho(Q)$ is an intersection point of D and E. The converse is also true.

Lemma 4.2. If q = [x, y, z] lies on both D and E, then Q = [x, y, z, 0] lies on all of C_i for $i \leq k$.

Hence, $Q \in C_i \cap C_j$ for $i \neq j$ if and only if $q \in D \cap E$.

Proof. By definition, $\rho^{-1}(q) = \{Q\}$ and $Q \in \mathscr{C}$. Since each C_i is mapped onto E, the intersection $\rho^{-1}(q) \cap C_i$ is nonempty. Hence $Q \in C_i$ as desired.

The last assertion follows immediately by Lemma 3.2.

4.3. Intersection multiplicities. Recall that we have the following equation

$$de^2 = m^2 k C^2 + m^2 \sum_{i \neq j} C_i \cdot C_j.$$

To give a lower bound for C^2 , we need to estimate the sum on the right-hand side. Moreover, by definition, we have

$$\sum_{i \neq j} C_i \cdot C_j = \sum_{\substack{i \neq j \\ Q \in C_i \cap C_j}} I_Q(C_i, C_j)$$

Hence, it suffices to compute the sum

$$\sum_{i \neq j} I_P(C_i, C_j)$$

where $P = [x_0, y_0, z_0, 0]$ is chosen above. Our main technical results are summarized as follows.

Proposition 4.4. With the notations above, we have

- (A) m = 1; in other words, \mathscr{C} is a reduced scheme.
- (B) The following equality holds:

$$\iota_P(\mathscr{C}) = (d-1)i_p(D, E) + d\mu_p(E) - d + 1.$$

(C) We have

$$\sum_{i \neq j} I_P(C_i, C_j) = (d-1)i_p(D, E) + d\mu_p(E) - k\mu_P(C) - (d-k).$$

and thus

$$\sum_{i \neq j} C_i \cdot C_j = d(d-1)e + d \sum_{p \in D \cap E} \mu_p(E) - k \sum_{p \in D \cap E} \mu_{\rho^{-1}(p)}(C) - (d-k) \# (D \cap E).$$

The proof will be presented in Section 6.

Now we use the above results to study the Bounded Negativity Conjecture for Fermat surfaces. It is clear that we only need to consider negative curves C. Then the curve \mathscr{C} necessarily has $k \geq 2$ irreducible components since otherwise C^2 would be nonnegative. Moreover, the number k divides d since the Galois action for ρ is a cyclic action of order d. To ensure that the polynomial h is v-general, we only need to assume $C^2 \neq 2 - d$; see the discussions in Section 3. Therefore, combining all of our discussions above, we get the following theorem.

Theorem 4.5. Let $X_d \subset \mathbb{P}^3$ be the Fermat surface defined by $x^d + y^d + z^d + w^d = 0$ for $d \geq 5$, and C a reduced and irreducible curve on X_d with $C^2 < 2 - d < 0$. Then (1) There exists an irreducible homogeneous polynomial q(x, y, z) which vanishes

on C.

Denote by $e = \deg g$. Define two curves E and D in \mathbb{P}^2 by

$$E: g = 0 \text{ and } D: x^d + y^d + z^d = 0.$$

(2) There exists $k \ge 2$ and k|d such that

$$C^{2} = \frac{d}{k}e^{2} - \frac{d(d-1)}{k}e - \frac{d}{k}\sum_{p\in D\cap E}\mu_{p}(E) + \sum_{[x,y,z]\in D\cap E}\mu_{[x,y,z,0]}(C) + \left(\frac{d}{k} - 1\right)\#(D\cap E).$$

In particular, when d is prime, then k = d and

$$C^{2} = e^{2} - (d-1)e - \sum_{p \in D \cap E} \mu_{p}(E) + \sum_{[x,y,z] \in D \cap E} \mu_{[x,y,z,0]}(C).$$

As a corollary, we get two sufficient conditions for the Bounded Negativity Conjecture to hold on Fermat surfaces.

Corollary 4.6. With the notations above, the Bounded Negativity Conjecture holds on the Fermat surface X_d if one of the following conditions is satisfied: (a)

$$\liminf_{e \to \infty} \inf_{E} \left(e^2 - (d-1)e - \sum_{p \in D \cap E} \mu_p(E) \right) \ge -\lambda$$

for some constant $\lambda > 0$ which may depend on d, where E ranges over all irreducible projective plane curves of degree e and D is the projective plane curve defined by $x^d + y^d + z^d = 0.$

(b)

$$\limsup_{e\to\infty}\sup_E \sup_E \frac{\sum\limits_{p\in D\cap E} \mu_p(E)}{e^2} < 1,$$

where again E ranges over all irreducible projective plane curves of degree e and D is as defined above.

Proof. By our computations before, it is clear that (a) implies the Bounded Negativity Conjecture. Indeed, if (a) holds, there exists $e_0 > d + 1$ such that

$$\inf_{E} \left(e^2 - (d-1)e - \sum_{p \in D \cap E} \mu_p(E) \right) \ge -(\lambda+1)$$

for $e \ge e_0$. On the other hand, the geometric genus g(E) of E is given by

$$g(E) = \frac{(e-1)(e-2)}{2} - \sum_{p \in E} \frac{1}{2} \left(\mu_p(E) + r_p - 1 \right)$$

by [12, Cor.7.1.3], where r_p is the number of local branches of E at p. It follows that

$$\sum_{p \in D \cap E} \mu_p(E) \le \sum_{p \in E} \mu_p(E) \le (e-1)(e-2)$$

Hence, for $e < e_0$, we have

$$e^{2} - (d-1)e - \sum_{p \in D \cap E} \mu_{p}(E) \geq e^{2} - (d-1)e - (e-1)(e-2)$$
$$= (4-d)e - 2$$
$$\geq -(d-4)e_{0} - 2.$$

Therefore, we may take

$$b_{X_d} = \max\{\lambda + 1, (d - 4)e_0 + 2\}.$$

The condition (b) implies (a), as can be seen as follows: if (b) holds, then for e sufficiently large, we have

$$e^{2} - (d-1)e - \sum_{p \in D \cap E} \mu_{p}(E) \ge e^{2} - (d-1)e - (1-\alpha)e^{2} = \alpha e^{2} - (d-1)e.$$

for some $\alpha > 0$. The desired result follows by noting that

$$\alpha e^2 - (d-1)e = \alpha \left(e - \frac{d-1}{2\alpha}\right)^2 - \frac{(d-1)^2}{4\alpha} \ge -\frac{(d-1)^2}{4\alpha}.$$

5. Discussions and Problems

In this section, we investigate some cases in which we are able to estimate the intersection number C^2 which is equal to

$$\frac{d}{k}e^2 - \frac{d(d-1)}{k}e - \frac{d}{k}\sum_{p\in D\cap E}\mu_p(E) + \sum_{[x,y,z]\in D\cap E}\mu_{[x,y,z,0]}(C) + \left(\frac{d}{k} - 1\right)\#(D\cap E).$$

This equality is from Theorem 4.5.

The first key obstacle for us to prove the Bounded Negativity Conjecture on X_d comes from the term

$$\sum_{p \in D \cap E} \mu_p(E)$$

which may be comparable to $e^2 - (d-1)e$ for e sufficiently large. Moreover, this problem arises from two possible causes: (i) $D \cap E$ contains many singular points of E; (ii) $\mu_p(E)$ is large for $p \in D \cap E$.

5.1. The case $\#(D \cap E) = de$. We first consider the case where $D \cap E$ contains as many points as possible, i.e., $\#(D \cap E) = de$. Then for each $p \in D \cap E$, D and Eintersect transversely at p by [6, Cor.2.6.8]. In particular, p is a smooth point of E, and hence $\mu_p(E) = 0$. Therefore,

$$\sum_{p \in D \cap E} \mu_p(E) = 0.$$

5.2. The case where $\mu_p(E)$ is large for some $p \in D \cap E$. In the extreme case, $\#(D \cap E) = 1$ and $D \cap E$ is the only singular point of E; moreover, E is a rational cuspidal curve and C is smooth. Then

$$\frac{d}{k}e^{2} - \frac{d(d-1)}{k}e - \frac{d}{k}\sum_{p\in D\cap E}\mu_{p}(E) + \sum_{[x,y,z]\in D\cap E}\mu_{[x,y,z,0]}(C) + \left(\frac{d}{k} - 1\right)\#(D\cap E)$$

$$= \frac{d}{k}e^{2} - \frac{d(d-1)}{k}e - \frac{d}{k}(e-1)(e-2) + \left(\frac{d}{k} - 1\right)$$

$$= \frac{d}{k}\left((4-d)e - 1\right) - 1,$$

which can be very negative for e sufficiently large.

Hence we propose the following problem:

Problem 5.3. Does there exist a sequence of irreducible homogeneous polynomials g_{e_i} of degrees e_i for a sequence of integers $e_i \to \infty$ such that the following conditions are satisfied?

- (i) The projective plane curve $V(g_{e_i})$ is a rational cuspidal curve with exactly one singular point lying on the curve $x^d + y^d + z^d = 0$; and
- (ii) The curve \mathscr{C}_{e_i} in \mathbb{P}^3 defined by $g_{e_i}(x, y, z) = 0, x^d + y^d + z^d + w^d = 0$ is a reducible curve with smooth irreducible components.

If the answer is positive, then taking an irreducible component C_{e_i} of \mathscr{C}_{e_i} , we will have

$$C_{e_i}^2 = \frac{d}{k_i} \left((4 - d)e_i - 1 \right) - 1 \to -\infty$$

where $2 \leq k_i \leq d$. This would disprove the Bounded Negativity Conjecture for X_d .

5.4. An invariant. From Corollary 4.6 (b), we are led to consider the following invariant

$$\limsup_{e \to \infty} \sup_{E} \frac{\sum_{p \in D \cap E} \mu_p(E)}{e^2}$$

More generally, given any reduced homogeneous polynomial f of degree d, we define

$$I_f = \limsup_{e \to \infty} \sup_{g \text{ irreducible}} \frac{\sum_{p \in V(f) \cap V(g)} \mu_p(V(g))}{e^2},$$

where V(g) denotes the projective plane curve defined by g.

Problem 5.5. (1) If $f = x^d + y^d + z^d$, is it true that $I_f < 1$?

- (2) More generally, if f defines a smooth projective curve in \mathbb{P}^2 , does it hold that $I_f < 1$?
- (3) Does the invariant I_f depend on the equation of f, or at least the degree d of f?

6. PUISEUX SERIES, RESULTANTS, AND INTERSECTION MULTIPLICITIES

In this section, we first recall basic notions concerning Puiseux series and resultants, and then use them to prove properties of intersection multiplicities of plane curve germs. Finally, we give the proof of Proposition 4.4.

6.1. **Puiseux series.** We follow the notations in [6, Sec.1.2, p.17]. The ring $\mathbb{C}[[u]]$ denotes the set of formal power series, and $\mathbb{C}\{u\}$ the analytic local ring of $(\mathbb{C}, 0)$. Denote by $\mathbb{C}((u))$ the field of fractions of $\mathbb{C}[[u]]$. A typical element of $\mathbb{C}((u))$ can be represented as

$$\sum_{i\geq r} a_i u^i,$$

for some $r \in \mathbb{Z}$ and $a_i \in \mathbb{C}$.

The field of fractionary power series, denoted by $\mathbb{C}\langle\langle u \rangle\rangle$ is the direct limit of the fields $\mathbb{C}((u^{\frac{1}{n}}))$, where the morphisms in the directed system are given by

$$\mathbb{C}((u^{\frac{1}{n}})) \to \mathbb{C}((u^{\frac{1}{n'}})), \quad n|n' \text{ and } \sum a_i u^{\frac{i}{n}} \mapsto \sum a_i u^{\frac{in'/n}{n'}}.$$

Let

$$s = \sum_{i \ge r} a_i u^{\frac{i}{n}}$$

be a fractionary power series, we define the *order* of s to be

$$o_u(s) = \begin{cases} \infty, & \text{if } s = 0; \\ \frac{\min\{i: a_i \neq 0\}}{n}, & \text{if } s \neq 0. \end{cases}$$

Fractionary power series s with $o_u(s) > 0$ will be called *Puiseux series*. After a simultaneous reduction of all the fractional exponents effectively appearing in s, we may assume that n and gcd $\{i : a_i \neq 0\}$ are coprime, and then we call n the polydromy order of s, denoted by $\nu(s)$. We will also use the following notation: if $n = \nu(s)$ and s can be written as

$$s(u) = \sum a_i u^{\frac{i}{n}},$$

then define

$$s_{,\alpha}(u) = \sum e^{2\pi\sqrt{-1}\alpha i/n} a_i u^{\frac{i}{n}}$$

for $\alpha = 1, ..., n$. Moreover, we say that s is a convergent fractionary power series if the formal power series $s(t^n)$ has a positive convergence radius.

Suppose $f \in \mathbb{C}[[u, v]]$ is a formal power series satisfying f(0, 0) = 0. We say a Puiseux series $s \in \mathbb{C}\langle\langle u \rangle\rangle$ is a v-root (or simply, root) of f is f(u, s(u)) = 0. Furthermore, we say that f is v-general of order k, if

$$f(0,v) = c_k v^k + c_{k+1} v^{k+1} + \cdots,$$

for some constant $c_k \neq 0$.

For convenience, for $s, s' \in \mathbb{C}\langle\langle u \rangle\rangle$, we use $s \simeq s'$ to denote the relation

$$s = (a + s'')s'$$

for some $a \in \mathbb{C}^*$ and s'' a Puiseux series. Note that if $s \simeq s'$, then $o_u(s) = o_u(s')$.

Some basic results which we will implicitly use in the theory of Puiseux series are summarized as follows; we refer to [6, Chap.1] for the proof.

Proposition 6.2. The following statements are true:

- (i) The field $\mathbb{C}\langle\langle u \rangle\rangle$ is algebraically closed.
- (ii) If $f \in \mathbb{C}\{u, v\}$ is a convergent power series and a Puiseux series $s \in \mathbb{C}\langle\langle u \rangle\rangle$ is a root of f, then s is convergent.
- (iii) If $f \in \mathbb{C}\{u, v\}$ is v-general of order b and a Puiseux series $s \in \mathbb{C}\langle\langle u \rangle\rangle$ is a root of f such that $\nu(s) = b$, then

$$\prod_{\alpha=1}^{b} (v - s_{,\alpha}(u)) \text{ divides } f \text{ in } \mathbb{C}\{u, v\}.$$

Moreover, if, in addition, f is irreducible, then

$$f = c(u, v) \prod_{\alpha=1}^{b} (v - s_{,\alpha}(u))$$

for an invertible element c(u, v) in $\mathbb{C}\{u, v\}$.

6.3. **Resultants.** We recall the definition and basic properties of resultants; all the results are well known. The interested reader may consult [5, Sec.1.4].

Let K be a field of characteristic zero, which will be taken as $\mathbb{C}\langle\langle u \rangle\rangle$ in our discussion about intersection multiplicities. Let f, g are two polynomials in K[v] of the following form:

$$f = a_0 v^d + a_1 v^{d-1} + \dots + a_d,$$

and

$$g = b_0 v^e + b_1 v^{e-1} + \dots + b_e,$$

with $a_0, b_0 \neq 0$. The resultant $\operatorname{Res}(f, g)$ is the determinant

$ a_0 $	a_1	a_2	• • •	a_d	0	•••	0
0	a_0	a_1	• • •	a_{d-1}	a_d	•••	0
:	÷	·	·	÷	·	·	:
0	0	0	a_0	• • •	• • •	a_{d-1}	a_d
b_0	b_1	b_2	• • •	b_e	0	•••	0
0	b_0	b_1	•••	b_{e-1}	b_e	•••	0
:	÷	·.	·	÷	·	·	:
0	0	0	b_0	•••	• • •	b_{e-1}	b_e

which has e rows of a_i and d rows of b_j .

Alternatively, if $\lambda_1, \dots, \lambda_d$ and μ_1, \dots, μ_e are respectively the roots of f, g in the algebraic closure of K, counted with multiplicities, then

$$\operatorname{Res}(f,g) = a_0^e b_0^d \prod_{\substack{1 \le i \le d \\ 1 \le j \le e}} (\lambda_i - \mu_j).$$

For the polynomial f, the discriminant Disc(f) is defined as

$$\operatorname{Disc}(f) = a_0^{2d-2} \prod_{i < j} (\lambda_i - \lambda_j)^2$$

Proposition 6.4. The following assertions are true:

(i)

$$\operatorname{Disc}(f) = (-1)^{d(d-1)/2} a_0^{2d-2} \prod_{i \neq j} (\lambda_i - \lambda_j) = \frac{(-1)^{d(d-1)/2}}{a_0} \operatorname{Res}\left(f, \frac{\partial f}{\partial v}\right).$$

- (ii) $\operatorname{Res}(f,g) = (-1)^{de} \operatorname{Res}(g,f).$
- (iii) If d = 0, then $\operatorname{Res}(f, g) = a_0^e$.
- (iv) If h is another polynomial in K[v], then

 $\operatorname{Res}(fg, h) = \operatorname{Res}(f, h)\operatorname{Res}(g, h).$

(v) If $a_0 = 1$, then for any $h \in K[v]$, we have

$$\operatorname{Res}(f,g) = \operatorname{Res}(f,g-fh).$$

(vi) If $f = f_1 \cdots f_k$, then

$$\operatorname{Disc}(f) = \left(\prod_{i=1}^{k} \operatorname{Disc}(f_i)\right) \left(\prod_{1 \le i < j \le k} \operatorname{Res}(f_i, f_j)^2\right).$$

6.5. Intersection multiplicities. Now we use the Puiseux series and resultants to study intersection multiplicities of plane curve germs. Suppose F, G are two reduced curve germs in $(\mathbb{C}^2, 0)$, defined by

$$F: f(u,v) = 0,$$

and

G: g(u, v) = 0.

By the Weierstrass preparation theorem (see [6, Thm.1.8.7]), we may assume that f, g are Weierstrass polynomials:

$$f = v^d + a_1(u)v^{d-1} + \dots + a_d(u),$$

and

$$g = v^e + b_1(u)v^{e-1} + \dots + b_e(u),$$

where $a_i, b_j \in \mathbb{C}\{u\}$ satisfy $a_i(0) = b_j(0) = 0$ for $1 \leq i \leq d, 1 \leq j \leq e$. For convenience, we will also regard a_i, b_j as elements of $\mathbb{C}\langle\langle u \rangle\rangle$ by the inclusion $\mathbb{C}\{u\} \subset \mathbb{C}\langle\langle u \rangle\rangle$.

The intersection multiplicity of F and G at 0, denoted by $i_0(F,G)$, which is equal to the intersection multiplicity of f and g at 0, denoted by $i_0(f,g)$, is defined as

$$i_0(F,G) = i_0(f,g) = \dim_{\mathbb{C}} \frac{\mathbb{C}\{u,v\}}{(f,g)}$$

The Milnor number of F at 0, denoted by $\mu_0(F)$ or $\mu_0(f)$, is defined as the intersection multiplicity at 0 of $\frac{\partial f}{\partial u}$ and $\frac{\partial f}{\partial v}$:

$$\mu_0(F) = i_0\left(\frac{\partial f}{\partial u}, \frac{\partial f}{\partial v}\right) = \dim_{\mathbb{C}} \frac{\mathbb{C}\{u, v\}}{\left(\frac{\partial f}{\partial u}, \frac{\partial f}{\partial v}\right)}$$

The following results are classical; the interested reader may consult [6, Chap.2] or [9, Chap.I, §3].

Proposition 6.6. (i) $i_0(f,g) < \infty$ if and only if F and G do not share a common *irreducible component.*

- (ii) If h is another element in $\mathbb{C}\{u,v\}$, h(0,0) = 0, then $i_0(fg,h) = i_0(f,h) + i_0(g,h)$.
- (iii) If F is irreducible and s = s(u) is a v-root of f such that $\nu(s) = d$, then

$$i_0(f,g) = do_u(g(u,s(u))) = \sum_{\alpha=1}^d o_u(g(u,s_{\alpha}(u)))$$

(iv) (Halphen's formula) If s_1, \dots, s_d and s'_1, \dots, s'_e respectively are the v-roots of f and g, then

$$i_0(f,g) = \sum_{\substack{1 \le i \le d \\ 1 \le j \le e}} o_u(s_i - s'_j).$$

As a corollary, we have

$$i_0(f,g) = o_u \left(\operatorname{Res}(f,g) \right)$$

(v) If $f = f_1 \cdots f_k$, then

$$\mu_0(f) = \sum_{i=1}^k \mu_0(f_i) + \sum_{i \neq j} i_0(f_i, f_j) - k + 1.$$

Now we establish the relationship between the Minor number and the discriminant.

Lemma 6.7. With the notations above, we have

$$o_u(\text{Disc}(f)) = \mu_0(f) + d - 1.$$

Proof. By [12, Lem.6.5.7, p.148], we have that the number

$$i_0\left(pf+qu\frac{\partial f}{\partial u},\frac{\partial f}{\partial v}\right)$$

is independent of (p,q) provided $p,q \ge 0$ and $(p,q) \ne (0,0)$. It follows that

$$o_u\left(\operatorname{Disc}(f)\right) = i_0\left(f, \frac{\partial f}{\partial v}\right) = i_0\left(u\frac{\partial f}{\partial u}, \frac{\partial f}{\partial v}\right) = i_0\left(u, \frac{\partial f}{\partial v}\right) + \mu_0(f).$$

Recall that $f = v^d + \sum a_i(u)v^{d-i}$. We get that

$$i_0\left(u,\frac{\partial f}{\partial v}\right) = i_0(u,v^{d-1}) = d-1.$$

Hence, we get the desired equality $o_u(\text{Disc}(f)) = \mu_0(f) + d - 1$.

6.8. **Proof of Proposition 4.4.** Before presenting the proof of Proposition 4.4, we first recall our settings. We are given a reduced and irreducible curve C on X_d , from which we obtain the following objects:

(1) The branched Galois cover

$$\rho: X_d \to \mathbb{P}^2, \quad [x, y, z, w] \mapsto [x, y, z].$$

(2) There exists a homogeneous polynomial g(x, y, z) of degree e such that

$$(g) = m(C_1 + \dots + C_k)_{\sharp}$$

with $C_i^2 = C^2$ for i = 1, ..., k, and $k \ge 2$. (3) The subscheme \mathscr{C} on X_d defined by g, i.e.,

$$\mathscr{C}: g(x, y, z) = 0$$

(4) The curves D and E in \mathbb{P}^2 defined as

 $D: x^d + y^d + z^d = 0$ and E: g(x, y, z) = 0.

In addition, D and E intersect at only finitely many points.

- (5) Fix an intersection point $P = [x_0, y_0, z_0, w_0]$ of different irreducible components of \mathscr{C} . We have $w_0 = 0$. We also assume $x_0 \neq 0$ and $z_0 \neq 0$, and denote $p = [x_0, y_0, z_0]$.
- (6) Introduce the affine coordinates

$$u = \frac{y}{x} - \frac{y_0}{x_0}, \qquad v = \frac{z}{x} - \frac{z_0}{x_0}, \qquad \omega = \frac{w}{x},$$

and then the scheme $\mathscr{C}: g = 0$ on X_d is defined in $\mathbb{C}^3 = \{x \neq 0\} \subset \mathbb{P}^3$ by

$$h(u, v) = 0, \quad l(u, v) + \omega^d = 0,$$

where

$$h(u, v) = g\left(1, u + \frac{y_0}{x_0}, v + \frac{z_0}{x_0}\right)$$

and

$$l(u, v) = 1 + \left(u + \frac{y_0}{x_0}\right)^d + \left(v + \frac{z_0}{x_0}\right)^d.$$

(7) Viewed as an element in $\mathbb{C}\{u, v\}$, the function h is v-general of order b.

Now we are ready to give the proof of Proposition of 4.4. To begin with, we first decompose h in $\mathbb{C}\{u, v\}$ into irreducible factors

$$h = h_0 h_1 \cdots h_r;$$

where h_1, \ldots, h_r are irreducible Weierstrass polynomial in $\mathbb{C}\{u\}[v]$ of degrees b_1, \ldots, b_r respectively, and h_0 is an invertible element in $\mathbb{C}\{u, v\}$. Recall that by assumption h is v-general of order b, hence $b = b_1 + \cdots + b_r$.

For $i \leq r$, let s_i be a Puiseux series which is a root of h_i ; without loss of generality, we assume $\nu(s_i) = b_i$. Then, by Proposition 6.2 (iii) we have

$$h_i = \prod_{\alpha=1}^{b_i} (v - s_{i,\alpha}(u)), \ i \le r.$$

(Recall that by our convention, if s_i is written as

$$s_i(u) = \sum a_k u^{\frac{k}{b_i}}$$

then

$$s_{i,\alpha}(u) = \sum e^{2\pi\sqrt{-1}\alpha k/b_i} a_k u^{\frac{k}{b_i}}$$

for $\alpha = 1, \dots, b_i$.) Set

$$\phi_i(u,\omega) = \prod_{\alpha=1}^{b_i} \left(\omega^d + l\left(u, s_{i,\alpha}(u)\right) \right), \ i \le r$$

and

$$\Phi(u,\omega)=\phi_1\cdots\phi_r.$$

Then Φ is a Weierstrass polynomial in $\mathbb{C}\{u\}[\omega]$ of degree bd and moreover, the scheme \mathscr{C} is locally defined by the equation $\Phi(u, \omega) = 0$ in the local coordinate system (u, ω) on X_d .

- (A) To show that \mathscr{C} is reduced, it is sufficient to show
- (a_1) For $i \leq r, \phi_i$ is reduced; and
- (a₂) For $1 \le i < j \le r$, ϕ_i and ϕ_j are coprime as elements in $\mathbb{C}\langle\langle u \rangle\rangle[\omega]$.

For (a_1) , by definition of the intersection multiplicity, we have

$$o_u(l(u, s_{i,\alpha}(u))) = \frac{i_0(h_i, l)}{b_i} \le \frac{i_p(D, E)}{b_i} < \infty.$$

In particular, $l(u, s_{i,\alpha}(u)) \neq 0$; and thus $\omega^d + l(u, s_{i,\alpha}(u))$ is reduced. Moreover, recall that

$$l(u, v) = 1 + \left(u + \frac{y_0}{x_0}\right)^d + \left(v + \frac{z_0}{x_0}\right)^d$$

hence, for $\alpha \neq \beta$,

$$l(u, s_{i,\alpha}(u)) - l(u, s_{i,\beta}(u))$$

$$= (s_{i,\alpha}(u) - s_{i,\beta}(u)) \left(d\left(\frac{z_0}{x_0}\right)^{d-1} + r(u) \right)$$

$$\simeq s_{i,\alpha}(u) - s_{i,\beta}(u)$$

where $\lim_{u\to 0} r(u) = 0$ and \simeq means equality up to multiplication by an invertible element. Since $\nu(s_i) = b_i$ and $\alpha \neq \beta$, we have

$$s_{i,\alpha}(u) - s_{i,\beta}(u) \neq 0,$$

and thus

$$gcd\left(\omega^{d}+l\left(u,s_{i,\alpha}(u)\right),\omega^{d}+l\left(u,s_{i,\beta}(u)\right)\right)=1.$$

Consequently, ϕ_i is reduced.

For (a_2) , it suffices to show that for any $i \neq j$ and for all $\alpha \leq b_i, \beta \leq b_j$,

$$l(u, s_{i,\alpha}(u)) \neq l(u, s_{j,\beta}(u)).$$

Similar to the discussions above, we have

$$l(u, s_{i,\alpha}(u)) - l(u, s_{j,\beta}(u)) \simeq s_{i,\alpha}(u) - s_{j,\beta}(u)$$

Recall that h_i and h_j are coprime, so they do not have common roots in $\mathbb{C}\langle\langle u \rangle\rangle$. Therefore $s_{i,\alpha}(u) - s_{j,\beta}(u) \neq 0$.

Remark 6.9. Proposition 4.4 (A) also follows from the fact that for all $q \in E \setminus (E \cap D)$, the inverse image $\rho^{-1}(q) \subset \mathscr{C}$ consists of exactly d points.

(B) To compute $\mu_P(\mathscr{C})$, we use Lemma 6.7. Then

$$\mu_P(\mathscr{C}) = o_u \left(\text{Disc}(\Phi) \right) - bd + 1.$$

Hence, we need to calculate $o_u(\text{Disc}(\Phi))$. By Proposition 6.4 (vi), we have from $\Phi = \phi_1 \cdots \phi_r$ the following equality

$$o_u(\operatorname{Disc}(\Phi)) = \sum_{i=1}^r o_u(\operatorname{Disc}(\phi_i)) + \sum_{1 \le i < j \le r} o_u(\operatorname{Res}(\phi_i, \phi_j)^2).$$

Moreover, for $i \leq r$,

$$o_{u} (\operatorname{Disc}(\phi_{i}))$$

$$= \sum_{\alpha=1}^{b_{i}} o_{u} \left(\operatorname{Disc} \left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right)\right)\right)$$

$$+ \sum_{1 \leq \alpha < \beta \leq b_{i}} o_{u} \left(\operatorname{Res} \left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right), \omega^{d} + l\left(u, s_{i,\beta}(u)\right)\right)^{2}\right).$$

We will prove the following results.

(b₁) For $i \leq r$, $\sum_{\alpha=1}^{b_i} o_u \left(\text{Disc} \left(\omega^d + l \left(u, s_{i,\alpha}(u) \right) \right) \right) = (d-1)i_0(h_i, l),$

$$(b_2) \text{ For } i \leq r,$$

$$\sum_{1 \leq \alpha < \beta \leq b_i} o_u \left(\operatorname{Res} \left(\omega^d + l \left(u, s_{i,\alpha}(u) \right), \omega^d + l \left(u, s_{i,\beta}(u) \right) \right)^2 \right)$$

$$= do_u \left(\operatorname{Disc}(h_i) \right)$$

$$= d \left(\mu_0(h_i) + b_i - 1 \right).$$

$$(b_3) \text{ For } 1 \leq i < j \leq r,$$

 (b_4) We have

$$o_u(\text{Disc}(\Phi)) = (d-1)i_p(D, E) + d(\mu_p(E) + b - 1).$$

 $o_u \left(\operatorname{Res}(\phi_i, \phi_j)^2 \right) = do_u \left(\operatorname{Res}(h_i, h_j)^2 \right).$

For (b_1) , we have

Disc
$$\left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right)\right) \simeq \operatorname{Res}\left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right), \omega^{d-1}\right)$$

$$\simeq \left(l\left(u, s_{i,\alpha}(u)\right)\right)^{d-1};$$

hence,

$$o_u \left(\text{Disc} \left(\omega^d + l \left(u, s_{i,\alpha}(u) \right) \right) \right) = (d-1) \left(l \left(u, s_{i,\alpha}(u) \right) \right)$$
$$= \frac{(d-1)i_0(h_i, l)}{b_i}.$$

Summing over $\alpha = 1, \ldots, b_i$ gives the desired equality.

For (b_2) , we have

$$\operatorname{Res} \left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right), \omega^{d} + l\left(u, s_{i,\beta}(u)\right) \right)$$

=
$$\operatorname{Res} \left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right), l\left(u, s_{i,\beta}(u)\right) - l\left(u, s_{i,\alpha}(u)\right) \right)$$

$$\simeq \left(l\left(u, s_{i,\beta}(u)\right) - l\left(u, s_{i,\alpha}(u)\right) \right)^{d}.$$

Recall that

$$l(u, s_{i,\beta}(u)) - l(u, s_{i,\alpha}(u)) \simeq s_{i,\beta}(u) - s_{i,\alpha}(u).$$

Hence,

$$\operatorname{Res}\left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right), \omega^{d} + l\left(u, s_{i,\beta}(u)\right)\right) \\ \simeq \left(s_{i,\beta}(u) - s_{i,\alpha}(u)\right)^{d}.$$

It follows that

$$\sum_{1 \le \alpha < \beta \le b_i} o_u \left(\operatorname{Res} \left(\omega^d + l \left(u, s_{i,\alpha}(u) \right), \omega^d + l \left(u, s_{i,\beta}(u) \right) \right)^2 \right)$$

= $d \sum_{1 \le \alpha < \beta \le b_i} o_u \left(s_{i,\beta}(u) - s_{i,\alpha}(u) \right)^2$
= $do_u \left(\operatorname{Disc}(h_i) \right)$
= $do_u \left(\mu_0(h_i) + b_i - 1 \right),$

where the last equality follows from Lemma 6.7.

For (b_3) , we have

$$\operatorname{Res}(\phi_{i}, \phi_{j})^{2}$$

$$= \prod_{\alpha=1}^{b_{i}} \prod_{\beta=1}^{b_{j}} \operatorname{Res}\left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right), \omega^{d} + l\left(u, s_{j,\beta}(u)\right)\right)^{2}$$

$$\simeq \prod_{\alpha=1}^{b_{i}} \prod_{\beta=1}^{b_{j}} \operatorname{Res}\left(\omega^{d} + l\left(u, s_{i,\alpha}(u)\right), l\left(u, s_{j,\beta}(u)\right) - l\left(u, s_{i,\alpha}(u)\right)\right)^{2}$$

$$\simeq \prod_{\alpha=1}^{b_{i}} \prod_{\beta=1}^{b_{j}} \left(l\left(u, s_{j,\beta}(u)\right) - l\left(u, s_{i,\alpha}(u)\right)\right)^{2d}.$$

Recall that

$$l(u, s_{j,\beta}(u)) - l(u, s_{i,\alpha}(u)) \simeq s_{j,\beta}(u) - s_{i,\alpha}(u)$$

hence,

$$\operatorname{Res}(\phi_i, \phi_j)^2 \simeq \prod_{\alpha=1}^{b_i} \prod_{\beta=1}^{b_j} (s_{j,\beta}(u) - s_{i,\alpha}(u))^{2d}$$
$$\simeq \operatorname{Res}(h_i, h_j)^{2d}.$$

It follows that

$$o_u \left(\operatorname{Res}(\phi_i, \phi_j)^2 \right) = do_u \left(\operatorname{Res}(h_i, h_j)^2 \right)$$

as desired.

For (b_4) , we combine $(b_1)-(b_3)$ and get

$$o_u (\text{Disc}(\Phi)) = \sum_{i=1}^r \left((d-1)i_0(h_i, l) + do_u (\text{Disc}(h_i)) \right) + \sum_{1 \le i < j \le r} \left(do_u \left(\text{Res}(h_i, h_j)^2 \right) \right)$$

= $(d-1) \sum_{i=1}^r i_0(h_i, l) + do_u (\text{Disc}(h)).$

Recall that $i_0(h, l) = i_p(E, D)$; and by Lemma 6.7, we have

 $o_u\left(\mathrm{Disc}(h)\right) = \mu_p(E) + b - 1.$

The equality in (b_4) follows immediately.

Now we are ready to finish the proof of (B). From (b_4) and the equality

$$o_u(\operatorname{Disc}(\Phi)) = \mu_P(\mathscr{C}) + bd - 1,$$

we get

$$\mu_P(\mathscr{C}) = (d-1)i_p(D, E) + d(\mu_p(E) + b - 1) - bd + 1$$

= $(d-1)i_p(D, E) + d\mu_p(E) - d + 1.$

(C) Since C_1, \ldots, C_k are the irreducible components of \mathscr{C} , we have

$$\mu_P(\mathscr{C}) = \sum_{i \neq j} i_P(C_i, C_j) + \sum_{i=1}^k \mu_P(C_i) - k + 1.$$

Note that P is a fixed point under the action of the Galois group for the branched cover ρ , and the Galois group acts transitively on $\{C_1, \ldots, C_k\}$, hence

 $\mu_P(C_i) = \mu_P(C)$, for all i = 1, ..., k.

It follows that

$$\sum_{i \neq j} i_P(C_i, C_j) = \mu_P(\mathscr{C}) - k\mu_P(C) + k - 1.$$

Using (B), we get

$$\sum_{i \neq j} i_P(C_i, C_j) = (d-1)i_p(D, E) + d\mu_p(E) - k\mu_P(C) - (d-k).$$

Summing over all possible intersection points and using Lemma 4.2, we obtain

$$\sum_{i \neq j} C_i \cdot C_j = (d-1)D \cdot E + d \sum_{p \in D \cap E} \mu_p(E) - k \sum_{p \in D \cap E} \mu_{\rho^{-1}(p)}(C) - (d-k) \# (D \cap E).$$

Note that $D \cdot E = de$, the equality in (C) follows immediately.

The proof of Proposition 4.4 is complete now.

References

- T. Bauer, C. Bocci, S. Cooper, S. Di Rocco, M. Dumnicki, B. Harbourne, K. Jabbusch, A. L. Knutsen, A. Küronya, R. Miranda, J. Roé, H. Schenck, T. Szemberg, and Z. Teitler, *Recent developments and open problems in linear series*, in *Contributions to Algebraic Geometry*, P. Pragacz, ed., EMS Ser. Congr. Rep., European Math. Soc., Zürich, 2012, pp. 93–140. Impanga Conference on Algebraic Geometry, Bedlewo, Poland, Jul 04-10, 2010. 1
- [2] T. Bauer, S. Di Rocco, B. Harbourne, J. Huizenga, A. Lundman, P. Pokora, and T. Szemberg, Bounded negativity and arrangements of lines, Int. Math. Res. Not. IMRN 19(2015), 9456– 9471. 1
- [3] T. Bauer, B. Harbourne, A. L. Knutsen, A. Küronya, S. Müller-Stach, X. Roulleau, and T. Szemberg, Negative curves on algebraic surfaces, Duke Math. J. 162(2013), 1877–1894. 1, 1, 1
- [4] T. Bauer, P. Pokora, and D. Schmitz, On the boundedness of the denominators in the Zariski decom position on surfaces, J. Reine Angew. Math. 733(2017), 251–259. 1
- [5] R. Benedetti, and J.-J. Risler, Real algebraic and semi-algebraic Sets, Hermann, 1990. 6.3
- [6] E. Casas-Alvero, Singularities of plane curves, Cambridge University Press, 2000. 5.1, 6.1, 6.5
- [7] C. Ciliberto, A.L. Knutsen, J. Lesieutre, V. Lozovanu, R. Miranda, Y. Mustopa, and D. Testa, A few questions about curves on surfaces, Rend. Circ. Mat. Palermo (2) 66 (2017), 195–204. 1
- [8] C. Galindo, F. Monserrat, and C.-J. Moreno-Ávila, On weighted bounded negativity for rational surfaces, arXiv:2408.05466. 1
- [9] G.-M. Greuel, C. Lossen, and E. Shustin, Introduction to singularities and deformations, Springer, 2007. 6.5
- [10] F. Hao, Weak bounded negativity conjecture, Proc. Amer. Math. Soc. 147 (2019), 3233–3238.
 1
- [11] R. Laface, and P. Pokora, Towards the weighted bounded negativity conjecture for blow-ups of algebraic surfaces, Manuscripta Math. 163 (2020), 361–373. 1
- [12] C.T.C. Wall, Singular points of plane curves, Cambridge University Press, 2004.
- 4.3, 6.5

HEFEI NATIONAL LABORATORY, HEFEI 230088, CHINA *Email address:* wzhj@ustc.edu.cn