# Theory and Applications of Kernel Stein Discrepancy on Riemannian Manifolds

Xiaoda Qu† and Baba C. Vemuri‡ †Department of Statistics ‡Department of CISE University of Florida

January 3, 2025

#### Abstract

Distributional comparison is a fundamental problem in statistical data analysis with numerous applications in a variety of fields including but not limited to Sciences and Engineering. Numerous methods exist for distributional comparison but Kernel Stein Discrepancy (KSD) has gained significant popularity in recent times. In this paper, we present a novel mathematically rigorous and consistent generalization of KSD to Riemannian manifolds. We first generalize the Stein's operator to Riemannian manifolds and use it to establish the Stein's Lemma on Riemannian manifolds. Then we define a novel Stein class and use it to develop what we call qualified kernels that are used to define the KSD in closed form on Rimeannian manifolds. We present several examples of our theory applied to commonly encountered Riemannian manifolds in applications namely, the Riemannian homogeneous spaces, for example, the n-sphere, the Grassmanian, Stiefel, the manifold of symmetric positive definite matrices and others. On these aforementioned manifolds, we consider a variety of distributions with intractable normalization constants and derive closed form expressions for the KSD and the minimum KSD estimator (mKSDE). Several theoretical properties of mKSDE are established and we present results of comparison between mKSDE and MLE on a widely popular example, the sphere.

# 1 Introduction

Data residing in curved spaces have recently received growing attention in numerous fields of Science and Engineering. To model their underlying curved geometry, it is natural to model the space in which they reside with known manifold geometries for example, (i) the Stiefel manifold,  $V_r(N)$ , commonly used to model the space of directional data in fields of computer vision and medical image analysis [\[10,](#page-26-0) [64\]](#page-29-0), dynamic system [\[7\]](#page-25-0) and rigid body motion [\[36,](#page-27-0) [50,](#page-28-0) [57\]](#page-28-1), (ii) Grassmann manifold  $\mathcal{G}_r(N)$  in signal processing [\[12,](#page-26-1) [46,](#page-28-2) [60\]](#page-28-3), shape analysis [\[23,](#page-26-2) [5,](#page-25-1) [68\]](#page-29-1) and image processing [\[13,](#page-26-3) [14,](#page-26-4) [59\]](#page-28-4), (iii) covariance matrices are modeled as points on the manifold of symmetric positive definite (SPD) matrices  $\mathcal{P}(N)$  in diffusion magnetic resonance imaging [\[4,](#page-25-2) [18,](#page-26-5) [65,](#page-29-2) [31\]](#page-27-1) and brain computer interfaces [\[9,](#page-25-3) [67\]](#page-29-3).

However, due to the lack of vector space structure, significant complications arise in the formulation and application of statistical methods to such curved spaces. e.g., the data points lying on manifolds can not be simply summed up, thus the notion of the classical arithmetic mean is not meaningful in general. Among all the challenges, the most significant one is the issue of normalization constant associated with probability distributions defined on the manifold-valued (M-valued) random variables, which arises in distributional comparison, parametric estimation and numerous practical applications. Even the simplest distribution on the simplest curved manifold, e.g., the von Mises Fisher distribution  $p(x) \propto \exp(\mu^\top x)$  on a sphere  $\mathbb{S}^{d-1}$ , has a normalization constant that is intractable. Furthermore, the KL-divergence  $KL(p, q) := \mathbb{E}_p[log \frac{p}{q}]$ , which is the most commonlyused loss function in parameter estimation, distributional comparison and neural network training, highly relies on the computation of normalization constant of  $p$  and  $q$ .

In practice, approximating these constants and their derivative with respect to the parameters of the distribution requires the use of numerical methods such as the gradient descent and/or its variants, which has been resorted to by many researchers in statistics, machine learning and robotics literature [\[21,](#page-26-6) [22,](#page-26-7) [29,](#page-27-2) [30\]](#page-27-3) but at the expense of a high computation cost. It would of course be more desirable to fully avoid computing this intractable constant and simultaneously achieve high accuracy in parameter estimation. In fact, this will be the main objective of this paper.

Kernel Stein Discrepancy (KSD), a normalization free loss function was first introduced by Liu et al. [\[37\]](#page-27-4) as a measure of goodness of fit and for model evaluation. KSD measures the difference between distributions by using a combination of the so called Stein's method and the well established reproducing kernels Hilbert space (RKHS) theory. KSD has since been extensively researched on, including various aspects within a general framework  $[35, 43, 44, 49]$  $[35, 43, 44, 49]$  $[35, 43, 44, 49]$  $[35, 43, 44, 49]$  $[35, 43, 44, 49]$  $[35, 43, 44, 49]$ , characterization of its scope  $[24, 25, 54]$  $[24, 25, 54]$  $[24, 25, 54]$  $[24, 25, 54]$  $[24, 25, 54]$ , exploration of its asymptotic properties relating to minimization  $[2, 48]$  $[2, 48]$ , its diverse applications [\[11,](#page-26-10) [37,](#page-27-4) [41\]](#page-27-6), and generalizations to manifolds [\[3,](#page-25-5) [53,](#page-28-10) [66\]](#page-29-4).

#### 1.1 Context

At its core, the KSD consists of an RKHS,  $\mathcal{H}_{\kappa}$ , defined by a kernel function  $\kappa$  on  $\mathbb{R}^d$ , known as the Stein's class. Additionally, it incorporates a Stein's operator  $S_p$ , dependent on the candidate distribution  $p$  but independent of its normalizing constant. The role of the Stein operator is to map elements from  $\mathcal{H}_{\kappa}$  to real integrable functions. These operators must satisfy *Stein's identity*, given by,  $\mathbb{E}_p[\mathcal{S}_p f] = 0$  for all  $f \in \mathcal{H}_{\kappa}$ . This leads us to the Stein pair defined as:

### Stein pair

A pair  $(S_p, \mathcal{H}_\kappa)$  consisting of a *Stein operator*  $S_p$  and a *Stein class*  $\mathcal{H}_\kappa$  satisfying *Stein's identity*, i.e.,  $\mathbb{E}_p[\mathcal{S}_p f] = 0$  for all  $f \in \mathcal{H}_\kappa$ , is called a *Stein pair.* 

With these foundational elements in place, the KSD between distributions  $p$  and  $q$  can be defined as follows:

<span id="page-1-1"></span>
$$
\text{KSD}(p, q) := \sup \left\{ \mathbb{E}_q[\mathcal{S}_p f] : f \in \mathcal{H}_\kappa, \|f\|_{\mathcal{H}_\kappa} \le 1 \right\}. \tag{1}
$$

# 1.1.1 KSD on  $\mathbb{R}^d$

Most of the existing works have focused on the KSD defined on  $\mathbb{R}^d$ . Let  $\mathcal{H}_\kappa$  be a RKHS on  $\mathbb{R}^d$ associated with kernel  $\kappa$ , and  $\mathcal{H}^d_{\kappa} := \mathcal{H}_{\kappa} \times \cdots \times \mathcal{H}_{\kappa}$  be the *d*-times product of  $\mathcal{H}_{\kappa}$ , equipped with the inner product  $\langle f, g \rangle_{\mathcal{H}_{\kappa}^d} = \sum_{l=1}^d \langle f_l, g_l \rangle_{\mathcal{H}_{\kappa}}$  for  $f = (f_1, \ldots, f_d)$  and g in  $\mathcal{H}_{\kappa}^d$ . Given a differentiable density p on  $\mathbb{R}^d$ , the most commonly adopted [\[2,](#page-25-4) [11,](#page-26-10) [25,](#page-26-9) [37,](#page-27-4) [41,](#page-27-6) [48\]](#page-28-9) Stein operator  $\mathcal{A}_p$  on  $\mathbb{R}^d$  is defined as

<span id="page-1-0"></span>
$$
\mathcal{A}_p: f \mapsto \sum_{l=1}^d \left[ \frac{\partial f_l}{\partial x^l} + f^l \frac{\partial}{\partial x^l} \log p \right], \quad f \in \mathcal{H}_\kappa^d. \tag{2}
$$

KSD is then obtained by substituting  $(2)$  into  $(1)$ , i.e.,

<span id="page-2-0"></span>
$$
\text{KSD}(p,q) := \sup \left\{ \mathbb{E}_q[\mathcal{A}_p f] : f \in \mathcal{H}_\kappa^d, \|f\|_{\mathcal{H}_\kappa^d} \le 1 \right\}. \tag{3}
$$

Clearly, one can easily see from the definition and Stein's identity that  $KSD(p, q) \geq 0$  and  $KSD(p, p)$ 0. In fact, as demonstrated in [\[2,](#page-25-4) [11,](#page-26-10) [25,](#page-26-9) [37,](#page-27-4) [41,](#page-27-6) [48\]](#page-28-9), if the kernel function k is  $C_0$ -universal [\[8\]](#page-25-6), then KSD will uniquely characterize  $p$ , i.e., for  $p$  and  $q$  that satisfy mild regularity conditions, there is  $KSD(p, q) = 0 \Leftrightarrow p = q$ . Notably, the computation of  $\mathcal{A}_p$  is independent of the normalizing constant of  $p$ , so is the associated KSD in  $(3)$ .

The minimum kernel Stein discrepancy estimator (MKSDE) is one of the most important application of KSD, which was first proposed in  $[2]$  and further researched on in  $[41, 48]$  $[41, 48]$ . The MKSDE minimizes the KSD between the empirical distribution of the samples and a parametrized family  $p_{\theta}$ , to acquire an estimate  $p_{\theta^*}$  of the underlying distribution from the samples. The MKSDE converges under mild regularity conditions, thus can serve as a normalization-free alternative to MLE.

#### 1.1.2 Existing generalizations to manifold

In contrast to the extensive work on KSD in  $\mathbb{R}^d$ , generalizations of KSD to Riemannian manifolds is scarce and the existing generalizations are somewhat restrictive as will be evident from the following discussion. In [\[3\]](#page-25-5), Barp et al. adopted the Stein's operator  $f \mapsto \Delta f + g(\nabla \log p, \nabla f)$  and the Sobolev space as their RKHS on compact manifolds. It is a significant challenge to identify a closed form kernel for such a large RKHS on a curved manifold. An alternative approach to tackle this challenge, not discussed in [\[3\]](#page-25-5), is to restrict a Sobolev-type kernel from  $\mathbb{R}^d$  to the manifold, as suggested in [\[19,](#page-26-11) Thm. 5]. However, this approach does not ensure a closed-form expression for the kernel. In [\[66\]](#page-29-4), Xu et al. adopted the same Stein's operator  $\mathcal{A}_p$  in [\(2\)](#page-1-0), replacing the coordinate  $x^i$  with the local coordinate chart on the manifold, and applying Stokes's theorem to show Stein's identity. However, there is no global chart on any compact manifold, e.g., sphere  $\mathbb{S}^{N-1}$ , Stiefel manifold  $\mathcal{V}_r(N)$ , Grassmann manifold  $\mathcal{G}_r(N)$  and many others. In order for Stein's identity based on the Stokes's theorem to hold in their method, the target density p must vanish outside the singular boundary of the chart on such compact manifolds. In [\[53\]](#page-28-10), Qu et al. also adopted [\(2\)](#page-1-0), but replaced the vector fields  $\frac{\partial}{\partial x^l}$  in [\(2\)](#page-1-0) with the left invariant vector fields utilizing the Lie groups structure, so as to circumvent the issue of local coordinates on Lie groups. However, there are many manifolds that are widely encountered in applications, including the sphere  $\mathbb{S}^{N-1}$ , Stiefel manifold  $V_r(N)$ , Grassmann manifold  $\mathcal{G}_r(N)$  and the manifold of symmetric positice definite matrices  $\mathcal{P}(N)$ that will be addressed in this work, which do not possess a Lie group structure.

#### 1.2 Our work and contributions

Our contributions in this work are itemized below.

• KSD on general Riemannian manifolds: In §[3,](#page-4-0) we propose a novel Stein operator on the general complete Riemannian manifold and study its properties. Unlike the previous works, it leads to a normalization-free KSD that is not only applicable to all complete Riemannian manifold but also independent of the choice of local coordinates. More importantly, our KSD does not require an excessively large RKHS to characterize the target distribution (see Thm. [3.5\)](#page-8-0) and thus its Stein class can be chosen as any one of the most widely-used RKHSs with easy-to-compute kernels (see e.g. Ex. [2.1\)](#page-4-1). This significantly increases its applicability to machine learning, engineering and other fields. In §[3.4,](#page-8-1) we show that the KSD can be further simplified on Riemannian homogeneous spaces utilizing isometry structure and killing fields. We will elaborate on these topics subsequently.

- MKSDE and its applications: In §[4,](#page-10-0) we introduce the MKSDE obtained by minimizing our novel KSD and its asymptotic properties. In §[4.4,](#page-13-0) we introduce the composite goodness of fit test, one of the most important application based on MKSDE. These results follow the same outline in our previous work [\[53\]](#page-28-10) on Lie groups, but we generalize the theory so that it is applicable to all complete Riemannian manifolds. In §[6,](#page-23-0) we present two applications of our KSD to one of the most widely-encountered manifolds in science and engineering namely, the Stiefel manifold  $\mathcal{V}_2(3)$ . Specifically, the first experiments [6.1](#page-23-1) will address the issue of the normalization constant that arises in MLE and how the estimation obtained using proposed normalization-free KSD yields far more accurate parameter estimates compared to MLE that uses approximations for the normalization constant. The second experient justifies the power of the composite goodness of fit test based on our MKSDE.
- Explicit closed forms: The most significant property of our KSD and MKSDE, is that they have closed forms on some of the most widely-encountered manifolds, including Stiefel manifold  $\mathcal{V}_r(N)$  (including the sphere  $\mathbb{S}^{N-1}$  and the rotation group  $SO(N)$  as  $\mathcal{V}_1(N) = \mathbb{S}^{N-1}$  and  $V_{N-1}(N) = SO(N)$ , Grassmann manifold  $G_r(N)$  and the manifold of symmetric positive definite matrices  $\mathcal{P}(N)$ . In §[5,](#page-13-1) we will compute the explicit form of our KSD and our MKSDE for the exponential family of distributions on these manifolds, which will facilitate the usage of our method in practice.

# 2 Mathematical Background

In this section, we will introduce several pivotal theorems that will be used subsequently in this work. To stay within the journal imposed page limits, definitions of several relevant concepts used throughout this paper, including the Riemannian manifold  $M$ , Riemannian metric  $g$ , Riemannian distance  $\rho$ , vector field D and its pointwise length  $|D| := \sqrt{g(D, D)}$ , local curves  $[\mathfrak{c}(t)]$ , volume measure  $\mu$ , divergence operator div, Riemannian gradient  $\nabla$ , reproducing kernel Hilbert space (RKHS)  $\kappa$ ,  $C_0$ -universality, and the notion of Bochner integral, are introduced in the supplementary material.

First three theorems will play fundamental roles in the construction of our Stein operator in §[3.1.](#page-4-2) The first theorem [\[20\]](#page-26-12) generalizes the classical Stokes's theorem, which ensures the Stein's identity holds. The second theorem  $[61, \text{Lem. } 4.34]$  demonstrates the connection between the differentiability of the kernel and the differentiability of the functions in its associated RKHS, which ensures all functions in the RKHS are differentiable so that the Stein operator is applicable. The third theorem [\[61,](#page-29-5) §A.5.4] captures the interchangeability between the Bochner integral and a continuous linear functional, which serves as the key to obtaining the closed form of KSD in Thm. [3.1.](#page-7-0)

<span id="page-3-0"></span>**Theorem 2.1** (Divergence theorem). Let  $M$  be a complete Riemannian manifold. For a locally Lipschitz continuous vector field  $D$  on  $M$  such that  $|D|$  and div  $D$  are both integrable w.r.t volume measure  $\mu$ , the following identity holds:  $\int_M \text{div } D d\mu = 0$ .

<span id="page-3-1"></span>**Theorem 2.2.** Suppose  $\kappa$  is a twice continuous differentiable kernel on manifold M, i.e.,  $\kappa \in$  $C^2(M\times M)$ , then  $\mathcal{H}_\kappa\subset C^1(M)$ . Furthermore, for a tangent vector  $D\in T_{x_0}M$  at  $x_0\in M$ , we have  $(D\kappa)_{x_0} \in \mathcal{H}_\kappa$  and  $Df(x_0) = \langle f, (D\kappa)_{x_0} \rangle_{\mathcal{H}_\kappa}$ . Here  $(D\kappa)_{x_0}$  represents the function obtained by letting D act on the first argument of  $\kappa$  and fix the first argument at  $x_0$ .

<span id="page-3-2"></span>**Theorem 2.3.** Let  $X : \Omega \to \mathcal{B}$  be a B-valued random variable  $(r.v.)$  from a probability space  $(\Omega, \mathcal{F}, P)$  to a Banach space B. We say X is Bochner integrable if  $\mathbb{E} \|X\| < +\infty$ , in which case there exists an unique  $b_0 \in \mathcal{B}$ , such that for any continuous linear functional f on  $\mathcal{B}$ ,  $f(b_0) = \mathbb{E}f(X)$ . We define the expectation of X to be  $\mathbb{E}X := b_0$ .

The next theorem, called Mercer's theorem [\[61,](#page-29-5) §4.5], ensures the spectral decomposition of a kernel  $\kappa$ , which is pivotal in the depiction of the approximating distribution of the empirical KSDs in Thm. [3.7.](#page-10-1)

<span id="page-4-3"></span>**Theorem 2.4** (Mercer's theorem). Suppose  $\kappa : \Omega \times \Omega \to \mathbb{R}$  is a symmetric semi-positive definite bivariate function on a  $\sigma$ -finite measure space  $(\Omega, \mathcal{F}, \nu)$  and satisfies  $\kappa \in L^2(\nu \times \nu)$ , then there exists a sequence of positive numbers  $\lambda_k$ ,  $k \geq 1$  and a sequence of orthonormal eigen-functions  $\phi_k \in L^2(\nu)$ such that

$$
\sum_{k=1}^{n} \lambda_k \phi(x)\phi(y) \to \kappa(x, y) \quad \text{in} \quad L^2(\nu \times \nu), \text{ as } n \to \infty.
$$

Then  $\{\lambda_k\}$  are said to be the eigenvalues of  $\kappa$ .

The next theorem [\[42,](#page-27-7) Thm. 17][\[56,](#page-28-11) Thm. 2] characterizes the radial kernels on  $\mathbb{R}^d$ , which are a class of kernels that satisfy our mathematical requirements perfectly and will be extensively used in §[5.](#page-13-1)

**Theorem 2.5.** Suppose  $\psi : [0, +\infty) \to \mathbb{R}$  is a continuous function. Then

- 1. The bivariate function  $\kappa(x, y) := \exp(-\psi(||x-y||_{\mathbb{R}^d}^2))$  for  $x, y \in \mathbb{R}^d$  is a kernel on  $\mathbb{R}^d$  if and only if there exists a finite Borel measure  $\lambda$  on  $[0, +\infty)$  such that  $\psi(t) := -\log \int_0^{+\infty} e^{-t\sigma} d\lambda(\sigma)$ ,  $t \ge$ 0. Such kernel  $\kappa$  is said to be a radial kernel on  $\mathbb{R}^d$ .
- 2. A radial kernel  $\kappa$  is  $C_0$ -universal on  $\mathbb{R}^d$  if  $\lambda$  is not concentrated at 0.

<span id="page-4-1"></span>Example 2.1 (Radial kernels). Following radial kernels  $[42]$  will be used in §[5:](#page-13-1)

- Gaussian kernel: $\kappa(x, y) = \exp(-\frac{\tau}{2} ||x y||_{\mathbb{R}^d}^2)$  for some  $\tau > 0$ .
- Inverse quadratic kernel:  $\kappa(x, y) = (\beta + ||x y||_{\mathbb{R}^d}^2)^{-\gamma}$  for some  $\beta, \gamma > 0$ .

# <span id="page-4-0"></span>3 KSD on Riemannian Manifolds

# <span id="page-4-2"></span>3.1 Stein operator on Riemannian Manifolds

First we seek to generalize the Stein pair  $(\mathcal{A}_p, \mathcal{H}_\kappa^d)$  on  $\mathbb{R}^d$  defined in Eq. [\(2\)](#page-1-0) to Riemannian manifolds and then use it to develop the KSD. It is straightforward to see that following assumptions must be maintained on manifolds.

Standing Assumptions 1,2 and 3

- 1. The Riemannian manifold  $M$  is complete and connected so that the generalized divergence theorem (Thm. [2.1\)](#page-3-0) holds.
- 2. The probability density p of the target distribution w.r.t the volume measure  $\mu$  on M is locally Lipschitz continuous so that the derivative of  $p$  is well-defined by Rademacher theorem [\[17,](#page-26-13) Thm. 3.1.6].
- **3.** The kernel function  $\kappa \in C^2(M \times M)$ , so that all functions in its associated RKHS  $\mathcal{H}_{\kappa}$  are continuously differentiable by Thm. [2.2.](#page-3-1)

As previously stated, there is no global chart on a curved manifold, thus the partial derivatives  $\{\frac{\partial}{\partial x^l}\}\$  do not generalize to Riemannian manifolds globally. To have corresponding global derivatives, we should resort to the vector fields  $D<sup>l</sup>$  on manifolds instead, in which case the resulting Stein operator maps f to  $\sum_{l=1}^{m} [D^{l} f_{l} + f_{l} D^{l} \log p]$ . However, one should note that the fact, div  $\frac{\partial}{\partial x^{l}} = 0$ plays an important role in the Stein operator  $\mathcal{A}_p$ , since

<span id="page-5-0"></span>
$$
\sum_{l=1}^{d} \operatorname{div} (fp \frac{\partial}{\partial x^{l}}) = p \sum_{l=1}^{d} \left( \frac{\partial f}{\partial x^{l}} + f \frac{\partial}{\partial x^{l}} \log p + f \operatorname{div} \frac{\partial}{\partial x^{l}} \right)
$$
  

$$
= p \mathcal{A}_{p} f + pf \sum_{l=1}^{d} \operatorname{div} \frac{\partial}{\partial x^{l}} = p \mathcal{A}_{p} f,
$$
 (4)

The first equality is based on the property [\[52,](#page-28-12) Exer. 2.5.5] of the divergence operator div( $fD$ ) =  $Df + f$  div D. This equation leads to Stein's identity for  $\mathcal{A}_p$  as

$$
\mathbb{E}_p[\mathcal{A}_p f] = \int_{\mathbb{R}^d} p \mathcal{A}_p f dx = \sum_{l=1}^d \int_{\mathbb{R}^d} \text{div}(fp \frac{\partial}{\partial x^l}) dx = 0,
$$

by divergence theorem (Thm. [2.1\)](#page-3-0).

The above derivation process falls apart on manifolds, since div  $D^l \neq 0$  in general and thus the last equality in Eq.[\(4\)](#page-5-0) fails to hold. To preserve the Stein's identity on manifolds, we keep the term in Eq.[\(4\)](#page-5-0) that contains div  $D<sup>l</sup>$ , and define the operator as:

Stein's operator on M

Definition 3.1 (Stein operator on M). Given a group of vector fields  $\{D^{l}\}_{l=1}^{m}$  on M, the *Stein* operator  $\mathcal{T}_p$  on M is defined as

<span id="page-5-1"></span>
$$
\mathcal{T}_p f = \sum_{l=1}^m \left[ D^l f_l + f_l D^l \log p + f_l \operatorname{div} D^l \right], \quad f \in \mathcal{H}_\kappa^m. \tag{5}
$$

Here,  $D^l \log p$  is set to 0 whenever  $p = 0$ .

Remark. In contrast to the Euclidean case, the number of vector fields m here is not compelled to equal the dimension  $d := \dim M$  of M, as long as Stein's identity holds. In fact, we will usually need more vector fields than in the case of Euclidean space for manifolds, to preserve the properties that the Stein operator on Euclidean space possesses, as we will elaborate in §[3.3.](#page-7-1)

*Example* 3.1 (Euclidean space). For the case  $M = \mathbb{R}^d$ , let  $D^l = \frac{\partial}{\partial x^l}$ ,  $1 \leq l \leq d$ . Since div  $\frac{\partial}{\partial x^l} = 0$ , then Stein operator  $\mathcal{T}_p$  on manifold [\(5\)](#page-5-1) will degenerate to the Stein operator  $\mathcal{A}_p$  on Euclidean space  $(2).$  $(2).$ 

*Example* 3.2 (Lie groups). The case where M is a Lie group was presented in [\[53\]](#page-28-10). Let  $D^l$ ,  $1 \leq l \leq d$ be the left invariant vector fields on G, then div  $D^l = D^l \Delta$ , where  $\Delta$  is the modular function. For illustrative examples, we refer the readers to our recent work [\[53\]](#page-28-10).

# 3.2 KSD on Riemannian Manifolds

With the Stein operator  $\mathcal{T}_p f$  in hand, we can now define the KSD on M as follows:

Definition 3.2 (KSD on Riemannian manifolds). Given the Stein operator  $\mathcal{T}_p f$  in [\(5\)](#page-5-1), the kernel Stein discrepancy (KSD) on M is defined by plugging  $\mathcal{T}_p$  into [\(1\)](#page-1-1), i.e.,

<span id="page-6-0"></span>
$$
\text{KSD}(p,q) := \sup \left\{ \mathbb{E}_q[\mathcal{T}_p f] : f \in \mathcal{H}_\kappa^m, \|f\|_{\mathcal{H}_\kappa^m} \le 1 \right\}. \tag{6}
$$

A remarkable property of KSD on  $\mathbb{R}^d$  [\[11,](#page-26-10) [37\]](#page-27-4), as well as the KSD on Lie groups [\[53\]](#page-28-10) is that it has a closed form represented by an integral, which facilitates its use in the practical applications. The KSD on Riemannian manifold defined in [\(6\)](#page-6-0) preserves this property, as we derive next.

For notational convenience, we specify each component  $\mathcal{T}_p^l$  of the Stein operator  $\mathcal{T}_p$  as  $\mathcal{T}_p^l: h \mapsto$  $D^l f + h D^l \log p + h \operatorname{div} D^l$  for  $h \in \mathcal{H}_{\kappa}$ . Thus,  $\mathcal{T}_p f = \sum_{l=1}^m \mathcal{T}_p^l f_l$ , for  $f = (f_1, \ldots, f_m) \in \mathcal{H}_{\kappa}^m$ . We let  $\mathcal{T}_{p}^{l} \kappa$  denote the bivariate function obtained by letting  $\mathcal{T}_{p}^{l}$  act on the first argument of  $\kappa$ , let  $(\mathcal{T}_{p}^{l} \kappa)_{x}$ represent the univariate function obtained by fixing the first argument of  $\mathcal{T}_p^l \kappa$  at x, and let  $(\vec{\mathcal{T}}_p \kappa)_x$ represent the vector-valued function  $((\mathcal{T}_p^1\kappa)_x,\ldots,(\mathcal{T}_p^m\kappa)_x).$ 

By Thm. [2.2,](#page-3-1) it is straightforward that  $(\mathcal{T}_p^l \kappa)_x \in \mathcal{H}_\kappa$  and thus  $(\vec{\mathcal{T}}_p \kappa)_x \in \mathcal{H}_\kappa^m$  for all x. Further-more, Thm. [2.2](#page-3-1) implies that  $\langle h, (\mathcal{T}_{p}^{l} \kappa)_{x} \rangle_{\mathcal{H}_{\kappa}} = \mathcal{T}_{p}^{l} h$  for all  $h \in \mathcal{H}_{\kappa}$ , thus if we substitute the function  $(\mathcal{T}_{p}^{l}\kappa)_{y}$  for h, we will have

<span id="page-6-1"></span>
$$
\kappa_p(x, y) := \langle (\vec{\mathcal{T}}_p \kappa)_y, (\vec{\mathcal{T}}_p \kappa)_x \rangle_{\mathcal{H}_\kappa^m} = \sum_{l=1}^m \langle (\mathcal{T}_p^l \kappa)_y, (\mathcal{T}_p^l \kappa)_x \rangle_{\mathcal{H}_\kappa}
$$

$$
= \sum_{l=1}^m \mathcal{T}_{p(x)}^l \mathcal{T}_{p(y)}^l \kappa = \sum_{l=1}^m \mathcal{T}_{p(y)}^l \mathcal{T}_{p(x)}^l \kappa. \tag{7}
$$

Here  $\mathcal{T}_{p(x)}^l$  and  $\mathcal{T}_{p(y)}^l$  represent the operators acting on the first argument x of  $\kappa$  and the second argument y of  $\kappa$  respectively, and  $\mathcal{T}_{p(x)}^l \mathcal{T}_{p(y)}^l \kappa$  represent the bivariate function by letting  $\mathcal{T}_{p(y)}^l$  act on y first and then let  $\mathcal{T}_{p(x)}^l$  act on x of  $\kappa$ . Note that  $\kappa_p(x, y)$  is semi-positive definite.

As  $x \mapsto (\vec{T}_{p} \kappa)_x$  is a continuous map from M to  $\mathcal{H}_{\kappa}^m$ , given a M-valued random variable X,  $(\vec{\mathcal{T}}_p \kappa)_X$  is a  $\mathcal{H}_\kappa^m$ -valued random variable. Note that  $\kappa_p(x,x) = \langle (\vec{\mathcal{T}}_p \kappa)_x, (\vec{\mathcal{T}}_p \kappa)_x \rangle_{\mathcal{H}_\kappa^m} = \|(\vec{\mathcal{T}}_p \kappa)_x\|_{\mathcal{H}_\kappa^m}^2$ thus if  $\sqrt{\kappa_p(X,X)}$  is integrable, then  $(\vec{\mathcal{T}}_p\kappa)_X$  is Bochner integrable. We denote its expectation by  $\mathbb{E}(\vec{\mathcal{T}}_{p}\kappa)_{X}$ . Given a M-valued r.v. X that follows the distribution q, we have

$$
\text{KSD}(p,q) = \sup_{\|f\|_{\mathcal{H}_\kappa^m} \le 1} \mathbb{E}[\mathcal{T}_p f(X)] = \sup_{\|f\|_{\mathcal{H}_\kappa^m} \le 1} \mathbb{E}_q \langle f, (\vec{\mathcal{T}}_p \kappa)_X \rangle_{\mathcal{H}_\kappa^m}
$$
  
Thm. 2.3 = 
$$
\sup_{\|f\|_{\mathcal{H}_\kappa^m} \le 1} \langle f, \mathbb{E}(\vec{\mathcal{T}}_p \kappa)_X \rangle_{\mathcal{H}_\kappa^m} = \|\mathbb{E}(\vec{\mathcal{T}}_p \kappa)_X\|_{\mathcal{H}_\kappa^m}.
$$

Let Y be another q-distributed M-valued r.v. independent with X. By Thm. [2.3,](#page-3-2) we have

$$
\|\mathbb{E}(\vec{\mathcal{T}}_{p}\kappa)_{X}\|_{\mathcal{H}_{\kappa}^{m}}^{2} = \langle \mathbb{E}(\vec{\mathcal{T}}_{p}\kappa)_{X}, \mathbb{E}(\vec{\mathcal{T}}_{p}\kappa)_{Y} \rangle_{\mathcal{H}_{\kappa}^{m}} = \mathbb{E}_{X} \mathbb{E}_{Y} \langle (\vec{\mathcal{T}}_{p}\kappa)_{X}, (\vec{\mathcal{T}}_{p}\kappa)_{Y} \rangle_{\mathcal{H}_{\kappa}^{m}} = \mathbb{E}_{X} \mathbb{E}_{Y} \kappa_{p}(X, Y) = \iint_{M \times M} \kappa_{p}(x, y) q(x) q(y) \mu(dx) \mu(dy).
$$
\n(8)

We summarize this result into following theorem:

#### KSD in Closed Form

<span id="page-7-0"></span>**Theorem 3.1** (Closed form). Suppose  $\sqrt{\kappa_p(x, x)}$  is q-integrable, then the KSD on M defined in Eq.  $(6)$  satisfies

<span id="page-7-2"></span>
$$
KSD2(p,q) = \iint \kappa_p(x,y)q(x)q(y)\mu(dx)\mu(dy).
$$
 (9)

Here  $\kappa_p(x, y)$  is the function introduced in [\(7\)](#page-6-1).

Stein's Identity on M

**Theorem 3.2** (Stein's identity). Suppose  $\sqrt{\kappa(x,x)} \sum_{l=1}^{m} |D^l|$  and  $\sqrt{\kappa_p(x,x)}$  are all pintegrable, then  $\mathbb{E}_p[\mathcal{T}_p f] = 0$  for all  $f \in \mathcal{H}_\kappa$ , that is,  $\text{KSD}(p, p) = 0$  or equivalently  $p = q \Rightarrow$  KSD $(p, q) = 0$ .

#### <span id="page-7-1"></span>3.3 Characterization of KSD

The Stein operator  $\mathcal{T}_p$  in [\(5\)](#page-5-1) satisfies Stein's identity, i.e.,  $\mathbb{E}_p[\mathcal{T}_p f] = 0$  for  $f \in \mathcal{H}_\kappa^m$ , which is equivalent to stating that,  $p = q \Rightarrow \text{KSD}(p, q) = 0$ . However, to serve as a loss functions in practice, the KSD should satisfy the reverse direction, i.e.,  $KSD(p,q) = 0 \Rightarrow p = q$  with mild regularity conditions on  $p$  and  $q$ , in which case we say the KSD characterizes  $p$ . The original Stein operator  $\mathcal{A}_p$  on  $\mathbb{R}^d$  in [\(2\)](#page-1-0) does distinguish p from all locally Lipschitz continuous q when the kernel function  $\kappa$  is *qualified*, which we now define precisely.

Let X be a locally compact Hausdorff topological space and  $C_0(\mathcal{X})$  be the space of all continuous functions on  $X$  that vanish at infinity.

Definition 3.3 (Qualified Kernel). A separately continuous bounded kernel  $\kappa$  on X is qualified if for any  $f \in C_0(\mathcal{X})$  and  $\epsilon > 0$ , there exists  $g \in \mathcal{H}_\kappa$  such that sup  $|f - g| < \epsilon$ .

Following theorem provides us with several qualified kernels that can be easily adopted in practice.

Kernel Qualification

**Theorem 3.3.** Suppose  $\psi : M \to \mathbb{R}^{d'}$  is a smooth embedding for some  $d' > d$ , and suppose κ is C<sub>0</sub>-universal on  $\mathbb{R}^{d'}$ , then the restriction of κ onto M, i.e.,  $\kappa|_M := \kappa(\psi(x), \psi(x'))$  for  $x, x' \in M$  is a qualified kernel on M.

We expect  $\mathcal{T}_p$  in [\(5\)](#page-5-1) to distinguish p from such q if  $\kappa$  is qualified. However, if we recall the process of derivation of  $\mathcal{A}_p$  on  $\mathbb{R}^d$  in the original literature [\[2,](#page-25-4) [11,](#page-26-10) [25,](#page-26-9) [37,](#page-27-4) [41,](#page-27-6) [48\]](#page-28-9), we discover that, intuitively, the component  $\mathcal{A}_p^l f(x) := \frac{\partial f}{\partial x^l}(x) + f(x) \frac{\partial}{\partial x^l} \log p(x)$  of each l in  $\mathcal{A}_p f$ , detects the difference between the slopes of log p and log q along the direction  $\frac{\partial}{\partial x^l}$  at x. Therefore,  $\{\frac{\partial}{\partial x^l}\}_{l=1}^d$  must be a basis at each point of  $\mathbb{R}^d$ , so that  $\mathcal{A}_p = \sum_{l=1}^d \mathcal{A}_p^l$  can detect the differences along all directions, so that we can conclude  $\frac{\partial}{\partial x^l} \log p(x) = \frac{\partial}{\partial x^l} \log q(x)$  for all  $1 \leq l \leq d$  after some reasoning. In addition, to further conclude  $p = q$ , we need the connectedness of  $\mathbb{R}^d$ . This motivates us to make the following additional assumptions:

Standing Assumption 4 and 5

- 4. M is connected.
- **5.** For each  $x \in M$ ,  $D_x^1, \ldots, D_x^m$  span the entire  $T_xM$ .

In this paper, we maintain the standing assumptions  $1,2,3,4,5$ .

<span id="page-8-2"></span>Theorem 3.4. There always exists a collection of vector fields that satisfies the standing assumption 5 on M.

Then we have the following characterization of KSD:

#### KSD Characterization

<span id="page-8-0"></span>**Theorem 3.5.** Suppose  $\kappa$  is a qualified kernel and q is a locally Lipschitz continuous density on M. If  $\sum_{l=1}^m |D^l|$ ,  $\sum_{l=1}^m |D^l \log(p/q)|$ ,  $\sqrt{\kappa_p(x,x)}$  are q-integrable, then  $p = q \iff$  $KSD(p, q) = 0.$ 

### <span id="page-8-1"></span>3.4 Stein operator on Riemannian Homogeneous Spaces

The definition of the Stein operator in [\(5\)](#page-5-1) does not only require one to find a group of basis that satisfy standing assumption 4, but also requires one to compute the divergence of these vector fields, which is usually challenging for a general vector field, e.g., the one obtained in the proof of Thm. [3.4.](#page-8-2) However, in practice, most of the commonly encountered manifolds are Riemannian Homogeneous spaces. In such spaces, we can select the vector fields  $D<sup>l</sup>$  as a special kind of vector fields, killing fields, to get around such computational issues.

To introduce the Riemannian Homogeneous space, we introduce the notion of isometry and group action first. An *isometry* of a Riemannian manifold  $M$  is a diffeomorphism from  $M$  onto itself that preserves the distance. The isometry group  $I(M)$  is the group of all isometries of M, which is a Lie group due to the Myers–Steenrod theorem  $[52, Thm. 5.6.19]$ . A group action of a Lie group G on a Riemannian manifold M is an continuous group homomorphism  $\Psi: G \to I(M)$  that maps each element g to some isometry  $\Psi_q$  on M. Conventionally, we omit symbol  $\Psi$ , and denote the group action by  $g.x := \Psi_{g}(x)$ . Note that  $e.x = x$ .

A Riemannian homogeneous space H, abbreviated as homogeneous space in this work, is a Riemannian manifold such that there exists a Lie group  $G$  that acts transitively on  $H$ , i.e., for each  $x, y \in H$ , there exists  $g \in G$  such that  $g.x = y$ .

Let e be the identity of G. For a tangent vector  $E \in T_eG$ , we take a local curve  $\mathfrak{e}(t)$  in the equivalence class of E. Note that  $e.x = x$ , thus  $e(t).x$  is a local curve on H at x, and thus corresponds to a tangent vector in  $T_xH$ . For each  $x \in H$ ,  $e(t)$ . x corresponds to a tangent vector, thus they form a vector field, denoted by  $K$ . Such a vector field  $K$  is said to be a killing field of  $G$  on  $H$ . This correspondence is linear, i.e., if  $E^1$  and  $E^2$  corresponds to  $K^1$  and  $K^2$  respectively, then  $aE^1 + bE^2$ corresponds to  $aK^1 + bK^2$ . Specifically, killing fields are *divergence-free*, i.e., div  $K = 0$ .

**Theorem 3.6.** Suppose Lie group  $G$  acts transitively on homogeneous space  $H$ . For a basis  $E^1,\ldots,E^m$  of  $T_eG$  (m = dim G), they correspond to a group of killing field  $K^1,\ldots,K^m$  in the way previously introduced. Then  $K^1, \ldots, K^m$  is a group of divergence-free vector fields on H that satisfies standing assumption 5.

In this case, the *Stein operator*  $\mathcal{T}_p$  becomes

Stein Operator on Homogeneous Spaces

$$
\mathcal{T}_p f \mapsto f \mapsto \sum_{l=1}^m \left[ K^l f_l + f_l K^l \log p \right], \quad f \in \mathcal{H}_\kappa^m. \tag{10}
$$

#### <span id="page-9-0"></span>3.5 Empirical kernel Stein discrepancy

The integral closed form [\(9\)](#page-7-2) is one of the most significant properties of KSD. However, it may not be computable in practice due to following commonly-encountered situations:

#### Only samples available

In practice, q may be accessible only via samples instead of its density being known. This is commonly encountered in parameter estimation problems where, it is common to use a parameterized density family  $p_{\theta}$  to approximate an unknown distribution q, merely from its samples.

#### Intractable Integral

Sometimes, we do have the form of the density q but the integral in  $(9)$  is intractable. For example, in the rotation tracking problem encountered in robotics  $[62]$ , one must approximate the posterior distribution of  $Q_k$  with a von Mises-Fisher distribution so that the tracking algorithm (Kalman filter) updates consistently lie in the same space. In a Bayesian fusion problem [\[34\]](#page-27-8), a similar situation arises when it is required to guarantee that the result of the fusion stays in the family.

These above two situations also arise in the KL-divergence setting, KL $(p, q) = \mathbb{E}_q[\log p] - \mathbb{E}_q[\log q]$ , and in practice, it is common to use the empirical KL-divergence. For example, suppose we aim to approximate an unknown density q with a density family  $p_{\alpha}$  using the KL-divergence, but only have samples  $x_i$  from q instead of its density. We then minimize the empirical KLdivergence  $n^{-1}\sum_i \log p_\alpha(x_i)$  as an alternative  $(\mathbb{E}_q[\log q]$  is constant w.r.t.  $\alpha$ ), which converges to  $\mathbb{E}_q[\log p]$  almost surely by the law of large number. The minimizer of the empirical KL-divergence  $n^{-1} \sum_{i} \log p_{\alpha}(x_i)$  is exactly the maximum likelihood estimator.

Analogously, the KSD has empirical versions, i.e., the  $U$ - and  $V$ -statistics:

<span id="page-9-1"></span>
$$
U_n(p) := \frac{1}{n(n-1)} \sum_{i \neq j} \kappa_p(x_i, x_j), \quad V_n(p) := \frac{1}{n^2} \sum_{i,j} \kappa_p(x_i, x_j), \tag{11}
$$

which can serve as the alternatives to the integral KSD in  $(9)$  when we only have samples from q. Several prior research works [\[11,](#page-26-10) [37,](#page-27-4) [66\]](#page-29-4) have developed a kernel Stein goodness of fit test based on the U-statistics.

If we do have the density of q but the integral in  $(9)$  is intractable, then we can draw samples from q with various sampling algorithms, e.g., Hamiltonion Monte Carlo or Metropolis-Hastings algorithms and adopt  $U_n$  or  $V_n$  as the measurement of the dissimilarity between p and q. Alternatively, if these sampling methods are hard to implement, we could use the importance sampling scheme to sample from another easy-to-sample density  $w$  and consider the following *weighted empirical KSD*:

<span id="page-10-2"></span>
$$
U_n^w(p) := \frac{1}{n(n-1)} \sum_{i \neq j} \kappa_p^w(x_i, x_j), \quad V_n^w(p) := \frac{1}{n^2} \sum_{i,j} \kappa_p^w(x_i, x_j), \tag{12}
$$

where  $\kappa_p^w$  is the weighted  $\kappa_p$  function given by

$$
\kappa_p^w(x, y) := \kappa_p(x, y) \frac{q(x)}{w(x)} \frac{q(y)}{w(y)}.
$$
\n(13)

If  $w = q$ , then  $\kappa_p^w$  will degenerate to  $\kappa_p$ , thus  $U_n^w$  and  $V_n^w$  will degenerate to  $U_n$  and  $V_n$ .

*Example* 3.3. Consider the Riemannian Gaussian distribution  $q \propto \exp\left(-\frac{\rho^2(x,\bar{x}_0)}{2}\right)$  on  $\mathbb{S}^{N-1}$  for some fixed  $\bar{x}_0 \in \mathbb{S}^{N-1}$ . Let the easy-to-sample distribution  $\omega$  be the uniform distribution on  $\mathbb{S}^{N-1}$ , then the weighted empirical KSDs between  $p$  and  $q$  are:

$$
U_n^w = \frac{1}{n(n-1)} \sum_{i \neq j} \kappa_p(x_i, x_j) q(x_i) q(x_j), \quad V_n^w = \frac{1}{n^2} \sum_{i,j} \kappa_p(x_i, x_j) q(x_i) q(x_j).
$$

Note that the empirical KSDs are proportional to the normalizing constant of q.

Similar to the empirical KL-divergence, the empirical KSD will converges to KSD almost surely, as stated in the next theorem:

## Asymptotic properties for empirical KSD

<span id="page-10-1"></span>**Theorem 3.7.** Suppose M is a Riemannian manifold and  $\kappa_p^w(x, x)$  is w-integrable. Given M-valued samples  $x_i \sim w$ , we have

1. 
$$
U_n^w(p), V_n^w(p) \xrightarrow{a.s.} \text{KSD}^2(p,q)
$$
 with the rate of  $O_p(n^{-1})$ ,

2. If  $p \neq q$ , then

$$
\sqrt{n}[U_n^w - \text{KSD}^2(p, q)] \xrightarrow{d} N(0, \tilde{\sigma}^2), \quad \sqrt{n}[V_n^w - \text{KSD}^2(p, q)] \xrightarrow{d} N(0, \tilde{\sigma}^2),
$$

where  $\tilde{\sigma}^2 := 4var_{x' \sim w}[\mathbb{E}_{x \sim w} \kappa_p^w(x, x')]$ .

3. If  $p = q$ , then  $nU_n^w$  $\stackrel{d.}{\rightarrow} \sum^{\infty}$  $k=1$  $\lambda_k (Z_k^2 - 1), \quad nV_n^w$  $\stackrel{d.}{\rightarrow} \sum^{\infty}$  $k=1$  $\lambda_k Z_k^2,$ where  $\lambda_k$  are the eigenvalues of  $\kappa_p^w(x, y)$  as introduced in Thm. [2.4](#page-4-3) and  $Z_k$  are i.i.d. standard Gaussian samples.

# <span id="page-10-0"></span>4 Minimum kernel Stein discrepancy estimator

Since KSD measures the dissimilarity between distributions, one may natually use it on distributional approximation. Suppose we want to approximate a density q with a density  $p_{\alpha}$ , then we define the global minimizer

$$
\hat{\alpha} := \operatorname{argmin} \operatorname{KSD}(p_{\alpha}, q) \tag{14}
$$

as the minimum kernel Stein discrepancy estimator (MKSDE). However, as we explained in §[3.5,](#page-9-0)  $KSD(p_{\alpha}, q)$  may not be computable in most of the situations in practice. Therefore, we minimize a suite of different empirical KSDs over  $\alpha$  based on various situations:

$$
U_n^w(\alpha) := \frac{1}{n(n-1)} \sum_{i \neq j} \kappa_\alpha^w(x_i, x_j), \quad V_n^w(\alpha) := \frac{1}{n^2} \sum_{i,j} \kappa_\alpha^w(x_i, x_j),
$$

where  $\kappa_\alpha^w(x, x') := \kappa_{p_\alpha}^w(x, x')$ . Here  $U_n^w$  and  $V_n^w$  accommodate the unweighted case  $w = q$ .

Although U-statistics is far more frequently mentioned in prior works  $[2, 11, 37, 66]$  $[2, 11, 37, 66]$  $[2, 11, 37, 66]$  $[2, 11, 37, 66]$  $[2, 11, 37, 66]$  $[2, 11, 37, 66]$  as it is an unbiased estimator of  $KSD<sup>2</sup>(p<sub>\alpha</sub>, q)$ , we notice that the V-statistics exhibits better stability for optimization, as explained in the next section studying the asymptotic properties of MKSDE.

#### <span id="page-11-2"></span>4.1 Asymptotic Properties of MKSDE

To obtain the asymptotic behavior of the MKSDE, we re-tag the index  $\alpha$  of the density family by θ, and assume that θ is from some topological space  $\Theta$ . We denote  $\text{KSD}(\theta) := \text{KSD}(p_{\theta}, q)$  and let  $\kappa_{\theta}^{w}(x, x') := \kappa_{p_{\theta}}^{w}(x, x')$ . Let  $\Theta_0 \subset \Theta$  be the set of best approximators  $\theta_0$ , i.e.,  $\text{KSD}(\theta_0) = \inf_{\theta} \text{KSD}(\theta)$ . With the additional topological structure on  $\Theta$ , we can establish stronger asymptotic results of  $U_n^w$ and  $V_n^w$ . The asymptotic results in this section are satisfied by both  $U_n^w$  and  $V_n^w$ . For notational convenience, we let  $W_n(\theta)$  denote whichever  $U_n^w(\theta)$  or  $V_n^w(\theta)$ .

Compact Convergence for KSD

<span id="page-11-0"></span>**Theorem 4.1.** If  $\kappa_{(\cdot)}^w(\cdot, \cdot)$  is jointly continuous and  $\sup_{\theta \in K} \kappa_{\theta}^w(x, x)$  is w-integrable for any compact  $K \subset \Theta$ , then  $W_n(\theta) \to \text{KSD}^2(\theta)$  compactly almost surely, i.e., the following event

 $W_n(\theta) \to \text{KSD}^2(\theta)$  uniformly on any compact  $K \subset \Theta$ 

almost surely happens. As a corollary,  $KSD(\cdot)$  is continuous if  $\Theta$  is locally compact.

Let  $\widehat{\Theta}_n$  be the set of MKSDE, i.e., global minimizers  $\widehat{\theta}_n$  of  $W_n$ , which is a random set. We can establish the strong consistency of MKSDE with Thm. [4.1](#page-11-0) in hand.

Strong Consistency of MKSDE

<span id="page-11-1"></span>**Theorem 4.2.** Suppose all the conditions in Thm. [4.1](#page-11-0) hold and suppose  $\Theta$  satisfies one of following three conditions:

- 1. Θ is compact;
- 2. Θ is a geodesic convex subset of a Riemannian manifold,  $W_n$  is convex on  $\Theta_1$ ,  $\Theta_0$  is non-empty, compact and  $\Theta_0 \subset \check{\Theta}_2$  (interior);
- 3.  $\Theta = \Theta_1 \times \Theta_2$ , where  $\Theta_1$  is compact, and for each fixed  $\theta_1 \in \Theta_1$ ,  $\{\theta_1\} \times \Theta_2$  and  $W_n(\theta_1, \cdot)$ satisfy the second condition.

Then  $\Theta_0$ ,  $\Theta_n$  are non-empty for large n and  $\sup_{\theta \in \widehat{\Theta}_n} \rho(\theta, \Theta_0) \to 0$  almost surely.

It is worth noting that if q is a member of the family  $p_{\theta}$ , then the global minimizer set  $\Theta_0$  is a singleton  $\{\theta_0\}$ . In such situation, the MKSDE always converges to the unique ground truth  $\theta_0$ . Moreover, if the parameter space  $\Theta := \Theta_1 \times \cdots \times \Theta_m$  is multi-dimensional, we combine all compact components into one compact parameter space, and hence the convex components as well, so that Thm. [4.1](#page-11-0) is still applicable.

Existing result in literature [\[2,](#page-25-4) Thm. 3.3] showed the consistency of MKSDE, assuming that the parameter space Θ is either compact or satisfies the conditions of convexity, which is not applicable to the case where there are compact and convex parameters simultaneously. For example, consider the Riemannian Gaussian distribution  $p \propto \exp(-\frac{\rho^2(x,\bar{x})}{2\sigma^2})$  on a compact manifold, if we redenote  $\varsigma := \sigma^{-2}$ , then  $\mu$  is a "compact" parameter and  $\varsigma$  is a "convex" parameter.

To establish the asymptotic normality of MKSDE, we assume that  $\Theta$  is a connected Riemannian manifold with the Riemannian logarithm map Log. We assume that  $\Theta_0$  and  $\Theta_n$  are non-empty for n large enough, and  $\hat{\theta}_n$  is a sequence of MKSDE that converges to one of the global minimizer  $\theta_0$  of  $KSD(\theta)$ . Additionally, we assume the following conditions:

(A1)  $\kappa_{\theta}^{w}(x, y)$  is jointly continuous, and twice continuously differentiable in  $\theta$ .

- (A2) there exists a compact neighborhood K of  $\theta_0$  such that  $\sup_{\theta \in K} \|\nabla \kappa_{\theta_0}^w(x, y)\|$  is  $w \times w$ -integrable.
- (A3)  $\|\nabla \kappa_{\theta_0}^w(x, y)\|^2$  and  $\|\mathcal{I}_{\theta_0}(x, y)\|$  are  $w \times w$ -integrable,  $\|\mathcal{I}_{\theta_0}(x, x)\|$  is w-integrable.
- (A4)  $\mathcal{I}_{\theta_0}(x, y)$  is equi-continuous at  $\theta_0$ .
- (A5)  $\Gamma := \frac{1}{2} \mathbb{E}_{x,y \sim w} [\mathcal{I}_{\theta_0}(x,y)]$  is invertible.

Here  $\nabla \kappa_{\theta}^{w}(x, y)$  represents the gradient of  $\kappa_{\theta}^{w}(x, y)$  w.r.t  $\theta$ , and  $\mathcal{I}_{\theta}(x, y)$  represents the Hessian of  $\kappa_{\theta}^{w}(x, y)$  w.r.t  $\theta$ . In addition, let Σ be the covariance matrix of the random vector  $\mathbb{E}_{Y \sim w}[\kappa_{\theta_0}^{w}(x, Y)].$ 

#### CLT for MKSDE

Theorem 4.3. Under assumption  $A1 \sim A5$ ,  $\sqrt{n} \text{Log}_{\theta_0}(\hat{\theta}_n) \stackrel{d}{\rightarrow} \mathcal{N}(0,\Gamma^{-1}\Sigma\Gamma^{-1})$ . In more practical situations, we have  $\sqrt{n}[\mathcal{I}(\hat{\theta}_n)]^{1/2} \text{Log}_{\hat{\theta}_n}(\theta_0) \stackrel{d}{\rightarrow} \mathcal{N}(0, I).$ 

### 4.2 MKSDE Composite goodness of Fit Test

KSD is commonly used to develop normalization-free goodness of fit tests [\[11,](#page-26-10) [37\]](#page-27-4), which test whether a group of samples can be well-modeled by a given distribution  $p$ . More precisely, if we denote by q the unknown underlying distribution of samples, then we aim to test  $H_0: p = q$  versus  $H_1: p \neq q$ . However, in many applications the candidate distribution for the given samples is usually not a specific distribution but a parameterized family  $p_{\theta}$ , while most existing methods only apply to a specific candidate, and requires testing individually for each member of  $p<sub>\theta</sub>$ . Recently, a *composite* goodness-of-fit test was developed to test whether a group of samples matches a family  $p_\theta$  [\[28\]](#page-27-9), i.e., test  $H_0$ :  $\exists \theta_0, p_{\theta_0} = q$  versus  $H_1 : \forall \theta, p_{\theta} \neq q$ , and was later generalized to all Lie groups in [\[53\]](#page-28-10). This however was valid for samples of distributions over Lie groups and not general Riemannian manifolds. In this section, we will generalize the composite goodness of fit test to all Riemannian manifold, develop a one-shot (direct) method, using the MKSDE obtained by minimizing [\(12\)](#page-10-2).

To implement the test, we assume the conditions in Thm. [4.2](#page-11-1) hold, so that  $\Theta_0$  and  $\widehat{\Theta}_n$  are nonempty. We also assume that  $\Theta_0$  is a singleton so that the MKSDE  $\hat{\theta}_n = \text{argmin wKSD}_n^2(\theta)$  converges to the unique ground truth  $\theta_0$ . We follows the notation in [4.1](#page-11-2) letting  $W_n$  denote whichever  $U_n^w$  or  $V_n^w$ .

Under the null hypothesis  $H_0$ ,  $nW_n(\theta_0)$  converges to  $\sum_k \lambda_k (Z_k^2 - 1)$  or  $\sum_k \lambda_k Z_k^2$  in distribution asymptotically by Thm. [3.7.](#page-10-1) Let  $\gamma_{1-\beta}$  be the  $(1-\beta)$ -quantile of  $\sum_k \lambda_k (Z_k^2 - 1)$  or  $\sum_{k=1}^{\infty} \lambda_k Z_k^2$  with significance level  $\beta$ . We reject  $H_0$  if  $nW_n(\hat{\theta}_n) \geq \gamma_{1-\beta}$ , as it implies  $nW_n(\theta_0) \geq nW_n(\hat{\theta}_n) \geq \gamma_{1-\beta}$ , since  $\hat{\theta}_n$  is the minimizer of  $W_n$ .

As there is no method in general to directly compute the infinite eigenvalues  $\lambda_k$ -s of  $\kappa_{\theta_0}^w$ , Gretton et al. [\[27\]](#page-27-10) introduced a method to approximate  $\lambda_k$ -s by the eigenvalues of the empirical matrix  $G_n^w(\theta_0) := n^{-1}(\kappa_{\theta_0}^w(x_i, x_j))_{ij}$ . Note that the ground truth  $\theta_0$  is unknown in our setting, but the minimizers  $\hat{\theta}_n$  converge to  $\theta_0$  almost surely under specific conditions by Thm. [4.2.](#page-11-1) Therefore, we may approximate  $\lambda_k$  by the eigenvalues of the empirical matrix  $G_n^w(\hat{\theta}_n) := n^{-1}(\kappa_{\hat{\theta}_n}^w(x_i, x_j))_{ij}$ . Let  $\hat{\lambda}_k$  be the eigenvalues of the matrix  $G_n^w(\hat{\theta}_n)$  (set  $\hat{\lambda}_k := 0$  for  $k > n$ ), then we have

<span id="page-13-0"></span>**Theorem 4.4.** Under the conditions in Thm.  $\{1.2, \sum_{k=1}^{\infty} (\hat{\lambda}_k - \lambda_k) Z_k^2 \} 0$  in probability.

Therefore, the  $\sum_{k=1}^{n} \hat{\lambda}_k (Z_k^2 - 1)$  and  $\sum_{k=1}^{n} \hat{\lambda}_k Z_k^2$  can serve as an empirical estimate of the asymptotic distribution  $\sum_{k=1}^{\infty} \lambda_k (Z_k^2 - 1)$  and  $\sum_{k=1}^{\infty} \lambda_k Z_k^2$ . The MKSDE goodness of fit test algorithm is summarized in the following algorithm block [1.](#page-13-2)

Algorithm 1: MKSDE goodness of fit test

<span id="page-13-2"></span>**Input**: population  $x_1, \ldots, x_n \sim w$ ; sample size *n*; number of generations *n'*; significance level  $\beta$ . **Test:**  $H_0$ :  $\exists \theta_0, p_{\theta_0} = q$  versus  $H_1$ :  $\forall \theta, p_{\theta} \neq q$ . Procedure:

- 1. Obtain minimizer  $\hat{\theta}_n$  of  $W_n(\theta)$  in [\(12\)](#page-10-2).
- 2. Obtain the eigenvalues  $\hat{\lambda}_1, \ldots, \hat{\lambda}_n$  of  $G_n^w(\hat{\theta}_n) := n^{-1}(\kappa_{\hat{\theta}_n}^w(x_i, x_j))_{ij}$ .
- 3. Sample  $Z_k^l \sim N(0, 1), 1 \leq k \leq n, 1 \leq l \leq n'$  independently.
- 4. Compute  $\gamma^l = \sum_{k=1}^n \hat{\lambda}_k [(Z_k^l)^2 1]$  or  $\sum_{k=1}^n \hat{\lambda}_k (Z_k^l)^2$ .
- 5. Determine estimation  $\hat{\gamma}_{1-\beta}$  of  $(1-\beta)$ -quantile using  $\gamma^1, \ldots, \gamma^{n'}$ .

**Output:** Reject  $H_0$  if  $nW_n(\hat{\theta}_n) > \hat{\gamma}_{1-\beta}$ .

# <span id="page-13-1"></span>5 Closed Form KSD on Homogeneous Spaces

As we can see from our earlier discussion, the calculation of  $\kappa_p$  function from [\(7\)](#page-6-1) plays an important role in the practical usage of KSD and MKSDE. Suppose  $H$  is a homogeneous space, and  $\{K^l\}_{l=1}^m$  is a killing field basis,  $\kappa$  is kernel function and p is a density on H. After some straightforward calculations, we obtain

<span id="page-13-3"></span>
$$
\kappa_p(x,y) = \sum_{l=1}^m \mathcal{T}_{p(y)}^l \mathcal{T}_{p(x)}^l \kappa = \kappa \cdot \sum_{l=1}^m \left[ K_x^l \log(p\kappa) \cdot K_y^l \log(p\kappa) + K_y^l K_x^l \log \kappa \right]. \tag{15}
$$

In this section, we will present a surprising result namely that, the formula [\(15\)](#page-13-3) of  $\kappa_p$  function can be further simplified into closed form expressions in most commonly-encountered homogeneous spaces, including:

• the Stiefel manifolds  $\mathcal{V}_r(N) := \{ X \in \mathbb{R}^{N \times r} : X^\top X = I_{r \times r} \}$ , inheriting the subspace topology from  $\mathbb{R}^{N\times r}$ , including the sphere  $\mathcal{V}_1(N) := \mathbb{S}^{N-1}$ , the special orthogonal group  $\mathcal{V}_{N-1}(N) =$ SO(N) and the orthogonal group  $\mathcal{V}_N(N) = O(N)$ .

- the Grassman manifold  $\mathcal{G}_r(N) := \{ X \in \mathbb{R}^{N \times N} : X^{\top} = X, X^2 = X, \text{tr}(X) = r \}$ , the space of all *r*-planes in  $\mathbb{R}^N$ , or equivalently, the space of orthogonal projections of  $\mathbb{R}^N$  into *r*-dimensional subspaces, or equivalently, the space of  $N$  by  $N$  idempotent symmetric matrices with rank  $r$ .
- $\mathcal{P}(N) : \{X \in \mathbb{R}^{N \times N} : X^{\top} = X, X \succ 0\}$ , the manifold of  $(N, N)$  symmetric positive definite (SPD) matrices, inheriting the subspace topology from  $\mathbb{R}^{N \times N}$ .

Even more surprisingly, the MKSDE obtained by minimizing the empirical KSDs [\(11\)](#page-9-1) and [\(12\)](#page-10-2), also has closed form expressions, if  $p_{\theta}$  is the exponential family given by

$$
p(x|\theta) \propto \exp(\theta^\top \zeta(x) + \eta(x)), \quad \theta \in \mathbb{R}^s
$$

where  $\zeta := (\zeta_1, \ldots, \zeta_s)^\top \in C^1(H, \mathbb{R}^s), \, \eta \in C^1(H, \mathbb{R})$  for some  $s \in \mathbb{N}_+$ . We plug this into [\(15\)](#page-13-3) and get

$$
\kappa_p(x, y) = \theta^{\top} \cdot \kappa \sum_{l=1}^m K_x^l \zeta \cdot K_y^l \zeta^{\top} \cdot \theta + \kappa \sum_{l=1}^m K_x^l \log(e^{\eta} \kappa) K_y^l \zeta^{\top} \theta + \kappa \sum_{l=1}^m K_y^l \log(e^{\eta} \kappa) K_x^l \zeta^{\top} \theta + c(x, y),
$$
\n(16)

where  $c(x, y)$  denote the terms independent of  $p_\theta$  thus independent of its parameter  $\theta$ . Note that  $K_x^l \zeta$  is a s-dimensional vector, thus  $K_x^l \zeta \cdot K_y^l \zeta^\top$  is the matrix  $(K_x^l \zeta_i \cdot K_y^l \zeta_j)_{ij}$ . Furthermore, we let

<span id="page-14-2"></span>
$$
Q(x,y) = \kappa \sum_{l=1}^{m} K_x^l \zeta \cdot K_y^l \zeta^\top, \quad b(x,y) = \kappa \sum_{l=1}^{m} K_x^l \log(e^{\eta} \kappa) \cdot K_y^l \zeta. \tag{17}
$$

Therefore, the empirical KSDs in [\(11\)](#page-9-1) and [\(12\)](#page-10-2) will be quadratic forms of  $\theta$ :

<span id="page-14-1"></span>
$$
U_n^w(\theta) = \theta^\top Q_u \theta + 2b_u \theta + c, \quad V_n^w(\theta) = \theta^\top Q_v \theta + 2b_v \theta + c,
$$
\n(18)

where

<span id="page-14-0"></span>
$$
Q_u := \sum_{i \neq j} \frac{Q(x_i, x_j)}{n(n-1)} \frac{q(x_i)q(x_j)}{w(x_i)w(x_j)}, \quad b_u := \sum_{i \neq j} \frac{b(x_i, x_j)}{n(n-1)} \frac{q(x_i)q(x_j)}{w(x_i)w(x_j)},
$$
  
\n
$$
Q_v := \sum_{i,j} \frac{Q(x_i, x_j)}{n^2} \frac{q(x_i)q(x_j)}{w(x_i)w(x_j)}, \quad b_v := \sum_{i,j} \frac{b(x_i, x_j)}{n^2} \frac{q(x_i)q(x_j)}{w(x_i)w(x_j)}.
$$
\n(19)

It is noteworthy that  $Q_u, Q_v$  are both symmetric. Furthermore,  $Q_v$  is always semi-positive definite, thus  $V_n(\theta)$  can always attain its minimum values for exponential family, while  $U_n$  can not. Additionally,  $Q_u$  and  $Q_v$  are not necessarily invertible, e.g.  $p_\theta$  is not identifiable. In such cases, the global minimum point can be represented by the Moore–Penrose inverse  $Q_u^+$  and  $Q_v^+$  of  $Q_u$  and  $Q_v$ . We summarize into the following theorem:

<span id="page-14-4"></span>**Theorem 5.1.**  $Q_u$ ,  $Q_v$  defined in [\(19\)](#page-14-0) are both symmetric, and  $Q_v$  is positive semi-definite. If further  $Q_u$  is positive semi-definite, then the global minimizer sets of  $U_n^w(\theta)$  and  $V_n^w(\theta)$  in [\(18\)](#page-14-1) can be represented by the Moore-Penrose inverse  $Q_u^+$ ,  $Q_v^+$  of  $Q_u$ ,  $Q_v$  as follows

<span id="page-14-3"></span>
$$
\operatorname{argmin} U_n^w(\theta) = \{-Q_u^+ b_u - (I - Q_u^+ Q_u)x : x \in \mathbb{R}^s\},
$$
  

$$
\operatorname{argmin} V_n^w(\theta) = \{-Q_v^+ b_v - (I - Q_v^+ Q_v)x : x \in \mathbb{R}^s\}.
$$
  
(20)

For the unweighted case, we just ignore the weighted ratio  $\frac{q(x_i)q(x_j)}{w(x_i)w(x_j)}$  in [\(19\)](#page-14-0).

To obtain the forms of MKSDE, it suffices to derive the closed form of  $Q(x, y)$  and  $b(x, y)$  in [\(17\)](#page-14-2) on different manifolds. In this section, we will explicitly calculate aforementioned closed forms of KSD and MKSDE on commonly-encountered homogeneous spaces, and provide multiple examples for specific families and specific choice of kernels.

#### 5.1 Matrix Algebra

In this section, since the sample space  $M$  is taken to be a matrix manifold, we first introduce some background matrix algebra along with convenient notation that will be used subsequently in this section. We refer the reader to [\[26\]](#page-27-11) and [\[38\]](#page-27-12) for more details on this topic.

#### Matrix Inner Product space

The Frobenius inner product

$$
\langle A, B \rangle_{\mathcal{F}} := \text{tr}(A^{\top}B) = \sum_{i=1}^{N} \sum_{j=1}^{r} a_{ij} b_{ij}, \quad \text{for } A = (a_{ij}) \in \mathbb{R}^{N \times r} \text{ and } B = (b_{ij}) \in \mathbb{R}^{N \times r},
$$

is usually considered to be the canonical inner product on  $\mathbb{R}^{N \times r}$ . The Frobenius norm is given by  $||A||_F := \sqrt{\langle A, A \rangle_F}$ . We let vec $(A)$  denote the vectorization of A obtained by stacking the columns, and define then following functions

$$
\mathscr{S}(X) = (A + A^{\top})/2, \quad \mathscr{A}(A) = (A - A^{\top})/2,
$$

be the symmetrization and skew-symmetrization of a squared matrix  $A \in \mathbb{R}^{N \times N}$ . Let  $\mathscr{S}(\mathbb{R}^{N \times N})$  and  $\mathscr{A}(\mathbb{R}^{N\times N})$  be the linear subspaces of all symmetric and skew-symmetric  $N\times N$  matrices respectively. It can be easily checked that they are the orthogonal complements of each other, and  $X = \mathscr{S}(X)$  +  $\mathscr{A}(X)$  is the orthogonal decomposition of X onto  $\mathscr{S}(\mathbb{R}^{N\times N})$  and  $\mathscr{A}(\mathbb{R}^{N\times N})$ .

Let  $E_{ij}$  be the matrix prescribed with all zeros everywhere except a 1 at the  $(i, j)$ <sup>th</sup> entry. Specifically, when  $r = 1$ , we set  $e_i$  be the  $N \times 1$  vector prescribed with all zeros except 1 at the i<sup>th</sup> element. Let  $\mathcal{E}_{ij} := \left(\frac{\sqrt{2}}{2}E_{ij} - \frac{\sqrt{2}}{2}E_{ji}\right)$ . Then following proposition is easy to verify: √

<span id="page-15-0"></span>**Proposition 5.1.**  $\{E_{ij}, 1 \le i, j \le N\}$  and  $\{\mathcal{E}_{ij}, 1 \le i < j \le N\}$  are orthonormal bases of  $\mathbb{R}^{N \times N}$ and  $\mathscr{A}(\mathbb{R}^{N\times N})$  respectively, and for any  $A, B \in \mathbb{R}^{N\times N}$  we have

$$
\sum_{i,j} \langle E_{ij}, A \rangle_{\mathcal{F}} \cdot \langle E_{ij}, B \rangle_{\mathcal{F}} = \langle A, B \rangle_{\mathcal{F}}, \quad \sum_{i < j} \langle \mathcal{E}_{ij}, A \rangle_{\mathcal{F}} \cdot \langle \mathcal{E}_{ij}, B \rangle_{\mathcal{F}} = \langle \mathscr{A}(A)^{\top}, \mathscr{A}(B) \rangle_{\mathcal{F}}.
$$

#### 5.1.1 Gradient of functions on a matrix manifold

Suppose M is a sub-manifold of  $\mathbb{R}^{N \times r}$ , i.e., a matrix manifold whose elements are  $N \times r$  matrices, and M is endowed with the Riemannian metric  $g_X(\cdot, \cdot)$  for  $X \in M$ .

For a real-valued function f on  $M$ , the Riemannian gradient of  $f$  at  $X$  is defined as the tangent vector  $\nabla_X^M f \in T_X M$ , such that  $D_X f = g_X(D_X, \nabla_X^M f)$  for all tangent vectors  $D_X \in T_X M$ . Analogously, since tangent space  $T_XM$  at each point  $X \in M$  is a linear subspace of  $\mathbb{R}^{N \times r}$ , we may define the Euclidean gradient of f w.r.t the Frobenius inner product  $\langle \cdot, \cdot \rangle_F$ , i.e., a matrix  $\nabla_X^{\mathbb{R}} f \in \mathbb{R}^{N \times r}$ such that  $D_X f = \langle \nabla^{\mathbb{R}}_X f, D_X \rangle_F$  for all  $D_X \in T_X M \subset \mathbb{R}^{\tilde{N} \times r}$ .

Note that  $\nabla^{\mathbb{R}}_X f$  is not necessarily an element in  $T_XM$ , thus the Euclidean gradient is not unique in general. For example, consider function  $f(x) = \mu^{\top} x$  for x on the sphere  $\mathbb{S}^{N-1}$ . Since the Riemannian

gradient must lie in  $T_x \mathbb{S}^{N-1}$ , we have  $\nabla_x^M f = \mu - (\mu^T x)x$ , the orthogonal projection of  $\mu$  onto the  $T_x\mathbb{S}^{N-1}$ . On the other hand, as  $x \perp T_x\mathbb{S}^{N-1}$ , the Euclidean gradient  $\nabla_x^{\mathbb{R}}f$  can be any vector with the form  $\mu - \alpha x, \alpha \in \mathbb{R}$ .

In this work, we address this notion only for computational and representational convenience, but it is not widely used in other works due to non-uniqueness. As different choices of  $\nabla^{\mathbb{R}}_{X} f$  will deliver the same value of  $\langle \nabla_X^{\mathbb{R}} f, D_X \rangle_F$  for any  $D_X \in T_XM$ , and they only appear inside the inner product bracket  $\langle \cdot, \cdot \rangle_F$  in this work, the non-uniqueness will not influence any specific calculation that follows in this section.

# 5.2 Stiefel Manifold  $\mathcal{V}_r(N)$  (Special Cases:  $\mathbb{S}^{N-1}$ ,  $\text{SO}(N)$  and  $\text{O}(N)$ )

We refer the readers to  $\left[1, 15\right]$  for a detailed discussion on the definition and geometry of the Stiefel manifold. It is known that  $O(N)$  acts transitively on  $\mathcal{V}_r(N)$  and the isometry corresponding to each  $O \in O(N)$  is  $X \mapsto O.X := OX$  for  $X \in V_r(N)$ . The tangent space of  $O(N)$  at identity is  $\mathfrak{o}(N) = \mathscr{A}(\mathbb{R}^{N \times N})$ , the space of all skew-symmetric  $N \times N$ -matrices, where  $\{\mathcal{E}_{ij} : 1 \leq i < j \leq N\}$ is an orthogonal basis as introduced in §[3.4.](#page-8-1) We take local curves  $O_{ij}(t)$  at identity corresponding to each  $\mathcal{E}_{ij}$ , i.e.,  $\frac{d}{dt}O_{ij}(t)|_{t=0} = \mathcal{E}_{ij}$ , then the killing field corresponding to each  $\mathcal{E}_{ij}$  is  $K_X^{ij} := \frac{d}{dt}O_{ij}(t)|_{t=0}X = \frac{d}{dt}O_{ij}(t)|_{t=0}X = \mathcal{E}_{ij}X$ , i.e., the vector field  $K^{ij}$  that assigns eac vector  $\mathcal{E}_{ij}X \in T_X\mathcal{V}_r(N)$ . We plug this into [\(15\)](#page-13-3) so that the  $k_p$  function on Stiefel manifolds equals

$$
\kappa_p(X,Y) = \kappa \cdot \sum_{i < j} K_X^{ij} \log(p\kappa) \cdot K_Y^{ij} \log(p\kappa) + \kappa \cdot \sum_{i < j} K_Y^{ij} K_X^{ij} \log \kappa.
$$

The first term can be written in a closed form:

$$
\kappa \cdot \sum_{i < j} K_X^{ij} \log(p\kappa) \cdot K_Y^{ij} \log(p\kappa) = \kappa \cdot \sum_{i < j} \text{tr}[X^\top \mathcal{E}_{ij}^\top \nabla_X^{\mathbb{R}} \log(p\kappa)] \cdot \text{tr}[Y^\top \mathcal{E}_{ij}^\top \nabla_Y^{\mathbb{R}} \log(p\kappa)]
$$
\n
$$
= \kappa \cdot \sum_{i < j} \langle \mathcal{E}_{ij}, \nabla_X^{\mathbb{R}} \log(p\kappa) X^\top \rangle_F \cdot \langle \mathcal{E}_{ij}, \nabla_Y^{\mathbb{R}} \log(p\kappa) Y^\top \rangle_F
$$
\nBased on Prop. 5.1

\n
$$
= \kappa \cdot \langle \mathscr{A}(\nabla_X^{\mathbb{R}} \log(p\kappa) X^\top), \mathscr{A}(\nabla_Y^{\mathbb{R}} \log(p\kappa) Y^\top) \rangle_F.
$$

aThe summation of  $K_Y^{ij} K_X^{ij} \log \kappa$  relies on the specific form of  $\kappa$  and thus has no closed form in general, which can not be further simplified. However, if  $\kappa$  is a radial kernel, i.e.,

$$
\kappa(X, Y) = \exp(-\psi(||X - Y||_{\mathcal{F}}^2)) = \exp(-\psi(2r - 2\operatorname{tr}[X^{\top}Y])), \quad X, Y \in \mathcal{V}_r(N),
$$

for some  $\psi \in C^2[0, +\infty)$ , then  $K_Y^{ij} K_X^{ij} \log \kappa$  equals

$$
K_Y^{ij} K_X^{ij} \log \kappa = -\frac{d^2}{dt dt'} \psi(2r - 2 \operatorname{tr}[X^\top O_{ij}^\top(t) O_{ij}(t')Y])|_{t,t'=0}
$$
  
= 
$$
\underbrace{2\psi'(\|X - Y\|_{\text{F}}^2) \cdot \operatorname{tr}[X^\top \mathcal{E}_{ij}^\top \mathcal{E}_{ij} Y]}_{(i)}
$$
  

$$
-4\psi''(\|X - Y\|_{\text{F}}^2) \cdot \operatorname{tr}[X^\top \mathcal{E}_{ij}^\top Y] \cdot \operatorname{tr}[X^\top \mathcal{E}_{ij} Y].
$$

Note that  $\mathcal{E}_{ij}^{\top} \mathcal{E}_{ij} = \frac{1}{2} E_{ii} + \frac{1}{2} E_{jj}$ , thus  $\sum_{i < j} \mathcal{E}_{ij}^{\top} \mathcal{E}_{ij} = \frac{N-1}{2} I$ . Therefore, the summation of (*i*) over  $i < j$  equals  $(N-1)\psi'(\|X-Y\|_{\text{F}}^2)\langle X,Y\rangle_{\text{F}}$ . The summation of  $(ii)$  follows the mechanic based on the Prop. [5.1,](#page-15-0) which equals  $4\psi''(\|X-Y\|_{\mathrm{F}}^2)\|\mathscr{A}(XY^{\top})\|_{\mathrm{F}}^2$ . We summarize into following theorem:

**Theorem 5.2.** Given density p and kernel κ, the  $\kappa_p$  function on  $\mathcal{V}_r(N)$  is given by

$$
\kappa_p(X, Y) = \langle \mathscr{A}(\nabla_X^{\mathbb{R}} \log(p\kappa) X^{\top}), \mathscr{A}(\nabla_Y^{\mathbb{R}} \log(p\kappa) Y^{\top}) \rangle_F \cdot \kappa(X, Y) + \cdot \sum_{i < j} K_Y^{ij} K_X^{ij} \log \kappa \cdot \kappa(X, Y).
$$
\n(21)

Furthermore, if  $\kappa(X,Y) = e^{-\psi(||X-Y||_F^2)}$  is a radial kernel for some  $\psi \in C^2[0,+\infty)$ , then

<span id="page-17-0"></span>
$$
\kappa_p(X, Y) = \kappa \cdot \langle \mathscr{A}(\nabla_X^{\mathbb{R}} \log(p\kappa) X^{\top}), \mathscr{A}(\nabla_Y^{\mathbb{R}} \log(p\kappa) Y^{\top}) \rangle_F \cdot \kappa(X, Y) + (N-1)\psi'(\|X - Y\|_F^2) \langle X, Y \rangle_F \cdot \kappa(X, Y) + 4\psi''(\|X - Y\|_F^2) \|\mathscr{A}(XY^{\top})\|_F^2 \cdot \kappa(X, Y).
$$
\n(22)

Next we explicitly calculate the closed form of MKSDE for the exponential family on  $\mathcal{V}_r(N)$ . It suffices to calculate the matrix  $Q(X,Y) = k \sum_{i < j} (K_X^{ij} \zeta_k \cdot K_Y^{ij} \zeta_l)_{kl}$  and the vector  $b(X,Y) =$  $\kappa \cdot \sum_{i < j} [K_X^{ij} \eta + K_X^{ij}] \log \kappa] K_Y^{ij} \zeta$  introduced in [\(17\)](#page-14-2). For  $Q(X, Y)$ , we have

$$
\sum_{i < j} K_X^{ij} \zeta_k \cdot K_Y^{ij} \zeta_l = \sum_{i < j} \text{tr}[(\mathcal{E}_{ij} X)^\top \nabla_X^{\mathbb{R}} \zeta_k] \cdot \text{tr}[(\mathcal{E}_{ij} Y)^\top \nabla_Y^{\mathbb{R}} \zeta_l]
$$
\n
$$
= \sum_{i < j} \langle \mathcal{E}_{ij}, \nabla_X^{\mathbb{R}} \zeta_k X^\top \rangle_F \cdot \langle \mathcal{E}_{ij}, \nabla_Y^{\mathbb{R}} \zeta_l Y^\top \rangle_F
$$
\nProp. 5.1 =  $\langle \mathcal{A}(\nabla_X^{\mathbb{R}} \zeta_k X^\top), \mathcal{A}(\nabla_Y^{\mathbb{R}} \zeta_l Y^\top) \rangle_F$ ,

Similarly, for  $b(X, Y)$ , we have

$$
\sum_{i < j} K_X^{ij} \log(e^{\eta} \kappa) \cdot K_Y^{ij} \zeta_k = \langle \mathscr{A}(\nabla_X^{\mathbb{R}} \log(e^{\eta} \kappa) X^{\top}), \mathscr{A}(\nabla_Y^{\mathbb{R}} \zeta_k Y^{\top}) \rangle_{\mathcal{F}}.
$$

To sum up, we have

**Theorem 5.3** (MKSDE for exponential family). Given the exponential family  $p_\theta \propto \exp(\theta^\top \zeta(X))$  $\eta(X)$ , let  $Q(X,Y)$  and  $b(X,Y)$  be the matrix and vector given by

$$
Q(X,Y)_{kl} = \langle \mathscr{A}(\nabla_X^{\mathbb{R}} \zeta_k X^{\top}), \mathscr{A}(\nabla_Y^{\mathbb{R}} \zeta_l Y^{\top}) \rangle_F \cdot \kappa(X,Y),
$$
  
\n
$$
b(X,Y)_k = \langle \mathscr{A}(\nabla_X^{\mathbb{R}} \log(e^{\eta} \kappa) X^{\top}), \mathscr{A}(\nabla_Y^{\mathbb{R}} \zeta_k Y^{\top}) \rangle_F \cdot \kappa(X,Y),
$$
\n(23)

Then the MKSDE can be computed via  $(20)$ .

Next, we calculate several examples for specific radial kernel  $\kappa$  and density p on Stiefel manifold. To obtain the KSD in [\(22\)](#page-17-0), it suffices to compute  $\nabla_X^{\mathbb{R}} \log \kappa$ ,  $\nabla_X^{\mathbb{R}} \log p$  and  $\psi'$  and  $\psi''$ . To obtain the MKSDE of exponential family  $p(X) \propto \exp(\theta^{\top} \zeta(X) + \eta(X))$  in [\(20\)](#page-14-3), we only need to compute  $\nabla^{\mathbb{R}}_X \zeta_i$ and  $\nabla_X^{\mathbb{R}} \eta$ , to construct the

### Commonly-used distributions

Most of the widely-used distribution families on Stiefel manifold  $\mathcal{V}_r(N)$  have intractable normalizing constant, including:

• Matrix Fisher (MF) family:  $p(X; F) \propto \exp[\text{tr}(F^{\top}X)]$  with parameter  $F \in \mathbb{R}^{N \times r}$ , which belongs to the exponential family since  $\log p = \text{tr}(F^{\top}X)$  is linear in F. We have  $\nabla_X^{\mathbb{R}} \log p = F$ and  $\overline{\nabla}_X^{\mathbb{R}} \eta = 0$ . Note that  $\zeta(X) = X$ , thus the Euclidean gradient of its  $(i, j)$ <sup>th</sup> component  $\zeta_{ij} = \operatorname{tr}[E_{ij}^{\top} X]$  is  $\nabla_X^{\mathbb{R}} \zeta_{ij} = E_{ij} \in \mathbb{R}^{N \times r}$ .

- Matrix Bingham (MB) family:  $p(X; A) \propto \exp[{\rm tr}(X^{\top}AX)]$  with parameter  $A \in \mathbb{R}^{N \times N}$ , which belongs to the exponential family since  $\log p = \text{tr}[(XX^\top)^\top A]$  is linear in A. We have  $\nabla_X^{\mathbb{R}} \log p = (A + \mathcal{S}(A))X$  and  $\nabla_X^{\mathbb{R}} \eta = 0$ . Note that  $\zeta(X) = XX^{\top}$ , thus its  $(i, j)^{\text{th}}$  component  $\zeta_{ij} = \text{tr}[E_{ij}^{\top} (XX^{\top})]$ . Here  $E_{ij} \in \mathbb{R}^{N \times N}$ . Therefore,  $\nabla^{\mathbb{R}}_X \zeta_{ij} = (E_{ij} + E_{ji})X$ . The MB family is not identifiable, since any  $A_1$ ,  $A_2$  such that  $\mathscr{S}(A_1) = \mathscr{S}(A_2)$  will correspond to the same distribution.
- Matrix Fisher-Bingham (MFB) family:  $p(X; A, F) \propto \exp[{\rm tr}(X^{\top}AX + F^{\top}X)]$  with parameter  $(A, F) \in \mathbb{R}^{N \times (N+r)}$ , combining A and F by row, which belongs to the exponential family since  $\log p = \text{tr}(X^\top AX + F^\top X)$  is linear in  $(A, F)$ . We have  $\nabla_X^{\mathbb{R}} \log p = (A + \mathcal{S}(A))X + F$ ,  $\nabla^{\mathbb{R}}_X \eta = 0$ . Note that  $\zeta(X) = (XX^{\top}, X)$ , thus  $\nabla^{\mathbb{R}}_X \zeta_{ij} = (E_{ij}^N + E_{ji}^N)X$  for  $1 \leq j \leq N$ , and  $\nabla^{\mathbb{R}}_X \zeta_{ij} = E^r_{ij}$  for  $N + 1 \leq j \leq N + r$ , where  $E^N_{ij} \in \mathbb{R}^{\bar{N} \times N}$  and  $E^r_{ij} \in \mathbb{R}^{\bar{N} \times r}$ . The MFB family is also non-identifiable.
- Riemannian Gaussian(RG) family:  $p(X) \propto \exp(-\frac{d^2(X,\bar{X})}{2\sigma^2})$  with parameters  $\bar{X} \in \mathcal{V}_r(N)$ ,  $\sigma > 0$ . The RG family is not a member of the exponential family. For Riemannian Gaussian family,  $\log p(X) = -\frac{1}{2\sigma^2}d^2(X,\bar{X})$ . The Riemannian gradient of  $d^2$  function is  $-2\log_X(\bar{X})$  [\[51\]](#page-28-13), the Riemannian gradient of  $\log p(X)$  is  $\sigma^{-2} \text{Log}_X(\bar{X})$ . Here Log is the Riemannian logarithm on  $V_r(N)$ , which can be computed numerically [\[70\]](#page-29-7). It is shown in [\[15,](#page-26-14) §2.4.1] that the Riemannian metric on  $\mathcal{V}_r(N)$  is given by  $\langle D_1, D_2 \rangle_X = \text{tr} \left( D_1^\top (I - \frac{1}{2} XX^\top) D_2 \right)$ , thus  $\nabla_X^{\mathbb{R}} \log p = \sigma^{-2} (I \frac{1}{2}XX^{\top}$ ) Log<sub>X</sub>  $\overline{X}$ .

#### Commonly-used kernels

The most widely-used kernel on  $\mathcal{V}_r(N)$  includes:

• Gaussian kernel:

$$
\kappa(X,Y) = \exp\left(-\frac{\tau}{2}||X - Y||_{\mathcal{F}}^2\right) \propto \exp(\tau \operatorname{tr}(X^{\top}Y)), \quad X, Y \in \mathcal{V}_r(N).
$$

Note that  $\nabla_X^{\mathbb{R}} \log \kappa = \tau Y$ ,  $\psi(x) = \frac{\tau}{2} x$ ,  $\psi'(x) = \frac{\tau}{2}$  and  $\psi''(x) = 0$ .

• Inverse quadratic kernel:

$$
\kappa(X, Y) = (\beta + \|X - Y\|_{\text{F}}^2)^{-\gamma}, \quad X, Y \in \mathcal{V}_r(N).
$$
  
Note that  $\nabla_X^{\mathbb{R}} \log \kappa = \frac{2\gamma}{\beta + \|X - Y\|_{\text{F}}^2} Y, \ \psi(x) = \gamma \log(\beta + x), \ \psi'(x) = \frac{\gamma}{\beta + x} \text{ and } \psi''(x) = -\frac{\gamma}{(\beta + x)^2}.$ 

### 5.3 Grassmann Manifold  $\mathcal{G}_r(N)$

We refer the readers to [\[15\]](#page-26-14) for the detailed definition and geometry of Grassmann manifold. It is known that  $O(N)$  acts transitively on  $\mathcal{G}_r(N)$ , and the isometry corresponding to each  $O \in O(N)$  is  $X \mapsto O.X = OXO^{\top}$ . The tangent space of  $O(N)$  at identity is  $\mathfrak{o}(N) = \mathscr{A}(\mathbb{R}^{N \times N})$ , the space of all skew-symmetric  $N \times N$ -matrices, where  $\{\mathcal{E}_{ij} : 1 \leq i < j \leq N\}$  is an orthogonal basis as introduced in §[3.4.](#page-8-1) We take local curves  $O_{ij}(t)$  at identity corresponding to each  $\mathcal{E}_{ij}$ , i.e.,  $\frac{d}{dt}O_{ij}(t)|_{t=0} = \mathcal{E}_{ij}$ , then the killing field corresponding to each  $\mathcal{E}_{ij}$  is  $K_X^{ij} := \frac{d}{dt} O_{ij}(t)|_{t=0}$ .  $X = \frac{d}{dt} O_{ij}(t) X O_{ij}^{\top}(t)|_{t=0} = \mathcal{E}_{ij} X - \mathcal{E}_{ij}$  $X\mathcal{E}_{ij}$ , i.e., the vector field  $K^{ij}$  that assigns each X with tangent vector  $\mathcal{E}_{ij}X - X\mathcal{E}_{ij} \in T_X\mathcal{G}_r(N)$ . We plug this into [\(15\)](#page-13-3) so that the  $\kappa_p$  function on  $\mathcal{G}_r(N)$  equals

$$
\kappa_p(X,Y) = \kappa \cdot \sum_{i < j} K_X^{ij} \log(p\kappa) \cdot K_Y^{ij} \log(p\kappa) + \kappa \cdot \sum_{i < j} K_Y^{ij} K_X^{ij} \log \kappa.
$$

Then the first term can be written in a closed form:

$$
\kappa \cdot \sum_{i < j} K_X^{ij} \log(p\kappa) \cdot K_Y^{ij} \log(p\kappa)
$$
\n
$$
= \kappa \cdot \sum_{i < j} \text{tr}[(\mathcal{E}_{ij}X - X\mathcal{E}_{ij})\nabla_X^{\mathbb{R}} \log(p\kappa)] \cdot \text{tr}[(\mathcal{E}_{ij}Y - Y\mathcal{E}_{ij})\nabla_Y^{\mathbb{R}} \log(p\kappa)]
$$
\n
$$
= 4\kappa \cdot \sum_{i < j} \langle \mathcal{E}_{ij}, \mathcal{S}(\nabla_X^{\mathbb{R}} \log(p\kappa))X \rangle_{\mathcal{F}} \cdot \langle \mathcal{E}_{ij}, \mathcal{S}(\nabla_Y^{\mathbb{R}} \log(p\kappa))Y \rangle_{\mathcal{F}}
$$
\n
$$
1 - 4\kappa \langle \mathcal{S}(\mathcal{G}(\nabla_X^{\mathbb{R}} \log(p\kappa)))X \rangle_{\mathcal{F}} \cdot \langle \mathcal{S}_{ij}, \mathcal{S}(\nabla_X^{\mathbb{R}} \log(p\kappa)))Y \rangle_{\mathcal{F}}
$$

Based on Prop.  $5.1 = 4\kappa \langle \mathscr{A}[\mathscr{S}(\nabla^{\mathbb{R}}_{X} \log(p\kappa))X], \mathscr{A}[\mathscr{S}(\nabla^{\mathbb{R}}_{Y} \log(p\kappa))Y]\rangle_{F}$  $5.1 = 4\kappa \langle \mathscr{A}[\mathscr{S}(\nabla^{\mathbb{R}}_{X} \log(p\kappa))X], \mathscr{A}[\mathscr{S}(\nabla^{\mathbb{R}}_{Y} \log(p\kappa))Y]\rangle_{F}$ .

The summation of  $K_Y^{ij} K_X^{ij} \log k$  relies on the specific form of  $\kappa$  and thus has no closed form in general, which can not be further simplified. However, if  $\kappa$  is a radial kernel, i.e.,

$$
\kappa(X,Y) = \exp(-\psi(||X - Y||_{\mathrm{F}}^2)) = \exp(-\psi(2r - 2\,\mathrm{tr}[XY])), \quad X, Y \in \mathcal{G}_r(N),
$$

for some  $\psi \in C^2[0, +\infty)$ . Then we have

$$
K_Y^{ij} K_X^{ij} \log \kappa = -\frac{d^2}{dt dt'} \psi(2r - 2 \operatorname{tr}[O_{ij}(t) X O_{ij}^\top(t) O_{ij}(t') Y O_{ij}^\top(t'))|_{t,t'=0}
$$
  

$$
= 2\psi'(\|X - Y\|_{\mathrm{F}}^2) \cdot \underbrace{\operatorname{tr}[(\mathcal{E}_{ij} X - X \mathcal{E}_{ij})(\mathcal{E}_{ij} Y - Y \mathcal{E}_{ij})]}_{(I)}
$$
  

$$
- 4\psi''(\|X - Y\|_{\mathrm{F}}^2) \cdot \underbrace{\langle Y X - XY, \mathcal{E}_{ij} \rangle_{\mathrm{F}} \cdot \langle XY - Y X, \mathcal{E}_{ij} \rangle_{\mathrm{F}}}_{(II)}.
$$

Due to Prop. [5.1,](#page-15-0) the summation of  $(II)$  over  $i < j$  equals  $-||XY - YX||_F^2$ , as we note that  $XY-YX$  is skew-symmetric itself. For  $(I)$ , note that  $(I) = \text{tr}[2X\mathcal{E}_{ij}Y\mathcal{E}_{ij} - (XY+YX)\mathcal{E}_{ij}\mathcal{E}_{ij}]$ . We set  $X = (x_{ij})$  and  $Y = (y_{ij})$  and then we have

$$
\sum_{i < j} 2 \operatorname{tr}[\mathcal{E}_{ij} X \mathcal{E}_{ij} Y] = \sum_{i < j} \operatorname{tr}[(E_{ji} - E_{ij}) X (E_{ji} - E_{ij}) Y]
$$
\n
$$
= \sum_{i < j} \operatorname{tr}[E_{ij} X E_{ij} Y + E_{ji} X E_{ji} Y - E_{ij} X E_{ji} Y - E_{ji} X E_{ij} Y]
$$
\n
$$
= \sum_{i < j} (x_{ji} y_{ji} + x_{ij} y_{ij} - x_{jj} y_{ii} - x_{ii} y_{jj})
$$
\n
$$
= \sum_{i \neq j} (x_{ij} y_{ij} - x_{ii} y_{jj}) = \sum_{i, j} (x_{ij} y_{ij} - x_{ii} y_{jj})
$$
\n
$$
= \operatorname{tr}(XY) - \operatorname{tr}(X) \operatorname{tr}(Y) = \langle X, Y \rangle_{F} - r^{2}.
$$

Furthermore, we have  $\sum_{i \leq j} tr[(XY + YX)\mathcal{E}_{ij}\mathcal{E}_{ij}] = (N-1)\langle X, Y \rangle_F$  since  $\sum_{i \leq j} \mathcal{E}_{ij}\mathcal{E}_{ij} = -\frac{N-1}{2}I_N$ . Therefore, the summation of (I) over  $i < j$  equals  $N\langle X, Y \rangle_F - r^2$ . We summarize into following theorem:

**Theorem 5.4.** Given density p and kernel k, the  $k_p$  function is given by

$$
\kappa_p(X, Y) = 4 \langle \mathscr{A}(\nabla_X^{\mathbb{R}} \log(p\kappa)X), \mathscr{A}(\nabla_Y^{\mathbb{R}} \log(p\kappa)Y) \rangle_F \kappa(X, Y) + \sum_{i < j} K_Y^{ij} K_X^{ij} \log \kappa \cdot \kappa(X, Y), \tag{24}
$$

where all Euclidean gradients are chosen to be symmetric. Furthermore, if  $\kappa$  is a radial kernel, i.e.,  $\kappa(X,Y) = e^{-\psi(||X-\tilde{Y}||_{\text{F}}^2)}$  for some  $\psi \in C^2[0,+\infty)$ , then the  $\kappa_p$  function equals

$$
\kappa_p(X, Y) = 4\langle \mathscr{A}(\nabla_X^{\mathbb{R}} \log(p\kappa)X), \mathscr{A}(\nabla_Y^{\mathbb{R}} \log(p\kappa)Y) \rangle_F \kappa(X, Y) + 2\psi'(\|X - Y\|_F^2) \cdot (N\langle X, Y \rangle_F - r^2) \kappa(X, Y) + 4\psi''(\|X - Y\|_F^2) \|XY - YX\|_F^2 \kappa(X, Y).
$$
\n(25)

Next we explicitly calculate the closed form of MKSDE for the exponential family on  $\mathcal{G}_r(N)$ . It suffices to calculate the matrix  $Q(X,Y) = \kappa \sum_{i < j} (K_X^{ij} \zeta_k \cdot K_Y^{ij} \zeta_l)_{kl}$  and the vector  $b(X,Y) =$  $\kappa \cdot \sum_{i < j} [K_X^{ij} \eta + K_X^{ij} \log \kappa] K_Y^{ij} \zeta$  introduced in [\(17\)](#page-14-2). For  $Q(X, Y)$ , we have

$$
\sum_{i < j} K_X^{ij} \zeta_k \cdot K_Y^{ij} \zeta_l = \sum_{i < j} \text{tr}[(\mathcal{E}_{ij} X - X \mathcal{E}_{ij}) \nabla_X^{\mathbb{R}} \zeta_k] \cdot \text{tr}[(\mathcal{E}_{ij} Y - Y \mathcal{E}_{ij}) \nabla_Y^{\mathbb{R}} \zeta_l]
$$
\n
$$
= \sum_{i < j} \langle \mathcal{E}_{ij}, \mathcal{S}(\nabla_X^{\mathbb{R}} \zeta_k) X \rangle_F \cdot \langle \mathcal{E}_{ij}, \mathcal{S}(\nabla_Y^{\mathbb{R}} \zeta_l) Y \rangle_F
$$
\nProp. 5.1 =  $\langle \mathcal{A}[\mathcal{S}(\nabla_X^{\mathbb{R}} \zeta_k) X], \mathcal{A}[\mathcal{S}(\nabla_Y^{\mathbb{R}} \zeta_l) Y] \rangle_F,$ 

Similarly, for  $b(X, Y)$ , we have

$$
\sum_{i < j} K_X^{ij} \log(e^n \kappa) \cdot K_Y^{ij} \zeta_k = \langle \mathscr{A}[\mathscr{S}(\nabla_X^{\mathbb{R}} \log(e^n \kappa))X], \mathscr{A}[\mathscr{S}(\nabla_Y^{\mathbb{R}} \zeta_k)Y] \rangle_F.
$$

To sum up, we have

**Theorem 5.5** (MKSDE for exponential family). Given the exponential family  $p_\theta \propto \exp(\theta^\top \zeta(X))$  $\eta(X)$ , let  $Q(X, Y)$  and  $b(X, Y)$  be the matrix and vector given by

$$
Q(X,Y)_{kl} = \kappa(X,Y) \cdot \langle \mathscr{A}[\mathscr{S}(\nabla^{\mathbb{R}}_{X}\zeta_{k})X], \mathscr{A}[\mathscr{S}(\nabla^{\mathbb{R}}_{Y}\zeta_{l})Y]\rangle_{F},
$$
  
\n
$$
b(X,Y)_{k} = \kappa(X,Y) \cdot \langle \mathscr{A}[\mathscr{S}(\nabla^{\mathbb{R}}_{X}\log(e^{\eta}\kappa))X], \mathscr{A}[\mathscr{S}(\nabla^{\mathbb{R}}_{Y}\zeta_{k})Y]\rangle_{F},
$$
\n(26)

Then the MKSDE can be computed via  $(20)$ .

Next, we calculate several examples for specific kernel  $\kappa$  and distribution p on Grassmann manifold.

#### Commonly-used distributions

Most of the widely-used distribution families on Grassmann manifold have intractable normalizing constant, including:

- Matrix Fisher (MF) family:  $p(X) \propto \exp[\text{tr}(F^{\top}X)]$  with parameter  $F \in \mathbb{R}^{N \times N}$ , which is a member of the exponential family since  $tr(F^{\top}X)$  is linear in F. We have  $\nabla_X^{\mathbb{R}} \log p = \mathscr{S}(F)$ and  $\nabla^{\mathbb{R}}_X \eta = 0$ . Note that  $\zeta(X) = X$ , thus the Euclidean gradient of its  $(i, j)$ <sup>th</sup> component  $\zeta_{ij} = \operatorname{tr}[E_{ij}^{\top} X]$  is  $\nabla_X^{\mathbb{R}} \zeta_{ij} = E_{ij} \in \mathbb{R}^{N \times N}$ .
- Riemannian Gaussian (RG) family:  $p(X) \propto \exp(-\frac{d^2(X,\bar{X})}{2\sigma^2})$  with parameters  $\bar{X} \in \mathcal{G}_r(N)$ ,  $\sigma > 0$ . The RG family is not a member of the exponential family. For Riemannian Gaussian family,  $\log p(X) = -\frac{1}{2\sigma^2}d^2(X,\bar{X})$ . The Riemannian gradient of  $d^2$  function is  $-2\log_X(\bar{X})[51]$  $-2\log_X(\bar{X})[51]$ , the Riemannian gradient of  $\log p(X)$  is  $\sigma^{-2} \text{Log}_X(\overline{X})$ . Here Log is the Riemannian logarithm on  $\mathcal{G}_r(N)$ , which can be computed numerically [\[70\]](#page-29-7). It is known [\[6,](#page-25-8) §2.4] that the Riemannian metric on  $\mathcal{G}_r(N)$  is given by  $\langle D_1, D_2 \rangle_X = \frac{1}{2} \text{tr}(D_1 D_2)$ , thus  $\nabla_X^{\mathbb{R}} \log p(X) = \frac{1}{2} \nabla_X^M \log p(X) = \frac{1}{2\sigma^2} \text{Log}_X(\bar{X})$ . The Riemannian logarithm Log can be computed numerically [\[6,](#page-25-8) §5.2].

• Matrix Bingham (MB) and Matrix Fisher-Bingham (MFB) family: The MB and MFB family will degenerate to the matrix Fisher family on  $\mathcal{G}_r(N)$ , as  $p(X) \propto \exp[\text{tr}(XAX + F^{\top}X)]$  $\exp[\text{tr}(AXX + FX)] = \exp[\text{tr}((A+F)X)].$ 

### Commonly-used kernels

The most widely-used kernels on  $\mathcal{G}_r(N)$  include

• Gaussian kernel:

$$
k(X,Y) = \exp\left(-\frac{\tau}{2}||X - Y||_{\mathcal{F}}^2\right) \propto \exp(\tau \operatorname{tr}(XY)), \quad X, Y \in \mathcal{G}_r(N).
$$

Note that  $\nabla_X^{\mathbb{R}} \log k = \tau Y$ ,  $\psi(x) = \frac{\tau x}{2}$ ,  $\psi'(x) = \frac{\tau}{2}$  and  $\psi''(x) = 0$ .

• Inverse quadratic kernel:

$$
k(X,Y) = (\beta + \|X - Y\|_{\text{F}}^2)^{-\gamma}, \quad X, Y \in \mathcal{G}_r(N).
$$
  
Note that  $\nabla_X^{\mathbb{R}} \log k = \frac{2\gamma}{\beta + \|X - Y\|_{\text{F}}^2} Y, \ \psi(x) = \gamma \log(\beta + x), \ \psi'(x) = \frac{\gamma}{\beta + x}$  and  $\psi''(x) = -\frac{\gamma}{(\beta + x)^2}$ .

### 5.4 Space of Symmetric Positive Definite Matrices  $\mathcal{P}(N)$

It is known that  $GL(N)$ , the general linear group of all non-singular matrices, acts transitively on  $\mathcal{P}(N)$ , where the isometry corresponding to each  $G \in GL(N)$  is  $X \mapsto G.X = G^{\top}XG$ . The tangent space of  $\text{GL}(N)$  at identity is  $\mathfrak{gl}(N) := \mathbb{R}^{N \times N}$ , where  $\{E_{ij} : 1 \le i, j \le N\}$  is an orthogonal basis, as introduced in §[3.4.](#page-8-1) We take local curves  $G_{ij}(t)$  of  $E_{ij}$  on  $\mathcal{P}(N)$  such that  $\frac{d}{dt}G_{ij}(t)|_{t=0} = E_{ij}$ , then the killing field corresponding to  $E_{ij}$  is  $K_X^{ij} := \frac{d}{dt}G(t).X|_{t=0} = \frac{d}{dt}G_{ij}(t)^{\top} X G_{ij}(t)|_{t=0} = E_{ji}X + X E_{ij}$ . We plug this into [\(15\)](#page-13-3) so that the  $\kappa_p$  function on  $\mathcal{P}(N)$  equals

$$
\kappa_p(X,Y) = \kappa \cdot \sum_{i < j} K_X^{ij} \log(p\kappa) \cdot K_Y^{ij} \log(p\kappa) + \kappa \cdot \sum_{i < j} K_Y^{ij} K_X^{ij} \log \kappa.
$$

The the first term can be written in a closed form:

$$
\kappa \cdot \sum_{i < j} K_X^{ij} \log(p\kappa) \cdot K_Y^{ij} \log(p\kappa)
$$
\n
$$
= \kappa \cdot \sum_{i < j} \text{tr}[(E_{ji}X + XE_{ij})\nabla_X^{\mathbb{R}} \log(p\kappa)] \cdot \text{tr}[(E_{ji}Y + YE_{ij})\nabla_Y^{\mathbb{R}} \log(p\kappa)]
$$
\n
$$
= 4\kappa \cdot \sum_{i < j} \langle E_{ij}, X \mathcal{S}(\nabla_X^{\mathbb{R}} \log(p\kappa)) \rangle_{\text{tr}} \cdot \langle E_{ij}, Y \mathcal{S}(\nabla_Y^{\mathbb{R}} \log(p\kappa)) \rangle_{\text{tr}}
$$
\n
$$
= 4\kappa \langle X \mathcal{S}(\nabla_X^{\mathbb{R}} \log(p\kappa)), Y \mathcal{S}(\nabla_Y^{\mathbb{R}} \log(p\kappa)) \rangle_{\text{F}}.
$$

The summation of  $K_Y^{ij} K_X^{ij} \log \kappa$  relies on the specific form of  $\kappa$  and thus can not be further simplified. However, if  $\kappa$  is a radial kernel, i.e.,

$$
\kappa(X, Y) = \exp(-\psi(||X - Y||_{\mathcal{F}}^2)), \quad X, Y \in \mathcal{P}(N),
$$

for some  $\psi \in C^2[0, +\infty)$ , then we have

$$
K_Y^{ij} K_X^{ij} \log \kappa = 2\psi'(\|X - Y\|_{\text{F}}^2) \operatorname{tr}[(E_{ji}X + XE_{ij})(E_{ji}Y + YE_{ij})]
$$
  
\n
$$
- 4\psi''(\|X - Y\|_{\text{F}}^2) \operatorname{tr}[(E_{ji}X + XE_{ij})(X - Y)] \operatorname{tr}[(E_{ji}Y + YE_{ij})(X - Y)]
$$
  
\n
$$
= 4\psi'(\|X - Y\|_{\text{F}}^2) \underbrace{\operatorname{tr}[XE_{ij}YE_{ij} + XE_{ij}E_{ji}Y]}_{(i)}
$$
  
\n
$$
+ 16\psi''(\|X - Y\|_{\text{F}}^2) \underbrace{\operatorname{tr}[(X - Y)XE_{ij}] \operatorname{tr}[(X - Y)YE_{ij}]}_{(ii)}.
$$

Denote  $X := (x_{ij})$  and  $Y = (y_{ij})$ . Note that  $tr[E_{ij} X E_{ij} Y] = x_{ji} y_{ij} = x_{ij} y_{ij}$  since X is symmetric. Also note that  $E_{ij}E_{ji} = E_{ii}$ . Therefore, the summation of (i) over i, j equals  $4(N +$  $1)\psi'(\|X-Y\|_{\text{F}}^2)\langle X,Y\rangle_{\text{F}}$ . The summations of  $(ii)$  follows from the Prop. [5.1,](#page-15-0) which equals  $16\psi'(\|X-Y\|_{\text{F}}^2)$  $Y\Vert_{\mathrm{F}}^2 \rangle \langle X(X-Y), Y(X-Y) \rangle_{\mathrm{F}}$ . To sum up,

**Theorem 5.6.** Given the kernel function  $\kappa$ , the  $\kappa_p$  function is,

$$
\kappa_p(X, Y) = 4\langle X \mathcal{S}(\nabla_X^{\mathbb{R}} \log(p\kappa)), Y \mathcal{S}(\nabla_Y^{\mathbb{R}} \log(p\kappa)) \rangle_F \cdot \kappa(X, Y) + \sum_{i,j} K_Y^{ij} K_Y^{ij} \log \kappa \cdot \kappa(X, Y).
$$
\n(27)

Furthermore, if  $\kappa$  is a radial kernel, i.e.,  $\kappa(X,Y) = e^{-\psi(||X-Y||_F^2)}$  for some  $\psi \in C^2[0,+\infty)$ , then we have,

$$
\kappa_p(X, Y) = 4\langle X \mathcal{S}(\nabla_X^{\mathbb{R}} \log(p\kappa)), Y \mathcal{S}(\nabla_Y^{\mathbb{R}} \log(p\kappa)) \rangle_F \cdot \kappa(X, Y) + 4(N+1)\langle X, Y \rangle_F \psi'(\|X - Y\|_F^2) \cdot \kappa(X, Y) + 16\psi''(\|X - Y\|_F^2) \langle X(X - Y), Y(X - Y) \rangle_F \cdot \kappa(X, Y).
$$
\n(28)

Next we explicitly calculate the closed form of MKSDE for the exponential family on  $\mathcal{G}_r(N)$ . It suffices to calculate the matrix  $Q(X,Y) = \kappa \sum_{i,j} (K_X^{ij} \zeta_k \cdot K_Y^{ij} \zeta_l)_{kl}$  and the vector  $b(X,Y) =$  $\kappa \cdot \sum_{i,j} [K_X^{ij} \eta + K_X^{ij} \log \kappa] K_Y^{ij} \zeta$  introduced in [\(17\)](#page-14-2). For  $Q(X, Y)$ , we have

$$
\sum_{i,j} K_X^{ij} \zeta_k \cdot K_Y^{ij} \zeta_l = \sum_{i,j} \text{tr}[E_{ji}X + XE_{ij}) \nabla_X^{\mathbb{R}} \zeta_k] \cdot \text{tr}[(E_{ji}Y - YE_{ij}) \nabla_Y^{\mathbb{R}} \zeta_l]
$$
  

$$
= \sum_{i,j} \langle E_{ij}, X \mathcal{S}(\nabla_X^{\mathbb{R}} \zeta_k) \rangle_{\mathcal{F}} \cdot \langle E_{ij}, Y \mathcal{S}(\nabla_Y^{\mathbb{R}} \zeta_l) \rangle_{\mathcal{F}}
$$
  
Prop. 5.1 =  $\langle X \mathcal{S}(\nabla_X^{\mathbb{R}} \zeta_k), Y \mathcal{S}(\nabla_Y^{\mathbb{R}} \zeta_l) \rangle_{\mathcal{F}}$ ,

Similarly, for  $b(X, Y)$ , we have

$$
\sum_{i < j} K_X^{ij} \log(e^{\eta} \kappa) \cdot K_Y^{ij} \zeta_k = \langle X \mathcal{S}(\nabla_X^{\mathbb{R}} \log(e^{\eta} \kappa)), Y \mathcal{S}(\nabla_Y^{\mathbb{R}} \zeta_k) \rangle_F.
$$

To sum up, we have

**Theorem 5.7** (MKSDE for exponential family). Given the exponential family  $p_\theta \propto \exp(\theta^\top \zeta(X))$  $\eta(X)$ , let  $Q(X,Y)$  and  $b(X,Y)$  be the matrix and vector given by

$$
Q(X,Y)_{kl} = \kappa(X,Y) \cdot \langle X \mathcal{S}(\nabla^{\mathbb{R}}_X \zeta_k), Y \mathcal{S}(\nabla^{\mathbb{R}}_Y \zeta_l) \rangle_F,
$$
  
\n
$$
b(X,Y)_k = \kappa(X,Y) \cdot \langle X \mathcal{S}(\nabla^{\mathbb{R}}_X \log(e^{\eta} \kappa)), Y \mathcal{S}(\nabla^{\mathbb{R}}_Y \zeta_k) \rangle_F,
$$
\n(29)

Then the MKSDE can be computed using  $(20)$ .

#### Commonly-used distributions

- Wishart family:  $p(X; V, r) \propto |X|^{(r-N+1)/2} \exp\left(-\frac{1}{2} \text{tr}[V^{-1}X]\right)$ , with parameter  $V \in \mathcal{P}(N)$ ,  $1 \leq N \leq r \in \mathbb{N}_+$ . The Wishart distribution is not a member of the exponential family, since the domain of V is  $\mathcal{P}(N)$ , which is not a vector space. Due to Jacobi's formula [\[38,](#page-27-12) Thm. 8.1], we have  $\nabla_X^{\mathbb{R}} \log p = -\frac{1}{2} V^{-1} + \frac{r - N + 1}{2} X^{-1}$ .
- Riemannian Gaussian family:  $p(X) \propto \exp\left(-\frac{d^2(X,\bar{X})}{2\sigma^2}\right)$  $\left(\frac{(X,X)}{2\sigma^2}\right)$  with parameters  $\bar{X} \in \mathcal{P}(N)$ ,  $\sigma > 0$ . The Riemannian Gaussian family is not a member of the exponential family. As shown in [\[45\]](#page-28-14), on  $\mathcal{P}(N)$ ,  $d^2(X,\bar{X}) = \text{tr}(\text{Log}^2(\bar{X}^{-1}X))$ . By [\[45,](#page-28-14) Prop. 2.1], we have

$$
\frac{d}{dt}d^{2}(X(t),\bar{X}) = \frac{d}{dt}\operatorname{tr}(\operatorname{Log}^{2}(\bar{X}^{-1}X(t))) = 2\operatorname{tr}\left[\operatorname{Log}(\bar{X}^{-1}X(t))X^{-1}(t)\frac{d}{dt}X(t)\right].
$$

Therefore,  $\nabla_X^{\mathbb{R}} \log p(X) = -\sigma^{-2} X^{-1} \text{Log}(\bar{X}^{-1} X).$ 

#### Commonly-used kernels

The most widely-used kernels on  $\mathcal{P}(N)$  include

• Gaussian kernel:

$$
\kappa(X,Y) = \exp\left(-\frac{\tau}{2}||X - Y||_{\mathcal{F}}^2\right), \quad X, Y \in \mathcal{P}(N).
$$

Note that  $\nabla_X^{\mathbb{R}} \log \kappa = \tau(Y - X), \psi(x) = \frac{\tau x}{2}, \psi'(x) = \frac{\tau}{2}, \psi''(x) = 0.$ 

• Inverse quadratic kernel:

$$
\kappa(X,Y) = (\beta + \|X - Y\|_{\mathcal{F}}^2)^{-\gamma}, \quad X, Y \in \mathcal{G}_r(N).
$$

Note that  $\nabla^{\mathbb{R}}_X \log \kappa = \frac{2\gamma(Y-X)}{\beta + \|X-Y\|}$  $\frac{2\gamma(Y-X)}{\beta+\|X-Y\|^2_{\text{tr}}}, \psi(x)=\gamma\log(\beta+x), \, \psi'(x)=\frac{\gamma}{\beta+x}, \, \psi''(x)=-\frac{\gamma}{(\beta+x)^2}.$ 

# <span id="page-23-0"></span>6 Experiments

In this section, we present two experiments to demonstrate the power of our kernel Stein method on manifolds. In the first experiments, we will compare the MLE and MKSDE in estimating the parameters of the matrix Fisher distribution on a Stiefel manifold  $\mathcal{V}_r(N)$ , illustrating the advantage of MKSDE over MLE due to the presence of the intractable normalizing constant in the MLE. In the second experiment, we validate the power of composite goodness of fit test using MKSDE by comparing the matrix Fisher distribution and matrix Bingham distribution on a Stiefel manifold. The kernel we choose in these experiments is the Gaussian kernel  $\kappa(X, Y) = \exp(\text{tr}(X^\top Y)), X, Y \in$  $V_r(N)$ . Code for computing the KSD, MKSDE and conducting the composite goodness of fit test is provided on GitHub at <https://github.com/cvgmi/KSD-on-Riemannian-Manifolds>.

# <span id="page-23-1"></span>6.1 MKSDE vs. MLE

The normalizing constant  $c(F)$  of the matrix Fisher distribution  $p(X) \propto \exp(\text{tr}(F^{\top}X))$  on a Stiefel manifold is an intractable hypergeometric function of F. The most widely-used classical method to compute the MLE of the matrix Fisher distribution on a Stiefel manifold utilize two direct approximate solutions, introduced in [\[40,](#page-27-13) §13.2.3]. In general, the first solution approximates the MLE well when  $F$  is small, while the second approximates the MLE relatively accurately when

<span id="page-24-0"></span>

Figure 1: Frobenius distances between the estimators and ground truth

 $F$  is large. However, it should be noted that both approximate solutions work poorly for mediumvalued  $F$ , and the second approximate solution involves solving a non-linear multivariate equation, which is burdened with relatively hight computational cost.

In this experiment, we obtain the samples from a matrix Fisher distribution  $p(X) \propto \exp(\text{tr}(F_0^{\top} X))$ with ground truth  $F_0$  on  $\mathcal{V}_r(N)$ , then compute the MLE,  $\hat{F}_{\text{MLE}}$ , the MKSDE  $\hat{F}_{\text{MKSDE-U}}$  and  $\hat{F}_{MKSDE-V}$  obtained by minimizing  $U_n^w$  and  $V_n^w$  in [\(12\)](#page-10-2) respectively. The figure [1](#page-24-0) shows the Frobenius distance between the ground truth  $F_0$  and estimators including  $\hat{F}_{\text{MLE}}, \hat{F}_{\text{MKSDE-U}}$  and  $\hat{F}_{\text{MKSDE-V}}$ with varying values of  $F_0$ . Here we set  $E_1 = (1,0;1,0;1,0) \in \mathbb{R}^{3 \times 2}$  and  $E_2 = (1,1;1,1;1,1) \in \mathbb{R}^{3 \times 2}$ and the value of  $F_0$  will vary in  $0.3 * E_1$ ,  $E_1$ ,  $5 * E_1$ ,  $0.3 * E_2$ ,  $E_2$  and  $5 * E_2$ . As depicted in figure [1,](#page-24-0) the approximate solution using the MLE worsens as  $F_0$  becomes larger. This experiment demonstrates the performance of MKSDE which is independent of the normalization constant.

#### 6.2 Composite goodness of fit test

In this experiment, we conduct the composite goodness of fit test presented in algorithm block [1](#page-13-2) to check whether a group of samples from a specific matrix Fisher distribution can be modeled by the matrix Bingham family. The table  $1$  depicts the  $p$  values of the composite goodness-of-fit test under different values of  $F$  and the number of samples  $n$ . Intuitively, an MF distribution with small  $F$  becomes nearly uniform, and an MF distribution with large  $F$  will concentrate around the dominant directions specified by F. Therefore, the MF distributions belong approximately to the MB family when  $A$  is small or large, but differ from the shape of the MB family for  $F$  in-between. This is consistent with the results in Table [1.](#page-25-9) In addition, the loss function corresponding to the V-statistic shows better stability in the context of optimization compared to the U-statistic, this is because the global minimizer of  $V_n^w$  always exists, as stated in Thm. [5.1.](#page-14-4)

<span id="page-25-9"></span>

(a) p-values of the U-stat					
number of samples	100	150	200	250	300
$F = 0.3 * E_1$	0.4670	0.3307	0.0582	0.0223	0.0034
$F=E_1$	0.3420	0.0713	0.0320	0.0018	0.0007
$F = 5 * E_1$	0.2624	0.1528	0.0139	0.0282	0.0214
(b) p-values of the V-stat					
number of samples	100	150	200	250	300
$F = 0.3 * E_1$	0.3923	0.1506	0.0348	0.0213	0.0028
$F=E_1$	0.0687	0.0045	0.0012	0.0001	0.0000
$F = 5 * E_1$	0.0202	0.0173	0.0008	0.0030	0.0024

Table 1: p-values of the composite goodness of fit test

# Acknowledgements

This research was in part funded by the NIH NINDS and NIA grant RO1NS121099 to Vemuri.

# References

- <span id="page-25-7"></span>[1] P-A Absil, Robert Mahony, and Rodolphe Sepulchre. Optimization algorithms on matrix manifolds. Princeton University Press, 2008.
- <span id="page-25-4"></span>[2] Alessandro Barp, Francois-Xavier Briol, Andrew Duncan, Mark Girolami, and Lester Mackey. Minimum Stein discrepancy estimators. Advances in Neural Information Processing Systems, 32, 2019.
- <span id="page-25-5"></span>[3] Alessandro Barp, Chris Oates, Emilio Porcu, Mark Girolami, et al. A Riemann-Stein kernel method. arXiv preprint arXiv:1810.04946, 2018.
- <span id="page-25-2"></span>[4] Peter J Basser, James Mattiello, and Denis LeBihan. Mr diffusion tensor spectroscopy and imaging. Biophysical journal, 66(1):259–267, 1994.
- <span id="page-25-1"></span>[5] Martin Bauer, Martins Bruveris, and Peter W. Michor. Overview of the geometries of shape spaces and diffeomorphism groups. Journal of Mathematical Imaging and Vision,  $50(1-2)$ ,  $2014$ .
- <span id="page-25-8"></span>[6] Thomas Bendokat, Ralf Zimmermann, and P-A Absil. A Grassmann manifold handbook: Basic geometry and computational aspects. arXiv preprint arXiv:2011.13699, 2020.
- <span id="page-25-0"></span>[7] Thomas J. Bridges and Sebastian Reich. Computing lyapunov exponents on a stiefel manifold. Physica D: Nonlinear Phenomena, 156(3):219–238, 2001.
- <span id="page-25-6"></span>[8] Claudio Carmeli, Ernesto De Vito, Alessandro Toigo, and Veronica Umanitá. Vector valued reproducing kernel hilbert spaces and universality. Analysis and Applications, 8(01):19–61, 2010.
- <span id="page-25-3"></span>[9] Igor Carrara, Bruno Aristimunha, Marie-Constance Corsi, Raphael Yokoingawa de Camargo, Sylvain Chevallier, and Theodore Papadopoulo. Geometric neural network based on phase space for bci-eeg decoding. Journal of Neural Engineering, 2024.
- <span id="page-26-0"></span>[10] Rudrasis Chakraborty, Monami Banerjee, and Baba C Vemuri. Statistics on the space of trajectories for longitudinal data analysis. In 2017 IEEE 14th international symposium on biomedical imaging (ISBI 2017), pages 999–1002. IEEE, 2017.
- <span id="page-26-10"></span>[11] Kacper Chwialkowski, Heiko Strathmann, and Arthur Gretton. A kernel test of goodness of fit. In International conference on machine learning, pages 2606–2615. PMLR, 2016.
- <span id="page-26-1"></span>[12] Wei Dai, Youjian Liu, and Brian Rider. Quantization bounds on grassmann manifolds and applications to mimo communications. IEEE Transactions on Information Theory, 54(3):1108– 1123, 2008.
- <span id="page-26-3"></span>[13] Ganggang Dong, Gangyao Kuang, Na Wang, and Wei Wang. Classification via sparse representation of steerable wavelet frames on grassmann manifold: Application to target recognition in sar image. IEEE Transactions on Image Processing, 26(6):2892–2904, 2017.
- <span id="page-26-4"></span>[14] Xiaowen Dong, Pascal Frossard, Pierre Vandergheynst, and Nikolai Nefedov. Clustering on multi-layer graphs via subspace analysis on grassmann manifolds. IEEE Transactions on signal processing, 62(4):905–918, 2013.
- <span id="page-26-14"></span>[15] Alan Edelman, Tomás A Arias, and Steven T Smith. The geometry of algorithms with orthogonality constraints. SIAM journal on Matrix Analysis and Applications, 20(2):303–353, 1998.
- <span id="page-26-15"></span>[16] Lawrence C Evans and Ronald F Garzepy. Measure Theory and Fine Properties of Functions. Routledge, 2018.
- <span id="page-26-13"></span>[17] Herbert Federer. Geometric measure theory. Springer, 1996.
- <span id="page-26-5"></span>[18] Thomas P Fletcher and Sarang Joshi. Riemannian geometry for the statistical analysis of diffusion tensor data. Signal Processing, 87(2):250–262, 2007.
- <span id="page-26-11"></span>[19] Edward Fuselier and Grady B Wright. Scattered data interpolation on embedded submanifolds with restricted positive definite kernels: Sobolev error estimates. SIAM Journal on Numerical Analysis, 50(3):1753–1776, 2012.
- <span id="page-26-12"></span>[20] Matthew P Gaffney. A special Stokes's theorem for complete riemannian manifolds. Annals of Mathematics, pages 140–145, 1954.
- <span id="page-26-6"></span>[21] Igor Gilitschenski, Roshni Sahoo, Wilko Schwarting, Alexander Amini, Sertac Karaman, and Daniela Rus. Deep orientation uncertainty learning based on a bingham loss. In International conference on learning representations, 2020.
- <span id="page-26-7"></span>[22] Jared Glover and Leslie Pack Kaelbling. Tracking the spin on a ping pong ball with the quaternion bingham filter. In 2014 IEEE international conference on robotics and automation (ICRA), pages 4133–4140. IEEE, 2014.
- <span id="page-26-2"></span>[23] Colin R Goodall and Kanti V Mardia. Projective shape analysis. Journal of Computational and Graphical Statistics, 8(2):143–168, 1999.
- <span id="page-26-8"></span>[24] Jackson Gorham and Lester Mackey. Measuring sample quality with stein's method. Advances in Neural Information Processing Systems, 28, 2015.
- <span id="page-26-9"></span>[25] Jackson Gorham and Lester Mackey. Measuring sample quality with kernels. In International Conference on Machine Learning, pages 1292–1301. PMLR, 2017.
- <span id="page-27-11"></span>[26] Alexander Graham. Kronecker Product and Matrix Calculus with Applications. Dover, 2018.
- <span id="page-27-10"></span>[27] Arthur Gretton, Kenji Fukumizu, Zaid Harchaoui, and Bharath K Sriperumbudur. A fast, consistent kernel two-sample test. Advances in Neural Information Processing Systems, 22, 2009.
- <span id="page-27-9"></span>[28] Oscar Key, Arthur Gretton, François-Xavier Briol, and Tamara Fernandez. Composite goodness-of-fit tests with kernels. arXiv preprint arXiv:2111.10275, 2021.
- <span id="page-27-2"></span>[29] Alfred Kume, Simon P Preston, and Andrew TA Wood. Saddlepoint approximations for the normalizing constant of fisher–bingham distributions on products of spheres and stiefel manifolds. Biometrika, 100(4):971–984, 2013.
- <span id="page-27-3"></span>[30] Alfred Kume and Tomonari Sei. On the exact maximum likelihood inference of fisher–bingham distributions using an adjusted holonomic gradient method. Statistics and Computing, 28:835– 847, 2018.
- <span id="page-27-1"></span>[31] Denis Le Bihan, Jean-François Mangin, Cyril Poupon, Chris A Clark, Sabina Pappata, Nicolas Molko, and Hughes Chabriat. Diffusion tensor imaging: concepts and applications. Journal of Magnetic Resonance Imaging: An Official Journal of the International Society for Magnetic Resonance in Medicine, 13(4):534–546, 2001.
- <span id="page-27-14"></span>[32] John M Lee. Riemannian Manifolds: An Introduction to Curvature. Springer Science & Business Media, 2006.
- <span id="page-27-15"></span>[33] John M Lee. *Introduction to Smooth Manifolds*. Springer, 2013.
- <span id="page-27-8"></span>[34] Taeyoung Lee. Bayesian attitude estimation with the matrix fisher distribution on so (3). IEEE Transactions on Automatic Control, 63(10):3377–3392, 2018.
- <span id="page-27-5"></span>[35] Christophe Ley, Gesine Reinert, and Yvik Swan. Stein's method for comparison of univariate distributions. 2017.
- <span id="page-27-0"></span>[36] Christophe Ley and Thomas Verdebout. Applied directional statistics: modern methods and case studies. CRC Press, 2018.
- <span id="page-27-4"></span>[37] Qiang Liu, Jason Lee, and Michael Jordan. A kernelized Stein discrepancy for goodness-of-fit tests. In International Conference on Machine Learning, pages 276–284. PMLR, 2016.
- <span id="page-27-12"></span>[38] Jan R Magnus and Heinz Neudecker. Matrix differential calculus with applications in statistics and econometrics. John Wiley & Sons, 2019.
- <span id="page-27-16"></span>[39] Jonathan H. Manton and Pierre-Olivier Amblard. 2015.
- <span id="page-27-13"></span>[40] Kanti V Mardia, Peter E Jupp, and KV Mardia. Directional statistics, volume 2. Wiley Online Library, 2000.
- <span id="page-27-6"></span>[41] Takuo Matsubara, Jeremias Knoblauch, François-Xavier Briol, and Chris J Oates. Robust generalised Bayesian inference for intractable likelihoods. Journal of the Royal Statistical Society Series B: Statistical Methodology, 84(3):997–1022, 2022.
- <span id="page-27-7"></span>[42] Charles A Micchelli, Yuesheng Xu, and Haizhang Zhang. Universal kernels. Journal of Machine Learning Research, 7(12), 2006.
- <span id="page-28-5"></span>[43] Guillaume Mijoule, Gesine Reinert, and Yvik Swan. Stein operators, kernels and discrepancies for multivariate continuous distributions. arXiv preprint arXiv:1806.03478, 2018.
- <span id="page-28-6"></span>[44] Guillaume Mijoule, Gesine Reinert, and Yvik Swan. Stein's density method for multivariate continuous distributions. arXiv preprint arXiv:2101.05079, 2021.
- <span id="page-28-14"></span>[45] Maher Moakher. A differential geometric approach to the geometric mean of symmetric positivedefinite matrices. SIAM journal on matrix analysis and applications, 26(3):735–747, 2005.
- <span id="page-28-2"></span>[46] Bishwarup Mondal, Satyaki Dutta, and Robert W Heath. Quantization on the grassmann manifold. IEEE Transactions on Signal Processing, 55(8):4208–4216, 2007.
- <span id="page-28-17"></span>[47] James R Munkres. Topology. (No Title), 2000.
- <span id="page-28-9"></span>[48] Chris Oates et al. Minimum kernel discrepancy estimators. arXiv preprint arXiv:2210.16357, 2022.
- <span id="page-28-7"></span>[49] Chris J Oates, Mark Girolami, and Nicolas Chopin. Control functionals for Monte Carlo integration. Journal of the Royal Statistical Society. Series B (Statistical Methodology), pages 695–718, 2017.
- <span id="page-28-0"></span>[50] Karim Oualkacha and Louis-Paul Rivest. On the estimation of an average rigid body motion. Biometrika, 99(3):585–598, 2012.
- <span id="page-28-13"></span>[51] Xavier Pennec. Hessian of the riemannian squared distance. Preprint. https://www-sop. inria. fr/members/Xavier. Pennec/AOS-DiffRiemannianLog. pdf, 2017.
- <span id="page-28-12"></span>[52] Peter Petersen. Riemannian Geometry. Springer, 2016.
- <span id="page-28-10"></span>[53] Xiaoda Qu, Xiran Fan, and Baba C. Vemuri. Kernel stein discrepancy on lie groups: Theory and applications. IEEE Transactions on Information Theory, 70(12):8961–8974, 2024.
- <span id="page-28-8"></span>[54] Xiaoda Qu and Baba C Vemuri. A Framework for Improving the Characterization Scope of Stein's Method on Riemannian Manifolds. arXiv preprint arXiv:2209.08424, 2022.
- <span id="page-28-16"></span>[55] Herman Rubin. Uniform convergence of random functions with applications to statistics. The Annals of Mathematical Statistics, pages 200–203, 1956.
- <span id="page-28-11"></span>[56] Isaac J Schoenberg. Metric spaces and completely monotone functions. Annals of Mathematics, 39(4):811–841, 1938.
- <span id="page-28-1"></span>[57] Jörn Schulz, Sungkyu Jung, Stephan Huckemann, Michael Pierrynowski, JS Marron, and Stephen M Pizer. Analysis of rotational deformations from directional data. Journal of Computational and Graphical Statistics, 24(2):539–560, 2015.
- <span id="page-28-15"></span>[58] Robert J Serfling. Approximation theorems of mathematical statistics. John Wiley & Sons, 2009.
- <span id="page-28-4"></span>[59] Krishan Sharma and Renu Rameshan. Image set classification using a distance-based kernel over affine grassmann manifold. IEEE transactions on neural networks and learning systems, 32(3):1082–1095, 2020.
- <span id="page-28-3"></span>[60] A. Srivastava. A bayesian approach to geometric subspace estimation. IEEE Transactions on Signal Processing, 48(5):1390–1400, 2000.
- <span id="page-29-5"></span>[61] Ingo Steinwart and Andreas Christmann. Support Vector Machines. Springer Science & Business Media, 2008.
- <span id="page-29-6"></span>[62] Sofia Suvorova, Stephen D Howard, and Bill Moran. Tracking rotations using maximum entropy distributions. IEEE Transactions on Aerospace and Electronic Systems, 57(5):2953–2968, 2021.
- <span id="page-29-9"></span>[63] Terence Tao. Topics in random matrix theory, volume 132. American Mathematical Society, 2023.
- <span id="page-29-0"></span>[64] Pavan Turaga, Ashok Veeraraghavan, and Rama Chellappa. Statistical analysis on stiefel and grassmann manifolds with applications in computer vision. In 2008 IEEE conference on computer vision and pattern recognition, pages 1–8. IEEE, 2008.
- <span id="page-29-2"></span>[65] Zhizhou Wang and B.C. Vemuri. Dti segmentation using an information theoretic tensor dissimilarity measure. IEEE Transactions on Medical Imaging, 24(10):1267–1277, 2005.
- <span id="page-29-4"></span>[66] Wenkai Xu and Takeru Matsuda. Interpretable Stein Goodness-of-fit Tests on Riemannian Manifold. In International Conference on Machine Learning, pages 11502–11513. PMLR, 2021.
- <span id="page-29-3"></span>[67] Xiaoqi Xu, Darrick Lee, Nicolas Drougard, and Raphaëlle N Roy. Signature methods for braincomputer interfaces. Scientific Reports, 13(1):21367, 2023.
- <span id="page-29-1"></span>[68] Ryoma Yataka, Kazuki Hirashima, and Masashi Shiraishi. Grassmann manifold flows for stable shape generation. Advances in Neural Information Processing Systems, 36:72377–72411, 2023.
- <span id="page-29-8"></span>[69] In-Kwon Yeo and Richard A Johnson. A uniform strong law of large numbers for u-statistics with application to transforming to near symmetry. Statistics  $\mathscr B$  probability letters, 51(1):63–69, 2001.
- <span id="page-29-7"></span>[70] Ralf Zimmermann. A matrix-algebraic algorithm for the riemannian logarithm on the stiefel manifold under the canonical metric. SIAM Journal on Matrix Analysis and Applications, 38(2):322–342, 2017.

# Supplement

Theory and Applications of Kernel Stein Discrepancy on Riemannian Manifolds

Xiaoda Qu, Baba C. Vemuri

In this supplement we provide the necessary mathematical backgrounds that were omitted in the main text and the proofs to our theorems.

# A Mathematical Background

In this section, we set up the notations and standing assumptions, and then briefly introduce several mathematical definitions and concepts that will be used in the rest of the paper. For a more comprehensive study of the background material presented here, we refer the readers to some standard text books on differential geometry [\[32,](#page-27-14) [33,](#page-27-15) [52\]](#page-28-12), reproducing kernel Hilbert spaces (RKHS) [\[8,](#page-25-6) [61\]](#page-29-5) and Bochner integral [\[61,](#page-29-5) §A.5.4].

#### Riemannian Geometry

We begin by introducing smooth manifolds followed by tangent and cotangent tensors, then, Riemannian metric and distance. This is followed by the volume measure, the divergence theorem, Lie groups, Killing fields and finally Riemannian homogeneous spaces.

#### Smooth Manifolds

A manifold M is a second countable Hausdorff space that is locally homeomorphic to an open set in  $\mathbb{R}^d$ . The number d is the dimension of M. Such local homeomorphisms are called *charts*. A smooth structure on  $M$  is assigned via a collection of charts that covers the manifold such that the composition of every pair of charts of the collection is smooth on  $\mathbb{R}^d$ . A manifold with a smooth structure is called a smooth manifold.

#### Tangent and Cotangent Vector

A local curve at a point  $x \in M$  is a smooth map  $\mathfrak{c} : (-\epsilon, \epsilon) \to M$  with  $\mathfrak{c}(0) = x$ . A tangent vector at x is an equivalence class of local curves at x with the same gradient  $\frac{d}{dt}(\xi \circ \mathfrak{c})(0)$  for some chart  $(U, \xi)$ . The tangent space  $T_xM$  is the n-dimensional vector space of tangent vectors at  $x \in M$ . The *cotangent space*  $T_x^*M$  is then the dual space of  $T_xM$ . For a tangent vector  $D_x \in T_xM$  and a smooth functions f,  $D_x f$  represents the directional derivative of f along  $D_x$ , defined via  $D_x f = \frac{d}{dt} f \circ \mathfrak{c}(0)$ where c is a local curve corresponding to  $D_x$ . A vector field D is a smooth assignment of vector fields to tangent spaces at each point, where  $D_x$  represents the vector assigned to tangent space at x. For a map  $f: M \to M'$  between two manifolds, the differential  $df_x$  of f at x is defined as the map  $df_x: T_xM \to T_{f(x)}M'$ ,  $[\mathfrak{c}(t)] \to [f(\mathfrak{c}(t))]$ , i.e, mapping the equivalence class of  $\mathfrak{c}(t)$ , which is a tangent vector in  $T_xM$ , to the equivalence class of  $f(\mathfrak{c}(t))$ , which is a tangent vector in  $T_{f(x)}M'$ .

#### Riemannian Metric and Distance

A Riemannian metric g is a smooth assignment of inner product to tangent spaces at each point  $x \in M$ . A Riemannian manifold M is a smooth manifold endowed with a Riemannian metric g. For a connected Riemannian manifold  $M$ , any two points  $x, y$  can be connected by a piecewise smooth curve  $\gamma : [0, 1] \to M$  with  $\gamma(0) = x, \gamma(1) = y$ . The length of the curve is then defined as  $L(\gamma) = \int_0^1 |\gamma'(t)| dt$ . The Riemannian distance d is then given by  $d(x, y) =$ inf  $\{L(\gamma):$  for all  $\gamma$  connecting  $x, y\}$ . A Riemannian manifold M is called *complete* if  $(M, d)$  is a complete metric space.

#### Volume Measure

The volume measure  $\mu$  on M can be defined as the n-dimensional Hausdorff measure or defined through integration of differential forms. These two definitions coincide. For details we refer the readers to [\[16,](#page-26-15) [33\]](#page-27-15). In this work, the readers are only required to know that the volume measure is the generalization of Euclidean Lebesgue measure to Riemannian manifolds, and commonly plays the role of a dominating measure of probability density functions on the manifolds.

#### Differential Operator

The Riemannian gradient operator  $\nabla$  maps a smooth function f to a smooth vector field  $\nabla f$ such that  $D(f) = q(D, \nabla f)$  for all smooth vector fields D. For the rather complicated definition of the divergence operator div, we refer the reader to  $[52, 82.1.3]$ . Very briefly, it maps a smooth vector field D to a smooth function div D. The divergence operator satisfies the property  $[52, \text{Exer. } 2.5.5]$  $\operatorname{div}(fD) = Df + f \operatorname{div} D.$ 

**Theorem A.1** (Divergence theorem). Let M be a complete Riemannian manifold. For a locally Lipschitz continuous vector field  $D$  on  $M$  such that  $|D|$  and div  $D$  are both integrable w.r.t volume measure  $\mu$ , the following identity holds:  $\int_M \text{div } D d\mu = 0$ .

*Proof.* See  $[20]$ .

#### Embedding

A smooth map  $\psi : M \to \mathbb{R}^{d'}$  for some  $d' \geq d$  is said to be an *smooth embedding* if  $d\psi_x$  is injective for all  $x \in M$  and  $\psi$  is an homeomorphism between M and  $\psi(M) \subset \mathbb{R}^{d'}$  with subspace topology. Whitney embedding theorem [\[33,](#page-27-15) Thm. 6.15] shows that every smooth manifold admits a smooth embedding into higher dimensional Euclidean space.

#### Reproducing Kernel Hilbert Space

In this section, we briefly introduce the reproducing kernel Hilbert space (RKHS) which will be needed later to define the kernel Stein discrepancy (KSD). For a detailed exposition on RKHS, we refer the readers to [\[39\]](#page-27-16) and [\[61,](#page-29-5) §4].

A kernel function, abbreviated as kernel in this work, on a set X, is a bivariate function  $\kappa$ :  $X \times X \to \mathbb{R}$  such that (i) symmetric, i.e.,  $\kappa(x, y) = \kappa(y, x)$  for all  $x, y \in X$ ; (ii) positive definite, i.e., the matrix  $(\kappa(x_i, x_j))_{i,j} \succeq 0$  for all  $x_i \in X$ ,  $1 \leq i \leq n$ . It is widely known that (see e.g., [\[61,](#page-29-5) Thm. 4.20 & Thm. 4.21]), for each kernel  $\kappa$  on X, there exists an unique Hilbert space  $\mathcal{H}_{\kappa}$ , whose elements are real-valued functions on X, such that for each fixed  $x_0 \in X$ , the uni-variate function  $\kappa_{x_0}(\cdot) := \kappa(x_0, \cdot)$  belongs to  $\mathcal{H}_{\kappa}$ , and satisfies  $f(x_0) := \langle f, \kappa_{x_0} \rangle_{\mathcal{H}_{\kappa}}$  for all  $f \in \mathcal{H}_{\kappa}$ . Specifically, note

that  $\kappa(x,y) = \langle \kappa_x, \kappa_y \rangle_{\mathcal{H}_\kappa}$ . The function space  $\mathcal{H}_\kappa$  is said to be the *reproducing kernel Hilbert space* (RKHS) associated with kernel  $\kappa$ .

If X is a locally compact topological space, we say a kernel  $\kappa$  on X is  $C_0$  if  $\mathcal{H}_{\kappa} \subset C_0(X)$ , i.e., the space of continuous functions that vanish at infinity. Furthermore, a  $C_0$  kernel  $\kappa$  is said to be  $C_0$ -universal if  $\mathcal{H}_\kappa$  is dense in  $C_0(X)$  under the uniform norm. One type of the most widely-used  $C_0$ -universal kernels is the *radial kernels*:

**Theorem A.2** (Radial kernels). Suppose  $\psi : [0, +\infty) \to \mathbb{R}$  is a continuous. Then

- 1. The bivariate function  $\kappa(x, y) := \exp(-\psi(\|x y\|_{\mathbb{R}^d}^2))$  for  $x, y \in \mathbb{R}^d$  is a kernel on  $\mathbb{R}^d$  iff there exists a finite Borel measure  $\lambda$  on  $[0, +\infty)$  such that  $\psi(t) := -\log \int_0^{+\infty} e^{-t\sigma} d\lambda(\sigma)$ ,  $t \ge 0$ . Such kernel  $\kappa$  is said to be a radial kernel on  $\mathbb{R}^d$ .
- 2. A radial kernel  $\kappa$  is  $C_0$ -universal on  $\mathbb{R}^d$  if  $\lambda$  is not concentrated at 0.

*Proof.* The first statement is a classical result proved in [\[56\]](#page-28-11). The second statement is proved in [\[42,](#page-27-7) Thm. 17].

Furthermore, if the topological space  $X$  is endowed with a differential structure, i.e., a smooth manifold, then the kernel  $\kappa$  and its associated RKHS  $\mathcal{H}_{\kappa}$  will satisfy following properties:

**Theorem A.3.** Suppose  $\kappa$  is a kernel on a manifold M, and  $\kappa \in C^2(M \times M)$ , i.e., twice continuous differentiable, then all  $f \in \mathcal{H}_{\kappa}$  is  $C^1$ . Furthermore, given a tangent vector  $D \in T_{x_0}M$  at  $x_0 \in M$ , we have  $(D\kappa)_{x_0} \in \mathcal{H}_{\kappa}$  and  $Df(x_0) = \langle f, (D\kappa)_{x_0} \rangle_{\mathcal{H}_{\kappa}}$ . Here  $(D\kappa)_{x_0}$  represents the function obtained by letting D act on the first argument of  $\kappa$  and fix the first argument at  $x_0$ .

*Proof.* See [\[61,](#page-29-5) Lem.  $4.34$ ].

#### Bochner Integral

Given a probability space  $(\Omega, \mathcal{F}, P)$  and a separable Banach space  $\mathcal{B}$ , a  $\mathcal{B}$ -valued map  $X : \Omega \to \mathcal{B}$ is said to be *measurable* if  $X^{-1}(B)$  is measurable for all Borel sets  $B \subset \mathcal{B}$ . A (measurable) step function is a map in the form of  $X = \sum_{i=1}^{m} 1_{A_i} x_i$  for some  $x_1, \ldots, x_n \in \mathcal{B}$  and  $A_1, \ldots, A_n \in \mathcal{F}$ . For each measurable B-valued map X, there exists a sequence of measurable step functions  $X_n$  such that  $||X_n - X||_B \to 0$  pointwisely. A measurable B-valued map X is Bochner P-integrable if there exists a sequence of step functions  $X_n = \sum_{i=1}^{m_n} \mathbf{1}_{A_{i,n}} x_{i,n}$  such that  $\lim_{n\to\infty} \mathbb{E} ||X_n - X|| = 0$ , then the expectation of X is defined as  $\mathbb{E}X = \lim_{n\to\infty} \sum_{i=1}^{m_n} P(A_{i,n})x_{i,n}$ . A measurable B-valued map is P-integrable if and only if  $\mathbb{E}||X|| < +\infty$ . For a detailed exposition on Bochner integral, we refer the readers to  $[61, §A.5.4]$ . In addition, we have

**Theorem A.4.** Suppose X is a P-integrable B-valued map and f is a continuous linear functional on  $\mathcal{B}$ , then  $f(\mathbb{E}X) = \mathbb{E}f(X)$ .

# B Proofs of Theorems (Original to this Work)

### B.1 Proof of theorem 3.2

**Theorem** (Stein's identity). Suppose  $\sqrt{\kappa(x,x)}$   $D^l$ ,  $l = 1, ..., m$  and  $\sqrt{\kappa_p(x,x)}$  are all p-integrable, then  $\mathbb{E}_p[\mathcal{T}_p f] = 0$  for all  $f \in \mathcal{H}_\kappa$ , that is,  $\text{KSD}(p, p) = 0$  or equivalently  $p = q \Rightarrow \text{KSD}(p, q) = 0$ .

*Proof.* By Cauchy-Schwarz inequality, for  $f \in \mathcal{H}_\kappa^m$ , we have

$$
|\mathcal{T}_{p}f(x)| = |\langle f, (\vec{\mathcal{T}}_{p}\kappa)_{x}\rangle_{\mathcal{H}_{\kappa}^{m}}| \leq ||f||_{\mathcal{H}_{\kappa}^{m}} \cdot ||(\vec{\mathcal{T}}_{p}\kappa)_{x}||_{\mathcal{H}_{\kappa}^{m}} = ||f||_{\mathcal{H}_{\kappa}^{m}} \cdot \sqrt{\kappa_{p}(x,x)},
$$
  

$$
|\sum_{l=1}^{m} f_{l}(x)D_{x}^{l}| \leq \sum_{l=1}^{m} |D_{x}^{l}| |\langle f_{l}, \kappa_{x}\rangle_{\mathcal{H}_{\kappa}}| \leq \sum_{l=1}^{m} |D_{x}^{l}| ||f_{l}||_{\mathcal{H}_{\kappa}} ||\kappa_{x}||_{\mathcal{H}_{\kappa}}
$$
  

$$
\leq \sum_{l=1}^{m} |D_{x}^{l}| ||f_{l}||_{\mathcal{H}_{\kappa}} \sqrt{\kappa(x,x)} \leq ||f||_{\mathcal{H}_{\kappa}^{m}} \sqrt{\kappa(x,x)} \sum_{l=1}^{m} |D_{x}^{l}|.
$$

Therefore,  $|\sum_{l=1}^m f_l D^l|$  and  $\mathcal{T}_p f$  are p-integrable for all  $f \in \mathcal{H}_\kappa^m$ , since  $\sqrt{\kappa(x,x)} \sum_{l=1}^m |D^l|$  and  $\sqrt{\kappa_p(x,x)}$  are p-integrable. By Thm. 1.1,  $\mathbb{E}_p[\mathcal{T}_p f] = \int_M \text{div}(\sum_{l=1}^d p f_l D^l) d\mu = 0.$ 

### B.2 Proof of theorem 3.3

**Theorem.** Suppose  $\psi : M \to \mathbb{R}^{d'}$  is a smooth embedding for some  $d' > d$ , and suppose  $\kappa$  is  $C_0$ . universal on  $\mathbb{R}^{d'}$ , then the restriction of  $\kappa$  onto M, i.e.,  $\kappa|_M := \kappa(\psi(x), \psi(x'))$  for  $x, x' \in M$  is a qualified kernel on M.

*Proof.* Since  $\psi : M \to \mathbb{R}^{d'}$  is a smooth embedding, it is a diffeomorphism between M and  $\psi(M) \subset$  $\mathbb{R}^{d'}$  inheriting the subspace topology from  $\mathbb{R}^{d'}$ . Therefore, without loss of generality, we assume M is a subspace of  $\mathbb{R}^{d'}$ . Let  $\overline{M}$  be the closure of M in  $\mathbb{R}^{d'}$ . Let  $\mathcal{H}_M$  and  $\mathcal{H}_{\overline{M}}$  be the RKHS-s generated by  $\kappa_M$  and  $\kappa_{\overline{M}}$  on M and  $\overline{M}$  respectively.

It is proved in [\[8,](#page-25-6) Cor. 3] that if  $\kappa$  is  $C_0$ -universal on  $\mathbb{R}^{d'}$ , then its restriction  $\kappa|_{\overline{M}}$  onto the closed subset  $\overline{M}$  is a  $C_0$ -universal kernel on  $\overline{M}$ . Note that any function  $h \in C_0(M)$  can be extended to a function  $\bar{h} \in C_0(\overline{M})$  (set  $\bar{f} = 0$  on  $\partial M$ ). Since  $\kappa_{\overline{M}}$  is  $C_0$ -universal on  $\overline{M}$ , for any  $\epsilon > 0$ , there exists  $g \in \mathcal{H}_{\overline{M}} \subset C_0(\overline{M})$  such that  $\sup_{x \in \overline{M}} |g(x) - \overline{h}(x)| < \epsilon$ . Furthermore, by [\[8,](#page-25-6) Cor. 2],  $g|_M$  is an element in  $\mathcal{H}_M$ , thus  $\sup_{x \in M} |g_M(x) - h(x)| \leq \sup_{x \in \overline{M}} |g(x) - h(x)| < \epsilon$ , which proves the theorem by the arbitrariness of h. by the arbitrariness of  $h$ .

### B.3 Proof of theorem 3.4

Theorem. There always exists a collection of vector fields that satisfies the standing assumption 5 on M.

*Proof.* By Whitney embedding theorem [\[33,](#page-27-15) Thm 6.15], there exists a smooth embedding  $\psi : M \to$  $\mathbb{R}^{2d+1}$ . Let  $D_x^l$ ,  $1 \leq l \leq 2d+1$ , be the orthonormal projection of tangent vector  $\left(\frac{\partial}{\partial x^l}\right)_x$  onto the tangent space  $T_x\psi(M)$  of  $\psi(M)$ . Then  $d(\psi^{-1})(D^l)$ ,  $l=1,\ldots, 2d+1$ , is a group of vector fields on M such that they span the entire tangent space at each point.  $\blacksquare$ 

# B.4 Proof of theorem 3.5

 $\sum_{l=1}^{m} |D^l|$ ,  $\sum_{l=1}^{m} |D^l \log(p/q)|$ ,  $\sqrt{\kappa_p(x,x)}$  are q-integrable, then  $p = q \iff \text{KSD}(p,q) = 0$ . **Theorem.** Suppose  $\kappa$  is a qualified kernel, and q is a locally Lipschitz continuous density on M. If

*Proof.* Since  $\kappa$  is qualified, it is bounded by some constant  $C > 0$ . Therefore, the forward direction

is straightforward from Thm. 3.2. Furthermore, note that

$$
\sqrt{\kappa_q(x,x)} = \sqrt{\sum_{l=1}^m \|(\mathcal{T}_q^l \kappa)_x\|_{\mathcal{H}_\kappa^m}^2} = \sqrt{\sum_{l=1}^m \|(\mathcal{T}_p^l \kappa)_x - D^l \log(p/q) \cdot \kappa_x\|_{\mathcal{H}_\kappa^m}^2}
$$
  
\n
$$
\leq \sqrt{\sum_{l=1}^m \|(\mathcal{T}_p^l \kappa)_x\|_{\mathcal{H}_\kappa^m}^2} + \sqrt{\sum_{l=1}^m \|D^l \log(p/q) \cdot \kappa_x\|_{\mathcal{H}_\kappa^m}^2}
$$
  
\n
$$
= \sqrt{\kappa_p(x,x)} + \sqrt{\sum_{l=1}^m |D^l \log(p/q)|^2} \cdot \sqrt{\kappa(x,x)}
$$
  
\n
$$
\leq \sqrt{\kappa_p(x,x)} + \sqrt{C} \cdot \sum_{l=1}^m |D^l \log(p/q)|,
$$

which implies  $\sqrt{\kappa_q}(x, x)$  is q-integrable since  $\sqrt{\kappa_p(x, x)}$  and  $D^l \log(p/q)$  are q-integrable. The condition  $KSD(p, q) = 0$  further implies  $\mathbb{E}_q[\mathcal{T}_p f] = 0$  for all  $f \in \mathcal{H}_\kappa^m$ . Therefore,

$$
\sum_{l} \mathbb{E}_{q}[f_{l}D^{l} \log(p/q)] = \mathbb{E}_{q}[\mathcal{T}_{p}f - \mathcal{T}_{q}f] = \mathbb{E}_{q}[\mathcal{T}_{p}f] = 0, \text{ for all } f \in \mathcal{H}_{\kappa}^{m}.
$$

which implies  $\mathbb{E}_q[gD^l \log(p/q)] = 0$  for all  $g \in C_0(M)$ , as  $\kappa$  is qualified, which further implies that  $D^l \log(p/q) = 0$  for all l by the integrability of  $D^l \log(p/q)$ . As we assume M is connected,  $\log(p/q)$ is constant and thus  $p = q$ .

### B.5 Proof of Theorem 3.6

**Theorem.** Suppose Lie group G acts transitively on homogeneous space H. For a basis  $E^1, \ldots, E^m$ of  $T_eG$  (m = dim G), they correspond to a group of killing field  $K^1,\ldots,K^m$  in the way previously introduced. Then  $K^1, \ldots, K^m$  is a group of divergence-free vector fields on H that satisfies standing assumption 5.

*Proof.* It suffices to show that: for all  $x \in H$  and  $Y \in T_xH$ , there exists  $E \in T_eG$  such that its corresponding killing field K such that  $K_x = Y$ .

For each  $x \in H$ , let  $\mathcal{C}_x$  be the stabilizer of G at x, i.e.,  $\mathcal{C}_x := \{g \in G : g.x = x\}$ , which is a Lie subgroup of G. Note that  $T_eC_x$  is a subspace of  $T_eG$ , then we take a subspace  $\mathscr{C}_x$  of  $T_eG$  such that  $T_eC_x \oplus C_x = T_eG$  and take a basis  $D^j$ ,  $1 \leq j \leq \dim C_x$ , of  $C_x$ . Note that this basis corresponds to a group of vector field  $K^j$  on H, which are linearly independent at x. It is known that H is diffeomorphic to  $G/C_x$  and  $\dim H = \dim G - \dim C_x$ , thus  $\dim H = \dim C_x$ . Thus  $K_x^j$  span the entire  $T_xH$ .

# B.6 Proof of theorem 3.7

**Theorem.** Suppose M is a Riemannian manifold and  $\kappa_p^w(x, x)$  is w-integrable. Given M-valued samples  $x_i \sim w$ , we have

- 1.  $U_n^w(p), V_n^w(p) \xrightarrow{a.s.} \text{KSD}^2(p,q)$  with the rate of  $O_p(n^{-1}),$
- 2. If  $p \neq q$ , then

$$
\sqrt{n}[U_n^w - \text{KSD}^2(p, q)] \xrightarrow{d} N(0, \tilde{\sigma}^2), \quad \sqrt{n}[V_n^w - \text{KSD}^2(p, q)] \xrightarrow{d} N(0, \tilde{\sigma}^2),
$$

where  $\tilde{\sigma}^2 := 4var_{x' \sim w}[\mathbb{E}_{x \sim w} \kappa_p^w(x, x')]$ .

3. If  $p = q$ , then

$$
nU_n^w \xrightarrow{d_\cdot} \sum_{k=1}^\infty \lambda_k (Z_k^2 - 1), \quad nV_n^w \xrightarrow{d_\cdot} \sum_{k=1}^\infty \lambda_k Z_k^2,
$$

where  $\lambda_k$  are the eigenvalues of  $\kappa_p^w(x, y)$  as introduced in Thm. 2.4 and  $Z_k$  are i.i.d. standard Gaussian samples.

*Proof.* Note that  $\kappa_p^w(x, y) := \langle \frac{q(x)}{w(x)} \rangle$  $\frac{q(x)}{w(x)}(\vec{\mathcal{T}}_p\kappa)_x, \frac{q(y)}{w(y)}$  $\frac{q(y)}{w(y)}(\vec{\mathcal{T}}_{p}\kappa)_{y}\rangle_{\mathcal{H}_{\kappa}^{m}}$  is a semi-positive definite function, thus we have  $\kappa_p^w(x, y)^2 \leq \kappa_p^w(x, x) \kappa_p^w(y, y)$ , thus  $\mathbb{E}_q[\kappa_p^w(x, x)] < +\infty$  implies that  $\mathbb{E}_{q \times q}[\kappa_p^w(x, y)^2] < +\infty$ . Then the results follow directly from the classical asymptotic results of U-statistics and V-statistics  $[58, §5 \text{ and } §6]$  $[58, §5 \text{ and } §6]$ .

# B.7 Proof of theorem 4.1

**Theorem.** If  $\kappa_{(\cdot)}^w(\cdot,\cdot)$  is jointly continuous and  $\sup_{\theta \in K} \kappa_{\theta}^w(x,x)$  is w-integrable for any compact  $K \subset \Theta$ , then  $W_n(\theta) \to \text{KSD}^2(\theta)$  compactly almost surely, i.e., the following event

 $W_n(\theta) \to \text{KSD}^2(\theta)$  uniformly on any  $K \subset \Theta$ 

almost surely happens. As a corollary, if  $\Theta$  is locally compact,  $\text{KSD}(\theta)$  is continuous on  $\Theta$ .

We establish a lemma that will be needed for the proof.

<span id="page-35-0"></span>**Lemma B.1.** Suppose X is a M-valued r.v. following distribution P. Suppose  $f(\theta, x)$  is continuous on  $\Theta \times M$  and  $\sup_{\theta \in K} f(\theta, X)$  is integrable for all compact  $K \subset \Theta$ . Then

$$
n^{-1} \sum_{i=1}^{n} f(\theta, x_i) \to F(\theta) := \mathbb{E}[f(\theta, X)].
$$

compactly P-almost surely, and thus  $F(\theta)$  is continuous.

Proof. The lemma relies on a classical results  $[55, Thm. 1]$  regarding the uniform convergence of random functions. It suffices to check the condition of equi-continuity. Take a sequence of compact sets  $M_n$  such that  $M = \bigcup M_n$ , then the join continuity of  $f(\cdot, \cdot)$  will implies the joint uniform continuity of  $f(\theta, x)$  on  $K \times M_n$ , which further implies equi-continuity.

Proof. Note that  $|V_n^w(\theta) - \frac{n-1}{n}U_n^w(\theta)| = n^{-2}\sum_i \kappa_{\theta}^w(x_i, x_i) \to 0$  compactly almost surely, since  $n^{-1}\sum_{i} \kappa_{\theta}^{w}(x_{i},x_{i}) \to \mathbb{E}_{w}[\kappa_{\theta}^{w}(x,x)]$  compactly almost surely by the Lem. [B.1.](#page-35-0) Therefore, it suffices to show the result for  $U_n^w(\theta)$ . We apply a result [\[69,](#page-29-8) Thm. 1] regarding the uniform convergence of U-statistic. We will test the conditions one by one next.

First, we show  $\sup_{\theta \in K} |\kappa^w_{\theta}(x, y)|$  is  $\omega \times \omega$ -integrable. Note that  $\kappa^w_{\theta}(\cdot, \cdot)$  is positive definite, thus

$$
\sup_{\theta \in K} |\kappa^w_{\theta}(x, y)| \leq \sup_{\theta \in K} \kappa^w_{\theta}(x, x)^{1/2} \cdot \sup_{\theta \in K} \kappa^w_{\theta}(y, y)^{1/2},
$$

By Jensen's inequality, the w-integrability of  $\sup_{\theta \in K} \kappa^w_{\theta}(x,x)$  will imply the w-integrability of  $\sup_{\theta \in K} \kappa_\theta^w(x, x)^{1/2}$ . Therefore,  $\sup_{\theta \in K} |\kappa_\theta^w(x, y)|$  is  $w \times w$ -integrable.

Second, we show that for a sequence of compact sets  $M_n$  such that  $M = \bigcup M_n$ ,  $\kappa_{\theta}^{w}(x, y)$  and  $\mathbb{E}_{y\sim w}[\kappa_\theta^w(x,y)]$  are equi-continuous in  $\theta$  on  $K\times M_n\times M_n$  and  $K\times M_n$ . The equi-continuity of  $\kappa_{(\cdot)}^w(x,y)$  is straightforward from the uniform continuity of  $\kappa_{\theta}^w(x,y)$  on  $K \times M_n \times M_n$ . It suffices to show the equi-continuity of  $\mathbb{E}_{y \sim w}[\kappa_{(\cdot)}^w(\cdot, y)].$ 

For any compact  $K \subset \Theta$ , and any fixed  $\theta_0 \in K$ ,  $x_0 \in M$ , we choose a compact neighborhood  $W \subset M$  of  $x_0$ , and note that for any  $\theta \in K$ ,  $x \in W$  and any  $y \in M$ 

$$
\kappa_{\theta}^{w}(x, y) \leq \sup_{\theta \in K, x \in W} [\kappa_{\theta}^{w}(x, x)]^{1/2} \cdot \sup_{\theta \in K} [\kappa_{\theta}^{w}(y, y)]^{1/2}.
$$

Note that the right hand side is independent of  $\theta \in K$  and  $x \in W$ , and is integrable in y as the condition states. Therefore, regardless of any small perturbation on  $\theta$  and x, the function  $\kappa_{\theta}^{w}(x, \cdot)$ will be uniformly bounded by some integrable function of y that is independent of  $\theta$  and x. Applying dominating convergence theorem, we conclude that  $\mathbb{E}_{y \sim w}[\kappa_{(\cdot)}^w(\cdot,y)]$  restricted on  $K \times G$  is jointly continuous at  $(\theta_0, x_0)$ , and thus continuous on  $K \times G$  by the arbitrariness of  $\theta_0$  and  $x_0$ . It is thus uniformly continuous on  $K \times G_n$  and hence the condition of equi-continuity holds. Therefore,  $U_n^w(\theta) \to \text{KSD}^2(\theta)$  compactly almost surely.

The continuity of  $W_n$  is straightforward since it is a finite summation of continuous functions. Since  $KSD<sup>2</sup>$  is the uniform limit of  $W_n$  on any compact set,  $KSD<sup>2</sup>$  is continuous on any compact set. Therefore,  $KSD^2$  is continuous as continuity is a local property. This completes the proof.

# B.8 Proof of theorem 4.2

Theorem. Suppose all the conditions in Thm. 4.1 hold and suppose Θ satisfies one of following three conditions:

- 1.  $\Theta$  is compact;
- 2. Θ is a geodesic convex subset of a Riemannian manifold,  $W_n$  is convex on  $\Theta$ ,  $\Theta_0$  is non-empty, compact and  $\Theta_0 \subset \Theta_2$  (interior);
- 3.  $\Theta = \Theta_1 \times \Theta_2$ , where  $\Theta_1$  is compact, and for each fixed  $\theta_1 \in \Theta_1$ ,  $\{\theta_1\} \times \Theta_2$  and  $W_n(\theta_1, \cdot)$  satisfy the second condition.

Then  $\Theta_0$ ,  $\Theta_n$  are non-empty for large n and  $\sup_{\theta \in \widehat{\Theta}_n} \rho(\theta, \Theta_0) \to 0$  almost surely.

We establish some lemmas that will be needed for the proof.

<span id="page-36-0"></span>**Lemma B.2.** Suppose f is convex on  $\mathbb{R}^d$  and  $C \subset \mathbb{R}^d$  is an open set. If there exists  $x_0 \in C$  such that  $f(x_0) < f(x)$  for all  $x \in \partial C$ , then  $\inf_{x \notin C} f(x) = \inf_{x \in \partial C} f$  and the set of global minimizers of f is non-empty and contained in C.

*Proof.* For all  $x' \notin C$ , there exists  $0 \le \beta < 1$  such that  $\beta x_0 + (1 - \beta)x' \in \partial C$ , and thus

$$
\beta f(x_0) + (1 - \beta)f(x') \ge f(\beta x_0 + (1 - \beta)x') > f(x_0) \Longrightarrow f(x') > f(x_0).
$$

Furthermore,

$$
f(x') \ge \beta f(x_0) + (1 - \beta)f(x') \ge f(\beta x_0 + (1 - \beta)x') \ge \inf_{x \in \partial C} f(x).
$$

This concludes the proof.

Note that  $\Theta$  is locally compact, thus  $W_n$  and  $\text{KSD}^2$  are continuous on  $\Theta$  by Thm. 4.3. Furthermore, as the pointwise limit preserves the convexity,  $KSD^2(\theta_1, \cdot)$  is also convex in  $\theta_2$  for all  $\theta_1 \in \Theta_1$ . Let  $m^*(\theta_1) := \inf_{\theta_2 \in \Theta_2} \text{KSD}^2(\theta_1, \theta_2)$ , and denote by  $\tilde{\Theta}_0(\theta_1)$  the set of minimizers of  $\text{KSD}^2(\theta_1, \cdot)$ . Then we establish the next lemma.

<span id="page-37-0"></span>**Lemma B.3.** For each  $\theta_1 \in \Theta_1$ , there exists a neighborhood  $V_1$  of  $\theta_1$  and a bounded open set  $V_2 \subset \Theta_2$ such that for any  $\theta_1 \in V_1$ ,  $\tilde{\Theta}_0(\theta_1) \subset V_2$ .

*Proof.* Take a bounded  $V_2$  such that  $\tilde{\Theta}_0(\bar{\theta}_1) \subset V_2$ , then since  $\text{KSD}^2(\bar{\theta}_1, \cdot) - m^*(\bar{\theta}_1)$  is continuous and positive on  $\partial V_2$ , let  $\epsilon := \inf_{\theta_2 \in \partial V_2} \text{KSD}^2(\bar{\theta}_1, \theta_2) - m^*(\bar{\theta}_1) > 0$ . Note that  $\{\bar{\theta}_1\} \times \partial V_2$  is a subset of the open set  $\{KSD^2(\cdot,\cdot) > \epsilon/2 + m^*(\bar{\theta}_1)\}\)$ , thus we apply tube lemma [\[47,](#page-28-17) Lem. 26.8] and get an open neighborhood  $V_1$  of  $\bar{\theta}_1$  such that  $V_1' \times \partial V_2 \subset {\text{KSD}^2(\cdot, \cdot)} > \epsilon/2 + m^*(\bar{\theta}_1)$ , which implies that  $\inf_{V_1' \times \partial V_2} \text{KSD}^2 > m^*(\bar{\theta}_1) + \epsilon/2$ . However, we choose a  $\tilde{\theta}_2 \in \tilde{\Theta}_1(\bar{\theta}_1) \subset V_2$ , there exists a neighborhood  $V''_1$  of  $\bar{\theta}_1$  such that  $\sup_{\theta_1 \in V''_1} \text{KSD}^2(\theta_1, \tilde{\theta}_2) < m^* + \epsilon/2$ , as  $\text{KSD}^2$  is continuous at  $(\bar{\theta}_1, \tilde{\theta}_2)$ . Therefore, for any  $\theta_1 \in V_1 := V_1' \cap V''$ , we have  $\text{KSD}^2(\theta_1, \tilde{\theta}_2) < \inf_{\theta_2 \in \partial V_2} \text{KSD}^2(\theta_1, \theta)$ . We now apply Lem. [B.2,](#page-36-0) resulting in the minimizer set  $\tilde{\Theta}_0(\theta_1)$  is contained in  $V_2$ .

Now we are ready to prove the theorem.

*Proof.* 1. We show that  $\Theta_0$  is non-empty.

Let  $m_0^* := \inf_{\theta_1 \in \Theta_1} m^*(\theta_1) = \inf_{\theta \in \Theta} \text{KSD}^2(\theta)$ . Take a sequence  $\theta_{1,n} \in \Theta_1$  such that  $m^*(\theta_{1,n}) \downarrow$  $m_0^*$ . As  $\Theta_1$  is compact,  $\theta_{1,n}$  has a convergent sub-sequence, still denoted as  $\theta_{1,n}$ , whose limit point is denoted as  $\theta_{1,0}$ . We apply Lem. [B.3](#page-37-0) to  $\tilde{\Theta}_0(\theta_{1,0})$ , thus there exists a compact neighborhood  $V_1 \times V_2$ of  ${\{\theta_{1,0}\}\times\tilde{\Theta}_0(\theta_{1,0})}$  such that for all  $\theta_1 \in V_1$ ,  $\tilde{\Theta}_0(\theta_1) \subset V_2$ . Since KSD<sup>2</sup> is continuous on the compact set  $\{\theta_{1,0}\}\times \tilde{\Theta}_0(\theta_{1,0})$  and the size of  $V_1\times V_2$  can be taken as small as needed, we can shrink  $V_1\times V_2$ such that  $\inf_{\theta \in V_1 \times V_2}$  KSD<sup>2</sup>( $\theta$ ) >  $m^*(\theta_{1,n}) - \epsilon$ . However, note that  $\theta_{1,n}$  will finally enter  $V_1$ , and from then on,  $\{\theta_{1,n}\}\times\tilde{\Theta}_0(\theta_{1,n})\subset V_2$ , which implies  $\inf_{V_1\times V_2}KSD^2(\theta)\leq m^*(\theta_{1,n})\downarrow m_0^*$ . Therefore,  $m^*(\theta_{1,n}) - \epsilon \le \inf_{V_1 \times V_2} \text{KSD}^2(\theta) \le m^*(\theta_{1,n}) + m_0^*$ , and thus  $m^*(\theta_{1,0}) = m_0^*$  as  $\epsilon$  is arbitrary. Therefore,  $\Theta_0$  is non-empty.

2. We show that  $\Theta_0$  is compact.

Suppose  ${B_\alpha} \subset \Theta$  is a collection of open balls such that  $\Theta_0 \subset \bigcup B_\alpha$ . For each  $\theta_1$ , the  $\theta_1$ -section  $\Theta_0^{\theta_1} := \{(\theta, \theta') \in \Theta_0 | \theta = \theta_1\}$  of  $\Theta_0$ , if non-empty, is exactly the set  $\tilde{\Theta}_0(\theta_1)$ . Since  $\Theta_0^{\theta_1} = \tilde{\Theta}_0(\theta_1)$  is compact, there exists a finite subcollection of balls  $B_i^{\theta_1}$ ,  $1 \leq i \leq n_{\theta_1}$  such that  $\{\theta_1\} \times \tilde{\Theta}_0(\theta_1) \subset \bigcup_i B_i^{\theta_1}$ . We apply Lem. [B.3](#page-37-0) to each  $\theta_1$ , and conclude there exists open neighborhoods  $V_1^{\theta_1}$  and  $V_2^{\theta_1}$  such that  $\{\theta_1\} \times \tilde{\Theta}_0(\theta_1) \subset V_1^{\theta_1} \times V_2^{\theta_1} \subset \bigcup_i B_i^{\theta_1}$ . If the section  $\Theta_0^{\tilde{\theta}_1}$  is empty, we prescribe  $V_1^{\theta_1}$  and  $V_2^{\theta_2}$  with arbitrary neighborhood such that  $\{\theta_1\} \times \tilde{\Theta}_0(\theta_1) \subset V_1^{\theta_1} \times$ cover of  $\Theta_1$ , thus there exists an open cover  $V_1^{\theta_{1,i}}$  for  $\theta_{1,i}$ ,  $1 \leq i \leq n$ . Note that  $B_j^{\theta_{1,i}}$ ,  $1 \leq i \leq n$ ,  $1 \leq j \leq n_{\theta_{1,i}}$  is a finite subcover of  $\Theta_0$ . Therefore,  $\Theta_0$  is compact.

3. We show that the minimizers  $\hat{\theta}_n \in \widehat{\Theta}_n$  will eventually enter a compact set uniformly.

Since  $\Theta_0$  is compact, we choose an open set  $K \subset \Theta_2$  with compact closure such that  $\Theta_0 \subset \Theta_1 \times K$ . Since KSD<sup>2</sup>  $-m_0^*$  is continuous and positive on the compact set  $\Theta_1 \times \partial K$ , let  $\epsilon := \inf_{\Theta_1 \times \partial K}$  KSD<sup>2</sup>( $\theta$ )  $m_0^* > 0$ . By Thm. 4.2,  $W_n \to \text{KSD}^2$  uniformly on  $\Theta_1 \times \bar{K}$ , which implies for large enough n,  $W_n > m_0^* + \epsilon/2$  uniformly on  $\Theta_1 \times \partial K$  and  $W_n < m_0^* + \epsilon/2$  uniformly on  $\Theta_0$ . Therefore, we can apply Lem. [B.2](#page-36-0) to each  $W_n(\theta_1, \cdot)$ , and conclude that  $\widehat{\Theta}_n \subset \Theta_1 \times K$  for large enough n.

4. We show the theorem.

Since the global minimizers of  $W_n$  will eventually enter the compact set  $\Theta_1 \times \overline{K}$ . We assume that  $\Theta_n$  is set of minimizers of  $W_n$  over  $\Theta_1 \times \overline{K}$  WLOG.

For each  $\delta > 0$ , let  $\Theta_{\delta} := \{ \theta \in \Theta_1 \times \overline{K} : d(\theta, \Theta_0) < \delta \}$  be the closed  $\delta$ -neighborhood of  $\Theta_0$  in  $\Theta_1 \times \bar{K}$ , and let  $\Theta_{\delta}^c := \Theta_1 \times \bar{K} - \Theta_{\delta}$ . Note that  $\Theta_{\delta}^c$  is compact, and  $KSD^2 - m^*0$  is positive on  $\Theta_{\delta}^c$ , thus we let  $\epsilon_{\delta} := \inf_{\Theta_{\delta}^c} \text{KSD}^2(\theta) - m_0^* > 0$ . For n large enough such that  $\widehat{\Theta}_n \subset \Theta_1 \times \overline{K}$  and  $\sup |W_n - \text{KSD}^2| < \epsilon_\delta/2$ , thus for each  $\hat{\theta}_n \in \widehat{\Theta}_n$ ,

$$
\textup{KSD}^2(\hat{\theta}_n)-m_0^*
$$

which implies  $\theta_n \in \Theta_\delta$ , thus proving the theorem.

### B.9 Proof of theorem 4.3

**Theorem.** Under assumption  $A1 \sim A5$ ,  $\sqrt{n} \text{Log}_{\theta_0}(\hat{\theta}_n) \stackrel{d}{\rightarrow} \mathcal{N}(0,\Gamma^{-1}\Sigma\Gamma^{-1})$ . In more practical situations, we have  $\sqrt{n}[\mathcal{I}(\hat{\theta}_n)]^{1/2} \text{Log}_{\hat{\theta}_n}(\theta_0) \stackrel{d}{\rightarrow} \mathcal{N}(0, I).$ 

Proof. Throughout history, the asymptotic normality of the minimizer of a loss function is always established via a similar strategy to the classical results regarding the asymptotic normality of MLE, e.g, the asymptotic normality of MKSDE established in [\[2,](#page-25-4) [48\]](#page-28-9).

In this work, since the parameter space may not be a vector space, we can not directly adopt their proof. However, as the MKSDE  $\hat{\theta}_n$  converges to the ground truth  $\theta_0$ , the sequence will finally enter a neighborhood of  $\theta_0$ . Since a manifold is locally Euclidean, we may adopt their proof in a local chart of the manifold. It is convenient to take the normal coordinates given by the exponential map  $Exp_{\theta_0}$  at  $\theta_0$ , as the Jacobian of the  $Exp_{\theta_0}$  is I at 0 and the geodesic connecting  $\hat{\theta}_n$  with  $\theta_0$ coincides with the line segments connecting  $Log_{\theta_0}(\theta)$  with 0 on the tangent space at  $\theta_0$ .

The rest of the proof is the same as the one given in  $[2, 48]$  $[2, 48]$ .

# B.10 Proof of theorem 4.4

**Theorem.** Under the conditions in Thm. 4.2 hold, then  $\sum_{l=1}^{\infty} (\hat{\lambda}_k - \lambda_k) Z_k^2 \to 0$  in probability.

Let  $\hat{\lambda}'_k$  be the eigenvalues of the Gram matrix  $G_n^w(\theta_0)$  (set  $\hat{\lambda}'_k := 0$  for  $k > n$ ). It is shown in [\[27\]](#page-27-10) that  $\sum_{k=1}^{\infty} (\hat{\lambda}'_k - \lambda_k) Z_k^2 \to 0$  in probability. Therefore, it suffices to show that  $\sum_{k=1}^{n} (\hat{\lambda}'_k - \hat{\lambda}_k) Z_k^2 \to 0$ in probability. The Weilandt-Hoffman inequality  $[63, Eq. (1.64)]$  states that

$$
\sum_{l=1}^{n} |\hat{\lambda}'_l - \hat{\lambda}_l| \leq ||G_n^w(\theta_0) - G_n^w(\hat{\theta}_n)||_1,
$$

where  $||A||_1 := \text{tr} \sqrt{A^{\top}A}$  represents the nuclear norm of a squared matrix A. We will show that  $||G_n^w(\theta_0) - G_n^w(\hat{\theta}_n)||_1 \to 0$  almost surely, which will imply that  $\sum_{k=1}^n (\hat{\lambda}'_k - \hat{\lambda}_k) Z_k^2$  converges to 0 in probability. The proof relies on following lemma:

<span id="page-38-0"></span>**Lemma B.4.** Suppose H is a Hilbert space and  $x_i, y_i \in H$ ,  $1 \leq i \leq n$  and  $\Sigma := (\langle x_i, y_j \rangle)_{ij}$ . Then  $||\Sigma||_1 \le ||\vec{x}|| \cdot ||\vec{y}||$ . Here  $||\vec{x}|| := \sqrt{\sum_{i=1}^n ||x_i||^2}$  and  $||\vec{y}||$  likewise. Consequently,

$$
\|(\langle x_i, x_j \rangle)_{ij} - (\langle y_i, y_j \rangle)_{ij}\|_1 \leq \|\vec{x} - \vec{y}\| \cdot (\|\vec{x}\| + \|\vec{y}\|)
$$

*Proof.* Let  $\mathcal{H}_0 := \text{span}\{x_1, \ldots, x_n, y_1, \ldots, y_n\}$  and  $m := \dim \mathcal{H}_0$ . Take an orthogonal basis  $\{e\}_{i=1}^m$  of  $\mathcal{H}_0$  and let  $x_i = \sum_{j=1}^m a_i^j e_j$  and  $y_i = \sum_{j=1}^m b_i^j e_j$  for  $1 \leq i \leq n$ , and let  $A := (a_i^j)_{ij}$ ,  $B := (b_i^j)_{ij}$ . Then  $\Sigma = AB^{\top}$ . Let  $\Sigma = PSQ^{\top}$  be the singular value decomposition of  $\Sigma$ , then apply the Cauchy-Schwartz inequality

$$
\|\Sigma\|_1 = \text{tr}(S) = \text{tr}(P^\top A B^\top Q) \le \|P^\top A\|_F \|B^\top Q\|_F = \|A\|_F \|B\|_F.
$$

The first part of the lemma follows from the fact that  $||A||_F = ||x||^2$  and  $||B||_F = ||y||^2$ . For the second part, just note that

$$
\begin{aligned} ||(\langle x_i, x_j \rangle)_{ij} - (\langle y_i, y_j \rangle)_{ij}||_1 &\leq ||(\langle x_i, x_j - y_j \rangle)_{ij}||_1 + ||(\langle x_i - y_i, y_j \rangle)_{ij}||_1 \\ &\leq ||\vec{x} - \vec{y}||(||\vec{x}|| + ||\vec{y}||). \end{aligned}
$$

■

Next we prove the theorem.

*Proof.* Recall the construction in §3.1, the vector  $\vec{\mathcal{A}}_p \kappa_x := (\mathcal{A}_p^1 \kappa_x, \dots, \mathcal{A}_p^d \kappa_x)^\top$  is an element in  $\mathcal{H}_{\kappa}^d$ , and

$$
\kappa_p^w(x_i, x_j) = \left\langle \frac{p(x_i)}{\omega(x_i)} \vec{\mathcal{A}}_p \kappa_{x_i}, \frac{p(x_j)}{\omega(x_j)} \vec{\mathcal{A}}_p \kappa_{x_j} \right\rangle_{\mathcal{H}_k^d}.
$$

For notational simplicity, let  $\phi_{\theta}(x) := \frac{p(x)}{\omega(x)} \vec{\mathcal{A}}_{p_{\theta}} \kappa_x \in \mathcal{H}_{\kappa}^d$ , then note that

$$
G_n^w(\theta_0) = n^{-1}(\langle \phi_{\theta_0}(x_i), \phi_{\theta_0}(x_j) \rangle)_{ij}, \quad G_n^w(\hat{\theta}_n) = n^{-1}(\langle \phi_{\hat{\theta}_n}(x_i), \phi_{\hat{\theta}_n}(x_j) \rangle)_{ij}.
$$

Apply the Lem. [B.4](#page-38-0) to get

$$
n||G_n^w(\theta_0) - G_n^w(\hat{\theta}_n)||_1 \le
$$
  

$$
\left(\sqrt{\sum_{i=1}^n ||\phi_{\hat{\theta}_n}(x_i)||^2} + \sqrt{\sum_{i=1}^n ||\phi_{\theta_0}(x_i)||^2}\right) \cdot \sqrt{\sum_{i=1}^n ||\phi_{\hat{\theta}_n}(x_i) - \phi_{\theta_0}(x_i)||^2}
$$

Let  $f(\theta, x) := ||\phi_{\theta}(x) - \phi_{\theta_0}(x)||^2$ , then we can rewrite above inequality as

$$
||G_n^w(\theta_0) - G_n^w(\hat{\theta}_n)||_1
$$
  
\n
$$
\leq \sqrt{n^{-1} \sum_{i=1}^n f(\hat{\theta}_n, x_i)} \left( \sqrt{n^{-1} \sum_{i=1}^n ||\phi_{\hat{\theta}_n}(x_i)||^2} + \sqrt{n^{-1} \sum_{i=1}^n ||\phi_{\theta_0}(x_i)||^2} \right)
$$
  
\n
$$
\leq \sqrt{n^{-1} \sum_{i=1}^n f(\hat{\theta}_n, x_i)} \left( \sqrt{n^{-1} \sum_{i=1}^n f(\hat{\theta}_n, x_i)} + 2 \sqrt{n^{-1} \sum_{i=1}^n ||\phi_{\theta_0}(x_i)||^2} \right).
$$

Clearly, by law of large numbers,

$$
n^{-1} \sum_{i=1}^{n} \|\phi_{\theta_0}(x_i)\|^2 = n^{-1} \sum_{i=1}^{n} \kappa_{\theta_0}^w(x_i, x_i) \xrightarrow{\omega - a.s} \int_G \kappa_{\theta_0}^w(x, x) w_{\theta_0}(x) \mu(dx) < +\infty,
$$

thus  $n^{-1} \sum_{i=1}^{n} ||\phi_{\theta_0}(x_i)||^2 = O_w(1)$ .

To show  $\|\tilde{G}_n^w(\theta_0) - G_n^w(\hat{\theta}_n)\| \to 0$   $\omega$ -a.s., it suffices to show  $n^{-1} \sum_{i=1}^n f(\hat{\theta}_n, x_i) \to 0$   $\omega$ -a.s.. Note that

$$
f(\theta, x) \le 2\|\phi_{\theta}(x)\|^2 + 2\|\phi_{\theta_0}(x)\|^2 = 2\kappa_{\theta}^{w}(x, x) + 2\kappa_{\theta_0}^{w}(x, x),
$$

thus  $\sup_{\theta \in K} f(\theta, x)$  is  $\omega$ -integrable for any compact set  $K \subset \Theta$ , as the conditions in Thm. 4.2 state. Therefore, we can apply Lem. [B.1](#page-35-0) to conclude that

$$
n^{-1} \sum_{i=1}^{n} f(\theta, x_i) \to F(\theta) := \int_{G} f(\theta, x) \omega(dx),
$$

compactly  $\omega$ -a.s. and  $F(\theta)$  is continuous. Note that  $f(\theta_0, x) = ||\phi_{\theta_0}(x) - \phi_{\theta_0}(x)||^2 = 0$ , thus  $F(\theta_0) = 0$ . Note that  $\hat{\theta}_n \to \theta_0$   $\omega$ -a.s., thus  $n^{-1} \sum_{i=1}^n f(\hat{\theta}_n, x_i) \to 0$   $\omega$ -a.s..

#### B.11 Proof of theorem 5.1

**Theorem.**  $Q_u$ ,  $Q_v$  defined in (19) are both symmetric, and  $Q_v$  is positive semi-definite. If further  $Q_u$  is positive semi-definite, then the global minimizer sets of  $U_n^w(\theta)$  and  $V_n^w(\theta)$  in (18) can be represented by the Moore-Penrose inverse  $Q_u^+$ ,  $Q_v^+$  of  $Q_u$ ,  $Q_v$  as follows

$$
\operatorname{argmin} U_n^w(\theta) = \{-Q_u^+ b_u - (I - Q_u^+ Q_u)x : x \in \mathbb{R}^s\},
$$
  

$$
\operatorname{argmin} V_n^w(\theta) = \{-Q_v^+ b_v - (I - Q_v^+ Q_v)x : x \in \mathbb{R}^s\}.
$$

For the unweighted case, we just ignore the weighted ratio  $\frac{q(x_i)q(x_j)}{w(x_i)w(x_j)}$  in (19).

Proof. Note that

$$
Q(x,y)^{\top} = \kappa(x,y) \sum_{l=1}^{m} (K_x^l \zeta \cdot K_y^l \zeta^{\top})^{\top} = \kappa(y,x) \sum_{l=1}^{m} K_y^l \zeta \cdot K_x^l \zeta^{\top} = Q(y,x),
$$

thus

$$
Q_u^{\top} = \sum_{i \neq j} \frac{Q(x_i, x_j)^{\top}}{n(n-1)} \frac{q(x_i)q(x_j)}{w(x_i)w(x_j)} = \sum_{j \neq i} \frac{Q(x_j, x_i)}{n(n-1)} \frac{q(x_j)q(x_i)}{w(x_j)w(x_i)} = Q_u,
$$

and  $Q_v$  is also symmetric by a parallel argument. Next we show  $Q_v$  is positive semi-definite. Given arbitrary fixed vector  $\theta_0 \in \mathbb{R}^s$ , we let  $\phi_x^l = \theta_0^\top \cdot K_x^l \zeta$ , which is a real-valued function on H. We let  $\mathbf{1}_n := (1, \ldots, 1)^\top \in \mathbb{R}^n$ ,  $\vec{\phi}^l := (\phi^l_{x_1}, \ldots, \phi^l_{x_n})^\top \in \mathbb{R}^n$  and further set

$$
\Phi = \sum_{l=1}^m \vec{\phi}^l \vec{\phi}^l \vec{\phi}^{l\top} \in \mathbb{R}^{n \times n}, \quad G_n := \left( \kappa(x_i, x_j) \frac{q(x_i)q(x_j)}{w(x_i)w(x_j)} \right)_{ij} \in \mathbb{R}^{n \times n}.
$$

Note that  $\Phi$  and  $G_n$  are both positive semi-definite, thus by Schur product theorem, their Hadamard product (elementwise product)  $G_n \circ \Phi$  is also positive semi-definite. Furthermore, since  $\theta^{\top} Q(x, y) \theta =$  $\kappa(x, y) \sum_{l=1}^{m} \phi_x^l \cdot \phi_y^l$ , we have

$$
\theta^{\top} Q_v \theta = n^{-2} \mathbf{1}_n^{\top} (G_n \circ \Phi) \mathbf{1}_n \ge 0,
$$

which implies  $Q_n$  is positive semi-definite by the arbitrariness of  $\theta$ . The rest of the theorem is straightforward.