Fiducial inference for partially identified parameters with applications to instrumental variable models

Yifan Cui^{*} Jan Hannig[†]

Abstract

In the past two decades, there has been a fast-growing literature on fiducial inference since it was first proposed by R. A. Fisher in the 1930s. However, most of the fiducial inference based methods and related approaches have been developed for point-identified models, i.e., statistical models where the parameters of interest are uniquely determined by the observed data and the model's assumptions. In this paper, we propose a novel fiducial approach for partially identified statistical models. As a leading example, we consider the instrumental variable model with a variety of causal assumptions and estimands. The proposed methods are illustrated through extensive simulations and a data analysis evaluating the effect of consuming Vitamin A supplementation on reducing mortality rates.

keywords: Causal Inference, Fiducial Inference, Instrumental Variable, Partial Identification, Uncertainty Quantification

^{*}Center for Data Science, Zhejiang University

[†]Department of Statistics and Operations Research, UNC-Chapel Hill

1 Introduction

Fiducial inference has its origins in a series of papers by R. A. Fisher (Fisher, 1922, 1930, 1933). The essential idea of the fiducial statement is switching the role of data and parameters to introduce meaningful distribution on the parameter space that summarizes knowledge about the unknown parameter without introducing any prior. In the past two decades, the fiducial inference, including its variants such as generalized fiducial inference (Hannig et al., 2016), extended fiducial inference (Liang et al., 2024), has been applied to a variety of subjects such as psychology (Liu and Hannig, 2017; Liu et al., 2019; Neupert et al., 2021), forensic science (Hannig et al., 2019; Hannig and Iyer, 2022), social network (Su et al., 2022), and reliability modeling (Chen et al., 2016; Cui and Hannig, 2019; Cui et al., 2023).

Fiducial inference has connections to several related approaches, including Dempster-Shafer theory (Dempster, 1968; Shafer, 1976), confidence distributions (Singh et al., 2005; Xie and Singh, 2013; Hjort and Schweder, 2018), inferential models (Martin and Liu, 2013, 2015; Liu and Martin, 2024), objective Bayesian inference (Berger et al., 2009, 2012) and the repro method (Xie and Wang, 2022, 2024; Hou et al., 2024). Many additional references are available for the interested reader in Xie and Singh (2013), Schweder and Hjort (2016), Cui and Xie (2023), and Murph et al. (2024).

While tremendous success has been seen in the area of the foundation of statistics, there are few papers on partially identified parameters. Partially identified models refer to statistical models where the parameters of interest are not uniquely determined by the available data and the models' assumptions. Instead of pinpointing a single value, these models yield a set of possible values (or bounds) for the parameters; for example, this arises when the data or assumptions are insufficient to fully identify the parameters, but they do constrain the range of feasible values.

Throughout, we consider instrumental variable methods in causal inference as a running example. Causal inference (Neyman, 1923; Rubin, 2005; Hernán and Robins,

2020; Ding, 2024; Wager, 2024) is critically important because it allows us to draw valid conclusions about causal relationships between variables, which is fundamental for decision-making. In observational studies and randomized experiments subject to non-compliance, unmeasured confounders might occur when there are variables that influence both the treatment and the outcome but are not included in the analysis. These unmeasured confounders can introduce bias and undermine the validity of causal inferences. Over the years, a prevailing strand of proposals has been proposed to mitigate unmeasured confounding bias, including instrumental variable (Imbens and Angrist, 1994; Angrist et al., 1996; Wang and Tchetgen Tchetgen, 2018), difference-in-differences (Abadie, 2005; Arkhangelsky et al., 2021; Ye et al., 2023), sensitivity analysis (Cornfield et al., 1959; Rosenbaum, 2002; VanderWeele and Ding, 2017; Zhao et al., 2019; Bonvini and Kennedy, 2022), negative controls (Miao et al., 2018; Shi et al., 2020), proximal causal inference (Tchetgen Tchetgen et al., 2024), and many other approaches.

An instrumental variable (IV) model (Imbens and Angrist, 1994; Angrist et al., 1996) is a powerful causal inference tool used to estimate causal relationships when there is potential endogeneity in the treatment variable. An IV is defined as a pretreatment variable that is independent of all unmeasured confounders, and does not have a direct causal effect on the outcome other than through the treatment. In a double-blind randomized trial, random assignment is a leading example of an ideal IV when patients fail to comply with their assigned treatment. The use of instrumental variables (IVs) is a celebrated approach to quantifying causal effects in observational studies or randomized trials with non-compliance. However, it might not be possible to uniquely identify the causal effect of a treatment using a valid IV.

A partially identified model (Balke and Pearl, 1994; Imbens and Manski, 2004; Chernozhukov et al., 2007, 2013; Richardson et al., 2015; Kaido et al., 2019) with instrumental variables (Robins, 1989; Manski, 1990; Balke and Pearl, 1997; Richardson and Robins, 2010;

Swanson et al., 2015, 2018) is used in statistics, econometrics, epidemiology, and social science to estimate causal effects when the relationship between treatment and outcome is not fully identifiable due to data limitations or violations of strong assumptions. In this paper, we propose a novel fiducial approach for partially identified statistical models. Specifically, we propose a novel acceptance sampler to sample from the derived fiducial distribution of the bounds. Upon obtaining fiducial samples, we use the samples to construct statistical inference, e.g., using the median of the samples as a point estimator, and using appropriate quantiles to construct confidence intervals for the lower and upper bounds, respectively.

The paper makes a number of significant contributions to both the fiducial and causal inference literature. First, we propose fiducial based acceptance sampling algorithm to quantify uncertainty of bounds for a variety of causal estimands under various assumptions by leveraging a binary IV. Second, we establish a novel Bernstein–von Mises theorem that verifies the frequentist validity of the proposed fiducial confidence intervals. As a consequence of the Bernstein–von Mises theorem, the proposed confidence intervals provide asymptotically correct coverage for the lower and upper bounds. Third, as a by-product, the acceptance rate of the proposed sampling algorithm is a natural estimator of fiducial probability of the observed data agreeing with the IV assumptions. Therefore, a high acceptance rate near 0 suggests that the IV assumptions are likely violated.

The remainder of the article is organized as follows. In Section 2, we briefly review instrumental variable models with a binary treatment, instrument, and outcome. In Section 3, we develop fiducial inference for partially identified causal estimand in instrumental variable models. In Section 4, we establish a novel Bernstein-von Mises theorem for our framework. Simulation studies are presented in Section 5. Section 6 describes a real data application on evaluating the effect of consuming Vitamin A supplementation on reducing mortality rates. Section 7 and Section 8 discuss a variety of causal estimands and assumptions under our framework. The article concludes with a discussion of future work in Section 9. Proofs are provided in the Appendix.

2 Preliminaries

We first introduce some notation. Let Y denote the outcome of interest and $A \in \{0, 1\}$ be a binary treatment indicator. There might be unmeasured confounders of the effect of A on Y which are not known apriori. Suppose also that one has observed a binary IV $Z \in \{0, 1\}$. Throughout we assume the observed data are independent and identically distributed realizations of (Y, A, Z). To simplify the presentation, we ignore covariates which can easily be incorporated into our framework.

Let A_z be a person's potential treatment under an intervention that sets IV to value z. Let $Y_{z,a}$ denote the potential outcome had, possibly contrary to fact, a person's IV and treatment value been set to z and a, respectively. Suppose that the following consistency assumption holds: $Y = Y_{Z,A}$ and $A = A_Z$.

Furthermore, we assume the following set of IV assumptions (Balke and Pearl, 1997; Swanson et al., 2018):

Assumption 1. (Exclusion restriction) $Y_{z,a} = Y_a$.

Assumption 2. (*IV independence*) $Z \perp \{Y_{a=0}, Y_{a=1}, A_{z=0}, A_{z=1}\}$.

Assumption 3. (IV positivity) 0 < P(Z = 1) < 1.

Assumption 1 assumes an individual-level exclusion restriction, i.e., there can be no individual direct causal effect of Z on Y not mediated by A. Assumption 2 ensures that the causal effect of Z on A and Y is unconfounded. Assumption 3 essentially states that a person has an opportunity to receive both IVs (Greenland, 2000; Hernán and Robins, 2006). Figure 1 gives an illustration of a valid IV.



Figure 1: A directed acyclic graph (DAG) for an IV, where U refers to unmeasured confounder.

Note that because of the independence of the complete data, the observed data can be summarized using two independent multinomials: $\mathbf{n}_z = (n_{z00}, n_{z01}, n_{z10}, n_{z11}) \sim$ Multinomial (n_z, \mathbf{q}_z) , where $z \in \{0, 1\}$, $\mathbf{q}_z = (q_{z00}, q_{z01}, q_{z10}, q_{z11}) \in \Delta^3$, the probability $q_{zay} = P(A = a, Y = y | Z = z) \ a, y \in \{0, 1\}$, and $\Delta^{k-1} = \{\mathbf{p} : \sum_{i=1}^k p_i = 1, p_i \ge 0, i = 1, \dots, k\}$ is a k - 1-dimensional probability simplex. Notice that the counts n_{zay} and conditional probabilities q_{zay} are indexed depending on outcomes they are associated with. In particular, the indexes z, a, y mean the following: z = 0, 1 correspond to the observed value of Z, a = 0, 1 matches the observed value of A, and y = 0, 1 is the observed value of Y.

Next, we introduce the following notation $p_{ij,kl}$ that will facilitate us describe the causal effect:

$$p_{ij,kl} = P(A_0 = i, A_1 = j, Y_0 = k, Y_1 = l), \quad i, j, k, l \in \{0, 1\}.$$

Naturally,

$$\sum p_{ij,kl} = 1,\tag{1}$$

$$p_{ij,kl} \ge 0. \tag{2}$$

The probability $p_{ij,kl}$ stands for the probability of selecting a person who when Z = 0observes A = i and when Z = 1 observes A = j, and at the same time if A = 0 observes Y = k, if A = 1 observes Y = l. Notice that some of these cannot be distinguished based on the observed data. For example, $p_{00,01}$ is the probability of never-takers

responding to a tr	eatment if	they	actually	took	it.	Table	1	shows	the	partition	of
population into 16	disjoint set	s.									

Never-taker	Prob	Always-taker	Prob	rob Complier		Defier	Prob
00,00	$p_{00,00}$	11,00	$p_{11,00}$	01,00	$p_{01,00}$	10,00	$p_{10,00}$
$00,\!01$	$p_{00,01}$	11,01	$p_{11,01}$	01,01	$p_{01,01}$	10,01	$p_{10,01}$
00,10	$p_{00,10}$	$11,\!10$	$p_{11,10}$	$01,\!10$	$p_{01,10}$	10,10	$p_{10,10}$
00,11	$p_{00,11}$	11,11	$p_{11,11}$	01,11	$p_{01,11}$	10,11	$p_{10,11}$

Table 1: A partition of population to 16 disjoint sets.

These probabilities are useful in describing many causal quantities of interest. For example, the celebrated average causal effect

$$E(Y_1 - Y_0) = P(Y_1 = 1) - P(Y_0 = 1)$$

= $p_{01,01} + p_{10,01} + p_{00,01} + p_{11,01} - p_{01,10} - p_{10,10} - p_{00,10} - p_{11,10}.$

We will derive fiducial based stochastic bounds on average causal effect and other quantities. To this end we first link these unobservable probabilities with probabilities of observable events:

Proposition 2.1. Under Assumptions 1, 2, and 3,

$$q_{000} = p_{00,01} + p_{00,00} + p_{01,01} + p_{01,00}$$

$$q_{001} = p_{00,10} + p_{00,11} + p_{01,11} + p_{01,10}$$

$$q_{010} = p_{11,10} + p_{11,00} + p_{10,10} + p_{10,00}$$

$$q_{011} = p_{11,01} + p_{11,11} + p_{10,11} + p_{10,01}$$

$$q_{100} = p_{00,01} + p_{00,00} + p_{10,01} + p_{10,00}$$

$$q_{101} = p_{00,10} + p_{00,11} + p_{10,11} + p_{10,10}$$

$$q_{110} = p_{11,10} + p_{11,00} + p_{01,10} + p_{01,00}$$

$$q_{111} = p_{11,01} + p_{11,11} + p_{01,11} + p_{01,01}$$
(3)

Note that in (3) the sum of the first four equations is 1 as is the sum of the last right equations.

3 Generalized fiducial Distribution (GFD)

3.1 GFD for Multinomial Distribution

In order to assess uncertainty due to the randomness of the observed data, we derive a GFD for the probabilities $p_{ij,kl}$. We will follow an argument introduced in Lawrence et al. (2009), see also Hannig et al. (2016) and Lawrence et al. (2024), that derived GFD for parameters of multinomial distribution using arguments from Dempster-Shafer theory (Dempster, 2008). Fix $z \in \{0, 1\}$. Lawrence et al. (2009) show that GFD for q_{zay} , is obtained by considering the random polygon determined by inequalities:

$$V_{z00}^* \le q_{z00}, \quad V_{z01}^* \le q_{z01}, \quad V_{z10}^* \le q_{z10}, \quad V_{z11}^* \le q_{z11},$$
 (4)

where $V_{zay}^* \sim \text{Beta}(n_{zay}, 1)$, $a, y \in \{0, 1\}$, are independent random variables, truncated to the event that a solution to (4) exists, i.e., $\mathcal{Z}_z = \{V_{z00}^* + V_{z01}^* + V_{z10}^* + V_{z11}^* \leq 1\}$. Next, Lawrence et al. (2009) show that the distribution of $V_{z00}^*, \ldots, V_{z11}^*$ truncated to \mathcal{Z}_z is the same as the unconditional distribution of $V_{z00}^*, \ldots, V_{z11}^*$ where $(V_z^*, V_{z00}^*, \ldots, V_{z11}^*)$ follows Dirichlet $(1, n_{z00}, \ldots, n_{z11})$. Consequently, the random set

$$S_{z}^{*} = \{\mathbf{q}_{z} \in \Delta^{3}; \text{ satisfying (4)}\}$$

$$= \{(V_{z}^{*} + V_{z00}^{*}, V_{z01}^{*}, V_{z10}^{*}, V_{z11}^{*}), (V_{z00}^{*}, V_{z}^{*} + V_{z01}^{*}, V_{z10}^{*}, V_{z11}^{*}), (V_{z00}^{*}, V_{z01}^{*}, V_{z11}^{*}, V_{z11}^{*}), (V_{z00}^{*}, V_{z01}^{*}, V_{z11}^{*}, V_{z11}^{*}), (V_{z00}^{*}, V_{z01}^{*}, V_{z10}^{*}, V_{z11}^{*}), (V_{z00}^{*}, V_{z01}^{*}, V_{z00}^{*}, V_{z11}^{*}), (V_{z00}^{*}, V_{z01}^{*}, V_{z10}^{*}, V_{z11}^{*})\}$$

Any vector q_{zay}^* selected from the random polygon can be viewed as a sample from the GFD distribution for q_{zay} . Moreover, for any set \mathcal{A} the $P^*(\mathcal{S}_z \cap \mathcal{A} \neq \emptyset)$ is called plausibility of \mathcal{A} and represents the fiducial belief that the set \mathcal{A} is feasible given the observed data.

3.2 GFD for partially identified parameters

Let us now derive the GFD for our problem. We have observed two independent multinomial vectors, with parameters that are related to each other by Proposition 2.1. Thus the joint GFD for this problem is sampled from the random polygon

$$V_{000}^{*} \leq p_{00,01} + p_{00,00} + p_{01,01} + p_{01,00}$$

$$V_{001}^{*} \leq p_{00,10} + p_{00,11} + p_{01,11} + p_{01,10}$$

$$V_{010}^{*} \leq p_{11,10} + p_{11,00} + p_{10,10} + p_{10,00}$$

$$V_{011}^{*} \leq p_{11,01} + p_{11,11} + p_{10,11} + p_{10,01}$$

$$V_{100}^{*} \leq p_{00,01} + p_{00,00} + p_{10,01} + p_{10,00}$$

$$V_{101}^{*} \leq p_{00,10} + p_{00,11} + p_{10,11} + p_{10,10}$$

$$V_{110}^{*} \leq p_{11,10} + p_{11,00} + p_{01,10} + p_{01,00}$$

$$V_{111}^{*} \leq p_{11,01} + p_{11,11} + p_{01,11} + p_{01,01},$$
(6)

where again $V_{zay}^* \sim \text{Beta}(n_{zay}, 1)$ are independent random variables truncated to the event that a solution of (1), (2), and (6) exists. Because of the form of the right hand of (6), there is no longer a closed form expression for the random polygon. However, we can nevertheless sample from it using a simple acceptance sampling algorithm.

To this end, notice that the inequalities in (6) can be divided into 2 groups of 4 inequalities, each group corresponding to a different multinomial observation. Thus following the ideas described above, we propose to generate $(V_z^*, V_{z00}^*, V_{z01}^*, V_{z10}^*, V_{z11}^*)$ from Dirichlet $(1, n_{z00}, n_{z01}, n_{z10}, n_{z11})$ independently for both z = 0, 1, and then accept the proposed combined $(V_{000}^*, \ldots, V_{111}^*)$ if there exist $p_{ij,kl}$ that satisfy (1), (2), (6). The details of the proposed acceptance sampler are described in Algorithm 1.

Alg	gorit	hm	1:	А	causal	acce	ptance	samp	ler
-----	-------	----	----	---	--------	------	--------	------	-----

Input: Dataset $(Z_i, A_i, Y_i), n_{\text{mcmc}}$

1 Attempts $\leftarrow 0$;

2 for $j \leftarrow 1$ to n_{mcmc} do

3	\mathbf{r}	epeat
4		$Attempts \leftarrow Attempts + 1;$
5		Generate $(V_z^*, V_{z00}^*, V_{z01}^*, V_{z10}^*, V_{z11}^*)$ from Dirichlet $(1, n_{z00}, n_{z01}, n_{z10}, n_{z11})$
		independently for both $z = 0, 1,;$
6		Check if the system (1) , (2) , (6) has a solution (by solving an LP problem);
7	u	ntil Feasible $(V_{000}^*, \dots, V_{111}^*)$ found;
8	v	$V_j^* \leftarrow (V_{000}^*, \dots, V_{111}^*);$
9 e	nd	

10 return Accepted fiducial samples \mathbf{V}_{j}^{*} , $j = 1, \ldots, n_{\text{mcmc}}$ and acceptance rate $\frac{n_{\text{mcmc}}}{\text{Attempts}}$;

Surprisingly, the acceptance probability of our algorithm has itself a fiducial interpretation. Define the following set of observable probabilities $\mathbf{q} = (q_{000}, \dots, q_{111})$:

 $\mathcal{F} = \{ \mathbf{q} \in \Delta^3 \times \Delta^3; \text{ so that there exists } p_{ij,kl} \text{ satisfying } (1), (2), (3) \}.$ (7)

Notice that Algorithm 1 accepts a proposed sample if and only if $(S_0 \times S_1) \cap \mathcal{F} \neq \emptyset$, where S_z are defined in (5). Thus, the acceptance rate is an estimator of the fiducial probability $P^*((S_0 \times S_1) \cap \mathcal{F} \neq \emptyset)$. Proposition 2.1 implies that for $\mathbf{q} \in \mathcal{F}$ it is feasible that the data was generated from an IV model, while for $\mathbf{q} \in \mathcal{F}^{\complement}$ it is not possible for the causal assumptions to be satisfied. Consequently, the acceptance rate of the algorithm is an estimator of fiducial probability of the observed data agreeing with the IV assumptions. In other words, high acceptance rate points toward high trust in feasibility of the IV assumption, while acceptance rate near 0 suggests that the IV assumptions are likely violated.

After generating fiducial samples \mathbf{V}_{j}^{*} , we then propose Algorithm 2 to obtain upper and lower fiducial samples for a targeted parameter. Algorithm 2: Generating upper and lower fiducial samples for a targeted estimand

(average treatment effect as a special case) **Input:** The fiducial samples $\mathbf{V}_{j}^{*}, j = 1, \dots, n_{\text{mcmc}}$.

1 for j = 1 to n_{mcmc} do

Solve $\mathbf{2}$ $u_j^* = \max_p f(p_{00,00}, \dots, p_{11,11})$ $= \max_{p} p_{01,01} + p_{10,01} + p_{00,01} + p_{11,01} - p_{01,10} - p_{10,10} - p_{00,10} - p_{11,10}$ subject to constraints (1), (2), (6);Solve 3 $l_j^* = \min_p f(p_{00,00}, \dots, p_{11,11})$ $= \min_{p} p_{01,01} + p_{10,01} + p_{00,01} + p_{11,01} - p_{01,10} - p_{10,10} - p_{00,10} - p_{11,10}$ subject to constraints (1), (2), (6);

- 4 end
- 5 return The upper and lower fiducial samples u_j^* and l_j^* , $j = 1, \ldots, n_{\text{mcmc}}$.

Based on the fiducial samples, we construct pointwise confidence intervals by finding intervals of a given fiducial probability. In particular, a 95% confidence interval is formed by taking the empirical 0.025 quantile of l_i^* as a lower limit and the empirical 0.975 quantile of u_j^* as an upper limit. We also propose to use the pointwise median of l_j^* and u_j^* as a point estimator for l_j^* and u_j^* , respectively.

4 Bernstein-von Mises theorem

In this section, we present a novel Bersntein-von Mises theorem. First, we provide the following lemma.

In order to fully appreciate the mode of convergence discussed below, notice that

there are two sources of randomness present. First is the usual randomness of the observed data, e.g., **N**. The second source of the randomness is the fiducial distribution defined for each fixed data set. We indicate this randomness by using stars, e.g., P^* , \mathbf{V}^* . We prefer this notation to using a conditioning symbol, as fiducial distribution was not obtained by the Bayes theorem. The main results use convergence in distribution in probability. In particular, $X^* \xrightarrow{\mathcal{D}} Y$ in probability if the Levy-Prokhorov distance $\nu(X^*, Y) \xrightarrow{P} 0$.

Lemma 4.1. Let $\mathbf{N}_n \in \mathbb{R}^d$ be $Multinomial(n, \mathbf{q}), \mathbf{q} \in \Delta^d$. Let $\mathbf{V}_n^* | \mathbf{N}_n = \mathbf{n}_n$ be $Dirichlet(1, \mathbf{n}_n)$. Then as $n \to \infty$

1. $\sqrt{n}(\mathbf{N}_n/n - \mathbf{q}) \xrightarrow{\mathcal{D}} N(\mathbf{0}, \Sigma), \text{ where } \Sigma = \operatorname{diag}(\mathbf{q}) - \mathbf{q}\mathbf{q}^{\top}.$ 2. $\sqrt{n}(\mathbf{V}_n^* - \mathbf{N}_n^0/n) \xrightarrow{\mathcal{D}} N(\mathbf{0}, \Sigma^0) \text{ in probability where}$ $\mathbf{N}_n^0 = \begin{pmatrix} 0\\ \mathbf{N}_n \end{pmatrix} \text{ and } \Sigma^0 = \begin{pmatrix} 0 & \mathbf{0}^{\top}\\ \mathbf{0} & \Sigma \end{pmatrix}.$

The following theorem validates our assertion that the acceptance rate can be used to assert whether the IV assumptions are satisfied. In what follows \mathcal{F}° is the interior of \mathcal{F} given by (7).

Theorem 4.1. Let $\mathbf{N}_{z} = (n_{z00}, \dots, n_{z11})$ be independent $Multinomial(n_{z}, \mathbf{q}_{z})$, with $\mathbf{q}_{z} = (q_{z00}, q_{z01}, q_{z10}, q_{z11})^{\top}$ and $z \in \{0, 1\}$. Given $\mathbf{N}_{z} = \mathbf{n}_{z}$, $z \in \{0, 1\}$, let $\mathbf{V}^{*} = (V_{z}^{*}, V_{z00}^{*}, \dots, V_{z11}^{*})$ be independent $Dirichlet(1, \mathbf{n}_{z}), z \in \{0, 1\}$. Then for $\mathbf{q} = (q_{000}, \dots, q_{111})$, as $n_{0}, n_{1} \to \infty$:

1. if $\mathbf{q} \in \mathcal{F}^{\circ}$, then $P^{*}((\mathcal{S}_{0} \times \mathcal{S}_{1}) \cap \mathcal{F} \neq \emptyset) \xrightarrow{P} 1;$ 2. if $\mathbf{q} \in (\mathcal{F}^{\complement})^{\circ}$, then $P^{*}((\mathcal{S}_{0} \times \mathcal{S}_{1}) \cap \mathcal{F} \neq \emptyset) \xrightarrow{P} 0;$ 3. if $\mathbf{q} \in \partial \mathcal{F}$ and $n_{z}/(n_{0} + n_{1}) \rightarrow \lambda_{z} \in (0, 1)$, then

$$\liminf P(P^*((\mathcal{S}_0 \times \mathcal{S}_1) \cap \mathcal{F} \neq \emptyset) \le u) \ge u \text{ for all } u \in (0,1).$$

Note that the feasibility set \mathcal{F} is polyhedral cone. If $\mathbf{q} \in \partial \mathcal{F}$ is additionally in the interior of a face, then the inequality in (C.1) becomes equality in the limit and the acceptance rate itself actually converges to the uniform distribution. Thus large values of the acceptance rate indicate that we are safely within the region where IV assumptions are satisfied, small values mean that the IV assumptions are likely violated while values in-between indicate we may be near the boundary.

Next we prove the Bernstein-von Mises for solutions of an optimization problem in Algorithm 2 or one of its variants. We will use notation introduced in the previous theorem.

Theorem 4.2. Let \mathbf{N}_z be independent $Multinomial(n_z, \mathbf{q}_z), z \in \{0, 1\}$. Let the conditional distribution $\mathbf{V}_z^* | \mathbf{N}_z = \mathbf{n}_z$ be independent $Dirichlet(1, \mathbf{n}_z), z \in \{0, 1\}$.

For each $\bar{\mathbf{q}} \in \mathcal{F}$ define $g(\bar{\mathbf{q}}) = \min f(p_{00,00}, \dots, p_{11,11})$ subject to (1), (2), (3), where f is the objective function determined by the estimand of interest. Assume that there is an open set \mathcal{Q} so that $\mathbf{q} \in \mathcal{Q} \subset \mathcal{F}$, and the function $g(\bar{\mathbf{q}})$ is continuously differentiable on \mathcal{Q} . Set $n_z/(n_0 + n_1) \rightarrow \lambda_z \in (0, 1)$. Then as $n_0, n_1 \rightarrow \infty$

1.
$$\sqrt{n_0 + n_1}(g(\mathbf{N}_0/n_0, \mathbf{N}_1/n_1) - g(\mathbf{q})) \xrightarrow{\mathcal{D}} N(0, \sigma^2), where$$

$$\sigma^2 = \sum_{z=0}^1 \lambda_z^{-1} \left(\sum_{a,y=0}^1 \left(\frac{\partial g(\mathbf{q})}{\partial q_{zay}} \right)^2 q_{zay} - \left(\sum_{a,y=0}^1 \frac{\partial g(\mathbf{q})}{\partial q_{zay}} q_{zay} \right)^2 \right).$$

2. Denote the random variable $G^* = \min f(p_{00,00}, \dots, p_{11,11})$ subject to (1), (2), (6).

$$\sqrt{n_0 + n_1}(G^* - g(\mathbf{N}_0/n_0, \mathbf{N}_1/n_1)) \xrightarrow{\mathcal{D}} N(0, \sigma^2)$$
 in probability.

Note that Theorem 4.2 also holds for $\max f(p_{00,00}, \ldots, p_{11,11})$. The consequence of this result is that we have a correct coverage for confidence intervals for the theoretical lower and upper bounds on the quantity of interest.

The assumption on differentiability of $g(\mathbf{q})$ is satisfied if the objective function f is continuously differentiable and the active constraints in (2) are the same for all $\mathbf{q} \in \mathcal{Q}$.

5 Simulation study

In this section, we examined the coverage and average length of 95% fiducial confidence intervals for the lower and upper bounds. We considered the following scenario from Cui (2021):

$$Pr(Y = 1|U, A) = U/16 + 1/5A + 1/15,$$

$$Pr(A = 1|U, Z) = U/16 + 2/5Z + 1/2,$$

$$U, Z \sim \text{Bernoulli}(0.5).$$

and a variant setting with no unmeasured confounding

$$\Pr(Y = 1|A) = 1/5A + 1/15,$$

 $\Pr(A = 1|Z) = 1/5Z + 1/5,$
 $Z \sim \text{Bernoulli}(0.5).$

We chose various sample sizes n = 25, 50, 100. Each scenario was simulated 200 times. The fiducial estimates were based on 1000 iterations. In Tables 2-5, LR denotes the error rate that the true parameter is less than the lower confidence limit; UR denotes the error rate that the true parameter is greater than the upper confidence limit. The two-sided error rate is obtained by adding the values in columns LR and UR, and the value around 5% in aggregate indicates achieving nominal coverage. WD is the average width of the confidence interval.

We also compare the proposed method with a Bayesian approach with Dirichlet prior (Loh and Richardson, 2016). We use the recommended two priors, Dirichlet(1,1,0,0,1,1,1,1) and Dirichlet(1/2,1/2,0,0,1,1,1,1) as given in Loh and Richardson (2016). As can be seen from the tables, the proposed fiducial confidence intervals maintain the aggregate coverage and have reasonable length. The proposed fiducial approach is comparable to other methods in Scenario 1 and outperforms other methods in Scenario 2. More importantly, we do not need to specify a particular prior.

	Fiducial			Dirie	chlet p	orior 1	Dirichlet prior 2			
	LR	UR	WD	LR	UR	WD	LR	UR	WD	
n = 25	0.5	4.0	0.61	2.5	0.5	0.59	2.5	1.0	0.62	
n = 50	1.5	3.5	0.46	3.0	2.0	0.45	4.0	2.5	0.46	
n=100	4.0	1.5	0.34	4.5	1.0	0.34	4.5	1.0	0.34	

Table 2: Error rates in percent and average width of 95% confidence intervals for the lower bound of Scenario 1. LR denotes the error rate that the true parameter is less than the lower confidence limit; UR denotes the error rate that the true parameter is greater than the upper confidence limit; WD is the average width of the confidence interval. Dirichlet priors 1 and 2 use Dirichlet(1,1,0,0,1,1,1,1) and Dirichlet(1/2,1/2,0,0,1,1,1,1), respectively.

	Fiducial			Dirie	chlet p	orior 1	Dirichlet prior 2			
	LR	UR	WD	LR	UR	WD	LR	UR	WD	
n=25	4.5	1.0	0.48	0	1.5	0.50	0.5	3.0	0.52	
n=50	4.5	1.0	0.36	0.5	1.5	0.37	1.0	3.0	0.37	
n=100	2.5	1.5	0.27	2.0	1.5	0.27	1.5	1.5	0.27	

Table 3: Error rates in percent and average width of 95% confidence intervals for the upper bound of Scenario 1. LR denotes the error rate that the true parameter is less than the lower confidence limit; UR denotes the error rate that the true parameter is greater than the upper confidence limit; WD is the average width of the confidence interval. Dirichlet priors 1 and 2 use Dirichlet(1,1,0,0,1,1,1,1) and Dirichlet(1/2,1/2,0,0,1,1,1,1), respectively.

6 Real data application

We considered the Vitamin A example from Balke and Pearl (1997) which is a classic illustration of how causal inference can be applied to estimate the effect of an interven-

	I	Fiduci	al	Dirie	chlet p	orior 1	Dirichlet prior 2			
	LR	UR	WD	LR	UR	WD	LR	UR	WD	
n = 25	2.0	1.0	0.50	2.5	0	0.50	7.5	0.5	0.50	
n=50	1.5	1.5	0.36	1.5	0.5	0.36	4.0	1.0	0.36	
n=100	3.5	1.5	0.27	4.0	1.5	0.27	6.0	0	0.27	

Table 4: Error rates in percent and average width of 95% confidence intervals for the lower bound of Scenario 2. LR denotes the error rate that the true parameter is less than the lower confidence limit; UR denotes the error rate that the true parameter is greater than the upper confidence limit; WD is the average width of the confidence interval. Dirichlet priors 1 and 2 use Dirichlet(1,1,0,0,1,1,1,1) and Dirichlet(1/2,1/2,0,0,1,1,1,1), respectively.

	Ι	Fiduci	al	Dirio	chlet p	rior 1	Dirichlet prior 2			
	LR	UR	WD	LR	UR	WD	LR	UR	WD	
n=25	3.0	0.5	0.49	0	6.0	0.51	0.5	4.5	0.53	
n=50	1.0	2.5	0.37	0	10.5	0.37	0.5	6.0	0.38	
n=100	2.0	2.5	0.27	0	5.0	0.27	0.5	4.5	0.27	

Table 5: Error rates in percent and average width of 95% confidence intervals for the upper bound of Scenario 2. LR denotes the error rate that the true parameter is less than the lower confidence limit; UR denotes the error rate that the true parameter is greater than the upper confidence limit; WD is the average width of the confidence interval. Dirichlet priors 1 and 2 use Dirichlet(1,1,0,0,1,1,1,1) and Dirichlet(1/2,1/2,0,0,1,1,1,1), respectively.

tion subject to unmeasured confounders. Balke and Pearl (1997) analyzed data from a randomized experiment, where 450 villages were randomly offered oral doses of vitamin A supplementation, with 221 assigned to the control group and 229 assigned to the treatment group. In the study, researchers are interested in the effect of consuming Vitamin A supplementation on reducing mortality rates.

The dataset contained 10231 individuals from villages assigned to vitamin A supplementation and 10919 untreated individuals which defines a natural binary IV Z. Balke and Pearl (1997) reported bounds on the average treatment effect, $-0.1946 \leq E(Y_1 - Y_0) \leq 0.0054$. Their conclusion is that the vitamin A supplementation, if uniformly administered, is capable of increasing mortality rate by as much as 19.46% and is incapable of reducing mortality rate by more than 5.4%.

We applied the proposed method to provide uncertainty quantification. The fiducial estimates were based on 10000 iterations. The fiducial point estimators for the bounds are the same as those reported by Balke and Pearl (1997). The proposed 95% confidence interval for the lower bound is (-0.2019, -0.1873) and the proposed 95% confidence interval for the upper bound is (0.0040, 0.0071). Our estimates suggest that both bounds are statistically significant.

7 Other causal estimands

7.1 Complier average causal effect

Given our framework, it is straightforward to see that the complier average causal effect (Imbens and Angrist, 1994; Angrist et al., 1996) is

$$E[Y_1 - Y_0 | A_1 > A_0] = \frac{p_{01,01} - p_{01,10}}{p_{01,00} + p_{01,01} + p_{01,10} + p_{01,11}}.$$

Therefore, one can replace u_j^* and l_j^* in Algorithm 2 with

$$u_j^* = \max_p \frac{p_{01,01} - p_{01,10}}{p_{01,00} + p_{01,01} + p_{01,10} + p_{01,11}},$$

and

$$l_j^* = \min_p \frac{p_{01,01} - p_{01,10}}{p_{01,00} + p_{01,01} + p_{01,10} + p_{01,10}}.$$

Similarly, the always-taker's, never-taker's, and defyer's average causal effect can also be obtained.

7.2 Nudge average treatment effect

Recently, Tchetgen Tchetgen (2024) proposes the so-called nudge average treatment effect, i.e., the IV estimand that recovers the average causal effect for the subgroup of units for whom the treatment is manipulable by the instrument. Given our framework, the nudge average treatment effect is

$$E[Y_1 - Y_0 | A_1 \neq A_0] = \frac{p_{01,01} + p_{10,01} - p_{01,10} - p_{10,10}}{p_{01,00} + p_{01,01} + p_{01,10} + p_{01,11} + p_{10,00} + p_{10,01} + p_{10,10} + p_{10,11}}$$

Therefore, one can replace u_j^* and l_j^* in Algorithm 2 with

$$u_j^* = \max_p \frac{p_{01,01} + p_{10,01} - p_{01,10} - p_{10,10}}{p_{01,00} + p_{01,01} + p_{01,10} + p_{01,11} + p_{10,00} + p_{10,01} + p_{10,10} + p_{10,11}}$$

and

$$l_j^* = \min_p \frac{p_{01,01} + p_{10,01} - p_{01,10} - p_{10,10}}{p_{01,00} + p_{01,01} + p_{01,10} + p_{01,11} + p_{10,00} + p_{10,01} + p_{10,10} + p_{10,11}}$$

8 Other causal assumptions

8.1 Monotonicity

Monotonicity (Imbens and Angrist, 1994; Angrist et al., 1996) is a commonly seen assumption in IV analysis, which ensures that the instrument affects treatment assignment in a consistent direction across all units. This assumption is important in settings like complier average treatment effect identification, where we are concerned with the average treatment effect among compliers.

Given our framework, monotonicity refers to $p_{10,00} = 0$, $p_{10,01} = 0$, $p_{10,10} = 0$, $p_{10,11} = 0$, i.e., there are no defiers. In Section E, we show that under monotonicity, the complier average causal effect is indeed point identified through our framework.

8.2 New drug assumption

Given our framework, the new drug assumption $A_0 = 0$ implies $p_{11,00} = 0, p_{11,01} = 0, p_{11,10} = 0, p_{10,10} = 0, p_{10,10} = 0, p_{10,11} = 0, p_{10,11} = 0$, i.e., there are neither always-takers nor defiers.

9 Discussion

In this paper, we proposed novel fiducial approaches for learning partially identified parameters in a statistical model. Using IV models as a running example, we proposed a novel acceptance sampler to sample from fiducial distribution. As a by-product, the acceptance rate of the proposed sampling algorithm provided a measure of whether the observed data agree with the IV assumptions. We established Bernstein-von Mises theorem for the proposed estimator to provide theoretical justification of our confidence intervals. We illustrated our framework via simulation studies and a real data application. Our paper contributes to the literature on both foundations of statistics and causal inference.

Several generalizations can be made following the proposed framework.

1) We focus on binary instruments, treatments, and outcomes in this paper. The idea can be extended to categorical variables.

2) We consider the randomization scheme of Z to be known. If it is not known, one can include this part into data generation mechanism to account for uncertainty.

3) It would be interesting to explore our framework in meditation analysis (VanderWeele, 2015; Steen and Vansteelandt, 2018) and survival analysis (Fleming and Harrington, 2013) under partial identification.

4) It would be of interest to extend our framework to other sets of IV assumptions such as stochastic monotonicity (Small et al., 2017), homogeneous instrument-treatment association (Wang and Tchetgen Tchetgen, 2018), homogeneous treatment-

outcome association (Wang and Tchetgen Tchetgen, 2018), no unmeasured common effect modifier (Cui and Tchetgen Tchetgen, 2021), and other causal estimands such as average treatment effect on the treated (Robins, 1989).

5) It would also be of interest to extend our framework to weak IVs (Zhao et al., 2020; Ye et al., 2021), invalid IVs (Small, 2007; Kang et al., 2016; Guo et al., 2018; Tchetgen Tchetgen et al., 2021; Ye et al., 2024; Kang et al., 2024; Dukes et al., 2024), and proxy variables (Miao et al., 2018; Cui et al., 2024; Ghassami et al., 2023).

Appendix

A Proof of Proposition 2.1

Proof. We first prove the first equation in (3). By definition, we know that

$$p_{00,01} + p_{00,00} + p_{01,01} + p_{01,00}$$

= $P(A_0 = 0, A_1 = 0, Y_0 = 0, Y_1 = 1) + P(A_0 = 0, A_1 = 0, Y_0 = 0, Y_1 = 0)$
+ $P(A_0 = 0, A_1 = 1, Y_0 = 0, Y_1 = 1) + P(A_0 = 0, A_1 = 1, Y_0 = 0, Y_1 = 0)$
= $P(A_0 = 0, Y_0 = 0, Y_1 = 1) + P(A_0 = 0, Y_0 = 0, Y_1 = 0)$
= $P(A_0 = 0, Y_0 = 0)$
= $P(A_0 = 0, Y_0 = 0|Z = 0)$ (by Assumption 2)
= $P(A = 0, Y = 0|Z = 0)$] = q_{000} .

The other equalities are proved analogously.

B Proof of Lemma 4.1

Proof. The first assertion follows immediately by the Central Limit Theorem.

For the second statement, let us first assume that $\mathbf{N}_n/n \to \mathbf{q}$ a.s. Conditionally on $\mathbf{N}_n = \mathbf{n}_n$, let $\mathbf{X}_n = (X_{n,0}, X_{n,1}, \dots, X_{n,d})^{\top}$ be independent $X_{n,i} \sim \Gamma(n_{n,i}, 1)$ random variables, with $n_{n,0} = 1$. Using a coupling we select a version of $\mathbf{X}_n/n \to \mathbf{q}^0$ whenever $\mathbf{n}_n/n \to \mathbf{q}$. A direct calculation using characteristic function shows that

$$n^{-1/2}(\mathbf{X}_n - \mathbf{N}_n^0) \xrightarrow{\mathcal{D}} N(\mathbf{0}, \operatorname{diag}(\mathbf{q}^0)) \text{ a.s., where } \mathbf{q}_0 = \begin{pmatrix} 0 \\ \mathbf{q} \end{pmatrix}.$$

Next, let $f(\mathbf{X}_n) = \mathbf{X}_n/(\mathbf{1}^{\top}\mathbf{X}_n) \sim \text{Dirichlet}(1, \mathbf{n}_n)$ and set $\mathbf{V}_n^* = f(\mathbf{X}_n)$. Using Taylor series we get

$$f(\mathbf{X}_n/n) = f(\mathbf{N}_n^0/n) - \int_0^1 Df(\mathbf{N}_n^0/n + v(\mathbf{X}_n/n - \mathbf{N}_n^0/n)) dv (\mathbf{X}_n/n - \mathbf{N}_n^0/n)$$

and consequently

$$\sqrt{n} \left(\mathbf{V}_n^* - \mathbf{N}_n^0 / n \right) = \int_0^1 Df(\mathbf{N}_n^0 / n + v(\mathbf{X}_n / n - \mathbf{N}_n^0 / n)) dv \ n^{-1/2} (\mathbf{X}_n - \mathbf{N}_n^0).$$
(B.1)

The integral on the right hand side of (B.1) converges to $I - \mathbf{q}^0 \mathbf{1}^\top$ a.s. and consequently $\sqrt{n} \left(\mathbf{V}_n^* - \mathbf{N}_n^0 / n \right) \xrightarrow{\mathcal{D}} N(\mathbf{0}, \Sigma^0)$ a.s.

Finally, if $\mathbf{N}_n/n \xrightarrow{P} \mathbf{q}$ only, the theorem follows by use of subsequence criterion for convergence in probability and the Prokhorov's metric between distribution of $\sqrt{n} \left(\mathbf{V}_n^* - \mathbf{N}_n^0/n \right)$ and $N(\mathbf{0}, \Sigma^0)$.

C Proof of Theorem 4.1

Proof. Lemma 4.1 and Equation (5) imply that $\sqrt{n_z} \operatorname{diam} S_z \xrightarrow{P} 0$ as $n_z \to \infty$. Moreover, for any open ball \mathcal{B} containing \mathbf{q} , $P^*(S_0 \times S_1 \subset \mathcal{B}) \xrightarrow{P} 1$. Statements 1 and 2 follow immediately.

Next, notice that \mathcal{F} is convex. If $\mathbf{q} \in \partial \mathcal{F}$ then there is a supporting hyperplane plane A and a corresponding normal vector \mathbf{v} pointing out of \mathcal{F} . Notice that

$$P^*((\mathcal{S}_0 \times \mathcal{S}_1) \cap \mathcal{F} = \emptyset) \ge P^*(\min \mathbf{v}^\top (\mathcal{S}_0 \times \mathcal{S}_1) > \mathbf{v}^\top \mathbf{q}).$$
(C.1)

Lemma 4.1 implies that

$$\sqrt{n_0 + n_1}((\mathbf{N}_0/n_0, \mathbf{N}_1/n_1) - \mathbf{q}) \xrightarrow{\mathcal{D}} \mathbf{Z} \sim N(\mathbf{0}, \Sigma),$$
 (C.2)

where

$$\Sigma = \begin{pmatrix} \lambda_0^{-1}(\operatorname{diag}(\mathbf{q}_0) - \mathbf{q}_0\mathbf{q}_0^{\top}) & 0\\ 0 & \lambda_1^{-1}(\operatorname{diag}(\mathbf{q}_1) - \mathbf{q}_1\mathbf{q}_1^{\top}) \end{pmatrix}$$

Moreover, $\mathbf{v}^{\top}\mathbf{Z}$ is a non-degenerate, mean zero, Gaussian random variable. Denote F(s) its distribution function.

Skorokhod representation theorem allows us to select sequence of the multinationals for which the convergence in (C.2) is a.s. For such a sequence

$$\sqrt{n_0 + n_1}((\mathcal{S}_0 \times \mathcal{S}_1) - (\mathbf{N}_0/n_0, \mathbf{N}_1/n_1)) \xrightarrow{\mathcal{D}} N(\mathbf{0}, \Sigma) \text{ a.s.}$$

Thus by subtracting $\mathbf{v}^{\top}(\mathbf{N}_0/n_0, \mathbf{N}_1/n_1)$ and multiplying by $\sqrt{n_0 + n_1}$ on both sides one gets

$$P^*(\min \mathbf{v}^{\top}(\mathcal{S}_0 \times \mathcal{S}_1) > \mathbf{v}^{\top}\mathbf{q}) \to F(-\mathbf{v}^{\top}\mathbf{Z}) \sim U(0,1) \text{ a.s}$$

The assertion follows.

D Proof of Theorem 4.2

Proof. Theorem 4.1 implies that the fiducial probability that G^* exist, i.e., \mathbf{V}^* is such that the optimization problem is feasible, goes to 1 in probability. Consequently, the result is a direct application of Lemma 4.1 and the delta method.

E Proof of Section 8.1

Proof. When there is no defier, $p_{10,00} = 0$, $p_{10,01} = 0$, $p_{10,10} = 0$, $p_{10,11} = 0$. Recall that we have

$$\begin{aligned} q_{000} &= (p_{00,01} + p_{00,00} + p_{01,01} + p_{01,00}) \\ q_{100} &= (p_{00,01} + p_{00,00} + p_{10,01} + p_{10,00}) \\ q_{010} &= (p_{11,10} + p_{11,00} + p_{10,10} + p_{10,00}) \\ q_{110} &= (p_{11,10} + p_{11,00} + p_{01,10} + p_{01,00}) \\ q_{001} &= (p_{00,10} + p_{00,11} + p_{01,11} + p_{01,10}) \\ q_{101} &= (p_{11,01} + p_{11,11} + p_{10,11} + p_{10,01}) \\ q_{111} &= (p_{11,01} + p_{11,11} + p_{01,11} + p_{01,01}), \end{aligned}$$

and the complier average causal effect is

$$E[Y_1 - Y_0 | A_1 > A_0]$$

= $\frac{p_{01,01} - p_{01,10}}{p_{01,00} + p_{01,01} + p_{01,10} + p_{01,11}}.$

By a simple calculation, we have

 $p_{01,01} - p_{01,10} = q_{000} - q_{110} - q_{100} + q_{010},$

and

 $p_{01,00} + p_{01,01} + p_{01,10} + p_{01,11} = (q_{000} + q_{110} + q_{001} + q_{111} - q_{100} - q_{010} - q_{101} - q_{011})/2,$

which completes the proof.

References

- Abadie, A. (2005), "Semiparametric difference-in-differences estimators," *The review* of economic studies, 72, 1–19.
- Angrist, J. D., Imbens, G. W., and Rubin, D. B. (1996), "Identification of causal effects using instrumental variables," *Journal of the American statistical Association*, 91, 444–455.
- Arkhangelsky, D., Athey, S., Hirshberg, D. A., Imbens, G. W., and Wager, S. (2021),"Synthetic difference-in-differences," *American Economic Review*, 111, 4088–4118.
- Balke, A. and Pearl, J. (1994), "Counterfactual probabilities: Computational methods, bounds and applications," in Uncertainty in artificial intelligence, Elsevier, pp. 46– 54.
- (1997), "Bounds on treatment effects from studies with imperfect compliance," Journal of the American statistical Association, 92, 1171–1176.
- Berger, J. O., Bernardo, J. M., and Sun, D. (2009), "The formal definition of reference priors," *The Annals of Statistics*, 37, 905–938.
- (2012), "Objective priors for discrete parameter spaces," Journal of the American Statistical Association, 107, 636–648.

- Bonvini, M. and Kennedy, E. H. (2022), "Sensitivity Analysis via the Proportion of Unmeasured Confounding," Journal of the American Statistical Association, 117, 1540–1550.
- Chen, P., Xu, A., and Ye, Z.-S. (2016), "Generalized fiducial inference for accelerated life tests with Weibull distribution and progressively type-II censoring," *IEEE Transactions on Reliability*, 65, 1737–1744.
- Chernozhukov, V., Hong, H., and Tamer, E. (2007), "Estimation and confidence regions for parameter sets in econometric models 1," *Econometrica*, 75, 1243–1284.
- Chernozhukov, V., Lee, S., and Rosen, A. M. (2013), "Intersection bounds: Estimation and inference," *Econometrica*, 81, 667–737.
- Cornfield, J., Haenszel, W., Hammond, E. C., Lilienfeld, A. M., Shimkin, M. B., and Wynder, E. L. (1959), "Smoking and lung cancer: recent evidence and a discussion of some questions," *Journal of the National Cancer institute*, 22, 173–203.
- Cui, Y. (2021), "Individualized Decision-Making Under Partial Identification: Three Perspectives, Two Optimality Results, and One Paradox," *Harvard Data Science Review*, 3, 1–19.
- Cui, Y. and Hannig, J. (2019), "Nonparametric generalized fiducial inference for survival functions under censoring (with discussions and rejoinder)," *Biometrika*, 106, 501–518.
- Cui, Y., Hannig, J., and Kosorok, M. (2023), "A unified nonparametric fiducial approach to interval-censored data," *Journal of the American Statistical Association*.
- Cui, Y., Pu, H., Shi, X., Miao, W., and Tchetgen Tchetgen, E. (2024), "Semiparametric proximal causal inference," *Journal of the American Statistical Association*, 119, 1348–1359.

- Cui, Y. and Tchetgen Tchetgen, E. (2021), "A semiparametric instrumental variable approach to optimal treatment regimes under endogeneity," *Journal of the American Statistical Association*, 116, 162–173.
- Cui, Y. and Xie, M.-g. (2023), "Confidence distribution and distribution estimation for modern statistical inference," in Springer Handbook of Engineering Statistics, Springer, pp. 575–592.
- Dempster, A. (1968), "Upper and Lower Probabilities Generated by a Random Closed Interval," The Annals of Mathematical Statistics, 39, 957–966.
- Dempster, A. P. (2008), "The Dempster-Shafer Calculus for Statisticians," International Journal of Approximate Reasoning, 48, 365–377.
- Ding, P. (2024), A first course in causal inference, CRC Press.
- Dukes, O., Richardson, D. B., Shahn, Z., Robins, J. M., and Tchetgen Tchetgen, E. J. (2024), "Using negative controls to identify causal effects with invalid instrumental variables," *Biometrika*, asae064.
- Fisher, R. A. (1922), "On the mathematical foundations of theoretical statistics," Philosophical Transactions of the Royal Society of London. Series A, 222, 309 – 368.
- (1930), "Inverse probability," *Proceedings of the Cambridge Philosophical Society*, xxvi, 528–535.
- (1933), "The concepts of inverse probability and fiducial probability referring to unknown parameters," *Proceedings of the Royal Society of London series A*, 139, 343–348.
- Fleming, T. R. and Harrington, D. P. (2013), Counting processes and survival analysis, vol. 625, John Wiley & Sons.

- Ghassami, A., Shpitser, I., and Tchetgen, E. T. (2023), "Partial identification of causal effects using proxy variables," *arXiv preprint arXiv:2304.04374*.
- Greenland, S. (2000), "An introduction to instrumental variables for epidemiologists," International journal of epidemiology, 29, 722–729.
- Guo, Z., Kang, H., Tony Cai, T., and Small, D. S. (2018), "Confidence intervals for causal effects with invalid instruments by using two-stage hard thresholding with voting," *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 80, 793–815.
- Hannig, J. and Iyer, H. (2022), "Testing for calibration discrepancy of reported likelihood ratios in forensic science," *Journal of the Royal Statistical Society Series A: Statistics in Society*, 185, 267–301.
- Hannig, J., Iyer, H., Lai, R. C., and Lee, T. C. (2016), "Generalized Fiducial Inference: A Review and New Results," *Journal of the American Statistical Association*, 111, 1346–1361.
- Hannig, J., Riman, S., Iyer, H., and Vallone, P. M. (2019), "Are reported likelihood ratios well calibrated?" Forensic Science International: Genetics Supplement Series, 7, 572–574.
- Hernán, M. A. and Robins, J. M. (2006), "Instruments for causal inference: an epidemiologist's dream?" *Epidemiology*, 17, 360–372.
- (2020), "Causal inference: What If,".
- Hjort, N. L. and Schweder, T. (2018), "Confidence distributions and related themes," Journal of Statistical Planning and Inference, 195, 1–13.
- Hou, X., Zhang, L., Wang, P., and Xie, M.-g. (2024), "Repro Samples Method for High-dimensional Logistic Model," arXiv preprint arXiv:2403.09984.

- Imbens, G. W. and Angrist, J. D. (1994), "Identification and Estimation of Local Average Treatment Effects," *Econometrica*, 62, 467–475.
- Imbens, G. W. and Manski, C. F. (2004), "Confidence intervals for partially identified parameters," *Econometrica*, 72, 1845–1857.
- Kaido, H., Molinari, F., and Stoye, J. (2019), "Confidence intervals for projections of partially identified parameters," *Econometrica*, 87, 1397–1432.
- Kang, H., Guo, Z., Liu, Z., and Small, D. (2024), "Identification and Inference with Invalid Instruments," Annual Review of Statistics and Its Application, 12.
- Kang, H., Zhang, A., Cai, T. T., and Small, D. S. (2016), "Instrumental variables estimation with some invalid instruments and its application to Mendelian randomization," *Journal of the American statistical Association*, 111, 132–144.
- Lawrence, E., Liu, C., Vander Wiel, S., and Zhang, J. (2009), "A new method for multinomial inference using Dempster-Shafer theory," Preprint.
- Lawrence, E. C., Murph, A. C., Wiel, S. A. V., and Liu, C. (2024), "A New Method for Multinomial Inference using Dempster-Shafer Theory," .
- Liang, F., Kim, S., and Sun, Y. (2024), "Extended fiducial inference: toward an automated process of statistical inference," *Journal of the Royal Statistical Society Series* B: Statistical Methodology, qkae082.
- Liu, C. and Martin, R. (2024), "Inferential models and possibility measures," in Handbook of Bayesian, Fiducial, and Frequentist Inference, Chapman and Hall/CRC, pp. 344–363.
- Liu, Y. and Hannig, J. (2017), "Generalized Fiducial Inference for Logistic Graded Response Models," *Psychometrika*, 82, 1097–1125.

- Liu, Y., Hannig, J., and Pal Majumder, A. (2019), "Second-Order Probability Matching Priors for the Person Parameter in Unidimensional IRT Models," *Psychometrika*, 84, 701–718.
- Loh, W. W. and Richardson, T. S. (2016), noncompliance: Causal Inference in the Presence of Treatment Noncompliance Under the Binary Instrumental Variable Model, r package version 0.2.2.
- Manski, C. F. (1990), "Nonparametric bounds on treatment effects," The American Economic Review, 80, 319–323.
- Martin, R. and Liu, C. (2013), "Inferential models: A framework for prior-free posterior probabilistic inference," *Journal of the American Statistical Association*, 108, 301 – 313.
- (2015), Inferential models: Reasoning with uncertainty, Chapman & Hall/CRC
 Monographs on Statistics & Applied Probability, CRC Press.
- Miao, W., Geng, Z., and Tchetgen Tchetgen, E. J. (2018), "Identifying causal effects with proxy variables of an unmeasured confounder," *Biometrika*, 105, 987–993.
- Murph, A. C., Hannig, J., and Williams, J. P. (2024), "Introduction to generalized fiducial inference," in *Handbook of Bayesian*, *Fiducial*, and *Frequentist Inference*, Chapman and Hall/CRC, pp. 276–299.
- Neupert, S. D., Growney, C. M., Zhu, X., Sorensen, J. K., Smith, E. L., and Hannig, J. (2021), "BFF: bayesian, fiducial, and frequentist analysis of cognitive engagement among cognitively impaired older adults," *Entropy*, 23, 428.
- Neyman, J. (1923), "Sur les applications de la theorie des probabilites aux experiences agricoles: Essai des principes (In Polish). English translation by DM Dabrowska and TP Speed (1990)," *Statistical Science*, 5, 465–480.

- Richardson, A., Hudgens, M. G., Gilbert, P. B., and Fine, J. P. (2015), "Nonparametric bounds and sensitivity analysis of treatment effects," *Statistical science: a review journal of the Institute of Mathematical Statistics*, 29, 596.
- Richardson, T. S. and Robins, J. M. (2010), "Analysis of the binary instrumental variable model," *Heuristics, Probability and Causality: A Tribute to Judea Pearl*, 25, 415–444.
- Robins, J. M. (1989), "The analysis of randomized and non-randomized AIDS treatment trials using a new approach to causal inference in longitudinal studies," *Health service research methodology: a focus on AIDS*, 113–159.
- Rosenbaum, P. R. (2002), Observational Studies, New York, NY: Springer New York, pp. 1–17.
- Rubin, D. B. (2005), "Causal inference using potential outcomes: Design, modeling, decisions," Journal of the American Statistical Association, 100, 322–331.
- Schweder, T. and Hjort, N. L. (2016), Confidence, likelihood, probability, vol. 41, Cambridge University Press.
- Shafer, G. (1976), A mathematical theory of evidence, Princeton university press Princeton.
- Shi, X., Miao, W., and Tchetgen, E. T. (2020), "A selective review of negative control methods in epidemiology," *Current epidemiology reports*, 7, 190–202.
- Singh, K., Xie, M., and Strawderman, W. E. (2005), "Combining information from independent sources through confidence distributions," *The Annals of Statistics*, 33, 159–183.

Small, D. S. (2007), "Sensitivity analysis for instrumental variables regression with

overidentifying restrictions," Journal of the American Statistical Association, 102, 1049–1058.

- Small, D. S., Tan, Z., Ramsahai, R. R., Lorch, S. A., and Brookhart, M. A. (2017), "Instrumental Variable Estimation with a Stochastic Monotonicity Assumption," *Statistical Science*, 32, 561 – 579.
- Steen, J. and Vansteelandt, S. (2018), "Mediation analysis," in Handbook of Graphical Models, CRC Press, pp. 405–438.
- Su, Y., Hannig, J., and Lee, T. C. (2022), "Uncertainty Quantification in Graphon Estimation Using Generalized Fiducial Inference," *IEEE Transactions on Signal and Information Processing over Networks*, 8, 597–609.
- Swanson, S. A., Hernán, M. A., Miller, M., Robins, J. M., and Richardson, T. S. (2018), "Partial identification of the average treatment effect using instrumental variables: review of methods for binary instruments, treatments, and outcomes," *Journal of* the American Statistical Association, 113, 933–947.
- Swanson, S. A., Holme, Ø., Løberg, M., Kalager, M., Bretthauer, M., Hoff, G., Aas, E., and Hernán, M. A. (2015), "Bounding the per-protocol effect in randomized trials: an application to colorectal cancer screening," *Trials*, 16, 1–11.
- Tchetgen Tchetgen, E., Sun, B., and Walter, S. (2021), "The GENIUS approach to robust Mendelian randomization inference," *Statistical Science*, 36, 443–464.
- Tchetgen Tchetgen, E. J. (2024), "The Nudge Average Treatment Effect," arXiv eprints, arXiv-2410.
- Tchetgen Tchetgen, E. J., Ying, A., Cui, Y., Shi, X., and Miao, W. (2024), "An introduction to proximal causal inference," *Statistical Science*, 39, 375–390.

- VanderWeele, T. J. (2015), Explanation in causal inference: methods for mediation and interaction, Oxford University Press.
- VanderWeele, T. J. and Ding, P. (2017), "Sensitivity analysis in observational research: introducing the E-value," Annals of internal medicine, 167, 268–274.
- Wager, S. (2024), "Causal Inference: A Statistical Learning Approach,".
- Wang, L. and Tchetgen Tchetgen, E. (2018), "Bounded, efficient and multiply robust estimation of average treatment effects using instrumental variables," *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 80, 531–550.
- Xie, M. and Singh, K. (2013), "Confidence Distribution, the Frequentist Distribution Estimator of a Parameter: A Review," *International Statistical Review*, 81, 3 – 39.
- Xie, M. and Wang, P. (2024), "Repro Samples Method for a Performance Guaranteed Inference in General and Irregular Inference Problems," arXiv preprint arXiv:2402.15004.
- Xie, M.-g. and Wang, P. (2022), "Repro samples method for finite-and large-sample inferences," *arXiv preprint arXiv:2206.06421*.
- Ye, T., Ertefaie, A., Flory, J., Hennessy, S., and Small, D. S. (2023), "Instrumented difference-in-differences," *Biometrics*, 79, 569–581.
- Ye, T., Liu, Z., Sun, B., and Tchetgen Tchetgen, E. (2024), "GENIUS-MAWII: For robust Mendelian randomization with many weak invalid instruments," *Journal of* the Royal Statistical Society Series B: Statistical Methodology, qkae024.
- Ye, T., Shao, J., and Kang, H. (2021), "Debiased inverse-variance weighted estimator in two-sample summary-data Mendelian randomization," *The Annals of statistics*, 49, 2079–2100.

- Zhao, Q., Small, D. S., and Bhattacharya, B. B. (2019), "Sensitivity analysis for inverse probability weighting estimators via the percentile bootstrap," Journal of the Royal Statistical Society Series B: Statistical Methodology, 81, 735–761.
- Zhao, Q., Wang, J., Hemani, G., Bowden, J., and Small, D. S. (2020), "Statistical inference in two-sample summary-data Mendelian randomization using robust adjusted profile score," *The Annals of Statistics*, 48, 1742 – 1769.