Linear-Quadratic Optimal Control for Mean-Field Stochastic Differential Equations in Infinite-Horizon with Regime Switching

Hongwei Mei^{*} Qingmeng Wei[†] and Jiongmin Yong[‡]

January 3, 2025

Abstract: This paper is concerned with stochastic linear quadratic (LQ, for short) optimal control problems in an infinite horizon with conditional mean-field term in a switching regime environment. The orthogonal decomposition introduced in [21] has been adopted. Desired algebraic Riccati equations (AREs, for short) and a system of backward stochastic differential equations (BSDEs, for short) in infinite time horizon with the coefficients depending on the Markov chain have been derived. The determination of closed-loop optimal strategy follows from the solvability of ARE and BSDE. Moreover, the solvability of BSDEs leads to a characterization of open-loop solvability of the optimal control problem.

Keywords: Linear-quadratic optimization problem, infinite-horizon, conditional mean-field, Markov switching, Riccati equations.

AMS Mathematics Subject Classification. 93E20, 49N10, 60F17.

1 Introduction

Let $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be a complete filtered probability space on which a one-dimensional standard Brownian motion $W(\cdot)$ is defined. On the same probability space, a Markov chain $\alpha(\cdot)$ is defined with a finite state space $\mathcal{M} = \{1, \dots, m_0\}$ and the generator $\Lambda = (\lambda_{\iota j})_{m_0 \times m_0}$, which is independent of $W(\cdot)$. Let \mathbb{F}^W and \mathbb{F}^{α} be the natural filtrations of $W(\cdot)$ and $\alpha(\cdot)$, augmented by all the \mathbb{P} -null sets, respectively. We define $\mathbb{F} = \mathbb{F}^W \vee \mathbb{F}^{\alpha}$ and denote the conditional expectation with respect to \mathbb{F}^{α} by $\mathbb{E}^{\alpha}_t[\xi] := \mathbb{E}[\xi|\mathcal{F}^{\alpha}_t]$.

We consider the following *n*-dimensional controlled mean-field stochastic differential equation (SDE, for short) with regime switching (governed by the Markov chain $\alpha(\cdot)$):

$$(1.1) \begin{cases} dX(t) = \left\{ A(\alpha(t))X(t) + \bar{A}(\alpha(t))\mathbb{E}_{t}^{\alpha}[X(t)] + B(\alpha(t))u(t) + \bar{B}(\alpha(t))\mathbb{E}_{t}^{\alpha}[u(t)] + b(t) \right\} dt \\ + \left\{ C(\alpha(t))X(t) + \bar{C}(\alpha(t))\mathbb{E}_{t}^{\alpha}[X(t)] + D(\alpha(t))u(t) + \bar{D}(\alpha(t))\mathbb{E}_{t}^{\alpha}[u(t)] + \sigma(t) \right\} dW(t), \quad t \ge s, \\ X(s) = \xi, \qquad \alpha(s) = \iota \in \mathcal{M}, \end{cases}$$

and the following cost functional

(1.2)
$$J^{\infty}(s,\iota,\xi;u(\cdot)) = \mathbb{E}\int_{s}^{\infty} f(t,\alpha(t),X(t),\mathbb{E}_{t}^{\alpha}[X(t)],u(t),\mathbb{E}_{t}^{\alpha}[u(t)])dt$$

^{*}Department of Mathematics and Statistics, Texas Tech University, Lubbock, TX 79409, USA; email: hongwei.mei@ttu.edu. This author is supported in part by Simons Travel Grant MP-TSM-00002835.

[†]School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China; email: weiqm100@nenu. edu.cn. This author is supported in part by the Natural Science Foundation of Jilin Province for Outstanding Young Talents (No. 20230101365JC), the National Key R&D Program of China (No. 2023YFA1009002), the National Natural Science Foundation of China (No. 12371443).

[‡]Department of Mathematics, University of Central Florida, Orlando, FL 32816, USA; email: jiongmin.yong@ucf.edu. This author is supported in part by NSF Grant DMS-2305475.

where

(1.3)
$$f(t,\iota,x,\bar{x},u,\bar{u}) = \frac{1}{2} \Big[\langle Q(\iota)x,x \rangle + 2\langle S(\iota)x,u \rangle + \langle R(\iota)u,u \rangle + \langle \bar{Q}(\iota)\bar{x},\bar{x} \rangle + 2\langle \bar{S}(\iota)\bar{x},\bar{u} \rangle + \langle \bar{R}(\iota)\bar{u},\bar{u} \rangle + 2\langle q(t),x \rangle + 2\langle \bar{q}(t),\bar{x} \rangle + 2\langle r(t),u \rangle + 2\langle \bar{r}(t),\bar{u} \rangle \Big].$$

Here all the coefficients $A(\cdot), \bar{A}(\cdot), B(\cdot), \bar{B}(\cdot), C(\cdot), \bar{C}(\cdot), D(\cdot), \bar{D}(\cdot)$ of the state equation (1.1) and the quadratic weight matrices $Q(\cdot), \bar{Q}(\cdot), S(\cdot), \bar{S}(\cdot), R(\cdot), \bar{R}(\cdot)$ of the cost functional (1.2) are some deterministic maps defined on \mathcal{M} ; the nonhomogeneous terms $b(\cdot), \sigma(\cdot)$ of the state equation (1.1) and the linear weighting terms $q(\cdot), r(\cdot)$ of the cost functional (1.2) are some integrable stochastic processes; and the linear mean-field weighting terms $\bar{q}(\cdot), \bar{r}(\cdot)$ of the cost functional (1.2) are some integrable stochastic processes on $[0, \infty)$. Due to $\alpha(\cdot)$ being a Markov chain, the coefficients (such as $A(\alpha(\cdot)))$ and weight functions (such as $Q(\alpha(\cdot)))$ are all of random. Our goal is to minimize the cost functional (1.2) subject to the state equation (1.1) over some set of admissible controls. Such a problem is referred to as a mean-field linear-quadratic optimal control problem (MF-LQ problem, for short), denoted by Problem (MF-LQ)^{∞}.

Linear quadratic (LQ, for short) problem has been extensively studied since the seminal works of Bellman-Glicksberg–Gross [2], Kalman [13] and Letov [16] appeared around 1960. There has been a vast amount of works on LQ control problems and their variations. Let us briefly mention a very small portion of the relevant works. For the classical theory of deterministic LQ problems, see Lee–Markus [15], Willems [36], Anderson-Moore [1], Wonham [38]. See also Bernhard [4], Zhang [40], Delfour [9], and Delfour-Sbarba [10] for a zero-sum differential game version. Study of stochastic LQ problems began with the works of Kushner [14] and Wonham [37] in the 1960s. See also McLane [20], Davis [8], Bensoussan [3], and so on, In 1998, Chen-Li-Zhou [5] found that for stochastic LQ problems, the weighting matrix of the control in the cost functional does not have to be positive definite, or even could be negative definite to some extent. This kind of problems have been termed to be *indefinite LQ problem* for convenience, initiating which trigged quite a few investigations. For stochastic coefficient case of such a problem, Chen-Yong [6, 7] studied the local solvability of backward stochastic differential Riccati equation. Tang [32, 33] solved the case of random coefficients with degenerate (positive semi-definite) control weight in the cost functional. The general theory for indefinite LQ problem with deterministic coefficients was completely established by Sun-Yong [28], Sun-Li-Yong [26] and Sun-Yong [29], under the framework of open and closed-loop solvability. See the book [30]. In 2021, Sun–Xiong–Yong [27] finally completely solved the general case of indefinite LQ problem with random coefficients. See Lü–Wang ([19]) for an infinite-dimensional version. In 2013, Yong [39] studied the problem with mean-field, followed by Huang-Li-Yong [12], Li-Sun-Yong [18], Sun [25], Wei-Yong-Yu [34], Sun-Yong [31], Li-Shi-Yong [17]. For Markov regime switching case, see Pham [22] and Zhang-Li-Xiong [41]. Moreover, the problem with coefficients being adapted to a martingale and with conditional mean-field interaction in a finite horizon has been studied by Mei–Wei–Yong [21]. This work can be regarded as a continuation of [21].

We now highlight the main features of the MF-LQ problem associated with (1.1) and (1.2), as well as the main contributions of the current paper.

(i) In order the cost functional associated with the state equation to be well-defined, we need to have a proper stabilizability of the state equation (1.1). Such a notion is technically new, due to two issues: (a) all the coefficients of the state equation and the quadratic weight functions in the cost functional depend on the Markov chain $\alpha(\cdot)$, so that they are all random in some special form, adapted to \mathbb{F}^{α} ; and (b) the state equation as well as the cost functional contains the conditional expectation terms $\mathbb{E}_t^{\alpha}[X(\cdot)]$ and $\mathbb{E}_t^{\alpha}[u(\cdot)]$. We will establish some reasonably easy-to-check conditions under which our Problem (MF-LQ)^{∞} is well-formulated.

(ii) Following the natural idea of completing squares, we derive the algebraic Riccati equations (AREs, for short) and the corresponding system of backward stochastic differential equations (BSDEs, for short) in an infinite horizon with the coefficients depending on the Markov chain. The well-posedness of AREs and BSDEs will determine the closed-loop optimal strategy.

(iii) For AREs, we use the results of the finite horizon LQ problem from [21]. Under the stabilizability condition, we are able to pass to the limits to get the so-called stabilizing solutions of AREs. For BSDEs, due to the fact that the stabilizability of our state equation is weaker than the constant coefficient system,

the result found in [30] cannot be applied directly. A new method is introduced to prove the well-posedness of such BSDEs.

(vi) Besides the Problem $(MF-LQ)^{\infty}$ to be cosed-loop solvable once the well-posedness of AREs and BSDEs have been established. We also obtained a characterization of open-loop solvability of the problem in terms of a forward-backward system, with the coupling by the stationarity conditions.

The rest of the paper is arranged as follows. In Section 2, some preliminaries are presented including the martingale measure and orthogonal decomposition. Stabilizability related notions and results are presented in Section 3. Some relevant notions and several immediate results are presented in Section 4. Closed-loop solvability results are proved in Section 5. In particular, Subsection 5.1 is devoted to the completing squares method. The solvability of AREs and BSDEs are carried out in Subsections 5.2 and 5.3, respectively. In Section 6, an open-loop solvability result is presented. Finally, some concluding remarks are made in Section 7.

2 Preliminaries

Throughout this paper, besides the notations introduced in the previous section, let \mathbb{S}^k (\mathbb{S}^k_+ , \mathbb{S}^k_{++}) be the set of all the $k \times k$ symmetric (positive semi-definite, positive definite) matrices, I be the identity matrix or operator in a suitable space, M^{\top} and $\mathscr{R}(M)$ be the transpose and range of a matrix M, respectively. We write M > N ($M \ge N$, resp.), meaning that M - N is positive definite (positive semi-definite, resp.).

For Euclidean space \mathbb{H} and $0 < T \leq \infty$, we define

Naturally, replacing \mathcal{F}_s , \mathbb{F} of $L^2_{\mathcal{F}_s}(\Omega; \mathbb{H})$, $L^2_{\mathbb{F}}(s, T; \mathbb{H})$ by \mathcal{F}^{α}_s , \mathbb{F}^{α} , we recognize the new spaces $L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{H})$, $L^2_{\mathbb{F}^{\alpha}}(s, T; \mathbb{H})$. Furthermore, the spaces $L^2_{\mathbb{F}_-}(s, T; \mathbb{H})$, $L^2_{\mathbb{F}^{\alpha}_-}(s, T; \mathbb{H})$ can also be identified by rewriting the measurability in $L^2_{\mathbb{F}}(s, T; \mathbb{H})$, $L^2_{\mathbb{F}^{\alpha}}(s, T; \mathbb{H})$ as \mathbb{F} -predictable and \mathbb{F}^{α} -predictable, resp.

Any $(s, \iota, \xi) \in \mathscr{D}$ is called an *admissible initial triple*. Also, we let

$$\mathscr{U}[s,\infty) = L^2_{\mathbb{F}}(s,\infty;\mathbb{R}^m), \qquad \mathscr{U}[s,T] = L^2_{\mathbb{F}}(s,T;\mathbb{R}^m).$$

Any $u(\cdot) \in \mathscr{U}[s, \infty)$ is called a *feasible control* on $[s, \infty)$ and any $u(\cdot) \in \mathscr{U}[s, T]$ is called a *feasible control* on [s, T]. We now introduce the following hypotheses.

- (A1) $A(\cdot), \bar{A}(\cdot), C(\cdot), \bar{C}(\cdot) : \mathcal{M} \to \mathbb{R}^{n \times n}, B(\cdot), \bar{B}(\cdot), D(\cdot), \bar{D}(\cdot) : \mathcal{M} \mapsto \mathbb{R}^{n \times m}.$
- (A2) $b(\cdot), \sigma(\cdot), q(\cdot) \in L^2_{\mathbb{F}}(0,\infty;\mathbb{R}^n), \bar{q}(\cdot) \in L^2_{\mathbb{F}^\alpha}(0,\infty;\mathbb{R}^n), r(\cdot) \in L^2_{\mathbb{F}}(0,\infty;\mathbb{R}^m), \bar{r}(\cdot) \in L^2_{\mathbb{F}^\alpha}(0,\infty;\mathbb{R}^m).$

(A3) The following hold: $Q(\cdot), \bar{Q}(\cdot) : \mathcal{M} \to \mathbb{S}^n_{++}, R(\cdot), \bar{R}(\cdot) : \mathcal{M} \to \mathbb{S}^m_{++}, S(\cdot), \bar{S}(\cdot) : \mathcal{M} \to \mathbb{R}^{m \times n}$, and for each $i \in \mathcal{M}$,

(2.1)
$$Q(i) - S(i)^{\top} R(i)^{-1} S(i) \in \mathbb{S}_{++}^{n}, Q(i) + \bar{Q}(i) - [S(i) + \bar{S}(i)]^{\top} [R(i) + \bar{R}(i)]^{-1} [S(i) + \bar{S}(i)] \in \mathbb{S}_{++}^{n}$$

Clearly, under (A1), all the coefficients of the state equation (1.1) are bounded. Thus, for any admissible initial triple $(s, \iota, \xi) \in \mathscr{D}$ and any feasible control $u(\cdot) \in \mathscr{U}[s, \infty)$, the state equation (1.1) admits a unique solution $X(\cdot) = X(\cdot; s, \iota, \xi, u(\cdot)) \in L^2_{\mathbb{F}}(s, T; \mathbb{R}^n)$ for any T > 0, but might not be in $L^2_{\mathbb{F}}(s, \infty; \mathbb{R}^n)$. Thus, $J^{\infty}(s, \iota, \xi; u(\cdot))$ might not be well-defined or finite under (A1)–(A2). We therefore introduce the set $\mathscr{U}_{ad}[s, \infty)$ as follows:

$$\mathscr{U}_{ad}[s,\infty) = \Big\{ u(\cdot) \in \mathscr{U}[0,\infty) \mid X(\cdot;s,\imath,\xi,u(\cdot)) \in L^2_{\mathbb{F}}(0,\infty;\mathbb{R}^n), \ \forall (s,\imath,\xi) \in \mathscr{D} \Big\}.$$

We will show later that $\mathcal{U}_{ad}[s,\infty) \neq \emptyset$ under a certain condition. Now, our LQ problem can be stated as follows.

Problem (MF-LQ)^{∞}. For $(s, i, \xi) \in \mathscr{D}$, find $\bar{u}(\cdot) \in \mathscr{U}_{ad}[s, \infty)$ such that

(2.2)
$$J^{\infty}(s,\iota,\xi;\bar{u}(\cdot)) = \inf_{u(\cdot)\in\mathscr{U}_{ad}[s,\infty)} J^{\infty}(s,\iota,\xi;u(\cdot)) \equiv V^{\iota}(s,\xi), \qquad \text{inf } \varnothing \equiv \infty.$$

Any $\bar{u}(\cdot) \in \mathscr{U}_{ad}[s, \infty)$ satisfying (2.2) is called an *open-loop optimal control*, the corresponding state process $\bar{X}(\cdot) = X(\cdot; s, i, \xi, \bar{u}(\cdot))$ is called an *open-loop optimal state process*, and $(\bar{X}(\cdot), \bar{u}(\cdot))$ is called an *open-loop optimal pair*. In this case, we say the above problem to be *open-loop solvable*.

In the case that $b(\cdot), \sigma(\cdot), q(\cdot), \bar{q}(\cdot), r(\cdot), \bar{r}(\cdot)$ are zero, the state equation (1.1) becomes

$$(2.3) \qquad \begin{cases} dX(t) = \left\{ A(\alpha(t))X(t) + \bar{A}(\alpha(t))\mathbb{E}_{t}^{\alpha}[X(t)] + B(\alpha(t))u(t) + \bar{B}(\alpha(t))\mathbb{E}_{t}^{\alpha}[u(t)] \right\} dt \\ + \left\{ C(\alpha(t))X(t) + \bar{C}(\alpha(t))\mathbb{E}_{t}^{\alpha}[X(t)] + D(\alpha(t))u(t) + \bar{D}(\alpha(t))\mathbb{E}_{t}^{\alpha}[u(t)] \right\} dW(t), \ t \ge s, \\ X(s) = \xi \in L^{2}_{\mathcal{F}_{s}}(\Omega; \mathbb{R}^{n}), \quad \alpha(s) = \iota \in \mathcal{M}, \end{cases}$$

which is called a *homogeneous system* and the cost functional becomes

(2.4)
$$J_0^{\infty}(s,\iota,\xi;u(\cdot)) = \mathbb{E}\int_s^{\infty} f_0(\alpha(t),X(t),\mathbb{E}_t^{\alpha}[X(t)],u(t),\mathbb{E}_t^{\alpha}[u(t)])dt$$

where

$$(2.5) \qquad f_0(\iota, x, \bar{x}, u, \bar{u}) = \frac{1}{2} \Big[\langle Q(\iota)x, x \rangle + 2 \langle S(\iota)x, u \rangle + \langle R(\iota)u, u \rangle + \langle \bar{Q}(\iota)\bar{x}, \bar{x} \rangle + 2 \langle \bar{S}(\iota)\bar{x}, \bar{u} \rangle + \langle \bar{R}(\iota)\bar{u}, \bar{u} \rangle \Big].$$

We call (2.4) a *purely quadratic* cost functional. For the case of (2.3) with (2.4), the corresponding LQ problem is said to be *homogeneous*, denoted by Problem (MF-LQ)₀^{∞}. In what follows, for convenience, we will denote the *homogeneous system* (2.3) by $[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$.

2.1 Orthogonal Decomposition

To achieve the stabilizability condition, we adopt the orthogonal decomposition constructed in [21] to derive an equivalent formulation for Problem $(MF-LQ)^{\infty}$.

For any $s \in [0,T), L^2_{\mathcal{F}_s}(\Omega;\mathbb{H})$ is a Hilbert space under the following inner product

$$\mathbb{E}\langle \xi, \eta \rangle \equiv \int_{\Omega} \langle \xi(\omega), \eta(\omega) \rangle d\mathbb{P}(\omega), \qquad \xi, \eta \in L^{2}_{\mathcal{F}_{s}}(\Omega; \mathbb{H})$$

The space $L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{H})$, with the same inner product as above, is a closed subspace of $L^2_{\mathcal{F}_s}(\Omega; \mathbb{H})$ and its orthogonal complement in $L^2_{\mathcal{F}_s}(\Omega; \mathbb{H})$ is given by

$$L^{2}_{\mathcal{F}^{\alpha}_{s}}(\Omega;\mathbb{H})^{\perp} := \left\{ \xi \in L^{2}_{\mathcal{F}_{s}}(\Omega;\mathbb{H}) \mid \mathbb{E}\langle \xi,\eta\rangle = 0, \quad \forall \eta \in L^{2}_{\mathcal{F}^{\alpha}_{s}}(\Omega;\mathbb{H}) \right\}.$$

For any $\xi \in L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{H})$, define

$$\Pi_s[\xi] := \mathbb{E}[\xi|\mathcal{F}_s^\alpha]$$

Similar to [21], we see that Π_s induces the following orthogonal decomposition:

$$\xi = \Pi_s[\xi] + (\xi - \Pi_s[\xi]) \in L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{H}) \oplus L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{H})^{\perp}$$

On the other hand, thanks to Lemma A.1 in [21], we have $\mathbb{E}_s^{\alpha}[\cdot] = \mathbb{E}^{\alpha}[\cdot] \equiv \mathbb{E}[\cdot|\mathcal{F}_{\infty}^{\alpha}]$. Therefore,

(2.6)
$$\Pi_s[\xi] = \mathbb{E}_s^{\alpha}[\xi] = \mathbb{E}^{\alpha}[\xi], \qquad \Pi_s^{\perp}[\xi] = \xi - \mathbb{E}_s^{\alpha}[\xi] = \xi - \mathbb{E}^{\alpha}[\xi], \qquad \forall \xi \in L^2_{\mathcal{F}_s}(\Omega; \mathbb{H}).$$

Based on the above, for any $0 \leq s < T$, we further define $\Pi : L^2_{\mathbb{F}}(s,T;\mathbb{H}) \to L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H})$ as follows:

(2.7)
$$\Pi[v(\cdot)](t) = \Pi_t[v(t)] \equiv \mathbb{E}_t^{\alpha}[v(t)] = \mathbb{E}^{\alpha}[v(t)], \quad \text{a.e. } t \in [s, T], \ \forall v(\cdot) \in L^2_{\mathbb{F}}(s, T; \mathbb{H}).$$

Note that for any $v(\cdot) \in L^2(s, T; \mathbb{H})$, $\Pi[v(\cdot)](t)$ is only defined for almost all $t \in [s, T]$, as a process on [s, T]. We now show that Π is the orthogonal projection from $L^2_{\mathbb{F}}(s, T; \mathbb{H})$ onto $L^2_{\mathbb{F}^{\alpha}}(s, T; \mathbb{H})$. In fact, first of all, if $v(\cdot) = \bar{v}(\cdot)$ in $L^2_{\mathbb{F}}(s, T; \mathbb{H})$, we get

$$\mathbb{E}\int_{s}^{T}\left|\Pi[v(\cdot)](t) - \Pi[\bar{v}(\cdot)](t)\right|^{2}dt \leq \mathbb{E}\int_{s}^{T}\left|v(t) - \bar{v}(t)\right|^{2}dt = 0,$$

which leads to $\Pi[v(\cdot)] = \Pi[\bar{v}(\cdot)]$ in $L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H})$. This means that Π is well-defined. Clearly, $\Pi^2 = \Pi$ and

$$\begin{split} \langle \Pi[v(\cdot)], \bar{v}(\cdot) \rangle &= \mathbb{E} \int_{s}^{T} \langle \mathbb{E}^{\alpha}[v(t)], \bar{v}(t) \rangle dt = \mathbb{E} \int_{s}^{T} \langle \mathbb{E}^{\alpha}[v(t)], \mathbb{E}^{\alpha}[\bar{v}(t)] \rangle dt \\ &= \mathbb{E} \int_{s}^{T} \langle v(t), \mathbb{E}^{\alpha}[\bar{v}(t)] \rangle dt = \langle v(\cdot), \Pi[\bar{v}(\cdot)] \rangle. \end{split}$$

Thus, Π is a self-adjoint idempotent, which means that Π is an orthogonal projection from $L^2_{\mathbb{F}}(s,T;\mathbb{H})$ onto $L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H})$. Next, we denote $\Pi^{\perp} := I - \Pi$, which is the orthogonal projection from $L^2_{\mathbb{F}}(s,T;\mathbb{H})$ onto $L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H})^{\perp}$, where

$$L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H})^{\perp} := \Big\{ v(\cdot) \in L^2_{\mathbb{F}}(s,T;\mathbb{H}) \mid \mathbb{E} \int_s^T \langle v(t), \bar{v}(t) \rangle dt = 0, \quad \forall \bar{v}(\cdot) \in L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H}) \Big\}.$$

Also

$$L^2_{\mathbb{F}^{\alpha}_{-}}(s,T;\mathbb{H})^{\perp} := \Big\{ v(\cdot) \in L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H})^{\perp} \mid v(\cdot) \text{ is } \mathbb{F}\text{-predictable } \Big\} = \Pi^{\perp} \Big(L^2_{\mathbb{F}_{-}}(s,T;\mathbb{H}) \Big).$$

From the above, we have the following orthogonal decompositions:

$$\begin{cases} L^2_{\mathbb{F}}(s,T;\mathbb{H}) = L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H})^{\perp} \oplus L^2_{\mathbb{F}^{\alpha}}(s,T;\mathbb{H}),\\ L^2_{\mathbb{F}_{-}}(s,T;\mathbb{H}) = L^2_{\mathbb{F}^{\alpha}_{-}}(s,T;\mathbb{H})^{\perp} \oplus L^2_{\mathbb{F}^{\alpha}_{-}}(s,T;\mathbb{H}),\\ M^2_{\mathbb{F}_{-}}(s,T;\mathbb{H}) = M^2_{\mathbb{F}^{\alpha}_{-}}(s,T;\mathbb{H})^{\perp} \oplus M^2_{\mathbb{F}^{\alpha}_{-}}(s,T;\mathbb{H}). \end{cases}$$

Here $M^2_{\mathbb{F}_-}(s,T;\mathbb{H})$ will be defined later in (2.17). More details on the above orthogonal decompositions can be found in [21]. We also note that all above arguments are valid for $T = \infty$. In the sequel, we will write

$$\Pi_1 = \Pi^\perp, \qquad \Pi_2 = \Pi.$$

2.2 An Equivalent Formulation of Problem $(MF-LQ)^{\infty}$

Denote $\theta_i = \prod_i [\theta]$ for $\theta = X, u, b, \sigma, \xi$. Applying the orthogonal projection \prod_2 to state equation (1.1), then subtracting from (1.1) leads to

$$(2.8) \begin{cases} dX_1(t) = \left\{ A_1(\alpha(t))X_1(t) + B_1(\alpha(t))u_1(t) + b_1(t) \right\} dt \\ + \left\{ C_1(\alpha(t))X_1(t) + C_2(\alpha(t))X_2(t) + D_1(\alpha(t))u_1(t) + D_2(\alpha(t))u_2(t) + \sigma(t) \right\} dW(t), \\ dX_2(t) = \left\{ A_2(\alpha(t))X_2(t) + B_2(\alpha(t))u_2(t) + b_2(t) \right\} dt, \quad t \in [s, \infty), \\ X_1(s) = \xi_1, \quad X_2(s) = \xi_2, \quad \alpha(s) = i. \end{cases}$$

Here

(2.9)
$$A_1(i) := A(i), \quad A_2(i) := A(i) + \bar{A}(i), \quad B_1(i) := B(i), \quad B_2(i) := B(i) + \bar{B}(i), \\ C_1(i) := C(i), \quad C_2(i) := C(i) + \bar{C}(i), \quad D_1(i) := D(i), \quad D_2(i) := D(i) + \bar{D}(i).$$

At the same time, the cost functional (1.2) can be (provided the integrals exist) written as

$$(2.10) \qquad J^{\infty}(s,\iota,\xi_{1},\xi_{2};u_{1}(\cdot),u_{2}(\cdot)) := J^{\infty}(s,\iota,\xi;u(\cdot))$$
$$= \frac{1}{2}\sum_{k=1}^{2} \mathbb{E} \int_{s}^{\infty} \left[\langle Q_{k}(\alpha(t))X_{k}(t),X_{k}(t)\rangle + 2\langle S_{k}(\alpha(t))X_{k}(t),u_{k}(t)\rangle + \langle R_{k}(\alpha(t))u_{k}(t),u_{k}(t)\rangle + 2\langle q_{k}(t),X_{k}(t)\rangle + 2\langle r_{k}(t),u_{k}(t)\rangle \right] dt,$$

with

(2.11)

$$Q_{1}(i) := Q(i), \quad Q_{2}(i) := Q(i) + \bar{Q}(i), \quad S_{1}(i) := S(i), \quad S_{2}(i) := S(i) + \bar{S}(i), \\
R_{1}(i) := R(i), \quad R_{2}(i) := R(i) + \bar{R}(i), \\
q_{1}(t) = q(t) - \Pi[q(t)], \quad q_{2}(t) = \Pi[q(t) + \bar{q}(t)], \\
r_{1}(t) = r(t) - \Pi[r(t)], \quad r_{2}(t) = \Pi[r(t) + \bar{r}(t)].$$

According to the above, Problem (MF-LQ)^{∞} can be equivalently formulated with the state equation (2.8) and the cost functional (2.10).

Now, for the homogeneous case, i.e., $b(\cdot), \sigma(\cdot), q(\cdot), \bar{q}(\cdot), r(\cdot), \bar{r}(\cdot)$ are all zero, (2.3) becomes

(2.12)
$$\begin{cases} dX_1(t) = \left\{ A_1(\alpha(t))X_1(t) + B_1(\alpha(t))u_1(t) \right\} dt \\ + \left\{ C_1(\alpha(t))X_1(t) + C_2(\alpha(t))X_2 + D_1(\alpha(t))u_1(t) + D_2(\alpha(t))u_2(t) \right\} dW(t), \\ dX_2(t) = \left\{ A_2(\alpha(t))X_2(t) + B_2(\alpha(t))u_2(t) \right\} dt, \quad t \in [s, \infty), \\ X_1(s) = \xi_1, \quad X_2(s) = \xi_2, \quad \alpha(s) = i, \end{cases}$$

and the cost functional (2.4) becomes (again, if the integrals exist)

$$(2.13) \qquad \begin{aligned} J_0^{\infty}(s,\iota,\xi_1,\xi_2;u_1(\cdot),u_2(\cdot)) &:= J_0^{\infty}(s,\iota,\xi;u(\cdot)) \\ &= \frac{1}{2} \sum_{k=1}^2 \mathbb{E} \int_s^{\infty} \Big[\langle Q_k(\alpha(t))X_k(t),X_k(t) \rangle + 2\langle S_k(\alpha(t))X_k(t),u_k(t) \rangle + \langle R_k(\alpha(t))u_k(t),u_k(t) \rangle \Big] dt. \end{aligned}$$

For convenience, we denote $[A_1, C_1, C_2; B_1, D_1, D_2]_1$ for the system (2.12) of $X_1(\cdot)$. Similarly, we denote $[A_2, B_2]_2$ the system (2.12) for $X_2(\cdot)$. The whole system is denoted by $\{[A_1, C_1, C_2; B_1, D_1, D_2]_1; [A_2, B_2]_2\}$. The corresponding LQ problem is denoted by Problem (MF-LQ)₀[∞].

Further, if the system is uncontrolled, then it becomes

(2.14)
$$\begin{cases} dX^{0}(t) = \left(A(\alpha(t))X^{0}(t) + \bar{A}(\alpha(t))\mathbb{E}_{t}^{\alpha}[X^{0}(t)]\right)dt + \left(C(\alpha(t))X^{0}(t) + \bar{C}(\alpha(t))\mathbb{E}_{t}^{\alpha}[X^{0}(t)]\right)dW(t), \\ t \in [s, \infty), \\ X^{0}(s) = \xi, \quad \alpha(s) = \iota, \end{cases}$$

whose solution is $X^0(\cdot) \equiv X^0(\cdot; s, \xi, i)$. We denote such a system simply by $[A, \overline{A}, C, \overline{C}] \equiv [A, \overline{A}, C, \overline{C}; 0, 0, 0, 0]$. Under our orthogonal decomposition, the above can also be written as

(2.15)
$$\begin{cases} dX_1^0(t) = A_1(\alpha(t))X_1^0(t)dt + [C_1(\alpha(t))X_1^0(t) + C_2(\alpha(t))X_2^0]dW(t), \\ dX_2^0(t) = A_2(\alpha(t))X_2^0(t)dt, \quad t \in [s,\infty), \\ X_1^0(s) = \xi_1, \quad X_2^0(s) = \xi_2, \quad \alpha(s) = \iota, \end{cases}$$

whose solution is $(X_1^0(\cdot;\xi_1,\xi_2,i), X_2^0(\cdot;s,\xi_1,\xi_2,i))$, where $A_k(\cdot), C_k(\cdot)$ (k = 1, 2) are given by (2.9). According to our notation, such a system is denoted by $\{[A_1, C_1, C_2, 0, 0, 0]_1; [A_2, 0]_2\} \equiv \{A_1, A_2, C_1, C_2\}$.

Remark 2.1. Note that condition (2.1) in (A3) can be written as

(2.16)
$$Q_k(i) - S_k(i)R_k(i)^{-1}S_k(i) \in \mathbb{S}_{++}^n, \quad i \in \mathcal{M}, \quad k = 1, 2.$$

Then the cost functional for Problem (MF-LQ)_0^{\infty} is uniformly convex on $\mathscr{U}_{ad}[s,\infty)$, i.e.

$$J_0^{\infty}(s,\iota,\xi_1,\xi_2;u_1(\cdot),u_2(\cdot)) \geqslant \varepsilon \sum_{k=1}^2 \mathbb{E} \int_s^{\infty} |u_k(t)|^2 dt, \qquad \forall u(\cdot) \in \mathscr{U}_{ad}[s,\infty),$$

for some $\varepsilon > 0$. Consequently, Problem (MF-LQ)^{∞} admits a unique open-loop optimal control in $\mathscr{U}_{ad}[s, \infty)$. One of the main focuses in this paper is to find a feedback representation of the open-loop optimal control.

2.3 Martingale Measure of $\alpha(\cdot)$

In this subsection, we will construct a martingale associated with the Markov chain. Such a martingale will be used in the BSDEs in the future.

Recall that $\mathcal{M} = \{1, \dots, m_0\}$, and $\alpha : [0, \infty) \times \Omega \to \mathcal{M}$ is a continuous-time Markov chain with the *transition probability*:

$$p_{\imath \jmath}(t,s) = \mathbb{P}\big(\alpha(t) = \jmath \mid \alpha(s) = \imath\big), \qquad \imath, \jmath \in \mathcal{M}, \quad 0 \leqslant s < t.$$

We assume that $\alpha(\cdot)$ is homogeneous, i.e., $p_{ij}(t,s) = p_{ij}(t-s)$, where

$$p_{ij}(r) = \mathbb{P}(\alpha(r) = j \mid \alpha(0) = i), \quad i, j \in \mathcal{M}, \quad r \ge 0.$$

Denote its *transition probability matrix* as follows:

$$\mathbf{P}(r) = \left(p_{ij}(r)\right)_{m_0 \times m_0}, \qquad r \ge 0$$

For any $j \in \mathcal{M} \setminus \{i\}$, let

$$\tau_{ij}^s = \inf\{r > 0 \mid \alpha(s+r) = j, \ \alpha(s-) = i\}$$

which follows an exponential distribution, i.e.,

$$\mathbb{P}(\tau_{ij}^s > r) = \int_r^\infty \lambda_{ij} e^{-\lambda_{ij}\theta} d\theta = e^{-\lambda_{ij}r}, \qquad r > 0, \quad j \neq i,$$

where λ_{ij} satisfies the so-called *q*-property:

$$\lambda_{ij} > 0, \quad i \neq j, \qquad \sum_{j=1}^{m_0} \lambda_{ij} = 0,$$

and we refer to $\mathbf{\Lambda} = (\lambda_{ij})$ as the generator of the Markov chain $\alpha(\cdot)$. Since $\alpha(\cdot)$ is stationary, one has that (for $0 \leq s < t < \infty$)

$$p_{ij}(t-s) = p_{ij}(s,t) = \delta_{ij} + \lambda_{ij}(t-s) + o(t-s) = \begin{cases} \lambda_{ij}(t-s) + o(t-s), & j \neq i, \\ 1 + \lambda_{ii}(t-s) + o(t-s), & j = i, \end{cases}$$

Next, for $i \neq j$, we define

$$\widetilde{M}_{ij}(t) = \sum_{0 < s \leq t} \mathbf{1}_{[\alpha(s_{-})=i]} \mathbf{1}_{[\alpha(s)=j]} \equiv \text{accumulative jump number from } i \text{ to } j \text{ in } (0, \langle \widetilde{M}_{ij} \rangle(t) = \int_{0}^{t} \lambda_{ij} \mathbf{1}_{[\alpha(s_{-})=i]} ds, \quad M_{ij}(t) = \widetilde{M}_{ij}(t) - \langle \widetilde{M}_{ij} \rangle(t), \qquad t \geq 0.$$

t],

Then, by [23], p.35, (21.12) Lemma (see also [11, 24]), $M_{ij}(\cdot)$ is a purely discontinuous and square-integrable martingale (with respect to \mathbb{F}^{α}). For convenience, we let

$$M_{ii}(t) = M_{ii}(t) = \langle M_{ii} \rangle(t) = 0, \qquad t \ge 0.$$

We call $\{M_{ij}(\cdot) \mid i, j \in \mathcal{M}\}$ the martingale measure of Markov chain $\alpha(\cdot)$.

Now we want to define the stochastic integral with respect to such a martingale measure. Introduce the following Hilbert spaces

$$(2.17) \qquad \begin{cases} M_{\mathbb{F}_{-}}^{2}(s,T;\mathbb{H}) = \left\{\varphi(\cdot) = (\varphi(\cdot,1),\cdots,\varphi(\cdot,m_{0})) \mid \varphi(\cdot,\imath) \text{ is } \mathbb{H}\text{-valued and } \mathbb{F}\text{-predictable} \\ \text{with } \sum_{\iota\neq\jmath} \mathbb{E}\int_{s}^{T} |\varphi(r,\jmath)|^{2} d\widetilde{M}_{\imath\jmath}(r) < \infty, \quad \forall \jmath \in \mathcal{M} \right\}, \\ M_{\mathbb{F}_{-}^{\alpha}}^{2}(s,T;\mathbb{H}) = \left\{\varphi(\cdot) \in M_{\mathbb{F}_{-}}^{2}(s,T;\mathbb{H}) \mid \varphi(\cdot) \text{ is } \mathbb{F}^{\alpha}\text{-predictable} \right\}. \end{cases}$$

It can be seen that $M^2_{\mathbb{F}^{\alpha}_{-}}(s,T;\mathbb{H}) \subset M^2_{\mathbb{F}_{-}}(s,T;\mathbb{H})$. For any $\varphi(\cdot) \in M^2_{\mathbb{F}_{-}}(s,T;\mathbb{H})$, we write

$$\int_{s}^{t} \varphi(r) dM(r) := \sum_{i \neq j} \int_{s}^{t} \varphi(r, j) \mathbf{1}_{[\alpha(r^{-})=i]} dM_{ij}(r),$$

which is a (local) martingale with quadratic variation

$$\mathbb{E}\Big(\int_{s}^{t}\varphi(r)dM(r)\Big)^{2} = \mathbb{E}\int_{s}^{t}\sum_{i\neq j}|\varphi(r,j)|^{2}\lambda_{ij}\mathbf{1}_{[\alpha(r)=i]}dr.$$

Note that for any map $\Sigma : \mathcal{M} \to \mathbb{H}$, we have

$$d\Sigma(\alpha(t)) = \sum_{\substack{j \neq \alpha(t^{-})}} \left([\Sigma(j) - \Sigma(\alpha(t^{-}))] d\widetilde{M}_{\alpha(t^{-})j} \right)$$

(2.18)
$$= \sum_{\substack{j \in \mathcal{M}}} \lambda_{\alpha(t^{-})j} \Sigma(j) dt + \sum_{\substack{j \in \mathcal{M}}} \left(\Sigma(j) - \Sigma(\alpha(t^{-})) \right) \left(d\widetilde{M}_{\alpha(t^{-})j} - \lambda_{\alpha(t^{-})j} dt \right)$$

$$\equiv \Lambda[\Sigma](\alpha(t)) dt + \sum_{\substack{j \in \mathcal{M}}} \left(\Sigma(j) - \Sigma(\alpha(t^{-})) \right) dM_{\alpha(t^{-})j},$$

where

(2.19)
$$\Lambda[\Sigma](i) = \sum_{j \in \mathcal{M}} \lambda_{ij} \Sigma(j), \qquad i \in \mathcal{M}.$$

Next, we also have the following product rule: for any maps $\Sigma_1, \Sigma_2 : \mathcal{M} \to \mathbb{H}$, by (2.18),

$$\begin{split} d\Big[\Sigma_1(\alpha(t))\Sigma_2(\alpha(t))\Big] &= \Big(d\Sigma_1(\alpha(t))\Big)\Sigma_2(\alpha(t)) + \Sigma_1(\alpha(t))d\Sigma_2(\alpha(t)) + \langle d\Sigma_1(\alpha(t)), d\Sigma_2(\alpha(t))\rangle \\ &= \Big[\Lambda[\Sigma_1](\alpha(t))\Sigma_2(\alpha(t)) + \Sigma_1(\alpha(t))\Lambda[\Sigma_2](\alpha(t)) + \sum_{j\in\mathcal{M}}\lambda_{\alpha(t)j}\Big(\Sigma_1(j) - \Sigma_1(\alpha(t))\Big)\Big(\Sigma_2(j) - \Sigma_2(\alpha(t))\Big)\Big]dt \\ &+ \sum_{j\in\mathcal{M}}\Big[\Big(\Sigma_1(j) - \Sigma_1(\alpha(t))\Big)\Sigma_2(\alpha(t)) + \Sigma_1(\alpha(t))\Big(\Sigma_2(j) - \Sigma_2(\alpha(t))\Big)\Big] \\ &+ \Big(\Sigma_1(j) - \Sigma_1(\alpha(t))\Big)\Big(\Sigma_2(j) - \Sigma_2(\alpha(t))\Big)\Big]dM_{\alpha(t^-)j}(t). \end{split}$$

3 Stability and Stabilizability

In this section, we consider the long-term behavior of the state process $X(\cdot)$ of the controlled state equation so that Problem (MF-LQ)^{∞} is well-defined. We first consider the uncontrolled homogeneous state equation (2.14) or (2.15). The following definitions will be necessary.

Definition 3.1. (i) System $[A, \overline{A}, C, \overline{C}]$ is said to be L^2 -exponentially stable if there are constants $K, \delta > 0$ such that for any $(s, \xi, i) \in \mathcal{D}$, the solution of (2.14) satisfies

(3.1)
$$\mathbb{E}|X^0(t;s,\xi,i)|^2 \leqslant Ke^{-\delta(t-s)}\mathbb{E}|\xi|^2, \qquad t \ge s,$$

or equivalently, system $\{A_1, A_2, C_1, C_2\}$ is said to be L^2 -exponentially stable, if there are $K, \delta > 0$ such that for any $(s, \xi_1, \xi_2, i) \in \mathcal{D}$, the solution of (2.15) satisfies

(3.2)
$$\sum_{k=1}^{2} |X_k^0(t; s, \xi_1, \xi_2, \imath)|^2 \leqslant K e^{-\delta(t-s)} \sum_{k=1}^{2} \mathbb{E} |\xi_k|^2, \qquad t \ge s.$$

(ii) System $[A, \overline{A}, C, \overline{C}]$ is said to be L^2 -integrable if for any $(s, \xi, i) \in \mathscr{D}$, the solution of (2.14) satisfies

(3.3)
$$X^{0}(\cdot; s, \xi, i) \in L^{2}_{\mathbb{F}}(s, \infty; \mathbb{R}^{n}),$$

or equivalently, system $\{A_1, A_2, C_1, C_2\}$ is said to be L^2 -integrable, if for any $(s, \xi_1, \xi_2, i) \in \mathcal{D}$, the solution of (2.15) satisfies

(3.4)
$$X_1^0(\cdot; s, \xi_1, \xi_2, i) \in L^2_{\mathbb{F}^\alpha}(s, \infty; \mathbb{R}^n)^\perp, \qquad X_2^0(\cdot; s, \xi_1, \xi_2, i) \in L^2_{\mathbb{F}^\alpha}(s, \infty; \mathbb{R}^n).$$

(iii) System $[A, \overline{A}, C, \overline{C}]$ is said to be *dissipative* if there exists a measurable function $P : \mathcal{M} \to \mathbb{S}^n_{++}$ such that for some $\varepsilon > 0$,

$$(3.5) \qquad \mathbb{E}\Big[\langle P(\alpha(t))X^{0}(t), X^{0}(t)\rangle\Big] \leqslant \mathbb{E}\Big[\langle P(\alpha(s))X^{0}(s), X^{0}(s)\rangle - \varepsilon \mathbb{E}\int_{s}^{t} \langle P(\alpha(r))X^{0}(r), X^{0}(r)\rangle dr\Big],\\ \forall 0 \leqslant s < t < \infty,$$

for any solution $X^0(\cdot)$ of (2.14); or equivalently, system $\{A_1, A_2, C_1, C_2\}$ is said to be *dissipative*, if there exist measurable functions $P_1, P_2 : \mathcal{M} \to \mathbb{S}^n_{++}$ such that for some $\varepsilon > 0$,

$$(3.6) \qquad \mathbb{E}\Big[\langle P_k(\alpha(t))X_k^0(t), X_k^0(t)\rangle\Big] \leqslant \mathbb{E}\Big[\langle P_k(\alpha(s))X_k^0(s), X_k^0(s)\rangle - \varepsilon \int_s^t \langle P_k(\alpha(r))X_k^0(r), X_k^0(r)\rangle dr\Big],\\ 0 \leqslant s < t < \infty, \quad k = 1, 2,$$

for any solution $(X_1^0(\cdot), X_2^0(\cdot))$ of (2.15).

Note that (3.5) roughly means the map $t \mapsto \mathbb{E}\langle P(\alpha(t))X^0(t), X^0(t)\rangle$ has a negative time-derivative if $X^0(t) \neq 0$. This amounts to saying that the quantity $\mathbb{E}\langle P(\alpha(t))X^0(t), X^0(t)\rangle$ is decreasing/dissipative as a map of t.

Next, we introduce the following system of algebraic Lyapunov-type inequalities:

(3.7)
$$\Lambda[P_1(\cdot)](i) + A_1(i)^\top P_1(i) + P_1(i)A_1(i) + C_1(i)^\top P_1(i)C_1(i) < 0, \Lambda[P_2(\cdot)](i) + A_2(i)^\top P_2(i) + P_2(i)A_2(i) < 0, \qquad i \in \mathcal{M}.$$

Here $A_i(\cdot), C_i(\cdot)$ (i = 1, 2) are given by (2.9), and at the same time, we also introduce the following:

(3.8)
$$\Lambda[P_1(\cdot)](i) + A_1(i)^\top P_1(i) + P_1(i)A_1(i) + C_1(i)^\top P_1(i)C_1(i) \leq -\varepsilon P_1(i), \quad i \in \mathcal{M}$$

$$\Lambda[P_2(\cdot)](i) + A_2(i)^\top P_2(i) + P_2(i)A_2(i) + C_2(i)^\top P_1(i)C_2(i) \leq -\varepsilon P_2(i), \quad i \in \mathcal{M},$$

where $\varepsilon > 0$. We have a simple proposition.

Proposition 3.2. Lyapunov type inequalities (3.7) admit a finite solution pair $(P_1(i), P_2(i))$ $(i \in \mathcal{M})$ if and only if so do (3.8).

Proof. Because $P_1(i) \in \mathbb{S}_{++}^n$, it can be easily seen that the first inequalities in (3.7) and (3.8) are equivalent. For the second parts, we notice that if the first inequality in (3.7) holds for P_1 , it also holds for λP_1 with any $\lambda > 0$. By the arbitrariness of $\lambda > 0$, the second inequalities in (3.7) and (3.8) are also equivalent. The proof is complete.

Now, we are ready to prove the following result.

Proposition 3.3. Let (A1) concerning $A, \overline{A}, C, \overline{C}$ hold. Then the following are equivalent:

- (i) System $[A, \overline{A}, C, \overline{C}]$ is L²-exponentially stable;
- (ii) System $[A, \overline{A}, C, \overline{C}]$ is L^2 -integrable;
- (iii) System $[A, \overline{A}, C, \overline{C}]$ is dissipative;

(iv) System (3.7) (or equivalently (3.8)) of Lyapunov inequalities admits a positive definite solution $(P_1(\cdot), P_2(\cdot))$.

Proof. The proof for (i) \Rightarrow (ii) and (iii) \Leftrightarrow (iv) is trivial. (ii) \Rightarrow (iv). Let $(\Psi_1^{s,\iota}(\cdot), \Psi_2^{s,\iota}(\cdot))$ solve the following SDE

(3.9)
$$\begin{cases} d\Psi_1^{s,\iota}(t) = A_1(\alpha(t))\Psi_1^{s,\iota}(t)dt + C_1(\alpha(t))\Psi_1^{s,\iota}(t)dW(t), \\ d\Psi_2^{s,\iota}(t) = A_2(\alpha(t))\Psi_2^{s,\iota}(t)dt, \quad t \in [s,\infty), \\ \Psi_1^{s,\iota}(s) = I, \quad \Psi_2^{s,\iota}(s) = I, \quad \alpha(s) = \iota. \end{cases}$$

Clearly, for $k = 1, 2, \Psi_k^{s,\iota}(\cdot)$ is L^2 -integrable and invertible. Thus, for $L_1 : \mathcal{M} \mapsto \mathbb{S}_{++}^n$, we may define

$$P_1(i) = \mathbb{E}\left[\int_s^\infty \Psi_1^{s,i}(t)^\top L_1(\alpha(t))\Psi_1^{s,i}(t)dt \mid \alpha(s) = i\right]$$
$$\equiv \mathbb{E}_{s,i}\left[\int_s^\infty \Psi_1^{s,i}(t)^\top L_1(\alpha(t))\Psi_1^{s,i}(t)dt\right] \in \mathbb{S}_{++}^n.$$

Next, for any given $(s, i, \xi_1, 0) \in \mathscr{D}$, applying Itô's formula for $t \mapsto \langle P_1(\alpha(t))\Psi_1(t)\xi_1, \Psi_1(t)\xi_1 \rangle$, we have

$$\mathbb{E}_{s,\iota} \langle P_1(\alpha(t_2)) \Psi_1^{s,\iota}(t_2) \xi_1, \Psi_1^{s,\iota}(t_2) \xi_1 \rangle - \mathbb{E}_{s,\iota} \langle P_1(\alpha(t_1)) \Psi_1^{s,\iota}(t_1) \xi_1, \Psi_1^{s,\iota}(t_1) \xi_1 \rangle$$

= $\mathbb{E}_{s,\iota} \int_{t_1}^{t_2} \left\langle \left(\Lambda[P_1] + A_1^\top P_1 + P_1 A_1 + C_1^\top P_1 C_1 \right) (\alpha(r)) \Psi_1^{s,\iota}(r) \xi_1, \Psi_1^{s,\iota}(r) \xi_1 \right\rangle dr.$

By the definition of $P_1(\cdot)$, (3.9), we get

$$\begin{split} \mathbb{E}_{s,\imath} \langle P_1(\alpha(t_1)) \Psi_1^{s,\imath}(t_1) \xi_1, \Psi_1^{s,\imath}(t_1) \xi_1 \rangle &- \mathbb{E}_{s,\imath} \langle P_1(\alpha(t_2)) \Psi_1^{s,\imath}(t_2) \xi_1, \Psi_1^{s,\imath}(t_2) \xi_1 \rangle \\ &= \mathbb{E}_{s,\imath} \Big\langle \Big[\int_{t_1}^{\infty} [\Psi_1^{t_1,\alpha(t_1)}(r)]^\top L_1(\alpha(r)) \Psi_1^{t_1,\alpha(t_1)}(r) dr \Big] \Psi_1^{s,\imath}(t_1) \xi_1, \Psi_1^{s,\imath}(t_1) \xi_1 \Big\rangle \\ &- \mathbb{E}_{s,\imath} \Big\langle \Big[\int_{t_2}^{\infty} [\Psi_1^{t_2,\alpha(t_2)}(r)]^\top L_1(\alpha(r)) \Psi_1^{t_2,\alpha(t_2)}(r) dr \Big] \Psi_1^{s,\imath}(t_2) \xi_1, \Psi_1^{s,\imath}(t_2) \xi_1 \Big\rangle \\ &= - \mathbb{E}_{s,\imath} \int_{t_2}^{t_1} \langle L_1(\alpha(r)) \Psi_1^{s,\imath}(r) \xi_1, \Psi_1^{s,\imath}(r) \xi_1 \rangle dr. \end{split}$$

Taking $L_1(\cdot) = I$ and noting that $\Psi_1^{s,\iota}(\cdot) > 0$, by the arbitrariness of $t_1 > t_2$, ι and x_1 , we have

$$(\Lambda[P_1] + A_1^{\top}P_1 + P_1A_1 + C_1^{\top}P_1C_1)(j) = -L_1(j) = -I < 0, \quad j \in \mathcal{M}.$$

Now let us construct P_2 . For some $L_2(\cdot) : \mathcal{M} \mapsto \mathbb{S}^n_{++}$, we define

$$P_2(\iota) := \mathbb{E}\Big[\int_s^\infty \Psi_2^{s,\iota}(t)^\top L_2(\alpha(t))\Psi_2^{s,\iota}(t)dt \Big| \alpha(s) = \iota\Big].$$

Similar to $P_1(\cdot)$ above, taking $L_2(\cdot) = I$ and one can steadily check that

$$(\Lambda[P_2] + A_2^\top P_2 + P_2 A_2)(j) = -I < 0,$$

Then (iv) is proved.

(iv) \Rightarrow (i). Applying Itô's formula, using the couple (P_1, P_2) in (3.8), one has (with $\alpha(t)$ being suppressed)

$$\frac{d}{dt}\mathbb{E}\sum_{k=1}^{2}\langle P_{k}(\alpha(t))X_{k}^{0}(t),X_{k}^{0}(t)\rangle = \sum_{k=1}^{2}\mathbb{E}\Big[\langle\Big(\Lambda[P_{k}(\cdot)]+P_{k}A_{k}+A_{k}^{\top}P_{k}+C_{k}^{\top}P_{1}C_{k}\Big)X_{k}^{0}(t),X_{k}^{0}(t)\rangle \\ \leqslant -\varepsilon\mathbb{E}\sum_{k=1}^{2}\langle P_{k}(\alpha(t))X_{k}^{0}(t),X_{k}^{0}(t)\rangle.$$

Grownwall's inequality yields

(3.10)
$$\mathbb{E}\sum_{k=1}^{2} \langle P_k(\alpha(t))X_k^0(t), X_k^0(t) \rangle \leqslant e^{-\varepsilon(t-s)} \mathbb{E}\sum_{k=1}^{2} \langle P_k(\iota)X_k^0(s), X_k^0(s) \rangle.$$

Then, because $P_k(\cdot) \in \mathbb{S}_{++}^n$, k = 1, 2, we get $\mathbb{E} \sum_{k=1}^2 |X_k^0(t)|^2 \leq K e^{-\varepsilon(t-s)} \sum_{k=1}^2 \mathbb{E} |\xi_k|^2$. The proof is complete.

Let us compare the following with (2.14):

(3.11)
$$\begin{cases} dX_{i}^{0}(t) = \left(A(i)X_{i}^{0}(t) + \bar{A}(i)\mathbb{E}_{t}^{\alpha}[X_{i}^{0}(t)]\right)dt + \left(C(i)X_{i}^{0}(t) + \bar{C}(i)\mathbb{E}_{t}^{\alpha}[X_{i}^{0}(t)]\right)dW(t), \quad t \ge s, \\ X_{i}^{0}(s) = \xi, \quad i \in \mathcal{M}. \end{cases}$$

The above are $m_0 = |\mathcal{M}|$ (the number of elements in \mathcal{M}) decoupled systems with $i \in \mathcal{M}$ as the parameter. Clearly, if system (3.11) is L^2 -exponentially stable for all $i \in \mathcal{M}$ so is (2.14). However, the contrary is not true. Here is a simple example.

Example 3.4. Consider the one-dimensional system with Markov chain-dependent coefficient:

(3.12)
$$\begin{cases} dX^{0}(t) = A(\alpha(t))X^{0}(t)dt, \\ X^{0}(s) = \xi. \end{cases}$$

Suppose the Markov chain $\alpha(\cdot)$ has two states $\mathcal{M} = \{1, 2\}$, with generator $\mathbf{\Lambda} = \begin{pmatrix} -10 & 10 \\ 1 & -1 \end{pmatrix}$, and A(1) = 1, A(2) = -1. Also consider systems

(3.13)
$$\begin{cases} dX_i^0(t) = A(i)X_i^0(t)dt, \\ X_i^0(s) = \xi, \end{cases} \quad i \in \mathcal{M}. \end{cases}$$

It is clear that when i = 1, the corresponding system is not stable. Now, we let $P_1, P_2 : \mathcal{M} \to (0, \infty)$ such that for k = 1, 2,

$$\begin{split} &\Lambda[P_k(\cdot)](1) + 2A(1)P_k(1) = \lambda_{11}P_k(1) + \lambda_{12}P_k(2) + 2P_k(1) = -8P_k(1) + 10P_k(2) < 0, \\ &\Lambda[P_k(\cdot)](2) + 2A(2)P_k(2) = \lambda_{21}P_k(1) + \lambda_{22}P_k(2) - 2P_k(2) = P_k(1) - 3P_k(2) < 0. \end{split}$$

These can be achieved if we select

$$0 < \frac{1}{3}P_k(1) < P_k(2) < \frac{4}{5}P_k(1).$$

Hence, we may take, say, $P_1(1) = P_2(1) = 1$, $P_1(2) = P_2(2) \in (\frac{1}{3}, \frac{4}{5})$ (for example, $P_1(2) = P_2(2) = \frac{1}{2}$). Thus, the system (3.12) is stable.

Next, we consider the controlled homogeneous SDE (2.12). For any $\Theta_1, \Theta_2 : \mathcal{M} \to \mathbb{R}^{m \times n}$, we consider the following homogeneous system:

(3.14)
$$\begin{cases} dX_1(t) = (A_1 + B_1\Theta_1)X_1(t)dt + [(C_1 + D_1\Theta_1)X_1(t) + (C_2 + D_2\Theta_2)X_2(t)]dW(t), \\ dX_2(t) = (A_2 + B_2\Theta_2)X_2(t)dt, \quad t \in [s, \infty), \\ X_1(s) = \xi_1, \ X_2(s) = \xi_2, \quad \alpha(s) = \iota. \end{cases}$$

This is the system (2.12) when the controls are taken to be the following:

$$u_k(t) = \Theta_k(\alpha(t))X_k(t), \qquad t \in [s,T], \quad k = 1, 2,$$

which are referred to as *feedback controls*. Thus, according to our notation, this system should be denoted by $\{A_1 + B_1\Theta_1, A_2 + B_2\Theta_2, C_1 + D_1\Theta_1, C_2 + D_2\Theta_2\}$.

Definition 3.5. System $[A, \bar{A}, C, \bar{C}; B, \bar{B}, D, \bar{D}]$ is said to be L^2 -stabilizable if there are (deterministic matrix functions) $\Theta_1, \Theta_2 : \mathcal{M} \to \mathbb{R}^{m \times n}$ such that system $\{A_1 + B_1\Theta_1, A_2 + B_2\Theta_2, C_1 + D_1\Theta_1, C_2 + D_2\Theta_2\}$ is L^2 -integrable. In this case, (Θ_1, Θ_2) is called an L^2 -stabilizer of $[A, \bar{A}, C, \bar{C}; B, \bar{B}, D, \bar{D}]$. We denote the set of all such stabilizers by $\mathbf{S}[A, \bar{A}, C, \bar{C}; B, \bar{B}, D, \bar{D}]$.

Definition 3.6. The maps $(\Theta_1, \Theta_2) : \mathcal{M} \mapsto \mathbb{R}^{m \times n} \times \mathbb{R}^{m \times n}$ is said to be a *(uniform)* L^2 -dissipative strategy of system $[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$ if there exist $P_1, P_2 : \mathcal{M} \mapsto \mathbb{S}^n_{++}$ such that, for any $j \in \mathcal{M}$,

$$(3.15) \quad \left(\Lambda[P_k] + (A_k + B_k \Theta_k)^\top P_k + P_k (A_k + B_k \Theta_k) + (C_k + D_k \Theta_k)^\top P_1 (C_k + D_k \Theta_k)\right)(j) < 0, \quad k = 1, 2.$$

By Proposition 3.3, the following proposition is immediate.

Proposition 3.7. Under (A1), the following are equivalent:

- (i) (Θ_1, Θ_2) is an L^2 -stabilizer of system $[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$.
- (ii) (Θ_1, Θ_2) is an L^2 -dissipative strategy of system $[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$.

The following is an example which shows that if the Markov chain has infinite-many states, then the above (i) does not imply (ii).

Example 3.8. Let the probability space have the decomposition

$$\Omega = \bigcup_{i=1}^{\infty} \Omega_i, \qquad \mathbb{P}(\Omega_i) = p_i > 0, \qquad \sum_{i=1}^{\infty} p_i = 1.$$

Let $\mathcal{M} = \{1, 2, 3, \dots\}$, and consider an infinite state Markov chain $\alpha : \Omega \to \mathcal{M}$ with

$$\alpha(t,\omega) = i, \qquad t \ge 0, \quad \omega \in \Omega_i, \quad i \ge 1.$$

We look at one-dimensional differential equation

$$dX(t) = [A(\alpha(t))X(t) + u(t)]dt, \qquad t \ge 0,$$

with

$$A(i) = -\lambda_i < 0, \qquad i \ge 1, \qquad \lambda_i \downarrow 0, \qquad \sum_{i=1}^{\infty} \frac{p_i}{\lambda_i} < \infty.$$

Then with $u(\cdot) = 0$,

$$X(t,\omega) = X(0) \sum_{i=1}^{\infty} e^{-\lambda_i t} \mathbf{1}_{\Omega_i}(\omega).$$

Thus

$$\int_0^\infty \mathbb{E} |X(t)|^2 dt = |X(0)|^2 \sum_{i=1}^\infty \int_0^\infty p_i e^{-2\lambda_i t} dt = \frac{|X(0)|^2}{2} \sum_{i=1}^\infty \frac{p_i}{\lambda_i} < \infty.$$

That is, 0 is an L^2 -stabilizer. However, (3.10) cannot be true for any $\varepsilon > 0$, which implies that 0 is not an L^2 -dissipative strategy.

Note that similar to Example 3.4, system $[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$ is L^2 -stabilizable does not mean that for all $i \in \mathcal{M}$, the following system is L^2 -stabilizable:

(3.16)
$$\begin{cases} dX_{i}(t) = \left\{ A(i)X_{i}(t) + \bar{A}(i)\mathbb{E}_{t}^{\alpha}[X_{i}(t)] + B(i)u(t) + \bar{B}(i)\mathbb{E}_{t}^{\alpha}[u(t)] \right\} dt \\ + \left\{ C(i)X_{i}(t) + \bar{C}(i)\mathbb{E}_{t}^{\alpha}[X_{i}(t)] + D(i)u(t) + \bar{D}(i)\mathbb{E}_{t}^{\alpha}[u(t)] \right\} dW(t), \ t \ge s, \\ X_{i}(s) = \xi \in L^{2}_{\mathcal{F}_{s}}(\Omega; \mathbb{R}^{n}), \quad i \in \mathcal{M}, \end{cases}$$

This will bring us some essential difficulties later on.

We now introduce the following hypotheses.

(A4) $\mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}] \neq \emptyset.$

,

 $(\mathbf{A4})'(0,0) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}], \text{ i.e., system } [A, \overline{A}, C, \overline{C}] \text{ is } L^2\text{-exponentially stable.}$

It is obvious that the condition $(\mathbf{A4})'$ is stronger than $(\mathbf{A4})$. Now, we consider state equation (2.8) with $[A, \overline{A}, C, \overline{C}]$ being L^2 -exponentially stable. It is known that in this case, the homogeneous equation has a solution in $L^2_{\mathbb{F}^{\alpha}}(s, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(s, \infty; \mathbb{R}^n)$. The following proposition says that this is true for nonhomogeneous systems.

Proposition 3.9. Under (A4)', for any $(s, \iota, \xi_1, \xi_2, u_1(\cdot), u_2(\cdot)) \in \mathscr{D} \times \mathscr{U}[s, \infty)$, (2.8) admits a unique solution $(X_1(\cdot), X_2(\cdot)) \in L^2_{\mathbb{F}^\alpha}(s, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^\alpha}(s, \infty; \mathbb{R}^n)$.

Proof. We notice the existence of a solution $(X_1(\cdot), X_2(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(s, T; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(s, T; \mathbb{R}^n)$ to SDE (2.8) for any T > 0 is trivial because of the linear structure. Now let us verify $(X_1(\cdot), X_2(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(s, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(s, \infty; \mathbb{R}^n)$. Under (A4)', system $\{A_1, A_2, C_1, C_2\}$ is L^2 -exponentially stable. Thus, we have P_1, P_2 :

 $\mathcal{M} \mapsto \mathbb{S}^n_{++}$ satisfy (3.8). Applying Ito's formula to $t \mapsto \sum_{k=1}^2 \langle P_k(\alpha(t)) X_k(t), X_k(t) \rangle$, it follows that

$$\begin{aligned} \frac{d}{dt} \mathbb{E} \sum_{k=1}^{2} \langle P_k(\alpha(t)) X_k(t), X_k(t) \rangle \\ &= \sum_{k=1}^{2} \left(\mathbb{E} \langle \Lambda[P_k] X_k, X_k \rangle + \mathbb{E} \langle P_k(A_k X_k + B_k u_k + b_k), X_k \rangle + \langle P_k X_k, A_k X_k + B_k u_k + b_k \rangle \right) \\ &\quad + \mathbb{E} \langle P_1(C_1 X_1 + C_2 X_2 + D_1 u_1 + D_2 u_2 + \sigma), C_1 X_1 + C_2 X_2 + D_1 u_1 + D_2 u_2 + \sigma \rangle \\ &\leqslant -\varepsilon \mathbb{E} \sum_{k=1}^{2} \langle P_k(\alpha(t)) X_k(t), X_k(t) \rangle + L \sum_{k=1}^{2} \left(\mathbb{E} \langle X_k, u_k \rangle + \mathbb{E} \langle X_k, b_k \rangle + \mathbb{E} \langle X_k, \sigma_k \rangle + \mathbb{E} |u_k|^2 + \mathbb{E} |\sigma_k|^2 \right) \\ &\leqslant -\frac{\varepsilon}{2} \mathbb{E} \sum_{k=1}^{2} \langle P_k(\alpha(t)) X_k(t), X_k(t) \rangle + \sum_{k=1}^{2} \left[\frac{L}{\varepsilon} \mathbb{E} |b_k|^2 + \left(L + \frac{L}{\varepsilon} \right) \mathbb{E} [|u_k|^2 + |\sigma_k|^2] \right]. \end{aligned}$$

In the above inequality, we have used the fact that the system is exponentially stable. Then Gronwall's inequality yields that

$$\mathbb{E}\sum_{k=1}^{2} \langle P_k(\alpha(t))X_k(t), X_k(t)\rangle \leqslant e^{-\frac{\varepsilon}{2}(t-s)} \mathbb{E}\sum_{k=1}^{2} \langle P_k(\iota)\xi_k, \xi_k\rangle + \left(L + \frac{L}{\varepsilon}\right) \int_s^t e^{-\frac{\varepsilon}{2}(t-r)} \mathbb{E}[|u(r)|^2 + |b(r)|^2 + |\sigma(r)|^2] dr.$$

Consequently,

$$\begin{split} &\int_{s}^{\infty} \mathbb{E} \sum_{k=1}^{2} \langle P_{k}(\alpha(t)) X_{k}(t), X_{k}(t) \rangle dt \leqslant \frac{2}{\varepsilon} \mathbb{E} \sum_{k=1}^{2} \langle P_{k}(\iota) \xi_{k}, \xi_{k} \rangle \\ &\quad + \left(L + \frac{L}{\varepsilon} \right) \int_{s}^{\infty} \int_{s}^{t} e^{-\frac{\varepsilon}{2}(t-r)} \mathbb{E}[|u(r)|^{2} + |b(r)|^{2} + |\sigma(r)|^{2}] dr dt \\ &\leqslant \frac{2}{\varepsilon} \mathbb{E} \sum_{k=1}^{2} \langle P_{k}(\iota) \xi_{k}, \xi_{k} \rangle + \frac{2}{\varepsilon} \left(L + \frac{L}{\varepsilon} \right) \int_{s}^{\infty} \mathbb{E}[|u(r)|^{2} + |b(r)|^{2} + |\sigma(r)|^{2}] dr < \infty. \end{split}$$

Note that $P_k(\cdot) \in \mathbb{S}^n_{++}$, k = 1, 2, it follows that $(X_1(\cdot), X_2(\cdot)) \in L^2_{\mathbb{F}^\alpha}(s, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^\alpha}(s, \infty; \mathbb{R}^n)$.

From the above proposition, we see that under (A4), one clearly has

(3.17)
$$\mathscr{U}_{ad}[s,\infty) \neq \varnothing.$$

Thus, under (A1)–(A4), Problem (MF-LQ)^{∞} is meaningful.

4 Some Notions for Problem $(MF-LQ)^{\infty}$

In this section, we introduce some relevant notions for Problem $(MF-LQ)^{\infty}$. First of all, the following definition is necessary.

Definition 4.1. Problem (MF-LQ)^{∞} is said to be *finite* at $(s, i, \xi) \in \mathscr{D}$ if $\mathscr{U}_{ad}[s, \infty) \neq \emptyset$, and

(4.1)
$$\inf_{u(\cdot)\in\mathscr{U}_{ad}[s,\infty)} J^{\infty}(s,\iota,\xi;u(\cdot)) > -\infty.$$

If the above is true for any $(s, i, \xi) \in \mathscr{D}$, we simply say that Problem (MF-LQ)^{∞} is *finite*.

Note that it is not very interesting if Problem (MF-LQ)^{∞} is finite at some $(s, i, \xi) \in \mathscr{D}$ with $\mathscr{U}_{ad}[s, \infty)$ being very small. We now present the following result.

Proposition 4.2. Let **(A1)**–(**A2)** hold. Let Problem (MF-LQ)^{∞} be finite at some $(s, i, \xi) \in \mathscr{D}$. Then $\mathscr{U}_{ad}[s, \infty)$ is a subspace of $\mathscr{U}[s, \infty)$, $J^{\infty}(s, i, \xi; 0)$ is finite, and $u(\cdot) \mapsto J^{\infty}(s, i, \xi; u(\cdot))$ is convex on $\mathscr{U}_{ad}[s, \infty)$.

Proof. Let $(s, i, \xi) \in \mathscr{D}$ such that (4.1) holds. Since the state equation is linear, we must have

(4.2)
$$X(t) \equiv X(t; s, \imath, \xi, u(\cdot)) = \mathcal{L}_0(t)\xi + \mathcal{L}_1[b(\cdot)](t) + \mathcal{L}_2[\sigma(\cdot)](t) + \mathcal{L}_3[u(\cdot)](t), \qquad t \ge s.$$

Here, \mathcal{L}_0 , etc. are (s, i)-dependent linear operators with proper domains and ranges. Then

$$(4.3) J^{\infty}(s,\imath,\xi;u(\cdot)) = \langle M_2u(\cdot),u(\cdot)\rangle + \langle M_1(\xi;b(\cdot),\sigma(\cdot),q(\cdot),r(\cdot)),u(\cdot)\rangle + J^{\infty}(s,\imath,\xi;0).$$

In the above, M_2 is a linear operator from $\mathscr{U}_{ad}[s,\infty)$ into itself, only depending on (s,i), $M_1 \in \mathscr{U}_{ad}[s,\infty)$ is linear in its arguments and depends on (s,i). Then we obtain that for any $\beta \in \mathbb{R}$,

$$J^{\infty}(s, i, \xi; \beta u(\cdot)) = \langle M_2[\beta u(\cdot)], \beta u(\cdot) \rangle + \langle M_1, \beta u(\cdot) \rangle + J^{\infty}(s, i, \xi; 0) \\ = \beta^2 \langle M_2 u(\cdot), u(\cdot) \rangle + \beta \langle M_1, u(\cdot) \rangle + J^{\infty}(s, i, \xi; 0).$$

Hence, $J^{\infty}(s, i, \xi; 0)$ is finite. Further, when $u(\cdot) \in \mathscr{U}_{ad}[s, \infty)$, one has $\beta u(\cdot) \in \mathscr{U}_{ad}[s, \infty)$, for any $\beta \in \mathbb{R}$. This together with the above implies

$$(4.4) M_2 \ge 0,$$

which gives the convexity of $u(\cdot) \mapsto J^{\infty}(s, \iota, \xi; u(\cdot))$ on $\mathscr{U}_{ad}[s, \infty)$. Now, by (4.4), we see that for $u(\cdot), v(\cdot) \in \mathscr{U}_{ad}[s, \infty)$, one has

$$\begin{aligned} 0 &\leqslant \langle M_2[\beta_1 u(\cdot) + \beta_2 v(\cdot)], \beta_1 u(\cdot) + \beta_2 v(\cdot)) = \beta_1^2 \langle M_2 u(\cdot), u(\cdot) \rangle + 2\beta_1 \beta_2 \langle M_2 u(\cdot), v(\cdot) \rangle + \beta_2^2 \langle M_2 v(\cdot). v(\cdot) \rangle \\ &\leqslant 2\beta_1^2 \langle M_2 u(\cdot), u(\cdot) \rangle + 2\beta_2^2 \langle M_2 v(\cdot), v(\cdot) \rangle < \infty. \end{aligned}$$

Hence, $\mathscr{U}_{ad}[s,\infty)$ is a subspace of $\mathscr{U}[s,\infty)$.

Further, we have the following result if some additional condition holds.

Proposition 4.3. Let (A1)-(A3) hold. Then the following hold:

(i) $u(\cdot) \mapsto J(s, i, \xi; u(\cdot))$ is non-negative on $\mathscr{U}[s, \infty)$, which implies Problem (MF-LQ)^{∞} to be finite.

(ii) $u(\cdot) \mapsto J(s, i, \xi; u(\cdot))$ is uniformly convex on $\mathscr{U}_{ad}[s, \infty)$, which implies the following: For given $(s, i, \xi) \in \mathscr{D}$, open-loop optimal control $\bar{u}(\cdot)$ uniquely exists and it is characterized by the unique solution of the following:

(4.5)
$$M_2 \bar{u} + M_1 = 0.$$

Proof. (i) is trivial. (ii) The unform convexity of $u(\cdot) \mapsto J(s, i, \xi; u(\cdot))$ can be proved similar to [30], p.37, Proposition 2.5.1. Then, by [30], p.7, Proposition 1.3.1, we get the implication.

Next, we assume (A4) holds. Picking any $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$, and $v(\cdot) \equiv v_1(\cdot) \oplus v_2(\cdot) \in \mathscr{U}[s, \infty)$, we call them a *closed-loop strategy*. Take

(4.6)
$$u_k^{\Theta_k, v_k}(t) \equiv u_k(t) = v_k(t) + \Theta_k(\alpha(t))X_k(t), \qquad t \ge 0,$$

which is called the *outcome* of $(\Theta_1(\cdot), \Theta_2(\cdot), v_1(\cdot), v_2(\cdot))$, where $X(\cdot) \equiv X_1(\cdot) \oplus X_2(\cdot)$ is the state process corresponding to the control $u(\cdot) = u_1(\cdot) \oplus u_2(\cdot)$. Any such a form control is called a *closed-loop control*. Under such a control, we have

$$(4.7) \begin{cases} dX_1(t) = \left\{ [A_1(\alpha(t)) + B_1(\alpha(t))\Theta_1(\alpha(t))]X_1(t) + B_1(\alpha(t))v_1(t) + b_1(t) \right\} dt \\ + \left\{ [C_1(\alpha(t)) + D_1(\alpha(t))\Theta_1(\alpha(t))]X_1(t) + [C_2(\alpha(t)) + D_2(\alpha(t))\Theta_2(\alpha(t))]X_2(t) + D_1(\alpha(t))v_1(t) + D_2(\alpha(t))v_2(t) + \sigma(t) \right\} dW(t), \\ dX_2(t) = \left\{ [A_2(\alpha(t)) + B_2(\alpha(t))\Theta_2(\alpha(t))]X_2(t) + B_2(\alpha(t))v_2(t) + b_2(t) \right\} dt, \quad t \in [s, \infty), \\ X_1(s) = \xi_1, \quad X_2(s) = \xi_2, \quad \alpha(s) = \iota. \end{cases}$$

From the above, we are ready to introduce the following definition.

Definition 4.4. A strategy $(\Theta_1^*(\cdot), \Theta_2^*(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$ and $(v_1^*(\cdot), v_2^*(\cdot)) \in \mathscr{U}[s, \infty)$ is said be *closed-loop optimal* at (s, i) if

(4.8)
$$J^{\infty}(s, \imath, \xi_1, \xi_2; u_1^{\Theta_1^*, v_1^*}(\cdot), u_2^{\Theta_2^*, v_2^*}(\cdot)) \leqslant J^{\infty}(s, \imath, \xi_1, \xi_2; u_1^{\Theta_1, v_1}(\cdot), u_2^{\Theta_2, v_2}(\cdot)),$$

for any $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \bar{A}, C, \bar{C}; B, \bar{B}, D, \bar{D}], (v_1(\cdot), v_2(\cdot)) \in \mathscr{U}[s, \infty) \text{ and } \xi \equiv (\xi_1, \xi_2) \in L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{R}^n)^{\perp} \times L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{R}^n).$ When this happens, we say that Problem (MF-LQ)^{\infty} is *closed-loop solvable* at (s, i).

Note that (4.8) is required to hold for all ξ . In the above case, $(\Theta_1^*(\cdot), \Theta_2^*(\cdot), v_1^*(\cdot), v_2^*(\cdot))$ (which is called a *closed-loop optimal strategy*) is optimal in the class of closed-loop strategies. Now, if $(\Theta_1^*(\cdot), \Theta_2^*(\cdot), v_1^*(\cdot), v_2^*(\cdot))$ is a closed-loop optimal, with $X^*(\cdot) = X_1^*(\cdot) \oplus X_2^*(\cdot) \in L^2_{\mathbb{F}}(s, \infty; \mathbb{R}^n)$ being the corresponding closed-loop optimal state process, then we may write (4.7) as (suppressing $\alpha(\cdot)$)

$$\begin{cases} dX_1^*(t) = \left\{ \begin{bmatrix} A_1 X_1^*(t) + B_1[v_1^*(t) + \Theta_1^* X_1^*(t)] + b_1(t) \right\} dt \\ + \left\{ C_1 X_1^*(t) + C_2 X_2^*(t) + D_1[v_1^*(t) + \Theta_1^* X_1^*(t)] + D_2[v_2^*(t) + \Theta_2^* X_2^*(t)] + \sigma(t) \right\} dW(t), \\ dX_2^*(t) = \left\{ A_2 X_2^*(t) + B_2[v_2^*(t) + \Theta_2^* X_2^*(t)] + b_2(t) \right\} dt, \quad t \in [s, \infty), \\ X_1^*(s) = \xi_1, \quad X_2^*(s) = \xi_2, \quad \alpha(s) = \iota. \end{cases}$$

Hence, by taking

$$u_k^*(t) = v_k^*(t) + \Theta_k^*(\alpha(t))X_k^*(t), \quad k = 1, 2,$$

we have $u_1^*(\cdot) \oplus u_2^*(\cdot) \in \mathscr{U}_{ad}[s, \infty)$. Likewise, for any $u_1(\cdot) \oplus u_2(\cdot) \in \mathscr{U}_{ad}[s, \infty)$, with $X(\cdot) = X_1(\cdot) \oplus X_2(\cdot) \in L^2_{\mathbb{F}}(s, \infty; \mathbb{R}^n)$ being the corresponding state process, we may write

(4.9)
$$\begin{cases} dX_1(t) = \left\{ [A_1 + B_1\Theta_1^*]X_1(t) + B_1[u_1(t) - \Theta_1^*X_1(t)] + b_1(t) \right\} dt \\ + \left\{ [C_1 + D_1\Theta_1^*]X_1(t) + [C_2 + D_2\Theta_2^*]X_2(t) \\ + D_1[u_1(t) - \Theta_1^*X_1(t)] + D_2[u_2(t) - \Theta_2^*X_2(t)] + \sigma(t) \right\} dW(t), \\ dX_2(t) = \left\{ [A_2 + B_2\Theta_2^*]X_2(t) + B_2[u_2(t) - \Theta_2^*X_2(t)] + b_2(t) \right\} dt, \quad t \in [s, \infty), \\ X_1(s) = \xi_1, \quad X_2(s) = \xi_2, \quad \alpha(s) = \iota. \end{cases}$$

Thus, $X(\cdot)$ is the state process under $(\Theta_1^*(\cdot), \Theta_2^*(\cdot), v_1(\cdot), v_2(\cdot))$ where

(4.10)
$$v_k(t) = u_k(t) - \Theta_k^*(\alpha(t))X_k(t), \quad k = 1, 2$$

Then,

$$\begin{split} J^{\infty}(s,\imath,\xi_{1},\xi_{2};u_{1}^{*}(\cdot),u_{2}^{*}(\cdot)) &= J^{\infty}(s,\imath,\xi_{1},\xi_{2};u_{1}^{\Theta_{1}^{*},v_{1}^{*}}(\cdot),u_{2}^{\Theta_{2}^{*},v_{2}^{*}}(\cdot)) \\ &\leqslant J^{\infty}(s,\imath,\xi_{1},\xi_{2};u_{1}^{\Theta_{1}^{*},v_{1}}(\cdot),u_{2}^{\Theta_{2}^{*},v_{2}}(\cdot)) = J(s,\imath,\xi_{1},\xi_{2};u_{1}(\cdot),u_{2}(\cdot)). \end{split}$$

Thus, $(u_1^*(\cdot), u_2^*(\cdot))$ is open-loop optimal. This means that if Problem (MF-LQ)^{∞} is closed-loop solvable, then it is open-loop solvable.

Now, let us assume (A4), and let $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$. Then, under the control (4.6), we have state equation (4.7) and the cost functional $J^{\infty}(s, i, \xi_1, \xi_2; u_1^{\Theta_1, v_1}(\cdot), u_2^{\Theta_2, v_2}(\cdot))$. The corresponding LQ problem is denoted by Problem (MF-LQ)^{$\infty, *$}. The following result will be useful below in deriving a characterization of open-loop solvability of Problem (MF-LQ)^{∞}.

Proposition 4.5. Let (A1)–(A4) hold and $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$. Then

(4.11)
$$\mathscr{U}_{ad}[s,\infty) = \left\{ u_1^{\Theta_1,v_1}(\cdot) \oplus u_2^{\Theta_2,v_2}(\cdot) \mid v_1(\cdot) \oplus v_2(\cdot) \in \mathscr{U}[s,\infty) \right\}.$$

Moreover, Problem (MF-LQ) is open-loop solvable at (s, i, ξ) if and only if so is Problem (MF-LQ)^{$\infty, *$}.

Proof. We first prove (4.11). Let $u(\cdot) = u_1(\cdot) \oplus u_2(\cdot) \in \mathscr{U}_{ad}[s, \infty)$. Since $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \bar{A}, C, \bar{C}; B, \bar{B}, D, \bar{D}]$, for any $u_1(\cdot) \oplus u_2(\cdot) \in \mathscr{U}_{ad}[s, \infty)$, we may write (2.8) as (4.9) replacing $(\Theta_1^*(\cdot), \Theta_2^*(\cdot))$ by $(\Theta_1(\cdot), \Theta_2(\cdot))$. Then $X(\cdot)$ is the state process under strategy $(\Theta_1(\cdot), \Theta_2(\cdot), v_1(\cdot), v_2(\cdot))$. This means the left-hand side of (4.11) is in the right-hand side. Conversely, for any $v_1(\cdot) \oplus v_2(\cdot) \in \mathscr{U}[s, \infty)$, let $X(\cdot)$ be the state process under $(\Theta_1(\cdot), \Theta_2(\cdot), v_1(\cdot), v_2(\cdot))$. Then we write

$$dX_{1}(t) = \left\{ [A_{1} + B_{1}\Theta]X_{1}(t) + v_{1}(t) + b_{1}(t) \right\} dt \\ + \left\{ [C_{1} + D_{1}\Theta_{1})X_{1}(t) + (C_{2} + D_{2}\Theta_{2})X_{2}(t) + D_{1}v_{1}(t) + D_{2}v_{2}(t) + \sigma(t) \right\} dW(t), \\ dX_{2}(t) = \left\{ (A_{2} + B_{2}\Theta_{2})X_{2}(t) + B_{2}v_{2}(t) + b_{2}(t) \right\} dt, \qquad t \in [s, \infty), \\ X_{1}(s) = \xi_{1}, \quad X_{2}(s) = \xi_{2}, \quad \alpha(s) = \iota.$$

Since $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$, the above system is L^2 -exponentially stable. Thus, $X(\cdot) = X_1(\cdot) \oplus X_2(\cdot) \in L^2_{\mathbb{F}}(s, \infty; \mathbb{R}^n)$. Then the right-hand side of (4.11) is in the left-hand side. Therefore, we have the equality.

Now, by a similar argument, we are able to show the equivalence of the open-loop solvability of Problems $(MF-LQ)^{\infty}$ and $(MF-LQ)^{\infty,*}$.

5 Closed-Loop Solvability

With all the above preparation, we are ready to state the main result of this paper concerning the closed-loop solvability of Problem $(MF-LQ)^{\infty}$.

Theorem 5.1. Suppose (A1)–(A4) hold. Then we have the following conclusions:

(i) There exists a unique solution $P_1, P_2 : \mathcal{M} \to \mathbb{S}^n_{++}$ to the following ARE:

(5.1)
$$\begin{cases} \Lambda[P_k(\cdot)](i) + P_k(i)A_k(i) + A_k(i)^\top P_k(i) + C_k(i)^\top P_1(i)C_k(i) + Q_k(i) \\ -[P_k(i)B_k(i) + C_k(i)^\top P_1(i)D_k(i) + S_k(i)^\top][R_k(i) + D_k(i)^\top P_1(i)D_k(i)]^{-1} \\ \cdot [B_k(i)^\top P_k(i) + D_k(i)^\top P_1(i)C_k(i) + S_k(i)] = 0, \\ R_k(i) + D_k(i)^\top P_1(i)D_k(i) > 0, \qquad k = 1, 2, \end{cases}$$

such that $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$ where

$$\Theta_k = -(R_k + D_k^\top P_1 D_k)^{-1} (B_k^\top P_k + D_k^\top P_1 C_k + S_k), \ k = 1, 2.$$

(ii) There exists a unique adapted solution $(Y_1(\cdot), Z(\cdot), Z_1^M(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}}(0, \infty; \mathbb{R}^n) \times M^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)^{\perp}$ and $(Y_2(\cdot), Z_2^M(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n) \times M^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)$ to the following BSDE

(5.2)
$$\begin{cases} dY_1 = -\left((A_1^{\Theta_1})^\top Y_1 + (C_1^{\Theta_1})^\top \Pi_1[Z] + P_1 b_1 + (C_1^{\Theta_1})^\top P_1 \sigma_1 + q_1 + \Theta_1^\top r_1\right) dt + Z dW + Z_1^M dM, \\ dY_2 = -\left((A_2^{\Theta_2})^\top Y_2 + (C_2^{\Theta_2})^\top \Pi_2[Z] + P_2 b_2 + (C_2^{\Theta_2})^\top P_1 \sigma_2 + q_2 + \Theta_2^\top r_2\right) dt + Z_2^M dM, \\ \lim_{t \to \infty} \mathbb{E}[|Y_1(t)|^2 + |Y_2(t)|^2] = 0. \end{cases}$$

Here $A_i^{\Theta_i} = A_i + B_i \Theta_i$ and $C_i^{\Theta_i} = C_i + D_i \Theta_i$.

(iii) The 4-tuple $(\Theta_1(\cdot), \Theta_2(\cdot), v_1(\cdot), v(\cdot))$ is the closed-loop optimal strategy of Problem (MF-LQ)^{∞}, where

(5.3)
$$v_k = -(R_k + D_k^{\dagger} P_1 D_k)^{-1} (B_k^{\dagger} Y_k + D_k^{\dagger} \Pi_k [Z] + D_k^{\dagger} P_1 \sigma_k + r_k), \qquad k = 1, 2.$$

Thus, the following outcome

(5.4)
$$\bar{u}_k = \Theta_k X_k - (R_k + D_k^\top P_1 D_k)^{-1} (B_k^\top Y_k + D_k^\top \Pi_k [Z] + D_k^\top P_1 \sigma_k + r_k), \qquad k = 1, 2,$$

is the optimal open-loop optimal control.

The above theorem will be proved in the following three subsections. First, we will present the derivation of the ARE (5.1) and BSDE (5.2) through completing the squares. It can also be seen that if both of the two equations admit some required solutions as stated in (i) and (ii), then $(\bar{u}_1(\cdot), \bar{u}_2(\cdot))$ in (iii) is the unique optimal control. Then we complete the proof by establishing the well-posedness of (5.1) and (5.2).

5.1 Completing the Squares

The purpose of this section is to approach Problem (MF-LQ)^{∞} by a classical method — completing the squares. In what follows, we suppress $\alpha(\cdot)$ and t. First of all, for any $P_1, P_2 : \mathcal{M} \to \mathbb{S}^n_{++}$, we have

$$\begin{split} & \mathbb{E}\Big[\langle P_{1}(\alpha(T))X_{1}(T), X_{1}(T)\rangle - \langle P_{1}(\alpha(s))\xi_{1}, \xi_{1}\rangle\Big] \\ &= \mathbb{E}\!\int_{s}^{T}\!\!\Big(\langle \big[\Lambda[P_{1}] + P_{1}A_{1} + A_{1}^{\top}P_{1}\big]X_{1}, X_{1}\rangle + 2\langle P_{1}(B_{1}u_{1} + b_{1}), X_{1}\rangle \\ &\quad + \langle P_{1}(C_{1}X_{1} + D_{1}u_{1} + \sigma_{1}), C_{1}X_{1} + D_{1}u_{1} + \sigma_{1}\rangle + \langle P_{1}(C_{2}X_{2} + D_{2}u_{2} + \sigma_{2}), C_{2}X_{2} + D_{2}u_{2} + \sigma_{2}\rangle\Big)dt \\ &= \mathbb{E}\int_{s}^{T}\Big(\langle \big\{\Lambda[P_{1}] + P_{1}A_{1} + A_{1}^{\top}P_{1} + C_{1}^{\top}P_{1}C_{1}\big\}X_{1}, X_{1}\rangle + \langle C_{2}^{\top}P_{1}C_{2}X_{2}, X_{2}\rangle \\ &\quad + \langle D_{1}^{\top}P_{1}D_{1}u_{1}, u_{1}\rangle + 2\langle u_{1}, (B_{1}^{\top}P_{1} + D_{1}^{\top}P_{1}C_{1})X_{1} + D_{1}^{\top}P_{1}\sigma_{1}\rangle + 2\langle P_{1}b_{1} + C_{1}^{\top}P_{1}\sigma_{1}, X_{1}\rangle + \langle P_{1}\sigma_{1}, \sigma_{1}\rangle \\ &\quad + \langle D_{2}^{\top}P_{1}D_{2}u_{2}, u_{2}\rangle + 2\langle u_{2}, D_{2}^{\top}P_{1}C_{2}X_{2} + D_{2}^{\top}P_{1}\sigma_{2}\rangle + 2\langle C_{2}^{\top}P_{1}\sigma_{2}, X_{2}\rangle + \langle P_{1}\sigma_{2}, \sigma_{2}\rangle\Big)dt, \end{split}$$

and

$$\mathbb{E}\Big[\langle P_{2}(\alpha(T))X_{2}(T), X_{2}(T)\rangle - \langle P_{2}(\alpha(s))\xi_{2}, \xi_{2}\rangle\Big]$$

= $\mathbb{E}\int_{s}^{T}\Big(\langle\{\Lambda[P_{2}] + P_{2}A_{2} + A_{2}^{\top}P_{2}\}X_{2}, X_{2}\rangle + 2\langle P_{2}[B_{2}u_{2} + b_{2}], X_{2}\rangle\Big)dt$
= $\mathbb{E}\int_{s}^{T}\Big(\langle\{\Lambda[P_{2}] + P_{2}A_{2} + A_{2}^{\top}P_{2}\}X_{2}, X_{2}\rangle + 2\langle u_{2}, B_{2}^{\top}P_{2}X_{2}\rangle + 2\langle P_{2}b_{2}, X_{2}\rangle\Big)dt.$

In the above, the cross terms are zero, for example,

$$\mathbb{E}\int_{s}^{\infty} \langle P_{1}C_{1}X_{1}, C_{2}X_{2} \rangle dt = \mathbb{E}\int_{s}^{\infty} \mathbb{E}_{t}^{\alpha} \langle P_{1}C_{1}X_{1}, C_{2}X_{2} \rangle dt = 0.$$

Next, let $(Y(\cdot), Z(\cdot), Z^M(\cdot)) \in L^2_{\mathbb{F}}(s, \infty; \mathbb{R}^n) \times L^2_{\mathbb{F}}(s, \infty; \mathbb{R}^n) \times L^2_{\mathbb{F}^\alpha}(s, \infty; \mathbb{R}^n)$ be the unique adapted solution of the following BSDE:

$$\begin{cases} dY(t) = \Gamma(t)dt + Z(t)dW(t) + Z^M(t)dM(t), & t \ge s, \\ \lim_{t \to \infty} \mathbb{E}|Y(t)|^2 = 0, \end{cases}$$

with $\Gamma(\cdot)$ being undetermined. Let

$$\begin{split} Y_{2}(\cdot) &= \Pi_{2}[Y(\cdot)], \quad Z_{2}(\cdot) = \Pi_{2}[Z(\cdot)], \quad Z_{2}^{M}(\cdot) = \Pi_{2}[Z^{M}(\cdot)], \quad \Gamma_{2}(t) = \Pi_{2}[\Gamma(\cdot)], \\ Y_{1}(\cdot) &= Y(\cdot) - Y_{2}(\cdot), \quad Z_{1}(\cdot) = Z(\cdot) - Z_{(}\cdot), \\ Z_{1}^{M}(\cdot) &= Z^{M}(\cdot) - Z_{2}^{M}(\cdot), \quad \Gamma_{1}(t) = \Gamma(t) - \Gamma_{2}(t). \end{split}$$

Applying Π_2 to the above BSDE, and set $Y_1(\cdot), Z_1^M(\cdot)$ and $\Gamma_1(\cdot)$, etc. as above, we have

$$dY_1 = \Gamma_1 dt + Z dW + Z_1^M dM, \qquad \lim_{t \to \infty} \mathbb{E}|Y_1(t)|^2 = 0,$$

$$dY_2 = \Gamma_2 dt + Z_2^M dM, \qquad \lim_{t \to \infty} \mathbb{E}|Y_2(t)|^2 = 0.$$

Then

$$\mathbb{E}\Big[\langle Y_1(T), X_1(T) \rangle - \langle Y_1(s), \xi_1 \rangle \Big]$$

= $\mathbb{E} \int_s^T \Big(\langle \Gamma_1, X_1 \rangle + \langle Y_1, A_1 X_1 + B_1 u_1 + b_1 \rangle + \langle Z, C_1 X_1 + C_2 X_2 + D_1 u_1 + D_2 u_2 + \sigma \rangle \Big) dt,$

 $\quad \text{and} \quad$

$$\mathbb{E}\Big[\langle Y_2(T), X_2(T) \rangle - \langle Y_2(s), \xi_2 \rangle\Big] = \mathbb{E}\int_s^T \Big(\langle \Gamma_2, X_2 \rangle + \langle Y_2, A_2X_2 + B_2u_2 + b_2 \rangle\Big) dt.$$

Then

$$\begin{split} J^{\infty}(s,\iota,\xi_{1},\xi_{2};u_{1}(\cdot),u_{2}(\cdot)) &:= J^{\infty}(s,\iota,\xi;u(\cdot)) \\ &= \frac{1}{2}\sum_{k=1}^{2} \mathbb{E} \int_{s}^{\infty} \Big[\langle Q_{k}X_{k},X_{k} \rangle + 2\langle S_{k}X_{k},u_{k} \rangle + \langle R_{k}u_{k},u_{k} \rangle + 2\langle q_{k},X_{k} \rangle + 2\langle r_{k},u_{k} \rangle \Big] dt \\ &= \frac{1}{2}\sum_{k=1}^{2} \mathbb{E} \Big\{ \langle P_{k}(\alpha(s))\xi_{k},\xi_{k} \rangle + 2\langle Y_{k}(s),\xi_{k} \rangle - \langle P_{k}(\alpha(T))X_{k}(T),X_{k}(T) \rangle - 2\langle Y_{k}(T),X_{k}(T) \rangle \\ &+ \int_{s}^{\infty} \Big(\langle Q_{k}X_{k},X_{k} \rangle + 2\langle S_{k}X_{k},u_{k} \rangle + \langle R_{k}u_{k},u_{k} \rangle + 2\langle q_{k},X_{k} \rangle + 2\langle r_{k},u_{k} \rangle \Big] dt \Big\} \\ &+ \frac{1}{2} \mathbb{E} \int_{s}^{T} \Big(\langle \{\Lambda[P_{1}] + P_{1}A_{1} + A_{1}^{\top}P_{1} + C_{1}^{\top}P_{1}C_{1}\}X_{1},X_{1} \rangle + \langle C_{2}^{\top}P_{1}C_{2}X_{2},X_{2} \rangle + \langle P_{1}\sigma_{1},\sigma_{1} \rangle \\ &+ \langle D_{1}^{\top}P_{1}D_{1}u_{1},u_{1} \rangle + 2\langle u_{1},(B_{1}^{\top}P_{1} + D_{1}^{\top}P_{1}C_{1})X_{1} + D_{1}^{\top}P_{1}\sigma_{1} \rangle + 2\langle P_{1}b_{1} + C_{1}^{\top}P_{1}\sigma_{1},X_{1} \rangle \\ &+ \langle D_{2}^{\top}P_{1}D_{2}u_{2},u_{2} \rangle + 2\langle u_{2},D_{2}^{\top}P_{1}C_{2}X_{2} + D_{2}^{\top}P_{1}\sigma_{2} \rangle + 2\langle C_{2}^{\top}P_{1}\sigma_{2},X_{2} \rangle + \langle P_{1}\sigma_{2},\sigma_{2} \rangle \\ &+ \langle \{\Lambda[P_{2}] + P_{2}A_{2} + A_{2}^{\top}P_{2}\}X_{2},X_{2} \rangle + 2\langle u_{2},B_{2}^{\top}P_{2}X_{2} \rangle + 2\langle P_{2}b_{2},X_{2} \rangle \\ &+ 2\langle \Gamma_{1},X_{1} \rangle + 2\langle Y_{1},A_{1}X_{1} + B_{1}u_{1} + b_{1} \rangle + 2\langle Z,C_{1}X_{1} + C_{2}X_{2} + D_{1}u_{1} + D_{2}u_{2} + \sigma \rangle \\ &+ 2\langle \Gamma_{2},X_{2} \rangle + 2\langle Y_{2},A_{2}X_{2} + B_{2}u_{2} + b_{2} \rangle \Big] dt. \end{split}$$

We assume that as $T \to \infty$, the above limits exist, in particular, $X_k(T) \to 0$ and $Y_k(T)$ is bounded. Then for the terms in $L^2_{\mathbb{F}^\alpha}(s,\infty;\mathbb{R}^n)$, we have

$$\begin{split} &\mathbb{E} \int_{s}^{\infty} \left(\langle Q_{1}X_{1}, X_{1} \rangle + 2 \langle S_{1}X_{1}, u_{1} \rangle + \langle R_{1}u_{1}, u_{1} \rangle + 2 \langle q_{1}, X_{1} \rangle + 2 \langle r_{1}, u_{1} \rangle \\ &+ \langle \{\Lambda[P_{1}] + P_{1}A_{1} + A_{1}^{\top}P_{1} + C_{1}^{\top}P_{1}C_{1}\}X_{1}, X_{1} \rangle + \langle P_{1}\sigma_{1}, \sigma_{1} \rangle \\ &+ \langle D_{1}^{\top}P_{1}D_{1}u_{1}, u_{1} \rangle + 2 \langle u_{1}, (B_{1}^{\top}P_{1} + D_{1}^{\top}P_{1}C_{1})X_{1} + D_{1}^{\top}P_{1}\sigma_{1} \rangle + 2 \langle P_{1}b_{1} + C_{1}^{\top}P_{1}\sigma_{1}, X_{1} \rangle \\ &+ 2 \langle \Gamma_{1}, X_{1} \rangle + 2 \langle Y_{1}, A_{1}X_{1} + B_{1}u_{1} + b_{1} \rangle + 2 \langle Z_{1}, C_{1}X_{1} + D_{1}u_{1} + \sigma_{1} \rangle \right) dt \\ &= \mathbb{E} \int_{s}^{\infty} \left(\langle \{\Lambda[P_{1}] + P_{1}A_{1} + A_{1}^{\top}P_{1} + C_{1}^{\top}P_{1}C_{1} + Q_{1}\}X_{1}, X_{1} \rangle \\ &+ 2 \langle \Gamma_{1} + A_{1}^{\top}Y_{1} + C_{1}^{\top}Z_{1} + P_{1}b_{1} + C_{1}^{\top}P_{1}\sigma_{1} + q_{1}, X_{1} \rangle + \langle (R_{1} + D_{1}^{\top}P_{1}D_{1})u_{1}, u_{1} \rangle \\ &+ 2 \langle u_{1}, (B_{1}^{\top}P_{1} + D_{1}^{\top}P_{1}C_{1} + S_{1})X_{1} + B_{1}^{\top}Y_{1} + D_{1}^{\top}Z_{1} + D_{1}^{\top}P_{1}\sigma_{1} + r_{1} \rangle \\ &+ 2 \langle Y_{1}, b_{1} \rangle + 2 \langle Z_{1}, \sigma_{1} \rangle + \langle P_{1}\sigma_{1}, \sigma_{1} \rangle \right) dt \\ &= \mathbb{E} \int_{s}^{\infty} \left(\langle \{\Lambda[P_{1}] + Q_{1}\}X_{1}, X_{1} \rangle + |\mathcal{R}_{1}^{\frac{1}{2}}[u_{1} + \mathcal{R}_{1}^{-1}(S_{1}X_{1} + B_{1}^{\top}Y_{1} + D_{1}^{\top}Z_{1} + D_{1}^{\top}P_{1}\sigma_{1} + r_{1})]|^{2} \\ &+ 2 \langle Y_{1}, b_{1} \rangle + 2 \langle Z_{1}, \sigma_{1} \rangle + \langle P_{1}\sigma_{1}, \sigma_{1} \rangle \right) dt \\ &= \mathbb{E} \int_{s}^{\infty} \left(\langle \{\Lambda[P_{1}] + Q_{1} - S_{1}^{\top}\mathcal{R}_{1}^{-1}S_{1}\}X_{1}, X_{1} \rangle + |\mathcal{R}_{1}^{\frac{1}{2}}[u_{1} - \Theta_{1}X_{1} + \mathcal{R}_{1}^{-1}(B_{1}^{\top}Y_{1} + D_{1}^{\top}Z_{1} + D_{1}^{\top}P_{1}\sigma_{1} + r_{1})]|^{2} \\ &+ 2 \langle \Gamma_{1} + (A_{1} - B_{1}\mathcal{R}^{-1}S_{1})^{\top}Y_{1} + (C_{1} - D_{1}\mathcal{R}_{1}^{-1}S_{1})^{\top}Z_{1} \right) \right]$$

$$\begin{split} &+P_{1}b_{1}+(C_{1}-D_{1}\mathcal{R}_{1}^{-1}\mathcal{S}_{1})^{\top}P_{1}\sigma_{1}+q_{1}-\mathcal{S}_{1}^{\top}\mathcal{R}_{1}^{-1}r_{1},X_{1}\rangle \\ &-\left|\mathcal{R}_{1}^{-\frac{1}{2}}(B_{1}^{\top}Y_{1}+D_{1}^{\top}Z_{1}+D_{1}^{\top}P_{1}\sigma_{1}+r_{1})\right|^{2}+2\langle Y_{1},b_{1}\rangle+2\langle Z_{1},\sigma_{1}\rangle+\langle P_{1}\sigma_{1},\sigma_{1}\rangle\Big)dt \\ &=\mathbb{E}\int_{s}^{\infty}\left(\langle\left\{\Lambda[P_{1}]+\mathcal{Q}_{1}-\mathcal{S}_{1}^{\top}\mathcal{R}_{1}^{-1}\mathcal{S}_{1}\right\}X_{1},X_{1}\rangle+\left|\mathcal{R}_{1}^{\frac{1}{2}}\left[u_{1}-\Theta_{1}X_{1}+\mathcal{R}_{1}^{-1}(B_{1}^{\top}Y_{1}+D_{1}^{\top}Z_{1}+D_{1}^{\top}P_{1}\sigma_{1}+r_{1})\right]\right|^{2} \\ &-2\langle\Gamma_{1}+(A_{1}^{\Theta_{1}})^{\top}Y_{1}+(C_{1}^{\Theta_{1}})^{\top}Z_{1}+P_{1}b_{1}+(C_{1}^{\Theta_{1}})^{\top}P_{1}\sigma_{1}+q_{1}+\Theta_{1}^{\top}r_{1},X_{1}\rangle \\ &-\left|\mathcal{R}_{1}^{-\frac{1}{2}}(B_{1}^{\top}Y_{1}+D_{1}^{\top}Z_{1}+D_{1}^{\top}P_{1}\sigma_{1}+r_{1})\right|^{2}+2\langle Y_{1},b_{1}\rangle+2\langle Z_{1},\sigma_{1}\rangle+\langle P_{1}\sigma_{1},\sigma_{1}\rangle\Big)dt, \end{split}$$

where

(5.5)
$$\begin{aligned} \mathcal{Q}_1 &= P_1 A_1 + A_1^\top P_1 + C_1^\top P_1 C_1 + Q_1, \quad \mathcal{S}_1 = B_1^\top P_1 + D_1^\top P C_1 + S_1, \quad \mathcal{R}_1 = R_1 + D_1^\top P_1 D_1, \\ \Theta_1 &= -\mathcal{R}_1^{-1} \mathcal{S}_1, \quad A_1^{\Theta_1} = A_1 + B_1 \Theta_1, \quad C_1^{\Theta_1} = C_1 + D_1 \Theta_1. \end{aligned}$$

Since $P_1(\cdot) \in \mathbb{S}^n_{++}$, we have $\mathcal{R}_1(\cdot) \in \mathbb{S}^m_{++}$. The terms involving X_2 become

$$\begin{split} & \mathbb{E} \int_{s}^{\infty} \left(\langle Q_{2}X_{2}, X_{2} \rangle + 2 \langle S_{2}X_{2}, u_{2} \rangle + \langle R_{2}u_{2}, u_{2} \rangle + 2 \langle q_{2}, X_{2} \rangle + 2 \langle r_{2}, u_{2} \rangle \\ & + \langle C_{2}^{T}P_{1}C_{2}X_{2}, X_{2} \rangle + \langle D_{2}^{T}P_{1}D_{2}u_{2}, u_{2} \rangle + 2 \langle u_{2}, D_{2}^{T}P_{1}\sigma_{2} \rangle + \langle P_{1}\sigma_{2}, \sigma_{2} \rangle \\ & + \langle \{\Lambda[P_{2}] + P_{2}A_{2} + A_{2}^{T}P_{2}\}X_{2}, X_{2} \rangle + 2 \langle u_{2}, (B_{2}^{T}P_{2} + D_{2}^{T}P_{1}C_{2})X_{2} \rangle + 2 \langle P_{2}b_{2}, X_{2} \rangle \\ & + 2 \langle Z_{2}, C_{2}X_{2} + D_{2}u_{2} + \sigma_{2} \rangle + 2 \langle \Gamma_{2}, X_{2} \rangle + 2 \langle Y_{2}, A_{2}X_{2} + B_{2}u_{2} + b_{2} \rangle \right) dt \\ & = \mathbb{E} \int_{s}^{\infty} \left(\langle \{\Lambda[P_{2}] + P_{2}A_{2} + A_{2}^{T}P_{2} + C_{2}^{T}P_{1}C_{2} + Q_{2}\}X_{2}, X_{2} \rangle \\ & + 2 \langle \Gamma_{2} + A_{2}^{T}Y_{2} + P_{2}b_{2} + C_{2}^{T}P_{1}\sigma_{2} + q_{2}, X_{2} \rangle \\ & + \langle (R_{2} + D_{2}^{T}P_{1}D_{2})u_{2}, u_{2} \rangle + 2 \langle u_{2}, (B_{2}^{T}P_{2} + D_{2}^{T}P_{1}C_{2} + S_{2})X_{2} + B_{2}^{T}Y_{2} + D_{2}^{T}Z_{2} + D_{2}^{T}P_{1}\sigma_{2} + r_{2} \rangle \\ & + 2 \langle Z_{2}, \sigma_{2} \rangle + 2 \langle Y_{2}, b_{2} \rangle + \langle P_{1}\sigma_{2}, \sigma_{2} \rangle \right) dt \\ & = \mathbb{E} \int_{s}^{\infty} \left(\langle \{\Lambda[P_{2}] + Q_{2}\}X_{2}, X_{2} \rangle + |\mathcal{R}_{2}^{\frac{1}{2}}[u_{2} + \mathcal{R}_{2}^{-1}(S_{2}X_{2} + B_{2}^{T}Y_{2} + D_{2}^{T}Z_{2} + D_{2}^{T}P_{1}\sigma_{2} + r_{2})]|^{2} \\ & + 2 \langle \Gamma_{2} + A_{2}^{T}Y_{2} + C_{2}^{T}Z_{2} + P_{2}b_{2} + C_{2}^{T}P_{1}\sigma_{2} + q_{2}, X_{2} \rangle - |\mathcal{R}_{2}^{-\frac{1}{2}}(S_{2}X_{2} + B_{2}^{T}Y_{2} + D_{2}^{T}Z_{2} + D_{2}^{T}P_{1}\sigma_{2} + r_{2})|^{2} \\ & + 2 \langle \Gamma_{2} + A_{2}^{T}Y_{2} + C_{2}^{T}Z_{2} + P_{2}b_{2} + C_{2}^{T}P_{1}\sigma_{2} + q_{2}, X_{2} \rangle - |\mathcal{R}_{2}^{-\frac{1}{2}}(S_{2}X_{2} + B_{2}^{T}Y_{2} + D_{2}^{T}Z_{2} + D_{2}^{T}P_{1}\sigma_{2} + r_{2})|^{2} \\ & + 2 \langle Y_{2}, b_{2} \rangle + 2 \langle Z_{2}, \sigma_{2} \rangle + \langle P_{1}\sigma_{2}, \sigma_{2} \rangle \right) dt \\ & = \mathbb{E} \int_{s}^{\infty} \left(\langle [\Lambda[P_{2}] + Q_{2} - S_{2}^{T}\mathcal{R}_{2}^{-1}S_{2}]X_{2}, X_{2} \rangle + |\mathcal{R}_{2}^{\frac{1}{2}}[u_{2} - \Theta_{2}X_{2} + \mathcal{R}_{2}^{-1}(B_{2}^{T}Y_{2} + D_{2}^{T}Z_{2} + D_{2}^{T}P_{1}\sigma_{2} + r_{2})] |^{2} \\ & - 2 \langle \Gamma_{2} + (A_{2}^{\Theta_{2}})^{T}Y_{2} + (C_{2}^{\Theta_{2}})^{T}Z_{2} + P_{2}b_{2} + (C_{2}^{\Theta_{2}})^{T}P_{1}\sigma_{2} + Q_{2} \otimes Z_{2}, \sigma_{2} \rangle + \langle P_{1}\sigma_{$$

where (compare with (5.5))

(5.6)
$$\begin{aligned} \mathcal{Q}_2 &= P_2 A_2 + A_2^\top P_2 + C_2^\top P_1 C_2 + Q_2, \quad \mathcal{S}_2 = B_2^\top P_2 + D_2^\top P_1 C_2 + S_2, \quad \mathcal{R}_2 = R_2 + D_2^\top P_1 D_2, \\ \Theta_2 &= -\mathcal{R}_2^{-1} \mathcal{S}_2, \quad A_2^{\Theta_2} = A_2 + B_2 \Theta_2, \quad C_2^{\Theta_2} = C_2 + D_2 \Theta_2. \end{aligned}$$

Similar to the above, we also have $\mathcal{R}_2(\cdot) \in \mathbb{S}^m_{++}$. Hence, in this case, one has

$$\begin{split} J^{\infty}(s,\imath,\xi_{1},\xi_{2};u_{1}(\cdot),u_{2}(\cdot)) &= \frac{1}{2}\mathbb{E}\sum_{k=1}^{2}\Big\{\langle P_{k}\xi_{k},\xi_{k}\rangle + 2\langle Y_{k}(s),\xi_{k}\rangle + \int_{s}^{\infty}\Big(\langle\big\{\Lambda[P_{k}]+\mathcal{Q}_{k}-\mathcal{S}_{k}^{\top}\mathcal{R}_{k}^{-1}\mathcal{S}_{k}\big\}X_{k},X_{k}\rangle \\ &+ \big|\mathcal{R}_{k}^{\frac{1}{2}}\big[u_{k}-\Theta_{k}X_{k}+\mathcal{R}_{k}^{-1}(B_{k}^{\top}Y_{k}+D_{k}^{\top}Z_{k}+D_{k}^{\top}P_{1}\sigma_{k}+r_{k})\big]\big|^{2} \\ &- 2\langle\Gamma_{k}+(A_{k}^{\Theta_{k}})^{\top}Y_{k}+(C_{2}^{\Theta_{k}})^{\top}Z_{k}+P_{k}b_{k}+(C_{k}^{\Theta_{k}})^{\top}P_{1}\sigma_{k}+q_{k}+\Theta_{k}^{\top}r_{k}),X_{k}\rangle \\ &- |\mathcal{R}_{k}^{-\frac{1}{2}}(B_{k}^{\top}Y_{k}+D_{k}^{\top}Z_{k}+D_{k}^{\top}P_{1}\sigma_{k}+r_{k})|^{2}+2\langle Y_{k},b_{k}\rangle+2\langle Z_{k},\sigma_{k}\rangle+\langle P_{1}\sigma_{k},\sigma_{k}\rangle\Big)dt\Big\}. \end{split}$$

Now, suppose we are able to do the following:

• The ARE

(5.7)
$$\Lambda[P_k](i) + \mathcal{Q}_k(i) - \mathcal{S}_k(i)^\top \mathcal{R}_k(i)^{-1} \mathcal{S}_k(i) = 0, \qquad \mathcal{R}_k(i) > 0, \qquad i \in \mathcal{M}, \quad k = 1, 2$$

admits a unique stabilizing solution $P_k(i)$, $i \in \mathcal{M}$, k = 1, 2, meaning that by defining $\Theta_k(\cdot)$ as in (5.5)–(5.6), one has

$$(\Theta_1, \Theta_2) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$$

This will make the quadratic terms of $X_1(\cdot)$ and $X_2(\cdot)$ in the completing-squares form of cost functional vanish.

• The following BSDE admits an adapted solution $(Y_1(\cdot), Y_2(\cdot), Z(\cdot), Z_1^M(\cdot), Z_2^M(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n) \times L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)$

(5.8)
$$\begin{cases} dY_1 = -\left((A_1^{\Theta_1})^\top Y_1 + (C_1^{\Theta_1})^\top \Pi_1[Z] + P_1 b_1 + (C_1^{\Theta_1})^\top P_1 \sigma_1 + q_1 + \Theta_1^\top r_1 \right) dt + Z dW + Z_1^M dM, \\ \lim_{t \to \infty} \mathbb{E} |Y_1(t)|^2 = 0, \end{cases}$$

and

(5.9)
$$\begin{cases} dY_2 = -\left((A_2^{\Theta_2})^\top Y_2 + (C_2^{\Theta_2})^\top \Pi_2[Z] + P_2 b_2 + (C_2^{\Theta_2})^\top P_1 \sigma_2 + q_2 + \Theta_2^\top r_2\right) dt + Z_2^M dM, \\ \lim_{t \to \infty} \mathbb{E}|Y_1(t)|^2 = 0. \end{cases}$$

This will make the weighting matrices of the linear terms of $X_1(\cdot)$ and $X_2(\cdot)$ in the completing-squares form of cost functional to be well-defined, and vanish. Note that if the ARE and the above BSDEs have the feature that the (closed-loop) system $\{A_1^{\Theta_1}, A_2^{\Theta_2}, C_1^{\Theta_1}, C_2^{\Theta_2}\}$ is L^2 -exponentially stable. Thus, as far as the well-posedness of BSDEs (5.2) is concerned, it is equivalent to assume that the system $[A, \overline{A}, C, \overline{C}]$ is already L^2 -exponentially stable, namely (A4)' holds. Hence, in the following subsections, we will assume (A4)' for convenience.

• One could choose

$$\bar{u}_k(t) = \Theta_k X_k - \mathcal{R}_k^{-1} (B_k^\top Y_k + D_k^\top \Pi_k[Z] + D_k^\top P_1 \sigma_k + r_k), \qquad k = 1, 2$$

This, together with others, will make the cost functional minimal.

In the above case, we have

$$J^{\infty}(s, \imath, \xi_1, \xi_2; \bar{u}_1(\cdot), \bar{u}_2(\cdot)) = \frac{1}{2} \mathbb{E} \sum_{k=1}^2 \Big\{ \langle P_k \xi_k, \xi_k \rangle + 2 \langle Y_k(s), \xi_k \rangle \\ - \int_s^{\infty} \Big(|\mathcal{R}_k^{-\frac{1}{2}} (B_k^\top Y_k + D_k^\top Z_k + D_k^\top P_1 \sigma_k + r_k)|^2 + 2 \langle Y_k, b_k \rangle + 2 \langle Z_k, \sigma_k \rangle + \langle P_1 \sigma_k, \sigma_k \rangle \Big) dt \Big\}.$$

From the above, we can see that $(\bar{u}_1(\cdot), \bar{u}_2(\cdot))$ is the unique optimal control Problem (MF-LQ)^{∞} if the well-posedness of the ARE (5.1) and BSDE (5.2) holds. The detailed proof will be presented in the next two sections, respectively.

5.2 Solvability of AREs

In this subsection, we will establish the solvability of the ARE (5.1), under proper conditions. The main idea is to consider the corresponding homogeneous LQ problem in finite time horizon [0, T], denoted by Problem (MF-LQ)₀^T. Under (A1)–(A3), by [21], Problem (MF-LQ)₀^T is closed-loop solvable so that the corresponding differential Riccati equation admits a solution $P_k^T(t, i)$ and then let $T \to \infty$ to get the limit $P_k^\infty(i)$ which will be the proper solution to the ARE (5.1), under (A4). Now, let us make the above precisely.

Theorem 5.2. Let **(A1)**–(**A4)** hold. Then, the ARE (5.1) admits a unique solution $(\widetilde{P}_1(i), \widetilde{P}_2(i))$, $i \in \mathcal{M}$, so that $(\widetilde{\Theta}_1(\cdot), \widetilde{\Theta}_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$, where

$$\widetilde{\Theta}_{k}(i) = -[R_{k}(i) + D_{k}(i)^{\top} \widetilde{P}_{1}(i) D_{k}(i)]^{-1} [B_{k}(i)^{\top} \widetilde{P}_{k}(i) + D_{k}(i)^{\top} \widetilde{P}_{1}(i) C_{k}(i) + S_{k}(i)], \qquad k = 1, 2.$$

Proof. First of all, we recall

(5.10)
$$J_0^{\infty}(s, i, \xi_1, \xi_2; u_1(\cdot), u_2(\cdot)) = \frac{1}{2} \sum_{k=1}^2 \mathbb{E} \int_s^{\infty} \left(\langle Q_k X_k^0, X_k^0 \rangle + 2 \langle S_k X_k^0, u_k \rangle + \langle R_k u_k, u_k \rangle \right) dt,$$

and for T > 0 large, we define

(5.11)
$$J_0^T(s, i, \xi_1, \xi_2; u_1(\cdot), u_2(\cdot)) = \frac{1}{2} \sum_{k=1}^2 \mathbb{E} \int_s^T \left(\langle Q_k X_k^0, X_k^0 \rangle + 2 \langle S_k X_k^0, u_k \rangle + \langle R_k u_k, u_k \rangle \right) dt,$$

where $(X_1^0(\cdot), X_2^0(\cdot)) = (X_1^0(\cdot; s, i, \xi_1, u_1, u_2), X_2^0(\cdot; s, i, \xi_2, u_2))$ is the solution of (2.12). By (A3), similar to [30], p.37, Proposition 2.5.1, we have

$$J_0^{\infty}(s, i, 0, 0; u_1(\cdot), u_2(\cdot)) \ge \delta \sum_{k=1}^2 \mathbb{E} \int_s^\infty |u_k(t)|^2 dt,$$

and

$$J_0^T(s, i, 0, 0; u_1(\cdot), u_2(\cdot)) \ge \delta \sum_{k=1}^2 \mathbb{E} \int_s^T |u_k(t)|^2 dt$$

for some $\delta > 0$. Hence, there exists $\widetilde{P}_k : \mathcal{M} \mapsto \mathbb{S}^n_{++}$ and $\widetilde{P}^T_k : [0,T] \times \mathcal{M} \to \mathbb{S}^n_{++}$ such that

$$V_0^{\infty}(s, i, \xi_1, \xi_2) = \inf_{(u_1(\cdot), u_2(\cdot)) \in \mathscr{U}_{ad}[s, \infty)} J_0^{\infty}(s, i, \xi_1, \xi_2; u_1(\cdot), u_2(\cdot)) = \sum_{k=1}^2 \mathbb{E} \langle \widetilde{P}_k(i) \xi_k, \xi_k \rangle,$$
$$V_0^T(s, i, \xi_1, \xi_2) = \inf_{(u_1(\cdot), u_2(\cdot)) \in \mathscr{U}[s, T]} J_0^T(t, i, \xi_1, \xi_2; u_1(\cdot), u_2(\cdot)) = \sum_{k=1}^2 \mathbb{E} \langle \widetilde{P}_k^T(t, i) \xi_k, \xi_k \rangle,$$

Moreover, the following Bellman principle of optimality holds:

$$\sum_{k=1}^{2} \mathbb{E} \langle \widetilde{P}_{k}^{T}(s,i)\xi_{i},\xi_{i} \rangle = \inf_{(u_{1}(\cdot),u_{2}(\cdot))\in\mathscr{U}[s,T]} \mathbb{E} \sum_{k=1}^{2} \left[\langle \widetilde{P}_{k}^{T}(t,\alpha(t))X_{i}^{0}(t),X_{i}^{0}(t) \rangle + \int_{s}^{t} \left(\langle Q_{i}(\alpha(r))X_{i}^{0},X_{i}^{0} \rangle + 2\langle S_{i}(\alpha(r))X_{i}^{0},u_{i} \rangle + \langle R_{i}(\alpha(r))u_{i},u_{i} \rangle \right) dr \right]$$

Now let $P_k^T(t) = \tilde{P}_k^T(t, \alpha(t))$. Similar to [21], for $k = 1, 2, P_k^T(\cdot)$ solves the following backward stochastic Riccati equation:

(5.12)
$$\begin{cases} dP_k^T + \zeta_k^T dM + \left[P_k^T A_k + A_k^\top P_k^T + C_k^\top P_1^T C_k + Q_k - (B_k^\top P_k^T + D_k^\top P_1^T C_k + S_k)^\top (R_k + D_k^\top P_1^T D_k)^{-1} (B_k^\top P_k^T + D_k^\top P_1^T C_k + S_k) \right] dt = 0, \\ P_k^T(T) = 0. \end{cases}$$

Now, let $(s, i, \xi_1, \xi_2) \in \mathscr{D}$ be given. For any $\varepsilon > 0$, there exists $(u_1^{\varepsilon}(\cdot), u_2^{\varepsilon}(\cdot)) \in \mathscr{U}_{ad}[s, \infty)$ such that

$$\begin{split} V_0^{\infty}(s, i, \xi_1, \xi_2) &\equiv \mathbb{E}[\langle \widetilde{P}_1(i)\xi_1, \xi_1 \rangle + \langle \widetilde{P}_2(i)\xi_2, \xi_2 \rangle] \\ \geqslant \frac{1}{2} \mathbb{E} \sum_{k=1}^2 \int_s^{\infty} \left(\langle Q_k X_k^{0,\varepsilon}, X_k^{0,\varepsilon} \rangle + 2 \langle S_k X_k^{0,\varepsilon}, u_k^{\varepsilon} \rangle + \langle R_k u_k^{\varepsilon}, u_k^{\varepsilon} \rangle \right) dt - \varepsilon \\ &= J_0^T(s, i, \xi_1, \xi_2; u_1^{\varepsilon}(\cdot), u_2^{\varepsilon}(\cdot)) + \mathbb{E} \sum_{k=1}^2 \int_T^{\infty} \left(\langle Q_k X_k^{0,\varepsilon}, X_k^{0,\varepsilon} \rangle + 2 \langle S_k X_k^{0,\varepsilon}, u_k^{\varepsilon} \rangle + \langle (R_k u_k^{\varepsilon}, u_k^{\varepsilon} \rangle \right) dt - \varepsilon, \end{split}$$

where

$$(X_1^{0,\varepsilon}, X_2^{0,\varepsilon}) = (X_1^0(\cdot; s, \imath, \xi_1; u_1^{\varepsilon}(\cdot), u_2^{\varepsilon}(\cdot)), X_2^0(\cdot; s, \imath, \xi_2; u_1^{\varepsilon}(\cdot), u_2^{\varepsilon}(\cdot))) \in L^2_{\mathbb{F}^{\alpha}}(s, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(s, \infty; \mathbb{R}^n).$$

Letting $T \to \infty$ and then $\varepsilon \to 0$, we have

$$\mathbb{E}[\langle \widetilde{P}_1(\imath)\xi_1,\xi_1\rangle + \langle \widetilde{P}_2(\imath)\xi_2,\xi_2\rangle] \geqslant \lim_{T \to \infty} \mathbb{E}[\langle P_1^T(s)\xi_1,\xi_1\rangle + \langle P_2^T(s)\xi_2,\xi_2\rangle].$$

Moreover, for any $(u_1(\cdot), u_2(\cdot)) \in \mathscr{U}[s, T]$,

$$J_0^T(s, i, \xi_1, \xi_2; u_1(\cdot) \oplus 0I_{(T,\infty)}, u_2(\cdot) \oplus 0I_{(T,\infty)}) = J_0^T(s, i, \xi_1, \xi_2; u_1(\cdot), u_2(\cdot)).$$

Therefore, for any $(s, i, \xi_1, \xi_2) \in \mathscr{D}$,

$$\mathbb{E}[\langle \widetilde{P}_{1}(i)\xi_{1},\xi_{1}\rangle + \langle \widetilde{P}_{2}(i)\xi_{2},\xi_{2}\rangle] = V_{0}^{\infty}(s,i,\xi_{1},\xi_{2}) \leqslant V_{0}^{T}(s,i,\xi_{1},\xi_{2}) = \mathbb{E}[\langle P_{1}^{T}(s)\xi_{1},\xi_{1}\rangle + \langle P_{2}^{T}(s)\xi_{2},\xi_{2}\rangle].$$

By the arbitrariness of $(\xi_1, \xi_2) \in L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{R}^n)^{\perp} \times L^2_{\mathcal{F}^{\alpha}_s}(\Omega; \mathbb{R}^n)$ and (5.12), it is necessary that

$$\lim_{T \to \infty} \widetilde{P}_k^T(s, i) = \lim_{T \to \infty} P_k^T(s) = \widetilde{P}_k(i) \quad \text{with} \quad R_k(i) + D_k(i)^\top \widetilde{P}_1(i) D_k(i) > 0, \ i \in \mathcal{M}.$$

We notice that $\lim_{T\to\infty} \widetilde{P}_k^T(t,i)$ is independent of t for any finite $t \ge s$. Therefore

$$\lim_{T \to \infty} P_k^T(t) = \widetilde{P}_k(\alpha(t))$$

Because $P_k^T(t)$ is decreasing in T, taking $T \to \infty$ in (5.12), we have

(5.13)
$$\widetilde{P}_{k}(\alpha(t)) + \int_{s}^{t} \left(\widetilde{P}_{k}A_{k} + A_{k}^{\top}\widetilde{P}_{k} + C_{k}^{\top}\widetilde{P}_{k}C_{k} + Q_{k} - (B_{k}^{\top}\widetilde{P}_{k} + D_{k}^{\top}\widetilde{P}_{1}C_{k} + S_{k})^{\top}(R_{k} + D_{\kappa}^{\top}\widetilde{P}_{1}D_{k})^{-1}(B_{k}^{\top}\widetilde{P}_{k} + D_{k}^{\top}\widetilde{P}_{1}C_{k} + S_{k}) \right) d\tau$$

with $\alpha(s) = \iota$ is a martingale. The Itô's formula on $\widetilde{P}_k(\alpha(t))$, together the arbitrariness of ι , yields that \widetilde{P}_k is the solution to (5.1).

Now let us verify $(\widetilde{\Theta}_1(\cdot), \widetilde{\Theta}_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$. Suppose that $(\widetilde{X}_1(\cdot), \widetilde{X}_2(\cdot), \widetilde{u}_1(\cdot), \widetilde{u}_2(\cdot))$ is the optimal pair of Problem (MF-LQ)₀^{∞}. Itô's formula yields that

$$\frac{d}{dt} \sum_{k=1}^{2} \mathbb{E}[\langle \widetilde{P}_{k}(\alpha(t))\widetilde{X}_{k}(t), \widetilde{X}_{k}(t) \rangle]$$

$$= \sum_{k=1}^{2} \mathbb{E}\Big[\left\langle \left(\Lambda[\widetilde{P}_{k}] + A_{k}^{\top}\widetilde{P}_{k} + \widetilde{P}_{k}A_{k} + C_{k}^{\top}\widetilde{P}_{1}C_{k} \right)\widetilde{X}_{k}, \widetilde{X}_{i} \right\rangle \\
+ \sum_{k=1}^{2} \mathbb{E}\Big[2 \langle (\widetilde{P}_{k}B_{k} + C_{k}^{\top}\widetilde{P}_{1}D_{k})\widetilde{u}_{k}, \widetilde{X}_{k} \rangle + \langle D_{k}^{\top}\widetilde{P}_{1}D_{k}\widetilde{u}_{k}, \widetilde{u}_{k} \rangle \Big] \\
= \sum_{k=1}^{2} \mathbb{E}\Big[\langle (R_{k} + D_{k}^{\top}\widetilde{P}_{1}D_{k})(\widetilde{u}_{k} - \widetilde{\Theta}_{k}\widetilde{X}_{k}), \widetilde{u}_{k} - \widetilde{\Theta}_{k}\widetilde{X}_{k} \rangle \Big] \\
- \sum_{k=1}^{2} \mathbb{E}\Big[\langle Q_{k}\widetilde{X}_{k}, \widetilde{X}_{k} \rangle + 2 \langle S_{k}\widetilde{u}_{k}, \widetilde{X}_{k} \rangle + \langle R_{k}\widetilde{u}_{k}, \widetilde{u}_{k} \rangle \Big].$$

By (5.1), long but straightforward calculation yields that

$$\mathbb{E}\sum_{k=1}^{2} \langle \widetilde{P}_{k}(i)\xi_{k},\xi_{k}\rangle = V_{0}^{\infty}(s,i,\xi_{1},\xi_{2}) = J_{0}^{\infty}(s,i,\xi_{1},\xi_{2};\widetilde{u}_{1}(\cdot),\widetilde{u}_{2}(\cdot))$$
$$= \mathbb{E}\sum_{k=1}^{2} \langle \widetilde{P}_{k}(i)\xi_{k},\xi_{k}\rangle + \mathbb{E}\sum_{k=1}^{2} \int_{s}^{\infty} |(R_{k}+D_{k}^{\top}\widetilde{P}_{1}D_{k})^{\frac{1}{2}}(\widetilde{u}_{k}(t)-\widetilde{\Theta}_{k}(\alpha(t))\widetilde{X}_{k}(t))|^{2}dt$$

This says that the optimal control $\widetilde{u}_k(\cdot) = \widetilde{\Theta}_k(\alpha(\cdot))\widetilde{X}_k(\cdot) \in \mathscr{U}_{ad}[s,\infty)$, i.e. $(\widetilde{X}_1(\cdot),\widetilde{X}_2(\cdot))$ solves the following

$$\begin{cases} d\widetilde{X}_1 = (A_1 + B_1\widetilde{\Theta}_1)\widetilde{X}_1 dt + \left((C_1 + D_1\widetilde{\Theta}_1)\widetilde{X}_1 + (C_2 + D_2\widetilde{\Theta}_2)\widetilde{X}_2 \right) dW(t) \\ d\widetilde{X}_2 = (A_2 + B_2\widetilde{\Theta}_2)\widetilde{X}_2, \quad t \in [s, \infty) \\ \widetilde{X}_1(s) = \xi_1, \ \widetilde{X}_2(s) = \xi_2, \ \alpha(s) = \iota. \end{cases}$$

Because $(\widetilde{X}_1(\cdot), \widetilde{X}_2(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(s, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(s, \infty; \mathbb{R}^n)$, it follows that $(\widetilde{\Theta}_1(\cdot), \widetilde{\Theta}_2(\cdot))$ is a stabilizer i.e., $(\widetilde{\Theta}_1(\cdot), \widetilde{\Theta}_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$. If $(\widetilde{P}_1, \widetilde{P}_2)$ is the solution of (5.1), we can actually conclude that

$$J_0^{\infty}(s,\iota,\xi_1,\xi_2;u_1(\cdot),u_2(\cdot)) \geqslant \sum_{k=1}^2 \mathbb{E}\langle \widetilde{P}_k(\iota)\xi_k,\xi_k\rangle.$$

The equality holds when $\widetilde{u}_i(\cdot) = \widetilde{\Theta}_k(\alpha(\cdot))\widetilde{X}_k(\cdot) \in \mathscr{U}_{ad}[s,\infty)$. Therefore

$$V_0^{\infty}(s, \imath, \xi_1, \xi_2) = \sum_{k=1}^2 \mathbb{E} \langle \widetilde{P}_k(\imath) \xi_k, \xi_k \rangle$$

which implies the uniqueness directly.

5.3 BSDE in Infinite Horizon

In this section, we are going to study the well-posedness of BSDE (5.2) in an infinite horizon, with the coefficients depending on the Markov chain. Such a result can be concluded from the following theorem directly.

Theorem 5.3. Under (A1)–(A4), for any $(\varphi_1(\cdot), \varphi_2(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)$, the following system of BSDEs on $[0, \infty)$ admits a unique adapted solution $(Y_1(\cdot), Z(\cdot), Z_1^M(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}}(0, \infty; \mathbb{R}^n) \times M^2_{\mathbb{F}^{\alpha}_{-}}(0, \infty; \mathbb{R}^n)^{\perp}$ and $(Y_2(\cdot), Z_2^M(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(0, \infty; \mathbb{R}^n) \times M^2_{\mathbb{F}^{\alpha}_{-}}(0, \infty; \mathbb{R}^n)$:

(5.14)
$$\begin{cases} dY_1(t) = -\left[A_1^{\Theta_1}(\alpha(t))^\top Y_1(t) + C_1^{\Theta_1}(\alpha(t))^\top \Pi_1[Z(t)] + \varphi_1(t)\right] dt + Z(t) dW(t) + Z_1^M(t) dM(t) \\ dY_2(t) = -\left[A_2^{\Theta_2}(\alpha(t))^\top Y_2(t) + C_2^{\Theta_2}(\alpha(t))^\top \Pi_2[Z(t)] + \varphi_2(t)\right] dt + Z_2^M(t) dM(t), \\ \lim_{t \to \infty} \mathbb{E}|Y_i(t)|^2 = 0, \qquad i = 1, 2. \end{cases}$$

To illustrate the non-triviality of the above result, let us recall the classical situation. The following is found in [30].

Proposition 5.4. Let [A, C] be L^2 -stable, i.e., there exists a $P \in \mathbb{S}^n_{++}$ such that

$$PA + A^{\top}P + C^{\top}PC \leqslant -I.$$

Then for any $\varphi(\cdot) \in L^2_{\mathbb{R}^W}(0,\infty;\mathbb{R}^n)$, the following BSDE

(5.15)
$$\begin{cases} dY(t) = -[A^{\top}Y(t) + C^{\top}Z(t) + \varphi(t)]dt + Z(t)dW(t), & t \in [0, \infty), \\ \lim_{t \to \infty} \mathbb{E}|Y(t)|^2 = 0. \end{cases}$$

admits a unique adapted solution $(Y(\cdot), Z(\cdot)) \in L^2_{\mathbb{F}^W}(\Omega; C([0, \infty); \mathbb{R}^n)) \times L^2_{\mathbb{F}^W}(0, \infty; \mathbb{R}^n)$, and the following holds:

$$\mathbb{E}\Big(\sup_{t\in[0,\infty)}|Y(t)|^2 + \int_0^\infty |Z(t)|^2 dt\Big) \leqslant K\mathbb{E}\int_0^\infty |\varphi(t)|^2 dt$$

From the above result, we see that if the following are L^2 -exponentially stable for all $i \in \mathcal{M}$:

(5.16)
$$\begin{cases} dX_1(t) = \{A_1(i)X_1(t) + B_1(i)u_1(t)\}dt \\ +\{C_1(i)X_1(t) + C_2(i)X_2 + D_1(i)u_1(t) + D_2(i)u_2(t)\}dW(t)\} \\ dX_2(t) = \{A_2(i)X_2(t) + B_2(i)u_2(t)\}dt, \quad t \in [s, \infty), \\ X_1(s) = \xi_1, \quad X_2(s) = \xi_2, \end{cases}$$

then similar to [30], we are able to get the unique solvability of (5.14). However, we have indicated in Section 3 that the L^2 -exponentially stabilizability of system $[A, \bar{A}, C, \bar{C}; B, \bar{B}, D, \bar{D}]$ does not imply that of (5.16). Thus, we need to find a new approach.

Note that BSDE (5.14) is associated with system (4.7) which is L^2 -exponentially stable if we take $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \overline{A}, C, \overline{C}; B, \overline{B}, D, \overline{D}]$. Thus, to investigate the solvability of BSDE (5.14), it suffices to assume the original system $[A, \overline{A}, C, \overline{C}]$ is L^2 -exponential stable. Hence, we only need to consider the solvability of the following BSDE:

(5.17)
$$\begin{cases} dY_1(t) = -\left[A_1(\alpha(t))^\top Y_1(t) + C_1(\alpha(t))^\top \Pi_1[Z(t)] + \varphi_1(t)\right] dt + Z(t) dW(t) + Z_1^M(t) dM(t) \\ dY_2(t) = -\left[A_2(\alpha(t))^\top Y_2(t) + C_2(\alpha(t))^\top \Pi_2[Z(t)] + \varphi_2(t)\right] dt + Z_2^M(t) dM(t), \\ \lim_{t \to \infty} \mathbb{E}|Y_i(t)|^2 = 0, \quad i = 1, 2. \end{cases}$$

Proof of Theorem 5.3. First of all, as we pointed out in Section 5.1, it suffices for us to look at the case that $[A, \overline{A}, C, \overline{C}]$ is L^2 -exponential stable, i.e., $(\mathbf{A4})'$ holds. In what follows, we keep this assumption and let $\Theta_1 = \Theta_2 = 0$. The rest of the proof will be divided into several steps.

Step 1. Since $[A, \overline{A}, C, \overline{C}]$ is L^2 -exponential stable, we have $P_1, P_2 : \mathcal{M} \to \mathbb{S}^n_{++}$ such that

(5.18)
$$\Lambda[P_k] + P_k A_k + A_k^\top P_k + C_k^\top P_1 C_k < 0, \qquad k = 1, 2.$$

Because each $P_k(i) \in \mathbb{S}_{++}^n$, it admits a unique square root $P_k(i)^{\frac{1}{2}} \in \mathbb{S}_{++}^n$. From (2.18), it can be seen that

$$dP_k(\alpha(t))^{\frac{1}{2}} = \Lambda[P_k^{\frac{1}{2}}](\alpha(t))dt + \sum_{i\neq j} E_i(i,j)\lambda_{ij}\mathbf{1}_{(\alpha(t^-)=i)}dM_{ij}(t)$$

with $E_i(i, j) = P_k(j)^{\frac{1}{2}} - P_k(i)^{\frac{1}{2}} \in \mathbb{S}^n$. Suppose that $(X_1^0(\cdot), X_2^0(\cdot))$ is the solution of (2.15), with s = 0. By (5.18), Itô's formula and Gronwall's inequality together yield that for some $\varepsilon > 0$,

$$\sum_{k=1}^{2} \mathbb{E} \langle P_k(\alpha(t)) X_k^0(t), X_k^0(t) \rangle \leqslant e^{-\varepsilon t} \sum_{k=1}^{2} \langle P_k(\alpha(0)) \xi_k, \xi_k \rangle,$$

which leads to

(5.19)
$$\lim_{t \to 0^+} \frac{1}{t} \Big(\sum_{k=1}^2 \mathbb{E} \langle P_k(\alpha(t)) X_k^0(t), X_k^0(t) \rangle - \sum_{k=1}^2 \langle P_k(\alpha(0)) \xi_k, \xi_k \rangle \Big) \leqslant -\varepsilon \sum_{k=1}^2 \langle P_k(\alpha(0)) \xi_k, \xi_k \rangle.$$

Let $\widetilde{X}_k(t) = P_k(\alpha(t))^{\frac{1}{2}} X_k^0(t)$. Then, Itô's formula implies that

(5.20)
$$\begin{cases} d\widetilde{X}_{1}(t) = \widetilde{A}_{1}(\alpha(t))\widetilde{X}_{1}(t)dt + \left[\widetilde{C}_{1}(\alpha(t))\widetilde{X}_{1}(t) + \widetilde{C}_{2}(\alpha(t))\widetilde{X}_{2}(t)\right]dW \\ + \sum_{i \neq j} \widetilde{E}_{1}(i,j)\widetilde{X}_{1}(t^{-})\lambda_{ij}\mathbf{1}_{(\alpha(t^{-})=i)}dM_{ij}(t), \\ d\widetilde{X}_{2}(t) = \widetilde{A}_{2}(\alpha(t))\widetilde{X}_{2}(t)dt + \sum_{i \neq j} \widetilde{E}_{2}(i,j)\widetilde{X}_{2}(t^{-})\lambda_{ij}\mathbf{1}_{(\alpha(t^{-})=i)}dM_{ij}(t), \quad t \in [0,\infty) \end{cases}$$

where

$$\widetilde{A}_{k}(i) = P_{k}(i)^{\frac{1}{2}}A_{k}(i)P_{k}(i)^{-\frac{1}{2}} + \Lambda[P_{k}^{\frac{1}{2}}](i)P_{k}(i)^{-\frac{1}{2}}, \quad \widetilde{E}_{k}(i,j) = E_{k}(i,j)P_{k}(i)^{-\frac{1}{2}},$$

$$\widetilde{C}_{1}(i) = P_{1}(i)^{\frac{1}{2}}C_{1}(i)P_{1}(i)^{-\frac{1}{2}}, \quad \widetilde{C}_{2}(i) = P_{1}(i)^{\frac{1}{2}}C_{2}(i)P_{2}(i)^{-\frac{1}{2}}.$$

Note that $\langle P_k(\alpha(t))X_k^0(t),X_k^0(t)\rangle = |\widetilde{X}_k(t)|^2$ and

$$\frac{d}{dt} \sum_{k=1}^{2} \mathbb{E} |\widetilde{X}_{k}(t)|^{2} = \sum_{k=1}^{2} \mathbb{E} \left\langle \left(\widetilde{A}_{k}(\alpha(t)) + \widetilde{A}_{k}(\alpha(t))^{\top} + \widetilde{C}_{k}(\alpha(t))^{\top} \widetilde{C}_{k}(\alpha(t)) + \sum_{j \neq \alpha(t)} \lambda_{\alpha(t)j} \widetilde{E}_{k}^{\top}(\alpha(t), j) \widetilde{E}_{k}(\alpha(t), j) \right) \widetilde{X}_{k}(t), \widetilde{X}_{k}(t) \right\rangle$$

Then (5.19) leads to

(5.21)
$$\widetilde{A}_{k}(i) + \widetilde{A}_{k}(i)^{\top} + \widetilde{C}_{k}(i)^{\top} \widetilde{C}_{k}(i) + \sum_{j \neq i} \lambda_{ij} \widetilde{E}_{k}(i,j)^{\top} \widetilde{E}_{k}(i,j) < 0, \quad i \in \mathcal{M}$$

Step 2. Suppose the following BSDE on $[0,\infty)$ admits a unique adapted solution $(\widetilde{Y}_1(\cdot),\widetilde{Z}(\cdot),\widetilde{Z}_1^M(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(0,\infty;\mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}^{\alpha}}(0,\infty;\mathbb{R}^n) \times M^2_{\mathbb{F}^{\alpha}_{-}}(0,\infty;\mathbb{R}^n)^{\perp}$ and $(\widetilde{Y}_2(\cdot),\widetilde{Z}_2^M(\cdot)) \in L^2_{\mathbb{F}^{\alpha}}(0,\infty;\mathbb{R}^n) \times M^2_{\mathbb{F}^{\alpha}_{-}}(0,\infty;\mathbb{R}^n)$:

$$(5.22) \quad \begin{cases} 0 = d\widetilde{Y}_{1}(t) + \widetilde{Z}(t)dW(t) + \widetilde{Z}_{1}^{M}(t)dM(t) \\ + \left[\widetilde{A}_{1}(\alpha(t))^{\top}\widetilde{Y}_{1}(t) - \widetilde{C}_{1}(\alpha(t))^{\top}\Pi_{1}[\widetilde{Z}(t)] + \widetilde{\varphi}_{1}(t) - \sum_{i \neq j}\widetilde{E}_{1}^{\top}(i,j)\widetilde{Z}_{1}^{M}(t,j)\lambda_{ij}I(\alpha(t)=i)\right]dt, \\ 0 = d\widetilde{Y}_{2}(t) + \widetilde{Z}_{2}^{M}(t)dM(t) \\ + \left[\widetilde{A}_{2}(\alpha(t))^{\top}\widetilde{Y}_{2}(t) - \widetilde{C}_{2}(\alpha(t))^{\top}\Pi_{2}[Z(t)] + \widetilde{\varphi}_{2}(t) - \sum_{i \neq j}\widetilde{E}_{2}^{\top}(i,j)\widetilde{Z}_{2}^{M}(t,j)\lambda_{ij}I(\alpha(t)=i)\right]dt, \\ \lim_{t \to \infty} \mathbb{E}|\widetilde{Y}_{i}(t)|^{2} = 0, \end{cases}$$

where $\tilde{\varphi}_k(t) = P_k(\alpha(t))^{-\frac{1}{2}} \varphi_k(t)$. Then, by letting $Y_k(t) = P_k(\alpha(t))^{\frac{1}{2}} \tilde{Y}_k(t)$, using Itô's formula, we have

$$\begin{split} dY_{1}(t) &= P_{1}(\alpha(t^{-}))^{\frac{1}{2}}d\widetilde{Y}_{1}(t) + \{d[P_{1}(\alpha(t))^{\frac{1}{2}}]\}\widetilde{Y}_{1}(t^{-}) - \sum_{i\neq j} E_{1}(i,j)\widetilde{Z}_{1}^{M}(t,j)\lambda_{ij}\mathbf{1}_{(\alpha(t)=i)}dt \\ &- \sum_{j\neq i} E_{1}(i,j)\widetilde{Z}_{1}^{M}(t,j)\mathbf{1}_{(\alpha(t^{-})=i)}dM_{ij}(t) \\ &= -P_{1}(\alpha(t^{-}))^{\frac{1}{2}}\Big[\widetilde{Z}(t)dW(t) + \sum_{j\neq i}\widetilde{Z}_{1}^{M}(t,j)\mathbf{1}_{(\alpha(t^{-})=i)}dM_{ij}(t) \\ &+ \Big(\widetilde{A}_{1}(\alpha(t))^{\top}\widetilde{Y}_{1}(t) - \widetilde{C}_{1}(\alpha(t))^{\top}\Pi_{1}[\widetilde{Z}(t)] + \widetilde{\varphi}_{1}(t)\Big)dt - \sum_{i\neq j}\widetilde{E}_{1}^{\top}(i,j)\widetilde{Z}_{1}^{M}(t,j)\lambda_{ij}\mathbf{1}_{(\alpha(t)=i)}dt\Big] \\ &+ \Big(\Lambda[P^{\frac{1}{2}}](\alpha(t))\widetilde{Y}_{1}(t)dt + \sum_{j\neq i} E_{1}(i,j)\mathbf{1}_{(\alpha(t^{-})=i)}\widetilde{Y}_{1}(t^{-})dM_{ij}(t)\Big) \\ &- \sum_{i\neq j} E_{1}(i,j)\widetilde{Z}_{1}^{M}(t,j)\lambda_{ij}\mathbf{1}_{(\alpha(t)=i)}dt - \sum_{j\neq i} E_{1}(i,j)\widetilde{Z}_{1}^{M}(t,j)\mathbf{1}_{(\alpha(t^{-})=i)}dM_{ij}(t) \\ &= -P_{1}(\alpha(t))^{\top}\widetilde{Z}(t)dW - \Big[A_{1}(\alpha(t))^{\top}Y_{1} - C_{1}(\alpha(t))^{\top}\Pi_{1}[P_{1}(\alpha(t)^{\frac{1}{2}}\widetilde{Z}(t)] + \varphi_{1}\Big]dt \\ &- \sum_{j\neq i} \Big(E_{1}(i,j)\widetilde{Z}_{1}^{M}(t,j) + P_{1}(\alpha(t^{-}))^{\frac{1}{2}}\widetilde{Z}_{1}^{M}(t,j) - E_{1}(i,j)\widetilde{Y}_{1}(t^{-})\Big)\mathbf{1}_{(\alpha(t^{-})=i)}dM_{ij}(t) \\ &= -ZdW - Z_{1}^{M}dM - \Big[A_{1}^{\top}Y_{1} - C_{1}^{\top}\Pi_{1}[Z] + \varphi_{1}\Big]dt. \end{split}$$

where $Z(t) = P_1(\alpha(t))^\top \widetilde{Z}(t)$ and $Z_1^M(t,j) = E_1(\alpha(t^-),j)\widetilde{Z}_1^M(t,j) + P_1(\alpha(t^-))^{\frac{1}{2}}\widetilde{Z}_1^M(t,j) - E_1(\alpha(t^-),j)\widetilde{Y}_1(t^-)$. Here we have used the symmetricity of $E_k(i,j)$. Similarly one can derive that

$$dY_2(t) = -Z_2^M(t)dM(t) - \left[A_2^\top Y_2(t) - C_2^\top \Pi_2[Z(t)] + \varphi_2(t)\right]dt.$$

Therefore we have constructed the required solution to (5.17). We also can see that $Y_k(t)$ and $\tilde{Y}_k(t)$ have one-to-one correspondence and therefore $Y_k(\cdot)$ is uniquely determined. The last limits are clear.

Step 3. Now it suffices to prove that (5.22) has an adapted solution. The proof is split into several substeps.

Substep 1. A priori estimate for (5.22) when $\tilde{\varphi}_1(\cdot) = 0$ and $\tilde{\varphi}_2(\cdot) = 0$. Itô's formula yields that

$$\begin{aligned} \frac{d}{dt} \mathbb{E} \sum_{i=1}^{2} |\tilde{Y}_{i}(t)|^{2} &= -\mathbb{E} \sum_{i=1}^{2} \langle \tilde{Y}_{i}(t), (\tilde{A}_{i}(\alpha(t)) + \tilde{A}_{i}^{\top}(\alpha(t))) \tilde{Y}_{i}(t) \rangle dt \\ &+ \mathbb{E} \sum_{i=1}^{2} \left(|\Pi_{i}[\tilde{Z}(t)]|^{2} + 2 \langle \tilde{C}_{i}^{\top}(\alpha(t)) \Pi_{i}[Z(t)], \tilde{Y}_{i}(t) \rangle \right) dt \\ &+ \sum_{i=1}^{2} \mathbb{E} \sum_{i \neq j} \left(|\tilde{Z}_{i}^{M}(t, j)|^{2} + 2 \langle \tilde{E}_{i}^{\top}(i, j) \tilde{Z}_{i}^{M}(t, j), Y_{i}(t) \rangle \right) \lambda_{ij} \mathbf{1}_{\{\alpha(t)=i\}} dt \\ \end{aligned}$$

$$(5.23) \qquad = -\mathbb{E} \sum_{i=1}^{2} \left\langle \tilde{Y}_{i}, \left(\tilde{A}_{i}(\alpha(t)) + \tilde{A}_{i}^{\top}(\alpha(t)) + \tilde{C}_{i}^{\top}(\alpha(t)) \tilde{C}_{i}(\alpha(t)) \right) \\ &+ \sum_{i \neq j} \tilde{E}_{i}^{\top}(\alpha(t), j) \tilde{E}_{i}(\alpha(t), j) \lambda_{ij} \mathbf{1}_{\{\alpha(t)=i\}} \right) \tilde{Y}_{i} \right\rangle \\ &+ \mathbb{E} \sum_{i=1}^{2} \left(|\Pi_{i}[\tilde{Z}(t)] + \tilde{C}_{i}(\alpha(t)) \tilde{Y}_{i}(t)|^{2} + \sum_{i \neq j} |\tilde{Z}_{i}^{M}(t, j) + \tilde{E}_{i}(\iota, j) \tilde{Y}_{i}(t)|^{2} \lambda_{ij} \mathbf{1}_{\{\alpha(t)=i\}} \right) \\ &\geqslant \mathbb{E} \sum_{i=1}^{2} \left(\varepsilon |\tilde{Y}_{i}(t)|^{2} + |\Pi_{i}[\tilde{Z}(t)] + \tilde{C}_{i}(\alpha(t)) \tilde{Y}_{i}(t)|^{2} + \sum_{i \neq j} |\tilde{Z}_{i}^{M}(t, j) + \tilde{E}_{i}(\iota, j) \tilde{Y}_{i}(t)|^{2} \lambda_{ij} \mathbf{1}_{\{\alpha(t)=i\}} \right) \end{aligned}$$

where we used (5.21) in the last step.

Substep 2. Consider the following BSDE on [0, T) for finite T > 0:

$$(5.24) \begin{cases} 0 = d\widetilde{Y}_{1}(t;T) + \widetilde{Z}(t;T)dW(t) + \widetilde{Z}_{1}^{M}(t;T)dM(t) \\ + \left[\widetilde{A}_{1}^{\top}(\alpha(t))\widetilde{Y}_{1}(t;T) - \widetilde{C}_{1}^{\top}(\alpha(t))\Pi_{1}[\widetilde{Z}(t;T)] + \widetilde{\varphi}_{1}(t) - \sum_{\iota \neq j}\widetilde{E}_{1}^{\top}(\iota,j)\widetilde{Z}_{1}^{M}(t,j;T)\lambda_{\iota j}\mathbf{1}_{\{\alpha(t)=\iota\}}\right]dt, \\ 0 = d\widetilde{Y}_{2}(t;T) + \widetilde{Z}_{2}^{M}(t;T)dM(t) \\ + \left[\widetilde{A}_{2}^{\top}(\alpha(t))\widetilde{Y}_{2}(t;T) - \widetilde{C}_{2}^{\top}(\alpha(t))\Pi_{2}[\widetilde{Z}(t;T)] + \widetilde{\varphi}_{2}(t) - \sum_{\iota \neq j}\widetilde{E}_{2}^{\top}(\iota,j)\widetilde{Z}_{2}^{M}(t,j;T)\lambda_{\iota j}\mathbf{1}_{\{\alpha(t)=\iota\}}\right]dt, \\ \widetilde{Y}_{1}(T;T) = \widetilde{Y}_{2}(T;T) = 0. \end{cases}$$

It is a linear BSDE (with mean-field term) on a finite horizon. Due to the orthogonal structure, it admits a unique solution such that

$$\begin{aligned} &(\widetilde{Y}_1(\cdot;T),\widetilde{Z}(\cdot;T),\widetilde{Z}_1^M(\cdot;T)) \in L^2_{\mathbb{F}^{\alpha}}(0,T;\mathbb{R}^n)^{\perp} \times L^2_{\mathbb{F}}(0,T,\mathbb{R}^n) \times M^2_{\mathbb{F}^{\alpha}_{-}}(0,T;\mathbb{R}^n)^{\perp},\\ &(\widetilde{Y}_2(\cdot;T),\widetilde{Z}_2^M(\cdot;T)) \in L^2_{\mathbb{F}^{\alpha}}(0,T;\mathbb{R}^n) \times M^2_{\mathbb{F}^{\alpha}_{-}}(0,T;\mathbb{R}^n). \end{aligned}$$

Applying Itô's formula to $\sum_{i=1}^{2} |\widetilde{Y}_{i}(t;T)|^{2}$, similar to (5.23), one has

(5.25)
$$\frac{d}{dt} \mathbb{E} \sum_{i=1}^{2} |\widetilde{Y}_{i}(t;T)|^{2} \geq \mathbb{E} \sum_{i=1}^{2} \frac{\varepsilon}{2} |\widetilde{Y}_{i}(t;T)|^{2} + |\Pi_{i}[\widetilde{Z}(t;T)] + \widetilde{C}_{i}(\alpha(t))\widetilde{Y}_{i}(t;T)|^{2} + \mathbb{E} \sum_{i\neq j} |\widetilde{Z}_{i}^{M}(t,j;T) + \widetilde{E}_{i}(\imath,j)\widetilde{Y}_{i}(t;T)|^{2} \lambda_{\imath j} \mathbf{1}_{\{\alpha(t)=\imath\}} - \frac{K}{\varepsilon} \mathbb{E} \sum_{i=1}^{2} |\widetilde{\varphi}_{i}(t)|^{2}.$$

Then Gronwall's inequality yields that

(5.26)
$$\mathbb{E}\sum_{i=1}^{2}|\widetilde{Y}_{i}(t;T)|^{2} \leqslant \frac{K}{\varepsilon}\sum_{i=1}^{2}\mathbb{E}\int_{t}^{T}e^{-\frac{\varepsilon}{2}(r-t)}|\widetilde{\varphi}_{i}(r)|^{2}dr$$

for some $\varepsilon > 0$ where K is an uniform constant. Using (5.23), we have, for $0 \leq t \leq T$,

$$\frac{d}{dt} \mathbb{E} \sum_{i=1}^{2} |\widetilde{Y}_{i}(t;T) - \widetilde{Y}_{i}(t;T+T_{0})|^{2} \ge \varepsilon \mathbb{E} \sum_{i=1}^{2} |\widetilde{Y}_{i}(t;T) - \widetilde{Y}_{i}(t;T+T_{0})|^{2} \\
+ \mathbb{E} \sum_{i=1}^{2} \left| \Pi_{i}[\widetilde{Z}(t;T)] - \Pi_{i}[\widetilde{Z}(t;T+T_{0})] + \widetilde{C}_{i} \Big(\widetilde{Y}_{i}(t;T) - \widetilde{Y}_{i}(t;T+T_{0}) \Big) \Big|^{2} \\
+ \sum_{i=1}^{2} \mathbb{E} \sum_{i\neq j} \left| \widetilde{Z}_{i}^{M}(t,j;T) - \widetilde{Z}_{i}^{M}(t,j;T+T_{0}) + \widetilde{E}_{i}(\iota,j) \Big(\widetilde{Y}_{i}(t;T) - \widetilde{Y}_{i}(t;T+T_{0}) \Big) \Big|^{2} \lambda_{\iota j} \mathbf{1}_{\{\alpha(t)=i\}}.$$

Substep 3. We now let $T \to \infty$. By (5.26) and (5.27), Gronwall's inequality yields that

$$\begin{split} & \mathbb{E}\sum_{i=1}^{2}|\widetilde{Y}_{i}(t;T+T_{0})-\widetilde{Y}_{i}(t;T)|^{2}\leqslant e^{\varepsilon(t-T)}\mathbb{E}\sum_{i=1}^{2}|\widetilde{Y}_{i}(T;T+T_{0})-\widetilde{Y}_{i}(T;T)|^{2} \\ &= e^{\varepsilon(t-T)}\mathbb{E}\sum_{i=1}^{2}|\widetilde{Y}_{i}(T;T+T_{0})|^{2}\leqslant \frac{K}{\varepsilon}e^{\varepsilon(t-T)}\sum_{i=1}^{2}\mathbb{E}\int_{T}^{T+T_{0}}e^{-\frac{\varepsilon}{2}(r-t)}|\widetilde{\varphi}_{i}(r)|^{2}dr \\ &\leqslant \frac{K}{\varepsilon}e^{\varepsilon(t-T)}\sum_{i=1}^{2}\mathbb{E}\int_{T}^{\infty}|\widetilde{\varphi}_{i}(r)|^{2}dr \to 0 \text{ as } T \to \infty. \end{split}$$

This concludes that $\widetilde{Y}_i(t;T)$ is a Cauchy sequence in $L^2_{\mathcal{F}_t}$ as $T \to \infty$ with a limit $\widetilde{Y}_i(t)$ for each finite $t \in [0,\infty)$. The above estimate also yields that $\widetilde{Y}_i(\cdot;T)$ converges to $\widetilde{Y}_i(\cdot)$ in $L^2_{\mathbb{F}}(0,T_1;\mathbb{R}^n)$ for any finite $T_1 > 0$. We also notice that $\lim_{t\to\infty} \mathbb{E}|\widetilde{Y}_i(t)|^2 = 0$ from (5.26) and $\widetilde{Y}_i(\cdot) \in L^2_{\mathbb{F}}(0,\infty;\mathbb{R}^n)$ because

$$\begin{split} & \mathbb{E}\sum_{i=1}^{2}\int_{0}^{\infty}|\widetilde{Y}_{i}(t)|^{2}dt \leqslant \frac{K}{\varepsilon}\mathbb{E}\sum_{i=1}^{2}\int_{0}^{\infty}\int_{t}^{\infty}e^{-\frac{\varepsilon}{2}(r-t)}|\widetilde{\varphi}_{i}(r)|^{2}drdt \\ & = \frac{K}{\varepsilon}\mathbb{E}\sum_{i=1}^{2}\int_{0}^{\infty}\int_{0}^{r}e^{-\frac{\varepsilon}{2}(r-t)}|\widetilde{\varphi}_{i}(r)|^{2}dtdr \leqslant K\mathbb{E}\sum_{i=1}^{2}\int_{0}^{\infty}|\widetilde{\varphi}_{i}(r)|^{2}dr < \infty \end{split}$$

where we used (5.26) in the first inequality. From (5.27), we see that $\widetilde{Z}(\cdot;T)$, $\widetilde{Z}_1^M(\cdot;T)$ and $\widetilde{Z}_2^M(\cdot;T)$ is a Cauchy sequence in $L^2_{\mathbb{F}}(0,T_1;\mathbb{R}^n)$, $M^2_{\mathbb{F}^\alpha}(0,T_1;\mathbb{R}^n)^{\perp}$ and $M^2_{\mathbb{F}^\alpha}(0,T_1;\mathbb{R}^n)$ respectively, for any $T_1 > 0$. Taking $T \to \infty$ in (5.24), we have

$$\begin{split} 0 &= \widetilde{Y}_{1}(t_{2}) - \widetilde{Y}_{1}(t_{1}) + \int_{t_{1}}^{t_{2}} \widetilde{Z}(t) dW(t) + \int_{t_{1}}^{t_{2}} \widetilde{Z}_{1}^{M}(t) dM(t) \\ &+ \int_{t_{1}}^{t_{2}} \left[\widetilde{A}_{1}^{\top}(\alpha(t)) \widetilde{Y}_{1}(t) - \widetilde{C}_{1}^{\top}(\alpha(t)) \Pi_{1}[\widetilde{Z}(t)] + \widetilde{\varphi}_{1}(t) - \sum_{i \neq j} \widetilde{E}_{1}^{\top}(i, j) \widetilde{Z}_{1}^{M}(t, j) \lambda_{ij} \mathbf{1}_{\{\alpha(t)=i\}} dt, \\ 0 &= \widetilde{Y}_{1}(t_{2}) - \widetilde{Y}_{1}(t_{1}) + \int_{t_{1}}^{t_{2}} \widetilde{Z}_{2}^{M}(t) dM(t) \\ &+ \int_{t_{1}}^{t_{2}} \left[\widetilde{A}_{2}^{\top}(\alpha(t)) \widetilde{Y}_{2}(t) - \widetilde{C}_{2}^{\top}(\alpha(t)) \Pi_{2}[\widetilde{Z}(t)] + \widetilde{\varphi}_{2}(t) - \sum_{i \neq j} \widetilde{E}_{2}^{\top}(i, j) \widetilde{Z}_{2}^{M}(t, j) \lambda_{ij} \mathbf{1}_{\{\alpha(t)=i\}} \right] dt, \end{split}$$

for any $t_2 > t_1 \ge 0$. This says we have solved the BSDE (5.24) on [0, T) for any T > 0. Moreover, we see that $\widetilde{Z}(\cdot)$ and $\widetilde{Z}^M(\cdot)$ are L^2 -integrable on $[0, \infty)$ from (5.25). Finally, the uniqueness can be concluded from (5.23) directly. The proof is complete.

6 Open-Loop Solvability

This section is concerned with the open-loop solvability of Problem (MF-LQ)^{∞} based on the solvability of the BSDE (5.2). First of all, under (A4), by taking $(\Theta_1(\cdot), \Theta_2(\cdot)) \in \mathbf{S}[A, \bar{A}, C, \bar{C}; B, \bar{B}, D, \bar{D}]$, we may

formulate Problem $(MF-LQ)^{\infty,*}$ whose open-loop solvability is equivalent to that of Problem $(MF-LQ)^{\infty}$. Thus to investigate the open-loop solvability of Problem $(MF-LQ)^{\infty}$, we need only to consider the case that (A4)' holds. Hence, in what follows, we will keep this assumption, without loss of generality. We now state and prove the following theorem.

Theorem 6.1. Let **(A1)**–(**A3)** and **(A4)'** hold. Then $(\bar{X}(\cdot), \bar{u}(\cdot))$ is an open-loop optimal pair at $(s, \iota, \xi) \in \mathscr{D}$ if and only if for some adapted and square integrable $(Y(\cdot), Z(\cdot), Z^M(\cdot))$, the following FBSDE holds

$$(6.1) \begin{cases} d\bar{X}_{1}(t) = \left\{A_{1}(\alpha(t))\bar{X}_{1}(t) + B_{1}(\alpha(t))\bar{u}_{1}(t) + b_{1}(t)\right\}dt \\ + \left\{C_{1}(\alpha(t))\bar{X}_{1}(t) + C_{2}(\alpha(t))\bar{X}_{2}(t) + D_{1}(\alpha(t))\bar{u}_{1}(t) + D_{2}(\alpha(t))\bar{u}_{2}(t) + \sigma(t)\right\}dW(t), \\ d\bar{X}_{2}(t) = \left\{A_{2}(\alpha(t))\bar{X}_{2}(t) + B_{2}(\alpha(t))\bar{u}_{2}(t) + b_{2}(t)\right\}dt, \quad t \in [s, \infty), \\ \bar{X}_{1}(s) = \xi_{1}, \quad \bar{X}_{2}(s) = \xi_{2}, \quad \alpha(s) = i, \\ dY_{1}(t) = -\left[A_{1}(\alpha(t))^{\top}Y_{1}(t) + C_{1}(\alpha(t))^{\top}\Pi_{1}[Z(t)] + Q_{1}(\alpha(t))\bar{X}_{1}(t) + S_{1}(\alpha(t))^{\top}\bar{u}_{1}(t) + q_{1}(t)\right]dt \\ + Z(t)dW(t) + Z_{1}^{M}(t)dM(t) \\ dY_{2}(t) = -\left[A_{2}(\alpha(t))^{\top}Y_{2}(t) + C_{2}(\alpha(t))^{\top}\Pi_{2}[Z(t)] + Q_{2}(\alpha(t))\bar{X}_{2}(t)) + S_{2}(\alpha(t))^{\top}\bar{u}_{2}(t) + q_{2}(t)\right]dt \\ + Z_{2}^{M}(t)dM(t), \\ \lim_{t \to \infty} \mathbb{E}\left[|Y_{1}(t)|^{2} + |Y_{2}(t)|^{2}\right] = 0, \end{cases}$$

and the following stationarity conditions hold

(6.2)
$$B_k(\alpha(t))^{\top} Y_k(t) + D_k(\alpha(t))^{\top} Z_k(t) + S_k(\alpha(t)) \bar{X}_k(t) + R_k(\alpha(t)) \bar{u}_k(t) + r_k(t) = 0, \qquad k = 1, 2.$$

The above is a coupled FBSDEs, with the coupling being through the stationarity conditions (6.2).

Proof. From Proposition 4.3, we know that it suffices to prove the necessary part. Let $\bar{u}(\cdot) \in \mathscr{U}_{ad}[s, \infty)$ be an open-loop optimal control with the corresponding state process $\bar{X}(\cdot)$. Then by its optimality, for any $u(\cdot) \in \mathscr{U}_{ad}[s, \infty)$ with $X(\cdot)$ being the corresponding state process, one has (see (2.10), suppressing $\alpha(\cdot)$)

$$0 \leq \lim_{\varepsilon \to 0} \frac{J^{\infty}(s, \imath, \xi; \bar{u}(\cdot) + \varepsilon u(\cdot)) - J^{\infty}(s, \imath, \xi; \bar{u}(\cdot))}{\varepsilon}$$
$$= \sum_{k=1}^{2} \mathbb{E} \int_{s}^{\infty} \left(\langle Q_{k} \bar{X}_{k}(t) + S_{k}^{\top} \bar{u}_{k}(t) + q_{k}(t), X_{k}(t) \rangle + \langle S_{k} \bar{X}_{k}(t) + R_{k} \bar{u}_{k}(t) + r_{k}(t), u_{k}(t) \rangle \right) dt$$

with

(6.3)
$$\begin{cases} dX_1(t) = \left\{ A_1(\alpha(t))X_1(t) + B_1(\alpha(t))u_1(t) \right\} dt \\ + \left\{ C_1(\alpha(t))X_1(t) + C_2(\alpha(t))X_2(t) + D_1(\alpha(t))u_1(t) + D_2(\alpha(t))u_2(t) \right\} dW(t) \\ dX_2(t) = \left\{ A_2(\alpha(t))X_2(t) + B_2(\alpha(t))u_2(t) \right\} dt, \quad t \in [s, \infty), \\ X_1(s) = 0, \quad X_2(s) = 0, \quad \alpha(s) = i. \end{cases}$$

Since system $[A, \overline{A}, C, \overline{C}]$ is L^2 -exponentially stable, by Theorem 5.3, we see that the BSDEs in (6.1) are solvable. Now, by Itô's formula, we have

$$\mathbb{E}\langle X_{1}(T), Y_{1}(t) \rangle = \mathbb{E} \int_{s}^{T} \left(\langle A_{1}X_{1} + B_{1}u_{1}, Y_{1} \rangle - \langle X_{1}, A_{1}^{\top}Y_{1} + C_{1}^{\top}Z_{1} + Q_{1}\bar{X}_{1} + S_{1}\bar{u}_{1} + q_{1} \rangle + \langle C_{1}X_{1} + C_{2}X_{2} + D_{1}u_{1} + D_{2}u_{2}, Z \rangle \right) dt$$
$$= \mathbb{E} \int_{s}^{T} \left(\langle u_{1}, B_{1}^{\top}Y_{1} \rangle - \langle X_{1}, Q_{1}\bar{X}_{1} + S_{1}^{\top}\bar{u}_{1} + q_{1} \rangle + \langle u_{1}, D_{1}^{\top}Z_{1} \rangle + \langle C_{2}X_{2} + D_{2}u_{2}, Z \rangle \right) dt.$$

and

$$\mathbb{E}\langle X_{2}(T), Y_{2}(T) \rangle = \mathbb{E} \int_{s}^{T} \left(\langle A_{2}X_{2} + B_{2}u_{2}, Y_{2} \rangle - \langle X_{2}, A_{2}^{\top}Y_{2} + C_{2}^{\top}Z_{2} + Q_{2}\bar{X}_{2} + S_{2}^{\top}\bar{u}_{2} + q_{2} \rangle \right) dt$$
$$= \mathbb{E} \int_{s}^{T} \left(\langle u_{2}, B_{2}^{\top}Y_{2} \rangle - \langle X_{2}, C_{2}^{\top}Z_{2} + Q_{2}\bar{X}_{2} + S_{2}^{\top}\bar{u}_{2} + q_{2} \rangle \right) dt.$$

Hence,

$$\mathbb{E}\Big[\langle X_1(T), Y_1(T)\rangle + \langle X_2(T), Y_2(T)\rangle\Big]$$

= $\mathbb{E}\int_s^T \Big(\langle u_1, B_1^\top Y_1 + D_1^\top Z_1\rangle + \langle u_2, B_2^\top Y_2 + D_2^\top Z_2\rangle$
 $-\langle X_1, Q_1 \bar{X}_1 + S_1^\top \bar{u}_1 + q_1\rangle - \langle X_2, Q_2 \bar{X}_2 + S_2^\top \bar{u}_2 + q_2\rangle\Big)dt$

Let $T \to \infty$, we get

$$\mathbb{E}\int_{s}^{\infty} \left(\langle X_{1}, Q_{1}\bar{X}_{1} + S_{1}^{\top}\bar{u}_{1} + q_{1} \rangle + \langle X_{2}, Q_{2}\bar{X}_{2} + S_{2}^{\top}\bar{u}_{2} + q_{2} \rangle \right) dt$$
$$= \mathbb{E}\int_{s}^{\infty} \left(\langle u_{1}, B_{1}^{\top}Y_{1} + D_{1}^{\top}Z_{1} \rangle + \langle u_{2}, B_{2}^{\top}Y_{2} + D_{2}^{\top}Z_{2} \rangle \right) dt.$$

Therefore,

$$0 = \sum_{k=1}^{2} \mathbb{E} \int_{s}^{\infty} \langle B_{k}^{\top} Y_{k} + D_{k}^{\top} Z_{k} + S_{k} \bar{X}_{k}(t) + R_{k} \bar{u}_{k}(t) + r_{k}(t), u_{k}(t) \rangle dt.$$

By the arbitrariness of $u_k(\cdot)$, our conclusion follows.

7 Concluding Remarks

In the paper, we have studied the linear quadratic optimal control problems in infinite-horizon for mean-field stochastic differential equations in a switching environment, called Problem $(MF-LQ)^{\infty}$. To deal with the mean-field terms involved, an orthogonal projection introduced in [21] is adopted which leads to a new linear optimal control problem on the product of two orthogonal spaces. Under some general assumptions, we find the closed-loop optimal strategy by means of AREs and a system of BSDEs. The solvability of the BSDEs also leads to the characterization of open-loop solvability of Problem $(MF-LQ)^{\infty}$ in terms of FBSDEs. The authors admit that the positive-definite conditions have been assumed, for the presentation of the current paper. We will report results without assuming these positive-definite conditions before long.

References

- B. O. D Anderson and J. B. Moore, *Linear Optimal Control*, Prentice-Hall Englewood Cliffs, New Jersey, 1971.
- [2] R. Bellman, I. Glicksberg, and O. Gross, Some Aspects of the Mathematical Theory of Control Processes, Rand Corporation, Santa Monica, California, 1958.
- [3] A. Bensoussan, Lecture on Stochastic Control, Lecture Notes in Math. Vol. 972, Springer-Verlag, Berlin, 1981.
- [4] P. Bernhard, Linear-quadratic, two-person, zero-sum differential games: necessary and sufficient conditions, J. Optim. Theory Appl., 27 (1979), 51–69.
- [5] S. Chen, X. Li and X. Y. Zhou, Stochastic linear quadratic regulators with indefinite weight costs, SIAM J. Control Optim., 36 (1998), 1695–1702.

- [6] S. Chen and J. Yong, Stochastic linear quadratic optimal control problems with random coefficients, Chin. Ann. Math., 21 B (2000), 323–338.
- [7] S. Chen and J. Yong, Stochastic linear quadratic optimal control problems, Appl. Math. Optim., 43 (2001), 21–45.
- [8] M. H. A. Davis, Linear Estimation and Stochastic Control, Chapman & Hall, London, 1977.
- M. C. Delfour, Linear quadratic differential games: saddle point and Riccati differential equations, SIAM J. Control Optim., 46 (2007), 750–774.
- [10] M. C. Delfour and O. D. Sbarba, Linear quadratic differential games: closed loop saddle points, SIAM J. Control Optim., 47 (2009), 3138–3166.
- [11] C. Donnelly and A. J. Heunis, Quadratic risk minimization in regime0switching model wit portfolio constraints, SIAM J. Control Optim., 50 (2012), 2431–2461.
- [12] J. Huang, X. Li, and J. Yong, A linear-quadratic optimal control problem for mean-field stochastic differential equations in infinite horizon, Math. Control Relat. Fields, 5 (2015), 97–139.
- [13] R. E. Kalman, Contributions to the theory of optimal control, Bol. Soc. Mat. Mexicana, 5 (1960), 102–119.
- [14] H. J. Kushner, Optimal stochastic control, IRE Trans. Auto. Control, 7 (1962), 120–122.
- [15] E. B. Lee and L. Markus, Foundations of Optimal Control Theory, John Wiley, New York, 1967.
- [16] A. M. Letov, Analytic design of regulators, Avtomat. i Telemekh., 21 (1960), 436–441; 561–568; 661–665; 22 (1961), 425–435; 23 (1962), 1405–1413 (in Russian).
- [17] X. Li, J. Shi, and J. Yong, Mean-field linear-quadratic stochastic differential games in an infinite horizon, ESAIM Control Optim. Calc. Var., 27 (2021), 81.
- [18] X. Li, J. Sun, and J. Yong, Mean-field stochastic linear quadratic optimal control problems: closed-loop solvability, Probability, Uncertainty and Quantitative Risk, 1 (2016), 2.
- [19] Q. Lü and T. Wang, Optimal feedback controls of stochastic linear quadratic control problems in infinite dimensions with random coefficients, J. Math. Pures Appl., 173 (2023), 195–242.
- [20] P. J. McLane, Optimal stochastic control of linear systems with state- and control-dependent disturbances, IEEE Trans. Auto. Control, 16 (1971), 793–798.
- [21] H. Mei, Q. Wei, and J. Yong, Linear-quadratic optimal control problem for mean-field stochastic differential equations with a type of random coefficients, Numer. Algebra, Control Optim., 14 (2024), 813–852.
- [22] H. Pham, Linear quadratic optimal control of conditional McKean-Vlasov equation with random coefficients and applications, Probab. Uncertain Quantstive Risk, 1 (2016), 7.
- [23] L. R. G. Rogers and D. Williams, Diffusions, Markov Processes and Martingales, Vol.2, 2nd ed., Cambridge Univ., Press, 2000.
- [24] E. J. Rolón Gutiérrez, S. L. Nguyen, and G. Yin, Markovian-switching systems: backward and forwardbackward stochastic differential equations, mean-field interactions, and nonzero-sum differential games, Appl. Math. Optim., 89 (2024), 33.
- [25] J. Sun, Mean-field stochastic linear quadratic optimal control problems: open-loop solvabilities, ESAIM Control Optim. Calc. Var., 23 (2017), 1099–1127.
- [26] J. Sun, X. Li, and J. Yong. Open-loop and closed-loop solvabilities for stochastic linear quadratic optimal control problems, SIAM J. Control Optim., 54 (2016), 2274–2308.

- [27] J. Sun, J. Xiong, and J. Yong, Indefinite stochastic linear-quadratic optimal control problems with random coefficients: closed-loop representation of open-loop optimal controls. The Annals of Applied Probability, 31 (2021), 460–499.
- [28] J. Sun and J. Yong, Linear quadratic stochastic differential games: Open-loop and closed-loop saddle points, SIAM J. Control Optim., 52 (2014), 4082–4121.
- [29] J. Sun and J. Yong, Stochastic linear quadratic optimal control problems in infinite horizon, Appl. Math. Optim., 78 (2018), 145–183,
- [30] J. Sun and J. Yong, Stochastic Linear-Quadratic Optimal Control Theory: Open-Loop and Closed-Loop Solutions, Springer Brief in Math., Springer, 2020.
- [31] J. Sun and J. Yong, Stochastic Linear-Quadratic Optimal Control Theory: Differential Games and Mean-Field Problems, SpringerBrief in Math., Springer, 2020.
- [32] S. Tang, General linear quadratic optimal stochastic control problems with random coefficients: Linear stochastic Hamilton systems and backward stochastic Riccati equations, SIAM J. Control Optim., 42 (2003), 53–75.
- [33] S. Tang, Dynamic programming for general linear quadratic optimal stochastic control with random coefficients, SIAM J. Control Optim., 53 (2015), 1082–1106.
- [34] Q. Wei, J. Yong, and Z. Yu, Linear quadratic optimal control problems with operator coefficients: openloop solutions, ESAIM COCV, 25 (2019). 17.
- [35] J. Wen, X. Li, J. Xiong, X. Zhang, Stochastic linear quadratic optimal control problems with random coefficients and Markovian regime switching system, arXiv:2202.04880.
- [36] J. C. Willems, Least squares stationary optimal control and the algebraic Riccati equation, IEEE Trans. Automat. Control, 16 (1971), 621–634.
- [37] W. M. Wonham, On a matrix Riccati equation of stochastic control, SIAM J. Control, 6 (1968), 681–697.
- [38] W. M. Wonham, Linear Multivariable Control: A Geometric Approach, 2nd Edition, Springer-Verlag, 1979.
- [39] J. Yong, Linear-quadratic optimal control problems for mean-field stochastic differential equations, SIAM J. Control Optim., 51(2013), 2809–2838.
- [40] P. Zhang, Some results on two-person zero-sum linear quadratic differential games, SIAM J. Control Optim., 43 (2005), 2157–2165.
- [41] X. Zhang, X. Li, and J. Xiong, Open-loop and closed-loop solvabilities for stochastic linear quadratic optimal control problems of Markovian regime-switching system, ESAIM: COCV, 27 (2021), 69.