

Partial versus total resetting for Lévy flights in d dimensions: similarities and discrepancies

Costantino Di Bello,¹ Aleksei Chechkin,^{1,2,3} Tomasz Grzywny,² Karol Szczypkowski,² Bartosz Trojan,² and Zbigniew Palmowski²

¹*University of Potsdam, Institute of Physics & Astronomy, 14476 Potsdam, Germany*

²*Faculty of Pure and Applied Mathematics, Wrocław University of Science and Technology, Wyb. Wyspiańskiego 27, 50-370 Wrocław, Poland*

³*German-Ukrainian Core of Excellence Max Planck Institute of Microstructure Physics, Weinberg 2, 06120 Halle (Saale), Germany*

(*Electronic mail: dibello@uni-potsdam.de)

(Dated: 3 January 2025)

While stochastic resetting (or total resetting) is less young and more established concept in stochastic processes, partial stochastic resetting (PSR) is a relatively new field. PSR means that, at random moments in time, a stochastic process gets multiplied by a factor between 0 and 1, thus approaching but not reaching the resetting position. In this paper, we present new results on PSR highlighting the main similarities and discrepancies with total resetting. Specifically, we consider both symmetric α -stable Lévy processes (Lévy flights) and Brownian motion with PSR in arbitrary d dimensions. We derive explicit expressions for the propagator and its stationary measure, and discuss in detail their asymptotic behavior. Interestingly, while approaching to stationarity, a dynamical phase transition occurs for the Brownian motion, but not for Lévy flights. We also analyze the behavior of the process around the resetting position and find significant differences between PSR and total resetting.

PACS numbers: 05.40.-a, 02.50.-r, 05.10.Gg

Consider an insurance company. At a random moment, a customer incurs an accident and gets a refund from the insurance. From the company's point of view, refunding someone corresponds to an instantaneous sudden loss of money. If we consider that the company's total wealth is aleatory, we could mathematically model it as a stochastic process with random jumps. In physical literature, we refer to these kinds of models as *stochastic processes with resets*. There has been a great interest in resetting processes in the last fifteen years. In this paper, we will consider a peculiar model of stochastic resetting named *partial resetting*. With partial resetting, we mean that the instantaneous jump in the stochastic process is proportional to the current value of the process. In the most studied total resetting mechanism, instead, the stochastic process gets instantaneously set to a specific value independently of any past value of the process. In this paper, we want to study how this kind of mechanism affects the behavior of generic Lévy flights in arbitrary d dimensions. We obtained, both analytically and numerically, that total and partial resetting, although being equivalent under many aspects, show some completely different behaviors.

I. INTRODUCTION

In the search for a hidden target, it is sometimes beneficial to restart from the beginning¹. This simple fact has inspired many researchers in the last decade. Although this idea can be found in some earlier works^{1,2}, we can with no doubt trace the birth of *stochastic resetting* with the seminal paper by Evans and Majumdar³. This simple yet powerful idea of restarting has been the spark that started a long series of works, not only in the field of stochastic processes. Many applications were found in computer science^{4,5}, physics⁶, finance⁷, and biology⁸. It was found in^{3,9,10} that a random searcher moving with resetting takes a significantly lower time to reach its target. After less than 15 years it is already difficult to mention all the relevant papers in the field, and some topical reviews¹¹⁻¹⁴ are already available. Recently, a new variant of the stochastic resetting model, sometimes called *partial stochastic resetting*¹⁵⁻²⁰, has been introduced in the physics literature. Considering a generic stochastic process X_t , partial stochastic resetting (PSR) consists of multiplying, at random times T_n , the current position of X_{T_n} for a certain factor $c \in [0, 1]$. This kind of model is not completely new: the *additive-increase multiplicative-decrease* (AIMD) model is analogous. AIMD was used to describe the transmission of data via internet using the transmission control protocol (TCP)²¹⁻²⁶. Nevertheless, the limit of AIMD is that it is a piecewise deterministic model, in the sense that the underlying process X_t , i.e. the process between consecutive partial resets, is deterministic. New works on PSR¹⁶⁻²⁰, instead, take into

account the stochasticity of X_t . The peculiarity of PSR of having sudden instantaneous crashes makes it a suitable model for actuarial sciences²⁷.

The scope of our paper is twofold. After a brief review of the known facts about PSR, we will first discuss new analytical results on multidimensional isotropic Lévy flights with PSR. Second, we will present all the features of PSR making a comparison with total resetting. As analytical derivations are heavily technical and accessible only to an extremely specialized audience, we preferred reporting them in a companion paper²⁸. Here we focus on the results and not on the technicalities of the derivation.

II. THE MODEL

Let us consider a d -dimensional process $X_t = (X_t^1, X_t^2, \dots, X_t^d)$, in which X_t^i are independent and identically distributed Lévy flights (LF) with stability index $\alpha \in (0, 2]$, that is α -stable Lévy processes. The case $\alpha = 2$ refers to the Brownian motion. We denote with X_t and Y_t the two processes respectively without and with partial resetting. At random times $(T_i)_{i \in \mathbb{N}}$ the stochastic process Y_t gets multiplied by a factor $c \in [0, 1]$, and we assume that inter-arrival times $T_i - T_{i-1}$ are independent and exponentially distributed with the same rate (inverse mean) $r \forall i \in \mathbb{N}$. In other words, for each infinitesimal time interval dt there is a probability $r dt$ that a partial resetting event occurs, moreover, between two inter-arrival times the two processes X_t Y_t coincide. Therefore, we can summarize the evolution of Y_t with the following equation

$$Y_t = \begin{cases} X_t, & \text{if } t < T_1, \\ cY_{T_n^-} + X_t - X_{T_n}, & \text{for } t \in [T_n, T_{n+1}), n \in \mathbb{N}. \end{cases} \quad (1)$$

A graphical representation of the evolution of the Y_t for total and partial resetting is available in figure 1 for two different one-dimensional Lévy flights.




FIG. 1. Comparison between total resetting (yellow line) and partial resetting (black line). Red lines highlight resetting events. In panel (a) we set $\alpha = 2$ (Brownian motion), while in panel (b) we set $\alpha = 1$ (Cauchy).

The propagator of the process X_t with initial condition $X_{t_0} = x_0$ is denoted by $p_0(x, t|x_0, t_0)$, where $x, x_0 \in \mathbb{R}^d$. Due to homogeneity in space and time of X_t we have $p_0(x, t|x_0, t_0) = p_0(x - x_0, t - t_0|0, 0) \equiv p_0(x - x_0, t - t_0)$.

As it was shown in¹⁸, that Y_t is also homogeneous in time, therefore we can set $t_0 = 0$ everywhere without losing generality. We recall that the propagator of a symmetric d -dimensional Lévy flight, in terms of its Fourier transform, reads

$$\hat{p}_0(k, t) \equiv \int_{\mathbb{R}^d} dx e^{ik \cdot x} p_0(x, t) = \exp[ix_0 \kappa - D_\alpha |k|^\alpha t] \quad (2)$$

where $\alpha \in]0, 2]$ is called stability index, while D_α is a scale parameter sometimes called *noise intensity*²⁹, clearly $k \in \mathbb{R}^d$. We are interested in the pdf $p_r(x, t|x_0)$ of the process Y_t . An analytical, yet cumbersome, expression for p_r in one dimension is already available in¹⁸. Nevertheless, the authors of¹⁸, apart from stating the analytical expression of p_r , did not discuss any feature of this propagator. Here, we would like first to generalize the expression of p_r to arbitrary d -dimensions, and second to analyze some interesting features of this pdf like moments, tails, approach to stationarity, and non-equilibrium steady state. A complete proof of the results that we present here is available in the companion paper²⁸. We prefer to split the work into two papers for multiple reasons. First, the complete analytical proof of our results is extremely long and requires advanced mathematical techniques which might be of interest to an extremely specialized mathematical audience, only. Second, we would like to present the main features of partial resetting, highlighting the similarities and discrepancies from total resetting. The paper is organized as follows. In section III we recapitulate the mathematical description of the model and all the known results already available in the literature. In section IV we present an alternative expression of the propagator obtained with the technique of splines. In section V we discuss in full detail the stationary distribution, its moments and its scaling for large $|x|$ in full details. In section VI we discuss the relaxation of the propagator to a stationary state and the emergence of a dynamical phase transition for Brownian motion. In section VII we analyze the behavior of the propagator around $|x| = 0$. In all the sections the features of partial resetting are compared to the ones of total resetting. A complete summary of all the features discussed, compared to the ones of total resetting, is available in Table I.

III. PRELIMINARIES

A. Useful notations

Let us first give a few notations that we are going to use in the entire manuscript. We denote with

$$m \equiv c^\alpha. \quad (3)$$

We will make extensive use of the q -Pochhammer symbols which are defined as

$$(a; q)_0 \equiv 1, \quad (a; q)_n \equiv \prod_{j=0}^{n-1} (1 - aq^j), \quad (4)$$

$$(a; q)_\infty \equiv \prod_{j=0}^{\infty} (1 - aq^j),$$

and q -Gamma function

$$\Gamma_q(x) \equiv (1 - q)^{1-x} \frac{(q; q)_\infty}{(q^x; q)_\infty}, \quad x \notin -\mathbb{N}. \quad (5)$$

B. Known results

Here, we recall some basics and known results on the PSR. Since in the whole manuscript we fix $x_0 = 0$, we will drop the dependence on x_0 in the propagator and we will write simply $p_r(x, t)$. The pantograph³⁰ Fokker-Planck equation (FPE) for this process in one dimension has been found in¹⁶. The generalization to many dimensions is straightforward and reads

$$\partial_t p_r(x, t) = D_\alpha \Delta^{\alpha/2} p_r(x, t) + \frac{r}{c} p_r\left(\frac{x}{c}, t\right) - r p_r(x, t). \quad (6)$$

Here $\Delta^{\alpha/2}$ is the fractional Laplacian, defined in terms of its Fourier transform

$$\mathcal{F}\left[(-\Delta)^{\alpha/2} f(x)\right](k) = |k|^\alpha \hat{f}(k), \quad (7)$$

$$p_r(x, t) = e^{-rt} \sum_{n=0}^{\infty} r^n \sum_{k=0}^n \frac{1}{(m^{-1}; m^{-1})_k (m; m)_{n-k}} \int_0^t \left((1 - \delta_{n0}) \frac{(t-t')^{n-1}}{(n-1)!} + \delta_{n0} \delta(t-t') \right) p_0(x, m^k t') dt', \quad (14)$$

where δ is the Dirac delta function (we used the same symbol for both discrete and continuous cases). The propagator for total resetting has, instead, a much simpler expression³¹

$$p_r(x, t) = e^{-rt} p_0(x, t) + r \int_0^t e^{-rt'} p_0(x, t') dt'. \quad (15)$$

where $|k|$ is the euclidean norm of the vector $k \in \mathbb{R}^d$, while $\hat{f}(k)$ denotes the Fourier transform of f , defined as

$$\hat{f}(k) = \int_{\mathbb{R}^d} dx e^{ik \cdot x} f(x). \quad (8)$$

In the case of total resetting ($c \rightarrow 0^+$), the FPE reads (see³¹)

$$\partial_t p_r(x, t) = D_\alpha \Delta^{\alpha/2} p_r(x, t) + r \delta(x) - r p_r(x, t). \quad (9)$$

It can be proved, using eq. (6), that for PSR the double Fourier-Laplace transform of the propagator, defined as

$$\hat{p}_r(k, s) \equiv \int_0^\infty e^{-st} \int_{\mathbb{R}^d} e^{ik \cdot x} p_r(x, t) dx dt, \quad s > 0, k \in \mathbb{R}^d, \quad (10)$$

reads

$$\hat{p}_r(k, s) = \sum_{n=0}^{\infty} r^n \prod_{j=0}^n \frac{1}{r + s + D_\alpha m^j |k|^\alpha}. \quad (11)$$

Equation (11) is completely analogous to eq. (16) in¹⁸ for $d = 1$. The only term depending on the dimension d is the norm of the vector $|k|$. As expected the propagator is isotropic, as it only depends on $|k|$.

The stationary distribution $p^{(s)}(x)$ can be computed taking the limit $t \rightarrow \infty$ in eq. (6), thus it solves

$$D_\alpha \Delta^{\alpha/2} p_r^{(s)}(x) + \frac{r}{c} p_r^{(s)}\left(\frac{x}{c}, t\right) - r p_r^{(s)}(x) = 0, \quad (12)$$

and its Fourier transform reads

$$\hat{p}_r(k) = \prod_{j=0}^{\infty} \frac{r}{r + D_\alpha m^j |k|^\alpha}. \quad (13)$$

Equation (13) is also a known result from¹⁸ (see eq. (27)) for the case $d = 1$.

An analytical inversion of the double transform in eq. (11) has been performed in¹⁸ (see eq. (23)). The expression for d dimensions reads

Formula (14) is rather cumbersome and difficult to handle. It is possible, in principle, to express the integral appearing in (14) in terms of hypergeometric functions (see eq. (36) in¹⁸). Nevertheless, for an efficient numerical implementation, it is useful to use a different series representation, which we will discuss in the next section. With such an efficient implementation, it will be possible to analyze the tails of the distribution with high accuracy

and to verify the existence of a dynamical phase transition.

We conclude this section by mentioning the stationary distribution, also found in¹⁸ for one-dimensional processes (see eq. (32)). It can be written in terms of Fox functions³² and reads

$$p_r^{(s)}(x) = \frac{1}{(c^\alpha; c^\alpha)_\infty} \sum_{n=0}^{\infty} \frac{1}{(c^{-\alpha}; c^{-\alpha})_n} \frac{1}{\alpha|x|} \times \\ \times H_{2,3}^{2,1} \left[\left(\frac{r}{D} \right)^{1/\alpha} \frac{|x|}{c^n} \left| \begin{matrix} (1, 1/\alpha), (1, 1/2) \\ (1, 1), (1, 1/\alpha), (1, 1/2) \end{matrix} \right. \right]. \quad (16)$$

In¹⁸ authors computed neither the moments nor the asymptotics. We will also expose our results for these two important features.

We do not report here the explicit formulae for total resetting, but they can be easily derived by taking the limit $c \rightarrow 0^+$ or $m \rightarrow 0^+$ (see references^{3,31,33}).

IV. ALTERNATIVE REPRESENTATION OF THE PROPAGATOR

It is well known that the propagator $p_r(x, t)$ also solves the last renewal equation¹⁸

$$p_r(y, t|x) = e^{-rt} p_0(y, t|x) + \\ + r \int_0^t \int_{\mathbb{R}^d} e^{-rs} p_0(z, s|x) p_r(y, t-s|cz) dz ds. \quad (17)$$

The meaning of the rhs of (17) is rather simple: either, with probability e^{-rt} , the random walk does not incur in any partial reset up to time t , and thus freely propagates from x to y with $p_0(y, t|x)$, or, with probability re^{-rs} , a partial resetting event occurs at some time $0 < s < t$. This second possibility means that in the time interval $[0, s[$ the random walk propagates unperturbed from x to some $z \in \mathbb{R}^d$ with propagator $p_0(z, s|x)$, then in the time interval $[s, t]$ the partial stochastic resetting process propagates from cz to y with $p_r(y, t-s|cz)$. Of course this term has to be integrated over all possible values of s and z . The formula for total resetting (15) can be retrieved by sending $c \rightarrow 0^+$ (see³).

As shown in the companion paper²⁸ (see section 3), a possible way to find the solution to the renewal equation is to define recursively the sequence $(P_n : n \in \mathbb{N})$ of splines via

$$P_1(u) = \frac{1}{1-m} \mathbb{1}_{[m,1]}(u), \quad (18)$$

$$P_{n+1}(u) = \max(u - m^{n+1}; 0)^n \int_u^1 \frac{P_n(v)}{(v - m^{n+1})^{n+1}} dv, \quad (19)$$

where $\mathbb{1}_{[m,1]}(u)$ is the indicator function of the interval $[m, 1]$, i.e. it is 1 if $u \in [m, 1]$ and 0 otherwise. Note that

each P_n is supported on $[m^n, 1]$. In²⁸ it was proved that the solutions (17) reads

$$p_r(x, t) = \int_0^\infty p_0(x, u) \mu_t(du), \quad x \in \mathbb{R}^d, \quad (20)$$

where μ_t is a measure defined as

$$\mu_t(du) = e^{-rt} \delta(t) du + e^{-rt} \sum_{j=1}^{\infty} r^j t^j P_j(u/t) \frac{du}{t}. \quad (21)$$

Notice that this solution is valid only for $c > 0$. In the case $c = 0$ (total resetting) the integral in eq.(20) does not converge. As stated before, we do not report here the proof, but we mention that it makes use of the scaling properties of Lévy flights³⁴. This alternative representation of the propagator has several advantages. The first is that it is possible to analyze the tails of the distribution, and thus to identify, in the case of Brownian motion with PSR, a dynamical phase transition. Second, in this way it is possible to avoid using special functions that are generally not numerically stable. A comparison between a simple Monte Carlo simulation and the aforementioned spline solution is available in Fig. 2.

In the next section we proceed with a discussion on the stationary distribution and its scaling for large $|x|$.




FIG. 2. Comparison in log scale between Monte Carlo sampling of positions (dots) with spline solution given in eqs. (20),(21) for two different values of α . In panel (a) we have $\alpha = 2$, in (b) $\alpha = 1$. Both simulations were made with parameters $d = 1$, $D = 1/2$, $t = 5$, $r = 1/2$, $c = 0.5$.

V. STATIONARY DISTRIBUTION

As to the stationary distribution, one can prove, using the same spline method (see²⁸ for details), that the stationary PDF $p_r^{(s)}$ for a d-dimensional process reads

$$p_r^{(s)}(x) = \frac{1}{(m; m)_\infty} \sum_{k=0}^{\infty} (-1)^k \frac{m^{\frac{1}{2}k(k-1)}}{(m; m)_k} \times \\ \times \int_0^\infty e^{-m^{-k} u} r^{d/\alpha} p_0(r^{1/\alpha} x, u) du, \quad (22)$$

where $x \in \mathbb{R}^d$. In the case of Brownian motion ($\alpha = 2$) the integral can be computed explicitly,

$$p_r^{(s)}(x) = \frac{1}{(2\pi)^{d/2}} \frac{(r/D)^{d/2}}{(m; m)_\infty} \sum_{k=0}^{\infty} (-1)^k \frac{m^{\frac{1}{2}k(k-1)}}{(m; m)_k} \times \\ \times \left(m^{\frac{k}{2}} \sqrt{\frac{r}{D}} |x| \right)^{1-d/2} K_{d/2-1} \left(\frac{\sqrt{r/D} |x|}{m^{k/2}} \right), \quad (23)$$

where $K_{d/2-1}$ is the modified Bessel function of the second type, and we simply set $D_2 = D$. For $d = 1$ in the case of Brownian motion we have

$$p_r^{(s)}(x) = \frac{1}{2} \frac{\sqrt{r/D}}{(m; m)_\infty} \sum_{k=0}^{\infty} (-1)^k \frac{m^{\frac{k^2}{2}}}{(m; m)_k} e^{-m^{-\frac{k}{2}} \sqrt{r/D} |x|} \quad (24)$$

which is in agreement with equation (37) of¹⁸, see also¹⁶.

A. Asymptotic tail of the propagator and the stationary measure

Recall that the Lévy measure of an isotropic d -dimensional Lévy flight with index $\alpha \in (0, 2)$ equals (see³⁴)

$$\nu(x) = \frac{2^\alpha \Gamma((d+\alpha)/2)}{\pi^{d/2} |\Gamma(-\alpha/2)|} |x|^{-d-\alpha}. \quad (25)$$

In²⁸ it was proved that in the case of the Lévy flight with index $0 < \alpha < 2$ we have

$$\lim_{\substack{|x| \rightarrow +\infty \\ t \rightarrow +\infty}} \frac{p_r(x, t)}{\nu(x)} = \frac{1}{r(1-m)} \quad (26)$$

and therefore

$$p_r^{(s)}(x) \sim \frac{1}{r(1-m)} \nu(x) \quad \text{as } |x| \rightarrow +\infty. \quad (27)$$

where we write $f(x) \sim g(x)$ if $\lim_{|x| \rightarrow +\infty} f(x)/g(x) = 1$.

The asymptotics (26) is a consequence of one big jump principle: the large values of $|x|$ are possible only when we have one big jump with regularly varying density (25) that should happens till time t .

On the other side, if X_t is a Brownian motion in \mathbb{R}^d , then from (23) it follows that

$$p_r^{(s)}(x) \sim \frac{1}{2} \frac{(r/D)^{(d+1)/4}}{(m; m)_\infty} (2\pi)^{-\frac{d-1}{2}} |x|^{-\frac{d-1}{2}} e^{-\sqrt{r/D} |x|} \\ \text{as } |x| \rightarrow +\infty. \quad (28)$$

We can observe that from eqs. (27)(28) it is possible to obtain the scaling for total resetting by sending $m \rightarrow 0$. In both cases, partial and total resetting exhibit the same scaling for large $|x|$.

B. Moments of stationary distribution

With a simple calculation, as usual available in²⁸, it can be proved that the fractional moments of the stationary distribution for all $-1 < \gamma < \alpha$

$$\int_{\mathbb{R}^d} |x|^\gamma p_r^{(s)}(x) dx = \langle |Y_\infty|^\gamma \rangle = \\ = \frac{\Gamma(\gamma/\alpha + 1)}{\Gamma_m(\gamma/\alpha + 1)} (1-m)^{-\gamma/\alpha} r^{-\gamma/\alpha} \langle |X_1|^\gamma \rangle. \quad (29)$$

Equation (29) is one of the new results of this paper. Additionally, if $d = 1$ and X_t is a symmetric α -stable process with $\alpha \in (0, 2)$, then for $-1 < \gamma < \alpha$ ³⁵,

$$\langle |X_1|^\gamma \rangle = \frac{2^\gamma \Gamma(\frac{1+\gamma}{2}) \Gamma(1 - \frac{\gamma}{\alpha})}{\sqrt{\pi} \Gamma(1 - \frac{\gamma}{2})}. \quad (30)$$

Similarly, if $d = 1$ and X_t is a Brownian motion B_t , then for $\gamma > -1$ one can see that

$$\langle |B_1|^\gamma \rangle = \frac{2^\gamma \Gamma(\frac{1+\gamma}{2})}{\sqrt{\pi}}. \quad (31)$$

In particular, for $k \in \mathbb{N}$ and one-dimensional Brownian motion,

$$\langle Y_\infty^{2k-1} \rangle = \int_{\mathbb{R}} x^{2k-1} p_r^{(s)}(x) dx = 0, \quad (32)$$

and

$$\langle Y_\infty^{2k} \rangle = \int_{\mathbb{R}} x^{2k} p_r^{(s)}(x) dx = r^{-k} \frac{(2k)!}{(m; m)_k}. \quad (33)$$

Hence in this case the variance of the stationary distribution Y_∞ equals

$$\langle Y_\infty^2 \rangle = \frac{2}{r(1-m)}. \quad (34)$$

The same result for total resetting could be obtained by taking the limit $m \rightarrow 0^+$. The results are summarized in Table I.

VI. NON-EQUILIBRIUM STATIONARY STATE

Partial resetting, similarly to total resetting, is characterized by the existence of a non-equilibrium steady state (NESS). Indeed, when a resetting event occurs the random walker is instantaneously transferred from some position x to cx . The way in which resetting is defined is such that the reversed jump, i.e. from cx to x , is not possible. This peculiarity is responsible for the presence of the so called *dynamical phase transition*³⁶. The concept is very well explained in³⁷ for the case of Brownian motion in $d = 1$ with total resetting. The idea is that

when t tends to $+\infty$ the propagator of the process can be written in the scaling form

$$p_r(x, t) \sim e^{-tI(x/t)} \quad (35)$$

where the function I is generally called *large deviation function* (LDF). If the LDF shows a discontinuity in one

$$p_r(x, t) = \begin{cases} p_r^{(s)}(x) (1 + \mathcal{O}(t^{-1})) & \text{for } y < y^* \\ e^{-rt} (4\pi Dt)^{-\frac{d}{2}} e^{-\frac{|x|^2}{4Dt}} \left[1 + \frac{4Dt^2}{|x|^2} \vartheta\left(\frac{4Dt^2}{|x|^2}\right) + \mathcal{O}\left(\frac{4Dt^2}{|x|^2}\right) \right] & \text{for } y > y^*, \end{cases} \quad (36)$$

where $y = \frac{|x|}{t}$ is constant, $y^* = 2\sqrt{Dr}$, $x \in \mathbb{R}^d$ and the function $\vartheta(z)$ is defined as

$$\vartheta(z) = \sum_{j=0}^{\infty} \frac{1}{(m; m)_{j+1}} z^j, \quad |z| < 1. \quad (37)$$

This suggests that the LDF in eq. (35) can be written as

$$I(y) = \begin{cases} \sqrt{\frac{r}{D}} |y| & \text{for } |y| < y^* \\ r + \frac{y^2}{4D} & \text{for } |y| > y^*, \end{cases} \quad (38)$$

which is precisely the same LDF obtained in³⁷ for the total resetting. Interestingly, the dynamical phase transition is not affected either by the dimension or by the value $c \in [0, 1[$. From this point of view, partial and total resetting are identical. A graphical representation of this result is presented in Figs. 3 and 4. The result is proved in²⁸ by making use of (20) and of the Stein method³⁸. The physical meaning of this transition is the following: first of all, the propagator $p_r(x, t)$ can be interpreted as $\langle \delta(X_t - x) \rangle$, i.e. we could think of starting N independent random walks and computing the histogram of positions at time t . Second, since in the outer region $|x| > \sqrt{4Dr}t$ the far asymptotics of the propagator is Gaussian, mainly trajectories that never experienced resetting will be the ones contributing to the density $p_r(x, t)$ of this region. Therefore, following³⁷, we can argue that in the thermodynamical limit $N \rightarrow \infty$ the phase transition corresponds to a separation between the trajectories that reset many times with respect to the ones that never reset.

The dynamical phase transition is, instead, absent in the case of $\alpha \neq 2$. This is evident in Figure 5. An intuitive explanation for this behavior is the following. In Ref.³⁹ it was shown that the propagator of a LF for any $t > 0$ and large x scales as

$$p_0(x, t) \sim t|x|^{-d-\alpha} \quad (39)$$

Therefore, substituting the previous equation into (20) we have that

$$p_r(x, t) \sim A(t)|x|^{-d-\alpha} \quad (40)$$

of its derivatives we are in the presence of a dynamical phase transition (we will come back later on the physical meaning of this kind of transition). In the literature on stochastic processes, the LDF is generally obtained either from the theory of large deviations or from saddle-point methods³⁷. In our companion paper²⁸ we were able to prove that for $\alpha = 2$ we have




FIG. 3. Brownian motion with PSR in $d = 1$. In panel (a) $p_r(x, t)$ is shown for two different values of time. The solid green curve represents the stationary distribution $p^{(s)}$, see (22). The two vertical dashed lines separate the inner region, in which stationarity has been achieved, from the transient region. In particular, the two red ones refer to $t = 10$, while the blue ones refer to $t = 15$. In panel (b) we again show the propagator $p_r(x, t)$, but with respect to the rescaled coordinate x/t . At variance with panel (a), in which we plotted the stationary distribution, here we represented the scaling $e^{-tI(x/t)}$. Close to $x = 0$, the higher order exponentials in eq. (24) are non-negligible, and this explains the poor agreement for small $|x|$. Both figures were obtained with $D = 0.5$, $c = 0.5$, $r = 0.5$.

where $A(t)$ is some function of time such that $\lim_{t \rightarrow \infty} A(t) = \text{const}$. Moreover, the stationary distribution exhibits the same decay (compare eq. (27)). Therefore, in the process of relaxation, the asymptotics of the propagator does not change, and a dynamical phase transition is absent for any d . The same argument applies in the case of LF with total resetting. Previous works, indeed, did not observe such a dynamical phase transition in $d = 1$ ^{33,40}.

VII. BOUNDEDNESS OF THE PROPAGATOR

As we have seen so far, there are no major differences between partial and total resetting. Scalings, tails, dynamical phase transitions are precisely the same for the two processes. Yet there is a feature, that we discuss in this section, that distinguishes between the two




FIG. 4. Large deviation function vs rescaled variable $y = x/rt$. All symbols represent the probability distribution for different values of time, the solid curve represents the large deviation function (38). The dashed line highlights the phase transition.




FIG. 5. Propagator of Lévy flight with PSR in the case $\alpha = 1$ (Cauchy). The symbols represent different values of time, while the solid line represents the stationary distribution given by eq. (22). In panel (b) a zoom of the far asymptotics of the distribution is shown. The parameters of the simulations are $D = 0.5$, $r = 0.5$, $c = 0.5$.

stochastic processes. It has been observed in³¹ that the propagator of a Brownian motion with total resetting in $d \geq 2$ dimensions is unbounded around $|x| = 0$, i.e. $\lim_{|x| \rightarrow 0} p_r(x, t) = +\infty$. The reason for this fact is rather easy: the propagator of a LF has the scaling property

$$p_0(x, t) = t^{-d/\alpha} p_0(x/t^{-1/\alpha}, 1) \quad (41)$$

which, in the case of total resetting, after substituting it into eq. (15) and taking the limit $|x| \rightarrow 0$ gives

$$\lim_{|x| \rightarrow 0} p_r(x, t) = \lim_{|x| \rightarrow 0} \left[e^{-rt} p_0(x, t) + r \int_0^t e^{-ru} u^{-d/\alpha} p_0(x/u^{-1/\alpha}, 1) du \right], \quad (42)$$

where the integral at the rhs diverges for $|x| \rightarrow 0$ whenever $d \geq \alpha$. On the other hand, for PSR after substitut-

ing the same scaling (41) into eq. (20) we get

$$\begin{aligned} \lim_{|x| \rightarrow 0} p_r(x, t) &= \lim_{|x| \rightarrow 0} \left[\int_0^\infty u^{-d/\alpha} p_0(x/u^{-1/\alpha}, 1) \mu_t(u) du \right] = \\ &= p_0(0, 1) \int_0^\infty u^{-d/\alpha} \mu_t(u) du \quad (43) \end{aligned}$$

where the integral in the last step does not diverge since all splines are identically equal to 0 close to $u = 0$ (see eqs. (19)). In Fig. 6 we plotted the propagator for several values of d and α .




FIG. 6. Figure showing boundedness of the propagator. The parameters of the simulations are $D = 0.5$, $r = 0.5$, $c = 0.5$, $t = 5$, $\alpha = 2$ (panel (a)), $\alpha = 1$ (panel (a)).

VIII. CONCLUSIONS

We have studied the behavior of partial stochastic resetting for symmetric Lévy flights in arbitrary space dimension and compare it with total resetting. Interestingly, the two processes both show the existence of a stationary distribution with the same asymptotics, the same approach to stationarity and the same phase transition in the case of Brownian motion. The only feature distinguishing the two processes is the behavior around $|x| = 0$. We have summarized all these features in Table I.

We hope our paper will open many perspectives around PSR. Future directions of research could be the study of the first-passage time distribution for Lévy flights⁴¹ with PSR, or the extension of our theory to asymmetric Lévy flights⁴² and to non-stable Lévy processes³⁴. Another interesting direction of the research could be studying the universal singularities of the models with resetting^{43,44}. We believe that our work can serve as a model for practical applications, e.g. in finance or in actuarial sciences.

TABLE I.

Comparison of α -stable processes, $\alpha \in]0, 2]$, $c \in [0, 1[$		
Process	α -stable with PSR ($c \neq 0$)	α -stable with total resetting ($c = 0$)
FPE: $\partial_t p_r(x, t) =$	$D_\alpha \Delta^{\alpha/2} p_r(x, t) + \frac{r}{c} p_r\left(\frac{r}{c}, t\right) - r p_r(x, t)$	$D_\alpha \Delta^{\alpha/2} p_r(x, t) + r \delta(x) - r p_r(x, t)$
$\hat{p}_r(k, s) =$	$\sum_{n=0}^{\infty} r^n \prod_{j=0}^n \frac{1}{r+s+ k ^{\alpha} c^{\alpha j}}$	$\frac{r+s}{s} \frac{1}{r+s+ k ^{\alpha}}$
$p_r(x, t) =$	See eq. (14)	$\lim_{c \rightarrow 0}$ in eq. (14)
Spline $p_r(x, t)$	See eqs. (18, 19, 20, 21)	Not available
$\alpha \neq 2$: $p_r(x, t) \sim$	$A(t) x ^{-d-\alpha}$ where $A(t) \rightarrow \text{const}$ when $t \rightarrow \infty$	$A(t) x ^{-d-\alpha}$
$\alpha = 2$: $p_r(x, t) \sim$	$e^{-tI(x/t)}$	$e^{-tI(x/t)}$
$\langle x ^\gamma p_r^{(s)}(x) \rangle =$	with $I(y) = \begin{cases} \sqrt{\frac{r}{D}} y & \text{if } y < y^* \\ r + \frac{y^2}{4D} & \text{if } y > y^* \end{cases}$ $\frac{\Gamma(\gamma/\alpha+1)}{\Gamma_m(\gamma/\alpha+1)} (1-m)^{\gamma/\alpha} r^{-\gamma/\alpha} \langle X_1^\gamma \rangle$ with $\langle X_1^\gamma \rangle = \frac{2^\gamma \Gamma\left(\frac{1+\gamma}{2}\right) \Gamma\left(1-\frac{\gamma}{\alpha}\right)}{\sqrt{\pi} \Gamma\left(1-\frac{\gamma}{2}\right)}$	$\Gamma(\gamma/\alpha + 1) r^{-\gamma/\alpha} \langle X_1^\gamma \rangle$
Boundedness	$p_r(x, t)$ bounded around $x = 0$	$p_r(x, t)$ not bounded around $x = 0$

ACKNOWLEDGMENTS

The authors thank Dr Amin Padash for useful discussions. This research is partially supported by the Polish National Science Centre under the grant No. 2023/49/B/ST1/00678. Aleksei Chechkin acknowledges support from the BMBF Project 01DK24006 PLASMA-SPIN-ENERGY.

- ¹A. Montanari and R. Zecchina, “Optimizing searches via rare events,” *Phys. Rev. Lett.* **88**, 178701 (2002).
- ²S. C. Manrubia and D. H. Zanette, “Stochastic multiplicative processes with reset events,” *Phys. Rev. E* **59**, 4945–4948 (1999).
- ³M. Evans and S. Majumdar, “Diffusion with stochastic resetting,” *Phys. Rev. Lett.* **106**, 160601 (2011).
- ⁴S. M. Maurer and B. A. Huberman, “Restart strategies and internet congestion,” *Journal of Economic Dynamics and Control* **25**, 641–654 (2001), agent-based Computational Economics (ACE).
- ⁵H. Tong, C. Faloutsos, and J. Y. Pan, “Random walk with restart: fast solutions and applications,” *Knowledge and Information Systems* **14**, 327–346 (2008).
- ⁶T. Sandev, V. Domazetoski, L. Kocarev, R. Metzler, and A. Chechkin, “Heterogeneous diffusion with stochastic resetting,” *Journal of Physics A: Mathematical and Theoretical* **55**, 074003 (2022).
- ⁷P. Jolakoski, A. Pal, T. Sandev, L. Kocarev, R. Metzler, and V. Stojkoski, “A first passage under resetting approach to income dynamics,” *Chaos, Solitons & Fractals* **175**, 113921 (2023).
- ⁸E. Roldan, A. Lisica, Sanchez-Taltavull, D., and S. Grill, “Stochastic resetting in backtrack recovery by RNA polymerases,” *Phys. Rev. E* **93**, 062411 (2016).
- ⁹M. R. Evans and S. N. Majumdar, “Diffusion with optimal resetting,” *Journal of Physics A: Mathematical and Theoretical* **44**, 435001 (2011).
- ¹⁰A. Chechkin and I. M. Sokolov, “Random search with resetting: A unified renewal approach,” *Phys. Rev. Lett.* **121**, 050601 (2018).
- ¹¹S. Gupta and A. M. Jayannavar, “Stochastic resetting: A (very) brief review,” *Frontiers in Physics* **10** (2022), 10.3389/fphy.2022.789097.
- ¹²A. Nagar and S. Gupta, “Stochastic resetting in interacting particle systems: a review,” *Journal of Physics A: Mathematical and Theoretical* **56**, 283001 (2023).
- ¹³A. Kundu and S. Reuveni, “Preface: stochastic resetting—theory and applications,” *Journal of Physics A: Mathematical and Theoretical* **57**, 060301 (2024).
- ¹⁴A. Pal, V. Stojkoski, and T. Sandev, “Random resetting in search problems,” in *Target Search Problems*, edited by D. Grebenkov, R. Metzler, and G. Oshanin (Springer Nature Switzerland, Cham, 2024) pp. 323–355.
- ¹⁵M. Dahlenburg, A. V. Chechkin, R. Schumer, and R. Metzler, “Stochastic resetting by a random amplitude,” *Phys. Rev. E* **103**, 052123 (2021).
- ¹⁶J. K. Pierce, “An advection-diffusion process with proportional resetting,” (2022), arXiv:2204.07215 [cond-mat.stat-mech].
- ¹⁷O. Tal-Friedman, Y. Roichman, and S. Reuveni, “Diffusion with partial resetting,” *Phys. Rev. E* **106**, Paper No. 054116, 13 (2022).
- ¹⁸C. Di Bello, A. Chechkin, A. Hartmann, Z. Palmowski, and R. Metzler, “Time-dependent probability density function for partial resetting dynamics,” *New J. Phys.* **25**, 082002 (2023).
- ¹⁹K. S. Olsen and D. Gupta, “Thermodynamic work of partial resetting,” *Journal of Physics A: Mathematical and Theoretical* **57**, 245001 (2024).
- ²⁰K. S. Olsen and H. Löwen, “Partial stochastic resetting with refractory periods,” *Journal of Physics A: Mathematical and Theoretical* **57**, 485001 (2024).
- ²¹V. Dumas, F. Guillemin, and P. Robert, “A Markovian analysis of additive-increase multiplicative-decrease algorithms,” *Adv. in Appl. Probab.* **34**, 85–111 (2002).
- ²²F. Guillemin, P. Robert, and B. Zwart, “AIMD algorithms and exponential functionals,” *Ann. Appl. Probab.* **14**, 90–117 (2004).
- ²³T. Ott, J. Kemperman, and M. Mathis, “The stationary behavior of ideal TCP congestion avoidance,” (1996), unpublished.
- ²⁴R. van der Hofstad, S. Kapodistria, Z. Palmowski, and S. Shneer, “Unified approach for solving exit problems for additive-increase and multiplicative-decrease processes,” *J. Appl. Probab.* **60**, 85–105 (2023).
- ²⁵A. Löpker and W. Stadje, “Hitting times and the running maximum of Markovian growth-collapse processes,” *J. Appl. Probab.* **48**, 295–312 (2011).
- ²⁶J. van Leeuwen, A. Löpker, and T. Ott, “TCP and iso-stationary transformations,” *Queueing Syst.* **63**, 459–475 (2009).
- ²⁷S. Asmussen and H. Albrecher, *Ruin Probabilities*, 2nd ed. (WORLD SCIENTIFIC, 2010) <https://www.worldscientific.com/doi/pdf/10.1142/7431>.
- ²⁸T. Grzywny, K. Szczypkowski, Z. Palmowski, and B. Trojan, “Stationary states for stable processes with partial resetting,” (2024), arXiv:2412.15626 [math.PR].
- ²⁹A. V. Chechkin, R. Metzler, J. Klafter, and V. Y. Gonchar, “Introduction to the theory of lévy flights,” in *Anomalous Transport* (John Wiley & Sons, Ltd, 2008) Chap. 5, pp. 129–162, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/9783527622979.ch5>.
- ³⁰L. Fox, D. F. Mayers, J. R. Ockendon, and A. B. Tayler, “On a Functional Differential Equation,” *IMA Journal of Applied Mathematics* **8**, 271–307 (1971), <https://academic.oup.com/imamat/article-pdf/8/3/271/1968911/8-3-271.pdf>.
- ³¹M. Evans and S. Majumdar, “Diffusion with resetting in arbitrary spatial dimension,” *J. Phys. A* **47**, 285001, 19 (2014).
- ³²J. E. J., “The h-function with applications in statistics and other disciplines,” *Technometrics* **21**, 392–393 (1979).
- ³³L. Kuśmierz and E. Gudowska-Nowak, “Optimal first-arrival times in lévy flights with resetting,” *Phys. Rev. E* **92**, 052127 (2015).
- ³⁴K. Sato, *Lévy Processes and Infinitely Divisible Distributions*, Cambridge studies in advanced mathematics (Cambridge University Press, 1999).
- ³⁵R. Metzler and J. Klafter, “The random walk’s guide to anomalous diffusion: a fractional dynamics approach,” *Physics Reports* **339**, 1–77 (2000).
- ³⁶H. Touchette, “Introduction to dynamical large deviations of markov processes,” *Physica A: Statistical Mechanics and its Applications* **504**, 5–19 (2018).
- ³⁷S. Majumdar, S. Sabhapandit, and G. Schehr, “Dynamical transition in the temporal relaxation of stochastic processes under resetting,” *Phys. Rev. E* **91**, 052131, 8 (2015).
- ³⁸E. M. Stein, *Singular Integrals and Differentiability Properties of Functions (PMS-30)* (Princeton University Press, 1970).
- ³⁹R. M. Blumenthal and R. Gettoor, “Some theorems on stable processes,” *Transactions of the American Mathematical Society* **95**, 263–273 (1960).
- ⁴⁰L. Kusmierz, S. N. Majumdar, S. Sabhapandit, and G. Schehr, “First order transition for the optimal search time of lévy flights with resetting,” *Phys. Rev. Lett.* **113**, 220602 (2014).
- ⁴¹T. Koren, M. A. Lomholt, A. V. Chechkin, J. Klafter, and R. Metzler, “Leapover lengths and first passage time statistics for lévy flights,” *Phys. Rev. Lett.* **99**, 160602 (2007).
- ⁴²A. Padash, T. Sandev, H. Kantz, R. Metzler, and A. V. Chechkin, “Asymmetric lévy flights are more efficient in random search,” *Fractal and Fractional* **6** (2022), 10.3390/fractalfract6050260.
- ⁴³A. L. Stella, A. Chechkin, and G. Teza, “Universal singularities of anomalous diffusion in the richardson class,” *Phys. Rev. E* **107**, 054118 (2023).
- ⁴⁴A. L. Stella, A. Chechkin, and G. Teza, “Anomalous dynamical scaling determines universal critical singularities,” *Phys. Rev. Lett.* **130**, 207104 (2023).