

The clock advance description of the Epidemic Type Aftershock Sequence (ETAS) model

Matthias Holschneider^{1*}

¹University of Potsdam, Institute of Mathematics

Key Points:

- ETAS model
- triggering of aftershocks
- microseismicity on all scales
- jump processes

arXiv:2501.01233v2 [physics.geo-ph] 5 Jan 2025

*DFG SFB 1294

Corresponding author: Matthias Holschneider, hols@uni-potsdam.de

Abstract

In this short paper we propose to extend the ETAS model to micro-seismic events. For that we interpret the triggered events in an ETAS model as individual local clock advances of an independent background process. The solution of the ETAS model thus becomes the sum of an infinite Markov chain of independent time adjusted background processes. This allows the incorporation of events at all scales. No artificial small magnitude cutoff is needed.

Plain language summary

A widely used model for the description of seismic activity is the ETAS model. It describes how starting from some background activity, earthquakes trigger new earthquakes giving rise to a full cascade of aftershocks. Until now, it was not possible to include microseismicity into the description due to the infinite number of such events arising in a finite time. In this paper I propose a shift of interpretation of the triggering mechanism. As I show, we may describe the triggering effect of a single earthquake as an additional independent, clock advanced background process. In this formulation, the ETAS cascade can be understood as a converging sequence of local clock advance functions and the microseismicity can be fully incorporated. A great advantage of this concept is, that the artificial minimal triggering magnitude can now be safely removed from the model and net effect of the micro-seismic activities in terms of macroscopic creep may be analyzed.

1 The ETAS model

The epidemic type aftershock model (ETAS) is usually written in terms of time varying intensity or frequency of a Poisson point process of events in time and magnitude (see (Ogata, 1988)). Here we shall limit ourselves to time dependent processes and do not consider the spatial aspect so that we have

$$\mu(t, m | \mathcal{H}_t) = 10^{a-bm} + \sum_{t_i \leq t} h(t - t_i) 10^{\kappa + \alpha m_i + a - bm}, \quad m_0 \leq m < m_{max}.$$

The notation \mathcal{H}_t stands for (the σ -algebra generated by) all events occurring before the time t .

The first term on the right-hand side is the background process. It is a Poisson point process in time \times magnitude with intensity

$$g(t, m) = 10^{a-bm}.$$

The term intensity (or frequency) refers to the fact that in any time \times magnitude region Ω the number of events N follows a Poisson distribution with frequency Λ

$$N \sim \frac{\Lambda^N}{N!} e^{-\Lambda},$$

and the rate Λ is simply

$$\Lambda = \int_{\Omega} g(m, t) dm dt.$$

The magnitudes are independent of the event times and are following a Gutenberg Richter distribution law

$$m | t_i \sim \frac{b \ln(10) 10^{-bm}}{10^{m_0} - 10^{m_{max}}}, \quad m_0 \leq m \leq m_{max}.$$

and the total Poisson rate of events with $m_0 < m < m_{max}$ per time is

$$\text{rate of } t_i = \int_{m_0}^{m_{\infty}} g(m) dm = \frac{1}{b \ln(10)} 10^a (10^{-bm_{max}} - 10^{-bm_0})$$

The kernel h that distributes the triggered seismicity to future times is assumed to be normalized

$$\int_0^\infty h(\tau) d\tau = 1.$$

It is commonly assumed to be of the Omori family

$$h(t) = \frac{C}{(1+t/c)^p}, \quad p > 1,$$

however, for our analysis this specific form does not play any role.

Then we have to following trigger function that describes the Poisson point process intensity of the aftershocks generated by an event at time τ and magnitude m'

$$t, m | \tau, m' \sim g(m) h(t - \tau) F(m')$$

with F the productivity function

$$F(m) = 10^{\kappa + \alpha m}.$$

We agree, that $m(t)$ denotes the formal catalog function

$$m(t) = m_i \text{ if } t = t_i, \quad m(t) = -\infty \text{ if no event at } t$$

With this notation we have $F(m(t)) = 0$ whenever there is no event at t We then can write the ETAS model more concisely as

$$\mu(t, m | \mathcal{H}_t) = g(m) + \sum_{\tau \leq t} g(m) F(m(\tau)) h(t - \tau)$$

The sum over all times $\tau \leq t$ is well-defined since only a finite number of times actually contribute, at least as long as $m_0 > 0$.

Then, the random process defined by the ETAS equation can be realized in two different ways.

The causal sequential picture. We simply follow the equation as stated. So, starting from a single random event time t_1 drawn from the appropriate exponential distribution (the inter-event time distribution of a Poisson process with constant time rate), we pick independently a magnitude m_1 from the Gutenberg Richter distribution. This will change the Poissonian rate for all upcoming events. We pick the next time point from the altered non-stationary Poisson rate, sample a magnitude and continue that way. For a finite smallest magnitude $m_0 > -\infty$, this strategy is valid, since with probability one, in every step the non-stationary Poisson rate of the future events is bounded, and thus the next event time is always strictly $t_{i+1} > t_i$ yielding an increasing sequence of times. So the ETAS model exists for some time at least. It may however happen, that the time intervals become shorter and shorter leading to a run-away effect in the seismicity in finite time. Under conditions linked to the branching ratio this however does not happen and the sequence of “next” events grows until it leaves $[0, T]$ and the whole catalog is generated in a finite time. This approach however fails if we want to include arbitrary small events, of which the frequency tends to ∞ . Indeed then, there is no “next” event anymore. It is like the rational numbers, where there isn’t any “next” number after $1/2$, say.

Generation picture. Instead, we could also use the additivity of the Poisson process to proceed in a non-causal way. The sum of two Poisson point processes with non-stationary rate $\lambda_1(t, m)$ and $\lambda_2(t, m)$ is again a Poisson process with rate $\lambda(t, m) = \lambda_1(t, m) + \lambda_2(t, m)$. This means we can either directly sample from λ or we sample from λ_1 and λ_2 independently and join the catalogs. This leads to the following strategy. In order to obtain a random realization of the ETAS model for $t \in [0, T]$ we start from a random catalog of background events defined through the background intensity $g(t, m)$ For each of

the finely many background events we sample from the triggered events, discarding all events with $t > T$, to obtain the generation 1 events, and so on. With probability one, for finite m_0 and a branching ratio < 1 this procedure comes to an end. Indeed, the total number of siblings of the Galton-Watson process is finite. For a branching ratio > 1 , the expected number of events triggered by each background event grows exponential with each generation. Note however, that even for a branching ratio > 1 , still with a finite probability all background events in $[0, T]$ only have a finite total number of siblings and again the ETAS summation processes comes to an end with probability > 0 .

It has been argued, and given the above description it seems reasonable, that the ETAS model undergoes a kind of phase transition, whenever the branching number (i.e. the expected number of earthquakes triggered by an earthquake) exceeds the threshold of 1 see e.g. (Helmstetter & Sornette, 2002). In (Nandan et al., 2021) the authors analyze the criticality of the earthquake generation process with the help of these characteristics. For that reason, in order to allow for a sub-critical branching ratio, until now, the lower threshold m_0 below which no earthquakes are triggered anymore had to be set to a finite positive value since otherwise, the branching number which is based on the expected number of siblings would be necessarily infinite. In this paper I want to show that if instead of looking at the number of earthquakes, we define a productivity ratio as a measure for the aftershock activity, the ETAS model can be defined easily to include all magnitudes down to $m_0 = -\infty$. Clearly the number of earthquakes when including micro-seismicity to arbitrary small magnitudes will go to ∞ however the total cumulated impact on the triggering capacity remains finite.

The lower threshold m_0 that is thought to be necessary for the ETAS model to be well-defined has to be distinguished from an observational threshold below which the earthquakes can not be detected anymore. This parameter m_0 however constitutes an ad hoc chosen parameter for the model. It is essentially impossible to estimate this value from data. It is rather unnatural to assume that the scaling law of the Gutenberg Richter distribution is not valid down to micro seismic events. Clearly at some very small scale the physics may change completely, but the assumed cut-off scales m_0 are way above the limit at which in laboratory experiments the fundamental scaling laws of earthquake physics can still be observed.

In this paper, I show, how the ETAS model formulated in the language of jump processes and Levy processes can be extended consistently to incorporate all scales of microseismicity. This opens the way to build statistically useful models that absorbs the unobserved microscale into a Brownian motion component of some micro-seismic Levy hum. The aim of this paper however is to lay down the basic mathematical tools from Levy type jump processes.

Although not strictly necessary mathematically we propose to describe the triggering of earthquakes by the time advance of some background type seismic activity. Thus, once we have understood the background seismicity including arbitrary small magnitude events, we can understand the ETAS process as an infinite superposition of time advanced back-ground processes. Each of these processes has infinite many events, however the time scales stretch in a geometric fashion making the sum of all these time scaled back-ground processes converge to a limit process.

We first show the principle for an ETAS model that depends only on time. In the last chapter we show how to extend the description to a space-time ETAS model.

2 The ETAS model in moment space

In (Kagan, 2002), Kagan has proposed to write the Gutenberg Richter law in Moment space in which case it becomes a Pareto distribution. Although it would be possible to do everything in the magnitude parametrization of the event size, we introduce

an alternative parametrization of the magnitude that is better suited for the analysis of the ETAS dynamics. We call it the productivity moment scale Y since it measures the magnitude of an earthquake in terms of its productivity to trigger other earthquakes. So we define

$$Y = F(m).$$

This is possible since F is strictly monotonically growing and thus no information is lost by replacing m with Y . From now on we set $m_0 = -\infty$. Then the background Poisson intensity in time $t \times$ productivity moment Y is stationary and can be computed as the change of variable formula for the density of a measure. It comes

$$t, Y \sim \psi(Y) = \frac{g(t, F^{-1}(Y))}{F'(F^{-1}(Y))} = \gamma Y^{-1-b/\alpha}, \quad 0 < Y \leq Y_{max}$$

We suppose that $\alpha > b$. This is necessary to ensure our construction to make sense for arbitrary small moments (see below). Since we assume the background to be stationary in time, the right-hand side is a function of Y only. The constant γ and the maximal productivity moment, Y_{max} , are given by

$$\gamma = \frac{10^{a+\kappa b/\alpha}}{\ln(10)\alpha}, \quad Y_{max} = F(m_{max}) = 10^{\kappa+\alpha m_{max}}.$$

The right hand can be understood as the symbol of a Levy jump process. Then, for any interval $I \subset (0, \infty)$, the integral $\int_I \psi(Y) dY$ would be the Poisson rate of jumps with jump size in I . We want to make this construction a little more explicit. Fix an arbitrary time interval $[0, T]$. Then consider a random catalog generated from the Poisson intensity $\eta(t, Y) = \psi(Y)$. Since the total integral is infinite

$$\int_0^T \int_0^\infty \eta(t, Y) dt dY = T \int_0^{Y_{max}} \psi(Y) dY = \infty,$$

the standard way of sampling from a non-homogeneous (in Y) Poisson point process does not apply since we can not sample from a Poisson distribution P_λ with parameter $\lambda = \infty$. However, we may decompose the magnitudes into disjoint dyadic intervals (see Fig 3)

$$I_k = (Y_{max}2^{-k+1}, Y_{max}2^{-k}], \quad \bigcup_{k=0}^{\infty} I_k = (0, Y_{max}].$$

Now for each I_k we may sample as usual. That is we compute the total rate of events in $[0, T] \times I_k$

$$N_k = \int_0^T \int_{I_k} \psi(Y) dY \sim 2^{kb/\alpha}.$$

We then draw a Poisson variable $L_k \sim P_{N_k}$. Given this random number of events, we distribute L_k events in time and moment according to the pdf with density

$$p(t, Y) = \frac{\psi(Y)}{TN_k}, \quad (t, Y) \in [0, T] \times I_k$$

This defines a random catalog Ξ_k with moments in $Y_k \in I_k$. Then, the collection of these sub-catalogs defines the total catalog

$$\Xi = \bigcup_k I_k.$$

It contains infinite many events. However, the events are still countable, since in any slab I_k we only have a finite number. Moreover, with probability one no two events share the same time point. Therefore, we may replace the catalog set Ξ also by the catalog function $Y(t)$ defined as follows. It is zero except for those points $t = \tau$ at which we have an event $(\tau, Y) \in \Xi$. Since there is at most one, we can define $Y(t)$ without ambiguity as the productivity moment of the event at t . Note that $Y(t)$ is not a measure, it is




Figure 1. The jump distribution can be cut into dyadic blocks.

a function that is zero almost everywhere. However, It may be associated with a measure via

$$\nu_Y = \sum_{\tau} Y(\tau) \delta_{\tau}$$

The sum is well-defined, since it actually runs only over a countable set only and the sum of the amplitudes of the delta function have a finite value in any finite time interval almost surely (see below). We now consider the cumulated productivity moments

$$C_Y(t) = \sum_{\tau \leq t} Y(\tau) = \chi_{[0, \infty)} * \nu_Y(t).$$

This defines a non-decreasing function which is right continuous but not necessarily left continuous. However, the left limits exist and thus the jumps are well-defined. These jumps happen at all event times

$$Y(t) = \Delta C_Y(t) = C_Y(t) - C_Y(t^-), \quad C_Y(t^-) = \lim_{\tau \rightarrow t, \tau < t} C_Y(\tau).$$

This so called cadlag property (continue à droite limit à gauche) allows us to map seismic catalogs containing arbitrary small events to such well-defined semi-continuous non-decreasing functions. Vice versa any such non-decreasing cadlag function defines a possibly infinite, not time-ordered, seismic catalog and a catalog function $Y(t)$. As a random process, the process $C_Y(t)$ has independent increments and all its finite increments are infinitely divisible (see e.g. (Doob, 1953)).

In order to actually compute (better approximate) such a function numerically we can again resort to the decomposition I_k . If $Y_k(t)$ and $C_{Y_k}(t)$ are the corresponding cat-




Figure 2. In the upper figure is depicted a random sample the Levy path for the first three moment blocks. The blue line is an approximation to the limit by using only the first 3 blocks.

alog functions and cumulated moment functions, we then have the following scaling relation

$$Y_{k+1}(t) \simeq 2^{-1}Y_k(2^{\alpha/b}t), \quad C_{Y_{k+1}}(t) \simeq 2^{-1}C_{Y_k}(2^{\alpha/b}t)$$

where the symbol \simeq means both process have the same distributions. Indeed, going from block k to block $k+1$ the jump-size is scaled by $1/2$ whereas the frequency increases by a factor $2^{\alpha/b} > 1$. Therefore, starting from the process Y^0 and C_{Y^0} we have the following Weierstrass fractal function like multi-scale expansion for the background process Y^B and C_{Y^B}

$$Y^B(t) \simeq \sum_{k=0}^{\infty} 2^{-k}Y_0(2^{k\alpha/b}t), \quad C_{Y^B}(t) \simeq \sum_{k=0}^{\infty} 2^{-k}C_{Y_0}(2^{k\alpha/b}t)$$

The terms in the sum are each independent, rescaled (in time and event/jump size) realizations of the common band-limited background process. In Figure 2 we have shown an example of this construction.

The average value of $C_{Y^B}(t)$ may be computed explicitly. By Campbell's theorem (see Appendix A) we have

$$\mathbb{E}(C_{Y^B}(t)) = \int_0^t \int_0^{Y_{max}} Y \psi(Y) dY = t\mathbb{E}(Y)$$

Therefore with probability one the growth rate of $C_{Y^B}(t)$ is given by the expectation of the productivity of the background process.

$$D = \lim_{t \rightarrow \infty} \frac{1}{t} C_{Y^B}(t) = \mathbb{E}(Y) = \mathbb{E}(F(m))$$

So explicitly

$$D = \int_{Y>0} Y \psi(Y) dY = \int g(m) F(m) dm = \gamma \int_0^{Y_{max}} Y^{-b/\alpha} = \frac{\gamma Y_{max}^{1-b/\alpha}}{1-b/\alpha}$$

or in terms of the magnitude limit m_{max}

$$D = \frac{10^{a+\kappa+(\alpha-b)m_{max}}}{\ln(10)(\alpha-b)}.$$

To ensure a finite average productivity coming from the small events, we need to assume

$$\alpha > b, \quad Y_{max} < \infty.$$

This same condition is also necessary and sufficient that all the previous computations have a well-defined meaning. In the next section we shall add the condition $D < 1$ to ensure the ETAS model to be well-defined.

In general a rate $\eta(t, Y)$ of jumps may be time dependent and the band limited construction can be adapted accordingly. For the ETAS model a simplification however occurs in case that the distribution of triggered event magnitudes does not depend on the magnitude of the triggering event. That is, we assume for simplicity that Bath's law does not hold. Then the triggered seismicity may be absorbed in a change of time as we shall see now. In order to include Bath's law, we would have to include additional filter operators, which we leave for now.

3 The clock advance picture

We now rewrite the triggering process using the language of Levy pure jump processes. For a single triggering event at τ and moment $Y(\tau)$ the next generation events triggered by this event are drawn from the rate

$$Y, t | \tau Y(\tau) \sim \eta(t, Y(\tau)) = \psi(Y) Y(\tau) h(t - \tau).$$

We now use the following fact that shows how amplitude modulation of the background process can be absorbed into time deformations.

Fact If a jump process has non-stationary jump size Poisson intensity $\eta(t, Y)$ that factorizes

$$\eta(t, Y) = f(t) \psi(Y)$$

then the Levy process of the cumulated Y satisfies

$$C_Y^\eta(t) = C_Y(F(t)), \quad F'(t) = f(t), \quad F(0) = 0$$

The equality means, the right and left side have the same distribution. This is intuitively clear since if locally the time is scaled by $F' = f$ its local frequency is multiplied by the same factor.

We introduce the elementary clock advance function H

$$H(t) = \int_0^t h(\tau) d\tau.$$

Then let $C_{YB}(t)$ be an independent process of cumulated background moments. The triggered events by the event at $\tau, Y(\tau)$ can therefore be identified with the clock modified background process

$$C_{YB}(Y(\tau)H(t - \tau)),$$

or on the catalog function itself

$$Y^B(Y(\tau)H(t - \tau))$$




Figure 3. The triggered events can be realized by modifying the local time of an independent background process. Here we illustrate this by an event of unit size $Y = 1$. The clock modified catalog is displayed horizontally whereas the original background catalog is along the vertical axis. Only the events colored in red will be in the locally-clock-advanced catalog.

If we have two events at τ_1 and τ_2 we have for the catalog

$$Y^{B,1}(Y(\tau_1)H(t - \tau_1)) + Y^{B,2}(Y(\tau_2)H(t - \tau_2))$$

with $Y^{B,i}$, $i = 1, 2$ two independent realizations of the background. Since the sum of two independent Poisson processes with two rates is the same as a single process with the sum of the rates we can the above process by time deforming a single realization of the background

$$Y^B(Y(\tau_1)H(t - \tau_1) + Y(\tau_2)H(t - \tau_2)).$$

We therefore can define a total clock transform function that is applied to the background process via

$$H^* = \sum_{\tau} Y^B(\tau)H(t - \tau) = H * \sum_{\tau} Y^B(\tau)\delta_{\tau} = h * C_{Y^B}(t) = H * \nu_{Y^B}(t).$$

The process of the first generation of triggered events is then

$$C_{Y^T}(t) = C_{Y^B}(H^*(t)).$$

Note that the local clock function H^* inherits the continuity from h , and it is strictly monotonically growing. It therefore defines a valid time.

This can be iterated to obtain the sequence of triggered aftershocks. If $Y^{T,k}(t)$ is the catalog function of the triggered events triggered at generation k , we define the clock advance function as

$$H^{*,k}(t) = h * C_{Y^{T,k}}(t) = H * \nu_{Y^{T,k}}(t)$$

and the triggered events and the corresponding Levy path at generation $k+1$ are then

$$Y^{T,k+1}(t) = Y^{B,k}(H^{*,k}(t)), \quad C_{Y^{T,k+1}} = C_{Y^{B,k}}(H^{*,k}(t))$$

with independent background process $Y^{B,k}(t)$. The ETAS catalog function / path would then be

$$Y^{ETAS}(t) = Y^B(t) + \sum_{k=1}^{\infty} Y^{T,k}(t), \quad C_{Y^{ETAS}}(t) = C_{Y^B}(t) + \sum_{k=1}^{\infty} C_{Y^{T,k}}(t).$$

Observe that the cumulated productivity moments on the right are not a Levy process, since the increments are not independent. However, they constitute still a non-decreasing, cadlag (continue à droite, limit à gauche) function. The jumps are precisely the events of the infinite catalog

$$Y^{ETAS}(t) = \Delta C_{Y^{ETAS}}(t).$$

We still have to show, that the sum of random functions actually converges almost-surely. This will be the case whenever the expected productivity is < 1

$$D = \mathbb{E}(Y) < 1 \Leftrightarrow \text{ETAS sum converges (with probability one)}$$

We only give a schematic proof here. For that note that the longtime slope of the background Levy path, $C_{Y^B}(t)$ is D , $C_{Y^B}(t)/t \rightarrow D$ for $t \rightarrow \infty$. Suppose $C_Y^k(t)$ has a slope of D_k

$$t^{-1}C_Y^k(t) \rightarrow D_k \quad t \rightarrow \infty$$

Then the clock advance function $H^{*,k}(t) = h * C_Y^k(t)$ has the same slope since convolution with h reproduces the slope

$$\int_0^t h(t-\tau)\tau d\tau = -H(t-\tau)\tau|_0^t + \int_0^t H(\tau-t)d\tau = \int_0^t H(\tau)d\tau \sim t \quad (t \gg 1)$$

Therefore the time in generation k is asymptotically rescaled by D_k . If this is applied to the background it produces a process with slope

$$t^{-1}C_{Y^B}(H^{*,k}(t)) \rightarrow DD_k \quad \Rightarrow D_{k+1} = DD_k = D^{k+1}.$$

By induction, we have a geometric series which can be summed to show that

$$\lim_{t \rightarrow \infty} \frac{1}{t} C_{Y^{ETAS}}(t) = \frac{D}{1-D}$$

Note however that this is a rather hand-wavy non-rigorous argument since it does not take into account the random nature of these slopes. We refer to a forthcoming paper on this subject where we use a martingale argument.

4 A Markov chain in infinite dimensional state space

It is interesting to note, that our algorithm can be cast into a Markov process with an infinite dimensional state space. Consider a sequence $C_B^k(t)$, of independent background processes. The state is given by a pair of $(E(t), W(t))$ of non-decreasing right-continuous functions. The transition process is defined through

$$E_{k+1}(t) = E_k(t) + W_k(t), \quad W_{k+1} = C_B^{k+1}(h * W_k(t))$$

Then, starting from $(0, C_B^0)$ this defines a Markov chain. Under the stability condition above we have that the ETAS sampling paths are obtained as a limit

$$\lim_{k \rightarrow \infty} E_k = Y^{ETAS}$$

5 Extension to spatial processes

On the level of Poisson intensity in time and trigger moment space the triggering dynamic can be written as

$$\eta^{k+1}(t, Y) = \sum_{\tau} Y^{T,k}(\tau) h(t - \tau) \psi(Y)$$

Sampling from this Poisson density can be achieved through time deforming a sample from an independent background sample $Y^{B,k}(\tau)$

$$Y^{T,k}(t) = Y^{B,k}(H^{*,k}(t))$$

This description can be extended to spatially distributed ETAS processes.

For this we need to consider space-time Point processes described through an intensity $\eta(x, t, Y)$ of the form

$$\eta(x, t, Y) = f(x, t) \psi(Y).$$

This factorization comes from our assumption, that the magnitude distribution and hence the production moment distribution of the triggered events is the same as the one of the background.

Instead of considering Poisson point processes in the space \times time \times moment space, we can again look at the cumulated moments. For that for a spatial domain Ω the cumulated Y of the events taking place in Ω produce a Levy jump process.

$$C_{\Omega, Y}(t) = \sum_{(x, \tau) | x \in \Omega, \tau \leq t} Y(\tau, x)$$

It can be realized through a clock advance of the background process. For this we define a spacial clock advance function of every space point x via

$$H(x, t) = \int_0^t f(x, \tau) d\tau.$$

Then, the clock advance in Ω is

$$H_{\Omega}(t) = \int_{\Omega} H(x, t) dx = \int_0^t \int_{\Omega} f(x, \tau) dx d\tau$$

The process of the time points of the events in Ω as described through η can be realized by clock modifying the one dimensional background process $C_Y^B(t)$ of section 3.

$$C_{\Omega, Y}(t) \sim C_{Y^B}(H_{\Omega}(t))$$

On the level of the catalog function over space \times time itself we can write for the clock advanced catalog

$$Y^{\eta}(x, t) = Y^B(x, H(x, t)).$$

We are actually considering here a Levy process valued measures (see e.g (Griffiths & Riedle, 2021)). In the spatial extension of the ETAS process, the seismicity is redistributed not only in time but also in space via a kernel, which may depend on Y . The conditional Poisson intensity reads

$$x, t, Y | x', \tau, Y' \sim \Gamma(x | x', Y') \psi(Y) Y' h(t - \tau) \quad (1)$$

We suppose that Γ is normalized

$$\int \Gamma(x | \dots) dx = 1 \quad (2)$$

so that Y maintains its meaning as productivity. Then the ETAS equations read

$$\mu(x, t, Y | \mathcal{H}_t) = \psi(Y) + \sum_{(x', \tau) | \tau \leq t} \Gamma(x | y', Y(x', \tau)) Y(x', \tau) h(t - \tau) \psi(Y)$$

The ETAS generation iteration is now defined as follows. Let $B^k(t, x)$, $k = 0, 1, \dots$ be independent background processes. We then consider the following Markov chain of pairs of processes (E, W) starting from $(0, B^0)$

$$E_{k+1}(x, t) = E_k(x, t) + W_k(x, t), \quad W_{k+1}(x, t) = B^{k+1}(x, H_k(x, t))$$

The Poissonian intensity of generation $k+1$ can be computed from the events at generation k as

$$\eta^{k+1}(x, t, Y) = f^{k+1}(x, t) \psi(Y),$$

with

$$f^{k+1}(x, t) = \sum_{x', \tau} \Gamma(x | x', W_k(x', \tau)) W_k(x', \tau) h(t - \tau)$$

The clock advance function that maps background events to triggered events reads

$$H_k(x, t) = \sum_{x', \tau} \Gamma(x | x', W_k(x', \tau)) W_k(x', \tau) H(t - \tau)$$

If this process converges, the resulting limit process of E_k , $k \rightarrow \infty$ is the ETAS process including microseismicity. Again, the condition for convergence is that the expected productivity is < 1 .

6 Conclusion and outlook

In this paper I have shown that the concept of Levy jump processes allows the extension of the ETAS model to arbitrary small earthquakes under the condition that the expected productivity remains < 1 . The productivity is measured in units of the background process and not in terms of the number of siblings of a triggering earthquake. The latter is infinite all the time if we include micro seismicity. This shows that the non-physical small magnitude cutoff can be removed in a consistent way. The consequences for the estimation of the ETAS parameters needs to be analyzed. It is clear, that the specific form of the Gutenberg Richter law does not play a role. We only need that the productivity function has a first moment. Therefore, for instance we may assume a completely scale-free Gutenberg Richter law $\sim 10^{-bm}$ without cutoff but with a two-scale productivity law

$$F(m) \sim 10^{\alpha-m}, \quad m < m^*, \quad \text{and} \quad F(m) \sim 10^{\alpha+m} \quad m \geq m^*$$

We only need

$$\alpha_- > b, \quad m < m^* \quad \text{and} \quad \alpha_+ < b \quad m > m^*,$$

to ensure a finite total productivity function. Adjusting the coupling constants we can achieve a mean moment productivity < 1 , which ensures the convergence of the ETAS iteration.

Acknowledgments

This work was financed through the SFB1294 CRC research grant.

Appendix A Campbell's theorem

We need the theorem of total expectation

$$\mathbb{E}(A) = \mathbb{E}_B \mathbb{E}_A(A|B),$$

Moreover we need Campbell's theorem. It states that for a point process with intensity Λ over some domain and a function g the random sum

$$G = \sum_x g(x)$$

has mean value

$$\mathbb{E}(G) = \int g(x)\Lambda(x)dx$$

Therefore, if Λ itself is a random process, we may apply the iterated expectation and it comes

$$\mathbb{E}(G) = \int g(x)\mathbb{E}(\Lambda(x))dx$$

References

- Doob, J. L. (1953). *Stochastic processes*. New York: John Wiley & Sons. (MR 15,445b. Zbl 0053.26802.)
- Griffiths, M., & Riedle, M. (2021, May). Modelling lévy space-time white noises. *Journal of the London Mathematical Society*, *104*(3), 1452–1474. doi: 10.1112/jlms.12465
- Helmstetter, A., & Sornette, D. (2002). Subcritical and supercritical regimes in epidemic models of earthquake aftershocks. *Journal of Geophysical Research: Solid Earth*, *107*(B10), ESE 10-1-ESE 10-21.
- Kagan, Y. Y. (2002, 03). Seismic moment distribution revisited: I. statistical results. *Geophysical Journal International*, *148*(3), 520-541. Retrieved from <https://doi.org/10.1046/j.1365-246x.2002.01594.x> doi: 10.1046/j.1365-246x.2002.01594.x
- Nandan, S., Ram, S. K., Ouillon, G., & Sornette, D. (2021, Mar). Is seismicity operating at a critical point? *Phys. Rev. Lett.*, *126*, 128501.
- Ogata, Y. (1988). *Journal of the American Statistical Association*, *83*(401), 9-27.