ABSTRACT REGULAR POLYTOPES OF FINITE IRREDUCIBLE COXETER GROUPS

MALCOLM HOONG WAI CHEN AND PETER ROWLEY

ABSTRACT. Here, for W the Coxeter group D_n where n > 4, it is proved that the maximal rank of an abstract regular polytope for W is n-1 if n is even and n if n is odd. Further it is shown that W has abstract regular polytopes of rank r for all r such that $3 \le r \le n-1$, if n is even, and $3 \le r \le n$, if n is odd. The possible ranks of abstract regular polytopes for the exceptional finite irreducible Coxeter groups are also determined.

1. INTRODUCTION

Finite Coxeter groups appear in many guises in the mathematics literature – as Weyl groups of semisimple Lie algebras, reflection groups, and automorphism groups of regular polytopes to mention a few. It is the last area that is of interest here. We recall that Sym(n+1), the symmetric group of degree n+1, also the Coxeter group of type A_n is the automorphism group of a regular *n*-simplex, while the Coxeter group of type B_n is the automorphism group of the *n*-cube and its dual, the *n*-cross polytope.

In this paper we examine abstract regular polytopes of the finite irreducible Coxeter groups W. As is documented in McMullen and Schulte [27], this is equivalent to investigating the C-strings of W, and we follow that approach. A *C-string* of a group G is a set of involutions $S = \{s_1, \ldots, s_r\}$ of G which generates the group. Additionally, setting $I = \{1, \ldots, r\}$ and, for $J \subseteq I, W_J = \langle s_j \mid j \in J \rangle$, they must satisfy

(i) for all $J, K \subseteq I, W_J \cap W_K = W_{J \cap K}$ (intersection property); and

(ii) $s_i s_j = s_j s_i$ for all $i, j \in I$ with $|i - j| \ge 2$ (string property).

We refer to r as the rank of the C-string, equivalently, the rank of the associated abstract regular polytope, and also call (G, S) a string C-group. The maximal rank for a C-string of W will be denoted by $r_{\max}(W)$. If W is either of type A_n or B_n , then their defining presentation in terms of fundamental (or simple) reflections yields a C-string of rank n (see Section 5.5 of [18] for the intersection property). Moreover $r_{\max}(W) = n$ when $W \cong A_n$, a consequence of a recent deep result by Whiston [30] which uses the classification of finite simple groups. For $W \cong B_n$, $r_{\max}(W) = n$ when n is even and $r_{\max}(W) = n + 1$ when n is odd. In the latter case the corresponding C-string is degenerate, meaning the string is disconnected. See Theorem 2.5(ii) for more details. If we insist on only considering non-degenerate strings, then we have $r_{\max}(W) = n$. On the other hand, $I_2(m)$, identified as Dih(2m) are also well known to only exhibit rank two polytopes given

²⁰²⁰ Mathematics Subject Classification. 52B11, 20B25, 20F55.

 $Key\ words\ and\ phrases.$ Abstract regular polytopes, string C-groups, finite irreducible Coxeter groups.

by the *m*-gon. Now in the remaining infinite family of finite irreducible Coxeter groups we have D_n whose Dynkin diagrams are not strings. Our first result concerns the maximal rank of C-strings for this family. Noting that D_4 has no C-strings, we have the following theorem.

Theorem 1.1. Suppose that W is the Coxeter group D_n with $n \ge 5$. If n is even, then $r_{\max}(W) = n - 1$ and if n is odd, then $r_{\max}(W) = n$.

The question as to the existence of abstract regular polytopes for intermediate rank values is the subject of the next theorem.

Theorem 1.2. Suppose that W is the Coxeter group D_n with $n \ge 5$. Then W has a C-string of rank r for all r with $3 \le r \le r_{\max}(W)$.

To complete the picture we consider the exceptional finite irreducible Coxeter groups.

Theorem 1.3. Suppose that W is an exceptional finite irreducible Coxeter group.

- (i) If W is one of $I_2(m)$, H_3 , H_4 , F_4 , then $r_{max}(W)$ is the Coxeter rank of W.
- (ii) If W is one of E_6, E_7, E_8 , then $r_{max}(W)$ is 5, 6, and 7, respectively.

The quest to classify string C-groups started from early experimental results by Hartley [16] and also the joint work of Leemans and Vauthier [26] which resulted in atlases of abstract regular polytopes for small groups. More efficient computer algorithms have also been developed which successfully enumerated the C-strings of some sporadic simple groups [17, 21, 22]. This experimental data has led to some interesting conjectures which were later proven theoretically. In particular, the possible ranks of C-strings has been investigated for some infinite families of almost simple groups, namely the Suzuki groups Sz(q) [19], Ree groups ${}^{2}G_{2}(q)$ [25], groups with socle PSL(2, q) [11, 23, 24], groups PSL(3, q) and PGL(3, q) [6], groups PSL(4, q) [4], symmetric groups [13], alternating groups [10, 14], orthogonal and symplectic groups [2, 3]. We refer the interested reader to a recent survey article by Leemans [20] for more details of these investigations.

Only a few family of groups are known to give rise to string C-groups of arbitrarily large rank. As already noted, the highest possible rank of a string C-group representation for $\operatorname{Sym}(n)$ is n-1, and for $n \geq 12$, the highest rank for $\operatorname{Alt}(n)$ is $\lfloor \frac{n-1}{2} \rfloor$ [10]. Furthermore, $\operatorname{Sym}(n)$ has C-strings of rank r for every $3 \leq r \leq n-1$ [13], and for $n \geq 12$, $\operatorname{Alt}(n)$ also has C-strings of rank r for every $3 \leq r \leq \lfloor \frac{n-1}{2} \rfloor$ [14]. Likewise for all integers $k, m \geq 2$, the orthogonal groups $O^{\pm}(2m, 2^k)$ and the symplectic groups $\operatorname{Sp}(2m, 2^k)$ have string C-group representations of rank r and 2m + 1 [3], respectively. Interestingly in [5], Brooksbank proved a 'rank reduction theorem' which can be applied on known string C-groups to obtain C-strings of the same group with smaller ranks, and consequently it was shown that $O^{\pm}(2m, 2^k)$ and $\operatorname{Sp}(2m, 2^k)$ have string C-group representations of rank n/2 for a transitive group of degree n see [15]. Finally, we mention a further study in which two interesting families of C-strings for B_n are constructed [28].

This paper is structured as follows. In Section 2, we state some preliminary results on Coxeter groups and string C-groups. Lemma 2.1 plays an important role as our investigations into $W \cong D_n$ are conducted in Sym(2n). Section 3 begins with Theorem 3.1 showing that $r_{\max}(W) < n$ when n is even. The remainder of

this section focuses on constructing various C-strings for W and then Lemmas 2.6 and 2.7 together with Theorem 2.8 are called upon. Our final section gives a census of abstract regular polytopes for the exceptional finite irreducible Coxeter groups.

2. Preliminaries

We recall that a *Coxeter group* is a group W with presentation $\langle s_1, \ldots, s_n | (s_i s_j)^{m_{ij}} \rangle$ where $m_{ii} = 1$ and $m_{ij} = m_{ji} \geq 2$ is a positive integer or infinity for every $1 \leq i < j \leq n$ (infinity being read as no relation). The *Coxeter rank* of W is n. Associated with this presentation is the *Coxeter diagram* where the nodes correspond to the s_i with a bond between s_i and s_j if $i \neq j$ and $m_{ij} > 2$. If this diagram is connected, then W is said to be an *irreducible* Coxeter group. Coxeter [12] (see also [18]) classified the finite irreducible Coxeter groups. There are four infinite families denoted A_n , B_n , D_n , $I_2(m)$, and six exceptional groups H_3 , H_4 , F_4 , E_6 , E_7 , E_8 (we shall blur the distinction between the group and its root system).

As already mentioned in Section 1, we shall focus on the Coxeter groups D_n and the exceptional groups. Suppose W is isomorphic to D_n , where n > 4. Then W = SN where $S \cong \text{Sym}(n)$ and N is a normal subgroup of W with N an elementary abelian subgroup of order 2^{n-1} . The subgroup N will sometimes be called the subgroup of (even) sign changes. In this paper we frequently view W as a subgroup of Sym(2n), as specified in our first lemma.

Lemma 2.1. [18, (2.10)] Let $\beta_0 = (1, n+1)(2, n+2)$ and $\beta_i = (i, i+1)(n+i, n+i+1)$ for every $1 \leq i \leq n-1$ be permutations in Sym(2n). Then $\langle \beta_0, \beta_1, \ldots, \beta_{n-1} \rangle$ is isomorphic to D_n .

In a similar fashion, we also have the well-known characterization of Sym(n).

Lemma 2.2. [18, (6.4)] Let G be a group with presentation $\langle s_1, \ldots, s_{n-1} | (s_i s_j)^{m_{ij}} \rangle$ where $m_{ii} = 1$, $m_{ij} = 2$ if $|i - j| \ge 2$ and $m_{ij} = 3$ if |i - j| = 1 for every $1 \le i, j \le n-1$. Then G is isomorphic to Sym(n).

Observe that in Lemma 2.1, $W \cong D_n$ is a transitive subgroup of Sym(2n). Also note, using the notation from Lemma 2.1 that $\langle \beta_1, \ldots, \beta_{n-1} \rangle \cong \text{Sym}(n)$, while products of an even number of transpositions of the form (i, n + i), for some $1 \le i \le n$ yields the normal elementary abelian subgroup of sign changes. The next lemma will be used frequently in Section 3.

Lemma 2.3. Suppose that $W \cong D_n$ with N being the subgroup of sign changes. If $H \leq W$ is such that W = HN and $H \cap N \leq Z(W)$, then W = H.

Proof. Since N is abelian and W = HN, $H \cap N \leq W$. The only non-trivial normal subgroup of W contained in N is Z(W) (if n is even) and N. Thus, as $H \cap N \nleq Z(W)$, $H \cap N = N$ which implies H = HN = W.

Now let G be a group generated by involutions s_1, \ldots, s_r with the ordering such that $s_i s_j = s_j s_i$ (or equivalently, $(s_i s_j)^2 = 1$) if $|i - j| \ge 2$ for every $i, j \in$ $I = \{1, \ldots, r\}$. Then $(G, \{s_1, \ldots, s_r\})$ is said to be a string group generated by involutions. Set $G_J = \langle s_j \mid j \in J \rangle$ for every $J \subseteq I$, and sometimes we may write G_{i_1,\ldots,i_k} in place of $G_{\{i_1,\ldots,i_k\}}$. Its Schläfli type is the sequence $\{p_1, \ldots, p_{r-1}\}$ where p_i is the order of $s_i s_{i+1}$ for every $1 \le i \le r-1$. If a symbol p_i appears k times in adjacent places in the sequence we may write p_i^k in place of writing $p_i k$ times. So a string group generated by involutions will be a string C-group if the intersection condition also holds.

A set R of elements of a group G is *independent* if for every $g \in R$, $g \notin \langle R \setminus \{g\} \rangle$. Let $\mu(G)$ denote the maximum size of an independent subset of G.

Theorem 2.4. Suppose that $G \cong \text{Sym}(m)$. Then

- (i) $\mu(G) \le m 1$.
- (ii) Assume that $m \ge 5$. If (G, S) is a string group generated by involutions and S is an independent set of size m - 1, then S is the set of Coxeter generators for G.

Proof. (i) This is a theorem of Whiston [30]. (ii) This follows from Cameron and Cara [8] which classifies independent sets of size m - 1 in Sym(m) when $m \ge 7$, while m = 5, 6 can be checked using MAGMA [1].

Theorem 2.4 is deployed in the proof of Theorem 3.1, as is our next theorem which gives an upper bound for the rank of a C-string of transitive permutation groups. This is due to Cameron, Fernandes, Leemans and Mixer and is as follows.

Theorem 2.5. [9, Theorem 1.2] Let G be a string C-group of rank r which is isomorphic to a transitive subgroup of Sym(n) other than Sym(n) or Alt(n). Then one of the following holds.

- (i) $r \le n/2$.
- (ii) $n \equiv 2 \pmod{4}$ and $G \cong B_n \cong C_2 \wr \operatorname{Sym}(n/2)$ with Schläfli type $\{2, 3^{n/2-2}, 4\}$.
- (iii) G is imprimitive and is one of the following.
 - (a) G = 6T9 with Schläfli type $\{3, 2, 3\}$.
 - (b) G = 6T11 with Schläfli type $\{2, 3, 3\}$.
 - (c) G = 6T11 with Schläfli type $\{2, 3, 4\}$.
 - (d) G = 8T45 with Schläfli type $\{3, 4, 4, 3\}$.

Here, nTj is the *j*-th conjugacy class among transitive subgroups of Sym(n) according to Butler and McKay [7].

(iv) G is primitive. In this case, n = 6 and G is obtained from the degree six permutation representation of $Sym(5) \cong PGL_2(5)$ and is the 4-simplex of Schläfli type $\{3,3,3\}$.

We will also make use of the following lemmas to verify that the generating set of involutions we construct indeed give us string C-groups.

Lemma 2.6. [27, 2E16(a) and 11A10] Let $(G, \{s_1, \ldots, s_r\})$ be a string group generated by involutions where $G_{1,\ldots,r-1}$ and $G_{2,\ldots,r}$ are string C-groups. Then G is a string C-group if either of the following conditions are satisfied.

- (i) $G_{1,...,r-1} \cap G_{2,...,r} = G_{2,...,r-1}$.
- (ii) $s_r \notin G_{1,\dots,r-1}$ and $G_{2,\dots,r-1}$ is maximal in $G_{2,\dots,r}$.

Lemma 2.7. [27, 2E16(b)] Let $(G, \{s_1, \ldots, s_r\})$ be a string group generated by involutions such that $G_{1,\ldots,r-1}$ is a string C-group and $G_{1,\ldots,r-1} \cap G_{k,\ldots,r} = G_{k,\ldots,r-1}$ for every $2 \le k \le r-1$. Then G is also a string C-group.

Once we have constructed a C-string of maximal rank, we will use the following rank reduction theorem by Brooksbank and Leemans to obtain C-strings of smaller ranks. **Theorem 2.8.** [5, Theorem 1.1 and Corollary 1.3] Let $(G, \{s_1, \ldots, s_n\})$ be a nondegenerate string C-group of rank $n \ge 4$ with Schläfli type $\{p_1, \ldots, p_{n-1}\}$. If $s_1 \in \langle s_1s_3, s_4 \rangle$, then $(G, \{s_2, s_1s_3, s_4, \ldots, s_n\})$ is a string C-group of rank n - 1. In particular, G has a C-string of rank n - i for every $0 \le i \le t$, where

 $t = \max\{j \in \{0, \dots, n-3\} \mid \text{for all } i \in \{0, \dots, j\}, p_{3+i} \text{ is odd}\}.$

3. C-Strings for D_n

Theorem 3.1. Let n be even with $n \ge 6$. If $W \cong D_n$, then $r_{\max}(W) < n$.

Proof. By Lemma 2.1 we may regard W as a subgroup of Sym(2n) and hence, appealing to Theorem 2.5, $r_{\max}(W) \leq n$. So we must show that W has no C-strings of rank n. Assuming that W has a C-string S with |S| = n, we seek a contradiction. Let $S = \{s_1, \ldots, s_n\}$ with $s_i s_j = s_j s_i$ for all $i, j \in I = \{1, \ldots, n\}$ with $|i - j| \geq 2$. Put $\overline{W} = W/N$ where N is the subgroup of W consisting of the even sign changes, and we use the bar convention. So $\overline{W} \cong \text{Sym}(n)$.

(3.1.1) For $i \in I, s_i \notin Z(W)$.

Suppose that $s_i \in Z(W)$ and set $H = \langle S \setminus \{s_i\} \rangle$. Then W/Z(W) = HZ(W)/Z(W)and therefore HZ(W) = W = HN with $H \cap N \leq Z(W)$. Thus W = H by Lemma 2.3, against S satisfying the intersection condition. This proves (3.1.1).

Let $T \subseteq S$ be such that \overline{T} is an independent generating set for \overline{W} . Hence $|\overline{T}| \leq n-1$ by Theorem 2.4(i). Further \overline{T} is a connected string in \overline{W} . For if not then $\overline{T} = \overline{T}_1 \cup \overline{T}_2$, $T_i \subseteq S$ with $[\overline{T}_1, \overline{T}_2] = 1$ and $\overline{T}_1 \neq \emptyset \neq \overline{T}_2$. But then $\langle \overline{T}_1 \rangle \langle \overline{T}_2 \rangle = \overline{W}$ with $[\langle \overline{T}_1 \rangle, \langle \overline{T}_2 \rangle] = 1$, contrary to $\overline{W} \cong \text{Sym}(n)$.

(3.1.2) $|\overline{T}| = n - 1$ or n - 2. Moreover, if $|\overline{T}| = n - 2$, then $\overline{T} = \{\overline{s}_2, \dots, \overline{s}_{n-1}\}$.

Suppose that $|\overline{T}| \leq n-3$. Then there exists $s \in S$ such that [T, s] = 1. From $\langle \overline{T} \rangle = \overline{W}$, we get $\overline{s} \in Z(\overline{W})$ and then, as $\overline{W} \cong \text{Sym}(n)$, we have $s \in N$. Hence, using (3.1.1) and Lemma 2.3, $W = \langle T \cup \{s\} \rangle$ which is impossible as S satisfies the intersection property. Thus $|\overline{T}| = n-1$ or n-2. A similar argument applies to show $\overline{T} = \{\overline{s}_2, \ldots, \overline{s}_{n-1}\}$ when $|\overline{T}| = n-2$.

 $(3.1.3) |\overline{T}| \neq n-1.$

Suppose that $|\overline{T}| = n - 1$. We may assume, as \overline{T} is connected, that $\overline{T} = \{\overline{s}_1, \ldots, \overline{s}_{n-1}\}$. Put $\overline{X} = \langle \overline{s}_1, \ldots, \overline{s}_{n-2} \rangle$. By Theorem 2.4(ii) \overline{T} consists of the Coxeter generators for \overline{W} . Thus $\overline{X} \cong \text{Sym}(n-1)$. Because $[s_n, s_i] = 1$ for $i = 1, \ldots, n-2, \ \overline{s}_n \in C_{\overline{W}}(\overline{X})$. Since $C_{\overline{W}}(\overline{X}) = 1, \ s_n \in N$. Now n-1 being odd implies $C_N(\overline{X}) = Z(W)$ giving $s_n \in Z(W)$, against (3.1.1). This rules out $|\overline{T}| = n-1$.

 $(3.1.4) |\overline{T}| \neq n-2.$

Assume that $|\overline{T}| = n - 2$. By (3.1.2) we have $\overline{T} = \{\overline{s}_2, \ldots, \overline{s}_{n-1}\}$. Put $X = \langle s_2, \ldots, s_{n-1} \rangle$ and $X_1 = \langle s_1, s_2, \ldots, s_{n-1} \rangle$. Then $W = XN = X_1N$ and, as $X \leq \langle s_1, s_2, \ldots, s_{n-1} \rangle$.

 $X_1, X_1 = X(X_1 \cap N)$. If $X_1 \cap N = 1$, then $X_1 \cong \text{Sym}(n)$ and so $X = X_1$. But this contradicts the intersection condition. Thus $X_1 \cap N \neq 1$. If $X_1 \cap N \neq Z(W)$, then Lemma 2.3 forces $X_1 = W$, also a contradiction. Hence $X_1 \cap N = Z(W)$ and consequently $X_1 = XZ(W)$. A similar argument applies for $X_n = \langle s_2, \ldots, s_{n-1}, s_n \rangle$ to also yield $X_n = XZ(W)$. But then

$$W = \langle s_1, \dots, s_n \rangle \le XZ(W) \ne W,$$

so proving (3.1.4).

Combining (3.1.2), (3.1.3) and (3.1.4) yields the desired contradiction, so establishing Theorem 3.1. $\hfill \Box$

In the remainder of this section we construct examples of C-strings for D_n . For every integer $n \ge 5$, we define t_1, \ldots, t_n in Sym(2n) as follows.

$$t_1 = \prod_{j=2}^{n} (j, n+j),$$

$$t_i = (i-1, i)(n+i-1, n+i) \text{ for every } 2 \le i \le n.$$

Lemma 3.2. For every $1 \le i, j \le n$, we have

$$order \ of \ t_i t_j = \begin{cases} 1, & if \ i = j, \\ 2, & if \ |i - j| \ge 2, \\ 3, & if \ |i - j| = 1 \ and \ \{i, j\} \neq \{1, 2\}, \\ 4, & if \ \{i, j\} = \{1, 2\}. \end{cases}$$

Proof. Each of the elements t_1, \ldots, t_n are involutions as they are defined as products of pairwise disjoint transpositions. Note that

$$t_1 t_2 = (1, 2, n+1, n+2) \prod_{j=3}^n (j, n+j),$$

$$t_i t_{i+1} = (i-1, i+1, i)(n+i-1, n+i+1, n+i) \text{ for every } 2 \le i \le n.$$

For every $2 \leq i, j \leq n$ with $|i - j| \geq 2$, t_i and t_j are disjoint, whereas for every $3 \leq k \leq n$, note that

$$t_1 t_k = (k-1, n+k)(k, n+k-1) \prod_{\substack{2 \le j \le n \\ j \notin \{k-1, k\}}} (j, n+j).$$

As such, $t_i t_j$ is a product of pairwise disjoint transpositions for every $1 \le i, j \le n$ with $|i - j| \ge 2$. The lemma follows from examining each of the products $t_i t_j$ in disjoint cycle notation.

Lemma 3.3. $\langle t_1, \ldots, t_n \rangle$ is a string C-group.

Proof. We prove by induction on k that $\langle t_1, \ldots, t_k \rangle$ is a string C-group for every $3 \leq k \leq n$. For the base case k = 3, it is easy to verify that $\langle t_1, t_2 \rangle \cap \langle t_2, t_3 \rangle = \langle t_2 \rangle$. So now suppose that k > 3 and that $\langle t_1, \ldots, t_{k-1} \rangle$ is a string C-group. Now by Lemmas 2.2 and 3.2, $\langle t_2, \ldots, t_k \rangle \cong \text{Sym}(k)$ is a string C-group and $\langle t_2, \ldots, t_{k-1} \rangle \cong \text{Sym}(k-1)$ which is maximal in $\langle t_2, \ldots, t_k \rangle$. We also have $t_k \notin \langle t_1, \ldots, t_{k-1} \rangle$ because t_k moves k-1 to k, which are in different $\langle t_1, \ldots, t_k \rangle$ -orbits. It follows from Lemma 2.6 that $\langle t_1, \ldots, t_k \rangle$ is a string C-group.

Lemma 3.4. Let n be odd and $n \ge 5$. Then $\langle t_1, \ldots, t_n \rangle$ is isomorphic to D_n .

Proof. Let $G = \langle t_1, \ldots, t_n \rangle$, $H = \langle \beta_0, \beta_1, \ldots, \beta_{n-1} \rangle$ where $\beta_0 = (1, n+1)(2, n+2)$ and $\beta_i = (i, i+1)(n+i, n+i+1)$ for every $1 \leq i \leq n-1$. Then $H \cong D_n$ by Lemma 2.1 and we want to show that G = H. Since $t_i = \beta_{i-1}$ for every $2 \leq i \leq n$, it suffices to show that $\beta_0 \in G$ and $t_1 \in H$.

We note that

$$t_1 t_2 = (1, 2, n+1, n+2) \prod_{j=3}^{n} (j, n+j),$$

whence $\beta_0 = (1, n + 1)(2, n + 2) = (t_1t_2)^2 \in G$. Now let $\alpha_{i,j} = (i, n + i)(j, n + j)$ and $\gamma_{i,j} = (i, j)(n + i, n + j)$ for every $2 \le i, j \le n$. Note that $\alpha_{2,3} = \beta_0^{\beta_2 \beta_1} \in H$. Now for $4 \le k \le n$, we have

$$\gamma_{2,k} = \beta_2 \prod_{j=4}^k \beta_{j-1}$$
 and $\gamma_{3,k} = \beta_2 \gamma_{2,k}$

It follows that for every $4 \leq i, j \leq n$, we have $\alpha_{i,j} = \alpha_{2,3}^{\gamma_{2,i}\gamma_{3,j}} \in H$ and hence, $t_1 = \alpha_{2,3}\alpha_{4,5} \cdots \alpha_{n-1,n} \in H$.

Theorem 3.5. Let n be odd and $n \ge 5$. Then $\{t_1, \ldots, t_n\}$ is a rank n C-string of D_n with Schläfli type $\{4, 3^{n-2}\}$. Therefore, $r_{\max}(D_n) = n$ and there is a rank r C-string of D_n for every $3 \le r \le n$.

Proof. The first assertion follows from Lemmas 3.2, 3.3 and 3.4. The second and third assertion follows from Theorem 2.5 and Theorem 2.8, respectively. \Box

To describe our C-strings, we will use an interesting graph called the string C-group permutation representation (CPR) graph, first introduced in [29]. Let $(G, \{s_1, \ldots, s_r\})$ be a string C-group identified as a permutation group of degree m. Then the *CPR graph* of G is an r-edge-labelled multigraph with vertex set $\{1, \ldots, m\}$ such that $\{i, j\}$ is an edge of label k if s_k moves i to j. Recall the notation for $W \cong D_n$ when $W \leq \text{Sym}(2n)$ as described in Section 2. That is, W = SN where $S \cong \text{Sym}(n)$ and N is the subgroup of even sign changes.

Theorem 3.6. [13, Lemma 21] Let $n \ge 5$ and $3 \le d \le n-2$. Then Sym(n) has a C-string of rank d with Schläffli type $\{3^{d-3}, 6, n-d+2\}$ whose CPR graph can be obtained by appending n-d vertices and edges with labels d-1 and d alternatively to the CPR graph of the d-simplex, as follows.

$$\bullet \quad 1 \quad \bullet \quad 2 \quad \bullet \quad \cdots \quad \bullet \quad d - 1 \quad \bullet \quad d \quad \bullet \quad d - 1 \quad \bullet \quad d \quad \bullet \quad \cdots \quad \bullet$$

Lemma 3.7. Let $G = \langle s_1, \ldots, s_r \rangle \leq W \leq \text{Sym}(2n)$ with $W \cong D_n$ be a string group generated by involutions. If $G_{1,\ldots,r-1} = S$ is a string C-group and $s_r \in N \setminus Z(G)$, then G = W and is also a string C-group.

Proof. By Lemma 2.3 we have G = W. Let $2 \le k \le r-1$. Since $s_r \in N$, $G_{k,\ldots,r} \le G_{k,\ldots,r-1}N$, and so $G_{k,\ldots,r} = G_{k,\ldots,r-1}(G_{k,\ldots,r} \cap N)$ by the Dedekind law. Then by Dedekind law again,

$$G_{1,...,r-1} \cap G_{k,...,r} = G_{1,...,r-1} \cap (G_{k,...,r-1}(G_{k,...,r} \cap N))$$

= $G_{k,...,r-1}(G_{1,...,r-1} \cap G_{k,...,r} \cap N)$
= $G_{k,...,r-1},$

as $G_{1,\ldots,r-1} \cap N = S \cap N = 1$. Thus the lemma follows from Lemma 2.7.

Let $n \ge 6$ be an even integer for the remainder of this section. We first construct rank three C-strings of D_n , defining t_1, t_2, t_3 in Sym(2n) as follows.

$$t_1 = (1, 2)(n + 1, n + 2)(n - 1, 2n - 1)(n, 2n),$$

$$t_2 = (2, 3)(4, 5) \cdots (n - 2, n - 1)(n + 2, n + 3)(n + 4, n + 5) \cdots (2n - 2, 2n - 1),$$

$$t_3 = (3, 4)(5, 6) \cdots (n - 1, n)(n + 3, n + 4)(n + 5, n + 6) \cdots (2n - 1, 2n).$$



Lemma 3.8. $\{t_1, t_2, t_3\}$ is a rank 3 C-string of D_n with Schläfli type $\{12, n-1\}$.

Proof. Each of the elements t_1, t_2, t_3 are involutions as they are defined as products of pairwise disjoint transpositions. Note that

$$t_1t_2 = (1,3,2)(n-1,2n-2,2n-1,n-2)(n,2n)$$

(4,5)...(n-4,n-3)(n+4,n+5)...(2n-4,2n-3),
$$t_2t_3 = (2,4,\ldots,n,n-1,n-3,\ldots,3)(n+2,n+4,\ldots,2n,2n-1,2n-3,\ldots,n+3),$$

$$t_1t_3 = (1,2)...(n-3,n-2)(n-1,2n)(n,2n-1).$$

Since $\langle t_1, t_2 \rangle \cong \text{Dih}(24)$, its elements are of the form $t_1^{\varepsilon}(t_1t_2)^k$ for some integers $\varepsilon \in \{0, 1\}$ and $0 \le k \le 11$. Also since $\langle t_2, t_3 \rangle \le S \cap \text{Stab}_{\text{Sym}(2n)}(1)$, we have

It follows that $\langle t_1, t_2 \rangle \cap \langle t_2, t_3 \rangle = \langle t_2 \rangle$ and so $\langle t_1, t_2, t_3 \rangle$ is a string C-group with Schläffli type $\{12, n-1\}$. We now show that $H = \langle t_1, t_2, t_3 \rangle$ is isomorphic to D_n . Note that $\overline{H} = S$ by Theorem 3.6 and so $HN = D_n$. Since

$$(n-2, 2n-2)(n-1, 2n-1) = (t_1t_2)^6 \in N$$

and $(n-2, 2n-2)(n-1, 2n-1) \notin Z(D_n)$. Then calling upon Lemma 2.3 yields $H = D_n$.

We now construct rank r C-strings of D_n for every $4 \le r \le n-1$ as follows. We use Theorem 3.6 to construct a rank r-1 C-string for the subgroup $S \cong \text{Sym}(n)$ of D_n , and then append an element in N. We note that the construction depends on the parity of r.

If r is odd, we define t_1, \ldots, t_r in Sym(2n) as follows.

8

If r is even, we define t_1, \ldots, t_r in Sym(2n) as follows.

$$t_{i} = (i, i+1)(n+i, n+i+1) \text{ for every } 1 \le i \le r-3,$$

$$t_{r-2} = (r-2, r-1)(r, r+1) \cdots (n-2, n-1)(n+r-2, n+r-1)$$

$$(n+r, n+r+1) \cdots (2n-2, 2n-1),$$

$$t_{r-1} = (r-1, r)(r+1, r+2) \cdots (n-1, n)(n+r-1, n+r) \cdots (2n-1, 2n),$$

$$t_{r} = (n-2, 2n-2)(n-1, 2n-1).$$

$$r-2 \qquad r-1 \qquad r-2 \qquad r-1 \qquad r-2 \qquad r-1$$

Lemma 3.9. Let n be even and $n \ge 6$. Then for every $4 \le r \le n-1$, $\{t_1, \ldots, t_r\}$ is a rank r C-string of D_n with Schläftli type $\{3^{r-4}, 6, n-r+3, 4\}$.

Proof. Each of the elements t_1, \ldots, t_r are involutions as they are defined as products of pairwise disjoint transpositions. Note that for every $1 \le k \le r-3$, we have

$$t_k t_r = \begin{cases} (k, k+1)(n+k, n+k+1)(n-1, 2n-1)(n, 2n) &, \text{ if } r \text{ is odd} \\ (k, k+1)(n+k, n+k+1)(n-2, 2n-2)(n-1, 2n-1) &, \text{ if } r \text{ is even} \end{cases},$$

which are products of pairwise disjoint transpositions because

$$k+1 \le (r-3)+1 \le ((n-1)-3)+1 = n-3 < n-2.$$

Likewise, if r is odd we have

$$t_{r-2}t_r = (n-1,2n)(n,2n-1)(r-2,r-1)(r,r+1)\cdots(n-3,n-2)$$

(n+r-2,n+r-1)(n+r,n+r+1)\cdots(2n-3,2n-2),

whereas if r is even we have

$$t_{r-2}t_r = (n-2, 2n-1)(n-1, 2n-2)(r-2, r-1)(r, r+1)\cdots(n-4, n-3)$$
$$(n+r-2, n+r-1)(n+r, n+r+1)\cdots(2n-4, 2n-3),$$

which are also products of pairwise disjoint transpositions. Finally, we have

$$t_{r-1}t_r = (n-1, n-2, 2n-1, 2n-2)(n, 2n)$$

(r-1, r)(r+1, r+2) · · · (n-4, n-3)
(n+r-1, n+r)(n+r+1, n+r+2) · · · (2n-4, 2n-3)

for r odd and we also have

$$t_{r-1}t_r = (n-2, n-3, 2n-2, 2n-3)(n-1, n, 2n-1, 2n)$$

(r-1,r)(r+1,r+2)...(n-5, n-4)
(n+r-1, n+r)(n+r+1, n+r+2)...(2n-5, 2n-4),

for r even. It follows from Lemma 3.7 that $\langle t_1, \ldots, t_r \rangle = D_n$ is a string C-group with Schläfli type $\{3^{r-4}, 6, n-r+3, 4\}$.

Theorem 3.10. Let n be even and $n \ge 6$. Then $r_{\max}(D_n) = n - 1$ and there is a rank r C-string of D_n for every $3 \le r \le n - 1$.

Proof. The theorem follows from Lemmas 3.8, 3.9 and Theorem 3.1. \Box

Together Theorems 3.1, 3.5 and 3.10 prove Theorems 1.1 and 1.2.

4. Exceptional Finite Irreducible Coxeter Groups

The following table lists the number of abstract regular polytopes (up to isomorphism and duality with the number of self-dual polytopes in brackets) for each exceptional Coxeter groups computed using MAGMA [1].

TABLE 1. Number of polytopes for each exceptional Coxeter group.

Group	Total	Rank 3	Rank 4	Rank 5	Rank 6	Rank 7	$\mathrm{Rank} \geq 8$
H_3	8(1)	8(1)	0	0	0	0	0
H_4	59(6)	45(2)	14(4)	0	0	0	0
F_4	5(1)	3(0)	2(1)	0	0	0	0
E_6	147(18)	87(12)	50(4)	10(2)	0	0	0
E_7	3662(10)	1577(10)	1525(0)	465(0)	95(0)	0	0
E_8	11689(142)	6746(117)	3584(22)	986(2)	310(0)	63(1)	0

References

- W. Bosma, J. Cannon, and C. Playoust, The Magma algebra system. I. The user language, J. Symbolic Comput. 24 (1997), no. 3–4, 235–265.
- [2] P. A. Brooksbank, On the ranks of string C-group representations for symplectic and orthogonal groups. Polytopes and Discrete Geometry, Contemp. Math. 764 (2021), 31–41.
- [3] P. A. Brooksbank, J. T. Ferrara, and D. Leemans, Orthogonal groups in characteristic 2 acting on polytopes of high rank, Discrete Comput. Geom. 63 (2020), no. 3, 656–669.
- [4] P. A. Brooksbank and D. Leemans, Polytopes of large rank for PSL(4, Fq), J. Algebra 452 (2016), 390–400.
- [5] P. A. Brooksbank and D. Leemans, Rank reduction of string C-group representations, Proc. Amer. Math. Soc. 147 (2019), no. 12, 5421–5426.
- [6] P. A. Brooksbank and D. A. Vicinsky, *Three-dimensional classical groups acting on polytopes*, Discrete Comput. Geom. 44 (2010), no. 3, 654–659.
- [7] G. Butler and J. McKay, The transitive groups of degree up to eleven, Comm. Algebra 11 (1983), no. 8, 863–911.

- [8] P. J. Cameron and P. Cara, Independent generating sets and geometries for symmetric groups, J. Algebra 258 (2002), no. 2, 641–650.
- [9] P. J. Cameron, M. E. Fernandes, D. Leemans, and M. Mixer, String C-groups as transitive subgroups of S_n, J. Algebra 447 (2016), 468–478.
- [10] P. J. Cameron, M. E. Fernandes, D. Leemans, and M. Mixer, *Highest rank of a polytope for* A_n, Proc. Lond. Math. Soc. (3) **115** (2017), no. 1, 135–176.
- [11] T. Connor, J. De Saedeleer, and D. Leemans, Almost simple groups with socle PSL(2,q) acting on abstract regular polytopes, J. Algebra 423 (2015), 550–558.
- [12] H. S. M. Coxeter, The complete enumeration of finite groups of the form $R_i^2 = (R_i R_j)^{k_{ij}} = 1$, J. Lond. Math. Soc. **10** (1935), no. 1, 21–25.
- [13] M. E. Fernandes and D. Leemans, Polytopes of high rank for the symmetric groups, Adv. Math. 228 (2011), no. 6, 3207–3222.
- [14] M. E. Fernandes and D. Leemans, String C-group representations of alternating groups, Ars Math. Contemp. 17 (2019), no. 1, 291-310.
- [15] M. E. Fernandes and C. A. Piedade, Regular polytopes of rank n/2 for transitive groups of degree n (2024), available at arXiv:2407.16003.
- [16] M. I. Hartley, An atlas of small regular abstract polytopes, Period. Math. Hungar. 53 (2006), no. 1–2, 149–156.
- [17] M. I. Hartley and A. Hulpke, *Polytopes derived from sporadic simple groups*, Contrib. Discrete Math. 5 (2010), no. 2, 106–118.
- [18] J. E. Humphreys, Reflection Groups and Coxeter Groups, Cambridge University Press, 1990.
- [19] D. Leemans, Almost simple groups of Suzuki type acting on polytopes, Proc. Amer. Math. Soc. 134 (2006), no. 12, 3649–3651.
- [20] D. Leemans, String C-group representations of almost simple groups: a survey. Polytopes and Discrete Geometry, Contemp. Math. 764 (2021), 157–178.
- [21] D. Leemans and M. Mixer, Algorithms for classifying regular polytopes with a fixed automorphism group, Contrib. Discrete Math. 7 (2012), no. 2, 105–118.
- [22] D. Leemans and J. Mulpas, The string C-group representations of the Suzuki, Rudvalis and O'Nan sporadic groups, Art. Discrete Appl. Math. 5 (2022), no. 3, Article #P3.09, 12 pp.
- [23] D. Leemans and E. Schulte, Groups of type L₂(q) acting on polytopes, Adv. Geom. 7 (2007), no. 4, 529–539.
- [24] D. Leemans and E. Schulte, Polytopes with groups of type PGL₂(q), Ars Math. Contemp. 2 (2009), no. 2, 163–171.
- [25] D. Leemans, E. Schulte, and H. Van Maldeghem, Groups of Ree type in characteristic 3 acting on polytopes, Ars Math. Contemp. 14 (2018), no. 2, 209–226.
- [26] D. Leemans and L. Vauthier, An atlas of abstract regular polytopes for small groups, Aequationes Math. 72 (2006), no. 3, 313–320.
- [27] P. McMullen and E. Schulte, Abstract Regular Polytopes, Cambridge University Press, 2002.
- [28] R. Nicolaides and P. Rowley, Two families of unravelled abstract regular polytopes in B_n , J. Group Theory **25** (2022), no. 6, 1133–1148.
- [29] D. Pellicer, CPR graphs and regular polytopes, European J. Combin. 29 (2008), no. 1, 59-71.
- [30] J. Whiston, Maximal independent generating sets of the symmetric group, J. Algebra 232 (2000), no. 1, 255–268.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANCHESTER, MANCHESTER, UNITED KINGDOM.

 $Email \ address: \verb"malcolmhoongwai.chen@postgrad.manchester.ac.uk"$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANCHESTER, MANCHESTER, UNITED KINGDOM.

Email address: peter.j.rowley@manchester.ac.uk