

# Reinforcement Learning for Respondent-Driven Sampling

Justin Weltz <sup>\*</sup>    Angela Yoon <sup>†</sup>    Yichi Zhang <sup>‡</sup>  
 Alexander Volfovsky <sup>§</sup>    Eric Laber <sup>¶</sup>

## Abstract

Respondent-driven sampling (RDS) is widely used to study hidden or hard-to-reach populations by incentivizing study participants to recruit their social connections. The success and efficiency of RDS can depend critically on the nature of the incentives, including their number, value, call to action, etc. Standard RDS uses an incentive structure that is set *a priori* and held fixed throughout the study. Thus, it does not make use of accumulating information on which incentives are effective and for whom. We propose a reinforcement learning (RL) based adaptive RDS study design in which the incentives are tailored over time to maximize cumulative utility during the study. We show that these designs are more efficient, cost-effective, and can generate new insights into the social structure of hidden populations. In addition, we develop methods for valid post-study inference which are non-trivial due to the adaptive sampling induced by RL as well as the complex dependencies among subjects due to latent (unobserved) social network structure. We provide asymptotic regret bounds and illustrate its finite sample behavior through a suite of simulation experiments.

---

<sup>\*</sup>Department of Statistical Science, Duke University, justin.weltz@duke.edu

<sup>†</sup>Department of Neurosurgery and Neurology, Duke University, angela.yoon@duke.edu

<sup>‡</sup>Department of Computer Science and Statistics, University of Rhode Island, yichizhang@uri.edu

<sup>§</sup>Department of Statistical Science, Duke University, alexander.volfovsky@duke.edu

<sup>¶</sup>Department of Statistical Science, Duke University, eric.laber@duke.edu

Social network data are at the forefront of healthcare research. It is widely acknowledged that a better understanding of community social structure can provide insights into the prevalence of mental illnesses such as depression, transmittable diseases such as HIV, syphilis, and COVID, and other conditions like obesity and type-II diabetes (Ma et al., 2007; Tabák et al., 2012; Eisenberg et al., 2013; Perry et al., 2018). Currently, available methodologies for studying the underlying social network in a community fall broadly into two categories: (1) complete community census (such as in the Framingham Heart Study, Mahmood et al., 2014), or (2) network sampling algorithms. The magnitude of modern healthcare needs makes complete censuses nearly unachievable (and generally impractical) as the relevant sampling frames are usually unknown. Consequently, network sampling techniques have proven invaluable (see Raifman et al., 2022, for a recent survey).

Respondent-driven sampling (RDS) is a network sampling algorithm based on participant referral which is frequently employed for surveillance in public health research (Heckathorn, 1997; Wejnert and Heckathorn, 2011; Heckathorn and Cameron, 2017). RDS begins with an initial group (usually a convenience sample) of individuals who are given a limited number of coupons and asked to recruit other members of the population of interest by giving them one of the coupons directly or by providing their contact information to the research team. Recipients of these initial coupons redeem them with study researchers; they are then compensated, interviewed, and given new coupons to recruit additional subjects. This process continues until a sufficient sample is generated, the study budget or duration is reached, or some other stopping criterion is met.

The attributes of the coupons, e.g., their number, value, call to action, expiration date, etc., can play a critical role in shaping the evolution of the RDS process. However, in standard RDS, researchers give each participant an identical coupon allocation which is typically based on convention rather than characteristics of the population under study (Goel and Salganik, 2010). This is inefficient in that coupon return rates, population coverage, and cost may be poor relative to what could be achieved with a tailored coupon allocation. Despite being widely recognized, attempts to address this issue have been

scarce. Lunagómez et al. (2018) proposed calibrating the number of coupons by simulating RDS under a parametric model and an informative prior, but they do not consider adapting coupon allocations to information accumulating during the study. McFall et al. (2021) ran a two-stage RDS study in which an initial RDS study was used to identify characteristics of participants likely to be successful recruiters, then in a subsequent RDS study, participants with these characteristics were given extra coupons (see VanOrsdale, 2023, for an extensive simulation study of such two-stage designs). Two-stage designs make use of interim data but include only a single adaptation step and require independent samples collected across two studies, which is often impractical.

We use reinforcement learning (RL, Sutton and Barto, 2018) to adapt coupon allocations over time within a single RDS study in such a way that some study objective is optimized. Example study objectives include maximizing information about disease prevalence and reaching as many individuals in the target population as possible under budget and time constraints. Our approach uses a Markov branching process (Sevast'yanov and Zubkov, 1974; Athreya et al., 2004) as a working model to guide coupon allocation during the study. However, for post-study inference, we do not assume that this model is correct. Instead, we develop a novel projection confidence set that provides valid finite sample coverage for a large class of functionals of the generative model, even when the true model is not identifiable under RDS. This is non-trivial as we are combining two procedures which are notorious for their inferential challenges: (i) RDS which is complicated by differential response probabilities, homophily, and other design effects (Goel and Salganik, 2010; Gile and Handcock, 2010; Tomas and Gile, 2011; Lu et al., 2012; Roch and Rohe, 2018; Rohe, 2019); and (ii) adaptive RL-based experimentation for which standard bootstrap or normality based inference procedures can fail to provide nominal coverage (Deshpande et al., 2018; Zhang et al., 2020; Bibaut et al., 2021; Zhan et al., 2023; Bibaut and Kallus, 2024). To address these complications, we develop valid post adaptive-experimentation inference procedures for  $M$ -estimators constructed from Markov decision processes (MDPs, Putterman, 1994), a result which is of independent interest. In simulation experiments, our proposed procedure, which we term RL-RDS,

significantly improves efficiency relative to static and two-stage RDS designs. Furthermore, the projection confidence sets deliver nominal coverage without being excessively conservative.

The contributions of this work are summarized as follows: (1) we develop the first principled framework for adaptive RDS using RL; (2) we show that a Markov branching process approximation to the RDS process is useful for guiding online adaptation; (3) we prove regret bounds for our RL algorithm under the Markov branching process model; and (4) we develop valid finite sample inference methods for adaptive-RDS without requiring identifiability.

In Section 1, we review RDS. In Section 2, we introduce our branching process approximation and RL-based adaptive coupon selection. In Section 2.1, we present regret bounds for the branching process approximation. In Section 3, we introduce our inference approach and prove that confidence regions constructed by this procedure achieve nominal coverage in finite samples and concentrate asymptotically. Lastly, in Section 4, we present a suite of simulation experiments comparing RL-RDS, static, and two-stage designs.

## 1 Setup and Notation

An RDS study recruits participants in epochs or waves. The initial epoch, which comprises individuals  $\mathcal{E}_0 \subseteq \mathbb{N}$ , is typically collected as a convenience sample. Generally, we use  $i$  to index interim study participants organized in epochs and  $v$  to index interim study participants ordered by arrival time. Each participant  $i \in \mathcal{E}_0$  receives an allocation of coupons  $\mathbf{A}_i \in \mathcal{A}$  to distribute among their social contacts. These coupons explain the study and encourage participation by offering an incentive. The data collected in the zeroth epoch are thus  $\mathcal{Z}_0 \triangleq \{(T_i, \mathbf{X}_i, Y_i, \mathbf{A}_i, C_i) : i \in \mathcal{E}_0\}$ , where  $T_i \in \mathbb{R}_+$  is the arrival time of participant  $i$ ,  $\mathbf{X}_i \in \mathcal{X} \subseteq \mathbb{R}^p$  are their covariates,  $Y_i \in \mathcal{Y} \subseteq [0, 1]$  is an outcome of interest,  $\mathbf{A}_i \in \mathcal{A}$  is their coupon allocation, and  $C_i \in \mathbb{R}_+$  is the cost of their recruitment. The set of individuals in epoch  $j \geq 1$  is denoted by  $\mathcal{E}_j \subseteq \mathbb{N} \setminus \{\cup_{j' < j} \mathcal{E}_{j'}\}$  and their

associated data are  $\mathcal{Z}_j \triangleq \{(R_i, T_i, \mathbf{X}_i, Y_i, \mathbf{A}_i, C_i) : i \in \mathcal{E}_j\}$ , where  $R_i \in \mathcal{E}_{j-1}$  denotes the recruiter of individual  $i \in \mathcal{E}_j$ . Individuals in each epoch are given coupons to recruit the next epoch until available resources are depleted or another stopping criterion is met; for concreteness, we assume that the study terminates the first time the total cost exceeds a fixed budget  $D$ . Figure (1) illustrates the evolution of an RDS sample.

We note that the covariates  $\mathbf{X}_i$  and the outcome  $Y_i$  may (though they need not) be measured simultaneously. Even if this is the case,  $Y_i$  is distinguished by its role in defining the adaptive RDS algorithm’s objective. As detailed below, we define an optimal coupon allocation strategy as one that maximizes the cumulative sum of the outcome across the sample. For example, in an RDS study of people who inject drugs, the outcome may be choosing to be tested for HIV, and participant attributes might include demographic information, PrEP use, history of STI testing, and attitudes and intentions related to risky behaviors (Risser et al., 2009). In an RDS study targeting colorectal cancer screening among non-utilizers of a healthcare system, participant attributes might include demographic information, family medical history, previous FIT or colonoscopy screening, and risk factors for colorectal cancer. The outcome might be a participant’s screening intention. Post-selection inference could focus on the distribution of the covariates, the outcome, or both (Cooks et al., 2022).

Let  $J$  denote the total number of epochs. We use an overline to represent history so that  $\bar{\mathcal{E}}_J \triangleq \bigcup_{j \leq J} \mathcal{E}_j$  are the complete sets of study participants, and  $\bar{\mathcal{Z}}_J \triangleq \bigcup_{j \leq J} \mathcal{Z}_j$  are the data that correspond to the members of  $\bar{\mathcal{E}}_J$ . For simplicity, we assume that subjects are processed sequentially; i.e., no two individuals arrive at exactly the same time. Therefore, the data may be equivalently represented as  $\mathcal{D}^\kappa \triangleq \{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v, C^v)\}_{v=1}^\kappa$ , in which individuals are indexed by their arrival times.

Under adaptive RDS, when an individual arrives in the study, they are assigned a coupon allocation based on accumulated information on all prior participants. Let  $\mathbf{H}^v \in \mathcal{H}^v$  denote the information available to researchers at the time the  $v^{\text{th}}$  subject is given their coupon allocation  $\mathbf{A}^v \in \mathcal{A}$ . For each  $\mathbf{H}^v = \mathbf{h}^v$ , let  $\psi^v(\mathbf{h}^v) \subseteq \mathcal{A}$  denote the set of allowable coupon combinations given  $\mathbf{h}^v$  (e.g., this set may be restricted to ensure budget

constraints are not violated). A deterministic allocation strategy  $\boldsymbol{\pi} = (\pi^1, \pi^2, \dots) \in \Pi$  is a sequence of functions such that  $\pi^v : \mathcal{H}^v \rightarrow \mathcal{A}$  and  $\pi^v(\mathbf{h}^v) \in \psi^v(\mathbf{h}^v)$  for all  $\mathbf{h}^v$ . The set  $\Pi$  can be restricted to exclude allocation strategies that are inherently unfair or harmful or to improve the tractability of finding the optimal allocation strategy. We define an optimal allocation strategy,  $\boldsymbol{\pi}^{\text{opt}} \in \Pi$ , as maximizing the expected value of the cumulative outcome across the RDS sample.

We formalize the optimal allocation strategy within the potential outcomes framework (Rubin, 1978; Splawa-Neyman et al., 1990). For each  $v \geq 2$ , let  $\mathbf{H}^{v*}(\bar{\mathbf{a}}^{v-1})$  denote the potential history under the sequence of coupon allocations  $\bar{\mathbf{a}}^{v-1} \triangleq (\mathbf{a}^1, \mathbf{a}^2, \dots, \mathbf{a}^{v-1})$ . For any deterministic allocation strategy  $\boldsymbol{\pi} \in \Pi$ , the potential history at time  $v \geq 2$  is

$$\mathbf{H}^{v*}(\boldsymbol{\pi}) \triangleq \sum_{\bar{\mathbf{a}}^{v-1}} \mathbf{H}^{v*}(\bar{\mathbf{a}}^{v-1}) \prod_{k=1}^{v-1} \mathbb{I}[\pi^k \{ \mathbf{H}^{k*}(\bar{\mathbf{a}}^{k-1}) \} = \mathbf{a}^k],$$

where we have defined  $\mathbf{H}^{1*}(\mathbf{a}^0) \equiv \mathbf{H}^1$  and  $\mathbb{I}(u)$  as an indicator of the event  $u$ . Similarly, for each individual  $v$ , let  $Y^{v*}(\boldsymbol{\pi})$  be the potential outcome and  $C^{v*}(\boldsymbol{\pi})$  the potential cost under  $\boldsymbol{\pi}$ . The potential number of participants under allocation strategy  $\boldsymbol{\pi}$  is thus

$$n^*(\boldsymbol{\pi}) \triangleq \inf \left\{ g : \sum_{v=1}^g C^{v*}(\boldsymbol{\pi}) > D \right\}.$$

We assume that there exists  $Q \in \mathbb{N}$  such that  $\sup_{\boldsymbol{\pi} \in \Pi} n^*(\boldsymbol{\pi}) \leq Q$  almost surely; in application, this incurs no loss in generality. Let  $\Delta^{v*}(\boldsymbol{\pi})$  be an indicator that the budget has not been exceeded when individual  $v$  enters the study under  $\boldsymbol{\pi}$ . For each  $n \in \mathbb{N}$ , define the history-value function of  $\boldsymbol{\pi}$  at  $\mathbf{h}^n$  as

$$V^n(\mathbf{h}^n, \boldsymbol{\pi}) \triangleq \mathbb{E} \left\{ \sum_{v=n}^Q \Delta^{v*}(\boldsymbol{\pi}) Y^{v*}(\boldsymbol{\pi}) \mid \mathbf{H}^n = \mathbf{h}^n \right\},$$

where this expectation is over the sampling process. An optimal allocation strategy,  $\boldsymbol{\pi}^{\text{opt}}$ , satisfies  $V^n(\mathbf{h}^n, \boldsymbol{\pi}^{\text{opt}}) \geq V^n(\mathbf{h}^n, \boldsymbol{\pi})$  for each  $n$ , all feasible  $\boldsymbol{\pi}$ , and  $\mathbf{h}^n \in \mathcal{H}^n$ .

We identify  $\boldsymbol{\pi}^{\text{opt}}$  in terms of the data-generating  $\boldsymbol{\pi}$  model by making a series of assumptions that are standard for sequential decision problems (Tsiatis et al., 2019). Define the

collection of all potential outcomes as

$$\mathcal{W} \triangleq \{ \mathbf{H}^{v*}(\bar{\mathbf{a}}^{v-1}), Y^{v*}(\bar{\mathbf{a}}^v), C^{v*}(\bar{\mathbf{a}}^v) : \mathbf{a}^v \in \psi^v \{ \mathbf{H}^{v*}(\bar{\mathbf{a}}^{v-1}) \} \}_{v \geq 1}.$$

**Assumption 1** (Strong ignorability). *For all  $v \in \mathbb{N}$ ,  $\mathcal{W} \perp \mathbf{A}^v | \mathbf{H}^v$ .*

**Assumption 2** (Consistency). *For all  $v \in \mathbb{N}$ ,  $\mathbf{H}^v = \mathbf{H}^{v*}(\bar{\mathbf{A}}^{v-1})$ ,  $Y^v = Y^{v*}(\bar{\mathbf{A}}^v)$ , and  $C^v = C^{*v}(\bar{\mathbf{A}}^v)$ ; i.e., the observed histories, outcomes, and costs are equal to their counterfactual counterparts under the coupon allocations actually given.*

**Assumption 3** (Positivity). *For all  $v \in \mathbb{N}$ ,  $\mathbf{h}^v \in \mathcal{H}^v$  and  $\mathbf{a} \in \psi^v(\mathbf{h}^v)$ , there exists  $\epsilon > 0$  such that  $\mathbb{P}(\mathbf{A}^v = \mathbf{a} | \mathbf{H}^v = \mathbf{h}^v) \geq \epsilon$ .*

Under these assumptions, one can express the likelihood for the counterfactual cumulative outcome under any coupon allocation strategy in terms of the data-generating model (Robins, 2004). However, the curvature of this likelihood may be zero or near zero in some regions of the parameter space even under simple network models (Crawford et al., 2018; Wetz et al., 2024). Consequently, estimators of parameters indexing the RDS process can be extremely volatile and online learning based on such estimators is similarly volatile, especially in small samples. To make online learning tractable, one must impose additional structure on the model. This could be done through an informative prior, though such priors are difficult to posit and have been shown to exert unacceptably large influence on the operating characteristics of resulting estimators (Wetz et al., 2024). Instead, we posit a simple branching process working model that is parsimonious and stable when fit to the RDS data as it accumulates, yet flexible enough to capture salient features of the RDS process for online learning. As noted previously, we do not require this model be correctly specified.

## 2 Reinforcement Learning for RDS

Most inferential techniques for RDS are based on simplifications of the RDS process (Heckathorn, 1997; Rohe, 2015; Crawford et al., 2018) or graph model (Gile and Hand-




Figure 1: RDS is a complex stochastic process that samples without replacement over a social network. The observed RDS sample is composed of coupon exchanges, illustrated by arrows ( $\rightarrow$ ). The unobserved connections between sample participants are represented as dashed lines ( $---$ ). The observed data resembles a branching process.

cock, 2015; Gile, 2011) because the network structure is often only weakly identifiable from the sample (Weltz et al., 2024). Consequently, these estimators can be biased and unstable. Instead of estimating the graph dynamics directly, we posit a partially controllable branching process as a working model for RDS. This model captures relationships between recruiters and recruits, and its tractable likelihood facilitates asymptotic regret guarantees and robust inference techniques. However, we emphasize that our inferential approach in Section 3 does not assume the branching process model is correct.

Let  $M^v$  be the number of recruits of participant  $v$ ,  $\underline{\mathbf{X}}^v = (\mathbf{X}_1^v, \dots, \mathbf{X}_{M^v}^v)$  the collection of their covariates,  $\underline{\mathbf{Y}}^v = (Y_1^v, \dots, Y_{M^v}^v)$  their response statuses, and  $\underline{\mathbf{T}}^v = (T_1^v, \dots, T_{M^v}^v)$  their response times. We write  $\underline{\mathbf{X}}^0, \underline{\mathbf{Y}}^0, \mathbf{A}^0$ , and  $\underline{\mathbf{T}}^0$  to denote the information collected in the initial sample, treating the first epoch as recruits of a fictional recruiter “zero.” We posit a branching process working model that factors as

$$\prod_{j=1}^J \prod_{v \in \mathcal{E}_j} f(\underline{\mathbf{Y}}^v | \underline{\mathbf{X}}^v, \underline{\mathbf{T}}^v, M^v, \mathbf{A}^v, \mathbf{H}^v) f(\underline{\mathbf{X}}^v | \underline{\mathbf{T}}^v, M^v, \mathbf{A}^v, \mathbf{H}^v) \times f(\underline{\mathbf{T}}^v | M^v, \mathbf{A}^v, \mathbf{H}^v) f(M^v | \mathbf{A}^v, \mathbf{H}^v) f(\mathbf{A}^v | \mathbf{H}^v), \quad (1)$$

where we define  $f(\mathbf{A}^0 | \mathbf{H}^0) = f(\mathbf{A}^0)$  and assume that  $C^v$  is a known function of  $(\mathbf{X}^v, \mathbf{A}^v, Y^v, \underline{\mathbf{T}}^v, \underline{\mathbf{X}}^v, \underline{\mathbf{Y}}^v)$ . According to this model, the recruits associated with recruiter  $v$  are not affected by the coupon allocation to individual  $v'$  (or recruits of  $v'$ ) if  $v' > v$ .

During the RDS process, our goal is to guide incentive allocation to maximize the expected cumulative outcome. This outcome is chosen to characterize study effectiveness, e.g., an indicator that a participant consents to disease screening. Online sequential decision making to optimize a cumulative objective fits naturally within the framework of reinforcement learning (RL; Sutton and Barto, 2018). Thompson sampling is an RL algorithm that has been shown to possess favorable theoretical properties and strong empirical performance on a wide range of problems (Chapelle and Li, 2011; Agrawal and Goyal, 2013; Gopalan et al., 2014; Gopalan and Mannor, 2015; Russo and Van Roy, 2016; Russo et al., 2018; Laber et al., 2018). However, it has never been studied as an approach to optimal allocation in RDS. This context is especially challenging due to budget constraints and complex dependence between study participants.

We consider a class of parametric models for the components of the branching process, specified in Equation (1), which we index by  $\beta = (\beta_y, \beta_t, \beta_x, \beta_m) \in \mathcal{B}$ , where  $\mathcal{B}$  is a bounded, open subset of  $\mathbb{R}^k$ . The contribution to the joint density from the  $v$ th participant is composed of the following components

$$\begin{aligned}
f(\underline{\mathbf{Y}}^v | \underline{\mathbf{X}}^v, \underline{\mathbf{T}}^v, M^v, \mathbf{A}^v, \mathbf{H}^v) &= f(\underline{\mathbf{Y}}^v | \underline{\mathbf{X}}^v, \underline{\mathbf{T}}^v, M^v, \mathbf{A}^v, \mathbf{H}^v; \beta_y), \\
f(\underline{\mathbf{X}}^v | \underline{\mathbf{T}}^v, M^v, \mathbf{A}^v, \mathbf{H}^v) &= f(\underline{\mathbf{X}}^v | \underline{\mathbf{T}}^v, M^v, \mathbf{A}^v, \mathbf{H}^v; \beta_x), \\
f(\underline{\mathbf{T}}^v | M^v, \mathbf{A}^v, \mathbf{H}^v) &= f(\underline{\mathbf{T}}^v | M^v, \mathbf{A}^v, \mathbf{H}^v; \beta_t), \\
f(M^v | \mathbf{A}^v, \mathbf{H}^v) &= f(M^v | \mathbf{A}^v, \mathbf{H}^v; \beta_m).
\end{aligned} \tag{2}$$

To approximate an optimal strategy after each participant's arrival, we use the branching process model to simulate potential futures starting from the current RDS sample. Recall that  $Q \in \mathbb{N}$  is defined such that  $\sup_{\pi} n^*(\pi) \leq Q$  almost surely. For any  $n \in \mathbb{N}$ ,  $\mathbf{h}^n \in \mathcal{H}^n$ ,  $\pi \in \Pi$ ,  $\beta \in \mathcal{B}$ , and  $B \in \mathbb{N}$ , we denote by  $\mathcal{K}^B(\mathbf{h}^n, \pi; \beta)$  a set of  $B$  trajectories of length  $Q - n$  simulated under  $\beta$ . For  $b \in B$ , the trajectory begins with the RDS information collected so far,  $\mathbf{H}_{b,\pi,\beta}^n = \mathbf{h}^n$ , and the coupons,  $\mathbf{A}_{b,\pi,\beta}^\ell$ , are assigned according to  $\pi^\ell(\mathbf{H}_{b,\pi,\beta}^\ell)$  for  $\ell = n, n + 1, \dots, Q$ , i.e.,

$$\mathcal{K}^B(\mathbf{h}^n, \pi; \beta) \triangleq \left\{ \left( Y_{b,\pi,\beta}^n, \mathbf{H}_{b,\pi,\beta}^n, \mathbf{A}_{b,\pi,\beta}^n, Y_{b,\pi,\beta}^{n+1}, \mathbf{H}_{b,\pi,\beta}^{n+1}, \mathbf{A}_{b,\pi,\beta}^{n+1}, \dots, Y_{b,\pi,\beta}^Q, \mathbf{H}_{b,\pi,\beta}^Q \right) \right\}_{b=1}^B.$$

These simulated trajectories will be used to approximate an optimal strategy,  $\boldsymbol{\pi}^{\text{opt}}$ . Under Assumptions 1-2, the history-value function reduces to

$$V^n(\mathbf{h}^n, \boldsymbol{\pi}; \boldsymbol{\beta}) = \mathbb{E}_{\boldsymbol{\beta}} \left\{ \sum_{v=n}^Q \Delta^v(\boldsymbol{\pi}) Y^v(\boldsymbol{\pi}) \mid \mathbf{H}^n = \mathbf{h}^n \right\}$$

for any  $\boldsymbol{\beta} \in \mathcal{B}$ ,  $n \in \mathbb{N}$ , history  $\mathbf{H}^n = \mathbf{h}^n$ , and  $\boldsymbol{\pi} \in \boldsymbol{\Pi}$ . Under Assumption 3, this expectation can be estimated by averaging cumulative reward over the simulated trajectories,

$$\widehat{V}_B^n(\mathbf{h}^n, \boldsymbol{\pi}; \boldsymbol{\beta}) = \frac{\sum_{b=1}^B \left( \sum_{v=n}^Q \Delta_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^v Y_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^v \right)}{B}.$$

The estimated optimal policy under  $\boldsymbol{\beta}$  upon observing  $\mathbf{H}^n = \mathbf{h}^n$  is thus  $\widehat{\boldsymbol{\pi}}_B^n(\mathbf{h}^n; \boldsymbol{\beta}) \in \arg \max_{\boldsymbol{\pi} \in \boldsymbol{\Pi}} \widehat{V}_B^n(\mathbf{h}^n, \boldsymbol{\pi}; \boldsymbol{\beta})$ .

Deciding between allocations that appear to be optimal given current estimated parameter values and those that might improve parameter estimates and thereby lead to better decisions in the future is a fundamental problem in reinforcement learning. This balance between information gain and optimization is commonly known as the exploration-exploitation trade-off in statistics and computer science (Berry and Fristedt, 1985; Sutton and Barto, 2018; Szepesvári, 2010; Slivkins et al., 2019; Lattimore and Szepesvári, 2020). We consider a variant of Thompson sampling with clipping (Zhang et al., 2020) to ensure sufficient exploration when determining coupon allocations.

At each decision point, our method samples a coupon allocation for the newest study participant approximately proportional to the probability that it is optimal. Let  $\widehat{\boldsymbol{\beta}}^n$  be an estimator of the parameters indexing the branching process model based on a sample of size  $n \in \mathbb{N}$ <sup>1</sup>, and  $\widehat{P}^n$  denote an estimator of the sampling distribution of  $\widehat{\boldsymbol{\beta}}^n$ . Given state  $\mathbf{H}^n = \mathbf{h}^n$  and  $\boldsymbol{\pi} \in \boldsymbol{\Pi}$ , let

$$\widehat{\rho}_B^n(\mathbf{h}^n, \boldsymbol{\pi}) = \int \mathbb{I} \left\{ \boldsymbol{\pi} \in \arg \max_{\boldsymbol{\pi} \in \boldsymbol{\Pi}} \widehat{V}_B^n(\mathbf{h}^n, \boldsymbol{\pi}; \boldsymbol{\beta}) \right\} d\widehat{P}^n(\boldsymbol{\beta})$$

---

<sup>1</sup>In application, a natural choice for this statistic (and the one we use for the simulations in Section 4) is the maximum likelihood estimator for the parameters indexing the branching process model; however the theory holds more generally.

be the estimated confidence that  $\boldsymbol{\pi}$  is the optimal policy. If  $\hat{\rho}_B^n(\mathbf{h}^n, \boldsymbol{\pi})$  is high, it means that  $\boldsymbol{\pi}$  maximizes the cumulative reward under the points in  $\hat{P}^n(\boldsymbol{\beta})$  that are most likely. Additionally, for  $\mathbf{a}^n \in \psi^n(\mathbf{h}^n)$ , let

$$\hat{\xi}_B^n(\mathbf{h}^n, \mathbf{a}^n) = \int_{\boldsymbol{\Pi}} \mathbb{I}\{\pi^n(\mathbf{h}^n) = \mathbf{a}^n\} d\hat{\rho}_B^n(\mathbf{h}^n, \boldsymbol{\pi})$$

be the estimated probability that  $\mathbf{a}^n$  is the optimal action. Thompson sampling with clipping at level  $\epsilon \in (0, 1)$  selects action  $\mathbf{A}^n = \mathbf{a}^n$  with probability proportional to  $\min\left[1 - \epsilon, \max\left\{\epsilon, \hat{\xi}_B^n(\mathbf{h}^n, \mathbf{a}^n)\right\}\right]$ . Lemma 11.1 in the Supplemental Materials shows that this clipping constraint satisfies the following assumption, which is needed for the theoretical developments in the next section.

**Assumption 4** (Bounded action selection probabilities). *For all  $v \in \mathbb{N}$ , assume there exists  $\rho_{\min}, \rho_{\max} \in \mathbb{R}^+$  such that  $\rho_{\min} \leq \sqrt{1/\mathbb{P}(\mathbf{A}^v|\mathbf{H}^v)} \leq \rho_{\max}$  with probability one.*

We provide justification for the proposed variant of Thompson Sampling with the branching process approximation in the following section.

## 2.1 Asymptotic Regret Bounds

In this section, we show that if the branching model is correctly specified, then Thompson Sampling with clipping achieves optimal regret. This result is both novel in its own right and justifies the use of Thompson Sampling for coupon allocation with our branching process approximation.

Recall that  $\mathcal{B}$  is a bounded, open subset of  $\mathbb{R}^k$  for  $k \in \mathbb{N}$ . We assume a parametric model indexed by fixed but unknown parameter  $\boldsymbol{\beta}^* \in \mathcal{B}$ . We construct a weighted maximum likelihood estimator of  $\boldsymbol{\beta}^*$  by extending the M-estimation approach described in Zhang et al. (2021) to general MDPs. Zhang et al. show that the maximum likelihood estimator (MLE) constructed from adaptively sampled data in a linear contextual bandit can converge to a non-normal limit when two arms have the same mean reward. A normal asymptotic limit is obtained by re-weighting the likelihood by a function of the propensity score in such a way that the asymptotic variance is stabilized (see also Deshpande et al.,

2018; Hadad et al., 2021; Zhan et al., 2023; Bibaut and Kallus, 2024; Zhan et al., 2024). In RL for RDS this re-weighting will depend on the coupon assignment distribution.

We consider an asymptotic regime based on complete generations (epochs) of study participants. For  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ , we restrict the information available before the coupon allocation to participant  $i$  to  $\bar{\mathcal{Z}}_{j-1}$ . Additionally, we do not use coupon allocation information associated with individuals in the same epoch as the current study participant. Define  $r(i)$  to be the rank of participant  $i$ 's arrival time; (i.e., if participant  $i$  were the third participant in the study, then  $r(i) = 3$ ). For  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ , we assume that the coupon allocation for participant  $i$  is based on historical information,  $\mathbf{H}_i$ , that is an element of sigma field

$$\sigma \left[ \{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v, C^v)\}_{v=1}^{r(i)} \cap (\bar{\mathcal{Z}}_{j-1} \setminus \{\mathbf{A}_i\}_{i \in \mathcal{E}_{j-1}}) \right]. \quad (3)$$

Define the complete information associated with the recruits of individual  $i$  as  $\mathbf{D}_i \triangleq \{\underline{\mathbf{Y}}_i, \underline{\mathbf{X}}_i, \underline{\mathbf{T}}_i, M_i, \mathbf{A}_i\} \in \mathcal{D}$ , and define the field associated with the first  $j$  generations as  $\mathcal{F}_j \triangleq \sigma(\bar{\mathcal{Z}}_j \setminus \{\mathbf{A}_i\}_{i \in \mathcal{E}_j})$ . In addition, under the branching process model, we assume that the individuals in generation  $\mathcal{Z}_j$  are conditionally independent given  $\mathcal{F}_{j-1}$  and  $\{\mathbf{A}_i\}_{i \in \mathcal{E}_{j-1}}$ , so that

$$\mathbb{P} \left( \{\mathbf{D}_i\}_{i \in \mathcal{E}_{j-1}} \mid \{\mathbf{A}_i\}_{i \in \mathcal{E}_{j-1}}, \mathcal{F}_{j-1} \right) = \prod_{i \in \mathcal{E}_{j-1}} \mathbb{P}(\mathbf{D}_i \mid \mathbf{A}_i, \mathcal{F}_{j-1}).$$

Define the ‘‘complete generation’’ likelihood as

$$\prod_{j=1}^J \mathbb{P}(\mathcal{Z}_j \mid \mathcal{F}_{j-1}) = \prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} \mathbb{P}(\{\underline{\mathbf{Y}}_i, \underline{\mathbf{X}}_i, \underline{\mathbf{T}}_i, M_i\} \mid \mathbf{A}_i, \mathcal{F}_{j-1}) \mathbb{P}(\mathbf{A}_i \mid \mathbf{H}_i), \quad (4)$$

where the equality follows from Equation (3) under which  $\mathbb{P}(\mathbf{A}_i \mid \mathcal{F}_{j-1}, \{\mathbf{D}_k\}_{k \in \mathcal{E}_{j-1}, k \neq i}) = \mathbb{P}(\mathbf{A}_i \mid \mathcal{F}_{j-1}) = \mathbb{P}(\mathbf{A}_i \mid \mathbf{H}_i)$ .

In our context, the estimated optimal policy need not converge to a fixed strategy because of the non-stationarity of the branching process we use to model RDS. Consequently, as anticipated by the literature on inference after adaptive sampling, the asymptotic behavior of the MLE based on (4) is difficult to characterize (Bibaut and Kallus,

2024). However, we are able to obtain a parametric rate of convergence after a suitable re-weighting of the likelihood. Let  $\hat{\pi}$  be the policy followed by the reinforcement algorithm, and  $\tilde{\pi}$  be the policy that samples from among the available coupon allocations with equal probability; i.e.,  $\mathbb{P}_{\tilde{\pi}}(\mathbf{a}|\mathbf{H}_i = \mathbf{h}_i) \triangleq 1/|\psi_i(\mathbf{h}_i)|$  if  $\mathbf{a} \in \psi_i(\mathbf{h}_i)$  and zero otherwise, and  $\mathbb{P}_{\hat{\pi}}(\mathbf{a}|\mathbf{H}_i = \mathbf{h}_i) \triangleq \mathbb{P}(\mathbf{A}_i = \mathbf{a}|\mathbf{H}_i = \mathbf{h}_i)$ . We use

$$W_i \triangleq \sqrt{\frac{\mathbb{P}_{\tilde{\pi}}(\mathbf{A}_i|\mathbf{H}_i)}{\mathbb{P}_{\hat{\pi}}(\mathbf{A}_i|\mathbf{H}_i)}},$$

as stabilizing weights in the log-likelihood. Our complete generation M-estimator maximizes the weighted log-likelihood

$$\hat{\beta}_J \triangleq \arg \max_{\beta \in \mathcal{B}} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i l_i(\beta),$$

where for  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ ,  $l_i(\beta) \triangleq l(\beta, \mathbf{D}_i) \triangleq \log \{\mathbb{P}(\mathbf{D}_i | \mathcal{F}_{j-1}; \beta)\}$ .

Below we state the technical assumptions under which we derive regret bounds for the estimated optimal policy. We verify that these assumptions hold in the working model described by Equation (6) in Section 2.2. Define the number of sample participants in generation  $j$  as  $\kappa_j \triangleq |\mathcal{E}_j|$  and the total number of individuals recruited before generation  $j$  as  $\bar{\kappa}_j \triangleq |\bar{\mathcal{E}}_{j-1}|$ . Additionally, note that the complete data-generating distribution is specified by the branching process parameter,  $\beta$ , and the reinforcement learning policy,  $\pi$ . The expectation taken with respect to this distribution is denoted  $\mathbb{E}_{\beta, \pi}$ .

**Assumption 5** (Branching asymptotics). *For all recruiters  $v \in \mathbb{N}$ , and their recruits  $j \in \{1, 2, \dots, M^v\}$ , there exists  $\alpha > 0$  such that  $T_j^v - T^v \geq \alpha$  with probability one. In addition, the number of coupons in an allocation is bounded above by  $L_*$ . Lastly, the branching process is super-critical; i.e., there exists a random variable  $\mathcal{I}$  such that for any  $\epsilon > 0$  there exists  $\delta > 0$  such that*

$$\lim_{j \rightarrow \infty} \kappa_j / m^j \rightarrow \mathcal{I} \quad \text{a.s., where } \mathbb{P}(\mathcal{I} \geq \delta) \geq 1 - \epsilon.$$

**Assumption 6** (Growing budget asymptotics). *The budget grows over time as follows. Let  $S^n$  denote the budget when individual  $n$  is recruited. For all  $n \in \mathbb{N}$ ,  $0 < S^n - \sum_{v=1}^n C^v < C^*$  with probability one for some fixed  $C^* \in \mathbb{R}^+$ .*

**Assumption 7** (Identifiability and differentiability). *The parameter indexing the branching process is identifiable. Additionally, for all  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ , the first two derivatives of  $l_i(\boldsymbol{\beta})$  with respect to any  $\boldsymbol{\beta} \in \mathcal{B}$  exist.*

**Assumption 8** (Moment conditions). *The parameter that indexes the true generative process,  $\boldsymbol{\beta}^* \in \mathcal{B}$ , is in the interior of  $\mathcal{B}$ . For all  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ , the first two moments of  $l_i(\boldsymbol{\beta}^*)$ ,  $\dot{l}_i(\boldsymbol{\beta}^*)$ , and  $\ddot{l}_i(\boldsymbol{\beta}^*)$  conditional on  $\mathcal{F}_{j-1}$  are bounded almost surely.*

**Assumption 9** (Lipschitz). *There exists a real-valued function  $g : \mathcal{D} \rightarrow \mathbb{R}$  such that for all  $j \in \mathbb{N}$ ,  $i \in \mathcal{E}_{j-1}$ , and  $\boldsymbol{\beta}, \boldsymbol{\beta}' \in \mathcal{B}$ ,*

$$|l_i(\boldsymbol{\beta}) - l_i(\boldsymbol{\beta}')| \leq g(\mathbf{D}_i) \|\boldsymbol{\beta} - \boldsymbol{\beta}'\|_2,$$

where  $\mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\}$  is bounded almost surely.

**Assumption 10** (Well-separated maximizer). *For any  $\epsilon > 0$ , there exists  $J_0 \in \mathbb{N}$  and  $\delta > 0$  such that for all  $J \geq J_0$ ,*

$$\inf_{\boldsymbol{\beta} \in \mathcal{B}: \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\|_2 > \epsilon} \left[ \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\tilde{\boldsymbol{\pi}}, \boldsymbol{\beta}^*} \{l_i(\boldsymbol{\beta}^*) - l_i(\boldsymbol{\beta}) | \mathcal{F}_{j-1}\} \right] \geq \delta \text{ a.s.}$$

Assumptions 5 and 6 specify the asymptotic regime for the branching process. Assumption 5 implies that  $J \rightarrow \infty$  and that the generation sizes are consistent with Galton-Watson processes (Athreya et al., 2004). Assumption 6 states that the budget grows in such a way that the remaining budget is always bounded; this avoids trivial solutions in which maximal resources are allocated at each time point. Assumptions 7-10 ensure that the log-likelihood is well-behaved. In Assumptions 7 and 8, we assume that the log-likelihood is identifiable, two times differentiable, and its components (and their derivatives) have finite second moments. Assumption 9 limits the complexity of the

log-likelihood function so that the weighted log-likelihood converges uniformly. Assumption 10 requires that  $\beta^*$  be a “well-separated” point of maximum and is a standard assumption for consistency; e.g., Theorem 5.7 of Van der Vaart (2000). Note that Assumptions 9 and 10 are unnecessary for consistency if the log-likelihood is concave.

Under Assumption 7, for any policy  $\pi \in \Pi$ ,  $\beta \in \mathcal{B}$ ,  $j \in \mathbb{N}$ , and  $i \in \mathcal{E}_{j-1}$ , it follows that  $\mathbb{E}_{\beta, \pi} \{ \dot{l}_i(\beta) \mid \mathbf{A}_i, \mathbf{H}_i \} = 0$ . Consequently,  $\sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\beta)$  is a martingale difference with respect to the filtration  $\{\mathcal{F}_j\}_{j \geq 1}$ ,

$$\mathbb{E}_{\beta, \pi} \left\{ \sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\beta) \mid \mathcal{F}_{j-1} \right\} = \mathbb{E}_{\beta, \pi} \left[ \sum_{i \in \mathcal{E}_{j-1}} W_i \mathbb{E}_{\beta, \pi} \{ \dot{l}_i(\beta) \mid \mathbf{A}_i, \mathcal{F}_{j-1} \} \mid \mathcal{F}_{j-1} \right] = 0.$$

We define the variance of the weighted score function conditioned on this filtration,

$$\eta_J \triangleq \sum_{j=1}^J \mathbb{E}_{\beta^*, \hat{\pi}} \left\{ \sum_{i \in \mathcal{E}_{j-1}} W_i^2 \dot{l}_i(\beta^*) \dot{l}_i(\beta^*)^\top \mid \mathcal{F}_{j-1} \right\} = \sum_{j=1}^J \mathbb{E}_{\beta^*, \hat{\pi}} \left\{ \sum_{i \in \mathcal{E}_{j-1}} \dot{l}_i(\beta^*) \dot{l}_i(\beta^*)^\top \mid \mathcal{F}_{j-1} \right\},$$

where the equality follows from

$$\begin{aligned} \mathbb{E}_{\beta^*, \hat{\pi}} \left\{ W_i^2 \dot{l}_i(\beta^*) \dot{l}_i(\beta^*)^\top \mid \mathcal{F}_{j-1} \right\} &= \int W_i^2 \dot{l}_i(\beta^*) \dot{l}_i(\beta^*)^\top f(\mathbf{D}_i \mid \mathbf{A}_i, \mathcal{F}_{j-1}; \beta^*) \mathbb{P}_{\hat{\pi}}(\mathbf{A}_i \mid \mathcal{F}_{j-1}) d\nu \\ &= \int \frac{\mathbb{P}_{\hat{\pi}}(\mathbf{A}_i \mid \mathbf{H}_i)}{\mathbb{P}_{\hat{\pi}}(\mathbf{A}_i \mid \mathbf{H}_i)} \dot{l}_i(\beta^*) \dot{l}_i(\beta^*)^\top f(\mathbf{D}_i \mid \mathbf{A}_i, \mathcal{F}_{j-1}; \beta^*) \mathbb{P}_{\hat{\pi}}(\mathbf{A}_i \mid \mathbf{H}_i) d\nu \\ &= \mathbb{E}_{\beta^*, \hat{\pi}} \left\{ \dot{l}_i(\beta^*) \dot{l}_i(\beta^*)^\top \mid \mathcal{F}_{j-1} \right\}. \end{aligned}$$

We see that re-weighting the outer product of the score function removes the dependence of the estimating equation’s conditional variance on the RL algorithm  $\hat{\pi}$ .

While  $\eta_J$  is a constant in the setting considered by Zhang et al. (2021), here it is a random variable. Consequently, characterizing the asymptotic behavior of  $\hat{\beta}_J$  requires new theory for martingale estimating functions constructed from controllable branching processes. For matrix  $A$ , let  $\sigma_{\min}(A)$  be its minimum eigenvalue. In addition, let  $I_k$  be the  $k$  dimensional identity matrix. To extend asymptotic theory for  $M$ -estimators collected under a contextual bandit to the more general setting of an MDP, we make use of the following standard assumptions.

**Assumption 11** (Martingale stabilizing variance). *As  $J \rightarrow \infty$ , there exists a sequence of constant (i.e., not random) positive definite matrices  $\{\Sigma_J\}_{J \geq 1}$  such that*

$$\Sigma_J^{-1/2} \eta_J \Sigma_J^{-1/2} \xrightarrow{p} U,$$

where  $U$  is a (random) positive definite matrix and  $U \succ 0$  with probability one.

**Assumption 12** (Information accumulation). *For some  $\delta > 0$ ,  $\liminf_{J \rightarrow \infty} \sigma_{\min} \{\eta_J / \bar{\kappa}_J\} \geq \delta$  with probability one.*

**Assumption 13** (Equicontinuity). *There exists an  $\epsilon_i > 0$  and a function  $f : \mathcal{D} \rightarrow \mathbb{R}$  such that for all  $j \in \mathbb{N}$ ,  $i \in \mathcal{E}_{j-1}$  and  $0 < \epsilon \leq \epsilon_i$ , there exists  $\delta_\epsilon$  such that*

$$\sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\|_2 \leq \delta_\epsilon} \|\ddot{l}_i(\beta) - \ddot{l}_i(\beta^*)\|_2 \leq \epsilon f(\mathbf{D}_i) \quad \text{a.s.},$$

and  $\mathbb{E}_{\beta^*, \bar{\pi}} \{f(\mathbf{D}_i) | \mathcal{F}_{j-1}\}$  is bounded almost surely.

Assumption 11 requires that the conditional variance of the log-likelihood components converges. This assumption is the crux of our extension from the contextual bandit framework of Zhang et al. (2021) to more general MDPs. It is also standard in asymptotic theory for martingales, e.g., Theorem 3.2 in Hall and Heyde (2014). Assumption 12 requires that information accumulates over time. Assumption 13 ensures the equicontinuity of the empirical Fisher information.

**Theorem 2.1.** *For  $\delta > 0$ , define the event  $E_{\mathcal{J}} = \{\mathcal{J} > \delta\}$ , where  $\mathcal{J}$  is defined in Assumption 5. Under Assumptions 1-13, as  $J \rightarrow \infty$ ,*

$$\widehat{\beta}_J - \beta^* = O_p(1/\sqrt{\bar{\kappa}_J}) \tag{5}$$

on event  $E_{\mathcal{J}}$ .

We consider the cumulative reward of policies implemented over study participants as they arrive at the study, represented by data  $\mathcal{D}^\kappa \triangleq \{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v, C^v)\}_{v=1}^\kappa$  for  $\kappa \in \mathbb{N}$ . Consequently, we require a Lipschitz condition for the log-likelihood of this

arrival process to establish asymptotic regret bounds. For  $v \in \mathbb{N}$ , define the log-likelihood components of the arrival process as

$$q^v(\boldsymbol{\beta}) \triangleq q\{\boldsymbol{\beta}, (R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v)\} \triangleq \log [\mathbb{P}\{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v) | \mathbf{H}^{v-1}, \mathbf{A}^{v-1}, \boldsymbol{\beta}\}].$$

**Assumption 14** (Lipschitz 2). *There exists a real-valued function  $e$  such that for all  $v \in \mathbb{N}$ , and  $\boldsymbol{\beta}, \boldsymbol{\beta}' \in \mathcal{B}$ ,*

$$|q^v(\boldsymbol{\beta}) - q^v(\boldsymbol{\beta}')| \leq e\{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v)\} \|\boldsymbol{\beta} - \boldsymbol{\beta}'\|_2,$$

where for any  $\boldsymbol{\pi} \in \Pi$ ,  $\mathbb{E}_{\boldsymbol{\beta}^*, \boldsymbol{\pi}} [e\{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v)\}^2]$  is bounded.

Define  $t_{\min} \geq 0$  as the minimum processing time for an individual and  $t_{\max} > 0$  as the coupon expiration time so that  $t_{\min} \leq T_j^v - T^v \leq t_{\max}$  for any  $v \in \mathbb{N}$  and  $j \in \{1, 2, \dots, M^v\}$ . For a sample of size  $\kappa$ , define  $J_\kappa$  as the last complete epoch induced by the expiration of coupons,  $J_\kappa \triangleq \max\{j : \forall i \in \mathcal{E}_j, T_i + t_{\max} < T^\kappa\}$ . Additionally, define  $n_J \triangleq \min\{\kappa : J_\kappa = J\}$  as the index of the first individual recruited after all the coupons associated with members of epoch  $J$  have expired.

**Theorem 2.2.** *For any  $\mathbf{h}^{n_J} \in \mathcal{H}^{n_J}$ , define  $\hat{\boldsymbol{\pi}}_J \in \arg \max_{\boldsymbol{\pi} \in \Pi} V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \hat{\boldsymbol{\beta}}_J)$ . For  $\delta > 0$ , define the event  $E_{\mathcal{J}} = \{\mathcal{J} > \delta\}$ , where  $\mathcal{J}$  is defined in Assumption 5. Under Assumptions 1-14 and for a fixed  $\mathbf{h}^{n_J} \in \mathcal{H}^{n_J}$ , it follows that*

$$V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}^{\text{opt}}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \hat{\boldsymbol{\pi}}_J; \boldsymbol{\beta}^*) = O_p(1/\sqrt{\kappa_J})$$

as  $J \rightarrow \infty$  on event  $E_{\mathcal{J}}$ .

The proofs of Theorems 2.1 and 2.2 are in the Supplemental Materials. Lemma 10.1 in the Supplementary Materials ensures that  $\lim_{\kappa \rightarrow \infty} J_\kappa \rightarrow \infty$  a.s. under Assumption 5, implying that inference based on the last complete epoch will achieve the asymptotic guaranties of Theorems 2.1 and 2.2. Note that  $\lim_{\kappa \rightarrow \infty} J_\kappa \rightarrow \infty$  a.s. implies  $n_J < \infty$  for all  $J \in \mathbb{N}$ .

## 2.2 A Branching Process Example

In this section, we provide an example of a working branching process model that might be used in the context of RL-RDS and illustrate how the assumptions used in Theorems (2.1) and (2.2) can be verified for this model. Define  $T_{i,l}$ ,  $\mathbf{X}_{i,l}$ ,  $Y_{i,l}$ , and  $A_{i,l}$  for  $l = 1, \dots, M_i$  as the arrival times, covariates, rewards, and coupon types associated with the potential recruits of recruiter  $i$  respectively. Define  $\mathbb{A} = \{-1, 1\}$  as the set of possible coupon types (these might reflect different calls to action for example), and  $\mathbf{Z}_{i,l} = (1, \mathbf{X}_{i,l}, \mathbf{X}_{i,l} \mathbb{I}(A_{i,l} = -1))$ .

We consider a working model of the form:

$$\begin{aligned} \mathbb{P}(M_i = m_i | \mathbf{H}_i, \mathbf{A}_i) &= \frac{\lambda^{m_i}/m_i!}{\sum_{\ell=0}^{|\mathbf{A}_i|} (\lambda^\ell/\ell!)}, m_i = 0, \dots, |\mathbf{A}_i|, \\ T_{i,l} - T_i | \mathbf{H}_i, \mathbf{A}_i, M_i &\sim \text{Truncated Exponential}(\zeta, t_{\min}, t_{\max}), l = 1, \dots, M_i, \\ \mathbf{X}_{i,l} | \mathbf{H}_i, U_{i,l}, \mathbf{A}_i, M_i &\sim \text{Normal}(\boldsymbol{\phi}_a + G_a \mathbf{X}_i, \Sigma_a), l = 1, \dots, M_i, a = A_{i,l}, \\ Y_{i,l} | \mathbf{H}_i, \mathbf{X}_i, U_{i,l}, \mathbf{A}_i, M_i &\sim \text{Bernoulli} \left\{ \frac{1}{1 + \exp(-\mathbf{Z}_{i,l}^\top \boldsymbol{\beta}_y)} \right\}, l = 1, \dots, M_i, \end{aligned} \quad (6)$$

where  $\lambda, \zeta \in \mathbb{R}$ ,  $\{G_a\}_{a \in \mathbb{A}} \in \mathbb{R}^{p \times p}$ ,  $\{\boldsymbol{\phi}_a\}_{a \in \mathbb{A}} \in \mathbb{R}^{p \times 1}$ ,  $\boldsymbol{\beta}_y \in \mathbb{R}^\ell$ , and  $\{\Sigma_a\}_{a \in \mathbb{A}} \in \mathbb{R}^{p \times p}$ . In the context of a randomized experiment, Assumption 1 (strong ignorability), Assumption 3 (positivity), and Assumption 6 (budget growth) can be ensured to hold by design. Thus, we assume them as a matter of course. Assumption 2 (consistency) is also assumed (this is standard as consistency is sometimes considered as an axiom rather than an assumption). We evaluate the assumptions that are sufficient for the convergence and regret results to hold for our working model in the Supplementary Materials. We verify them under the model conditions:

(C1) the same number of coupons, defined as  $L \in \mathbb{N}$ , are given to each participant,  $t_{\min} > 0$ , and  $\lambda$  is such that

$$\lambda \in \left\{ \lambda : \sum_{m_i=1}^L m_i \frac{\lambda^{m_i}/m_i!}{\sum_{\ell=0}^L (\lambda^\ell/\ell!)} > 1 \right\};$$

(C2) for each  $i \in \mathbb{N}$ ,  $A_{i,l}$  is constant across  $l \in \{1, 2, \dots, M_i\}$ ;

(C3) the set of coupon allocations available for each participant is constant throughout the study;

(C4)  $\mathcal{X}$  is compact;

(C5)  $\mathcal{B} \subseteq \mathbb{R}^q$  is convex and compact, and  $\beta^* \in \mathcal{B}$  is an interior point of  $\mathcal{B}$ ;

(C6)  $\frac{1}{|\mathbb{A}|} \sum_{a \in \mathbb{A}} \log \|G_a\|_2 < 0$ , recalling that  $\|\cdot\|$  is the spectral norm.

Under Conditions (C1)-(C6), we show that there exists a random variable  $\mathcal{J}$  s.t.

$$\mathcal{J} = \lim_{j \rightarrow \infty} m^{-j} \kappa_j \quad \text{a.s.}, \quad \mathbb{P}(\mathcal{J} \geq \delta) \geq \epsilon. \quad (7)$$

where  $\mathbb{E}(\mathcal{J}) = 1$  (Athreya et al., 2004). Consequently, event  $E_{\mathcal{J}}$  is well defined. Condition (C6) ensures that the auto-regressive covariate process is not explosive.

**Theorem 2.3.** *Assume that Conditions (C1)-(C6) hold as well as Assumptions 1-3 and 6. Then, under the working model given above, the conclusions of Theorems (2.1) and (2.2) hold.*

The proof of the preceding result involves deriving a new weak law of large numbers for Galton-Watson branching processes to verify Assumption 11. Theorem 2.3 implies that for the posited model, policy-search with Thompson sampling attains favorable regret bounds as  $J \rightarrow \infty$ . Because Equation 7 and Condition (C1) satisfy Assumption 5, we know that Lemma 10.1 is also satisfied under the assumptions made in Theorem 2.3. In the next section, we discuss inference given an RL-RDS sample without assuming an underlying branching model.

### 3 Inference for RL-RDS

In this section, we derive valid inference for functionals of the population network model. To account for the underlying social network and the idiosyncrasies of RDS, we do not

assume that data-generating model is a branching process. Instead, we consider a dynamic network model indexed by  $\theta^* \in \Theta$  for which we derive asymptotic confidence sets. Projections of these regions are then used to conduct inference for functionals of the data-generating model, e.g., disease prevalence, rate of risky behavior, attitudes toward public health services, etc.

We recall that the data-generating model (i.e.,  $\theta^*$ ) need not be identifiable under an RDS sampling scheme. Nevertheless, it is still possible to obtain valid confidence intervals by inverting a test (e.g., see Robins, 2004; Laber and Murphy, 2011). Our test is based on the likelihood ratio for the covariate distribution in the branching process working model. We reiterate that this is only used to construct a test and that we are not assuming that this model is correct. Let  $\ell_\kappa^\theta(\beta_{\mathbf{x}})$  be the log-likelihood<sup>2</sup> of the branching process covariate model for a collection of  $\kappa$  subjects sampled under the RDS process when  $\theta \in \Theta$  is the true parameter. Define the MLE of the working model at  $\theta$  as

$$\widehat{\beta}_{\mathbf{x}}^\kappa(\theta) \in \arg \max_{\beta_{\mathbf{x}} \in \mathcal{B}} \ell_\kappa^\theta(\beta_{\mathbf{x}}).$$

At the true parameter,  $\theta^*$ , we let  $\ell_\kappa \triangleq \ell_\kappa^{\theta^*}$  and  $\widehat{\beta}_{\mathbf{x}}^\kappa \triangleq \widehat{\beta}_{\mathbf{x}}^\kappa(\theta^*)$ .

Recall that  $\mathcal{B}$  is a bounded, open subset of  $\mathbb{R}^k$ . Let  $s : \Theta \rightarrow \mathcal{B}$  be a fixed function. Our confidence region is based on the distribution of the proximity of  $\widehat{\beta}_{\mathbf{x}}^\kappa(\theta)$  to  $s(\theta)$ . In our application, we choose  $s(\theta)$  to be an asymptotic limit of  $\widehat{\beta}_{\mathbf{x}}^\kappa(\theta)$  though other choices are possible and may be more appropriate in other contexts; hence, we let  $s$  be arbitrary. Given  $s$ , define the sampling distribution of the log-likelihood ratio statistic at  $\theta$  as

$$-2 \left[ \ell_\kappa^\theta \{s(\theta)\} - \ell_\kappa^\theta \left\{ \widehat{\beta}_{\mathbf{x}}^\kappa(\theta) \right\} \right] \sim P_\kappa^\theta(s),$$

and the  $1 - \alpha$  quantile of  $P_\kappa^\theta(s)$  as  $\gamma_{1-\alpha, \kappa}^\theta(s)$ . A confidence region for  $\theta^*$  is

$$\Gamma_{1-\alpha, \kappa}(s) = \left\{ \theta : -2 \left[ \ell_\kappa \{s(\theta)\} - \ell_\kappa \left\{ \widehat{\beta}_{\mathbf{x}}^\kappa \right\} \right] \leq \gamma_{1-\alpha, \kappa}^\theta(s) \right\}. \quad (8)$$

---

<sup>2</sup>Note that in this inference procedure, we use the complete sample  $v \in \{1, 2, \dots, \kappa\}$  because we are no longer restricted to epoch structured data by the branching process.

For  $\boldsymbol{\theta} \in \Theta$ , we sample from  $P_\kappa^\boldsymbol{\theta}(s)$  by simulating a dataset of size  $\kappa$  under RDS at  $\boldsymbol{\theta}$  and calculate  $-2 \left[ \ell_\kappa^\boldsymbol{\theta} \{s(\boldsymbol{\theta})\} - \ell_\kappa^\boldsymbol{\theta} \left\{ \widehat{\boldsymbol{\beta}}_\mathbf{x}^\kappa(\boldsymbol{\theta}) \right\} \right]$  given this data. We can approximate  $\gamma_{1-\alpha, \kappa}^\boldsymbol{\theta}(s)$  to arbitrary precision by generating a large number of draws from  $P_\kappa^\boldsymbol{\theta}(s)$ . That  $\Gamma_{1-\alpha, \kappa}(s)$  achieves nominal coverage is easily verified as, by construction, we have

$$\mathbb{P} \{ \boldsymbol{\theta}^* \in \Gamma_{1-\alpha, \kappa}(s) \} = \mathbb{P} \left\{ -2 \left[ \ell_\kappa \{s(\boldsymbol{\theta}^*)\} - \ell_\kappa \left\{ \widehat{\boldsymbol{\beta}}_\mathbf{x}^\kappa \right\} \right] \leq \gamma_{1-\alpha, \kappa}^{\boldsymbol{\theta}^*}(s) \right\} = 1 - \alpha.$$

This equality holds in finite samples regardless of the function  $s$ .

The preceding result shows that  $\Gamma_{1-\alpha, \kappa}(s)$  achieves nominal coverage, we now consider another key attribute of this interval, its asymptotic concentration. To do this, we make the following additional assumptions. First, we assume that the average log-likelihood for each  $\boldsymbol{\beta}_\mathbf{x} \in \mathcal{B}$  converges to a finite limit.

**Assumption 15** (Pointwise convergence). *For any  $\boldsymbol{\theta} \in \Theta$  and for each  $\boldsymbol{\beta}_\mathbf{x} \in \mathcal{B}$ ,  $\ell_\kappa^\boldsymbol{\theta}(\boldsymbol{\beta}_\mathbf{x})/\kappa$  converges almost surely to a finite limit  $\bar{\ell}^\boldsymbol{\theta}(\boldsymbol{\beta}_\mathbf{x})$  as  $\kappa \rightarrow \infty$ . Define  $\bar{\boldsymbol{\beta}}_\mathbf{x}(\boldsymbol{\theta}) = \arg \max_{\boldsymbol{\beta}_\mathbf{x} \in \mathcal{B}} \bar{\ell}^\boldsymbol{\theta}(\boldsymbol{\beta}_\mathbf{x})$ , then  $\bar{\boldsymbol{\beta}}_\mathbf{x}(\boldsymbol{\theta})$  is finite almost surely.*

Under this assumption, the asymptotic limit of  $\widehat{\boldsymbol{\beta}}_\mathbf{x}^\kappa(\boldsymbol{\theta})$  is well-defined. We suggest using this limit,  $\bar{\boldsymbol{\beta}}_\mathbf{x}(\boldsymbol{\theta})$ , as the function  $s(\boldsymbol{\theta})$ . Let  $\Gamma_{1-\alpha, \kappa}$  denote the confidence set for  $\boldsymbol{\theta}^*$  under this choice. We also assume that the average log-likelihood over a compact set stays strictly concave asymptotically.

**Assumption 16.** *For any  $\boldsymbol{\theta} \in \Theta$ , the log-likelihood of the working model,  $\ell_\kappa^\boldsymbol{\theta}$ , is concave for every  $\kappa \in \mathbb{N}$ . Additionally,  $\ell_\kappa^\boldsymbol{\theta}(\boldsymbol{\beta}_\mathbf{x})/\kappa = O_p(1)$  as  $\kappa \rightarrow \infty$ . Lastly, for any  $\boldsymbol{\theta} \in \Theta$ , and any compact set  $\mathcal{B}' \subseteq \mathcal{B}$ , there exists  $\delta_{\mathcal{B}', \boldsymbol{\theta}} > 0$  such that*

$$\lim_{\kappa \rightarrow \infty} \sup_{\boldsymbol{\beta}_\mathbf{x} \in \mathcal{B}'} \sigma_{\min} \left\{ -\ell_\kappa^\boldsymbol{\theta}(\boldsymbol{\beta}_\mathbf{x})/\kappa \right\} \geq \delta_{\mathcal{B}', \boldsymbol{\theta}} \quad \text{a.s.}$$

Define the equivalence class

$$\Theta^* = \left\{ \boldsymbol{\theta} \in \Theta : \bar{\boldsymbol{\beta}}_\mathbf{x}(\boldsymbol{\theta}) = \bar{\boldsymbol{\beta}}_\mathbf{x}(\boldsymbol{\theta}^*) \right\}$$

as the set of network models that have the same asymptotic branching process parameter

as the true network model. If there is a unique  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$  for each  $\boldsymbol{\theta} \in \Theta$ , then  $\Theta^* = \{\boldsymbol{\theta}^*\}$ . Otherwise, the size of  $\Theta^*$  can be thought of as the “price” our inference approach pays for using a working model to perform inference on  $\boldsymbol{\theta}^*$  instead of the true data-generating model. In Theorem 3.1, we show that Assumptions 15-16 with Conditions (C2) and (C4) imply that  $\Gamma_{1-\alpha,\kappa}$  concentrates around  $\Theta^*$  as  $\kappa \rightarrow \infty$ .

**Theorem 3.1.** *Under Assumptions 1-3, 15-16,  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$  is unique and*

$$\widehat{\beta}_{\mathbf{x}}^{\kappa}(\boldsymbol{\theta}) \xrightarrow{p} \bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$$

as  $\kappa \rightarrow \infty$ . Additionally, for any  $\boldsymbol{\theta} \notin \Theta^*$ ,

$$\lim_{\kappa \rightarrow \infty} \mathbb{P}(\boldsymbol{\theta} \in \Gamma_{1-\alpha,\kappa}) \rightarrow 0.$$

The proof of Theorem 3.1 is in the Supplemental Materials. If we use the branching process in Equation 6 as the working model, we can substitute Assumption 17, Condition (2), and Condition (4) for Assumption 16 in Theorem 3.1.

**Assumption 17.** *For any  $n \in \mathbb{N}$ , define fields  $\mathcal{F}^n = \sigma[\{R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v, C^v\}_{v=1}^n]$ . For any  $\boldsymbol{\theta} \in \Theta$ , there exists a positive definite  $\Delta$ ,  $\delta > 0$ , and  $N \in \mathbb{N}$  such that  $\forall \mathbf{a} \in \mathcal{A}$  and  $n \geq N$ ,*

$$\mathbb{E}_{\boldsymbol{\theta}} \{ \mathbf{X}^{R^n} \mathbf{X}^{T^n} \mathbb{I}(\mathbf{A}^{R^n} = \mathbf{a}) \mid \mathcal{F}^{n-1} \} \succeq \Delta, \quad \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{A}^{R^n} = \mathbf{a} \mid \mathcal{F}^{n-1}) \geq \delta.$$

We note that setting  $s(\boldsymbol{\theta}) = \bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$  necessitates approximating  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$  for an arbitrary  $\boldsymbol{\theta} \in \Theta$ . Algorithm 1 in the Supplemental Materials describes our method of approximation as well as the full procedure for constructing  $\Gamma_{1-\alpha,\kappa}$  when  $s(\boldsymbol{\theta}) = \bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$ .

## 4 RL-RDS Simulations

We conducted a series of simulation experiments to evaluate the operating characteristics of RL-RDS. To allow comparisons with the two-stage procedure proposed by McFall

et al. (2021) (see also VanOrsdale (2023)), we consider the setting in which the goal is to recruit the largest subset of people in a hidden population with a given binary trait, e.g., undiagnosed HIV. The outcome is thus an indicator of this trait. We estimate the optimal policy,  $\boldsymbol{\pi}^{\text{opt}}$ , using RL-RDS. After collecting the sample, we use the confidence set derived in Section 3 to construct projection intervals for parameters indexing the target population’s network and covariate models.

The hidden population network is generated as follows. Given population size  $N$ , each population member  $v \in \{1, 2, \dots, N\}$  is assigned attributes  $\mathbf{X}^v \stackrel{\text{iid}}{\sim} \text{Normal}(\boldsymbol{\mu}, \Sigma)$ , where  $\boldsymbol{\mu} \in \mathbb{R}^p$ , and  $\Sigma \in \mathbb{R}^{p \times p}$ . Define  $\mathcal{A}_N \in \{0, 1\}^{N \times N}$  to be the adjacency matrix representing links between the population members. We construct the adjacency matrix using a latent distance network model, which is known to be flexible and projective (Spencer and Shalizi, 2017). This model specifies the probability of a connection between individuals  $i$  and  $j$  as

$$\mathbb{P} \left\{ (\mathcal{A}_N)_{i,j} = 1 \mid \mathbf{X}^i = \mathbf{x}^i, \mathbf{X}^j = \mathbf{x}^j \right\} = 1 / \left[ 1 + \exp \left\{ - (\rho_0 - \rho_1 \|\mathbf{x}^i - \mathbf{x}^j\|_2) \right\} \right],$$

where  $\rho_0, \rho_1 \in \mathbb{R}$ .

To characterize the evolution of RDS, for each  $v \in \{1, 2, \dots, N\}$ , define

$$\mathcal{N}^v = \{j \in \{1, 2, \dots, N\} \setminus \{v\} : (\mathcal{A}_N)_{v,j} = 1\}$$

as the neighborhood of  $v$ . Label the set of individual  $v$ ’s potential recruits as  $\mathcal{M}_1^v$ ; i.e., the un-recruited members of  $\mathcal{N}^v$  when  $v$  is recruited. Furthermore, define  $\mathcal{M}_2^v \subseteq \mathcal{M}_1^v$  as the coupon-constrained set of potential recruits for participant  $v$ . If  $|\mathcal{M}_1^v| \leq |\mathbf{A}^v|$ , then  $\mathcal{M}_2^v \equiv \mathcal{M}_1^v$ . If  $|\mathcal{M}_1^v| > |\mathbf{A}^v|$ , then sample  $|\mathbf{A}^v|$  recruits from  $\mathcal{M}_1^v$  according to probabilities  $\mathbf{p}^v = u(\mathbf{A}^v, \underline{\mathbf{X}}^v)$ , where  $u : \mathcal{A} \times \mathcal{X}^{|\mathcal{M}_1^v|} \rightarrow [0, 1]^{|\mathcal{M}_1^v|}$ , and label them  $\mathcal{M}_2^v$ . This allows for the neighbor selection process to depend on the characteristics of the neighbors and the coupon allocation type. When individual  $v$  is recruited, we assign arrival times,  $T_j^v$ , to the edges between recruiter,  $v$ , and hidden population members,  $j \in \mathcal{M}_2^v$ , such that  $T_j^v - T^v \sim \text{Truncated Exponential}(\zeta, t_{\min}, t_{\max})$ . At time  $T_j^v$ , participant  $j$  enters the

study (if they have not been previously recruited) and edge  $\{v, j\}$  is recorded. The reward is independently drawn for recruit  $j$  of participant  $v$  according to  $Y_j^v | \mathbf{X}_j^v, A_j^v \sim \text{Bernoulli} [1 / \{1 + \exp(-\mathbf{Z}_j^{v\top} \boldsymbol{\beta}_y)\}]$ . We note that the complete data-generating process is parameterized by  $\boldsymbol{\theta} \triangleq (\rho_0, \rho_1, \boldsymbol{\mu}, \Sigma, \boldsymbol{\beta}_y, \zeta, u, t_{\min}, t_{\max})$ .

## 4.1 Policies

We evaluate the performance of RL-RDS against a suite of alternative strategies. At each step, the researcher can choose from a finite selection of coupon types. The fixed allocation policies (i.e., those that give the same coupon allocation type to all participants) represent the current standard in RDS. The train-and-implement policy (aka, explore-than-exploit) mimics the procedure used by McFall et al. (2021), which determines an incentive strategy using a pilot study. We describe each strategy below.

1. **Fixed** offers a fixed coupon allocation  $\mathbf{a} \in \mathcal{A}$  to every study participant. If  $\mathbf{a} \notin \psi^v(\mathbf{h}^v)$ , then pick a random coupon allocation from  $\psi^v(\mathbf{h}^v)$  to give to the  $v^{\text{th}}$  study participant.
2. **Random** offers a random element of  $\psi^v(\mathbf{h}^v)$  to the  $v^{\text{th}}$  study participant.
3. **Train-and-Implement** uses half of the budget for a “pilot study,” in which the the Random policy is used to assign coupon allocations. It then conducts policy search using the pilot study data to estimate the branching process working model. This estimated policy (without updating) is then used to determine coupon allocations for the remainder of the budget.
4. **RL-RDS** uses the Random policy to assign coupon allocations to participants in a short “warm-up” period (50 participants in the simulations below). Then, it performs policy search with Thompson sampling for the remainder of the budget.

To conduct RL-RDS, we use the following space of policies,  $\Pi$ . Define  $\alpha_0 \in \mathbb{R}$ ,  $\boldsymbol{\alpha}_1 \in \mathbb{R}^p$ , and  $\boldsymbol{\alpha} = (\alpha_0, \boldsymbol{\alpha}_1)$ . For  $n \in \mathbb{N}$  and state  $\mathbf{h}^n \in \mathcal{H}^n$ , we consider policies of the

form  $\boldsymbol{\pi}(\mathbf{h}^n) = \{\pi^n(\mathbf{h}^n), \pi^{n+1}(\mathbf{h}^{n+1}), \dots\}$  such that for  $v \geq n$ ,

$$\pi^v(\mathbf{h}^v) = \pi^v(\mathbf{h}^v, \boldsymbol{\alpha}) = \pi^v(\mathbf{x}^v, \boldsymbol{\alpha}) = g^v \left[ \frac{1}{1 + \exp\{-\alpha_0 + \mathbf{x}^{v\top} \boldsymbol{\alpha}_1\}} \right], \quad (9)$$

where  $g^v : (0, 1) \rightarrow \phi^v(\mathbf{h}^v)$  maps a continuous score (dependent on the participant's covariates) to a coupon allocation,  $\mathbf{a}^v \in \psi^v(\mathbf{h}^v)$ . We first draw  $\widehat{\boldsymbol{\beta}}^n$  from the generalized bootstrap estimator of the sampling distribution of the MLE (Chatterjee and Bose, 2005). We then generate synthetic data sets,

$$\mathcal{K}^B(\mathbf{h}^n, \boldsymbol{\pi}; \boldsymbol{\beta}) \triangleq \left\{ \left( Y_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^n, \mathbf{H}_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^n, \mathbf{A}_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^n, Y_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^{n+1}, \mathbf{H}_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^{n+1}, \mathbf{A}_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^{n+1}, \dots, Y_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^Q, \mathbf{H}_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^Q \right) \right\}_{b=1}^B,$$

and calculate

$$\widehat{V}_B^n(\mathbf{h}^n, \boldsymbol{\pi}; \boldsymbol{\beta}) = \frac{\sum_{b=1}^B \left( \sum_{v=n}^Q \Delta_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^v Y_{b,\boldsymbol{\pi},\boldsymbol{\beta}}^v \right)}{B}.$$

for each  $\boldsymbol{\pi} \in \Pi$ . To determine the coupon allocation for the current study participant, we set  $\widehat{\boldsymbol{\alpha}}_B^n = \arg \max_{\boldsymbol{\alpha}} \widehat{V}_B^n(\mathbf{h}^n, \boldsymbol{\alpha}; \widehat{\boldsymbol{\beta}}^n)$ , and assign  $\mathbf{a}^n = \pi^n(\mathbf{x}^n, \widehat{\boldsymbol{\alpha}}_B^n)$ . Alternatively, we could use a grid approximation to  $\Pi$  or a gradient descent method to approximate  $\widehat{\boldsymbol{\alpha}}_B^n$ .

## 4.2 Results

In the following simulations, we set the hidden population size to  $N = 5,000$ . The recruitment process begins with an initial sample of 25 individuals randomly drawn from the population,  $|\mathcal{E}_0| = 25$ . The graph model is defined by  $\rho_0^* = 0$ ,  $\boldsymbol{\mu}^* = (1, 1, 1)$ , and  $\Sigma^* = \text{diag}\{(10, 10, 10)\}$ . We vary  $\rho_1^* \in \{1, 2\}$ , to compare simulation experiments in dense and sparse network settings respectively. The researchers have access to three types of coupon allocations  $\mathcal{A} \triangleq \{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$  that correspond to three coupon types  $\mathbb{A} \triangleq \{a_1, a_2, a_3\}$ . Each allocation has 5 coupons. We found that limiting the number of coupons given to each study participant in the pilot study and the warm-up period of the T&I and the RLRDS policies respectively allows us to observe the effects of the learned policies earlier in the sampling process. Consequently, we give two coupons to individuals in the pilot and warm-up period, and increase the allotment to 5 coupons afterwards. Additionally,

$C^v \equiv 1$  for all  $v \in \mathbb{N}$ . For recruit  $j$  of participant  $v$ , define the reward model components as  $\mathbf{Z}_j^v \triangleq \{1, \mathbf{X}_j^v, \mathbb{I}(A_j^v = a_2)\mathbf{X}_j^v, \mathbb{I}(A_j^v = a_3)\mathbf{X}_j^v\}$  and  $\boldsymbol{\beta}_y^* \triangleq (-1, 3\mathbf{k}, -3\mathbf{k}, -6\mathbf{k})$ , where  $\mathbf{k} \triangleq (1, -1, -1)$ . Furthermore, we define the neighbor selection probability distribution as  $\mathbf{p}^v = u(\mathbf{A}^v, \underline{\mathbf{X}}^v)$  such that

$$p_j^v = \begin{cases} \frac{\|\mathbf{X}^v - \mathbf{X}_j^v\|_2^2}{\sum_{j \in \mathcal{M}_1^v} \|\mathbf{X}^v - \mathbf{X}_j^v\|_2^2}, & \text{if } \mathbf{A}^v = \mathbf{a}_2, \\ 1/|\mathcal{M}_1^v|, & \text{otherwise,} \end{cases}$$

where  $\mathcal{M}_1^v$  is the set of participant  $v$ 's potential recruits (as defined in the previous section). This implies that recruiters with the second coupon type will be more likely to recruit neighbors that are different from them. The basic policy objective is clear: we want to ensure that coupon type 1,  $\mathbf{a}_1$ , is used to recruit individuals with covariates that satisfy  $\mathbf{X}^{v\top} \mathbf{k} > 0$ , and coupon type 3,  $\mathbf{a}_3$ , is used to recruit individuals with covariates that satisfy  $\mathbf{X}^{v\top} \mathbf{k} < 0$ . We also wish to recruit individuals with large  $|\mathbf{X}^{v\top} \mathbf{k}|$  because this will increase the probability of the desired outcome contingent on the correct coupon type being awarded. Consequently, it is possible that coupon type 2,  $\mathbf{a}_2$ , is optimal for participant  $v$  if  $|\mathbf{X}^{v\top} \mathbf{k}|$  is low because it increases the likelihood of observing high values in the future. We define the policy space by specifying the function  $g^v$  from (9) as

$$g^v(z) = \begin{cases} \mathbf{a}_1, & \text{if } z > 0.66, \\ \mathbf{a}_2, & \text{if } 0.66 > z \geq 0.33, \\ \mathbf{a}_3, & \text{if } 0.33 > z. \end{cases}$$

This policy space implies that correctly assigning coupons 1 or 3 will depend on the sign of the  $\boldsymbol{\alpha}$  components. The frequency of coupon 2 allocation will be determined by the magnitude of  $\boldsymbol{\alpha}$ . This structure makes finding an optimal policy computationally feasible while maintaining sufficient difficulty to showcase the strength of RL-RDS. Lastly, we set  $\zeta = 1$ ,  $t_{\min} = 0$ , and  $t_{\max} = 3$ .

Figure 2 illustrates the estimated value of policies in both the sparse and dense network settings. It indicates that RL-RDS outperforms all competitor policies by a significant

margin in each regime. In the Supplemental Materials, we test RL-RDS under two additional simulation paradigms. We vary the value/cost and number of coupons given to RDS participants in these experiments. In these contexts, we introduce another branching process working model that incorporates the incentive value in the covariate and reward models (while reducing their dimensionality). This model is described by Equation 36 in the Supplemental Materials. Figures 4 and 5 confirm that RL-RDS outperforms all competitor policies by a significant margin under multiple graph density and coupon allocation settings.

We construct 95% confidence regions for the full network model,  $(\rho_0^*, \rho_1^*, \boldsymbol{\mu}^*, \Sigma^*)$ , with a variety of inference procedures. In these experiments, we assume that  $\Sigma^*$  is diagonal and  $q$  is known to reduce the computational burden of the inference methods. We also add an additional network setting,  $\rho_1^* = 0.5$  (corresponding to a higher density network). Table 1 depicts coverage results for these confidence regions. The table indicates that our simulation-based inference (SBI) technique, which is described by Algorithm 1 in the Supplemental Materials, achieves nominal coverage or greater across all graph settings. In the sparse setting, bootstrapping with the log-likelihood ratio (BS LLR) and approximate bayesian computation (ABC) achieve nominal coverage as well. However, as the graph becomes denser, ABC and BS LLR’s coverage decreases while SBI’s coverage stays above 95%; i.e. the comparison methods fail to adequately quantify uncertainty while our method succeeds. The details of ABC, BS LLR, and bootstrapping with the wald interval (BS WI) are described in the Supplemental Materials.

Table 2 reports 95% projection interval coverage results for each dimension of the average covariate value; i.e.,  $\nu_i$  for  $i \in \{1, 2, 3\}$ . We see that the projection intervals associated with every inference method have high coverage except for BS WI. This is unsurprising since we conduct inference calibrated to have 95% coverage over all network parameters simultaneously. Encouragingly, our method, SBI, has a smaller interval length than contenders that achieve nominal coverage (BS LLR and ABC).




Figure 2: This figure compares the estimated cumulative reward of each policy with 90% Monte Carlo confidence intervals over multiple sample sizes and graph densities.

Table 1: Simultaneous Network Parameter Coverage

Method	Graph Density	Coverage
<b>SBI</b>	Low	0.975
<b>SBI</b>	Medium	0.962
<b>SBI</b>	High	0.979
ABC	Low	0.950
ABC	Medium	0.749
ABC	High	0.393
BS LLR	Low	0.945
BS LLR	Medium	0.937
BS LLR	High	0.908
BS WI	Low	0.526
BS WI	Medium	0.480
BS WI	High	0.508

These are simultaneous coverage rates for the network parameters over 250 simulations for three graph settings.

Table 2: Covariate Distribution Mean Intervals

Method	Dim.	Coverage	Interval
<b>SBI</b>	1	0.99	2.78
<b>SBI</b>	2	0.99	2.71
<b>SBI</b>	3	1.00	2.72
ABC	1	1.00	3.04
ABC	2	1.00	3.23
ABC	3	1.00	3.09
BS LLR	1	0.98	2.86
BS LLR	2	0.98	2.82
BS LLR	3	0.97	2.84
BS WI	1	0.88	2.15
BS WI	2	0.87	2.15
BS WI	3	0.89	2.15

These are coverage rates and interval lengths averaged over graph settings for the mean of each dimension of the covariate distribution over 250 simulations.

## 5 Discussion

We showed that RL-driven adaptive RDS can lead to dramatic improvements in the effectiveness of RDS. In the course of deriving an RL strategy, we: (i) showed that a branching process approximation to RDS is useful for guiding coupon selection without having to fully model the underlying system dynamics; (ii) extended asymptotic theory for adaptively sampled M-estimators to Markov Decision Processes; and (iii) developed novel methods for post adaptive-RDS inference that remain valid even when the true model is not identifiable.

This work is the first to consider fully adaptive RDS methods using RL. Consequently, there are a number of open problems and interesting directions for future work. One such direction is the development of a regret bound that explicitly depends on the quality of the branching process approximation. While branching processes have been used extensively to model epidemics evolving on networks, the quality of these approximations remains an important open question. Another direction for future work regards statistical efficiency. As noted in the introduction, it is possible to fold measures of information gain into participant outcomes to improve power in post-study analyses. However, how best to do this for various network models is not clear. Finally, as we considered projective dynamic

graph models, exploring whether there is a more general class of models to which the asymptotic theory still applies is yet another potential area for future research.

## References

- Shipra Agrawal and Navin Goyal. Further optimal regret bounds for thompson sampling. In *Artificial intelligence and statistics*, pages 99–107, 2013.
- Krishna B Athreya, Peter E Ney, and PE Ney. *Branching processes*. Courier Corporation, 2004.
- Donald A Berry and Bert Fristedt. *Bandit problems: sequential allocation of experiments (Monographs on statistics and applied probability)*. Springer, 1985.
- Aurélien Bibaut and Nathan Kallus. Demistifying inference after adaptive experiments. *arXiv preprint arXiv:2405.01281*, 2024.
- Aurélien Bibaut, Maria Dimakopoulou, Nathan Kallus, Antoine Chambaz, and Mark van Der Laan. Post-contextual-bandit inference. *Advances in neural information processing systems*, 34:28548–28559, 2021.
- Lawrence D Brown. *Fundamentals of statistical exponential families: with applications in statistical decision theory*. Ims, 1986.
- Olivier Chapelle and Lihong Li. An empirical evaluation of thompson sampling. In *Advances in neural information processing systems*, pages 2249–2257, 2011.
- Snigdhanu Chatterjee and Arup Bose. Generalized bootstrap for estimating equations. *The Annals of Statistics*, 33(1):414 – 436, 2005. doi: 10.1214/009053604000000904. URL <https://doi.org/10.1214/009053604000000904>.
- Eric J Cooks, Kyle A Duke, Jordan M Neil, Melissa J Vilaro, Danyell Wilson-Howard, Francois Modave, Thomas J George, Folakemi T Odedina, Benjamin C Lok, Peter Carek, et al. Telehealth and racial disparities in colorectal cancer screening: A pilot study of how virtual clinician characteristics influence screening intentions. *Journal of clinical and translational science*, 6(1):e48, 2022.

- Forrest W Crawford, Jiacheng Wu, and Robert Heimer. Hidden population size estimation from respondent-driven sampling: a network approach. *Journal of the American Statistical Association*, 113(522):755–766, 2018.
- Jean-François Delmas and Laurence Marsalle. Detection of cellular aging in a galton-watson process. *Stochastic Processes and their Applications*, 120(12):2495–2519, 2010.
- Yash Deshpande, Lester Mackey, Vasilis Syrgkanis, and Matt Taddy. Accurate inference for adaptive linear models. In *International Conference on Machine Learning*, pages 1194–1203. PMLR, 2018.
- Daniel Eisenberg, Ezra Golberstein, Janis L Whitlock, and Marilyn F Downs. Social contagion of mental health: evidence from college roommates. *Health economics*, 22(8):965–986, 2013.
- Pak Wing Fong, Wai Keung Li, CW Yau, and Chun Shan Wong. On a mixture vector autoregressive model. *Canadian Journal of Statistics*, 35(1):135–150, 2007.
- Christian Francq and J-M Zakoian. Stationarity of multivariate markov-switching arma models. *Journal of Econometrics*, 102(2):339–364, 2001.
- Krista J Gile. Improved inference for respondent-driven sampling data with application to hiv prevalence estimation. *Journal of the American Statistical Association*, 106(493):135–146, 2011.
- Krista J Gile and Mark S Handcock. 7. respondent-driven sampling: An assessment of current methodology. *Sociological methodology*, 40(1):285–327, 2010.
- Krista J Gile and Mark S Handcock. Network model-assisted inference from respondent-driven sampling data. *Journal of the Royal Statistical Society. Series A, (Statistics in Society)*, 178(3):619, 2015.
- Sharad Goel and Matthew J Salganik. Assessing respondent-driven sampling. *Proceedings of the National Academy of Sciences*, 107(15):6743–6747, 2010.

- Aditya Gopalan and Shie Mannor. Thompson sampling for learning parameterized markov decision processes. In *Conference on Learning Theory*, pages 861–898, 2015.
- Aditya Gopalan, Shie Mannor, and Yishay Mansour. Thompson sampling for complex online problems. In *International Conference on Machine Learning*, pages 100–108, 2014.
- Vitor Hadad, David A Hirshberg, Ruohan Zhan, Stefan Wager, and Susan Athey. Confidence intervals for policy evaluation in adaptive experiments. *Proceedings of the national academy of sciences*, 118(15):e2014602118, 2021.
- Peter Hall and Christopher C Heyde. *Martingale limit theory and its application*. Academic press, 2014.
- James D Hamilton. A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica: Journal of the econometric society*, pages 357–384, 1989.
- Douglas D Heckathorn. Respondent-driven sampling: a new approach to the study of hidden populations. *Social problems*, 44(2):174–199, 1997.
- Douglas D Heckathorn and Christopher J Cameron. Network sampling: From snowball and multiplicity to respondent-driven sampling. *Annual review of sociology*, 43:101–119, 2017.
- Eric B Laber and Susan A Murphy. Adaptive confidence intervals for the test error in classification. *Journal of the American Statistical Association*, 106(495):904–913, 2011.
- Eric B Laber, Nick J Meyer, Brian J Reich, Krishna Pacifici, Jaime A Collazo, and John M Drake. Optimal treatment allocations in space and time for on-line control of an emerging infectious disease. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 67(4):743–789, 2018.
- Tor Lattimore and Csaba Szepesvári. *Bandit algorithms*. Cambridge University Press, 2020.

- Xin Lu, Linus Bengtsson, Tom Britton, Martin Camitz, Beom Jun Kim, Anna Thorson, and Fredrik Liljeros. The sensitivity of respondent-driven sampling. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 175(1):191–216, 2012.
- Simón Lunagómez, Marios Papamichalis, Patrick J Wolfe, and Edoardo M Airoidi. Evaluating and optimizing network sampling designs: Decision theory and information theory perspectives. *arXiv preprint arXiv:1811.07829*, 2018.
- Xiaoyan Ma, Qiyun Zhang, Xiong He, Weidong Sun, Hai Yue, Sanny Chen, H Fisher Raymond, Yang Li, Min Xu, Hui Du, et al. Trends in prevalence of hiv, syphilis, hepatitis c, hepatitis b, and sexual risk behavior among men who have sex with men: results of 3 consecutive respondent-driven sampling surveys in beijing, 2004 through 2006. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 45(5):581–587, 2007.
- Syed S Mahmood, Daniel Levy, Ramachandran S Vasani, and Thomas J Wang. The framingham heart study and the epidemiology of cardiovascular disease: a historical perspective. *The lancet*, 383(9921):999–1008, 2014.
- Allison M McFall, Bryan Lau, Carl Latkin, Aylur K Srikrishnan, Santhanam Anand, Canjeevaram K Vasudevan, Shruti H Mehta, and Sunil S Solomon. Optimizing respondent-driven sampling to find undiagnosed hiv-infected people who inject drugs. *AIDS*, 35(3):485–494, 2021.
- Whitney K Newey and Daniel McFadden. Chapter 36 large sample estimation and hypothesis testing. volume 4 of handbook of econometrics. *Elsevier*, 12:2111–2245, 1994.
- Brea L Perry, Bernice A Pescosolido, and Stephen P Borgatti. *Egocentric network analysis: Foundations, methods, and models*, volume 44. Cambridge university press, 2018.
- Martin L Putterman. Markov decision processes. *John Wiley and Sons, New York*, 1994.
- Sarah Raifman, Michelle A DeVost, Jean C Digitale, Yea-Hung Chen, and Meghan D Morris. Respondent-driven sampling: a sampling method for hard-to-reach populations and beyond. *Current Epidemiology Reports*, 9(1):38–47, 2022.

- JMH Risser, P Padgett, M Wolverton, and WL Risser. Relationship between heterosexual anal sex, injection drug use and hiv infection among black men and women. *International journal of STD & AIDS*, 20(5):310–314, 2009.
- J.M. Robins. Optimal structural nested models for optimal sequential decisions. In *Proceedings of the Second Seattle Symposium in Biostatistics: Analysis of Correlated Data*, 2004.
- Sebastien Roch and Karl Rohe. Generalized least squares can overcome the critical threshold in respondent-driven sampling. *Proceedings of the National Academy of Sciences*, 115(41):10299–10304, 2018.
- Karl Rohe. Network driven sampling; a critical threshold for design effects. *arXiv preprint arXiv:1505.05461*, 2015.
- Karl Rohe. A critical threshold for design effects in network sampling. *Annals of Statistics*, 47(1):556–582, 2019.
- D.B. Rubin. Bayesian inference for causal effects: The role of randomization. *The Annals of Statistics*, 6(1):34–58, 1978.
- Daniel Russo and Benjamin Van Roy. An information-theoretic analysis of thompson sampling. *The Journal of Machine Learning Research*, 17(1):2442–2471, 2016.
- Daniel J Russo, Benjamin Van Roy, Abbas Kazerouni, Ian Osband, and Zheng Wen. A tutorial on thompson sampling. *Foundations and Trends® in Machine Learning*, 11(1):1–96, 2018.
- Boris A Sevast’yanov and Andreĭ M Zubkov. Controlled branching processes. *Theory of Probability & Its Applications*, 19(1):14–24, 1974.
- Aleksandrs Slivkins et al. Introduction to multi-armed bandits. *Foundations and Trends® in Machine Learning*, 12(1-2):1–286, 2019.
- Neil A Spencer and Cosma Rohilla Shalizi. Projective, sparse, and learnable latent position network models. *arXiv preprint arXiv:1709.09702*, 2017.

- Jerzy Splawa-Neyman, DM Dabrowska, TP Speed, et al. On the application of probability theory to agricultural experiments. essay on principles. section 9. *Statistical Science*, 5 (4):465–472, 1990.
- Robert Stelzer. On markov-switching arma processes—stationarity, existence of moments, and geometric ergodicity. *Econometric Theory*, 25(1):43–62, 2009.
- R.S. Sutton and A.G. Barto. *Reinforcement Learning: An Introduction*. The MIT Press, 2018.
- Csaba Szepesvári. Algorithms for reinforcement learning. *Synthesis Lectures on Artificial Intelligence and Machine Learning*, 4(1):1–103, 2010.
- Adam G Tabák, Christian Herder, Wolfgang Rathmann, Eric J Brunner, and Mika Kivimäki. Prediabetes: a high-risk state for diabetes development. *The Lancet*, 379 (9833):2279–2290, 2012.
- Amber Tomas and Krista J Gile. The effect of differential recruitment, non-response and non-recruitment on estimators for respondent-driven sampling. *Electronic Journal of Statistics*, 5:899–934, 2011.
- Anastasios A Tsiatis, Marie Davidian, Shannon T Holloway, and Eric B Laber. *Dynamic Treatment Regimes: Statistical Methods for Precision Medicine*. CRC press, 2019.
- Aad W Van der Vaart. *Asymptotic statistics*, volume 3. Cambridge university press, 2000.
- Josey VanOrsdale. *Adaptive Respondent Driven Sampling of Social Networks: A Simulation-Based Study Using Machine Learning*. PhD thesis, The University of Nebraska-Lincoln, 2023.
- Cyprian Wejnert and Douglas Heckathorn. Respondent-driven sampling: operational procedures, evolution of estimators, and topics for future research. *The SAGE handbook of innovation in social research methods*. London: SAGE Publications, Ltd, pages 473–97, 2011.

Justin David Naggar Wertz, Eric Laber, and Alexander Volfovsky. Hidden population estimation with indirect inference and auxiliary information. In *The 40th Conference on Uncertainty in Artificial Intelligence*, 2024.

Ruohan Zhan, Zhimei Ren, Susan Athey, and Zhengyuan Zhou. Policy learning with adaptively collected data. *Management Science*, 2023.

Ruohan Zhan, Zhimei Ren, Susan Athey, and Zhengyuan Zhou. Policy learning with adaptively collected data. *Management Science*, 70(8):5270–5297, 2024.

Kelly Zhang, Lucas Janson, and Susan Murphy. Statistical inference with m-estimators on adaptively collected data. *Advances in neural information processing systems*, 34: 7460–7471, 2021.

Kelly W Zhang, Lucas Janson, and Susan A Murphy. Inference for batched bandits. *arXiv preprint arXiv:2002.03217*, 2020.

# Supplemental Materials

## 6 Glossary

$i$ (subscript)	index for interim study participants organized in epochs
$v$ (superscript)	index for interim study participants ordered by arrival times
$\mathbf{A}_i$	coupon allocation given to $i$
$T_i$	arrival time of $i$
$\mathbf{X}_i$	covariate of $i$
$Y_i$	outcome of interest for $i$
$C_i$	cost of recruitment for $i$
$R_i$	recruiter of $i$
$M_i$	number of recruits of $i$
$\underline{\mathbf{X}}_i, \underline{\mathbf{Y}}_i, \underline{\mathbf{T}}_i$	collection of covariates, response statuses, and response times associated with the recruits of participant $i$
$\mathbf{D}_i$	complete information associate with the recruits of individual $i$
$\mathbf{H}^v$	information available to researchers at the time $v$ is given their coupon allocation
$\mathcal{E}_j, \bar{\mathcal{E}}_j$	epoch $j$ and all epochs up to epoch $j$ respectively
$\kappa_j, \bar{\kappa}_j$	size of epoch $j$ and size of all epochs up to epoch $j - 1$ respectively
$\mathcal{Z}_j, \bar{\mathcal{Z}}_j$	data associated with epoch $j$ and data associated with all epochs up to epoch $j$ respectively
$\mathcal{F}_j$	sigma field associated with the first $j$ epochs without the coupon allocation information associated with individuals in the $j^{\text{th}}$ epoch
$\mathcal{D}^\kappa$	data indexed by arrival time (first $\kappa$ individuals)
$\boldsymbol{\pi}$	deterministic coupon allocation strategy
$\boldsymbol{\pi}^{\text{opt}}$	optimal coupon allocation strategy
$\psi^v(\mathbf{h}^v)$	set of allowable coupon allocations given $\mathbf{h}^v$
$V^v(\mathbf{h}^v, \boldsymbol{\pi})$	history-value function of $\boldsymbol{\pi}$ at $\mathbf{h}^v$
$\hat{\xi}_B^v(\mathbf{h}^v, \mathbf{a}^v)$	estimated probability that $\mathbf{a}^v$ is the optimal action given $\mathbf{h}^v$
$\boldsymbol{\beta} \in \mathcal{B}$	parameter that indexes the branching process
$W_i$	stabilizing weights for log-likelihood
$l_i(\boldsymbol{\beta})$	log-likelihood associated with $\mathbf{D}_i$ conditional on $\mathcal{F}_{j-1}$ , where $i \in \mathcal{E}_{j-1}$
$\hat{\boldsymbol{\beta}}_J$	M-estimator of weighted log-likelihood based on complete epoch information up to epoch $J$
$\mathcal{I}$	asymptotic event associated with super-critical branching processes
$q^v(\boldsymbol{\beta})$	log-likelihood component associated with $v$ of the arrival process data, $\mathcal{D}^\kappa$
$\boldsymbol{\theta} \in \Theta$	parameter that indexes the dynamic network model
$\ell_\kappa^\theta(\boldsymbol{\beta}_\mathbf{x})$	log-likelihood of the branching process covariate model for a collection of $\kappa$ subjects sampled under the RDS process when $\boldsymbol{\theta}$ is the true parameter
$\bar{\boldsymbol{\beta}}_\mathbf{x}(\boldsymbol{\theta})$	maximizes limit of $\ell_\kappa^\theta(\boldsymbol{\beta}_\mathbf{x})/\kappa$
$\Gamma_{1-\alpha, \kappa}(\bar{\boldsymbol{\beta}}_\mathbf{x})$	confidence region for $\boldsymbol{\theta}^*$ based on $\bar{\boldsymbol{\beta}}_\mathbf{x}$

Table 3: Table of notation

## 7 Inference Algorithms

Algorithm 1 generates an approximation of the confidence interval described in Section 3. Recall that we need to approximate  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$  for an arbitrary  $\boldsymbol{\theta} \in \Theta$ . We do this by first generating a large network for  $\boldsymbol{\theta}$  and simulating an RDS study of size  $K \gg \kappa$  over this network. Using this simulated sample, we maximize the log-likelihood of the branching process working model to obtain  $\widehat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta})$ .

---

**Algorithm 1:** SBI Confidence Set

---

- 1 **Input:** Set of parameter values  $\Theta_S \subseteq \Theta$ ; the MLE of the working model,  $\widehat{\beta}_{\mathbf{x}}^{\kappa}(\boldsymbol{\theta}^*)$ ;  
 $K$ , the sample size for the approximation of  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$ ; and  $B$ , the number of  
simulated branching processes used to approximate the test statistic distribution
  - 2 **for**  $\boldsymbol{\theta} \in \Theta_S$  **do**
  - 3     Generate a large network from  $\boldsymbol{\theta}$  (of size greater than  $K$ ), and simulate an  
RDS study of size  $K$  over this graph
  - 4     Use the branching process working model with this sample to find  $\widehat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta})$
  - 5     Simulate  $B$  branching processes of size  $\kappa$  over graphs generated according to  $\boldsymbol{\theta}$
  - 6     For  $b \in \{1, 2, \dots, B\}$ , calculate  $\widehat{\beta}_{\mathbf{x},b}^{\kappa}(\boldsymbol{\theta})$
  - 7     Construct a set of test statistic values
$$\widehat{P}_{\kappa}^{\boldsymbol{\theta}}(\bar{\beta}_{\mathbf{x}}) = \left\{ -2 \left[ \ell_{\kappa}^{\boldsymbol{\theta}} \left\{ \widehat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta}) \right\} - \ell_{\kappa}^{\boldsymbol{\theta}} \left\{ \widehat{\beta}_{\mathbf{x},b}^{\kappa}(\boldsymbol{\theta}) \right\} \right], b \in \{1, 2, \dots, B\} \right\}$$
  - 8     Determine the  $1 - \alpha$  upper quantile of  $\widehat{P}_{\kappa}^{\boldsymbol{\theta}}(\bar{\beta}_{\mathbf{x}})$  and label this  $\widehat{\gamma}_{1-\alpha,\kappa}^{\boldsymbol{\theta}}(\bar{\beta}_{\mathbf{x}})$
  - 9     If  $-2 \left[ \ell_{\kappa}^{\boldsymbol{\theta}^*} \left\{ \widehat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta}) \right\} - \ell_{\kappa}^{\boldsymbol{\theta}^*} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa}(\boldsymbol{\theta}^*) \right\} \right] \leq \widehat{\gamma}_{1-\alpha,\kappa}^{\boldsymbol{\theta}}(\bar{\beta}_{\mathbf{x}})$ , then include  $\boldsymbol{\theta}$  in  $\Gamma_{1-\alpha,\kappa}$
  - 10 **end**
  - 11 **Output:**  $\Gamma_{1-\alpha,\kappa}$
-

We compare our inference approach to a series of natural alternatives. Algorithm 2 is inspired by approximate Bayesian computation (ABC).

---

**Algorithm 2:** ABC Confidence Set

---

- 1 **Input:** Set of parameter values  $\Theta_S \subseteq \Theta$ ;  $K$ , the sample size for the approximation of  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$ ; and  $B$ , the number of bootstrap samples
  - 2 Take  $B$  bootstrapped samples using a generalized bootstrap for estimating equations as described in Chatterjee and Bose (2005) and find the MLE for every sample, label this set  $\{\hat{\beta}_{\mathbf{x},b}^{\kappa}\}_{b=1}^B$
  - 3 For each  $\boldsymbol{\theta} \in \Theta_S$ , generate a large network consistent with  $\boldsymbol{\theta}$  (of size greater than  $K$ ), and simulate an RDS study of size  $K$  over this graph
  - 4 Use the log-likelihood of the branching process working model with this sample to find  $\hat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta})$
  - 5 **Output:**  $\Gamma_{1-\alpha,\kappa} = \{\hat{\boldsymbol{\theta}}_b = \arg \min_{\boldsymbol{\theta} \in \Theta} \|\hat{\beta}_{\mathbf{x},b}^{\kappa} - \hat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta})\|^2, b \in \{1, 2, \dots, B\}\}$
- 

Algorithm 3 describes the bootstrap with Wald interval (BS WI) and the bootstrap with likelihood ratio (BS LLR) approaches. It uses a generalized bootstrap for estimating equations as described in Chatterjee and Bose (2005).

---

**Algorithm 3:** Bootstrap Confidence Set
 

---

- 1 **Input:** Set of parameter values  $\Theta_S \subseteq \Theta$ ; the MLE of the working model,  $\widehat{\beta}_{\mathbf{x}}^\kappa$ ;  $K$ ,  
the sample size for the approximation of  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$ ; and  $B$ , the number of  
bootstrapped samples
  - 2 Take  $B$  bootstrapped samples using a generalized bootstrap for estimating  
equations as described in Chatterjee and Bose (2005)
  - 3 For  $b \in \{1, 2, \dots, B\}$ , estimate the MLE,  $\widehat{\beta}_{\mathbf{x},b}^\kappa$ , for each bootstrap from the  
log-likelihood  $\ell_{\kappa,b}$
  - 4 Construct a set of test statistic the following manner:
  - 5 **if LLR then**
  - 6      $\widehat{P}_\kappa = \left\{ -2 \left[ \ell_{\kappa,b}(\widehat{\beta}_{\mathbf{x}}^\kappa) - \ell_{\kappa,b}(\widehat{\beta}_{\mathbf{x},b}^\kappa) \right], b \in \{1, 2, \dots, B\} \right\}$
  - 7 **else**
  - 8     Calculate  $\widehat{\Sigma}_B^\kappa = \sum_{b=1}^B \left\{ \widehat{\beta}_{\mathbf{x},b}^\kappa - \widehat{\beta}_{\mathbf{x}}^\kappa \right\} \left\{ \widehat{\beta}_{\mathbf{x},b}^\kappa - \widehat{\beta}_{\mathbf{x}}^\kappa \right\}^\top / B$   
        $\widehat{P}_\kappa = \left\{ \left\{ \widehat{\beta}_{\mathbf{x},b}^\kappa - \widehat{\beta}_{\mathbf{x}}^\kappa \right\}^\top \widehat{\Sigma}_B^\kappa \left\{ \widehat{\beta}_{\mathbf{x},b}^\kappa - \widehat{\beta}_{\mathbf{x}}^\kappa \right\} \right\}$
  - 9 **end**
  - 10 Determine a  $1 - \alpha$  upper quantile of  $\widehat{P}_\kappa$  and label this  $\widehat{\gamma}_{1-\alpha,\kappa}$
  - 11 **for**  $\boldsymbol{\theta} \in \Theta_S$  **do**
  - 12     Generate a large network consistent with  $\boldsymbol{\theta}$  (of size greater than  $K$ ), and  
simulate an RDS study of size  $K$  over this graph
  - 13     Use the log-likelihood of the branching process working model with this  
sample to find  $\widehat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta})$
  - 14     **if LLR then**
  - 15         If  $-2 \left[ \ell_\kappa \left\{ \widehat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta}) \right\} - \ell_\kappa \left\{ \widehat{\beta}_{\mathbf{x}}^\kappa \right\} \right] \leq \widehat{\gamma}_{1-\alpha,\kappa}$ , then include  $\boldsymbol{\theta}$  in  $\Gamma_{1-\alpha,\kappa}$
  - 16     **else**
  - 17         If  $\left\{ \widehat{\beta}_{\mathbf{x}}^\kappa - \widehat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta}) \right\}^\top \widehat{\Sigma}_B^\kappa \left\{ \widehat{\beta}_{\mathbf{x}}^\kappa - \widehat{\beta}_{\mathbf{x}}^K(\boldsymbol{\theta}) \right\} \leq \widehat{\gamma}_{1-\alpha,\kappa}$ , then include  $\boldsymbol{\theta}$  in  $\Gamma_{1-\alpha,\kappa}$
  - 18     **end**
  - 19 **end**
  - 20 **Output:**  $\Gamma_{1-\alpha,\kappa}$
-

## 8 Proof of Theorem 2.1

### 8.1 Supporting Martingale Limit Theory

Theorems 8.1-8.3 establish useful martingale laws of large numbers.

**Theorem 8.1** (Theorem 2.19 from Hall and Heyde (2014)). *Let  $\{X_n, n \geq 1\}$  be a sequence of random variables and  $\{\mathcal{F}_n, n \geq 1\}$  be an increasing sequence of  $\sigma$ -fields such that  $X_n$  is  $\mathcal{F}^n$  measurable for each  $n$ . Let  $X$  be a r.v. and  $c$  a constant such  $\mathbb{E}|X| < \infty$ ,  $\mathbb{E}(|X| \log^+ |X|) < \infty$ , and  $\mathbb{P}(|X_n| > x) \leq c\mathbb{P}(|X| > x)$  for each  $x \geq 0$  and  $n \geq 1$  (note that if  $X_n$  is bounded, this condition is satisfied). Then*

$$\frac{1}{n} \sum_{i=1}^n \{X_i - \mathbb{E}(X_i | \mathcal{F}_{i-1})\} \rightarrow 0$$

*a.s. as  $n \rightarrow \infty$ .*

**Theorem 8.2** (Theorem 2.17 from Hall and Heyde (2014)). *Let  $\{S_n = \sum_{i=1}^n X_i, \mathcal{F}_n, n \geq 1\}$  be a martingale, and let  $1 \leq p \leq 2$ . Then  $S_n$  converges a.s. on the set  $\{\sum_{i=1}^{\infty} \mathbb{E}(|X_i|^p | \mathcal{F}_{i-1}) < \infty\}$ .*

**Theorem 8.3** (Theorem 2.18 from Hall and Heyde (2014)). *Let  $\{S_n = \sum_{i=1}^n X_i, \mathcal{F}_n, n \geq 1\}$  be a martingale and  $\{U_n, n \geq 1\}$  a non-decreasing sequence of positive random variables such that  $U_n$  is  $\mathcal{F}_{n-1}$ -measurable for each  $n$ . If  $1 \leq p \leq 2$ , then*

$$\lim_{n \rightarrow \infty} U_n^{-1} S_n = 0 \text{ a.s.}$$

*on the set  $\{\lim_{n \rightarrow \infty} U_n = \infty, \sum_{i=1}^{\infty} U_i^{-p} \mathbb{E}(|X_i|^p | \mathcal{F}_{i-1}) < \infty\}$ .*

We also state the Toeplitz Lemma from Hall and Heyde (2014).

**Lemma 8.1.** *For  $i \geq 1$ , if  $x_i$  are real numbers,  $a_i$  are positive numbers, and  $b_n = \sum_{i=1}^n a_i$  diverges, then  $x_n \rightarrow x$  ensures that*

$$b_n^{-1} \sum_{i=1}^n a_i x_i \rightarrow x.$$

Lastly, we state a strong law of large numbers specific to the branching process model developed in Section 2. Assume the historical information and epoch setup of Section 2 for the following Theorem.

**Theorem 8.4.** *For function  $z : \mathcal{D} \rightarrow \mathbb{R}$ , assume for any  $j \in \mathbb{N}$ ,  $i \in \mathcal{E}_{j-1}$ ,  $\beta \in \mathcal{B}$ , and  $B < \infty$  that*

$$\mathbb{E}_{\beta, \hat{\pi}} \{z^2(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\} < B. \quad (10)$$

*Additionally, assume that for  $j \in \mathbb{N}$ ,  $\{\mathbf{D}_i\}_{i \in \mathcal{E}_{j-1}}$  are conditionally independent given  $\mathcal{F}_{j-1}$ .*

*For  $\delta > 0$ , define the event  $E_{\mathcal{J}} = \{\mathcal{J} > \delta\}$ , where  $\mathcal{J}$  is defined in Assumption 5. Then,*

$$\lim_{J \rightarrow \infty} \frac{1}{\bar{\kappa}_J} \left[ \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i z(\mathbf{D}_i) - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\} \right] \rightarrow 0$$

*on the event  $E_{\mathcal{J}}$ .*

*Proof.* We first consider the sum

$$\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i z(\mathbf{D}_i) - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\}.$$

This is clearly a martingale sequence with respect to fields  $\{\mathcal{F}_j\}_{j=1}^J$  and martingale differences

$$\left[ \sum_{i \in \mathcal{E}_{j-1}} W_i z(\mathbf{D}_i) - \mathbb{E}_{\beta, \hat{\pi}} \left\{ \sum_{i \in \mathcal{E}_{j-1}} W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1} \right\} \right]_{j=1}^J.$$

Consequently, we can invoke Theorem 8.3, implying that

$$\lim_{J \rightarrow \infty} \frac{1}{\bar{\kappa}_J} \left[ \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i z(\mathbf{D}_i) - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\} \right] = 0$$

on the set

$$\left\{ \begin{aligned} & \lim_{J \rightarrow \infty} \bar{\kappa}_J \rightarrow \infty, \\ & \lim_{J \rightarrow \infty} \sum_{j=1}^J \frac{1}{\bar{\kappa}_j^2} \mathbb{E}_{\beta, \hat{\pi}} \left( \left[ \sum_{i \in \mathcal{E}_{j-1}} W_i z(\mathbf{D}_i) - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) | \mathcal{F}_{j-1}\} \right]^2 \mid \mathcal{F}_{j-1} \right) < \infty \end{aligned} \right\}.$$

We examine

$$\begin{aligned} & \sum_{j=1}^J \frac{1}{\bar{\kappa}_j^2} \mathbb{E}_{\beta, \hat{\pi}} \left( \left[ \sum_{i \in \mathcal{E}_{j-1}} W_i z(\mathbf{D}_i) - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) | \mathcal{F}_{j-1}\} \right]^2 \mid \mathcal{F}_{j-1} \right) \stackrel{(a)}{=} \\ & \sum_{j=1}^J \frac{1}{\bar{\kappa}_j^2} \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta, \hat{\pi}} \left( [W_i z(\mathbf{D}_i) - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) | \mathcal{F}_{j-1}\}]^2 \mid \mathcal{F}_{j-1} \right) = \\ & \sum_{j=1}^J \frac{1}{\bar{\kappa}_j^2} \sum_{i \in \mathcal{E}_{j-1}} \left( \mathbb{E}_{\beta, \hat{\pi}} \{W_i^2 z^2(\mathbf{D}_i) | \mathcal{F}_{j-1}\} - \right. \\ & \quad \left. \mathbb{E}_{\beta, \hat{\pi}} [\mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) | \mathcal{F}_{j-1}\}^2 | \mathcal{F}_{j-1}] \right) \leq \\ & \sum_{j=1}^J \frac{1}{\bar{\kappa}_j^2} \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta, \hat{\pi}} (W_i^2 z^2(\mathbf{D}_i) | \mathcal{F}_{j-1}) = \\ & \sum_{j=1}^J \frac{1}{\bar{\kappa}_j^2} \sum_{i \in \mathcal{E}_{j-1}} \int W_i^2 z^2(\mathbf{D}_i) f(\mathbf{D}_i | \mathbf{A}_i, \mathcal{F}_{j-1}) \mathbb{P}_{\hat{\pi}}(\mathbf{A}_i | \mathcal{F}_{j-1}) d\nu \stackrel{(b)}{=} \\ & \sum_{j=1}^J \frac{1}{\bar{\kappa}_j^2} \sum_{i \in \mathcal{E}_{j-1}} \int \frac{\mathbb{P}_{\hat{\pi}}(\mathbf{A}_i | \mathbf{H}_i)}{\mathbb{P}_{\hat{\pi}}(\mathbf{A}_i | \mathbf{H}_i)} z^2(\mathbf{D}_i) f(\mathbf{D}_i | \mathbf{A}_i, \mathcal{F}_{j-1}) \mathbb{P}_{\hat{\pi}}(\mathbf{A}_i | \mathbf{H}_i) d\nu = \\ & \sum_{j=1}^J \frac{1}{\bar{\kappa}_j^2} \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta, \hat{\pi}} (z^2(\mathbf{D}_i) | \mathcal{F}_{j-1}) \stackrel{(c)}{\leq} \\ & \sum_{j=1}^J \frac{\kappa_j}{\bar{\kappa}_j^2} B \stackrel{(d)}{\leq} \\ & \sum_{j=1}^J \frac{1}{\kappa_j} B. \end{aligned}$$

Equality (a) follows from the fact that for  $i > g$ ,

$$\begin{aligned}
& \mathbb{E}_{\beta, \hat{\pi}} ([W_i z(\mathbf{D}_i) - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\}] [W_g z(\mathbf{D}_g) - \mathbb{E}_{\beta, \hat{\pi}} \{W_g z(\mathbf{D}_g) \mid \mathcal{F}_{j-1}\}] \mid \mathcal{F}_{j-1}) \\
&= \mathbb{E}_{\beta, \hat{\pi}} \left( [\mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}, \mathbf{D}_g\} - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\}] \times \right. \\
&\quad \left. [W_g z(\mathbf{D}_g) - \mathbb{E}_{\beta, \hat{\pi}} \{W_g z(\mathbf{D}_g) \mid \mathcal{F}_{j-1}\}] \mid \mathcal{F}_{j-1} \right) \\
&\stackrel{(i)}{=} \mathbb{E}_{\beta, \hat{\pi}} \left( [\mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\} - \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\}] \times \right. \\
&\quad \left. [W_g z(\mathbf{D}_g) - \mathbb{E}_{\beta, \hat{\pi}} \{W_g z(\mathbf{D}_g) \mid \mathcal{F}_{j-1}\}] \mid \mathcal{F}_{j-1} \right) \\
&= 0.
\end{aligned}$$

Equality (i) follows because

$$\mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}, \mathbf{D}_g\} = \mathbb{E}_{\beta, \hat{\pi}} \{W_i z(\mathbf{D}_i) \mid \mathcal{F}_{j-1}\}.$$

Equality (b) follows from the fact that  $\mathbb{P}_{\hat{\pi}}(\mathbf{A}_i \mid \mathcal{F}_{j-1}) = \mathbb{P}_{\hat{\pi}}(\mathbf{A}_i \mid \mathbf{H}_i)$  and the definition of  $W_i$ . Inequality (c) follows from Equation 10. Additionally, we know

$$\begin{aligned}
\sum_{j=1}^J \frac{1}{\kappa_j} B &= \sum_{j=1}^J \frac{1}{m^{j-1}} \frac{m^{j-1}}{\kappa_j} B \\
&= \left( \sum_{j=1}^J \frac{1}{m^{j-1}} \right) \left( \sum_{j=1}^J \frac{1}{m^{j-1}} \right)^{-1} \sum_{j=1}^J \frac{1}{m^{j-1}} \frac{m^{j-1}}{\kappa_j} B \\
&\stackrel{\text{a.s.}}{\rightarrow} \left( \sum_{j=1}^J \frac{1}{m^{j-1}} \right) \mathcal{I}^{-1} B
\end{aligned}$$

by Lemma 8.1 and Assumption 5. Because we assume event  $E_{\mathcal{I}}$ , we know that  $\mathcal{I}^{-1} < \frac{1}{\delta}$ .

Consequently,

$$\sum_{j=1}^J \frac{1}{\kappa_j} B \stackrel{\text{a.s.}}{\rightarrow} \frac{1}{1 - 1/m} \mathcal{I}^{-1} B < \frac{B}{\delta(1 - 1/m)} < \infty.$$

■

## 8.2 Consistency

First, we show asymptotic consistency of  $\widehat{\beta}_J$ . We follow the proof of Zhang et al. (2021), making slight changes to account for a more general context. We define  $e_i(\beta) \triangleq W_i l_i(\beta)$ . By the definition of  $\widehat{\beta}_J$ ,

$$\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} e_i(\widehat{\beta}_J) = \sup_{\beta \in \mathcal{B}} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} e_i(\beta) \geq \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} e_i(\beta^*). \quad (11)$$

Note that  $\|\widehat{\beta}_J - \beta^*\| > \epsilon > 0$  implies that

$$\sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} e_i(\beta) = \sup_{\beta \in \mathcal{B}} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} e_i(\beta). \quad (12)$$

Define  $\mathbb{P}_{\beta^*, \widehat{\pi}}$  as the probability distribution associated with the complete data-generating distribution, where  $\beta^*$  is the generative parameter associated with the branching process and  $\widehat{\pi}$  is the policy followed during data collection. Equations 11 and 12 imply the following inequality,

$$\begin{aligned} \mathbb{P}_{\beta^*, \widehat{\pi}} \left( \|\widehat{\beta}_J - \beta^*\| > \epsilon \right) &\leq \mathbb{P}_{\beta^*, \widehat{\pi}} \left\{ \sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} e_i(\beta) \geq \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} e_i(\beta^*) \right\} \\ &= \mathbb{P}_{\beta^*, \widehat{\pi}} \left[ \sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i l_i(\beta) - \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i l_i(\beta^*) \geq 0 \right] \\ &= \mathbb{P}_{\beta^*, \widehat{\pi}} \left[ \sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \left\{ \left( \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \left[ W_i l_i(\beta) - \mathbb{E}_{\beta^*, \widehat{\pi}} \{W_i l_i(\beta) | \mathcal{F}_{j-1}\} + \right. \right. \right. \right. \\ &\quad \left. \left. \left. \mathbb{E}_{\beta^*, \widehat{\pi}} \{W_i l_i(\beta) | \mathcal{F}_{j-1}\} \right] \right) \right. \\ &\quad \left. - \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i l_i(\beta^*) - \mathbb{E}_{\beta^*, \widehat{\pi}} \{W_i l_i(\beta^*) | \mathcal{F}_{j-1}\} + \mathbb{E}_{\beta^*, \widehat{\pi}} \{W_i l_i(\beta^*) | \mathcal{F}_{j-1}\}] \geq 0 \right\} \right]. \end{aligned}$$

By the triangle inequality,

$$\begin{aligned}
&\leq \mathbb{P}_{\beta^*, \hat{\pi}} \left\{ \underbrace{\sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \left( \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i l_i(\beta) - \mathbb{E}_{\beta^*, \hat{\pi}} \{W_i l_i(\beta) | \mathcal{F}_{j-1}\}] \right)}_{(a)} \right. \\
&\quad + \underbrace{\sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \left\{ \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} (\mathbb{E}_{\beta^*, \hat{\pi}} [W_i \{l_i(\beta) - l_i(\beta^*)\} | \mathcal{F}_{j-1}]) \right\}}_{(b)} \\
&\quad \left. - \underbrace{\frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i l_i(\beta^*) - \mathbb{E}_{\beta^*, \hat{\pi}} \{W_i l_i(\beta^*) | \mathcal{F}_{j-1}\}] \geq 0 \right\}}_{(c)}. \tag{13}
\end{aligned}$$

We first analyze quantity (c), which is a martingale by construction. By Assumption 8 (the moments condition), we know that there exists  $B_{l^2} \in \mathbb{R}^+$  such that  $\mathbb{E}_{\beta^*, \hat{\pi}} \{l_i^2(\beta^*) | \mathcal{F}_{j-1}\} \leq B_{l^2}$ . Consequently, by Theorem 8.4,  $\lim_{J \rightarrow \infty} |(c)| \rightarrow 0$  almost surely. We use a uniform martingale strong law of large numbers, Lemma 8.2, to prove that  $\lim_{J \rightarrow \infty} |(a)| \rightarrow 0$  almost surely.

Therefore, it is sufficient for consistency to show that there exists  $\delta' > 0$  such that

$$\lim_{J \rightarrow \infty} \sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \left\{ \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} (\mathbb{E}_{\beta^*, \hat{\pi}} [W_i \{l_i(\beta) - l_i(\beta^*)\} | \mathcal{F}_{j-1}]) \right\} \leq -\delta' \text{ a.s.}$$

By the law of iterated expectations,

$$\begin{aligned}
&\sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \left\{ \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} (\mathbb{E}_{\beta^*, \hat{\pi}} [W_i \{l_i(\beta) - l_i(\beta^*)\} | \mathcal{F}_{j-1}]) \right\} = \\
&\sup_{\beta \in \mathcal{B}: \|\beta - \beta^*\| > \epsilon} \left\{ \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \left( \int_{\mathcal{A}_i} \mathbb{P}(\mathbf{A}_i = \mathbf{a} | \mathcal{F}_{j-1}) \times \right. \right. \\
&\quad \left. \left. W_i \mathbb{E}_{\beta^*} \{l_i(\beta) - l_i(\beta^*) | \mathbf{A}_i = \mathbf{a}, \mathcal{F}_{j-1}\} d\mathbf{a} \right) \right\}. \quad (*)
\end{aligned}$$

First, by the restriction of information explained in Section 2, we know that

$\mathbb{P}(\mathbf{A}_i = \mathbf{a} | \mathcal{F}_{j-1}) = \mathbb{P}_{\hat{\pi}}(\mathbf{a} | H_i)$ . For  $\rho_{\min}, \rho_{\max} \in \mathbb{R}$ , we know that  $\rho_{\min} \leq W_i \leq \rho_{\max}$  by Assumption 4. Because  $\mathbb{E}_{\beta^*} \{l_i(\beta) - l_i(\beta^*) | \mathbf{A}_i = \mathbf{a}_i, \mathcal{F}_{j-1}\} \leq 0$  with probability 1, we

find that

$$\begin{aligned}
(*) &\leq \sup_{\boldsymbol{\beta} \in \mathcal{B}: \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\| > \epsilon} \left\{ \frac{1}{\rho_{\max} \bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \left( \int_{\mathcal{A}_i} \mathbb{P}_{\tilde{\boldsymbol{\pi}}}(\mathbf{a} \mid \mathbf{H}_i) \times \right. \right. \\
&\quad \left. \left. W_i^2 \mathbb{E}_{\boldsymbol{\beta}^*} \{l_i(\boldsymbol{\beta}) - l_i(\boldsymbol{\beta}^*) \mid \mathbf{A}_i = \mathbf{a}, \mathcal{F}_{j-1}\} d\mathbf{a} \right) \right\} \\
&= \sup_{\boldsymbol{\beta} \in \mathcal{B}: \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\| > \epsilon} \left\{ \frac{1}{\rho_{\max} \bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \left( \int_{\mathcal{A}_i} \mathbb{P}_{\tilde{\boldsymbol{\pi}}}(\mathbf{a} \mid \mathbf{H}_i) \times \right. \right. \\
&\quad \left. \left. \mathbb{E}_{\boldsymbol{\beta}^*} \{l_i(\boldsymbol{\beta}) - l_i(\boldsymbol{\beta}^*) \mid \mathbf{A}_i = \mathbf{a}, \mathcal{F}_{j-1}\} d\mathbf{a} \right) \right\} \\
&= \sup_{\boldsymbol{\beta} \in \mathcal{B}: \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\| > \epsilon} \left[ \frac{1}{\rho_{\max} \bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{l_i(\boldsymbol{\beta}) - l_i(\boldsymbol{\beta}^*) \mid \mathcal{F}_{j-1}\} \right].
\end{aligned}$$

From Assumption 10, it follows that

$$\frac{1}{\rho_{\max}} \lim_{J \rightarrow \infty} \sup_{\boldsymbol{\beta} \in \mathcal{B}: \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\| > \epsilon} \left[ \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{l_i(\boldsymbol{\beta}) - l_i(\boldsymbol{\beta}^*) \mid \mathcal{F}_{j-1}\} \right] \leq -\delta \frac{1}{\rho_{\max}} \text{ a.s.}$$

Thus, Equation 13 holds with  $\delta' \triangleq \delta \frac{1}{\rho_{\max}}$ .

### 8.3 Convergence Rate

First, we will prove that

$$\Sigma_J^{1/2} (\hat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^*) = O_p(1). \tag{14}$$

Define the weighted log-likelihood at  $\boldsymbol{\beta} \in \mathcal{B}$  as

$$\mathcal{M}_J(\boldsymbol{\beta}) \triangleq \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i l_i(\boldsymbol{\beta}).$$

We begin with a Taylor series expansion of  $\dot{\mathcal{M}}_J$  between  $\hat{\boldsymbol{\beta}}_J$  and  $\boldsymbol{\beta}^*$ ,

$$0 = \dot{\mathcal{M}}_J(\boldsymbol{\beta}^*) + \ddot{\mathcal{M}}_J(\bar{\boldsymbol{\beta}}^J) (\hat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^*),$$

where  $\bar{\beta}^J$  is between  $\widehat{\beta}_J$  and  $\beta^*$ . Multiplying and dividing by  $\Sigma_J^{-1/2}$ , we find that

$$-\Sigma_J^{-1/2} \dot{\mathcal{M}}_J(\beta^*) = \Sigma_J^{-1/2} \left\{ \ddot{\mathcal{M}}_J(\bar{\beta}^J) \right\} \Sigma_J^{-1/2} \Sigma_J^{1/2} \left( \widehat{\beta}_J - \beta^* \right).$$

If we show that

$$\Sigma_J^{-1/2} \dot{\mathcal{M}}_J(\beta^*) = O_p(1), \quad (15)$$

and

$$\sigma_{\min} \left( \Sigma_J^{-1/2} \ddot{\mathcal{M}}_J(\bar{\beta}^J) \Sigma_J^{-1/2} \right)^{-1} = O_p(1), \quad (16)$$

then Equation 14 follows. Consequently, we will divide the proof of Equation 14 into proofs of Equations 15 and 16. This proof only relies on Assumptions 1-8 and 11.

### 8.3.1 Proof of Equation 15

We first prove that

$$\Sigma_J^{-1/2} \dot{\mathcal{M}}_J(\beta^*) = O_p(1).$$

Note that

$$\Sigma_J^{-1/2} \dot{\mathcal{M}}_J(\beta^*) = \Sigma_J^{-1/2} \left\{ \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\beta^*) \right\}.$$

Define  $\mathbf{c}_g \in \mathbb{R}^k$  as a standard basis vector with 1 in the  $g^{\text{th}}$  position and 0 elsewhere. We know that for any  $g \in \{1, 2, \dots, k\}$ ,

$$\left\{ \mathbf{c}_g^\top \Sigma_J^{-1/2} \sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\beta^*) \right\}_{j=1}^J$$

is a martingale difference sequence with respect to fields  $\{\mathcal{F}_j\}_{j=1}^J$ . For any  $j \in \mathbb{N}$  and  $\boldsymbol{\pi} \in \Pi$ ,

$$\begin{aligned} \mathbb{E}_{\boldsymbol{\beta}^*, \boldsymbol{\pi}} \left\{ \mathbf{c}_g^\top \sum_{i \in \mathcal{E}_{j-1}} W_i l_i(\boldsymbol{\beta}^*) | \mathcal{F}_{j-1} \right\} &= \mathbf{c}_g^\top \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \boldsymbol{\pi}} \left\{ W_i l_i(\boldsymbol{\beta}^*) | \mathcal{F}_{j-1} \right\} \\ &= \mathbb{E}_{\boldsymbol{\beta}^*, \boldsymbol{\pi}} \left[ \mathbf{c}_g^\top \sum_{i \in \mathcal{E}_{j-1}} W_i \mathbb{E}_{\boldsymbol{\beta}^*} \left\{ l_i(\boldsymbol{\beta}^*) | \mathcal{F}_{j-1}, \mathbf{A}_i \right\} | \mathcal{F}_{j-1} \right] \\ &= 0. \end{aligned}$$

We now apply Theorem 8.2 (Theorem 2.15 from Hall and Heyde (2014)) to this martingale difference sequence. By this theorem,

$$\sum_{j=1}^J \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left[ \left\{ \mathbf{c}_g^\top \boldsymbol{\Sigma}_J^{-1/2} \sum_{i \in \mathcal{E}_{j-1}} W_i l_i(\boldsymbol{\beta}^*) \right\}^2 | \mathcal{F}_{j-1} \right] \xrightarrow{p} \mathbf{c}^\top U \mathbf{c}.$$

is sufficient for

$$\sum_{j=1}^J \mathbf{c}_g^\top \boldsymbol{\Sigma}_J^{-1/2} \sum_{i \in \mathcal{E}_{j-1}} W_i l_i(\boldsymbol{\beta}^*)$$

to converge almost surely. We find that

$$\begin{aligned}
& \sum_{j=1}^J \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left[ \left\{ \mathbf{c}_g^\top \Sigma_J^{-1/2} \sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\boldsymbol{\beta}^*) \right\}^2 \middle| \mathcal{F}_{j-1} \right] \\
&= \sum_{j=1}^J \left[ \left\{ \mathbf{c}_g^\top \Sigma_J^{-1/2} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left\{ \left( \sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\boldsymbol{\beta}^*) \right) \left( \sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\boldsymbol{\beta}^*)^\top \right) \middle| \mathcal{F}_{j-1} \right\} \Sigma_J^{-1/2} \mathbf{c}_g \right\} \right] \\
&\stackrel{(a)}{=} \mathbf{c}_g^\top \Sigma_J^{-1/2} \left\{ \sum_{j=1}^J \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left[ \sum_{i \in \mathcal{E}_{j-1}} \left\{ W_i^2 \dot{l}_i(\boldsymbol{\beta}^*) \dot{l}_i(\boldsymbol{\beta}^*)^\top \right\} \middle| \mathcal{F}_{j-1} \right] \right\} \Sigma_J^{-1/2} \mathbf{c}_g \\
&\stackrel{(b)}{=} \mathbf{c}_g^\top \Sigma_J^{-1/2} \left\{ \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left\{ \frac{\mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i)}{\mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i)} \dot{l}_i(\boldsymbol{\beta}^*) \dot{l}_i(\boldsymbol{\beta}^*)^\top \middle| \mathcal{F}_{j-1} \right\} \right\} \Sigma_J^{-1/2} \mathbf{c}_g \\
&= \mathbf{c}_g^\top \Sigma_J^{-1/2} \left\{ \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \int \frac{\mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i | \mathbf{H}_i)}{\mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i | \mathbf{H}_i)} \dot{l}_i(\boldsymbol{\beta}^*) \dot{l}_i(\boldsymbol{\beta}^*)^\top \times \right. \\
&\quad \left. f(\mathbf{D}_i | \mathbf{A}_i, \mathcal{F}_{j-1}; \boldsymbol{\beta}^*) \mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i | \mathcal{F}_{j-1}) d\nu \right\} \Sigma_J^{-1/2} \mathbf{c}_g \\
&\stackrel{(c)}{=} \mathbf{c}_g^\top \Sigma_J^{-1/2} \left\{ \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \int \frac{\mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i | \mathbf{H}_i)}{\mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i | \mathbf{H}_i)} \dot{l}_i(\boldsymbol{\beta}^*) \dot{l}_i(\boldsymbol{\beta}^*)^\top \times \right. \\
&\quad \left. f(\mathbf{D}_i | \mathbf{A}_i, \mathcal{F}_{j-1}; \boldsymbol{\beta}^*) \mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i | \mathbf{H}_i) d\nu \right\} \Sigma_J^{-1/2} \mathbf{c}_g \\
&= \mathbf{c}_g^\top \Sigma_J^{-1/2} \left\{ \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \int \dot{l}_i(\boldsymbol{\beta}^*) \dot{l}_i(\boldsymbol{\beta}^*)^\top f(\mathbf{D}_i | \mathbf{A}_i, \mathcal{F}_{j-1}; \boldsymbol{\beta}^*) \mathbb{P}_{\hat{\boldsymbol{\pi}}}(\mathbf{A}_i | \mathbf{H}_i) d\nu \right\} \Sigma_J^{-1/2} \mathbf{c}_g \\
&= \mathbf{c}_g^\top \Sigma_J^{-1/2} \left[ \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left\{ \dot{l}_i(\boldsymbol{\beta}^*) \dot{l}_i(\boldsymbol{\beta}^*)^\top \middle| \mathcal{F}_{j-1} \right\} \right] \Sigma_J^{-1/2} \mathbf{c}_g \\
&\stackrel{(d)}{=} \mathbf{c}_g^\top \Sigma_J^{-1/2} \eta_J \Sigma_J^{-1/2} \mathbf{c}_g.
\end{aligned}$$

Equality (a) follows from the law of iterated expectations applied to the cross terms of the sum (as explained in the proof of Theorem 8.4); equality (b) follows from the definition of  $W_i$ ; equality (c) follows from the definition of historical information,  $\mathbf{H}_i$ ; and equality (d) follows from the definition of  $\eta_J$ . By Assumption 11,

$$\mathbf{c}_g^\top \Sigma_J^{-1/2} \eta_J \Sigma_J^{-1/2} \mathbf{c}_g \xrightarrow{p} \mathbf{c}_g^\top U \mathbf{c}_g.$$

By Theorem 8.2, this implies that

$$\mathbf{c}_g^\top \Sigma_J^{-1/2} \sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\boldsymbol{\beta}^*)$$

converges a.s. to a random variable, which we label  $V_g$  for  $g \in \{1, 2, \dots, k\}$ . Labeling  $\mathbf{V} = (V_1, V_2, \dots, V_k)$ , we observe that

$$\lim_{J \rightarrow \infty} \Sigma_J^{-1/2} \sum_{i \in \mathcal{E}_{j-1}} W_i \dot{l}_i(\boldsymbol{\beta}^*) \xrightarrow{p} \mathbf{V}.$$

This verifies equation 15.

### 8.3.2 Proof of Equation 16

First, note that the differentiability and moment conditions of the log-likelihood (Assumptions 7 and 8) imply that

$$\eta_J = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left\{ \dot{l}_i(\boldsymbol{\beta}^*) \dot{l}_i(\boldsymbol{\beta}^*)^\top \mid \mathcal{F}_{j-1} \right\} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left\{ -\ddot{l}_i(\boldsymbol{\beta}^*) \mid \mathcal{F}_{j-1} \right\}.$$

Define

$$\alpha_J \triangleq \sum_{j=1}^J -\mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left\{ \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}^*) \mid \mathcal{F}_{j-1} \right\}.$$

By Lemmas 11.2 and 11.3 and the fact that  $\rho_{\min} \leq W_i \leq \rho_{\max}$ ,

$$\begin{aligned}
\sigma_{\min} \left( \Sigma_J^{-1/2} \left\{ -\ddot{\mathcal{M}}_J(\bar{\beta}^J) \right\} \Sigma_J^{-1/2} \right) &\geq \sigma_{\min} \left( \Sigma_J^{-1/2} \left\{ -\ddot{\mathcal{M}}_J(\bar{\beta}^J) + \ddot{\mathcal{M}}_J(\beta^*) \right\} \Sigma_J^{-1/2} \right) + \\
&\quad \sigma_{\min} \left( \Sigma_J^{-1/2} \left\{ -\ddot{\mathcal{M}}_J(\beta^*) - \alpha_J \right\} \Sigma_J^{-1/2} \right) + \\
&\quad \sigma_{\min} \left( \Sigma_J^{-1/2} \alpha_J \Sigma_J^{-1/2} \right) \\
&\geq \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} \sigma_{\min} \left( \frac{\ddot{\mathcal{M}}_J(\bar{\beta}^J) - \ddot{\mathcal{M}}_J(\beta^*)}{\bar{\kappa}_J} \right) \times \\
&\quad \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} + \\
&\quad \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} \sigma_{\min} \left\{ \frac{-\ddot{\mathcal{M}}_J(\beta^*) - \alpha_J}{\bar{\kappa}_J} \right\} \times \\
&\quad \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} + \\
&\quad \sigma_{\min} \left( \Sigma_J^{-1/2} \alpha_J \Sigma_J^{-1/2} \right) \\
&\stackrel{(a)}{\geq} \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} \sigma_{\min} \left\{ \frac{\ddot{\mathcal{M}}_J(\bar{\beta}^J) - \ddot{\mathcal{M}}_J(\beta^*)}{\bar{\kappa}_J} \right\} \times \\
&\quad \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} + \\
&\quad \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} \sigma_{\min} \left\{ \frac{-\ddot{\mathcal{M}}_J(\beta^*) - \alpha_J}{\bar{\kappa}_J} \right\} \times \\
&\quad \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} + \\
&\quad \frac{1}{\rho_{\max}} \sigma_{\min} \left( \Sigma_J^{-1/2} \eta_J \Sigma_J^{-1/2} \right).
\end{aligned}$$

By Assumption 11,  $\sigma_{\min} \left( \Sigma_J^{-1/2} \eta_J \Sigma_J^{-1/2} \right)^{-1} = O_p(1)$  as  $J \rightarrow \infty$ . Inequality (a) will be verified by Property (4) below. In sum, it is sufficient to establish four properties:

1.  $\sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} \leq \left\| (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\|_2 = O_p(1)$
2.  $\left\| \left\{ \ddot{\mathcal{M}}_J(\beta^*) - \ddot{\mathcal{M}}_J(\bar{\beta}^J) \right\} / \bar{\kappa}_J \right\|_2 = o_p(1)$ .
3.  $\left\| \left\{ -\ddot{\mathcal{M}}_J(\beta^*) - \alpha_J \right\} / \bar{\kappa}_J \right\|_2 = o_p(1)$
4.  $\alpha_J \succeq \frac{1}{\rho_{\max}} \eta_J$

Properties (1)-(3) ensure that

$$\begin{aligned} \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} \sigma_{\min} \left\{ \frac{\ddot{\mathcal{M}}_J(\bar{\beta}^J) - \ddot{\mathcal{M}}_J(\beta^*)}{\bar{\kappa}_J} \right\} \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} &= o_p(1), \text{ and} \\ \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} \sigma_{\min} \left\{ \frac{-\ddot{\mathcal{M}}_J(\beta^*) - \alpha_J}{\bar{\kappa}_J} \right\} \sigma_{\min} \left\{ (\Sigma_J / \bar{\kappa}_J)^{-1/2} \right\} &= o_p(1). \end{aligned}$$

This allows us to conclude that

$$\begin{aligned} \sigma_{\min} \left( \Sigma_J^{-1/2} \left\{ -\ddot{\mathcal{M}}_J(\bar{\beta}^J) \right\} \Sigma_J^{-1/2} \right)^{-1} &\leq \left[ \frac{1}{\rho_{\max}} \sigma_{\min} \left\{ \Sigma_J^{-1/2} \eta_J \Sigma_J^{-1/2} \right\} + o_p(1) + o_p(1) \right]^{-1} \\ &= O_p(1). \end{aligned}$$

**We begin with Property (1).** By Assumption 12, there exists  $\epsilon > 0$  such that

$$\begin{aligned} \sigma_{\min}(\eta_J / \bar{\kappa}_J) &\geq \epsilon + o_p(1) \Rightarrow \\ \bar{\kappa}_J \|\eta_J^{-1}\|_2 &\leq 1/\epsilon + o_p(1). \end{aligned}$$

By Assumption 11,  $\left\| \Sigma_J^{-1/2} \eta_J \Sigma_J^{-1/2} \right\|_2 = O_p(1)$ . Consequently, by the sub-multiplicativity of the spectral norm,

$$\begin{aligned} \bar{\kappa}_J \|\Sigma_J^{-1}\|_2 &= \bar{\kappa}_J \|\Sigma_J^{-1} \eta_J \eta_J^{-1}\|_2 \\ &\leq \bar{\kappa}_J \|\Sigma_J^{-1} \eta_J\|_2 \|\eta_J^{-1}\|_2 \\ &\leq O_p(1) \left\{ \frac{1}{\epsilon} + o_p(1) \right\} \end{aligned}$$

as  $J \rightarrow \infty$ . Note that the last equality follows because the eigenvalues of  $\Sigma_J^{-1} \eta_J$  are the same as  $\Sigma_J^{-1/2} \eta_J \Sigma_J^{-1/2}$ .

**We now prove Property (2).** We analyze

$$\begin{aligned} \left\| \frac{\ddot{\mathcal{M}}_J(\beta^*) - \ddot{\mathcal{M}}_J(\bar{\beta}^J)}{\bar{\kappa}_J} \right\|_2 &= \left\| \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \left\{ \ddot{l}_i(\beta^*) - \ddot{l}_i(\bar{\beta}^J) \right\}}{\bar{\kappa}_J} \right\|_2 \\ &\leq \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \|\ddot{l}_i(\beta^*) - \ddot{l}_i(\bar{\beta}^J)\|_2}{\bar{\kappa}_J}, \end{aligned}$$

where the last inequality follows from the triangle inequality. We wish to show that for any  $\epsilon > 0$ ,

$$\lim_{J \rightarrow \infty} \mathbb{P} \left( \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \|\ddot{l}_i(\boldsymbol{\beta}^*) - \ddot{l}_i(\bar{\boldsymbol{\beta}}^J)\|_2}{\bar{\kappa}_J} > \epsilon \right) = 0. \quad (17)$$

Recall that Assumption 13 implies there exists an  $\epsilon_j > 0$  and a function  $f : \mathcal{D} \rightarrow \mathbb{R}$  such that for all  $0 < \epsilon^* \leq \epsilon_j$ , there exists  $\delta_{\epsilon^*}$  such that

$$\sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}, \boldsymbol{\beta} \in \mathcal{B}: \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\|_2 \leq \delta_{\epsilon^*}} \|\ddot{l}_i(\boldsymbol{\beta}) - \ddot{l}_i(\boldsymbol{\beta}^*)\|_2 \leq f(\mathbf{D}_i) \epsilon^* \quad \text{a.s.},$$

and  $\mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{f^2(\mathbf{D}_i) | \mathcal{F}_{j-1}\}$  is bounded almost surely. Because  $\bar{\boldsymbol{\beta}}^J$  is consistent for  $\boldsymbol{\beta}^*$ , we know that for any  $\delta_{\epsilon^*}$ ,  $\lim_{J \rightarrow \infty} \|\bar{\boldsymbol{\beta}}^J - \boldsymbol{\beta}^*\| \leq \delta_{\epsilon^*}$  a.s. Consequently,

$$\begin{aligned} & \lim_{J \rightarrow \infty} \mathbb{P} \left( \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \|\ddot{l}_i(\boldsymbol{\beta}^*) - \ddot{l}_i(\bar{\boldsymbol{\beta}}^J)\|_2}{\bar{\kappa}_J} > \epsilon \right) \\ & \leq \lim_{J \rightarrow \infty} \mathbb{P} \left( \epsilon^* \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i f(\mathbf{D}_i)}{\bar{\kappa}_J} > \epsilon \right). \end{aligned}$$

We now analyze the quantity

$$\begin{aligned} \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i f(\mathbf{D}_i) &= \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i f(\mathbf{D}_i) - \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{W_i f(\mathbf{D}_i) | \mathcal{F}_{j-1}\} \\ &+ \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{W_i f(\mathbf{D}_i) | \mathcal{F}_{j-1}\}. \end{aligned}$$

We know that

$$\frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i f(\mathbf{D}_i) - \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{W_i f(\mathbf{D}_i) | \mathcal{F}_{j-1}\} = o_p(1) \quad (18)$$

by Theorem 8.4 because Assumption 13 states that  $\mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{f^2(\mathbf{D}_i) | \mathcal{F}_{j-1}\}$  is bounded almost surely. Therefore,

$$\frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i f(\mathbf{D}_i) = o_p(1) + \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{W_i f(\mathbf{D}_i) | \mathcal{F}_{j-1}\}.$$

By Assumption 4, we know that  $\rho_{\min} \leq W_i \leq \rho_{\max}$ . Because  $f(\mathbf{D}_i)$  is a positive function, we can conclude that

$$\begin{aligned} \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i f(\mathbf{D}_i) &\leq o_p(1) + \frac{1}{\rho_{\min} \bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{W_i^2 f(\mathbf{D}_i) | \mathcal{F}_{j-1}\} \\ &\leq o_p(1) + B_f / \rho_{\min}. \end{aligned}$$

Defining  $\epsilon^* \triangleq \epsilon \rho_{\min} / (2B_f)$ , we find that

$$\begin{aligned} &\lim_{J \rightarrow \infty} \mathbb{P} \left( \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \|\ddot{l}(\boldsymbol{\beta}^*) - \ddot{l}(\bar{\boldsymbol{\beta}}^J)\|_2}{\bar{\kappa}_J} > \epsilon \right) \\ &\leq \lim_{J \rightarrow \infty} \mathbb{P} \left( \epsilon \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i f(\mathbf{D}_i)}{\bar{\kappa}_J} > \epsilon \right) \\ &\leq \lim_{J \rightarrow \infty} \mathbb{P} \left( \epsilon \rho_{\min} / (2B_f) \right. \\ &\quad \left. \{o_p(1) + B_f / \rho_{\min}\} > \epsilon \right) \\ &\leq \lim_{J \rightarrow \infty} \mathbb{P} (o_p(1) > \epsilon/2) \rightarrow 0. \end{aligned}$$

Consequently, Property (2) is satisfied.

**We now prove Property (3).** We verify the following condition,

$$\lim_{J \rightarrow \infty} \left\| \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} -W_i \ddot{l}(\boldsymbol{\beta}^*) - \sum_{j=1}^J \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left\{ \sum_{i \in \mathcal{E}_{j-1}} -W_i \ddot{l}(\boldsymbol{\beta}^*) \mid \mathcal{F}_{j-1} \right\}}{\bar{\kappa}_J} \right\|_2 = 0 \text{ a.s.}$$

By Assumption 8 (the moments assumption) and Assumption 4, for any  $p, q \in [k]$ , there exists  $B_{j_2}$  such that for any  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ ,

$$\mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left\{ [\ddot{l}_i(\boldsymbol{\beta}^*)]_{p,q}^2 \mid \mathcal{F}_{j-1} \right\} \leq B_{j_2} \text{ a.s.}$$

Consequently, we know that

$$\left\| \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} -W_i \ddot{l}(\boldsymbol{\beta}^*) - \alpha_J}{\bar{\kappa}_J} \right\| = o_p(1)$$

by Theorem 8.4 (component-wise).

**Lastly, we prove Property (4).** First, we find that

$$\begin{aligned}\alpha_J &\triangleq - \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \hat{\pi}} \left\{ W_i \ddot{l}(\beta^*) \mid \mathcal{F}_{j-1} \right\} \\ &= \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \hat{\pi}} \left[ W_i \mathbb{E}_{\beta^*} \left\{ -\ddot{l}(\beta^*) \mid \mathbf{A}_i, \mathcal{F}_{j-1} \right\} \mid \mathcal{F}_{j-1} \right].\end{aligned}$$

We know  $\mathbb{E}_{\beta^*} \left\{ -\ddot{l}(\beta^*) \mid \mathbf{A}_i, \mathcal{F}_{j-1} \right\} \succeq 0$  (because this is equivalent to the variance of the score function), and therefore we can express

$$\begin{aligned}&\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \hat{\pi}} \left[ W_i \mathbb{E}_{\beta^*} \left\{ -\ddot{l}(\beta^*) \mid \mathbf{A}_i, \mathcal{F}_{j-1} \right\} \mid \mathcal{F}_{j-1} \right] \\ &\succeq \frac{1}{\rho_{\max}} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \hat{\pi}} \left[ W_i^2 \mathbb{E}_{\beta^*} \left\{ -\ddot{l}(\beta^*) \mid \mathbf{A}_i, \mathcal{F}_{j-1} \right\} \mid \mathcal{F}_{j-1} \right] \\ &= \frac{1}{\rho_{\max}} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \hat{\pi}} \left[ \mathbb{E}_{\beta^*} \left\{ -\ddot{l}(\beta^*) \mid \mathbf{A}_i, \mathcal{F}_{j-1} \right\} \mid \mathcal{F}_{j-1} \right] \\ &= -\frac{1}{\rho_{\max}} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \hat{\pi}} \left\{ \ddot{l}(\beta^*) \mid \mathcal{F}_{j-1} \right\} \\ &= \frac{1}{\rho_{\max}} \eta_J.\end{aligned}$$

This satisfies Property (4). We have verified Equation 16.

Equation 14,

$$\Sigma_J^{1/2} \left( \hat{\beta}_J - \beta^* \right) = O_p(1),$$

follows from Equations 15 and 16. Equation 14 is used to conclude that

$$\left( \hat{\beta}_J - \beta^* \right) = O_p(1/\sqrt{\kappa_J})$$

by showing

$$\begin{aligned}
\left\| \sqrt{\bar{\kappa}_J} \left( \widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^* \right) \right\|_2 &= \left\| \bar{\kappa}_J^{-1/2} \Sigma_J^{-1/2} \Sigma_J^{1/2} \left( \widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^* \right) \right\|_2 \\
&\leq \left\| \bar{\kappa}_J^{-1/2} \Sigma_J^{-1/2} \right\|_2 \left\| \Sigma_J^{1/2} \left( \widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^* \right) \right\|_2 \\
&= O_p(1).
\end{aligned}$$

The last equality follows from Property (1) and Equation 14.

**Lemma 8.2.** *Let  $e_i(\boldsymbol{\beta}) \triangleq W_i l_i(\boldsymbol{\beta})$ . Under Assumptions 1-13,*

$$\sup_{\boldsymbol{\beta} \in \mathcal{B}} \left( \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [e_i(\boldsymbol{\beta}) - \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{e_i(\boldsymbol{\beta}) | \mathcal{F}_{j-1}\}] \right) = o_p(1).$$

Lemma 8.2 is a martingale uniform law of large numbers, and its proof mirrors Lemma 2 of Zhang et al. (2021).

*Proof.* Like Zhang et al. (2021), we begin by establishing a finite bracketing number for the log-likelihood function based on Assumption 9 and the fact that  $\mathcal{B}$  is bounded.

**Finite Bracketing Number.** Let  $\delta > 0$ . We construct a set  $B_\delta$  that is made up of pairs of functions  $(b, u)$  and satisfies the following criteria.

1. We denote  $l_i(\boldsymbol{\beta})$  as  $l(\boldsymbol{\beta}, \mathbf{D}_i)$  to emphasize its dependence on data  $\mathbf{D}_i \in \mathcal{D}$ . We show that for any  $\boldsymbol{\beta} \in \mathcal{B}$ , we can find functions  $(b, u) \in B_\delta$  such that  $b(\mathbf{D}_i) \leq l(\boldsymbol{\beta}, \mathbf{D}_i) \leq u(\mathbf{D}_i)$  and

$$\sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{u(\mathbf{D}_i) - b(\mathbf{D}_i) | \mathcal{F}_{j-1}\} \leq \delta \quad \text{a.s.}$$

2. There are a finite number of pairs in  $B_\delta$ ,  $|B_\delta| < \infty$ .
3. For any  $(b, u) \in B_\delta$ , there exists  $m_g < \infty$  that does not depend on  $\delta$  such that

$$\sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{b(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \leq m_g \quad \text{a.s. and}$$

$$\sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{u(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \leq m_g \quad \text{a.s.}$$

We now construct  $B_\delta$ . Create a grid over  $\mathcal{B}$  with a meshwidth of  $\lambda > 0$ , and let the points in this grid be the set  $G_\lambda \subseteq \mathcal{B}$ . By construction, this implies that for any  $\boldsymbol{\beta} \in \mathcal{B}$ , we can find  $\boldsymbol{\beta}' \in G_\lambda$  such that  $\|\boldsymbol{\beta}' - \boldsymbol{\beta}\| \leq \lambda$ .

By Assumption 9, we know that for any  $\boldsymbol{\beta}, \boldsymbol{\beta}' \in \mathcal{B}$ ,  $j \in \mathbb{N}$ , and  $i \in \mathcal{E}_{j-1}$ , there exists function  $g$  and constant  $m_g < \infty$  such that  $|l(\boldsymbol{\beta}, \mathbf{D}_i) - l(\boldsymbol{\beta}', \mathbf{D}_i)| \leq g(\mathbf{D}_i)\|\boldsymbol{\beta} - \boldsymbol{\beta}'\|_2$  and

$$\mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \leq m_g \quad \text{a.s.} \quad (19)$$

We specify  $B_\delta = \{(l(\boldsymbol{\beta}, \mathbf{D}_i) - \lambda g(\mathbf{D}_i), l(\boldsymbol{\beta}, \mathbf{D}_i) + \lambda g(\mathbf{D}_i)) : \boldsymbol{\beta} \in G_\lambda\}$  and show that this set satisfies properties (1)-(3) for a certain value of  $\lambda$ . Note that because  $\mathcal{B}$  is bounded, property (2) is satisfied because the number of points in  $G_\lambda$  is finite. To show that (1) holds for  $B_\delta$ , recall that for any  $\boldsymbol{\beta} \in \mathcal{B}$ , we can find  $\boldsymbol{\beta}' \in G_\lambda$  such that  $\|\boldsymbol{\beta} - \boldsymbol{\beta}'\| \leq \lambda$ . Define  $\boldsymbol{\beta}'_\lambda(\boldsymbol{\beta}) \triangleq \min_{\boldsymbol{\beta}' \in G_\lambda} \|\boldsymbol{\beta} - \boldsymbol{\beta}'\|$ . By Assumption 9, this implies that

$$|l(\boldsymbol{\beta}, \mathbf{D}_i) - l\{\boldsymbol{\beta}'_\lambda(\boldsymbol{\beta}), \mathbf{D}_i\}| \leq g(\mathbf{D}_i)\|\boldsymbol{\beta} - \boldsymbol{\beta}'_\lambda(\boldsymbol{\beta})\|_2 \leq g(\mathbf{D}_i)\lambda.$$

Therefore,

$$l\{\boldsymbol{\beta}'_\lambda(\boldsymbol{\beta}), \mathbf{D}_i\} - \lambda g(\mathbf{D}_i) \leq l(\boldsymbol{\beta}, \mathbf{D}_i) \leq l\{\boldsymbol{\beta}'_\lambda(\boldsymbol{\beta}), \mathbf{D}_i\} + \lambda g(\mathbf{D}_i),$$

implying that there exists  $(b, u) \in B_\delta$  such that  $b(\mathbf{D}_i) \leq l(\boldsymbol{\beta}, \mathbf{D}_i) \leq u(\mathbf{D}_i)$  for all  $\mathbf{D}_i \in \mathcal{D}$ .

Additionally, note that for any  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ ,

$$\begin{aligned} & \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left[ \left| l\{\boldsymbol{\beta}'_\lambda(\boldsymbol{\beta}), \mathbf{D}_i\} + \lambda g(\mathbf{D}_i) - [l\{\boldsymbol{\beta}'_\lambda(\boldsymbol{\beta}), \mathbf{D}_i\} - \lambda g(\mathbf{D}_i)] \right| \middle| \mathcal{F}_{j-1} \right] \\ &= 2\lambda \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} (|g(\mathbf{D}_i)| | \mathcal{F}_{j-1}) \leq 2\lambda \sqrt{m_g} \quad \text{a.s.} \end{aligned}$$

The inequality above holds by Jensen's inequality,

$$\mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{|g(\mathbf{D}_i)| | \mathcal{F}_{j-1}\} \leq \sqrt{\mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\}} \quad \text{a.s.,}$$

and Equation 19. We conclude that (1) holds for our choice of  $B_\delta$  by letting the meshwidth

$$\lambda \triangleq \delta / (2\sqrt{m_g}).$$

Lastly, we show that (3) holds. Note that for any  $\beta \in \mathcal{B}$ ,

$$\begin{aligned} & \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} [\{l(\beta, \mathbf{D}_i) + \lambda g(\mathbf{D}_i)\}^2 | \mathcal{F}_{j-1}] \\ & \leq 2 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{l(\beta, \mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} + 2\lambda^2 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \quad \text{a.s.} \end{aligned}$$

by the triangle inequality and the fact that for any  $p, q \in \mathbb{R}$ ,

$$\begin{aligned} (p + q)^2 &= p^2 + 2pq + q^2 \\ &= 2p^2 + 2q^2 - p^2 + 2pq - q^2 \\ &= 2p^2 + 2q^2 - (p - q)^2 \\ &\leq 2p^2 + 2q^2. \end{aligned}$$

By the same logic,

$$\begin{aligned} \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} [\{l(\beta, \mathbf{D}_i) - \lambda g(\mathbf{D}_i)\}^2 | \mathcal{F}_{j-1}] &\leq 2 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{l(\beta, \mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \\ &\quad + 2\lambda^2 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \quad \text{a.s.} \end{aligned}$$

Because  $l(\mathbf{D}_i, \beta) = l(\mathbf{D}_i, \beta) - l(\mathbf{D}_i, \beta^*) + l(\mathbf{D}_i, \beta^*)$ , we upper bound

$$\begin{aligned} & 2 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{l(\beta, \mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} + 2\lambda^2 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \\ & \leq 4 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{l(\beta^*, \mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} + 4 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} [\{l(\beta, \mathbf{D}_i) - l(\beta^*, \mathbf{D}_i)\}^2 | \mathcal{F}_{j-1}] \\ & \quad + 2\lambda^2 \sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \quad \text{a.s.} \end{aligned}$$

Note that  $\sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \tilde{\pi}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \leq m_g$  a.s., and  $\mathbb{E}_{\beta^*, \tilde{\pi}} \{l(\beta^*, \mathbf{D}_i)^2 | \mathcal{F}_{j-1}\}$  is bounded by Assumption 8.

By Assumption 9, for any  $\boldsymbol{\beta} \in \mathcal{B}$ ,

$$\begin{aligned} |l(\boldsymbol{\beta}, \mathbf{D}_i) - l(\boldsymbol{\beta}^*, \mathbf{D}_i)| &\leq g(\mathbf{D}_i) \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\|_2 \Rightarrow \\ \{l(\boldsymbol{\beta}, \mathbf{D}_i) - l(\boldsymbol{\beta}^*, \mathbf{D}_i)\}^2 &\leq g(\mathbf{D}_i)^2 \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\|_2^2 \Rightarrow \\ \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left[ \{l(\boldsymbol{\beta}, \mathbf{D}_i) - l(\boldsymbol{\beta}^*, \mathbf{D}_i)\}^2 | \mathcal{F}_{j-1} \right] &\leq \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \|\boldsymbol{\beta} - \boldsymbol{\beta}^*\|_2^2 \text{ a.s.} \end{aligned}$$

$\|\boldsymbol{\beta} - \boldsymbol{\beta}^*\|_2^2$  is bounded because  $\mathcal{B}$  is bounded, and  $\sup_{j \in \mathbb{N}, i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{g(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \leq m_g$  a.s. We conclude that property (3) holds. We now use  $B_\delta$  in the main argument for the proof of uniform convergence.

**Main Argument:** We now show that for any  $\epsilon > 0$ ,

$$\mathbb{P}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left\{ \sup_{\boldsymbol{\beta} \in \mathcal{B}} \left( \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [e_i(\boldsymbol{\beta}) - \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{e_i(\boldsymbol{\beta}) | \mathcal{F}_{j-1}\}] \right) > \epsilon \right\} = o(1). \quad (20)$$

Let  $\delta > 0$  (we will choose  $\delta$  later). Let  $B_\delta$  be the set of pairs of functions as constructed earlier. Note that by property (1) of  $B_\delta$ , we get the following upper bound,

$$\begin{aligned} &\sup_{\boldsymbol{\beta} \in \mathcal{B}} \left( \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i l_i(\boldsymbol{\beta}) - \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{W_i l_i(\boldsymbol{\beta}) | \mathcal{F}_{j-1}\}] \right) \\ &\leq \max_{(b, u) \in B_\delta} \left( \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i u(\mathbf{D}_i) - \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{W_i b(\mathbf{D}_i) | \mathcal{F}_{j-1}\}] \right) \text{ a.s. } (*) \end{aligned}$$

By subtracting and adding  $\mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{W_i u(\mathbf{D}_i) | \mathcal{F}_{j-1}\}$ , and using the triangle inequality, we find that

$$\begin{aligned} (*) &\leq \max_{(b, u) \in B_\delta} \left[ \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} [W_i \{u(\mathbf{D}_i) - b(\mathbf{D}_i)\} | \mathcal{F}_{j-1}] \right] + \\ &\quad \max_{(b, u) \in B_\delta} \left( \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i u(\mathbf{D}_i) - \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \{W_i u(\mathbf{D}_i) | \mathcal{F}_{j-1}\}] \right) \text{ a.s.} \end{aligned}$$

By Assumption 4 and the fact that  $u(\mathbf{D}_i) - b(\mathbf{D}_i) \geq 0$ ,

$$\begin{aligned} \mathbb{E}_{\beta^*, \hat{\pi}} [W_i \{u(\mathbf{D}_i) - b(\mathbf{D}_i) | \mathcal{F}_{j-1}\}] &\leq \frac{1}{\rho_{\min}} \mathbb{E}_{\beta^*, \hat{\pi}} [W_i^2 \{u(\mathbf{D}_i) - b(\mathbf{D}_i) | \mathcal{F}_{j-1}\}] \\ &= \frac{1}{\rho_{\min}} \mathbb{E}_{\beta^*, \hat{\pi}} \{u(\mathbf{D}_i) - b(\mathbf{D}_i) | \mathcal{F}_{j-1}\} \\ &\leq \frac{1}{\rho_{\min}} \delta. \end{aligned}$$

The last inequality holds by property (1) of  $B_\delta$ . Because  $\max_{i \in [n]} a_i \leq \sum_{i=1}^n |a_i|$ ,

$$\begin{aligned} &\sup_{\beta \in \mathcal{B}} \left( \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i l_i(\beta) - \mathbb{E}_{\beta^*, \hat{\pi}} \{W_i l_i(\beta) | \mathcal{F}_{j-1}\}] \right) \\ &\leq \frac{1}{\rho_{\min}} \delta + \sum_{(b, u) \in B_\delta} \left| \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i u(\mathbf{D}_i) - \mathbb{E}_{\beta^*, \hat{\pi}} \{W_i u(\mathbf{D}_i) | \mathcal{F}_{j-1}\}] \right|. \end{aligned}$$

By property (3) of  $B_\delta$ , we know that

$$\mathbb{E}_{\beta^*, \hat{\pi}} \{W_i^2 u(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} = \mathbb{E}_{\beta^*, \hat{\pi}} \{u(\mathbf{D}_i)^2 | \mathcal{F}_{j-1}\} \leq m_g.$$

Therefore, by Theorem 8.4, for any  $(b, u) \in B_\delta$ ,

$$\left| \frac{1}{\bar{\kappa}_J} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} [W_i u(\mathbf{D}_i) - \mathbb{E}_{\beta^*, \hat{\pi}} \{W_i u(\mathbf{D}_i) | \mathcal{F}_{j-1}\}] \right| = o_p(1).$$

Because  $|B_\delta| < \infty$  by Property (2), the convergence holds for all  $(b, u) \in B_\delta$  simultaneously, so

$$(*) \leq \frac{1}{\rho_{\min}} \delta + o_p(1).$$

Equation 20 is satisfied by choosing  $\delta = \rho_{\min} \epsilon / 2$ . ■

## 9 Proof of Theorem 2.2

We establish asymptotic regret bounds for RL-RDS by proving Theorem 2.2. The following consequence of Markov's inequality will be useful.

**Lemma 9.1.** Let  $\{X_n\}_{n \geq 1}$  be a sequence of random variables where  $X_n \in \mathcal{X} \subset \mathbb{R}^p$  and  $\|\cdot\|$  is an arbitrary norm on  $\mathbb{R}^p$ . If  $\mathbb{E}\|X_n\| = O(1)$  for all  $n \geq 1$ , then  $\{X_n\}_{n \geq 1} = O_p(1)$ .

*Proof.* If  $\mathbb{E}\|X_n\| = O(1)$  for all  $n \geq 1$ ,  $\exists M > 0$  such that  $\forall n \geq 1$ ,

$$\mathbb{E}\|X_n\| \leq M.$$

We need to show that given  $\epsilon$ ,  $\exists V_\epsilon$  such that  $P(\|X_n\| \geq V_\epsilon) \leq \epsilon$  for all  $n \geq 1$ . We know that

$$\mathbb{P}(\|X_n\| \geq V_\epsilon) \leq \frac{\mathbb{E}\|X_n\|}{V_\epsilon} \leq \frac{M}{V_\epsilon}$$

by the Markov Inequality. Choosing  $V_\epsilon = M/\epsilon$  gives the desired result.  $\blacksquare$

In Assumption 6, we define  $S^n$  as the budget left when individual  $n$  is recruited, where  $S^n$  satisfies  $0 < S^n - \sum_{v=1}^n C^v < C^*$  for  $C^* \in \mathbb{R}^+$  and each  $n \in \mathbb{N}$ . We define an upper bound on the potential number of additional recruits under allocation strategy  $\boldsymbol{\pi}$  at any state  $\mathbf{h}^{n_J}$  for  $J \in \mathbb{N}$  as

$$n^*(\boldsymbol{\pi}, \mathbf{h}^k) \triangleq \inf \left\{ n : \sum_{v=k}^n C^{v*}(\boldsymbol{\pi}, \mathbf{h}^k) > C^* \right\}.$$

We assume that there exists  $S^* \in \mathbb{N}$  such that for any  $\mathbf{h}^{n_J} \in \mathcal{H}^{n_J}$ ,  $\sup_{\boldsymbol{\pi} \in \Pi} n^*(\boldsymbol{\pi}, \mathbf{h}^{n_J}) \leq S^*$  a.s.

*Proof of Theorem 2.2.*

$$\begin{aligned} V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) &= V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J) + V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J) \\ &\leq V^{n_J}(\mathbf{h}^{n_J}, \widehat{\boldsymbol{\pi}}_J; \widehat{\boldsymbol{\beta}}_J) + V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J) \\ &= V^{n_J}(\mathbf{h}^{n_J}, \widehat{\boldsymbol{\pi}}_J; \boldsymbol{\beta}^*) + V^{n_J}(\mathbf{h}^{n_J}, \widehat{\boldsymbol{\pi}}_J; \widehat{\boldsymbol{\beta}}_J) - V^{n_J}(\mathbf{h}^{n_J}, \widehat{\boldsymbol{\pi}}_J; \boldsymbol{\beta}^*) + \\ &\quad V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J) \\ &\leq V^{n_J}(\mathbf{h}^{n_J}, \widehat{\boldsymbol{\pi}}_J; \boldsymbol{\beta}^*) + 2 \sup_{\boldsymbol{\pi} \in \Pi} \left| V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J) \right|, \end{aligned}$$

where the first inequality follows from the definition of  $\widehat{\boldsymbol{\pi}}_J$ . The upper bound above does

not depend on  $\boldsymbol{\pi}$ , and so it holds for  $\boldsymbol{\pi}^{\text{opt}}$ ,

$$V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}^{\text{opt}}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \widehat{\boldsymbol{\pi}}_J; \boldsymbol{\beta}^*) \leq 2 \sup_{\boldsymbol{\pi} \in \Pi} |V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J)|.$$

We know that  $V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}^{\text{opt}}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \widehat{\boldsymbol{\pi}}_J; \boldsymbol{\beta}^*) \geq 0$  because  $\boldsymbol{\pi}^{\text{opt}} \in \arg \max_{\boldsymbol{\pi} \in \Pi} V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*)$ ,

so

$$|V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}^{\text{opt}}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \widehat{\boldsymbol{\pi}}_J; \boldsymbol{\beta}^*)| \leq 2 \sup_{\boldsymbol{\pi} \in \Pi} |V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J)|.$$

We now show that  $\sup_{\boldsymbol{\pi} \in \Pi} |V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J)| = O_p(1/\sqrt{\kappa_J})$ , which proves the desired result. We expand  $V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J)$  as follows. Recall that  $\Delta^v$  is an indicator that the budget has not been exceeded when individual  $v$  enters the study. Note that  $\exists S^*$  such that we can express

$$\begin{aligned} & V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J) = \\ & \int \left( \sum_{k=1}^{S^*} \Delta^k y^k \right) \left\{ \prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \boldsymbol{\beta}^*) \pi(\mathbf{a}^m | \mathbf{h}^m) \right\} d\lambda(\mathbf{h}^{S^*}) - \\ & \int \left( \sum_{k=1}^{S^*} \Delta^k y^k \right) \left\{ \prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \widehat{\boldsymbol{\beta}}_J) \pi(\mathbf{a}^m | \mathbf{h}^m) \right\} d\lambda(\mathbf{h}^{S^*}), \end{aligned} \quad (21)$$

where  $\mathbf{h}^1 = \mathbf{h}^{n_J}$ . Refactoring this expression and applying the definition of the log-likelihood results in

$$\begin{aligned} & V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \widehat{\boldsymbol{\beta}}_J) = \\ & \int \left( \sum_{k=1}^{S^*} \Delta^k y^k \right) \left\{ 1 - \frac{\prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \widehat{\boldsymbol{\beta}}_J) \pi(\mathbf{a}^m | \mathbf{h}^m)}{\prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \boldsymbol{\beta}^*) \pi(\mathbf{a}^m | \mathbf{h}^m)} \right\} \times \\ & \prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \boldsymbol{\beta}^*) \pi(\mathbf{a}^m | \mathbf{h}^m) d\lambda(\mathbf{h}^{S^*}) = \\ & \int \left( \sum_{k=1}^{S^*} \Delta^k y^k \right) \left[ 1 - \exp \left\{ \ell_{S^*}(\widehat{\boldsymbol{\beta}}_J) - \ell_{S^*}(\boldsymbol{\beta}^*) \right\} \right] \times \\ & \left\{ \prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \boldsymbol{\beta}^*) \pi(\mathbf{a}^m | \mathbf{h}^m) \right\} d\lambda(\mathbf{h}^{S^*}). \end{aligned}$$

By Assumption 14, we know that

$$\begin{aligned} \sqrt{\kappa_J} \left| \ell_{S^*}(\widehat{\boldsymbol{\beta}}_J) - \ell_{S^*}(\boldsymbol{\beta}^*) \right| &\leq \sqrt{\kappa_J} \sum_{v=1}^{S^*} |q^v(\widehat{\boldsymbol{\beta}}_J) - q^v(\boldsymbol{\beta}^*)| \\ &\leq \left[ \sum_{v=1}^{S^*} |e\{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v)\}| \right] \left\| \sqrt{\kappa_J} (\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^*) \right\|_2. \end{aligned}$$

We know  $\sqrt{\kappa_J}(\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^*) = O_p(1)$  by Theorem 2.1. Labeling  $e^v \triangleq e\{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v)\}$  and noting that  $e^v$  is strictly positive, we also know that for any  $\boldsymbol{\pi} \in \Pi$ ,

$$\mathbb{E}_{\boldsymbol{\beta}^*, \boldsymbol{\pi}}(|e^v|) = O(1) \quad (22)$$

by Jensen's Inequality. By Lemma 9.1, this implies that  $\sum_{v=1}^{S^*} |e^v| = O_p(1)$ . We conclude that

$$\sqrt{\kappa_J} \left| \ell_{S^*}(\widehat{\boldsymbol{\beta}}_J) - \ell_{S^*}(\boldsymbol{\beta}^*) \right| = O_p(1) \Rightarrow \left| \ell_{S^*}(\widehat{\boldsymbol{\beta}}_J) - \ell_{S^*}(\boldsymbol{\beta}^*) \right| = O_p(1/\sqrt{\kappa_J}).$$

By Taylor expansion,

$$1 - \exp \left\{ \ell_{S^*}(\widehat{\boldsymbol{\beta}}_J) - \ell_{S^*}(\boldsymbol{\beta}^*) \right\} = \ell_{S^*}(\widehat{\boldsymbol{\beta}}_J) - \ell_{S^*}(\boldsymbol{\beta}^*) + o_p(1/\sqrt{\kappa_J}). \quad (23)$$

Because  $Y^v \in [0, 1]$  for all  $v \in \mathbb{N}$ ,

$$\left| \sum_{k=1}^{S^*} Y^k \Delta^k \right| = \sum_{k=1}^{S^*} |Y^k \Delta^k| \leq S^*,$$

and

$$\begin{aligned} |(21)| &\leq S^* \times \left| \ell_{S^*}(\widehat{\boldsymbol{\beta}}_J) - \ell_{S^*}(\boldsymbol{\beta}^*) + o_p(1/\sqrt{\kappa_J}) \right| \\ &\times \left\{ \prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \boldsymbol{\beta}^*) \pi(\mathbf{a}^m | \mathbf{h}^m) \right\} d\lambda(\mathbf{h}^{S^*}). \end{aligned} \quad (24)$$

Leveraging equations (22), (23), and (24) as well as the Cauchy-Schwartz Inequality, we

know that as  $J \rightarrow \infty$

$$\begin{aligned}
|(21)| &\leq S^* \times \frac{1}{\sqrt{\bar{\kappa}_J}} \left\| \sqrt{\bar{\kappa}_J} \left\{ \hat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^* \right\} \right\|_2 \left\{ \sum_{v=1}^{S^*} |e^v| \right\} \\
&\quad \times \left\{ \prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \boldsymbol{\beta}^*) \pi(\mathbf{a}^m | \mathbf{h}^m) \right\} d\lambda(\mathbf{h}^{S^*}) + \\
& o_p(1/\sqrt{\bar{\kappa}_J}) \times \frac{1}{\sqrt{\bar{\kappa}_J}} \left\| \sqrt{\bar{\kappa}_J} \left\{ \hat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^* \right\} \right\|_2 \left\{ \sum_{v=1}^{S^*} |e^v| \right\} \\
&\quad \times \left\{ \prod_{m=1}^{S^*} f(\mathbf{h}^{m+1} | \mathbf{h}^m, \mathbf{a}^m; \boldsymbol{\beta}^*) \pi(\mathbf{a}^m | \mathbf{h}^m) \right\} d\lambda(\mathbf{h}^{S^*}) \\
&\leq S^* \times \frac{1}{\sqrt{\bar{\kappa}_J}} \left\| \sqrt{\bar{\kappa}_J} \left\{ \hat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^* \right\} \right\|_2 \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left\{ \sum_{v=1}^{S^*} |e^v| \right\} + \\
&\quad o_p(1/\sqrt{\bar{\kappa}_J}) \times \frac{1}{\sqrt{\bar{\kappa}_J}} \left\| \sqrt{\bar{\kappa}_J} \left\{ \hat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^* \right\} \right\|_2 \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left\{ \sum_{v=1}^{S^*} |e^v| \right\}.
\end{aligned}$$

This follows because  $\sqrt{\bar{\kappa}_J}(\hat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^*) = O_p(1)$ , and  $\mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left( \sum_{v=1}^{S^*} |e^v| \right) = O(1)$ .

Therefore,  $|V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \boldsymbol{\beta}^*) - V^{n_J}(\mathbf{h}^{n_J}, \boldsymbol{\pi}; \hat{\boldsymbol{\beta}}_J)| = O_p(1/\sqrt{\bar{\kappa}_J})$  as  $J \rightarrow \infty$ . This proves the result. ■

## 10 Epoch Asymptotics

As mentioned in Section 2.1 in the main text, we observe the branching process data in order of arrival time,  $\mathcal{D}^\kappa \triangleq \{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v, C^v)\}_{v=1}^\kappa$ . Define  $J_\kappa$  as the last complete epoch induced by coupon expiration,  $J_\kappa \triangleq \max\{j : \forall i \in \mathcal{E}_j, T_i + t_{\max} < T^\kappa\}$ . Lemma 10.1 ensures that  $\lim_{\kappa \rightarrow \infty} J_\kappa \rightarrow \infty$  a.s. under Assumption 5, implying that inference based on the last complete epoch will achieve the asymptotic guarantees of Theorem 2.3.

**Lemma 10.1.** *Assume that the working model specified in Equation 6 is the true generative model. Under Assumption 5,*

$$J_\kappa \rightarrow \infty \text{ a.s.}$$

*Proof.* First, we show that

$$\lim_{\kappa \rightarrow \infty} T^\kappa = \infty.$$

Define  $V_S$  as the seed sample set. Under Assumption 5, the number of coupons in a coupon allocation is upper bounded by  $L_*$ . It is sufficient to show that for any  $g \in \mathbb{N}$ ,  $T^{|V_S|L_*^{g-1}+1} \geq gt_{\min}$ .

We employ a proof by induction. First, we show the base case. Suppose  $g = 1$ . We know that  $T^{|V_S|+1} \geq t_{\min}$  by Assumption 5. We now assume that for any  $g \in \mathbb{N}$ ,

$$T^{|V_S|*L_*^{g-1}+1} \geq gt_{\min}.$$

Note that  $t_{\min} > 0$ . Consequently, an upper bound on the number of active coupons in the branching process at time  $T^{|V_S|*L_*^{g-1}}$  is

$$|V_S|L_*^{g-1} * L_* = |V_S|L_*^g.$$

Note that any study participant recruited by a new member of the sample (an individual recruited after the first  $|V_S|L_*^{g-1}$  study participants) will have a recruitment time greater than  $(g+1)t_{\min}$ ; i.e., for any  $\kappa$  such that  $R^\kappa > |V_S|L_*^{g-1}$ ,  $T^\kappa \geq (g+1)t_{\min}$ . Because  $|V_S|L_*^g - |V_S|L_*^{g-1}$  is an upper bound on the number of active coupons at time  $T^{|V_S|L_*^{g-1}}$ , we know that  $R^{|V_S|L_*^g} \geq |V_S|L_*^{g-1}$ . Consequently,

$$T^{|V_S|L_*^g+1} > (g+1)t_{\min}.$$

We conclude that

$$\lim_{\kappa \rightarrow \infty} T^\kappa = \infty$$

by induction. Now we establish that

$$\lim_{\kappa \rightarrow \infty} J_\kappa = \infty.$$

Recall that  $\bar{\kappa}_J \triangleq |\bar{\mathcal{E}}_J|$ . Additionally, recall that  $|\mathbf{A}^v| \leq L_*$  for all  $v \in N$ . Consequently,

for all  $j \in \mathbb{N}$ ,  $|\bar{\mathcal{E}}_j| \leq L_*^j < \infty$ . This implies that  $\max_{i \in \bar{\mathcal{E}}_j} T_i \leq t_{\max} L_*^j$ . Consequently, for any  $j \in \mathbb{N}$ ,

$$1 = P \left( \lim_{\kappa \rightarrow \infty} \{t_{\max} L_*^j + t_{\max} < T^\kappa\} \right) \leq P \left( \lim_{\kappa \rightarrow \infty} \{\forall i \in \bar{\mathcal{E}}_j, T_i + t_{\max} < T^\kappa\} \right).$$

We conclude that  $\lim_{\kappa \rightarrow \infty} J_\kappa = \infty$ . ■

## 11 Proof of Theorem 2.3

In this section, we prove Theorem 2.3 for the branching model described in Equation 6. Under Conditions (C1)-(C6), we verify Assumptions 4, 5, and 7-13. This will allow us to invoke Theorems 2.1 and 2.2 for the branching model described by Equation 6.

**General notation for this section.** For matrix  $A \in \mathbb{R}^{n \times n}$ ,  $\sigma_{\min}(A)$  and  $\sigma_{\max}(A)$  are the minimum and maximum singular values of  $A$  respectively. Define  $\text{vec}(A)$  as the vectorization of  $A$ . For a symmetric matrix  $X \in \mathbb{R}^{d \times d}$ , define  $\text{vech}(X)$  as the vectorization of the lower-triangular elements of  $X$ . Label  $I_d \in \mathbb{R}^{d \times d}$  as the  $d$ -dimensional identity matrix.

### 11.1 The Branching Models Derivatives

Before we begin verifying assumptions, it will be useful to calculate the derivatives of the log-likelihood of the branching process specified in Equation 6. First, we define the data

$$\mathbf{D}_i = (\underline{\mathbf{Y}}_i, \underline{\mathbf{X}}_i, \underline{\mathbf{T}}_i, M_i, \mathbf{A}_i).$$

For example,  $\mathbf{X}_i$  is the covariate vector of recruiter  $i$ , and  $\underline{\mathbf{X}}_i = (\mathbf{X}_{i,1}, \mathbf{X}_{i,2}, \dots, \mathbf{X}_{i,M_i})$  are the recruits of recruiter  $i$ . The complete likelihood for the branching process is

$$\begin{aligned}
\mathcal{L}_J(\boldsymbol{\beta}) &\triangleq \\
\mathcal{L}_J(\boldsymbol{\beta}_y) \mathcal{L}_J(\{\boldsymbol{\phi}_a, G_a, \Sigma_a\}_{a \in \mathbb{A}}) \mathcal{L}_J(\zeta, \lambda) &\triangleq \\
\prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} \prod_{l=1}^{M_i} \left[ \frac{1}{1 + \exp(-\mathbf{Z}_{i,l}^\top \boldsymbol{\beta}_y)} \right]^{Y_{i,l}} \left[ \frac{1}{1 + \exp(\mathbf{Z}_{i,l}^\top \boldsymbol{\beta}_y)} \right]^{1-Y_{i,l}} &\times \\
\prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} \prod_{l=1}^{M_i} (2\pi)^{-p/2} \left| \sum_{a \in \mathbb{A}} \Sigma_a \mathbb{I}(A_{i,l} = a) \right|^{-1/2} &\times \\
\exp \left[ -\frac{1}{2} \sum_{a \in \mathbb{A}} \{(\mathbf{X}_{i,l} - \boldsymbol{\phi}_a - G_a \mathbf{X}_i)^\top \Sigma_a^{-1} (\mathbf{X}_{i,l} - \boldsymbol{\phi}_a - G_a \mathbf{X}_i)\} \mathbb{I}(A_{i,l} = a) \right] & \\
\prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} M_i! \left[ \frac{\prod_{l=1}^{M_i} \zeta e^{-\zeta U_{i,l}}}{[e^{-\zeta t_{\min}} - e^{-\zeta t_{\max}}]^{M_i}} \right] \frac{\lambda^{M_i}/M_i!}{\sum_{\ell=k}^{|\mathbf{A}^v|} (\lambda^\ell/\ell!)} &.
\end{aligned} \tag{25}$$

Lastly, we rewrite the logarithm of the part of the likelihood that involves  $\boldsymbol{\beta}_t = \zeta$  and  $\boldsymbol{\beta}_m = \lambda$  as

$$\begin{aligned}
\ell_J(\zeta, \lambda) &\triangleq \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l=1}^{M_i} \{ \log(\zeta) - \zeta U_{i,l} - \log(e^{-\zeta t_{\min}} - e^{-\zeta t_{\max}}) \} + \\
&\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \left[ M_i \log(\lambda) - \log \left\{ \sum_{\ell=k}^{|\mathbf{A}^v|} (\lambda^\ell/\ell!) \right\} \right].
\end{aligned} \tag{26}$$

We will need the hessian of the log-likelihood of this branching process for the proofs that follow. Recall that  $\mathbb{A}$  is the set of possible coupon types. We note that each coupon allocation is a set of identical coupons, implying that the sets  $\mathcal{A}$  and  $\mathbb{A}$  have a one-to-one correspondence. Consequently, for  $\mathbf{a} \in \mathcal{A}$ , there exists  $a \in \mathbb{A}$  such that

$$\mathbb{I}(\mathbf{A}_i = \mathbf{a}) = \mathbb{I}(A_{i,l} = a)$$

for every  $l \in \{1, 2, \dots, M_i\}$ . Consequently, we can represent the complete branching

process likelihood as

$$\begin{aligned}
& \mathcal{L}_J(\boldsymbol{\beta}) \triangleq \\
& \mathcal{L}_J(\boldsymbol{\beta}_y) \mathcal{L}_J(\{\boldsymbol{\phi}_{\mathbf{a}}, G_{\mathbf{a}}, \Sigma_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) \mathcal{L}_J(\zeta, \lambda) \triangleq \\
& \prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} \prod_{l=1}^{M_i} \left[ \frac{1}{1 + \exp(-\mathbf{Z}_{i,l}^\top \boldsymbol{\beta}_y)} \right]^{Y_{i,l}} \left[ \frac{1}{1 + \exp(\mathbf{Z}_{i,l}^\top \boldsymbol{\beta}_y)} \right]^{1-Y_{i,l}} \times \\
& \prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} \prod_{l=1}^{M_i} (2\pi)^{-p/2} \left| \sum_{\mathbf{a} \in \mathcal{A}} \Sigma_{\mathbf{a}} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \right|^{-1/2} \times \\
& \exp \left[ -\frac{1}{2} \sum_{\mathbf{a} \in \mathcal{A}} \{(\mathbf{X}_{i,l} - \boldsymbol{\phi}_{\mathbf{a}} - G_{\mathbf{a}} \mathbf{X}_i)^\top \Sigma_{\mathbf{a}}^{-1} (\mathbf{X}_{i,l} - \boldsymbol{\phi}_{\mathbf{a}} - G_{\mathbf{a}} \mathbf{X}_i)\} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \right] \\
& \prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} M_i! \left[ \frac{\prod_{l=1}^{M_i} \zeta e^{-\zeta U_{i,l}}}{[e^{-\zeta t_{\min}} - e^{-\zeta t_{\max}}]^{M_i}} \right] \frac{\lambda^{M_i} / M_i!}{\sum_{\ell=k}^{|\mathbf{A}^v|} (\lambda^\ell / \ell!)}.
\end{aligned} \tag{27}$$

We will use this likelihood for the proofs that follow.

**The Hessian for the log-likelihood of the covariate model.** Define  $\mathbf{X}_i^* = (1, \mathbf{X}_i)$

and  $G_{\mathbf{a}}^\dagger = (\boldsymbol{\phi}_{\mathbf{a}}^\top, G_{\mathbf{a}}^\top)^\top$ . The likelihood of the covariate model parameter,

$\boldsymbol{\beta}_{\mathbf{x}} = \{G_{\mathbf{a}}^\dagger, \Sigma_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}$ , is

$$\begin{aligned}
\mathcal{L}_J(\{G_{\mathbf{a}}^\dagger, \Sigma_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) & \triangleq \prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} \prod_{l=1}^{M_i} (2\pi)^{-p/2} \left| \sum_{\mathbf{a} \in \mathcal{A}} \Sigma_{\mathbf{a}} \mathbb{I}(c = \mathbf{a}) \right|^{-1/2} \times \\
& \exp \left[ -\frac{1}{2} \sum_{\mathbf{a} \in \mathcal{A}} \{(\mathbf{X}_{i,l} - G_{\mathbf{a}}^\dagger \mathbf{X}_i^*)^\top \Sigma_{\mathbf{a}}^{-1} (\mathbf{X}_{i,l} - G_{\mathbf{a}}^\dagger \mathbf{X}_i^*)\} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \right] \\
& = \prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} \prod_{l=1}^{M_i} (2\pi)^{-p/2} \left| \sum_{\mathbf{a} \in \mathcal{A}} \Sigma_{\mathbf{a}} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \right|^{-1/2} \times \\
& \exp \left[ -\frac{1}{2} \sum_{\mathbf{a} \in \mathcal{A}} \left\{ \mathbf{X}_{i,l}^\top \Sigma_{\mathbf{a}}^{-1} \mathbf{X}_{i,l} - 2\mathbf{X}_i^{*\top} G_{\mathbf{a}}^\dagger{}^\top \Sigma_{\mathbf{a}}^{-1} \mathbf{X}_{i,l} + \right. \right. \\
& \quad \left. \left. \mathbf{X}_i^{*\top} G_{\mathbf{a}}^\dagger{}^\top \Sigma_{\mathbf{a}}^{-1} G_{\mathbf{a}}^\dagger \mathbf{X}_i^* \right\} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \right].
\end{aligned}$$

The equality above simply follows from distributing. For  $\mathbf{a} \in \mathcal{A}$ , we reparameterize  $\Omega_{\mathbf{a}} = -\frac{1}{2}\Sigma_{\mathbf{a}}^{-1}$  and  $\Gamma_{\mathbf{a}} = \Sigma_{\mathbf{a}}^{-1}G_{\mathbf{a}}^{\dagger}$ .

$$\begin{aligned} \mathcal{L}_J(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) &= \prod_{j=1}^J \prod_{i \in \mathcal{E}_{j-1}} \prod_{l=1}^{M_i} (2\pi)^{-p/2} |\sum_{\mathbf{a} \in \mathcal{A}} -2\Omega_{\mathbf{a}} \mathbb{I}(\mathbf{A}_i = \mathbf{a})|^{1/2} \times \\ &\quad \exp \left[ \sum_{\mathbf{a} \in \mathcal{A}} \left\{ \mathbf{X}_{i,l}^{\top} \Omega_{\mathbf{a}} \mathbf{X}_{i,l} + \mathbf{X}_i^{*\top} \Gamma_{\mathbf{a}}^{\top} \mathbf{X}_{i,l} + \right. \right. \\ &\quad \left. \left. \frac{1}{4} \mathbf{X}_i^{*\top} \Gamma_{\mathbf{a}}^{\top} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{X}_i^* \right\} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \right]. \end{aligned}$$

The reparameterized log-likelihood is

$$\begin{aligned} \ell_J(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) &\triangleq \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l=1}^{M_i} \sum_{\mathbf{a} \in \mathcal{A}} \left[ (-p/2) \log(2\pi) + \frac{1}{2} \log | -2\Omega_{\mathbf{a}} | + \right. \\ &\quad \left. \mathbf{X}_{i,l}^{\top} \Omega_{\mathbf{a}} \mathbf{X}_{i,l} + \mathbf{X}_i^{*\top} \Gamma_{\mathbf{a}}^{\top} \mathbf{X}_{i,l} + \right. \\ &\quad \left. \frac{1}{4} \mathbf{X}_i^{*\top} \Gamma_{\mathbf{a}}^{\top} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{X}_i^* \right] \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \\ &= \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l=1}^{M_i} \sum_{\mathbf{a} \in \mathcal{A}} \left[ (-p/2) \log(2\pi) + \frac{1}{2} \log | -2\Omega_{\mathbf{a}} | + \right. \\ &\quad \left. \text{tr}(\Omega_{\mathbf{a}} \mathbf{X}_{i,l} \mathbf{X}_{i,l}^{\top}) + \text{tr}(\Gamma_{\mathbf{a}}^{\top} \mathbf{X}_{i,l} \mathbf{X}_i^{*\top}) + \right. \\ &\quad \left. \text{tr}\left(\frac{1}{4} \Gamma_{\mathbf{a}}^{\top} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{X}_i^* \mathbf{X}_i^{*\top}\right) \right] \mathbb{I}(\mathbf{A}_i = \mathbf{a}), \end{aligned}$$

where the second equality follows from rearranging terms and using the properties of the trace operator. For  $\mathbf{a} \in \mathcal{A}$ , define  $n_{\mathbf{a}} \triangleq \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} M_i \mathbb{I}(\mathbf{A}_i = \mathbf{a})$  and  $\mathbf{V}_{\mathbf{a}}^J \in \mathbb{R}^{(p+1) \times (p+1)}$  such that  $\mathbf{V}_{\mathbf{a}}^J \triangleq \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} M_i \mathbf{X}_i^* \mathbf{X}_i^{*\top} \mathbb{I}(\mathbf{A}_i = \mathbf{a})$ . We apply the differential operator

two times and find

$$\begin{aligned}
\mathbf{d}^2 \ell_J (\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) &= \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l=1}^{M_i} \sum_{\mathbf{a} \in \mathcal{A}} -\mathbf{d}^2 \left\{ -\frac{1}{2} \log | -2\Omega_{\mathbf{a}} | - \right. \\
&\quad \left. \frac{1}{4} \text{tr} \left( \Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{X}_i^* \mathbf{X}_i^{*\top} \right) \right\} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \\
&= -\mathbf{d} \left\{ -\frac{n_{\mathbf{a}}}{2} \text{tr} (\Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}}) - \frac{1}{4} \text{tr} (2\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^J) + \right. \\
&\quad \left. \frac{1}{4} \text{tr} (\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^J) \right\} \\
&= -\left\{ \frac{n_{\mathbf{a}}}{2} \text{tr} (\Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}}) - \right. \\
&\quad \frac{1}{2} \text{tr} (\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^J) - \\
&\quad \frac{1}{2} \text{tr} (\mathbf{d}\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^J) + \\
&\quad \text{tr} (\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^J) + \\
&\quad \left. \frac{1}{4} \text{tr} (\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^J) \right\}.
\end{aligned}$$

We observe that for  $\mathbf{a}, \mathbf{a}' \in \mathcal{A}$  and  $\mathbf{a} \neq \mathbf{a}'$ ,

$$\frac{\partial \ell_J (\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}})}{\partial (\text{vec}(\Gamma_{\mathbf{a}}), \text{vec}(\Omega_{\mathbf{a}})) \partial (\text{vec}(\Gamma_{\mathbf{a}'}) , \text{vec}(\Omega_{\mathbf{a}'}))^\top} = [0]_{(p+1)^2 \times p^2}.$$

We express

$$\begin{aligned}
\ddot{\ell}_J (\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) &= \\
&= -\text{diag} \left[ \left\{ \left( \begin{array}{cc} \mathbf{V}_{\mathbf{a}}^J \otimes \Sigma_{\mathbf{a}} & 2(\mathbf{V}_{\mathbf{a}}^J G_{\mathbf{a}}^{*\top} \otimes \Sigma_{\mathbf{a}}) \\ 2(G_{\mathbf{a}}^* \mathbf{V}_{\mathbf{a}}^J \otimes \Sigma_{\mathbf{a}}) & 4(G_{\mathbf{a}}^* \mathbf{V}_{\mathbf{a}}^J G_{\mathbf{a}}^{*\top} \otimes \Sigma_{\mathbf{a}}) + 2n_{\mathbf{a}}(\Sigma_{\mathbf{a}} \otimes \Sigma_{\mathbf{a}}) \end{array} \right) \right\}_{\mathbf{a} \in \mathcal{A}} \right] \\
&= -\text{diag} \left[ \left\{ \left( \begin{array}{cc} \mathbf{V}_{\mathbf{a}}^J & 2\mathbf{V}_{\mathbf{a}}^J G_{\mathbf{a}}^{*\top} \\ 2G_{\mathbf{a}}^* \mathbf{V}_{\mathbf{a}}^J & 4G_{\mathbf{a}}^* \mathbf{V}_{\mathbf{a}}^J G_{\mathbf{a}}^{*\top} + 2n_{\mathbf{a}} \Sigma_{\mathbf{a}} \end{array} \right) \otimes \Sigma_{\mathbf{a}} \right\}_{\mathbf{a} \in \mathcal{A}} \right].
\end{aligned}$$

The second equality follows from properties of Kronecker products.

**The Hessian for the log-likelihood of the reward model.** The estimating equation for the reward model parameter,  $\beta_y$ , is

$$\frac{\partial \ell_J(\beta_y)}{\partial \beta_y} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l=1}^{M_i} \left[ Y_{i,l} \mathbf{Z}_{i,l} - \left\{ \frac{1}{1 + \exp(-\mathbf{Z}_{i,l}^\top \beta_y)} * \mathbf{Z}_{i,l} \right\} \right].$$

The derivative of the estimating equation (the hessian of the log-likelihood) is

$$\frac{\partial^2 \ell_J(\beta_y)}{\partial \beta_y \partial \beta_y^\top} = - \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l=1}^{M_i} \mathbf{Z}_{i,l} \mathbf{Z}_{i,l}^\top \left\{ \frac{1}{1 + \exp(-\mathbf{Z}_{i,l}^\top \beta_y)} \right\} \left\{ \frac{1}{1 + \exp(\mathbf{Z}_{i,l}^\top \beta_y)} \right\}.$$

**The Hessian for the log-likelihood of the arrival model.** Appealing to Equation 26, the estimating equation for the arrival model parameter,  $\beta_t = \zeta$ , is

$$\frac{\partial \ell_J(\zeta)}{\partial \zeta} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l=1}^{M_i} \frac{1}{\zeta} - U_{i,l} - \frac{t_{\min} e^{-\zeta t_{\min}} - t_{\max} e^{-\zeta t_{\max}}}{e^{-\zeta t_{\min}} - e^{-\zeta t_{\max}}}.$$

The derivative of the estimating equation (the hessian of the log-likelihood) is

$$\frac{\partial^2 \ell_J(\zeta)}{\partial \zeta^2} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} -M_i \left( \frac{1}{\zeta^2} + \frac{t_{\min} e^{-\zeta t_{\min}} - t_{\max}^2 e^{-\zeta t_{\max}}}{[e^{-\zeta t_{\min}} - e^{-\zeta t_{\max}}]^2} \right).$$

**The Hessian for the log-likelihood of the family model.** We reparameterize the log-likelihood in Equation 26, where  $\tau = \log(\lambda)$ :

$$\ell_J(\tau) = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \left[ M_i \tau - \log \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} (e^{\tau \ell} / \ell!) \right\} \right].$$

The score function is

$$\frac{\partial \ell_J(\tau)}{\partial \tau} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \left[ M_i - \tau \frac{\sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell} / \ell!}{\sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell!} \right].$$

This makes the Hessian of the log-likelihood

$$\frac{\partial^2 \ell_J(\tau)}{\partial \tau^2} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} -\tau^2 \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau \ell} / \ell! \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell} / \ell! \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\}^2}.$$

## 11.2 Verification of Assumptions 4-5 and 7-8

### 11.2.1 Assumption 4

**Assumption 4** is satisfied by Thompson sampling with a clipping constraint. We show this in Lemma 11.1.

**Lemma 11.1.** *For any  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ , assume  $\mathbf{A}_i$  is assigned by RL-RDS with a clipping constraint at level  $\epsilon_1 \in (0, 1)$  (as specified in Section 2). Additionally, assume Conditions (C1)-(C6). There exists  $\epsilon_{\min}, \epsilon_{\max} > 0$  such that for all  $\mathbf{a} \in \mathcal{A}$ ,*

$$\epsilon_{\min} \leq \mathbb{P}(\mathbf{A}_i = \mathbf{a} | \mathbf{H}_i) \leq \epsilon_{\max} \quad \text{w.p. 1.}$$

This implies that for  $\rho_{\min} = \sqrt{\epsilon_{\min}}$  and  $\rho_{\max} = \sqrt{\epsilon_{\max}}$ ,

$$\rho_{\min} \leq \mathbf{W}_i \leq \rho_{\max} \quad \text{w.p. 1.}$$

*Proof.* Note that by Condition (C3), the set of possible coupon allocations is constant for the duration of the study. For any  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ , Thompson sampling with clipping at level  $\epsilon_1 \in (0, 1)$  selects allocation  $\mathbf{a} \in \mathcal{A}$  with probability

$$\mathbb{P}(\mathbf{A}_i = \mathbf{a}) = \frac{p_i^{\mathbf{a}}}{\sum_{\mathbf{a}' \in \mathcal{A}} p_i^{\mathbf{a}'}},$$

where  $p_i^{\mathbf{a}} \triangleq \min \left[ 1 - \epsilon_1, \max \left\{ \epsilon_1, \widehat{\xi}_B^i(\mathbf{H}_i, \mathbf{a}) \right\} \right]$  (refer to Section 2 for the definition of  $\widehat{\xi}_B^i$ ).

This implies that

$$\mathbb{P}(\mathbf{A}_i = \mathbf{a}) = \frac{p_i^{\mathbf{a}}}{\sum_{\mathbf{a}' \in \mathcal{A}} p_i^{\mathbf{a}'}} \geq \frac{\epsilon_1}{(1 - \epsilon_1)|\mathcal{A}|}.$$

Consequently, for any  $\mathbf{a} \in \mathcal{A}$ ,

$$\frac{1 - \epsilon_1}{1 + (|\mathcal{A}| - 2)\epsilon_1} \geq \mathbb{P}(\mathbf{A}_i = \mathbf{a}).$$

The inequality follows from the fact that  $1 - \epsilon_1 \geq p_i^{\mathbf{a}} \geq \epsilon_1$ . We define

$$\epsilon_{\min} \triangleq \epsilon_1 / \{(1 - \epsilon_1)|\mathcal{A}|\}, \quad \epsilon_{\max} \triangleq \frac{1 - \epsilon_1}{1 + (|\mathcal{A}| - 2)\epsilon_1}.$$

■

### 11.2.2 Assumption 5

**Assumption 5** is satisfied by Equation 7 (in Section 2.1 of the main text) and Condition (C1). See the verification of Assumption 11 for a further discussion of Equation 7.

### 11.2.3 Assumptions 7 and 8

**Assumption 7** is verified by the derivatives calculated in Section 11.1 and the fact that each component of this branching model is a full rank exponential family. Consequently, the natural parameters are identifiable (Brown, 1986). **Assumption 8** follows from the score and Hessian functions calculated in Section 11.1. Observe that the likelihood and Hessian components are continuous and finite in the data for a given parameter value. Consequently, the fact that the data are bounded implies that for any  $j \in \mathbb{N}$  and  $i \in \mathcal{E}_{j-1}$ ,  $\dot{l}_i(\boldsymbol{\beta})$  and  $\ddot{l}_i(\boldsymbol{\beta})$  are bounded (by the extreme value theorem). This implies that the conditional expectations of these functions are bounded and **Assumption 8** is satisfied.

## 11.3 Verification of Assumptions 9 and 11-14

We now verify Assumptions 9 and 11-14.

### 11.3.1 Note on Assumption 10

Note that Assumption 10 is only used in the proof of consistency in Section 8. It is unnecessary because the log-likelihood of the branching process specified in Equation 6 is concave. We will provide another proof of consistency in Section 11.4 that does not use Assumption 10.

### 11.3.2 Supporting Lemmas

Lemmas 11.2 and 11.3 are useful properties of positive semi-definite matrices.

**Lemma 11.2.** *Define positive semi-definite matrices  $A, B \in \mathbb{R}^{d \times d}$ . It follows that*

$$\sigma_{\min}(A + B) \geq \sigma_{\min}(A) + \sigma_{\min}(B).$$

*Proof.*

$$\begin{aligned} \sigma_{\min}(A + B) &= \min_{x \neq 0} \frac{x^\top (A + B)x}{x^\top x} \\ &= \min_{x \neq 0} \left( \frac{x^\top Ax}{x^\top x} + \frac{x^\top Bx}{x^\top x} \right) \\ &\geq \min_{x \neq 0} \frac{x^\top Ax}{x^\top x} + \min_{x \neq 0} \frac{x^\top Bx}{x^\top x} \\ &= \sigma_{\min}(A) + \sigma_{\min}(B). \end{aligned}$$

The first line follows from the definition of the minimum singular value for positive semi-definite matrices, and the second from distributing. For any functions  $f$  and  $g$ ,  $\min_{x \in \mathcal{X}} \{f(x) + g(x)\} \geq \min_{x \in \mathcal{X}} f(x) + \min_{x \in \mathcal{X}} g(x)$ . Consequently, line 3 is true. Line 4 applies the definition of the minimum singular value again. ■

**Lemma 11.3.** *Let  $A, B \in \mathbb{R}^{d \times d}$  be positive semi-definite matrices. It follows that*

$$\sigma_{\min}(A)\sigma_{\min}(B) \leq \sigma_{\min}(AB).$$

*Proof.* First, we assume that matrices  $A$  and  $B$  are invertible. By the sub-multiplicativity

of the spectral norm, we know that

$$\sigma_{\max}(AB) \leq \sigma_{\max}(A)\sigma_{\max}(B).$$

Consequently,

$$\begin{aligned} \sigma_{\max}\{(AB)^{-1}\} &\leq \sigma_{\max}(A^{-1})\sigma_{\max}(B^{-1}) \Rightarrow \\ \sigma_{\min}(AB)^{-1} &\leq \sigma_{\min}(A)^{-1}\sigma_{\min}(B)^{-1} \Rightarrow \\ \sigma_{\min}(A)\sigma_{\min}(B) &\leq \sigma_{\min}(AB). \end{aligned}$$

The first line applies the sub-multiplicativity of the spectral norm. For any matrix  $A$ ,  $\sigma_{\max}(A^{-1}) = \sigma_{\min}(A)^{-1}$ . Consequently, line 2 is true.

If matrices  $A$  and  $B$  are not invertible, then  $\sigma_{\min}(A) = 0$ ,  $\sigma_{\min}(B) = 0$ , and  $\sigma_{\min}(AB) = 0$ . Therefore, the result follows trivially. ■

**Lemma 11.4.** *Let  $M \in \mathbb{R}^{d \times d}$  be a matrix such that*

$$M = \begin{pmatrix} A & B \\ B^\top & C \end{pmatrix}$$

where  $A \in \mathbb{R}^{d_1 \times d_1}$  and  $C \in \mathbb{R}^{d_2 \times d_2}$  are symmetric and  $d_1 + d_2 = d$ .  $M$  is positive definite if and only if  $A$  is positive definite and  $C - B^\top A^{-1}B$  is positive definite. Additionally,

$$\sigma_{\min}(M) \geq \sigma_{\min}(A)\sigma_{\min}(C - B^\top A^{-1}B).$$

*Proof.* Define  $I_{d_1} \in \mathbb{R}^{d_1 \times d_1}$  and  $I_{d_2} \in \mathbb{R}^{d_2 \times d_2}$  as the  $d_1$  and  $d_2$ -dimensional identity matrices

respectively. We express  $M$  as

$$\begin{aligned} M &= \begin{pmatrix} A & B \\ B^\top & C \end{pmatrix} \\ &= \begin{pmatrix} I_{d_1} & 0 \\ B^\top A^{-1} & I_{d_2} \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & C - B^\top A^{-1} B \end{pmatrix} \begin{pmatrix} I_{d_1} & A^{-1} B \\ 0 & I_{d_2} \end{pmatrix}. \end{aligned}$$

Because

$$\begin{pmatrix} I_{d_1} & 0 \\ B^\top A^{-1} & I_{d_2} \end{pmatrix}$$

and its transpose are invertible, we know that  $M$  is positive definite if and only if

$$Q = \begin{pmatrix} A & 0 \\ 0 & C - B^\top A^{-1} B \end{pmatrix}$$

is positive definite. The matrix  $Q$  is block diagonal, so it is positive definite if and only if  $A$  and  $C - B^\top A^{-1} B$  are positive definite. Additionally, by Lemma 11.3,

$$\begin{aligned} \sigma_{\min}(M) &\geq \sigma_{\min} \left\{ \begin{pmatrix} I_{d_1} & 0 \\ B^\top A^{-1} & I_{d_2} \end{pmatrix} \right\} \sigma_{\min} \left\{ \begin{pmatrix} A & 0 \\ 0 & C - B^\top A^{-1} B \end{pmatrix} \right\} \times \\ &\quad \sigma_{\min} \left\{ \begin{pmatrix} I_{d_1} & A^{-1} B \\ 0 & I_{d_2} \end{pmatrix} \right\} \\ &\geq \sigma_{\min}(A) \sigma_{\min}(C - B^\top A^{-1} B) \sigma_{\min}(I_{d_1})^2 \sigma_{\min}(I_{d_2})^2 \\ &= \sigma_{\min}(A) \sigma_{\min}(C - B^\top A^{-1} B) \end{aligned}$$

■

### 11.3.3 Assumptions 9 and 14

We verify **Assumption 9** first. We write  $\dot{l}_i(\boldsymbol{\beta}^*) = \dot{l}(\boldsymbol{\beta}^*, \mathbf{D}_i)$  as a function  $\dot{l} : \mathcal{B} \times \mathcal{D} \rightarrow \mathbb{R}$ . From Section 11.2, we know that  $\dot{l}$  is continuous over both  $\mathcal{B}$  and  $\mathcal{D}$ . Because  $\mathcal{X}$  is compact (and the other data types are bounded by definition), we know that  $\mathcal{D}$  is compact. By the extreme value theorem, we can define  $\gamma < \infty$ ,

$$\max_{\mathbf{d} \in \mathcal{D}} \max_{\boldsymbol{\beta} \in \mathcal{B}} \left\| \dot{l}(\boldsymbol{\beta}, \mathbf{d}) \right\|_2 \leq \gamma.$$

$\mathcal{B}$  is a convex subset of  $\mathbb{R}^q$ . Consequently, by the mean value theorem, we know that for all  $j \in \mathbb{N}$ ,  $i \in \mathcal{E}_{j-1}$ , and  $\boldsymbol{\beta}, \boldsymbol{\beta}' \in \mathcal{B}$ ,

$$|l_i(\boldsymbol{\beta}) - l_i(\boldsymbol{\beta}')| \leq |\dot{l}_i(\bar{\boldsymbol{\beta}})| \|\boldsymbol{\beta} - \boldsymbol{\beta}'\|_2,$$

for some  $\bar{\boldsymbol{\beta}} = t\boldsymbol{\beta} + (1-t)\boldsymbol{\beta}'$  where  $t \in [0, 1]$ . Because  $|\dot{l}_i(\bar{\boldsymbol{\beta}})| = |\dot{l}(\bar{\boldsymbol{\beta}}, \mathbf{D}_i)| \leq \gamma$ , we know that

$$|l_i(\boldsymbol{\beta}) - l_i(\boldsymbol{\beta}')| \leq \gamma \|\boldsymbol{\beta} - \boldsymbol{\beta}'\|_2$$

and **Assumptions 9** is satisfied.

The likelihood of the arrival process implied by data  $\mathcal{D}^\kappa = \{(R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v)\}_{v=1}^\kappa$  is simply a censored (or integrated) version of the complete generation likelihood of Section 11.2. Consequently, write  $\dot{q}^v(\boldsymbol{\beta}) = \dot{q}(\boldsymbol{\beta}, \mathbf{D}^v)$  (the components of this likelihood as defined in Section 2) as a function  $\dot{q} : \mathcal{B} \times \mathcal{D} \rightarrow \mathbb{R}$ , we know that  $\dot{q}$  is continuous over both  $\mathcal{B}$  and  $\mathcal{D}$ . Therefore, **Assumption 14** is satisfied by the logic presented in the previous paragraph.

### 11.3.4 Assumption 11

In this section, we verify **Assumption 11**. To do this, we introduce the branching process in a manner consistent with Delmas and Marsalle (2010). Most of the following setup is taken directly from their paper; we reproduce it here for the reader's convenience. We modify the background and theory where necessary to generalize their proofs to more

than two possible recruits (or cells in their case).

Let  $\mathbb{G}_0 = \emptyset$ ,  $\mathbb{G}_j = \{1, \dots, L\}^j$  for  $j \in \mathbb{N}$ ,  $\mathbb{T}_r = \cup_{0 \leq j \leq r} \mathbb{G}_j$ . The set  $\mathbb{G}_j$  contains all possible recruits in the  $j$ -th generation. Note that for participant  $i \in \mathbb{G}_j$ ,  $j = |i|$  denotes the generation of  $i$ . Additionally, for  $i, j \in \mathbb{T}$ , we can represent the concatenation of their positions as  $ij = (i, j)$ . For recruit  $i \in \mathbb{T}$ , we still denote  $\mathbf{X}_i \in \mathbb{R}^p$  as an individual's covariates as specified in the model described by Equation 6. We define the transition kernel implied by this covariate process over the branching tree as follows. Let  $(E, \mathcal{E})$  be a measurable space, and let  $P$  be a probability kernel on  $E \times \mathcal{E}^L$  with values in  $[0, 1]$ . This implies that  $P(\cdot, A)$  is measurable for all  $A \in \mathcal{E}^L$  and  $P(\mathbf{x}, \cdot)$  is a probability measure on  $(E^L, \mathcal{E}^L)$ . Additionally, for any real-valued function  $g$  defined on  $E^L$  and  $\mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_L)$ , we set

$$Pg(\mathbf{x}) = \int_{E^L} g(\mathbf{q})P(\mathbf{x}, d\mathbf{q}).$$

We call a stochastic process indexed by  $\mathbb{T}$ ,  $\{\mathbf{X}_i, i \in \mathbb{T}\}$ , a fixed branching Markov chain on a measurable space  $(E, \mathcal{E})$  with initial distribution  $\nu$  and probability kernel  $P$  if:

- $\mathbf{X}_\emptyset$  is distributed as  $\nu$
- It satisfies a Markov property: for any measurable real-valued bounded functions  $(g_i, i \in \mathbb{T})$  defined on  $E^L$ , we have that for all  $j \geq 0$

$$\mathbb{E} \left[ \prod_{i \in \mathbb{G}_j} g_i(\mathbf{X}_{i_1}, \mathbf{X}_{i_2}, \dots, \mathbf{X}_{i_L}) \mid \sigma(\mathbf{X}_k, k \in \mathbb{T}_j) \right] = \prod_{i \in \mathbb{G}_j} Pg_i(\mathbf{X}_i).$$

We consider the metric measurable space  $(S, \mathcal{S})$ , and add a cemetery point to  $S$ ,  $\delta$ . Let  $\bar{S} = \mathbb{R}^p \cup \{\delta\}$  and  $\bar{\mathcal{S}}$  be the  $\sigma$ -field generated by  $\mathcal{S}$  and  $\{\delta\}$ . Let  $P^*$  be a probability kernel defined on  $\bar{S} \times \bar{S}^L$  such that

$$P^*(\delta, \{(\delta, \delta, \dots, \delta)\}) = 1. \tag{28}$$

Note that this condition means that  $\delta$  is an absorbing state.

The covariate process specified in Equation 6,  $\{\mathbf{X}_i, i \in \mathbb{T}^*\}$  with  $\mathbb{T}^* = \{i \in \mathbb{T} : \mathbf{X}_i \neq \delta\}$  is a fixed branching Markov chain on  $(\bar{S}, \bar{S})$  with  $P^*$  satisfying Equation 28. Note that the recruit covariate distributions are independently and identically distributed. In other words, for  $V \triangleq (V_1 \times V_2 \times \cdots \times V_L) \in \mathcal{S}^L$  and an induced kernel  $P_1$ ,

$$\int_V P(\mathbf{x}, \mathbf{q}) d\mathbf{q} = \int_{V^1} \int_{V^2} \cdots \int_{V^L} P_1(\mathbf{x}, d\mathbf{q}_1) P_1(\mathbf{x}, d\mathbf{q}_2) \cdots P_1(\mathbf{x}, d\mathbf{q}_L) d\mathbf{q}_1 d\mathbf{q}_2 \cdots d\mathbf{q}_L.$$

Additionally, the family size distribution is only dependent on the total number of recruits. This fact, along with the i.i.d. property described above, implies that, for  $\bar{V} = (\bar{V}^1, \bar{V}^2, \dots, \bar{V}^L) \in \bar{S}^L$ ,

$$\begin{aligned} \int_{\bar{V}} P^*(\mathbf{x}, \mathbf{q}) d\mathbf{q} &= \int_{V^1} \int_{V^2} \cdots \int_{V^L} P^*(\mathbf{x}, \mathbf{q}) d\mathbf{q} \\ &= \int_{V^{z(1)}} \int_{V^{z(2)}} \cdots \int_{V^{z(L)}} P^*(\mathbf{x}, \mathbf{q}) d\mathbf{q} \end{aligned}$$

for any permutation  $z$  of  $\{1, 2, \dots, L\}$ . We will call this property the “exchangeability” of  $P^*$ . Lastly, the process is “spatially homogeneous;” i.e., for  $\bar{V} \in \{S, \delta\}^L$  and  $\mathbf{x}_1, \mathbf{x}_2 \in S$ ,

$$P^*(\mathbf{x}_1, \bar{V}) = P^*(\mathbf{x}_2, \bar{V}).$$

In other words, the family size distribution does not depend on the covariate value of the parent. By Condition (C1), the Galton-Watson (GW) tree implied by this process is super-critical.

We now cite some well known results from the branching process literature (Athreya et al., 2004). For any subset  $O \subset \mathbb{T}$ , let

$$O^* = O \cap \mathbb{T}^* = \{o \in O, \mathbf{X}_k \neq \delta\}$$

be the subset of  $O$  that are “realized” recruits. Labeling  $Z_j \triangleq |\mathbb{G}_j^*|$ ,  $\{Z_j, j \in \mathbb{N}\}$  is a GW

process. For  $j \geq 0$ , we know

$$\mathbb{E}(|\mathbb{G}_j^*|) = m^j.$$

And, for  $r \geq 0$ , we know

$$\mathbb{E}(|\mathbb{T}_r^*|) = \sum_{j=0}^r \mathbb{E}(|\mathbb{G}_j^*|) = \sum_{j=0}^r m^j = \frac{m^{r+1} - 1}{m - 1}.$$

Additionally, there exists a random variable  $\mathcal{I}$  s.t.

$$\mathcal{I} = \lim_{j \rightarrow \infty} m^{-j} |\mathbb{G}_j^*| \quad \text{a.s. and in } L^2 \quad (29)$$

where  $\mathbb{E}(\mathcal{I}) = 1$ .

We now define a series of useful sub-probability kernels. The first is on  $S \times \mathcal{S}^2$  such that

$$P_2^* = P^* \left\{ \cdot, (\cdot \cap S^2) \times \bar{S}^{L-2} \right\}.$$

Note that by the exchangeability of  $P^*$ , for any  $i, j \in \{0, 1, \dots, L-2\}$  such that  $i + j \leq L-2$ ,

$$P_2^* = P^* \left\{ \cdot, (\cdot \cap S^2) \times \bar{S}^{L-2} \right\} = P^* \left\{ \cdot, \bar{S}^i \times (\cdot \cap S) \times \bar{S}^j \times (\cdot \cap S) \times \bar{S}^{L-2-i-j} \right\}.$$

The second sub-probability kernel is defined on  $S \times \mathcal{S}$  in the following manner,

$$P_1^* = P^* \left\{ \cdot, (\cdot \cap S) \times \bar{S}^{L-1} \right\}.$$

By the same exchangeability property, we know that for any  $0 \leq i \leq L-1$ ,

$$P_1^* = P^* \left\{ \cdot, (\cdot \cap S) \times \bar{S}^{L-1} \right\} = P^* \left\{ \cdot, \bar{S}^i \times (\cdot \cap S) \times \bar{S}^{L-1-i} \right\}.$$

We introduce an auxiliary Markov chain. Let  $\{\mathbf{K}_j, j \in \mathbb{N}\}$  be a Markov chain on  $S$

with  $\mathbf{K}_0$  distributed as  $\mathbf{X}^\theta$  and transition kernel

$$Q = \frac{L}{m} P_1^*.$$

The distribution of  $\mathbf{K}_j$  corresponds to the distribution of  $\mathbf{X}_I$  conditional on  $\{I \in \mathbb{T}^*\}$ , where  $I$  is chosen at random from  $\mathbb{G}_j$ .

We explore the nature of transition kernel  $Q$  as it will be useful in theory later. We define  $P_1$  as a probability kernel on  $S \times \mathcal{S}$  that represents the marginal of  $P$  (note the use of  $P$  instead of  $P^*$  here). Consequently, it represents the dynamics of a single recruiter, recruit covariate process (with the guarantee that the recruit exists). Formally,  $P_1(\cdot, V)$  is measurable for  $V \in \mathcal{S}$ , where

$$P_1(\cdot, V) = P(\cdot, V \times S^{L-1}),$$

and  $P_1(\mathbf{x}, \cdot)$  is a probability measurable on  $(S, \mathcal{S})$  such that

$$P_1(\mathbf{x}, \cdot) = P(\mathbf{x}, \cdot \times S^{L-1}).$$

We can also define  $P_1$  as the sub-probability kernel  $P_1^*$  re-normalized over  $S$ . We see that for  $\mathbf{x} \in S$ ,

$$P_1(\mathbf{x}, \cdot) = P_1^*(\mathbf{x}, \cdot | \cdot \in S) = \frac{P_1^*(\mathbf{x}, \cdot)}{P_1^*(\mathbf{x}, S)},$$

or in other words  $P_1 = P_1^*/P_1^*(\mathbf{x}, S)$ . We now show that

$$Q = \frac{L}{m} P_1^*(\mathbf{x}, S) = P_1^*/P_1^*(\mathbf{x}, S) = P_1,$$

where  $Q$  is defined as the transition kernel for the auxiliary Markov chain  $(\mathbf{K}_j, j \in \mathbb{N})$  above.

**Lemma 11.5.** *Let  $Q$  be defined as above, then*

$$Q = \frac{L}{m} P_1^* = P_1^*/P_1^*(\mathbf{x}, S) = P_1$$

*Proof.* To prove the following statement, we need to show that

$$\frac{m}{L} = P_1^*(\mathbf{x}, S).$$

Due to the spatial homogeneity of the Markov process, we can define  $p_\gamma$  for  $\gamma \in \{0, 1, \dots, L\}$  as the probability that  $\gamma$  people are recruited (regardless of the recruiter's covariates). We evaluate  $P_1^*(\mathbf{x}, S)$  in terms of these probabilities (keeping in mind that the recruits are identically distributed),

$$\begin{aligned} P_1^*(\mathbf{x}, S) &= \frac{p_1}{L!/\{(L-1)!1!\}} + \frac{(L-1)!/\{(L-2)!1!\} p_2}{L!/\{(L-2)!2!\}} + \\ &\quad \frac{(L-1)!/\{(L-3)!2!\} p_3}{L!/\{(L-3)!3!\}} + \dots + p_L \\ &= \sum_{i=1}^L \frac{i p_i}{L} \\ &= \frac{m}{L}. \end{aligned}$$

■

If  $(E, \mathcal{E})$  is a metric measurable space, then define  $\mathcal{B}_b(E)$  (resp.  $\mathcal{B}_+(E)$ ) to be the set of bounded (resp. non-negative) real-valued measurable functions on  $E$ . The set  $\mathcal{C}_b(E)$  (resp.  $\mathcal{C}_+(E)$ ) denotes the set of bounded (resp. non-negative) real-valued continuous functions defined on  $E$ . For a finite measure  $\lambda$  on  $(E, \mathcal{E})$  and  $f \in \mathcal{B}_b(E) \cup \mathcal{B}_+(E)$  we shall write

$$\langle \lambda, f \rangle = \int f(\mathbf{x}) d\lambda(\mathbf{x}).$$

Additionally, we write  $\mathbb{E}_{\mathbf{x}}$  when  $\mathbf{X}_\emptyset = \mathbf{x}$ . We end this preamble with the definition of ergodicity for the Markov chain  $\{\mathbf{K}_j, j \in \mathbb{N}\}$ .

**Definition 11.1.** *The Markov chain  $\{\mathbf{K}_j, j \in \mathbb{N}\}$  is ergodic if there exists a probability measure  $\mu$  on  $(S, \mathcal{S})$  such that for all  $f \in \mathcal{C}_b(S)$  and all  $\mathbf{x} \in S$ ,*

$$\lim_{j \rightarrow \infty} \mathbb{E}_{\mathbf{x}} \{f(\mathbf{K}_j)\} = \langle \mu, f \rangle.$$

Before tackling the strong law of large numbers for the branching process covariate model, we reproduce a helpful Lemma from Delmas and Marsalle (2010) with slight adaptations.

**Lemma 11.6.** For  $f \in \mathcal{B}_b(S) \cup \mathcal{B}_+(S)$ ,

$$\begin{aligned} \mathbb{E} \{f(\mathbf{K}_j)\} &\stackrel{(a)}{=} m^{-j} \sum_{i \in \mathbb{G}_j} \mathbb{E} \{f(\mathbf{X}_i) \mathbb{I}(i \in \mathbb{T}^*)\} \stackrel{(b)}{=} \frac{\sum_{i \in \mathbb{G}_j} f(\mathbf{X}_i) \mathbb{I}(i \in \mathbb{T}^*)}{\sum_{i \in \mathbb{G}_j} \mathbb{P}(i \in \mathbb{T}^*)} \\ &= \mathbb{E} \{f(\mathbf{X}_I) \mid I \in \mathbb{T}^*\}. \end{aligned}$$

where  $I$  is a uniform random variable on  $\mathbb{G}_j$  independent of  $\mathbf{X}$ .

*Proof.* We consider equality (a). Recall that  $\mathbf{K}_0$  has distribution  $\nu$ . For  $i \in \mathbb{G}_j$ , we know

$$\mathbb{E} \{f(\mathbf{X}_i) \mathbb{I}(i \in \mathbb{T}^*)\} = \mathbb{E} \{f(\mathbf{X}_i) \mathbb{I}(\mathbf{X}_i \neq \delta)\} = \langle \nu, (P_1^*)^j f \rangle,$$

following from Equation 28 and the definition of  $P_1^*$ . Consequently,

$$\begin{aligned} \sum_{i \in \mathbb{G}_j} \mathbb{E} \{f(\mathbf{X}_i) \mathbb{I}(i \in \mathbb{T}^*)\} &= \sum_{i \in \{1, 2, \dots, L\}^j} \langle \nu, (P_1^*)^j f \rangle = \langle \nu, (LP_1^*)^j f \rangle = m^j \langle \nu, Q^j f \rangle \\ &= m^j \mathbb{E} \{f(\mathbf{K}_j)\}. \end{aligned}$$

This gives the first equality. Then take  $f = 1$  in the previous equality to get  $m^j = \sum_{i \in \mathbb{G}_j} \mathbb{P}(i \in \mathbb{T}^*)$  and equality (b). To prove Equality (c), we show by the law of total probability that

$$\begin{aligned} \mathbb{E} \{f(\mathbf{X}_I) \mid I \in \mathbb{T}^*\} &= \frac{f(\mathbf{X}_I) \mathbb{I}(I \in \mathbb{T}^*)}{\mathbb{P}(I \in \mathbb{T}^*)} \\ &= \frac{\sum_{i \in \mathbb{G}_j} f(\mathbf{X}_i) \mathbb{I}(i \in \mathbb{T}^*) \mathbb{P}(I = i)}{\sum_{i \in \mathbb{G}_j} \mathbb{P}(i \in \mathbb{T}^*) \mathbb{P}(I = i)} \\ &= \frac{\sum_{i \in \mathbb{G}_j} f(\mathbf{X}_i) \mathbb{I}(i \in \mathbb{T}^*)}{\sum_{i \in \mathbb{G}_j} \mathbb{P}(i \in \mathbb{T}^*)}. \end{aligned}$$

■

We recall that  $\nu$  denotes the distribution of  $\mathbf{X}_0$ . Any function  $f$  defined on  $S$  is

extended to  $\bar{S}$  by  $f(\delta) = 0$ . Let  $F$  be a vector subspace of  $\mathcal{B}(S)$  s.t.

1.  $F$  contains the constants
2.  $F^2 \triangleq \{f^2 : f \in F\} \subset F$
3.  $F \otimes F \subset L^1(P(\mathbf{x}, \cdot))$  for all  $x \in S$  and  $P(f_0 \otimes f_1) \in F$  for all  $f_0, f_1 \in F$
4. For  $\delta \in [0, 1]$ ,  $F \subset L^1(P_1^*(\mathbf{x}, \cdot))$  for all  $\mathbf{x} \in S$  and  $P_1^*(f) \in F$  for all  $f \in F$
5. There exists a probability measure  $\mu$  on  $(S, \mathcal{S})$  such that  $F \subset L^1(\mu)$  and  $\lim_{j \rightarrow \infty} \mathbb{E}_{\mathbf{x}} \{f(\mathbf{K}_j)\} = \langle \mu, f \rangle$  for all  $\mathbf{x} \in S$  and  $f \in F$
6. For all  $f \in F$ , there exists  $g \in F$  such that for all  $j \in \mathbb{N}$ ,  $|Q^j f| \leq g$ .
7.  $F \subset L^1(\nu)$

By convention, a function defined on  $\bar{S}$  is said to belong to  $F$  if its restriction to  $S$  belongs to  $F$ . **Note that if  $\mathbf{K}_j$  is ergodic and  $\mathbf{x} \rightarrow P^*g$  is continuous on  $S$  for all  $g \in \mathcal{C}_b(\bar{S}^2)$  then the set  $F = \mathcal{C}_b(S)$  fulfills Properties (1)-(7). Additionally, the stipulation that  $\mathbf{x} \rightarrow P^*g$  is continuous on  $S$  for all  $g \in \mathcal{C}_b(\bar{S}^2)$  is satisfied if  $\mathbf{K}_j$  has a continuous density (because integrals of continuous functions are continuous).**

---

**The Covariate Process of Equation 6.** First, note that any continuous  $f$  is automatically bounded since  $\mathcal{X}$  is bounded. Additionally,  $P^*g$  is an integral over a continuous density, so we know that  $\mathbf{x} \rightarrow P^*g$  is continuous on  $S$  for all  $g \in \mathcal{C}_b(\bar{S}^2)$ . Consequently, in order for  $F$  to satisfy Properties (1)-(7) above for the covariate process specified in Equation 6, we simply require that  $F$  is the space of continuous functions and  $\mathbf{K}_j$  is Ergodic.

Assume that the probability distributions (and expectations) of this section are with respect to the true branching parameters,  $\beta^*$ , and the uniform stabilizing policy,  $\tilde{\pi}$ . We now examine the covariate process specified in Equation 6

under  $\tilde{\pi}$ , in which the coupon allocations are assigned uniformly at random. Define  $\mathcal{R}^p$  as the Borel  $\sigma$ -algebra generated by  $\mathbb{R}^p$ . For  $\mathbf{x} \in \mathbb{R}^p$  and  $B \subset \mathcal{R}^p$ , we know that

$$P_1(\mathbf{x}, B) = Q(\mathbf{x}, B) = \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \text{Normal}(\boldsymbol{\phi}_{\mathbf{a}} + G_{\mathbf{a}}\mathbf{x}, \Sigma_{\mathbf{a}})$$

Consequently, the auxiliary Markov chain  $(\mathbf{K}_j, j \in \mathbb{N})$  is

$$f(\mathbf{K}_j \mid \mathbf{K}_{j-1} = \mathbf{k}_{j-1}) = \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \text{Normal}(\boldsymbol{\phi}_{\mathbf{a}} + G_{\mathbf{a}}\mathbf{k}_{j-1}, \Sigma_{\mathbf{a}}).$$

This is a Markov-Switching Autoregressive Model (Hamilton, 1989; Francq and Zakoian, 2001; Stelzer, 2009). According to Fong et al. (2007), Condition (C6), which states that

$$\frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \log \|G_{\mathbf{a}}\|_2 < 0,$$

implies that the  $(\mathbf{K}_j, j \in \mathbb{N})$  is Ergodic.

We label the stationary distribution for  $\mathbf{K}_j$  as  $\mu$  such that for all  $\mathbf{x} \in S$ ,

$$\lim_{j \rightarrow \infty} \mathbb{E}_{\mathbf{x}} \{f(\mathbf{K}_j)\} = \langle \mu, f \rangle.$$

We lower bound the variance (with the law of total variation) of the stationary distribution,  $\mu$ ,

$$\text{Var}_{\mu}(\mathbf{K}_j) \succeq \text{Var}_{\mu} \{\mathbb{E}(\mathbf{K}_j \mid \mathbf{K}_{j-1})\} \succeq \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \Sigma_{\mathbf{a}}.$$

Define  $\Sigma_{\mathbf{k}} \triangleq \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \Sigma_{\mathbf{a}}$ .

We now prove a Weak Law of Large numbers over the function class  $F$ . We again follow Delmas and Marsalle (2010) with slight additions. Because  $(\mathbf{K}_j, j \in \mathbb{N})$  is Ergodic and  $\mathbf{K}_j$  has a continuous density for all  $j \in \mathbb{N}$ , we only need  $F = \mathcal{C}_b(S)$  to satisfy Properties (1)-(7).

**Theorem 11.1.** *Let  $\{\mathbf{X}_i, i \in \mathbb{T}^*\}$  be the covariate process specified in Equation 6. Let*

$F = C_b(S)$ , which implies it satisfies Properties (1)-(7). Then,

$$\left\{ \frac{1}{m^n} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i), j \in \mathbb{N} \right\} \rightarrow \langle \mu, f \rangle_{\mathcal{S}}$$

in  $L^2$  as  $j \rightarrow \infty$ , where  $\mathcal{S}$  is defined by Equation 29.

*Proof.* We first assume that  $\langle \mu, f \rangle = 0$ . We have

$$\begin{aligned} \left\| \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) \right\|_2^2 &= \mathbb{E} \left[ \left( \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) \mathbb{I}(i \in \mathbb{T}^*) \right)^2 \right] \\ &= \sum_{i \in \mathbb{G}_j^*} \mathbb{E} \{ f^2(\mathbf{X}_i) \mathbb{I}(i \in \mathbb{T}^*) \} + B_j \\ &= m^j \mathbb{E} \{ f^2(\mathbf{K}_j) \} + B_j, \end{aligned}$$

with  $B_j = \sum_{(p,q) \in \mathbb{G}_j^2, i \neq j} \mathbb{E} [f(\mathbf{X}_p)f(\mathbf{X}_q)\mathbb{I}\{(p,q) \in \mathbb{T}^{*2}\}]$ . We used Lemma 11.6 for the last equality. For  $p, q \in \mathbb{G}_j^*$ , define  $p \wedge q$  as the most recent common ancestor of  $p$  and  $q$ . We compute  $B_q$  by decomposing the sum according to  $k = p \wedge q$ :  $B_j = \sum_{r=0}^{j-1} \sum_{k \in \mathbb{G}_r} C_k$  where

$$C_k = \sum_{(p,q) \in \mathbb{G}_j^2, p \wedge q = k} \mathbb{E} [f(\mathbf{X}_p)f(\mathbf{X}_q)\mathbb{I}\{(p,q) \in \mathbb{T}^{*2}\}].$$

If  $|k| = q - 1$ , we get

$$\begin{aligned} C_k &= \sum_{(p,q) \in \mathbb{G}_1^2, p \wedge q = \emptyset} \mathbb{E} \left( \mathbb{E}_{\mathbf{X}_k} [f(\mathbf{X}_{kp})f(\mathbf{X}_{kq})\mathbb{I}\{(kp,kq) \in \mathbb{T}^{*2}\}] \mathbb{I}(k \in \mathbb{T}^*) \right) \\ &= L(L-1) \mathbb{E} \{ P_2^*(f \otimes f)(\mathbf{X}_k) \mathbb{I}(k \in \mathbb{T}^*) \} \end{aligned}$$

If  $|k| < q - 1$ , we set  $r = |k|$  and derive

$$\begin{aligned}
C_k &= \sum_{(w,z) \in \mathbb{G}_1^2, w \wedge z = \emptyset} \sum_{(p,q) \in \mathbb{G}_{j-r-1}^2} \mathbb{E} \left[ \mathbb{E}_{\mathbf{X}_{kw}} \{f(\mathbf{X}_{kwp}) \mathbb{I}(kwp \in \mathbb{T}^*)\} \times \right. \\
&\quad \left. \mathbb{E}_{\mathbf{X}_{kz}} \{f(\mathbf{X}_{kzq}) \mathbb{I}(kzq \in \mathbb{T}^*)\} \mathbb{I}(kw \in \mathbb{T}^*, kz \in \mathbb{T}^*) \right] \\
&= \sum_{(w,z) \in \mathbb{G}_1^2, w \wedge z = \emptyset} \mathbb{E} \left( \left[ \sum_{p \in \mathbb{G}_{j-r-1}} \mathbb{E}_{\mathbf{X}_{kw}} \{f(\mathbf{X}_{kwp}) \mathbb{I}(kwp \in \mathbb{T}^*)\} \right] \times \right. \\
&\quad \left. \left[ \sum_{q \in \mathbb{G}_{j-r-1}} \mathbb{E}_{\mathbf{X}_{kz}} \{f(\mathbf{X}_{kzq}) \mathbb{I}(kzq \in \mathbb{T}^*)\} \right] \mathbb{I}(kw \in \mathbb{T}^*, kz \in \mathbb{T}^*) \right) \\
&\stackrel{(a)}{=} m^{2(j-r-1)} \sum_{(w,z) \in \mathbb{G}_1^2, w \wedge z = \emptyset} \mathbb{E} \left[ \mathbb{E}_{\mathbf{X}_{kw}} \{f(\mathbf{K}_{j-r-1})\} \times \right. \\
&\quad \left. \mathbb{E}_{\mathbf{X}_{kz}} \{f(\mathbf{K}_{j-r-1})\} \mathbb{I}(kw \in \mathbb{T}^*, kz \in \mathbb{T}^*) \right] \\
&\stackrel{(b)}{=} m^{2(j-r-1)} \sum_{(w,z) \in \mathbb{G}_1^2, w \wedge z = \emptyset} \mathbb{E} [Q^{j-r-1} f(\mathbf{X}_{kw}) Q^{j-r-1} f(\mathbf{X}_{kz}) \mathbb{I}(kw \in \mathbb{T}^*, kz \in \mathbb{T}^*)] \\
&= m^{2(j-r-1)} \sum_{(w,z) \in \mathbb{G}_1^2, w \wedge z = \emptyset} \mathbb{E} [(Q^{j-r-1} f \otimes Q^{j-r-1} f)(\mathbf{X}_{kw}, \mathbf{X}_{kz}) \mathbb{I}(kw \in \mathbb{T}^*, kz \in \mathbb{T}^*)] \\
&\stackrel{(c)}{=} m^{2(j-r-1)} \sum_{(w,z) \in \mathbb{G}_1^2, w \wedge z = \emptyset} \mathbb{E} [P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f)(\mathbf{X}_k) \mathbb{I}(k \in \mathbb{T}^*)] \\
&= L(L-1) m^{2(j-r-1)} \mathbb{E} [P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f)(\mathbf{X}_k) \mathbb{I}(k \in \mathbb{T}^*)]
\end{aligned}$$

Equality (a) follows from Lemma 11.6 and (b) through the definition of  $Q$ . Equality (c) follows from the definition of  $P_2^*$ . Consequently, by Lemma 11.6,

$$\begin{aligned}
B_j &= L(L-1) \sum_{r=0}^{j-1} m^{2(j-r-1)} \sum_{k \in \mathbb{G}_r} \mathbb{E} [P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f)(\mathbf{X}_k) \mathbb{I}(k \in \mathbb{T}^*)] \\
&\stackrel{(a)}{=} L(L-1) \sum_{r=0}^{j-1} m^{2(j-r-1)} m^r \langle \nu, Q^r P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f) \rangle \\
&= L(L-1) \sum_{r=0}^{j-1} m^{2j-r-2} \langle \nu, Q^r P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f) \rangle,
\end{aligned}$$

where equality (a) follows from Lemma 11.6. Therefore, we find that

$$\begin{aligned} \left\| m^{-j} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) \right\|_2^2 &= m^{-2j} \left\| \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) \right\|_2^2 \\ &= m^{-j} \mathbb{E} \{ f^2(\mathbf{K}_j) \} + \\ &\quad L(L-1)m^{-2} \sum_{r=0}^{j-1} m^{-r} \langle \nu, Q^r P^* (Q^{j-r-1} f \otimes Q^{j-r-1} f) \rangle \end{aligned}$$

Because  $f \in F$ , properties (2), (4), and (6) imply that  $E \{ f^2(\mathbf{K}_j) \} < \infty$  for any  $j \in \mathbb{N}$  and

$$\lim_{j \rightarrow \infty} m^{-j} \mathbb{E} \{ f^2(\mathbf{K}_j) \} = 0.$$

Properties (3), (4), and (5) along with  $\langle \mu, f \rangle = 0$  imply that  $P(Q^{j-r-1} f \otimes Q^{j-r-1} f)$  converges to 0 as  $j \rightarrow \infty$  (with  $r$  fixed) and is bounded uniformly in  $j > r$  by a function in  $F$ . Thus, properties (5) and (6) imply that  $\langle \nu, Q^r P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f) \rangle$  converges to 0 as  $j \rightarrow \infty$  (with  $r$  fixed) and is bounded uniformly in  $j > r$  by a fixed constant, say  $C$ .

For any  $\epsilon > 0$ , we can choose  $r_0$  such that  $\sum_{r > r_0} m^{-r} C \leq \epsilon/2$ . Additionally, choose  $j_0 > r_0$  such that for  $j \geq j_0$  and  $r \leq r_0$ , we have

$$|\langle \nu, Q^r P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f) \rangle| \leq \epsilon/(2r_0).$$

We get that for all  $j \geq j_0$

$$\sum_{r=0}^{j-1} m^{-r} |\langle \nu, Q^r P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f) \rangle| \leq \sum_{r=0}^{r_0} \epsilon/(2r_0) + \sum_{r > r_0}^{j-1} m^{-r} C \leq \epsilon/2 + \epsilon/2 = \epsilon.$$

Therefore,

$$\lim_{j \rightarrow \infty} \sum_{r=0}^{j-1} m^{-r} \langle \nu, Q^r P_2^* (Q^{j-r-1} f \otimes Q^{j-r-1} f) \rangle = 0.$$

We can now conclude that if  $\langle \mu, f \rangle = 0$ , then

$$\lim_{j \rightarrow \infty} \left\| m^{-j} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) \right\|_2 = 0.$$

We consider the case when  $\langle \mu, f \rangle$  is arbitrary. For any function  $f \in F$ , we can construct  $g = f - \langle \mu, f \rangle$ . We decompose as follows,

$$m^{-j} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) = m^{-j} \sum_{i \in \mathbb{G}_j^*} g(\mathbf{X}_i) + \langle \mu, f \rangle m^{-j} |\mathbb{G}_j^*|.$$

Because  $g \in F$  and  $\langle \mu, g \rangle = 0$ , we know that

$$\lim_{j \rightarrow \infty} \left\| m^{-j} \sum_{i \in \mathbb{G}_j^*} g(\mathbf{X}_i) \right\|_2 = 0.$$

Because  $(m^{-j} |\mathbb{G}_j^*|, j \geq 1)$  converges to  $L^2$  (and a.s.) to  $\mathcal{I}$ , we can conclude that

$$m^{-j} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) \xrightarrow{L^2} \langle \mu, f \rangle \mathcal{I}$$

as  $j \rightarrow \infty$ . ■

We now prove a similar result for the average over all individuals in  $\mathbb{T}_r^*$ , the first  $r$  generations. We set  $t_r \triangleq \mathbb{E}(|\mathbb{T}_r^*|)$  and recall that

$$t_r = \frac{m^{r+1} - 1}{m - 1}.$$

We now state a lemma that is a direct consequences of Lemma 8.1.

**Lemma 11.7.** *Let  $(v_r, r \in \mathbb{N})$  be a sequence of real numbers converging to  $a \in \mathbb{R}_+$ , and  $m$  be a real number such that  $m > 1$ . Let*

$$w_r = \sum_{j=0}^r m^{j-r-1} v_j.$$

*Then the sequence  $(w_r, r \in \mathbb{N})$  converges to  $a/(m - 1)$ .*

*Proof.* First we rearrange the sum  $w_r = \sum_{j=0}^r m^{j-r-1} v_j$ ,

$$w_r = \sum_{j=0}^r m^{j-r-1} v_j = \sum_{j=0}^r m^{-j-1} v_{r-j} = \frac{1}{m} \left( \sum_{j=0}^r m^{-j} \right) \left( \sum_{j=0}^r m^{-j} \right)^{-1} \sum_{j=0}^r m^{-j} v_{r-j}.$$

By Lemma 8.1,

$$\lim_{r \rightarrow \infty} \left( \sum_{j=0}^r m^{-j} \right)^{-1} \sum_{j=0}^r m^{-j} v_{r-j} = a.$$

By the formula for a geometric series,

$$\lim_{r \rightarrow \infty} \sum_{j=0}^r m^{-j} = \frac{1}{1 - \frac{1}{m}}.$$

Consequently,

$$\lim_{r \rightarrow \infty} w_r = \frac{1}{m} \left( \sum_{j=0}^r m^{-j} \right) \left( \sum_{j=0}^r m^{-j} \right)^{-1} \sum_{j=0}^r m^{-j} v_{r-j} = \frac{1}{m} \frac{1}{1 - \frac{1}{m}} a = \frac{a}{m-1}.$$

■

We now state the weak law of large numbers when averaging over the complete tree  $\mathbb{T}_r^*$  as  $r \rightarrow \infty$ .

**Theorem 11.2.** *Let  $\{\mathbf{X}_i, i \in \mathbb{T}^*\}$  be the covariate process specified in Equation 28. Let  $F$  satisfy (i)-(vi) and  $f \in F$ . Then,*

$$\left\{ \frac{1}{t_r} \sum_{i \in \mathbb{T}_r^*} f(\mathbf{X}_i), r \in \mathbb{N} \right\} \rightarrow \langle \mu, f \rangle_{\mathcal{S}}$$

in  $L^2$  as  $r \rightarrow \infty$ , where  $\mathcal{S}$  is defined by Equation 29.

*Proof.* Recalling that

$$t_r = \frac{m^{r+1} - 1}{m - 1}$$

we know that

$$\begin{aligned}
\left\| \frac{1}{t_r} \sum_{i \in \mathbb{T}_r^*} f(\mathbf{X}_i) - \langle \mu, f \rangle_{\mathcal{J}} \right\|_2 &= \left\| \sum_{j=0}^r \frac{m^j}{t_r} \left( \frac{1}{m^j} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) - \langle \mu, f \rangle_{\mathcal{J}} \right) \right\|_2 \\
&\leq \sum_{j=0}^r \frac{m^j}{t_r} \left\| \frac{1}{m^j} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) - \langle \mu, f \rangle_{\mathcal{J}} \right\|_2 \\
&= m^{r+1} \frac{m-1}{m^{r+1}-1} \sum_{j=0}^r m^{j-r-1} \left\| \frac{1}{m^j} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) - \langle \mu, f \rangle_{\mathcal{J}} \right\|_2 \\
&= \frac{m-1}{1-m^{-r-1}} \sum_{j=0}^r m^{j-r-1} \left\| \frac{1}{m^j} \sum_{i \in \mathbb{G}_j^*} f(\mathbf{X}_i) - \langle \mu, f \rangle_{\mathcal{J}} \right\|_2
\end{aligned}$$

Consequently,

$$\left\| \frac{1}{t_r} \sum_{i \in \mathbb{T}_r^*} f(\mathbf{X}_i) - \langle \mu, f \rangle_{\mathcal{J}} \right\|_2 \rightarrow 0$$

as  $r \rightarrow \infty$  by Theorem 11.1 and Lemma 11.7. ■

**Note that convergence in  $L^2$  implies convergence in probability, so we have proven a weak law of large numbers (WLLN) for the branching covariate process.**

---

Now that we have proven a WLLN for the branching covariate process implied by Equation 6, we return to the task of verifying Assumption 11. First, recall that

$$m = \sum_{m^v=1}^L m^v \frac{\lambda^{m^v}/m^v!}{\sum_{\ell=0}^{|\mathbf{A}^v|} (\lambda^\ell/\ell!)}$$

Additionally, define  $\Sigma_J = t_{J-1}$ . Note that the differentiability and moment conditions of the log-likelihood imply that

$$\eta_J = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \bar{\pi}} \left\{ \dot{l}_i(\beta^*) \dot{l}_i(\beta^*)^\top \mid \mathcal{F}_{j-1} \right\} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\beta^*, \bar{\pi}} \left\{ -\ddot{l}_i(\beta^*) \mid \mathcal{F}_{j-1} \right\}.$$

Consequently, we will use the Hessians calculated in Section 11.2 in the following theory.

We divide  $\eta_J$  into components corresponding to the different models that compose the

branching process,  $\eta_J = \text{diag}(\eta_J^{\mathbf{x}}, \eta_J^y, \eta_J^m, \eta_J^t)$ .

First, we analyze the covariate model component,  $\eta_J^{\mathbf{x}}$ . We define quantities

$$\mathbf{V}_j^{\mathbf{a}} \triangleq \sum_{i \in \mathcal{E}_{j-1}} M_i \mathbf{X}_i^* \mathbf{X}_i^{*\top} \mathbb{I}(\mathbf{A}_i = \mathbf{a}), \quad n_j^{\mathbf{a}} \triangleq \sum_{i \in \mathcal{E}_{j-1}} M_i \mathbb{I}(\mathbf{A}_i = \mathbf{a}),$$

$$\eta_J^{\mathbf{x}} \triangleq \sum_{j=1}^J \text{diag} \left[ \mathbb{E}_{\beta^*, \tilde{\pi}} \left\{ \left( \begin{array}{cc} \mathbf{V}_j^{\mathbf{a}} & 2\mathbf{V}_j^{\mathbf{a}} \mathbf{G}_{\mathbf{a}}^{*\top} \\ 2\mathbf{G}_{\mathbf{a}}^* \mathbf{V}_j^{\mathbf{a}} & 4\mathbf{G}_{\mathbf{a}}^* \mathbf{V}_j^{\mathbf{a}} \mathbf{G}_{\mathbf{a}}^{*\top} + 2n_j^{\mathbf{a}} \Sigma_{\mathbf{a}}^* \end{array} \right) \otimes \Sigma_{\mathbf{a}}^* \mid \mathcal{F}_{j-1} \right\}_{\mathbf{a} \in \mathcal{A}} \right].$$

We define

$$Q_{j,\mathbf{a}}^{\mathbf{x}} \triangleq \mathbb{E} \left\{ \sum_{i \in \mathcal{E}_{j-1}} M_i \mathbf{X}_i^* \mathbf{X}_i^{*\top} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \mid \mathcal{F}_{j-1} \right\} = \frac{1}{|\mathcal{A}|} \sum_{i \in \mathcal{E}_{j-1}} m \mathbf{X}_i^* \mathbf{X}_i^{*\top},$$

and

$$Q_{j,\mathbf{a}}^{\mathbf{a}} \triangleq \mathbb{E} \left\{ \sum_{i \in \mathcal{E}_{j-1}} M_i \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \mid \mathcal{F}_{j-1} \right\} = \frac{1}{|\mathcal{A}|} m \kappa_{j-1}.$$

Recall that

$$t_J = \frac{m^{J+1} - 1}{m - 1}.$$

Then

$$\begin{aligned} \Sigma_J^{-1/2} \eta_J^{\mathbf{x}} \Sigma_J^{-1/2} &= \\ \sum_{j=1}^J \Sigma_J^{-1/2} \text{diag} \left[ \left\{ \left( \begin{array}{cc} Q_{j,\mathbf{a}}^{\mathbf{x}} & 2Q_{j,\mathbf{a}}^{\mathbf{x}} \mathbf{G}_{\mathbf{a}}^{*\top} \\ 2\mathbf{G}_{\mathbf{a}}^* Q_{j,\mathbf{a}}^{\mathbf{x}} & 4\mathbf{G}_{\mathbf{a}}^* Q_{j,\mathbf{a}}^{\mathbf{x}} \mathbf{G}_{\mathbf{a}}^{*\top} + 2Q_{j,\mathbf{a}}^{\mathbf{a}} \Sigma_{\mathbf{a}}^* \end{array} \right) \otimes \Sigma_{\mathbf{a}}^* \right\}_{\mathbf{a} \in \mathcal{A}} \right] \Sigma_J^{-1/2} &= \\ \sum_{j=1}^J \text{diag} \left[ \left\{ \left( \begin{array}{cc} Q_{j,\mathbf{a}}^{\mathbf{x}} & 2Q_{j,\mathbf{a}}^{\mathbf{x}} \mathbf{G}_{\mathbf{a}}^{*\top} \\ 2\mathbf{G}_{\mathbf{a}}^* Q_{j,\mathbf{a}}^{\mathbf{x}} & 4\mathbf{G}_{\mathbf{a}}^* Q_{j,\mathbf{a}}^{\mathbf{x}} \mathbf{G}_{\mathbf{a}}^{*\top} + 2Q_{j,\mathbf{a}}^{\mathbf{a}} \Sigma_{\mathbf{a}}^* \end{array} \right) \otimes \Sigma_{\mathbf{a}}^* \right\}_{\mathbf{a} \in \mathcal{A}} \right] / t_{J-1}. \end{aligned}$$

We know that

$$\begin{aligned} Q_{j,\mathbf{a}}^{\mathbf{x}}/t_{J-1} &= \frac{1}{|\mathcal{A}|} \sum_{i \in \mathcal{E}_{j-1}} m \mathbf{X}_i^* \mathbf{X}_i^{*\top} / t_{J-1} \\ &= \frac{1}{|\mathcal{A}|} m \begin{pmatrix} \kappa_{j-1}/t_{J-1} & \sum_{i \in \mathcal{E}_{j-1}} \mathbf{X}_i^\top / t_{J-1} \\ \sum_{i \in \mathcal{E}_{j-1}} \mathbf{X}_i / t_{J-1} & \sum_{i \in \mathcal{E}_{j-1}} \mathbf{X}_i \mathbf{X}_i^\top / t_{J-1} \end{pmatrix}. \end{aligned}$$

Note that  $\mathbf{X}_i$  and  $\mathbf{X}_i \mathbf{X}_i^\top$  are both continuous, bounded functions of  $\mathbf{X}_i$  because  $\mathcal{X}$  is bounded (as previously mentioned). By Theorem 11.2, as  $J \rightarrow \infty$ ,

$$\begin{pmatrix} \sum_{j=1}^J \kappa_{j-1}/t_{J-1} & \sum_{i \in \mathcal{E}_{j-1}} \mathbf{X}_i^\top / t_{J-1} \\ \sum_{i \in \mathcal{E}_{j-1}} \mathbf{X}_i / t_{J-1} & \sum_{i \in \mathcal{E}_{j-1}} \mathbf{X}_i \mathbf{X}_i^\top / t_{J-1} \end{pmatrix} \xrightarrow{p} \mathcal{I} C_*^{\mathbf{x}},$$

where  $C_*^{\mathbf{x}}$  is a constant matrix. Additionally,

$$Q_{j,\mathbf{a}}^{\mathbf{a}} = \frac{1}{|\mathcal{A}|} m \kappa_{j-1} / t_{J-1} \xrightarrow{p} \frac{m}{|\mathcal{A}|} \mathcal{I}.$$

Consequently, we know that

$$\begin{aligned} &\Sigma_J^{-1/2} \eta_J^{\mathbf{x}} \Sigma_J^{-1/2} = \\ &\sum_{j=1}^J \text{diag} \left[ \mathbb{E}_{\beta^*, \tilde{\pi}} \left\{ \begin{pmatrix} \mathbf{V}_j^{\mathbf{a}} & 2\mathbf{V}_j^{\mathbf{a}} G_{\mathbf{a}}^{*\top} \\ 2G_{\mathbf{a}}^* \mathbf{V}_j^{\mathbf{a}} & 4G_{\mathbf{a}}^* \mathbf{V}_j^{\mathbf{a}} G_{\mathbf{a}}^{*\top} + 2n_j^{\mathbf{a}} \Sigma_{\mathbf{a}}^* \end{pmatrix} \otimes \Sigma_{\mathbf{a}}^* \mid \mathcal{F}_{j-1} \right\}_{\mathbf{a} \in \mathcal{A}} \right] / t_{J-1} \xrightarrow{p} C_*^{\mathbf{x}} \mathcal{I} \end{aligned}$$

as  $J \rightarrow \infty$ .

**Lemma 11.8.** *Under conditions (1)-(6), for any covariate parameters  $\beta_{\mathbf{x}} = \{\Sigma_{\mathbf{a}}, G_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}} \in \mathcal{B}$  and some  $\delta_{\beta_{\mathbf{x}}} > 0$ ,*

$$\begin{aligned} &\sigma_{\min} \left( \sum_{j=1}^J \mathbb{E}_{\beta^*, \tilde{\pi}} \left[ \text{diag} \left\{ \begin{pmatrix} \mathbf{V}_j^{\mathbf{a}} & 2\mathbf{V}_j^{\mathbf{a}} G_{\mathbf{a}}^\top \\ 2G_{\mathbf{a}} \mathbf{V}_j^{\mathbf{a}} & 4G_{\mathbf{a}} \mathbf{V}_j^{\mathbf{a}} G_{\mathbf{a}}^\top + 2n_j^{\mathbf{a}} \Sigma_{\mathbf{a}} \end{pmatrix} \otimes \Sigma_{\mathbf{a}} \right\}_{\mathbf{a} \in \mathcal{A}} / t_{J-1} \mid \mathcal{F}_{j-1} \right] \right) \\ &\geq \delta_{\beta_{\mathbf{x}}} \quad \text{w.p. 1} \end{aligned}$$

*Proof.* By Lemma 11.4, we know that

$$\begin{aligned} & \lim_{J \rightarrow \infty} \sigma_{\min} \left\{ \sum_{j=1}^J \left\{ \begin{pmatrix} Q_{j,\mathbf{a}}^{\mathbf{x}} & 2Q_{j,\mathbf{a}}^{\mathbf{x}} G_{\mathbf{a}}^{\top} \\ 2G_{\mathbf{a}} Q_{j,\mathbf{a}}^{\mathbf{x}} & 4G_{\mathbf{a}} Q_{j,\mathbf{a}}^{\mathbf{x}} G_{\mathbf{a}}^{\top} + 2Q_{j,\mathbf{a}}^{\mathbf{a}} \Sigma_{\mathbf{a}} \end{pmatrix} \otimes \Sigma_{\mathbf{a}} \right\} / t_{J-1} \right\} \geq \\ & \lim_{J \rightarrow \infty} \sigma_{\min} \left( \sum_{j=1}^J Q_{j,\mathbf{a}}^{\mathbf{x}} / t_{J-1} \right) \sigma_{\min} \left( \sum_{j=1}^J 2Q_{j,\mathbf{a}}^{\mathbf{a}} \Sigma_{\mathbf{a}} / t_{J-1} \right) \end{aligned}$$

because

$$\lim_{J \rightarrow \infty} \sum_{j=1}^J \left\{ 4G_{\mathbf{a}} Q_{j,\mathbf{a}}^{\mathbf{x}} G_{\mathbf{a}}^{\top} + 2Q_{j,\mathbf{a}}^{\mathbf{a}} \Sigma_{\mathbf{a}} - 4G_{\mathbf{a}} Q_{j,\mathbf{a}}^{\mathbf{x}} Q_{j,\mathbf{a}}^{\mathbf{x}^{-1}} Q_{j,\mathbf{a}}^{\mathbf{x}} G_{\mathbf{a}}^{\top} \right\} / t_{J-1} = \lim_{J \rightarrow \infty} \sum_{j=1}^J 2Q_{j,\mathbf{a}}^{\mathbf{a}} \Sigma_{\mathbf{a}} / t_{J-1}.$$

We know that  $\lim_{J \rightarrow \infty} \sum_{j=1}^J 2Q_{j,\mathbf{a}}^{\mathbf{a}} \Sigma_{\mathbf{a}} / t_{J-1} = 2m_{\mathcal{I}} \Sigma_{\mathbf{a}} / |\mathcal{A}|$  a.s., so we only need to prove that

$$\sum_{j=1}^J Q_{j,\mathbf{a}}^{\mathbf{x}} / t_{J-1} \xrightarrow{p} \frac{m_{\mathcal{I}}}{|\mathcal{A}|} \begin{pmatrix} 1 & \mathbb{E}_{\mu}(\mathbf{K}_i)^{\top} \\ \mathbb{E}_{\mu}(\mathbf{K}_i) & \mathbb{E}_{\mu}(\mathbf{K}_i \mathbf{K}_i^{\top}) \end{pmatrix}$$

is positive definite. By a second application of Lemma 11.4,

$$\begin{aligned} \frac{m_{\mathcal{I}}}{|\mathcal{A}|} \sigma_{\min} \left\{ \begin{pmatrix} 1 & \mathbb{E}_{\mu}(\mathbf{K}_i)^{\top} \\ \mathbb{E}_{\mu}(\mathbf{K}_i) & \mathbb{E}_{\mu}(\mathbf{K}_i \mathbf{K}_i^{\top}) \end{pmatrix} \right\} \frac{m_{\mathcal{I}}}{|\mathcal{A}|} & \geq \sigma_{\min}(1) \times \\ & \sigma_{\min} \left\{ \mathbb{E}_{\mu}(\mathbf{K}_i \mathbf{K}_i^{\top}) - \mathbb{E}_{\mu}(\mathbf{K}_i) \mathbb{E}_{\mu}(\mathbf{K}_i)^{\top} \right\} \\ & = \frac{m_{\mathcal{I}}}{|\mathcal{A}|} \sigma_{\min} \{ \text{Var}_{\mu}(\mathbf{K}_i) \} \\ & \succeq \frac{m_{\mathcal{I}}}{|\mathcal{A}|} \sigma_{\min}(\Sigma_{\mathbf{k}}). \end{aligned}$$

Note that there exists  $\delta_*$  such that  $\mathcal{I} > \delta_*$  by assumption. Using the properties of kronecker products, we know that defining

$$\epsilon_{\mathbf{a}} \triangleq 2 \frac{m^2 \delta_*^2}{|\mathcal{A}|^2} \sigma_{\min}(\Sigma_{\mathbf{k}}) \sigma_{\min}(\Sigma_{\mathbf{a}})^2,$$

and

$$\delta_{\beta \mathbf{x}} \triangleq \prod_{\mathbf{a} \in \mathcal{A}} \epsilon_{\mathbf{a}}$$

completes the proof. ■

By Lemma 11.8,  $\sigma_{\min}(\mathcal{I}C_*^{\mathbf{x}}) \geq \delta_{\beta_{\mathbf{x}}}$ . Consequently, Assumption 11 is satisfied for the covariate model.

Next, we analyze the reward model component,  $\eta_J^y$ .

$$\frac{\partial^2 \ell_J(\beta_y)}{\partial \beta_y \partial \beta_y^\top} = - \sum_{j=1}^J \sum_{i \in \mathcal{E}_j} \mathbf{Z}_i \mathbf{Z}_i^\top \left\{ \frac{1}{1 + \exp(-\mathbf{Z}_i^\top \beta_y)} \right\} \left\{ \frac{1}{1 + \exp(\mathbf{Z}_i^\top \beta_y)} \right\}.$$

Remember that  $\mathbf{Z}_i = \sum_{\mathbf{a} \in \mathcal{A}} h_{\mathbf{a}}(\mathbf{X}_i) \mathbb{I}(\mathbf{A}_{r^i} = \mathbf{a})$ , where  $h_{\mathbf{a}}$  is a continuous function in  $\mathbf{X}_i$  that depends on  $\mathbf{a} \in \mathcal{A}$ . Consequently,

$$\begin{aligned} \frac{\partial^2 \ell_J(\beta_y)}{\partial \beta_y \partial \beta_y^\top} &= - \sum_{j=1}^J \sum_{i \in \mathcal{E}_j} \sum_{\mathbf{a} \in \mathcal{A}} h_{\mathbf{a}}(\mathbf{X}_i) h_{\mathbf{a}}(\mathbf{X}_i)^\top \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_i)^\top \beta_y)} \right\} \times \\ &\quad \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_i)^\top \beta_y)} \right\} \mathbb{I}(\mathbf{A}_{r^i} = \mathbf{a}). \end{aligned}$$

We now express

$$\begin{aligned}
\eta_J^y &= \sum_{j=1}^J \mathbb{E} \left\{ \sum_{i \in \mathcal{E}_j} \sum_{\mathbf{a} \in \mathcal{A}} h_{\mathbf{a}}(\mathbf{X}_i) h_{\mathbf{a}}(\mathbf{X}_i)^\top \times \right. \\
&\quad \left. \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_i)^\top \boldsymbol{\beta}_y^*)} \right\} \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_i)^\top \boldsymbol{\beta}_y^*)} \right\} \times \right. \\
&\quad \left. \mathbb{I}(\mathbf{A}_{r^i} = \mathbf{a}) \mid \mathcal{F}_{j-1} \right\} \\
&= \sum_{j=1}^J \mathbb{E} \left\{ \sum_{i \in \mathcal{E}_j} \sum_{\mathbf{a} \in \mathcal{A}} \sum_{l \in \mathbb{G}_1} \mathbb{I}(il \in \mathbb{T}^*) h_{\mathbf{a}}(\mathbf{X}_i) h_{\mathbf{a}}(\mathbf{X}_i)^\top \times \right. \\
&\quad \left. \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_i)^\top \boldsymbol{\beta}_y^*)} \right\} \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_i)^\top \boldsymbol{\beta}_y^*)} \right\} \times \right. \\
&\quad \left. \mathbb{I}(\mathbf{A}_{r^i} = \mathbf{a}) \mid \mathcal{F}_{j-1} \right\} \\
&= \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l \in \mathbb{G}_1} \mathbb{E} \left\{ \mathbb{I}(il \in \mathbb{T}^*) \mathbb{E} \left\{ \sum_{\mathbf{a} \in \mathcal{A}} h_{\mathbf{a}}(\mathbf{X}_{il}) h_{\mathbf{a}}(\mathbf{X}_{il})^\top \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y^*)} \right\} \times \right. \right. \\
&\quad \left. \left. \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y^*)} \right\} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \mid \mathbf{X}_i, il \in \mathbb{T}^* \right\} \mid \mathbf{X}_i \right\} \\
&= \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l \in \mathbb{G}_1} \mathbb{E} \left\{ \mathbb{I}(il \in \mathbb{T}^*) \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a}' \in \mathcal{A}} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ h_{\mathbf{a}}(\mathbf{X}_{il}) h_{\mathbf{a}}(\mathbf{X}_{il})^\top \right. \right. \\
&\quad \left. \left. \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y^*)} \right\} \times \right. \right. \\
&\quad \left. \left. \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y^*)} \right\} \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \mid \mathbf{A}_i = \mathbf{a}', \mathbf{X}_i, il \in \mathbb{T}^* \right\} \mid \mathbf{X}_i \right\} \\
&= \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l \in \mathbb{G}_1} \mathbb{E} \left\{ \mathbb{I}(il \in \mathbb{T}^*) \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ h_{\mathbf{a}}(\mathbf{X}_{il}) h_{\mathbf{a}}(\mathbf{X}_{il})^\top \right. \right. \\
&\quad \left. \left. \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y^*)} \right\} \times \right. \right. \\
&\quad \left. \left. \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y^*)} \right\} \mid \mathbf{A}_i = \mathbf{a}, \mathbf{X}_i, il \in \mathbb{T}^* \right\} \mid \mathbf{X}_i \right\}
\end{aligned}$$

We label

$$f(\mathbf{X}_{il}) = \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ h_{\mathbf{a}}(\mathbf{X}_{il}) h_{\mathbf{a}}(\mathbf{X}_{il})^\top \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y^*)} \right\} \times \right. \\ \left. \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y^*)} \right\} \mid \mathbf{A}_i = \mathbf{a}, \mathbf{X}_i \right\},$$

where  $f$  is continuous because  $h_{\mathbf{a}}$  is continuous (and it is a composition of continuous functions). Consequently,

$$\eta_J^y = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \sum_{l \in \mathbb{G}_1} \mathbb{E} \{ \mathbb{I}(il \in \mathbb{T}^*) f(\mathbf{X}_{il}) \mid \mathbf{X}_i \} \\ = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} LP_1^* \{ f(\mathbf{X}_i) \}.$$

Lastly, we can define  $f^*(\mathbf{X}_i) = LP_1^* \{ f(\mathbf{X}_i) \}$  and

$$\eta_J^y = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} f^*(\mathbf{X}_i).$$

Note that  $f^*$  is continuous and bounded since compositions and integrals of continuous functions are continuous. It is now clear that

$$\eta_J^y / t_{J-1} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} f^*(\mathbf{X}_i) / t_{J-1} \xrightarrow{p} \mathcal{I} C_*^y,$$

where  $C_*^y$  is a constant matrix.

**Lemma 11.9.** *Under conditions (1)-(6), for any  $\boldsymbol{\beta}_y \in \mathcal{B}$ , there exists  $\delta_{\boldsymbol{\beta}_y}$  such that*

$$\lim_{J \rightarrow \infty} \sigma_{\min} \left( \sum_{j=1}^J \mathbb{E} \left\{ \sum_{i \in \mathcal{E}_j} \sum_{\mathbf{a} \in \mathcal{A}} h_{\mathbf{a}}(\mathbf{X}_i) h_{\mathbf{a}}(\mathbf{X}_i)^\top \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_i)^\top \boldsymbol{\beta}_y)} \right\} \times \right. \right. \\ \left. \left. \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_i)^\top \boldsymbol{\beta}_y)} \right\} \times \right. \right. \\ \left. \left. \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \mid \mathcal{F}_{j-1} \right\} / t_{J-1} \right) \geq \delta_{\boldsymbol{\beta}_y} \quad \text{w.p. } 1$$

*Proof.* Label

$$\sigma_y = \lim_{J \rightarrow \infty} \sigma_{\min} \left( \sum_{j=1}^J \mathbb{E} \left\{ \sum_{i \in \mathcal{E}_j} \sum_{\mathbf{a} \in \mathcal{A}} h_{\mathbf{a}}(\mathbf{X}_i) h_{\mathbf{a}}(\mathbf{X}_i)^\top \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_i)^\top \boldsymbol{\beta}_y)} \right\} \times \right. \right. \\ \left. \left. \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_i)^\top \boldsymbol{\beta}_y)} \right\} \times \right. \right. \\ \left. \left. \mathbb{I}(\mathbf{A}_i = \mathbf{a}) \mid \mathcal{F}_{j-1} \right\} \right).$$

We start with analyzing the quantity

$$\sum_{l \in \mathbb{G}_1} \mathbb{E} \left\{ \mathbb{I}(il \in \mathbb{T}^*) \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ h_{\mathbf{a}}(\mathbf{X}_{il}) h_{\mathbf{a}}(\mathbf{X}_{il})^\top \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y)} \right\} \times \right. \right. \\ \left. \left. \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{X}_{il})^\top \boldsymbol{\beta}_y)} \right\} \mid il \in \mathbb{T}^*, \mathbf{A}_i = \mathbf{a}, \mathbf{X}_i \right\} \mid \mathbf{X}_i \right\},$$

Because  $\mathcal{X}$  is compact and  $h_{\mathbf{a}}$  is continuous, we know that

$$c_{\exp} \triangleq \min_{\mathbf{x} \in \mathcal{X}} \min_{\mathbf{a} \in \mathcal{A}} \left\{ \frac{1}{1 + \exp(-h_{\mathbf{a}}(\mathbf{x})^\top \boldsymbol{\beta}_y)} \right\} \left\{ \frac{1}{1 + \exp(h_{\mathbf{a}}(\mathbf{x})^\top \boldsymbol{\beta}_y)} \right\},$$

where  $c_{\exp} > 0$  by the extreme value theorem. Consequently,

$$\succ c_{\exp} \sum_{l \in \mathbb{G}_1} \mathbb{E} \left\{ \mathbb{I}(il \in \mathbb{T}^*) \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ h_{\mathbf{a}}(\mathbf{X}_{il}) h_{\mathbf{a}}(\mathbf{X}_{il})^\top \mid \mathbf{A}_i = \mathbf{a}, \mathbf{X}_i, il \in \mathbb{T}^* \right\} \mid \mathbf{X}_i \right\}.$$

To progress we need to be more explicit about  $h_{\mathbf{a}}$ . In our model  $Z_i$  is an interaction between  $X_i$  and an indicator of the coupon type given to participant  $i$ . We demonstrate the positive definite property for a paradigm where we have two coupon types  $\mathcal{A} = \{\mathbf{a}, \mathbf{a}'\}$ , making  $\mathbf{Z}_i = (1, \mathbf{X}_i, \mathbf{X}_i \mathbb{I}(\mathbf{A}_i = \mathbf{a}))$ . However, this logic extends to a larger number of

coupon types. We find that

$$\begin{aligned}
& \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ \mathbf{Z}_{il} \mathbf{Z}_{il}^\top \mid il \in \mathbb{T}^*, \mathbf{A}_i = \mathbf{a}, \mathbf{X}_i \right\} \\
&= \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ \mathbf{Z}_{il} \mathbf{Z}_{il}^\top \mid \mathbf{A}_i = \mathbf{a}, \mathbf{X}_i \right\} \\
&= \mathbb{E} \left\{ \begin{pmatrix} 1 & \mathbf{X}_{il}^\top & 0 \\ \mathbf{X}_{il}^\top & \mathbf{X}_{il} \mathbf{X}_{il}^\top & 0 \\ 0 & 0 & 0 \end{pmatrix} \mid \mathbf{X}_i, \mathbf{A}_i = \mathbf{a}' \right\} \\
&\quad + \mathbb{E} \left\{ \begin{pmatrix} 1 & \mathbf{X}_{il}^\top & \mathbf{X}_{il}^\top \\ \mathbf{X}_{il}^\top & \mathbf{X}_{il} \mathbf{X}_{il}^\top & \mathbf{X}_{il} \mathbf{X}_{il}^\top \\ \mathbf{X}_{il}^\top & \mathbf{X}_{il} \mathbf{X}_{il}^\top & \mathbf{X}_{il} \mathbf{X}_{il}^\top \end{pmatrix} \mid \mathbf{X}_i, \mathbf{A}_i = \mathbf{a} \right\}.
\end{aligned}$$

We label  $\mu_{\mathbf{a}, \mathbf{x}} = \mathbb{E} \{ \mathbf{X}_{il} \mid \mathbf{X}_i = \mathbf{x}, \mathbf{A}_i = \mathbf{a} \}$  and  $\Delta_{\mathbf{a}} = \mathbb{E} \{ \mathbf{X}_{il} \mathbf{X}_{il}^\top \mid \mathbf{X}_i = \mathbf{x}, \mathbf{A}_i = \mathbf{a} \}$ . Note that  $\Delta_{\mathbf{a}}$  does not depend on  $\mathbf{X}_i$ . We find that

$$\begin{aligned}
\Delta_0 &\triangleq \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ \mathbf{Z}_{il} \mathbf{Z}_{il}^\top \mid \mathbf{A}_i = \mathbf{a}, \mathbf{X}_i = \mathbf{x} \right\} = \begin{pmatrix} 1 & \mu_{\mathbf{a}', \mathbf{x}}^\top & 0 \\ \mu_{\mathbf{a}', \mathbf{x}} & \Delta_{\mathbf{a}'} & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 1 & \mu_{\mathbf{a}, \mathbf{x}}^\top & \mu_{\mathbf{a}, \mathbf{x}}^\top \\ \mu_{\mathbf{a}, \mathbf{x}} & \Delta_{\mathbf{a}} & \Delta_{\mathbf{a}} \\ \mu_{\mathbf{a}, \mathbf{x}} & \Delta_{\mathbf{a}} & \Delta_{\mathbf{a}} \end{pmatrix} \\
&= \begin{pmatrix} 2 & \mu_{\mathbf{a}', \mathbf{x}}^\top + \mu_{\mathbf{a}, \mathbf{x}}^\top & \mu_{\mathbf{a}, \mathbf{x}}^\top \\ \mu_{\mathbf{a}', \mathbf{x}} + \mu_{\mathbf{a}, \mathbf{x}} & \Delta_{\mathbf{a}'} + \Delta_{\mathbf{a}} & \Delta_{\mathbf{a}} \\ \mu_{\mathbf{a}, \mathbf{x}} & \Delta_{\mathbf{a}} & \Delta_{\mathbf{a}} \end{pmatrix}.
\end{aligned}$$

By Lemma 11.4, we know that  $\sigma_{\min}(\Delta_0) \geq 2\sigma_{\min}(\Delta_1)$ , where

$$\begin{aligned}\Delta_1 &\triangleq \begin{pmatrix} \Sigma_{\mathbf{a}} + \Sigma_{\mathbf{a}'} + \frac{(\mu_{\mathbf{a},\mathbf{x}} - \mu_{\mathbf{a}',\mathbf{x}})(\mu_{\mathbf{a},\mathbf{x}} - \mu_{\mathbf{a}',\mathbf{x}})^\top}{2} & \Sigma_{\mathbf{a}} + \frac{(\mu_{\mathbf{a},\mathbf{x}} - \mu_{\mathbf{a}',\mathbf{x}})\mu_{\mathbf{a},\mathbf{x}}^\top}{2} \\ \Sigma_{\mathbf{a}} + \frac{\mu_{\mathbf{a},\mathbf{x}}(\mu_{\mathbf{a},\mathbf{x}} - \mu_{\mathbf{a}',\mathbf{x}})^\top}{2} & \Sigma_{\mathbf{a}} + \frac{\mu_{\mathbf{a},\mathbf{x}}\mu_{\mathbf{a},\mathbf{x}}^\top}{2} \end{pmatrix} \\ &= \begin{pmatrix} \Sigma_{\mathbf{a}} + \Sigma_{\mathbf{a}'} & \Sigma_{\mathbf{a}} \\ \Sigma_{\mathbf{a}} & \Sigma_{\mathbf{a}} \end{pmatrix} + \begin{pmatrix} \frac{(\mu_{\mathbf{a},\mathbf{x}} - \mu_{\mathbf{a}',\mathbf{x}})(\mu_{\mathbf{a},\mathbf{x}} - \mu_{\mathbf{a}',\mathbf{x}})^\top}{2} & \frac{(\mu_{\mathbf{a},\mathbf{x}} - \mu_{\mathbf{a}',\mathbf{x}})\mu_{\mathbf{a},\mathbf{x}}^\top}{2} \\ \frac{\mu_{\mathbf{a},\mathbf{x}}(\mu_{\mathbf{a},\mathbf{x}} - \mu_{\mathbf{a}',\mathbf{x}})^\top}{2} & \frac{\mu_{\mathbf{a},\mathbf{x}}\mu_{\mathbf{a},\mathbf{x}}^\top}{2} \end{pmatrix} \\ &\succcurlyeq \begin{pmatrix} \Sigma_{\mathbf{a}} + \Sigma_{\mathbf{a}'} & \Sigma_{\mathbf{a}} \\ \Sigma_{\mathbf{a}} & \Sigma_{\mathbf{a}} \end{pmatrix}.\end{aligned}$$

We label

$$\Delta_2 \triangleq \begin{pmatrix} \Sigma_{\mathbf{a}} + \Sigma_{\mathbf{a}'} & \Sigma_{\mathbf{a}} \\ \Sigma_{\mathbf{a}} & \Sigma_{\mathbf{a}} \end{pmatrix}.$$

By Lemma 11.4 again, we find that

$$\begin{aligned}\sigma_{\min}\{\Delta_2\} &\geq \sigma_{\min}(\Sigma_{\mathbf{a}})\sigma_{\min}(\Sigma_{\mathbf{a}} + \Sigma_{\mathbf{a}'} - \Sigma_{\mathbf{a}}\Sigma_{\mathbf{a}}^{-1}\Sigma_{\mathbf{a}}) \\ &= \sigma_{\min}(\Sigma_{\mathbf{a}})\sigma_{\min}(\Sigma_{\mathbf{a}'}) > 0.\end{aligned}$$

We can conclude that

$$\begin{aligned}&c_{\text{exp}} \sum_{l \in \mathbb{G}_1} \mathbb{E} \left\{ \mathbb{I}(il \in \mathbb{T}^*) \frac{1}{|\mathcal{A}|} \sum_{\mathbf{a} \in \mathcal{A}} \mathbb{E} \left\{ h_{\mathbf{a}}(\mathbf{X}_{il}) h_{\mathbf{a}}(\mathbf{X}_{il})^\top \mid \mathbf{A}_i = \mathbf{a}, \mathbf{X}_i \right\} \mid \mathbf{X}_i \right\} \\ &\succcurlyeq c_{\text{exp}} \sum_{l \in \mathbb{G}_1} \mathbb{E} \left\{ \mathbb{I}(il \in \mathbb{T}^*) \Delta_2 \mid \mathbf{X}_i \right\} \\ &\succcurlyeq c_{\text{exp}} m \Delta_2.\end{aligned}$$

Therefore,

$$\begin{aligned}\sigma_y &\geq \lim_{J \rightarrow \infty} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \frac{1}{t_{J-1}} c_{\text{exp}} m \sigma_{\min}(\Delta_2) \\ &\geq \lim_{J \rightarrow \infty} \frac{\bar{\kappa}_J}{t_{J-1}} c_{\text{exp}} m \sigma_{\min}(\Delta_2) \\ &= \mathcal{I} c_{\text{exp}} m \sigma_{\min}(\Delta_2).\end{aligned}$$

Note that there exists  $\delta_*$  such that  $\mathcal{J} > \delta_*$  by assumption. Defining

$$\delta_{\beta_y} \triangleq \delta^* c_{\text{exp}} m \sigma_{\min}(\Delta_2)$$

completes the proof. ■

We conclude that  $\sigma_{\min}(\mathcal{J}C_*^y) \geq \delta_{\beta_y}^*$  and we can conclude **Assumption 11** is satisfied for the reward model component of the branching process.

Next, we analyze the time model component,  $\eta_J^t$ . The Hessian for this part of the model is

$$\frac{\partial^2 \ell_J(\zeta)}{\partial \zeta^2} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} -M_i \left( \frac{1}{\zeta^2} + \frac{ae^{-\zeta a} - b^2 e^{-\zeta b}}{[e^{-\zeta a} - e^{-\zeta b}]^2} \right).$$

We get that

$$\begin{aligned} \lim_{J \rightarrow \infty} \eta_J^t / t_{J-1} &= \lim_{J \rightarrow \infty} \bar{\kappa}_J / t_{J-1} \left( \frac{1}{\zeta^{*2}} + \frac{ae^{-\zeta^* a} - b^2 e^{-\zeta^* b}}{[e^{-\zeta^* a} - e^{-\zeta^* b}]^2} \right) m \\ &= \mathcal{J} \left( \frac{1}{\zeta^{*2}} + \frac{ae^{-\zeta^* a} - b^2 e^{-\zeta^* b}}{[e^{-\zeta^* a} - e^{-\zeta^* b}]^2} \right) m \\ &> 0. \end{aligned}$$

Where the last inequality follows because there exists  $\delta_*$  such that  $\mathcal{J} > \delta_*$  by assumption. We can conclude **Assumption 11** is satisfied for the time model component of the branching process.

Lastly, we analyze the family size model component,  $\eta_J^m$ .

$$\frac{\partial^2 \ell_J(\tau)}{\partial \tau^2} = \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} -\tau^2 \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau \ell} / \ell! \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell} / \ell! \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\}^2}.$$

We find

$$\begin{aligned}
& \lim_{J \rightarrow \infty} \eta_J / t_{J-1} \\
&= \lim_{J \rightarrow \infty} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \tau^{*2} \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau^* \ell / \ell!} \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau^* \ell / \ell!} \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau^* \ell / \ell!} \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau^* \ell / \ell!} \right\}^2} / t_{J-1} \\
&= \lim_{J \rightarrow \infty} \frac{\bar{\kappa}_J}{t_{J-1}} \tau^{*2} \frac{\left\{ \sum_{\ell=k}^L e^{\tau^* \ell / \ell!} \right\} \left\{ \sum_{\ell=k}^L \ell^2 e^{\tau^* \ell / \ell!} \right\} - \left\{ \sum_{\ell=k}^L \ell e^{\tau^* \ell / \ell!} \right\}^2}{\left\{ \sum_{\ell=k}^L e^{\tau^* \ell / \ell!} \right\}^2} \\
&= \mathcal{I} \tau^{*2} \frac{\left\{ \sum_{\ell=k}^L e^{\tau^* \ell / \ell!} \right\} \left\{ \sum_{\ell=k}^L \ell^2 e^{\tau^* \ell / \ell!} \right\} - \left\{ \sum_{\ell=k}^L \ell e^{\tau^* \ell / \ell!} \right\}^2}{\left\{ \sum_{\ell=k}^L e^{\tau^* \ell / \ell!} \right\}^2}
\end{aligned}$$

Note that there exists  $\delta_*$  such that  $\mathcal{I} > \delta_*$  by assumption. We can conclude **Assumption 11** is satisfied for the family model component of the branching process.

In conclusion, we have confirmed that

$$\lim_{J \rightarrow \infty} \eta_J / t_{J-1} = \mathcal{I} \mathbb{C}$$

where  $\mathcal{I} \mathbb{C} > 0$  w.p. 1 on the event  $E_{\mathcal{I}}$ . We establish one more lemma that will be useful in the consistency proof of Section 11.4.

**Lemma 11.10.** *Recall in the model described in Equation 6,  $\beta_t \triangleq \zeta$  and  $\beta_m \triangleq \tau$ . Under conditions (1)-(6), for any  $\zeta, \tau \in \mathcal{B}$ , there exists  $\delta_{\beta_t}, \delta_{\beta_m} > 0$ ,*

$$\begin{aligned}
& \lim_{J \rightarrow \infty} \sum_{j=1}^J \mathbb{E} \left[ \sum_{i \in \mathcal{E}_{j-1}} M_i \left( \frac{1}{\zeta^2} + \frac{a e^{-\zeta a} - b^2 e^{-\zeta b}}{[e^{-\zeta a} - e^{-\zeta b}]^2} \right) \mid \mathcal{F}_{j-1} \right] / t_{J-1} \geq \delta_{\beta_t} \text{ a.s.} \\
& \lim_{J \rightarrow \infty} \sum_{j=1}^J \mathbb{E} \left\{ \sum_{i \in \mathcal{E}_{j-1}} \tau^2 \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell / \ell!} \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau \ell / \ell!} \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell / \ell!} \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell / \ell!} \right\}^2} \mid \mathcal{F}_{j-1} \right\} / t_{J-1} \\
& \geq \delta_{\beta_m} \text{ a.s.}
\end{aligned}$$

on event  $E_{\mathcal{I}}$ .

*Proof.* For any  $\zeta \in \mathcal{B}$ ,

$$\begin{aligned}
& \lim_{J \rightarrow \infty} \sum_{j=1}^J \mathbb{E} \left[ \sum_{i \in \mathcal{E}_{j-1}} M_i \left( \frac{1}{\zeta^2} + \frac{ae^{-\zeta a} - b^2 e^{-\zeta b}}{[e^{-\zeta a} - e^{-\zeta b}]^2} \right) \mid \mathcal{F}_{j-1} \right] / t_{J-1} = \\
& \lim_{J \rightarrow \infty} (\bar{\kappa}_J / t_{J-1}) m \left( \frac{1}{\zeta^2} + \frac{ae^{-\zeta a} - b^2 e^{-\zeta b}}{[e^{-\zeta a} - e^{-\zeta b}]^2} \right) = \\
& \mathcal{I} m \left( \frac{1}{\zeta^2} + \frac{ae^{-\zeta a} - b^2 e^{-\zeta b}}{[e^{-\zeta a} - e^{-\zeta b}]^2} \right) \geq \\
& \delta^* m \left( \frac{1}{\zeta^2} + \frac{ae^{-\zeta a} - b^2 e^{-\zeta b}}{[e^{-\zeta a} - e^{-\zeta b}]^2} \right) \text{ a.s.}
\end{aligned}$$

where the last inequality follows from the fact that there exists  $\delta_*$  such that  $\mathcal{I} > \delta_*$  by assumption. Defining

$$\delta_{\beta_t} \triangleq \delta^* m \left( \frac{1}{\zeta^2} + \frac{ae^{-\zeta a} - b^2 e^{-\zeta b}}{[e^{-\zeta a} - e^{-\zeta b}]^2} \right)$$

completes the first part of the lemma.

For any  $\tau \in \mathcal{B}$ , we find that

$$\begin{aligned}
& \lim_{J \rightarrow \infty} \sum_{j=1}^J \mathbb{E} \left\{ \sum_{i \in \mathcal{E}_{j-1}} \tau^2 \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau \ell} / \ell! \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell} / \ell! \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\}^2} \mid \mathcal{F}_{j-1} \right\} / t_{J-1} \geq \\
& \lim_{J \rightarrow \infty} \frac{\bar{\kappa}_J}{\tau_{J-1}} \tau^2 \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau \ell} / \ell! \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell} / \ell! \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\}^2} \geq \\
& \mathcal{I} \tau^2 \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau \ell} / \ell! \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell} / \ell! \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\}^2} \geq \\
& \delta^* \tau^2 \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau \ell} / \ell! \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell} / \ell! \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\}^2} \text{ a.s.}
\end{aligned}$$

where the last inequality follows from the fact that there exists  $\delta_*$  such that  $\mathcal{I} > \delta_*$  by assumption. Defining

$$\delta_{\beta_m} \triangleq \delta^* \tau^2 \frac{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\} \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell^2 e^{\tau \ell} / \ell! \right\} - \left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} \ell e^{\tau \ell} / \ell! \right\}^2}{\left\{ \sum_{\ell=k}^{|\mathbf{A}_i|} e^{\tau \ell} / \ell! \right\}^2}$$

completes the proof of the lemma. ■

### 11.3.5 Assumption 12

**Assumption 12** follows from the fact that  $\Sigma_J = t_{J-1}$ , verification of Assumption 11 and Lemmas 11.8-11.10,

$$\lim_{J \rightarrow \infty} \eta_J / t_{J-1} = \mathcal{C}$$

where  $\mathcal{C} \succ 0$  w.p. 1. Consequently,

$$\frac{\eta_J}{\bar{\kappa}_J} = \frac{\eta_J}{t_{J-1}} \frac{t_{J-1}}{\bar{\kappa}_J} \xrightarrow{p} \mathcal{C} \mathcal{J} \mathcal{J}^{-1} = \mathcal{C} \succ 0$$

by the continuous mapping theorem. We conclude that **Assumption 12** is satisfied.

### 11.3.6 Assumption 13

We prove **Assumption 13**. Set any  $\epsilon_i > 0$ . We conceptualize  $\ddot{l}_i(\beta^*)$  as a function  $\ddot{\lambda} : \mathcal{B} \times \mathcal{D} \rightarrow \mathbb{R}^{k \times k}$ . From Section 11.2, we know that  $\ddot{\lambda}$  is continuous over both  $\mathcal{B}$  and  $\mathcal{D}$ . Because  $\mathcal{X}$  is compact, we know that  $\mathcal{D}$  is compact. Consequently,  $\ddot{\lambda}$  is uniformly continuous in  $\mathcal{B} \times \mathcal{D}$  (since  $\mathcal{B}$  is compact too). Therefore, for any  $0 < \epsilon \leq \epsilon_i$ , we can find  $\delta_\epsilon$  such that for any  $\beta, \beta'$  and  $\mathbf{d}, \mathbf{d}'$  such

$$\sup_{\|\beta - \beta'\|_2 + \|\mathbf{d} - \mathbf{d}'\|_2 \leq \delta_\epsilon} \left\| \ddot{\lambda}(\beta, \mathbf{d}) - \ddot{\lambda}(\beta', \mathbf{d}') \right\|_2 \leq \epsilon.$$

Since  $\{\beta, \mathbf{d}, \mathbf{d}' : \|\beta - \beta^*\|_2 \leq \delta_\epsilon, \|\mathbf{d} - \mathbf{d}'\|_2 = 0\} \subseteq \{\beta, \beta', \mathbf{d}, \mathbf{d}' : \|\beta - \beta'\|_2 + \|\mathbf{d} - \mathbf{d}'\|_2 \leq \delta_\epsilon\}$ , we know that

$$\sup_{\beta : \|\beta - \beta^*\|_2 \leq \delta_\epsilon} \sup_{\mathbf{d} \in \mathcal{D}} \left\| \ddot{\lambda}(\beta, \mathbf{d}) - \ddot{\lambda}(\beta^*, \mathbf{d}) \right\|_2 \leq \epsilon.$$

**Assumption 13** is satisfied.

## 11.4 Consistency Proof

Note that we do not need Assumptions 9 and 10 in the proof of consistency for the branching process described in Equation 6. We begin with a finite Taylor series expansion,

$$0 = \dot{\mathcal{M}}_J(\boldsymbol{\beta}^*) + \ddot{\mathcal{M}}_J(\bar{\boldsymbol{\beta}}^J)(\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^*),$$

where  $\bar{\boldsymbol{\beta}}^J$  is between  $\widehat{\boldsymbol{\beta}}_J$  and  $\boldsymbol{\beta}^*$ . By the concavity of exponential families (Brown, 1986) – this can also be observed in the proofs of Lemmas 11.8-11.10 – we know that for all  $J \in \mathbb{N}$  and  $\boldsymbol{\beta} \in \mathcal{B}$ ,  $\ddot{\mathcal{M}}_J(\boldsymbol{\beta})/t_{J-1}$  is invertible. Consequently,

$$\begin{aligned} -\dot{\mathcal{M}}_J(\boldsymbol{\beta}^*)/t_{J-1} &= \left\{ \ddot{\mathcal{M}}_J(\bar{\boldsymbol{\beta}}^J)/t_{J-1} \right\} (\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^*) \Rightarrow \\ &-\left\{ \ddot{\mathcal{M}}_J(\bar{\boldsymbol{\beta}}^J)/t_{J-1} \right\}^{-1} \dot{\mathcal{M}}_J(\boldsymbol{\beta}^*)/t_{J-1} = \widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^*. \end{aligned}$$

Under Assumptions 1-8 and Assumption 11, we know that

$$\dot{\mathcal{M}}_J(\boldsymbol{\beta}^*)/t_{J-1}^{1/2} = O_p(1)$$

by Section 8.3.1. Consequently,

$$\dot{\mathcal{M}}_J(\boldsymbol{\beta}^*)/t_{J-1} = o_p(1).$$

Therefore, we only need to show

$$\left\{ -\ddot{\mathcal{M}}_J(\bar{\boldsymbol{\beta}}^J)/t_{J-1} \right\}^{-1} = O_p(1)$$

to prove that

$$\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}^* = o_p(1).$$

To do this, we show that for any  $\boldsymbol{\beta} \in \mathcal{B}$ ,

$$\sigma_{\min} \left\{ -\ddot{\mathcal{M}}_J(\boldsymbol{\beta})/t_{J-1} \right\} \geq \epsilon_{\boldsymbol{\beta}}^*/\rho_{\max} + o_p(1). \quad (30)$$

This implies that

$$\left\| \left\{ -\ddot{\mathcal{M}}_J(\boldsymbol{\beta})/t_{J-1} \right\}^{-1} \right\|_2 \leq \rho_{\max}/\epsilon_{\boldsymbol{\beta}}^* + o_p(1).$$

To show Equation 30, we first define

$$\begin{aligned} \nu_J(\boldsymbol{\beta}) &= - \sum_{j=1}^J \mathbb{E}_{\boldsymbol{\beta}^*, \tilde{\boldsymbol{\pi}}} \left\{ \sum_{i \in \mathcal{E}_{j-1}} \ddot{l}_i(\boldsymbol{\beta}) \mid \mathcal{F}_{j-1} \right\}, \\ \alpha_J(\boldsymbol{\beta}) &= - \sum_{j=1}^J \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left\{ \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \mid \mathcal{F}_{j-1} \right\} \end{aligned}$$

By Lemmas 11.8-11.10, we know that (on the event  $E_{\mathcal{I}}$ ) for any  $\boldsymbol{\beta} \in \mathcal{B}$  and

$$\epsilon_{\boldsymbol{\beta}}^* = \min \left\{ \delta_{\boldsymbol{\beta}_m^*}, \delta_{\boldsymbol{\beta}_t^*}, \delta_{\boldsymbol{\beta}_{\mathbf{x}}^*}, \delta_{\boldsymbol{\beta}_y^*} \right\} \text{ (where } \delta_{\boldsymbol{\beta}_m^*}, \delta_{\boldsymbol{\beta}_t^*}, \delta_{\boldsymbol{\beta}_{\mathbf{x}}^*}, \delta_{\boldsymbol{\beta}_y^*} \text{ are defined in Lemmas 11.8-11.10),}$$

$$\lim_{J \rightarrow \infty} \nu_J(\boldsymbol{\beta})/t_{J-1} \geq \epsilon_{\boldsymbol{\beta}}^*. \quad (31)$$

By Lemmas 11.2 and 11.3,

$$\begin{aligned} \lim_{J \rightarrow \infty} \sigma_{\min} \left\{ \ddot{\mathcal{M}}_J(\boldsymbol{\beta})/t_{J-1} \right\} &\geq \frac{1}{\rho_{\max}} \lim_{J \rightarrow \infty} \sigma_{\min} \left\{ \left\{ - \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \right\} \alpha_J(\boldsymbol{\beta})^{-1} \nu_J(\boldsymbol{\beta})/t_{J-1} \right\} \\ &\geq \frac{1}{\rho_{\max}} \lim_{J \rightarrow \infty} \sigma_{\min} \left\{ \left\{ - \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \right\} \alpha_J(\boldsymbol{\beta})^{-1} \right\} \sigma_{\min} \{ \nu_J(\boldsymbol{\beta})/t_{J-1} \} \\ &= \frac{1}{\rho_{\max}} \lim_{J \rightarrow \infty} \sigma_{\min} \left\{ \left\{ - \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \right\} \alpha_J(\boldsymbol{\beta})^{-1} - I + I \right\} \sigma_{\min} \{ \nu_J(\boldsymbol{\beta})/t_{J-1} \} \\ &\geq \frac{1}{\rho_{\max}} \lim_{J \rightarrow \infty} \left( \sigma_{\min} \left\{ \left\{ - \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \right\} \alpha_J(\boldsymbol{\beta})^{-1} - I \right\} + 1 \right) \sigma_{\min} \{ \nu_J(\boldsymbol{\beta})/t_{J-1} \}. \end{aligned}$$

The first inequality follows from the inequality

$$\alpha_J \succ \frac{1}{\rho_{\max}} \nu_J. \quad (32)$$

Equation 32 follows from the fact that  $l(\boldsymbol{\beta})$  is concave (because each component of the branching process is a member of a full exponential family – this can also be observed in

the proofs of Lemmas 11.8-11.10),  $-\ddot{l}(\boldsymbol{\beta}) \succeq 0$ , so for every  $\boldsymbol{\beta} \in \mathcal{B}$ ,

$$\begin{aligned} \alpha_J &\triangleq \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left\{ W_i(-\ddot{l}(\boldsymbol{\beta})) \mid \mathcal{F}_{j-1} \right\} \\ &\succeq \frac{1}{\rho_{\max}} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left\{ W_i^2(-\ddot{l}(\boldsymbol{\beta})) \mid \mathcal{F}_{j-1} \right\} \\ &= \frac{1}{\rho_{\max}} \sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} \mathbb{E}_{\boldsymbol{\beta}^*, \hat{\boldsymbol{\pi}}} \left\{ (-\ddot{l}(\boldsymbol{\beta})) \mid \mathcal{F}_{j-1} \right\}. \end{aligned}$$

Additionally, from Property (3) of Section 8, we know that

$$\lim_{J \rightarrow \infty} \left\| \frac{\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) - \alpha_J(\boldsymbol{\beta})}{\bar{\kappa}_J} \right\|_2 = 0 \quad \text{a.s.} \quad (33)$$

We can extend this to

$$\begin{aligned} &\left\| \left\{ -\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \right\} \alpha_J(\boldsymbol{\beta})^{-1} - I \right\|_2 \\ &= \left\| \left\{ \left\{ -\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \right\} - \alpha_J(\boldsymbol{\beta}) \right\} \alpha_J(\boldsymbol{\beta})^{-1} \right\|_2 \\ &\leq \left\| \left\{ \left\{ -\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \right\} - \alpha_J(\boldsymbol{\beta}) \right\} \right\|_2 \|\alpha_J(\boldsymbol{\beta})^{-1}\|_2 \\ &\leq \rho_{\max} \left\| \frac{\left\{ \left\{ -\sum_{j=1}^J \sum_{i \in \mathcal{E}_{j-1}} W_i \ddot{l}_i(\boldsymbol{\beta}) \right\} - \alpha_J(\boldsymbol{\beta}) \right\}}{\bar{\kappa}_J} \right\|_2 \frac{\bar{\kappa}_J}{t_{J-1}} \|\nu_J(\boldsymbol{\beta})^{-1}\|_2 t_{J-1} \\ &= o_p(1) \{O_p(1) + o_p(1)\} \left\{ \frac{1}{\epsilon_{\boldsymbol{\beta}}^*} + o_p(1) \right\} \\ &= o_p(1). \end{aligned}$$

The third to last equality follows from Equation 32. The second to last equality follows from Equation 31, Equation 33, and the fact that

$$\lim_{J \rightarrow \infty} \frac{\bar{\kappa}_J}{t_{J-1}} = \mathcal{I} \quad \text{a.s.}$$

Consequently, we know that

$$\sigma_{\min} \left\{ -\ddot{\mathcal{M}}_J(\boldsymbol{\beta})/t_{J-1} \right\} \geq \epsilon_{\boldsymbol{\beta}}^*/\rho_{\max} + o_p(1).$$

This implies that

$$t_{J-1} \left\| \ddot{\mathcal{M}}_J(\boldsymbol{\beta})^{-1} \right\|_2 \leq \rho_{\max}/\epsilon_{\boldsymbol{\beta}}^* + o_p(1).$$

Note that  $\left\| \ddot{\mathcal{M}}_J(\boldsymbol{\beta})^{-1} \right\|_2$  is a continuous function of  $\boldsymbol{\beta}$  by Section 11.1. Because  $\mathcal{B}$  is compact, we know that

$$\sup_{\boldsymbol{\beta} \in \mathcal{B}} t_{J-1} \left\| \ddot{\mathcal{M}}_J(\boldsymbol{\beta})^{-1} \right\|_2 \leq \sup_{\boldsymbol{\beta} \in \mathcal{B}} \rho_{\max}/\epsilon_{\boldsymbol{\beta}}^* + o_p(1) < \infty$$

by the extreme value theorem. Consequently, we know that

$$t_{J-1} \left\| \ddot{\mathcal{M}}_J(\bar{\boldsymbol{\beta}}_J)^{-1} \right\|_2 = O_p(1).$$

This demonstrates that  $\widehat{\boldsymbol{\beta}}_J$  is consistent.

## 11.5 Proof of Theorem 2.3

We have proved consistency and verified Assumptions 5-8 and Assumptions 11-14. This proves Theorems 2.1 and 2.2. Theorem 2.3 follows from Theorems 2.1 and 2.2.

# 12 Generalized RDS Inference

## 12.1 Hessian of the Covariate Model

**The Hessian for the log-likelihood of the covariate model.** Define  $\mathbf{X}_*^v \triangleq (1, \mathbf{X}^v)$  and  $G_{\mathbf{a}}^\dagger = (\boldsymbol{\phi}_{\mathbf{a}}^\top, G_{\mathbf{a}}^\top)$ . Define  $\mathbb{A}$  to be the set of possible coupon types. We note that each coupon allocation is a set of identical coupons, implying that the sets  $\mathcal{A}$  and  $\mathbb{A}$  have a

one to one correspondence. Consequently, for  $\mathbf{a} \in \mathcal{A}$ , there exists  $a \in \mathbb{A}$  such that

$$\mathbb{I}(\mathbf{A}_i = \mathbf{a}) = \mathbb{I}(A_{i,l} = a)$$

for every  $i \in \mathbb{N}$  and  $l \in \{1, 2, \dots, M_i\}$ . Consequently, we can represent the complete branching process likelihood as

$$\begin{aligned} \mathcal{L}_\kappa^\theta \left( \{G_{\mathbf{a}}^\dagger, \Sigma_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}} \right) &\triangleq \prod_{v=1}^{\kappa} (2\pi)^{-p/2} \left| \sum_{\mathbf{a} \in \mathcal{A}} \Sigma_{\mathbf{a}} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \right|^{-1/2} \times \\ &\exp \left[ -\frac{1}{2} \sum_{\mathbf{a} \in \mathcal{A}} \{ (\mathbf{X}^v - G_{\mathbf{a}}^\dagger \mathbf{X}_*^{R^v})^\top \Sigma_{\mathbf{a}}^{-1} (\mathbf{X}^v - G_{\mathbf{a}}^\dagger \mathbf{X}_*^{R^v}) \} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \right] \\ &= \prod_{v=1}^{\kappa} (2\pi)^{-p/2} \left| \sum_{\mathbf{a} \in \mathcal{A}} \Sigma_{\mathbf{a}} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \right|^{-1/2} \times \\ &\exp \left[ -\frac{1}{2} \sum_{\mathbf{a} \in \mathcal{A}} \left\{ \mathbf{X}^{v\top} \Sigma_{\mathbf{a}}^{-1} \mathbf{X}^v - 2 \mathbf{X}_*^{R^v\top} G_{\mathbf{a}}^{\dagger\top} \Sigma_{\mathbf{a}}^{-1} \mathbf{X}^v + \right. \right. \\ &\quad \left. \left. \mathbf{X}_*^{R^v\top} G_{\mathbf{a}}^{\dagger\top} \Sigma_{\mathbf{a}}^{-1} G_{\mathbf{a}}^\dagger \mathbf{X}_*^{R^v} \right\} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \right]. \end{aligned}$$

We will use this likelihood for the proofs that follow.

For  $\mathbf{a} \in \mathcal{A}$ , we reparameterize  $\Omega_{\mathbf{a}} = -\frac{1}{2} \Sigma_{\mathbf{a}}^{-1}$  and  $\Gamma_{\mathbf{a}} = \Sigma_{\mathbf{a}}^{-1} G_{\mathbf{a}}^\dagger$ .

$$\begin{aligned} \mathcal{L}_\kappa^\theta \left( \{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}} \right) &= \prod_{v=1}^{\kappa} (2\pi)^{-p/2} \left| \sum_{\mathbf{a} \in \mathcal{A}} -2\Omega_{\mathbf{a}} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \right|^{1/2} \times \\ &\exp \left[ \sum_{\mathbf{a} \in \mathcal{A}} \left\{ \mathbf{X}^{v\top} \Omega_{\mathbf{a}} \mathbf{X}^v + \mathbf{X}_*^{R^v\top} \Gamma_{\mathbf{a}}^\top \mathbf{X}^v + \right. \right. \\ &\quad \left. \left. \frac{1}{4} \mathbf{X}_*^{R^v\top} \Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{X}_*^{R^v} \right\} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \right]. \end{aligned}$$

The reparameterized log-likelihood is

$$\begin{aligned}
\ell_\kappa^\theta(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) &= \sum_{v=1}^{\kappa} \sum_{\mathbf{a} \in \mathcal{A}} \left[ (-p/2) \log(2\pi) + \frac{1}{2} \log | -2\Omega_{\mathbf{a}} | + \right. \\
&\quad \mathbf{X}^{v\top} \Omega_{\mathbf{a}} \mathbf{X}^v + \mathbf{X}_*^{R^v\top} \Gamma_{\mathbf{a}}^\top \mathbf{X}^v + \\
&\quad \left. \frac{1}{4} \mathbf{X}_*^{R^v\top} \Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{X}_*^{R^v} \right] \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \\
&= \sum_{v=1}^{\kappa} \sum_{\mathbf{a} \in \mathcal{A}} \left[ (-p/2) \log(2\pi) + \frac{1}{2} \log | -2\Omega_{\mathbf{a}} | + \right. \\
&\quad \text{tr}(\Omega_{\mathbf{a}} \mathbf{X}^v \mathbf{X}^{v\top}) + \text{tr}(\Gamma_{\mathbf{a}}^\top \mathbf{X}^v \mathbf{X}_*^{R^v\top}) + \\
&\quad \left. \text{tr}\left(\frac{1}{4} \Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{X}_*^{R^v} \mathbf{X}_*^{R^v\top}\right) \right] \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}),
\end{aligned}$$

where the second equality follows from rearranging terms and using the properties of the trace operator. For  $\mathbf{a} \in \mathcal{A}$ , define  $\kappa_{\mathbf{a}} = \sum_{v=1}^{\kappa} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a})$  and  $\mathbf{V}_{\mathbf{a}}^\kappa \in \mathbb{R}^{(p+1) \times (p+1)}$  such that  $\mathbf{V}_{\mathbf{a}}^\kappa = \sum_{v=1}^{\kappa} \mathbf{X}_*^{R^v} \mathbf{X}_*^{R^v\top} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a})$ . We apply the differential operator two times and find

$$\begin{aligned}
\mathbf{d}^2 \ell_\kappa^\theta(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) &= \sum_{v=1}^{\kappa} \sum_{\mathbf{a} \in \mathcal{A}} -\mathbf{d}^2 \left\{ -\frac{1}{2} \log | -2\Omega_{\mathbf{a}} | - \right. \\
&\quad \left. \frac{1}{4} \text{tr}\left(\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{X}_*^{R^v} \mathbf{X}_*^{R^v\top}\right) \right\} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \\
&= -\mathbf{d} \left\{ -\frac{\kappa_{\mathbf{a}}}{2} \text{tr}(\Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}}) - \frac{1}{4} \text{tr}(2\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^\kappa) + \right. \\
&\quad \left. \frac{1}{4} \text{tr}(\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^\kappa) \right\} \\
&= -\left\{ \frac{\kappa_{\mathbf{a}}}{2} \text{tr}(\Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}}) - \right. \\
&\quad \frac{1}{2} \text{tr}(\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^\kappa) - \\
&\quad \frac{1}{2} \text{tr}(\mathbf{d}\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^\kappa) + \\
&\quad \text{tr}(\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^\kappa) + \\
&\quad \left. \frac{1}{4} \text{tr}(\Gamma_{\mathbf{a}}^\top \Omega_{\mathbf{a}}^{-1} \mathbf{d}\Omega_{\mathbf{a}} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}} \mathbf{V}_{\mathbf{a}}^\kappa) \right\}.
\end{aligned}$$

We observe that for  $\mathbf{a}, \mathbf{a}' \in \mathcal{A}$  and  $\mathbf{a} \neq \mathbf{a}'$ ,

$$\frac{\partial \ell_\kappa^\theta(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}})}{\partial (\text{vec}(\Gamma_{\mathbf{a}}), \text{vec}(\Omega_{\mathbf{a}})) \partial (\text{vec}(\Gamma_{\mathbf{a}'}) , \text{vec}(\Omega_{\mathbf{a}'}))} = [0]_{(p+1)^2 \times p^2}.$$

We express

$$\begin{aligned} & \ddot{\ell}_\kappa^\theta(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) \\ &= -\text{diag} \left[ \left\{ \begin{pmatrix} \mathbf{V}_{\mathbf{a}}^\kappa \otimes \Sigma_{\mathbf{a}} & 2(\mathbf{V}_{\mathbf{a}}^\kappa \mathbf{G}_{\mathbf{a}}^{\dagger \top} \otimes \Sigma_{\mathbf{a}}) \\ 2(\mathbf{G}_{\mathbf{a}}^\dagger \mathbf{V}_{\mathbf{a}}^\kappa \otimes \Sigma_{\mathbf{a}}) & 4(\mathbf{G}_{\mathbf{a}}^\dagger \mathbf{V}_{\mathbf{a}}^\kappa \mathbf{G}_{\mathbf{a}}^{\dagger \top} \otimes \Sigma_{\mathbf{a}}) + 2\kappa_{\mathbf{a}}(\Sigma_{\mathbf{a}} \otimes \Sigma_{\mathbf{a}}) \end{pmatrix} \right\}_{\mathbf{a} \in \mathcal{A}} \right] \\ &= -\text{diag} \left[ \left\{ \begin{pmatrix} \mathbf{V}_{\mathbf{a}}^\kappa & 2\mathbf{V}_{\mathbf{a}}^\kappa \mathbf{G}_{\mathbf{a}}^{\dagger \top} \\ 2\mathbf{G}_{\mathbf{a}}^\dagger \mathbf{V}_{\mathbf{a}}^\kappa & 4\mathbf{G}_{\mathbf{a}}^\dagger \mathbf{V}_{\mathbf{a}}^\kappa \mathbf{G}_{\mathbf{a}}^{\dagger \top} + 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}} \end{pmatrix} \otimes \Sigma_{\mathbf{a}} \right\}_{\mathbf{a} \in \mathcal{A}} \right]. \end{aligned}$$

The second equality follows from properties of kronecker products.

## 12.2 Consistency of an M-estimator

We establish the conditions for consistency of an M-estimator under a sequence of concave estimating equations. This will be leveraged in Section 12.3.

**Lemma 12.1** (Theorem 2.7 from Newey and McFadden (1994)). *If there is a function  $Q_0(\theta)$  and sequence of functions  $\{\widehat{Q}_n(\theta)\}_{n \geq 1}$  such that*

1.  $Q_0(\theta)$  is uniquely maximized at  $\theta_0$ .
2.  $\theta_0$  is an element of the interior of a convex set  $\Theta$ .
3. For all  $n \geq 1$ ,  $\widehat{Q}_n(\theta)$  is concave.
4.  $\widehat{Q}_n(\theta) \xrightarrow{p} Q_0(\theta)$  for all  $\theta \in \Theta$ .

Then,  $\widehat{\theta}_n = \arg \max_{\theta} \widehat{Q}_n(\theta)$  exists with probability approaching one, and  $\widehat{\theta}_n \xrightarrow{p} \theta_0$ .

## 12.3 Generalized RDS Inference Proof

We prove consistency of the MLE in this section. First, we establish some helpful lemmas.

**Lemma 12.2.** *Under Assumptions 1-3, the log-likelihood for the branching process specified in Equation 6 is concave.*

*Proof.* Recall that for  $\mathbf{a} \in \mathcal{A}$ ,  $\Omega_{\mathbf{a}} \triangleq -\frac{1}{2}\Sigma_{\mathbf{a}}^{-1}$  and  $\Gamma_{\mathbf{a}} \triangleq \Sigma_{\mathbf{a}}^{-1}G_{\mathbf{a}}^{\dagger}$ . For  $\mathbf{a} \in \mathcal{A}$ , define  $\kappa_{\mathbf{a}} \triangleq \sum_{v=1}^{\kappa} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a})$  and  $\mathbf{V}_{\mathbf{a}}^{\kappa} \in \mathbb{R}^{(p+1) \times (p+1)}$  such that  $\mathbf{V}_{\mathbf{a}}^{\kappa} \triangleq \sum_{v=1}^{\kappa} (1, \mathbf{X}^{r^v}) (1, \mathbf{X}^{r^v})^{\top} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a})$ . From Section 12.1, we know that

$$\ddot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) = -\text{diag} \left[ \left\{ \begin{pmatrix} \mathbf{V}_{\mathbf{a}}^{\kappa} & 2\mathbf{V}_{\mathbf{a}}^{\kappa}G_{\mathbf{a}}^{\dagger \top} \\ 2G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} & 4G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} G_{\mathbf{a}}^{\dagger \top} + 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}} \end{pmatrix} \otimes \Sigma_{\mathbf{a}} \right\}_{\mathbf{a} \in \mathcal{A}} \right].$$

We begin by analyzing the quantity,

$$\mathbb{M}^{\kappa} \triangleq \begin{pmatrix} \mathbf{V}_{\mathbf{a}}^{\kappa} & 2\mathbf{V}_{\mathbf{a}}^{\kappa}G_{\mathbf{a}}^{\dagger \top} \\ 2G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} & 4G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} G_{\mathbf{a}}^{\dagger \top} + 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}} \end{pmatrix}.$$

By Lemma 11.4, we know that  $\mathbb{M}^{\kappa}$  is positive semi-definite since  $\mathbf{V}_{\mathbf{a}}^{\kappa}$  and  $4G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} G_{\mathbf{a}}^{\dagger \top} + 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}} - 4G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} G_{\mathbf{a}}^{\dagger \top} = 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}}$  are positive semi-definite. Consequently, because  $\mathbb{M}^{\kappa}$  and  $\Sigma_{\mathbf{a}}$  are positive semi-definite regardless of  $\mathbf{a} \in \mathcal{A}$ ,

$$0 \succeq \ddot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}).$$

We conclude that  $\ddot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) / \kappa$  is concave for all  $\kappa \in \mathbb{N}$ . ■

We now show that the Hessian of the log-likelihood is negative definite almost surely under Assumption 17.

**Lemma 12.3.** *Under Assumptions 1-3, 6, 17, and Conditions (C2) and (C4),, assume that the true underlying model for RDS is indexed by  $\boldsymbol{\theta}$ , and the working model is the branching process specified in Equation 6. Define the MLE of the working model covariate distribution as  $\widehat{\boldsymbol{\beta}}_{\mathbf{x}}^{\kappa}(\boldsymbol{\theta}) = \arg \max_{\boldsymbol{\beta}_{\mathbf{x}} \in \mathcal{B}} \ell_{\kappa}^{\boldsymbol{\theta}}(\boldsymbol{\beta}_{\mathbf{x}})$ . For any compact set  $\mathcal{B}' \subset \mathcal{B}$ , there exists  $\delta_{\boldsymbol{\theta}, \mathcal{B}'} > 0$  such that*

$$\liminf_{\kappa \rightarrow \infty} \inf_{\boldsymbol{\beta}_{\mathbf{x}} \in \mathcal{B}'} \sigma_{\min} \left( -\ddot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\boldsymbol{\beta}_{\mathbf{x}}) / \kappa \right) \geq \delta_{\boldsymbol{\theta}, \mathcal{B}'} \quad \text{a.s.}$$

*Proof.* Recall that for  $\mathbf{a} \in \mathcal{A}$ ,  $\Omega_{\mathbf{a}} \triangleq -\frac{1}{2}\Sigma_{\mathbf{a}}^{-1}$  and  $\Gamma_{\mathbf{a}} \triangleq \Sigma_{\mathbf{a}}^{-1}G_{\mathbf{a}}^{\dagger}$ . For  $\mathbf{a} \in \mathcal{A}$ , define  $\kappa_{\mathbf{a}} \triangleq \sum_{v=1}^{\kappa} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a})$  and  $\mathbf{V}_{\mathbf{a}}^{\kappa} \in \mathbb{R}^{(p+1) \times (p+1)}$  such that  $\mathbf{V}_{\mathbf{a}}^{\kappa} \triangleq \sum_{v=1}^{\kappa} (1, \mathbf{X}^{R^v})(1, \mathbf{X}^{R^v})^{\top} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a})$ . From Section 12.1, we know that

$$\ddot{\ell}_{\kappa}^{\theta}(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) = -\text{diag} \left[ \left\{ \left( \begin{array}{cc} \mathbf{V}_{\mathbf{a}}^{\kappa} & 2\mathbf{V}_{\mathbf{a}}^{\kappa}G_{\mathbf{a}}^{\dagger \top} \\ 2G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} & 4G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} G_{\mathbf{a}}^{\dagger \top} + 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}} \end{array} \right) \otimes \Sigma_{\mathbf{a}} \right\}_{\mathbf{a} \in \mathcal{A}} \right].$$

We begin by analyzing the quantity,

$$\mathbb{M}^{\kappa} \triangleq \begin{pmatrix} \mathbf{V}_{\mathbf{a}}^{\kappa} & 2\mathbf{V}_{\mathbf{a}}^{\kappa}G_{\mathbf{a}}^{\dagger \top} \\ 2G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} & 4G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} G_{\mathbf{a}}^{\dagger \top} + 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}} \end{pmatrix}.$$

By Lemma 11.4, we know that  $\mathbb{M}^{\kappa}$  is positive semi-definite since  $\mathbf{V}_{\mathbf{a}}^{\kappa}$  and  $4G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} G_{\mathbf{a}}^{\dagger \top} + 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}} - 4G_{\mathbf{a}}^{\dagger} \mathbf{V}_{\mathbf{a}}^{\kappa} \mathbf{V}_{\mathbf{a}}^{\kappa -1} \mathbf{V}_{\mathbf{a}}^{\kappa} G_{\mathbf{a}}^{\dagger \top} = 2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}}$  are positive semi-definite. Consequently, because  $\mathbb{M}^{\kappa}$  and  $\Sigma_{\mathbf{a}}$  are positive semi-definite regardless of  $\mathbf{a} \in \mathcal{A}$ ,

$$0 \succeq \ddot{\ell}_{\kappa}^{\theta}(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}).$$

We now show that  $-\ddot{\ell}_{\kappa}^{\theta}$  is asymptotically **positive definite**. By Lemma 11.4,

$$\sigma_{\min}(\mathbb{M}^{\kappa}/\kappa) \geq \sigma_{\min}(\mathbf{V}_{\mathbf{a}}^{\kappa}/\kappa) \sigma_{\min}(2\kappa_{\mathbf{a}} \Sigma_{\mathbf{a}}/\kappa).$$

To characterize the convexity of  $\ddot{\ell}_{\kappa}^{\theta}(\{\Gamma_{\mathbf{a}}^{\dagger}, \Sigma_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}})/\kappa$  as  $\kappa \rightarrow \infty$ , we first analyze

$$\begin{aligned} \lim_{\kappa \rightarrow \infty} \mathbf{V}_{\mathbf{a}}^{\kappa}/\kappa &= \lim_{\kappa \rightarrow \infty} \frac{1}{\kappa} \sum_{v=1}^{\kappa} (1, \mathbf{X}^{R^v})(1, \mathbf{X}^{R^v})^{\top} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \\ &= \lim_{\kappa \rightarrow \infty} \frac{1}{\kappa} \sum_{v=1}^{\kappa} \begin{pmatrix} 1 & \mathbf{X}^{R^v \top} \\ \mathbf{X}^{R^v} & \mathbf{X}^{R^v} \mathbf{X}^{R^v \top} \end{pmatrix} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}). \end{aligned}$$

Define fields  $\mathcal{F}^{\kappa} = \sigma[\{R^v, T^v, \mathbf{X}^v, Y^v, \mathbf{A}^v, C^v\}_{v=1}^{\kappa}]$  and quantity

$$\Delta^v \triangleq \mathbb{E} \{ (1, \mathbf{X}^{R^v})(1, \mathbf{X}^{R^v})^{\top} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \mid \mathcal{F}^{v-1} \}.$$

By Lemma 11.4,

$$\Delta^v \succ \mathbb{E} \left\{ \mathbf{X}^{R^v} \mathbf{X}^{R^v \top} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \mid \mathcal{F}^{v-1} \right\}.$$

Additionally, by Assumption 17, there exists  $N$  such that for  $\kappa \geq N$ ,  $\Delta^\kappa \succeq \Delta \succ 0$ .

We know that  $\forall v \in \mathbb{N}$ , every entry of  $\mathbf{X}^v \mathbf{X}^{v \top}$  is bounded because  $\mathcal{X}$  is compact. By Theorem 8.1,

$$\lim_{\kappa \rightarrow \infty} \left\{ \sum_{v=1}^{\kappa} (1, \mathbf{X}^{R^v}) (1, \mathbf{X}^{R^v})^\top \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) - \sum_{v=1}^{\kappa} \Delta^v \right\} / \kappa = 0 \quad \text{a.s.}$$

There exists  $N \in \mathbb{N}$  such that

$$\begin{aligned} & \lim_{\kappa \rightarrow \infty} \frac{1}{\kappa} \sum_{v=1}^{\kappa} (1, \mathbf{X}^{R^v}) (1, \mathbf{X}^{R^v})^\top \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \\ &= \lim_{\kappa \rightarrow \infty} \left\{ \sum_{v=1}^{\kappa} (1, \mathbf{X}^{R^v}) (1, \mathbf{X}^{R^v})^\top \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) - \sum_{v=1}^{\kappa} \Delta^v \right\} / \kappa \\ & \quad + \lim_{\kappa \rightarrow \infty} \sum_{v=1}^{\kappa} \Delta^v / \kappa \\ &= \lim_{\kappa \rightarrow \infty} \sum_{v=1}^{\kappa} \Delta^v / \kappa \quad \text{a.s.} \\ &= \lim_{\kappa \rightarrow \infty} \sum_{v=N}^{\kappa} \Delta^v / \kappa + \sum_{v=1}^N \Delta^v / N \quad \text{a.s.} \\ &\succeq \lim_{\kappa \rightarrow \infty} \frac{\kappa - N}{\kappa} \Delta \\ &\succeq \Delta \end{aligned}$$

Consequently, by the continuous mapping theorem

$$\lim_{\kappa \rightarrow \infty} \sigma_{\min} \left\{ \frac{1}{\kappa} \sum_{v=1}^{\kappa} (1, \mathbf{X}^{R^v}) (1, \mathbf{X}^{R^v})^\top \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) \right\} \geq \sigma_{\min}(\Delta).$$

Defining  $\epsilon_1 \triangleq \sigma_{\min}(\Delta) > 0$ , we conclude that  $\lim_{\kappa \rightarrow \infty} \sigma_{\min}(\mathbf{V}_{\mathbf{a}}^\kappa / \kappa) \geq \epsilon_1$  a.s. Next, we

analyze  $2\kappa_{\mathbf{a}}\Sigma_{\mathbf{a}}/\kappa$ . By Assumption 17 and Theorem 8.1,

$$\begin{aligned}
\lim_{\kappa \rightarrow \infty} \kappa_{\mathbf{a}}/\kappa &= \lim_{\kappa \rightarrow \infty} \left\{ \sum_{v=1}^{\kappa} \mathbb{I}(\mathbf{A}^{R^v} = \mathbf{a}) - \mathbb{P}(\mathbf{A}^{R^v} = \mathbf{a} \mid \mathcal{F}^{\kappa-1}) \right\} / \kappa + \sum_{v=1}^{\kappa} \mathbb{P}(\mathbf{A}^{R^v} = \mathbf{a} \mid \mathcal{F}^{\kappa-1}) / \kappa \\
&\geq \lim_{\kappa \rightarrow \infty} \sum_{v=1}^{\kappa} \mathbb{P}(\mathbf{A}^{R^v} = \mathbf{a} \mid \mathcal{F}^{\kappa-1}) / \kappa \\
&\geq \lim_{\kappa \rightarrow \infty} \sum_{v=N}^{\kappa} \mathbb{P}(\mathbf{A}^{R^v} = \mathbf{a} \mid \mathcal{F}^{\kappa-1}) / \kappa \\
&\geq \delta \quad \text{a.s.}
\end{aligned}$$

Consequently,  $\lim_{\kappa \rightarrow \infty} \kappa_{\mathbf{a}}/\kappa \geq \delta$  a.s. as  $\kappa \rightarrow \infty$ .

Therefore,

$$\lim_{\kappa \rightarrow \infty} \sigma_{\min}(2\kappa_{\mathbf{a}}\Sigma_{\mathbf{a}}/\kappa) \geq 2\sigma_{\min}(\Sigma_{\mathbf{a}})\delta \quad \text{a.s.}$$

Because the minimum eigenvalue of a matrix is a continuous function, by the extreme value theorem, we know that for some  $\alpha > 0$ ,

$$\min_{\mathbf{a} \in \mathcal{A}} \min_{\Sigma_{\mathbf{a}} \in \mathcal{B}'} \sigma_{\min}(\Sigma_{\mathbf{a}}) \geq \alpha.$$

Defining  $\epsilon_2 \triangleq 2\alpha\delta\epsilon_1$ , we find that

$$\lim_{\kappa \rightarrow \infty} \sigma_{\min}(\mathbb{M}^{\kappa}/\kappa) \geq \epsilon_2 \quad \text{a.s.}$$

Defining  $\epsilon_3 \triangleq \epsilon_2\alpha$ , for any  $\mathbf{a} \in \mathcal{A}$ , we find that

$$\begin{aligned}
\lim_{\kappa \rightarrow \infty} \sigma_{\min} \left\{ -\ell_{\kappa}^{\theta}(\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}) \right\} / \kappa &= \sigma_{\min}(\mathbb{M}^{\kappa} \otimes \Sigma_{\mathbf{a}}) \\
&= \sigma_{\min}(\mathbb{M}^{\kappa}) \sigma_{\min}(\Sigma_{\mathbf{a}}) \\
&\geq \epsilon_3 \quad \text{a.s.}
\end{aligned}$$

Line 2 follows because for square matrices  $A$  and  $B$ ,  $\sigma_{\min}(A \otimes B) = \sigma_{\min}(A)\sigma_{\min}(B)$ . In

conclusion,

$$\begin{aligned} \lim_{\kappa \rightarrow \infty} \sigma_{\min} \left\{ \dot{\ell}_{\kappa}^{\boldsymbol{\theta}} (\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) / \kappa \right\} &= \lim_{\kappa \rightarrow \infty} \prod_{\mathbf{a} \in \mathcal{A}} \sigma_{\min} \left\{ \dot{\ell}_{\kappa}^{\boldsymbol{\theta}} (\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}) \right\} / \kappa \\ &\leq -\epsilon_3^{|\mathcal{A}|} \quad \text{a.s.} \\ &< 0 \quad \text{a.s.,} \end{aligned}$$

where the first line follows from Lemma 11.4 (where the off-diagonal blocks are zero). We complete the proof by defining  $\delta_{\boldsymbol{\theta}, \mathcal{B}'} \triangleq \epsilon_3$ . ■

We verify the last part of Assumption 16.

**Lemma 12.4.** *Under Assumptions 1-3, 6, 17, and Conditions (C2) and (C4), assume that the true underlying model for RDS is indexed by  $\boldsymbol{\theta}$ , and the working model is the branching process specified in Equation 6. For any  $\boldsymbol{\beta}_{\mathbf{x}} \in \mathcal{B}$ ,*

$$\dot{\ell}_{\kappa}^{\boldsymbol{\theta}} (\boldsymbol{\beta}_{\mathbf{x}}) / \kappa = O_p(1)$$

as  $\kappa \rightarrow \infty$ .

*Proof.* Write

$$\dot{\ell}_{\kappa}^{\boldsymbol{\theta}} (\boldsymbol{\beta}_{\mathbf{x}}) = \sum_{v=1}^{\kappa} q^v (\boldsymbol{\beta}_{\mathbf{x}}) = \sum_{v=1}^{\kappa} q (\boldsymbol{\beta}_{\mathbf{x}}, \{Y^v, \mathbf{X}^v, T^v, R^v, \mathbf{A}^v\}).$$

We see from Section 12.1 that the function  $q$  is continuous over the compact space  $\mathcal{Y} \times \mathcal{X} \times [t_{\min}, t_{\max}] \times \{1, 2, \dots, v\} \times \mathcal{A}$  (which defines the data space). Consequently, for any given value of  $\boldsymbol{\beta}_{\mathbf{x}}$ ,  $q$  has a maximum over possible data values. Call this maximum  $q^*(\boldsymbol{\beta}_{\mathbf{x}})$ . Consequently,

$$\dot{\ell}_{\kappa}^{\boldsymbol{\theta}} (\boldsymbol{\beta}_{\mathbf{x}}) / \kappa = \sum_{v=1}^{\kappa} q (\boldsymbol{\beta}_{\mathbf{x}}, \{Y^v, \mathbf{X}^v, T^v, R^v, \mathbf{A}^v\}) \leq \sum_{v=1}^{\kappa} q^* / \kappa = q^*(\boldsymbol{\beta}_{\mathbf{x}}).$$

The lemma follows. ■

Lemmas 12.2-12.4 verify Assumption 16 under the working model specified by Equation 6, Assumptions 1-3, 6, 17, and Conditions (C2) and (C4). We now prove Theorem 3.1 under Assumption 16 (which suffices to prove Theorem 3.1 under both assumption sets).

*Proof of Theorem 3.1. Proof of consistency.* Because  $\mathcal{B}$  is open, we know that  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$  is an element of the interior of  $\mathcal{B}$ . Consequently, there exists  $\gamma^* > 0$  such that

$$N_{\gamma^*} \triangleq \{\beta_{\mathbf{x}} : \|\beta_{\mathbf{x}} - \bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})\|_2 \leq \gamma^*\}$$

is a compact subset of  $\mathcal{B}$ .

Under Assumption 16, we know that  $\ell_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}})/\kappa$  is concave for all  $\kappa \in \mathbb{N}$ , and that for  $\gamma^* > 0$  there exists a  $\delta > 0$  such that

$$\lim_{\kappa \rightarrow \infty} \inf_{\beta_{\mathbf{x}} \in N_{\gamma^*}} \sigma_{\min} \left( -\dot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}})/\kappa \right) \geq \delta \text{ a.s.} \quad (34)$$

We now show that for  $\beta_{\mathbf{x}} \in N_{\gamma^*}$ ,  $\ell_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}})/\kappa$  is strictly concave asymptotically. For any  $\beta_{\mathbf{x}}, \beta'_{\mathbf{x}} \in N_{\gamma^*}$  and  $t \in [0, 1]$ , we set  $\beta_{\mathbf{x}}^{\dagger} = t\beta_{\mathbf{x}} + (1-t)\beta'_{\mathbf{x}}$ . For  $\beta''_{\mathbf{x}} \in \{\alpha\beta_{\mathbf{x}} + (1-\alpha)\beta_{\mathbf{x}}^{\dagger} : \alpha \in [0, 1]\}$ ,  $\beta'''_{\mathbf{x}} \in \{\alpha\beta_{\mathbf{x}}^{\dagger} + (1-\alpha)\beta'_{\mathbf{x}} : \alpha \in [0, 1]\}$ , and an  $\epsilon \in (0, \delta)$ , there exists  $N \in \mathbb{N}$  such that  $\forall \kappa \geq N$  (by Taylor expansion),

$$\begin{aligned} \ell_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}})/\kappa &= \left\{ \ell_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}}^{\dagger}) + \dot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}}^{\dagger})^{\top} (\beta_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger}) + \frac{1}{2} (\beta_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger})^{\top} \ddot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\beta''_{\mathbf{x}}) (\beta_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger}) \right\} / \kappa \\ &\leq \ell_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}}^{\dagger}) / \kappa + \dot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}}^{\dagger})^{\top} (\beta_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger}) / \kappa - \frac{1}{2} \|\beta_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger}\|_2^2 \epsilon \text{ a.s.,} \end{aligned}$$

and

$$\begin{aligned} \ell_{\kappa}^{\boldsymbol{\theta}}(\beta'_{\mathbf{x}})/\kappa &= \left\{ \ell_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}}^{\dagger}) + \dot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}}^{\dagger})^{\top} (\beta'_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger}) + \frac{1}{2} (\beta'_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger})^{\top} \ddot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\beta'''_{\mathbf{x}}) (\beta'_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger}) \right\} / \kappa \\ &\leq \ell_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}}^{\dagger}) / \kappa + \dot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}}^{\dagger})^{\top} (\beta'_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger}) / \kappa - \frac{1}{2} \|\beta'_{\mathbf{x}} - \beta_{\mathbf{x}}^{\dagger}\|_2^2 \epsilon \text{ a.s.} \end{aligned}$$

Both of these inequalities follow because (1) for any  $x \in \mathbb{R}^d$  and semi-positive definite matrix  $A \in \mathbb{R}^{d \times d}$ , we know that  $x^{\top} A x \geq \sigma_{\min}(A) x^{\top} x$  and (2) we have a lower bound on  $-\ddot{\ell}_{\kappa}^{\boldsymbol{\theta}}(\beta_{\mathbf{x}})/\kappa$  for any  $\beta_{\mathbf{x}} \in N_{\gamma^*}$  by Equation 34.

From these two expressions, we find that  $\forall \kappa \geq N$ ,

$$\begin{aligned}
t\ell_\kappa^\theta(\beta_{\mathbf{x}})/\kappa + (1-t)\ell_\kappa^\theta(\beta'_{\mathbf{x}})/\kappa &\leq \ell_\kappa^\theta(\beta_{\mathbf{x}}^\dagger)/\kappa - \\
&\quad \frac{1}{2} \left\{ t\|\beta_{\mathbf{x}} - \beta_{\mathbf{x}}^\dagger\|_2^2 + (1-t)\|\beta'_{\mathbf{x}} - \beta_{\mathbf{x}}^\dagger\|_2^2 \right\} \in \text{a.s.} \\
&= \ell_\kappa^\theta(\beta_{\mathbf{x}}^\dagger)/\kappa - \frac{1}{2} \left\{ t(1-t)^2\|\beta_{\mathbf{x}} - \beta'_{\mathbf{x}}\|_2^2 + \right. \\
&\quad \left. (1-t)t^2\|\beta_{\mathbf{x}} - \beta'_{\mathbf{x}}\|_2^2 \right\} \in \text{a.s.} \\
&= \ell_\kappa^\theta(\beta_{\mathbf{x}}^\dagger)/\kappa - \frac{1}{2}t(1-t)\|\beta_{\mathbf{x}} - \beta'_{\mathbf{x}}\|_2^2 \in \text{a.s.}
\end{aligned}$$

Lines 1, 2, and 3 follow from distributing, combining like terms, and refactoring (as well as the inequalities stated above). By point-wise convergence, this implies that

$$t\bar{\ell}^\theta(\beta_{\mathbf{x}}) + (1-t)\bar{\ell}^\theta(\beta'_{\mathbf{x}}) \leq \bar{\ell}^\theta(\beta_{\mathbf{x}}^\dagger) - \frac{1}{2} \{t(1-t)\|\beta'_{\mathbf{x}} - \beta_{\mathbf{x}}\|_2^2\} \in \text{a.s.}$$

Consequently, we know that  $\bar{\ell}^\theta$  is strictly concave a.s. over the compact set  $N_{\gamma^*}$ ; i.e., for any  $\beta_{\mathbf{x}}, \beta'_{\mathbf{x}} \in N_{\gamma^*}$  and  $t \in (0, 1)$ ,

$$t\bar{\ell}^\theta(\beta_{\mathbf{x}}) + (1-t)\bar{\ell}^\theta(\beta'_{\mathbf{x}}) < \bar{\ell}^\theta(t\beta_{\mathbf{x}} + (1-t)\beta'_{\mathbf{x}}).$$

Lastly, we show that strict concavity over  $N_{\gamma^*}$  implies that  $\bar{\beta}_{\mathbf{x}}(\theta) \in \arg \max_{\beta_{\mathbf{x}} \in \mathcal{B}} \bar{\ell}^\theta(\beta_{\mathbf{x}})$  is unique. For a proof by contradiction, assume that  $\beta_{\mathbf{x}} \in \mathcal{B}$  such that  $\beta_{\mathbf{x}} \neq \bar{\beta}_{\mathbf{x}}(\theta)$  and  $\beta_{\mathbf{x}} \in \arg \max_{\beta_{\mathbf{x}} \in \mathcal{B}} \bar{\ell}^\theta(\beta_{\mathbf{x}})$  as well. By concavity, for any  $t \in (0, 1)$ ,

$$\begin{aligned}
\bar{\ell}^\theta\{\bar{\beta}_{\mathbf{x}}(\theta)\} &= t\bar{\ell}^\theta\{\bar{\beta}_{\mathbf{x}}(\theta)\} + (1-t)\bar{\ell}^\theta(\beta_{\mathbf{x}}) \\
&\leq \bar{\ell}^\theta\{t\bar{\beta}_{\mathbf{x}}(\theta) + (1-t)\beta_{\mathbf{x}}\}.
\end{aligned}$$

Because  $\bar{\beta}_{\mathbf{x}}(\theta)$  and  $\beta_{\mathbf{x}}$  are maxima, this implies that for any  $t \in (0, 1)$ ,

$$\bar{\ell}^\theta\{\bar{\beta}_{\mathbf{x}}(\theta)\} = \bar{\ell}^\theta(\beta_{\mathbf{x}}) = \bar{\ell}^\theta\{t\bar{\beta}_{\mathbf{x}}(\theta) + (1-t)\beta_{\mathbf{x}}\}.$$

For any  $\beta_{\mathbf{x}} \notin N_{\gamma^*}$ , we find that for  $t \geq 1 - \gamma^*/\|\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) - \beta_{\mathbf{x}}\|_2$ ,

$$\begin{aligned} \|\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) - (t\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) + (1-t)\beta_{\mathbf{x}})\|_2 &= \|(1-t)\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) - (1-t)\beta_{\mathbf{x}}\|_2 \\ &= (1-t)\|\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) - \beta_{\mathbf{x}}\|_2 \\ &\leq \gamma^*. \end{aligned}$$

We now pick any  $t^* = \max[1 - \gamma^*/\{2\|\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) - \beta_{\mathbf{x}}\|_2\}, 0]$  and label  $\beta'_{\mathbf{x}} = t^*\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) + (1-t^*)\beta_{\mathbf{x}}$ . We know that  $\beta'_{\mathbf{x}} \in N_{\gamma^*}$ , and  $\beta'_{\mathbf{x}} \in \arg \max_{\beta_{\mathbf{x}} \in \mathcal{B}} \bar{\ell}^{\theta}(\beta_{\mathbf{x}})$ . Additionally,

$$\begin{aligned} \bar{\ell}^{\theta} \{\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})\} &= \frac{1}{2} \bar{\ell}^{\theta} \{\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})\} + \frac{1}{2} \bar{\ell}^{\theta}(\beta'_{\mathbf{x}}) \\ &< \bar{\ell}^{\theta} \left\{ \frac{1}{2} \bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) + \frac{1}{2} \beta'_{\mathbf{x}} \right\}, \end{aligned} \tag{35}$$

where the last inequality is strict because  $\bar{\ell}^{\theta}$  is strictly concave over  $N_{\gamma^*}$ . Equation 35 is a **contradiction** because  $(\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta}) + \beta'_{\mathbf{x}})/2$  has a higher log-likelihood than  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$ . Consequently,  $\bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$  must be unique.

We now know that  $\bar{\ell}^{\theta}$  has a unique maximizer and  $\ell_{\kappa}^{\theta}(\beta_{\mathbf{x}})/\kappa$  is concave for all  $\kappa \in \mathbb{N}$ . By Lemma 12.1,  $\widehat{\beta}_{\mathbf{x}}^{\kappa}(\boldsymbol{\theta}) \xrightarrow{P} \bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$ .

**Ridge regression penalty.** We also prove consistency under a ridge regression penalty.

The log-likelihood with a ridge regression component parameterized by  $\alpha \in \mathbb{R}^+$  is

$$\ell_{\kappa}^{\theta, \alpha}(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) = \ell_{\kappa}^{\theta}(\{\Gamma_{\mathbf{a}}, \Omega_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{A}}) - \sum_{\mathbf{a} \in \mathcal{A}} \frac{\alpha}{2} \text{tr}(\Gamma_{\mathbf{a}}^{\top} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}}).$$

Define  $\widehat{\beta}_{\mathbf{x}}^{\kappa}(\boldsymbol{\theta}, \alpha) = \arg \max_{\beta_{\mathbf{x}} \in \mathcal{B}} \ell_{\kappa}^{\theta, \alpha}(\beta_{\mathbf{x}})$ . We want to prove that  $\widehat{\beta}_{\mathbf{x}}^{\kappa}(\boldsymbol{\theta}, \alpha) \xrightarrow{P} \bar{\beta}_{\mathbf{x}}(\boldsymbol{\theta})$  as  $\kappa \rightarrow \infty$ . Since  $-\Gamma_{\mathbf{a}}^{\top} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}}$  is negative semi-definite, we know that  $-\sum_{\mathbf{a} \in \mathcal{A}} \frac{\alpha}{2} \text{tr}(\Gamma_{\mathbf{a}}^{\top} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}})$  is concave. We conclude that  $\ell_{\kappa}^{\alpha}$  is concave for all  $\kappa \in \mathbb{N}$  since it is the sum of two concave functions. Additionally, since

$$\lim_{\kappa \rightarrow \infty} \sum_{\mathbf{a} \in \mathcal{A}} \frac{\alpha}{2} \text{tr}(\Gamma_{\mathbf{a}}^{\top} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}}) / \kappa = 0,$$

we know that for any  $\beta_{\mathbf{x}} \in \mathcal{B}$ ,

$$\lim_{\kappa \rightarrow \infty} \ell_{\kappa}^{\alpha}(\beta_{\mathbf{x}}) / \kappa = \lim_{\kappa \rightarrow \infty} \left[ \ell_{\kappa}(\beta_{\mathbf{x}}) / \kappa - \sum_{\mathbf{a} \in \mathcal{A}} \frac{\alpha}{2} \text{tr}(\Gamma_{\mathbf{a}}^{\top} \Omega_{\mathbf{a}}^{-1} \Gamma_{\mathbf{a}}) / \kappa \right] = \bar{\ell}^{\theta}(\beta_{\mathbf{x}}).$$

Using Lemma 12.1, we conclude that  $\widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta, \alpha) \xrightarrow{p} \bar{\beta}_{\mathbf{x}}(\theta)$ .

**Concentration.** We now prove the concentration statement of Theorem 3.1: for any  $\theta \notin \Theta^*$ ,

$$\lim_{\kappa \rightarrow \infty} \mathbb{P} \{ \theta \in \Gamma_{1-\alpha, \kappa} \} \rightarrow 0.$$

Recall that  $\Theta^* = \{ \theta \in \Theta : \bar{\beta}(\theta) = \bar{\beta}(\theta^*) \}$ . For a given  $\theta \in \Theta$ , we define the sampling distribution of the log-likelihood ratio statistic as  $P_{\kappa}^{\theta}(\bar{\beta}_{\mathbf{x}})$  such that

$$-2 \left[ \ell_{\kappa}^{\theta} \{ \bar{\beta}_{\mathbf{x}}(\theta) \} - \ell_{\kappa}^{\theta} \{ \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) \} \right] \sim P_{\kappa}^{\theta}(\bar{\beta}_{\mathbf{x}}).$$

Furthermore, define the  $1 - \alpha$  quantile of  $P_{\kappa}^{\theta}(\bar{\beta}_{\mathbf{x}})$  as  $\gamma_{1-\alpha, \kappa}^{\theta}(\bar{\beta}_{\mathbf{x}})$  and the confidence set in question as

$$\Gamma_{1-\alpha, \kappa} = \left\{ \theta : -2 \left[ \ell_{\kappa} \{ \bar{\beta}_{\mathbf{x}}(\theta) \} - \ell_{\kappa} \{ \widehat{\beta}_{\mathbf{x}}^{\kappa} \} \right] \leq \gamma_{1-\alpha, \kappa}^{\theta}(\bar{\beta}_{\mathbf{x}}) \right\}.$$

We know that  $\widehat{\beta}_{\mathbf{x}}^{\kappa}$  is consistent for  $\bar{\beta}_{\mathbf{x}}(\theta^*)$ . Therefore, we know that for any  $\epsilon_0 > 0$  such that  $\{ \beta_{\mathbf{x}} : \|\beta_{\mathbf{x}} - \bar{\beta}_{\mathbf{x}}(\theta^*)\|_2 \leq \epsilon_0 \} \subset \mathcal{B}$ , there exists  $N_1 \in \mathbb{N}$  such that for all  $\kappa \geq N_1$ ,  $\|\widehat{\beta}_{\mathbf{x}}^{\kappa} - \bar{\beta}_{\mathbf{x}}(\theta^*)\| \leq \epsilon_0$  almost surely.

Define the compact set  $\mathcal{B}_c^{\theta} = \bar{\beta}_{\mathbf{x}}(\theta) \cup \{ \beta_{\mathbf{x}} : \|\beta_{\mathbf{x}} - \bar{\beta}_{\mathbf{x}}(\theta^*)\|_2 \leq \epsilon_0 \}$ . Additionally, for any  $\beta_{\mathbf{x}}^p, \beta_{\mathbf{x}}^g \in \mathcal{B}_c^{\theta}$ , define  $\mathcal{B}_*^{\theta} = \{ \beta_{\mathbf{x}} \in \{ \alpha \beta_{\mathbf{x}}^p + (1 - \alpha) \beta_{\mathbf{x}}^g : \alpha \in [0, 1] \} \}$ . Note that  $\mathcal{B}_*^{\theta}$  is compact because it is the image of a continuous function on a compact set in Euclidean space.

Under Assumption 16, we know that there exists an  $N_2 \in \mathbb{N}$  such that for an  $\epsilon_{\theta} > 0$

and any  $\kappa \geq N_2$

$$\inf_{\beta_{\mathbf{x}} \in \mathcal{B}_{*}^{\theta}} \sigma_{\min} \left\{ -\ddot{\ell}_{\kappa}(\beta_{\mathbf{x}}) / \kappa \right\} \geq \epsilon_{\theta} \quad \text{a.s.}$$

Consequently, for all  $\kappa \geq \max\{N_1, N_2\}$ , any  $\theta \in \Theta$ , and a  $\beta'_{\mathbf{x}} \in \left\{ \alpha \widehat{\beta}_{\mathbf{x}}^{\kappa} + (1 - \alpha) \bar{\beta}_{\mathbf{x}}(\theta) : \alpha \in [0, 1] \right\}$ ,

$$\begin{aligned} & -2 \left[ \ell_{\kappa} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\} - \ell_{\kappa} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\} \right] / \kappa \\ &= 2 \left[ \ell_{\kappa} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\} - \ell_{\kappa} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\} \right] / \kappa \\ &= 2 \left[ \ell_{\kappa} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\} - \ell_{\kappa} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\} - \dot{\ell}_{\kappa} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\}^{\top} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\} - \right. \\ & \quad \left. \frac{1}{2} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\}^{\top} \ddot{\ell}_{\kappa} \left\{ \beta'_{\mathbf{x}} \right\} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\} \right] / \kappa \\ &= \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\}^{\top} \left\{ -\ddot{\ell}_{\kappa}(\beta'_{\mathbf{x}}) / \kappa \right\} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\} \\ &\geq \epsilon_{\theta} \left\| \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\|_2^2 \quad \text{a.s.} \end{aligned}$$

Line 3 follows from an exact Taylor expansion. Line 4 follows from the fact that  $\dot{\ell}_{\kappa} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa} \right\} = 0$ . Line 5 follows from the fact that  $\beta'_{\mathbf{x}} \in \mathcal{B}_{*}^{\theta}$ .

For  $\beta''_{\mathbf{x}} \in \left\{ \alpha \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) + (1 - \alpha) \bar{\beta}_{\mathbf{x}}(\theta) : \alpha \in [0, 1] \right\}$ , we now upper bound

$$\begin{aligned} 0 &\leq -2 \left[ \ell_{\kappa}^{\theta} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\} - \ell_{\kappa}^{\theta} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) \right\} \right] / \kappa \\ &= 2 \left[ \ell_{\kappa}^{\theta} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) \right\} - \ell_{\kappa}^{\theta} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\} \right] / \kappa \\ &= 2 \left[ \ell_{\kappa}^{\theta} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\} - \ell_{\kappa}^{\theta} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\} + \dot{\ell}_{\kappa}^{\theta} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\}^{\top} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) - \bar{\beta}_{\mathbf{x}}(\theta) \right\} + \right. \\ & \quad \left. \frac{1}{2} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) \right\}^{\top} \ddot{\ell}_{\kappa}^{\theta} \left\{ \beta''_{\mathbf{x}} \right\} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) \right\} \right] / \kappa \\ &= 2 \left[ \dot{\ell}_{\kappa}^{\theta} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\} / \kappa \right]^{\top} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) - \bar{\beta}_{\mathbf{x}}(\theta) \right\} - \\ & \quad \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) \right\}^{\top} \left[ -\ddot{\ell}_{\kappa}^{\theta} \left\{ \beta''_{\mathbf{x}} \right\} / \kappa \right] \left\{ \bar{\beta}_{\mathbf{x}}(\theta) - \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) \right\} \\ &\leq 2 \left[ \dot{\ell}_{\kappa}^{\theta} \left\{ \bar{\beta}_{\mathbf{x}}(\theta) \right\} / \kappa \right]^{\top} \left\{ \widehat{\beta}_{\mathbf{x}}^{\kappa}(\theta) - \bar{\beta}_{\mathbf{x}}(\theta) \right\} \quad \text{a.s.} \end{aligned}$$

Lines 1, 2, and 3 follow from simple algebra. Line 4 follows from the fact that we know from the same logic as Lemma 12.3 that  $\ell_{\kappa}^{\theta}$  is concave for any  $\theta \in \Theta$ .

By Assumption 16, we know that  $\ell_\kappa^\theta \{\bar{\beta}_x(\theta)\} / \kappa = O_p(1)$  as  $\kappa \rightarrow \infty$ . Because  $\lim_{\kappa \rightarrow \infty} \left\| \bar{\beta}_x(\theta) - \widehat{\beta}_x^\kappa(\theta) \right\|_2^2 = 0$  a.s., we know that

$$\lim_{\kappa \rightarrow \infty} \left( 2 \left[ \ell_\kappa^\theta \{\bar{\beta}_x(\theta)\} / \kappa \right]^\top \left\{ \bar{\beta}_x(\theta) - \widehat{\beta}_x^\kappa(\theta) \right\} \right) = 0. \quad \text{a.s.}$$

Therefore, we know that  $\lim_{\kappa \rightarrow \infty} \gamma_{1-\alpha, \kappa}^\theta(\bar{\beta}_x) / \kappa = 0$  a.s. We find that

$$\begin{aligned} \Gamma_{1-\alpha, \kappa} &\triangleq \left\{ \theta : -2 \left[ \ell_\kappa \{\bar{\beta}_x(\theta)\} - \ell_\kappa \{\widehat{\beta}_x^\kappa\} \right] \leq \gamma_{1-\alpha, \kappa}^\theta(\bar{\beta}_x) \right\} \\ &= \left\{ \theta : -2 \left[ \ell_\kappa \{\bar{\beta}_x(\theta)\} - \ell_\kappa \{\widehat{\beta}_x^\kappa\} \right] / \kappa \leq \gamma_{1-\alpha, \kappa}^\theta(\bar{\beta}_x) / \kappa \right\} \\ &\subseteq \left\{ \theta : \epsilon_\theta \left\| \bar{\beta}_x(\theta) - \widehat{\beta}_x^\kappa \right\|_2^2 \leq \gamma_{1-\alpha, \kappa}^\theta(\bar{\beta}_x) / \kappa \right\} \quad \text{a.s.} \end{aligned}$$

Line 2 follows by dividing both sides of the inequality by  $\kappa$ . Line 3 follows because for any  $\theta \in \Theta$ ,

$$-2 \left[ \ell_\kappa \{\bar{\beta}_x(\theta)\} - \ell_\kappa \{\widehat{\beta}_x^\kappa\} \right] / \kappa \geq \epsilon_\theta \left\| \bar{\beta}_x(\theta) - \widehat{\beta}_x^\kappa \right\|_2^2 \quad \text{a.s.}$$

as  $\kappa \rightarrow \infty$ . Because  $\lim_{\kappa \rightarrow \infty} \widehat{\beta}_x^\kappa = \bar{\beta}_x(\theta^*)$ , we know that if  $\bar{\beta}_x(\theta^*) \neq \bar{\beta}_x(\theta)$ ,

$$\epsilon_\theta \left\| \bar{\beta}_x(\theta) - \widehat{\beta}_x^\kappa \right\|_2^2 \rightarrow \epsilon_\theta \left\| \bar{\beta}_x(\theta) - \bar{\beta}_x(\theta^*) \right\|_2^2 > 0.$$

Additionally,  $\gamma_{1-\alpha, \kappa}^\theta(\bar{\beta}_x) / \kappa \rightarrow 0$ . Consequently, for any  $\theta \notin \Theta^*$ ,

$$\lim_{\kappa \rightarrow \infty} \mathbb{P}(\theta \in \Gamma_{1-\alpha, \kappa}) \leq \lim_{\kappa \rightarrow \infty} \mathbb{P} \left( \epsilon_\theta \left\| \bar{\beta}_x(\theta) - \widehat{\beta}_x^\kappa \right\|_2^2 \leq \gamma_{1-\alpha, \kappa}^\theta(\bar{\beta}_x) / \kappa \right) \rightarrow 0.$$

■

## 13 Branching Model Paradigm for RL-RDS

In this section, we conduct a series of simulation experiments to evaluate the operating characteristics of RL-RDS when the branching process is the true generative model. To allow comparisons with the two-stage procedure proposed by McFall et al. (2021) (see

also VanOrsdale (2023)), we consider the setting in which the goal is to recruit the largest subset of people in a hidden population with a given binary trait, e.g., undiagnosed HIV. The outcome is thus an indicator of this trait. We estimate the optimal policy,  $\boldsymbol{\pi}^{\text{opt}}$ , using RL-RDS. In this paradigm, we assume that the branching process described in Section 2.2 is the true generative model. We restate this model here for convenience.

Recall that  $T_{i,l}$ ,  $\mathbf{X}_{i,l}$ ,  $Y_{i,l}$ , and  $A_{i,l}$  for  $l = 1, \dots, M_i$  are the arrival times, covariates, rewards, and coupon types associated with the potential recruits of recruiter  $i$  respectively. Treating  $A_{i,l}$  as a factor, let  $\mathbf{Z}_{i,l} \in \mathcal{Z} \subseteq \mathbb{R}^\ell$  be a row in the model matrix of a model that includes a main effect,  $\mathbf{X}_{i,l}$  and its interaction with  $A_{i,l}$ . For example, if  $\mathcal{A} = \{-1, 1\}$ , then  $\mathbf{Z}_{i,l} = (1, \mathbf{X}_{i,l}, \mathbf{X}_{i,l} \mathbb{I}(A_{i,l} = -1))$ . Define  $\mathbb{A}$  as the set of possible coupon types.

We consider a working model of the form (same as Equation 6):

$$\begin{aligned} \mathbb{P}(M_i = m_i | \mathbf{H}_i, \mathbf{A}_i) &= \frac{\lambda^{m_i} / m_i!}{\sum_{\ell=0}^{|\mathbf{A}_i|} (\lambda^\ell / \ell!)}, m_i = 0, \dots, |\mathbf{A}_i|, \\ T_{i,l} - T_i | \mathbf{H}_i, \mathbf{A}_i, M_i &\sim \text{Truncated Exponential}(\zeta, t_{\min}, t_{\max}), l = 1, \dots, M_i, \\ \mathbf{X}_{i,l} | \mathbf{H}_i, U_{i,l}, \mathbf{A}_i, M_i &\sim \text{Normal}(\boldsymbol{\phi}_a + G_a \mathbf{X}_i, \Sigma_a), l = 1, \dots, M_i, a = a_{i,l} \\ Y_{i,l} | \mathbf{H}_i, \mathbf{X}_i, U_{i,l}, \mathbf{A}_i, M_i &\sim \text{Bernoulli} \left\{ \frac{1}{1 + \exp(-\mathbf{Z}_{i,l}^\top \boldsymbol{\beta}_y)} \right\}, j = 1, \dots, M_i, \end{aligned}$$

where  $\lambda, \zeta \in \mathbb{R}$ ,  $\{G_a\}_{a \in \mathbb{A}} \in \mathbb{R}^{p \times p}$ ,  $\{\boldsymbol{\phi}_a\}_{a \in \mathbb{A}} \in \mathbb{R}^{p \times 1}$ ,  $\boldsymbol{\beta}_y \in \mathbb{R}^\ell$ , and  $\{\Sigma_a\}_{a \in \mathbb{A}} \in \mathbb{R}^{p \times p}$ . We draw the covariates of the initial sample,  $\mathcal{E}_0$ , from a multivariate normal distribution,  $\text{Normal}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ .

### 13.1 Policies

We evaluate the performance of RL-RDS against a suite of alternative strategies. At each step, the researcher can choose from a finite selection of coupon types. The fixed allocation policies (i.e., those that give the same coupon allocation type to all participants) represent the current standard in RDS. The train and implement policy mimics the procedure used by McFall et al. (2021), which determines an incentive strategy using a pilot study. We describe each policy below.

1. **Fixed** offers a fixed coupon allocation  $\mathbf{a} \in \mathcal{A}$  to every study participant. If  $\mathbf{a} \notin \psi^v(\mathbf{h}^v)$ , then pick a random coupon allocation from  $\psi^v(\mathbf{h}^v)$  to give to the  $v^{\text{th}}$  study participant.
2. **Random** offers a random element of  $\psi^v(\mathbf{h}^v)$  to the  $v^{\text{th}}$  study participant.
3. **Train and Implement** uses half of the budget for a “pilot study,” in which the the Random policy is used to assign coupon allocations. It then conducts policy search using the pilot study data to estimate the branching process working model. This estimated policy (without updating) is used to determine coupon allocations for the remainder of the budget.
4. **RL-RDS** uses the Random policy to assign coupon allocations to participants in a short “warm-up” period (50 participants in the simulations below). Then, it performs policy search with Thompson sampling as outlined in Section 2 for the remainder of the budget.

To conduct RL-RDS, we establish a reasonable space of policies,  $\Pi$ . Define  $\alpha_0 \in \mathbb{R}$ ,  $\alpha_1 \in \mathbb{R}^p$ , and  $\alpha = (\alpha_0, \alpha_1)$ . For  $n \in \mathbb{N}$  and state  $\mathbf{h}^n \in \mathcal{H}^n$ , we consider policies of the form  $\pi(\mathbf{h}^n) = \{\pi^n(\mathbf{h}^n), \pi^{n+1}(\mathbf{h}^{n+1}), \dots\}$  such that for  $v \geq n$ ,

$$\pi^v(\mathbf{h}^v) = \pi^v(\mathbf{h}^v, \alpha) = \pi^v(\mathbf{x}^v, \alpha) = g^v \left[ \frac{1}{1 + \exp\{-(\alpha_0 + \mathbf{x}^{v\top} \alpha_1)\}} \right],$$

where  $g^v : (0, 1) \rightarrow \phi^v(\mathbf{h}^v)$  maps a continuous score (dependent on the participant’s covariates) to a coupon allocation,  $\mathbf{a}^v \in \psi^v(\mathbf{h}^v)$ . We first draw  $\widehat{\beta}^n$  from the MLE sampling distribution using a generalized bootstrap for estimating equations (Chatterjee and Bose, 2005). We then generate synthetic data sets,

$$\mathcal{K}^B(\mathbf{h}^n, \pi; \widehat{\beta}^n) = \left\{ \left( \mathbf{h}_b^n, A_b^n, Y_b^n, \mathbf{h}_b^{n+1}, A_b^{n+1}, Y_b^{n+1}, \dots, \mathbf{h}_b^Q, Y_b^Q \right) \right\}_{b=1}^B,$$

and calculate

$$\widehat{V}_B^n(\mathbf{h}^n, \pi; \widehat{\beta}^n) = \frac{\sum_{b=1}^B \{\sum_{v \geq n} \Delta_b^v Y_b^v\}}{B}$$

for each  $\boldsymbol{\pi} \in \Pi$  (or each  $\boldsymbol{\pi}$  in a grid approximation of  $\Pi$ ). To determine the coupon allocation given to the latest study participant, we set  $\widehat{\boldsymbol{\alpha}}_B^n = \arg \max_{\boldsymbol{\alpha}} \widehat{V}_B^n(\mathbf{h}^n, \boldsymbol{\alpha}; \widehat{\boldsymbol{\beta}}^n)$ , and assign  $\mathbf{a}^n = \pi^n(\mathbf{x}^n, \widehat{\boldsymbol{\alpha}}_B^n)$ .

## 13.2 Results

In the following simulations, we set the hidden population size to  $N = 5,000$ . The recruitment process begins with an initial sample of 25 individuals randomly drawn from the population,  $|\mathcal{E}_0| = 25$  with  $\boldsymbol{\mu}^* = (1, 1, 1)^\top$  and  $\boldsymbol{\Sigma}^* = \text{diag}(1/8)$ . We define

$$G_{a_1} = \begin{pmatrix} \phi_1 & 0 & 0 \\ 0 & \phi_1 & 0 \\ 0 & 0 & \phi_1 \end{pmatrix}, G_{a_2} = \begin{pmatrix} \phi_1 * 0.975 & 0 & 0 \\ 0 & \phi_1 * 0.975 & 0 \\ 0 & 0 & \phi_1 * 0.975 \end{pmatrix}$$

$$G_{a_3} = \begin{pmatrix} \phi_1 * 0.95 & 0 & 0 \\ 0 & \phi_1 * 0.95 & 0 \\ 0 & 0 & \phi_1 * 0.95 \end{pmatrix}, \boldsymbol{\phi}_{a_1} = \boldsymbol{\phi}_{a_2} = \boldsymbol{\phi}_{a_3} = \boldsymbol{\phi}_0,$$

where we set  $\boldsymbol{\phi}_0 = (0.05, 0.05, 0.05)$  and  $\phi_1 = 0.95$  in our weak correlation setting, and we set  $\boldsymbol{\phi}_0 = (0.025, 0.025, 0.025)$  and  $\phi_1 = 0.975$  in our strong correlation setting. The researchers have access to three types of coupon allocations  $\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ , and each allocation has 5 coupons. We found that limiting the number of coupons given to each study participant in the pilot study and the warm-up period of the T&I and the RLRDS policies respectively allows us to observe the effects of the learned policies earlier in the sampling process. Consequently, we give two coupons to individuals in the pilot and warm-up period, and increase the allotment to 5 coupons afterwards. Additionally,  $C^v \equiv 1$  for all  $v \in \mathbb{N}$ . Define the reward model components as  $\mathbf{Z}_j^v \triangleq \{1, \mathbf{X}_j^v, \mathbb{I}(\mathbf{A}_j^v = \mathbf{a}_2)\mathbf{X}_j^v, \mathbb{I}(\mathbf{A}_j^v = \mathbf{a}_3)\mathbf{X}_j^v\}$  and  $\boldsymbol{\beta}_y^* \triangleq (-1, 3\mathbf{k}, -3\mathbf{k}, -6\mathbf{k})$ , where  $\mathbf{k} \triangleq (1, -1, -1)$ . The basic policy objective is clear: we want to ensure that coupon type 1,  $\mathbf{a}_1$ , is used to recruit individuals with covariates that satisfy  $\mathbf{X}^{v\top} \mathbf{k} > 0$ , and coupon type 3,  $\mathbf{a}_3$ , is used to recruit individuals with covariates




Figure 3: This figure compares the estimated cumulative reward of each policy with 90% Monte Carlo confidence intervals over multiple sample sizes and graph densities.

that satisfy  $\mathbf{X}^v \top \mathbf{k} < 0$ . We define the policy space by specifying the function  $g^v$ ,

$$g^v(z) = \begin{cases} \mathbf{a}_1, & \text{if } z > 0.66, \\ \mathbf{a}_2, & \text{if } 0.66 > z \geq 0.33, \\ \mathbf{a}_3, & \text{if } 0.33 > z. \end{cases}$$

This policy space implies that correctly assigning coupons 1 or 3 will depend on the sign of the  $\boldsymbol{\alpha}$  components. The frequency of coupon 2 allocation will be determined by the magnitude of  $\boldsymbol{\alpha}$ . This structure makes finding an optimal policy computationally feasible while maintaining sufficient difficulty to showcase the strength of RL-RDS. Lastly, we set  $\lambda = 3$ ,  $\zeta = 0.5$ ,  $t_{\min} = 0$ , and  $t_{\max} = 3$ .

Figure 3 illustrates the estimated value of policies in both the sparse and dense network settings. It indicates that RL-RDS outperforms all competitor policies by a significant margin in each regime. Note that the train and implement (T&I) method does not adapt its policy after the initial sample, causing the margin between RL-RDS and this strategy to increase for larger budgets.

## 14 Alternative Graph Model Paradigms for RL-RDS

The setup of this section mirrors the setup of Section 4. We conduct a series of simulation experiments to evaluate the operating characteristics of RL-RDS when the graph process of Section 4 is the true generative model. Here, we evaluate the performance of RL-RDS under two additional simulation settings. In both of these paradigms, the coupon “type” is held fixed.

1. Researchers can vary the coupon value while the number of coupons is fixed.
2. Researchers can vary the number of coupons while the coupon value is fixed.

The fixed allocation policies (i.e., those that give the same coupon allocation type to all participants) represent the current standard in RDS. The train and implement policy mimics the procedure used by McFall et al. (2021), which determines an incentive strategy using a pilot study. We describe each policy below.

1. **Fixed (Min)** offers  $\min_{a \in \psi^v(\mathbf{h}^v)}$  to the  $v^{\text{th}}$  study participant.
2. **Fixed (Half)** offers  $\lfloor \max_{a \in \psi^v(\mathbf{h}^v)} / 2 \rfloor$  to the  $v^{\text{th}}$  study participant.
3. **Fixed (Max)** offers  $\max_{a \in \psi^v(\mathbf{h}^v)}$  to the  $v^{\text{th}}$  study participant.
4. **Random** offers a random element of  $\psi^v(\mathbf{h}^v)$  to the  $v^{\text{th}}$  study participant.
5. **Train and Implement** uses half of the budget for a “pilot study,” in which the the Random policy is used to assign coupon allocations. It then conducts policy search using the pilot study data to estimate the branching process working model. This estimated policy (without updating) is used to determine coupon allocations for the remainder of the budget.
6. **RL-RDS** uses the Random policy to assign coupon allocations to participants in a short “warm-up” period (50 participants in the simulations below). Then, it performs policy search with Thompson sampling as outlined in Section 2 for the remainder of the budget.

The working model we use for inference in the experiments in this section involves the value of the coupons explicitly and reduces the dimensionality of the covariate and reward models. The possible coupons will have values between 0 and 1,  $\mathbb{A} \subseteq [0, 1]$ .

$T_{i,l}, \mathbf{X}_{i,l}, Y_{i,l}$  and  $A_{i,l}$  for  $l = 1, \dots, M_i$  are the arrival times, covariates, rewards, and incentive values associated with the potential recruits of study participant  $i$  respectively.

$$\begin{aligned}
P(M_i = m_i) &= \frac{\lambda^{m_i}/m_i!}{\sum_{\ell=0}^{|\mathbf{A}_i|} (\lambda^\ell/\ell!)}, m_i = 0, \dots, |\mathbf{A}_i|, \\
U_{i,l} \triangleq T_{i,l} - T_i &\sim \text{TruncatedExponential}(\zeta_0 + \zeta_1 A_{i,l}, b), l = 1, \dots, M_i, \\
\mathbf{X}_{i,l} &\sim \text{Normal}[\boldsymbol{\phi}_0 + \phi_1 \mathbf{X}_{i,l}, \{\Omega_0 + (1 - A_{i,l}) \Omega_1\}^{-1}], l = 1, \dots, M_i, \\
Y_{i,l} &\sim \text{Bernoulli}\left\{\frac{1}{1 + \exp(-\mathbf{Z}_{i,l}^\top \boldsymbol{\beta}_y)}\right\}, l = 1, \dots, M_i,
\end{aligned} \tag{36}$$

where the incentive,  $A_{i,l} \in [0, 1]$ , is coded so that a value of 0 encodes the minimal incentive and 1 encodes the maximum incentive. For simplicity (and to align with common study constraints), we assume that the incentive allocation strategy is such that  $A_{i,l}$  is constant across  $l$ . Lastly, we assume that there is an upper bound on the number of coupons that can be given to a single participant; i.e., there exists  $L \in \mathbb{N}$  such that  $\forall i \in \mathbb{N}, |\mathbf{A}_i| \leq L$ .

## 14.1 Results

In all following simulations, the hidden population size is  $N = 5,000$ . The recruitment process begins with an initial sample of 25 individuals randomly drawn from the population,  $|\mathcal{E}_0| = 25$ . The graph model is defined by  $\psi_0^* = 0$  and  $\psi_1^* \in \{0.5, 2\}$  to compare between simulation experiments in dense and sparse network settings respectively. We specify the covariate distribution with  $\boldsymbol{\mu}^* = (1, 1, 1)$  and  $\Sigma^* = 10 * I_3$ . We define the arrival time distribution with  $\zeta_0^* = 0.5$  and  $\zeta_1^* = 6$ .  $\zeta_1^*$  controls the relationship between the incentive offered to a potential recruit and their arrival time, which effects whether we observe them before the end of the process. Consequently, for large  $\zeta_1^*$ , there is a high positive association between the incentive assigned to a recruiter and the likelihood of

observing their recruits.

We switch to superscript indexing for study participants when discussing policy estimation to emphasize that the assignment of coupon allocations happens upon the arrival of a study participant. In the first simulation setting, we set the coupon package size to 5 for all recruits and vary the value of the coupons. For the recruits  $j \in \{1, 2, \dots, M^v\}$  of participant  $v$ , we define  $\mathbf{Z}_j^v \triangleq (1, A_j^v, \mathbf{X}_j^v, A_j^v * \mathbf{X}_j^v)$ . We make  $\boldsymbol{\beta}_y^* \in \mathbb{R}^{2p+2}$  sparse, setting it equal to  $\boldsymbol{\beta}_y^* = (-4, 0, 0, 0, 0, 3, 0, 0)$ . This makes the basic policy objective clear: give high incentives to study participants who are likely to recruit individuals with a particular characteristic in order to maximize cumulative utility. We define the possible incentive values as  $\{0.1, 0.2, 0.3, \dots, 1\}$  and the policy space for participant  $v$  as

$$\begin{aligned} \pi^v(\mathbf{H}^v) &\triangleq \mathbf{A}^v, \text{ where for } A_j^v \in \mathbf{A}^v, \\ A_j^v &= \frac{1}{10} \left[ 10 \times \frac{1}{1 + \exp[-(\alpha_0 + \mathbf{x}^{v\top} \boldsymbol{\alpha}_1)]} + \frac{1}{2} \right], \text{ for } j \in 1, \dots, M^v. \end{aligned}$$

Therefore, the policy is determined by  $\boldsymbol{\alpha}_1 \in \mathbb{R}^p$  and  $\alpha_0 \in \mathbb{R}$ .

In the second simulation context, we hold the coupon value constant and allow the researcher to vary the size of  $\mathbf{A}_i$ . We make  $\mathbf{Z}^v = (1, \mathbf{X}^v)$  and set  $\boldsymbol{\beta}_y = (-1, 2, -2, -2)$ . This provides a strong signal to prioritize policies that recruit individuals who exhibit the covariate pattern:  $X_1^v > 0, X_2^v < 0, X_3^v < 0$ . In this context, we make the possible coupon package sizes  $\{1, 2, 3, \dots, 7\}$ , and specify the policy space as

$$\pi^v(\mathbf{h}^v) \triangleq |\mathbf{A}^v| = \left[ 7 \times \frac{1}{1 + \exp\{-(\alpha_0 + \mathbf{x}^{v\top} \boldsymbol{\alpha}_1)\}} + \frac{1}{2} \right].$$

Again, the policy is determined by  $\boldsymbol{\alpha}_1 \in \mathbb{R}^p$  and  $\alpha_0 \in \mathbb{R}$ .

Figures 4 and 5 contain the estimated value of policies in both simulation settings. They indicate that RL-RDS outperforms all competitor policies by a significant margin in each regime. Additionally, the effect sizes increase slightly as the network becomes sparser and the similarities between neighbors become stronger (because there is more “signal” for the polices to leverage). Lastly, note that the train and implement method does not adapt its policy after its “pilot study,” causing the margin between RL-RDS




Figure 4: This figure compares the estimated cumulative reward of each policy with 90% Monte Carlo confidence intervals in simulation setting 1.

and this strategy to generally increase in the larger budget setting.




Figure 5: This figure compares the estimated cumulative reward of each policy with 90% Monte Carlo confidence intervals in simulation setting 2.