Nonparametric estimation of a factorizable density using diffusion models

Hyeok Kyu Kwon¹, Dongha Kim², Ilsang Ohn³, and Minwoo Chae¹

 1 Department of Industrial and Management Engineering Pohang University of Science and Technology

²Department of Statistics, Sungshin Women's University ³Department of Statistics, Inha University

Abstract

In recent years, diffusion models, and more generally score-based deep generative models, have achieved remarkable success in various applications, including image and audio generation. In this paper, we view diffusion models as an implicit approach to nonparametric density estimation and study them within a statistical framework to analyze their surprising performance. A key challenge in high-dimensional statistical inference is leveraging low-dimensional structures inherent in the data to mitigate the curse of dimensionality. We assume that the underlying density exhibits a low-dimensional structure by factorizing into low-dimensional components, a property common in examples such as Bayesian networks and Markov random fields. Under suitable assumptions, we demonstrate that an implicit density estimator constructed from diffusion models adapts to the factorization structure and achieves the minimax optimal rate with respect to the total variation distance. In constructing the estimator, we design a sparse weight-sharing neural network architecture, where sparsity and weight-sharing are key features of practical architectures such as convolutional neural networks and recurrent neural networks.

Keywords: Bayesian network, diffusion model, factorizable density, Markov random field, minimax optimality, score-based generative model, weight-sharing neural network.

1 Introduction

Suppose we have observations X^1, \ldots, X^n , which are independent and identically distributed Ddimensional random variables following an unknown distribution P_0 with density p_0 . Inference of the unknown P_0 (or its density p_0) is a fundamental task in unsupervised learning, and various methodologies and related theories have been developed over the past few decades (e.g. [Hastie et al.,](#page-23-0) [2009,](#page-23-0) [Tsybakov,](#page-26-0) [2008,](#page-26-0) Giné and Nickl, [2016\)](#page-23-1). For large D , however, inferring high-dimensional distributions becomes prohibitively difficult due to the well-known phenomenon called the curse of dimensionality.

Even when the dimension D is large, real-world data often exhibit various low-dimensional structures. In such cases, one can leverage this structural information in statistical inference to effectively mitigate the curse of dimensionality. Sparsity [\(Hastie et al.,](#page-23-2) [2015\)](#page-23-2) of high-dimensional parameters is one of the most important low-dimensional structures in the statistical literature. Additive [\(Hastie](#page-23-3) [and Tibshirani,](#page-23-3) [1990\)](#page-23-3) and manifold structures [\(Genovese et al.,](#page-23-4) [2012\)](#page-23-4), where data are assumed to lie close to a low-dimensional manifold within an ambient space, are also well-studied in nonparametric contexts.

When information about the underlying low-dimensional structure is available, it is often possible to construct statistical procedures that perform substantially better than those that ignore this structural information. In real data analysis, however, knowledge about whether the data under consideration possesses a specific structure is often unavailable, except in very specific circumstances. As a result, statistical procedures that can adapt to various structures are preferred. Here, adaptation roughly means that a procedure performs as well as one that explicitly leverages the structural information. For instance, deep neural networks (DNNs) exhibit such adaptive properties in various structural function estimation problems [\(Imaizumi and Fukumizu,](#page-24-0) [2022,](#page-24-0) [Schmidt-Hieber,](#page-26-1) [2020,](#page-26-1) [Tang](#page-26-2) [and Yang,](#page-26-2) [2024,](#page-26-2) [Chae et al.,](#page-22-0) [2023\)](#page-22-0).

In this paper, we focus on a specific low-dimensional structure that accommodates a broad family of probability distributions. Although this structure is well-known in statistical communities, nonparametric adaptive statistical procedures for it have rarely been studied in the literature. Specifically, we assume that the density function p_0 is factorizable as

$$
p_0(\mathbf{x}) = \prod_{I \in \mathcal{I}} g_I(\mathbf{x}_I), \tag{1.1}
$$

where $\mathcal{I} \subseteq 2^{[D]}$ is a collection of index sets, $\mathbf{x}_I = (x_i)_{i \in I}$, and each g_I is a |I|-variate function. Here, $[D] = \{1, \ldots, D\}$, and |I| denotes the cardinality of I. Note that such factorizable densities are commonly found in graphical model contexts [\(Liu and Lafferty,](#page-25-0) [2019\)](#page-25-0), such as Bayesian networks and Markov random fields. In particular, the conditional independence structure induced by undirected graphical models (Markov random fields) is highly suitable for modeling images, where spatially close pixels tend to be strongly correlated, while those far apart exhibit weak correlations [\(Ji,](#page-24-1) [2020,](#page-24-1) [Vandermeulen et al.,](#page-27-0) [2024b,](#page-27-0)[a\)](#page-27-1). See Section [4](#page-7-0) for further details on these examples.

If the density function p_0 belongs to a β -Hölder class, one can construct an estimator converging to p_0 as the sample size n tends to infinity with the rate of $n^{-\beta/(D+2\beta)}$ with respect to the total variation distance. This is a well-known minimax optimal rate for a class of β -Hölder densities [\(Tsybakov,](#page-26-0) [2008,](#page-26-0) Giné and Nickl, 2016). If we reduce the density class to factorizable densities as in (1.1) for a given I , classical nonparametric theory states that the corresponding minimax optimal rate becomes $n^{-\beta/(d+2\beta)}$, where $d = \max_{I \in \mathcal{I}} |I|$ is the *effective dimension* corresponding to the worst component function. (Here, we assumed that D is fixed and all component functions have the same degree of smoothness β .) Once we know p_0 is factorizable as in [\(1.1\)](#page-1-0), it is not difficult to construct an estimator for p_0 with the rate $n^{-\beta/(d+2\beta)}$ under suitable technical assumptions. It is challenging, however, to construct an estimator that is adaptive to the factorization structure. To the best of our knowledge, (theoretically) adaptive estimators, not necessarily achieving the optimal rate, have only been considered in very recent articles [\(Bos and Schmidt-Hieber,](#page-22-1) [2024,](#page-22-1) [Vandermeulen et al.,](#page-27-0) [2024b,](#page-27-0)[a\)](#page-27-1).

Diffusion models [\(Sohl-Dickstein et al.,](#page-26-3) [2015,](#page-26-3) [Ho et al.,](#page-23-5) [2020,](#page-23-5) [Song et al.,](#page-26-4) [2020,](#page-26-4) [2021\)](#page-26-5) are scorebased generative models that have demonstrated remarkable success in recent years across many applications. Here, score-based means that they model the score function, which is the gradients of the log density. Even compared to modern deep generative models such as variational autoencoders (VAEs) [\(Kingma and Welling,](#page-24-2) [2014,](#page-24-2) [Rezende et al.,](#page-26-6) [2014\)](#page-26-6), generative adversarial networks (GANs) [\(Goodfellow et al.,](#page-23-6) [2014,](#page-23-6) [Arjovsky et al.,](#page-22-2) [2017,](#page-22-2) [Mroueh et al.,](#page-25-1) [2018\)](#page-25-1), and normalizing flows [\(Dinh et al.,](#page-23-7) [2015,](#page-23-7) [Rezende and Mohamed,](#page-25-2) [2015\)](#page-25-2), diffusion models have achieved state-of-the-art performance in several domains, including images [\(Rombach et al.,](#page-26-7) [2022,](#page-26-7) [Dhariwal and Nichol,](#page-23-8) [2021\)](#page-23-8), videos [\(Ho](#page-23-9) [et al.,](#page-23-9) [2022\)](#page-23-9), and audio [\(Kong et al.,](#page-24-3) [2021\)](#page-24-3).

Diffusion models operate via two complementary stochastic processes. First, a forward diffusion process gradually transforms the initial distribution P_0 into a simple noise distribution. At each time step of this forward process, the score function of the marginal density is estimated by minimizing a suitable score matching objective. Second, a reverse process is defined as the time-reversal of the forward process. To generate samples, one begins with the noise distribution and approximates the reverse process by plugging in the estimated score function. Further details are provided in Section [2.](#page-3-0)

Despite the remarkable empirical success of diffusion models, their statistical properties remain largely unexplored. Although diffusion models do not produce an explicit density estimator, one can define an implicit density estimator from the generative process. A primary goal of this paper is to prove that this implicit density estimator is adaptive to the factorization structure [\(1.1\)](#page-1-0) and achieves the minimax optimal convergence rate $n^{-\beta/(d+2\beta)}$, up to a logarithmic factor, with respect to the total variation distance (Theorem [5.2\)](#page-15-0).

To obtain an optimal estimator, we only need to carefully choose a network architecture. Notably, we consider *sparse weight-sharing neural networks* to construct an optimal estimator. Although sparse weight-sharing networks, such as convolutional neural networks (CNNs; [LeCun et al.,](#page-24-4) [1989,](#page-24-4) [Krizhevsky et al.,](#page-24-5) [2012\)](#page-24-5) and recurrent neural networks [\(Rumelhart et al.,](#page-26-8) [1986,](#page-26-8) [Sutskever et al.,](#page-26-9) [2014\)](#page-26-9), are widely used in practical applications, their theoretical benefits have not been thoroughly studied in the literature. To the best of our knowledge, there are only a few articles showing that CNNs perform as well as vanilla feedforward neural networks [\(Petersen and Voigtlaender,](#page-25-3) [2020,](#page-25-3) [Oono and Suzuki,](#page-25-4) [2019,](#page-25-4) [Yang et al.,](#page-27-2) [2024,](#page-27-2) [Fang and Cheng,](#page-23-10) [2023\)](#page-23-10). While we are not aware whether the optimality of the estimator remains valid if weight-sharing networks are replaced by vanilla feedforward neural networks, weight-sharing plays a key role in our construction to reduce model complexity.

Statistical theories for diffusion models have been studied in several recent articles. [Oko et al.](#page-25-5) [\(2023\)](#page-25-5) proved that the implicit density estimator from the diffusion model is minimax optimal within the nonparametric smooth density estimation framework, using total variation and Wasserstein distances as evaluation metrics. Subsequently, [Zhang et al.](#page-27-3) [\(2024\)](#page-27-3) and [Wibisono et al.](#page-27-4) [\(2024\)](#page-27-4) relaxed certain technical assumptions in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5). Although these papers introduced several interesting mathematical techniques for handling diffusion models, they did not address the issue of the curse of dimensionality. To tackle this issue, [Tang and Yang](#page-26-2) [\(2024\)](#page-26-2) demonstrated that the estimator from the diffusion model is minimax optimal with respect to the Wasserstein metric under the smooth manifold assumption. Under a similar regime, [Azangulov et al.](#page-22-3) [\(2024\)](#page-22-3) established tighter upper bounds for the convergence rate in terms of the ambient dimension D. While the manifold structure is an interesting low-dimensional structure, an optimal estimator adaptive to this structure can also be constructed using methods other than diffusion models [\(Tang and Yang,](#page-26-10) 2023 , Stéphanovitch et al., [2024\)](#page-26-11).

Building on these existing theories, our results provide new insights into diffusion models, demonstrating their adaptivity to novel low-dimensional structures. In particular, while various interesting statistical theories have been developed for VAEs [\(Kwon and Chae,](#page-24-6) [2024,](#page-24-6) [Chae et al.,](#page-22-0) [2023\)](#page-22-0) and GANs [\(Liang,](#page-24-7) [2021,](#page-24-7) [Uppal et al.,](#page-26-12) [2019,](#page-26-12) [Chae,](#page-22-4) [2022,](#page-22-4) Stéphanovitch et al., [2024,](#page-26-11) [Tang and Yang,](#page-26-10) [2023,](#page-26-10) [Puchkin et al.,](#page-25-6) [2024\)](#page-25-6), the factorization structure (1.1) , which is closely related to the conditional independence structure of directed and undirected graphs, has not been explored in the literature on deep generative models. While the estimators proposed in [Bos and Schmidt-Hieber](#page-22-1) [\(2024\)](#page-22-1), [Vandermeulen](#page-27-0) [et al.](#page-27-0) [\(2024b,](#page-27-0)[a\)](#page-27-1) are adaptive to the factorization structure, diffusion models are not only adaptive to this structure but also to other structures discussed above, making them significantly more practical alternatives.

The remainder of this paper is organized as follows. In Section [2,](#page-3-0) we introduce diffusion models and define our implicit density estimator. Section [3](#page-5-0) presents the class of weight-sharing networks, while Section [4](#page-7-0) details the main assumption—the factorization assumption. Our main theoretical results are provided in Section [5.](#page-9-0) In Section [6,](#page-17-0) we discuss the benefits of diffusion models compared to the vanilla score matching estimator. We present small experimental results in Section [7](#page-18-0) and conclude with discussions in Section [8.](#page-20-0) All proofs are provided in the Appendix.

1.1 Notations and definitions

Vectors are denoted using boldface notation. For a multi-index $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_D)^\top \in (\mathbb{Z}_{\geq 0})^D$, denote D^γ the mixed partial derivative operator $\partial^{\gamma} \setminus \partial x_1^{\gamma_1} \cdots \partial x_D^{\gamma_D}$, where $\gamma = \sum_{i=1}^D \gamma_i$. For any $\beta, K > 0$, let $\mathcal{H}_{D}^{\beta,K}(A)$ be the class of every real-valued function g on $A\subseteq\mathbb{R}^D$ such that

$$
\sum_{\gamma \leq \lfloor \beta \rfloor} \sup_{{\bf x} \in A} |(D^{\gamma} g)({\bf x})| + \sum_{\gamma = \lfloor \beta \rfloor} \sup_{{\bf x}, {\bf y} \in A \atop {\bf x} \neq {\bf y}} \frac{|(D^{\gamma} g)({\bf x}) - (D^{\gamma} g)({\bf y})|}{\|{\bf x} - {\bf y}\|_{\infty}^{\beta - \lfloor \beta \rfloor}} \leq K,
$$

where $\lfloor \beta \rfloor$ denotes the largest integer strictly smaller than β . We often denote $\mathcal{H}_{D}^{\beta,K}(A)$ as $\mathcal{H}^{\beta,K}(A)$ when the dimension is obvious from the contexts. For a vector **x**, we denote the ℓ^p -norm, $1 \leq p \leq \infty$, and the number of nonzero elements as $\|\mathbf{x}\|_p$ and $\|\mathbf{x}\|_0$, respectively. Let $\phi_{\sigma,D}$ be the density function of the multivariate normal distribution $\mathcal{N}(\mathbf{0}_D, \sigma^2 \mathbb{I}_D)$, where $\mathbf{0}_D$ and \mathbb{I}_D are D-dimensional zero vector and identity matrix, respectively. For simplicity, we often denote $\phi_{\sigma,D}$ as ϕ_{σ} when the dimension is obvious from the contexts. The notation $a \leq b$ means that $a \leq Cb$, where C is some constant that is non-important in the given context. Similarly, $a \approx b$ implies that $a \lesssim b$ and $b \lesssim a$. Finally, the notation $C = C(A_1, \ldots, A_n)$ means that the constant C depends only on A_1, \ldots, A_n .

2 Diffusion models

In this section, we provide a brief introduction to the diffusion model proposed in [Song et al.](#page-26-5) [\(2021\)](#page-26-5) and define the estimator studied in our main results. Let $(\mathbf{X}_t)_{t\geq 0}$ be the process satisfying the stochastic differential equation (SDE)

$$
d\mathbf{X}_t = -\alpha_t \mathbf{X}_t dt + \sqrt{2\alpha_t} dB_t, \quad \mathbf{X}_0 \sim P_0,
$$
\n(2.1)

where $(\mathbf{B}_t)_{t\geq 0}$ is a standard D-dimensional Brownian motion and $t\mapsto \alpha_t : [0,\infty) \to [0,\infty)$ is a (known) Borel measurable function. The stochastic process (X_t) is often referred to as a time-inhomogeneous Ornstein-Uhlenbeck (OU) process, and has been studied in [Song et al.](#page-26-5) [\(2021\)](#page-26-5), [Chen et al.](#page-23-11) [\(2023b\)](#page-23-11). For the OU process (2.1) , the transition kernel is explicitly given as Gaussian. Specifically, the conditional distribution of \mathbf{X}_t given $\mathbf{X}_0 = \mathbf{x}_0$ is $\mathcal{N}(\mu_t \mathbf{x}_0, \sigma_t^2 \mathbb{I}_D)$, where $\mu_t = \exp(-\int_0^t \alpha_s \, ds)$ and $\sigma_t^2 = 1 - \mu_t^2$. We denote this conditional distribution and the corresponding density as $P_t(\cdot | \mathbf{x}_0)$ and $p_t(\cdot | \mathbf{x}_0)$, respectively. We also denote P_t and p_t as the marginal distribution and density of \mathbf{X}_t , respectively. Hence, we have

$$
p_t(\mathbf{x}) = \int \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}P_0(\mathbf{y}) = \int \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) p_0(\mathbf{y}) \mathrm{d}\mathbf{y}.
$$

Note that p_t converges very quickly to the standard Gaussian density as $t \to \infty$; see [Bakry et al.](#page-22-5) [\(2014\)](#page-22-5) for a rigorous statement. Let $f_0(x, t) = \nabla \log p_t(x)$. Note that the map $x \mapsto f_0(x, t)$ is the score function corresponding to the marginal density p_t . As a convention, we also call f_0 a score function.

For a given (non-random) $T > 0$, let $(Y_t)_{t \in [0,\overline{T})}$ be the reverse-time process defined as $Y_t =$ $\mathbf{X}_{\overline{T}-t}$. Then, it is well-known [\(Anderson,](#page-22-6) [1982\)](#page-22-6) that $(\mathbf{Y}_t)_{t\in[0,\overline{T})}$ is also a diffusion process under mild assumptions. More specifically, once

$$
\int_s^{\overline{T}} \mathbb{E}\left[p_t(\mathbf{X}_t) + 2\alpha_t \left\|\nabla \log p_t(\mathbf{X}_t)\right\|_2^2 \right] dt < \infty \quad \forall s > 0
$$

and the map $t \mapsto \alpha_t$ is bounded above, we have

$$
d\mathbf{Y}_{t} = \left[\alpha_{\overline{T}-t}\mathbf{Y}_{t} + 2\alpha_{\overline{T}-t}\nabla\log p_{\overline{T}-t}(\mathbf{Y}_{t})\right]dt + \sqrt{2\alpha_{\overline{T}-t}}d\mathbf{B}_{t}
$$

\n
$$
= \left[\alpha_{\overline{T}-t}\mathbf{Y}_{t} + 2\alpha_{\overline{T}-t}\mathbf{f}_{0}(\mathbf{Y}_{t},\overline{T}-t)\right]dt + \sqrt{2\alpha_{\overline{T}-t}}d\mathbf{B}_{t}, \quad \mathbf{Y}_{0} \sim P_{\overline{T}},
$$
\n(2.2)

see Threom 2.1 of [Haussmann and Pardoux](#page-23-12) [\(1986\)](#page-23-12). Note that the Brownian motions in [\(2.1\)](#page-3-1) and (2.2) are not identical. However, we use the same notation B_t to denote a standard Brownian motion as a convention throughout the paper.

Once we have an estimator $\hat{\mathbf{f}}$ for the score function \mathbf{f}_0 , one can simulate the reverse process starting from a standard Gaussian to obtain samples from the estimated distribution. The score function can be estimated via the score matching (Hyvärinen, [2005\)](#page-23-13) or its scalable variations [\(Vincent,](#page-27-5) [2011,](#page-27-5) [Song](#page-26-4) [et al.,](#page-26-4) [2020,](#page-26-4) [Yu et al.,](#page-27-6) [2022\)](#page-27-6).

Let F be a class of functions $(\mathbf{x}, t) \mapsto \mathbf{f}(\mathbf{x}, t)$ used to model the score function \mathbf{f}_0 . A detailed description of the class $\mathcal F$ in our theory is provided in Section [3.](#page-5-0) At the population level, the best approximator to f_0 in F can be defined as the solution to the following optimization problem

$$
\underset{\mathbf{f}\in\mathcal{F}}{\text{minimize}} \int_0^{\overline{T}} \lambda_t \mathbb{E}\left[\|\mathbf{f}(\mathbf{X}_t, t) - \mathbf{f}_0(\mathbf{X}_t, t)\|_2^2\right] dt
$$
\n
$$
\iff \underset{\mathbf{f}\in\mathcal{F}}{\text{minimize}} \int_0^{\overline{T}} \lambda_t \mathbb{E}\left[\|\mathbf{f}(\mathbf{X}_t, t) - \nabla \log p_t(\mathbf{X}_t)\|_2^2\right] dt,
$$
\n(2.3)

where $\lambda_t \geq 0$ is a weight. Based on the well-known fact [\(Vincent,](#page-27-5) [2011\)](#page-27-5) that

$$
\mathbb{E}\left[\|\mathbf{f}(\mathbf{X}_t,t)-\nabla \log p_t(\mathbf{X}_t)\|_2^2\right] = \mathbb{E}\left[\|\mathbf{f}(\mathbf{X}_t,t)-\nabla \log p_t(\mathbf{X}_t\mid \mathbf{X}_0)\|_2^2\right] + C_t,
$$

where C_t is a constant depending only on t and f_0 and

$$
\nabla \log p_t(\mathbf{x}_t \mid \mathbf{x}_0) = \frac{\partial \log p_t(\mathbf{x}_t \mid \mathbf{x}_0)}{\partial \mathbf{x}_t} = -\frac{\mathbf{x}_t - \mu_t \mathbf{x}_0}{\sigma_t^2},
$$

the minimization problem (2.3) can be equivalently written as

$$
\text{minimize}_{\mathbf{f} \in \mathcal{F}} \int_0^{\overline{T}} \lambda_t \mathbb{E} \left[\| \mathbf{f}(\mathbf{X}_t, t) - \nabla \log p_t(\mathbf{X}_t \mid \mathbf{X}_0) \|_2^2 \right] dt
$$

$$
\iff \text{minimize}_{\mathbf{f} \in \mathcal{F}} \int_0^{\overline{T}} \lambda_t \mathbb{E} \left[\mathbb{E} \left(\left\| \mathbf{f}(\mathbf{X}_t, t) + \frac{\mathbf{X}_t - \mu_t \mathbf{X}_0}{\sigma_t^2} \right\|_2^2 \mid \mathbf{X}_0 \right) \right] dt.
$$

In practice, λ_t is set to zero for sufficiently small t to avoid potential singularity issues. This leads to the following ERM (empirical risk minimization) estimator

$$
\widehat{\mathbf{f}} \in \underset{\mathbf{f} \in \mathcal{F}}{\operatorname{argmin}} \frac{1}{n} \sum_{i=1}^{n} \ell_{\mathbf{f}}(\mathbf{X}^{i}),\tag{2.4}
$$

where

$$
\ell_{\mathbf{f}}(\mathbf{x}) = \int_{\underline{T}}^{\overline{T}} \lambda_t \mathbb{E} \left[\left\| \mathbf{f}(\mathbf{X}_t, t) + \frac{\mathbf{X}_t - \mu_t \mathbf{X}_0}{\sigma_t^2} \right\|_2^2 \mid \mathbf{X}_0 = \mathbf{x} \right] dt \tag{2.5}
$$

is the loss function and $\underline{T} > 0$ is a sufficiently small number.

Let $(\mathbf{Y}_t)_{t \in [0, \overline{T} - \underline{T}]}$ be the solution to the SDE

$$
\mathrm{d}\widehat{\mathbf{Y}}_t = \left[\alpha_{\overline{T}-t} \widehat{\mathbf{Y}}_t + 2\alpha_{\overline{T}-t} \widehat{\mathbf{f}}\left(\widehat{\mathbf{Y}}_t, \overline{T}-t\right) \right] \mathrm{d}t + \sqrt{2\alpha_{\overline{T}-t}} \mathrm{d}\mathbf{B}_t, \quad \widehat{\mathbf{Y}}_0 \sim \mathcal{N}(\mathbf{0}_D, \mathbb{I}_D) \tag{2.6}
$$

and $\mathbf{X}_t = \mathbf{Y}_{T-t}$. Let P_t be the marginal distribution of \mathbf{X}_t and \hat{p}_t be the corresponding Lebesgue density. Also, let $P = P_T$ and $\hat{p} = \hat{p}_T$. The existence of \hat{p}_t is guaranteed under mild assumptions; see [Bogachev et al.](#page-22-7) [\(2011\)](#page-22-7) for details. Although \hat{p}_t is only defined implicitly through the SDE [\(2.6\)](#page-5-1), it is a function of data, hence an estimator for the unknown density p_t . Since we expect that $p_t \approx p_0$ for sufficiently small t, \hat{p} can serve as an estimator for p_0 .

Remark 2.1. Note that the loss function (2.5) involves integrals (with respect to \mathbf{X}_t and t) that are not directly tractable. In practice, a slightly different loss function with augmented variables is considered for computational tractability [\(Sohl-Dickstein et al.,](#page-26-3) [2015,](#page-26-3) [Song and Ermon,](#page-26-13) [2019\)](#page-26-13). Specifically, with a slight abuse of notation, define the loss function

$$
\ell_{\mathbf{f}}(\mathbf{x}_0, \mathbf{x}_t, t) = \left\| \mathbf{f}(\mathbf{x}_t, t) + \frac{\mathbf{x}_t - \mu_t \mathbf{x}_0}{\sigma_t^2} \right\|_2^2.
$$

By regarding T as a random variable independent of the stochastic process (X_t) and supported on $[T, T]$ with the density proportional to λ_t , we have

$$
\mathbb{E}\ell_{\mathbf{f}}(\mathbf{X}_0,\mathbf{X}_T,T)=\mathbb{E}\ell_{\mathbf{f}}(\mathbf{X}_0).
$$

Therefore, although the loss function [\(2.6\)](#page-5-1) itself is not directly tractable, one can approximate the solution to the minimization problem (2.4) using stochastic gradient methods.

Remark 2.2. The target estimator in our theoretical study in Section [5](#page-9-0) is \hat{p} as defined above, which is the density of $\mathbf{X}_{\underline{T}} = \mathbf{Y}_{\overline{T} - \underline{T}}$. In practice, samples from the estimated distribution are generated by numerically solving the SDE (2.6) using methods such as Euler-Maruyama discretization [\(Kloeden](#page-24-8) [and Platen,](#page-24-8) [2011,](#page-24-8) [Song et al.,](#page-26-5) [2021\)](#page-26-5), starting from an initial sample drawn from the standard normal distribution. Hence, more delicate statistical theory should incorporate these discretization errors. As an independent line of work, there are various articles studying discretization errors in diffusion models [\(Oko et al.,](#page-25-5) [2023,](#page-25-5) [Chen et al.,](#page-23-11) [2023b,](#page-23-11) [Nakano,](#page-25-7) [2024,](#page-25-7) [Bortoli,](#page-22-8) [2022,](#page-22-8) [Benton et al.,](#page-22-9) [2024,](#page-22-9) [Chen](#page-22-10) [et al.,](#page-22-10) [2023a,](#page-22-10) [Li et al.,](#page-24-9) [2024\)](#page-24-9). Combining our statistical theory given in Section [5](#page-9-0) with these works, the main results remain valid if the SDE is discretized with a sufficiently fine time partition. For additional details, we refer to Section 5.3 of [Oko et al.](#page-25-5) [\(2023\)](#page-25-5).

3 Weight-sharing neural networks

In this section, we define neural networks that are used as a function class $\mathcal F$ to model the score function f_0 described in Section [2.](#page-3-0) Instead of vanilla feedforward neural networks, we consider sparse weight-sharing architectures, which are widely used in practical applications. By incorporating such sparsity and weight-sharing structures into the network architecture that models the score function, one can substantially reduce the model complexity (often expressed in terms of metric entropy; see Lemma [C.2](#page-88-0)), ultimately leading to a reduction in estimation error.

For a positive integer m, let $\rho_m : \mathbb{R}^m \to \mathbb{R}^m$ be the (coordinatewise) ReLU activation function defined as

$$
\rho_m(\mathbf{x}) = (\max\{x_1, 0\}, \dots, \max\{x_m, 0\})^\top
$$

for $\mathbf{x} = (x_1, \ldots, x_m)^\top$. For simplicity, we often denote ρ_m as ρ . For $L \in \mathbb{N}_{\geq 2}$, $\mathbf{d} = (d_1, \ldots, d_{L+1}) \in$ $\mathbb{N}^{L+1}, s \in \mathbb{N}, M > 0, m = (m_1, \ldots, m_{L-1}) \in \mathbb{N}^{L-1}$ and $\mathcal{P}_{m} = ((\mathcal{Q}_i, \mathcal{R}_i))_{i \in [L-1]},$ where $\mathcal{Q}_i =$ $(Q_i^{(j)}$ $\binom{(j)}{i}_{j\in[m_i]}$ is a collection of $d_i \times d_i$ permutation matrices and $\mathcal{R}_i = (R_i^{(j)})$ $(i^{(j)})_{j\in[m_i]}$ is a collection of $d_{i+1} \times d_{i+1}$ permutation matrices, let $\mathcal{F}_{\text{WSNN}} = \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\text{m}})$ be the class of functions $\mathbf{f} : \mathbb{R}^{d_1} \to \mathbb{R}^{d_{L+1}}$ of the form

$$
\mathbf{f}(\mathbf{x}) = W_L \left(\mathbf{f}_{L-1} \circ \cdots \circ \mathbf{f}_1 \right) (\mathbf{x}) + \mathbf{b}_L,
$$

$$
\mathbf{f}_i(\cdot) = \rho \left(\sum_{j=1}^{m_i} R_i^{(j)} \left(W_i Q_i^{(j)} \cdot + \mathbf{b}_i \right) \right)
$$
(3.1)

with $W_i \in \mathbb{R}^{d_{i+1} \times d_i}$ and $\mathbf{b}_i \in \mathbb{R}^{d_{i+1}}$ satisfying

$$
\max_{1 \leq i \leq L} \{ \max\left(||W_i||_{\infty}, ||\mathbf{b}_i||_{\infty} \right) \} \leq M, \quad \sum_{i=1}^L ||W_i||_0 + ||\mathbf{b}_i||_0 \leq s.
$$

Here, $||W_i||_{\infty}$ and $||W_i||_0$ denote the entrywise maximum norm and the number of nonzero elements of the matrix W_i , respectively.

The network [\(3.1\)](#page-6-0) includes vanilla feedforward neural networks as special cases. For example, if $m = (1, \ldots, 1)$ and all matrices in \mathcal{P}_m are identity matrices, the class \mathcal{F}_{WSNN} reduces to the usual class of sparse networks considered in the literature. In this case, we often denote $\mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\text{m}})$ as $\mathcal{F}_{NN}(L, \mathbf{d}, s, M)$.

The network [\(3.1\)](#page-6-0) is designed to incorporate sparsity and weight-sharing into the architecture. As a simple example, consider a weight matrix $W \in \mathbb{R}^{3d \times 4d}$ with the following block structure, where each block has the same size of $d \times d$:

$$
W = \begin{bmatrix} W_0 & 0_{d,d} & 0_{d,d} & W_0 \\ 0_{d,d} & 0_{d,d} & W_0 & 0_{d,d} \\ 0_{d,d} & W_0 & 0_{d,d} & W_0 \end{bmatrix}.
$$

Here, $0_{d,d}$ denotes the $d \times d$ zero-matrix. Two important features of W are that it is sparse, in the sense that many elements of W are exactly zero, and that the sub-matrix W_0 is shared across different rows and columns. The formula (3.1) is one way to effectively represent neural networks with sparse weight-sharing matrices like W. For example, it is straightforward to construct $3d \times 3d$ permutation matrices $R^{(j)}$ and $4d \times 4d$ permutation matrices $Q^{(j)}$, for $j \leq 5$, that satisfy

$$
W = \sum_{j=1}^{5} R^{(j)} \begin{bmatrix} W_0 & 0_{d,3d} \\ 0_{2d,d} & 0_{2d,3d} \end{bmatrix} Q^{(j)}.
$$

The number of nonzero elements of W is $5d^2$, but we can express it with a smaller sparsity of d^2 by weight-sharing architecture.

Sparse weight-sharing matrices are used in many practically important architectures, such as convolutional neural networks [\(LeCun et al.,](#page-24-4) [1989,](#page-24-4) [Krizhevsky et al.,](#page-24-5) [2012\)](#page-24-5) and recurrent neural networks [\(Rumelhart et al.,](#page-26-8) [1986,](#page-26-8) [Sutskever et al.,](#page-26-9) [2014\)](#page-26-9); see also [Zhang et al.](#page-27-7) [\(2021\)](#page-27-7), [Jagtap et al.](#page-24-10) [\(2022\)](#page-24-10) for additional examples. Note that CNN architectures are frequently adopted in diffusion models [\(Sohl-Dickstein et al.,](#page-26-3) [2015,](#page-26-3) [Ho et al.,](#page-23-5) [2020,](#page-23-5) [Ronneberger et al.,](#page-26-14) [2015\)](#page-26-14).

As an illustrative example, consider a convolution operation with an input image **x** of size $d_1 = s_1^2$, a filter of size $s = s_0^2$, and an output image y of size $d_2 = (s_1 - s_0 + 1)^2$, as illustrated in Figure [1.](#page-8-0) Let $\mathbf{w} = (w_1, \dots, w_s)^\top$ be the vectorized version of the filter, and let $\widetilde{W} \in \mathbb{R}^{d_2 \times d_1}$ be the weight matrix corresponding to the convolution operation, such that $y = \widetilde{W}x$. One can observe that \widetilde{W} is a sparse weight-sharing matrix and can be represented in the form of [\(3.1\)](#page-6-0). To see this, note that each row of \widetilde{W} can be obtained by permuting the vector $(\mathbf{w}^\top, 0_{1,d_1-s}) \in \mathbb{R}^{d_1}$. Thus, for $j \in [d_2]$, there exists a $d_1 \times d_1$ permutation matrix $Q^{(j)}$ such that $y_j = (\mathbf{w}^\top, 0_{1,d_1-s})Q^{(j)}\mathbf{x}$. Also, for $j \in [d_2]$, let $R^{(j)}$ be the $d_2 \times d_2$ permutation matrix that swaps the first and the jth row when it is left-multiplied by a matrix. Then, we can express the matrix \widetilde{W} in the form of [\(3.1\)](#page-6-0) by

$$
\widetilde{W} = \sum_{j=1}^{d_2} R^{(j)} W Q^{(j)}, \quad W = \begin{pmatrix} \mathbf{w}^{\top} & 0_{1, d_1 - s} \\ 0_{d_2 - 1, s} & 0_{d_2 - 1, d_1 - s} \end{pmatrix} \in \mathbb{R}^{d_2 \times d_1}.
$$

One may additionally incorporate padding and stride operations [\(Paszke et al.,](#page-25-8) [2019\)](#page-25-8) into the convolution operation described in Figure [1.](#page-8-0) The corresponding weight matrix, with these additional operations, can also be expressed in the form of (3.1) by carefully selecting the permutation matrices.

4 Factorizable densities

4.1 Factrorization assumption

In this section, we introduce the low-dimensional assumption considered in our main results and provide some well-known examples. Formally, we consider the following factorization assumption.

(F) There exists a set $\mathcal{I} \subseteq 2^{[D]}$ and functions $g_I : \mathbb{R}^{|I|} \to \mathbb{R}$ for each $I \in \mathcal{I}$ such that

$$
p_0(\mathbf{x}) = \prod_{I \in \mathcal{I}} g_I(\mathbf{x}_I), \quad \forall \mathbf{x} \in \mathbb{R}^D.
$$

For a density p_0 satisfying (F) , let $d = \max_{I \in \mathcal{I}} |I|$ denote the largest number of variables that any g_I depends on. We refer to d as the effective dimension corresponding to the factorizable density p_0 . As a simple example, if p_0 is the density of a random vector $\mathbf{X} = (X_1, \ldots, X_D)$ and each component of **X** is mutually independent, then $d = 1$. In the following subsections, we present examples based on conditional independence structures, which are often represented using graphical models.

4.2 Example: Bayesian networks

A Bayesian network is a random vector $\mathbf{X} = (X_1, \ldots, X_D)$ whose conditional independence structure can be represented by a directed acyclic graph (DAG) with the vertex set $\{1, \ldots, D\}$. For a Bayesian network, each variable X_i is conditionally independent of all other variables given its parent variables

	$\overline{0}$								
	$0\quad 0\quad 0$ $0 \quad 0$	$\overline{0}$ $0 \quad 0 \quad 0 \quad 0$	$\overline{}$ $\overline{}$ $\overline{0}$	$\overline{\mathbf{0}}$					$(w_1 \ w_2 \ 0 \ 0 \ w_3 \ w_4 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0$ $0 \quad w_1 \ w_2 \quad 0 \quad 0 \quad w_3 \ w_4 \quad 0 \quad 0$ 0 0 w_1 w_2 0 0 w_3 w_4 0 0 0 0 0 0 0 0 0 $0 \t 0 \t 0 \t 0 \t w_1 \t w_2 \t 0 \t 0 \t w_3 \t w_4 \t 0 \t 0 \t 0 \t 0 \t 0 \t 0$ $0\quad 0\quad 0\quad 0\quad 0\quad 0\quad w_1\ w_2\quad 0\quad 0\quad w_3\ w_4\quad 0\quad 0\quad 0\quad 0\quad 0\quad 0$ $0 \t 0 \t 0 \t 0 \t 0 \t 0 \t w_1 \t w_2 \t 0 \t 0 \t w_3 \t w_4 \t 0 \t 0 \t 0 \t 0$ $0 \t 0 \t 0 \t w_1 \t w_2 \t 0 \t 0 \t w_3 \t w_4 \t 0 \t 0$ $0 \t 0 \t 0 \t w_1 \t w_2 \t 0 \t 0 \t w_3 \t w_4 \t 0$ 0 0 0 0 0 w_1 w_2 0 0 w_3 w_4

Figure 1: Example of a 2-dimensional convolution operation with an input $\mathbf{x} \in \mathbb{R}^{16}$, a filter vector $\mathbf{w} \in \mathbb{R}^4$ and output $\mathbf{y} \in \mathbb{R}^9$. The operation can be represented as a matrix multiplication [\(Goodfellow](#page-23-14) [et al.,](#page-23-14) [2016\)](#page-23-14), given by $y = \widetilde{W}x$.

 $\mathbf{X}_{pa(i)} = (X_j)_{j \in pa(i)}$, where $pa(i)$ denotes the set of parent indices of vertex i. Accordingly, the density $p_0(\cdot)$ of a Bayesian network **X** factorizes as

$$
p_0(\mathbf{x}) = \prod_{i=1}^D p_i(x_i \mid \mathbf{x}_{pa(i)}) = \prod_{i=1}^D g_i(x_i, \mathbf{x}_{pa(i)}),
$$

where $p_i(\cdot | \mathbf{x}_{pa(i)})$ is the conditional density of X_i given $\mathbf{X}_{pa(i)} = \mathbf{x}_{pa(i)}$ and $g_i(x_i, \mathbf{x}_{pa(i)}) = p_i(x_i)$ $\mathbf{x}_{pa(i)}$). Hence, p_0 satisfies the assumption (F) with $\mathcal{I} = \{\{i\} \cup pa(i) : i \in [D]\}\$ and $d = 1 +$ $\max_{i \in [D]} |pa(i)|$, see Figure [2\(a\)](#page-9-1) for an illustrative example.

4.3 Example: Markov random fields

A Markov random field over an undirected graph G with vertex set $\{1, \ldots, D\}$ is a random vector $\mathbf{X} = (X_1, \ldots, X_D)$ with the property that each variable X_i is conditionally independent of all other variables given its neighbors, often referred to as the local Markov property [\(Lauritzen,](#page-24-11) [1996\)](#page-24-11). If the density $p_0(\cdot)$ of **X** is strictly positive, the local Markov property holds if and only if

$$
p_0(\mathbf{x}) = \prod_{C \in \mathcal{C}} g_C(\mathbf{x}_C) \tag{4.1}
$$

(a) A Bayesian network with $d = 4$ (b) A Markov random field with $d = 3$

Figure 2: Examples of directed and undirected graphical model structures for a 7-dimensional random vector. In both cases, the effective dimension d is strictly less than $D = 7$.

Figure 3: An image of the digit '0' from the MNIST dataset [\(LeCun et al.,](#page-24-12) [1998\)](#page-24-12), along with two possible undirected graph structures for MNIST. For each pixel, a larger neighborhood may be considered depending on the degree of spatial correlations.

for some functions g_C , where C denotes the set of all (maximal) cliques in the graph, as stated by the celebrated Hammersley-Clifford theorem [\(Hammersley and Clifford,](#page-23-15) [1971,](#page-23-15) [Lauritzen,](#page-24-11) [1996\)](#page-24-11). Here, a clique is a fully connected subset of the vertex set in a graph, and the factors g_C are referred to as potential functions in Markov random fields. Therefore, the assumption (F) holds with $\mathcal{I} = \mathcal{C}$ and $d = \max_{C \in \mathcal{C}} |C|$, where d represents the maximum number of vertices in the (maximal) cliques, see Figure [2\(b\)](#page-9-2) for an illustrative example.

Note that images consist of pixels with strong spatial correlations. It is, therefore, natural to assume that each pixel is conditionally independent of all other pixels given the pixels in its neighborhood. This makes the local Markov property particularly suitable for image data. For example, one might consider a graphical model structure, such as in Figure $3(b)$, which has a very small d (e.g., $d = 2$ in this example).

5 Main results

In this section, we present the main results of the paper. We first state the assumptions on p_0 required for the main results. Next, we provide approximation results for the score function f_0 using a weight-sharing network introduced in Section [3.](#page-5-0) Based on these approximation results, we derive

the convergence rate of the diffusion estimator \hat{p} in Section [2,](#page-3-0) which is minimax optimal up to a logarithmic factor.

5.1 Assumptions

For given data $\mathbf{X}^1, \ldots, \mathbf{X}^n$, let \widehat{p} and (\widehat{p}_t) be defined as in Section [2.](#page-3-0) Note that the estimators \widehat{p} and (\widehat{p}_t) depend only on the non-random quantities (α_t) , (λ_t) , \overline{T} , \underline{T} (which may depend on the sample size n), and the architecture $\mathcal{F} = \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\text{m}})$. Recall that α_t is the negative drift coefficient of the forward diffusion (2.1) , λ_t is the weight for the loss function (2.5) , and $[\underline{T},T]$ is the interval defining both the loss function and the estimator \hat{p} . Throughout the paper, we assume the following without explicit restatement:

- 1. $\mathbf{X}^1, \ldots, \mathbf{X}^n$ are i.i.d. from p_0 , supported on $[-1, 1]^D$.
- 2. $\sup_{\gamma \in \mathbb{N}} |D^{\gamma} \alpha_t| \leq 1$ for all $t \geq 0$ and $\underline{\tau} \leq \alpha_t \leq \overline{\tau}$ for constants $\underline{\tau}, \overline{\tau} > 0$.
- 3. $\lambda_t = 1$ for all $t > T$.

Note that the standard OU process, corresponding to $\alpha_t = 1$, satisfies the requirement for α_t . We also note that, with additional details, our main results can be generalized to broader choices of forward diffusion processes and weights λ_t . In addition to these basic assumptions, we will require the following additional assumptions:

(S) The factorization assumption (F) is satisfied and there exist constants $\beta, K > 0$ such that $p_0 \in \mathcal{H}^{\beta,K}([-1,1]^D)$ and $g_I \in \mathcal{H}^{\beta,K}([-1,1]^{|I|})$ for every $I \in \mathcal{I}$.

- (L) There exists a constant $\tau_1 > 0$ such that $p_0(\mathbf{x}) \geq \tau_1$ for every $\mathbf{x} \in [-1, 1]^D$.
- (B) In addition to (S), there exists a constant $\tau_2 \in (0,1)$ such that

$$
\sup_{\gamma \in \mathbb{N}^D} \sup_{\mathbf{x}: 1-\tau_2 \le ||\mathbf{x}||_{\infty} \le 1} |D^{\gamma} p_0(\mathbf{x})| \le K.
$$

Note that Assumptions (L) and (B) are also considered in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5). The assumption (L) on the lower bound of p_0 is required for technical reasons, though we believe it might be eliminated with additional technical developments. However, as our proof is already sufficiently lengthy, we do not explore these further technical details in the present paper. Recent works, such as [Zhang et al.](#page-27-3) (2024) and [Wibisono et al.](#page-27-4) (2024) , do not assume a lower bound on p_0 , but their results rely on a kernel density estimator, which is significantly different from the practical use of diffusion models.

Although p_0 is supported on the compact set $[-1,1]^D$, the support of p_t is \mathbb{R}^D for every $t > 0$. Furthermore, p_t and \hat{p}_t are infinitely differentiable on \mathbb{R}^D for $t > 0$. Accordingly, the difference between p_0 and \hat{p} , when regarded as densities over \mathbb{R}^D , can be large near the boundary of $[-1, 1]^D$. To address this issue, [Oko et al.](#page-25-5) [\(2023\)](#page-25-5) introduced an additional technical assumption that p_0 is infinitely smooth near the boundary, which substantially reduces the approximation error at the boundary. We adopt this strategy, leading to the assumption (B) . We also note that our main result, Theorem [5.2,](#page-15-0) remains valid if the constant τ_2 in assumption (B) is replaced by $(\log n)^{-\tau_{\rm bd}}$, where $\tau_{\rm bd}$ is an arbitrarily large constant.

The factorization assumption (F) , combined with the smooth components assumption (S) , forms the key structural assumption for our main results. Note that our results can be easily extended to the case where each factor function possesses a different level of smoothness. The factorization assumption with smooth nonparametric components has been investigated in the statistical literature under the framework of nonparametric graphical models, specifically in the Markov random fields. [Liu](#page-25-9) [et al.](#page-25-9) $(2011, 2012)$ $(2011, 2012)$ $(2011, 2012)$, Györfi et al. (2023) focused on undirected acyclic graphs (forests), where d is at most 2, and employed kernel methods. With this simple graph structure, [Liu et al.](#page-25-9) [\(2011\)](#page-25-9) developed a consistent graph selection method, while [Liu et al.](#page-25-10) [\(2012\)](#page-25-10) constructed a minimax optimal density estimator for the special case of $\beta = 2$. Further advancements for the case $\beta = 1$ were studied in Györfi et al. (2023) .

Nonparametric statistical theory for general undirected graph structures has been studied in some recent articles. In the case of $\beta = 1$, [Vandermeulen et al.](#page-27-1) [\(2024a](#page-27-1)[,b\)](#page-27-0) proposed estimators whose convergence rates do not depend on the data dimension D. More specifically, [Vandermeulen et al.](#page-27-1) $(2024a)$ introduced a novel quantity called the graph resilience r and derived a convergence rate of $n^{-1/(r+2)}$ (up to a logarithmic factor) with respect to the total variation distance. They showed that this quantity satisfies $d \leq r \leq D$, meaning their rate is optimal only in special cases where $d = r$. Notably, r can be much larger than d , for example, when the graph is a tree. [Vandermeulen et al.](#page-27-0) [\(2024b\)](#page-27-0) studied a more tractable DNN-based estimator with a convergence rate of $n^{-1/(d+4)}$, which is sub-optimal.

[Bos and Schmidt-Hieber](#page-22-1) [\(2024\)](#page-22-1) considered a slightly more general structure than the factorization assumption (F) , using a different type of estimator. Specifically, they assumed that p_0 has a composite structure with smooth component functions. It is well known that deep neural networks can adapt to composite structures in nonparametric function estimation; see [Schmidt-Hieber](#page-26-1) [\(2020\)](#page-26-1), [Bauer and](#page-22-11) [Kohler](#page-22-11) [\(2019\)](#page-22-11), [Kohler and Langer](#page-24-13) [\(2021\)](#page-24-13). [Bos and Schmidt-Hieber](#page-22-1) [\(2024\)](#page-22-1) transformed the density estimation problem into a nonparametric regression problem and then constructed a density estimator. With this approach, they achieved a convergence rate of $n^{-\beta/(d+2\beta)} \vee n^{-\beta/D}$ (up to a logarithmic factor). While this rate improves upon existing results, it is optimal only when $D \leq 2\beta + d$. Note that our main results can also be extended to the composite structure considered in [Bos and Schmidt-](#page-22-1)[Hieber](#page-22-1) [\(2024\)](#page-22-1) without significant difficulty.

5.2 Approximation theory

Theorem [5.1](#page-11-0) below presents the approximation results for the map $(\mathbf{x}, t) \mapsto \mathbf{f}_0(\mathbf{x}, t) = \nabla \log p_t(\mathbf{x})$ using weight-sharing neural networks, which serves as the key technical component of our main results.

Theorem 5.1. Suppose the density function p_0 satisfies the assumptions (S), (L), and (B). Let τ_{\min} and τ_{max} be constants with

$$
\tau_{\min} \ge \frac{4\beta}{d(\beta \wedge 1)} \vee \frac{1}{3D} \quad and \quad \tau_{\max} \ge \frac{2D}{\overline{\tau}}.
$$

Then, for every $m \geq C_2$, there exist a collection of permutation matrices $\mathcal{P}_{\mathbf{m}} = ((\mathcal{Q}_i, \mathcal{R}_i))_{i \in [L-1]}$ and a class of weight-sharing neural networks $\mathcal{F}_{\text{WSNN}} = \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\mathbf{m}})$ with

$$
L \le C_1 (\log m)^6 \log \log m, \quad \|\mathbf{d}\|_{\infty} \le C_1 m^{D+1} (\log m)^3,
$$

$$
s \le C_1 m (\log m)^5 \log \log m, \quad M \le \exp(C_1 {\log m})^6),
$$

$$
\|\mathbf{m}\|_{\infty} \le C_1 m^D
$$

such that

$$
\inf_{\mathbf{f}\in\mathcal{F}_{\text{WSNN}}\cap\mathcal{F}_{\infty}}\int_{\underline{T}}^{\overline{T}}\int_{\mathbb{R}^D}\|\mathbf{f}_{0}(\mathbf{x},t)-\mathbf{f}(\mathbf{x},t)\|_2^2p_t(\mathbf{x})\,\mathrm{d}\mathbf{x}\mathrm{d}t\leq C_1m^{-\frac{2\beta}{d}}(\log m)^{4(D+\beta\wedge 1)-3},
$$

where $\underline{T} = m^{-\tau_{\min}}, \overline{T} = \tau_{\max} \log m$ and

$$
\mathcal{F}_{\infty} = \bigg\{ \mathbf{f} : \|\mathbf{f}(\mathbf{x},t)\|_{\infty} \le \frac{C_1 \sqrt{\log m}}{\sigma_t} \quad \forall \mathbf{x} \in \mathbb{R}^D, t > 0 \bigg\}.
$$

Here, C_1 and C_2 are constants depending only on $(\beta, d, D, K, \tau_{\min}, \tau_{\max}, \tau_1, \tau_2, \overline{\tau}, \underline{\tau})$.

From the proof of Theorem [5.1,](#page-11-0) it can be observed that the class of permutation matrices $\mathcal{P}_{\mathbf{m}}$ can be chosen such that it depends only on m and β/d . Later, we select m based solely on β/d , meaning that the choice of \mathcal{P}_{m} ultimately depends only on β/d .

The proof of Theorem [5.1](#page-11-0) is provided in the Appendix. Here, we present an overview of the key ideas behind the proof. Note that $\nabla \log p_t(\mathbf{x}) = \nabla p_t(\mathbf{x})/p_t(\mathbf{x})$, and the division operation can be approximated by DNNs very efficiently, provided that the denominator is not too small. Since the ideas behind approximating the maps $(\mathbf{x}, t) \mapsto p_t(\mathbf{x})$ and $(\mathbf{x}, t) \mapsto \nabla p_t(\mathbf{x})$ are similar, we only present the key idea for approximating $(x, t) \mapsto p_t(x)$. For convenience, we use the informal notation $a \leq_{\text{log}} b$ to indicate that a is less than b up to a poly-logarithmic factor, such as $\log n$, $(\log m)^2$, or $\log \sigma_t^{-1}$. Similarly, we use the notation \approx_{log} to correspond to \approx .

For a given (sufficiently large) positive integer m , which roughly corresponds to the order of the number of nonzero network parameters, we will construct DNN approximators for the map $(\mathbf{x}, t) \mapsto$ $p_t(\mathbf{x})$ in four regions and combine them. These four regions for (\mathbf{x}, t) can be roughly defined as follows:

- (R1) (Outside of near-support) $\|\mathbf{x}\|_{\infty} \mu_t \gtrsim \sigma_t \sqrt{\mathbf{x}}$ $\overline{\log m}$
- (R2) (large t) $\|\mathbf{x}\|_{\infty} \mu_t \lesssim \sigma_t \sqrt{\log m}$ and $t \gtrsim m^{-(2-\delta)/D}$ for some $\delta > 0$
- (R3) (Boundary of near-support) $t \lesssim m^{-(2-\delta)/D}$ and $-\{\log(1/\sigma_t)\}^{-3/2} \lesssim ||\mathbf{x}||_{\infty} \mu_t \lesssim \sigma_t \sqrt{\frac{2\delta}{n}}$ $\overline{\log m}$
- (R4) (Interior of near-support) $t \lesssim m^{-(2-\delta)/D}$ and $\|\mathbf{x}\|_{\infty} \mu_t \lesssim -\{\log(1/\sigma_t)\}^{-3/2}$

Note that the maps $t \mapsto \mu_t$, $t \mapsto \sigma_t$, and $\mathbf{x} \mapsto \|\mathbf{x}\|_{\infty}$ can be approximated by DNNs very efficiently. Therefore, once we can approximate the map $(\mathbf{x}, t) \mapsto p_t(\mathbf{x})$ in each of the four regions, it is not difficult to combine them into a single function over the entire region.

In region (R1), p_t is nearly zero due to the sub-Gaussianity of p_t , making it easy to approximate. In region (R2), t is sufficiently large, and thus the map $\mathbf{x} \mapsto p_t(\mathbf{x})$ is much smoother than $\mathbf{x} \mapsto p_0(\mathbf{x})$. This smoother property enables the construction of a DNN with a moderate number of nonzero parameters, as in Lemma B.7 of [Oko et al.](#page-25-5) [\(2023\)](#page-25-5); see Proposition [B.3](#page-40-0) for details. Similarly, in region (R3), the map $\mathbf{x} \mapsto p_t(\mathbf{x})$ is very smooth due to the assumption (B), allowing us to construct a DNN with the desired approximation properties, similar to Lemmas B.2-B.5 of [Oko et al.](#page-25-5) [\(2023\)](#page-25-5); see Proposition [B.2](#page-40-1) for details.

The main challenge in the proof of Theorem [5.1](#page-11-0) lies in the approximation in region (R4). Note that $p_t(\mathbf{x}) \gtrsim 1$ in region (R4), and that

$$
p_t(\mathbf{x}) = \int_{\|\mathbf{z}\|_{\infty} \le 1} p_0(\mathbf{z}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{z}) d\mathbf{z}
$$

=
$$
\int_{\|\mathbf{x} + \sigma_t \mathbf{y}\|_{\infty} \le 1} \mu_t^{-D} p_0\left(\frac{\mathbf{x} + \sigma_t \mathbf{y}}{\mu_t}\right) \phi_1(\mathbf{y}) d\mathbf{y}.
$$
 (5.1)

To approximate the right-hand side of (5.1) , we first approximate it by a finite sum via a quadrature method, and then approximate the sum using a weight-sharing neural network.

To grasp the idea of approximation, it suffices to consider the approximation of a general function

$$
\mathbf{x} \mapsto \int_{[-1,1]^D} g(\mathbf{x}, \mathbf{y}) \mathrm{d}\mathbf{y},\tag{5.2}
$$

defined through a D-dimensional integral. Here, g is a function such that for each x , the map $y \mapsto g(x, y)$ belongs to $\mathcal{H}^{\beta,K}([-1, 1]^D)$, where $K > 0$ is a constant independent of x. We provide an idea for constructing a weight-sharing neural network to approximate the map [\(5.2\)](#page-13-0) with an error of $\epsilon \leq_{\log} m^{-\beta/d}$. It is well-known from numerical analysis [\(Novak,](#page-25-11) [1988\)](#page-25-11) that, to achieve an approximation error of ϵ for every function in $\mathcal{H}^{\beta,K}([-1,1]^D)$ using the Gauss–Legendre quadrature method, at least $O(\epsilon^{-D/\beta})$ quadrature points are necessary. Hence, [\(5.2\)](#page-13-0) can be approximated by a finite sum with $O(\epsilon^{-D/\beta})$ summands. However, to approximate this $O(\epsilon^{-D/\beta})$ -term summation using DNNs, we would need at least $O(\epsilon^{-D/\beta})$ network parameters (up to a poly-logarithmic factor), which results in a very large estimation error. To overcome this difficulty, instead of applying a single D-dimensional quadrature method, we apply a 1-dimensional m -point quadrature method D times to approximate the D-dimensional integral [\(5.2\)](#page-13-0). Specifically, let $(w_j)_{j\in[m]}$ and $(y_j)_{j\in[m]}$ be the m-point quadrature weights and nodes for 1-dimensional integrals over the interval $[-1, 1]$, that is,

$$
\left| \int_{-1}^{1} h(y) dy - \sum_{j=1}^{m} w_j h(y_j) \right| \lesssim m^{-\beta}, \quad \forall h \in \mathcal{H}^{\beta, K}([-1, 1]);
$$

see Lemma [B.1](#page-38-0) for details. Then, we can easily see that

$$
\left| \int_{[-1,1]^D} g(\mathbf{x}, \mathbf{y}) \mathrm{d} \mathbf{y} - \sum_{\mathbf{j} \in [m]^D} w_{\mathbf{j}} g(\mathbf{x}, \mathbf{y}_{\mathbf{j}}) \right| \lesssim m^{-\beta}
$$

where $w_j = \prod_{k=1}^D w_{j_k}$ and $\mathbf{y_j} = (y_{j_1}, \dots, y_{j_D})$. (We slightly abuse the notation for weights.) We next approximate the map

$$
\mathbf{x} \mapsto \sum_{\mathbf{j} \in [m]^{D}} w_{\mathbf{j}} g\left(\mathbf{x}, \mathbf{y}_{\mathbf{j}}\right) \tag{5.3}
$$

using weight-sharing neural networks. Although the summation in (5.3) consists of m^D terms and resembles a m^D -point, D-dimensional quadrature, it can be approximated by weight-sharing DNNs much more efficiently than a standard m^D -point, D-dimensional quadrature approximation. The key ingredients are the approximations of the following two maps:

$$
(w_1, \dots, w_m) \mapsto (w_j)_{j \in [m]^D} : \mathbb{R}^m \to \mathbb{R}^{m^D}
$$

$$
(\mathbf{x}, y_1, \dots, y_m) \mapsto (g(\mathbf{x}, \mathbf{y}_j))_{j \in [m]^D} : \mathbb{R}^{D+m} \to \mathbb{R}^{m^D}.
$$
 (5.4)

Although the w_j 's are distinct, each w_j is represented as a product of D terms from the m distinct values w_1, \ldots, w_m . Therefore, to approximate the first map of (5.4) , we only need to approximate the multiplication operation $(x, y) \mapsto xy$ and apply it multiple times. Note that multiplication can be approximated by DNNs very efficiently [\(Schmidt-Hieber,](#page-26-1) [2020\)](#page-26-1). With an additional trick, the repeated application of multiplication can be represented as a DNN of the form (3.1) , with a suitable choice of permutation matrices. Roughly speaking, to achieve an approximation error of $\epsilon \approx_{\log} m^{-\beta/d}$ for this

Figure 4: An illustration of why weight-sharing helps reduce model complexity: At some middle layers of the network, we need to approximate a map with inputs **x** and $y_j, j \in [m]^D$, and outputs $g_j = g(\mathbf{x}, \mathbf{y}_j), j \in [m]^D$. Since a single function g are approximated for m^D instances, leaving all network parameters as free parameters is inefficient. Weight-sharing can significantly reduce the number of distinct parameters. In this illustration, all edges with the same color share the same weight parameters.

map, we only need $O(\log m)$ distinct network parameters, which is the same as for approximating a single multiplication operation.

Similarly, weight sharing is crucial for approximating the second map in [\(5.4\)](#page-13-2). Since the function q are approximated for m^D instances, weight-sharing networks help reduce the number of distinct network parameters, see Figure [4](#page-14-0) for an illustration. The number of parameters required for a single evaluation of q with an approximation error of $m^{-\beta/d}$ depends on the structure of q. In our case, $O(m)$ parameters (up to a poly-logarithmic factor) are sufficient, due to the factorization property of p_0 .

By combining the results above, we can construct weight-sharing neural networks with $O(m)$ distinct parameters (up to a poly-logarithmic factor) to approximate [\(5.2\)](#page-13-0) with an error of $m^{-\beta/d}$.

Returning to the problem of approximating (5.1) , a key difference between (5.1) and (5.2) lies in the range of the integral, which depends on (x, t) . In particular, the diameter of the range also varies with t. As a result, we must use different quadrature weights and nodes for each pair (\mathbf{x}, t) . However, these quadrature weights and nodes can be expressed as (very) smooth functions of (\mathbf{x}, t) , making them easily approximated by deep neural networks. Full proofs, including additional technical details, are provided in the Appendix.

5.3 Convergence rate

The total variation distance between two Borel probability measures P and Q on \mathbb{R}^D is defined as

$$
d_{\mathrm{TV}}(P,Q) = \sup_A |P(A) - Q(A)|,
$$

where the supremum is taken over every Borel subset A of \mathbb{R}^D . We often denote $d_{\text{TV}}(P,Q)$ as $d_{\text{TV}}(p,q)$, where p and q are Lebesgue densities of P and Q, respectively. The following theorem provides the convergence rate of \hat{p} with respect to the total variation, which is our main result. Recall the definitions of the estimators $\hat{\mathbf{f}}$ and $\hat{\rho}$ from Section [2.](#page-3-0)

Theorem 5.2. Suppose that p_0 satisfies (S), (L), and (B). Let τ_{\min} and τ_{\max} be constants with

$$
\tau_{\min} \ge \frac{2\beta + d}{d} \left(\frac{4\beta}{d(\beta \wedge 1)} \vee \frac{1}{3D} \right) \quad \text{and} \quad \tau_{\max} \ge \frac{\beta}{\underline{\tau}(2\beta + d)} \vee \frac{2D(2\beta + d)}{d\overline{\tau}}.
$$

Let $\underline{T} = n^{-\tau_{\min}}$ and $\overline{T} = \tau_{\max} \log n$. Then, for every $n \geq C_4$, there exist a collection of permutation matrices $\mathcal{P}_{\mathbf{m}} = ((\mathcal{Q}_i, \mathcal{R}_i))_{i \in [L-1]}$ and a class of weight-sharing neural networks $\mathcal{F}_{\text{WSNN}} =$ $\mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\text{m}})$ with

$$
L \leq C_3 (\log n)^6 \log \log n, \quad ||\mathbf{d}||_{\infty} \leq C_3 n^{\frac{d(D+1)}{2\beta+d}} (\log n)^3,
$$

\n
$$
s \leq C_3 n^{\frac{d}{2\beta+d}} (\log n)^5 \log \log n, \quad M \leq \exp(C_3 {\log n}^6),
$$

\n
$$
\|\mathbf{m}\|_{\infty} \leq C_3 n^{\frac{dD}{2\beta+d}}
$$

satisfying

$$
\mathbb{E}\left[\int_{\underline{T}}^{\overline{T}} \int_{\mathbb{R}^D} \left\|\widehat{\mathbf{f}}(\mathbf{x},t) - \mathbf{f}_0(\mathbf{x},t)\right\|_2^2 p_t(\mathbf{x}) \mathrm{d}\mathbf{x} \mathrm{d}t\right] \le \epsilon_n^2
$$

and

$$
\mathbb{E}\left[d_{\text{TV}}\left(p_0, \hat{p}\right)\right] \leq \epsilon_n,
$$

where

$$
\epsilon_n = C_3 n^{-\frac{\beta}{2\beta + d}} \left\{ (\log n)^{2(D + \beta \wedge 1) - 3/2} + (\log n)^{10} \right\}
$$

and C_3, C_4 are constants depending only on $(\beta, d, D, K, \tau_{\min}, \tau_{\max}, \tau_1, \tau_2, \overline{\tau}, \underline{\tau})$.

Note that $\bar{\tau}$ and $\underline{\tau}$ can be treated as known constants. For example, if $\alpha_t = 1$ for all t, both constants can be set to 1. The constants $\tau_{\rm min}$ and $\tau_{\rm max}$ can also be chosen to depend solely on the single quantity β/d , ignoring their dependence on the known quantities $(D, \overline{\tau}, \underline{\tau})$. As discussed earlier, the class \mathcal{P}_{m} of permutation matrices can likewise be chosen to depend only on β/d . Similarly, if τ_1 , τ_2 , and K are treated as known constants, the hyperparameters $(L, \mathbf{d}, s, M, \mathcal{P}_m)$ defining the neural network $\mathcal{F}_{\text{WSNN}}$ can also be selected to depend solely on β/d . (Although this is not explicitly stated in Theorem [5.2,](#page-15-0) it can be deduced from the proof.) Therefore, the estimators \hat{f} and \hat{p} ultimately depend only on β/d .

In this sense, \hat{f} and \hat{p} are adaptive to the factorization structure because their construction does not rely on utilizing the structural information. We do not aim in this paper to construct a fully adaptive estimator, in the sense of estimators that do not depend on β/d . Although the architectures in Theorem [5.2,](#page-15-0) including the class \mathcal{P}_{m} of permutation matrices and the hyperparameters (L, d, s, M) , can be chosen to depend solely on β/d , in practice, much more complex architectures are often used, and hyperparameters are carefully tuned based on extensive experimental work.

Note that the convergence rate in Theorem [5.2](#page-15-0) is minimax-optimal up to a logarithm factor over the class of factorizable densities. Specifically, for $\beta, K > 0$ and $D, d \in \mathbb{N}$ with $d \leq D$, let

$$
\mathcal{G}(\beta, D, d, K) = \left\{ g_0 \in \mathcal{H}^{\beta, K}([-1, 1]^D) : g_0(\mathbf{x}) = \prod_{I \in \mathcal{I}} g_I(\mathbf{x}_I),
$$

$$
g_I \in \mathcal{H}^{\beta, K}([-1, 1]^{|I|}), \max_{I \in \mathcal{I}} |I| = d, \ \mathcal{I} \subseteq 2^{[D]}\right\}
$$

be the class of factorizable densities with smooth component functions. Then, we have

$$
\inf_{\widehat{p}} \sup_{p_0 \in \mathcal{G}(\beta, D, d, K)} \mathbb{E}[d_{\mathrm{TV}}(p_0, \widehat{p})] \gtrsim n^{-\beta/(2\beta + d)},
$$

where the infimum is taken over all estimators. The proof of this lower bound is straightforward, given the well-known result that the minimax rate for estimating a d-dimensional density in $\mathcal{H}^{\beta,K}([-1,1]^d)$ is $n^{-\beta/(2\beta+d)}$ (Giné and Nickl, [2016\)](#page-23-1).

The proof of Theorem [5.2](#page-15-0) is provided in the Appendix. Here, we present an overview of the key ideas behind the proof. Recall that $\hat{p} = \hat{p}_T$. By the triangle inequality, we have

$$
\mathbb{E}\left[d_{\text{TV}}(p_0,\hat{p})\right] \le d_{\text{TV}}(p_0,p_{\underline{T}}) + \mathbb{E}\left[d_{\text{TV}}(p_{\underline{T}},\hat{p}_{\underline{T}})\right].\tag{5.5}
$$

The first term in the right-hand side of (5.5) scales as a polynomial order in $\underline{T} = n^{-\tau_{\min}}$, thus we can control the error by choosing a large constant $\tau_{\rm min}$; see Lemma [C.1](#page-87-0) for details. The second term is the total variation distance between the distributions of $\mathbf{X}_{\underline{T}} = \mathbf{Y}_{\overline{T}-\underline{T}}$ and $\mathbf{X}_{\underline{T}} = \mathbf{Y}_{\overline{T}-\underline{T}}$. Note that the two processes $(\mathbf{Y}_t)_{t\in[0,\overline{T}]}$ and $(\mathbf{Y}_t)_{t\in[0,\overline{T}]}$ differ only in their initial distributions and drift functions. Hence, based on well-known results, we can bound the total variation distance by controlling each difference separately as follows:

$$
d_{\text{TV}}(p_{\overline{T}}, \widehat{p}_{\overline{T}})
$$
\n
$$
\leq d_{\text{TV}}(P_{\overline{T}}, \mathcal{N}(\mathbf{0}_D, \mathbb{I}_D)) + \int_{\overline{T}}^{\overline{T}} \int_{\mathbb{R}^D} \alpha_t^{-1} \left\| \widehat{\mathbf{f}}(\mathbf{x}, t) - \mathbf{f}_0(\mathbf{x}, t) \right\|_2^2 p_t(\mathbf{x}) \, \mathrm{d}\mathbf{x} \, \mathrm{d}t,\tag{5.6}
$$

see Remark 2.3 of [Bogachev et al.](#page-22-12) [\(2016\)](#page-22-12). Both terms on the right-hand side of [\(5.6\)](#page-16-1) correspond to the differences between the initial distributions and the drift functions, respectively. The first term can be easily controlled because $P_{\overline{T}}$ converges exponentially fast to $\mathcal{N}(\mathbf{0}_D, \mathbb{I}_D)$ as $\overline{T} = \tau_{\max} \log n$ increases. Thus, we can control the error by choosing a large constant τ_{max} .

The second term on the right-hand side of (5.6) represents the risk of the empirical risk minimizer \hat{f} . There is a substantial body of literature introducing techniques to bound the risk of empirical risk minimizers (e.g., [van der Vaart and Wellner,](#page-27-8) [1996,](#page-27-8) [Geer,](#page-23-17) [2000,](#page-23-17) [Wainwright,](#page-27-9) [2019\)](#page-27-9); see also Theorem C.4 of [Oko et al.](#page-25-5) [\(2023\)](#page-25-5) for the specific loss function considered in our paper. Technically, the risk can be decomposed into two terms: the approximation error and the estimation error, often referred to as the bias-variance decomposition. We have already bounded the approximation error in Section [5.2.](#page-11-1) To bound the estimation error, the key is to control the metric entropy of the weight-sharing networks $\mathcal{F}_{\text{WSNN}}$; see Lemma [C.2](#page-88-0) for details.

Remark 5.3. In (\mathbf{S}) , we assume that all factors g_I have the same smoothness level. This assumption can be relaxed, allowing each q_I to have a different level of smoothness. Specifically, suppose that for each $I \in \mathcal{I}$, we have $g_I \in \mathcal{H}^{\beta_I, K}([-1, 1]^{|I|})$ for some $\beta_I > 0$. Similarly to Theorem [5.1,](#page-11-0) an approximation error of ϵ for p_0 can be achieved using $O(\epsilon^{-d_*/\beta_*})$ network parameters (up to a polylogarithmic factor), where

$$
I_* = \underset{I \in \mathcal{I}}{\operatorname{argmin}} \frac{\beta_I}{|I|}, \quad \beta_* = \beta_{I_*}, \quad d_* = |I_*|.
$$

Based on this approximation result, one can show that

$$
\mathbb{E}\left[d_{\text{TV}}\left(p_0, \widehat{p}_{\underline{T}}\right)\right] \lesssim_{\log} n^{-\frac{\beta_*}{2\beta_* + d_*}}
$$

with a carefully chosen network architecture. The set-up of different smoothness levels includes the case of [Liu et al.](#page-25-12) [\(2007\)](#page-25-12), who considered a density of the form $p_0(\mathbf{x}) = g_I(\mathbf{x}_I)g_0(\mathbf{x})$, where g_0 is very smooth and $I \subseteq [D]$. Under the assumption that g_I has continuous second-order derivatives, they proposed a density estimator that achieves a convergence rate of $O(n^{-2/(4+|I|)+\epsilon})$ for any $\epsilon > 0$.

6 Sub-optimality of a vanilla score matching estimator

One of the main technical difficulties in our results in Section [5](#page-9-0) arises from the fact that p_t is no longer factorizable for $t > 0$. In practice, a key component of the success of score-based generative models and diffusion models lies in jointly modeling infinitely many score functions using deep neural networks via the map $(\mathbf{x}, t) \mapsto \mathbf{f}(\mathbf{x}, t)$ [\(Song and Ermon,](#page-26-13) [2019,](#page-26-13) [Song et al.,](#page-26-5) [2021\)](#page-26-5). Note that early works on the score estimation have focused on estimating the single score function $\mathbf{x} \mapsto \nabla \log p_0(\mathbf{x})$ via the score matching loss

$$
\widetilde{\ell}_{\mathbf{f}}(\mathbf{x}) = \mathrm{tr}(\nabla \mathbf{f}(\mathbf{x})) + \frac{1}{2} \left\| \mathbf{f}(\mathbf{x}) \right\|_2^2,
$$

which is based on the fact that

$$
\frac{1}{2}\mathbb{E}\left[\|\mathbf{f}(\mathbf{X}_0)-\nabla \log p_0(\mathbf{X}_0)\|_2^2\right] = \mathbb{E}\left[\mathrm{tr}\left(\nabla \mathbf{f}(\mathbf{X}_0)\right) + \frac{1}{2}\left\|\mathbf{f}(\mathbf{X}_0)\right\|_2^2\right] + C
$$

under mild assumptions, where C is a constant depending only on p_0 (Hyvärinen, [2005\)](#page-23-13).

For a given class F of score functions, let \hat{f}_{VS} be the corresponding empirical risk minimizer, that is,

$$
\widehat{\mathbf{f}}_{\text{VS}} \in \operatorname*{argmin}_{\mathbf{f} \in \mathcal{F}} \frac{1}{n} \sum_{i=1}^{n} \left[\operatorname{tr} \left(\nabla \mathbf{f}(\mathbf{X}^{i}) \right) + \frac{1}{2} \left\| \mathbf{f}(\mathbf{X}^{i}) \right\|_{2}^{2} \right].
$$

The corresponding density estimator can be defined via the Langevin diffusion. Specifically, let $(\mathbf{Z}_t)_{t>0}$ be the solution to the following Langevin equation:

$$
d\mathbf{Z}_t = -\nabla \log p_0(\mathbf{Z}_t) dt + \sqrt{2} dB_t, \quad \mathbf{Z}_0 \sim \mathcal{N}(\mathbf{0}_D, \mathbb{I}_D).
$$

Then, under mild assumptions, the distribution of \mathbf{Z}_t converges to p_0 as $t \to \infty$. The convergence speed can be exponentially fast under certain conditions on p_0 , such as when p_0 satisfies a Poincaré inequality or a log-Sobolev inequality [\(Bakry et al.,](#page-22-5) [2014\)](#page-22-5). Hence, one can define a density estimator \widehat{p}_{VS} as the limit distribution of the Langevin equation, with the true score function replaced by \widehat{f}_{VS} . We refer to $\hat{\mathbf{f}}_{\text{VS}}$ and \hat{p}_{VS} as vanilla score matching estimators for the score and density functions.

Note that vanilla score matching estimators are rarely used in modern large-scale generative problems. One reason is that the trace map $tr(\nabla f(\mathbf{x}))$ is computationally challenging to handle in high-dimensional (large D) problems, such as image generation tasks. Therefore, one may raise an important question: if the computation of \hat{f}_{VS} is tractable, would it perform well? More theoretically, one might ask whether \hat{f}_{VS} can achieve the optimal convergence rate.

Note that \hat{f}_{VS} is an empirical risk minimizer (or M-estimator); hence, one can use tools from standard empirical risk minimization theory to derive a convergence rate for \widehat{f}_{VS} . Although we do not provide specific proof in the present paper, one can expect the rate

$$
\mathbb{E}\left[\left\|\widehat{\mathbf{f}}_{\text{VS}}(\mathbf{X}_0) - \nabla \log p_0(\mathbf{X}_0)\right\|_2^2\right] \lesssim_{\log} n^{-\frac{\beta-1}{2(\beta-1)+d}} \tag{6.1}
$$

under assumptions similar to those in Theorem [5.2.](#page-15-0) While the rate in (6.1) is strictly slower than the rate in Theorem [5.2,](#page-15-0) it is minimax optimal for estimating the score function $\mathbf{x} \mapsto \nabla \log p_0(\mathbf{x})$ (or almost equivalently, the density derivative, under certain assumptions); see [Stone](#page-26-15) [\(1982\)](#page-26-15), [Singh](#page-26-16) [\(1977\)](#page-26-16), [Shen and Ghosal](#page-26-17) [\(2017\)](#page-26-17), [Yoo and Ghosal](#page-27-10) [\(2016\)](#page-27-10). The slower rate, compared to Theorem [5.2,](#page-15-0) arises from the fact that $\mathbf{x} \mapsto \nabla \log p_0(\mathbf{x})$ is only $(\beta - 1)$ -smooth, which is less smooth than the density $\mathbf{x} \mapsto p_0(\mathbf{x}).$

From the convergence rate of the score function estimator \hat{f}_{VS} , one can derive the same convergence rate with respect to the total variation for the corresponding density estimator \hat{p}_{VS} via [\(Girsanov,](#page-23-18) [1960,](#page-23-18) [Le Gall,](#page-24-14) [2016\)](#page-24-14); see also Remark 2.3 of [Bogachev et al.](#page-22-12) [\(2016\)](#page-22-12). While the rate $n^{-\frac{\beta-1}{2(\beta-1)+d}}$ is optimal for estimating the score function, the optimal rate for density estimation with respect to the total variation is strictly faster. Therefore, when vanilla score matching is viewed as an implicit density estimation method, its optimality would not be guaranteed.

7 Experimental analysis

Through a small simulation study, we empirically demonstrate in this section that the diffusion model effectively learns the true data distribution by comparing its performance to standard nonparametric procedures. While the empirical success of diffusion models is well-known in applications such as image generation, their performance in analyzing data from a smooth density is relatively less explored. Through simulations presented in this section, we show that diffusion models perform well in analyzing data from a smooth density. In particular, we anticipate that their performance is substantially better when the effective dimension d , derived from the factorization property, is small compared to the data dimension D. To evaluate this, we perform experiments using three types of simulated datasets and compare the results against conventional methods.

7.1 Dataset descriptions

Following the notation in the main results section, we denote the true density as p_0 , which can be expressed as $p_0(\mathbf{x}) = \prod_{I \in \mathcal{I}} g_I(\mathbf{x}_I)$. We analyze three types of true data distributions based on the value of the effective dimension d: (1) $d = 1$, (2) $d = 2$, and (3) $d = D$. To impose an image-like spatial structure, we organize the data into a grid with width and height of K, such that $D = K^2$. For each case, we consider various values of K , ranging from 3 to 15.

(Case 1) For the first case, we use the standard multivariate Gaussian density for p_0 , hence $d = 1$.

(Case 2) For the second case, we consider multivariate Gaussian densities with a non-identity covariance matrix, ensuring that the effective dimension is $d = 2$. To mimic the structure of conventional image data, as shown in Figure $3(b)$, we enforce a covariance structure where each element is conditionally dependent only on its four neighboring elements: those directly above, below, to the left, and to the right. This structure can be easily achieved by appropriately designing the corresponding precision matrix.

(Case 3) Finally, as for the third case, we consider finite mixtures of Gaussians of the form $p_0(\mathbf{x}) = \sum_{m=1}^M \pi_m \phi_1(\mathbf{x} - \boldsymbol{\mu}_m)$, where M is the number of mixture components, and π_m and $\boldsymbol{\mu}_m$ represent the mixing proportion and the mean vector of each component, respectively. For simplicity, we set $\pi_m = 1/M$ and generate μ_m from a standard multivariate Gaussian distribution. Since the

Figure 5: BPD values for Case 1. (From left to right: $K = 3, 5, 10, \text{ and } 15$.)

mixture density is not factorizable, we have $d = D$ in this case. We consider three values for M: 3, 5, and 8.

7.2 Learning algorithms and implementation details

For the diffusion model, we consider denoising diffusion probabilistic modeling (DDPM, [Ho et al.,](#page-23-5) [2020\)](#page-23-5), which corresponds to the standard OU process ($\alpha_t = 1$) and is one of the most widely used diffusion models.

DDPM is implemented using its public GitHub repository. The original architecture, with 37M parameters, is excessively large for the simulated datasets considered in this study. Therefore, we reduce its size to 1.7M parameters by modifying configuration options, such as the number of channels. We train DDPM using the Adam optimization algorithm [\(Kingma and Ba,](#page-24-15) [2015\)](#page-24-15) by 2,000 epochs, with a mini-batch size of 128 and a learning rate of 10−³ . We use Pytorch framework to run our algorithm using a single NVIDIA TITAN XP GPU.

As a baseline approach, we consider the kernel density estimation (KDE) method, using two types of kernel functions: Gaussian (KDE-G) and uniform (KDE-U) densities. Both KDE implementations are carried out using Scikit-learn module in Python.

7.3 Performance measure

We evaluate the performance of each method by calculating the negative log-likelihood normalized by the dimension, with a log base of 2, often referred to as bits per dimension (BPD) in image generation contexts. BPD normalizes the likelihood by the data dimensions, making it independent of the dimensionality of the data. It is important to note that evaluating the likelihood in diffusion models is not straightforward because we do not have an explicit density estimator. Since p_0 is known and easy to evaluate in the simulation, we use it to evaluate the performance of each method. Specifically, for the generated samples x_1^g $\mathbf{X}_1^g, \ldots, \mathbf{X}_{n'}^g$ of size n' , we use the following evaluation measure:

$$
BPD = -\frac{1}{D} \sum_{i=1}^{n'} \log_2 p_0(\mathbf{x}_i^g).
$$

For all cases considered, we set $n' = 3000$. In each experiment, we report the averaged BPD results over three repetitions with random data and random parameter initializations.

7.4 Performance results

For each case, we vary the size of the training dataset, n, from 100 to $20,000$ and compare the BPD of the trained DDPM with two KDE methods (KDE-G and KDE-U). The results for Case 1 and Case 2 are summarized in Figures [5](#page-19-0) and [6,](#page-20-1) respectively. In both cases, the performance of the diffusion model

Figure 6: BPD values for Case 2. (From left to right: $K = 3, 5, 10, \text{ and } 15$.)

Figure 7: BPD values for Case 3. (From left to right: $K = 3, 5, 10,$ and 15. From top to bottom: $M = 3, 5, \text{ and } 8.$

is substantially better than that of the two KDE methods for sufficiently large n , regardless of the data dimensions considered. It is worth noting that DDPM exhibits relatively unstable performance for small values of n , but as n increases, it effectively learns the data distribution. Furthermore, there appears to be a tendency for the discrepancy in BPD values between DDPM and the two KDE methods to grow as D increases.

The results of Case 3, with $M = 3, 5, 8$, are shown in Figure [7.](#page-20-2) Similar to Cases 1 and 2, the performance of the diffusion model is significantly better than that of the KDE methods for sufficiently large n . Interestingly, the margins in BPD values between DDPM and the other methods are relatively smaller compared to Cases 1 and 2. This phenomenon aligns with our theoretical results, which suggest that DDPM is particularly effective at learning data distributions with a small effective dimension, d. Although $d = D$ for Case 3, we hypothesize that the superior performance of the diffusion model might be attributed to its ability to adapt to other low-dimensional structures not covered in the existing theory.

8 Discussion

We have demonstrated that an estimator constructed from the diffusion model is adaptive to the factorization structure and achieves the minimax optimal convergence rates. In this section, we discuss some future directions related to our work.

Firstly, we believe that our analysis can be extended to high-dimensional settings where the data

dimension D diverges as the sample size tends to infinity. In this case, the convergence rate would depend on additional quantities such as D and $|\mathcal{I}|$. An important future task would be to characterize upper bounds for D , which might depend on the structure of p_0 , to guarantee statistical consistency. Although this generalization is a natural extension for statisticians, the techniques required, such as sharp approximation theory, present significant challenges.

Secondly, while we assumed that P_0 possesses a Lebesgue density, this assumption might be eliminated. For general probability distributions supported on the cube $[0, 1]^D$, the minimax optimal rate with respect to the Wasserstein distance is $n^{-1/D}$ for $D > 2$. If we restrict the class to distributions supported on a d -dimensional space (not necessarily a smooth manifold), the optimal rate improves to $n^{-1/d}$ [\(Weed and Bach,](#page-27-11) [2019\)](#page-27-11). It would be interesting to investigate whether an estimator constructed from the diffusion model achieves this optimal rate. Further structural assumptions could be considered through conditional independence in directed and undirected graphs, which might replace the factorization assumption (F) for densities.

Finally, recall that a key component in constructing an optimal estimator is the use of weightsharing neural networks to approximate functions defined through high-dimensional integrals of the form [\(5.2\)](#page-13-0). Recently, physics-informed neural networks (PINNs) have demonstrated remarkable success in modeling solutions to partial differential equations (PDEs) [\(Karniadakis et al.,](#page-24-16) [2021,](#page-24-16) [Raissi](#page-25-13) [et al.,](#page-25-13) [2019\)](#page-25-13). Notably, many solutions to PDEs, such as the heat equation and the Poisson equation, can be expressed as integrals of the form above [\(Courant and Hilbert,](#page-23-19) [2008\)](#page-23-19). This suggests that weight-sharing networks could serve as a promising architecture for theoretical analysis of PINNs.

References

- Anderson, B. D. (1982). Reverse-time diffusion equation models. Stochastic Processes and their Applications, 12(3):313–326.
- Arjovsky, M., Chintala, S., and Bottou, L. (2017). Wasserstein generative adversarial networks. In Proc. ICML, pages 214–223.
- Arnold, V. I. (2004). Lectures on Partial Differential Equations. Springer, Berlin.
- Azangulov, I., Deligiannidis, G., and Rousseau, J. (2024). Convergence of diffusion models under the manifold hypothesis in high-dimensions. ArXiv:2409.18804.
- Bakry, D., Gentil, I., and Ledoux, M. (2014). Analysis and Geometry of Markov Diffusion Operators. Springer, Cham.
- Bauer, B. and Kohler, M. (2019). On deep learning as a remedy for the curse of dimensionality in nonparametric regression. Ann. Statist., 47(4):2261–2285.
- Benton, J., Bortoli, V. D., Doucet, A., and Deligiannidis, G. (2024). Nearly d-linear convergence bounds for diffusion models via stochastic localization. In Proc. ICLR, pages 1–21.
- Bogachev, V., Röckner, M., and Shaposhnikov, S. (2011). On uniqueness problems related to the Fokker–Planck–Kolmogorov equation for measures. J. Math. Sci., 179(1):7–47.
- Bogachev, V. I., Krylov, N. V., Röckner, M., and Shaposhnikov, S. V. (2022). Fokker–Planck– Kolmogorov Equations. American Mathematical Society.
- Bogachev, V. I., Röckner, M., and Shaposhnikov, S. V. (2016). Distances between transition probabilities of diffusions and applications to nonlinear Fokker–Planck–Kolmogorov equations. J. Funct. Anal., 271(5):1262–1300.
- Bortoli, V. D. (2022). Convergence of denoising diffusion models under the manifold hypothesis. Transact. Mach. Learn. Res., pages 1–42.
- Bos, T. and Schmidt-Hieber, J. (2024). A supervised deep learning method for nonparametric density estimation. *Electron. J. Stat.*, 18(2):5601–5658.
- Bris, C. L. and Lions, P.-L. (2008). Existence and uniqueness of solutions to Fokker–Planck type equations with irregular coefficients. Commun. Partial. Differ. Equ., $33(7)$:1272–1317.
- Burden, R. L. and Faires, J. D. (2010). Numerical Analysis. Cengage Learning.
- Chae, M. (2022). Rates of convergence for nonparametric estimation of singular distributions using generative adversarial networks. ArXiv:2202.02890.
- Chae, M., Kim, D., Kim, Y., and Lin, L. (2023). A likelihood approach to nonparametric estimation of a singular distribution using deep generative models. J. Mach. Learn. Res., 24(77):1–42.
- Chen, S., Chewi, S., Lee, H., Li, Y., Lu, J., and Salim, A. (2023a). The probability flow ode is provably fast. In Proc. NeurIPS, pages 1–24.
- Chen, S., Chewi, S., Li, J., Li, Y., Salim, A., and Zhang, A. (2023b). Sampling is as easy as learning the score: theory for diffusion models with minimal data assumptions. In Proc. ICLR, pages 1–28.
- Courant, R. and Hilbert, D. (2008). Methods of Mathematical Physics, Volume 1. John Wiley & Sons.
- Dhariwal, P. and Nichol, A. (2021). Diffusion models beat gans on image synthesis. In Proc. NeurIPS, pages 8780–8794.
- Dinh, L., Krueger, D., and Bengio, Y. (2015). Nice: Non-linear independent components estimation. In Proc. ICLR, pages 1–13.
- Fang, Z. and Cheng, G. (2023). Optimal convergence rates of deep convolutional neural networks: Additive ridge functions. Transact. Mach. Learn. Res., pages 1–20.
- Geer, S. A. (2000). Empirical Processes in M-estimation. Cambridge University Press.
- Genovese, C. R., Perone-Pacifico, M., Verdinelli, I., and Wasserman, L. (2012). Minimax manifold estimation. J. Mach. Learn. Res., 13(1):1263–1291.
- Giné, E. and Nickl, R. (2016). Mathematical Foundations of Infinite-Dimensional Statistical Models. Cambridge University Press.
- Girsanov, I. V. (1960). On transforming a certain class of stochastic processes by absolutely continuous substitution of measures. Theory of Probability $\mathcal B$ Its Applications, 5(3):285–301.
- Goodfellow, I., Bengio, Y., and Courville, A. (2016). Deep Learning. MIT Press.
- Goodfellow, I., Pouget-Abadie, J., Mirza, M., Xu, B., Warde-Farley, D., Ozair, S., Courville, A., and Bengio, Y. (2014). Generative adversarial nets. In Proc. NIPS, pages 2672–2680.
- Györfi, L., Kontorovich, A., and Weiss, R. (2023). Tree density estimation. IEEE Trans. Inform. Theory, 69(2):1168–1176.
- Hammersley, J. M. and Clifford, P. (1971). Markov fields on finite graphs and lattices. Unpublished manuscript.
- Hastie, T., Tibshirani, R., and Friedman, J. H. (2009). The Elements of Statistical Learning: Data Mining, Inference, and Prediction. Springer, New York.
- Hastie, T., Tibshirani, R., and Wainwright, M. (2015). Statistical Learning with Sparsity. CRC Press.
- Hastie, T. J. and Tibshirani, R. J. (1990). Generalized Additive Models. Chapman and Hall/CRC.
- Haussmann, U. G. and Pardoux, E. (1986). Time reversal of diffusions. Ann. Probab., 14(4):1188–1205.
- Ho, J., Jain, A., and Abbeel, P. (2020). Denoising diffusion probabilistic models. In Proc. NeurIPS, pages 6840–6851.
- Ho, J., Salimans, T., Gritsenko, A., Chan, W., Norouzi, M., and Fleet, D. J. (2022). Video diffusion models. In Proc. NeurIPS, pages 8633–8646.
- Hyvärinen, A. (2005). Estimation of non-normalized statistical models by score matching. J. Mach. Learn. Res., 6(24):695–709.
- Imaizumi, M. and Fukumizu, K. (2022). Advantage of deep neural networks for estimating functions with singularity on hypersurfaces. J. Mach. Learn. Res., 23(111):1-54.
- Indritz, J. (1961). An inequality for hermite polynomials. *Proc. Amer. Math. Soc.*, 12(6):981–983.
- Jagtap, A. D., Shin, Y., Kawaguchi, K., and Karniadakis, G. E. (2022). Deep kronecker neural networks: A general framework for neural networks with adaptive activation functions. Neurocomputing, 468:165–180.
- Ji, Q. (2020). Probabilistic Graphical Models for Computer Vision. Academic Press.
- Karniadakis, G. E., Kevrekidis, I. G., Lu, L., Perdikaris, P., Wang, S., and Yang, L. (2021). Physicsinformed machine learning. Nature Reviews Physics, 3(6):422–440.
- Kingma, D. P. and Ba, J. (2015). Adam: A method for stochastic optimization. In Proc. ICLR, pages 1–15.
- Kingma, D. P. and Welling, M. (2014). Auto-encoding variational Bayes. In Proc. ICLR, pages 1–14.
- Kloeden, P. and Platen, E. (2011). Numerical Solution of Stochastic Differential Equations. Springer, Berlin.
- Kohler, M. and Langer, S. (2021). On the rate of convergence of fully connected deep neural network regression estimates. Ann. Statist., 49(4):2231–2249.
- Kong, Z., Ping, W., Huang, J., Zhao, K., and Catanzaro, B. (2021). Diffwave: A versatile diffusion model for audio synthesis. In Proc. ICLR, pages 1–17.
- Krizhevsky, A., Sutskever, I., and Hinton, G. E. (2012). Imagenet classification with deep convolutional neural networks. In Proc. NIPS, pages 1–9.
- Kwon, H. K. and Chae, M. (2024). Minimax optimal density estimation using a shallow generative model with a one-dimensional latent variable. In Proc. AISTATS, pages 469–477.
- Lauritzen, S. L. (1996). Graphical Models. Oxford University Press.
- Le Gall, J.-F. (2016). *Brownian Motion, Martingales, and Stochastic Calculus.* Springer, Cham.
- LeCun, Y., Boser, B., Denker, J., Henderson, D., Howard, R., Hubbard, W., and Jackel, L. (1989). Handwritten digit recognition with a back-propagation network. In Proc. NIPS, pages 396–404.
- LeCun, Y., Bottou, L., Bengio, Y., and Haffner, P. (1998). Gradient-based learning applied to document recognition. Proc. IEEE, 86(11):2278–2324.
- Li, G., Wei, Y., Chen, Y., and Chi, Y. (2024). Towards non-asymptotic convergence for diffusion-based generative models. In Proc. ICLR, pages 1–12.
- Liang, T. (2021). How well generative adversarial networks learn distributions. J. Mach. Learn. Res., $22(228):1-41.$
- Liu, H. and Lafferty, J. (2019). Nonparametric graphical models. In Maathuis, M., Drton, M., Lauritzen, S., and Wainwright, M., editors, Handbook of Graphical Models, pages 309–324. CRC Press.
- Liu, H., Lafferty, J., and Wasserman, L. (2007). Sparse nonparametric density estimation in high dimensions using the rodeo. In Proc. AISTATS, pages 283–290.
- Liu, H., Wasserman, L., and Lafferty, J. (2012). Exponential concentration for mutual information estimation with application to forests. In Proc. NIPS, pages 1–9.
- Liu, H., Xu, M., Gu, H., Gupta, A., Lafferty, J., and Wasserman, L. (2011). Forest density estimation. J. Mach. Learn. Res., 12(25):907–951.
- Mroueh, Y., Li, C.-L., Sercu, T., Raj, A., and Cheng, Y. (2018). Sobolev gan. In Proc. ICLR, pages 1–27.
- Nakada, R. and Imaizumi, M. (2020). Adaptive approximation and generalization of deep neural network with intrinsic dimensionality. J. Mach. Learn. Res., 21(174):1–38.
- Nakano, Y. (2024). Convergence of the denoising diffusion probabilistic models. $ArXiv:2406.01320$.
- Novak, E. (1988). Deterministic and Stochastic Error Bounds in Numerical Analysis. Springer, Berlin.
- Oko, K., Akiyama, S., and Suzuki, T. (2023). Diffusion models are minimax optimal distribution estimators. In Proc. ICML, pages 26517–26582.
- Oono, K. and Suzuki, T. (2019). Approximation and non-parametric estimation of resnet-type convolutional neural networks. In Proc. ICML, pages 4922–4931.
- Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z., Gimelshein, N., Antiga, L., Desmaison, A., Kopf, A., Yang, E., DeVito, Z., Raison, M., Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., Bai, J., and Chintala, S. (2019). PyTorch: An imperative style, highperformance deep learning library . In Proc. NeurIPS, pages 8024–8035.
- Pavliotis, G. A. (2014). Stochastic Processes and Applications: Diffusion Processes, the Fokker–Planck and Langevin Equations. Springer, New York.
- Petersen, P. and Voigtlaender, F. (2020). Equivalence of approximation by convolutional neural networks and fully-connected networks. Proc. Amer. Math. Soc., 148(4):1567–1581.
- Puchkin, N., Samsonov, S., Belomestny, D., Moulines, E., and Naumov, A. (2024). Rates of convergence for density estimation with generative adversarial networks. J. Mach. Learn. Res., 25(29):1– 47.
- Raissi, M., Perdikaris, P., and Karniadakis, G. E. (2019). Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. J. Comput. Phys., 378:686–707.
- Rezende, D. and Mohamed, S. (2015). Variational inference with normalizing flows. In Proc. ICML, pages 1530–1538.
- Rezende, D. J., Mohamed, S., and Wierstra, D. (2014). Stochastic backpropagation and approximate inference in deep generative models. In Proc. ICML, pages 1278–1286.
- Rombach, R., Blattmann, A., Lorenz, D., Esser, P., and Ommer, B. (2022). High-resolution image synthesis with latent diffusion models. In Proc. CVPR, pages 10684–10695.
- Ronneberger, O., Fischer, P., and Brox, T. (2015). U-net: Convolutional networks for biomedical image segmentation. In Proc. MICCAI, pages 234–241.
- Rumelhart, D. E., Hinton, G. E., and Williams, R. J. (1986). Learning representations by backpropagating errors. Nature, 323(6088):533–536.
- Schmidt-Hieber, J. (2020). Nonparametric regression using deep neural networks with ReLU activation function. Ann. Statist., 48(4):1875–1897.
- Shen, W. and Ghosal, S. (2017). Posterior contraction rates of density derivative estimation. Sankhya A, 79(2):336–354.
- Singh, R. S. (1977). Improvement on some known nonparametric uniformly consistent estimators of derivatives of a density. Ann. Statist., 5(2):394–399.
- Sohl-Dickstein, J., Weiss, E., Maheswaranathan, N., and Ganguli, S. (2015). Deep unsupervised learning using nonequilibrium thermodynamics. In Proc. ICML, pages 2256–2265.
- Song, Y. and Ermon, S. (2019). Generative modeling by estimating gradients of the data distribution. In Proc. NeurIPS, pages 1–13.
- Song, Y., Garg, S., Shi, J., and Ermon, S. (2020). Sliced score matching: A scalable approach to density and score estimation. In Proc. UAI, pages 574–584.
- Song, Y., Sohl-Dickstein, J., Kingma, D. P., Kumar, A., Ermon, S., and Poole, B. (2021). Score-based generative modeling through stochastic differential equations. In Proc. ICLR, pages 1–36.
- Stéphanovitch, A., Aamari, E., and Levrard, C. (2024). Wasserstein generative adversarial networks are minimax optimal distribution estimators. Ann. Statist., 52(5):2167–2193.
- Stone, C. J. (1982). Optimal global rates of convergence for nonparametric regression. Ann. Statist., 10(5):1040–1053.
- Sutskever, I., Vinyals, O., and Le, Q. V. (2014). Sequence to sequence learning with neural networks. In Proc. NIPS, pages 1–9.
- Tang, R. and Yang, Y. (2023). Minimax rate of distribution estimation on unknown submanifold under adversarial losses. Ann. Statist., 51(3):1282–1308.
- Tang, R. and Yang, Y. (2024). Adaptivity of diffusion models to manifold structures. In Proc. AISTATS, pages 1648–1656.
- Tsybakov, A. B. (2008). Introduction to Nonparametric Estimation. Springer, New York.
- Uppal, A., Singh, S., and Póczos, B. (2019). Nonparametric density estimation and convergence of GANs under Besov IPM losses. In Proc. NeurIPS, pages 9089–9100.
- van der Vaart, A. W. and Wellner, J. A. (1996). Weak Convergence and Empirical Processes. Springer, New York.
- Vandermeulen, R., Tai, W. M., and Aragam, B. (2024a). Breaking the curse of dimensionality in structured density estimation. To appear in Proc. NeurIPS.
- Vandermeulen, R., Tai, W. M., and Aragam, B. (2024b). Dimension-independent rates for structured neural density estimation. ArXiv:2411.15095.
- Vincent, P. (2011). A connection between score matching and denoising autoencoders. Neural Comput., 23(7):1661–1674.
- Wainwright, M. J. (2019). *High-Dimensional Statistics: A Non-Asymptotic Viewpoint.* Cambridge University Press.
- Weed, J. and Bach, F. (2019). Sharp asymptotic and finite-sample rates of convergence of empirical measures in Wasserstein distance. Bernoulli, 25(4A):2620–2648.
- Wibisono, A., Wu, Y., and Yang, K. Y. (2024). Optimal score estimation via empirical bayes smoothing. In Proc. COLT, pages 4958–4991.
- Yang, Y., Feng, H., and Zhou, D.-X. (2024). On the rates of convergence for learning with convolutional neural networks. ArXiv:2403.16459.
- Yoo, W. W. and Ghosal, S. (2016). Supremum norm posterior contraction and credible sets for nonparametric multivariate regression. Ann. Statist., 44(3):1069–1102.
- Yu, S., Drton, M., and Shojaie, A. (2022). Generalized score matching for general domains. *Informa*tion and Inference: A Journal of the IMA, 11(2):739–780.
- Zhang, A., Tay, Y., Zhang, S., Chan, A., Luu, A. T., Hui, S., and Fu, J. (2021). Beyond fully-connected layers with quaternions: Parameterization of hypercomplex multiplications with $1/n$ parameters. In Proc. ICLR, pages 1–13.
- Zhang, K., Yin, H., Liang, F., and Liu, J. (2024). Minimax optimality of score-based diffusion models: Beyond the density lower bound assumptions. In Proc. ICML, pages 60134–60178.

Appendix

A Auxililary Lemmas

This section provides auxiliary lemmas for proving the main theorems.

A.1 Several bounds regarding $p_t(\mathbf{x})$

In this subsection, we present several lemmas that bound $p_t(\mathbf{x})$, $\nabla \log p_t(\mathbf{x})$, and the derivatives of $p_t(\mathbf{x})$.

Lemma A.1 (Upper and lower bounds for $p_t(\mathbf{x})$). Let $K, \tau_1 > 0$ be given and suppose the true density p_0 satisfies that $\tau_1 \leq p_0(\mathbf{x}) \leq K$ for any $\mathbf{x} \in [-1,1]^D$. Then, there exists a constant $C_{S,1} =$ $C_{S,1}(D, K, \tau_1)$ such that

$$
C_{S,1}^{-1} \exp\left(-\frac{D\{(\|\mathbf{x}\|_{\infty} - \mu_t) \vee 0\}^2}{\sigma_t^2}\right) \le p_t(\mathbf{x}) \le C_{S,1} \exp\left(-\frac{\{(\|\mathbf{x}\|_{\infty} - \mu_t) \vee 0\}^2}{2\sigma_t^2}\right)
$$

for every $\mathbf{x} \in \mathbb{R}^D$ and $t \geq 0$.

Proof. This is a re-statement of Lemma A.2 in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5).

Lemma A.2 (Boundedness of score function). Let $K, \tau_1 > 0$ be given and suppose the true density p₀ satisfies that $\tau_1 \leq p_0(\mathbf{x}) \leq K$ for any $\mathbf{x} \in [-1,1]^D$. Then, there exists a positive constant $C_{S,2} =$ $C_{S,2}(D, K, \tau_1, \overline{\tau}, \underline{\tau})$ such that

$$
\|\nabla \log p_t(\mathbf{x})\|_2 \le \frac{C_{S,2}}{\sigma_t} \left(\frac{\|\mathbf{x}\|_{\infty} - \mu_t}{\sigma_t} \vee 1\right)
$$

for every $\mathbf{x} \in \mathbb{R}^D$ and $t \geq 0$.

Proof. This is a re-statement of Lemma A.3 in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5).

Lemma A.3 (Boundedness of derivatives). Let $K > 0$ be given and suppose the true density p_0 satisfies that $p_0(\mathbf{x}) \leq K$ for any $\mathbf{x} \in [-1,1]^D$. For any $\mathbf{k} \in \mathbb{Z}_{\geq 0}^D$, there exists a positive constant $C_{S,3} = C_{S,3}(D, K, \mathbf{k}, \overline{\tau}, \underline{\tau})$ such that

$$
\left|(\mathbf{D}^{\mathbf{k}} p_t)(\mathbf{x})\right| \le \frac{C_{S,3}}{\sigma_t^k}
$$

for every $\mathbf{x} \in \mathbb{R}^D$ and $t \geq 0$.

Proof. This is a re-statement of Lemma A.3 in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5), where $k = ||\mathbf{k}||_1$. \Box

A.2 Basic approximation results for neural networks

In this subsection, we present fundamental approximation results for using ReLU networks to approximate elementary functions. We also define the concatenation and parallelization in weight-sharing networks; see Lemma [A.8.](#page-30-0)

Lemma A.4 (Concatenation). Let $K \in \mathbb{N}, \{d_1, \ldots, d_{K+1}\} \subseteq \mathbb{N}$ be given. Consider $L^{(k)} \in \mathbb{N}_{\geq 2}$, $s^{(k)}$, $M^{(k)} > 0$ and $\mathbf{d}^{(k)} = (d_1^{(k)})$ $a_1^{(k)}, \ldots, a_{L^{(k)}}^{(k)})^{\top} \in \mathbb{N}^{L^{(k)}}$ with $d_1^{(k)} = d_k$ and $d_{L^{(k)}}^{(k)} = d_{k+1}$ for $k \in [K]$. For any neural networks f_1,\ldots,f_K with $f_k \in \mathcal{F}_{NN}(L^{(k)},\mathbf{d}^{(k)},s^{(k)},M^{(k)}), k \in [K]$, there exists a neural network $f \in \mathcal{F}_{NN}(L, d, s, M)$ with

$$
L = \sum_{k=1}^{K} L^{(k)}, \quad \|\mathbf{d}\|_{\infty} \le 2 \max_{k \in [K]} \|\mathbf{d}^{(k)}\|_{\infty}, \quad s \le 2 \sum_{k=1}^{K} s^{(k)}, \quad M = \max_{k \in [K]} M^{(k)}
$$

such that $f(\mathbf{x}) = (f_K \circ \cdots \circ f_1)(\mathbf{x})$ for any $\mathbf{x} \in \mathbb{R}^{d_1}$.

Proof. This is a re-statement of Remark 13 in [Nakada and Imaizumi](#page-25-14) [\(2020\)](#page-25-14).

Lemma A.5 (Parallelization). Let $K \in \mathbb{N}$ be given. Consider $L^{(k)} \in \mathbb{N}_{\geq 2}$, $s^{(k)}$, $M^{(k)} > 0$, $\mathbf{d}^{(k)} =$ $(d_1^{(k)}$ $\{a_1^{(k)}, \ldots, a_{L^{(k)}}^{(k)}\}^{\top} \in \mathbb{N}^{L^{(k)}}$ for $k \in [K]$. For any neural networks f_1, \ldots, f_K with

$$
f_k \in \mathcal{F}_{\text{NN}}(L^{(k)}, \mathbf{d}^{(k)}, s^{(k)}, M^{(k)}), \quad k \in [K],
$$

there exists a neural network $f \in \mathcal{F}_{NN}(L, \mathbf{d}, s, M)$ with

$$
L = \max_{k \in [K]} L^{(k)}, \quad ||\mathbf{d}||_{\infty} \le 2 \sum_{k=1}^{K} ||\mathbf{d}^{(k)}||_{\infty},
$$

$$
s \le 2 \sum_{k=1}^{K} \left(s^{(k)} + L d_{L^{(k)}}^{(k)} \right), \quad M \le \left(\max_{k \in [K]} M^{(k)} \right) \vee 1
$$

 \Box

such that

$$
f(\mathbf{x}) = \left(f_1(\mathbf{x}^{(1)}), \dots, f_K(\mathbf{x}^{(K)})\right) \in \mathbb{R}^{d_{L^{(1)}}^{(1)} + \dots + d_{L^{(K)}}^{(K)}}
$$

 $for \mathbf{x} = (\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(K)}) \in \mathbb{R}^{d_1^{(1)} + \dots + d_1^{(K)}}$. If $L^{(1)} = \dots = L^{(K)} = \tilde{L}$ with $\tilde{L} \in \mathbb{N}_{\geq 2}$, (L, \mathbf{d}, s, M) also satisfies

$$
L = \widetilde{L}, \quad \|\mathbf{d}\|_{\infty} \le \sum_{k=1}^{K} \|\mathbf{d}^{(k)}\|_{\infty}, \quad s \le \sum_{k=1}^{(K)} s^{(k)}, \quad M \le \max_{k \in [K]} M^{(k)}.
$$

Proof. This is a re-statement of Lemma F.3 in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5).

Lemma A.6 (Linear function). Let $W \in \mathbb{R}^{d_2 \times d_1}$, $\mathbf{b} \in \mathbb{R}^{d_2}$ be given with $d_1, d_2 \in \mathbb{N}$. There exists a neural networks $f_{lin} \in \mathcal{F}_{NN}(L, \mathbf{d}, s, M)$ with

$$
L = 2, \quad d = (d_1, 2d_2, d_2)T, \quad s = 2||W||_0 + 2||b||_0 + 2d_2, \quad M = \max\{||W||_{\infty}, ||b||_{\infty}, 1\}
$$

such that $f_{\text{lin}}(\mathbf{x}) = W\mathbf{x} + \mathbf{b}$ for any $\mathbf{x} \in \mathbb{R}^{d_1}$.

Proof. Note that $Wx + b = \rho(Wx + b) - \rho(-Wx - b)$ for any $x \in \mathbb{R}^{d_1}$. Let $f_{lin}(\cdot) = W_2 \rho(W_1 \cdot b_1) + b_2$ with

$$
W_1 = \left(W^\top, -W^\top\right)^\top \in \mathbb{R}^{2d_2 \times d_1}, \quad \mathbf{b}_1 = \left(\mathbf{b}^\top, -\mathbf{b}^\top\right)^\top \in \mathbb{R}^{2d_2},
$$

$$
W_2 = (\mathbb{I}_{d_2}, -\mathbb{I}_{d_2}) \in \mathbb{R}^{d_2 \times 2d_2}, \quad \mathbf{b}_2 = \mathbf{0}_{d_2} \in \mathbb{R}^{d_2},
$$

where \mathbb{I}_{d_2} denotes the $d_2 \times d_2$ identity matrix. Then, the assertion is followed by a simple calculation.

Lemma A.7 (Identity function). For any $L \geq \mathbb{N}_{\geq 2}$ and $m \in \mathbb{N}$, there exists a neural network $f_{\text{id}}^{(m,L)} \in \mathcal{F}_{\text{NN}}(L, \mathbf{d}, s, M)$ with

$$
\mathbf{d} = (m, 2m, ..., 2m, m)^{\top}, \quad s = 2mL, \quad M = 1
$$

such that $f_{\text{id}}^{(m,L)}(\mathbf{x}) = \mathbf{x}$ for any $\mathbf{x} \in \mathbb{R}^m$.

Proof. This is a re-statement of Lemma F.2 in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5).

The following lemma provides the concatenation and parallelization of two neural networks, where only one network has shared weight.

Lemma A.8 (Concatenation and parallelization of weight-sharing network). Consider the class of weight-sharing neural networks $\mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\text{m}})$ and neural networks $\mathcal{F}_{\text{NN}}(\widetilde{L}, \widetilde{\mathbf{d}}, \widetilde{s}, \widetilde{M})$. For any neural networks $f \in \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\mathbf{m}})$ and $\tilde{f} \in \mathcal{F}_{\text{NN}}(\tilde{L}, \tilde{\mathbf{d}}, \tilde{s}, \tilde{M})$, there exists a neural network $f_{\rm cc} \in \mathcal{F}_{\rm WSNN}(L_{\rm cc}, \mathbf{d}_{\rm cc}, s_{\rm cc}, M_{\rm cc}, \mathbf{m}_{\rm cc}, \mathcal{P}_{\rm cc})$ with

$$
L_{\rm cc} = L + \tilde{L}, \quad \|\mathbf{d}_{\rm cc}\|_{\infty} \le 2(\|\mathbf{d}\|_{\infty} \vee \|\mathbf{d}\|_{\infty}), \quad s_{\rm cc} \le 2s + 2\tilde{s}, \quad M_{\rm cc} = M \vee \tilde{M},
$$

 $\|\mathbf{m}_{cc}\|_{\infty} = \|\mathbf{m}\|_{\infty}$ and the set of permutation matrices \mathcal{P}_{cc} such that $f_{cc}(\mathbf{x}) = (\tilde{f} \circ f)(\mathbf{x})$ for any $\mathbf{x} \in \mathbb{R}^{d_1}$. Also, there exists a neural network $f_{\text{pr}} \in \mathcal{F}_{\text{NN}}(L_{\text{pr}}, \mathbf{d}_{\text{pr}}, s_{\text{pr}}, M_{\text{pr}}, \mathbf{m}_{\text{pr}}, \mathcal{P}_{\text{pr}})$ with

$$
L_{\text{pr}} = L \vee \bar{L}, \quad \|\mathbf{d}_{\text{pr}}\|_{\infty} \le 2\|\mathbf{d}\|_{\infty} + 2\|\mathbf{d}\|_{\infty},
$$

$$
s_{\text{pr}} \le 2s + 2\tilde{s} + 2(L \vee \tilde{L})(d_L + \tilde{d}_{\tilde{L}}), \quad M_{\text{pr}} = \max(M, \widetilde{M}, 1),
$$

 \Box

 \Box

 $\|\mathbf{m}_{\text{pr}}\|_{\infty} = \|\mathbf{m}\|_{\infty}$ and the set of permutation matrices \mathcal{P}_{pr} such that

$$
f_{\mathrm{pr}}(\mathbf{x}) = \left(f(\mathbf{x}_1), \widetilde{f}(\mathbf{x}_2)\right) \in \mathbb{R}^{d_L + \widetilde{d}_{\widetilde{L}}}
$$

for any $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2) \in \mathbb{R}^{d_1 + d_1}$.

Proof. The first assertion can be easily derived from Remark 13 of [Nakada and Imaizumi](#page-25-14) [\(2020\)](#page-25-14) with

$$
\mathbf{m}_{cc} = (\mathbf{m}, 1, \dots, 1) \in \mathbb{N}^{L_{cc}-1} \quad \text{and} \quad \mathcal{P}_{cc} = \mathcal{P} \cup \{Q_l, \mathcal{R}_l\}_{l \in \{L, \dots, L_{cc}-1\}},
$$

where Q_l and R_l are the set of $d_l \times d_l$ and $d_{l+1} \times d_{l+1}$ identity matrix, respectively.

For the second part, let $\{W_l, \mathbf{b}_l\}_{l \in [L]}$ and $\{W_l, \mathbf{b}_l\}_{l \in [\tilde{L}]}$ be the parameter matrices of f and f, respectively. If $L = L$, let $\mathbf{d}_{pr} = (d_1 + d_1, \ldots, d_{L+1} + d_{L+1})$ and $\mathbf{m}_{pr} = \mathbf{m}$. Also, for each $l \in [L]$, let $\mathcal{Q}_{\text{pr},l}$ and $\mathcal{R}_{\text{pr},l}$ be the set of permutation matrices of the form

$$
\begin{pmatrix} Q_l^{(j)} & \mathbf{0} \\ \mathbf{0} & \mathbb{I}_{\widetilde{d}_l} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} R_l^{(j)} & \mathbf{0} \\ \mathbf{0} & \mathbb{I}_{\widetilde{d}_{l+1}} \end{pmatrix}
$$

with $j \in [m_l]$, respectively. Then, the assertion follows with

$$
W_{\mathrm{pr},l} = \begin{pmatrix} W_l & \mathbf{0} \\ \mathbf{0} & m_l^{-1} \widetilde{W}_l \end{pmatrix} \quad \text{and} \quad \mathbf{b}_{\mathrm{pr},l} = \begin{pmatrix} \mathbf{b}_l \\ m_l^{-1} \widetilde{\mathbf{b}}_l \end{pmatrix}, \quad l \in [L-1],
$$

and $\mathcal{P}_{pr} = {\mathcal{Q}_{pr,l}, \mathcal{R}_{pr,l}}_{l \in [L-1]},$ where ${W_{pr,l}, \mathbf{b}_{pr,l}}_{l \in [L]}$ are the parameter matrices of f_{pr} . If $L = \widetilde{L} + 1$, consider a neural network **f** with L-layer and parameter matrices $\{\overline{W}_l, \mathbf{b}_l\}_{l \in [L]}$, where $\overline{W}_l = W_l, \mathbf{b}_l = \widetilde{\mathcal{L}}$ \mathbf{b}_l for $l \in [L-1]$, and

$$
\overline{W}_{\widetilde{L}} = \begin{pmatrix} \widetilde{W}_{\widetilde{L}} \\ -\widetilde{W}_{\widetilde{L}} \end{pmatrix}, \quad \overline{\mathbf{b}}_{\widetilde{L}} = \begin{pmatrix} \widetilde{\mathbf{b}}_{\widetilde{L}} \\ -\widetilde{\mathbf{b}}_{\widetilde{L}} \end{pmatrix}, \quad \overline{W}_{\widetilde{L}+1} = \begin{pmatrix} \mathbb{I}_{\widetilde{d}_{\widetilde{L}+1}} \\ -\mathbb{I}_{\widetilde{d}_{\widetilde{L}+1}} \end{pmatrix}, \quad \overline{\mathbf{b}}_{\widetilde{L}+1} = \mathbf{0}_{2 \widetilde{d}_{\widetilde{L}+1}}.
$$

We then apply the results for the case of parallelization between same layer network. If $L > \tilde{L} + 1$, consider a weight-sharing neural network $\bar{\mathbf{f}} = \mathbf{f}_{\text{id}}^{(d_{\tilde{L}+1}, L-L)} \circ \tilde{\mathbf{f}}$ with L-layer, where $\mathbf{f}_{\text{id}}^{(d_{\tilde{L}+1}, L-L)}$ is the neural network in Lemma [A.7.](#page-30-1) We then apply the results for the case of parallelization between same layer network.

If $L = \tilde{L} - 1$, consider a weight-sharing neural network \overline{f} with \tilde{L} -layer, parameter matrices $\{W_l, \mathbf{b}_l\}_{l \in [\tilde{L}]}$ with $W_l = W_l$, $\mathbf{b}_l = \mathbf{b}_l$ for $l \in [L-1]$, and

$$
\overline{W}_L = \begin{pmatrix} W_L \\ -W_L \end{pmatrix}, \quad \overline{\mathbf{b}}_L = \begin{pmatrix} \mathbf{b}_L \\ -\mathbf{b}_L \end{pmatrix}, \quad \overline{W}_{L+1} = \begin{pmatrix} \mathbb{I}_{d_{L+1}} \\ -\mathbb{I}_{d_{L+1}} \end{pmatrix}, \quad \overline{\mathbf{b}}_{L+1} = \mathbf{0}_{2d_{L+1}},
$$

 $m_L = 1$ and the set of permutation matrices $\mathcal{P} \cup \{Q_L, R_L\}$, where Q_L and R_L are the set of $d_L \times d_L$ and $2d_{L+1} \times 2d_{L+1}$ identity matrix, respectively. We then apply the results for the case of parallelization between same layer network. If $L < \widetilde{L} - 1$, consider a weight-sharing neural network $\overline{f} = f_{\text{id}}^{(d_{L+1}, L-L)} \circ f_{\text{id}}$ with \tilde{L} -layer and the set of permutation matrices $\mathcal{P} \cup \{\mathcal{Q}_l, \mathcal{R}_l\}_{L \leq l \leq \tilde{L}-1}$, where $\mathbf{f}_{\text{id}}^{(d_{L+1}, L-L)}$ is the neural network in Lemma [A.7](#page-30-1) and for each $l \in \{L, \ldots, \widetilde{L} - 1\}$, \mathcal{Q}_l and \mathcal{R}_l are the set of $d_l \times d_l$ and $d_{l+1} \times d_{l+1}$ identity matrix, respectively. We then apply the results for the case of parallelization between same layer network.

Lemma A.9 (Multiplication). Let $m \geq 2, C \geq 1, 0 < \tilde{\epsilon} \leq 1$ be given. For any $\epsilon > 0$, there exists a positive constant $C_{N,1}$ and a neural network $f_{\text{mult}} \in \mathcal{F}_{\text{NN}}(L, \mathbf{d}, s, M)$ with

$$
L \leq C_{N,1} \log m \{ \log(1/\epsilon) + m \log C \}, \quad \mathbf{d} = (m, 48m, \dots, 48m, 1)^\top,
$$

$$
s \leq C_{N,1} m \{ \log(1/\epsilon) + \log C \}, \quad M = C^m
$$

such that

$$
\left|f_{\text{mult}}(\widetilde{\mathbf{x}}) - \prod_{i=1}^{m} x_i\right| \le \epsilon + mC^{m-1}\widetilde{\epsilon}, \quad \forall \mathbf{x} \in [-C, C]^m, \widetilde{\mathbf{x}} \in \mathbb{R}^m \text{ with } \|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty} \le \widetilde{\epsilon},
$$

 $||f_{\text{mult}}||_{\infty} \leq C^m$ and $f_{\text{mult}}(\widetilde{\mathbf{x}}) = 0$ if $0 \in {\{\widetilde{x}_1, \ldots, \widetilde{x}_m\}}$.

Proof. This is a re-statement of Lemma F.6 in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5).

Lemma A.10 (Clipping function). Let $\underline{\mathbf{b}} = (\underline{b}_1, \ldots, \underline{b}_m), \overline{\mathbf{b}} = (\overline{b}_1, \ldots, \overline{b}_m) \in \mathbb{R}^m$ be given with $m \in \mathbb{N}$ and $\underline{b}_i \leq \overline{b}_i$ for all $i \in [m]$. Then, there exists a neural network

$$
f_{\text{clip}}^{(\underline{\mathbf{b}}, \overline{\mathbf{b}})} \in \mathcal{F}_{\text{NN}}(2, (m, 2m, m)^{\top}, 7m, \|\underline{\mathbf{b}}\|_{\infty} \vee \|\overline{\mathbf{b}}\|_{\infty})
$$

such that

$$
f_{\text{clip}}^{(\mathbf{b},\mathbf{b})}(\mathbf{x}) = (\bar{b}_1 \wedge \{x_1 \vee \underline{b}_1\}, \ldots, \bar{b}_m \wedge \{x_m \vee \underline{b}_m\}) \in \mathbb{R}^m
$$

for $\mathbf{x} = (x_1, \ldots, x_m) \in \mathbb{R}^m$.

Proof. This is a re-statement of Lemma F.4 in [Oko et al.](#page-25-5) [\(2023\)](#page-25-5).

Lemma A.11 (Logarithm function). For any $0 < \epsilon < 1/4$, there exists a positive constant $C_{N,2}$ and a neural network $f_{\text{log}} \in \mathcal{F}_{\text{NN}}(L, \mathbf{d}, s, M)$ with

$$
L \leq C_{N,2} \{ \log(1/\epsilon) \}^2 \log \log(1/\epsilon), \quad \|\mathbf{d}\|_{\infty} \leq C_{N,2} \{ \log(1/\epsilon) \}^3
$$

$$
s \leq C_{N,2} \{ \log(1/\epsilon) \}^5 \log \log(1/\epsilon), \quad M \leq \exp\left(8 \{ \log(1/\epsilon) \}^2\right)
$$

such that

$$
\left|\log x - f_{\log}(\widetilde{x})\right| \le \epsilon + \frac{|x - \widetilde{x}|}{\epsilon}
$$

for $x \in [\epsilon, 1/\epsilon]$ and $\widetilde{x} \in \mathbb{R}$.

Proof. Let $0 < \epsilon < 1/4$, $\delta = 1/8$ and $D_1 = \lfloor \frac{2 \log(1/\epsilon)}{\log(1+\delta)} - 1 \rfloor + 1$. Then, $[\epsilon, 1/\epsilon] \subseteq \bigcup_{i=1}^{D_1} [\underline{T}_i, \overline{T}_i]$, where $\underline{T}_i = (1+\delta)^{i-1}\epsilon$ and $\overline{T}_i = (1+\delta)^{i+1}\epsilon$ for $i \in [D_1]$. Let $D_2 = \lfloor \frac{\log(1/\epsilon)}{\log 2} \rfloor + 1$. For any $i \in [D_1]$ and $x \in [\underline{T}_i, \overline{T}_i]$, Taylor's theorem yields that

$$
\log x = P_i(x) + \frac{(-1)^{D_2 - 1}(x - \underline{T}_i)^{D_2}}{D_2\{\underline{T}_i + \xi(\overline{T}_i - \underline{T}_i)\}^{D_2}}
$$

for a suitable $\xi \in [0,1]$, where

$$
P_i(x) = \log \underline{T}_i + \frac{x - \underline{T}_i}{\underline{T}_i} + \sum_{k=2}^{D_2 - 1} \frac{(-1)^{k-1} (x - \underline{T}_i)^k}{k \underline{T}_i^k}.
$$

 \Box

Since $x - \underline{T}_i \leq \overline{T}_i - \underline{T}_i = \underline{T}_i(\delta^2 + 2\delta)$ and $\underline{T}_i \leq \underline{T}_i + \xi(\overline{T}_i - \underline{T}_i)$, it follows that

$$
|\log x - P_i(x)| \le \frac{1}{D_2} \left(\frac{\overline{T}_i - \underline{T}_i}{\underline{T}_i}\right)^{D_2} = \frac{\left(\delta^2 + 2\delta\right)^{D_2}}{D_2} \le \frac{2^{-D_2}}{D_2} \le \frac{\epsilon}{D_2}, \quad i \in [D_1]
$$
 (A.1)

for $x \in [\underline{T}_i, T_i]$. Let N_1 be a constant in Lemma [A.9.](#page-32-0) For $k \geq 2$, there exists a neural network $f_{\text{mult}}^{(k)} \in \mathcal{F}_{\text{NN}}(L_{\text{mult}}^{(k)}, \mathbf{d}_{\text{mult}}^{(k)}, s_{\text{mult}}^{(k)}, M_{\text{mult}}^{(k)})$ with

$$
L_{\text{mult}}^{(k)} \le N_1(k + D_2) \log k \{ \log(1/\epsilon) + \log D_2 \}, \quad \mathbf{d}_{\text{mult}}^{(k)} = (k, 48k, \dots, 48k, 1)^\top,
$$

\n
$$
s_{\text{mult}}^{(k)} \le N_1 k (k + D_2) \{ \log(1/\epsilon) + \log D_2 \}, \quad M_{\text{mult}}^{(k)} = \epsilon^{-k}
$$
\n(A.2)

such that $|f_{\text{mult}}^{(k)}(x_1,\ldots,x_k)-\prod_{i=1}^k x_i|\leq \epsilon^{D_2}/D_2$ for any $x_1,\ldots,x_k\in [-\epsilon^{-1},\epsilon^{-1}]$. For any $k\geq 1$ and $i \in [D_1]$, Lemma [A.6](#page-30-2) implies that there exists a neural network $f_{\text{lin}}^{(i,k)} \in \mathcal{F}_{\text{NN}}(2, (1, 2k, k)^\top, 6k, \underline{T}_i)$ such that $f_{\text{lin}}^{(i,k)}(x) = (x - \underline{T}_i, \dots, x - \underline{T}_i)^\top \in \mathbb{R}^k$ for any $x \in \mathbb{R}$. Combining Lemma [A.4](#page-29-1) with the last display, it follows that $f_{\text{pow}}^{(i,k)} = f_{\text{mult}}^{(k)} \circ f_{\text{lin}}^{(i,k)} \in \mathcal{F}_{\text{NN}}(L_{\text{pow}}^{(i,k)}, \mathbf{d}_{\text{pow}}^{(i,k)}, s_{\text{pow}}^{(i,k)}, M_{\text{pow}}^{(i,k)})$ for $i \in [D_1], k \geq 2$ with

$$
L_{\text{pow}}^{(i,k)} = L_{\text{mult}}^{(k)} + 2, \quad \|\mathbf{d}_{\text{pow}}^{(i,k)}\|_{\infty} \le 96k, \quad s_{\text{pow}}^{(i,k)} = 2s_{\text{mult}}^{(k)} + 12k, \quad M_{\text{pow}}^{(i,k)} = \underline{T}_i \vee M_{\text{mult}}^{(k)}
$$

and

$$
\left| f_{\text{pow}}^{(i,k)}(x) - (x - \underline{T}_i)^k \right| \le \frac{\epsilon^{D_2}}{D_2},
$$

for $x \in [\epsilon, \epsilon^{-1}]$. Consider functions $f_1, \ldots, f_{D_1} : \mathbb{R} \to \mathbb{R}$ such that

$$
f_i(\cdot) = \log \underline{T}_i + \frac{f_{\text{lin}}^{(i,1)}(\cdot)}{\underline{T}_i} + \sum_{k=2}^{D_2 - 1} \frac{(-1)^{k-1} f_{\text{pow}}^{(i,k)}(\cdot)}{k \underline{T}_i^k}, \quad i \in [D_1].
$$

Since $f_i - \log \underline{T}_i$ is a linear combination of $f_{\text{lin}}^{(i,1)}, f_{\text{pow}}^{(i,2)}, \dots, f_{\text{pow}}^{(i,D_2-1)}$ for $i \in [D_1]$, Lemma [A.4,](#page-29-1) Lemma [A.5](#page-29-2) and Lemma [A.6](#page-30-2) implies that $f_i \in \mathcal{F}_{NN}(L^{(i)}, \mathbf{d}^{(i)}, s^{(i)}, M^{(i)})$ with

$$
L^{(i)} \le L_{\text{pow}}^{(i, D_2 - 1)} + 2 \le D_3 D_2 \log D_2 \{ \log(1/\epsilon) + \log D_2 \},
$$

\n
$$
\|\mathbf{d}^{(i)}\|_{\infty} \le 2 \max \left(2 \sum_{k=2}^{D_2 - 1} \|\mathbf{d}_{\text{pow}}^{(i,k)}\|_{\infty} + 4, D_2 - 1 \right) \le D_3 D_2^2,
$$

\n
$$
s^{(i)} \le 2 \left\{ \sum_{k=2}^{D_2 - 1} \left(s_{\text{pow}}^{(i,k)} + L_{\text{pow}}^{(i, D_2 - 1)} + 2 \right) + L_{\text{pow}}^{(i, D_2 - 1)} + 8 \right\} + 2D_2 + 2
$$

\n
$$
\le D_3 D_2^3 \log D_2 \{ \log(1/\epsilon) + \log D_2 \},
$$

\n
$$
M^{(i)} \le \left(\max_{k \in [D_2 - 1]} M_{\text{pow}}^{(i,k)} \right) \vee \underline{T}_i^{-D_2 + 1} \le \epsilon^{-D_2}
$$

for a large enough constant $D_3 = D_3(N_1)$. Then,

$$
|P_i(x) - f_i(x)| \le \sum_{k=2}^{D_2 - 1} \frac{|f_{\text{pow}}^{(i,k)}(x) - (x - \underline{T}_i)^k|}{k \underline{T}_i^k}
$$

$$
\le \frac{D_2 - 2}{2\epsilon^{D_2 - 1}} \max_{2 \le k \le D_2 - 1} \left| f_{\text{pow}}^{(i,k)}(x) - (x - \underline{T}_i)^k \right| \le \frac{\epsilon}{2}, \quad i \in [D_1]
$$

for $x \in [\epsilon, \epsilon^{-1}]$, where the first inequality holds because $\underline{T}_i \geq \epsilon$. Combining $(A.1)$ with the last display, we have

$$
|\log x - f_i(x)| \le \left(\frac{1}{2} + \frac{1}{D_2}\right)\epsilon, \quad i \in [D_1]
$$
\n(A.4)

fo $x \in [\underline{T}_i, \overline{T}_i]$. Consider functions $f_{\text{swit}}^{(1)}, \ldots, f_{\text{swit}}^{(D_1)} : \mathbb{R} \to [0, 1]$ such that

$$
f_{\text{swit}}^{(1)}(\cdot) = \frac{1}{\overline{T}_1 - \underline{T}_2} \rho \left(-f_{\text{clip}}^{(\underline{T}_2, \overline{T}_1)}(\cdot) + \overline{T}_1 \right),
$$

\n
$$
f_{\text{swit}}^{(i)}(\cdot) = \frac{1}{\overline{T}_{i-1} - \underline{T}_i} \rho \left(f_{\text{clip}}^{(\underline{T}_i, \overline{T}_{i-1})}(\cdot) - \underline{T}_i \right) - \frac{1}{\overline{T}_i - \underline{T}_{i+1}} \rho \left(f_{\text{clip}}^{(\underline{T}_{i+1}, \overline{T}_i)}(\cdot) - \underline{T}_{i+1} \right),
$$

\n
$$
f_{\text{swit}}^{(D_1)}(\cdot) = \frac{1}{\overline{T}_{D_1 - 1} - \underline{T}_{D_1}} \rho \left(f_{\text{clip}}^{(\underline{T}_{D_1}, \overline{T}_{D_1 - 1})}(\cdot) - \underline{T}_{D_1} \right), \quad 2 \leq i \leq D_1 - 1,
$$

where $f_{\text{clip}}^{(\mathcal{I}_i, T_{i-1})} \in \mathcal{F}_{\text{NN}}(2, (1, 2, 1)^{\top}, 7, 8\epsilon^{-1})$ denotes the neural network in Lemma [A.10.](#page-32-1) Note that $\sum_{i=1}^{D_1} f_{s \text{wit}}^{(i)}(x) = 1$ for $x \in \mathbb{R}$ and $f_{s \text{wit}}^{(i)}(x) = 0$ for $x \in \bigcup_{j=1}^{D_1} [\underline{T}_j, \overline{T}_j] \setminus [\underline{T}_i, \overline{T}_i], i \in [D_1]$. Consider a function $f: \mathbb{R} \to \mathbb{R}$ such that $f(\cdot) = \sum_{i=1}^{D_1} f_{\text{mult}}^{(2)}(f_{\text{switch}}^{(i)}(\cdot), f_i(\cdot))$. Since $|f_{\text{mult}}^{(2)}(x_1, x_2) - x_1x_2| \le \epsilon^{D_2}/D_2$ for any $x_1, x_2 \in [-\epsilon^{-1}, \epsilon^{-1}]$, we have

$$
|\log x - f(x)| \leq \left| \log x - \sum_{i=1}^{D_1} f_{\text{switch}}^{(i)}(x) f_i(x) \right| + \frac{D_1 \epsilon^{D_2}}{D_2}
$$

=
$$
\left| \sum_{i=1}^{D_1} f_{\text{switch}}^{(i)}(x) \{ \log x - f_i(x) \} \right| + \frac{D_1 \epsilon^{D_2}}{D_2}
$$

$$
\leq \sum_{i=1}^{D_1} f_{\text{switch}}^{(i)}(x) |\log x - f_i(x)| + \frac{D_1 \epsilon^{D_2}}{D_2} \leq \left(\frac{1}{2} + \frac{1}{D_2} \right) \epsilon \sum_{i=1}^{D_1} f_{\text{switch}}^{(i)}(x) + \frac{D_1 \epsilon^{D_2}}{D_2}
$$

=
$$
\left(\frac{1}{2} + \frac{1 + D_1 \epsilon^{D_2 - 1}}{D_2} \right) \epsilon \leq \epsilon
$$

for $x \in \bigcup_{i=1}^{D_1} [\underline{T}_i, \overline{T}_i]$, where the second inequality holds by $(A.4)$. Combining $(A.2)$ and $(A.3)$ with Lemma [A.5](#page-29-2) and Lemma [A.4,](#page-29-1) we have $f_{\text{mult}}^{(2)}(f_{\text{switch}}^{(i)}(\cdot), f_i(\cdot)) \in \mathcal{F}_{\text{NN}}(\widetilde{L}^{(i)}, \widetilde{\mathbf{d}}^{(i)}, \widetilde{s}^{(i)}, \widetilde{M}^{(i)})$ for $i \in [D_1]$ with

$$
\widetilde{L}^{(i)} \le (L^{(i)} \vee 2) + L_{\text{mult}}^{(2)} \le D_4 D_2 \log D_2 \{ \log(1/\epsilon) + \log D_2 \}
$$

$$
\|\widetilde{\mathbf{d}}^{(i)}\|_{\infty} \le 2 \max \left(2 \|\mathbf{d}^{(i)}\|_{\infty} + 4, \|\mathbf{d}_{\text{mult}}^{(2)}\|_{\infty} \right) \le D_4 D_2^2
$$

$$
\widetilde{s}^{(i)} \le 4s^{(i)} + 4(L^{(i)} \vee 2) + 2s_{\text{mult}}^{(2)} + 28 \le D_4 D_2^3 \log D_2 \{ \log(1/\epsilon) + \log D_2 \}
$$

$$
\widetilde{M}^{(i)} \le \max \left(8\epsilon^{-1}, M^{(i)}, M_{\text{mult}}^{(2)}, 1 \right) \le \epsilon^{-D_2},
$$

where $D_4 = D_4(D_3)$ is a large enough constant. Let $f_{\text{clip}}^{(\epsilon, \epsilon^{-1})} \in \mathcal{F}_{\text{NN}}(2, (1, 2, 1)^{\top}, 7, \epsilon^{-1})$ be the nueral network in Lemma [A.10.](#page-32-1) Since f is a linear combination of $f_{\text{mult}}^{(2)}(f_{\text{switch}}^{(i)}(\cdot), f_i(\cdot))$ for each $i \in [D_1]$, Lemma [A.4,](#page-29-1) Lemma [A.5](#page-29-2) and Lemma [A.6](#page-30-2) implies that $f \circ f_{\text{clip}}^{(\epsilon, \epsilon^{-1})} \in \mathcal{F}_{\text{NN}}(L, \mathbf{d}, s, M)$ with

$$
L \leq \max_{i \in [D_1]} \widetilde{L}^{(i)} + 4 \leq D_5 \{ \log(1/\epsilon) \}^2 \log \log(1/\epsilon)
$$

$$
\|\mathbf{d}\|_{\infty} \leq 2 \max \left(2 \sum_{i=1}^{D_1} \|\widetilde{\mathbf{d}}^{(i)}\|_{\infty}, 4D_1 \right) \leq D_5 \{ \log(1/\epsilon) \}^3
$$

$$
s \leq 2 \sum_{i=1}^{D_1} \left(\widetilde{s}^{(i)} + \max_{j \in [D_1]} \widetilde{L}^{(j)} \right) + 4D_1 + 18 \leq D_5 \{ \log(1/\epsilon) \}^5 \log \log(1/\epsilon)
$$

$$
M \leq \max_{i \in [D_1]} \widetilde{M}^{(i)} \vee \epsilon^{-1} \leq \exp \left(8 \{ \log(1/\epsilon) \}^2 \right)
$$

for large enough constant $D_5 = D_5(D_4)$. Note that $|(f \circ f_{\text{clip}}^{(\epsilon,\epsilon^{-1})})(\tilde{x}) - \log x| \le |(f \circ f_{\text{clip}}^{(\epsilon,\epsilon^{-1})})(\tilde{x}) \log(\epsilon^{-1} \wedge {\{\tilde{x} \vee \epsilon\}})| + |\log(\epsilon^{-1} \wedge {\{\tilde{x} \vee \epsilon\}}) - \log x| \le \epsilon + \epsilon^{-1}|x - \tilde{x}| \text{ for any } x \in [\epsilon, \epsilon^{-1}] \text{ and } \tilde{x} \in \mathbb{R}.$ Then, the assertion follows by re-defining the constant.

Lemma A.12 (Negative exponential function). For any $0 < \epsilon < 2^{-4e+2}$, there exists a positive constant $C_{N,3}$ and a neural network $f_{\text{exp}} \in \mathcal{F}_{\text{NN}}(L, \mathbf{d}, s, M)$ with

$$
L \leq C_{N,3} \log(1/\epsilon) \log \log(1/\epsilon), \quad \|\mathbf{d}\|_{\infty} \leq C_{N,3} \{\log(1/\epsilon)\}^3
$$

$$
s \leq C_{N,3} \{\log(1/\epsilon)\}^4, \quad M \leq C_{N,3}\epsilon^{-1}
$$

such that

$$
\left|e^{-x} - f_{\exp}(\widetilde{x})\right| \le \epsilon + |x - \widetilde{x}|
$$

for any $x > 0$ and $\widetilde{x} \in \mathbb{R}$.

Proof. Let $0 < \epsilon < 2^{-4e+2}$, $D_1 = \lfloor \log(4/\epsilon) \rfloor + 1$, $D_2 = \lfloor \log(4/\epsilon) / \log 2 \rfloor + 1$ and $\underline{T}_i = i - 1$, $\overline{T}_i = i + 1$ for $i \in [D_1]$. Then, Taylor's theorem yields that for any $i \in [D_1]$ and $x \in [\underline{T}_i, T_i]$,

$$
e^{-x} = e^{-\underline{T}_i} e^{-(x-\underline{T}_i)} = e^{-\underline{T}_i} \left\{ P_i(x) + \frac{(-1)^{D_2} e^{-\xi(x-\underline{T}_i)} (x - \underline{T}_i)^{D_2}}{D_2!} \right\}
$$

for a suitable $\xi \in [0,1]$, where

$$
P_i(x) = 1 - (x - \underline{T}_i) + \sum_{k=2}^{D_2 - 1} \frac{(-1)^k (x - \underline{T}_i)^k}{k!}.
$$

Since $0 \leq x - \underline{T}_i \leq 2$ and $\underline{T}_i \geq 0$, it follows that

$$
\left|e^{-x} - e^{-\underline{T}_i} P_i(x)\right| \le \frac{2^{D_2}}{D_2!} \le \left(\frac{2e}{D_2}\right)^{D_2} \le \left(\frac{1}{2}\right)^{D_2} \le \frac{\epsilon}{4},\tag{A.5}
$$

where the second inequality holds because $k! \geq k^k e^{-k}$ for any $k \in \mathbb{N}$. Let N_1 be a constant in Lemma [A.9.](#page-32-0) For $k \geq 2$, there exists a neural network $f_{\text{mult}}^{(k)} \in \mathcal{F}_{\text{NN}}(L_{\text{mult}}^{(k)}, \mathbf{d}_{\text{mult}}^{(k)}, s_{\text{mult}}^{(k)}, M_{\text{mult}}^{(k)})$ with

$$
L_{\text{mult}}^{(k)} \le N_1 \log k \{ \log(4D_2/\epsilon^2) + k \log 2 \}, \quad \mathbf{d}_{\text{mult}}^{(k)} = (k, 48k, \dots, 48k, 1)^\top,
$$

\n
$$
s_{\text{mult}}^{(k)} \le N_1 k \{ \log(4D_2/\epsilon^2) + \log 2 \}, \quad M_{\text{mult}}^{(k)} = 2^k
$$
 (A.6)

such that $|f_{\text{mult}}^{(k)}(x_1,\ldots,x_k)-\prod_{i=1}^k x_i|\leq \epsilon^2/(4D_2)$ for any $x_1,\ldots,x_k\in[-2,2]$. For any $k\geq 1$ and $i \in [D_1]$, Lemma [A.6](#page-30-2) implies that there exists a neural network $f_{\text{lin}}^{(i,k)} \in \mathcal{F}_{\text{NN}}(2, (1, 2k, k)^\top, 6k, \underline{T}_i)$ such that $f_{\text{lin}}^{(i,k)}(x) = (x - \underline{T}_i, \dots, x - \underline{T}_i)^\top \in \mathbb{R}^k$ for any $x \in \mathbb{R}$. Combining Lemma [A.4](#page-29-1) with the last display, it follows that $f_{\text{pow}}^{(i,k)} = f_{\text{mult}}^{(k)} \circ f_{\text{lin}}^{(i,k)} \in \mathcal{F}_{\text{NN}}(L_{\text{pow}}^{(i,k)}, \mathbf{d}_{\text{pow}}^{(i,k)}, s_{\text{pow}}^{(i,k)}, M_{\text{pow}}^{(i,k)})$ for $i \in [D_1], k \geq 2$ with

$$
L_{\text{pow}}^{(i,k)} = L_{\text{mult}}^{(k)} + 2, \quad \|\mathbf{d}_{\text{pow}}^{(i,k)}\|_{\infty} \le 96k, \quad s_{\text{pow}}^{(i,k)} = 2s_{\text{mult}}^{(k)} + 12k, \quad M_{\text{pow}}^{(i,k)} = \underline{T}_i \vee M_{\text{mult}}^{(k)}
$$

and

$$
\left| f_{\text{pow}}^{(i,k)}(x) - (x - \underline{T}_i)^k \right| \le \frac{\epsilon^2}{4D_2}
$$

for $x \in [\underline{T}_i, \overline{T}_i]$. Consider functions $f_1, \ldots, f_{D_1} : \mathbb{R} \to \mathbb{R}$ such that

$$
f_i(\cdot) = 1 - f_{\text{lin}}^{(i,1)}(\cdot) + \sum_{k=2}^{D_2 - 1} \frac{(-1)^k f_{\text{pow}}^{(i,k)}(\cdot)}{k!}, \quad i \in [D_1].
$$
Since f_i-1 is a linear combination of $f_{\text{lin}}^{(i,1)}, f_{\text{pow}}^{(i,2)}, \ldots, f_{\text{pow}}^{(i,D_2-1)}$, Lemma [A.4,](#page-29-0) Lemma [A.5](#page-29-1) and Lemma [A.6](#page-30-0) implies that $f_i \in \mathcal{F}_{NN}(L^{(i)}, \mathbf{d}^{(i)}, s^{(i)}, M^{(i)})$ with

$$
L^{(i)} \le L_{\text{pow}}^{(i, D_2 - 1)} + 2 \le D_3 \log D_2 \{ \log(1/\epsilon) + D_2 \}
$$

\n
$$
\|\mathbf{d}^{(i)}\|_{\infty} \le 2 \max \left(2 \sum_{k=2}^{D_2 - 1} \|\mathbf{d}_{\text{pow}}^{(i,k)}\|_{\infty} + 4, D_2 - 1 \right) \le D_3 D_2^2
$$

\n
$$
s^{(i)} \le 2 \left\{ \sum_{k=2}^{D_2 - 1} \left(s_{\text{pow}}^{(i,k)} + L_{\text{pow}}^{(i, D_2 - 1)} + 2 \right) + L_{\text{pow}}^{(i, D_2 - 1)} + 8 \right\} + 2D_2 + 2
$$

\n
$$
\le D_3 D_2^2 \{ \log(1/\epsilon) + D_2 \}
$$

\n
$$
M^{(i)} \le \left(\max_{k \in [D_2 - 1]} M_{\text{pow}}^{(i,k)} \right) \vee 1 \le (D_1 + 1) \vee 2^{D_2 - 1}
$$
\n(A.7)

for a large enough constant $D_3 = D_3(N_1)$. Then,

$$
|P_i(x) - f_i(x)| \le \sum_{k=2}^{D_2 - 1} \frac{|f_{\text{pow}}^{(i,k)}(x) - (x - \underline{T}_i)^k|}{k!}
$$

$$
\le D_2 \max_{2 \le k \le D_2 - 1} \left| f_{\text{pow}}^{(i,k)}(x) - (x - \underline{T}_i)^k \right| \le \frac{\epsilon^2}{4}, \quad i \in [D_1]
$$

for $x \in [\underline{T}_i, \overline{T}_i]$. Combining $(A.5)$ with the last display, we have

$$
\left|e^{-x} - e^{-x} f_i(x)\right| \le \left|e^{-x} - e^{-x} f_i(x)\right| + |P_i(x) - f_i(x)| \le \frac{\epsilon}{4} + \frac{\epsilon^2}{4} \le \frac{\epsilon}{2}, \quad i \in [D_1]
$$
\n(A.8)

 $\text{for } x \in [\underline{T}_i, \overline{T}_i], \text{ where the first inequality holds because } e^{-\underline{T}_i} \leq 1. \text{ Consider functions } f_{\text{switch}}^{(1)}, \ldots, f_{\text{switch}}^{(D_1+1)}:$ $\mathbb{R} \to [0, 1]$ such that

$$
f_{\text{switch}}^{(1)}(\cdot) = \frac{1}{\overline{T}_1 - \underline{T}_2} \rho \left(-f_{\text{clip}}^{(\underline{T}_2, \overline{T}_1)}(\cdot) + \overline{T}_1 \right),
$$

\n
$$
f_{\text{switch}}^{(i)}(\cdot) = \frac{1}{\overline{T}_{i-1} - \underline{T}_i} \rho \left(f_{\text{clip}}^{(\underline{T}_i, \overline{T}_{i-1})}(\cdot) - \underline{T}_i \right) - \frac{1}{\overline{T}_i - \underline{T}_{i+1}} \rho \left(f_{\text{clip}}^{(\underline{T}_{i+1}, \overline{T}_i)}(\cdot) - \underline{T}_{i+1} \right),
$$

\n
$$
2 \leq i \leq D_1,
$$

\n
$$
f_{\text{switch}}^{(D_1+1)}(\cdot) = \frac{1}{\overline{T}_{D_1} - \underline{T}_{D_1+1}} \rho \left(f_{\text{clip}}^{(\underline{T}_{D_1+1}, \overline{T}_{D_1})}(\cdot) - \underline{T}_{D_1+1} \right),
$$

where $\underline{T}_{D_1+1} = D_1$ and $f_{\text{clip}}^{(\underline{T}_i, T_{i-1})} \in \mathcal{F}_{\text{NN}}(2, (1, 2, 1)^{\top}, 7, D_1)$ denotes the neural network in Lemma [A.10.](#page-32-0) Note that $\sum_{i=1}^{D_1+1} f_{\text{switch}}^{(i)}(x) = 1$ for $x \in \mathbb{R}$, and $f_{\text{switch}}^{(i)}(x) = 0$ for $x \in [0, \underline{T}_i] \cup [\overline{T}_i, \infty), i \in$ $[D_1]$ and $f_{\text{switch}}^{(D_1+1)}(x) = 0$ for $x \leq \underline{T}_{D_1+1}$. Consider a function $f : \mathbb{R} \to \mathbb{R}$ such that $f(\cdot) =$ $\sum_{i=1}^{D_1} e^{-T_i} f_{\text{mult}}^{(2)}(f_{\text{switch}}^{(i)}(\cdot), f_i(\cdot))$. Since $|f_{\text{mult}}^{(2)}(x_1, x_2) - x_1 x_2| \leq \epsilon^2/(4D_2)$ for any $x_1, x_2 \in [-2, 2]$, we have

$$
|e^{-x} - f(x)| \leq |e^{-x} - \sum_{i=1}^{D_1} e^{-\underline{T}_i} f_{\text{switch}}^{(i)}(x) f_i(x)| + \frac{D_1 \epsilon^2}{4D_2}
$$

\n
$$
= \left| \sum_{i=1}^{D_1} f_{\text{switch}}^{(i)}(x) \{ e^{-x} - e^{-\underline{T}_i} f_i(x) \} + f_{\text{switch}}^{(D_1+1)}(x) e^{-x} \right| + \frac{D_1 \epsilon^2}{4D_2},
$$

\n
$$
\leq \sum_{i=1}^{D_1} f_{\text{switch}}^{(i)}(x) |e^{-x} - e^{-\underline{T}_i} f_i(x)| + \frac{\epsilon}{4} + \frac{D_1 \epsilon^2}{4D_2}
$$

\n
$$
\leq \frac{\epsilon}{2} \sum_{i=1}^{D_1} f_{\text{switch}}^{(i)}(x) + \frac{\epsilon}{4} + \frac{D_1 \epsilon^2}{4D_2} \leq \left(\frac{3}{4} + \frac{(D_1 + 1)\epsilon}{4D_2} \right) \epsilon \leq \epsilon
$$

for $x \in [0,\infty)$, where the second inequality holds because $|f_{\text{switch}}^{(D_1+1)}(x)e^{-x}| \leq e^{-x} \leq \epsilon/4$ for $x \geq \underline{T}_{D_1+1}$ and the third inequality holds by $(A.8)$. Combining $(A.6)$ and $(A.7)$ with Lemma [A.5](#page-29-1) and Lemma [A.4,](#page-29-0) we have $f_{\text{mult}}^{(2)}(f_{\text{switch}}^{(i)}(\cdot), f_i(\cdot)) \in \mathcal{F}_{\text{NN}}(\widetilde{L}^{(i)}, \widetilde{\mathbf{d}}^{(i)}, \widetilde{s}^{(i)}, \widetilde{M}^{(i)})$ for $i \in [D_1]$ with

$$
\widetilde{L}^{(i)} \le (L^{(i)} \vee 2) + L_{\text{mult}}^{(2)} \le D_4 \log D_2 \{ \log(1/\epsilon) + \log D_2 \}
$$

$$
\|\widetilde{\mathbf{d}}^{(i)}\|_{\infty} \le 2 \max \left(2 \|\mathbf{d}^{(i)}\|_{\infty} + 4, \|\mathbf{d}_{\text{mult}}^{(2)}\|_{\infty} \right) \le D_4 D_2^2
$$

$$
\widetilde{s}^{(i)} \le 4s^{(i)} + 4(L^{(i)} \vee 2) + 2s_{\text{mult}}^{(2)} + 28 \le D_4 D_2^2 \{ \log(1/\epsilon) + D_2 \}
$$

$$
\widetilde{M}^{(i)} \le \max \left(D_1, M^{(i)}, M_{\text{mult}}^{(2)}, 1 \right),
$$

where $D_4 = D_4(D_3)$ is a large enough constant. Since f is a linear combination of $f_{\text{mult}}^{(2)}(f_{\text{switch}}^{(i)}(\cdot), f_i(\cdot))$ for each $i \in [D_1]$, Lemma [A.4,](#page-29-0) Lemma [A.5](#page-29-1) and Lemma [A.6](#page-30-0) implies that $f \circ \rho_1 \in \mathcal{F}_{NN}(L, \mathbf{d}, s, M)$ with

$$
L \leq \max_{i \in [D_1]} \widetilde{L}^{(i)} + 3 \leq D_5 \log(1/\epsilon) \log \log(1/\epsilon)
$$

$$
\|\mathbf{d}\|_{\infty} \leq 2 \max \left(2 \sum_{i=1}^{D_1} \|\widetilde{\mathbf{d}}^{(i)}\|_{\infty}, 4D_1 \right) \leq D_5 \{\log(1/\epsilon)\}^3
$$

$$
s \leq 2 \sum_{i=1}^{D_1} \left(\widetilde{s}^{(i)} + \max_{j \in [D_1]} \widetilde{L}^{(j)} \right) + 4D_1 + 4 \leq D_5 \{\log(1/\epsilon)\}^4
$$

$$
M \leq \max_{i \in [D_1]} \left(\widetilde{M}^{(i)} \vee e^{-\underline{T}_i} \right) \leq D_5 \epsilon^{-1}
$$

for large enough constant $D_5 = D_5(D_4)$. Note that $|(f \circ \rho_1)(\tilde{x}) - e^{-x}| \le |(f \circ \rho_1)(\tilde{x}) - e^{-(\tilde{x} \vee 0)}|$ $|e^{-(\tilde{x} \vee 0)} - e^{-x}| \leq \epsilon + |x - \tilde{x}|$ for any $x \geq 0$ and $\tilde{x} \in \mathbb{R}$. Then, the assertion follows by re-defining the constant.

Lemma A.13 (μ_t and σ_t). For any $0 < \epsilon < 1/2$, there exists a positive constant $C_{N,4} = C_{N,4}(\tau, \overline{\tau})$ and neural networks $f_{\mu} \in \mathcal{F}_{NN}(L_{\mu}, \mathbf{d}_{\mu}, s_{\mu}, M_{\mu}), f_{\sigma} \in \mathcal{F}_{NN}(L_{\sigma}, \mathbf{d}_{\sigma}, s_{\sigma}, M_{\sigma})$ with

$$
L_{\mu}, L_{\sigma} \leq C_{N,4} \{ \log(1/\epsilon) \}^2, \quad \|\mathbf{d}_{\mu}\|_{\infty}, \|\mathbf{d}_{\sigma}\|_{\infty} \leq C_{N,4} \{ \log(1/\epsilon) \}^2
$$

$$
s_{\mu}, s_{\sigma} \leq C_{N,4} \{ \log(1/\epsilon) \}^3, \quad M_{\mu}, M_{\sigma} \leq C_{N,4} \log(1/\epsilon)
$$

such that

$$
|\mu_{t_1} - f_\mu(t_1)| \le \epsilon \qquad \text{and} \qquad |\sigma_{t_2} - f_\sigma(t_2)| \le \epsilon
$$

for any $t_1 \geq 0$ and $t_2 \geq \epsilon$.

Proof. This is a re-statement of Lemma B.1 in [Oko et al.](#page-25-0) [\(2023\)](#page-25-0).

Lemma A.14 (Reciprocal function). For any $0 < \epsilon < 1$, there exists a positive constant $C_{N,5}$ and a neural network $f_{\text{rec}} \in \mathcal{F}_{\text{NN}}(L, \mathbf{d}, s, M)$ with

$$
L \le C_{N,5} \{ \log(1/\epsilon) \}^2, \quad \|\mathbf{d}\|_{\infty} \le C_{N,5} \{ \log(1/\epsilon) \}^3
$$

$$
s \le C_{N,5} \{ \log(1/\epsilon) \}^4, \quad M \le C_{N,5} \epsilon^{-2}
$$

such that

$$
\left|\frac{1}{x} - f_{\text{rec}}(\widetilde{x})\right| \le \epsilon + \frac{|x - \widetilde{x}|}{\epsilon^2}
$$

for any $x \in [\epsilon, 1/\epsilon]$ and $\widetilde{x} \in \mathbb{R}$.

Proof. This is a re-statement of Lemma F.7 in [Oko et al.](#page-25-0) [\(2023\)](#page-25-0).

38

 \Box

 \Box

B Proofs for the score function approximation

In this section, we provide the proof of Theorem [5.1.](#page-11-0) We begin by outlining the crucial lemmas and propositions.

For $n \in \mathbb{N}$, let P_n be the Legendre polynomial of degree n defined as

$$
P_n(x) = \left(\frac{1}{2^n n!}\right) \frac{d^n}{dx^n} (x^2 - 1)^n
$$

for $x \in \mathbb{R}$. It is well-known (page 114 of [Arnold](#page-22-0) [\(2004\)](#page-22-0)) that equation $P_n = 0$ has n distinct roots $\widetilde{x}_1^{(n)}$ $\widetilde{x}_1^{(n)}, \ldots, \widetilde{x}_n^{(n)}$ satisfying $-1 < \widetilde{x}_1^{(n)} < \cdots < \widetilde{x}_n^{(n)} < 1$. Let $\{\widetilde{w}_1^{(n)}\}$ $\{n_1^{(n)}, \ldots, \tilde{w}_n^{(n)}\}$ be the Gauss-Legendre quadrature weights, that is,

$$
\widetilde{w}_j^{(n)} = \begin{cases} \displaystyle\int_{-1}^1 \prod_{\substack{k=1 \\ k \neq j}}^n \left(\frac{x - \widetilde{x}_k^{(n)}}{\widetilde{x}_j^{(n)} - \widetilde{x}_k^{(n)}} \right) dx, & \text{if } n \ge 2\\ 2, & \text{if } n = 1. \end{cases}
$$

Let n_β be the largest integer strictly smaller than $\beta \vee 2$. For simplicity, we denote $(\tilde{x}_1^{(n_\beta)})$ $\widetilde{x}^{(n_\beta)}_{1},\ldots,\widetilde{x}^{(n_\beta)}_{n_\beta})$ and $(\widetilde{w}_1^{(n_\beta)}$ $(\alpha_{\beta}), \ldots, \widetilde{\mathcal{W}}_{n_{\beta}}^{(n_{\beta})})$ as $(\widetilde{x}_1, \ldots, \widetilde{x}_{n_{\beta}})$ and $(\widetilde{w}_1, \ldots, \widetilde{w}_{n_{\beta}})$, respectively. The following lemma provides an error bound for the m-points quadrature rule to approximate a one-dimensional integral.

Lemma B.1 (1-dimensional m-point quadrature rule). Let $A < B$ and $\beta, K > 0$ be given. For every $m \in n_\beta \mathbb{N}$, there exists $(w_i, x_i)_{i \in [m]}$ with $w_i > 0$ and $x_i \in (A, B)$ such that

$$
\left| \int_A^B g(x) dx - \sum_{i=1}^m w_i g(x_i) \right| \leq \left\{ \frac{n_\beta{}^\beta}{2^{\beta - \lfloor \beta \rfloor} \lfloor \beta \rfloor!} \right\} K(B-A)^\beta m^{-\beta},
$$

for every $g \in \mathcal{H}^{\beta, K}_1([A, B]).$ More specifically, one can choose

$$
w_i = \frac{(B-A)n_\beta}{2m}\widetilde{w}_{i-\lfloor i/n_\beta \rfloor},
$$

$$
x_i = A + \frac{(B-A)n_\beta}{2m} \left\{ \widetilde{x}_{i-\lfloor i/n_\beta \rfloor} + 2\lfloor i/n_\beta \rfloor + 1 \right\}.
$$

Let ϕ be the *one*-dimensional standard normal density. The following lemma provides a bound for the Hölder-norm of a function multiplied by ϕ and its derivative ϕ' .

Lemma B.2 (Preservation of Hölder continuity). Let $\beta, K > 0, A \subseteq \mathbb{R}$ be given and $g \in \mathcal{H}_1^{\beta, K}(A)$. Then, there exists a positive constant $C_{G,1} = C_{G,1}(\beta)$ such that $g\phi \in \mathcal{H}_1^{\beta, KC_{G,1}}(A)$ and $g\phi' \in$ $\mathcal{H}^{\beta,KC_{G,1}}_1$ $l_1^{p,n\cup G,1}(A).$

For $\mu, \sigma > 0$, define $p_{\mu,\sigma}(\cdot)$ as

$$
p_{\mu,\sigma}(\mathbf{x}) = \int_{\|\mathbf{y}\|_{\infty} \leq 1} p_0(\mathbf{y}) \phi_{\sigma}(\mathbf{x} - \mu \mathbf{y}) \mathrm{d}\mathbf{y} = \mu^{-D} \int_{\|\mathbf{x} + \sigma \mathbf{y}\|_{\infty} \leq \mu} p_0\left(\frac{\mathbf{x} + \sigma \mathbf{y}}{\mu}\right) \prod_{i=1}^D \phi(y_i) \mathrm{d}\mathbf{y}.
$$

Since p_0 is β -smooth under the (Smoothness) assumption, we can approximate the integral using the quadrature method with Lemma $B.1$ and Lemma $B.2$. Note however that \bf{y} in RHS ranges over a large set for small σ , and the error bound given in Lemma [B.1](#page-38-0) depends polynomially on the size of the interval. Since the tail of ϕ decays very quickly, one can control the numerical error as in the following lemma.

Lemma B.3 (Quadrature rule for $p_{\mu,\sigma}(\mathbf{x})$ and $\nabla p_{\mu,\sigma}(\mathbf{x})$). Let $\beta, K > 0$ be given and suppose that true density p_0 belongs to $\mathcal{H}^{\beta,K}([-1,1]^D)$. For τ_{bd} , τ_{tail} , $\mu, \sigma > 0$, $m \in n_\beta \mathbb{N}$, $i \in [D]$, $j \in [m]$ and $\mathbf{x} = (x_1, \dots, x_D)^\top \in \mathbb{R}^D$, let

$$
y_j^{(i)} = 2\sqrt{2\tau_{\text{tail}}} \{ \log(1/\sigma) \}^{\tau_{\text{bd}} + \frac{1}{2}} \left\{ -x_i - \mu + \frac{n_\beta \mu}{m} \left(\tilde{x}_{j - \lfloor j/n_\beta \rfloor} + 2 \lfloor j/n_\beta \rfloor + 1 \right) \right\},
$$

$$
w_j = 2\sqrt{2\tau_{\text{tail}}} n_\beta \tilde{w}_{j - \lfloor j/n_\beta \rfloor} \{ \log(1/\sigma) \}^{\tau_{\text{bd}} + \frac{1}{2}}.
$$

For $\mathbf{j} = (j_1, \ldots, j_D)^\top \in [m]^D$, let $\widetilde{\mathbf{y}}_{\mathbf{j}} = (y_{j_1}^{(1)})$ $y_{j_1}^{(1)}, \ldots, y_{j_D}^{(D)}$ $\binom{D}{j_D}^{\top} \in \mathbb{R}^D$. Then,

$$
\left\| \frac{\mathbf{x} + \sigma \tilde{\mathbf{y}}_{\mathbf{j}}}{\mu} \right\|_{\infty} \leq 1 - \frac{\{\log(1/\sigma)\}^{-\tau_{\text{bd}}}}{2}, \qquad \mathbf{j} \in [m]^D,
$$
\n
$$
\mu^D \left\| p_{\mu,\sigma}(\mathbf{x}) - \frac{1}{m^D} \sum_{\mathbf{j} \in [m]^D} \left\{ \prod_{i=1}^D w_{j_i} \phi(y_{j_i}^{(i)}) \right\} p_0 \left(\frac{\mathbf{x} + \sigma \tilde{\mathbf{y}}_{\mathbf{j}}}{\mu} \right) \right\| \leq \epsilon \qquad and
$$
\n
$$
\mu^D \left\| \sigma \nabla p_{\mu,\sigma}(\mathbf{x}) - \frac{1}{m^D} \sum_{\mathbf{j} \in [m]^D} \tilde{\mathbf{y}}_{\mathbf{j}} \left\{ \prod_{i=1}^D w_{j_i} \phi(y_{j_i}^{(i)}) \right\} p_0 \left(\frac{\mathbf{x} + \sigma \tilde{\mathbf{y}}_{\mathbf{j}}}{\mu} \right) \right\|_{\infty} \leq \epsilon
$$

 $\text{for every } ||\mathbf{x}||_{\infty} \leq \mu - \mu \{ \log(1/\sigma) \}^{-\tau_{\text{bd}}}, \mu \in [1/2, 1], \sigma \in (0, \widetilde{C}_2], \text{ where } \widetilde{C}_1 = \widetilde{C}_1(\beta, D, \tau_{\text{tail}}), \widetilde{C}_2 = \widetilde{C}_2(\beta, D, \tau_{\text{tail}}), \widetilde{C}_1 = \widetilde{C}_2(\beta, D, \tau_{\text{tail}}), \widetilde{C}_2 = \widetilde{C}_2(\beta, D, \tau_{\text{tail}}), \widetilde{C}_2 = \widetilde{C}_$ $\widetilde{C}_2(\beta, D, \tau_{\text{bd}}, \tau_{\text{tail}})$ and

$$
\epsilon = \widetilde{C}_1 K \left(\sigma^{\tau_{\rm tail}} + m^{-\beta} \{ \log(1/\sigma) \}^{(\tau_{\rm bd} + \frac{1}{2})\beta} \right) \{ \log(1/\sigma) \}^{(\tau_{\rm bd} + \frac{1}{2})(D-1)}.
$$

We can approximate the maps $(\mathbf{x}, t) \mapsto p_t(\mathbf{x})$ and $(\mathbf{x}, t) \mapsto \nabla p_t(\mathbf{x})$ using deep ReLU networks by replacing (μ, σ) in Lemma [B.3](#page-39-0) with (μ_t, σ_t) . As discussed in Section [5.2,](#page-11-1) a weight-sharing network is used to reduce the number of distinct network parameters. The approximation result is provided in the following proposition.

Proposition B.1 (Approximation at the interior of near-support). Suppose the true density p_0 verifies the assumption (S) and

$$
\tau_{\text{bd}} \in 1/2 + \mathbb{N}, \quad \tau_{\text{tail}} > 0, \quad \tau_{\text{min}} \ge \frac{4\beta}{d(\beta \wedge 1)}.
$$

Then, for every $m \geq C_5$, there exists a class of permutation matrices $\mathcal{P} = \{Q_i, \mathcal{R}_i\}_{i \in [L-1]}$ and weightsharing network $f \in \mathcal{F}_{WSNN}(L, d, s, M, P_m)$ with

$$
L \leq \widetilde{C}_3(\log m)^2 \log \log m, \quad \|\mathbf{d}\|_{\infty} \leq \widetilde{C}_3 m (\log m)^3,
$$

$$
s \leq \widetilde{C}_3 m (\log m)^5 \log \log m, \quad M \leq \exp \left(\widetilde{C}_3 {\log m}^2\right),
$$

$$
\|\mathbf{m}\|_{\infty} \leq \widetilde{C}_3 m^D
$$

satisfying

$$
\left\| \begin{pmatrix} \sigma_t \nabla p_t(\mathbf{x}) \\ p_t(\mathbf{x}) \end{pmatrix} - \mathbf{f}(\mathbf{x}, t) \right\|_{\infty} \le \widetilde{C}_4 \left(\log m \right)^{(\tau_{\rm bd} + \frac{1}{2})(D-1)} \left\{ t^{\frac{\tau_{\rm tail}}{2}} + m^{-\frac{\beta}{d}} \left(\log m \right)^{(\tau_{\rm bd} + \frac{1}{2})(\beta \vee 1)} \right\}
$$

for every $\mathbf{x} \in \mathbb{R}^D$ with $\|\mathbf{x}\|_{\infty} \leq \mu_t - \mu_t \{\log(1/\sigma_t)\}^{-\tau_{\text{bd}}}$ and $m^{-\tau_{\min}} \leq t \leq \overline{\tau}^{-1}(\widetilde{C}_2^2 \wedge 1/2).$

Here, $\widetilde{C}_2 = \widetilde{C}_2(\beta, D, \tau_{\text{bd}}, \tau_{\text{tail}})$ be the constant in Lemma [B.3,](#page-39-0) $\widetilde{C}_3 = \widetilde{C}_3(\beta, d, D, K, \overline{\tau}, \underline{\tau}, \tau_{\text{bd}}, \tau_{\text{tail}}, \tau_{\text{min}})$, $\widetilde{C}_4=\widetilde{C}_4(\beta,d,D,K,\underline{\tau},\tau_\mathrm{bd},\tau_\mathrm{tail},\tau_\mathrm{min}),\ and\ \widetilde{C}_5=\widetilde{C}_5(\beta,d,\underline{\tau},\tau_\mathrm{bd},\tau_\mathrm{tail},\tau_\mathrm{min}).$

As $t \to 0$, p_t is not lower bounded near the boundary of the support of p_0 due to the lower bound condition, making the approximation of $\nabla \log p_t$ challenging. With the assumption (\mathbf{B}) , p_0 is infinitely smooth so one can approximate p_0 efficiently with local polynomials by applying Taylor's theorem in the low-density region. Since a Gaussian density can also be efficiently approximated with local polynomials, one can calculate the integral in p_t closed form, and approximate the output with vanilla feedforward neural networks. The following proposition provides the approximation result, and our main proof strategy follows the proofs of Lemma B.2 - Lemma B.5 from [Oko et al.](#page-25-0) [\(2023\)](#page-25-0), with modifications for simplification.

Proposition B.2 (Approximation at the boundary of near-support). Let $K, \tau_{\text{bd}}, \tau_{\text{x}} > 0$, $0 < \tau_{\text{t}} < 1$, $0<\widetilde{\tau}_{\rm bd}<\tau_{\rm bd}$ be given and suppose the true density p_0 satisfies that $\|p_0\|_\infty\le K$. Then, for $0<\delta\le\widetilde{C}_8$ and p_0 satisfying

$$
\sup_{\alpha \in \mathbb{N}^D} \sup_{1 - \{\log(1/\delta)\}^{-\tilde{\tau}_{\text{bd}}} \le ||\mathbf{x}||_{\infty} \le 1} |(\mathbf{D}^{\alpha} p_0)(\mathbf{x})| \le K,
$$

there exists a network $f \in \mathcal{F}_{NN}(L, d, s, M)$ with

$$
L \leq \widetilde{C}_6 \{ \log(1/\delta) \}^4, \quad \|\mathbf{d}\|_{\infty} \leq \widetilde{C}_6 \{ \log(1/\delta) \}^{7+D\widetilde{\tau}_{\text{bd}}+D},
$$

$$
s \leq \widetilde{C}_6 \{ \log(1/\delta) \}^{11+D\widetilde{\tau}_{\text{bd}}+D}, \quad M \leq \exp\left(\widetilde{C}_6 \{ \log(1/\delta) \}^2\right),
$$

satisfying

$$
\left\| \begin{pmatrix} \sigma_t \nabla p_t(\mathbf{x}) \\ p_t(\mathbf{x}) \end{pmatrix} - \mathbf{f}(\mathbf{x}, t) \right\|_{\infty} \le \widetilde{C}_7 \delta \{\log(1/\delta)\}^D
$$

for every $\mathbf{x} \in \mathbb{R}^D$ with $\mu_t - \tau_{\mathbf{x}} \{ \log(1/\sigma_t) \}^{-\tau_{\text{bd}}} \leq ||\mathbf{x}||_{\infty} \leq \mu_t + \tau_{\mathbf{x}} \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \delta^{\tau_t}$.

Here, $\widetilde{C}_6 = \widetilde{C}_6(D, K, \overline{\tau}, \underline{\tau}, \tau_x), \ \widetilde{C}_7 = \widetilde{C}_7(D, K, \underline{\tau}), \ \widetilde{C}_8 = \widetilde{C}_8(D, \overline{\tau}, \tau_{\text{bd}}, \tau_x, \tau_t, \widetilde{\tau}_{\text{bd}})$ are positive constants.

For $t_* \geq 0$ and $t > 0$, we have

$$
p_{t_*+t}(\mathbf{x}) = \int_{\mathbb{R}^D} p_{t_*}(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}\mathbf{y}, \quad \mathbf{x} \in \mathbb{R}^D
$$

due to the Markov property of the process $(X_t)_{t\geq 0}$. Note that the map $x \mapsto p_{t_*}(x)$ is infinitely differentiable and its norm is bounded as $\|\mathcal{D}^{\mathbf{k}}p_{t*}(\cdot)\|_{\infty} \lesssim t_*^{-k/2}$ for any $\mathbf{k} \in \mathbb{N}^D$; see Lemma [A.3\)](#page-29-2). Then, one can approximate p_{t*} with a local Taylor expansion, yielding an error $O(m^{-k/D}t_*^{-k/2})$ using grid points bounded by $O(m)$, both up to a poly-logarithmic factor. Similar to the proof of Proposition [B.2,](#page-40-0) one can approximate the map $(\mathbf{x}, t) \mapsto p_{t_*+t}(\mathbf{x})$ with vanilla feedforward neural networks. The following proposition provides the approximation result, and our main proof strategy follows the proof of Lemma B.7 from [Oko et al.](#page-25-0) [\(2023\)](#page-25-0), with modifications for simplification.

Proposition B.3 (Approximation for large t). Let $K, \tau_1, \tau_x > 0, \tau_{sm} \in \mathbb{N}, \tau_{low} \in (0,1)$ be given and suppose the true density p_0 satisfies that $\tau_1 \leq p_0(\mathbf{x}) \leq K$ for any $\mathbf{x} \in [-1,1]^D$. Then, for $m \geq \widetilde{C}_{11}$, there exists a neural network $f \in \mathcal{F}_{NN}(L, d, s, M)$ with

$$
L \leq \widetilde{C}_9(\log m)^4, \quad \|\mathbf{d}\|_{\infty} \leq \widetilde{C}_9 m (\log m)^9,
$$

$$
s \leq \widetilde{C}_9 m (\log m)^{11}, \quad M \leq \exp(\widetilde{C}_9(\log m)^2)
$$

such that

$$
\left\| \begin{pmatrix} \sigma_t \nabla p_{t_*+t}(\mathbf{x}) \\ p_{t_*+t}(\mathbf{x}) \end{pmatrix} - \mathbf{f}(\mathbf{x},t) \right\|_{\infty} \le \widetilde{C}_{10} m^{-\frac{\tau_{\text{low}} \tau_{\text{sm}} - (D+1-\tau_{\text{low}})D}{D(1+D)}} (\log m)^{D(\frac{\tau_{\text{sm}}}{2}+1)}
$$

for every $\mathbf{x} \in \mathbb{R}^D$ with $\|\mathbf{x}\|_{\infty} \leq \mu_t + \tau_{\mathbf{x}} \sigma_t \sqrt{\log(1/\delta)}$, $\delta \leq t \leq \overline{\tau}^{-1} \log(1/\delta)$, where

$$
t_* = m^{-\frac{2-2\tau_{\text{low}}}{D}}
$$
 and $\delta = m^{-\frac{\tau_{\text{low}}\tau_{\text{sm}}+D+1-\tau_{\text{low}}}{D(1+D)}}$.

Here, $\widetilde{C}_9, \widetilde{C}_{10}, \widetilde{C}_{11}$ are positive constants depending on $(D, K, \overline{\tau}, \underline{\tau}, \tau_x, \tau_{\rm sm}, \tau_{\rm low})$.

B.1 Proofs of Lemma [B.1](#page-38-0) to [B.3](#page-39-0)

In this subsection, we provide the proof of Lemma [B.1,](#page-38-0) Lemma [B.2,](#page-38-1) and Lemma [B.3.](#page-39-0)

By the definition, we have $\exp(-\overline{\tau}t) \leq \mu_t \leq \exp(-\underline{\tau}t)$ and $1 - \exp(-2\underline{\tau}t) \leq \sigma_t^2 \leq 1 - \exp(-2\overline{\tau}t)$ for $t \geq 0$. Since $x/2 \leq 1 - e^{-x}$ for $0 \leq x \leq 1$ and $1 - e^{-x} \leq x$ for $x \geq 0$, we have

$$
\frac{1}{2} \le 1 - \overline{\tau}t \le \mu_t \le 1 - \frac{\tau t}{2} \le 1 \quad \text{and} \quad \sqrt{\tau t} \le \sigma_t \le \sqrt{2\overline{\tau}t} \tag{B.1}
$$

for any $0 \le t \le (2\overline{\tau})^{-1}$, which is widely used in the following proofs.

B.1.1 Proof of Lemma [B.1](#page-38-0)

Proof. Let $m_0 = \frac{2m}{(B-A)}$ $\frac{2m}{(B-A)n_{\beta}}$ and consider a function $g \in \mathcal{H}^{\beta,K}_1([A,B])$. Simple calculation yields that

$$
\int_{A}^{B} g(x) dx = \sum_{i=1}^{\frac{m_0(B-A)}{2}} \int_{A+\frac{2(i-1)}{m_0}}^{A+\frac{2i}{m_0}} g(x) dx = \sum_{i=1}^{\frac{m_0(B-A)}{2}} \int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} g_i(x) dx,
$$
\n(B.2)

where $g_i(x) = g(x + A + \frac{2i-1}{m_0})$ $\frac{i(i-1)}{m_0}$). For each $i \in \{1, \ldots, \frac{m_0(B-A)}{2}$ $\frac{B-A}{2}$, let L_i be the Lagrange interpolating polynomial of degree $n_{\beta} - 1$ that agrees with the function g_i at knots $\{\widetilde{x}_1/m_0, \ldots, \widetilde{x}_{n_{\beta}}/m_0\}$, defined as

$$
L_i(x) = \sum_{j=1}^{n_{\beta}} g_i(\widetilde{x}_j/m_0) l_j(x),
$$

where

$$
l_j(x) = \begin{cases} \prod_{k=1}^{[\beta]} \frac{m_0 x - \widetilde{x}_k}{\widetilde{x}_j - \widetilde{x}_k}, & \text{if } \beta > 2\\ \prod_{k \neq j}^{k=1} \frac{1}{\widetilde{x}_j - \widetilde{x}_k}, & \text{if } 0 < \beta \leq 2. \end{cases}
$$

for $j \in [n_\beta]$. Note that $L_i(\widetilde{x}_j/m_0) = g_i(\widetilde{x}_j/m_0)$ for any $j \in [n_\beta]$. If $0 < \beta \le 1$, $\tilde{x}_1 = 0$ and $L_i(x) = g_i(0)$. Since $g_i \in \mathcal{H}_1^{\beta,K}([-1/m_0, 1/m_0])$, we have

$$
\left| \int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} \left\{ g_i(x) - L_i(x) \right\} dx \right| \leq \int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} |g_i(x) - L_i(x)| dx
$$
\n
$$
\leq \int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} K|x|^{\beta} dx \leq \frac{2}{m_0} K m_0^{-\beta} = 2K m_0^{-(\beta+1)}.
$$
\n(B.3)

If $\beta > 1$, fix $\widetilde{x}_0 \in [-1, 1]$ satisfying $\widetilde{x}_0 \neq \widetilde{x}_j$ for $j \in [[\beta]]$. Consider a function $h : [-1/m_0, 1/m_0] \to \mathbb{R}$ such that

$$
h(x) = g_i(x) - L_i(x) - \left\{ g_i\left(\frac{\widetilde{x}_0}{m_0}\right) - L_i\left(\frac{\widetilde{x}_0}{m_0}\right) \right\} \prod_{j=1}^{|\beta|} \left(\frac{m_0 x - \widetilde{x}_j}{\widetilde{x}_0 - \widetilde{x}_j} \right).
$$

Then, $h(\tilde{x}_i/m_0) = 0$ for $j \in \{0, \ldots, |\beta|\}$ and g is | β |-times differentiable on $(-1/m_0, 1/m_0)$. Gen-eralized Rolle's Theorem (see Theorem 1.10 of [Burden and Faires](#page-22-1) [\(2010\)](#page-22-1)) implies that there exists a constant $\xi_{\tilde{x}_0} \in (-1/m_0, 1/m_0)$ such that $(D^{[\beta]}h)(\xi_{\tilde{x}_0}) = 0$. Since L_i is the polynomial of degree less than $\lfloor \beta \rfloor$ and $(D^{\lfloor \beta \rfloor} L_i)(\xi_{\widetilde{x}_0}) = 0$, a simple calculation yields that

$$
g_i\left(\frac{\widetilde{x}_0}{m_0}\right) = L_i\left(\frac{\widetilde{x}_0}{m_0}\right) + \frac{(\mathcal{D}^{\lfloor \beta \rfloor} f)(\xi_{\widetilde{x}_0})}{\lfloor \beta \rfloor!} \prod_{j=1}^{\lfloor \beta \rfloor} \left(\frac{\widetilde{x}_0 - \widetilde{x}_j}{m_0}\right).
$$

Note that $g_i(\tilde{x}_j/m_0) = L_i(\tilde{x}_j/m_0)$ for $j \in [\lfloor \beta \rfloor]$. Combining with the last display, there exists a function $\xi : [-1/m_0, 1/m_0] \to (-1/m_0, 1/m_0)$ such that

$$
g_i(x) = L_i(x) + \frac{(\mathcal{D}^{\lfloor \beta \rfloor} f)(\xi(x))}{\lfloor \beta \rfloor!} \prod_{j=1}^{\lfloor \beta \rfloor} \left(x - \frac{\widetilde{x}_j}{m_0} \right), \quad x \in \left[-\frac{1}{m_0}, \frac{1}{m_0} \right]
$$

,

where $\xi(x) = 0$ for $x \in \{\frac{\tilde{x}_1}{m_0}, \ldots, \frac{\tilde{x}_{|\beta|}}{m_0}\}$. For $x \in [-1/m_0, 1/m_0]$, we have

$$
\begin{split} \left| g_i(x) - L_i(x) - \frac{(\mathcal{D}^{\lfloor \beta \rfloor} g)(0)}{\lfloor \beta \rfloor !} \prod_{j=1}^{\lfloor \beta \rfloor} \left(x - \frac{\widetilde{x}_j}{m_0} \right) \right| \\ & = \left| \left\{ \frac{(\mathcal{D}^{\lfloor \beta \rfloor} g)(\xi(x)) - (\mathcal{D}^{\lfloor \beta \rfloor} g)(0)}{\lfloor \beta \rfloor !} \right\} \prod_{j=1}^{\lfloor \beta \rfloor} \left(x - \frac{\widetilde{x}_j}{m_0} \right) \right| \\ & \leq \frac{K |\xi(x)|^{\beta - \lfloor \beta \rfloor}}{\lfloor \beta \rfloor !} \prod_{j=1}^{\lfloor \beta \rfloor} \left(\frac{2}{m_0} \right) \leq \frac{K 2^{\lfloor \beta \rfloor}}{\lfloor \beta \rfloor !} m_0^{-\beta}, \end{split}
$$

where the first inequality holds because $g_i \in \mathcal{H}_1^{\beta,K}([-1/m_0, 1/m_0])$. Since $\{\tilde{x}_1, \ldots, \tilde{x}_{\lfloor \beta \rfloor}\}\$ are the roots of the Legendre polynomial, its orthogonality implies that $\int_{-1/m_0}^{1/m_0} \prod_{j=1}^{[\beta]} (x - \frac{\tilde{x}_j}{m_0}) dx = 0$. Combining with the last display, it follows that

$$
\left| \int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} \left\{ g_i(x) - L_i(x) \right\} dx \right|
$$
\n
$$
= \left| \int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} \left\{ g_i(x) - L_i(x) - \frac{(D^{[\beta]}g)(0)}{[\beta]!} \prod_{j=1}^{[\beta]} \left(x - \frac{\tilde{x}_j}{m_0} \right) \right\} dx \right|
$$
\n
$$
\leq \int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} \left| g_i(x) - L_i(x) - \frac{(D^{[\beta]}g)(0)}{[\beta]!} \prod_{j=1}^{[\beta]} \left(x - \frac{\tilde{x}_j}{m_0} \right) \right| dx \leq \frac{K2^{[\beta]+1}}{[\beta]!} m_0^{-(\beta+1)}.
$$
\n(B.4)

A simple calculation yields that

$$
\int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} L_i(x) = \sum_{j=1}^{n_\beta} g_i\left(\frac{\tilde{x}_j}{m_0}\right) \left\{ \int_{-\frac{1}{m_0}}^{\frac{1}{m_0}} l_j(x) dx \right\} = \sum_{j=1}^{n_\beta} \frac{\tilde{w}_j}{m_0} g_i\left(\frac{\tilde{x}_j}{m_0}\right).
$$

Combining $(B.2)$, $(B.3)$ and $(B.4)$ with the last display, we have

$$
\left| \int_{A}^{B} g(x) dx - \sum_{i=1}^{\frac{m_{0}(B-A)}{2}} \sum_{j=1}^{n_{\beta}} \frac{\widetilde{w}_{j}}{m_{0}} g_{i} \left(\frac{\widetilde{x}_{j}}{m_{0}} \right) \right| \leq \sum_{i=1}^{\frac{m_{0}(B-A)}{2}} \left| \int_{-\frac{1}{m_{0}}}^{\frac{1}{m_{0}}} \left\{ g_{i}(x) - L_{i}(x) \right\} dx \right|
$$

$$
\leq \frac{K(B-A)2^{\lfloor \beta \rfloor}}{\lfloor \beta \rfloor!} m_{0}^{-\beta} = \frac{(\lfloor \beta \rfloor \vee 1)^{\beta}}{2^{\beta-\lfloor \beta \rfloor} \lfloor \beta \rfloor!} m^{-\beta}.
$$

Then, the assertion follows because $\sum_{i=1}^{m} w_i g(x_i) = \sum_{i=1}^{m_0 (B-A)/2} \sum_{j=1}^{n_\beta} \frac{\tilde{w}_j}{m_0} g_i \left(\frac{\tilde{x}_j}{m_0} \right)$. \Box

B.1.2 Proof of Lemma [B.2](#page-38-1)

Proof. For any $n \in \mathbb{Z}_{\geq 0}$, it is well-known (see [\(Indritz,](#page-24-0) [1961\)](#page-24-0)) that

$$
\left|\frac{\mathrm{d}^n}{\mathrm{d}x^n}e^{-x^2}\right| \le \sqrt{2^n n!}e^{-\frac{x^2}{2}}, \quad x \in \mathbb{R}
$$

and moreover,

$$
\|\mathcal{D}^n\phi\|_{\infty} = \frac{1}{\sqrt{2\pi}} \left\| \frac{\mathrm{d}^n}{\mathrm{d}x^n} e^{-\frac{x^2}{2}} \right\|_{\infty} \le \frac{\sqrt{n!}}{\sqrt{2\pi}} \left\| e^{-\frac{x^2}{4}} \right\|_{\infty} \le \frac{\sqrt{n!}}{\sqrt{2\pi}},\tag{B.5}
$$

where the first inequality holds by the chain rule. Then,

$$
\sum_{\alpha=0}^{[\beta]} ||D^{\alpha}(g\phi)||_{\infty} = \sum_{\alpha=0}^{[\beta]} \left\| \sum_{r=0}^{\alpha} {\alpha \choose r} (D^r g)(D^{\alpha-r}\phi) \right\|_{\infty}
$$

\n
$$
\leq \sum_{\alpha=0}^{[\beta]} \sum_{r=0}^{\alpha} {\alpha \choose r} ||D^r g||_{\infty} ||D^{\alpha-r}\phi||_{\infty} \leq \sum_{\alpha=0}^{[\beta]} \sum_{r=0}^{\alpha} \alpha^{\alpha} ||D^r g||_{\infty} \frac{\sqrt{(\alpha-r)!}}{\sqrt{2\pi}}
$$

\n
$$
\leq \sum_{\alpha=0}^{[\beta]} K\alpha^{\alpha} \frac{\sqrt{\alpha!}}{\sqrt{2\pi}} \leq K (|\beta|+1)^{[\beta]+1} \frac{\sqrt{[\beta]!}}{\sqrt{2\pi}}.
$$
\n(B.6)

Similarly, we have

$$
\sum_{\alpha=0}^{\lfloor \beta \rfloor} \left\| \mathcal{D}^{\alpha}(g\phi') \right\|_{\infty} \le \sum_{\alpha=0}^{\lfloor \beta \rfloor} \sum_{r=0}^{\alpha} {\binom{\alpha}{r}} \left\| \mathcal{D}^r g \right\|_{\infty} \left\| \mathcal{D}^{\alpha-r+1}\phi \right\|_{\infty} \le K \left(\lfloor \beta \rfloor + 1 \right)^{n_{\beta}+1} \frac{\sqrt{(\lfloor \beta \rfloor + 1)!}}{\sqrt{2\pi}}. \tag{B.7}
$$

For any differentiable function $h:A\subseteq\mathbb{R}\to\mathbb{R}$ and $0<\gamma\leq 1,$ we have

$$
\sup_{\substack{x,y\in A\\x\neq y}}\frac{|h(x)-h(y)|}{|x-y|^{\gamma}} \leq \sup_{\substack{x,y\in A\\x\neq y\\|x-y|\leq 1}}\frac{|h(x)-h(y)|}{|x-y|^{\gamma}} + \sup_{\substack{x,y\in A\\|x-y|\geq 1}}\frac{|h(x)-h(y)|}{|x-y|\geq 1}
$$
\n
$$
\leq \sup_{\substack{x,y\in A\\x\neq y\\|x-y|\leq 1}}\frac{|h(x)-h(y)|}{|x-y|} + \sup_{\substack{x,y\in A\\|x-y|\geq 1}}|h(x)-h(y)| \leq ||h'||_{\infty} + 2||h||_{\infty}.
$$
\n(B.8)

If $\beta > 1$, it follows that

$$
\frac{\left|(\mathbf{D}^{[\beta]}(g\phi))(x) - (\mathbf{D}^{[\beta]}(g\phi))(y)\right|}{|x - y|^{\beta - [\beta]}} \\
= \frac{\left|\sum_{\alpha=0}^{[\beta]}\binom{[\beta]}{ \alpha}\left\{(\mathbf{D}^{\alpha}g)(x)(\mathbf{D}^{[\beta]-\alpha}\phi)(x) - (\mathbf{D}^{\alpha}g)(y)(\mathbf{D}^{[\beta]-\alpha}\phi)(y)\right\}\right|}{|x - y|^{\beta - [\beta]}} \\
\leq \sum_{\alpha=0}^{[\beta]} \binom{[\beta]}{ \alpha}\left|(\mathbf{D}^{\alpha}g)(x)\right| \left(\frac{\left|(\mathbf{D}^{[\beta]-\alpha}\phi)(x) - (\mathbf{D}^{[\beta]-\alpha}\phi)(y)\right|}{|x - y|^{\beta - [\beta]}}\right) \\
+ \sum_{\alpha=0}^{[\beta]} \binom{[\beta]}{ \alpha}\left|(\mathbf{D}^{[\beta]-\alpha}\phi)(y)\right| \left(\frac{\left|(\mathbf{D}^{\alpha}g)(x) - (\mathbf{D}^{\alpha}g)(y)\right|}{|x - y|^{\beta - [\beta]}}\right) \\
\leq \sum_{\alpha=0}^{[\beta]} \binom{[\beta]}{ \alpha}\|D^{\alpha}g\|_{\infty} \left(\left\|\mathbf{D}^{[\beta]-\alpha+1}\phi\right\|_{\infty} + 2\left\|\mathbf{D}^{[\beta]-\alpha}\phi\right\|_{\infty}\right) \\
+ \sum_{\alpha=0}^{[\beta]-1} \binom{[\beta]}{ \alpha}\|D^{[\beta]-\alpha}\phi\|_{\infty} \left(\left\|\mathbf{D}^{\alpha+1}g\right\|_{\infty} + 2\left\|\mathbf{D}^{\alpha}g\right\|_{\infty}\right) \\
+ \|\phi\|_{\infty} \left(\frac{\left|(\mathbf{D}^{[\beta]}g)(x) - (\mathbf{D}^{[\beta]}g)(y)\right|}{|x - y|^{\beta - [\beta]}}\right)
$$

for any $x, y \in A$ with $x \neq y$, where the last inequality holds by [\(B.8\)](#page-43-0). Combining [\(B.5\)](#page-43-1) with the last display, we have

$$
\sup_{\substack{x,y\in A \\ x\neq y}} \frac{|(D^{[\beta]}(g\phi))(x) - (D^{[\beta]}(g\phi))(y)|}{|x - y|^{\beta - [\beta]}} \n\leq \sum_{\alpha=0}^{[\beta]} {\binom{\beta}{\alpha}} ||D^{\alpha}g||_{\infty} (\left\| D^{[\beta]-\alpha+1}\phi \right\|_{\infty} + 2 \left\| D^{[\beta]-\alpha}\phi \right\|_{\infty}) \n+ \sum_{\alpha=0}^{[\beta]-1} {\binom{\beta}{\alpha}} ||D^{[\beta]-\alpha}\phi||_{\infty} (\left\| D^{\alpha+1}g \right\|_{\infty} + 2 \left\| D^{\alpha}g \right\|_{\infty}) \n+ \|\phi\|_{\infty} {\left(\sup_{x,y\in A} \frac{|(D^{[\beta]}g)(x) - (D^{[\beta]}g)(y)|}{|x - y|^{\beta - [\beta]}} \right)} \n\leq \sum_{\alpha=0}^{[\beta]} [\beta]^{\alpha} ||D^{\alpha}g||_{\infty} {\left(\frac{\sqrt{([\beta]-\alpha+1)!}}{\sqrt{2\pi}} + \frac{2\sqrt{([\beta]-\alpha)!}}{\sqrt{2\pi}} \right)} \n+ \sum_{\alpha=0}^{[\beta]-1} [\beta]^{\alpha} (\left\| D^{\alpha+1}g \right\|_{\infty} + 2 \left\| D^{\alpha}g \right\|_{\infty}) {\left(\frac{\sqrt{([\beta]-\alpha)!}}{\sqrt{2\pi}} \right)} \n+ \frac{1}{\sqrt{2\pi}} {\left(\sup_{x,y\in A} \frac{|(D^{[\beta]}g)(x) - (D^{[\beta]}g)(y)|}{|x - y|^{\beta - [\beta]}} \right)}.
$$

Moreover, the last display is bounded by

$$
\leq 3\lfloor\beta\rfloor^{\lfloor\beta\rfloor} \left(\frac{\sqrt{(\lfloor\beta\rfloor+1)!}}{\sqrt{2\pi}}\right) \left(\sum_{\alpha=0}^{\lfloor\beta\rfloor} \|\mathcal{D}^{\alpha}g\|_{\infty}\right) \n+ 3\lfloor\beta\rfloor^{\lfloor\beta\rfloor-1} \left(\frac{\sqrt{\lfloor\beta\rfloor!}}{\sqrt{2\pi}}\right) \left(\sum_{\alpha=0}^{\lfloor\beta\rfloor} \|\mathcal{D}^{\alpha}g\|_{\infty} + \sup_{\substack{x,y\in A \\ x\neq y}} \frac{|(\mathcal{D}^{\lfloor\beta\rfloor}g)(x) - (\mathcal{D}^{\lfloor\beta\rfloor}g)(y)|}{|x-y|^{\beta-\lfloor\beta\rfloor}}\right) \n\leq 6K\lfloor\beta\rfloor^{\lfloor\beta\rfloor} \left(\frac{\sqrt{(\lfloor\beta\rfloor+1)!}}{\sqrt{2\pi}}\right).
$$
\n(B.9)

Similarly, we have

$$
\sup_{\substack{x,y\in A \\ x\neq y}} \frac{|(D^{|\beta]}(g\phi'))(x) - (D^{|\beta]}(g\phi'))(y)|}{|x - y|^{\beta - |\beta|}}
$$
\n
$$
\leq \sum_{\alpha=0}^{|\beta|} {\binom{|\beta|}{\alpha}} ||D^{\alpha}g||_{\infty} (||D^{|\beta|- \alpha + 2}\phi||_{\infty} + 2 ||D^{|\beta|- \alpha + 1}\phi||_{\infty})
$$
\n
$$
+ \sum_{\alpha=0}^{|\beta|-1} {\binom{|\beta|}{\alpha}} ||D^{|\beta|- \alpha + 1}\phi||_{\infty} (||D^{\alpha+1}g||_{\infty} + 2 ||D^{\alpha}g||_{\infty})
$$
\n
$$
+ ||\phi'||_{\infty} \left(\sup_{x,y\in A} \frac{|(D^{|\beta|}g)(x) - (D^{|\beta|}g)(y)|}{|x - y|^{\beta - |\beta|}} \right)
$$
\n
$$
\leq \sum_{\alpha=0}^{|\beta|} n_{\beta}^{\alpha} ||D^{\alpha}g||_{\infty} \left(\frac{\sqrt{([\beta]-\alpha+2)!}}{\sqrt{2\pi}} + \frac{2\sqrt{([\beta]-\alpha+1)!}}{\sqrt{2\pi}} \right)
$$
\n
$$
+ \sum_{\alpha=0}^{|\beta|-1} |\beta|^{\alpha} (||D^{\alpha+1}g||_{\infty} + 2 ||D^{\alpha}g||_{\infty}) \left(\frac{\sqrt{([\beta]-\alpha+1)!}}{\sqrt{2\pi}} \right)
$$
\n
$$
+ \frac{1}{\sqrt{2\pi}} \left(\sup_{\substack{x,y\in A \\ x\neq y}} \frac{|(D^{|\beta|}g)(x) - (D^{|\beta|}g)(y)|}{|x - y|^{\beta - |\beta|}} \right)
$$
\n
$$
\leq 6K|\beta|^{|\beta|} \left(\frac{\sqrt{([\beta]+2)!}}{\sqrt{2\pi}} \right).
$$
\n(6)

If $0 < \beta \leq 1,$ we have

$$
\sup_{\substack{x,y\in A\\x\neq y}}\frac{|(g\phi)(x) - (g\phi)(y)|}{|x - y|^{\beta}} \le \sup_{\substack{x,y\in A\\x\neq y}}\frac{|g(x)\phi(x) - g(x)\phi(y) + g(x)\phi(y) - g(y)\phi(y)|}{|x - y|^{\beta}}
$$

$$
\le ||g||_{\infty} \left(\sup_{\substack{x,y\in A\\x\neq y}}\frac{|\phi(x) - \phi(y)|}{|x - y|^{\beta}}\right) + ||\phi||_{\infty} \left(\sup_{\substack{x,y\in A\\x\neq y}}\frac{|g(x) - g(y)|}{|x - y|^{\beta}}\right)
$$

$$
\le \left(||g||_{\infty} + \sup_{\substack{x,y\in A\\x\neq y}}\frac{|g(x) - g(y)|}{|x - y|^{\beta}}\right) \left(||\phi'||_{\infty} + 2 ||\phi||_{\infty}\right) \le \frac{3K}{\sqrt{2\pi}}
$$

and

$$
\sup_{\substack{x,y\in A\\x\neq y}}\frac{|(g\phi')(x)-(g\phi')(y)|}{|x-y|^\beta}\leq \left(\|g\|_\infty+\sup_{\substack{x,y\in A\\x\neq y}}\frac{|g(x)-g(y)|}{|x-y|^\beta}\right)\left(\|\phi''\|_\infty+2\|\phi'\|_\infty\right)
$$

$$
\leq \frac{3\sqrt{2}K}{\sqrt{2\pi}},
$$

where the second and third inequality holds by $(B.8)$ and $(B.5)$, respectively. Combining $(B.6)$, $(B.7)$, [\(B.9\)](#page-45-0), [\(B.10\)](#page-45-1) with the last display, we have $g\phi \in \mathcal{H}_1^{\beta,KD_1}(A)$ and $g\phi' \in \mathcal{H}_1^{\beta,KD_1}$, where

$$
D_1 = (\lfloor \beta \rfloor + 1)^{\lfloor \beta \rfloor + 1} \frac{\sqrt{(\lfloor \beta \rfloor + 1)!}}{\sqrt{2\pi}} + 6\lfloor \beta \rfloor^{\lfloor \beta \rfloor} \left(\frac{\sqrt{(\lfloor \beta \rfloor + 2)!}}{\sqrt{2\pi}} \right).
$$

The assertion follows by re-defining the constant.

B.1.3 Proof of Lemma [B.3](#page-39-0)

Proof. Consider τ_{bd} , $\tau_{\text{tail}} > 0$, $m \in n_{\beta} \mathbb{N}$ and real-valued D-dimensional vectors $\mathbf{x} = (x_1, \dots, x_D)^\top$, $\mathbf{y} =$ $(y_1, \ldots, y_D)^\top$ such that $\|\mathbf{x}\|_{\infty} \leq \mu - \mu \{ \log(1/\sigma) \}^{-\tau_{\text{bd}}}$ and $\|\mathbf{x} + \sigma \mathbf{y}\|_{\infty} \leq \mu$. Let $\mathbf{y}_{\lt i+1} = (y_1, \ldots, y_i)^\top \in$ \mathbb{R}^i and $\mathbf{y}_{>D-i} = (y_{D-i+1}, \ldots, y_D)^\top \in \mathbb{R}^i$ for $i \in [D]$. For $y \in \mathbb{R}$ and each i, denote $(\mathbf{y}_{\leq i}, y, \mathbf{y}_{>i})$ as a D-dimensional vector that is identical to y except for the *i*-th component, which is replaced by y.

Let
$$
D_{\sigma} = 2\sqrt{2\tau_{tail}} \{ \log(1/\sigma) \}^{(\tau_{\text{bd}} + \frac{1}{2})}
$$
 and fix $i \in [D]$. Then,

$$
-x_i - \mu \le -\mu {\log(1/\sigma)}^{-\tau_{\text{bd}}} < 0 \quad \text{and} \quad -x_i + \mu \ge \mu {\log(1/\sigma)}^{-\tau_{\text{bd}}} > 0.
$$

Moreover, $(-x_i - \mu)/\sigma < (-x_i - \mu)D_\sigma < 0$ and $(-x_i + \mu)/\sigma > (-x_i + \mu)D_\sigma > 0$ for small enough σ so that $0 < D_{\sigma} \leq (2\sigma)^{-1}$. Consider a one-dimensional real-valued function g_i such that

$$
g_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) = p_0\left(\frac{\mathbf{x} + \sigma(\mathbf{y}_{< i}, y, \mathbf{y}_{> i})}{\mu}\right) \phi(y), \quad y \in [(-x_i - \mu)/\sigma, (-x_i + \mu)/\sigma].
$$

Then,

$$
\left| \int_{\frac{-x_i-\mu}{2}}^{\frac{-x_i+\mu}{\sigma}} g_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy - \int_{(-x_i-\mu)D_{\sigma}}^{(-x_i+\mu)D_{\sigma}} g_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy \right|
$$
\n
$$
= \int_{\frac{-x_i-\mu}{\sigma}}^{(-x_i-\mu)D_{\sigma}} g_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy + \int_{(-x_i+\mu)D_{\sigma}}^{\frac{-x_i+\mu}{\sigma}} g_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy
$$
\n
$$
\leq K \int_{-\infty}^{(-x_i-\mu)D_{\sigma}} \phi(y) dy + K \int_{(-x_i+\mu)D_{\sigma}}^{\infty} \phi(y) dy
$$
\n
$$
\leq K \exp\left(-\frac{(-x_i-\mu)^2 D_{\sigma}^2}{2}\right) + K \exp\left(-\frac{(-x_i+\mu)^2 D_{\sigma}^2}{2}\right)
$$
\n
$$
\leq 2K \exp(-\tau_{\text{tail}} \log(1/\sigma)) = 2K\sigma^{\tau_{\text{tail}}},
$$
\n(B.11)

where the second inequality holds by the tail probability of the standard normal distribution. Let $C_{G,1} = C_{G,1}(\beta)$ be the constant in Lemma [B.2.](#page-38-1) Since $\sigma/\mu < 1$ for $\sigma < 2$, Lemma [B.2](#page-38-1) implies that $g_i \in \mathcal{H}_1^{\beta, KC_{G,1}}([(-x_i-\mu)D_{\sigma}, (-x_i+\mu)D_{\sigma}])$. Moreover, Lemma [B.1](#page-38-0) implies that

$$
\left| \int_{(-x_i - \mu)D_{\sigma}}^{(-x_i + \mu)D_{\sigma}} g_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy - \sum_{j=1}^m \widetilde{v}_j g_i \left(\widetilde{y}_j^{(i)}; \mathbf{y}_{< i}, \mathbf{y}_{> i} \right) \right|
$$
\n
$$
\leq \left\{ \frac{2^{\lfloor \beta \rfloor} n_{\beta}^{\beta}}{\lfloor \beta \rfloor!} \right\} C_{G,1} K \mu^{\beta} D_{\sigma}^{\beta} m^{-\beta}, \tag{B.12}
$$

 \Box

where

$$
\begin{split} &\widetilde{y}_j^{(i)} = (-x_i - \mu)D_\sigma + \frac{\mu D_\sigma n_\beta}{m}\left\{\widetilde{x}_{j-\lfloor\frac{j}{n_\beta}\rfloor} + 2\left\lfloor\frac{j}{n_\beta}\right\rfloor + 1\right\},\\ &\widetilde{v}_j = \frac{\mu D_\sigma n_\beta}{m}\widetilde{w}_{j-\lfloor\frac{j}{n_\beta}\rfloor}, \end{split}
$$

and $(-x_i - \mu)D_{\sigma} < \tilde{y}_j^{(i)} < (-x_i + \mu)D_{\sigma}$ for $j \in [m]$. Combining [\(B.11\)](#page-46-0) and [\(B.12\)](#page-46-1), we have

$$
\left| \int_{\frac{-x_i-\mu}{\sigma}}^{\frac{-x_i+\mu}{\sigma}} p_0\left(\frac{\mathbf{x} + \sigma(\mathbf{y}_{< i}, y, \mathbf{y}_{> i})}{\mu}\right) \phi(y) dy - \sum_{j=1}^m \widetilde{v}_j g_i\left(\widetilde{y}_j^{(i)}; \mathbf{y}_{< i}, \mathbf{y}_{> i}\right) \right| \le \epsilon, \tag{B.13}
$$

where $\epsilon = 2K\sigma^{\tau_{\text{tail}}} + 2^{\lfloor \beta \rfloor} n_{\beta}{}^{\beta} C_{G,1} K \mu^{\beta} D_{\sigma}^{\beta} m^{-\beta} / \lfloor \beta \rfloor!$. Since $|x_i| \leq \mu - \mu \{ \log(1/\sigma) \}^{-\tau_{\text{bd}}}$ and $\sigma D_{\sigma} \leq 1/2$, we have

$$
\frac{x_i}{2\mu} - \frac{1}{2} \le \frac{x_i + \sigma D_{\sigma}(-x_i - \mu)}{\mu} \le \frac{x_i + \sigma \widetilde{y}_j^{(i)}}{\mu} \le \frac{x_i + \sigma D_{\sigma}(-x_i + \mu)}{\mu} \le \frac{x_i}{2\mu} + \frac{1}{2}
$$

and

$$
\left|\frac{x_i + \sigma \widetilde{y}_j^{(i)}}{\mu}\right| \le 1 - \frac{\left\{\log(1/\sigma)\right\}^{-\tau_{\text{bd}}}}{2} < 1.
$$

Consider $F_{(j_1)}, \ldots, F_{(j_1,\ldots,j_D)}$ for $j_1, \ldots, j_D \in [m]$, defined as

$$
F_{(j_1,\ldots,j_{k-1})}
$$
\n
$$
= \int_{\|\mathbf{x}_{>k}+\sigma\mathbf{y}_{>k}\|_{\infty}\leq\mu} \left\{ \int_{\frac{-x_k-\mu}{\sigma}}^{\frac{-x_k+\mu}{\sigma}} p_0\left(\frac{\mathbf{x}+\sigma\left(\tilde{y}_{j_1}^{(1)},\ldots,\tilde{y}_{j_{k-1}}^{(k-1)},y,\mathbf{y}_{>k}^\top\right)^\top}{\mu}\right) \phi(y) \mathrm{d}y \right\} \prod_{i=k+1}^D \phi(y_i) \mathrm{d}\mathbf{y}_{>k},
$$

for $k \in \{2, ..., D-1\},\$

$$
F_{(j_1,\ldots,j_{D-1})} = \int_{\frac{-x_D+\mu}{\sigma}}^{\frac{-x_D+\mu}{\sigma}} p_0 \left(\frac{\mathbf{x} + \sigma \left(\tilde{y}_{j_1}^{(1)}, \ldots, \tilde{y}_{j_{D-1}}^{(D-1)}, y \right)^\top}{\mu} \right) \phi(y) \mathrm{d}y, \text{ and}
$$

$$
F_{(j_1,\ldots,j_D)} = p_0 \left(\frac{\mathbf{x} + \sigma \left(\tilde{y}_{j_1}^{(1)}, \ldots, \tilde{y}_{j_D}^{(D)} \right)^\top}{\mu} \right).
$$

For any $k \in \{2, \ldots, D-1\}$ and $j_1, \ldots, j_D \in [m]$, we have

$$
\left| F_{(j_1,\ldots,j_{k-1})} - \sum_{j_k=1}^m \widetilde{v}_{j_k} \phi \left(\widetilde{y}_{j_k}^{(k)} \right) F_{(j_1,\ldots,j_k)} \right|
$$
\n
$$
\leq \epsilon \int_{\|\mathbf{x}_{>k} + \sigma \mathbf{y}_{>k}\|_{\infty} \leq \mu} \left\{ \prod_{i=k+1}^D \phi(y_i) \right\} d\mathbf{y}_{>k} \leq \epsilon
$$
\n(B.14)

and

$$
\left| F_{(j_1,\ldots,j_{D-1})} - \sum_{j_D=1}^m \widetilde{v}_{j_D} \phi \left(\widetilde{y}_{j_D}^{(D)} \right) F_{(j_1,\ldots,j_D)} \right| \le \epsilon,
$$

where the first and last inequality holds by $(B.13)$. Note that

$$
\mu^D p_{\mu,\sigma}(\mathbf{x}) = \int_{\|\mathbf{x}_{>1} + \sigma \mathbf{y}_{>1}\|_{\infty} \leq \mu} \left[\int_{\frac{-x_1 - \mu}{\sigma}}^{\frac{-x_1 + \mu}{\sigma}} \left\{ p_0 \left(\frac{\mathbf{x} + \sigma(y, \mathbf{y}_{>1}^{\top})^{\top}}{\mu} \right) \phi(y) \right\} dy \right] \prod_{i=2}^D \phi(y_i) dy_{>1}.
$$

Then, we also have

$$
\left|\mu^D p_{\mu,\sigma}(\mathbf{x}) - \sum_{j_1=1}^m \widetilde{v}_{j_1} \phi\left(\widetilde{y}_{j_1}^{(1)}\right) F_{(j_1)}\right| \leq \epsilon \int_{\|\mathbf{x}_{>1} + \sigma \mathbf{y}_{>1}\|_{\infty} \leq \mu} \left\{ \prod_{i=2}^D \phi(y_i) \right\} d\mathbf{y}_{>1} \leq \epsilon,
$$

where the first inequality holds by $(B.13)$. Combining $(B.14)$ with the last display, we have

$$
\left| \mu^{D} p_{\mu,\sigma}(\mathbf{x}) - \sum_{j_{1},...,j_{D}=1}^{m} \prod_{k=1}^{D} \left\{ \tilde{v}_{j_{k}} \phi \left(\tilde{y}_{j_{k}}^{(k)} \right) \right\} F_{(j_{1},...,j_{D})} \right|
$$
\n
$$
\leq \left| \mu^{D} p_{\mu,\sigma}(\mathbf{x}) - \sum_{j_{1}=1}^{m} \tilde{v}_{j_{1}} \phi \left(\tilde{y}_{j_{1}}^{(1)} \right) F_{(j_{1})} \right|
$$
\n
$$
+ \sum_{i=2}^{D} \left| \sum_{j_{1},...,j_{i-1}=1}^{m} \prod_{k=1}^{i-1} \left\{ \tilde{v}_{j_{k}} \phi \left(\tilde{y}_{j_{k}}^{(k)} \right) \right\} \left\{ F_{(j_{1},...,j_{i-1})} - \sum_{j_{i}=1}^{m} \tilde{v}_{j_{i}} \phi \left(\tilde{y}_{j_{i}}^{(i)} \right) F_{(j_{1},...,j_{i})} \right\} \right|
$$
\n
$$
\leq \epsilon \left(1 + \sum_{i=2}^{D} \sum_{j_{1},...,j_{i-1}=1}^{m} \left| \prod_{k=1}^{i-1} \left\{ \tilde{v}_{j_{k}} \phi \left(\tilde{y}_{j_{k}}^{(k)} \right) \right\} \right|
$$
\n
$$
\leq \epsilon \left(1 + \sum_{i=2}^{D} \sum_{j_{1},...,j_{i-1}=1}^{m} \left| \frac{\prod_{k=1}^{i-1} \tilde{v}_{j_{k}}}{(2\pi)^{\frac{i-1}{2}}} \right|,
$$
\n(B.15)

where the last inequality holds because $|\phi| \leq 1/2$ 2π. For each $j \in [m]$, we have

$$
|\widetilde{v}_j| \leq \left\{\frac{\mu D_\sigma n_\beta}{m}\right\} \max\left(|\widetilde{w}_1|,\ldots,|\widetilde{w}_{n_\beta}|\right) = \frac{D_1 \{\log(1/\sigma)\}^{(\tau_{\text{bd}} + \frac{1}{2})}}{m},
$$

where the last inequality holds because $\mu \leq 1$ and $D_1 = 2\sqrt{2\tau_{\text{tail}}}n_{\beta}\max(|\tilde{w}_1|,\ldots,|\tilde{w}_{n_{\beta}}|)$. Then,

$$
\begin{split} & 1 + \sum_{i=2}^D \sum_{j_1,\ldots,j_{i-1}=1}^m \frac{|\prod_{k=1}^{i-1} \widetilde{v}_{j_k}|}{(2\pi)^{\frac{i-1}{2}}} \leq 1 + \sum_{i=2}^D \sum_{j_1,\ldots,j_{i-1}=1}^m \left(\frac{D_1 \{ \log(1/\sigma) \}^{(\tau_{\rm bd} + \frac{1}{2})}}{m\sqrt{2\pi}} \right)^{i-1} \\ & = 1 + \sum_{i=2}^D \left(\frac{D_1 \{ \log(1/\sigma) \}^{(\tau_{\rm bd} + \frac{1}{2})}}{\sqrt{2\pi}} \right)^{i-1} \leq D \left(\frac{D_1 \{ \log(1/\sigma) \}^{(\tau_{\rm bd} + \frac{1}{2})}}{\sqrt{2\pi}} \right)^{D-1}, \end{split}
$$

where the last inequality holds for small enough σ so that $D_1\{\log(1/\sigma)\}^{(\tau_{\text{bd}}+\frac{1}{2})} \geq \sqrt{\sigma}$ 2π . Also, there exists a constant $D_2=D_2(\beta,\tau_{\mathrm{tail}},C_{G,1})$ such that

$$
\epsilon \leq D_2 K \left(\sigma^{\tau_{\text{tail}}} + m^{-\beta} \{ \log(1/\sigma) \}^{(\tau_{\text{bd}} + \frac{1}{2})\beta} \right).
$$

Hence,

$$
\left| \mu^{D} p_{\mu,\sigma}(\mathbf{x}) - \sum_{j_{1},\dots,j_{D}=1}^{m} \prod_{k=1}^{D} \left\{ \tilde{v}_{j_{k}} \phi \left(\tilde{y}_{j_{k}}^{(k)} \right) \right\} F_{(j_{1},\dots,j_{D})} \right|
$$
\n
$$
\leq \epsilon \left(1 + \sum_{i=2}^{D} \sum_{j_{1},\dots,j_{i-1}=1}^{m} \frac{|\prod_{k=1}^{i-1} \tilde{v}_{j_{k}}|}{(2\pi)^{\frac{i-1}{2}}} \right)
$$
\n
$$
\leq D_{3} K \left(\sigma^{\tau_{\text{tail}}} + m^{-\beta} \{ \log(1/\sigma) \}^{(\tau_{\text{bd}} + \frac{1}{2})\beta} \right) \{ \log(1/\sigma) \}^{(\tau_{\text{bd}} + \frac{1}{2})(D-1)},
$$
\n(B.16)

where $D_3 = DD_2(D_1)$ √ $\overline{2\pi})^{D-1}.$

Note that

$$
\nabla p_t(\mathbf{x}) = \int_{\|\mathbf{y}\|_{\infty} \leq 1} \left(\frac{\mu \mathbf{y} - \mathbf{x}}{\sigma^2}\right) p_0(\mathbf{y}) \phi_{\sigma}(\mathbf{x} - \mu \mathbf{y}) d\mathbf{y}
$$

= $\sigma^{-1} \mu^{-D} \int_{\|\mathbf{x} + \sigma \mathbf{y}\|_{\infty} \leq \mu} (y_1, \dots, y_D)^{\top} p_0\left(\frac{\mathbf{x} + \sigma \mathbf{y}}{\mu}\right) \prod_{i=1}^D \phi(y_i) d\mathbf{y}.$

For $i \in [D]$, consider a one-dimensional real-valued function \tilde{g}_i such that

$$
\widetilde{g}_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) = y g_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}), \quad y \in [(-x_i - \mu)/\sigma, (-x_i + \mu)/\sigma].
$$

Then,

$$
\left| \int_{\frac{-x_i-\mu}{2}}^{\frac{-x_i+\mu}{2}} \widetilde{g}_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy \int_{(-x_i-\mu)D_{\sigma}}^{(-x_i+\mu)D_{\sigma}} \widetilde{g}_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy \right|
$$
\n
$$
= \int_{(x_i+\mu)D_{\sigma}}^{\frac{x_i+\mu}{2}} \widetilde{g}_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy + \int_{(-x_i+\mu)D_{\sigma}}^{\frac{-x_i+\mu}{2}} \widetilde{g}_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy
$$
\n
$$
\leq K \int_{(x_i+\mu)D_{\sigma}}^{\infty} y \phi(y) dy + K \int_{(-x_i+\mu)D_{\sigma}}^{\infty} y \phi(y) dy
$$
\n
$$
= K \exp \left(-\frac{(x_i+\mu)^2 D_{\sigma}^2}{2} \right) + K \exp \left(-\frac{(-x_i+\mu)^2 D_{\sigma}^2}{2} \right)
$$
\n
$$
\leq 2K \exp \left(-\tau_{\text{tail}} \log(1/\sigma) \right) = 2K \sigma^{\tau_{\text{tail}}},
$$
\n(B.17)

where the first equality holds because

$$
-x_i - \mu \le -\{\log(1/\sigma)\}^{-\tau_{\text{bd}}}/2 < 0 \quad \text{and} \quad -x_i + \mu \ge 2^{-1}\{\log(1/\sigma)\}^{-\tau_{\text{bd}}} > 0.
$$

Since $\phi'(y) = -y\phi(y)$ for $y \in \mathbb{R}$ and $\sigma/\mu < 1$, Lemma [B.2](#page-38-1) implies that $\widetilde{g}_i \in \mathcal{H}_1^{\beta, KC_{G,1}}((-x_i - \sigma_i)^2)$ μ) D_{σ} , $(-x_i + \mu)D_{\sigma}$]). Moreover, Lemma [B.1](#page-38-0) implies that

$$
\left| \int_{(-x_i - \mu)D_{\sigma}}^{(-x_i + \mu)D_{\sigma}} \tilde{g}_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) dy - \sum_{j=1}^m \tilde{v}_j \tilde{y}_j^{(i)} g_i \left(\tilde{y}_j^{(i)}; \mathbf{y}_{< i}, \mathbf{y}_{> i} \right) \right|
$$

$$
\leq \left\{ \frac{2^{\lfloor \beta \rfloor} n_{\beta}^{\beta}}{\lfloor \beta \rfloor!} \right\} C_{G,1} K \mu^{\beta} D_{\sigma}^{\beta} m^{-\beta}
$$

because $\widetilde{g}_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i}) = y g_i(y; \mathbf{y}_{< i}, \mathbf{y}_{> i})$. Combining [\(B.17\)](#page-49-0) with the last display, we have

$$
\left| \int_{\frac{-x_i-\mu}{\sigma}}^{\frac{-x_i+\mu}{\sigma}} p_0\left(\frac{\mathbf{x} + \sigma(\mathbf{y}_{< i}, y, \mathbf{y}_{> i})}{\mu}\right) y\phi(y) \mathrm{d}y - \sum_{j=1}^m \widetilde{v}_j \widetilde{y}_j^{(i)} g_i\left(\widetilde{y}_j^{(i)}; \mathbf{y}_{< i}, \mathbf{y}_{> i}\right) \right| \le \epsilon. \tag{B.18}
$$

Combining $(B.13)$ with a simple calculation, we have

$$
\left| \int_{\|\mathbf{x}_{-i} + \sigma \mathbf{y}_{-i}\|_{\infty} \leq \mu} p_0\left(\frac{\mathbf{x} + \sigma(\mathbf{y}_{\leq i}, y, \mathbf{y}_{>i})}{\mu}\right) \prod_{\substack{k=1 \ k \neq i}}^D \phi(y_k) \mathrm{d} \mathbf{y}_{\leq i} \n- \sum_{\substack{j_1, \dots, j_D = 1 \\ \text{w/o } j_i}}^m \prod_{\substack{k=1 \ k \neq i}}^D \left\{\widetilde{v}_k \phi\left(\widetilde{y}_{j_k}^{(k)}\right)\right\} p_0\left(\frac{\mathbf{x} + \sigma\left(\widetilde{y}_{j_1}^{(1)}, \dots, \widetilde{y}_{j_{i-1}}^{(i-1)}, y, \widetilde{y}_{j_{i+1}}^{(i+1)}, \dots, \widetilde{y}_{j_D}^{(D)}\right)^{\top}}{\mu}\right) \right| \n\leq \epsilon \left(1 + \sum_{h=1}^{D-1} \sum_{\substack{j_1, \dots, j_h = 1 \\ \text{w/o } j_i}}^m \prod_{\substack{k=1 \ k \neq i}}^h \left\{\left|\widetilde{v}_k\right| \phi\left(\widetilde{y}_{j_k}^{(k)}\right)\right\}\right)
$$

for any $y \in \mathbb{R}$ satisfying $|x_i + \sigma y| \leq \mu$, where $\sum_{j_1,\dots,j_D=1}^m w_{j_0} j_i$ denotes the summation over

$$
1 \leq j_1, \ldots, j_{i-1}, j_{i+1}, \ldots, j_D \leq m.
$$

Note that

$$
\mu^{D} \sigma(\nabla p_{t}(\mathbf{x}))_{i} = \int_{\|\mathbf{x}+\sigma\mathbf{y}\|_{\infty} \leq \mu} y_{i} p_{0} \left(\frac{\mathbf{x}+\sigma\mathbf{y}}{\mu}\right) \prod_{k=1}^{D} \phi(y_{k}) d\mathbf{y}
$$

$$
= \int_{\frac{-x_{i}+\mu}{\sigma}}^{-\frac{x_{i}+\mu}{\sigma}} \left\{ \int_{\|\mathbf{x}_{-i}+\sigma\mathbf{y}_{-i}\|_{\infty} \leq \mu} p_{0} \left(\frac{\mathbf{x}+\sigma\mathbf{y}}{\mu}\right) \prod_{\substack{k=1 \ k \neq i}}^{D} \phi(y_{k}) d\mathbf{y}_{
$$

Combining $(B.18)$ with the three last displays, we have

$$
\left| \int_{\|\mathbf{x}+\sigma\mathbf{y}\|_{\infty} \leq \mu} y_i p_0\left(\frac{\mathbf{x}+\sigma\mathbf{y}}{\mu}\right) \prod_{k=1}^D \phi(y_k) d\mathbf{y} - \sum_{j_1,\ldots,j_D=1}^m \tilde{y}_{j_i}^{(i)} \prod_{k=1}^D \left\{ \tilde{v}_{j_k} \phi\left(\tilde{y}_{j_k}^{(k)}\right) \right\} F_{(j_1,\ldots,j_D)} \right|
$$

\n
$$
\leq \epsilon \left(1 + \sum_{h=1}^{D-1} \sum_{\substack{j_1,\ldots,j_h=1 \\ j_1,\ldots,j_h=1}}^m \prod_{\substack{k=1 \\ k \neq i}}^h \left\{ |\tilde{v}_k| \phi\left(\tilde{y}_{j_k}^{(k)}\right) \right\} + \sum_{\substack{j_1,\ldots,j_D=1 \\ j_1,\ldots,j_D=1}}^m \prod_{\substack{k=1 \\ k \neq i}}^D \left\{ |\tilde{v}_k| \phi\left(\tilde{y}_{j_k}^{(k)}\right) \right\} \right) \int_{\frac{-x_i + \mu}{\sigma}}^{\frac{-x_i + \mu}{\sigma}} |y| \phi(y) dy
$$

\n
$$
\leq \frac{2\epsilon}{\sqrt{2\pi}} \left(1 + \sum_{h=1}^D \sum_{j_1,\ldots,j_h=1}^m \prod_{\substack{k=1 \\ j_1,\ldots,j_h=1}}^D \frac{|\tilde{v}_k|}{\sqrt{2\pi}} \right),
$$

where the last inequality holds because $\int_{-\infty}^{\infty} |y| \phi(y) = 2 \int_{0}^{\infty} y \phi(y) dy = \frac{2}{\sqrt{2}}$ $\frac{2}{2\pi}$ and $|\phi| \leq 1/$ √ 2π. Combining with $(B.16)$, the last display is bounded by

$$
\frac{2D_3K}{\sqrt{2\pi}} \left(\sigma^{\tau_{\rm tail}} + m^{-\beta} {\log(1/\sigma)}^{\{\tau_{\rm bd} + \frac{1}{2})\beta} \right) {\{\log(1/\sigma)\}}^{(\tau_{\rm bd} + \frac{1}{2})(D-1)},
$$

and the assertion follows by re-defining constants.

B.2 Proof of Proposition [B.1](#page-39-1)

Proof. Let $\delta > 0$ be a small enough value as described below. There exists neural networks $f_{\mu} \in$ $\mathcal{F}_{NN}(L_{\mu}, \mathbf{d}_{\mu}, s_{\mu}, M_{\mu}), f_{\sigma} \in \mathcal{F}_{NN}(L_{\sigma}, \mathbf{d}_{\sigma}, s_{\sigma}, M_{\sigma})$ with

$$
L_{\mu}, L_{\sigma} \leq C_{N,4} \{ \log(1/\delta) \}^2, \quad \|\mathbf{d}_{\mu}\|_{\infty}, \|\mathbf{d}_{\sigma}\|_{\infty} \leq C_{N,4} \{ \log(1/\delta) \}^2
$$

$$
s_{\mu}, s_{\sigma} \leq C_{N,4} \{ \log(1/\delta) \}^3, \quad M_{\mu}, M_{\sigma} \leq C_{N,4} \log(1/\delta)
$$

such that

$$
|\mu_t - f_\mu(t)| \le \delta \qquad \text{and} \qquad |\sigma_t - f_\sigma(t)| \le \delta \tag{B.19}
$$

for $t \geq \delta$, where $C_{N,4}$ is the constant in Lemma [A.13.](#page-37-0) for any $0 \leq t \leq (2\overline{\tau})^{-1}$.

Since $\log(1/x) = -\log x$ for any $x > 0$, Lemma [A.11](#page-32-1) implies that there exists a positive constant $D_1 = D_1(\underline{\tau})$ and neural network $f_{\log} \in \mathcal{F}_{\text{NN}}(L_{\log}, \mathbf{d}_{\log}, s_{\log}, M_{\log})$ with

$$
L_{\log} \le D_1 \{ \log(1/\delta) \}^2 \log \log(1/\delta), \quad \|\mathbf{d}_{\log}\|_{\infty} \le D_1 \{ \log(1/\delta) \}^3
$$

$$
s_{\log} \le D_1 \{ \log(1/\delta) \}^5 \log \log(1/\delta), \quad M_{\log} \le \exp(D_1 \{ \log(1/\delta) \}^2)
$$

 \Box

such that $|\log(1/x) - f_{\log}(\tilde{x})| \leq \sqrt{\tau \delta}/2 + (2/\sqrt{\tau \delta})$ $\sqrt{\underline{\tau}\delta}|x-\widetilde{x}|$ for $\sqrt{\underline{\tau}\delta}/2 \leq x \leq 0$ $\sqrt{\underline{\tau}\delta}/2$ ⁻¹ and $\widetilde{x} \in \mathbb{R}$. Combining with [\(B.19\)](#page-50-0), we have

$$
|\log(1/\sigma_t) - f_{\log}(f_{\sigma}(t))| \le \frac{2\sqrt{\delta}}{\sqrt{\mathcal{I}}} + \frac{\sqrt{\mathcal{I}\delta}}{2} = \left(\frac{2}{\sqrt{\mathcal{I}}} + \frac{\sqrt{\mathcal{I}}}{2}\right)\sqrt{\delta}
$$
(B.20)

for $\delta \leq t \leq (2\overline{\tau})^{-1}$. Lemma [A.9](#page-32-2) implies that for $k \geq 2$, there exists a neural network $f_{\text{mult}}^{(k)} \in$ $\mathcal{F}_{\rm NN}(L_{\rm mult}^{(k)},\mathbf{d}_{\rm mult}^{(k)},s_{\rm mult}^{(k)},M_{\rm mult}^{(k)})$ with

$$
L_{\text{mult}}^{(k)} \leq C_{N,1}(k\tau_{\text{bd}} + 1) \log k \log(1/\delta), \quad \|\mathbf{d}_{\text{mult}}^{(k)}\|_{\infty} = 48k,
$$

$$
s_{\text{mult}}^{(k)} \leq C_{N,1}k(\tau_{\text{bd}} + 1) \log(1/\delta), \quad M_{\text{mult}}^{(k)} = {\log(1/\delta)}^{k\tau_{\text{bd}}}
$$

such that

$$
\left| f_{\text{mult}}^{(k)}(\widetilde{x}_1, \dots, \widetilde{x}_k) - \prod_{i=1}^k x_i \right| \le \delta + k \{ \log(1/\delta) \}^{(k-1)\tau_{\text{bd}} \widetilde{\epsilon}} \tag{B.21}
$$

for any $\mathbf{x} = (x_1, \dots, x_k) \in \mathbb{R}^k$ with $\|\mathbf{x}\|_{\infty} \leq {\log(1/\delta)}^{\tau_{\text{bd}}}$ and $\widetilde{\mathbf{x}} = (\widetilde{x}_1, \dots, \widetilde{x}_k) \in \mathbb{R}^k$ with $\|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty} \leq \widetilde{\mathbf{x}}$ $\widetilde{\epsilon}$, where $0 < \widetilde{\epsilon} \leq 1$ and $C_{N,1}$ is the constant in Lemma [A.9.](#page-32-2) Combining Lemma [A.6](#page-30-0) and Lemma [A.4](#page-29-0) with the last display, there exists a neural network $f_{\text{pow}}^{(k)} \in \mathcal{F}_{\text{NN}}(L_{\text{pow}}^{(k)}, \mathbf{d}_{\text{pow}}^{(k)}, s_{\text{pow}}^{(k)}, M_{\text{pow}}^{(k)})$ with

$$
L_{\text{pow}}^{(k)} = L_{\text{mult}}^{(k)} + 2, \quad \|\mathbf{d}_{\text{pow}}^{(k)}\|_{\infty} \le 96k, \quad s_{\text{pow}}^{(k)} \le 2s_{\text{mult}}^{(k)} + 8k, \quad M_{\text{pow}}^{(k)} = M_{\text{mult}}^{(k)} \vee 1
$$

such that

$$
\left| f_{\text{pow}}^{(k)}(\widetilde{x}) - x^k \right| \le \delta + k \{ \log(1/\delta) \}^{(k-1)\tau_{\text{bd}}}\widetilde{\epsilon}
$$
\n(B.22)

for any $|x| \leq {\log(1/\delta)}^{\tau_{\text{bd}}}$ and $\tilde{x} \in \mathbb{R}$ with $|x - \tilde{x}| \leq \tilde{\epsilon}$. For $\delta \leq t \leq (2\overline{\tau})^{-1}$, we have

$$
|\log(1/\sigma_t)| \le \log(1/\sqrt{\tau \delta}) \le \log(1/\delta), \tag{B.23}
$$

where the first inequality holds by $(B.1 \text{ and the last inequality holds with small enough } \delta$ $(B.1 \text{ and the last inequality holds with small enough } \delta$. Combining $(B.20)$ with $(B.22)$, it follows that

$$
\begin{aligned}\n\left| \left\{ \log(1/\sigma_t) \right\}^{\tau_{\rm bd} + \frac{1}{2}} - f_{\rm pow}^{(\tau_{\rm bd} + \frac{1}{2})} \left(f_{\rm log} \left(f_{\sigma}(t) \right) \right) \right| \\
&\leq \delta + \left(\tau_{\rm bd} + \frac{1}{2} \right) \left(\frac{2}{\sqrt{\underline{\tau}}} + \frac{\sqrt{\underline{\tau}}}{2} \right) \sqrt{\delta} \left\{ \log(1/\delta) \right\}^{(\tau_{\rm bd} - \frac{1}{2}) \tau_{\rm bd}} \\
&\leq D_2 \sqrt{\delta} \left\{ \log(1/\delta) \right\}^{\tau_{\rm bd}^2 - \frac{\tau_{\rm bd}}{2}}\n\end{aligned} \tag{B.24}
$$

for $\delta \le t \le (2\overline{\tau})^{-1}$, where $D_2 = 1 + (\tau_{\text{bd}} + \frac{1}{2})$ $(\frac{1}{\sqrt{2}})(\frac{2}{\sqrt{2}} +$ $\sqrt{τ}$ $\frac{\sqrt{2}}{2}$). Combining [\(B.21\)](#page-51-2) with the last two displays, we have

$$
\left| \{ \log(1/\sigma_t) \}^{\tau_{\rm bd} + \frac{1}{2}} x - f_{\rm mult}^{(2)} \left(f_{\rm pow}^{(\tau_{\rm bd} + \frac{1}{2})} (f_{\rm log} (f_{\sigma}(t))), x \right) \right|
$$

$$
\leq \delta + 2D_2 \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\rm bd}^2 + \frac{\tau_{\rm bd}}{2}} \leq D_3 \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\rm bd}^2 + \frac{\tau_{\rm bd}}{2}}
$$

for $\delta \leq t \leq (2\overline{\tau})^{-1}$ and $|x| \leq 1$, where $D_3 = 1+2D_2$. Let $m \in n_\beta\mathbb{N}$ be a large enough value as described below. Then, consider functions $g_y^{(1)}, \ldots, g_y^{(m)} : [-1,1] \times [0,\infty) \to \mathbb{R}$ and $f_y^{(1)}, \ldots, f_y^{(m)} : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ such that

$$
g_{\mathbf{y}}^{(j)}(x,t)
$$

= $2\sqrt{2\tau_{\text{tail}}} \{\log(1/\sigma_t)\}^{\tau_{\text{bd}} + \frac{1}{2}} \left\{-x - \mu_t + \frac{\mu_t n_\beta}{m} \left(\tilde{x}_{j-\lfloor \frac{j}{n_\beta}\rfloor} + 2 \left\lfloor \frac{j}{n_\beta} \right\rfloor + 1\right)\right\}, \quad j \in [m]$

for $x \in [-1, 1], t \in [0, \infty)$ and

$$
f_{\mathbf{y}}^{(j)}(x,t) = 2\sqrt{2\tau_{\text{tail}}} \left\{ f_{\text{mult}}^{(2)} \left(f_{\text{pow}}^{(\tau_{\text{bd}} + \frac{1}{2})} \left(f_{\text{log}} \left(f_{\sigma}(t) \right) \right), x \right) - f_{\text{mult}}^{(2)} \left(f_{\text{pow}}^{(\tau_{\text{bd}} + \frac{1}{2})} \left(f_{\text{log}} \left(f_{\sigma}(t) \right) \right), f_{\mu}(t) \right) \right\} + \frac{n_{\beta}}{m} \left(\tilde{x}_{j - \lfloor \frac{j}{n_{\beta}} \rfloor} + 2 \left\lfloor \frac{j}{n_{\beta}} \right\rfloor + 1 \right) f_{\text{mult}}^{(2)} \left(f_{\text{pow}}^{(\tau_{\text{bd}} + \frac{1}{2})} \left(f_{\text{log}} \left(f_{\sigma}(t) \right) \right), f_{\mu}(t) \right) \right\}, \quad j \in [m]
$$

for $x, t \in \mathbb{R}$, where $\{(\tilde{x}_j, \tilde{w}_j) : j \in [n_\beta]\}$ are the constants in Lemma [B.3.](#page-39-0) Then,

$$
\begin{aligned}\n\left| g_y^{(j)}(x,t) - f_y^{(j)}(x,t) \right| \\
&\le 2\sqrt{2\tau_{\text{tail}}} \left\{ D_3 + \frac{D_3 n_\beta}{m} \left(\left| \tilde{x}_{j-\lfloor \frac{j}{n_\beta} \rfloor} \right| + 2 \left\lfloor \frac{j}{n_\beta} \right\rfloor + 1 \right) + 1 \right\} \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + \frac{\tau_{\text{bd}}}{2}}\n\end{aligned}
$$

and

$$
\left| g_y^{(j)}(x,t) \right| \le 2\sqrt{2\tau_{\text{tail}}} \left\{ 2 + \frac{2n_\beta}{m} \left(\left| \tilde{x}_{j-\lfloor \frac{j}{n_\beta} \rfloor} \right| + 2\left\lfloor \frac{j}{n_\beta} \right\rfloor + 1 \right) \right\} \left\{ \log(1/\delta) \right\}^{\tau_{\text{bd}} + \frac{1}{2}}
$$

for $|x| \leq 1$ and $\delta \leq t \leq (4\overline{\tau})^{-1}$, where the last inequality holds by [\(B.23\)](#page-51-3). Then, there exists a constant $D_4 = D_4(\beta, \tau_{tail}, \tau_{bd}, D_3)$ such that

$$
\left| f_{y}^{(j)}(x,t) \right| \leq \left| g_{y}^{(j)}(x,t) - f_{y}^{(j)}(x,t) \right| + \left| g_{y}^{(j)}(x,t) \right| \leq D_{4} \{ \log(1/\delta) \}^{\tau_{\text{bd}} + \frac{1}{2}}, \text{ and}
$$
\n
$$
\left| g_{y}^{(j)}(x,t) - f_{y}^{(j)}(x,t) \right| \leq D_{4} \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + \frac{\tau_{\text{bd}}}{2}} \tag{B.25}
$$

for $|x| \leq 1$ and $\delta \leq t \leq (2\overline{\tau})^{-1}$ with small enough δ so that $\sqrt{\delta} \{\log(1/\delta)\}^{\tau_{\rm bd}^2 + \frac{\tau_{\rm bd}}{2}} \leq {\log(1/\delta)}\}^{\tau_{\rm bd}+ \frac{1}{2}}$. Lemma [A.4](#page-29-0) implies that $f_{\text{pow}}^{(\tau_{\text{bd}} + \frac{1}{2})} \circ f_{\text{log}} \circ f_{\sigma} \in \mathcal{F}_{\text{NN}}(L_{\text{pl}\sigma}, d_{\text{pl}\sigma}, s_{\text{pl}\sigma}, M_{\text{pl}\sigma})$ with

$$
L_{\text{pl}\sigma} = L_{\text{pow}}^{(\tau_{\text{bd}} + \frac{1}{2})} + L_{\text{log}} + L_{\sigma} \leq D_5 \{ \log(1/\delta) \}^2 \log \log(1/\delta),
$$

$$
\|\mathbf{d}_{\text{pl}\sigma}\|_{\infty} \leq 2 \max \left(\|\mathbf{d}_{\text{pow}}^{(\tau_{\text{bd}} + \frac{1}{2})}\|_{\infty}, \|\mathbf{d}_{\text{log}}\|_{\infty}, \|\mathbf{d}_{\sigma}\|_{\infty} \right) \leq D_5 \{ \log(1/\delta) \}^3,
$$

$$
s_{\text{pl}\sigma} \leq 2 \left(s_{\text{pow}}^{(\tau_{\text{bd}} + \frac{1}{2})} + s_{\text{log}} + s_{\sigma} \right) \leq D_5 \{ \log(1/\delta) \}^5 \log \log(1/\delta),
$$

$$
M_{\text{pl}\sigma} = \max \left(M_{\text{pow}}^{(\tau_{\text{bd}} + \frac{1}{2})}, M_{\text{log}}, M_{\sigma} \right) \leq \exp \left(D_7 \{ \log(1/\delta) \}^2 \right),
$$

where $D_5 = D_5(\tau_{\text{bd}}, C_{N,1}, C_{N,4}, D_1)$. Then, Lemma [A.4,](#page-29-0) Lemma [A.5](#page-29-1) and Lemma [A.6](#page-30-0) imply that $f_{\mathbf{y}}^{(j)} \in \mathcal{F}_{\text{NN}}(L_{\mathbf{x}}, \mathbf{d}_{\mathbf{x}}, s_{\mathbf{x}}, M_{\mathbf{x}}^{(j)})$ for $j \in [m]$ with

$$
L_{\rm x} \le D_6 \{\log(1/\delta)\}^2 \log \log(1/\delta), \quad \|\mathbf{d}_{\rm x}\|_{\infty} \le D_6 \{\log(1/\delta)\}^3,
$$

$$
s_{\rm x} \le D_6 \{\log(1/\delta)\}^5 \log \log(1/\delta), \quad M_{\rm x}^{(j)} \le \exp\left(D_6 \{\log(1/\delta)\}^2\right),
$$
 (B.26)

where $D_6 = D_6(\beta, \tau_{tail}, C_{N,1}, C_{N,4}, D_3, D_5)$. Let $C_{N,5}$ be the constant in Lemma [A.14.](#page-37-1) Then, there exists a neural network $f_{\text{rec}} \in \mathcal{F}_{\text{NN}}(L_{\text{rec}}, \mathbf{d}_{\text{rec}}, s_{\text{rec}}, M_{\text{rec}})$ with

$$
L_{\text{rec}} \leq C_{N,5} \{ \log(1/\delta) \}^2, \quad \|\mathbf{d}_{\text{rec}}\|_{\infty} \leq C_{N,5} \{ \log(1/\delta) \}^3
$$

$$
s_{\text{rec}} \leq C_{N,5} \{ \log(1/\delta) \}^4, \quad M_{\text{rec}} \leq C_{N,5} \delta^{-2}
$$

such that $|x^{-1} - f_{\text{rec}}(x)| \le \delta$ for $x \in [\delta, 1/\delta]$. Combining with [\(B.19\)](#page-50-0), we have

$$
\left| \frac{1}{\mu_t} - f_{\text{rec}}(f_{\mu}(t)) \right| \le \left| \frac{1}{\mu_t} - \frac{1}{f_{\mu}(t)} \right| + \left| \frac{1}{f_{\mu}(t)} - f_{\text{rec}}(f_{\mu}(t)) \right|
$$
\n
$$
\le (\mu_t \wedge f_{\mu}(t))^{-2} |\mu_t - f_{\mu}(t)| + \delta \le 17\delta
$$
\n(B.27)

for $\delta \leq t \leq (2\overline{\tau})^{-1}$, where the second inequality holds because $1/4 \leq 1/2 - \delta \leq f_{\mu}(t)$ with $\delta \leq 1/4$. Consider functions $\tilde{f}_y^{(1)}, \ldots, \tilde{f}_y^{(m)} : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ such that

$$
\widetilde{f}_{y}^{(j)}(x,t) = f_{\text{mult}}^{(2)}\left(f_{\text{rec}}(f_{\mu}(t)), x + f_{\text{mult}}^{(2)}\left(f_{y}^{(j)}(x,t), f_{\sigma}(t)\right)\right), \quad j \in [m]
$$

for $x, t \in \mathbb{R}$. Then, Lemma [A.4,](#page-29-0) Lemma [A.5,](#page-29-1) Lemma [A.6](#page-30-0) and Lemma [A.7](#page-30-1) imply that $\tilde{f}_{y}^{(j)} \in$ $\mathcal{F}_{\text{NN}}(\widetilde{L}_{\text{x}}, \widetilde{\mathbf{d}}_{\text{x}}, \widetilde{s}_{\text{x}}, \widetilde{M}_{\text{x}}^{(j)})$ for $j \in [m]$ with

$$
\widetilde{L}_{\mathbf{x}} \le D_7 \{ \log(1/\delta) \}^2 \log \log(1/\delta), \quad \|\widetilde{\mathbf{d}}_{\mathbf{x}}\|_{\infty} \le D_7 \{ \log(1/\delta) \}^3,
$$
\n
$$
\widetilde{s}_{\mathbf{x}} \le D_7 \{ \log(1/\delta) \}^5 \log \log(1/\delta), \quad \widetilde{M}_{\mathbf{x}}^{(j)} \le \exp\left(D_7 \{ \log(1/\delta) \}^2 \right),
$$
\n(B.28)

where $D_7 = D_7(C_{N,4}, C_{N,5}, D_3, D_6)$. Note that $|\sigma_t|, |g_y^{(j)}(x, t)| \leq {\log(1/\delta)}^{\tau_{\text{bd}}}$ for $|x| \leq 1$ and $\delta \leq t \leq$ $(2\overline{\tau})^{-1}$ with small enough δ , due to the [\(B.1\)](#page-41-2) and [\(B.25\)](#page-52-0). Then,

$$
\left| \sigma_t g_{\mathbf{y}}^{(j)}(x,t) - f_{\text{mult}}^{(2)} \left(f_{\mathbf{y}}^{(j)}(x,t), f_{\sigma}(t) \right) \right| \le \delta + 2D_4 \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + \frac{3\tau_{\text{bd}}}{2}}
$$

for $x \in [0,1]$ and $\delta \le t \le (2\overline{\tau})^{-1}$, where the inequality holds by combining [\(B.19\)](#page-50-0) and [\(B.25\)](#page-52-0) with [\(B.21\)](#page-51-2). Also, $|x + \sigma_t g_y^{(j)}(x, t)|, |\mu_t^{-1}| \leq {\log(1/\delta)}^{\tau_{\text{bd}}}$ for $|x| \leq 1$ and $\delta \leq t \leq (2\overline{\tau})^{-1}$ with small enough δ, due to the $(B.1)$ and $(B.25)$. Combining $(B.27)$ and $(B.21)$ with the last display, we have

$$
\begin{aligned}\n\left| \frac{x + \sigma_t g_y^{(j)}(x, t)}{\mu_t} - \tilde{f}_y^{(j)}(x, t) \right| \\
&\leq \delta + 2\{\log(1/\delta)\}^{\tau_{\text{bd}}} \max\left(17\delta, \delta + 2D_4 \sqrt{\delta} \{\log(1/\delta)\}^{\tau_{\text{bd}}^2 + \frac{3\tau_{\text{bd}}}{2}}\right) \\
&\leq D_8 \sqrt{\delta} \{\log(1/\delta)\}^{\tau_{\text{bd}}^2 + \frac{5\tau_{\text{bd}}}{2}}\n\end{aligned} \tag{B.29}
$$

for $x \in [0,1]$ and $\delta \le t \le (2\overline{\tau})^{-1}$, where $D_8 = 35 + 2D_4$. Consider functions $g_{\rm w}^{(1)}, \ldots, g_{\rm w}^{(m)} : [0,\infty) \to \mathbb{R}$ and $f_{\mathbf{w}}^{(1)}, \ldots, f_{\mathbf{w}}^{(m)} : \mathbb{R} \to \mathbb{R}$ such that

$$
g_{\rm w}^{(j)}(t) = 2\sqrt{2\tau_{\rm tail}} n_\beta \widetilde{w}_{j-\lfloor \frac{j}{n_\beta}\rfloor} {\log(1/\sigma_t)}^{\tau_{\rm bd}+\frac{1}{2}}
$$

for $t \in [0, \infty)$ and

$$
f_{\rm w}^{(j)}(t) = 2\sqrt{2\tau_{\rm tail}} n_\beta \tilde{w}_{j-\lfloor \frac{j}{n_\beta} \rfloor} f_{\rm pow}^{(\tau_{\rm bd} + \frac{1}{2})} \left(f_{\rm log} \left(f_\sigma(t)\right)\right)
$$

for $t \in \mathbb{R}$. By [\(B.24\)](#page-51-4), we have

$$
\left| g_{\mathbf{w}}^{(j)}(t) - f_{\mathbf{w}}^{(j)}(t) \right| \le D_9 \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\rm bd}^2 - \frac{\tau_{\rm bd}}{2}},\tag{B.30}
$$

for $\delta \le t \le (2\overline{\tau})^{-1}$, where $D_9 = 2\sqrt{2\tau_{\text{tail}}}D_2n_\beta \max_{j\in[n_\beta]}\widetilde{w}_j$. Also, Lemma [A.4](#page-29-0) and Lemma [A.5](#page-29-1) implies that $f_{\mathbf{w}}^{(j)} \in \mathcal{F}_{\text{NN}}(L_{\mathbf{w}}, \mathbf{d}_{\mathbf{w}}, s_{\mathbf{w}}, M_{\mathbf{w}}^{(j)})$ for $j \in [m]$ with

$$
L_{\rm w} \leq L_{\rm mult}^{(2)} + L_{\rm pl\sigma} \vee L_{\mu} \leq D_{10} \{ \log(1/\delta) \}^2 \log \log(1/\delta),
$$

\n
$$
\|\mathbf{d}_{\rm w}\|_{\infty} \leq 4 \left\{ \|\mathbf{d}_{\rm mult}^{(2)}\|_{\infty} \vee (\|\mathbf{d}_{\rm pl\sigma}\|_{\infty} \vee \|\mathbf{d}_{\mu}\|_{\infty}) \right\} \leq D_{10} \{\log(1/\delta) \}^3,
$$

\n
$$
s_{\rm w} \leq 4 \left\{ s_{\rm mult}^{(2)} + 2 \left(L_{\rm pl\sigma} \vee L_{\mu} \right) + s_{\rm pl\sigma} + s_{\mu} \right\} \leq D_{10} \{ \log(1/\delta) \}^5 \log \log(1/\delta)
$$

\n
$$
M_{\rm w}^{(j)} \leq \max \left\{ 2\sqrt{2\tau_{\rm tail}} n_{\beta} \max_{j \in [n_{\beta}]} \tilde{w}_{j}, M_{\rm mult}^{(2)}, M_{\rm pl\sigma}, M_{\mu}, 1 \right\} \leq \exp \left(D_{10} \{ \log(1/\delta) \}^2 \right),
$$
\n(B.31)

where $D_{10} = D_{10}(\beta, \tau_{\text{tail}}, C_{N,1}, C_{N,4}, D_5)$. Consider a function $\mathbf{f}_{\text{pre}} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}^{3mD}$ such that

$$
\begin{aligned} (\mathbf{f}_{\text{pre}}(\mathbf{x},t))_{3m(i-1)+3(j-1)+1} &= f_{\mathbf{y}}^{(j)}(x_i,t), \quad (\mathbf{f}_{\text{pre}}(\mathbf{x},t))_{3m(i-1)+3(j-1)+2} = \widetilde{f}_{\mathbf{y}}^{(j)}(x_i,t),\\ (\mathbf{f}_{\text{pre}}(\mathbf{x},t))_{3m(i-1)+3(j-1)+3} &= f_{\mathbf{w}}^{(j)}(t), \quad i \in [D], j \in [m] \end{aligned}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Combining Lemma [A.5](#page-29-1) with [\(B.26\)](#page-52-2), [\(B.28\)](#page-53-0) and [\(B.31\)](#page-53-1), we have $\mathbf{f}_{\text{pre}} \in$ $\mathcal{F}_{\text{NN}}(L_{\text{pre}}, \mathbf{d}_{\text{pre}}, s_{\text{pre}}, M_{\text{pre}})$ with

$$
L_{\text{pre}} \leq \max\left(L_{\mathbf{x}}, \widetilde{L}_{\mathbf{x}}, L_{\mathbf{w}}\right) \leq D_{11}\{\log(1/\delta)\}^{2} \log \log(1/\delta),
$$

$$
\|\mathbf{d}_{\text{pre}}\|_{\infty} \leq 2\left(mD\|\mathbf{d}_{\mathbf{x}}\|_{\infty} + mD\|\widetilde{\mathbf{d}}_{\mathbf{x}}\|_{\infty} + mD\|\mathbf{d}_{\mathbf{w}}\|_{\infty}\right) \leq D_{11}m\{\log(1/\delta)\}^{3},
$$

$$
s_{\text{pre}} \leq 2\left\{3mD\max\left(L_{\mathbf{x}}, \widetilde{L}_{\mathbf{x}}, L_{\mathbf{w}}\right) + mDs_{\mathbf{x}} + mD\widetilde{s}_{\mathbf{x}} + mDs_{\mathbf{w}}\right\}
$$

$$
\leq D_{11}m\{\log(1/\delta)\}^{5} \log \log(1/\delta),
$$

$$
M_{\text{pre}} \leq \max\left(M_{\mathbf{x}}, \widetilde{M}_{\mathbf{x}}, M_{\mathbf{w}}, 1\right) \leq \exp\left(D_{11}\{\log(1/\delta)\}^{2}\right),
$$

where $D_{11} = D_{11}(D, D_6, D_7, D_{10}).$

The assumption (S) implies that $p_0 = g_2 \circ \mathbf{g}_1$ for functions $\mathbf{g}_1 : [-1,1]^D \to [-K,K]^{\mathcal{|I|}}$ and g_2 : $[-K, K]^{\vert\mathcal{I}\vert} \to \mathbb{R}$, where $\mathbf{g}_1 = (g_{11}, \ldots, g_{1\vert\mathcal{I}\vert})$ with $g_{1i} \in \mathcal{H}^{\beta, K}([-1, 1]^{\vert I \vert}), I \in \mathcal{I}$ and $g_2(x_1, \ldots, x_{\vert\mathcal{I}\vert}) =$ $\prod_{i=1}^{\lvert \mathcal{I} \rvert} x_i$ for $x_1, \ldots, x_{\lvert \mathcal{I} \rvert} \in [-K, K]$. A simple calculation yields that $g_2 \in \mathcal{H}^{\gamma, \widetilde{K}}([-K, K]^{|\mathcal{I}|})$ with $\widetilde{K} = (2K)^{|\mathcal{I}|}$ for any $\gamma \geq |\mathcal{I}| + 1$. Since $|\mathcal{I}| \leq 2^D$, Lemma 5 of [Chae et al.](#page-22-2) [\(2023\)](#page-22-2) implies that there exists neural networks $f_{p_0} \in \mathcal{F}_{\text{NN}}(L_{p_0}, \mathbf{d}_{p_0}, s_{p_0}, M_{p_0})$ with

$$
L_{p_0} \le D_{12} \log m, \quad \|\mathbf{d}_{p_0}\|_{\infty} \le D_{12} m, \quad s_{p_0} \le D_{12} m \log m, \quad M_{p_0} \le 1
$$

such that $|p_0(\mathbf{x}) - f_{p_0}(\mathbf{x})| \leq m^{-\frac{\beta}{d}}$ for $\|\mathbf{x}\|_{\infty} \leq 1$, where $D_{12} = D_{12}(\beta, d, D, K)$. Since $p_0 \in$ $\mathcal{H}^{\beta,K}([-1,1]^D)$, we have $|p_0(\mathbf{x}) - p_0(\widetilde{\mathbf{x}})| \leq K D \|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty}^{\beta \wedge 1}$ for any $\mathbf{x}, \widetilde{\mathbf{x}} \in [-1,1]^D$. Let \widetilde{C}_2 be the constant in Lemma [B.3.](#page-39-0) Then, we have

$$
\left|\frac{x + \sigma_t g_y^{(j)}(x, t)}{\mu_t}\right| \le 1 - \frac{\{\log(1/\sigma_t)\}^{-\tau_{\text{bd}}}}{2} < 1
$$

for $|x| \leq \mu_t - \mu_t \{ \log(1/\sigma_t) \}^{-\tau_{\text{bd}}}$ and $\delta \leq t \leq \overline{\tau}^{-1}(\tilde{C}_2^2 \wedge 1/4)$. Combining with [\(B.29\)](#page-53-2), we have

$$
\left| p_0 \left(\frac{\mathbf{x} + \sigma_t \left(g_y^{(j_1)}(x_1, t), \dots, g_y^{(j_D)}(x_D, t) \right)^{\top}}{\mu_t} \right) - f_{p_0} \left(\tilde{f}_y^{(j_1)}(x_1, t), \dots, \tilde{f}_y^{(j_D)}(x_D, t) \right) \right|
$$
\n
$$
\leq KDD_8^{\beta \wedge 1} \delta^{\frac{\beta \wedge 1}{2}} \{ \log(1/\delta) \}^{(\beta \wedge 1)(\tau_{\text{bd}}^2 + \frac{5\tau_{\text{bd}}}{2})} + m^{-\frac{\beta}{d}} \stackrel{\text{def}}{=} \epsilon_{p_0}, \quad j_1, \dots, j_D \in [m]
$$
\n(B.32)

for $\|\mathbf{x}\|_{\infty} \leq \mu_t - \mu_t \{\log(1/\sigma_t)\}^{-\tau_{\text{bd}}}$ and $\delta \leq t \leq \overline{\tau}^{-1}(\widetilde{C}_2^2 \wedge 1/4)$. Let $C_{N,3}$ be the constant in Lemma [A.12.](#page-35-2) Then, there exists a neural network $f_{\text{exp}} \in \mathcal{F}(L_{\text{exp}}, \mathbf{d}_{\text{exp}}, s_{\text{exp}}, M_{\text{exp}})$ with

$$
L_{\text{exp}} \leq C_{N,3} \log(1/\delta) \log \log(1/\delta), \quad \|\mathbf{d}_{\text{exp}}\|_{\infty} \leq C_{N,3} \{\log(1/\delta)\}^3,
$$

$$
s_{\text{exp}} \leq C_{N,3} \{\log(1/\delta)\}^4, \quad M_{\text{exp}} \leq C_{N,3} \delta^{-1},
$$

such that $|e^{-x} - f_{\text{exp}}(\tilde{x})| \le \delta + |x - \tilde{x}|$ for any $x \ge 0$ and $\tilde{x} \in \mathbb{R}$. Consider a function $f_{\phi}: \mathbb{R}^D \to \mathbb{R}$ such that

$$
f_{\phi}(\mathbf{x}) = (2\pi)^{-\frac{D}{2}} f_{\exp}\left(-\sum_{i=1}^{D} \frac{f_{\text{pow}}^{(2)}(x_i)}{2}\right)
$$

for $\mathbf{x} = (x_1, \ldots, x_D) \in \mathbb{R}^D$. Then, Lemma [A.4,](#page-29-0) Lemma [A.5](#page-29-1) and Lemma [A.6](#page-30-0) imply that $f_\phi \in$ $\mathcal{F}_{\text{NN}}(L_{\phi}, \mathbf{d}_{\phi}, s_{\phi}, M_{\phi})$ with

$$
L_{\phi} \le D_{13} \log(1/\delta) \log \log(1/\delta), \quad \|\mathbf{d}_{\phi}\|_{\infty} \le D_{13} \{\log(1/\delta)\}^{3}
$$

$$
s_{\phi} \le D_{13} \{\log(1/\delta)\}^{4}, \quad M_{\phi} \le D_{13} \delta^{-1},
$$

where $D_{13} = D_{13}(D, C_{N,1}, C_{N,3})$. Combining [\(B.25\)](#page-52-0) with [\(B.22\)](#page-51-1), it follows that

$$
\left| \prod_{i=1}^{D} \phi \left(g_{y}^{(j_i)}(x_i, t) \right) - f_{\phi} \left(f_{y}^{(j_1)}(x_1, t), \dots, f_{y}^{(j_D)}(x_D, t) \right) \right|
$$
\n
$$
\leq (2\pi)^{-\frac{D}{2}} \left[\delta + \left| \sum_{i=1}^{D} \frac{\left\{ g_{y}^{(j_i)}(x_i, t) \right\}^2 - f_{\text{pow}}^{(2)} \left(f_{y}^{(j_i)}(x_i, t) \right)}{2} \right| \right]
$$
\n
$$
\leq (2\pi)^{-\frac{D}{2}} \left[\delta + \frac{1}{2} \sum_{i=1}^{D} \left(\delta + 2 \{ \log(1/\delta) \}^{\tau_{\text{bd}}} \left| g_{y}^{(j_i)}(x_i, t) - f_{y}^{(j_i)}(x_i, t) \right| \right) \right]
$$
\n
$$
\leq D_{14} \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + \frac{3\tau_{\text{bd}}}{2}} \stackrel{\text{def}}{=} \epsilon_{\phi}, \quad j_1, \dots, j_D \in [m]
$$
\n(B.33)

for $\|\mathbf{x}\|_{\infty} \leq 1$ and $\delta \leq t \leq (2\overline{\tau})^{-1}$, where $D_{14} = D_{14}(D, D_4)$. Combining [\(B.30\)](#page-53-3) with [\(B.21\)](#page-51-2), we have

$$
\left| \prod_{i=1}^{D} g_{w}^{(j_i)}(t) - f_{\text{mult}}^{(D)} \left(f_{w}^{(j_1)}(t), \dots, f_{w}^{(j_D)}(t) \right) \right|
$$
\n
$$
\leq \delta + DD_{11} \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + (D - \frac{3}{2})\tau_{\text{bd}}} \leq D_{15} \sqrt{\delta} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + (D - \frac{3}{2})\tau_{\text{bd}}} \stackrel{\text{def}}{=} \epsilon_{w}
$$
\n(B.34)

for $\delta \leq t \leq (2\overline{\tau})^{-1}$ with small enough δ so that $|g_{\rm w}^{(j_i)}(t)| \leq {\log(1/\delta)}^{\tau_{\rm bd}}$ for each $j_i \in [m]$, where $D_{15} = 1 + DD_{11}.$

Note that $|p_0(\mathbf{x})| \le K$ for any $\|\mathbf{x}\|_{\infty} \le 1$, $|\phi(x)| \le 1/$ $\sqrt{2\pi}$ for any $x \in \mathbb{R}$ and $|g_{\mathbf{y}}^{(j)}(\mathbf{x},t)|, |g_{\mathbf{w}}^{(j)}(t)| \leq$ $D_{16}\{\log(1/\delta)\}^{\tau_{\rm bd}+\frac{1}{2}}$ for $j \in [m], ||\mathbf{x}||_{\infty} \leq 1, \delta \leq t \leq (4\overline{\tau})^{-1}$, where $D_{16} = 2\sqrt{2\tau_{\rm tail}}(2+n_{\beta})$ max $_{j\in [n_{\beta}]}$ \widetilde{w}_{j} . Let $\tilde{f}_{\text{mult}}^{(2)} \in \mathcal{F}_{\text{NN}}(\tilde{L}_{\text{mult}}^{(2)}, \tilde{\mathbf{d}}_{\text{mult}}^{(2)}, \tilde{s}_{\text{mult}}^{(2)}, \widetilde{M}_{\text{mult}}^{(2)})$ be the neural network in Lemma [A.9,](#page-32-2) with

$$
\widetilde{L}_{\text{mult}}^{(2)} \leq C_{N,1} \log 2 \left\{ (2D\tau_{\text{bd}} + D + 2) \log(1/\delta) + D \log D_{16} \right\},
$$

$$
\|\widetilde{\mathbf{d}}_{\text{mult}}^{(2)}\|_{\infty} \leq 96,
$$

$$
\widetilde{s}_{\text{mult}}^{(2)} \leq 2C_{N,1} \left\{ (D\tau_{\text{bd}} + D/2 + 1) \log(1/\delta) + D \log D_{16} \right\},
$$

$$
\widetilde{M}_{\text{mult}}^{(2)} = D_{16}^{2D} \left\{ \log(1/\delta) \right\}^{2D(\tau_{\text{bd}} + \frac{1}{2})}
$$

such that

$$
|\widetilde{f}_{\text{mult}}^{(2)}(\widetilde{\mathbf{x}}) - x_1 x_2| \le \delta + 2D_{16}^D \{ \log(1/\delta) \}^{D(\tau_{\text{bd}} + \frac{1}{2})} \widetilde{\epsilon}
$$

for all $\|\mathbf{x}\|_{\infty} \leq \{D_{11} \log(1/\delta)\}^{D(\tau_{\text{bd}} + \frac{1}{2})}, \tilde{\mathbf{x}} \in \mathbb{R}^2 \text{ with } \|\mathbf{x} - \tilde{\mathbf{x}}\|_{\infty} \leq \tilde{\epsilon}.$ Also, let

$$
\overline{f}_{\text{mult}}^{(3)} \in \mathcal{F}_{\text{NN}}(\overline{L}_{\text{mult}}^{(3)}, \overline{\mathbf{d}}_{\text{mult}}^{(3)}, \overline{s}_{\text{mult}}^{(3)}, \overline{M}_{\text{mult}}^{(3)})
$$

be the neural network in Lemma [A.9](#page-32-2) with

$$
\overline{L}_{\text{mult}}^{(3)} \leq C_{N,1} \log 3 \left\{ \log(1/\delta) + 3 \log(K \vee 1) \right\}, \quad \|\overline{\mathbf{d}}_{\text{mult}}^{(3)}\|_{\infty} \leq 144, \n\overline{s}_{\text{mult}}^{(3)} \leq 3C_{N,1} \left\{ \log(1/\delta) + \log(K \vee 1) \right\}, \quad \overline{M}_{\text{mult}}^{(3)} = K^3 \vee 1
$$

such that

$$
|\overline{f}_{\text{mult}}^{(3)}(\widetilde{\mathbf{x}}) - x_1 x_2 x_3| \le \delta + 3(K^2 \vee 1)\widetilde{\epsilon}
$$

for all $\|\mathbf{x}\|_{\infty} \leq K \vee 1$, $\widetilde{\mathbf{x}} \in \mathbb{R}^3$ with $\|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty} \leq \widetilde{\epsilon}$. Consider functions $\mathbf{f}_{\text{main}} : \mathbb{R}^D \times \mathbb{R}^D \times \mathbb{R}^D \to \mathbb{R}^{D+1}$ such that

$$
\begin{aligned} \left(\mathbf{f}_{\text{main}}(\mathbf{x}, \widetilde{\mathbf{x}}, \mathbf{w})\right)_i &= \widetilde{f}_{\text{mult}}^{(2)}\left(\overline{f}_{\text{mult}}^{(3)}\left(x_i, f_{p_0}(\widetilde{\mathbf{x}}), f_{\phi}(\mathbf{x})\right), f_{\text{mult}}^{(D)}(\mathbf{w})\right), \quad i \in [D],\\ \left(\mathbf{f}_{\text{main}}(\mathbf{x}, \widetilde{\mathbf{x}}, \mathbf{w})\right)_{D+1} &= \widetilde{f}_{\text{mult}}^{(2)}\left(\overline{f}_{\text{mult}}^{(3)}\left(1, f_{p_0}(\widetilde{\mathbf{x}}), f_{\phi}(\mathbf{x})\right), f_{\text{mult}}^{(D)}(\mathbf{w})\right) \end{aligned}
$$

for $\mathbf{x} = (x_1, \dots, x_D) \in \mathbb{R}^D$ and $\widetilde{\mathbf{x}}, \mathbf{w} \in \mathbb{R}^D$. Lemma [A.4,](#page-29-0) Lemma [A.5](#page-29-1) and Lemma [A.7](#page-30-1) implies that $\mathbf{f}_{\text{main}} \in \mathcal{F}_{\text{NN}}(L_{\text{main}}, \mathbf{d}_{\text{main}}, s_{\text{main}}, M_{\text{main}})$ with

$$
L_{\text{main}} \le D_{17} \left[\log m + \log(1/\delta) \log \log(1/\delta) \right], \quad \|\mathbf{d}_{\text{main}}\|_{\infty} \le D_{17} \left[m + \{\log(1/\delta)\}^{3} \right]
$$

$$
s_{\text{main}} \le D_{17} \left[m \log m + \{\log(1/\delta)\}^{4} \right], \quad M_{\text{main}} \le D_{17} \delta^{-1},
$$

where $D_{17} = D_{17}(D, K, C_{N,1}, D_{10}, D_{12}, D_{13}, D_{16})$. For $j_1, \ldots, j_D \in [m]$, consider a function $\mathbf{f}^{(j_1, \ldots, j_D)}$: $\mathbb{R}^D \times \mathbb{R} \to \mathbb{R}^{3D}$ such that

$$
\mathbf{f}^{(j_1,\ldots,j_D)}(\mathbf{x},t) = (f_{y}^{(j_1)}(x_1,t),\ldots,f_{y}^{(j_D)}(x_D,t),\tilde{f}_{y}^{(j_1)}(x_1,t),\ldots,\tilde{f}_{y}^{(j_D)}(x_D,t),f_{w}^{(j_1)}(t),\ldots,f_{w}^{(j_D)}(t))^\top.
$$

By $(B.32)$, $(B.33)$ and $(B.34)$, we have

$$
\left| p_0 \left(\frac{\mathbf{x} + \sigma_t \left(g_y^{(j_1)}(x_1, t), \dots, g_y^{(j_D)}(x_D, t) \right)^{\top}}{\mu_t} \right) \prod_{i=1}^D \phi \left(g_y^{(j_i)}(x_i, t) \right) g_w^{(j_i)}(t) - \left(\mathbf{f}_{\text{main}} \left(\mathbf{f}^{(j_1, \dots, j_D)}(\mathbf{x}, t) \right) \right)_{D+1} \right|
$$

$$
\leq \delta + 2D_{16}^D \{ \log(1/\delta) \}^{D(\tau_{\text{bd}} + \frac{1}{2})} \max \left\{ \delta + 3(K^2 \vee 1) \left(\epsilon_{p_0} \vee \epsilon_{\phi} \right), \epsilon_{w} \right\}
$$

$$
\leq D_{18} \{ \log(1/\delta) \}^{D(\tau_{\text{bd}} + \frac{1}{2})} \left[m^{-\frac{\beta}{d}} + \delta^{\frac{\beta \wedge 1}{2}} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + (D + \frac{3}{2}) \tau_{\text{bd}}} \right]
$$

and

$$
\begin{split}\n&\left| g_{\mathbf{y}}^{(j_k)}(x_k, t) p_0 \left(\frac{\mathbf{x} + \sigma_t \left(g_{\mathbf{y}}^{(j_1)}(x_1, t), \ldots, g_{\mathbf{y}}^{(j_D)}(x_D, t) \right)^{\top}}{\mu_t} \right) \prod_{i=1}^D \phi \left(g_{\mathbf{y}}^{(j_i)}(x_i, t) \right) g_{\mathbf{w}}^{(j_i)}(t) \right. \\
&\left. - \left(\mathbf{f}_{\text{main}} \left(\mathbf{f}^{(j_1, \ldots, j_D)}(\mathbf{x}, t) \right) \right)_k \right| \\
&\leq \delta + 2D_{16}^D \{ \log(1/\delta) \}^{D(\tau_{\text{bd}} + \frac{1}{2})} \max \left\{ \delta + 3(K^2 \vee 1) \left(\epsilon_{p_0} \vee \epsilon_{\phi} \right), \epsilon_{\mathbf{w}} \right\} \\
&\leq D_{18} \{ \log(1/\delta) \}^{D(\tau_{\text{bd}} + \frac{1}{2})} \left[m^{-\frac{\beta}{d}} + \delta^{\frac{\beta \wedge 1}{2}} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + (D + \frac{3}{2}) \tau_{\text{bd}}} \right], \quad k \in [D], j_1, \ldots, j_D \in [m] \\
\text{for } ||\mathbf{x}||_{\infty} \leq \mu_t - \mu_t \{ \log(1/\sigma_t) \}^{-\tau_{\text{bd}}} \text{ and } \delta \leq t \leq \overline{\tau}^{-1} (\widetilde{C}_2^2 \wedge 1/4), \text{ where}\n\end{split}
$$

$$
D_{18} = D_{18}(\beta, D, K, D_8, D_{14}, D_{15}, D_{16}).
$$

Let \widetilde{C}_1 be the constant in Lemma [B.3.](#page-39-0) It follows that

$$
\mu_t^D \left\| \frac{1}{m^D} \sum_{j_1, \dots, j_D = 1}^m \mathbf{f}_{\text{main}} \left(\mathbf{f}^{(j_1, \dots, j_D)}(\mathbf{x}, t) \right) - \left(\frac{\sigma_t \nabla p_t(\mathbf{x})}{p_t(\mathbf{x})} \right) \right\|_{\infty}
$$

\n
$$
\leq \widetilde{C}_1 K \{ \log(1/\sigma_t) \}^{(\tau_{\text{bd}} + \frac{1}{2})(D-1)} \left(\sigma_t^{\tau_{\text{tail}}} + m^{-\beta} \{ \log(1/\sigma_t) \}^{(\tau_{\text{bd}} + \frac{1}{2})\beta} \right)
$$

\n
$$
+ D_{18} \{ \log(1/\delta) \}^{D(\tau_{\text{bd}} + \frac{1}{2})} \left[m^{-\frac{\beta}{d}} + \delta^{\frac{\beta \wedge 1}{2}} \{ \log(1/\delta) \}^{\tau_{\text{bd}}^2 + (D + \frac{3}{2})\tau_{\text{bd}}} \right]
$$

for $\delta \leq t \leq \overline{\tau}^{-1}(\widetilde{C}_{2}^{2} \wedge 1/2)$. Let $m = n_{\beta}[\widetilde{m}]$ and $\delta = \widetilde{m}^{-\tau_{\min}}$ with large enough $\widetilde{m} > 0$. Since $\tau_{\min} \ge \frac{4\beta}{d(\beta\wedge 1)}$, we have $\delta^{\frac{\beta\wedge 1}{2}}\{\log(1/\delta)\}^{\tau_{\rm bd}^2 + (D+\frac{3}{2})\tau_{\rm bd}} \le \tilde{m}^{-\frac{\beta}{d}}$ for large enough \tilde{m} . Then,

$$
\left\| \frac{1}{m^D} \sum_{j_1, \dots, j_D = 1}^m \mathbf{f}_{\text{main}} \left(\mathbf{f}^{(j_1, \dots, j_D)}(\mathbf{x}, t) \right) - \left(\frac{\sigma_t \nabla p_t(\mathbf{x})}{p_t(\mathbf{x})} \right) \right\|_{\infty} \le D_{19} \left(\log \widetilde{m} \right)^{(\tau_{\text{bd}} + \frac{1}{2})(D-1)} \left\{ t^{\frac{\tau_{\text{tail}}}{2}} + \widetilde{m}^{-\frac{\beta}{d}} \left(\log \widetilde{m} \right)^{(\tau_{\text{bd}} + \frac{1}{2})(\beta \vee 1)} \right\},\tag{B.35}
$$

where $D_{19} = D_{19}(\beta, K, D, d, \tau_{\text{bd}}, \tau_{\text{tail}}, \tau_{\text{min}}, \overline{\tau}, \underline{\tau}, \widetilde{C}_1, D_{18}).$

Recall that $f_{\text{pre}} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}^{3mD}$ and $f_{\text{main}} : \mathbb{R}^{3D} \to \mathbb{R}^{2D+1}$. Consider a function $\widetilde{f}_{\text{pre}} : \mathbb{R}^D \times \mathbb{R} \to$ \mathbb{R}^{6mD} such that

$$
\left(\widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)\right)_{1:3mD} = \rho\left(\mathbf{f}_{\text{pre}}(\mathbf{x},t)\right), \quad \left(\widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)\right)_{3mD+1:6mD} = \rho\left(-\mathbf{f}_{\text{pre}}(\mathbf{x},t)\right)
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$, where $\mathbf{z}_{n_1:n_2} \in \mathbb{R}^{n_2-n_1+1}$ denotes the subvector of \mathbf{z} from n_1 -th component to n₂-th component. Let $W_1, \mathbf{b}_1, \ldots, W_{L_{\text{main}}}$, $\mathbf{b}_{L_{\text{main}}}$ be the weight matrices and shift vectors of the neural network f_{main} , where $d_{\text{main}} = (d_1, \ldots, d_{L_{\text{main}}+1})$ with $d_1 = 3D$, $d_{L_{\text{main}}+1} = 2D + 1$ and $W_l \in$ $\mathbb{R}^{d_{l+1} \times d_l}$, $\mathbf{b}_l \in \mathbb{R}^{d_{l+1}}$ for $l \in [L_{\text{main}}]$. Let $\widetilde{\mathbf{d}} = (\widetilde{d}_1, \dots, \widetilde{d}_{L_{\text{main}}+2})$, where $\widetilde{d}_1 = 2md_1$, $\widetilde{d}_l = m^D d_l$ for $l \in \{2,\ldots,L_{\text{main}}\}, \tilde{d}_{L_{\text{main}}+1} = 2m^D d_{L_{\text{main}}+1}$ and $\tilde{d}_{L_{\text{main}}+2} = d_{L_{\text{main}}+1}$. Let

$$
\widetilde{W}_1, \widetilde{\mathbf{b}}_1, \ldots, \widetilde{W}_{L_{\text{main}}}, \widetilde{\mathbf{b}}_{L_{\text{main}}}, \text{ and } \widetilde{W}_{L_{\text{main}}+1}
$$

be the block-sparse matrices and vectors, defined as

$$
\widetilde{W}_{1} = \begin{pmatrix} W_{1} & -W_{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \end{pmatrix} \in \mathbb{R}^{\widetilde{d}_{2} \times \widetilde{d}_{1}}, \qquad \widetilde{\mathbf{b}}_{1} = \begin{pmatrix} \mathbf{b}_{1} \\ \mathbf{0} \end{pmatrix} \in \mathbb{R}^{\widetilde{d}_{2}},
$$
\n
$$
\widetilde{W}_{l} = \begin{pmatrix} W_{l} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \in \mathbb{R}^{\widetilde{d}_{l+1} \times \widetilde{d}_{l}}, \qquad \widetilde{\mathbf{b}}_{l} = \begin{pmatrix} \mathbf{b}_{l} \\ \mathbf{0} \end{pmatrix} \in \mathbb{R}^{\widetilde{d}_{l+1}}, \quad \text{for } l \in \{2, \ldots, L_{\text{main}} - 1\},
$$
\n
$$
\widetilde{W}_{L_{\text{main}}} = \begin{pmatrix} W_{L_{\text{main}}} & \mathbf{0} \\ -W_{L_{\text{main}}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \in \mathbb{R}^{\widetilde{d}_{L_{\text{main}}} + 1 \times \widetilde{d}_{L_{\text{main}}}}, \qquad \widetilde{\mathbf{b}}_{L_{\text{main}}} = \begin{pmatrix} \mathbf{b}_{L_{\text{main}}} \\ -\mathbf{b}_{L_{\text{main}}} \\ \mathbf{0} \end{pmatrix} \in \mathbb{R}^{\widetilde{d}_{L_{\text{main}}} + 1}
$$

and

$$
\widetilde{W}_{L_{\text{main}}+1} \in \mathbb{R}^{\widetilde{d}_{L_{\text{main}}+2} \times \widetilde{d}_{L_{\text{main}}+1}},
$$
\n
$$
\left(\widetilde{W}_{L_{\text{main}}+1}\right)_{ij} = \begin{cases}\n1 & \text{if } i \in [\widetilde{d}_{L_{\text{main}}+2}], j = 1, \\
-1 & \text{if } i \in [\widetilde{d}_{L_{\text{main}}+2}], j = m^D d_{L_{\text{main}}+1} + 1, \\
0 & \text{otherwise.} \n\end{cases}
$$

For any $j_1, \ldots, j_D \in [m]$, consider a $\tilde{d}_1 \times \tilde{d}_1$ permutation matrix $Q_1^{(j_1, \ldots, j_D)}$ such that

$$
\left(Q_1^{(j_1,\ldots,j_D)}\mathbf{y}\right)_{D(k-1)+i} = y_{3m(i-1)+3(j_i-1)+k},
$$
\n
$$
\left(Q_1^{(j_1,\ldots,j_D)}\mathbf{y}\right)_{D(k-1)+i+3D} = y_{3m(i-1)+3(j_i-1)+k+3mD}, \quad k \in \{1,2,3\}, i \in [D],
$$

where $\mathbf{y} = (y_1, \dots, y_{\tilde{d}_1})^\top \in \mathbb{R}^{\tilde{d}_1}$. Then, the first 2d₁-components of $Q_1^{(j_1, \dots, j_D)} \tilde{\mathbf{f}}_{\text{pre}}(\mathbf{x}, t)$ are $\left(\rho\left(\mathbf{f}^{(j_1,...,j_D)}(\mathbf{x},t)\right)^\top,\rho\left(-\mathbf{f}^{(j_1,...,j_D)}(\mathbf{x},t)\right)^\top\right).$

Since $x \vee 0 - (-x \vee 0) = x$ for any $x \in \mathbb{R}$, we have

$$
\left(\widetilde{W}_1 Q_1^{(j_1,\ldots,j_D)}\widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)+\widetilde{\mathbf{b}}_1\right)_{1:d_2} = W_1 \mathbf{f}^{(j_1,\ldots,j_D)}(\mathbf{x},t)+\mathbf{b}_1,
$$

$$
\left(\widetilde{W}_1 Q_1^{(j_1,\ldots,j_D)}\widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)+\widetilde{\mathbf{b}}_1\right)_{d_2+1:\widetilde{d}_2} = \mathbf{0}, \quad j_1,\ldots,j_D \in [m]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. For any $j_1, \ldots, j_D \in [m]$ and $l \in [L_{\text{main}} - 1]$, consider a $\widetilde{d}_{l+1} \times \widetilde{d}_{l+1}$ permutation matrix $R_l^{(j_1,...,j_D)}$ $\binom{(J_1,\ldots,J_D)}{l}$ such that

$$
\left(R_l^{(j_1,\ldots,j_D)}\mathbf{y}\right)_{d_{l+1}\sum_{i=1}^D m^{i-1}(j_i-1)+1:d_{l+1}\sum_{i=1}^D m^{i-1}(j_i-1)+d_{l+1}}=\mathbf{y}_{1:d_{l+1}},\quad \mathbf{y}\in\mathbb{R}^{\widetilde{d}_{l+1}}.
$$

Combining with the last two display, we have

$$
\left(R_1^{(j_1,...,j_D)}\left(\widetilde{W}_1Q_1^{(j_1,...,j_D)}\widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)+\widetilde{\mathbf{b}}_1\right)\right)_{d_2\sum_{i=1}^D m^{i-1}(j_i-1)+1:d_2\sum_{i=1}^D m^{i-1}(j_i-1)+d_2}
$$
\n
$$
=W_1\mathbf{f}^{(j_1,...,j_D)}(\mathbf{x},t)+\mathbf{b}_1,
$$
\n
$$
\left(R_1^{(j_1,...,j_D)}\left(\widetilde{W}_1Q_1^{(j_1,...,j_D)}\widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)+\widetilde{\mathbf{b}}_1\right)\right)_{1:d_2\sum_{i=1}^D m^{i-1}(j_i-1)}=\mathbf{0},
$$
\n
$$
\left(R_1^{(j_1,...,j_D)}\left(\widetilde{W}_1Q_1^{(j_1,...,j_D)}\widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)+\widetilde{\mathbf{b}}_1\right)\right)_{d_2\sum_{i=1}^D m^{i-1}(j_i-1)+d_2+1:\widetilde{d}_2}=\mathbf{0}, \quad j_1,\ldots,j_D\in[m]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Therefore,

$$
\left(\sum_{j_1,\dots,j_D=1}^m R_1^{(j_1,\dots,j_D)}\left(\widetilde{W}_1 Q_1^{(j_1,\dots,j_D)}\widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)+\widetilde{\mathbf{b}}_1\right)\right)_{d_2 \sum_{i=1}^D m^{i-1}(\widetilde{j}_i-1)+1:d_2 \sum_{i=1}^D m^{i-1}(\widetilde{j}_i-1)+d_2}
$$
\n
$$
= W_1 \mathbf{f}^{(j_1,\dots,j_D)}(\mathbf{x},t)+\mathbf{b}_1, \quad \widetilde{j}_1,\dots,\widetilde{j}_D \in [m]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Also, for any $j_1, \ldots, j_D \in [m]$ and $l \in \{2, \ldots, L_{\text{main}}\}$, consider a $\tilde{d}_l \times \tilde{d}_l$ permutation matrix $Q_l^{(j_1,...,j_D)}$ $\binom{(J_1,\ldots,J_D)}{l}$ such that

$$
(Q_l^{(j_1,...,j_D)}\mathbf{y})_{1:d_l} = \mathbf{y}_{d_l \sum_{i=1}^D m^{i-1}(j_i-1)+1:d_l \sum_{i=1}^D m^{i-1}(j_i-1)+d_l}, \quad \mathbf{y} \in \mathbb{R}^{\widetilde{d}_l}.
$$

For any $j_1, \ldots, j_D \in [m]$, consider $\tilde{d}_{L_{\text{main}}} \times \tilde{d}_{L_{\text{main}}}$ permutation matrices $R_{L_{\text{main}}}^{(j_1, \ldots, j_D)}$ $(L_{\text{main}}^{(j_1,...,j_D)} \text{ and } Q_{L_{\text{main}}+1}^{(j_1,...,j_D)}$ $L_{\text{main}}+1$ such that

$$
\begin{split} \left(R^{(j_1,...,j_D)}_{L_{\text{main}}} \mathbf{y}\right)_{2d_{L_{\text{main}}}+1} &\sum_{i=1}^D m^{i-1}(j_i-1)+1:2d_{L_{\text{main}}}+1} \sum_{i=1}^D m^{i-1}(j_i-1)+2d_{L_{\text{main}}}+1} = \mathbf{y}_{1:2d_{L_{\text{main}}}+1},\\ \left(Q^{(j_1,...,j_D)}_{L_{\text{main}}} \mathbf{y}\right)_{1:2d_{L_{\text{main}}}+1} &= \mathbf{y}_{2d_{L_{\text{main}}}+1} \sum_{i=1}^D m^{i-1}(j_i-1)+1:2d_{L_{\text{main}}}+1} \sum_{i=1}^D m^{i-1}(j_i-1)+2d_{L_{\text{main}}}+1} \end{split}
$$

for $\mathbf{y} \in \mathbb{R}^{d_{L_{\text{main}}}+1}$. For $l \in [L_{\text{main}}]$, consider a function $\widetilde{\mathbf{f}}_l : \mathbb{R}^{\widetilde{d}_l} \to \mathbb{R}^{\widetilde{d}_{l+1}}$ such that

$$
\widetilde{\mathbf{f}}_l(\mathbf{x}) = \rho \left(\sum_{j_1,\dots,j_D=1}^m R_l^{(j_1,\dots,j_D)} \left(\widetilde{W}_l Q_l^{(j_1,\dots,j_D)} \mathbf{x} + \widetilde{\mathbf{b}}_l \right) \right).
$$

For any $\widetilde{j}_1, \ldots, \widetilde{j}_D \in [m]$, a simple calculation yields that

$$
\left(\widetilde{\mathbf{f}}_{L_{\text{main}}-1} \circ \cdots \circ \widetilde{\mathbf{f}}_1 \circ \widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)\right)_{d_{L_{\text{main}}}\sum_{i=1}^D m^{i-1}(\widetilde{j}_i-1)+1:d_{L_{\text{main}}}\sum_{i=1}^D m^{i-1}(\widetilde{j}_i-1)+d_{L_{\text{main}}}}}
$$
\n
$$
= \rho \left(W_{L_{\text{main}}-1} \cdot + \mathbf{b}_{L_{\text{main}}-1}\right) \circ \cdots \circ \rho \left(W_1 \cdot + \mathbf{b}_1\right) \circ \mathbf{f}^{(j_1,\dots,j_D)}(\mathbf{x},t),
$$

and

$$
\begin{split}\n\left(\tilde{\mathbf{f}}_{L_{\text{main}}}\circ\cdots\tilde{\mathbf{f}}_{1}\circ\tilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)\right)_{2d_{L_{\text{main}}+1}\sum_{i=1}^{D}m^{i-1}(\tilde{j}_{i}-1)+1:2d_{L_{\text{main}}+1}\sum_{i=1}^{D}m^{i-1}(\tilde{j}_{i}-1)+d_{L_{\text{main}}+1}} \\
&= \rho\left(W_{L_{\text{main}}}\cdot+\mathbf{b}_{L_{\text{main}}}\right)\circ\rho\left(W_{L_{\text{main}}-1}\cdot+\mathbf{b}_{L_{\text{main}}-1}\right)\circ\cdots\circ\rho\left(W_{1}\cdot+\mathbf{b}_{1}\right)\circ\mathbf{f}^{(j_{1},\ldots,j_{D})}(\mathbf{x},t), \\
\left(\tilde{\mathbf{f}}_{L_{\text{main}}}\circ\cdots\circ\tilde{\mathbf{f}}_{1}\circ\tilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t)\right)_{2d_{L_{\text{main}}+1}\sum_{i=1}^{D}m^{i-1}(\tilde{j}_{i}-1)+d_{L_{\text{main}}+1}+1:2d_{L_{\text{main}}+1}\sum_{i=1}^{D}m^{i-1}(\tilde{j}_{i}-1)+2d_{L_{\text{main}}+1}} \\
&= \rho\left(-W_{L_{\text{main}}}\cdot-\mathbf{b}_{L_{\text{main}}}\right)\circ\rho\left(W_{L_{\text{main}}-1}\cdot+\mathbf{b}_{L_{\text{main}}-1}\right)\circ\cdots\circ\rho\left(W_{1}\cdot+\mathbf{b}_{1}\right)\circ\mathbf{f}^{(j_{1},\ldots,j_{D})}(\mathbf{x},t)\n\end{split}
$$

for $\mathbf{x} \in \mathbb{R}^D, t \in \mathbb{R}$. Consider a vector-valued function $\mathbf{f} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}^{D+1}$ such that

$$
\mathbf{f}(\mathbf{x},t) = m^{-D} \rho \left(\sum_{j_1,\dots,j_D=1}^m \widetilde{W}_{L_{\text{main}}+1} Q_{L_{\text{main}}+1}^{(j_1,\dots,j_D)} \widetilde{\mathbf{f}}_{L_{\text{main}}} \circ \cdots \circ \widetilde{\mathbf{f}}_1 \circ \widetilde{\mathbf{f}}_{\text{pre}}(\mathbf{x},t) \right)
$$

Combining [\(B.35\)](#page-57-0) with the last two displays, we have

$$
\left\| \mathbf{f}(\mathbf{x},t) - \begin{pmatrix} \sigma_t \nabla p_t(\mathbf{x}) \\ p_t(\mathbf{x}) \end{pmatrix} \right\|_{\infty} \tag{B.36}
$$
\n
$$
\leq D_{19} \left(\log \widetilde{m} \right)^{(\tau_{\rm bd} + \frac{1}{2})(D-1)} \left\{ t^{\frac{\tau_{\rm tail}}{2}} + \widetilde{m}^{-\frac{\beta}{d}} \left(\log \widetilde{m} \right)^{(\tau_{\rm bd} + \frac{1}{2})(\beta \vee 1)} \right\}
$$

 $\text{for } ||\mathbf{x}||_{\infty} \leq \mu_t - \mu_t \{ \log(1/\sigma_t) \}^{\tau_{\text{bd}} + \frac{1}{2}} \text{ and } \widetilde{m}^{-\tau_{\text{min}}} \leq t \leq \overline{\tau}^{-1} (\widetilde{C}_2^2 \wedge 1/2).$ Note that $||\widetilde{W}||_0 = ||W_l||_0$ $\|\mathbf{b}_{l}\|_{0} = \|\mathbf{b}_{l}\|_{0}$ for $l \in \{2,\ldots,L_{\text{main}}-1\}$, $\|W_{L_{\text{main}}}\|_{0} = 2\|W_{L_{\text{main}}}\|_{0}$, $\|\mathbf{b}_{L_{\text{main}}}\|_{0} = 2\|\mathbf{b}_{L_{\text{main}}}\|_{0}$ and $\|\widetilde{W}_{L_{\text{main}}+1}\|_0 = 2d_{L_{\text{main}}+1}$. Let $L = L_{\text{pre}} + L_{\text{main}} + 2$ and $\mathbf{d} = (d_1, \ldots, d_{L+1}) \in \mathbb{N}^{L+1}$ with

$$
(d_1,\ldots,d_{L+1})=(d_{\text{pre}}^{(1)},\ldots,d_{\text{pre}}^{(L_{\text{pre}})},\widetilde{d}_1,\ldots,\widetilde{d}_{L_{\text{main}}+2},d_{L_{\text{main}}+1}),
$$

where $\mathbf{d}_{pre} = (d_{pre}^{(1)}, \ldots, d_{pre}^{(L_{pre}+1)})$. For $1 \leq i \leq L_{pre}$, let \mathcal{Q}_i and \mathcal{R}_i be the set $d_i \times d_i$ and $d_{i+1} \times d_{i+1}$ identity matrix, respectively. For $L_{pre} < i \leq L - 1$, let

$$
Q_i = \left\{ Q_{i-L_{\text{pre}}}^{(j_1,\dots,j_D)} : j_1,\dots,j_D \in [m] \right\} \text{ and } \mathcal{R}_i = \left\{ R_{i-L_{\text{pre}}}^{(j_1,\dots,j_D)} : j_1,\dots,j_D \in [m] \right\},\
$$

where $R_{L_{\text{main}}+1}^{(j_1,\ldots,j_D)}$ is a $\tilde{d}_{L_{\text{main}}+1} \times \tilde{d}_{L_{\text{main}}+1}$ identity matrix. Let $\mathbf{m} = (m_1,\ldots,m_{L-1})$ with $m_i = 1$ for $i \leq L_{\text{pre}}$ and $m_i = m^D$ for $i > L_{\text{pre}}$, and $\mathcal{P} = \{ \mathcal{Q}_l, \mathcal{R}_l \}_{l \in [L-1]}$. Then, $\mathbf{f} \in \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathbf{m}, \mathcal{P})$ with

$$
s \le 2s_{\text{pre}} + 2s_{\text{main}} + 4d_{L_{\text{main}}+1} \quad \text{and} \quad M = \max\left(M_{\text{pre}}, M_{\text{main}}, 1, m^{-D}\right).
$$

Recall that $m = n_\beta \lfloor \widetilde{m} \rfloor$ and $\delta = \widetilde{m}^{-\tau_{\min}}$. Thus,

$$
L \le D_{20} (\log \widetilde{m})^2 \log \log \widetilde{m}, \quad \|\mathbf{d}\|_{\infty} \le D_{20} \widetilde{m} (\log \widetilde{m})^3,
$$

$$
s \le D_{20} \widetilde{m} (\log \widetilde{m})^5 \log \log \widetilde{m}^{D+1}, \quad M \le \exp \left(D_{20} {\log \widetilde{m} }^2 \right),
$$

where $D_{20} = D_{20}(\tau_{\text{min}}, D_{11}, D_{17}, n_{\beta})$. Combining [\(B.36\)](#page-59-0) with the last display, the assertion follows by re-defining the constants. \Box

B.3 Proof of Proposition [B.2](#page-40-0)

Proof. Let $\tau_{tail} = 2 \vee \sqrt{(D+3)/(2e)}$. Given small enough $\delta > 0$ as described below, we have

$$
\int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_{\infty}\geq\tau_{\text{tail}}\sigma_t\sqrt{\log(1/\delta)}} p_0(\mathbf{y})\phi_{\sigma_t}(\mathbf{x}-\mu_t\mathbf{y})\mathrm{d}\mathbf{y}
$$
\n
$$
\leq K \int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_{\infty}\leq\tau_{\text{tail}}\sigma_t\sqrt{\log(1/\delta)}} \phi_{\sigma_t}(\mathbf{x}-\mu_t\mathbf{y})\mathrm{d}\mathbf{y} = K\mu_t^{-D} \int_{\|\mathbf{z}\|_{\infty}\geq\tau_{\text{tail}}\sqrt{\log(1/\delta)}} \phi_1(\mathbf{z})\mathrm{d}\mathbf{z}
$$
\n
$$
\leq K\mu_t^{-D} \sum_{i=1}^D \int_{|z_i|\geq\tau_{\text{tail}}\sqrt{\log(1/\delta)}} \phi(z_i)\mathrm{d}z_i \leq 2KD\mu_t^{-D} \delta^{\tau_{\text{tail}}^2/2}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t > 0$, where the last inequality holds by the tail probability of the standard normal distribution. Also, for $i \in [D]$, we have

$$
\left| \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \geq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y_i - x_i}{\sigma_t} \right) p_0(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} \right|
$$
\n
$$
\leq K \mu_t^{-D} \int_{\|\mathbf{z}\|_{\infty} \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)}} |z_i| \phi_1(\mathbf{z}) d\mathbf{z} \leq K \mu_t^{-D} \sum_{j=1}^D \int_{|z_j| \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)}} |z_i| \phi_1(\mathbf{z}) d\mathbf{z}
$$
\n
$$
= K \mu_t^{-D} \left\{ (D - 1) \mathbb{E}[|Z|] \int_{|z| \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)}} \phi(z) d\mathbf{z} + \int_{|z| \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)}} |z| \phi(z) d\mathbf{z} \right\}
$$
\n
$$
\leq K \mu_t^{-D} \left\{ 2(D - 1) \sqrt{2/\pi} \delta^{\tau_{\text{tail}}^2/2} + \sqrt{\mathbb{E}[Z^2]} \sqrt{\mathbb{P}\left(|Z| \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right)} \right\}
$$
\n
$$
\leq K \mu_t^{-D} \left\{ 2(D - 1) \sqrt{2/\pi} + \sqrt{2} \right\} \delta^{\tau_{\text{tail}}^2/4},
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t > 0$, where Z denotes the one-dimensional standard normal random variable and the second inequality holds by the Cauchy-Schwarz inequality. Since $\mu_t^{-D} \leq 2^D$ for $0 \leq t \leq (2\overline{\tau})^{-1}$,

$$
\left| p_t(\mathbf{x}) - \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} p_0(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}\mathbf{y} \right| \leq K D 2^{D+1} \delta^{\tau_{\text{tail}}^2/2},
$$
\n
$$
\left\| \sigma_t \nabla p_t(\mathbf{x}) - \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t \mathbf{y} - \mathbf{x}}{\sigma_t} \right) p_0(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}\mathbf{y} \right\|_{\infty} (B.37)
$$
\n
$$
\leq K 2^D \left\{ 2(D-1) \sqrt{2/\pi} + \sqrt{2} \right\} \delta^{\tau_{\text{tail}}^2/4}
$$

 $\text{for } \mathbf{x} \in \mathbb{R}^D \text{ and } 0 < t \leq (2\overline{\tau})^{-1}. \text{ For } 0 < t \leq (2\overline{\tau})^{-1} \text{ and } \mathbf{x}, \mathbf{y} \in \mathbb{R}^D \text{ with } ||\mathbf{x}||_{\infty} \geq \mu_t - \tau_{\mathbf{x}} \{ \log(1/\sigma_t) \}^{-\tau_{\text{bd}}}$ and $\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}$, we have

$$
\|\mathbf{y}\|_{\infty} \ge \frac{\|\mathbf{x}\|_{\infty} - \|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty}}{\mu_t} \ge 1 - \left(\frac{\tau_{\mathbf{x}} \{\log(1/\sigma_t)\}^{-\tau_{\rm bd}} + \tau_{\rm tail} \sigma_t \sqrt{\log(1/\delta)}}{\mu_t}\right)
$$

$$
\ge 1 - 2\left[\tau_{\mathbf{x}} \left\{\log(1/\sqrt{2\overline{\tau}t})\right\}^{-\tau_{\rm bd}} + \tau_{\rm tail} \sqrt{2\overline{\tau}t \log(1/\delta)}\right],
$$

where the last inequality holds by [\(B.1\)](#page-41-2). For $0 < t \leq \delta^{\tau_t}$ and small enough δ so that $\delta^{\tau_t} \leq (2\overline{\tau})^{-1}$, the last display is bounded by

$$
1 - 2\left[\tau_x \left\{ \log(1/\sqrt{2\overline{\tau} \delta^{\tau_t}}) \right\}^{-\tau_{\rm bd}} + \tau_{\rm tail} \sqrt{2\overline{\tau} \delta^{\tau_t}} \sqrt{\log(1/\delta)} \right].
$$

Moreover, the last display is lower bounded by $1 - 2D_1$ for small enough δ , where

$$
D_1 = \left(2\left[2\left\{\log(1/\delta)\right\}^{\widetilde{\tau}_{\text{bd}}} \vee 4\right] + 2\right)^{-1}.
$$

Then, $D_1 \leq (\log(1/\delta))^{-\tilde{\tau}_{\text{bd}}}/2) \wedge (1/4)$ and $D_1^{-1} \in 2\mathbb{N}$. Let $\mathbf{y}^{(1)}, \ldots, \mathbf{y}^{(D_2)} \in \mathbb{R}^D$ be distinct vectors satisfying that

$$
\left\{\mathbf{y}^{(1)},\ldots,\mathbf{y}^{(D_2)}\right\} = \left\{D_1(n_1,\ldots,n_D)^\top : n_i \in \mathbb{Z}, i \in [D]\right\} \cap \left\{\mathbf{y} \in \mathbb{R}^D : \|\mathbf{y}\|_{\infty} = 1 - D_1\right\},\
$$

where $D_2 = (2/D_1 - 1)^D - (2/D_1 - 3)^D$. Let $\mathcal{Y}_i = \{ \mathbf{y} \in \mathbb{R}^D : ||\mathbf{y} - \mathbf{y}^{(i)}||_{\infty} \le D_1 \}$ for $i \in [D_2]$. For any $i \in [D_2]$ and $\mathbf{y} \in \mathcal{Y}_i$, we have $1 - 2D_1 \le ||\mathbf{y}||_{\infty} \le 1$ and $1 - 2D_1 \le ||\mathbf{y}^{(i)}||_{\infty} \le 1$. Assume that the density p_0 satisfies

$$
\sup_{\mathbf{\alpha}\in\mathbb{N}^D} \sup_{1-\{\log(1/\delta)\}^{-\tilde{\tau}_{\text{bd}}}\leq\|\mathbf{x}\|_{\infty}\leq1} |(\mathbf{D}^{\mathbf{\alpha}}p_0(\mathbf{x}))|\leq K.
$$

With small enough δ so that $D_3 = \lfloor \log_{e/4}(1/\delta) \rfloor + 1 \geq 2$, Taylor's theorem for multivariate function implies that

$$
p_0(\mathbf{y}) = \sum_{0 \le k \, \le D_3} \frac{(D^k p_0)(\mathbf{y}^{(i)})}{\mathbf{k}!} (\mathbf{y} - \mathbf{y}^{(i)})^{\mathbf{k}} + \sum_{k = D_3} \frac{(D^k p_0)(\xi \mathbf{y}^{(i)} + (1 - \xi)\mathbf{y})}{\mathbf{k}!} (\mathbf{y} - \mathbf{y}^{(i)})^{\mathbf{k}}
$$

for a suitable $\xi \in [0,1]$ and $i \in [D_2], \mathbf{y} \in \mathcal{Y}_i$, where $(\mathbf{y} - \mathbf{y}^{(i)})^k = \prod_{j=1}^D (y_j - y_j^{(i)})$ $(j^{(i)}_j)^{k_j}$, $\mathbf{k}! = \prod_{j=1}^D k_j!$ and $k = ||\mathbf{k}||_1$. A simple calculation yields that

$$
\left| p_0(\mathbf{y}) - \sum_{0 \le k \,

$$
\le \sum_{k = D_3} K \prod_{j=1}^D \left(\frac{eD_1}{k_j}\right)^{k_j} \le K(D_3 + 1)^D \left(\frac{e}{4}\right)^{D_3} \le K3^D \delta \left\{\frac{\log(1/\delta)}{\log(e/4)}\right\}^D
$$
$$

for $\mathbf{y} \in \mathbb{R}^D$ and $i \in [D_2]$ because $k! \geq k^k e^{-k}$ for any $k \in \mathbb{Z}_{\geq 0}$ and $0 < D_1 \leq 1/4$. Since $\mathcal{Y}_1, \ldots, \mathcal{Y}_{D_2}$ are mutually disjoint except on a set of Lebesgue measure zero and $\bigcup_{i=1}^{D_2} \mathcal{Y}_i = \{ \mathbf{y} \in \mathbb{R}^D : 1 - 2D_1 \leq \mathcal{Y}_i \}$ $||\mathbf{y}||_{\infty} \leq 1$, we have

$$
\int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_{\infty}\leq\tau_{\text{tail}}\sigma_t\sqrt{\log(1/\delta)}} g(\mathbf{y}) d\mathbf{y} = \sum_{i=1}^{D_2} \int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_{\infty}\leq\tau_{\text{tail}}\sigma_t\sqrt{\log(1/\delta)}} g(\mathbf{y}) d\mathbf{y}
$$
\n
$$
= \int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_{\infty}\leq\tau_{\text{tail}}\sigma_t\sqrt{\log(1/\delta)}} \sum_{i=1}^{D_2} g(\mathbf{y}) \cdot 1\{\|\mathbf{y}-\mathbf{y}^{(i)}\|_{\infty}\leq D_1\} d\mathbf{y}
$$

for any continuous function $g : \mathbb{R}^D \to \mathbb{R}$, $\|\mathbf{x}\|_{\infty} \geq \mu_t - \tau_x {\log(1/\sigma_t)}^{-\tau_{\text{bd}}}$ and $0 < t \leq \delta^{\tau_t}$. Combining [\(B.37\)](#page-60-0) with the last two displays, we have

> $\overline{}$ $\overline{}$ \cdot $\overline{}$ $\overline{}$ $\overline{}$

$$
\left| p_t(\mathbf{x}) - \sum_{i=1}^{D_2} \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \le \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left\{ \sum_{0 \le k, < D_3} \frac{(D^k p_0)(\mathbf{y}^{(i)})}{k!} (\mathbf{y} - \mathbf{y}^{(i)})^k \right\} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}\mathbf{y} \right\}
$$
\n
$$
\le K D 2^{D+1} \delta^{\tau_{\text{tail}}^2/2} + K 3^D \delta \left\{ \frac{\log(1/\delta)}{\log(e/4)} \right\}^D
$$
\n
$$
\cdot \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \le \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \sum_{i=1}^{D_2} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \cdot 1 \{ \|\mathbf{y} - \mathbf{y}^{(i)}\|_{\infty} \le D_1 \} \mathrm{d}\mathbf{y}
$$
\n
$$
= K D 2^{D+1} \delta^{\tau_{\text{tail}}^2/2} + K 3^D \delta \left\{ \frac{\log(1/\delta)}{\log(e/4)} \right\}^D \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \le \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}\mathbf{y}
$$

for $\|\mathbf{x}\|_{\infty} \geq \mu_t - \tau_{\mathbf{x}} \{\log(1/\sigma_t)\}^{-\tau_{\text{bd}}}$ and $0 < t \leq \delta^{\tau_t}$. Moreover, the last display is bounded by

$$
KD2^{D+1}\delta^{\tau_{\text{tail}}^2/2} + K6^D \delta \left\{ \frac{\log(1/\delta)}{\log(e/4)} \right\}^D
$$
 (B.38)

because $\int_{\mathbb{R}^D} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} = \mu_t^{-D}$ and $\mu_t^{-D} \leq 2^D$ by [\(B.1\)](#page-41-2). Also, we have

$$
\left\|\sigma_{t}\nabla p_{t}(\mathbf{x}) - \sum_{i=1}^{D_{2}} \int_{\|\mathbf{x} - \mu_{t}\mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} \frac{\|\mathbf{y} - \mathbf{y}^{(i)}\|_{\infty} \leq D_{1}}{\|\mathbf{x} - \mu_{t}\mathbf{y}\|_{\infty} \leq D_{2}} \right\}
$$
\n
$$
\left\{\sum_{0 \leq k_{\cdot} < D_{3}} \frac{(D^{k}p_{0})(\mathbf{y}^{(i)})}{k!} \left(\frac{\mu_{t}\mathbf{y} - \mathbf{x}}{\sigma_{t}}\right) (\mathbf{y} - \mathbf{y}^{(i)})^{k} \right\} \phi_{\sigma_{t}}(\mathbf{x} - \mu_{t}\mathbf{y}) d\mathbf{y} \right\|_{\infty}
$$
\n
$$
\leq K 2^{D} \left\{ 2(D - 1)\sqrt{2/\pi} + \sqrt{2} \right\} \delta^{\tau_{\text{tail}}^{2}/4} + K 3^{D} \delta \left\{ \frac{\log(1/\delta)}{\log(e/4)} \right\}^{D}
$$
\n
$$
\cdot \int_{\|\mathbf{x} - \mu_{t}\mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} \sum_{i=1}^{D_{2}} \left\| \frac{\mu_{t}\mathbf{y} - \mathbf{x}}{\sigma_{t}} \right\|_{\infty} \phi_{\sigma_{t}}(\mathbf{x} - \mu_{t}\mathbf{y}) \cdot 1 \{\|\mathbf{y} - \mathbf{y}^{(i)}\|_{\infty} \leq D_{1}\} d\mathbf{y}
$$

for $\|\mathbf{x}\|_{\infty} \geq \mu_t - \tau_{\mathbf{x}} \{\log(1/\sigma_t)\}^{-\tau_{\text{bd}}}$ and $0 < t \leq \delta^{\tau_t}$. Moreover, the last display is bounded by

$$
K2^{D} \left\{ 2(D-1)\sqrt{2/\pi} + \sqrt{2} \right\} \delta^{\tau_{\text{tail}}^{2}/4} + 2K\tau_{\text{tail}}6^{D} \left\{ \log(e/4) \right\}^{-D} \delta \left\{ \log(1/\delta) \right\}^{D+1/2}
$$
(B.39)

because $\int_{\mathbb{R}^D} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} = \mu_t^{-D}$ and $\mu_t^{-D} \leq 2^D$ by [\(B.1\)](#page-41-2). For $x, y \in \mathbb{R}$ and $t > 0$, Taylor's theorem yields that

$$
\left|\exp\left(-\frac{(x-\mu_ty)^2}{2\sigma_t^2}\right)-\sum_{l=0}^{D_4-1}\frac{1}{l!}\left(-\frac{(x-\mu_ty)^2}{2\sigma_t^2}\right)^l\right|\leq \frac{1}{D_4!}\left(\frac{(x-\mu_ty)^2}{2\sigma_t^2}\right)^{D_4},
$$

where $D_4 = \lfloor 2e\tau_{\text{tail}}^2 \log(1/\delta) \rfloor + 1$ with small enough δ so that $D_4 \geq 1$. For $|x - \mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}$ and $t \geq 0$, the last display is further bounded by

$$
\left(\frac{e\tau_{\text{tail}}^2\log(1/\delta)}{D_4}\right)^{D_4} \le 2^{-D_4} \le \delta^{2e\tau_{\text{tail}}^2\log 2} \le \delta^{e\tau_{\text{tail}}^2},
$$

where the last inequality holds because $1/2 \le \log 2$ and $0 < \delta < 1$. Then,

$$
\left|\int_{|x-\mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y - x}{\sigma_t}\right)^m (y - \tilde{y})^k \exp\left(-\frac{(x - \mu_t y)^2}{2\sigma_t^2}\right) dy \right|
$$

$$
- \sum_{l=0}^{D_4 - 1} \frac{1}{l!} \int_{|x-\mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y - x}{\sigma_t}\right)^m (y - \tilde{y})^k \left(-\frac{(x - \mu_t y)^2}{2\sigma_t^2}\right)^l dy \right|
$$

$$
\leq \delta^{e\tau_{\text{tail}}^2} \int_{|x-\mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left|\frac{\mu_t y - x}{\sigma_t}\right|^m |y - \tilde{y}|^k dy
$$

$$
\leq \delta^{e\tau_{\text{tail}}^2} \int_{|y| \leq 1} D_1^k \left\{\tau_{\text{tail}} \sqrt{\log(1/\delta)}\right\}^m dy \leq \delta^{e\tau_{\text{tail}}^2} \left\{\tau_{\text{tail}} \sqrt{\log(1/\delta)}\right\}^m
$$

for any $x, \tilde{y} \in \mathbb{R}$ with $|\tilde{y}| = 1 - D_1, t \ge 0, k \in \mathbb{Z}_{\ge 0}$ and $m \in \{0, 1\}$, where the last inequality holds because $D_1 < 1$. Moreover,

$$
\left| \sum_{l=0}^{D_4-1} \frac{1}{l!} \int_{|x-\mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y - x}{\sigma_t} \right)^m (y - \tilde{y})^k \left(-\frac{(x - \mu_t y)^2}{2\sigma_t^2} \right)^l dy \right|
$$

$$
\leq \left(1 + \delta^{e\tau_{\text{tail}}^2} \right) \left\{ \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right\}^m
$$

for any $x, \widetilde{y} \in \mathbb{R}$ with $|\widetilde{y}| = 1 - D_1, t \ge 0, k \in \mathbb{Z}_{\ge 0}$ and $m \in \{0, 1\}$ because

$$
\left| \int_{|x-\mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y - x}{\sigma_t} \right)^m (y - \tilde{y})^k \exp\left(-\frac{(x - \mu_t y)^2}{2\sigma_t^2} \right) dy \right|
$$

$$
\leq D_1^k \{ \tau_{\text{tail}} \sqrt{\log(1/\delta)} \}^m \int_{|y| \leq 1} 1 dy \leq \left\{ \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right\}^m.
$$

Note that $\left|\prod_{j=1}^D x_j - \prod_{j=1}^D \widetilde{x}_j\right| \le DC^{D-1} \|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty}$ for any $\mathbf{x}, \widetilde{\mathbf{x}} \in [-C, C]^D$. Since the last two displays are bounded by $2\{\tau_{\text{tail}}\sqrt{\log(1/\delta)}\}^m$, we have

$$
\left| \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y_h - x_h}{\sigma_t} \right)^m (\mathbf{y} - \mathbf{y}^{(i)})^k \phi_{\sigma_t} (\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} - \right|
$$
\n
$$
\prod_{j=1}^D \sum_{l=0}^{D_4 - 1} \frac{1}{l! \sqrt{2\pi \sigma_t}} \int_{\|\mathbf{x}_j - \mu_t y_j\| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y_h - x_h}{\sigma_t} \right)^{m \cdot 1} (y_j - y_j^{(i)})^{k_j} \left(- \frac{(x_j - \mu_t y_j)^2}{2\sigma_t^2} \right)^l dy_j
$$
\n
$$
\leq (2\pi \sigma_t^2)^{-\frac{D}{2}} D 2^{D-1} \delta^{e \tau_{\text{tail}}^2} \left\{ \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right\}^{mD} \tag{B.40}
$$

for $\mathbf{x} \in \mathbb{R}^D, t \geq 0, h \in [D], i \in [D_2], j \in [D], \mathbf{k} \in \mathbb{Z}_{\geq 0}^D$ and $m \in \{0,1\}$ with small enough δ so that $\tau_{\text{tail}}\sqrt{\log(1/\delta)} > 1$. With $m = 0$ in the last display, the last integral satisfies that

$$
\frac{1}{l!\sqrt{2\pi\sigma_t}}\int_{|x_j - \mu_t y_j| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(y_j - y_j^{(i)}\right)^{k_j} \left(-\frac{(x_j - \mu_t y_j)^2}{2\sigma_t^2}\right)^l dy_j
$$
\n
$$
= \frac{1}{l!\sqrt{2\pi}} \int_{\substack{|z_j| \leq \tau_{\text{tail}} \sqrt{\log(1/\delta)} \\ |\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)}| \leq D_1}} \left(\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)}\right)^{k_j} z_j^{2l} (-2)^{-l} dz_j
$$
\n
$$
= \frac{(-2)^{-l}}{l!\sqrt{2\pi}} \int_{\substack{|z_j| \leq \tau_{\text{tail}} \sqrt{\log(1/\delta)} \\ |\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)}| \leq D_1}} \sum_{r_j=0}^{k_j} {k_j \choose r_j} \left(\mu_t^{-1} \sigma_t\right)^{r_j} \left(\mu_t^{-1} x_j - y_j^{(i)}\right)^{k_j - r_j} z_j^{r_j + 2l} dz_j
$$
\n
$$
= \frac{(-2)^{-l} \mu_t^{-k_j}}{l!\sqrt{2\pi}} \sum_{r_j=0}^{k_j} {k_j \choose r_j} \sigma_t^{r_j} \left(x_j - \mu_t y_j^{(i)}\right)^{k_j - r_j} \left(\frac{\overline{z}_{i,j}^{r_j + 2l + 1} - \underline{z}_{i,j}^{r_j + 2l + 1}}{r_j + 2l + 1}\right) \stackrel{\text{def}}{=} P_{i,j,k_j,l}(x_j, t),
$$

where

$$
\overline{z}_{i,j} = \min \left(\max \left(\frac{\mu_t(y_j^{(i)} + D_1) - x_j}{\sigma_t}, -\tau_{\text{tail}} \sqrt{\log(1/\delta)} \right), \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right),
$$

$$
\underline{z}_{i,j} = \min \left(\max \left(\frac{\mu_t(y_j^{(i)} - D_1) - x_j}{\sigma_t}, -\tau_{\text{tail}} \sqrt{\log(1/\delta)} \right), \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right).
$$

Combining $(B.1)$, $(B.38)$ and $(B.40)$ with the last two displays, we have

$$
|p_t(\mathbf{x}) - g_t(\mathbf{x})|
$$

\n
$$
\leq K D 2^{D+1} \delta^{\tau_{\text{tail}}^2/2} + K 6^D \delta \left\{ \frac{\log(1/\delta)}{\log(e/4)} \right\}^D
$$

\n
$$
+ \sum_{i=1}^{D_2} \sum_{0 \leq k \leq D_3} \left| \frac{(\mathbf{D}^k p_0)(\mathbf{y}^{(i)})}{\mathbf{k}!} \right| (2\pi \sigma_t^2)^{-\frac{D}{2}} D 2^{D-1} \delta^{e\tau_{\text{tail}}^2}
$$

\n
$$
\leq D_5 \left[\delta^{\tau_{\text{tail}}^2/2} + \delta \{ \log(1/\delta) \}^D + \delta^{e\tau_{\text{tail}}^2 - D/2} \{ \log(1/\delta) \}^{D\tau_{\text{bd}} + D} \right]
$$

for $\|\mathbf{x}\|_{\infty} \geq \mu_t - \tau_{\mathbf{x}} \{\log(1/\sigma_t)\}^{-\tau_{\text{bd}}}$ and $\delta \leq t \leq \delta^{\tau_t}$, where $D_5 = D_5(D, K, \tau, \tau_{\text{tail}})$ and $g_t : \mathbb{R}^D \to \mathbb{R}$ is a function such that

$$
g_t(\mathbf{x}) = \sum_{i=1}^{D_2} \sum_{0 \le k \, < D_3} \left\{ \frac{(\mathbf{D}^k p_0)(\mathbf{y}^{(i)})}{\mathbf{k}!} \right\} \prod_{j=1}^D \sum_{l=0}^{D_4 - 1} P_{i,j,k_j,l}(x_j,t), \quad \mathbf{x} \in \mathbb{R}^D.
$$

Since $\tau_{\text{tail}} \geq 2 \vee \sqrt{(D+3)/(2e)}$, we have

$$
|p_t(\mathbf{x}) - g_t(\mathbf{x})| \le 3D_5 \delta \{ \log(1/\delta) \}^D
$$
\n(B.41)

for small enough δ . Similarly, with $m = 1$ and $h = j$ in [\(B.40\)](#page-63-0), the last integral in (B.40) satisfies that

$$
\frac{1}{l!\sqrt{2\pi\sigma_t}}\int_{|x_j - \mu_t y_j| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y_j - x_j}{\sigma_t} \right) (y_j - y_j^{(i)})^{k_j} \left(-\frac{(x_j - \mu_t y_j)^2}{2\sigma_t^2} \right)^l dy_j
$$
\n
$$
= \frac{1}{l!\sqrt{2\pi}} \int_{\substack{|z_j| \leq \tau_{\text{tail}} \sqrt{\log(1/\delta)} \\ |\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)}| \leq D_1}} \left(\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)} \right)^{k_j} z_j^{2l+1} (-2)^{-l} dz_j
$$
\n
$$
= \frac{(-2)^{-l} \mu_t^{-k_j}}{l!\sqrt{2\pi}} \sum_{r_j=0}^{k_j} {k_j \choose r_j} \sigma_t^{r_j} (x_j - \mu_t y_j^{(i)})^{k_j - r_j} \left(\frac{\overline{z}_{i,j}^{r_j+2l+2} - \overline{z}_{i,j}^{r_j+2l+2}}{r_j + 2l + 2} \right) \stackrel{\text{def}}{=} \widetilde{P}_{i,j,k_j,l}(x_j,t).
$$

Combining $(B.1)$, $(B.39)$ and $(B.40)$ with the last display, we have

$$
\begin{split}\n&\left|\sigma_{t}\left(\nabla p_{t}(\mathbf{x})\right)_{h} - \tilde{g}_{t}^{(h)}(\mathbf{x})\right| \\
&\leq K2^{D} \left\{2(D-1)\sqrt{2/\pi} + \sqrt{2}\right\} \delta^{\tau_{\text{tail}}^{2}/4} + 2K\tau_{\text{tail}}6^{D} \{\log(e/4)\}^{-D} \delta \{\log(1/\delta)\}^{D+1/2} \\
&+ \sum_{i=1}^{D_{2}} \sum_{0 \leq k \, < D_{3}} \left|\frac{(D^{k}p_{0})(\mathbf{y}^{(i)})}{\mathbf{k}!}\right| (2\pi\sigma_{t}^{2})^{-\frac{D}{2}} D2^{D-1} \delta^{e\tau_{\text{tail}}^{2}} \left\{\tau_{\text{tail}}\sqrt{\log(1/\delta)}\right\}^{D} \\
&\leq D_{6} \left[\delta^{\tau_{\text{tail}}^{2}/4} + \delta \{\log(1/\delta)\}^{D+1/2} + \delta^{e\tau_{\text{tail}}^{2} - D/2} \{\log(1/\delta)\}^{D(\widetilde{\tau}_{\text{bd}}+3/2)}\right], \quad h \in [D]\n\end{split}
$$

for $\|\mathbf{x}\|_{\infty} \geq \mu_t - \tau_{\mathbf{x}} \{ \log(1/\sigma_t) \}^{-\tau_{\text{bd}}}$ and $\delta \leq t \leq \delta^{\tau_t}$, where $D_6 = D_6(D, K, \tau_{\text{tail}}, \tau)$ and $\widetilde{g}_t^{(h)}$ $t^{(h)}_t:\mathbb{R}^D\rightarrow$ $\mathbb{R}, h \in [D]$ is a function such that

$$
\widetilde{g}_t^{(h)}(\mathbf{x}) = \sum_{i=1}^{D_2} \sum_{0 \le k \lt D_3} \left\{ \frac{(\mathbf{D}^k p_0)(\mathbf{y}^{(i)})}{\mathbf{k}!} \right\} \left\{ \prod_{\substack{j=1 \ j \ne h}}^{D} \sum_{l=0}^{D_4-1} P_{i,j,k_j,l}(x_j,t) \right\} \left\{ \sum_{l=0}^{D_4-1} \widetilde{P}_{i,h,k_h,l}(x_h,t) \right\}.
$$

Since $\tau_{\text{tail}} \geq 2 \vee \sqrt{(D+3)/(2e)}$, we have

$$
\left|\sigma_t\left(\nabla p_t(\mathbf{x})\right)_h - \widetilde{g}_t^{(h)}(\mathbf{x})\right| \le 3D_6 \delta \{\log(1/\delta)\}^D, \quad h \in [D] \tag{B.42}
$$

for small enough δ .

Let $0 < \tilde{\delta} < \delta$ be a small enough value as described below. With $\tilde{\delta}^2 < 1/2$, Lemma [A.13](#page-37-0) implies that there exist neural networks $f_{\mu} \in \mathcal{F}_{NN}(L_{\mu}, \mathbf{d}_{\mu}, s_{\mu}, M_{\mu}), f_{\sigma} \in \mathcal{F}_{NN}(L_{\sigma}, \mathbf{d}_{\sigma}, s_{\sigma}, M_{\sigma})$ with

$$
L_{\mu}, L_{\sigma} \leq C_{N,4} \{ \log(1/\tilde{\delta}) \}^2, \quad \|\mathbf{d}_{\mu}\|_{\infty}, \|\mathbf{d}_{\sigma}\|_{\infty} \leq C_{N,4} \{ \log(1/\tilde{\delta}) \}^2
$$

$$
s_{\mu}, s_{\sigma} \leq C_{N,4} \{ \log(1/\tilde{\delta}) \}^3, \quad M_{\mu}, M_{\sigma} \leq C_{N,4} \log(1/\tilde{\delta})
$$

such that $|\mu_t - f_\mu(t)| \leq \tilde{\delta}$ and $|\sigma_t - f_\sigma(t)| \leq \tilde{\delta}$ for $t \geq \tilde{\delta}$, where $C_{N,4}$ is the constant in Lemma [A.13.](#page-37-0) Also, Lemma [A.14](#page-37-1) implies that there exist a neural network $f_{\text{rec}} \in \mathcal{F}_{\text{NN}}(L_{\text{rec}}, \mathbf{d}_{\text{rec}}, s_{\text{rec}}, M_{\text{rec}})$ with

$$
L_{\text{rec}} \leq C_{N,5} \{ \log(1/\tilde{\delta}) \}^2, \quad \|\mathbf{d}_{\text{rec}}\|_{\infty} \leq C_{N,5} \{ \log(1/\tilde{\delta}) \}^3
$$

$$
s_{\text{rec}} \leq C_{N,5} \{ \log(1/\tilde{\delta}) \}^4, \quad M_{\text{rec}} \leq C_{N,5} \tilde{\delta}^{-2}
$$

such that $|1/x - f_{\text{rec}}(x)| \leq \tilde{\delta}$ for any $x \in [\tilde{\delta}, 1/\tilde{\delta}],$ where $C_{N,5}$ is the constant in Lemma [A.14.](#page-37-1) Since $\sigma_t - \widetilde{\delta} \le f_\sigma(t) \le \sigma_t + \widetilde{\delta}$ for $t \ge \widetilde{\delta}$ and $\sqrt{\tau t} \le \sigma_t \le 1$ for $t \ge \delta$, we have $\delta \le f_\sigma(t) \le 2$ for $t \ge \delta$ with small enough δ so that $\delta \leq$ √ $\overline{\tau} \delta - \delta$ and $\delta \leq 1$. A simple calculation yields that

$$
|1/\sigma_t - f_{\text{rec}}(f_\sigma(t))| \le |1/\sigma_t - 1/f_\sigma(t)| + |1/f_\sigma(t) - f_{\text{rec}}(f_\sigma(t))|
$$

$$
\le \{\sigma_t \wedge f_\sigma(t)\}^{-2} |\sigma_t - f_\sigma(t)| + \tilde{\delta} \le (1 + \delta^{-2})\tilde{\delta}
$$
 (B.43)

for $t \geq \delta$. Lemma [A.9](#page-32-2) implies that there exists a neural network

$$
\widetilde{f}_{\text{mult}}^{(k)} \in \mathcal{F}_{\text{NN}}(\widetilde{L}_{\text{mult}}^{(k)}, \widetilde{\mathbf{d}}_{\text{mult}}^{(k)}, \widetilde{s}_{\text{mult}}^{(k)}, \widetilde{M}_{\text{mult}}^{(k)}), \quad k \ge 2
$$

with

$$
\widetilde{L}_{\text{mult}}^{(k)} \leq C_{N,1} \log k \{ \log(1/\widetilde{\delta}) + \log(1/\delta) \}, \quad \widetilde{d}_{\text{mult}}^{(k)} = (k, 48k, \dots, 48k, 1)^\top,
$$
\n
$$
\widetilde{s}_{\text{mult}}^{(2)} \leq C_{N,1} k \{ \log(1/\widetilde{\delta}) + \log(1/\delta) \}, \quad \widetilde{M}_{\text{mult}}^{(k)} = \delta^{-k}
$$

such that

$$
\left| \widetilde{f}_{\text{mult}}^{(k)}(\widetilde{x}_1, \dots, \widetilde{x}_k) - \prod_{i=1}^k x_i \right| \le \widetilde{\delta} + k \delta^{-(k-1)} \widetilde{\epsilon}
$$
\n(B.44)

for any $\mathbf{x} = (x_1, \dots, x_k) \in \mathbb{R}^k$ with $\|\mathbf{x}\|_{\infty} \leq \delta^{-1}$ and $\widetilde{\mathbf{x}} = (\widetilde{x}_1, \dots, \widetilde{x}_k) \in \mathbb{R}^k$ with $\|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty} \leq \widetilde{\epsilon}$, where $0 < \tilde{\epsilon} \leq 1$ and $C_{N,1}$ is the constant in Lemma [A.9.](#page-32-2) Let

$$
f_{\text{clip}} \in \mathcal{F}_{\text{NN}}(2, (1, 2, 1)^{\top}, 7, \tau_{\text{tail}} \sqrt{\log(1/\delta)})
$$

be the neural network in Lemma [A.10](#page-32-0) such that $f_{\text{clip}}(x) = (x \vee -\tau_{\text{tail}} \sqrt{\log(1/\delta)}) \wedge \tau_{\text{tail}} \sqrt{\log(1/\delta)}$ for $x \in \mathbb{R}$. For $i \in [D_1]$ and $j \in [D]$, consider functions $\overline{f}_{i,j}, \underline{f}_{i,j} : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ such that

$$
\overline{f}_{i,j}(x,t) = f_{\text{clip}}\left(\tilde{f}_{\text{mult}}^{(2)}\left(f_{\text{rec}}\left(f_{\sigma}(t)\right), \{y_j^{(i)} + D_1\} f_{\mu}(t) - x\right)\right),
$$
\n
$$
\underline{f}_{i,j}(x,t) = f_{\text{clip}}\left(\tilde{f}_{\text{mult}}^{(2)}\left(f_{\text{rec}}\left(f_{\sigma}(t)\right), \{y_j^{(i)} - D_1\} f_{\mu}(t) - x\right)\right),
$$

for $x, t \in \mathbb{R}$. Note that $|y_j^{(i)} + D_1| \le 1$ and $|y_j^{(i)} - D_1| \le 1$ for $i \in [D_1]$. Combining [\(B.43\)](#page-65-0) and [\(B.44\)](#page-65-1) with the last display, both $|\overline{z}_{i,j} - \overline{f}_{i,j}(x,t)|$ and $|\overline{z}_{i,j} - \overline{f}_{i,j}(x,t)|$ are bounded by

$$
\tilde{\delta} + 2\delta^{-1}(1+\delta^{-2})\tilde{\delta} \le 5\delta^{-3}\tilde{\delta}
$$
\n(B.45)

 $\text{for } |x| \leq \mu_t + \tau_{\mathbf{x}} \sigma_t \sqrt{\log(1/\delta)} \text{ and } \delta \leq t \leq \delta^{\tau_t} \text{ with small enough } \delta \text{ so that } |(y_j^{(i)} + D_1)\mu_t - x|,$ $|(y_j^{(i)} - D_1)\mu_t - x|$ and $1/\sigma_t$ are upper bounded by δ^{-1} . Since $\mu_t - \tilde{\delta} \le f_\mu(t) \le \mu_t + \tilde{\delta}$ for $t \ge \tilde{\delta}$, we have $1/4 \leq 1/2 - \tilde{\delta} \leq f_{\mu}(t) \leq 1 + \tilde{\delta} \leq 2$ and $1/4 \leq \mu_t \leq 2$ for $\delta \leq t \leq D_1$ with small enough $\tilde{\delta}$ by [\(B.1\)](#page-41-2). A simple calculation yields that

$$
|1/\mu_t - f_{\text{rec}}(f_\mu(t))| \le |1/\mu_t - 1/f_\mu(t)| + |1/f_\mu(t) - f_{\text{rec}}(f_\mu(t))|
$$

$$
\le \{\mu_t \wedge f_\mu(t)\}^{-1}|\mu_t - f_\mu(t)| + \tilde{\delta} \le 17\tilde{\delta}
$$
 (B.46)

for $\delta \leq t \leq \delta^{\tau_t}$. For any $i \in [D_2], j \in [D], k \in \{0, \ldots, D_3-1\}$, consider functions $f_{i,j,k}, \widetilde{f}_{i,j,k} : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ such that

$$
f_{i,j,k}(x,t) = \sum_{l=0}^{D_4-1} \sum_{r=0}^k {k \choose r} \left\{ \frac{(-2)^{-l}}{k!l!\sqrt{2\pi}(r+2l+1)} \right\} \left\{ \overline{f}_{i,j,k,l,r,r+2l+1} - \underline{f}_{i,j,k,l,r,r+2l+1} \right\},
$$

$$
\widetilde{f}_{i,j,k}(x,t) = \sum_{l=0}^{D_4-1} \sum_{r=0}^k {k \choose r} \left\{ \frac{(-2)^{-l}}{k!l!\sqrt{2\pi}(r+2l+2)} \right\} \left\{ \overline{f}_{i,j,k,l,r,r+2l+2} - \underline{f}_{i,j,k,l,r,r+2l+2} \right\},
$$

for $x, t \in \mathbb{R}$, where

$$
\overline{f}_{i,j,k,l,r,s} = \tilde{f}_{\text{mult}}^{(2k+s)} \left(f_{\text{rec}} \left(f_{\mu}(t) \right) \cdot \mathbf{1}_k, f_{\sigma}(t) \cdot \mathbf{1}_r, \left\{ x - f_{\mu}(t) y_j^{(i)} \right\} \cdot \mathbf{1}_{k-r}, \overline{f}_{i,j}(x,t) \cdot \mathbf{1}_s \right),
$$
\n
$$
\underline{f}_{i,j,k,l,r,s} = \tilde{f}_{\text{mult}}^{(2k+s)} \left(f_{\text{rec}} \left(f_{\mu}(t) \right) \cdot \mathbf{1}_k, f_{\sigma}(t) \cdot \mathbf{1}_r, \left\{ x - f_{\mu}(t) y_j^{(i)} \right\} \cdot \mathbf{1}_{k-r}, \underline{f}_{i,j}(x,t) \cdot \mathbf{1}_s \right),
$$

for $s \in \{r+2l+1, r+2l+2\}$. Combining [\(B.46\)](#page-65-2), [\(B.43\)](#page-65-0), [\(B.45\)](#page-65-3) and [\(B.44\)](#page-65-1) with the last two displays, the definition of $P_{i,j,k,l}(x,t)$ and $\widetilde{P}_{i,j,k,l}(x,t)$ implies that

$$
\begin{aligned}\n\left| \sum_{l=0}^{D_4 - 1} \frac{P_{i,j,k,l}(x,t)}{k!} - f_{i,j,k}(x,t) \right| \\
&\le \sum_{l=0}^{D_4 - 1} \sum_{r=0}^k {k \choose r} \left\{ \frac{2^{-l+1}}{k!l!\sqrt{2\pi}(r+2l+1)} \right\} \left\{ \widetilde{\delta} + 5(2k+r+2l+1)\delta^{-2k-r-2l-3} \widetilde{\delta} \right\}\n\end{aligned}
$$

and

$$
\left| \sum_{l=0}^{D_4-1} \frac{\widetilde{P}_{i,j,k,l}(x,t)}{k!} - \widetilde{f}_{i,j,k}(x,t) \right|
$$

$$
\leq \sum_{l=0}^{D_4-1} \sum_{r=0}^k {k \choose r} \left\{ \frac{2^{-l+1}}{k!l!\sqrt{2\pi}(r+2l+2)} \right\} \left\{ \widetilde{\delta} + 5(2k+r+2l+2)\delta^{-2k-r-2l-4} \widetilde{\delta} \right\}
$$

 $\text{for } i \in [D_2], j \in [D], k \in \{0, \ldots, D_3 - 1\}, |\mathbf{x}| \leq \mu_t + \tau_{\mathbf{x}} \sigma_t \sqrt{\log(1/\delta)} \text{ and } \delta \leq t \leq \delta^{\tau_t} \text{ with small enough }$ δ so that $\sigma_t, \mu_t^{-1}, |x - y_j^{(i)}|$ $\int_{j}^{(i)} |, \tau_{\text{tail}} \sqrt{\log(1/\delta)}$ are all upper bounded by δ^{-1} . Since $\sum_{r=0}^{k} {k \choose r}$ $\binom{k}{r} = 2^k$ and $2^{k}/k! \leq 2$ for all $k \in \mathbb{N}$, the last two displays are bounded by

$$
\sum_{l=0}^{D_4-1} \left(\frac{2^{-l+2}}{l!\sqrt{2\pi}(2l+1)} \right) \left\{ 1 + 5(3D_4 + 2l-1)\delta^{-3D_3 - 2l-1} \right\} \tilde{\delta} \le \delta^{-D_7 \log(1/\delta)} \tilde{\delta},\tag{B.47}
$$

where $D_7 = D_7(\tau_{tail})$. For any $i \in [D_2], j \in [D], k \in \{0, ..., D_3 - 1\}, l \in \{0, ..., D_4 - 1\}, |x| \le$ $\mu_t + \tau_{\mathbf{x}} \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \delta^{\tau_t}$, we have

$$
\left|\frac{P_{i,j,k,l}(x,t)}{k!}\right| \leq \left\{\frac{2^{-l-k+1}}{k!l!\sqrt{2\pi}(r+2l+1)}\sum_{r=0}^{k} \binom{k}{r}\right\} \left\{2\mu_t + \tau_{\mathbf{x}}\sigma_t\sqrt{\log(1/\delta)}\right\}^k \left\{\tau_{\text{tail}}\sqrt{\log(1/\delta)}\right\}^{k+2l+1}
$$

and

$$
\left|\frac{\widetilde{P}_{i,j,k,l}(x,t)}{k!}\right|\leq\left\{\frac{2^{-l-k+1}}{k!l!\sqrt{2\pi}(r+2l+1)}\sum_{r=0}^{k}\binom{k}{r}\right\}\left\{2\mu_{t}+\tau_{\text{x}}\sigma_{t}\sqrt{\log(1/\delta)}\right\}^{k}\left\{\tau_{\text{tail}}\sqrt{\log(1/\delta)}\right\}^{k+2l+2}
$$

by the definition of $P_{i,j,k,l}(x)$ and $\widetilde{P}_{i,j,k,l}(x)$. Since $\sum_{r=0}^{k} {k \choose r}$ $r_k^{(k)} = 2^k$ and $2^k/k! \leq 2$ for all $k \in \mathbb{N}$, the last two displays are bounded by $\{D_8 \log(1/\delta)\}^{k+l+1}$, where $D_8 = D_8(\tau_{\text{tail}}, \tau_{\text{x}})$. Then,

$$
\left| \sum_{l=0}^{D_4-1} \frac{P_{i,j,k,l}(x)}{k!} \right| \le D_4 \{ D_8 \log(1/\delta) \}^{D_3+D_4-1} \le \{ \log(1/\delta) \}^{D_9 \log(1/\delta)}
$$
 and

$$
\left| \sum_{l=0}^{D_4-1} \frac{\widetilde{P}_{i,j,k,l}(x)}{k!} \right| \le D_4 \{ D_8 \log(1/\delta) \}^{D_3+D_4-1} \le \{ \log(1/\delta) \}^{D_9 \log(1/\delta)}
$$

for $i \in [D_2], j \in [D], k \in \{0, \ldots, D_3 - 1\}, |x| \leq \mu_t + \tau_x \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \delta^{\tau_t}$, where $D_9 =$ $D_9(\tau_{\text{tail}}, D_8)$. Consider functions $f, \tilde{f}^{(1)}, \ldots, \tilde{f}^{(D)} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}$ such that

$$
f(\mathbf{x},t) = \sum_{i=1}^{D_2} \sum_{0 \le k \,
$$
\widetilde{f}^{(h)}(\mathbf{x},t) = \sum_{i=1}^{D_2} \sum_{0 \le k \,
$$
$$

where $f_{\text{mult}}^{(D)} \in \mathcal{F}_{\text{NN}}(L_{\text{mult}}^{(D)}, \mathbf{d}_{\text{mult}}^{(D)}, s_{\text{mult}}^{(D)}, M_{\text{mult}}^{(D)})$ is the neural network in Lemma [A.9](#page-32-2) with $L_{\text{mult}}^{(D)} \leq C_{N,1} \log D[\log(1/\tilde{\delta}) + DD_9 \log(1/\delta) \log \log(1/\delta)], \quad d_{\text{mult}}^{(D)} = (D, 48D, \dots, 48D, 1)^{\top},$ $s_{\text{mult}}^{(D)} \leq C_{N,1} D[\log(1/\tilde{\delta}) + D_9 \log(1/\delta) \log \log(1/\delta)], \quad M_{\text{mult}}^{(D)} = {\log(1/\delta)}^{DD_9 \log(1/\delta)}$

such that

$$
\left| f_{\text{mult}}^{(D)}(\widetilde{x}_1,\ldots,\widetilde{x}_D) - \prod_{i=1}^D x_i \right| \le \widetilde{\delta} + D\{\log(1/\delta)\}^{(D-1)D_9 \log(1/\delta)} \widetilde{\epsilon}
$$

for any $\mathbf{x} = (x_1, \ldots, x_D) \in \mathbb{R}^D$ with $\|\mathbf{x}\|_{\infty} \leq {\log(1/\delta)}^{\int_{\mathcal{D}} \log(1/\delta)}$ and $\widetilde{\mathbf{x}} = (\widetilde{x}_1, \ldots, \widetilde{x}_D) \in \mathbb{R}^D$ with $\|\mathbf{x}-\widetilde{\mathbf{x}}\|_{\infty} \leq \widetilde{\epsilon}$. Combining [\(B.47\)](#page-66-0) with the last display, we have

$$
|f(\mathbf{x},t) - g_t(\mathbf{x})|
$$

\n
$$
\leq \sum_{i=1}^{D_2} \sum_{0 \leq k \cdot < D_3} |(D^k p_0)(\mathbf{y}^{(i)})| \left\{ 1 + D\delta^{-D_7 \log(1/\delta)} \{ \log(1/\delta) \}^{(D-1)D_9 \log(1/\delta)} \right\} \tilde{\delta}
$$

\n
$$
\leq K D_2 D_3^D \left\{ 1 + D\delta^{-D_7 \log(1/\delta)} \{ \log(1/\delta) \}^{(D-1)D_9 \log(1/\delta)} \right\} \tilde{\delta} \leq \delta^{-D_{10} \log(1/\delta)} \tilde{\delta}
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \tau_{\mathbf{x}} \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq D_1$, where $D_{10} = D_{10}(D, K, \widetilde{\tau}_{\text{bd}}, D_7, D_9)$ is a large enough constant. Similarly, we have

$$
\left|\widetilde{f}^{(h)}(\mathbf{x},t)-\widetilde{g}_t^{(h)}(\mathbf{x})\right|\leq \delta^{-D_{10}\log(1/\delta)}\widetilde{\delta},\quad h\in[D]
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \tau_x \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \delta^{\tau_t}$. Let $\tilde{\delta} = \delta^{D_{10} \log(1/\delta) + 1}$. Combining [\(B.41\)](#page-64-0) and [\(B.42\)](#page-64-1) with the last two displays, we have

$$
|p_t(\mathbf{x}) - f(\mathbf{x}, t)| \le \delta + 3D_5 \delta \{ \log(1/\delta) \}^D \le (1 + 3D_5) \delta \{ \log(1/\delta) \}^D \quad \text{and}
$$

$$
\left| \sigma_t \left(\nabla p_t(\mathbf{x}) \right)_h - \tilde{f}^{(h)}(\mathbf{x}, t) \right| \le \delta + 3D_6 \delta \{ \log(1/\delta) \}^D \le (1 + 3D_6) \delta \{ \log(1/\delta) \}^D, \quad h \in [D]
$$

for $\mu_t - \tau_x {\log(1/\sigma_t)}^{-\tau_{\text{bd}}} \leq ||\mathbf{x}||_{\infty} \leq \mu_t + \tau_x \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \delta^{\tau_t}$. Note that

$$
D_2 \le D_{13} {\log(1/\delta)}^{D\widetilde{\tau}_{\text{bd}}}, \quad D_3 \le D_{13} \log(1/\delta), \quad D_4 \le D_{13} \log(1/\delta),
$$

where $D_{13} = D_{13}(D, \tau_{\text{tail}})$. Consider a function $f : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}^{D+1}$ such that

$$
\mathbf{f}(\mathbf{x},t) = (f(\mathbf{x},t), \widetilde{f}^{(1)}(\mathbf{x},t), \dots, \widetilde{f}^{(D)}(\mathbf{x},t))^\top
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Lemma [A.4,](#page-29-0) Lemma [A.5,](#page-29-1) Lemma [A.6](#page-30-0) and Lemma [A.7](#page-30-1) implies that $\mathbf{f} \in$ $\mathcal{F}_{NN}(L, \mathbf{d}, s, m)$ with

$$
L \le D_{11} \{ \log(1/\delta) \}^4, \quad \|\mathbf{d}\|_{\infty} \le D_{11} \{ \log(1/\delta) \}^{7+D\widetilde{\tau}_{\text{bd}}+D},
$$

$$
s \le D_{11} \{ \log(1/\delta) \}^{11+D\widetilde{\tau}_{\text{bd}}+D}, \quad M \le \exp(D_{11} \{ \log(1/\delta) \}^2),
$$

where $D_{11} = D_{11}(D, K, \tau_{tail}, C_{N,1}, C_{N,4}, C_{N,5}, D_9, D_{13})$. The assertion follows by re-defining the constants. \Box

B.4 Proof of Proposition [B.3](#page-40-1)

Proof. Let

$$
\tau_{\rm t} = \overline{\tau}^{-1}
$$
 and $\tau_{\rm tail} = \left\{ 4(D\overline{\tau}\tau_{\rm t} + 1) \vee \left(\frac{D+1}{e} \right) \right\}^{\frac{1}{2}}$.

Let $t_* > 0$ and $0 < \delta < 1$ be small enough values as described below. By the Markov property of $(\mathbf{X}_t)_{t\geq0}$, we have

$$
p_{t_*+t}(\mathbf{x}) = \int_{\mathbb{R}^D} p_{t_*}(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d} \mathbf{y}
$$

for any $\mathbf{x} \in \mathbb{R}^D$ and $t \geq 0$. Let $C_{S,1} = C_{S,1}(D,K,\tau_1)$ be the constant in Lemma [A.1.](#page-28-0) Since $|p_{t_*}(\mathbf{x})| \leq C_{S,1}$ for any $\mathbf{x} \in \mathbb{R}^D$, we have

$$
\int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_{\infty}\geq\tau_{\text{tail}}\sigma_t\sqrt{\log(1/\delta)}} p_{t_*}(\mathbf{y})\phi_{\sigma_t}(\mathbf{x}-\mu_t\mathbf{y})\mathrm{d}\mathbf{y} \n\leq C_{S,1} \int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_{\infty}\geq\tau_{\text{tail}}\sigma_t\sqrt{\log(1/\delta)}} \phi_{\sigma_t}(\mathbf{x}-\mu_t\mathbf{y})\mathrm{d}\mathbf{y} = C_{S,1}\mu_t^{-D} \int_{\|\mathbf{z}\|_{\infty}\geq\tau_{\text{tail}}\sqrt{\log(1/\delta)}} \phi_1(\mathbf{z})\mathrm{d}\mathbf{z} \n\leq C_{S,1}\mu_t^{-D} \sum_{i=1}^D \int_{|z_i|\geq\tau_{\text{tail}}\sqrt{\log(1/\delta)}} \phi(z_i)\mathrm{d}z_i \leq 2C_{S,1}D\mu_t^{-D}\delta^{\frac{\tau_{\text{tail}}^2}{2}}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \geq 0$, where the last inequality holds by the tail probability of the standard normal distribution. Also,

$$
\left| \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \geq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y_i - x_i}{\sigma_t} \right) p_{t_*}(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} \right|
$$
\n
$$
\leq C_{S,1} \mu_t^{-D} \int_{\|\mathbf{z}\|_{\infty} \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)}} |z_i| \phi_1(\mathbf{z}) d\mathbf{z} \leq C_{S,1} \mu_t^{-D} \sum_{j=1}^D \int_{|z_i| \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)}} |z_i| \phi_1(\mathbf{z}) d\mathbf{z}
$$
\n
$$
= C_{S,1} \mu_t^{-D} \left\{ (D-1) \mathbb{E}[|Z|] \int_{|z| \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)}} \phi(z) dz + \int_{|z| \geq \tau_{\text{tail}} \sqrt{\log(1/\delta)}} |z| \phi(z) dz \right\}
$$
\n
$$
\leq 2C_{S,1} \mu_t^{-D} \left\{ (D-1) \sqrt{2/\pi} \delta^{\tau_{\text{tail}}^2/2} + \sqrt{\mathbb{E}[Z^2]} \delta^{\tau_{\text{tail}}^2/4} \right\}
$$
\n
$$
\leq 2C_{S,1} D \mu_t^{-D} \delta^{\frac{\tau_{\text{tail}}^2}{4}}, \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \geq 0$, where Z denotes the one-dimensional standard normal random variable and the second inequality holds by the Cauchy-Schwarz inequality. Note that $\exp(-\overline{\tau}t) \leq \mu_t \leq \exp(-\underline{\tau}t)$ for $t \geq 0$. Then,

$$
\left| p_{t_{*}+t}(\mathbf{x}) - \int_{\|\mathbf{x} - \mu_{t}\mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} p_{t_{*}}(\mathbf{y}) \phi_{\sigma_{t}}(\mathbf{x} - \mu_{t}\mathbf{y}) d\mathbf{y} \right| \leq 2C_{S,1} D \delta^{\frac{\tau_{\text{tail}}^{2}}{2} - D\overline{\tau}\tau_{t}},
$$
\n
$$
\left\| \sigma_{t} \nabla p_{t_{*}+t}(\mathbf{x}) - \int_{\|\mathbf{x} - \mu_{t}\mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} \left(\frac{\mu_{t}\mathbf{y} - \mathbf{x}}{\sigma_{t}} \right)^{\top} p_{t_{*}}(\mathbf{y}) \phi_{\sigma_{t}}(\mathbf{x} - \mu_{t}\mathbf{y}) d\mathbf{y} \right\|_{\infty}
$$
\n
$$
\leq 2C_{S,1} D \delta^{\frac{\tau_{\text{tail}}^{2}}{4} - D\overline{\tau}\tau_{t}} \tag{B.48}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $0 \le t \le \tau_t \log(1/\delta)$. Lemma [A.1](#page-28-0) implies that $p_{t_*}(\mathbf{x}) \le C_{S,1} \delta^{\tau_{\text{tail}}^2/2}$ for $\|\mathbf{x}\|_{\infty} \ge$ $\mu_{t_*} + \tau_{\text{tail}} \sigma_{t_*} \sqrt{\log(1/\delta)}$. Then,

$$
\begin{aligned} &\left|\int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_\infty \leq \tau_{\mathrm{tail}} \sigma_t \sqrt{\log(1/\delta)}} p_{t*}(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x}-\mu_t\mathbf{y}) \mathrm{d}\mathbf{y} \right| \\ &\|\mathbf{y}\|_\infty \geq &\mu_{t*} + \tau_{\mathrm{tail}} \sigma_{t*} \sqrt{\log(1/\delta)} \\ &\leq 2C_{S,1} \delta^{\frac{\tau_{\mathrm{tail}}^2}{2}} \int_{\|\mathbf{x}-\mu_t\mathbf{y}\|_\infty \leq \tau_{\mathrm{tail}} \sigma_t \sqrt{\log(1/\delta)}} \phi_{\sigma_t}(\mathbf{x}-\mu_t\mathbf{y}) \mathrm{d}\mathbf{y} \\ &\|\mathbf{y}\|_\infty \geq &\mu_{t*} + \tau_{\mathrm{tail}} \sigma_{t*} \sqrt{\log(1/\delta)} \\ &\leq C_{S,1} \mu_t^{-D} \delta^{\frac{\tau_{\mathrm{tail}}^2}{2}} \int_{\mathbb{R}^D} \phi_1(\mathbf{z}) \mathrm{d}\mathbf{z} \leq C_{S,1} \delta^{\frac{\tau_{\mathrm{tail}}^2}{2} - D\overline{\tau}\tau_t} \end{aligned}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $0 \le t \le \tau_t \log(1/\delta)$, where the last inequaltiy holds because $\mu_t^{-D} \le \exp(D\overline{\tau}t)$. Also,

$$
\left| \int_{\substack{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)} \\ \|\mathbf{y}\|_{\infty} \geq \mu_{t_*} + \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y_i - x_i}{\sigma_t} \right) p_{t_*}(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} \right|
$$

$$
\leq C_{S,1} \delta^{\frac{\tau_{\text{tail}}^2}{2}} \mu_t^{-D} \int_{\mathbb{R}^D} |z_i| \phi_1(\mathbf{z}) d\mathbf{z} \leq \left(\sqrt{\frac{2}{\pi}} \right) C_{S,1} \delta^{\frac{\tau_{\text{tail}}^2}{2} - D\overline{\tau}\tau_t}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $0 \le t \le \tau_t \log(1/\delta)$. Combining with [\(B.48\)](#page-68-0), we have

$$
\left| p_{t_{*}+t}(\mathbf{x}) - \int_{\|\mathbf{x} - \mu_{t}\mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} p_{t_{*}}(\mathbf{y}) \phi_{\sigma_{t}}(\mathbf{x} - \mu_{t}\mathbf{y}) d\mathbf{y} \right|
$$
\n
$$
\leq C_{S,1}(2D+1) \delta^{\frac{\tau_{\text{tail}}^{2}}{2} - D\tau_{\text{te}}} \tag{B.49}
$$

and

$$
\begin{aligned}\n\left\|\sigma_{t_*+t} \nabla p_t(\mathbf{x}) - \int_{\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t \mathbf{y} - \mathbf{x}}{\sigma_t} \right)^{\top} p_{t_*}(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}\mathbf{y} \right\|_{\infty} \\
\leq C_{S,1} (2D + \sqrt{2/\pi}) \delta^{\frac{\tau_{\text{tail}}^2 - D\tau_{\tau_t}}{4}}\n\end{aligned}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $0 \le t \le \tau_t \log(1/\delta)$. Let $m_* \in \mathbb{N}_{\ge 2}$ be a large enough value as described below and $\mathbf{y}^{(1)}, \ldots, \mathbf{y}^{(D_1)} \in \mathbb{R}^D$ be distinct vectors satisfying that

$$
\left\{\mathbf{y}^{(1)},\ldots,\mathbf{y}^{(D_1)}\right\} = \left\{\tau_*(n_1,\ldots,n_D)^\top : n_i \in \mathbb{Z}, i \in [D]\right\} \cap \left\{\mathbf{y} \in \mathbb{R}^D : ||\mathbf{y}||_{\infty} \leq (m_*-1)\tau_*\right\},\
$$

where $\tau_* = {\mu_{t_*}} + \tau_{\text{tail}} \sigma_{t_*} \sqrt{\log(1/\delta)} \} / m_*$ and $D_1 = (2m_* - 1)^D$. Let $\mathcal{Y}_i = {\mathbf{y} \in \mathbb{R}^D : \|\mathbf{y} - \mathbf{y}^{(i)}\|_{\infty} \leq \tau_* }$ for $i \in [D_1]$. Taylor's theorem for multivariate function implies that

$$
p_{t_{*}}(\mathbf{y}) = \sum_{0 \leq k \, < \, \tau_{\text{sm}}} \frac{(\mathbf{D}^{\mathbf{k}} p_{t_{*}})(\mathbf{y}^{(i)})}{\mathbf{k}!} (\mathbf{y} - \mathbf{y}^{(i)})^{\mathbf{k}} + \sum_{k = \tau_{\text{sm}}} \frac{(\mathbf{D}^{\mathbf{k}} p_{t_{*}})(\xi \mathbf{y}^{(i)} + (1 - \xi)\mathbf{y})}{\mathbf{k}!} (\mathbf{y} - \mathbf{y}^{(i)})^{\mathbf{k}}
$$

for a suitable $\xi \in [0,1]$ and $i \in [D_1], \mathbf{y} \in \mathbb{R}^D$, where $(\mathbf{y} - \mathbf{y}^{(i)})^k = \prod_{j=1}^D (y_j - y_j^{(i)})$ $(\binom{i}{j} k_j, \; \mathbf{k}! \, = \, \prod_{j=1}^D k_j!$ and $k = ||\mathbf{k}||_1$. Combining with Lemma [A.3,](#page-29-2) we have

$$
\left| p_{t*}(\mathbf{y}) - \sum_{0 \leq k \, < \tau_{\text{sm}}} \frac{(D^k p_{t*})(\mathbf{y}^{(i)})}{k!} (\mathbf{y} - \mathbf{y}^{(i)})^k \right| \cdot 1\{\|\mathbf{y} - \mathbf{y}^{(i)}\|_{\infty} \leq \tau_*\}
$$
\n
$$
\leq \sum_{k = \tau_{\text{sm}}} C_{S,3} \sigma_{t*}^{-\tau_{\text{sm}}} \prod_{j=1}^D \left(\frac{e\tau_*}{k_j}\right)^{k_j} \leq C_{S,3}(\tau_{\text{sm}} + 1)^D \left(\frac{e\tau_*}{\sigma_{t*}}\right)^{\tau_{\text{sm}}}
$$

for $\mathbf{y} \in \mathbb{R}^D$ and $i \in [D_1]$ because $k! \geq k^k e^{-k}$ for any $k \in \mathbb{Z}_{\geq 0}$, where $C_{S,3} = C_{S,3}(D, K, \tau_{sm}, \overline{\tau}, \underline{\tau})$ is the constant in Lemma [A.3.](#page-29-2) Since $\mathcal{Y}_1, \ldots, \mathcal{Y}_{D_1}$ are mutually disjoint except on a set of Lebesgue measure zero and $\bigcup_{i=1}^{D_1} \mathcal{Y}_i = \{ \mathbf{y} \in \mathbb{R}^D : ||\mathbf{y}||_{\infty} \leq \mu_{t_*} + \tau_{\text{tail}} \sigma_{t_*} \sqrt{\log(1/\delta)} \},\$ we have

$$
\int_{\|\mathbf{y}\|_{\infty} \leq \mu_{t_{*}} + \tau_{\text{tail}} \sigma_{t_{*}} \sqrt{\log(1/\delta)}} g(\mathbf{y}) d\mathbf{y} = \sum_{i=1}^{D_{1}} \int_{\|\mathbf{y} - \mathbf{y}^{(i)}\|_{\infty} \leq \tau_{*}} g(\mathbf{y}) d\mathbf{y}
$$
\n
$$
= \int_{\|\mathbf{y}\|_{\infty} \leq \mu_{t_{*}} + \tau_{\text{tail}} \sigma_{t_{*}} \sqrt{\log(1/\delta)}} \sum_{i=1}^{D_{1}} g(\mathbf{y}) \cdot 1\{\|\mathbf{y} - \mathbf{y}^{(i)}\|_{\infty} \leq \tau_{*}\} d\mathbf{y}
$$

for any continuous function $g : \mathbb{R}^D \to \mathbb{R}$. Combining [\(B.49\)](#page-69-0) with the last two displays, we have

$$
\left| p_{t_{*}+t}(\mathbf{x}) - \sum_{i=1}^{D_{1}} \int_{\|\mathbf{x}-\mu_{t}\mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} \left\{ \sum_{0 \leq k_{*} < \tau_{\text{sm}}} \frac{(D^{k} p_{t_{*}})(\mathbf{y}^{(i)})}{k!} (\mathbf{y}-\mathbf{y}^{(i)})^{k} \right\} \phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y}) d\mathbf{y} \right|
$$
\n
$$
\leq C_{S,1}(2D+1) \delta^{\frac{\tau_{\text{tail}}^{2}}{2} - D\overline{\tau}_{\tau_{t}}} + C_{S,3}(\tau_{\text{sm}}+1)^{D} \left(\frac{e\tau_{*}}{\sigma_{t_{*}}} \right)^{\tau_{\text{sm}}} \int_{\|\mathbf{x}-\mu_{t}\mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} \sum_{i=1}^{D_{1}} \phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y}) \cdot 1 \{ \|\mathbf{y}-\mathbf{y}^{(i)}\|_{\infty} \leq \tau_{*} \} d\mathbf{y}
$$
\n
$$
\leq C_{S,1}(2D+1) \delta^{\frac{\tau_{\text{tail}}^{2}}{2} - D\overline{\tau}_{\tau_{t}}} + C_{S,3}(\tau_{\text{sm}}+1)^{D} \left(\frac{e\tau_{*}}{\sigma_{t_{*}}} \right)^{\tau_{\text{sm}}} \delta^{-D\overline{\tau}_{\tau_{t}}} \tag{B.50}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $0 \le t \le \tau_t \log(1/\delta)$, where the last inequality holds because $\int_{\mathbb{R}^D} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} = \mu_t^{-D}$ and $\mu_t^{-D} \leq \exp(D\overline{\tau}t) \leq \delta^{-D\overline{\tau}\tau_t}$. Also, we have

$$
\left\|\sigma_{t}\nabla p_{t_{*}+t}(\mathbf{x})-\sum_{i=1}^{D_{1}}\int_{\|\mathbf{x}-\mu_{t}\mathbf{y}\|_{\infty}\leq\tau_{\text{tail}}\sigma_{t}\sqrt{\log(1/\delta)}}\|\mathbf{y}-\mathbf{y}^{(i)}\|_{\infty}\leq\tau_{*}\right\}
$$
\n
$$
\left\{\sum_{0\leq k,\leq\tau_{\text{sm}}} \frac{(D^{k}p_{t_{*}})(\mathbf{y}^{(i)})}{k!} \left(\frac{\mu_{t}\mathbf{y}-\mathbf{x}}{\sigma_{t}}\right)^{\top}(\mathbf{y}-\mathbf{y}^{(i)})^{k}\right\}\phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y})\mathrm{d}\mathbf{y}\right\|_{\infty}
$$
\n
$$
\leq C_{S,1}(2D+\sqrt{2/\pi})\delta^{\frac{\tau_{\text{tail}}^{2}}{4}-D\overline{\tau}\tau_{t}}+C_{S,3}(\tau_{\text{sm}}+1)^{D}\left(\frac{e\tau_{*}}{\sigma_{t_{*}}}\right)^{\tau_{\text{sm}}} \tag{B.51}
$$
\n
$$
\cdot\int_{\|\mathbf{x}-\mu_{t}\mathbf{y}\|_{\infty}\leq\tau_{\text{tail}}\sigma_{t}\sqrt{\log(1/\delta)}}\sum_{i=1}^{D_{1}}\left\|\frac{\mu_{t}\mathbf{y}-\mathbf{x}}{\sigma_{t}}\right\|_{\infty}\phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y})\cdot1\{\|\mathbf{y}-\mathbf{y}^{(i)}\|_{\infty}\leq\tau_{*}\}\mathrm{d}\mathbf{y}
$$
\n
$$
\leq C_{S,1}(2D+\sqrt{2/\pi})\delta^{\frac{\tau_{\text{tail}}^{2}}{4}-D\overline{\tau}\tau_{t}}+C_{S,3}\tau_{\text{tail}}(\tau_{\text{sm}}+1)^{D}\left(\frac{e\tau_{*}}{\sigma_{t_{*}}}\right)^{\tau_{\text{sm}}}\delta^{-D\overline{\tau}\tau_{t}}\sqrt{\log(1/\delta)}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $0 \le t \le \tau_t \log(1/\delta)$, where the last inequality holds because $\int_{\mathbb{R}^D} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} = \mu_t^{-D}$ and $\mu_t^{-D} \leq \exp(D\overline{\tau}t) \leq \delta^{-D\overline{\tau}\tau_t}$.

For $x, y \in \mathbb{R}$ and $t > 0$, Taylor's theorem yields that

$$
\left|\exp\left(-\frac{(x-\mu_t y)^2}{2\sigma_t^2}\right)-\sum_{l=0}^{D_2-1}\frac{1}{l!}\left(-\frac{(x-\mu_t y)^2}{2\sigma_t^2}\right)^l\right|\leq \frac{1}{D_2!}\left(\frac{(x-\mu_t y)^2}{2\sigma_t^2}\right)^{D_2},
$$

where $D_2 = \lfloor 2e\tau_{\text{tail}}^2 \log(1/\delta) \rfloor + 1$ with small enough δ so that $D_2 \geq 1$. For $|x - \mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}$ and $t \geq 0$, the last display is further bounded by

$$
\left(\frac{e\tau_{\text{tail}}^2\log(1/\delta)}{D_2}\right)^{D_2} \le 2^{-D_2} \le \delta^{2e\tau_{\text{tail}}^2\log 2} \le \delta^{e\tau_{\text{tail}}^2},
$$

where the last inequality holds because $1/2 \le \log 2$ and $0 < \delta < 1$. Then,

$$
\begin{aligned} &\left|\int_{|x-\mu_t y| \leq \tau_{\mathrm{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y - x}{\sigma_t}\right)^u (y - \tilde{y})^k \exp\left(-\frac{(x - \mu_t y)^2}{2\sigma_t^2}\right) \mathrm{d}y \right. \\ & \left. - \sum_{l=0}^{D_2-1} \frac{1}{l!} \int_{|x-\mu_t y| \leq \tau_{\mathrm{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y - x}{\sigma_t}\right)^u (y - \tilde{y})^k \left(-\frac{(x - \mu_t y)^2}{2\sigma_t^2}\right)^l \mathrm{d}y \right| \\ & \leq \delta^{e\tau_{\mathrm{tail}}^2} \int_{|x-\mu_t y| \leq \tau_{\mathrm{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left|\frac{\mu_t y - x}{\sigma_t}\right|^u |y - \tilde{y}|^k \mathrm{d}y \\ & \leq \delta^{e\tau_{\mathrm{tail}}^2} \int_{|y| \leq m_* \tau_*} \left\{\tau_{\mathrm{tail}} \sqrt{\log(1/\delta)}\right\}^u \tau_*^k \mathrm{d}y = m_* \tau_*^{k+1} \left\{\tau_{\mathrm{tail}} \sqrt{\log(1/\delta)}\right\}^u \delta^{e\tau_{\mathrm{tail}}^2} \end{aligned}
$$

for any $x, \tilde{y} \in \mathbb{R}$ with $|\tilde{y}| = (m-1)\tau_*, t \geq 0, k \in \mathbb{Z}_{\geq 0}$ and $u \in \{0,1\}$. Note that $\mu_{t_*} \leq 1 - \underline{\tau}t_*/2 \leq 1$ and $\sqrt{\underline{\tau}t_*} \leq \sigma_{t_*} \leq$ √ $\overline{2\overline{\tau}t_*} \leq 1$ with $t_* \leq (2\overline{\tau})^{-1}$. For $k \leq \tau_{\rm sm}$, the last display is bounded by

$$
(1 + \tau_{\text{tail}})^{k+1} m_{\ast}^{-k} \delta^{e\tau_{\text{tail}}^2} \{ \log(1/\delta) \}^{\frac{k+1}{2}} \left\{ \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right\}^{u}
$$

$$
\leq \tau_{\text{tail}} (1 + \tau_{\text{tail}})^{\tau_{\text{sm}} + 1} \delta^{e\tau_{\text{tail}}^2} \{ \log(1/\delta) \}^{\frac{\tau_{\text{sm}}}{2} + 1}
$$

because $m_* \tau_* = \mu_{t_*} + \sigma_{t_*} \sqrt{\log(1/\delta)}$ and $\mu_{t_*} + \sigma_{t_*} \sqrt{\log(1/\delta)} \leq (1 + \tau_{tail}) \sqrt{\log(1/\delta)}$ with small enough δ so that $\delta \leq 1/e$ and $\tau_{\text{tail}}\sqrt{\log(1/\delta)} \geq 1$. Moreover,

$$
\left| \sum_{l=0}^{D_2-1} \frac{1}{l!} \int_{|x-\mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y - x}{\sigma_t} \right)^u (y - \tilde{y})^k \left(-\frac{(x - \mu_t y)^2}{2\sigma_t^2} \right)^l dy \right|
$$

$$
\leq \tau_{\text{tail}} \left(1 + \delta^{e\tau_{\text{tail}}^2} \right) (1 + \tau_{\text{tail}})^{\tau_{\text{sm}} + 1} {\log(1/\delta)}^{\frac{\tau_{\text{sm}}}{2} + 1}
$$

for any $x, \tilde{y} \in \mathbb{R}$ with $|\tilde{y}| = (m_{*}-1)\tau_{*}, t \geq 0, 0 \leq k \leq \tau_{\text{sm}}$ and $u \in \{0, 1\}$ because

$$
\left| \int_{|x-\mu_t y| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y - x}{\sigma_t} \right)^u (y - \tilde{y})^k \exp\left(-\frac{(x - \mu_t y)^2}{2\sigma_t^2} \right) dy \right|
$$

$$
\leq \int_{|y| \leq m_* \tau_*} \left\{ \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right\}^u \tau_*^k dy \leq m_* \tau_*^{k+1} \tau_{\text{tail}} \sqrt{\log(1/\delta)}
$$

$$
\leq \tau_{\text{tail}} (1 + \tau_{\text{tail}})^{\tau_{\text{sm}} + 1} \left\{ \log(1/\delta) \right\}^{\frac{\tau_{\text{sm}}}{2} + 1}.
$$

Note that $\left|\prod_{j=1}^D x_j - \prod_{j=1}^D \widetilde{x}_j\right| \le DC^{D-1} \|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty}$ for any $\mathbf{x}, \widetilde{\mathbf{x}} \in [-C, C]^D$. Since the last two displays are bounded by $2\tau_{tail}(1+\tau_{tail})^{\tau_{sm}+1}\{\log(1/\delta)\}^{\frac{\tau_{sm}}{2}+1}$, we have

$$
\left| \int_{\|\mathbf{x}-\mu_{t}\mathbf{y}\|_{\infty} \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} \left(\frac{\mu_{t} y_{h} - x_{h}}{\sigma_{t}} \right)^{u} (\mathbf{y} - \mathbf{y}^{(i)})^{k} \phi_{\sigma_{t}} (\mathbf{x} - \mu_{t}\mathbf{y}) d\mathbf{y} - \right|
$$
\n
$$
\prod_{j=1}^{D} \sum_{l=0}^{D_{2}-1} \frac{1}{l! \sqrt{2\pi\sigma_{t}}} \int_{|x_{j} - \mu_{t}y_{j}| \leq \tau_{\text{tail}} \sigma_{t} \sqrt{\log(1/\delta)}} \left(\frac{\mu_{t} y_{h} - x_{h}}{\sigma_{t}} \right)^{u \cdot 1\{h=j\}} (y_{j} - y_{j}^{(i)})^{k_{j}} \left(-\frac{(x_{j} - \mu_{t}y_{j})^{2}}{2\sigma_{t}^{2}} \right)^{l} dy_{j}
$$
\n
$$
\leq (2\pi\sigma_{t}^{2})^{-\frac{D}{2}} D 2^{D-1} \tau_{\text{tail}}^{D} (1 + \tau_{\text{tail}})^{D(\tau_{\text{sm}}+1)} \delta^{e\tau_{\text{tail}}^{2}} {\log(1/\delta)}^{D(\frac{\tau_{\text{sm}}}{2}+1)} \tag{B.52}
$$
for $\mathbf{x} \in \mathbb{R}^D, t > 0, h \in [D], i \in [D], j \in [D], \mathbf{k} \in \mathbb{Z}_{\geq 0}^D$ and $u \in \{0, 1\}$ with $k \leq \tau_{\text{sm}}$. With $u = 0$ in the last display, the second integral satisfies that

$$
\frac{1}{l!\sqrt{2\pi\sigma_t}}\int_{|x_j - \mu_t y_j| \leq \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}} \left(y_j - y_j^{(i)}\right)^{k_j} \left(-\frac{(x_j - \mu_t y_j)^2}{2\sigma_t^2}\right)^l dy_j
$$
\n
$$
= \frac{1}{l!\sqrt{2\pi}} \int_{\substack{|z_j| \leq \tau_{\text{tail}} \sqrt{\log(1/\delta)} \\ |\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)}| \leq \tau_*}} \left(\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)}\right)^{k_j} z_j^{2l} (-2)^{-l} dz_j
$$
\n
$$
= \frac{(-2)^{-l}}{l!\sqrt{2\pi}} \int_{\substack{|z_j| \leq \tau_{\text{tail}} \sqrt{\log(1/\delta)} \\ |\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)}| \leq \tau_*}} \sum_{r_j=0}^{k_j} {k_j \choose r_j} \left(\mu_t^{-1} \sigma_t\right)^{r_j} \left(\mu_t^{-1} x_j - y_j^{(i)}\right)^{k_j - r_j} z_j^{r_j + 2l} dz_j
$$
\n
$$
= \frac{(-2)^{-l} \mu_t^{-k_j}}{l!\sqrt{2\pi}} \sum_{r_j=0}^{k_j} {k_j \choose r_j} \sigma_t^{r_j} \left(x_j - \mu_t y_j^{(i)}\right)^{k_j - r_j} \left(\frac{\overline{z}_{i,j}^{r_j + 2l + 1} - \underline{z}_{i,j}^{r_j + 2l + 1}}{r_j + 2l + 1}\right) \stackrel{\text{def}}{=} P_{i,j,k_j,l}(x_j, t),
$$

where

$$
\overline{z}_{i,j} = \min \left(\max \left(\frac{\mu_t(y_j^{(i)} + \tau_*) - x_j}{\sigma_t}, -\tau_{\text{tail}} \sqrt{\log(1/\delta)} \right), \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right),
$$

$$
\underline{z}_{i,j} = \min \left(\max \left(\frac{\mu_t(y_j^{(i)} - \tau_*) - x_j}{\sigma_t}, -\tau_{\text{tail}} \sqrt{\log(1/\delta)} \right), \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right).
$$

Combining $(B.50)$ and $(B.52)$ with the last two displays, we have

$$
|p_{t_{*}+t}(\mathbf{x}) - g_{t_{*},t}(\mathbf{x})|
$$

\n
$$
\leq C_{S,1}(2D+1)\delta^{\frac{r_{\text{fail}}^2}{2} - D\overline{\tau}\tau_{\text{t}}} + C_{S,3}(\tau_{\text{sm}}+1)^D \left(\frac{e\tau_{*}}{\sigma_{t_{*}}}\right)^{\tau_{\text{sm}}} \delta^{-D\overline{\tau}\tau_{\text{t}}}
$$

\n
$$
+ \sum_{i=1}^{D_{1}} \sum_{0 \leq k \leq \tau_{\text{sm}}} \left| \frac{(\mathbf{D}^{\mathbf{k}} p_{t_{*}})(\mathbf{y}^{(i)})}{\mathbf{k}!} \right| (2\pi\sigma_{t}^2)^{-\frac{D}{2}} D2^{D-1} \tau_{\text{tail}}^D (1+\tau_{\text{tail}})^{D(\tau_{\text{sm}}+1)} \delta^{e\tau_{\text{tail}}^2} {\{\log(1/\delta)\}}^{D(\frac{\tau_{\text{sm}}}{2}+1)}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $0 < t \leq \tau_t \log(1/\delta)$, where $g_{t*,t} : \mathbb{R}^D \to \mathbb{R}$ is a function such that

$$
g_{t_{*},t}(\mathbf{x}) = \sum_{i=1}^{D_1} \sum_{0 \leq k < \tau_{\text{sm}}} \left\{ \frac{(\mathbf{D}^{\mathbf{k}} p_{t*})(\mathbf{y}^{(i)})}{\mathbf{k}!} \right\} \prod_{j=1}^{D} \sum_{l=0}^{D_2-1} P_{i,j,k_j,l}(x_j,t)
$$

for $\mathbf{x} \in \mathbb{R}^D$. Let

$$
m_* = \left[m^{1/D} + 1 \right]
$$
 and $t_* = m^{-\frac{2-2\eta_{ow}}{D}}$

with large enough $m \in \mathbb{R}$ so that $t_* \leq (2\overline{\tau})^{-1}$ and $m_* \in \mathbb{N}_{\geq 2}$. Since $\sigma_t^2 \geq 1 - \exp(-2\underline{\tau}t)$ for $t \geq 0$ and $1 - \exp(-x) \ge x/2$ for $0 \le x \le 1$, we have $\sigma_t^{-D} \le (\tau \delta)^{-D/2}$ for $t \ge \delta$ with $\delta \le \tau^{-1}$. Then,

$$
|p_{t_{*}+t}(\mathbf{x}) - g_{t_{*},t}(\mathbf{x})|
$$

\n
$$
\leq D_3 \left[\delta^{\frac{D\tau_{\text{tail}}^2 - D\overline{\tau}\tau_{\text{t}}}{2} + m_{*}^{-\tau_{\text{sm}}} t_{*}^{-\frac{\tau_{\text{sm}}}{2}} \delta^{-D\overline{\tau}\tau_{\text{t}}} \{ \log(1/\delta) \}^{\frac{\tau_{\text{sm}}}{2}} + m_{*}^{D} t_{*}^{-\frac{\tau_{\text{sm}}}{2}} \delta^{e\tau_{\text{tail}}^2 - D} \{ \log(1/\delta) \}^{D(\frac{\tau_{\text{sm}}}{2}+1)} \right]
$$

\n
$$
\leq D_3 \left[\delta^{\frac{D\tau_{\text{tail}}^2 - D\overline{\tau}\tau_{\text{t}}}{2} + m_{*}^{-\frac{\tau_{\text{low}}\tau_{\text{sm}}}{D}} \delta^{-D\overline{\tau}\tau_{\text{t}}} \{ \log(1/\delta) \}^{\frac{\tau_{\text{sm}}}{2}} + 2^D m^{\frac{D+1-\tau_{\text{low}}}{D}} \delta^{e\tau_{\text{tail}}^2 - D} \{ \log(1/\delta) \}^{D(\frac{\tau_{\text{sm}}}{2}+1)} \right]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $\delta \le t \le \tau_t \log(1/\delta)$, where $D_3 = D_3(D, \tau_{sm}, \tau, \tau_{tail}, C_{S,1}, C_{S,3})$ and the last inequality holds because $m^{1/D} \le m_* \le 2m^{1/D}$. Since $\tau_{\text{tail}}^2 = 4(D\overline{\tau}\tau_{\text{t}} + 1) \vee (D + 1)/e$, we have

$$
\frac{\tau_{\text{tail}}^2}{2} - D\overline{\tau}\tau_t \ge 1 \quad \text{and} \quad e\tau_{\text{tail}}^2 - D \ge 1.
$$

Then,

$$
|p_{t_*+t}(\mathbf{x}) - g_{t_*,t}(\mathbf{x})|
$$

\n
$$
\leq D_3 \left[\delta + m^{-\frac{\gamma_{\text{low}} \tau_{\text{sm}}}{D}} \delta^{-D\overline{\tau}\tau_t} \{ \log(1/\delta) \}^{\frac{\tau_{\text{sm}}}{2}} + 2^D m^{\frac{D+1-\tau_{\text{low}}}{D}} \delta \{ \log(1/\delta) \}^{D(\frac{\tau_{\text{sm}}}{2}+1)} \right]
$$
\n(B.53)

for $\mathbf{x} \in \mathbb{R}^D$ and $\delta \le t \le \tau_t \log(1/\delta)$. Similarly, with $u = 1$ and $h = j$ in [\(B.52\)](#page-71-0), the last integral in [\(B.52\)](#page-71-0) satisfies that

$$
\frac{1}{l!\sqrt{2\pi\sigma_t}}\int_{|x_j - \mu_t y_j| \leq \tau_{\text{tail}}\sigma_t \sqrt{\log(1/\delta)}} \left(\frac{\mu_t y_j - x_j}{\sigma_t} \right) (y_j - y_j^{(i)})^{k_j} \left(-\frac{(x_j - \mu_t y_j)^2}{2\sigma_t^2} \right)^l dy_j
$$
\n
$$
= \frac{1}{l!\sqrt{2\pi}} \int_{\substack{|z_j| \leq \tau_{\text{tail}} \sqrt{\log(1/\delta)} \\ |\mu_t^{-1}\sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)}| \leq \tilde{\tau}_*}} \left(\mu_t^{-1} \sigma_t z_j + \mu_t^{-1} x_j - y_j^{(i)} \right)^{k_j} z_j^{2l+1} (-2)^{-l} dz_j
$$
\n
$$
= \frac{(-2)^{-l} \mu_t^{-k_j}}{l!\sqrt{2\pi}} \sum_{r_j=0}^{k_j} {k_j \choose r_j} \sigma_t^{r_j} \left(x_j - \mu_t y_j^{(i)} \right)^{k_j - r_j} \left(\frac{\overline{z}_{i,j}^{r_j + 2l + 2} - \underline{z}_{i,j}^{r_j + 2l + 2}}{r_j + 2l + 2} \right) \stackrel{\text{def}}{=} \tilde{P}_{i,j,k_j,l}(x_j,t).
$$

Combining $(B.51)$ and $(B.52)$ with the last display, we have

$$
\begin{split} &\left|\sigma_{t}\left(\nabla p_{t_{*}+t}(\mathbf{x})\right)_{h}-\widetilde{g}_{t_{*},t}^{(h)}(\mathbf{x})\right| \\ &\leq C_{S,1}(2D+\sqrt{2/\pi})\delta^{\frac{\tau_{\rm tail}^{2}}{4}-D\overline{\tau}\tau_{\rm t}}+C_{S,3}\tau_{\rm tail}(\tau_{\rm sm}+1)^{D}\left(\frac{e\tau_{*}}{\sigma_{t_{*}}}\right)^{\tau_{\rm sm}}\delta^{-D\overline{\tau}\tau_{\rm t}}\sqrt{\log(1/\delta)} \\ &+\sum_{i=1}^{D_{1}}\sum_{0\leq k.<\tau_{\rm sm}}\left|\frac{(\mathbf{D}^{\mathbf{k}}p_{0})(\mathbf{y}^{(i)})}{\mathbf{k}!}\right|\left(2\pi\sigma_{t}^{2}\right)^{-\frac{D}{2}}D2^{D-1}\tau_{\rm tail}^{D}(1+\tau_{\rm tail})^{D(\tau_{\rm sm}+1)}\delta^{e\tau_{\rm tail}^{2}}\{\log(1/\delta)\}^{D(\frac{\tau_{\rm sm}}{2}+1)} \\ &\leq D_{4}\left[\delta^{\frac{\tau_{\rm tail}^{2}}{4}-D\overline{\tau}\tau_{\rm t}}+m_{*}^{-\tau_{\rm sm}}t_{*}^{-\frac{\tau_{\rm sm}}{2}}\delta^{-D\overline{\tau}\tau_{\rm t}}\{\log(1/\delta)\}^{\frac{\tau_{\rm sm}+1}{2}}+m_{*}^{D}t_{*}^{-\frac{\tau_{\rm sm}}{2}}\delta^{e\tau_{\rm tail}^{2}-D}\{\log(1/\delta)\}^{D(\frac{\tau_{\rm sm}}{2}+1)}\right] \\ &=D_{4}\left[\delta^{\frac{\tau_{\rm tail}^{2}}{4}-D\overline{\tau}\tau_{\rm t}}+m^{-\frac{\tau_{\rm low}\tau_{\rm sm}}{D}}\delta^{-D\overline{\tau}\tau_{\rm t}}\{\log(1/\delta)\}^{\frac{\tau_{\rm sm}+1}{2}}+2^{D}m^{\frac{D+1-\tau_{\rm low}}{D}}\delta^{e\tau_{\rm tail}^{2}-D}\{\log(1/\delta)\}^{D(\frac{\tau_{\rm sm}}{2}+1)}\right] \end{split}
$$

for $h \in [D]$, $\mathbf{x} \in \mathbb{R}^D$ and $\delta \le t \le \tau_t \log(1/\delta)$, where $D_4 = D_4(D, \tau_{sm}, \tau, \tau_{tail}, C_{S,1}, C_{S,3})$ and $\tilde{g}_{t_*,t}^{(h)}$: $\mathbb{R}^D \to \mathbb{R}, h \in [D]$ is a function such that

$$
\widetilde{g}_{t_*,t}^{(h)}(\mathbf{x}) = \sum_{i=1}^{D_1} \sum_{0 \leq k \, < \tau_{\text{sm}}} \left\{ \frac{(\mathbf{D}^k p_{t_*})(\mathbf{y}^{(i)})}{\mathbf{k}!} \right\} \left\{ \prod_{\substack{j=1 \ j \neq h}}^{D} \sum_{l=0}^{D_2-1} P_{i,j,k_j,l}(x_j,t) \right\} \left\{ \sum_{l=0}^{D_2-1} \widetilde{P}_{i,h,k_h,l}(x_h,t) \right\}.
$$

Since $\tau_{\text{tail}}^2 = 4(D\overline{\tau}\tau_t + 1) \vee (D+1)/e$, we have

$$
\frac{\tau_{\text{tail}}^2}{4} - D\overline{\tau}\tau_t \ge 1 \quad \text{and} \quad e\tau_{\text{tail}}^2 - D \ge 1.
$$

Then,

$$
\left|\sigma_{t}\left(\nabla p_{t_{*}+t}(\mathbf{x})\right)_{h} - \widetilde{g}_{t_{*},t}^{(h)}(\mathbf{x})\right|
$$
\n
$$
\leq D_{4}\left[\delta + m^{-\frac{\tau_{\text{low}}\tau_{\text{sm}}}{D}}\delta^{-D\overline{\tau}\tau_{t}}\left\{\log(1/\delta)\right\}^{\frac{\tau_{\text{sm}}+1}{2}} + 2^{D}m^{\frac{D+1-\tau_{\text{low}}}{D}}\delta\left\{\log(1/\delta)\right\}^{D(\frac{\tau_{\text{sm}}}{2}+1)}\right], \quad h \in [D] \tag{B.54}
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $\delta \le t \le \tau_t \log(1/\delta)$. Let

$$
\delta = m^{-\frac{\tau_{\text{low}}\tau_{\text{sm}}+D+1-\tau_{\text{low}}}{D(1+D\overline{\tau}\tau_{\text{t}})}}
$$

for large enough $m \in \mathbb{R}$. Combining [\(B.53\)](#page-73-0) and [\(B.54\)](#page-73-1) with the last display, we have

$$
|p_{t_{*}+t}(\mathbf{x}) - g_{t_{*},t}(\mathbf{x})| \le D_5 m^{-\frac{\tau_{\text{low}}\tau_{\text{sm}} - (D + 1 - \tau_{\text{low}})D\tau_{\tau_{t}}}{D(1 + D\overline{\tau}_{\tau_{t}})}} (\log m)^{D(\frac{\tau_{\text{sm}}}{2} + 1)},
$$

$$
\left| \sigma_t \left(\nabla p_{t_{*}+t}(\mathbf{x}) \right)_h - \tilde{g}_{t_{*},t}^{(h)}(\mathbf{x}) \right| \le D_5 m^{-\frac{\tau_{\text{low}}\tau_{\text{sm}} - (D + 1 - \tau_{\text{low}})D\tau_{\tau_{t}}}{D(1 + D\overline{\tau}_{\tau_{t}})}} (\log m)^{D(\frac{\tau_{\text{sm}}}{2} + 1)}, \quad h \in [D]
$$
\n(B.55)

for $\mathbf{x} \in \mathbb{R}^D$ and $\delta \le t \le \tau_t \log(1/\delta)$, where $D_5 = D_5(D, \overline{\tau}, \tau_t, \tau_{sm}, \tau_{low}, D_3, D_4)$.

Let $0 < \tilde{\delta} < \delta$ be a small enough value as described below. With $\tilde{\delta}^2 < 1/2$, Lemma [A.13](#page-37-0) implies that there exist neural networks $f_{\mu} \in \mathcal{F}_{NN}(L_{\mu}, \mathbf{d}_{\mu}, s_{\mu}, M_{\mu}), f_{\sigma} \in \mathcal{F}_{NN}(L_{\sigma}, \mathbf{d}_{\sigma}, s_{\sigma}, M_{\sigma})$ with

$$
L_{\mu}, L_{\sigma} \leq C_{N,4} \{ \log(1/\tilde{\delta}) \}^2, \quad \|\mathbf{d}_{\mu}\|_{\infty}, \|\mathbf{d}_{\sigma}\|_{\infty} \leq C_{N,4} \{ \log(1/\tilde{\delta}) \}^2
$$

\n
$$
s_{\mu}, s_{\sigma} \leq C_{N,4} \{ \log(1/\tilde{\delta}) \}^3, \quad M_{\mu}, M_{\sigma} \leq C_{N,4} \log(1/\tilde{\delta})
$$
\n(B.56)

such that $|\mu_t - f_\mu(t)| \leq \tilde{\delta}$ for $t \geq 0$ and $|\sigma_t - f_\sigma(t)| \leq \tilde{\delta}$ for $t \geq \tilde{\delta}$, where $C_{N,4}$ is the constant in Lemma [A.13.](#page-37-0) Also, Lemma [A.14](#page-37-1) implies that there exist a neural network $f_{\text{rec}} \in \mathcal{F}_{\text{NN}}(L_{\text{rec}}, \mathbf{d}_{\text{rec}}, s_{\text{rec}}, M_{\text{rec}})$ with

$$
L_{\text{rec}} \leq C_{N,5} \{ \log(1/\tilde{\delta}) \}^2, \quad \|\mathbf{d}_{\text{rec}}\|_{\infty} \leq C_{N,5} \{ \log(1/\tilde{\delta}) \}^3
$$

$$
s_{\text{rec}} \leq C_{N,5} \{ \log(1/\tilde{\delta}) \}^4, \quad M_{\text{rec}} \leq C_{N,5} \tilde{\delta}^{-2}
$$

such that $|1/x - f_{\text{rec}}(x)| \leq \tilde{\delta}$ for any $x \in [\tilde{\delta}, 1/\tilde{\delta}],$ where $C_{N,5}$ is the constant in Lemma [A.14.](#page-37-1) Since $\sigma_t - \tilde{\delta} \le f_\sigma(t) \le \sigma_t + \tilde{\delta}$ for $t \ge \tilde{\delta}$ and $\sqrt{\tau \delta} \le \sigma_t \le 1$ for $t \ge \delta$, we have $\delta \le f_\sigma(t) \le 2$ for $t \ge \delta$ with small enough δ so that $\delta \leq$ √ $\overline{\tau\delta} - \delta$ and $\delta \leq 1$. Then,

$$
|1/\sigma_t - f_{\text{rec}}(f_\sigma(t))| \le |1/\sigma_t - 1/f_\sigma(t)| + |1/f_\sigma(t) - f_{\text{rec}}(f_\sigma(t))|
$$

\n
$$
\le \{\sigma_t \wedge f_\sigma(t)\}^{-2} |\sigma_t - f_\sigma(t)| + \tilde{\delta} \le (1 + \delta^{-2})\tilde{\delta}
$$
\n(B.57)

for $t \geq \delta$. Since $\mu_t - \tilde{\delta} \leq f_\mu(t) \leq \mu_t + \tilde{\delta}$ for $t \geq \tilde{\delta}$ and $\delta^{\overline{\tau}\tau_t} \leq \mu_t \leq 1$ for $0 \leq t \leq \tau_t \log(1/\delta)$, we have $\delta^{\overline{\tau}\tau_{\rm t}}/2 \leq f_{\mu}(t) \leq 2$ for $\delta \leq t \leq \tau_{\rm t} \log(1/\delta)$ with small enough $\tilde{\delta}$ so that $\tilde{\delta} \leq \delta^{\overline{\tau}\tau_{\rm t}}/2$ and $\tilde{\delta} \leq 1$. Then,

$$
|1/\mu_t - f_{\text{rec}}(f_{\mu}(t))| \le |1/\mu_t - 1/f_{\mu}(t)| + |1/f_{\mu}(t) - f_{\text{rec}}(f_{\mu}(t))|
$$

$$
\le \{\mu_t \wedge f_{\mu}(t)\}^{-2}|\mu_t - f_{\mu}(t)| + \tilde{\delta} \le (1 + \delta^{-2\overline{\tau}\tau_t}/4)\tilde{\delta}
$$
 (B.58)

for $\delta \leq t \leq \tau_t \log(1/\delta)$. Lemma [A.9](#page-32-0) implies that for $k \geq 2$, there exists a neural network $\tilde{f}_{\text{mult}}^{(k)} \in$ $\mathcal{F}_{\text{NN}}(\widetilde{L}_{\text{mult}}^{(k)}, \widetilde{\mathbf{d}}_{\text{mult}}^{(k)}, \widetilde{s}_{\text{mult}}^{(k)}, \widetilde{M}_{\text{mult}}^{(k)})$ with

$$
\widetilde{L}_{\text{mult}}^{(k)} \leq C_{N,1} \log k \{ \log(1/\widetilde{\delta}) + \overline{\tau}\tau_t \log(1/\delta) \}, \quad \widetilde{d}_{\text{mult}}^{(k)} = (k, 48k, \dots, 48k, 1)^\top,
$$
\n
$$
\widetilde{s}_{\text{mult}}^{(2)} \leq C_{N,1} k \{ \log(1/\widetilde{\delta}) + \overline{\tau}\tau_t \log(1/\delta) \}, \quad \widetilde{M}_{\text{mult}}^{(k)} = \delta^{-\overline{\tau}\tau_t k}
$$

such that

$$
\left| \widetilde{f}_{\text{mult}}^{(k)}(\widetilde{x}_1, \dots, \widetilde{x}_k) - \prod_{i=1}^k x_i \right| \le \widetilde{\delta} + k \delta^{-\overline{\tau}\tau_t(k-1)} \widetilde{\epsilon}
$$
\n(B.59)

for any $\mathbf{x} = (x_1, \dots, x_k) \in \mathbb{R}^k$ with $\|\mathbf{x}\|_{\infty} \leq \delta^{-1}$ and $\widetilde{\mathbf{x}} = (\widetilde{x}_1, \dots, \widetilde{x}_k) \in \mathbb{R}^k$ with $\|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty} \leq \widetilde{\epsilon}$, where $0 < \tilde{\epsilon} \leq 1$ and $C_{N,1}$ is the constant in Lemma [A.9.](#page-32-0) Let

$$
f_{\text{clip}} \in \mathcal{F}_{\text{NN}}(2, (1, 2, 1)^{\top}, 7, \tau_{\text{tail}} \sqrt{\log(1/\delta)})
$$

be the neural network in Lemma [A.10](#page-32-1) such that $f_{\text{clip}}(x) = (x \vee - \tau_{\text{tail}} \sqrt{\log(1/\delta)}) \wedge \tau_{\text{tail}} \sqrt{\log(1/\delta)}$ for $x \in \mathbb{R}$. For $i \in [D_1]$ and $j \in [D]$, consider functions $\overline{f}_{i,j}, \underline{f}_{i,j} : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ such that

$$
\begin{split} \overline{f}_{i,j}(x,t) &= f_{\text{clip}}\left(\widetilde{f}_{\text{mult}}^{(2)}\left(f_{\text{rec}}\left(f_{\sigma}(t)\right), \{y_j^{(i)} + \tau_*\} f_{\mu}(t) - x\right)\right), \\ \underline{f}_{i,j}(x,t) &= f_{\text{clip}}\left(\widetilde{f}_{\text{mult}}^{(2)}\left(f_{\text{rec}}\left(f_{\sigma}(t)\right), \{y_j^{(i)} - \tau_*\} f_{\mu}(t) - x\right)\right), \end{split}
$$

 $\text{for } x, t \in \mathbb{R}$. Note that both $|y_j^{(i)} + \tau_*|$ and $|y_j^{(i)} - \tau_*|$ are upper bounded by $m_*\tau_* \leq (1 + \tau_{\text{tail}})\sqrt{\log(1/\delta)}$. Then, for $|x| \leq 1 + \tau_x \sqrt{\log(1/\delta)}$, both $|(y_j^{(i)} + \tau_*)\mu_t - x|$ and $|(y_j^{(i)} - \tau_*)\mu_t - x|$ are upper bounded by $\delta^{-\overline{\tau}_{\tau_t}}$ with small enough δ . Combining [\(B.56\)](#page-74-0), [\(B.57\)](#page-74-1) and [\(B.59\)](#page-74-2) with the last display, both $|\overline{z}_{i,j} - f_{i,j}(x,t)|$ and $|\underline{z}_{i,j} - \underline{f}_{i,j}(x,t)|$ are bounded by

$$
\widetilde{\delta} + 2\delta^{-\overline{\tau}\tau_t} (1 + \delta^{-2\overline{\tau}\tau_t}/4) \widetilde{\delta} \le 5\delta^{-3\overline{\tau}\tau_t} \widetilde{\delta},\tag{B.60}
$$

for $|x| \leq \mu_t + \tau_x \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \tau_t \log(1/\delta)$. Since $\mu_t - \tilde{\delta}^2 \leq f_\mu(t) \leq \mu_t + \tilde{\delta}^2$ for $t \geq \tilde{\delta}^2$, we have $1/4 \leq 1/2 - \tilde{\delta}^2 \leq f_\mu(t) \leq 1 + \tilde{\delta}^2 \leq 2$ and $1/4 \leq \mu_t \leq 2$ for $\delta \leq t \leq D_1$ with small enough $\tilde{\delta}$ by [\(B.1\)](#page-41-0). For any $i \in [D_2], j \in [D], k \in \{0, \ldots, \tau_{sm} - 1\}$, consider functions $f_{i,j,k}, \tilde{f}_{i,j,k} : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ such that

$$
f_{i,j,k}(x,t) = \sum_{l=0}^{D_2-1} \sum_{r=0}^k {k \choose r} \left\{ \frac{(-2)^{-l}}{k!l!\sqrt{2\pi}(r+2l+1)} \right\} \left\{ \overline{f}_{i,j,k,l,r,r+2l+1} - \underline{f}_{i,j,k,l,r,r+2l+1} \right\},
$$

$$
\widetilde{f}_{i,j,k}(x,t) = \sum_{l=0}^{D_2-1} \sum_{r=0}^k {k \choose r} \left\{ \frac{(-2)^{-l}}{k!l!\sqrt{2\pi}(r+2l+2)} \right\} \left\{ \overline{f}_{i,j,k,l,r,r+2l+2} - \underline{f}_{i,j,k,l,r,r+2l+2} \right\},
$$

for $x, t \in \mathbb{R}$, where

$$
\overline{f}_{i,j,k,l,r,s} = \tilde{f}_{\text{mult}}^{(2k+s)} \left(f_{\text{rec}} \left(f_{\mu}(t) \right) \cdot \mathbf{1}_k, f_{\sigma}(t) \cdot \mathbf{1}_r, \left\{ x - y_j^{(i)} f_{\mu}(t) \right\} \cdot \mathbf{1}_{k-r}, \overline{f}_{i,j}(x, t) \cdot \mathbf{1}_s \right),
$$
\n
$$
\underline{f}_{i,j,k,l,r,s} = \tilde{f}_{\text{mult}}^{(2k+s)} \left(f_{\text{rec}} \left(f_{\mu}(t) \right) \cdot \mathbf{1}_k, f_{\sigma}(t) \cdot \mathbf{1}_r, \left\{ x - y_j^{(i)} f_{\mu}(t) \right\} \cdot \mathbf{1}_{k-r}, \underline{f}_{i,j}(x, t) \cdot \mathbf{1}_s \right),
$$

for $s \in \{r+2l+1, r+2l+2\}$. Combining [\(B.56\)](#page-74-0), [\(B.58\)](#page-74-3), [\(B.60\)](#page-75-0) and [\(B.59\)](#page-74-2) with the last two displays, we have

$$
\begin{aligned}\n\left| \sum_{l=0}^{D_2-1} \frac{P_{i,j,k,l}(x,t)}{k!} - f_{i,j,k}(x,t) \right| \\
\leq \sum_{l=0}^{D_2-1} \sum_{r=0}^k {k \choose r} \left\{ \frac{2^{-l+1}}{k!l!\sqrt{2\pi}(2r+r+2l+1)} \right\} \left\{ \widetilde{\delta} + 5(2k+r+2l+1)\delta^{-(2k+r+2l+3)\overline{\tau}r_t} \widetilde{\delta} \right\}\n\end{aligned}
$$

and

$$
\left| \sum_{l=0}^{D_2-1} \frac{P_{i,j,k,l}(x,t)}{k!} - f_{i,j,k}(x,t) \right|
$$

\n
$$
\leq \sum_{l=0}^{D_2-1} \sum_{r=0}^k {k \choose r} \left\{ \frac{2^{-l+1}}{k!l!\sqrt{2\pi}(r+2l+2)} \right\} \left\{ \widetilde{\delta} + 5(2k+r+2l+2)\delta^{-(2k+r+2l+4)\overline{\tau}_{\tau_k}} \widetilde{\delta} \right\}
$$

for $i \in [D_1], j \in [D], k \in \{0, \ldots, \tau_{sm} - 1\}, |x| \leq \mu_t + \tau_x \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \tau_t \log(1/\delta)$ with small enough δ so that σ_t , $|x - \mu_t y_j^{(i)}|$ $\int_{j}^{(i)} |, \tau_{\text{tail}} \sqrt{\log(1/\delta)}$ are bounded by $\delta^{-\overline{\tau}\tau_{\text{t}}}$. Since $\sum_{r=0}^{k} {k \choose r}$ $\binom{k}{r} = 2^k$ and $2^{k}/k! \leq 2$ for all $k \in \mathbb{N}$, the last two displays are bounded by

$$
\sum_{l=0}^{D_2-1} \left(\frac{2^{-l+2}}{l!\sqrt{2\pi}(2l+1)} \right) \left\{ 1 + 5(3\tau_{\rm sm} + 2l-1)\delta^{-(3\tau_{\rm sm} + 2l+1)\bar{\tau}\tau_{\rm t}} \right\} \tilde{\delta} \leq \delta^{-D_6 \log(1/\delta)} \tilde{\delta},\tag{B.61}
$$

where $D_6 = D_6(\bar{\tau}, \tau_{tail}, \tau_{sm}, \tau_t)$. For any $i \in [D_1], j \in [D], k \in \{0, \ldots, \tau_{sm}-1\}, l \in \{0, \ldots, D_2-1\}, |x| \leq$ $\mu_t + \tau_x \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \tau_t \log(1/\delta)$, we have

$$
\left|\frac{P_{i,j,k,l}(x,t)}{k!}\right| \leq \left\{\frac{2^{-l+1}\mu_t^{-k}}{k!l!\sqrt{2\pi}(r+2l+1)}\sum_{r=0}^k \binom{k}{r}\right\} \left\{(2+\tau_x+\tau_{\text{tail}})\sqrt{\log(1/\delta)}\right\}^k \left\{\tau_{\text{tail}}\sqrt{\log(1/\delta)}\right\}^{k+2l+1}
$$

and

$$
\left| \frac{\widetilde{P}_{i,j,k,l}(x,t)}{k!} \right| \leq \left\{ \frac{2^{-l+1} \mu_t^{-k}}{k! l! \sqrt{2\pi} (r+2l+2)} \sum_{r=0}^k \binom{k}{r} \right\} \left\{ (2+\tau_x+\tau_{\text{tail}}) \sqrt{\log(1/\delta)} \right\}^k \left\{ \tau_{\text{tail}} \sqrt{\log(1/\delta)} \right\}^{k+2l+2}
$$

because $\mu_t \leq 1, \sigma_t \leq 1$, and $y_j^{(i)} \leq m_* \tau_* \leq (1 + \tau_{\text{tail}}) \sqrt{\log(1/\delta)}$. Since $\sum_{r=0}^k {k \choose r}$ $r \choose r = 2^k$ and $2^k/k! \leq 2$ for all $k \in \mathbb{N}$, the last two displays are bounded by

$$
\left(\frac{4}{\sqrt{2\pi}}\right)\delta^{-k\overline{\tau}\tau_{\rm t}}\left\{(2+\tau_{\rm x}+\tau_{\rm tail})^2\tau_{\rm tail}^2\log(1/\delta)\right\}^{k+l+1}.
$$

Then, both $|\sum_{l=0}^{D_2-1} P_{i,j,k,l}(x,t)/k!|$ and $|\sum_{l=0}^{D_2-1} \widetilde{P}_{i,j,k,l}(x,t)/k!|$ are upper bounded by

$$
D_2\left(\frac{4}{\sqrt{2\pi}}\right)\delta^{-(\tau_{\rm sm}-1)\bar{\tau}\tau_{\rm t}}\left\{(2+\tau_{\rm x}+\tau_{\rm tail})^2\tau_{\rm tail}^2\log(1/\delta)\right\}^{\tau_{\rm sm}+D_2-1}\leq \{\log(1/\delta)\}^{D_7\log(1/\delta)}
$$

for $i \in [D_1], j \in [D], k \in \{0, \ldots, \tau_{sm} - 1\}, |x| \leq \mu_t + \tau_x \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \tau_t \log(1/\delta)$, where $D_7 = D_7(\tau_{\text{tail}}, \tau_{\text{sm}}, \tau_{\text{x}}, \tau_{\text{t}}, \overline{\tau})$. Consider functions $f_{t_*}, \widetilde{f}_{t_*}^{(1)}, \ldots, \widetilde{f}_{t_*}^{(D)} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}$ such that

$$
f_{t*}(\mathbf{x},t) = \sum_{i=1}^{D_1} \sum_{0 \le k.<\tau_{\text{sm}}} \left\{ (\mathbf{D}^k p_{t*})(\mathbf{y}^{(i)}) \right\} f_{\text{mult}}^{(D)}(f_{i,1,k_1}(x_1,t),\ldots,f_{i,D,k_D}(x_D)) \text{ and}
$$

$$
\tilde{f}_{t*}^{(h)}(\mathbf{x},t) = \sum_{i=1}^{D_1} \sum_{0 \le k.<\tau_{\text{sm}}} \left\{ (\mathbf{D}^k p_{t*})(\mathbf{y}^{(i)}) \right\} f_{\text{mult}}^{(D)}\left(\tilde{f}_{i,h,k_h}(x_h),\underbrace{f_{i,1,k_1}(x_1,t),\ldots,f_{i,D,k_D}(x_D)}_{\text{without }f_{i,h,k_h}(x_h,t)} \right),
$$

where $f_{\text{mult}}^{(D)} \in \mathcal{F}_{\text{NN}}(L_{\text{mult}}^{(D)}, \mathbf{d}_{\text{mult}}^{(D)}, s_{\text{mult}}^{(D)}, M_{\text{mult}}^{(D)})$ is the neural network in Lemma [A.9](#page-32-0) with $L_{\text{mult}}^{(D)} \leq C_{N,1} \log D[\log(1/\tilde{\delta}) + DD_7 \log(1/\delta) \log \log(1/\delta)], \quad d_{\text{mult}}^{(D)} = (D, 48D, \dots, 48D, 1)^{\top},$ $s_{\text{mult}}^{(D)} \leq C_{N,1} D[\log(1/\tilde{\delta}) + D_7 \log(1/\delta) \log \log(1/\delta)], \quad M_{\text{mult}}^{(D)} = {\log(1/\delta)}^{\text{DD}_7 \log(1/\delta)}$

such that

$$
\left| \widetilde{f}_{\text{mult}}^{(D)}(\widetilde{x}_1,\ldots,\widetilde{x}_D) - \prod_{i=1}^D x_i \right| \le \widetilde{\delta} + D\{\log(1/\delta)\}^{(D-1)D_7 \log(1/\delta)} \widetilde{\epsilon}
$$

for any $\mathbf{x} = (x_1, \dots, x_D) \in \mathbb{R}^D$ with $\|\mathbf{x}\|_{\infty} \leq {\log(1/\delta)}^{\text{DD}_7 \log(1/\delta)}$ and $\widetilde{\mathbf{x}} = (\widetilde{x}_1, \dots, \widetilde{x}_D) \in \mathbb{R}^D$ with $\|\mathbf{x} - \tilde{\mathbf{x}}\|_{\infty} \leq \tilde{\epsilon}$. Combining [\(B.61\)](#page-75-1) with the last display, we have

$$
|f_{t*}(\mathbf{x},t) - g_t(\mathbf{x})|
$$

\n
$$
\leq \sum_{i=1}^{D_1} \sum_{0 \leq k \cdot \langle \tau_{sm}} |(D^k p_{t*})(\mathbf{y}^{(i)})| \left\{ 1 + D\delta^{-D_6 \log(1/\delta)} \{ \log(1/\delta) \}^{(D-1)D_7 \log(1/\delta)} \right\} \tilde{\delta}
$$

\n
$$
\leq D_1 \tau_{sm}^D C_{S,3} \sigma_{t*}^{-\tau_{sm}} \left\{ 1 + D\delta^{-D_6 \log(1/\delta)} \{ \log(1/\delta) \}^{(D-1)D_7 \log(1/\delta)} \right\} \tilde{\delta}
$$

\n
$$
\leq m^{\frac{\tau_{sm}(1-\tau_{low})+D}{D}} \delta^{-D_8 \log(1/\delta)} \tilde{\delta}
$$

for $||\mathbf{x}||_{\infty} \leq \mu_t + \tau_{\mathbf{x}} \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \tau_t \sqrt{\log(1/\delta)}$, where $D_8 = D_8(D, \tau, \tau_{\rm sm}, C_{S,3}, D_6, D_7)$. Similarly, we have

$$
\left| \tilde{f}_{t_*}^{(h)}(\mathbf{x},t) - \tilde{g}_t^{(h)}(\mathbf{x}) \right|
$$
\n
$$
\leq \sum_{i=1}^{D_1} \sum_{\substack{0 \leq k.<\tau_{\text{sm}}} } \left| \left(D^k p_{t_*} \right) (\mathbf{y}^{(i)}) \right| \left\{ 1 + D\delta^{-D_6 \log(1/\delta)} \{ \log(1/\delta) \}^{(D-1)D_7 \log(1/\delta)} \right\} \tilde{\delta}
$$
\n
$$
\leq m^{\frac{\tau_{\text{sm}}(1-\tau_{\text{low}})+D}{D}} \delta^{-D_8 \log(1/\delta)} \tilde{\delta}, \quad h \in [D]
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \tau_{\mathbf{x}} \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \tau_{\mathbf{t}} \sqrt{\log(1/\delta)}$. Let

$$
\widetilde{\delta} = \delta^{D_8 \log(1/\delta)} m^{-\frac{\tau_{\rm sm} + D}{D}}
$$

with large enough m. Combining $(B.55)$ with the second last display, we have

$$
|p_{t_*+t}(\mathbf{x}) - f_{t_*}(\mathbf{x}, t)| \le (1 + D_5) m^{-\frac{\tau_{\text{low}} \tau_{\text{sm}} - (D + 1 - \tau_{\text{low}}) D \overline{\tau}_{\tau_t}}{D(1 + D \overline{\tau}_{\tau_t})}} (\log m)^{D(\frac{\tau_{\text{sm}}}{2} + 1)},
$$

$$
\left| \sigma_t \left(\nabla p_{t_*+t}(\mathbf{x}) \right)_h - \tilde{f}_{t_*}^{(h)}(\mathbf{x}, t) \right| \le (1 + D_5) m^{-\frac{\tau_{\text{low}} \tau_{\text{sm}} - (D + 1 - \tau_{\text{low}}) D \overline{\tau}_{\tau_t}}{D(1 + D \overline{\tau}_{\tau_t})}} (\log m)^{D(\frac{\tau_{\text{sm}}}{2} + 1)}, \quad h \in [D]
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \tau_{\text{tail}} \sigma_t \sqrt{\log(1/\delta)}$ and $\delta \leq t \leq \tau_t \sqrt{\log(1/\delta)}$. Consider a function $\mathbf{f}_{t_*}: \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}^{D+1}$ such that $\mathbf{f}_{t_*}(\mathbf{x},t) = (f_{t_*}(\mathbf{x},t), \tilde{f}_{t_*}^{(1)}(\mathbf{x},t), \dots, \tilde{f}_{t_*}^{(D)}(\mathbf{x},t))^{\top}$ for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Lemma [A.4,](#page-29-0) Lemma [A.5,](#page-29-1) Lemma [A.6](#page-30-0) and Lemma [A.7](#page-30-1) implies that $f_{t_*} \in \mathcal{F}_{NN}(L, d, s, m)$ with

$$
L \le D_9(\log m)^4, \quad \|\mathbf{d}\|_{\infty} \le D_9 m(\log m)^9,
$$

$$
s \le D_9 m(\log m)^9, \quad M \le \exp(D_9(\log m)^2),
$$

where $D_9 = D_9(D, \overline{\tau}, \underline{\tau}, \tau_t, \tau_x, \tau_{sm}, \tau_{low}, \tau_{tail}, C_{S,3}, C_{N,4}, C_{N,5}, D_7, D_8)$ is a large enough constant. The assertion follows by re-defining the constants. \Box

B.5 Proof of Theorem [5.1](#page-11-0)

In this subsection, we provide the proof of Theorem [5.1](#page-11-0) by combining Propositions [B.1](#page-39-0) to [B.3.](#page-40-0)

Proof of Theorem [5.1.](#page-11-0) Let $m > 0$ be a large enough value as described below and

$$
D_1 = \sqrt{\frac{8\beta}{d}} \vee \sqrt{2\tau_{\min} + \frac{4\beta}{d}}.
$$

We will approximate $\nabla \log p_t(\mathbf{x})$ for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le \tau_{\max} \log m$ by neural networks by dividing the analysis into the following three cases:

- 1. (Interior of near-support) $\|\mathbf{x}\|_{\infty} \leq \mu_t {\log(1/\sigma_t)}^{-3/2}$ and $m^{-\tau_{\min}} \leq t \leq 3m^{-\frac{1}{2D}}$
- 2. (Boundary of near-support) $\mu_t \tau_{\min}^{3/2} \{ (4D)^{3/2} + 3 \} \{ \log(1/\sigma_t) \}^{-3/2} \le ||\mathbf{x}||_{\infty} \le \mu_t + \sigma_t D_1 \sqrt{\frac{\mu_t}{\sigma_t}}$ $\overline{\log m}$ and $m^{-\tau_{\min}} \leq t \leq 3m^{-\frac{1}{2D}}$
- 3. (large t) $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $2m^{-\frac{1}{2D}} \le t \le \tau_{\max} \log m$.

Then, we combine the networks into a single network and derive the approximation error over the entire region $(\mathbf{x}, t) \in \mathbb{R}^D \times [m^{-\tau_{\min}}, \tau_{\max} \log m].$

B.5.1 Interior of near-support

Let
$$
\tau_{tail}^{(1)} = 4D\beta/d
$$
 and $\tau_{bd}^{(1)} = 3/2$. Let
\n
$$
\widetilde{C}_3 = \widetilde{C}_3(\beta, d, D, K, \overline{\tau}, \underline{\tau}, \tau_{bd}^{(1)}, \tau_{tail}^{(1)}, \tau_{min}),
$$
\n
$$
\widetilde{C}_4 = \widetilde{C}_4(\beta, d, D, K, \underline{\tau}, \tau_{bd}^{(1)}, \tau_{tail}^{(1)}, \tau_{min}),
$$
\n
$$
\widetilde{C}_5 = \widetilde{C}_5(\beta, d, \underline{\tau}, \tau_{bd}^{(1)}, \tau_{tail}^{(1)}, \tau_{min})
$$

be the constants in Proposition [B.1,](#page-39-0) where (τ_{tail}, τ_{bd}) is replaced by $(\tau_{tail}^{(1)}, \tau_{bd}^{(1)})$. Also, let \widetilde{C}_2 = $\widetilde{C}_2(\beta, D, \tau_{\text{bd}}^{(1)}, \tau_{\text{tail}}^{(1)})$ be the constant in Lemma [B.3,](#page-39-1) where $(\tau_{\text{tail}}, \tau_{\text{bd}})$ is replaced by $(\tau_{\text{tail}}^{(1)}, \tau_{\text{bd}}^{(1)})$. For large enough m so that $m \geq \tilde{C}_5$ and $3m^{-1/(2D)} \leq \overline{\tau}^{-1}(\tilde{C}_2^2 \wedge 1/2)$, Proposition [B.1](#page-39-0) implies that there exists a class of permutation matrices $\mathcal{P}^{(1)} = \{ \mathcal{Q}_i^{(1)}, \mathcal{R}_i^{(1)} \}$ $\{i^{(1)}\}_{i\in[L-1]}$ and weight-sharing neural networks

$$
\mathbf{f}^{(1)} = (f_1^{(1)}, \dots, f_{D+1}^{(1)})^\top \in \mathcal{F}_{\text{NN}}(L^{(1)}, \mathbf{d}^{(1)}, s^{(1)}, M^{(1)}, \mathbf{m}^{(1)}, \mathcal{P}^{(1)})
$$

with

$$
L^{(1)} \leq \widetilde{C}_3 (\log m)^2 \log \log m, \quad \|\mathbf{d}^{(1)}\|_{\infty} \leq \widetilde{C}_3 m^{D+1} (\log m)^3,
$$

$$
s^{(1)} \leq \widetilde{C}_3 m (\log m)^5 \log \log m, \quad M^{(1)} \leq \exp(\widetilde{C}_3 {\log m}^2),
$$

$$
\|\mathbf{m}^{(1)}\|_{\infty} = m^D
$$

satisfying

$$
\left\| \begin{pmatrix} \sigma_t \nabla p_t(\mathbf{x}) \\ p_t(\mathbf{x}) \end{pmatrix} - \mathbf{f}^{(1)}(\mathbf{x}, t) \right\|_{\infty} \le \widetilde{C}_4 \left(3^{\frac{4D\beta}{d}} + 1 \right) m^{-\frac{\beta}{d}} (\log m)^{2(D-1+\beta\vee 1)} \stackrel{\text{def}}{=} \epsilon_1
$$

 $\|x\|_{\infty} \leq \mu_t - {\log(1/\sigma_t)}^{-3/2}$ and $m^{-\tau_{\min}} \leq t \leq 3m^{-\frac{1}{2D}}$ because $\mu_t \leq 1$. Note that $C_{S,1}^{-1} \leq p_t(\mathbf{x}) \leq t$ $C_{S,1}$ for $\|\mathbf{x}\|_{\infty} \leq \mu_t$, $t \geq 0$, and $\|\sigma_t \nabla p_t(\mathbf{x})\|_{\infty} \leq C_{S,3}$ for $\mathbf{x} \in \mathbb{R}^D$, $t \geq 0$, where $C_{S,1} = C_{S,1}(D, K, \tau_1)$ and $C_{S,3} = C_{S,3}(D, K, \overline{\tau}, \underline{\tau})$ are the constants in Lemma [A.1](#page-28-0) and Lemma [A.3,](#page-29-2) respectively. Also, $C_{S,1}^{-1}$ $S_{S,1}^{-1/2}$ ≤ $p_t(\mathbf{x}) - \epsilon_1$ ≤ $f_{D+1}^{(1)}(\mathbf{x}, t)$ ≤ $p_t(\mathbf{x}) + \epsilon_1$ ≤ 2C_{S,1} for $\|\mathbf{x}\|_{\infty}$ ≤ $\mu_t - \mu_t \{\log(1/\sigma_t)\}^{-3/2}$ and $m^{-\tau_{\min}}$ ≤ $t \leq 3m^{-\frac{1}{2D}}$ with large enough m so that $\epsilon_1 \leq C_{S,1}^{-1}$ $\mathcal{L}_{S,1}^{-1}/2$. Let $f_{\text{rec}}^{(\text{in})} \in \mathcal{F}_{\text{NN}}(L_{\text{rec}}^{(\text{in})}, \mathbf{d}_{\text{rec}}^{(\text{in})}, s_{\text{rec}}^{(\text{in})}, M_{\text{rec}}^{(\text{in})})$ be the neural networks in Lemma [A.14](#page-37-1) with

$$
L_{\text{rec}}^{(\text{in})} \leq C_{N,5} \{ \beta \log m/d \}^2, \quad \|\mathbf{d}_{\text{rec}}^{(\text{in})}\|_{\infty} \leq C_{N,5} \{ \beta \log m/d \}^3,
$$

$$
s_{\text{rec}}^{(\text{in})} \leq C_{N,5} \{ \beta \log m/d \}^4, \quad M_{\text{rec}}^{(\text{in})} \leq C_{N,5} m^{\frac{2\beta}{d}}
$$

such that $|1/x - f_{\text{rec}}^{(\text{in})}(x)| \le m^{-\beta/d}$ for $x \in [m^{-\beta/d}, m^{\beta/d}]$. For large enough m so that $m^{-\beta/d} \le C_{S,1}^{-1}$ $\binom{r-1}{S,1}/2,$

$$
\left|\frac{1}{p_t(\mathbf{x})} - f_{\text{rec}}^{(\text{in})} \left(f_{D+1}^{(1)}(\mathbf{x}, t) \right) \right| \le \left|\frac{1}{p_t(\mathbf{x})} - \frac{1}{f_{D+1}^{(1)}(\mathbf{x}, t)} \right| + \left|\frac{1}{f_{D+1}^{(1)}(\mathbf{x}, t)} - f_{\text{rec}}^{(\text{in})} \left(f_{D+1}^{(1)}(\mathbf{x}, t) \right) \right|
$$

$$
\le \{p_t(\mathbf{x}) \wedge f_{D+1}^{(1)}(\mathbf{x}, t)\}^{-2} \epsilon_1 + m^{-\frac{\beta}{d}} \le (4C_{S,1}^2 + 1)\epsilon_1
$$

for $||\mathbf{x}||_{\infty} \leq \mu_t - {\log(1/\sigma_t)}^{-3/2}$ and $m^{-\tau_{\min}} \leq t \leq 3m^{-\frac{1}{2D}}$. Let

$$
f_{\text{mult}}^{(\text{in})} \in \mathcal{F}_{\text{NN}}(L_{\text{mult}}^{(\text{in})}, \mathbf{d}_{\text{mult}}^{(\text{in})}, s_{\text{mult}}^{(\text{in})}, M_{\text{mult}}^{(\text{in})})
$$

be the neural networks in Lemma [A.9](#page-32-0) with

$$
L_{\text{mult}}^{(\text{in})} \leq C_{N,1} \log 2 \{ \beta \log m/d + 2 \log(C_{S,1} \vee C_{S,3}) + 2 \log 2 \}, \quad \mathbf{d}_{\text{mult}}^{(\text{in})} = (2, 96, \dots, 96, 1)^{\top},
$$

$$
s_{\text{mult}}^{(\text{in})} \leq C_{N,1} 2 \{ \beta \log m/d + \log(C_{S,1} \vee C_{S,3}) + \log 2 \}, \quad M_{\text{mult}}^{(\text{in})} = (C_{S,1} \vee C_{S,3})^2
$$

such that $|f_{\text{mult}}^{(\text{in})}(\tilde{\mathbf{x}}) - x_1x_2| \leq m^{-\beta/d} + 2(C_{S,1} \vee C_{S,3})\tilde{\epsilon}$ for all $\|\mathbf{x}\|_{\infty} \leq 2(C_{S,1} \vee C_{S,3}), \tilde{\mathbf{x}} \in \mathbb{R}^2$ with $\|\mathbf{x} - \tilde{\mathbf{x}}\|_{\infty} \leq \tilde{\epsilon}$ and $|f_{\text{mult}}^{(\text{in})}(\mathbf{x})| \leq (C_{S,1} \vee C_{S,3})^2$ for all $\mathbf{x} \in \mathbb{R}^2$, where $C_{N,1}$ is the constant in Lemma [A.9.](#page-32-0) Then,

$$
\left| \frac{\sigma_t(\nabla p_t(\mathbf{x}))_i}{p_t(\mathbf{x})} - f_{\text{mult}}^{(\text{in})} \left(f_i^{(1)}(\mathbf{x}, t), f_{\text{rec}}^{(\text{in})} \left(f_{D+1}^{(1)}(\mathbf{x}, t) \right) \right) \right|
$$
\n
$$
\leq m^{-\frac{\beta}{d}} + 2(C_{S,1} \vee C_{S,3}) (4C_{S,1}^2 + 1)\epsilon_1 \leq D_2 m^{-\frac{\beta}{d}} (\log m)^{2(D-1+\beta \vee 1)}, \quad i \in [D]
$$
\n(B.62)

for $||\mathbf{x}||_{\infty} \leq \mu_t - {\log(1/\sigma_t)}^{-3/2}$ and $m^{-\tau_{\min}} \leq t \leq 3m^{-\frac{1}{2D}}$, where $D_2 = D_2(\beta, D, C_{S,1}, C_{S,3}, \tilde{C}_4)$. Let $0 < \delta \leq T = m^{-\tau_{\min}}$ be a small enough value as described below. With $\delta < 1/2$, Lemma [A.13](#page-37-0) implies that there exist neural networks $f_{\sigma} \in \mathcal{F}_{NN}(L_{\sigma}, \mathbf{d}_{\sigma}, s_{\sigma}, M_{\sigma})$ with

$$
L_{\sigma} \leq C_{N,4} \{ \log(1/\delta) \}^2, \quad \|\mathbf{d}_{\sigma}\|_{\infty} \leq C_{N,4} \{ \log(1/\delta) \}^2
$$

$$
s_{\sigma} \leq C_{N,4} \{ \log(1/\delta) \}^3, \quad M_{\sigma} \leq C_{N,4} \log(1/\delta)
$$
 (B.63)

such that $|\sigma_t - f_\sigma(t)| \leq \delta$ for $t \geq \delta$, where $C_{N,4}$ is the constant in Lemma [A.13.](#page-37-0) Also, Lemma [A.14](#page-37-1) implies that there exists a neural network $f_{\text{rec}} \in \mathcal{F}_{\text{NN}}(L_{\text{rec}}, \mathbf{d}_{\text{rec}}, s_{\text{rec}}, M_{\text{rec}})$ with

$$
L_{\text{rec}} \leq C_{N,5} \{ \log(1/\delta) \}^2, \quad \|\mathbf{d}_{\text{rec}}\|_{\infty} \leq C_{N,5} \{ \log(1/\delta) \}^3
$$

$$
s_{\text{rec}} \leq C_{N,5} \{ \log(1/\delta) \}^4, \quad M_{\text{rec}} \leq C_{N,5} \delta^{-2}
$$

such that $|1/x - f_{\text{rec}}(x)| \leq \delta$ for any $x \in [\delta, 1/\delta]$. Since $\sigma_t - \delta \leq f_{\sigma}(t) \leq \sigma_t + \delta$ for $t \geq \delta$ and √ $\overline{\tau T}\leq \sigma_t\leq 1$ for $t\geq \underline{T}$, we have $\underline{T}\leq f_\sigma(t)\leq 2$ for $t\geq \underline{T}$ with large enough m so that $\underline{T}\leq$ √ $\overline{\tau}\overline{T}-\underline{T}$ and $\underline{T} \leq 1$. Then,

$$
|1/\sigma_t - f_{\text{rec}}(f_\sigma(t))| \le |1/\sigma_t - 1/f_\sigma(t)| + |1/f_\sigma(t) - f_{\text{rec}}(f_\sigma(t))|
$$

\n
$$
\le \{\sigma_t \wedge f_\sigma(t)\}^{-2} |\sigma_t - f_\sigma(t)| + \delta \le (1 + \underline{T}^{-2})\delta = (1 + m^{2\tau_{\text{min}}})\delta
$$
\n(B.64)

for $t \geq T$. Lemma [A.9](#page-32-0) implies that there exists a neural network

$$
f_{\text{mult}} \in \mathcal{F}_{\text{NN}}(L_{\text{mult}}, \mathbf{d}_{\text{mult}}, s_{\text{mult}}, M_{\text{mult}})
$$

with

$$
L_{\text{mult}} \leq C_{N,1} \log 2 \{ \log(1/\delta) + 3DD_1^2 \log m \}, \quad \mathbf{d}_{\text{mult}} = (2, 96, \dots, 96, 1)^\top,
$$

$$
s_{\text{mult}} \leq C_{N,1} 2 \{ \log(1/\delta) + 3DD_1^2 \log m \}, \quad M_{\text{mult}} = m^{6DD_1^2}
$$

such that

$$
|f_{\text{mult}}(\tilde{x}_1, \tilde{x}_2) - x_1 x_2| \le \delta + 2m^{3DD_1^2} \tilde{\epsilon}
$$
 (B.65)

for any $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2$ with $\|\mathbf{x}\|_{\infty} \leq m^{3DD_1^2}$ and $\tilde{\mathbf{x}} = (\tilde{x}_1, \tilde{x}_2) \in \mathbb{R}^k$ with $\|\mathbf{x} - \tilde{\mathbf{x}}\|_{\infty} \leq \tilde{\epsilon}$, where $0 < \tilde{\epsilon} \le 1$. Consider functions $\tilde{f}_1^{(1)}, \ldots, \tilde{f}_D^{(1)} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}$ such that

$$
\widetilde{f}_i^{(1)}(\mathbf{x},t) = f_{\text{mult}}\left(f_{\text{mult}}^{(\text{in})}\left(f_i^{(1)}(\mathbf{x},t), f_{\text{rec}}^{(\text{in})}\left(f_{D+1}^{(1)}(\mathbf{x},t)\right)\right), f_{\text{rec}}(f_{\sigma}(t))\right), \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. For large enough m so that $m^{3DD_1^2} \geq 4(C_{S,1} \vee C_{S,3})^2$, we have

$$
\left| \sigma_t^{-1} f_{\text{mult}}^{(\text{in})} \left(f_i^{(1)}(\mathbf{x}, t), f_{\text{rec}}^{(\text{in})} \left(f_{D+1}^{(1)}(\mathbf{x}, t) \right) \right) - \tilde{f}_i^{(1)}(\mathbf{x}, t) \right|
$$

$$
\leq \delta + 2m^{3DD_1^2} (1 + m^{2\tau_{\text{min}}})\delta \leq 5m^{5DD_1^2} \delta, \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \geq \underline{T}$ because $D_1^2 \geq \tau_{\min}$. Combining [\(B.62\)](#page-79-0) with the last display, we have

$$
\begin{aligned}\n&\left| (\nabla \log p_t(\mathbf{x}))_i - \tilde{f}_i^{(1)}(\mathbf{x}, t) \right| \sigma_t \\
&\leq \left| \frac{\sigma_t(\nabla p_t(\mathbf{x}))_i}{p_t(\mathbf{x})} - f_{\text{mult}}^{(\text{in})} \left(f_i^{(1)}(\mathbf{x}, t), f_{\text{rec}}^{(\text{in})} \left(f_{D+1}^{(1)}(\mathbf{x}, t) \right) \right) \right| + 4m^{5DD_1^2} \delta \sigma_t\n\end{aligned} \tag{B.66}
$$
\n
$$
\leq D_2 m^{-\frac{\beta}{d}} (\log m)^{2(D-1+\beta \vee 1)} + 4m^{5DD_1^2} \delta, \quad i \in [D]
$$

for $||\mathbf{x}||_{\infty} \leq \mu_t - {\log(1/\sigma_t)}^{-3/2}$ and $m^{-\tau_{\min}} \leq t \leq 3m^{-\frac{1}{2D}}$.

B.5.2 Boundary of near-support

Let $\delta^{(2)} = m^{-(3DD_1^2 + 2\beta/d)}$ and

$$
\tau_{\rm bd}^{(2)} = 3/2, \quad \widetilde{\tau}_{\rm bd}^{(2)} = 1,
$$
\n
$$
\tau_{\rm x}^{(2)} = \left\{ \left(3DD_1^2 + \frac{2\beta}{d} \right)^{-\frac{1}{2}} \vee 1 \right\} \left[D_1 \vee \tau_{\rm min}^{\frac{3}{2}} \left\{ (4D)^{\frac{3}{2}} + 3 \right\} \right],
$$
\n
$$
\tau_{\rm t}^{(2)} = \left\{ 3D \left(3DD_1^2 + \frac{2\beta}{d} \right) \right\}^{-1} \wedge \frac{1}{2}.
$$

Let $\widetilde{C}_6 = \widetilde{C}_6(D, K, \overline{\tau}, \underline{\tau}, \tau_x^{(2)}), \ \widetilde{C}_7 = \widetilde{C}_7(D, K, \underline{\tau}), \ \widetilde{C}_8 = \widetilde{C}_8(D, \overline{\tau}, \tau_{\rm bd}^{(2)}, \tau_x^{(2)}, \tau_t^{(2)})$ $(\tau_t^{(2)}, \tilde{\tau}_{\text{bd}}^{(2)})$ be the constants in Proposition [B.2,](#page-40-1) where $(\tau_{bd}, \tilde{\tau}_{bd}, \tau_x, \tau_t)$ is replaced by $(\tau_{bd}^{(2)}, \tilde{\tau}_{bd}^{(2)}, \tau_x^{(2)}, \tau_t^{(2)})$ $t_t^{(2)}$). For large enough m, we have

$$
\delta^{(2)} \leq \widetilde{C}_8, \quad 3m^{-\frac{1}{2D}} \leq \left\{ \delta^{(2)} \right\}^{\tau_{\rm t}^{(2)}}, \quad \left\{ \log \left(1/\delta^{(2)} \right) \right\}^{-\widetilde{\tau}_{\rm bd}^{(2)}} \leq \tau_2.
$$

Also, a simple calculation yields that

$$
\delta^{(2)} \le m^{-\tau_{\min}}, \quad \tau_{\mathbf{x}}^{(2)} \ge \tau_{\min}^{\frac{3}{2}} \left\{ (4D)^{\frac{3}{2}} + 3 \right\}, \quad D_1 \sqrt{\log m} \le \tau_{\mathbf{x}}^{(2)} \sqrt{\log \left(1/\delta^{(2)} \right)}.
$$

Then, Proposition [B.2](#page-40-1) implies that that there exists a neural networks

$$
\mathbf{f}^{(2)} = (f_1^{(2)}, \dots, f_{D+1}^{(2)})^\top \in \mathcal{F}(L^{(2)}, \mathbf{d}^{(2)}, s^{(2)}, M^{(2)})
$$

with

$$
L \leq \widetilde{C}_6 \left\{ \log \left(1/\delta^{(2)} \right) \right\}^4, \quad \|\mathbf{d}\|_{\infty} \leq \widetilde{C}_6 \left\{ \log \left(1/\delta^{(2)} \right) \right\}^{7+2D},
$$

$$
s \leq \widetilde{C}_6 \left\{ \log \left(1/\delta^{(2)} \right) \right\}^{11+2D}, \quad M \leq \exp \left(\widetilde{C}_6 \left\{ \log \left(1/\delta^{(2)} \right) \right\}^2 \right),
$$

such that

$$
\left\| \begin{pmatrix} \sigma_t \nabla p_t(\mathbf{x}) \\ p_t(\mathbf{x}) \end{pmatrix} - \mathbf{f}^{(2)}(\mathbf{x}, t) \right\|_{\infty} \le \widetilde{C}_7 \delta^{(2)} \left\{ \log \left(1/\delta^{(2)} \right) \right\}^D \stackrel{\text{def}}{=} \epsilon_2
$$

for $\mu_t - \tau_{\min}^{3/2} \{ (4D)^{3/2} + 3 \} \{ \log(1/\sigma_t) \}^{-3/2} \le ||\mathbf{x}||_{\infty} \le \mu_t + \sigma_t D_1 \sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le 3m^{-\frac{1}{2D}}$. Lemma [A.1](#page-28-0) implies that $C_{S,1}^{-1}m^{-DD_1^2} \leq p_t(\mathbf{x}) \leq C_{S,1}$ for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ √ $\overline{\log m}$ and $t \geq 0$. Note that $C_{S,1}^{-1}m^{-DD_1^2}/2 \le p_t(\mathbf{x}) - \epsilon_2 \le f_{D+1}^{(2)}(\mathbf{x}, t) \le p_t(\mathbf{x}) + \epsilon_2 \le 2C_{S,1}$ for $\|\mathbf{x}\|_{\infty} \le \mu_t + \sigma_t D_1$ √ $\overline{\log m}$ and $m^{-\tau_{\min}} \leq t \leq 3m^{-\frac{1}{2D}}$ with large enough m so that $\epsilon_2 \leq C_{S,1}^{-1}m^{-DD_1^2}/2$. Let

$$
f_{\text{rec}}^{(\text{bd})} \in \mathcal{F}_{\text{NN}}(L_{\text{rec}}^{(\text{bd})}, \mathbf{d}_{\text{rec}}^{(\text{bd})}, s_{\text{rec}}^{(\text{bd})}, M_{\text{rec}}^{(\text{bd})})
$$

be the neural networks in Lemma [A.14](#page-37-1) with

$$
L_{\text{rec}}^{(\text{bd})} \leq C_{N,5} \{ DD_1^2 \log m + \log(2C_{S,1}) \}^2, \quad \|\mathbf{d}_{\text{rec}}^{(\text{bd})}\|_{\infty} \leq C_{N,5} \{ DD_1^2 \log m + \log(2C_{S,1}) \}^3,
$$

$$
s_{\text{rec}}^{(\text{bd})} \leq C_{N,5} \{ DD_1^2 \log m + \log(2C_{S,1}) \}^4, \quad M_{\text{rec}}^{(\text{bd})} \leq 4C_{N,5} C_{S,1} m^{2DD_1^2}
$$

such that $|1/x - f_{\text{rec}}^{(\text{bd})}(x)| \leq C_{S,1}^{-1}m^{-DD_1^2}/2$ for $x \in [C_{S,1}^{-1}m^{-DD_1^2}/2, 2C_{S,1}m^{DD_1^2}]$. Then,

$$
\left| \frac{1}{p_t(\mathbf{x})} - f_{\text{rec}}^{(\text{bd})} \left(f_{D+1}^{(2)}(\mathbf{x}, t) \right) \right| \le \left| \frac{1}{p_t(\mathbf{x})} - \frac{1}{f_{D+1}^{(2)}(\mathbf{x}, t)} \right| + \left| \frac{1}{f_{D+1}^{(2)}(\mathbf{x}, t)} - f_{\text{rec}}^{(\text{bd})} \left(f_{D+1}^{(2)}(\mathbf{x}, t) \right) \right|
$$

\n
$$
\le \{ p_t(\mathbf{x}) \wedge f_{D+1}^{(2)}(\mathbf{x}, t) \}^{-2} \epsilon_2 + C_{S,1}^{-1} m^{-DD_1^2} / 2 \le (4C_{S,1}^2 + C_{S,1}^{-1} / 2) \epsilon_2 m^{2DD_1^2}
$$

for $\mu_t - \tau_{\min}^{3/2} \{ (4D)^{3/2} + 3 \} \{ \log(1/\sigma_t) \}^{-3/2} \le ||\mathbf{x}||_{\infty} \le \mu_t + \sigma_t D_1 \sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le 3m^{-\frac{1}{2D}}$. Let $f_{\text{mult}}^{(\text{bd})} \in \mathcal{F}_{\text{NN}}(L_{\text{mult}}^{(\text{bd})}, \mathbf{d}_{\text{mult}}^{(\text{bd})}, s_{\text{mult}}^{(\text{bd})}, M_{\text{mult}}^{(\text{bd})})$

be the neural networks in Lemma [A.9](#page-32-0) with

$$
L_{\text{mult}}^{\text{(bd)}} \leq C_{N,1} \log 2\{(2\beta/d + 1 + 2DD_1^2) \log m + 2 \log(C_{S,1} \vee C_{S,3}) + 2 \log 2\},
$$

\n
$$
\mathbf{d}_{\text{mult}}^{\text{(bd)}} = (2, 96, \dots, 96, 1)^{\top},
$$

\n
$$
s_{\text{mult}}^{\text{(bd)}} \leq C_{N,1} 2\{(2\beta/d + 1 + DD_1^2) \log m + \log(C_{S,1} \vee C_{S,3}) + \log 2\},
$$

\n
$$
M_{\text{mult}}^{\text{(bd)}} = (C_{S,1} \vee C_{S,3})^2 m^{2DD_1^2}
$$

such that $|f_{\text{mult}}^{(\text{in})}(\tilde{\mathbf{x}}) - x_1x_2| \le m^{-2\beta/d-1} + 2(C_{S,1} \vee C_{S,3})m^{DD_1^2}\tilde{\epsilon}$ for all $\|\mathbf{x}\|_{\infty} \le (C_{S,1} \vee C_{S,3})m^{DD_1^2}$, $\widetilde{\mathbf{x}} \in \mathbb{R}^2$ with $\|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty} \leq \widetilde{\epsilon}$ and $|f_{\text{mult}}^{(\text{bd})}(\mathbf{x})| \leq (C_{S,1} \vee C_{S,3})^2 m^{2DD_1^2}$ for all $\mathbf{x} \in \mathbb{R}^2$. Then,

$$
\left| \frac{\sigma_t(\nabla p_t(\mathbf{x}))_i}{p_t(\mathbf{x})} - f_{\text{mult}}^{\text{(bd)}} \left(f_i^{(2)}(\mathbf{x}, t), f_{\text{rec}}^{\text{(bd)}} \left(f_{D+1}^{(2)}(\mathbf{x}, t) \right) \right) \right|
$$
\n
$$
\leq m^{-2\beta/d-1} + 2(C_{S,1} \vee C_{S,3}) (4C_{S,1}^2 + C_{S,1}^{-1}/2) \epsilon_2 m^{3DD_1^2} \leq D_3 m^{-2\beta/d} (\log m)^D, \quad i \in [D]
$$
\n(B.67)

for $\mu_t - \tau_{\min}^{3/2} \{ (4D)^{3/2} + 3 \} \{ \log(1/\sigma_t) \}^{-3/2} \le ||\mathbf{x}||_{\infty} \le \mu_t + \sigma_t D_1 \sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le 3m^{-\frac{1}{2D}},$ where $D_3 = D_3(\beta, d, D, \widetilde{C}_7, C_{S,1}, C_{S,3}, D_1)$. Consider functions $\widetilde{f}_1^{(2)}, \ldots, \widetilde{f}_D^{(2)} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}$ such that

$$
\widetilde{f}_i^{(2)}(\mathbf{x},t) = f_{\text{mult}}\left(f_{\text{mult}}^{(\text{bd})}\left(f_i^{(2)}(\mathbf{x},t), f_{\text{rec}}^{(\text{bd})}\left(f_{D+1}^{(2)}(\mathbf{x},t)\right)\right), f_{\text{rec}}(f_{\sigma}(t))\right), \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. For large enough m so that $m^{3DD_1^2} \geq (C_{S,1} \vee C_{S,3})^2 m^{2DD_1^2}$, [\(B.64\)](#page-79-1) and [\(B.65\)](#page-79-2) implies that

$$
\left| \sigma_t^{-1} f_{\text{mult}}^{(\text{bd})} \left(f_i^{(2)}(\mathbf{x}, t), f_{\text{rec}}^{(\text{bd})} \left(f_{D+1}^{(2)}(\mathbf{x}, t) \right) \right) - \tilde{f}_i^{(2)}(\mathbf{x}, t) \right|
$$

$$
\leq \delta + 2m^{3DD_1^2} (1 + m^{2\tau_{\text{min}}})\delta \leq 5m^{5DD_1^2} \delta, \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \geq m^{-\tau_{\min}}$. Combining [\(B.67\)](#page-81-0) with the last display, we have

$$
\begin{split}\n&\left| (\nabla \log p_t(\mathbf{x}))_i - \tilde{f}_i^{(2)}(\mathbf{x}, t) \right| \sigma_t \\
&\leq \left| \frac{\sigma_t(\nabla p_t(\mathbf{x}))_i}{p_t(\mathbf{x})} - f_{\text{mult}}^{(\text{bd})} \left(f_i^{(2)}(\mathbf{x}, t), f_{\text{rec}}^{(\text{bd})} \left(f_{D+1}^{(2)}(\mathbf{x}, t) \right) \right) \right| + 5m^{5DD_1^2} \delta \sigma_t \\
&\leq D_3 m^{-2\beta/d} (\log m)^D + 5m^{5DD_1^2} \delta \leq D_3 m^{-\beta/d} + 5m^{5DD_1^2} \delta, \quad i \in [D]\n\end{split} \tag{B.68}
$$

for $\mu_t - \tau_{\min}^{3/2} \{ (4D)^{3/2} + 3 \} \{ \log(1/\sigma_t) \}^{-3/2} \le ||\mathbf{x}||_{\infty} \le \mu_t + \sigma_t D_1 \sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le 3m^{-\frac{1}{2D}},$ where the last ienquality holds for large enough m so that $(\log m)^D \leq m^{\beta/d}$.

B.5.3 Large t

Let
$$
m^{(3)} = \sqrt{m}
$$
, $\tau_{\text{low}}^{(3)} = 1/2$,
\n
$$
\tau_{\text{sm}}^{(3)} = \left[\left\{ \left(\frac{2D}{\tau_{\text{low}}^{(3)}} \right) (1+D) \left(\frac{2\beta}{d} + \frac{1}{4D} + 3DD_1^2 \right) + D \left(\frac{D+1-\tau_{\text{low}}^{(3)}}{\tau_{\text{low}}^{(3)}} \right) \right\}
$$
\n
$$
\vee \left\{ D(1+D) \left(\frac{\{\tau_{\text{max}}\overline{\tau}\} \vee \{1/2D\}}{\tau_{\text{low}}^{(3)}} \right) - \frac{D+1-\tau_{\text{low}}}{\tau_{\text{low}}^{(3)}} \right\} \right] + 1
$$
\n
$$
\tau_{\text{x}}^{(3)} = D_1 \left\{ \frac{2D(1+D)}{\tau_{\text{low}}^{(3)}\tau_{\text{sm}}^{(3)} + D + 1 - \tau_{\text{low}}^{(3)}} \right\}^{\frac{1}{2}}.
$$

Also, let

$$
t_* = \left\{ m^{(3)} \right\}^{-\frac{2 - 2\tau_{\text{low}}^{(3)}}{D}}, \quad \delta^{(3)} = \left\{ m^{(3)} \right\}^{-\frac{\tau_{\text{low}}^{(3)} \tau_{\text{sm}}^{(3)} + D + 1 - \tau_{\text{low}}^{(3)}}{D(1+D)}},
$$

and $\widetilde{C}_9, \widetilde{C}_{10}, \widetilde{C}_{11}$ be the constants in Proposition [B.3](#page-40-0) depending on $(D, K, \overline{\tau}, \underline{\tau}, \tau_x^{(3)}, \tau_{\rm sm}^{(3)}, \tau_{\rm low}^{(3)})$, where $(\tau_x, \tau_{sm}, \tau_{low})$ is replaced by $(\tau_x^{(3)}, \tau_{sm}^{(3)}, \tau_{low}^{(3)})$. A simple calculation yields that

$$
t_{*} = m^{-\frac{1}{2D}}, \quad \left\{ m^{(3)} \right\}^{-\frac{\tau_{\text{low}}^{(3)} \tau_{\text{sm}}^{(3)} - (D + 1 - \tau_{\text{low}}^{(3)})D}{D(1 + D)}} \leq m^{-\frac{2\beta}{d} - \frac{1}{4D} - 3DD_{1}^{2}},
$$

$$
\tau_{\text{x}}^{(3)} \sqrt{\log(1/\delta^{(3)})} = D_{1} \sqrt{\log m}, \quad \delta^{(3)} \leq m^{-\frac{1}{2D}}, \quad \tau_{\text{max}} \log m \leq \overline{\tau}^{-1} \log(1/\delta^{(3)}).
$$

Also, for large enough m , we have

$$
m^{(3)} \geq \widetilde{C}_{11}
$$
 and $\left\{\log m^{(3)}\right\}^{\frac{D\tau_{\rm sm}^{(3)}}{2} + D} \leq m^{\frac{\beta}{d}}$.

Then, Proposition [B.3](#page-40-0) implies that that there exists a neural networks

$$
\mathbf{f}^{(3)} = (f_1^{(3)}, \dots, f_{D+1}^{(3)})^\top \in \mathcal{F}(L^{(3)}, \mathbf{d}^{(3)}, s^{(3)}, M^{(3)})
$$

with

$$
L^{(3)} \leq \widetilde{C}_9(\log m/2)^4, \quad \|\mathbf{d}^{(3)}\|_{\infty} \leq \widetilde{C}_9\sqrt{m}(\log m/2)^9,
$$

$$
s^{(3)} \leq \widetilde{C}_9\sqrt{m}(\log m/2)^{11}, \quad M^{(3)} \leq \exp(\widetilde{C}_9{\log m/2})^2),
$$

such that

$$
\left\| \begin{pmatrix} \sigma_{t-t*} \nabla p_t(\mathbf{x}) \\ p_t(\mathbf{x}) \end{pmatrix} - \mathbf{f}^{(3)}(\mathbf{x}, t - t_*) \right\|_{\infty} \le \widetilde{C}_{10} m^{-\frac{\beta}{d} - \frac{1}{4D} - 3DD_1^2} \stackrel{\text{def}}{=} \epsilon_3
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $2m^{-\frac{1}{2D}} \leq t \leq \tau_{\max} \log m$, where $t_* = m^{-1/(2D)}$. Note that $C_{S,1}^{-1}m^{-DD_1^2}/2 \leq p_t(\mathbf{x}) - \epsilon_3 \leq f_{D+1}^{(3)}(\mathbf{x},t) \leq p_t(\mathbf{x}) + \epsilon_3 \leq 2C_{S,1}$ for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ √ $\overline{\log m}$ and $2m^{-\frac{1}{2D}} \le t \le \tau_{\max} \log m$ with large enough m so that $\epsilon_3 \le C_{S,1}^{-1} m^{-DD_1^2}/2$. Then,

$$
\left| \frac{1}{p_t(\mathbf{x})} - f_{\text{rec}}^{(\text{bd})} \left(f_{D+1}^{(3)}(\mathbf{x}, t - t_*) \right) \right|
$$
\n
$$
\leq \left| \frac{1}{p_t(\mathbf{x})} - \frac{1}{f_{D+1}^{(3)}(\mathbf{x}, t - t_*)} \right| + \left| \frac{1}{f_{D+1}^{(3)}(\mathbf{x}, t - t_*)} - f_{\text{rec}}^{(\text{bd})} \left(f_{D+1}^{(3)}(\mathbf{x}, t - t_*) \right) \right|
$$
\n
$$
\leq \{ p_t(\mathbf{x}) \wedge f_{D+1}^{(3)}(\mathbf{x}, t - t_*) \}^{-2} \epsilon_3 + C_{S,1}^{-1} m^{-DD_1^2} / 2 \leq (4C_{S,1}^2 + C_{S,1}^{-1}/2) \epsilon_3 m^{2DD_1^2}
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $2m^{-\frac{1}{2D}} \le t \le \tau_{\max} \log m$. Then,

$$
\left| \frac{\sigma_{t-t_*}(\nabla p_t(\mathbf{x}))_i}{p_t(\mathbf{x})} - f_{\text{mult}}^{\text{(bd)}} \left(f_i^{(3)}(\mathbf{x}, t - t_*) , f_{\text{rec}}^{\text{(bd)}} \left(f_{D+1}^{(3)}(\mathbf{x}, t - t_*) \right) \right) \right|
$$
\n
$$
\leq m^{-2\beta/d-1} + 2(C_{S,1} \vee C_{S,3}) (4C_{S,1}^2 + C_{S,1}^{-1}/2) \epsilon_3 m^{3DD_1^2} \leq D_4 m^{-\beta/d-1/(4D)} \tag{B.69}
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $2m^{-\frac{1}{2D}} \le t \le \tau_{\max} \log m$. where $D_4 = D_4(\widetilde{C}_{10}, C_{S,1}, C_{S,3})$. Consider functions $\tilde{f}_1^{(3)}, \ldots, \tilde{f}_D^{(3)} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}$ such that

$$
\widetilde{f}_i^{(3)}(\mathbf{x},t) = f_{\text{mult}}\left(f_{\text{mult}}^{(\text{bd})}\left(f_i^{(3)}(\mathbf{x},t-t_*), f_{\text{rec}}^{(\text{in})}\left(f_{D+1}^{(3)}(\mathbf{x},t-t_*)\right)\right), f_{\text{rec}}(f_{\sigma}(t-t_*))\right), \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Combining with $(B.64)$ and $(B.65)$, we have

$$
\left| \sigma_{t-t_*}^{-1} f_{\text{mult}}^{(\text{bd})} \left(f_i^{(3)}(\mathbf{x}, t - t_*) , f_{\text{rec}}^{(\text{bd})} \left(f_{D+1}^{(3)}(\mathbf{x}, t - t_*) , \right) \right) - \tilde{f}_i^{(3)}(\mathbf{x}, t) \right|
$$

$$
\leq \delta + 2m^{3DD_1^2} (1 + m^{2\tau_{\text{min}}}) \delta \leq 5m^{5DD_1^2} \delta, \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \ge 2m^{-1/(2D)}$. Note that $|\sigma_t/\sigma_{t-t_*}| \le |\sigma_{t-t_*}^{-1}| \le (2\tau)^{-1/2}m^{1/(4D)}$ for $t \ge 2m^{-1/(2D)}$. Combining $(B.69)$ with the last display, we have

$$
\begin{aligned} &\left| (\nabla \log p_t(\mathbf{x}))_i - \tilde{f}_i^{(3)}(\mathbf{x}, t) \right| \sigma_{t-t_*} \\ &\leq \left| \frac{\sigma_{t-t_*}(\nabla p_t(\mathbf{x}))_i}{p_t(\mathbf{x})} - f_{\text{mult}}^{(\text{bd})} \left(f_i^{(3)}(\mathbf{x}, t - t_*), f_{\text{rec}}^{(\text{bd})} \left(f_{D+1}^{(3)}(\mathbf{x}, t - t_*) \right) \right) \right| + 5m^{5DD_1^2} \delta \sigma_{t-t_*} \\ &\leq D_3 m^{-\beta/d - 1/(4D)} + 5m^{5DD_1^2} \delta, \quad i \in [D] \end{aligned}
$$

and

$$
\begin{aligned} &\left| (\nabla \log p_t(\mathbf{x}))_i - \tilde{f}_i^{(3)}(\mathbf{x}, t) \right| \sigma_t \\ &\leq \left| (\nabla \log p_t(\mathbf{x}))_i - \tilde{f}_i^{(3)}(\mathbf{x}, t) \right| \sigma_{t-t_*}(2\underline{\tau})^{-1/2} m^{1/(4D)} \\ &\leq D_3(2\underline{\tau})^{-1/2} \sqrt{\underline{\tau}} m^{-\beta/d} + 5(2\underline{\tau})^{-1/2} \sqrt{\underline{\tau}} m^{5DD_1^2 + 1/(4D)} \delta, \quad i \in [D] \end{aligned}
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $2m^{-\frac{1}{2D}} \le t \le \tau_{\max} \log m$. Let $\delta = m^{-5D}D_1^2 - 1/(4D) - \beta/d$. Then, there exists a positive constant $D_5 = D_5(\underline{\tau}, D_2, D_3, D_4)$ such that [\(B.66\)](#page-80-0), [\(B.68\)](#page-81-1) and the last display are bounded by

$$
D_5 m^{-\frac{\beta}{d}} (\log m)^{2(D-1+\beta \wedge 1)} \stackrel{\text{def}}{=} \epsilon_4 \tag{B.70}
$$

B.5.4 Combining into a single function

For large enough m so that $3m^{-1/(2D)} \leq (2\overline{\tau})^{-1}$, we have

$$
\left\{\log(1/\sqrt{\tau}) + \tau_{\min}\log m/2\right\}^{-3/2} \le \left\{\log(1/\sigma_t)\right\}^{-3/2} \le \left\{\log(1/\sqrt{6\tau}) + \log m/(4D)\right\}^{-3/2}
$$

for $m^{-\tau_{\min}} \leq t \leq 3m^{-1/(2D)}$ because $\sqrt{\tau t} \leq \sigma_t \leq$ √ $\overline{2\overline{\tau}t}$ for $0 \le t \le (2\overline{\tau})^{-1}$. For large enough m so that $\log(1/\sqrt{\tau}) \leq \tau_{\min} \log m/2$, it follows that

$$
\tau_{\min}^{-3/2} (\log m)^{-3/2} \le {\log(1/\sigma_t)}^{-3/2} \le (4D)^{3/2} (\log m)^{-3/2}
$$

for $m^{-\tau_{\min}} \leq t \leq 3m^{-1/(2D)}$. Then,

$$
\mu_t - \tau_{\min}^{3/2} \{ (4D)^{3/2} + 3 \} \{ \log(1/\sigma_t) \}^{-3/2} < \mu_t - \overline{x} < \mu_t - \underline{x} < \mu_t - \{ \log(1/\sigma_t) \}^{-3/2} \tag{B.71}
$$

for $m^{-\tau_{\min}} \leq t \leq 3m^{-1/(2D)}$, where

$$
\overline{x} = \{(4D)^{3/2} + 2\} (\log m)^{-3/2}
$$
 and $\underline{x} = \{(4D)^{3/2} + 1\} (\log m)^{-3/2}$.

Consider piecewise linear functions $f_{\text{swi}}^{(1)}$ $f_{\text{swit},x}^{(1)}, f_{\text{swit},x}^{(2)} : \mathbb{R} \to [0,1]$ such that

$$
f_{\text{swit,x}}^{(1)}(x) = (\log m)^{3/2} \rho \left(-f_{\text{clip}}^{(x,\overline{x})}(x) + \overline{x} \right) = \frac{1}{\overline{x} - \underline{x}} \max \left(-(x \vee \underline{x}) \wedge \overline{x} + \overline{x}, 0 \right),
$$

$$
f_{\text{swit,x}}^{(2)}(x) = (\log m)^{3/2} \rho \left(f_{\text{clip}}^{(x,\overline{x})}(x) - \underline{x} \right) = \frac{1}{\overline{x} - \underline{x}} \max \left((x \vee \underline{x}) \wedge \overline{x} - \underline{x}, 0 \right),
$$

for $x \in \mathbb{R}$, where $f_{\text{clip}}^{(\underline{x}, \overline{x})} \in \mathcal{F}_{\text{NN}}(2, (1, 2, 1)^{\top}, 7, \overline{x} \vee (\log m)^{3/2})$ is the neural network in Lemma [A.10.](#page-32-1) Note that $f_{\text{swi}}^{(1)}$ $f_{\text{swit},x}^{(1)}(x) + f_{\text{swi}}^{(2)}$ $f_{\text{swit},x}^{(2)}(x) = 1$ for $x \in \mathbb{R}$, and $f_{\text{swit}}^{(1)}$ $s_{\text{wit},x}^{(1)}(x) = 0$ for $x \geq \overline{x}$ and $f_{\text{swi}}^{(2)}$ $s_{\text{wit},x}(x) = 0$ for $x \leq \underline{x}$. Combining with [\(B.70\)](#page-83-1) and [\(B.71\)](#page-83-2), we have

$$
\begin{aligned} &\left| f_{\text{switch},x}^{(1)} \left(\mu_t - f_{\text{max}}(\mathbf{x}) \right) \widetilde{f}_i^{(1)}(\mathbf{x}, t) + f_{\text{switch},x}^{(2)} \left(\mu_t - f_{\text{max}}(\mathbf{x}) \right) \widetilde{f}_i^{(2)}(\mathbf{x}, t) - (\nabla \log p_t(\mathbf{x}))_i \right| \\ &\leq \epsilon_4 / \sigma_t, \quad i \in [D] \end{aligned}
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le 3m^{-\frac{1}{2D}}$, where $f_{\max}: \mathbb{R}^D \to \mathbb{R}$ is a function such that

$$
f_{\max}(\mathbf{x}) = \|\mathbf{x}\|_{\infty} = \rho(\cdots \rho(\rho(\rho(x_1 - x_2) + x_2 - x_3) + x_3 - x_4)\cdots + x_{D-1} - x_D) + x_D.
$$

Lemma [A.13](#page-37-0) implies that there exist neural networks $f_{\mu} \in \mathcal{F}_{NN}(L_{\mu}, \mathbf{d}_{\mu}, s_{\mu}, M_{\mu})$ with

$$
L_{\mu} \leq C_{N,4} \{ \log(1/\delta) \}^2, \quad, \|\mathbf{d}_{\mu}\|_{\infty} \leq C_{N,4} \{ \log(1/\delta) \}^2
$$

$$
s_{\mu} \leq C_{N,4} \{ \log(1/\delta) \}^3, \quad M_{\mu} \leq C_{N,4} \log(1/\delta)
$$

such that $|\mu_t - f_{\mu}(t)| \leq \delta$ for $t \geq 0$. Since $f_{\text{swi}}^{(1)}$ $f_{\text{swit},x}^{(1)}$ and $f_{\text{swi}}^{(2)}$ $\sum_{\text{swit},x}^{(2)}$ are $(\log m)^{\tau_{\text{bd}}}\text{-Lipschitz continuous},$ $|f_{\text{swi}}^{(i)}\rangle$ $s_{\text{switch},\textbf{x}}^{(i)}(\mu_t - \|\textbf{x}\|_\infty) - f_{\text{swi}}^{(i)}$ $\mathcal{S}_{\text{swit},\mathbf{x}}^{(i)}(f_{\mu}(t)-\|\mathbf{x}\|_{\infty})|\leq \delta(\log m)^{\tau_{\text{bd}}}$ for each $i\in\{1,2\},\mathbf{x}\in\mathbb{R}^D$ and $t\geq 0$. For $i \in [D]$, consider a function $f_i^{(x)}$ $f_i^{(x)} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}$ such that

$$
f_i^{(\mathbf{x})}(\mathbf{x},t)
$$

= $f_{\text{mult}}(f_{\text{swit},\mathbf{x}}^{(1)}(f_{\mu}(t) - f_{\text{max}}(\mathbf{x})), \tilde{f}_i^{(1)}(\mathbf{x},t)) + f_{\text{mult}}(f_{\text{swit},\mathbf{x}}^{(2)}(f_{\mu}(t) - f_{\text{max}}(\mathbf{x})), \tilde{f}_i^{(2)}(\mathbf{x},t))$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Combining with [\(B.65\)](#page-79-2), we have

$$
\begin{aligned} \left| f_i^{(\mathbf{x})}(\mathbf{x},t) - (\nabla \log p_t(\mathbf{x}))_i \right| \\ &\leq \epsilon_4/\sigma_t + 2\delta + 4m^{3DD_1^2}\delta(\log m)^{\tau_{\rm bd}} \leq \epsilon_4/\sigma_t + 6m^{3DD_1^2}\delta(\log m)^{\tau_{\rm bd}}, \quad i \in [D] \end{aligned} \tag{B.72}
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le 3m^{-\frac{1}{2D}}$. Similarly, let $\underline{t} = 2m^{-\frac{1}{2D}}$ and $\overline{t} = 3m^{-\frac{1}{2D}}$, consider piecewise linear functions $f_{\text{swi}}^{(1)}$ $f_{\text{switch}}^{(1)}, f_{\text{switch}}^{(2)} : \mathbb{R} \to [0, 1]$ such that

$$
f_{\text{switch},t}^{(1)}(t) = m^{\frac{1}{2D}} \rho \left(-f_{\text{clip}}^{(t,\bar{t})}(t) + \bar{t} \right) = \frac{1}{\bar{t} - \underline{t}} \max \left(-(t \vee \underline{t}) \wedge \bar{t} + \bar{t}, 0 \right),
$$

$$
f_{\text{switch},t}^{(2)}(t) = m^{\frac{1}{2D}} \rho \left(f_{\text{clip}}^{(t,\bar{t})}(t) - \underline{t} \right) = \frac{1}{\bar{t} - \underline{t}} \max \left((t \vee \underline{t}) \wedge \bar{t} - \underline{t}, 0 \right),
$$

where $f_{\text{clip}}^{(\underline{t},\overline{t})} \in \mathcal{F}_{\text{NN}}(2,(1,2,1)^{\top},7,\overline{t} \vee m^{\frac{1}{2D}})$ is the neural network in Lemma [A.10.](#page-32-1) Note that $f_{\text{swi}}^{(1)}$ $s_{\text{wit,t}}^{(1)}(t) +$ $f_{\text{swi}}^{(2)}$ $s_{\text{wit,t}}^{(2)}(t) = 1$ for $t \in \mathbb{R}$, and $f_{\text{swit}}^{(1)}$ $f_{\text{switch}}^{(1)}(t) = 0$ for $t \geq \overline{t}$ and $f_{\text{swi}}^{(2)}$ $s_{\text{switch}}^{(2)}(t) = 0$ for $t \leq \underline{t}$. Combining with $(B.70)$ and $(B.72)$, we have

$$
\left| f_{\text{switch}}^{(1)}(t) f_i^{(\mathbf{x})}(\mathbf{x}, t) + f_{\text{switch}}^{(2)}(t) \tilde{f}_i^{(3)}(\mathbf{x}, t) - (\nabla \log p_t(\mathbf{x}))_i \right|
$$

$$
\leq \epsilon_4 / \sigma_t + 6m^{3DD_1^2} \delta(\log m)^{\tau_{\text{bd}}}, \quad i \in [D]
$$

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le \tau_{\max} \log m$. Consider real-valued functions $f_1^{(x,t)}$ $f_{1}^{(x,t)},...,f_{D}^{(x,t)}$ such that

$$
f_i^{(\mathbf{x},t)}(\mathbf{x},t) = f_{\text{mult}}\left(f_{\text{switch},t}^{(1)}(t), f_i^{(\mathbf{x})}(\mathbf{x},t)\right) + f_{\text{mult}}\left(f_{\text{switch},t}^{(2)}(t), \tilde{f}_i^{(3)}(\mathbf{x},t)\right), \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Combining with [\(B.65\)](#page-79-2), we have

$$
\left| f_i^{(x,t)}(\mathbf{x},t) - (\nabla \log p_t(\mathbf{x}))_i \right| \le \epsilon_4/\sigma_t + 6m^{3DD_1^2} \delta(\log m)^{\tau_{\text{bd}}} + 4\delta
$$
\n
$$
\le \epsilon_4/\sigma_t + 10m^{3DD_1^2} \delta(\log m)^{\tau_{\text{bd}}} \le 11\epsilon_4/\sigma_t, \quad i \in [D]
$$
\n(B.73)

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le \tau_{\max} \log m$, where the last inequality holds with large enough m so that $m^{3DD_1^2}\delta(\log m)^{\tau_{\text{bd}}} \leq \epsilon_4$. Since $\|\mathbf{x}\|_{\infty} \leq \|\mathbf{x}\|_2$ for any $\mathbf{x} \in \mathbb{R}^D$, Lemma [A.2](#page-29-3) implies that $\sigma_t \|\nabla \log p_t(\mathbf{x})\|_{\infty} \leq C_{S,2}D_1$ √ $\overline{\log m}$ for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ √ $\overline{\log m}$ and $t \ge 0$ with large enough m so that D_1 √ $\overline{\log m} \ge 1$, where $C_{S,2} = C_{S,2}(D, K, \tau_1, \overline{\tau}, \underline{\tau})$ is the constant in Lemma [A.2.](#page-29-3) Combining with the last display, we have

$$
\sigma_t \left| f_i^{(\mathbf{x},t)}(\mathbf{x},t) \right| \le 11\epsilon_4 + C_{S,2}D_1\sqrt{\log m} \le D_5\sqrt{\log m}, \quad i \in [D]
$$

for $||\mathbf{x}||_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le \tau_{\max} \log m$ with large enough m so that $\epsilon_4 \le \sqrt{\frac{m}{\epsilon}}$ $\overline{\log m},$ where $D_5 = D_5(\beta, d, D, C_{S,2})$. Consider a function $\mathbf{f} = (f_1, \ldots, f_D)^\top : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}^D$ such that

$$
f_i(\mathbf{x}, t) = f_{\text{clip}}^{(-D_5 \sqrt{\log m}, D_5 \sqrt{\log m})} \left(f_i^{(\mathbf{x}, t)}(\mathbf{x}, t) \right),
$$

=
$$
\left(f_i^{(\mathbf{x}, t)}(\mathbf{x}, t) \vee -D_5 \sqrt{\log m} \right) \wedge D_5 \sqrt{\log m}, \quad i \in [D]
$$

for $\mathbf{x} \in \mathbb{R}^D$ and $t \in \mathbb{R}$, where $f_{\text{clip}}^{(-D_5\sqrt{\log m}, D_5\sqrt{\log m})} \in \mathcal{F}_{\text{NN}}(2, (1, 2, 1)^T, 7, D_5\sqrt{\log m})$ $\overline{\log m}$ is the neural network in Lemma [A.10.](#page-32-1) Since $0 < \sigma_t \leq 1$ for $t > 0$, we have $f_i(\mathbf{x}, t) = f_i^{(\mathbf{x}, t)}$ $i^{(X,t)}(x, t)$ for $i \in [D],$ $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le \tau_{\max} \log m$. Combining with [\(B.73\)](#page-85-0),

$$
\sigma_t \| \mathbf{f}(\mathbf{x}, t) - \nabla \log p_t(\mathbf{x}) \|_{\infty} \le 11 D_5 m^{-\frac{\beta}{d}} (\log m)^{(\tau_{\text{bd}} + \frac{1}{2})(D - 1 + \beta \wedge 1)}
$$
(B.74)

for $\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1$ $\sqrt{\log m}$ and $m^{-\tau_{\min}} \le t \le \tau_{\max} \log m$.

B.5.5 Outside of near-support

Note that

$$
\int_{\underline{T}}^{\overline{T}} \int_{\mathbb{R}^D} \|\nabla \log p_t(\mathbf{x}) - \mathbf{f}(\mathbf{x}, t)\|_2^2 p_t(\mathbf{x}) d\mathbf{x} \n\leq \int_{\underline{T}}^{\overline{T}} \int_{\|\mathbf{x}\|_{\infty} \leq \mu_t + \sigma_t D_1 \sqrt{\log m}} \|\nabla \log p_t(\mathbf{x}) - \mathbf{f}(\mathbf{x}, t)\|_2^2 p_t(\mathbf{x}) d\mathbf{x} \n+ \int_{\underline{T}}^{\overline{T}} \int_{\|\mathbf{x}\|_{\infty} \geq \mu_t + \sigma_t D_1 \sqrt{\log m}} \|\nabla \log p_t(\mathbf{x}) - \mathbf{f}(\mathbf{x}, t)\|_2^2 p_t(\mathbf{x}) d\mathbf{x}
$$

For $t \geq 0$,

$$
\int_{\|\mathbf{x}\|_{\infty}\geq\mu_{t}+D_{1}\sqrt{\log m}}p_{t}(\mathbf{x})\mathrm{d}\mathbf{x} = \int_{\|\mathbf{x}\|_{\infty}\geq\mu_{t}+\sigma_{t}D_{1}\sqrt{\log m}}\int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y})\mathrm{d}\mathbf{y}\mathrm{d}\mathbf{x}
$$
\n
$$
= \int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\int_{\|\sigma_{t}\mathbf{z}+\mu_{t}\mathbf{y}\|_{\infty}\geq\mu_{t}+\sigma_{t}D_{1}\sqrt{\log m}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\mathrm{d}\mathbf{y}
$$
\n
$$
\leq \int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\sum_{i=1}^{D}\int_{|\sigma_{t}z_{i}+\mu_{t}y_{i}|\geq\mu_{t}+\sigma_{t}D_{1}\sqrt{\log m}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\mathrm{d}\mathbf{y}
$$
\n
$$
\leq \int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\sum_{i=1}^{D}\int_{|z_{i}|\geq D_{1}\sqrt{\log m}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\mathrm{d}\mathbf{y} = D\int_{|z|\geq D_{1}\sqrt{\log m}}\phi(z)\mathrm{d}z,
$$

where the last inequality holds because $|y_i| \leq 1$. By the tail probability of the standard normal distribution, the last display is bounded by

$$
2Dm^{-D_1^2/2}.\t\t(B.75)
$$

Let $C_{S,2} = C_{S,2}(D, K, \tau_1, \overline{\tau}, \underline{\tau})$ be the constant in Lemma [A.2.](#page-29-3) Then, $\|\nabla \log p_t(\mathbf{x})\|_2 \leq C_{S,2}(\|\mathbf{x}\|_{\infty} \mu_t$)/ σ_t^2 for $\|\mathbf{x}\|_{\infty} \geq \mu_t + D_1$ √ $\overline{\log m}$ with large enough m so that D_1 √ $\overline{\log m} \geq 1$. Combining with the last display, we have

$$
\int_{\|\mathbf{x}\|_{\infty}\geq\mu_t+D_1\sqrt{\log m}} \|\nabla \log p_t(\mathbf{x})\|_2^2 p_t(\mathbf{x}) d\mathbf{x}
$$
\n
$$
\leq 2C_{S,2}^2 \sigma_t^{-4} \int_{\|\mathbf{x}\|_{\infty}\geq\mu_t+D_1\sqrt{\log m}} \left(\|\mathbf{x}\|_{\infty}^2 + \mu_t^2\right) p_t(\mathbf{x}) d\mathbf{x}
$$
\n
$$
\leq 2C_{S,2}^2 \sigma_t^{-4} \int_{\|\mathbf{x}\|_{\infty}\geq\mu_t+D_1\sqrt{\log m}} \|\mathbf{x}\|_2^2 p_t(\mathbf{x}) d\mathbf{x} + 2DC_{S,2}^2 \sigma_t^{-4} \mu_t^2 m^{-D_1^2/2}
$$

for $t \geq 0$. A simple calculation yields that

$$
\int_{\|\mathbf{x}\|_{\infty}\geq\mu_{t}+D_{1}\sqrt{\log m}}\|\mathbf{x}\|_{2}^{2}p_{t}(\mathbf{x})\mathrm{d}\mathbf{x}\n=\sum_{i=1}^{D}\int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\int_{\|\mathbf{x}\|_{\infty}\geq\mu_{t}+\sigma_{t}D_{1}\sqrt{\log m}}x_{i}^{2}\phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y})\mathrm{d}\mathbf{x}\mathrm{d}\mathbf{y}\n=\sum_{i=1}^{D}\int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\int_{\|\sigma_{t}\mathbf{z}+\mu_{t}\mathbf{y}\|_{\infty}\geq\mu_{t}+\sigma_{t}D_{1}\sqrt{\log m}}(\sigma_{t}z_{i}+\mu_{t}y_{i})^{2}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\mathrm{d}\mathbf{y}\n\leq \sum_{i=1}^{D}\int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\int_{\|\sigma_{t}\mathbf{z}+\mu_{t}\mathbf{y}\|_{\infty}\geq\mu_{t}+\sigma_{t}D_{1}\sqrt{\log m}}2(\sigma_{t}^{2}z_{i}^{2}+\mu_{t}^{2}y_{i}^{2})\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\mathrm{d}\mathbf{y}\n\leq \sum_{i=1}^{D}\int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\sum_{j=1}^{D}\int_{|\sigma_{t}z_{j}+\mu_{t}y_{j}|\geq\mu_{t}+\sigma_{t}D_{1}\sqrt{\log m}}2(\sigma_{t}^{2}z_{i}^{2}+\mu_{t}^{2}y_{i}^{2})\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\mathrm{d}\mathbf{y}
$$

for $t \geq 0$. Since $|y_j| \leq 1$ in the last integral, the last display is bounded by

$$
\sum_{i=1}^{D} \int_{\|\mathbf{y}\|_{\infty} \leq 1} p_0(\mathbf{y}) \sum_{j=1}^{D} \int_{|z_j| \geq D_1 \sqrt{\log m}} 2(\sigma_t^2 z_i^2 + \mu_t^2 y_i^2) \phi_1(\mathbf{z}) \, d\mathbf{z} \, d\mathbf{y}
$$

$$
\leq 2\sigma_t^2 \sum_{i=1}^{D} \sum_{j=1}^{D} \int_{|z_j| \geq D_1 \sqrt{\log m}} z_i^2 \phi_1(\mathbf{z}) \, d\mathbf{z} + 4D^2 \mu_t^2 m^{-D_1^2/2},
$$

where the last inequality holds by $(B.75)$. Furthermore,

 \overline{a}

$$
\sum_{i=1}^{D} \sum_{j=1}^{D} \int_{|z_j| \ge D_1 \sqrt{\log m}} z_i^2 \phi_1(\mathbf{z}) d\mathbf{z}
$$
\n
$$
= \sum_{i=1}^{D} \left\{ (D-1) \mathbb{E}[Z^2] \int_{|z| \ge D_1 \sqrt{\log m}} \phi(z) dz + \int_{|z| \ge D_1 \sqrt{\log m}} z^2 \phi(z) dz \right\}
$$
\n
$$
\le 2D \left\{ (D-1)m^{-D_1^2/2} + \sqrt{\mathbb{E}[Z^4]} m^{-D_1^2/4} \right\} \le 2D(D + \sqrt{3} - 1)m^{-D_1^2/4},
$$

where the first inequality holds by the Cauchy-Schwarz inequality. Hence, there exists a constant $D_6 = D_6(D, \underline{\tau}, C_{S,2})$ such that

$$
\int_{\|\mathbf{x}\|_{\infty}\geq\mu_t+D_1\sqrt{\log m}} \|\nabla \log p_t(\mathbf{x})\|_2^2 p_t(\mathbf{x}) d\mathbf{x} \leq D_6 \sigma_t^{-2} m^{-2\beta/d}
$$

for $t \geq m^{-\tau_{\min}}$ because $\mu_t \leq 1$ and $\sigma_t \geq \sqrt{\tau m^{-\tau_{\min}}}$. Combining with [\(B.75\)](#page-86-0), we have

$$
\int_{\|\mathbf{x}\|_{\infty}\geq\mu_t+D_1\sqrt{\log m}} \|\nabla \log p_t(\mathbf{x}) - \mathbf{f}(\mathbf{x},t)\|_2^2 p_t(\mathbf{x}) d\mathbf{x} \n\leq \int_{\|\mathbf{x}\|_{\infty}\geq\mu_t+D_1\sqrt{\log m}} 2\|\nabla \log p_t(\mathbf{x})\|_2^2 p_t(\mathbf{x}) d\mathbf{x} + \int_{\|\mathbf{x}\|_{\infty}\geq\mu_t+D_1\sqrt{\log m}} 2\|\mathbf{f}(\mathbf{x},t)\|_2^2 p_t(\mathbf{x}) d\mathbf{x} \n\leq 2D_6\sigma_t^{-2}m^{-2\beta/d} + 2D^2D_5m^{-D_1^2/2}\log m \n\leq \sigma_t^{-2} \left(2D_6m^{-2\beta/d} + 2D^2D_5m^{-2\beta/d}\log m\right)
$$

for $t \geq m^{-\tau_{\min}}$, where the second inequality holds because $\|\mathbf{f}(\mathbf{x},t)\|_{\infty} \leq D_5\sqrt{\log m}$ for $\mathbf{x} \in \mathbb{R}^D, t \in \mathbb{R}$. Combining with [\(B.74\)](#page-85-1), we have

$$
\sigma_t^2 \int_{\mathbb{R}^D} \|\nabla \log p_t(\mathbf{x}) - \mathbf{f}(\mathbf{x}, t)\|_2^2 p_t(\mathbf{x}) d\mathbf{x}
$$

\n
$$
\leq D11^2 D_5^2 m^{-\frac{2\beta}{d}} (\log m)^{(2\tau_{\text{bd}}+1)(D-1+\beta\wedge 1)} + 2D_6 m^{-\frac{2\beta}{d}} + 2D^2 D_5 m^{-D_1^2/2} \log m
$$

\n
$$
\leq D_7 m^{-\frac{2\beta}{d}} (\log m)^{(2\tau_{\text{bd}}+1)(D-1+\beta\wedge 1)}
$$

for $m^{-\tau_{\min}} \le t \le \tau_{\max} \log m$, where $D_7 = D_7(D, D_5, D_6)$. Since $\sigma_t^2 \ge \tau t$ for $t \ge 0$, we have

$$
\int_{\underline{T}}^{\overline{T}} \int_{\mathbb{R}^D} \|\nabla \log p_t(\mathbf{x}) - \mathbf{f}(\mathbf{x}, t)\|_2^2 p_t(\mathbf{x}) \, \mathrm{d}\mathbf{x} \, \mathrm{d}t
$$
\n
$$
\leq D_7 \underline{\tau}^{-1} m^{-\frac{2\beta}{d}} (\log m)^{(2\tau_{\text{bd}}+1)(D-1+\beta \wedge 1)} (\log \overline{T} - \log \underline{T})
$$
\n
$$
\leq D_8 m^{-\frac{2\beta}{d}} (\log m)^{(2\tau_{\text{bd}}+1)(D-1+\beta \wedge 1)+1},
$$

where $D_8 = D_8(\tau, \tau_{\text{max}}, \tau_{\text{min}}, D_7)$. Lemma [A.4,](#page-29-0) Lemma [A.5,](#page-29-1) Lemma [A.6,](#page-30-0) Lemma [A.7](#page-30-1) and Lemma [A.8](#page-30-2) implies that $f \in \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, m, M, \mathcal{P})$ with

$$
L \le D_9(\log m)^6 \log \log m, \quad \|\mathbf{d}\|_{\infty} \le D_9 m (\log m)^3,
$$

$$
s \le D_9 m^{D+1} (\log m)^5 \log \log m, \quad M \le \exp(D_9 {\log m})^6),
$$

 $\|\mathbf{m}\|_{\infty} \leq D_9 m^D$ and the set of permutation matrices \mathcal{P} , where

$$
D_9 = D_6(\beta, d, D, \tau_{\min}, \widetilde{C}_3, \widetilde{C}_6, \widetilde{C}_9, C_{N,1}, C_{N,4}, C_{N,5}, C_{S,1}, D_1)
$$

is a large enough constant. The assertion follows by re-defining the constants.

C Proofs for the convergence rate

In this section, we provide the proof of Theorem [5.2.](#page-15-0) We begin by outlining the cruical lemmas and a proposition.

Lemma C.1 (Error bound for small t). Let $\beta, K > 0$ be given and suppose the true density p_0 belongs to $\mathcal{H}^{\beta,K}([-1,1]^D)$. Then, there exist positive constants $\widetilde{C}_{12} = \widetilde{C}_{12}(\beta, D, K, \overline{\tau}, \underline{\tau})$ and $\widetilde{C}_{13} = \widetilde{C}_{13}(\overline{\tau}, \underline{\tau})$ such that

$$
\int_{\mathbb{R}^D} |p_0(\mathbf{x}) - p_t(\mathbf{x})| \, \mathrm{d}\mathbf{x} \leq \widetilde{C}_{12} \left\{ t \log(1/t) \right\}^{\frac{\beta \wedge 1}{2}}
$$

for $0 \leq t \leq \widetilde{C}_{13}$.

 \Box

For any function $f: \mathbb{R}^{n_1} \to \mathbb{R}^{n_2}, n_1, n_2 \in \mathbb{N}$ and $C > 0$, denote $\|\cdot\|_{L^{\infty}([-C,C]^{n_1})}$ as the sup-norm over $[-C, C]^{n_1}$, defined as

$$
\|\mathbf{f}\|_{L^{\infty}([-C,C]^{n_1})} = \sup_{\mathbf{x}\in[-C,C]^{n_1}} \|\mathbf{f}(\mathbf{x})\|_{\infty}.
$$

The following lemma provides a covering number of $\mathcal{F}_{\text{WSNN}}$. Our main proof strategy follows the proof of Lemma 5 from [Schmidt-Hieber](#page-26-0) [\(2020\)](#page-26-0), with modifications for weight-sharing networks.

Lemma C.2 (Covering number of $\mathcal{F}_{\text{WSNN}}$). Let $C, \delta > 0$ be given. For the class of weight-sharing neural networks $\mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\text{m}})$, we have

$$
\log N\left(\delta, \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\mathbf{m}}), \|\cdot\|_{L^{\infty}([-C,C]^{d_1})}\right) \leq (s+1)\log \left(\frac{4L^2 \|\mathbf{d}\|_{\infty}^2 \{\|\mathbf{m}\|_{\infty} \|\mathbf{d}\|_{\infty} (M \vee 1)\}^L (L + C + 2)}{\delta}\right).
$$

For any function class F, consisting of functions $\mathbf{f} : \mathbb{R}^D \times \mathbb{R} \to \mathbb{R}^D$, and $\underline{T}, \overline{T}, F, m > 0$ with $\underline{T} < \overline{T}$, define a class $\mathcal{L}^{\mathcal{I},T}(\mathcal{F},F,m)$ as

$$
\mathcal{L}^{\underline{T},\overline{T}}(\mathcal{F},F,m)=\left\{\ell_{\mathbf{f}}(\cdot): \mathbf{f}\in\mathcal{F} \quad \text{and} \quad \|\mathbf{f}(\mathbf{x},t)\|_{\infty}\leq F\sqrt{\log m}/\sigma_t, \quad \forall \mathbf{x}\in\mathbb{R}^D, t>0\right\},\
$$

where the loss function $\ell_f(\cdot)$ is defined in [\(2.5\)](#page-5-0) depending on (T, \overline{T}) . The following proposition provides a covering number of $\mathcal{L}^{T,T}$. Our main proof strategy follows the proof of Lemma C.2 from [Oko et al.](#page-25-0) [\(2023\)](#page-25-0), with modifications for weight-sharing networks.

Proposition C.1 (Covering number for the class of loss functions). Let F, $\tau_{\text{tail}}, \tau_{\text{min}}, \tau_{\text{max}} > 0, m > e$ be given. For the class of weight-sharing neural networks $\mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\text{m}})$, there exists a positive constant $\widetilde{C}_{14} = \widetilde{C}_{14}(D, \underline{\tau}, \tau_{\text{tail}}, \tau_{\text{min}}, \tau_{\text{max}}, F)$ such that

$$
\log N\left(\widetilde{C}_{14}m^{-\tau_{\rm tail}^2/4}\{\log m\}^2, \mathcal{L}^{\underline{T},\overline{T}}\left(\mathcal{F}_{\rm WSNN}\left(L,\mathbf{d},s,M,\mathcal{P}_{\mathbf{m}}\right),F,m\right),\|\cdot\|_{L^{\infty}([-1,1]^D)}\right) \leq (s+1)\log\left(\frac{4L^2\|\mathbf{d}\|_{\infty}^2\{\|\mathbf{m}\|_{\infty}\|\mathbf{d}\|_{\infty}(M\vee 1)\}^L\left\{L+(2+\tau_{\rm tail}\sqrt{\log m})\vee\tau_{\rm max}\sqrt{\log m}+2\right\}}{m^{-\tau_{\rm tail}^2/4}}\right),
$$

where $T = m^{-\tau_{\min}}$ and $\overline{T} = \tau_{\max} \log m$. Also, for any $\mathbf{f} \in \mathcal{F}(L, \mathbf{d}, s, M, \mathcal{P}_{\mathbf{m}})$ with $\|\mathbf{f}(\mathbf{x}, t)\|_{\infty} \leq$ F $\sqrt{\log m}/\sigma_t$ for $\mathbf{x} \in \mathbb{R}^D, t > 0$, we have

$$
\ell_{\mathbf{f}}(\mathbf{x}) \le \widetilde{C}_{15}(\log m)^2, \quad \mathbf{x} \in [-1, 1]^D,
$$

where $\widetilde{C}_{15} = \widetilde{C}_{15}(D, \underline{\tau}, \tau_{\min}, \tau_{\max}, F)$. In addition, for large enough m so that

$$
\widetilde{C}_{14}m^{-\tau_{\text{tail}}^2/4}(\log m)^2 \le DF^2/6 \quad \text{and} \quad \overline{T}-\underline{T} > 1,
$$

we have

$$
N\left(\widetilde{C}_{14}m^{-\tau_{\text{tail}}^2/4}\{\log m\}^2, \mathcal{L}^{\underline{T},\overline{T}}\left(\mathcal{F}_{\text{WSNN}}\left(L,\mathbf{d},s,M,\mathcal{P}_{\mathbf{m}}\right),F,m\right),\|\cdot\|_{L^{\infty}([-1,1]^D)}\right)\geq 3.
$$

C.1 Proof of Lemma [C.1](#page-87-0)

Proof. Note that p_0 is continuous and supported on $[-1, 1]^D$. Thus, for $t \ge 0$,

$$
\int_{\mathbb{R}^D} |p_0(\mathbf{x}) - p_t(\mathbf{x})| \, \mathrm{d}\mathbf{x} = \int_{\|\mathbf{x}\|_{\infty} \leq 1} |p_0(\mathbf{x}) - p_t(\mathbf{x})| \, \mathrm{d}\mathbf{x} + \int_{\|\mathbf{x}\|_{\infty} \geq 1} p_t(\mathbf{x}) \mathrm{d}\mathbf{x}.
$$

We will derive error bounds for each integral on the RHS.

Note that $p_0 \in \mathcal{H}^{\beta,K}([-1,1]^D)$. If $\beta \leq 1$, $|p_0(\mathbf{x}) - p_0(\mathbf{y})| \leq K \|\mathbf{x} - \mathbf{y}\|_{\infty}^{\beta}$ for any $\mathbf{x}, \mathbf{y} \in [-1,1]^D$ by the definition of $\mathcal{H}^{\beta,K}$. If $\beta > 1$, Mean value theorem implies that $|p_0(\mathbf{x}) - p_0(\mathbf{y})| \le K \|\mathbf{x} - \mathbf{y}\|_2$ for any $\mathbf{x}, \mathbf{y} \in [-1, 1]^D$. Combining two cases, we have

$$
|p_0(\mathbf{x})-p_0(\mathbf{y})| \leq K D \|\mathbf{x}-\mathbf{y}\|_{\infty}^{\beta \wedge 1}, \quad \mathbf{x}, \mathbf{y} \in [-1,1]^D.
$$

Then, for any $t \geq 0$,

$$
\int_{\|\mathbf{x}\|_{\infty}\leq 1} |p_0(\mathbf{x}) - p_t(\mathbf{x})| \, \mathrm{d}\mathbf{x} = \int_{\|\mathbf{x}\|_{\infty}\leq 1} \left| \int_{\|\mathbf{y}\|_{\infty}\leq 1} \left\{ p_0(\mathbf{x}) - p_0(\mathbf{y}) \right\} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}\mathbf{y} \right| \mathrm{d}\mathbf{x} \leq K D \int_{\|\mathbf{x}\|_{\infty}\leq 1} \int_{\|\mathbf{y}\|_{\infty}\leq 1} \|\mathbf{x} - \mathbf{y}\|_{\infty}^{\beta \wedge 1} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d}\mathbf{y} \mathrm{d}\mathbf{x}.
$$

For any $\delta > 0$, the last display is further bounded by

$$
KD\delta^{\beta \wedge 1} + KD \int_{\|\mathbf{x}\|_{\infty} \le 1} \int_{\|\mathbf{x}\|_{\infty} \le 1} \|\mathbf{x} - \mathbf{y}\|_{\infty}^{\beta \wedge 1} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d} \mathbf{y} \mathrm{d} \mathbf{x}
$$

$$
\le KD\delta^{\beta \wedge 1} + KD2^{\beta \wedge 1} \int_{\|\mathbf{x}\|_{\infty} \le 1} \int_{\|\mathbf{x}\|_{\infty} \le 1} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) \mathrm{d} \mathbf{y} \mathrm{d} \mathbf{x}.
$$
 (C.1)

Since $1 - \mu_t \geq 0$, we have $\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \geq \|\mathbf{x} - \mathbf{y}\|_{\infty} - (1 - \mu_t) \|\mathbf{y}\|_{\infty} \geq \delta - (1 - \mu_t)$ for $\|\mathbf{x} - \mathbf{y}\|_{\infty} \geq \delta$ and $||y||_{\infty} \leq 1$. Then, a simple calculation yields that

$$
\begin{aligned} &\int_{\|\mathbf{x}\|_{\infty}\leq 1}\int_{\|\mathbf{x}\|_{\infty}\leq 1}\phi_{\sigma t}(\mathbf{x}-\mu_{t}\mathbf{y})\mathrm{d}\mathbf{y}\mathrm{d}\mathbf{x}\leq \int_{\|\mathbf{x}\|_{\infty}\leq 1}\int_{\|\mathbf{x}-\mu_{t}\mathbf{y}\|_{\infty}\geq \delta-(1-\mu_{t})}\phi_{\sigma t}(\mathbf{x}-\mu_{t}\mathbf{y})\mathrm{d}\mathbf{y}\mathrm{d}\mathbf{x}\\ &=\mu_{t}^{-D}\int_{\|\mathbf{x}\|_{\infty}\leq 1}\int_{\|\mathbf{z}\|_{\infty}\geq \frac{\delta-(1-\mu_{t})}{\sigma_{t}}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\mathrm{d}\mathbf{x}=2^{D}\mu_{t}^{-D}\int_{\|\mathbf{z}\|_{\infty}\geq \frac{\delta-(1-\mu_{t})}{\sigma_{t}}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\\ &\leq 2^{D}\mu_{t}^{-D}\sum_{i=1}^{D}\int_{|z_{i}|\geq \frac{\delta-(1-\mu_{t})}{\sigma_{t}}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\leq D2^{D+1}\mu_{t}^{-D}\exp\left(-\frac{(\delta-(1-\mu_{t}))^{2}}{2\sigma_{t}^{2}}\right) \end{aligned}
$$

for $\delta \geq 1 - \mu_t$, where the last inequality holds by the tail probability of the standard normal distribution. By $(B.1)$, we have

$$
\frac{1}{2} \le 1 - \overline{\tau}t \le \mu_t \le 1 - \frac{\tau t}{2} \quad \text{and} \quad \sqrt{\tau t} \le \sigma_t \le \sqrt{2\overline{\tau}t}
$$

for any $0 \le t \le (2\overline{\tau})^{-1}$. Let $\delta = 1 - \mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)}$. Combining with [\(C.1\)](#page-89-0), we have

$$
\int_{\|\mathbf{x}\|_{\infty}\leq 1} |p_0(\mathbf{x}) - p_t(\mathbf{x})| \, \mathrm{d}\mathbf{x} \leq K D \delta^{\beta \wedge 1} + K D^2 2^{2D+1+\beta \wedge 1} \exp\left(-\frac{(\delta - (1 - \mu_t))^2}{2\sigma_t^2}\right)
$$
\n
$$
\leq K D \left(\overline{\tau} t + 2\sqrt{2\overline{\tau} t} \log(1/\sqrt{\underline{\tau} t})\right)^{\beta \wedge 1} + K D^2 2^{2D+1+\beta \wedge 1} \left(\sqrt{2\overline{\tau} t}\right)^2
$$
\n
$$
\leq D_1 \left\{ t \log(1/t) \right\}^{\frac{\beta \wedge 1}{2}}, \quad 0 \leq t \leq (2\overline{\tau})^{-1} \wedge 1,
$$

where $D_1 = D_1(\beta, D, K, \overline{\tau}, \underline{\tau}).$

Note that $\|\mathbf{x} - \mu_t \mathbf{y}\|_{\infty} \geq \|\mathbf{x}\|_{\infty} - \mu_t \|\mathbf{y}\|_{\infty} \geq 2\sigma_t \sqrt{\log(1/\sigma_t)}$ for $\|\mathbf{x}\|_{\infty} \geq \mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)}$ and ∥y∥[∞] ≤ 1. Then,

$$
\int_{\|\mathbf{x}\|_{\infty}\geq\mu_{t}+2\sigma_{t}\sqrt{\log(1/\sigma_{t})}}p_{t}(\mathbf{x})\mathrm{d}\mathbf{x} = \int_{\|\mathbf{x}\|_{\infty}\geq\mu_{t}+2\sigma_{t}\sqrt{\log(1/\sigma_{t})}}\int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y})\mathrm{d}\mathbf{y}\mathrm{d}\mathbf{x}
$$
\n
$$
= \int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\int_{\|\mathbf{x}\|_{\infty}\geq\mu_{t}+2\sigma_{t}\sqrt{\log(1/\sigma_{t})}}\phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y})\mathrm{d}\mathbf{x}\mathrm{d}\mathbf{y}
$$
\n
$$
\leq \int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\int_{\|\mathbf{x}-\mu_{t}\mathbf{y}\|_{\infty}\geq2\sigma_{t}\sqrt{\log(1/\sigma_{t})}}\phi_{\sigma_{t}}(\mathbf{x}-\mu_{t}\mathbf{y})\mathrm{d}\mathbf{x}\mathrm{d}\mathbf{y}
$$
\n
$$
= \mu_{t}^{-D}\int_{\|\mathbf{y}\|_{\infty}\leq1}p_{0}(\mathbf{y})\int_{\|\mathbf{z}\|_{\infty}\geq2\sqrt{\log(1/\sigma_{t})}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\mathrm{d}\mathbf{y} = \mu_{t}^{-D}\int_{\|\mathbf{z}\|_{\infty}\geq2\sqrt{\log(1/\sigma_{t})}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}
$$
\n
$$
\leq \mu_{t}^{-D}\sum_{i=1}^{D}\int_{|z_{i}|\geq2\sqrt{\log(1/\sigma_{t})}}\phi_{1}(\mathbf{z})\mathrm{d}\mathbf{z}\leq2D\mu_{t}^{-D}\sigma_{t}^{2}\leq D2^{D+1}\overline{\tau}t, \quad 0\leq t\leq(2\overline{\tau})^{-1},
$$

where the third inequality holds by the tail probability of the standard normal distribution. A simple calculation yields that

$$
\int_{1 \leq ||\mathbf{x}||_{\infty} \leq \mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)}} p_t(\mathbf{x}) d\mathbf{x}
$$
\n
$$
= \int_{1 \leq ||\mathbf{x}||_{\infty} \leq \mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)}} \int_{||\mathbf{y}||_{\infty} \leq 1} p_0(\mathbf{y}) \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} d\mathbf{x}
$$
\n
$$
\leq \int_{1 \leq ||\mathbf{x}||_{\infty} \leq \mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)}} K \int_{\mathbb{R}^D} \phi_{\sigma_t}(\mathbf{x} - \mu_t \mathbf{y}) d\mathbf{y} d\mathbf{x} = K \left| \left(\mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)} \right)^D - 1 \right|
$$
\n
$$
\leq K \sum_{k=1}^D {D \choose k} \left| \mu_t - 1 + 2\sigma_t \sqrt{\log(1/\sigma_t)} \right|^k, \quad t \geq 0.
$$

Note that

$$
\left|\mu_t - 1 + 2\sigma_t \sqrt{\log(1/\sigma_t)}\right| \le |\mu_t - 1| + \left|2\sigma_t \sqrt{\log(1/\sigma_t)}\right|
$$

$$
\le \overline{\tau}t + 2\sqrt{2\overline{\tau}t\log(1/\sqrt{\underline{\tau}t})} \le D_2\sqrt{t\log(1/t)}, \quad 0 \le t \le (2\overline{\tau})^{-1} \wedge 1,
$$

where $D_2 = D_2(\bar{\tau}, \bar{\tau})$. Let $D_3 = D_3(D_2, \bar{\tau})$ be a small enough constant so that $D_3 \leq (2\bar{\tau})^{-1} \wedge 1$ and $D_2\sqrt{t\log(1/t)} \leq 1$. Combining with the last two displays, we have

$$
\int_{1 \leq ||\mathbf{x}||_{\infty} \leq \mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)}} p_t(\mathbf{x}) d\mathbf{x} \leq K D^{D+1} D_2 \sqrt{t \log(1/t)}, \quad 0 \leq t \leq D_3.
$$

Therefore,

$$
\int_{\mathbb{R}^D} |p_0(\mathbf{x}) - p_t(\mathbf{x})| \, d\mathbf{x} = \int_{\|\mathbf{x}\|_{\infty} \le 1} |p_0(\mathbf{x}) - p_t(\mathbf{x})| \, d\mathbf{x} + \int_{\|\mathbf{x}\|_{\infty} \ge 1} p_t(\mathbf{x}) d\mathbf{x} \n\le D_1 \left\{ t \log(1/t) \right\}^{\frac{\beta \wedge 1}{2}} + \int_{1 \le \|\mathbf{x}\|_{\infty} \le \mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)}} p_t(\mathbf{x}) d\mathbf{x} + \int_{\|\mathbf{x}\|_{\infty} \ge \mu_t + 2\sigma_t \sqrt{\log(1/\sigma_t)}} p_t(\mathbf{x}) d\mathbf{x} \n\le D_1 \left\{ t \log(1/t) \right\}^{\frac{\beta \wedge 1}{2}} + KD^{D+1} D_2 \sqrt{t \log(1/t)} + D2^{D+1} \overline{\tau}t \n\le D_4 \left\{ t \log(1/t) \right\}^{\frac{\beta \wedge 1}{2}}, \quad 0 \le t \le D_3,
$$

where $D_4 = D_1 + D2^{D+1} \overline{\tau} + K D^{D+1} D_2$. The assertion follows by re-defining the constants. \Box

C.2 Proof of Lemma [C.2](#page-88-0)

Proof. For neural networks $f, f \in \mathcal{F}_{WSNN}(L, d, s, M, \mathcal{P}_m)$, let $\{W_l, \mathbf{b}_l\}_{l \in [L]}$ and $\{W_l, \mathbf{b}_l\}_{l \in [L]}$ be the parameter matrices of f and \tilde{f} , repsectively. For $l \in [L-1]$, let

$$
\mathbf{f}_l(\cdot) = \rho \left(\sum_{j=1}^{m_l} R_l^{(j)} \left(W_l Q_l^{(j)} \cdot + \mathbf{b}_l \right) \right) \quad \text{and} \quad \widetilde{\mathbf{f}}_l(\cdot) = \rho \left(\sum_{j=1}^{m_l} R_l^{(j)} \left(\widetilde{W}_l Q_l^{(j)} \cdot + \widetilde{\mathbf{b}}_l \right) \right).
$$

Given $\epsilon > 0$, assume that all parameter values of f and \tilde{f} are at most ϵ away from each other. Then,

$$
\|\mathbf{f}_{l}(\mathbf{x}) - \mathbf{f}_{l}(\widetilde{\mathbf{x}})\|_{\infty} \leq \left\| \sum_{j=1}^{m_{l}} R_{l}^{(j)} W_{l} Q_{l}^{(j)}(\mathbf{x} - \widetilde{\mathbf{x}}) \right\|_{\infty}
$$

\n
$$
\leq \sum_{j=1}^{m_{l}} \left\| W_{l} Q_{l}^{(j)}(\mathbf{x} - \widetilde{\mathbf{x}}) \right\|_{\infty} \leq \sum_{j=1}^{m_{l}} \left\{ \|W_{l}\|_{\infty} \cdot \left\| Q_{l}^{(j)}(\mathbf{x} - \widetilde{\mathbf{x}}) \right\|_{\infty} \right\}
$$

\n
$$
\leq m_{l} d_{l} M \|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty}, \quad l \in [L - 1], \ \mathbf{x}, \widetilde{\mathbf{x}} \in \mathbb{R}^{d_{l}}.
$$

because for $\mathbf{z} \in \mathbb{R}^{n_2}$, $n_1, n_2 \in \mathbb{N}$, we have $||W\mathbf{z}||_{\infty} \leq n_2 ||W||_{\infty} ||\mathbf{z}||_{\infty}$ for any matrix $W \in \mathbb{R}^{n_1 \times n_2}$ and $||Qz||_{\infty} = ||z||_{\infty}$ for any $n_1 \times n_1$ permutation matrix Q. A simple calculation yields that

$$
\|(\mathbf{f}_{L-1} \circ \cdots \circ \mathbf{f}_l)(\mathbf{x}) - (\mathbf{f}_{L-1} \circ \cdots \circ \mathbf{f}_l)(\widetilde{\mathbf{x}})\|_{\infty} \n\leq \left(\prod_{i=l}^{L-1} m_i d_i M\right) \|\mathbf{x} - \widetilde{\mathbf{x}}\|_{\infty}, \quad l \in [L-1], \mathbf{x}, \widetilde{\mathbf{x}} \in \mathbb{R}^{d_l}.
$$
\n(C.2)

Similarly, we have

$$
\left\| \mathbf{f}_{l}(\mathbf{x}) - \widetilde{\mathbf{f}}_{l}(\mathbf{x}) \right\|_{\infty} \leq \left\| \sum_{j=1}^{m_{l}} R_{l}^{(j)} \left(\left(W_{l} - \widetilde{W}_{l} \right) Q_{l}^{(j)} \mathbf{x} + \mathbf{b}_{l} - \widetilde{\mathbf{b}}_{l} \right) \right\|_{\infty} \n\leq \sum_{j=1}^{m_{l}} \left\| \left(W_{l} - \widetilde{W}_{l} \right) Q_{l}^{(j)} \mathbf{x} + \left(\mathbf{b}_{l} - \widetilde{\mathbf{b}}_{l} \right) \right\|_{\infty} \leq \sum_{j=1}^{m_{l}} \left\{ \left\| W_{l} - \widetilde{W}_{l} \right\|_{\infty} \cdot \left\| Q_{l}^{(j)} \mathbf{x} \right\|_{\infty} + \left\| \mathbf{b}_{l} - \widetilde{\mathbf{b}}_{l} \right\|_{\infty} \right\} \n\leq m_{l} \epsilon (1 + d_{l} \|\mathbf{x}\|_{\infty}) \leq m_{l} d_{l} \epsilon (1 + \|\mathbf{x}\|_{\infty}), \quad l \in [L - 1], \ \mathbf{x} \in \mathbb{R}^{d_{l}}.
$$

It follows that

$$
\left\| \left(\mathbf{f}_{l+1} \circ \widetilde{\mathbf{f}}_l \circ \cdots \circ \widetilde{\mathbf{f}}_1 \right) (\mathbf{x}) - \left(\widetilde{\mathbf{f}}_{l+1} \circ \widetilde{\mathbf{f}}_l \circ \cdots \circ \widetilde{\mathbf{f}}_1 \right) (\mathbf{x}) \right\|_{\infty} \n\leq m_{l+1} d_{l+1} \epsilon \left\{ 1 + \left\| \left(\widetilde{\mathbf{f}}_l \circ \cdots \circ \widetilde{\mathbf{f}}_1 \right) (\mathbf{x}) \right\|_{\infty} \right\} \n\leq m_{l+1} d_{l+1} \epsilon \left\{ 1 + \left(1 \vee \prod_{i=1}^{l} m_i d_i M \right) (l + \|\mathbf{x}\|_{\infty}) \right\} \n\leq m_{l+1} d_{l+1} \epsilon \left(1 \vee \prod_{i=1}^{l} m_i d_i M \right) (l + 1 + \|\mathbf{x}\|_{\infty}), \quad l \in [L-2], \ \mathbf{x} \in \mathbb{R}^{d_1},
$$
\n(C.3)

where the second inequality holds because

$$
\begin{aligned}\n\left\|\widetilde{\mathbf{f}}_i(\mathbf{z})\right\|_{\infty} &\leq \sum_{j=1}^{m_i} \left\|R_i^{(j)}\left(\widetilde{W}_i Q_i^{(j)} \mathbf{z} + \widetilde{\mathbf{b}}_i\right)\right\| \\
&\leq m_i M \left(1 + d_i \|\mathbf{z}\|_{\infty}\right) \leq m_i d_i M \left(1 + \|\mathbf{z}\|_{\infty}\right), \quad i \in [l], \mathbf{z} \in \mathbb{R}^{d_i}.\n\end{aligned}
$$

Let

$$
\mathbf{f}^{(l)}(\cdot) = \left(\mathbf{f}_{L-1} \circ \cdots \circ \mathbf{f}_{l+1} \circ \widetilde{\mathbf{f}}_l \circ \cdots \widetilde{\mathbf{f}}_1\right)(\cdot), \quad l \in [L-2]
$$

$$
\mathbf{f}^{(0)}(\cdot) = \left(\mathbf{f}_{L-1} \circ \cdots \mathbf{f}_1\right)(\cdot) \quad \text{and} \quad \mathbf{f}^{(L-1)}(\cdot) = \left(\widetilde{\mathbf{f}}_{L-1} \circ \cdots \circ \widetilde{\mathbf{f}}_1\right)(\cdot).
$$

Combining with $(C.2)$ and $(C.3)$, we have

$$
\left\| \mathbf{f}^{(l)}(\mathbf{x}) - \mathbf{f}^{(l+1)}(\mathbf{x}) \right\|_{\infty}
$$
\n
$$
\leq \left(\prod_{i=l+2}^{L-1} m_i d_i M \right) \left\{ m_{l+1} d_{l+1} \epsilon \left(1 \vee \prod_{i=1}^{l} m_i d_i M \right) (l+1 + ||\mathbf{x}||_{\infty}) \right\}
$$
\n
$$
\leq (M \vee 1)^{-1} \left\{ \prod_{i=1}^{L-1} m_i d_i (M \vee 1) \right\} \epsilon (l+1 + ||\mathbf{x}||_{\infty}), \quad l \in [L-3], \mathbf{x} \in \mathbb{R}^{d_1}
$$

and

$$
\left\| \mathbf{f}^{(L-2)}(\mathbf{x}) - \mathbf{f}^{(L-1)}(\mathbf{x}) \right\|_{\infty} \le m_{L-1} d_{L-1} \epsilon \left\{ 1 + \left(1 \vee \prod_{i=1}^{L-2} m_i d_i M \right) (L - 2 + ||\mathbf{x}||_{\infty}) \right\}
$$

$$
\le (M \vee 1)^{-1} \left\{ \prod_{i=1}^{L-1} m_i d_i (M \vee 1) \right\} \epsilon (L - 1 + ||\mathbf{x}||_{\infty}), \quad \mathbf{x} \in \mathbb{R}^{d_1}
$$

A simple calculation yields that

$$
\left\| \mathbf{f}(\mathbf{x}) - \widetilde{\mathbf{f}}(\mathbf{x}) \right\|_{\infty} \leq \sum_{l=0}^{L-2} \left\| W_L \left(\mathbf{f}^{(l)}(\mathbf{x}) - \mathbf{f}^{(l+1)}(\mathbf{x}) \right) \right\| + \left\| \left(W_L - \widetilde{W}_L \right) \mathbf{f}^{(L-1)}(\mathbf{x}) - \left(\mathbf{b}_L - \widetilde{\mathbf{b}}_L \right) \right\|_{\infty} \n\leq d_L \left\| W_L \right\|_{\infty} \sum_{l=0}^{L-2} \left\| \mathbf{f}^{(l)}(\mathbf{x}) - \mathbf{f}^{(l+1)}(\mathbf{x}) \right\| + d_L \left\| W_L - \widetilde{W}_L \right\|_{\infty} \cdot \left\| \mathbf{f}^{(L-1)}(\mathbf{x}) \right\|_{\infty} + \left\| \mathbf{b}_L - \widetilde{\mathbf{b}}_L \right\|_{\infty} \n\leq d_L M (M \vee 1)^{-1} \left\{ \prod_{i=1}^{L-1} m_i d_i (M \vee 1) \right\} \epsilon \sum_{l=0}^{L-2} (l+1 + \|\mathbf{x}\|_{\infty}) + d_L \epsilon \left\{ \prod_{i=1}^{L-1} m_i d_i M \right\} (1 + \|\mathbf{x}\|_{\infty}) + \epsilon \n\leq \epsilon d_L \left\{ \prod_{i=1}^{L-1} m_i d_i (M \vee 1) \right\} \left[\frac{L(L-1)}{2} + L \|\mathbf{x}\|_{\infty} + 2 \right], \quad \mathbf{x} \in \mathbb{R}^{d_1}.
$$

For $\mathbf{x} \in [-C, C]^{d_1}$, the last display is bounded by

$$
\epsilon d_L L \left\{ \prod_{i=1}^{L-1} m_i d_i (M \vee 1) \right\} (L + C + 2) \stackrel{\text{def}}{=} \delta.
$$

The total number of parameters in **f** is $T \stackrel{\text{def}}{=} \sum_{i=1}^{L} (d_i + 1)d_{i+1}$ and there are $\binom{T}{s}$ s^T combinations to pick s non-zero paramters. Since $T \leq 2L \|\mathbf{d}\|_{\infty}^2$ and $\binom{T}{s}$ $\binom{T}{s} \leq (2L \|\mathbf{d}\|_{\infty})^s$, we have

$$
N\left(\delta, \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\mathbf{m}}), \|\cdot\|_{L^{\infty}[-C,C]^{d_1}}\right) \leq \sum_{s_0=1}^{s} {T \choose s_0} N\left(\epsilon, [-M, M]^{s_0}, \|\cdot\|_{\infty}\right)
$$

$$
\leq s \left(2L \|\mathbf{d}\|_{\infty}^2\right)^s \left(\frac{2M}{\epsilon}\right)^s = s \left(\frac{4ML^2 \|\mathbf{d}\|_{\infty}^2 d_L \left\{\prod_{i=1}^{L-1} m_i d_i(M \vee 1)\right\}(L+C+2)}{\delta}\right)^s.
$$

For $\delta < 1$, the last display is bounded by

$$
\left(\frac{4L^2\|\mathbf{d}\|_{\infty}^2 \left\{\|\mathbf{m}\|_{\infty}\|\mathbf{d}\|_{\infty}(M\vee 1)\right\}^L (L+C+2)}{\delta}\right)^{s+1}.
$$

The assertion follows by taking the logarithm.

 \Box

C.3 Proof of Proposition [C.1](#page-88-1)

Proof. Fix $\widetilde{\mathbf{x}} = (\widetilde{x}_1, \dots, \widetilde{x}_D)^\top \in [-1, 1]^D$, $F > 0$ and $m > e$. Note that $p_t(\cdot | \widetilde{\mathbf{x}}) = \phi_{\sigma_t}(\cdot - \mu_t \widetilde{\mathbf{x}})$ for any $t \geq 0$. Then,

$$
\int_{\|\mathbf{y}\|_{\infty}\geq\mu_{t}+\tau_{\text{tail}}\sigma_{t}\sqrt{\log m}} p_{t}(\mathbf{y} \mid \widetilde{\mathbf{x}}) d\mathbf{y} \n\leq \sum_{i=1}^{D} \int_{|y_{i}| \geq \mu_{t}\widetilde{x}_{i}+\tau_{\text{tail}}\sigma_{t}\sqrt{\log m}} \phi_{\sigma_{t}}(\mathbf{y}-\mu_{t}\widetilde{\mathbf{x}}) d\mathbf{y} = D \int_{|z| \geq \tau_{\text{tail}}\sqrt{\log m}} \phi(z) dz, \quad t \geq 0,
$$

where ϕ denotes the one-dimensional standard normal density. By the tail probability of normal distribution, the last display is bounded by $2Dm^{-\tau_{\text{tail}}^2/2}$. Note that $\nabla \log p_t(\cdot | \tilde{\mathbf{x}}) = -(-\mu_t \tilde{\mathbf{x}})/\sigma_t^2$ for $t \geq 0$. Then,

$$
\int_{\|\mathbf{y}\|_{\infty}\geq\mu_{t}+\tau_{\text{tail}}\sigma_{t}\sqrt{\log m}} \|\nabla \log p_{t}(\mathbf{y} \mid \widetilde{\mathbf{x}})\|_{2}^{2} p_{t}(\mathbf{y} \mid \widetilde{\mathbf{x}}) d\mathbf{y} \n\leq \sum_{i=1}^{D} \int_{|y_{i}| \geq \mu_{t}\widetilde{x}_{i}+\tau_{\text{tail}}\sigma_{t}\sqrt{\log m}} \sum_{j=1}^{D} \left(\frac{y_{j}-\mu_{t}\widetilde{x}_{j}}{\sigma_{t}^{2}}\right)^{2} \phi_{\sigma_{t}}(\mathbf{y}-\mu_{t}\widetilde{\mathbf{x}}) d\mathbf{y} \n= \sum_{i=1}^{D} \sum_{j=1}^{D} \sigma_{t}^{-2} \int_{|z_{i}| \geq \tau_{\text{tail}}\sqrt{\log m}} z_{j}^{2} \phi_{1}(\mathbf{z}) d\mathbf{z}, \quad t \geq 0,
$$

where $\mathbf{y} = (y_1, \ldots, y_D)$ and $\mathbf{z} = (z_1, \ldots, z_D)$. By the Cauchy-Schwarz inequality, we have

$$
\int_{|z| \ge \tau_{\text{tail}} \sqrt{\log m}} z^2 \phi(z) dz \le 2\sqrt{\mathbb{E}[Z^4]} m^{-\tau_{\text{tail}}^2/4} = 2\sqrt{3}m^{-\tau_{\text{tail}}^2/4},
$$

where Z denotes the one-dimensional standard normal random variable. Combining with the last two displays, we have

$$
\int_{\|\mathbf{y}\|_{\infty}\geq\mu_{t}+\tau_{\text{tail}}\sigma_{t}\sqrt{\log m}} \|\nabla \log p_{t}(\mathbf{y} \mid \widetilde{\mathbf{x}})\|_{2}^{2} p_{t}(\mathbf{y} \mid \widetilde{\mathbf{x}}) d\mathbf{y}
$$
\n
$$
\leq D\sigma_{t}^{-2} \left\{ (D-1)\mathbb{E}[Z^{2}] \int_{|z|\geq\tau_{\text{tail}}\sqrt{\log m}} \phi(z) dz + \int_{|z|\geq\tau_{\text{tail}}\sqrt{\log m}} z^{2} \phi(z) dz \right\}
$$
\n
$$
\leq 2D(D+\sqrt{3}-1)\sigma_{t}^{-2} m^{-\tau_{\text{tail}}^{2}/4}, \quad t \geq 0.
$$

For any function $f \in \mathcal{F}(L, \mathbf{d}, s, M, \mathcal{P}_m)$ with $\|\mathbf{f}(\cdot, t)\|_{\infty} \leq F \sigma_t^{-1}$ √ $\overline{\log m}$ for $t \geq 0$, we have

$$
\begin{split}\n&\left|\ell_{\mathbf{f}}(\mathbf{x}) - \int_{\mathcal{I}}^{\overline{T}} \int_{\|\mathbf{y}\|_{\infty} \leq \mu_{t} + \tau_{\mathrm{tail}} \sigma_{t} \sqrt{\log m}} \|\mathbf{f}(\mathbf{y}, t) - \nabla \log p_{t}(\mathbf{y} \mid \mathbf{x})\|_{2}^{2} p_{t}(\mathbf{y} \mid \mathbf{x}) \mathrm{d}\mathbf{y} \mathrm{d}t\right| \\
&= \left|\int_{\mathcal{I}}^{\overline{T}} \int_{\|\mathbf{y}\|_{\infty} \geq \mu_{t} + \tau_{\mathrm{tail}} \sigma_{t} \sqrt{\log m}} \|\mathbf{f}(\mathbf{y}, t) - \nabla \log p_{t}(\mathbf{y} \mid \mathbf{x})\|_{2}^{2} p_{t}(\mathbf{y} \mid \mathbf{x}) \mathrm{d}\mathbf{y} \mathrm{d}t\right| \\
&\leq \int_{\mathcal{I}}^{\overline{T}} D \|\mathbf{f}(\cdot, t)\|_{\infty}^{2} \int_{\|\mathbf{y}\|_{\infty} \geq \mu_{t} + \tau_{\mathrm{tail}} \sigma_{t} \sqrt{\log m}} p_{t}(\mathbf{y} \mid \mathbf{x}) \mathrm{d}\mathbf{y} + 2D(D + \sqrt{3} - 1) \sigma_{t}^{-2} m^{-\tau_{\mathrm{tail}}^{2}/4} \mathrm{d}t \\
&\leq D \left\{ 2F^{2} D m^{-\tau_{\mathrm{tail}}^{2}/2} \log m + 2(D + \sqrt{3} - 1) m^{-\tau_{\mathrm{tail}}^{2}/4} \right\} \int_{\overline{T}}^{\overline{T}} \sigma_{t}^{-2} \mathrm{d}t, \quad \mathbf{x} \in [-1, 1]^{D}.\n\end{split}
$$

By [\(B.1\)](#page-41-0), $\sigma_t \geq \sqrt{\tau t}$ for $t \geq 0$. Then, the last display is bounded by

$$
D_1 m^{-\tau_{\text{tail}}^2/4} \{ \log m \}^2, \tag{C.4}
$$

where $D_1 = D_1(D, \underline{\tau}, \tau_{\min}, \tau_{\max}, F)$.

Let $\epsilon = m^{-\tau_{\text{tail}}^2/4}$ and $C = (2+\tau_{\text{tail}}\sqrt{\log m})\sqrt{\tau_{\text{max}}\sqrt{\log m}}$. Consider functions $\mathbf{f}_1, \mathbf{f}_2 \in \mathcal{F}(L, \mathbf{d}, s, M, \mathcal{P}_{\text{m}})$ such that $||\mathbf{f}_1(\cdot) - \mathbf{f}_2(\cdot)||_{L^{\infty}([-C,C]^{D+1})} \leq \epsilon$. For $\mathbf{y} \in [-C, C]^D, t \geq 0$, we have

$$
\begin{aligned}\n\|\mathbf{f}_1(\mathbf{y},t) - \nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}})\|_2^2 &- \|\mathbf{f}_2(\mathbf{y},t) - \nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}})\|_2^2 \\
&= \sum_{i=1}^D \left[\left\{ (\mathbf{f}_1(\mathbf{y},t))_i - (\nabla \log p_t(\mathbf{x}))_i \right\}^2 - \left\{ (\mathbf{f}_2(\mathbf{y},t))_i - (\nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}))_i \right\}^2 \right] \\
&= \sum_{i=1}^D \left\{ (\mathbf{f}_1(\mathbf{y},t))_i - (\mathbf{f}_2(\mathbf{y},t))_i \right\} \left\{ (\mathbf{f}_1(\mathbf{y},t))_i + (\mathbf{f}_2(\mathbf{y},t))_i - 2(\nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}))_i \right\}.\n\end{aligned}
$$

Thus,

$$
\begin{aligned}\n&\left|\|\mathbf{f}_1(\mathbf{y},t)-\nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}})\|_2^2 - \|\mathbf{f}_2(\mathbf{y},t)-\nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}})\|_2^2\right| \\
&\leq \epsilon \sigma_t^{-2} \left(2FD\sqrt{\log m} + 2\|\mathbf{y} - \mu_t \widetilde{\mathbf{x}}\|_1\right), \quad \mathbf{y} \in [-C,C]^D, t \geq 0.\n\end{aligned}
$$

Since $\mu_t \leq 1$ and $\sigma_t \leq 1$ for $t \geq 0$, we have

$$
\int_{\underline{T}}^{\overline{T}} \int_{\|\mathbf{y}\|_{\infty} \leq \mu_t + \tau_{\text{tail}} \sigma_t \sqrt{\log m}} \left| \|\mathbf{f}_1(\mathbf{y}, t) - \nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}) \|_2^2 - \|\mathbf{f}_2(\mathbf{y}, t) - \nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}) \|_2^2 \right| p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}) \mathrm{d}\mathbf{y} \mathrm{d}t
$$
\n
$$
\leq \int_{\underline{T}}^{\overline{T}} \int_{\|\mathbf{y}\|_{\infty} \leq \mu_t + \tau_{\text{tail}} \sigma_t \sqrt{\log m}} \epsilon \sigma_t^{-2} \left(2CD \sqrt{\log m} + 2\|\mathbf{y} - \mu_t \widetilde{\mathbf{x}} \|_1 \right) p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}) \mathrm{d}\mathbf{y} \mathrm{d}t
$$
\n
$$
\leq 2\underline{\tau} FD \epsilon \sqrt{\log m} \left(\log \overline{T} - \log \underline{T} \right) + 2D\epsilon \int_{\underline{T}}^{\overline{T}} \sigma_t^{-2} \left(2\mu_t + \tau_{\text{tail}} \sigma_t \sqrt{\log m} \right) \mathrm{d}t.
$$

Moreover, the last display is bounded by

$$
2\tau F D\epsilon \sqrt{\log m} \left(\log \overline{T} - \log \underline{T} \right) + 2\tau D\epsilon \left(2 + \tau_{\text{tail}} \sqrt{\log m} \right) \left(\log \overline{T} - \log \underline{T} \right) \le D_2 \epsilon \{\log m\}^{3/2},
$$

where $D_2 = D_2(D, \underline{\tau}, \tau_{\text{tail}}, \tau_{\text{min}}, \tau_{\text{max}}, F)$.

Combining with $(C.4)$, we have

$$
\left| l_{\mathbf{f}_1}^{\overline{T},\overline{T}}(\mathbf{x}) - l_{\mathbf{f}_2}^{\overline{T},\overline{T}}(\mathbf{x}) \right| \le 2D_1 m^{-\tau_{\rm tail}^2/4} {\{\log m\}}^2 + 2D_2 \epsilon {\{\log m\}}^{3/2}
$$

$$
\le D_3 m^{-\tau_{\rm tail}^2/4} {\{\log m\}}^2, \quad \mathbf{x} \in [-1,1]^D.
$$

where $D_3 = (2D_1 + 2D_2)$. Then,

$$
\log N\left(D_3 m^{-\tau_{\rm tail}^2/4} {\log m} \right)^2, \mathcal{L}^{\underline{T}, \overline{T}}\left(\mathcal{F}\left(L, \mathbf{d}, s, M, \mathcal{P}_m\right), F, m\right), \|\cdot\|_{L^{\infty}([-1,1]^D)}\right) \leq \log N\left(m^{-\tau_{\rm tail}^2/4}, \mathcal{F}\left(L, \mathbf{d}, s, M, \mathcal{P}_m\right), \|\cdot\|_{L^{\infty}([-C, C]^{D+1})}\right).
$$

Combining with Lemma [C.2,](#page-88-0) the last display is bounded by

$$
(s+1)\log\left(\frac{4 L^2\|\mathbf{d}\|_\infty^2 \left\{\|\mathbf{m}\|_\infty \|\mathbf{d}\|_\infty (M\vee 1)\right\}^L\left\{L+\left(2+\tau_{\mathrm{tail}}\sqrt{\log m}\right)\vee\tau_{\mathrm{max}}\sqrt{\log m}+2\right\}}{m^{-\tau_{\mathrm{tail}}^2/4}}\right).
$$

The first assertion follows by re-defining the constant.

For any function $f \in \mathcal{F}(L, \mathbf{d}, s, M, \mathcal{P}_m)$ with $\|\mathbf{f}(\cdot, t)\|_{\infty} \leq F \sigma_t^{-1}$ √ $\overline{\log m}$ for $t \geq 0$,

$$
\ell_{\mathbf{f}}(\mathbf{x})
$$
\n
$$
\leq 2D \int_{\underline{T}}^{\overline{T}} \|\mathbf{f}(\cdot,t)\|_{\infty}^{2} \int_{\mathbb{R}^{D}} \phi_{\sigma_{t}}(\mathbf{y}-\mu_{t}\mathbf{x}) \mathrm{d}\mathbf{y} \mathrm{d}t + 2 \int_{\underline{T}}^{\overline{T}} \int_{\mathbb{R}^{D}} \left\|\frac{\mathbf{y}-\mu_{t}\mathbf{x}}{\sigma_{t}^{2}}\right\|_{2}^{2} \phi_{\sigma_{t}}(\mathbf{y}-\mu_{t}\mathbf{x}) \mathrm{d}\mathbf{y} \mathrm{d}t
$$
\n
$$
\leq 2DF \log m \int_{\underline{T}}^{\overline{T}} \sigma_{t}^{-2} \mathrm{d}t + 2 \left(\int_{\underline{T}}^{\overline{T}} \sigma_{t}^{-2} \mathrm{d}t\right) \left(\int_{\mathbb{R}^{D}} \|\mathbf{z}\|_{2}^{2} \phi_{1}(\mathbf{z}) \mathrm{d}\mathbf{z}\right)
$$
\n
$$
\leq 2DF \underline{\tau} \log m \left(\log \overline{T} - \log \underline{T}\right) + 2D \underline{\tau} \left(\log \overline{T} - \log \underline{T}\right) \leq D_{4} \{\log m\}^{2}, \quad \mathbf{x} \in [-1,1]^{D},
$$

where $D_4 = D_4(D, \underline{\tau}, \tau_{\min}, \tau_{\max}, F)$. The second assertion follows by re-defining the constant.

Consider vector-valued constant functions $\tilde{f}_1, \tilde{f}_2, \tilde{f}_3$ such that $f_i(y, t) = (Fi/3, \ldots, Fi/3)^\top \in \mathbb{R}^D$ for $i \in \{1, 2, 3\}, \mathbf{y} \in \mathbb{R}^D$ and $t \in \mathbb{R}$. Then,

$$
\left\| \widetilde{\mathbf{f}}_i(\mathbf{y}, t) - \nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}) \right\|_2^2 - \left\| \widetilde{\mathbf{f}}_j(\mathbf{y}, t) - \nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}) \right\|_2^2
$$

\n
$$
= \sum_{k=1}^D \left\{ F(i - j)/3 \right\} \left\{ F(i + j)/3 - 2(\nabla \log p_t(\mathbf{y} \mid \widetilde{\mathbf{x}}))_k \right\}
$$

\n
$$
= \left\{ F(i - j)/3 \right\} \left\{ DF(i + j)/3 + 2 \sum_{k=1}^D \frac{y_k - \mu_t \widetilde{x}_k}{\sigma_t^2} \right\}, \quad i, j \in \{1, 2, 3\}
$$

for $\mathbf{y} = (y_1, \dots, y_D)^\top \in \mathbb{R}^D$ and $t > 0$. It follows that

$$
\ell_{\tilde{f}_i}(\mathbf{x}) - \ell_{\tilde{f}_j}(\mathbf{x})
$$
\n
$$
= \int_{\underline{T}}^{\overline{T}} \int_{\mathbb{R}^D} \{F(i-j)/3\} \left\{ DF(i+j)/3 + 2 \sum_{k=1}^D \frac{y_k - \mu_t \tilde{x}_k}{\sigma_t^2} \right\} p_t(\mathbf{y} \mid \mathbf{x}) \, \mathrm{d}\mathbf{y} \, \mathrm{d}t
$$
\n
$$
= \{ F(i-j)/3 \} \left\{ (\overline{T} - \underline{T}) DF(i+j)/3 + 2 \int_{\underline{T}}^{\overline{T}} \int_{\mathbb{R}^D} \sigma_t^{-1} \left(\sum_{k=1}^D z_k \right) \phi_1(\mathbf{z}) \, \mathrm{d}\mathbf{z} \, \mathrm{d}t \right\}
$$
\n
$$
= \left(\tau_{\text{max}} \log m - m^{-\tau_{\text{min}}} \right) DF^2 \left(i^2 - j^2 \right) / 9, \quad i \in \{1, 2, 3\}
$$

for $\mathbf{x} \in \mathbb{R}^D$. For large enough m so that $\tau_{\max} \log m - m^{-\tau_{\min}} > 1$, we have

$$
\left|\ell_{\widetilde{\mathbf{f}}_i}(\mathbf{x}) - \ell_{\widetilde{\mathbf{f}}_j}(\mathbf{x})\right| > DF^2/3, \quad i, j \in \{1, 2, 3\}, \ i \neq j.
$$

Thus, the following packing number (See [van der Vaart and Wellner](#page-27-0) [\(1996\)](#page-27-0) for the definition)

$$
D\left(DF^2/3, \mathcal{L}^{\underline{T}, \overline{T}}\left(\mathcal{F}\left(L, \mathbf{d}, s, M, \mathcal{P}_m\right), F, m\right), \|\cdot\|_{L^\infty([-1,1]^D)}\right)
$$

is lower bounded by 3. Combining with the inequality between pakcing number and covering number, we have

$$
N\left(DF^2/6, \mathcal{L}^{\underline{T}, \overline{T}}\left(\mathcal{F}\left(L, \mathbf{d}, s, M, \mathcal{P}_{\mathbf{m}}\right), F, m\right), \|\cdot\|_{L^{\infty}([-1,1]^D)}\right) \geq 3.
$$

For large enough m so that $D_3m^{-\tau_{\rm tail}^2/4}\{\log m\}^2 \le DF^2/6$, the last assertion follows by re-defining \Box the constants.

C.4 Proof of Theorem [5.2](#page-15-0)

In this subsection, we provide the proof of Theorem [5.2](#page-15-0) with auxiliary lemmas. For two probability measures P and Q on $\mathcal{X} \subseteq \mathbb{R}^D$, the Kullback-Leibler (KL) divergence is defined as

$$
KL(P, Q) = \begin{cases} \int_{\mathcal{X}} \log \frac{dP}{dQ} dP, & \text{if } P \ll Q\\ \infty, & \text{else.} \end{cases}
$$

We often denote $KL(P,Q)$ as $KL(p,q)$, where p and q are densities of P and Q, respectively. Hereafter, $C = C(\text{all})$ means that C is a constant depending on $(\beta, d, D, K, \tau_{\text{min}}, \tau_{\text{max}}, \tau_1, \tau_2, \overline{\tau}, \tau)$.

Proof of Theorem [5.2.](#page-15-0) Let $\tilde{\tau}_{min} = \tau_{min} d/(2\beta + d)$ and $\tilde{\tau}_{max} = \tau_{max} d/(2\beta + d)$. Let C_1, C_2 be the constants in Theorem [5.1](#page-11-0) depending on $(\beta, d, D, K, \tilde{\tau}_{min}, \tilde{\tau}_{max}, \tau_1, \tau_2, \overline{\tau}, \underline{\tau})$. Then, by replacing m in Theorem [5.1](#page-11-0) with $n^{d/(2\beta+d)}$, there exists a class of permutation matrices $\mathcal{P}_{\mathbf{m}}$ and a class of weight-sharing neural networks $\widetilde{\mathcal{F}}_{\text{WSNN}} = \mathcal{F}_{\text{WSNN}}(L, \mathbf{d}, s, M, \mathcal{P}_{\mathbf{m}})$ with

$$
L \le D_1 (\log n)^6 \log \log n, \quad \|\mathbf{d}\|_{\infty} \le D_1 n^{\frac{d(D+1)}{2\beta+d}} (\log n)^3,
$$

\n
$$
s \le D_1 n^{\frac{d}{2\beta+d}} (\log n)^5 \log \log n, \quad M \le \exp(D_1 {\log n})^6),
$$

\n
$$
\|\mathbf{m}\|_{\infty} \le D_1 n^{\frac{dD}{2\beta+d}}
$$

satisfying

$$
\inf_{\mathbf{f}\in\widetilde{\mathcal{F}}_{\text{WSNN}}\cap\mathcal{F}_{\infty}}\int_{\underline{T}}^{\overline{T}}\int_{\mathbb{R}^D}||\mathbf{f}(\mathbf{x},t)-\nabla\log p_t(\mathbf{x})||_2^2 p_t(\mathbf{x})\,d\mathbf{x}dt
$$
\n(C.5)\n
$$
\leq C_1 \left(\frac{d}{2\beta+d}\right)^{4(D-1+\beta\wedge 1)+1}n^{-\frac{2\beta}{2\beta+d}}(\log n)^{4(D-1+\beta\wedge 1)+1},
$$

for large enough *n* so that $n^{d/(2\beta+d)} \geq C_2$, where $D_1 = D_1(\beta, d, C_1)$ and

$$
\mathcal{F}_{\infty} = \left\{ \|\mathbf{f}(\mathbf{x},t)\|_{\infty} \leq C_1 \sqrt{d/(2\beta + d)} \sigma_t^{-1} \sqrt{\log n} \quad \forall \mathbf{x} \in \mathbb{R}^D, t > 0 \right\}.
$$

Let $\hat{\mathbf{f}}$ be an empirical risk minimizer over the class $\widetilde{\mathcal{F}}_{\text{WSNN}}$, defined as in [\(2.4\)](#page-4-0). By Triangle inequality, we have

$$
\mathbb{E}\left[d_{\rm TV}(P_0, \hat{P}_{\underline{T}})\right] \le d_{\rm TV}(P_0, P_{\underline{T}}) + \mathbb{E}\left[d_{\rm TV}(P_{\underline{T}}, \hat{P}_{\underline{T}})\right].\tag{C.6}
$$

We proceed to control each term on the RHS separately.

Let $\widetilde{C}_{12} = \widetilde{C}_{12}(\beta, D, K, \overline{\tau}, \underline{\tau})$ and $\widetilde{C}_{13}(\overline{\tau}, \underline{\tau})$ be the constants in Lemma [C.1.](#page-87-0) For large enough n so that $T \n\leq \widetilde{C}_{13}$, it follows that

$$
d_{\mathrm{TV}}(P_0, P_{\underline{T}}) = \frac{1}{2} \int_{\mathbb{R}^D} \left| p_0(\mathbf{x}) - p_{\underline{T}}(\mathbf{x}) \right| d\mathbf{x}
$$

\$\leq \widetilde{C}_{12} \left\{ \underline{T} \log(1/\underline{T}) \right\}^{\frac{\tau_{\min}(\beta \wedge 1)}{2}} / 2 = D_2 n^{-\frac{\tau_{\min}(\beta \wedge 1)}{2}} (\log n)^{\frac{\beta \wedge 1}{2}},

where $D_2 = \tilde{C}_{12} \tau_{\min}^{(\beta \wedge 1)/2} / 2$. Since $\tau_{\min} \ge \frac{2\beta}{(2\beta + d)(\beta \wedge 1)}$, the last display is bounded by

$$
D_2 n^{-\frac{\beta}{2\beta+d}} (\log n)^{\frac{\beta \wedge 1}{2}}.\tag{C.7}
$$

Consider functions $q_1, q_2 : \mathbb{R}^D \times [0, \overline{T} - \underline{T}] \to \mathbb{R}$ such that $q_1(\mathbf{x}, t) = p_{\overline{T}-t}(\mathbf{x})$ and $q_2(\mathbf{x}, t) = \widehat{p}_{\overline{T}-t}(\mathbf{x})$. Then, each q_1 and q_2 satisfy the corresponding well-known Fokker-Planck equation [\(Bris and Lions,](#page-22-0) [2008,](#page-22-0) [Bogachev et al.,](#page-22-1) [2022,](#page-22-1) [Pavliotis,](#page-25-1) [2014\)](#page-25-1) :

$$
\frac{\partial}{\partial t}q_1(\mathbf{x},t) = -\sum_{i=1}^D \frac{\partial}{\partial x_i} [\mathbf{b}_1(\mathbf{x},t)q_1(\mathbf{x},t)] + \sum_{i=1}^D \sum_{j=1}^D \frac{\partial^2}{\partial x_i \partial x_j} [a_1(t)q_1(\mathbf{x},t)]
$$

$$
\frac{\partial}{\partial t}q_2(\mathbf{x},t) = -\sum_{i=1}^D \frac{\partial}{\partial x_i} [\mathbf{b}_1(\mathbf{x},t)q_2(\mathbf{x},t)] + \sum_{i=1}^D \sum_{j=1}^D \frac{\partial^2}{\partial x_i \partial x_j} [a_2(t)q_2(\mathbf{x},t)],
$$

where

$$
\mathbf{b}_1(\mathbf{x}, t) = \alpha_{\overline{T}-t}\mathbf{x} + 2\alpha_{\overline{T}-t}\nabla \log p_{\overline{T}-t}(\mathbf{x}), \quad a_1(t) = \alpha_{\overline{T}-t},
$$

$$
\mathbf{b}_2(\mathbf{x}, t) = \alpha_{\overline{T}-t}\mathbf{x} + 2\alpha_{\overline{T}-t}\hat{\mathbf{f}}(\mathbf{x}, \overline{T}-t), \quad a_2(t) = \alpha_{\overline{T}-t}.
$$

Following the Remark 2.3 of [Bogachev et al.](#page-22-2) [\(2016\)](#page-22-2), we have

$$
d_{\text{TV}}(P_{\underline{T}}, \hat{P}_{\underline{T}}) = d_{\text{TV}}(q_1(\cdot, \overline{T} - \underline{T}), q_2(\cdot, \overline{T} - \underline{T}))
$$

\n
$$
\leq d_{\text{TV}}(q_1(\cdot, 0), q_2(\cdot, 0)) + \int_{\underline{T}}^{\overline{T}} \int_{\mathbb{R}^D} \alpha_t^{-1} \left\| \hat{\mathbf{f}}(\mathbf{x}, t) - \nabla \log p_t(\mathbf{x}) \right\|_2^2 p_t(\mathbf{x}) d\mathbf{x} dt.
$$
 (C.8)

By Pinsker's inequality, we have

$$
d_{\text{TV}}\left(q_1(\cdot,0), q_2(\cdot,0)\right) = d_{\text{TV}}\left(p_{\overline{T}}, \phi_1\right) \le \sqrt{\text{KL}\left(p_{\overline{T}}, \phi_1\right)/2} \tag{C.9}
$$

Since KL divergence is convex in its first argument, Jensen's inequality implies that

KL
$$
(p_{\overline{T}}, \phi_1) \leq \int_{\mathbb{R}^D} \text{KL} \left(\mathcal{N}(\mu_{\overline{T}} \mathbf{y}, \sigma_{\overline{T}} \mathbb{I}_D), \mathcal{N}(\mathbf{0}_D, \mathbb{I}_D) \right) dP_0(\mathbf{y}).
$$

because $p_{\overline{T}}(\cdot) = \int_{\mathbb{R}^D} \phi_{\sigma_{\overline{T}}}(\cdot - \mu_{\overline{T}} \mathbf{y}) \mathrm{d}P_0(\mathbf{y})$. KL divergence between two D-dimensional Gaussian random variables is known as

KL
$$
(\mathcal{N}(\mu_1, \Sigma_1), \mathcal{N}(\mu_2, \Sigma_2))
$$

= $\frac{1}{2} \left[\log \left(\frac{\det(\Sigma_2)}{\det(\Sigma_1)} \right) - D + (\mu_1 - \mu_2)^\top \Sigma_2^{-1} (\mu_1 - \mu_2) + \text{tr} (\Sigma_2^{-1} \Sigma_1) \right].$

Using that, we have

$$
\begin{split} &\text{KL}\left(p_{\overline{T}}, \phi_1\right) \le \int_{\mathbb{R}^D} \frac{1}{2} \left(\mu_{\overline{T}}^2 \|\mathbf{y}\|_2^2 + D\sigma_{\overline{T}} - D - D \log \sigma_{\overline{T}}\right) \mathrm{d}P_0(\mathbf{y}) \\ &= \frac{1}{2} \left(\mu_{\overline{T}}^2 \mathbb{E}_{P_0} \left[\|\mathbf{X}_0\|_2^2\right] + D\sigma_{\overline{T}} - D - D \log \sigma_{\overline{T}}\right) \\ &\le \frac{D}{2} \left\{\mu_{\overline{T}}^2 + \left|\sqrt{1 - \mu_{\overline{T}}^2} - 1\right| + \left|\log\left(1 - \mu_{\overline{T}}^2\right)/2\right|\right\}, \end{split}
$$

where the last inequality holds because $\mathbb{E}[\|\mathbf{X}_0\|_2^2] \le D$ and $\mu_{\overline{7}}^2$ $rac{2}{T}+\sigma \frac{2}{T}$ $\frac{2}{T} = 1$. Since $\mu_{\overline{T}} \leq \exp(-\underline{\tau}T)$ and $log(1 + x) \leq x$ for $x > -1$, the last display is bounded by

$$
\frac{D}{2}\left\{\mu_{\overline{T}}^2+\frac{\mu_{\overline{T}}^2}{\sqrt{1-\mu_{\overline{T}}^2+1}}+\frac{\mu_{\overline{T}}^2}{2}\right\}\leq \frac{5D\mu_{\overline{T}}^2}{4}\leq \left(\frac{5D}{4}\right)\exp\left(-2\underline{\tau}\tau_{\max}\log n\right)\leq \left(\frac{5D}{4}\right)n^{-\frac{2\beta}{2\beta+d}},
$$

where the last inequality holds because $\tau_{\text{max}} \geq \frac{\beta}{\tau(2\beta)}$ $\frac{\beta}{\mathcal{I}(2\beta+d)}$. Combining with [\(C.9\)](#page-97-0), we have

$$
d_{\text{TV}}(q_1(\cdot,0), q_2(\cdot,0)) = d_{\text{TV}}(p_{\overline{T}}, \phi_1) \le \sqrt{5D/8}n^{-\frac{\beta}{2\beta+d}}.\tag{C.10}
$$

Let

$$
\widetilde{\tau}_{\text{tail}} = \sqrt{8\beta/(2\beta + d)}
$$
 and $\widetilde{F} = C_1 \sqrt{d/(2\beta + d)}$.

Let $\widetilde{C}_{14} = \widetilde{C}_{14}(D, \underline{\tau}, \widetilde{\tau}_{\text{tail}}, \tau_{\text{min}}, \tau_{\text{max}}, \widetilde{F})$ and $\widetilde{C}_{15} = \widetilde{C}_{15}(D, \underline{\tau}, \widetilde{\tau}_{\text{tail}}, \tau_{\text{min}}, \tau_{\text{max}}, \widetilde{F})$ be the constants in Proposition [C.1.](#page-88-1) Then, by replacing m in Proposition [C.1](#page-88-1) with n , it follows that

$$
\log 3
$$
\n
$$
\leq \log N \left(\widetilde{C}_{14} n^{-2\beta/(2\beta+d)} \{ \log n \}^2, \mathcal{L}^{\mathcal{I}, \overline{T}} \left(\widetilde{\mathcal{F}}_{\text{WSNN}}, \widetilde{F}, n \right), \| \cdot \|_{L^{\infty}([-1,1]^D)} \right)
$$
\n
$$
\leq D_3 n^{\frac{d}{2\beta+d}} (\log n)^{17} (\log \log n)^2
$$

and

$$
\sup_{\substack{\mathbf{f}\in\widetilde{\mathcal{F}}_{\mathrm{WSNN}}\cap\mathcal{F}_{\infty}\\ \mathbf{y}\in[-1,1]^D}}\|\ell_{\mathbf{f}}(\mathbf{y})\|_{\infty}\leq\widetilde{C}_{15}(\log n)^2
$$

for large enough n so that $\widetilde{C}_{14}n^{-\widetilde{\tau}^2_{\text{tail}}/4}(\log n)^2 \leq D\widetilde{F}^2/6$ and $\overline{T}-\underline{T}>1$, where $D_3=D_3(\beta,d,D,\tau_{\text{max}},D_1)$. Combining the last two displays and $(C.5)$ with Theorem C.4 of [Oko et al.](#page-25-0) (2023) , we have

$$
\mathbb{E}\left[\int_{\underline{T}}^{\overline{T}}\int_{\mathbb{R}^D}\alpha_t^{-1}\left\|\widehat{\mathbf{f}}(\mathbf{x},t)-\nabla\log p_t(\mathbf{x})\right\|_2^2p_t(\mathbf{x})\mathrm{d}\mathbf{x}\mathrm{d}t\right]
$$

\n
$$
\leq \underline{\tau}^{-1}\mathbb{E}\left[\int_{\underline{T}}^{\overline{T}}\int_{\mathbb{R}^D}\left\|\widehat{\mathbf{f}}(\mathbf{x},t)-\nabla\log p_t(\mathbf{x})\right\|_2^2p_t(\mathbf{x})\mathrm{d}\mathbf{x}\mathrm{d}t\right]
$$

\n
$$
\leq D_4n^{-\frac{2\beta}{2\beta+d}}\left\{(\log n)^{4(D-1+\beta\wedge 1)+1}+(\log n)^{19}(\log\log n)^2\right\},
$$

where $D_4 = D_4(\beta, d, D, \tau_{\text{bd}}, \tau, C_1, \tilde{C}_{14}, \tilde{C}_{15}, D_3)$. Combining [\(C.8\)](#page-97-1) and [\(C.10\)](#page-97-2) with the last display, we have

$$
\mathbb{E}\left[d_{\text{TV}}\left(P_{\underline{T}}, \widehat{P}_{\underline{T}}\right)\right] \le D_5 n^{-\frac{\beta}{2\beta+d}} \left\{ (\log n)^{2(D-1+\beta\wedge 1)+1/2} + (\log n)^{10} \right\},\tag{C.11}
$$

where $D_5 = D_5(D, D_4)$ Combining [\(C.6\)](#page-96-1) and [\(C.7\)](#page-96-2) with the last display, we have

$$
\mathbb{E}\left[d_{\text{TV}}\left(P_0, \widehat{P}_{\underline{T}}\right)\right] \le D_6 n^{-\frac{\beta}{2\beta+d}} \left\{ (\log n)^{2(D-1+\beta\wedge 1)+1/2} + (\log n)^{10} \right\},\,
$$

where $D_6 = D_6(D_2, D_5)$. The assertion follows by re-defining the constants.

 \Box