# Bifurcation Analysis of Predator-Prey System using Conformable Fractional Order Discretization

January 7, 2025

Muhammad Rafaqat<sup>1</sup>, Abubakar Masha<sup>2</sup>, Nauman Ahmed<sup>1,3</sup>, Ali Raza<sup>4</sup>, Wojciech Sumelka<sup>5</sup>

<sup>1</sup> Department of Mathematics and Statistics, University of Lahore, Pakistan

<sup>2</sup> Department of Mathematics and Computer Science, Borno State University, Nigeria

<sup>3</sup> Department of Computer Science and Mathematics, Lebanese American University, Beirut, Lebanon <sup>4</sup> Department of Physical Analysis, University of Chenab, Gujrat, Pakistan<sup>5</sup> Institute of Structural

Analysis, Poznan University of Technology, Poznan, Poland

E.mail: <sup>5</sup> wojciech.sumelka@put.poznan.pl

#### 1 Abstract

In this paper, conformal fractional order discretization [\[31,](#page-14-0) [35,](#page-14-1) [36\]](#page-14-2) is used to analyze bifurcation analysis and stability of a predator-prey system. A continuous model has been discretized into a discrete one while preserving the fractional-order dynamics. This allows us to look more closely at the stability properties of the system and bifurcation phenomena, including period-doubling and Neimark-Sacker bifurcation. Through numerical and theoretical methods, this research investigated how the modification in system parameters affects the overall dynamics, which may have implications for ecological management and conservation strategies.

Keywords. Predator-Prey model; Fractional differential equations; Bifurcation; Stability results; simulations

# 2 Introduction

The fractional-order predator-prey system [\[1,](#page-12-0) [2\]](#page-12-1) is a development of the classical predator-prey model, such as the Lotka-Voltera system that describes the interaction between two species of predator and prey. In conventional models, ordinary differential equations with derivatives of integer order are typically used to illustrate population dynamics. The use of fractional calculus that deals with the derivatives and integrals of non-integer order gives rise to the fractional order predator-prey system [\[4\]](#page-13-0). The classical Lotka-Volterra predator-prey model is represented by the following set of ordinary differential equations (ODEs):

<span id="page-0-0"></span>
$$
\begin{aligned}\n\frac{dP}{dt} &= \alpha P - \beta PH, \\
\frac{dH}{dt} &= \delta PH - \gamma H,\n\end{aligned} \tag{2.1}
$$

where:

- $P$  represents the population of prey (e.g., rabbits),
- $H$  represents the population of predators (e.g., foxes),

•  $\alpha$ ,  $\beta$ ,  $\delta$ , and  $\gamma$  are parameters representing the birth rate of prey, predation rate, natural death rate of prey, and death rate of predators, respectively.

To introduce fractional-order derivatives into this model, one replaces the integer-order derivatives with fractional-order derivatives. For example, the fractional-order predator-prey model can be represented as follows:

$$
\begin{aligned}\n\frac{d^{\alpha}P}{dt^{\alpha}} &= \alpha P - \beta PH, \\
\frac{d^{\alpha}H}{dt^{\alpha}} &= \delta PH - \gamma H,\n\end{aligned} \tag{2.2}
$$

Here,  $\frac{d^{\alpha}}{dt^{\alpha}}$  denotes the fractional derivative of order  $\alpha$ , which introduces memory effects and non-local interactions into the system, allowing for a more accurate representation of real-world ecological dynamics. It has been noted that the specific form of the fractional derivative depends on the chosen fractional calculus approach (e.g., Riemann-Liouville, Caputo, etc.), and the choice of order  $\alpha$  affects the dynamics of the system.

Recently, some researchers have explored the applications of fractional order differential equations in various biological scenarios, such as ecological systems with delays [\[15,](#page-13-1) [16,](#page-13-2) [17\]](#page-13-3), epidemiological systems rooted in control  $[18]$ , and ecological systems involving diffusion  $[19, 25]$  $[19, 25]$  $[19, 25]$ , among others. These equations also find utility in a range of scientific and engineering disciplines  $[20, 23, 24]$  $[20, 23, 24]$  $[20, 23, 24]$  $[20, 23, 24]$  $[20, 23, 24]$ . Furthermore, in  $[21, 22]$  $[21, 22]$  $[21, 22]$  author discusses the Hopf bifurcation and chaos control in a fractional-order modified hybrid optimal system and fractionalorder hyperchaotic system. Recent research delves into topics like approximating solutions for non-linear fractional-order differential population models [\[27,](#page-14-6) [28\]](#page-14-7). We refer the reader [\[26\]](#page-14-8) while others investigate the qualitative dynamics of nonlinear interactions within biological systems  $[29, 30]$  $[29, 30]$  $[29, 30]$ . For more details on the bifurcation we refer the reader to [\[32,](#page-14-11) [33\]](#page-14-12).

This study investigates the stability analysis of a discrete-time predator-prey model using fractional-order discretization, clarifying the complexities of ecological dynamics through a lens that transcends traditional boundaries. By utilizing the power of fractional calculus, the study aims to shed more light on the intricate dynamics of predator-prey interactions, revealing new perspectives and enriching the understanding of ecological systems. Through rigorous analysis and simulation, the study seeks to clarify the underlying mechanisms governing system stability and resilience, paving the way for more informed conservation and management strategies in the face of environmental change.

In this paper, we study the following Lotka-Volterra systems

<span id="page-1-0"></span>
$$
\begin{cases}\n x_{n+1} = x_n e^{(r(1-x_n) - by_n)} \frac{h^{\alpha}}{\alpha} \\
 y_{n+1} = y_n e^{(bx_n - d)} \frac{h^{\alpha}}{\alpha}\n\end{cases}
$$
\n(2.3)

where  $r, b, d, h > 0$  and  $0 < \alpha \leq 1$  The Jacobian of the system  $(2.3)$  is  $J(E^*) =$ 

$$
\begin{pmatrix}\n e^{-\frac{h^{\alpha}(r(-1+x_n)+by_n)}{\alpha}} \left( \frac{-h^{\alpha}rx_n+\alpha}{\alpha} \right) & -e^{-\frac{h^{\alpha}(r(-1+x_n)+by_n)}{\alpha}} \left( \frac{bh^{\alpha}x_n}{\alpha} \right) \\
 \frac{h^{\alpha}(-d+bx_n)}{c} \left( \frac{bh^{\alpha}y_n}{\alpha} \right) & e^{-\frac{h^{\alpha}(-d+bx_n)}{\alpha}}\n\end{pmatrix}
$$
\n(2.4)

Section Existance and Stability of fixed points

**Lemma 2.1.** The system [\(2.3\)](#page-1-0) has the following positive fixed points if  $0 < \alpha \leq 1$  and  $b > d$ 

$$
E^* = \left(\frac{d}{b}, \frac{(b-d)r}{b^2}\right)
$$

The Jacobian of the system  $(2.3)$  at any point  $\left(\frac{d}{dx}\right)^{2}$  $\frac{d}{b}$ ,  $\frac{(b-d)r}{b^2}$  $b^2$ can be evaluated as

<span id="page-2-0"></span>
$$
\begin{pmatrix}\n1 - \frac{drh^{\alpha}}{b\alpha} & -\frac{dh^{\alpha}}{\alpha} \\
\frac{h^{\alpha}(b-d)r}{b\alpha} & 1\n\end{pmatrix}
$$
\n(2.5)

Next, to discuss the stability of the fixed point of  $(2.3)$ , we follow Lemma 1 from  $[37, 24]$  $[37, 24]$  $[37, 24]$ , therefore the characteristic equation of the Jacobian matrix  $(2.5)$  is the following:

$$
\lambda^2 + p\lambda + q = 0 \tag{2.6}
$$

$$
p = -2 + \frac{drh^{\alpha}}{b\alpha}
$$
  

$$
q = \frac{-drh^{\alpha}(dh^{\alpha} + \alpha) + b(h^{2\alpha}dr + \alpha^{2})}{b\alpha^{2}}
$$

**Theorem 2.2.** The following conditions hold about the stability of the positive fixed point  $E^*$ 

1. The positive fixed point  $E^*$  is sink if  $d < b$  and  $bh^{\alpha} < h^{\alpha}d + \alpha$  and one of the following condition holds

\n- $$
h^{\alpha} \geq \frac{2\alpha}{b-d}
$$
\n- $r < \frac{4b\alpha^2}{cd(-bh^{\alpha} + h^{\alpha}d + 2\alpha)}$
\n

- 2. The positive fixed point  $E^*$  is Saddle if the following condition holds
	- $d < b$ ,  $bh^{\alpha} < h^{\alpha}d + 2\alpha$  and  $b(h^{2\alpha}dr + 4\alpha^2) < h^{\alpha}dr(h^{\alpha}d + 2\alpha)$
- 3. The positive fixed point  $E^*$  is Source if  $d < b$ ,  $h^{\alpha}d + \alpha < bh^{\alpha}$  and one of the following condition are hold

\n- $$
h^{\alpha} \geq \frac{2\alpha}{b-d}
$$
\n- $r < \frac{4b\alpha^2}{h^{\alpha}d(-bh^{\alpha} + dh^{\alpha} + 2\alpha)}$
\n

4.  $E^*$  is non-hyperbolic and observed a flip bifurcation if the following conditions are hold

\n- $$
h^{\alpha} > \frac{2\alpha}{r}
$$
 and  $b = \frac{h^{\alpha} dr (h^{\alpha} d + 2\alpha)}{dh^{2\alpha} r + 4\alpha^2}$
\n- $\frac{h^{\alpha} dr}{b \alpha} \neq 2, 4$
\n

5. The positive fixed points  $E^*$  are complex if the following conditions are hold

\n- $$
0 < \frac{h^{\alpha} dr}{b \alpha} < 4
$$
\n- $b = d + \frac{\alpha}{h^{\alpha}}$
\n

### 3 Neimark Sacker Bifurcation and Flip Bifurcation

Now, we investigate the Neimark-Sacker and Flip bifurcation at the positive fixed point  $E^*$ Let us define a set

 $A_{NS} = \left\{ (\alpha, b, h, d, r) : \alpha, b, h, d, r > 0, 0 < \alpha \leq 1 \text{ and } b = d + \frac{\alpha}{4\sigma^2} \right\}$  $\frac{\alpha}{h^{\alpha}}$  and  $0 < \frac{h^{\alpha} dr}{b \alpha}$  $\frac{\alpha}{b\alpha} < 4$ Let b be a bifurcation parameter. The Neimark-Sacker bifurcations vary in a small neighborhood of  $A_{NS}$ . Choose  $\epsilon$  as a small perturbation parameter, where ( $|\epsilon| \ll 1$ ), then the perturbation of system [\(2.3\)](#page-1-0) is as follows:

$$
\begin{cases}\nx_{n+1} = x_n e^{(r(1-x_n) - (b+\epsilon)y_n)} \frac{h^\alpha}{\alpha} \\
y_{n+1} = y_n e^{(bx_n - d)} \frac{h^\alpha}{\alpha}\n\end{cases}
$$
\n(3.1)

where  $r, b, d, h > 0$  and  $0 < \alpha \leq 1$ 

Let us define a translation  $u_n = x_n - \frac{d}{dx}$  $\frac{d}{b}$  and  $v_n = y_n - \frac{(b-d)r}{b^2}$  $\frac{a}{b^2}$  that translate the positive equilibrium point of the system [\(2.3\)](#page-1-0) to the origin and we have the following

<span id="page-3-1"></span>
$$
\begin{cases}\nu_{n+1} = C_1 u_n + C_2 v_n + C_3 u_n^2 + C_4 v_n^2 + C_5 u_n v_n + C_6 u_n^2 v_n + C_7 u_n v_n^2 + C_8 u_n^3 + C_9 v_n^3 + O((|u_n| + |v_n|)^4) \\
v_{n+1} = D_1 u_n + D_2 v_n + D_3 u_n^2 + D_5 u_n v_n + D_6 u_n^2 v_n + D_8 u_n^3 + O((|u_n| + |v_n|)^4)\n\end{cases} (3.2)
$$

where 
$$
C_i
$$
's and  $D_i$ 's are given below  
\n
$$
C_1 = \frac{-h^{2\alpha}dr + d\alpha h^{\alpha} + \alpha^2 + \alpha h^{\alpha}\epsilon}{\alpha(dh^{\alpha} + \alpha + \epsilon h^{\alpha})}, C_2 = -\frac{h^{\alpha}d}{\alpha}, C_3 = \frac{rh^{\alpha}(h^{2\alpha}dr - 2d\alpha h^{\alpha} - 2\alpha^2 - 2\alpha\epsilon h^{\alpha})}{2\alpha^2(dh^{\alpha} + \alpha + \epsilon h^{\alpha})},
$$
\n
$$
C_4 = \frac{dh^{\alpha}(dh^{\alpha} + \alpha + \epsilon h^{\alpha})}{2\alpha^2}, C_5 = \frac{-h^{2\alpha}dr + d\alpha h^{\alpha} + \alpha^2 + \alpha\epsilon h^{\alpha}}{\alpha^2},
$$
\n
$$
C_6 = -\frac{rh^{\alpha}(h^{2\alpha}dr - 2d\alpha h^{\alpha} - 2\alpha^2 - 2\alpha\epsilon h^{\alpha})}{2\alpha^3}, C_7 = \frac{(dh^{\alpha} + \alpha + \epsilon h^{\alpha})\left(-h^{2\alpha}dr + d\alpha h^{\alpha} + \alpha^2 + \alpha\epsilon h^{\alpha}\right)}{2\alpha^3},
$$
\n
$$
C_8 = -\frac{(h^{2\alpha}r^2\left(h^{2\alpha}dr - 3d\alpha h^{\alpha} - 3\alpha^2 - 3\alpha\epsilon h^{\alpha}\right))}{6\alpha^3(dh^{\alpha} + \alpha + \epsilon h^{\alpha})}, C_9 = -\frac{(dh^{\alpha}(dh^{\alpha} + \alpha + h^{\alpha}\epsilon)^2)}{6\alpha^3},
$$
\n
$$
D_1 = \frac{h^{\alpha}r(\alpha + h^{\alpha}\epsilon)}{\alpha(h^{\alpha}d + \alpha + h^{\alpha}\epsilon)}, D_2 = 1, D_3 = \frac{h^{\alpha}r(\alpha + h^{\alpha}\epsilon)}{2\alpha^2}, D_5 = \frac{(h^{\alpha}d + \alpha + h^{\alpha}\epsilon)}{\alpha},
$$
\n
$$
D_6 = \frac{(h^{\alpha}d + \alpha + h^{\alpha}\epsilon)^2}{2\alpha^2}, D_8 = \frac{h^{\alpha}r(\alpha + h^{\alpha}\epsilon)(h^{\alpha}d + \alpha + h^{\alpha}\epsilon)}{6\alpha^3},
$$
\nLet,

<span id="page-3-0"></span>
$$
\lambda^2 - p(\epsilon)\lambda + q(\lambda) = 0 \tag{3.3}
$$

be the characteristic equation of the above system, where  $p(\epsilon)$  and  $q(\epsilon)$  are given by the following  $h^{2\alpha}dr - 2\alpha^2 - 2ch^{\alpha}\alpha(d+\epsilon)$ 

$$
p(\epsilon) = -\frac{\alpha}{\alpha(\alpha + h^{\alpha}(d + \epsilon))}
$$
  
\n
$$
q(\epsilon) = \frac{\alpha^3 + h^{3\alpha} dr \epsilon + h^{\alpha} \alpha^2 (d + \epsilon)}{\alpha^2 (\alpha + h^{\alpha}(d + \epsilon))}
$$
  
\nsince  $(\alpha, b, c, d, r, h) \in A_{NS}$ , then  $|\lambda_1|$ 

since  $(\alpha, b, c, d, r, h) \in A_{NS}$ , then  $|\lambda_1|=1$  and  $|\lambda_2|=1$ , where  $\lambda_1$  and  $\lambda_2$  are solutions of  $(3.3)$  and are given by

$$
\lambda_{1,2} = \frac{p(\epsilon)}{2} \pm i \frac{\sqrt{4q(\epsilon) - p(\epsilon)^2}}{2}
$$

Since  $|\lambda_1| = |\lambda_2| = \sqrt{q(\epsilon)}$  and at  $\epsilon = 0$ , we have the following result  $\frac{d|\lambda_1|}{d\epsilon} = \frac{d|\lambda_2|}{d\epsilon} = \frac{h^{3\alpha} dr}{2\alpha^2 (h^{\alpha} d \frac{n}{2\alpha^2(h^{\alpha}d+\alpha)}\neq 0$  and  $p(0) = \frac{-h^{2\alpha}dr + 2h^{\alpha}d\alpha + 2\alpha^2}{(1\alpha)^{1+\alpha}}$  $\frac{\alpha(h^{\alpha}d + \alpha)}{\alpha(h^{\alpha}d + \alpha)} \neq -2, 2$ and moreover

1.  $p(0) \neq 1$  if  $d \neq \frac{\alpha^2}{12\alpha}$  $h^{2\alpha}r - h^{\alpha}\alpha$ 

2. 
$$
p(0) \neq 0
$$
 if  $d \neq \frac{2\alpha^2}{h^{2\alpha}r - 2h^{\alpha}\alpha}$ 

which is equivalent to say that  $\lambda^m, \overline{\lambda}^m \neq 1$ ,  $(m = 1, 2, 3, 4)$  if the above two conditions hold. Let us define a transformation

$$
\begin{pmatrix}\nu_n \\ v_n \end{pmatrix} = \begin{pmatrix}\n-\frac{h^{\alpha}d}{\alpha} & 0 \\
\frac{h^2d}{2h^{\alpha}d\alpha} & -\frac{1}{2}\sqrt{\frac{h^{2\alpha}dr(-h^{2\alpha}dr + 4h^{\alpha}d\alpha + 4\alpha^2)}{\alpha^2(h^{\alpha}d + \alpha)^2}}\n\end{pmatrix} \begin{pmatrix}\ne_n \\ f_n \end{pmatrix}
$$
\nwe get the following canonical form of the system (3.2)

we get the following canonical form of the system [\(3.2\)](#page-3-1)

$$
\begin{pmatrix}\ne_{n+1} \\
f_{n+1}\n\end{pmatrix} = \begin{pmatrix}\n\frac{-h^{2\alpha}dr + 2h^{\alpha}d\alpha + 2\alpha^{2}}{2\alpha(h^{\alpha}d + \alpha)} & \frac{1}{2}\sqrt{\frac{(h^{\alpha})^{2}dr \left(-(h^{\alpha})^{2}dr + 4h^{\alpha}d\alpha + 4\alpha^{2}\right)}{\alpha^{2}(h^{\alpha}d + \alpha)^{2}}}\n\frac{1}{2}\sqrt{\frac{h^{2\alpha}dr \left(-(h^{\alpha})^{2}dr + 4h^{\alpha}d\alpha + 4\alpha^{2}\right)}{\alpha^{2}(h^{\alpha}d + \alpha)^{2}}}\n\frac{-(h^{\alpha})^{2}dr + 2h^{\alpha}d\alpha + 2\alpha^{2}}{2\alpha(h^{\alpha}d + \alpha)}\n\end{pmatrix}
$$
\n
$$
\begin{pmatrix}\ne_{n}\\f_{n}\n\end{pmatrix} + \begin{pmatrix}\ne_{n}\\f(e_{n},f_{n})\\G(e_{n},f_{n})\n\end{pmatrix}
$$

Where  $F(e_n, f_n)$  and  $G(e_n, f_n)$  are given by

$$
\begin{cases}\nF(e_n, f_n) = C_{11}e_n^2 + C_{12}f_n^2 + C_{13}e_nf_n + C_{14}e_n^2f_n + C_{15}e_nf_n^2 + C_{16}e_n^3 + C_{17}f_n^3 + O((|e_n|+|f_n|)^4) \\
G(e_n, f_n) = D_{11}e_n^2 + D_{12}f_n^2 + D_{13}e_nf_n + D_{14}e_n^2f_n + D_{15}e_nf_n^2 + D_{16}e_n^3 + D_{17}f_n^3 + O((|e_n|+|f_n|)^4)\n\end{cases}
$$
\n(3.4)

and  $C'_{ij}s$  and  $D'_{i,j}s$  are given by

$$
C_{11} = -\frac{h^{2\alpha} dr (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2})}{8(\alpha^{3}(h^{\alpha} d + \alpha))},
$$
\n
$$
C_{12} = \frac{h^{2\alpha} dr (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2})}{8\alpha^{3}(h^{\alpha} d + \alpha)},
$$
\n
$$
C_{13} = -\frac{h^{2\alpha} dr - 2h^{\alpha} d\alpha - 2\alpha^{2}}{2(\alpha(h^{\alpha} d + \alpha))} - \frac{\left(\sqrt{-\frac{h^{2\alpha} dr (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2})}{\alpha^{2}(h^{\alpha} d + \alpha)^{2}}}(h^{2\alpha} dr - 2h^{\alpha} d\alpha - 2\alpha^{2})\right)}{4\alpha^{2}},
$$
\n
$$
C_{14} = -\frac{h^{2\alpha} dr (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2})}{8\alpha^{3}(h^{\alpha} d + \alpha)} - \frac{\left(h^{2\alpha} dr (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2})\sqrt{-\frac{h^{2\alpha} dr (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2})}{\alpha^{2}(h^{\alpha} d + \alpha)^{2}}}\right)}{16\alpha^{4}},
$$
\n
$$
C_{15} = \frac{h^{2\alpha} dr (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2}) (h^{2\alpha} dr - 2h^{\alpha} d\alpha - 2\alpha^{2})}{16\alpha^{5}(h^{\alpha} d + \alpha)},
$$
\n
$$
C_{16} = -\frac{h^{4\alpha} d^{2} r^{2} (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 6\alpha^{2})}{48(\alpha^{5}(h^{\alpha} d + \alpha))},
$$
\n
$$
C_{17} = -\frac{h^{4\alpha} d^{2} r^{2} (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2}) \sqrt{-\frac{h^{2\alpha} dr (h^{2\alpha} dr - 4h^{\alpha} d\alpha - 4\alpha^{2})}{\alpha^{2}(
$$

$$
D_{16} = -\frac{\alpha \left( \alpha \left( \alpha \right) \right) \left( \alpha \right) \left( \
$$

To examine the Neimark-Sacker bifurcation, let us consider the first Lyapunov exponent, derived as follows:

$$
a = \left[ -Re\left( \frac{(1-2\lambda)\overline{\lambda}^2}{1-\lambda} \xi_{20}\xi_{11} - \frac{1}{2}|\xi_{11}|^2 - |\xi_{02}|^2 + Re(\overline{\lambda}\xi_{21}) \right) \right]_{\epsilon=0}
$$

where

$$
\xi_{20} = \frac{1}{8} \left[ (F_{e_ne_n} - F_{f_nf_n} + 2G_{e_nf_n}) + i(G_{e_ne_n} - G_{f_nf_n} - 2F_{e_nf_n}) \right].
$$
\n
$$
\xi_{11} = \frac{1}{4} \left[ (F_{e_ne_n} + F_{f_nf_n}) + i(G_{e_ne_n} + G_{f_nf_n}) \right].
$$
\n
$$
\xi_{02} = \frac{1}{8} \left[ (F_{e_ne_n} - F_{f_nf_n} - 2G_{e_nf_n}) + i(G_{e_ne_n} - G_{f_nf_n} + 2F_{e_nf_n}) \right].
$$
\n
$$
\xi_{21} = \frac{1}{16} \left[ (F_{e_ne_ne_n} + F_{e_nf_nf_n} + G_{e_ne_nf_n} + G_{f_nf_nf_n}) + i(G_{e_ne_ne_n} + G_{e_nf_nf_n} - F_{e_ne_nf_n} - F_{f_nf_nf_n}) \right].
$$

**Theorem 3.1.** Let  $(\alpha, b, h, d, r) \in A_{NS}$  and  $b \neq 0$ , then the system  $(2.3)$  observes a Neimark-Sacker bifurcation at the positive fixed point E<sup>∗</sup> when the bifurcation parameter varies in a small neighbourhood of b. Furthermore, when b is negative, an attracting invariant curve emanates from E<sup>∗</sup> . When b is positive, a repelling invariant curve emanates from  $E^*$ .

Let us define a set

$$
A_{PD} = \left\{ (\alpha, b, d, h) : \alpha, b, d, h > 0, 0 < \alpha \le 1, h^{\alpha} > \frac{2\alpha}{r}, b = \frac{h^{\alpha} dr (h^{\alpha} d + 2\alpha)}{dh^{2\alpha} r + 4\alpha^{2}} \text{ and } \frac{h^{\alpha} dr}{b\alpha} \ne 4 \right\}
$$

The system  $(2.3)$  observes a period-doubling bifurcation within a domain  $A_{PD}$ . Let b be a bifurcation parameter and  $\epsilon^*(|\epsilon^*| \ll 1)$  be a perturbation parameter, then the perturbation of the system [\(2.3\)](#page-1-0) is the following:

<span id="page-6-0"></span>
$$
\begin{cases}\nx_{n+1} = x_n e^{(r(1-x_n) - (b+\epsilon^*)y_n)} \frac{h^{\alpha}}{\alpha} \\
y_{n+1} = y_n e^{((b+\epsilon^*)x_n - d)} \frac{h^{\alpha}}{\alpha}\n\end{cases}
$$
\n(3.5)

Let  $u_n = x_n - \frac{d}{l}$  $\frac{d}{b}$  and  $v_n = y_n - \frac{(b-d)r}{b^2}$  $\frac{u}{b^2}$ , then the positive fixed point  $E^*$  transform the system [\(3.5\)](#page-6-0) into origin and we get the following:

$$
\begin{pmatrix}\nu_{n+1} \\
v_{n+1}\n\end{pmatrix} = \begin{pmatrix}\nc_{11}u_n + c_{12}v_n + c_{13}u_n^2 + c_{14}v_n^2 + c_{15}u_nv_n + c_{16}u_n\epsilon^* + c_{18}u_n^3 + c_{16}u_n\epsilon^* + c_{18}u_n\epsilon^* + c_{18}u_n\epsilon^* + O((|u_n| + |v_n| + \epsilon^*)^4) \\
c_{20}u_n^2v_n + c_{21}u_n^2\epsilon^* + c_{22}u_nv_n\epsilon^* + O((|u_n| + |v_n| + \epsilon^*)^4) \\
d_{11}u_n + d_{12}v_n + d_{13}u_n^2 + d_{15}u_nv_n + d_{18}u_n^3 + d_{20}u_n^2v_n + d_{18}u_n\epsilon^* + O((|u_n| + |v_n| + \epsilon^*)^4)\n\end{pmatrix}
$$
\n(3.6)

where  $c_{ij}$  and  $d_{ij}$  are given by

$$
c_{11} = \frac{-h^{2\alpha}dr + h^{\alpha}d\alpha - 2\alpha^{2}}{\alpha(dh^{\alpha} + 2\alpha)}, \ c_{12} = -\frac{dh^{\alpha}}{\alpha}, \ c_{13} = \frac{d\left(h^{3\alpha}r^{2} - 2h^{2\alpha}r\alpha\right)}{2\alpha^{2}(dh^{\alpha} + 2\alpha)},
$$
  
\n
$$
c_{14} = \frac{h^{3\alpha}d^{2}(drh^{\alpha} + 2r\alpha)}{2\alpha^{2}(h^{2\alpha}dr + 4\alpha^{2})}, \ c_{15} = \frac{h^{2\alpha}\left(h^{2\alpha}d^{2}r^{2} - (h^{\alpha})d^{2}r\alpha + 2dr\alpha^{2}\right)}{\alpha^{2}(h^{2\alpha}dr + 4\alpha^{2})},
$$
  
\n
$$
c_{16} = \frac{\left(h^{4\alpha}d^{2}r^{2} + 8h^{2\alpha}dr\alpha^{2} + 16\alpha^{4}\right)}{drh^{\alpha}\alpha(dh^{\alpha} + 2\alpha)^{2}},
$$
  
\n
$$
c_{18} = -\frac{h^{2\alpha}r^{2}\left(h^{2\alpha}dr - 3dh^{\alpha}\alpha - 2\alpha^{2}\right)}{6\left(\alpha^{3}(dh^{\alpha} + 2\alpha)\right)}, \ c_{20} = -\frac{h^{4\alpha}r^{2}\left(h^{\alpha}d^{2}r - 2d^{2}\alpha\right)}{2\left(\alpha^{3}\left(h^{2\alpha}dr + 4\alpha^{2}\right)\right)}
$$
  
\n
$$
c_{21} = \frac{\left(-h^{4\alpha}d^{2}r^{2} - 8h^{2\alpha}dr\alpha^{2} - 16\alpha^{4}\right)}{2d\alpha^{2}(dh^{\alpha} + 2\alpha)^{2}}, \ c_{22} = -\frac{h^{\alpha}}{\alpha}, \ d_{11} = \frac{2(h^{\alpha}r - 2\alpha)}{h^{\alpha}d + 2\alpha}
$$
  
\n
$$
d_{12} = 1, \ d_{13} = \frac{h^{2\alpha}dr(rr - 2\alpha)}{\alpha(h^{2\alpha}dr + 4\alpha^{2})}, \ d_{15} = \frac{h^{2\alpha}dr(h^{\alpha}d + 2\alpha)}{\alpha(h^{2\alpha}dr + 4\alpha^{2})},
$$
  
\n

Construct an invertible matrix

$$
T = \begin{pmatrix} -\frac{h^{\alpha}d + 2\alpha}{h^{\alpha}r - 2\alpha} & -\frac{h^{\alpha}d}{2\alpha} \\ 1 & 1 \end{pmatrix}
$$

and define a transformation

$$
\left(\begin{array}{c} e_n \\ f_n \end{array}\right) = T \left(\begin{array}{c} u_n \\ v_n \end{array}\right)
$$

we get the following

<span id="page-7-0"></span>
$$
\begin{pmatrix} e_{n+1} \\ f_{n+1} \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & \frac{-h^{2\alpha}dr + 3h^{\alpha}d\alpha + 2\alpha^2}{\alpha(h^{\alpha}d + 2\alpha)} \end{pmatrix} \begin{pmatrix} e_n \\ f_n \end{pmatrix} + \begin{pmatrix} F(e_n, f_n, \epsilon^*) \\ G(e_n, f_n, \epsilon^*) \end{pmatrix}
$$
(3.7)

where  $F(e_n, f_n, \epsilon^*)$  and  $G(e_n, f_n, \epsilon^*)$  are defined by  $F(e_n, f_n, \epsilon^*) = c_1 e_n^2 + c_2 e_n f_n + c_3 f_n^2 + c_4 \epsilon^* + c_5 e_n \epsilon^* + c_6 e_n^2 \epsilon^* + c_7 f_n \epsilon^* + c_8 e_n f_n \epsilon^* + c_9 f_n^2 \epsilon^* \quad + O((|e_n| + |f_n| + |\epsilon^*|)^4)$  $G(e_n, f_n, \epsilon^*) = d_1 e_n^2 + d_2 e_n f_n + d_3 f_n^2 + d_4 \epsilon^* + d_5 e_n \epsilon^* + d_6 e_n^2 \epsilon^* + d_7 f_n \epsilon^* + d_8 e_n f_n \epsilon^* + b_9 f_n^2 \epsilon^*$ +  $O((|e_n|+|f_n|+|\epsilon^*|)^4)$ where  $c_i$  and  $d_i$  are given by

$$
\begin{array}{rcl} c_2&=&\frac{h^{4\alpha}d^3r+4h^{3\alpha}d^2r\alpha+4h^{2\alpha}dr\alpha^2}{2\alpha^2(h^{2\alpha}dr+4\alpha^2)},\\ c_3&=&\frac{h^{5\alpha}d^3r(4+r)(h^{\alpha}r-2\alpha)}{4\alpha^3(dh^{\alpha}r+2\alpha)(h^{\alpha}r+4\alpha\alpha^2)},\quad c_4=\frac{(rh^{\alpha}-2\alpha)\left(h^{4\alpha}d^2r^2+8h^{2\alpha}dr\alpha^2+16\alpha^4\right)}{2r\alpha^2(h^{\alpha}d+2\alpha)\left(-h^{2\alpha}dr+4h^{\alpha}d\alpha+4\alpha^2\right)},\\ c_5&=&\frac{(h^{\alpha}d^2r^2+8h^{2\alpha}dr\alpha^2+16\alpha^4)}{h^{\alpha}dr\alpha(-h^{2\alpha}dr+4\alpha^2)},\quad c_6&=&\frac{(h^{\alpha}d-2\alpha)(h^{\alpha}d+2\alpha)^2(h^{2\alpha}dr+4\alpha^2)}{2r\alpha^2(h^{\alpha}d+r+2\alpha^2)}\\ c_7&=&\frac{(h^{\alpha}r-2\alpha)\left(h^{4\alpha}d^2r^2+8h^{2\alpha}dr\alpha^2+16\alpha^4\right)}{2r\alpha^2(h^{\alpha}d+2\alpha)\left(-\left(h^{\alpha})^2dr+4h^{\alpha}d\alpha+4\alpha^2\right)}\\ &&c_8&=&-\frac{(h^{\alpha}r-2\alpha)(h^{\alpha}d^3r+4h^{\alpha}d\alpha+4\alpha^2)}{2\left(\alpha^3(h^{\alpha}d+2\alpha)\left(-h^{2\alpha}dr+4h^{\alpha}d\alpha^2+4h^{3\alpha}dr\alpha^3-16h^{2\alpha}d\alpha^4-16h^{\alpha}\alpha^5\right)}\\ &&c_9&=&\frac{(h^{\alpha}r-2\alpha)(h^{\alpha}d^3r+4h^{\alpha}d\alpha^2r^2\alpha+8h^{\alpha}d^2r\alpha^2-12h^{3\alpha}d^2\alpha^3-4h^{3\alpha}dr\alpha^3-16h^{2\alpha}d\alpha^4-16h^{\alpha}\alpha^5)}{2\left(\alpha^3(h^{\alpha}d+2\alpha)\left(-h^{2\alpha}dr+4h^{\alpha}d
$$

Next we determine the center manifold  $W<sup>c</sup>(0,0,0)$  of the system  $(3.7)$  at the fixed point  $(0,0)$  in a small neighborhood of  $\epsilon^* = 0$ . Let

$$
W^{c}(0,0) = \{(e_n, f_n, \epsilon^*) \in \mathbb{R} : f_n = W(u_n, \epsilon^*) = h_0 \epsilon^* + h_1 e_n^2 + h_2 e_n \epsilon^* + h_3 \epsilon^{*2} + O(|e_n| + \epsilon^*)^3\}
$$
(3.8)

$$
f_{n+1} = W(e_{n+1}, \epsilon^*)
$$
  
\n
$$
f_{n+1} = W(-e_n + F(e_n, f_n, \epsilon^*), \epsilon^*)
$$
  
\n
$$
f_{n+1} = W(-e_n + F(e_n, W(e_n, \epsilon^*), \epsilon^*) , \epsilon^*)
$$
  
\n
$$
f_{n+1} = W(-e_n + F(e_n, h_0 \epsilon^* + h_1 e_n^2 + h_2 e_n \epsilon^* + h_3 \epsilon^* + O(|e_n| + |\epsilon^*|)^3, \epsilon^*), \epsilon^*)
$$
  
\n
$$
f_{n+1} = W(-e_n + c_1 e_n^2 + c_2 e_n (h_0 \epsilon^*)) + c_3 h_0^2 \epsilon^* + c_4 \epsilon^* + c_5 e_n \epsilon^* + C_7 h_0 \epsilon^* O(|e_n| + |\epsilon^*|)^3, \epsilon^*
$$
  
\n
$$
f_{n+1} = W(-e_n + c_4 \epsilon^* + c_1 \epsilon_n^2 + (c_5 + c_2 h_0) \epsilon^* e_n + h_0 (c_3 h_0 + c_7) \epsilon^* + O(|e_n|, \epsilon^*)^3), \epsilon^*
$$
  
\n
$$
f_{n+1} = h_0 \epsilon^* + h_1 (e_n^2 + c_4^2 + \epsilon^* - 2c_4 e_n \epsilon^*) + h_2 (-e_n \epsilon^* + c_4 \epsilon^*) + h_3 \epsilon^* + O(|e_n|, \epsilon^*)^3
$$
  
\n
$$
f_{n+1} = h_0 \epsilon^* + h_1 e_n^2 + (h_1 c_4^2 + h_3 + h_2 c_4) \epsilon^* + (2h_1, c_4 + h_2) e_n \epsilon^* + O(|e_n| + |\epsilon^*|)^3
$$

Also

$$
f_{n+1} = \alpha_1 f_n + G(e_n, f_n, \epsilon^*), \text{ where } \alpha_1 = \frac{-h^{2\alpha} dq + 3h^{\alpha} d\alpha + 2\alpha^2}{\alpha(h^{\alpha} d + 2\alpha)}
$$
  
\n
$$
f_{n+1} = \alpha_1 f_n + G(e_n, w(e_n \epsilon^*), \epsilon^*)
$$
  
\n
$$
f_{n+1} = \alpha_1 W(e_n, \epsilon^*) + G(e_n, h_0 \epsilon^* + h_1 e_n^2 + h_2 e_n \epsilon^* + h_3 \epsilon^* + O(|e_n| + \epsilon^*)^3, \epsilon^*)
$$
  
\n
$$
f_{n+1} = \alpha_1 (h_0 \epsilon^* + h_1 e_n^2 + h_2 e_n \epsilon^* + h_3 \epsilon^*) + d_1 e_n^2 + d_2 e_n (h_0 \epsilon^*) + d_3 (h_0^2 \epsilon^*) + d_4 \epsilon^* + d_5 e_n \epsilon^* + O(3) + d_7 h_0 \epsilon^*
$$
  
\n
$$
f_{n+1} = (d_4 + \alpha_1 h_2 + d_2 h_0) e_n \epsilon^* + (d_7 h_0 + h_3 \alpha_1 + d_3 h_0^2) \epsilon^* + O(3).
$$

compare coefficients we get the following

$$
d_4 + \alpha_1 h_0 = h_0 \to h_0 = \frac{d_4}{1 - \alpha_1},
$$
  
\n
$$
h_1 = \alpha_1 h_1 + d_1 \to h_1 = \frac{d_1}{1 - \alpha_2},
$$
  
\n
$$
-2h_1 c_4 - h_2 = \alpha_1 h_2 + d_2 h_0 + d_5 \to -2h_1 c_4 - d_2 h_0 - d_5 = \alpha_1 h_2 + h_2 \to h_2 = \frac{-2h_1 c_4 - d_2 h_0 - d_5}{1 + \alpha_1}
$$
  
\n
$$
h_3 = h_1 c_4^2 + h_2 c_4 = d_7 h_0 + h_3 \alpha_1 + d_3 h_0^2 \to h_3 = \frac{d_3 h_0^2 + d_7 h_0 - h_1 c_4^2 - h_2 c_4}{1 - \alpha_1}
$$
  
\nThus the center manifold can be approximated as

Thus the center manifold can be approximated as

$$
f_n = h_0 \epsilon^* + h_1 e_n^2 + h_2 e_n \epsilon^* + h_3 \epsilon^{*^2} + O(|e_n| + \epsilon^*)^3
$$

where  $h_0$ ,  $h_1$ ,  $h_2$  and  $h_3$  are given above. Substitute this in  $e_{n+1} = -e_n + F(e_n, f_n, \epsilon^*),$ we get the following equation that will describe the dynamics of the given system

$$
e_{n+1} = -e_n + c_1e_n^2 + c_2e_n(h_0e^* + h_1e_n^2 + h_2e_ne^* + h_3e^{*2}) + c_3(h_0e^{*2} + 2h_0e^*e_n^2) + c_4e^* + c_5e_ne^* + c_6e_n^2e^* + c_8e_ne^*(h_0e^*) + c_9e^*(h_0^2e^{*2}) + O(|e_n| + |e^*|)^4
$$
  
\n
$$
e_{n+1} = -e_n + c_1e_n^2 + c_4e^* + c_5e_ne^* + c_6e_n^2e^* + c_2h_0e_ne^* + c_3h_0e_ne^{*2} + c_3h_0^2e^{*2} + c_9h_0^2e^{*3} + c_7h_1e_n^2e^*
$$
  
\n
$$
+ 2c_3h_0h_1e_n^2e^* + c_2h_2e_n^2e^* + ...O(4)
$$

Let us define the right-hand side of the above equation by  $\mathbb{R}^n \Phi$ , then the following are holds about the flip bifurcation at (0,0)

$$
\beta_1 = \left( \frac{\partial^2 \Phi}{\partial e_n \partial \epsilon^*} + \frac{1}{2} \frac{\partial \Phi}{\partial \epsilon^*} \frac{\partial^2 \Phi}{\partial e_n^2} \right) \neq 0
$$
  

$$
\beta_2 = \left( \frac{1}{6} \frac{\partial^3 \Phi}{\partial e_n^3} + \left( \frac{1}{2} \frac{\partial^2 \Phi}{\partial e_n^2} \right)^2 \right) \neq 0
$$

# 4 Chaos Control

In this Section, we apply the hybrid control strategy to control the chaos generated by the period-doubling and Neimark-Sacker bifurcation in the system [\(2.3\)](#page-1-0).

Consider the following controlled system:

<span id="page-9-0"></span>
$$
\begin{cases}\n x_{n+1} = x_n e^{(r(1-x_n) - by_n)} \frac{h^{\alpha}}{\alpha} + (1-\beta)x_n \\
 y_{n+1} = y_n e^{(bx_n - d)} \frac{h^{\alpha}}{\alpha} + (1-\beta)y_n\n\end{cases}
$$
\n(4.1)

where  $0 < \beta < 1$  is the control parameter for the hybrid control method. It is interesting to note that the fixed points for  $(2.3)$  and  $(4.1)$  are the same. The Jacobian of the system  $(4.1)$  at the positive fixed point  $E^*$  is given by:

<span id="page-9-1"></span>
$$
J(E^*) = \begin{bmatrix} 2 - \frac{dh^\alpha r}{b\alpha} - \beta & -\frac{dh^\alpha}{\alpha} \\ \frac{(b-d)h^\alpha r}{b\alpha} & 2 - \beta \end{bmatrix}
$$
(4.2)

The trace and determinant of [\(4.2\)](#page-9-1) are given below:

$$
T = \text{trace}(J) = 4 - \frac{dh^{\alpha}r}{b\alpha} - 2\beta
$$
  

$$
D = \det(J) = \frac{(b-d)dh^{2\alpha}r + \alpha (dh^{\alpha}r + b\alpha(-2+\beta))(-2+\beta)}{b\alpha^{2}},
$$

The Jury condition states that the fixed point  $E^*$  of the controlled system  $(4.1)$  is stable if the following conditions are met:

$$
\begin{cases}\nT + D + 1 > 0 \\
-T + D + 1 > 0 \\
D - 1 < 0\n\end{cases} \tag{4.3}
$$

## 5 Numerical Examples

**Example 5.1.** For fixed values of parameters  $\alpha = 0.7$ ,  $h = 0.3$ ,  $d = 2$ ,  $r = 1$  and  $b = 3.62597$ , the system [\(2.3\)](#page-1-0) has the fixed point (0.551576, 0.12367). The Jacobian matrix of the system [\(2.3\)](#page-1-0) has the following eigen values  $\lambda_1 = 0.830386 + 0.557189I$  and  $\lambda = 0.830386 - 0.557189I$ . Since  $|\lambda_1| = 1$  and  $|\lambda_2| = 1$ , so the system [\(2.3\)](#page-1-0) observe a Neimark-Sacker bifurcation. For the bifurcation parameter  $b \in [3, 4]$  and  $x_0 = 0.5$ ,  $y_0 = 0.1$ , the figure [\(1\)](#page-9-2) describes the bifurcation of the system [\(2.3\)](#page-1-0) and the figure [\(5\)](#page-12-2) describes the bifurcation of the controlled system  $(4.1)$ . Figure  $(3)$  represents the graph of maximum Lyapunov exponent, and figure  $(4)$  is the phase portrait for different values of b for the system [\(2.3\)](#page-1-0).



<span id="page-9-2"></span>Figure 1: Bifurcation Diagrams for the System[\(2.3\)](#page-1-0)



Figure 2: Bifurcation Diagrams for the Controlled System [\(4.1\)](#page-9-0)



<span id="page-10-0"></span>Figure 3: Graph of the Maximum Lyapunov Exponent for the System [\(2.3\)](#page-1-0)



Figure 4: Phase Portrait for the System [\(2.3\)](#page-1-0)

<span id="page-11-0"></span>Example 5.2. For  $\alpha = 0.5$ ,  $h = 1$ ,  $r = 2$ ,  $d = 2.5$ ,  $d = 2.91667$ ,  $x_0 = 0.8$  and  $y_0 = 0.09$  then the system [\(2.3\)](#page-1-0) has a positive fixed point (0.857143, 0.0979592). The Jacobian of the system [\(2.3\)](#page-1-0) at this fixed point has the eigenvalues  $\lambda_1 = -1$  and  $\lambda_2 = -0.428571$ . Hence, system [\(2.3\)](#page-1-0) observes a period doubling bifurcation.



Figure 5: Bifurcation Diagrams for the System [\(2.3\)](#page-1-0)

# <span id="page-12-2"></span>6 Conclusion

In [\[34\]](#page-14-14), the author uses the forward Euler scheme to discretize the system [\(2.1\)](#page-0-0) and then investigates the complex dynamics of the discrete-time predator-prey system  $(2.1)$  in the closed first quadrant  $R_+^2$ . He observed that the system undergoes the flip bifurcation and Hopf bifurcation in the interior of the  $R_+^2$  using the centre manifold theorem and bifurcation theory. He also presented the numerical simulation not only to illustrate the theoretical analysis but also to exhibit the complex dynamic behaviours, such as the period-5,6,9,10,14,18,20,25 orbits, a cascade of period-doubling bifurcation in period-2,4,8, quasi-periodic orbits and the chaotic sets. However, these results reveal much better results than those of the continuous system. In our study, we use the piecewise constant argument technique, conformal fractional case, to discretize the system  $(2.1)$  and then investigate its complex dynamics. We also presented numerical examples to illustrate our theoretical work. We observe that it gives more accurate results than those in [\[34\]](#page-14-14). Graphical illustrations are also presented to support numerical work.

Acknowledgement This research was funded in whole by the National Science Centre, Poland, grant number 2023/51/B/ST8/01062. For the purpose of Open Access, the authors have applied a CC-BY public copyright license to any Author Accepted Manuscript (AAM) version arising from this submission.

Declaration of Conflicts of Interest The authors affirm they have no conflicts of interest to disclose concerning the current study.

Author Contribution Statement All authors contributed equally. Also, all authors reviewed the manuscript.

Availability of Data and Materials The datasets analyzed during the current study are available from the corresponding author upon reasonable request.

# References

- <span id="page-12-0"></span>[1] Kilbas, A. A., Srivastava, H. M., & Trujillo, J. J. (2006). Theory and applications of fractional differential equations (Vol. 204). Elsevier.
- <span id="page-12-1"></span>[2] Atangana, A., & G´omez-Aguilar, J. F. (2022). Application of Fractional calculus Modelling in Dynamics and Chaos. Solitons & Fractals, CRC Press, Taylor and Francis Group LLC.https://doi.org/10.1201/9781003006244
- [3] El-Sayed, A. M. A., & Arafa, A. A. M. (2013). Fractional-order prey-predator model with modified Leslie-Gower and Holling-type II schemes. Journal of Theoretical Biology, 318, 20-28.
- <span id="page-13-0"></span>[4] Baleanu, D., Magin, R. L., & Bhalekar, S. (2018). Fractional differential equations and models: Applications and numerical methods. Springer.
- [5] Pathak, S. D., & Das, S. (2016). Mathematical modeling and analysis of fractional order prey-predator system. Applied Mathematics and Computation, 276, 258-267.
- [6] Delavari, H., & Zarezadeh, M. (2019). Chaos and Fractional-Order Systems: Analysis and Design. CRC Press.
- [7] Li, C., & Guo, L. (2017). Fractional-order predator-prey system with modified Leslie-Gower and Holling-type II schemes. Chaos, Solitons & Fractals, 104, 667-673.
- [8] Podlubny, I. (1999). Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications. Academic Press.
- [9] Almeida, R., Pooseh, S., & Torres., D.F.M (2015). Computational methods in the Fractional Calculus of Variations, Imperial College Press. https://doi.org/10.1142/p991
- [10] Diethelm, K., & Ford, N. J. (2002). Analysis of fractional differential equations. Journal of Mathematical Analysis and Applications, 265(2), 229-248.
- [11] Caputo, M. (1967). Linear models of dissipation whose Q is almost frequency independent-II. Geophysical Journal International, 13(5), 529-539.
- [12] May, R. M. (1974). Stability and complexity in model ecosystems. Princeton University Press.
- [13] Grimm, V., & Railsback, S. F. (2005). Individual-based modeling and ecology. Princeton University Press.
- [14] Das, S., & Gupta, P.K. (2011). A mathematical model on fractional Lotka-Volterra equations. Journal of Theoretical Biology, 277, 1-6.
- <span id="page-13-1"></span>[15] Rihan, F.A., Lakshmanan, S., Hashish, A.H., Rakkiyappan, R., & Ahmed, E. (2015). Fractional-order delayed predator-prey systems with Holling type-II functional response. Non-linear Dynamics, 80, 777- 789.
- <span id="page-13-2"></span>[16] Chen, S., Wei, J., & Zhang, X. (2019). Bifurcation analysis for a delayed diffusive logistic population model in the advective heterogeneous environment. Journal of Dynamics and Differential Equations. Advance online publication. https://doi.org/10.1007/s10884-019-09739-0
- <span id="page-13-3"></span>[17] Abdeljawad, T., & Alzabut, J. (2018). On Riemann-Liouville fractional q-difference equations and their application to retarded logistic type model. Mathematical Methods in the Applied Sciences, 41(18), 8953-8962.
- <span id="page-13-4"></span>[18] Cao, X., Datta, A., Al Basir, F., & Roy, P.K. (2016). Fractional-order model of the disease psoriasis: A control based mathematical approach. Journal of Systems Science and Complexity, 29, 1-20.
- <span id="page-13-5"></span>[19] Li, H., Zhang, L., Hu, C., Jiang, Y., & Teng, Z. (2017). Dynamic analysis of a fractional-order singlespecies model with diffusion. Non-linear Analysis: Modelling and Control, 22(3), 303-316.
- <span id="page-13-6"></span>[20] Gutierrez-Vega, J.C. (2007). Fractionalization of optical beams: I. Planar analysis. Optics Letters, 32(11), 1521-1523.
- <span id="page-13-7"></span>[21] Abdelouahab, M.S., Hamri, N.E., & Wang, J. (2011). Hopf bifurcation and chaos in fractional-order modified hybrid optical system. Non-linear Dynamics, 69, 275-284.
- <span id="page-13-8"></span>[22] Li, X., & Wu, R. (2014). Hopf bifurcation analysis of a new commensurate fractional-order hyperchaotic system. Non-linear Dynamics, 78, 279-288.
- <span id="page-14-4"></span>[23] Tavazoei, M.S., Haeri, M., Attari, M., Bolouki, S., & Siami, M. (2009). More details on analysis of fractional-order Van der Pol oscillator. Journal of Vibration and Control, 15(6), 803-819.
- <span id="page-14-5"></span>[24] Grigorenko, I., & Grigorenko, E. (2003). Chaotic dynamics of the fractional Lorenz system. Physical Review Letters, 91, 034-101.
- <span id="page-14-3"></span>[25] Mohammad, S.T., & Mohammad, H. (2007). A necessary condition for double scroll attractor existence in fractional-order systems. Physics Letters A, 367, 102-113.
- <span id="page-14-8"></span>[26] Das, S. (2011). Introduction to Fractional Calculus for Scientists and Engineers. Springer, New York.
- <span id="page-14-6"></span>[27] Das, S., & Gupta, P.K. (2011). A mathematical model on fractional Lotka-Volterra equations. Journal of Theoretical Biology, 277, 1-6.
- <span id="page-14-7"></span>[28] Mondal, S., Bairagi, N., & Lahiri, A. (2017). A fractional calculus approach to Rosenzweig-MacArthur predator-prey model and its solution. Journal of Modern Methods in Numerical Mathematics, 8(1-2), 66-76.
- <span id="page-14-9"></span>[29] Li, H.L., Zhang, L., Hu, C., Jiang, Y.L., & Teng, Z. (2017). Dynamical analysis of a fractional-order predator-prey model incorporating a prey refuge. Journal of Applied Mathematics and Computation, 54(1-2), 435-449.
- <span id="page-14-10"></span>[30] Mondal, S., Bairagi, N., & N'Guerekata, G.M. (2019). Global stability of a Leslie-Gower-type fractional order tritrophic food chain model. Fractional Differential Calculus, 9(1), 149-161.
- <span id="page-14-0"></span>[31] Khalil, R., Al Horani, M., Yousef, A., & Sababheh, M. (2014). A new definition of fractional derivative. Journal of Computational and Applied Mathematics, 264, 65-70.
- <span id="page-14-11"></span>[32] Tarasov, V. E. (2011). Fractional dynamics: applications of fractional calculus to dynamics of particles, fields and media. Springer.
- <span id="page-14-12"></span>[33] Kuznetsov, Y. A. (2004). Elements of Applied Bifurcation Theory. Springer Science & Business Media.
- <span id="page-14-14"></span>[34] Xiaoli Liu, Dongmei Xiao (2007). Complex dynamic behaviors of a discrete time predator prey system. Chaos Solitons & Fractals, 32, 80-94.
- <span id="page-14-1"></span>[35] Rahimi, Z., Sumelka, W. and Yang, XJ. (2017). A new fractional nonlocal model and its application in free vibration of Timoshenko and Euler-Bernoulli beams. Eur. Phys. J. Plus 132, 479.
- <span id="page-14-2"></span>[36] Rahimi, Z., Sumelka, W. and Yang, X.-J. (2017). Linear and non-linear free vibration of nano beams based on a new fractional non-local theory. Engineering Computations, Vol. 34 No. 5, pp. 1754-1770.
- <span id="page-14-13"></span>[37] Qianqian Cui, Qiang Zhang, Zhipeng Qiu, Zengyun Hu, Complex dynamics of a discrete-time predatorprey system with Holling IV functional response, Chaos, Solitons & Fractals, Volume 87, 2016, Pages 158-171