PHASE TRANSITION FOR UNREGULARIZED M-ESTIMATION IN SINGLE INDEX MODEL

PIERRE C. BELLEC AND TAKUYA KORIYAMA

Abstract. This note studies phase transitions for the existence of unregularized M-estimators under proportional asymptotics where the sample size n and feature dimension p grow proportionally with $n/p \to \delta \in (0,\infty)$. We study the existence of M-estimators in single-index models where the response y_i depends on covariates $x_i \sim N(0, I_p)$ through an unknown index $w \in \mathbb{R}^p$ and an unknown link function. An explicit expression is derived for the critical threshold δ_{∞} that determines the phase transition for the existence of the M-estimator, generalizing the results of Candès and Sur [\[5\]](#page-8-0) for binary logistic regression to other single-index linear models.

Furthermore, we investigate the existence of a solution to the nonlinear system of equations governing the asymptotic behavior of the M-estimator when it exists. The existence of solution to this system for $\delta > \delta_{\infty}$ remain largely unproven outside the global null in binary logistic regression. We address this gap with a proof that the system admits a solution if and only if $\delta > \delta_{\infty}$, providing a comprehensive theoretical foundation for proportional asymptotic results that require as a prerequisite the existence of a solution to the system.

1. INTRODUCTION

Let $(x_i, y_i)_{i \in [n]}$ be a sample of i.i.d. observations where $x_i \in \mathbb{R}^p$ and follows the normal distribution $x_i \sim N(0, I_p)$. We consider responses $y_i \in \mathcal{Y}$ where $\mathcal{Y} \subset \mathbb{R}$ and assume a singleindex model of the form

$$
\mathbb{P}(y_i \leq t \mid \boldsymbol{x}_i) = F(t, \boldsymbol{x}_i^{\top} \boldsymbol{w}), \qquad \forall t \in \mathbb{R}
$$

where $F: \mathbb{R} \times \mathbb{R} \to [0, 1]$ is an unknown deterministic function and $w \in \mathbb{R}^p$ is an unknown index with $\|\boldsymbol{w}\|=1$. This includes generalized linear models, such as the Poisson model (in which $\mathcal{Y} = \{0, 1, 2, \dots\}$ or Binary logistic model (in which $\mathcal{Y} = \{-1, 1\}$ for instance).

An unregularized M-estimator is fit to observed data $(y_i, x_i)_{i \in [n]}$ by the minimization problem

(1.1)
$$
\inf_{\boldsymbol{b}\in\mathbb{R}^p}\sum_{i=1}^n\ell_{y_i}(\boldsymbol{x}_i^\top\boldsymbol{b})
$$

P.C. BELLEC: RUTGERS UNIVERSITY. T. KORIYAMA: UNIVERSITY OF CHICAGO. E-mail address: pierre.bellec@rutgers.edu, tkoriyam@uchicago.edu. Date: January 7, 2025.

where $\ell_y(\cdot)$ is convex for every $y \in \mathcal{Y}$. If the infimum is achieved, we say the M-estimator exists and denote it by \boldsymbol{b} , i.e.,

$$
\hat{\boldsymbol{b}} \in \mathop{\mathrm{argmin}}_{\boldsymbol{b} \in \mathbb{R}^p} \sum_{i=1}^n \ell_{y_i}(\boldsymbol{x}_i^{\top} \boldsymbol{b}).
$$

In this paper we focus on a high dimensional regime where the sample size and feature grow proportionally as

$$
\frac{n}{p} \to \delta \in (1, +\infty)
$$

Here the constant δ quantifies sample size per dimensions. In this proportional asymptotic regime, under logistic regression model with binary response $\mathcal{Y} = \{-1, 1\}$ and with $\ell_y(t) =$ $log(1 + exp(-yt))$ the logistic loss, the seminal work of [\[5\]](#page-8-0) establishes that the existence of the M-estimator undergoes a sharp phase transition at a critical threshold δ_{∞} :

- if $\delta > \delta_{\infty}$ then the M-estimator exists with high-probability (i.e., the infimum in [\(1.1\)](#page-0-0) is attained), while
- if $\delta < \delta_{\infty}$ then the M-estimator does not exist (i.e., the infimum is not attained) with high-probability.

If $\delta > \delta_{\infty}$, the behavior of the unregularized M-estimator $\hat{\boldsymbol{b}}$, including the limit in probability of $\hat{\mathbf{b}}^{\top}\mathbf{w}$ and the limit in probability of $\|(I_p - \mathbf{w}\mathbf{w}^{\top})\hat{\mathbf{b}}\|_2$, is characterized [\[9\]](#page-8-1) by the nonlinear system with three unknowns $(\gamma, a, \sigma) \in \mathbb{R}_{>0} \times \mathbb{R} \times \mathbb{R}_{>0}$:

(1.2)
\n
$$
\gamma^{-2}\delta^{-1}\sigma^2 = \mathbb{E}[\ell'_Y(\text{prox}[\gamma\ell_Y](aU + \sigma G))^2],
$$
\n
$$
0 = \mathbb{E}[U\ell'_Y(\text{prox}[\gamma\ell_Y](aU + \sigma G))],
$$
\n
$$
\sigma(1 - \delta^{-1}) = \mathbb{E}[G \text{ prox}[\gamma\ell_Y](aU + \sigma G)],
$$

where $G \sim N(0, 1)$ is independent of (U, Y) , and (U, Y) has the same distribution as $(\mathbf{x}_i^{\top} \mathbf{w}, y_i)$. In particular $U \sim N(0, 1)$. The phase transition result of Candès and Sur [\[5\]](#page-8-0) for Gaussian design and binary logistic regression has been extended by [\[11\]](#page-9-0) to elliptic covariate distributions and general binary response models. Han and Ren [\[6\]](#page-8-2) extended the phase transition of the logistic regression model with Gaussian covariates to the constrained minimization setting where the infimum in (1.1) is restricted to a closed convex cone.

Results such as [\[9,](#page-8-1) [8\]](#page-8-3) studying the behavior of the M-estimator on the side of the phase transition where it exists with high-probability assume that the nonlinear system [\(1.2\)](#page-1-0) admits a unique solution. Under the global null in binary logistic regression, i.e., when x_i is independent of y_i , [\[10\]](#page-9-1) establishes that $\delta_{\infty} = 2$ and that the system [\(1.2\)](#page-1-0) admits a unique solution if and only if $\delta > 2$. Beyond the global null, it was observed in [\[9\]](#page-8-1) that the system [\(1.2\)](#page-1-0) can be solved numerically if $\delta > \delta_{\infty}$ (where δ_{∞} is characterized in [\[5\]](#page-8-0)), and that if the solution exists, it is unique [\[9,](#page-8-1) Remark 2 of the supplement]. However to our knowledge there is no proof yet that the system admits a solution for $\delta > \delta_{\infty}$ except under the global null (see [\[9,](#page-8-1) discussion after eq. (16) of the supplement]).

The goal of the present paper is twofold:

- Characterize the critical threshold δ_{∞} for Gaussian covariates beyond binary response models, for instance the Poisson model.
- Prove that the system [\(1.2\)](#page-1-0) admits a unique solution if and only if $\delta > \delta_{\infty}$.

2. Main result

Let us introduce the three examples of interest that our assumptions will cover.

Example 2.1 (Binary logistic regression). Here, labels in $\mathcal{Y} = \{-1, 1\}$ and the loss function

$$
\ell_y(t) = \log(1 + \exp(-yt)).
$$

Example 2.2 (Logistic regression with repeated measurements). Let $q \geq 2$. Here, $\mathcal{Y} =$ $\{0, 1, \ldots, q\}$ and the loss function, corresponding to a binomial regression models with q throws and a sigmoid link function for the probability, is

$$
\ell_y(t) = q \log(1 + \exp(t)) - yt.
$$

If $q = 1$, this is equivalent to binary logistic regression by renaming $\{1,0\}$ to $\{1,-1\}$.

Example 2.3 (Poisson regression). If the labels are in $\mathcal{Y} = \{0, 1, 2, 3, \dots\} = \mathbb{N}$, the nonnegative likelihood of Poisson generalized linear model leads to the loss function

$$
\ell_y(t) = \exp(t) - yt.
$$

The phenomenon of the phase transition for the existence of the M-estimator results from lack of coercivity of some of the loss functions ℓ_{y_i} appearing in the optimization problem [\(1.1\)](#page-0-0). For instance,

- In the binary logistic regression case (Example [2.1\)](#page-2-0), the loss ℓ_{y_i} is not coercive for all y_i : it is increasing for $y_i = -1$ and decreasing for $y_i = 1$.
- For binomial logistic regression with $q \ge 2$ measurements, (Example [2.2\)](#page-2-1), the loss ℓ_{y_i} is coercive if $y_i \in \{1, ..., q-1\}$, increasing for $y_i = 0$ and decreasing for $y_i = q$.
- For Poisson regression, (Example [2.3\)](#page-2-2), the loss ℓ_{y_i} is coercive for $y_i \geq 1$ and increasing for $y_i = 0$.

The values of y_i leading to a coercive, increasing or decreasing loss $\ell_{y_i}(\cdot)$ and the distribution of $(\boldsymbol{x}_i^{\top} \boldsymbol{w}, y_i)$ will determine the critical threshold δ_{∞} . In order to study δ_{∞} , it will be thus useful to introduce the following notation: For a random variable Y valued in \mathcal{Y} , we define the events $\Omega_{\vee}(Y), \Omega_{\nearrow}(Y), \Omega_{\nwarrow}(Y)$ by

(2.1)
$$
\Omega_{\vee}(Y) = \{\ell_Y(\cdot) \text{ is coercive}\},
$$

$$
\Omega_{\nearrow}(Y) = \{\ell_Y \text{ is strictly increasing}\},
$$

$$
\Omega_{\searrow}(Y) = \{\ell_Y \text{ is strictly decreasing}\}.
$$

FIGURE 1. Three examples of loss functions.

We will assume that ℓ_Y is strictly convex in our working assumptions, so if ℓ_Y is not coercive it must be either strictly increasing or strictly decreasing. We first state an assumption that prevents the problem from being trivial.

Assumption 1. The loss ℓ_y is strictly convex for every $y \in \mathcal{Y}$, and the law of Y satisfies

$$
\mathbb{E}\Big[U^2\big(I\{\Omega_\vee(Y)\} + I\{\Omega_\searrow(Y), U > 0\} + I\{\Omega_\nearrow(Y), U < 0\}\big)\Big] > 0,
$$

$$
\mathbb{E}\Big[U^2\big(I\{\Omega_\vee(Y)\} + I\{\Omega_\searrow(Y), U < 0\} + I\{\Omega_\nearrow(Y), U > 0\}\big)\Big] > 0.
$$

This prevents the problem from being trivial in the following sense. Consider $p = 1$, and write $x_i = U_i$ (which is now scalar valued). Assume that the first line in Assumption [1](#page-3-0) is 0. The minimization problem (3.1) becomes

$$
\inf_{a \in \mathbb{R}} \sum_{i=1}^n \ell_{y_i}(aU_i).
$$

Since $\mathbb{P}(U^2 > 0) = 1$ by $U \sim N(0, 1)$, this implies $\mathbb{P}(\Omega_V(Y)) = 0$, so for each of the *n* terms, the loss $\ell_{y_i}(\cdot)$ is not coercive. In fact, each term is increasing in a, because $U_i > 0 \Rightarrow \Omega_{\neq}(y_i)$ and $U_i < 0 \Rightarrow \Omega_{\searrow}(y_i)$ by the first line in Assumption [1,](#page-3-0) so in this case $a \mapsto \sum_{i=1}^n \ell_{y_i}(aU_i)$ is increasing with probability one and the infimum is never attained. Similarly, if the second line in Assumption [1](#page-3-0) is zero, then $a \mapsto \sum_{i=1}^n \ell_{y_i}(aU_i)$ is decreasing and the infimum is never attained.

In conclusion, Assumption [1](#page-3-0) merely assumes that for $p = 1$, the infimum is achieved, i.e., the M-estimator exists. If the M-estimator does not exist for $p = 1$ then it will not exist for $p > 1$ $p > 1$ either, so Assumption 1 is required to avoid this trivial case that the M-estimator does not exist for all $p \geq 1$. The remaining of our working assumptions are given below.

Assumption 2. The loss ℓ_Y satisfies the following:

- (1) For all $y \in \mathcal{Y}$, $\ell_y : \mathbb{R} \to \mathbb{R}$ is C^1 , strictly convex, and not constant.
- (2) $\mathbb{E}[\inf_u \ell_Y(u)] < +\infty$ and $\mathbb{E}[\ell_Y(G)] < +\infty$ where $G \sim N(0, 1)$ independent of Y. $(3) \mathbb{P}(\Omega_{\vee}) < 1.$
- (4) There exists a positive constant b and $\sigma(Y)$ -measurable positive random variable $D(Y)$ satisfying $\mathbb{E}[D(Y)] < +\infty$ and $\mathbb{E}[D^2(Y)\Omega_{\vee}] < +\infty$ such that

$$
\forall u \in \mathbb{R}, \quad \ell_Y(u) \ge -D(Y) + \frac{1}{b} \times \begin{cases} u & \text{under } \Omega_{\nearrow} \\ |u| & \text{under } \Omega_{\vee} \\ -u & \text{under } \Omega_{\searrow} \end{cases}
$$

Beyond Assumption [1,](#page-3-0) Assumption [2](#page-3-1) requires differentiability of the loss (item 1) mild integrability conditions (item 2), requires that ℓ_{y_i} is not always coercive (item 3) and the requirement that in the directions where ℓ_Y diverges to $+\infty$, it does so at least as fast as an affine function with slope $1/b$ and squared integrable intercept $D(Y)$.

We define the critical threshold $\delta_{\infty} \in (0, +\infty]$ by

$$
\frac{1}{\delta_{\infty}} := \inf_{t \in \mathbb{R}} \Big(\mathbb{E} \Big[\Big(G + Ut \Big)^2 I \{ \Omega_{\vee}(Y) \} \Big] + \mathbb{E} \Big[\Big(G + Ut \Big)^2 I \{ \Omega_{\nearrow}(Y) \} \Big] + \mathbb{E} \Big[\Big(G + Ut \Big)^2 I \{ \Omega_{\searrow}(Y) \} \Big] \Big)
$$

where the positive part of any real a is denoted $a_{+} = \max(0, a)$ and the negative part $a_{-} =$ max(-a, 0). When the inf_{t∈R}(\cdots) = 0, δ_{∞} is interpreted as δ_{∞} = + ∞ . Above, the infimum over $t \in \mathbb{R}$ always admits a minimizer $t_* \in \mathbb{R}$ under Assumption [1](#page-3-0) (see Lemma [A.1\)](#page-9-2).

First, we claim that the threshold δ_{∞} characterizes the phase transition whether the Mestimator exists or not with high-probability under the proportional regime $n/p \to \delta$.

Theorem 2.4. As $n, p \to +\infty$ with $n/p \to \delta$, we have

$$
\mathbb{P}\Big(\text{the }M\text{-estimator exists, i.e., the inf }(1.1) \text{ is attained}\Big) \to \begin{cases} 1 & \delta > \delta_{\infty}, \\ 0 & \delta < \delta_{\infty}. \end{cases}
$$

The proof of Theorem [2.4](#page-4-0) is based on a common argument based on conic geometry and Gaussian kinematic formula given by [\[1\]](#page-7-0). This use of the kinematic formula of [\[1\]](#page-7-0) is similar to the argument in [\[5\]](#page-8-0) for the binary logistic regression model.

A more technical question, that requires an investigation beyond the application of the kinematic formula of [\[1\]](#page-7-0), is whether the critical threshold also characterizes the phase transition regarding the existence of solution to the nonlinear system [\(2.2\)](#page-4-1). A formal proof of existence of a unique solution to (2.2) is important, as it is required to leverage the Convex Gaussian Minmax Theorem (CGMT) of $[12]$. For instance, the works $[8, 7]$ $[8, 7]$ $[8, 7]$ which apply the CGMT in generalized linear models to study **b** assume in their theorems that the system (1.2) admits a unique solution.

Theorem 2.5.

6 PHASE TRANSITION FOR UNREGULARIZED M-ESTIMATION IN SINGLE INDEX MODEL

- If $\delta \leq \delta_{\infty}$, the system of nonlinear equations [\(1.2\)](#page-1-0) has no solution.
- If $\delta > \delta_{\infty}$, the system of nonlinear equations [\(1.2\)](#page-1-0) has a unique solution.

To our knowledge, a proof of this relationship between the critical threshold δ_{∞} and the existence of a solution to the system [\(1.2\)](#page-1-0) is new, even in the case of binary logistic regression (the global null case was handled in [\[10\]](#page-9-1), however outisde the global null case this phenomenon was observed numerically in [\[9\]](#page-8-1) without proof—see the discussion after eq. (16) of the supplement of $[9]$).

The tools we use to obtain this fact are based on the existence of a solution to a convex minimization problem in an infinite-dimensional Hilbert space which is the focus of the next section.

3. Infinite dimensional optimization problem

In this section, we define the mathematical objects at the heart of the proof of Theorem [2.5](#page-4-2) and outline the proof strategy. The key is the analysis of an infinite-dimensional convex optimization problem that is in a dual relationship with the nonlinear system.

The notation and setup of this infinite-dimensional convex optimization problem is heavily inspired by $[3]$, which studies the existence of solutions to systems of a similar nature to (1.2) in robust regression. In the robust regression setup of [\[3\]](#page-7-1) where coercive and Lipschitz loss functions $\ell_{y_i}(\cdot)$ are considered, the corresponding system has always a unique solution and the M-estimator always exists: there is no phase transition. A novelty of the present paper is to explain how these tools, in particular the infinite-dimensional optimization below, can be used to predict the phase transition for the existence of a minimizer in (1.1) and the phase transition for the existence of solutions to the system (1.2) .

To begin with, let us consider the almost sure equivalent classes $\mathcal H$ of squared integrable measurable functions of (G, U, Y)

$$
\mathcal{H} = \{v : \mathbb{R}^3 \to \mathbb{R} : \mathbb{E}[v(G, U, Y)^2] < +\infty\},\
$$

where $G \sim N(0, 1)$ and independent of (U, Y) . Here (U, Y) is equal in distribution to $(y_i, \mathbf{x}_i^{\top} \mathbf{w})$. Almost sure equivalence classes of $\mathcal H$ form a Hilbert space equipped with the usual inner product $\langle u, v \rangle := \mathbb{E}[u(G, U, Y)v(G, U, Y)].$ We will sometimes refer to H itself as the Hilbert space, in this case we implicitly identify random variables $v(G, U, Y)$ that are equal almost surely. Now we define two functions $\mathcal G$ and $\mathcal L$ as follows:

$$
\mathcal{G}: \mathcal{H} \to \mathbb{R}, \qquad v \mapsto ||v|| - \mathbb{E}[vG]/\sqrt{1 - \delta^{-1}}
$$

$$
\mathcal{L}: \mathbb{R} \times \mathcal{H} \to \mathbb{R} \cup \{+\infty\}, \qquad (a, v) \mapsto \begin{cases} \mathbb{E}[\ell_Y(aU + v)] & \text{if } \mathbb{E}[|\ell_Y(aU + v)|] < +\infty \\ +\infty & \text{otherwise} \end{cases}
$$

Here, $\mathcal L$ is a proper lower semicontinuous convex function, while $\mathcal G$ is a Lipschitz, finite valued, and convex function (See Lemma [B.1](#page-15-0) and Lemma [B.2\)](#page-15-1). With these functions $(\mathcal{L}, \mathcal{G})$, we claim that the system of nonlinear equations [\(1.2\)](#page-1-0) admits a unique solution if and only if the following infinite-dimensional convex optimization problem over $\mathbb{R} \times \mathcal{H}$

(3.1)
$$
\min_{(a,v)\in\mathbb{R}\times\mathcal{H}} \mathcal{L}(a,v) \quad \text{subject to} \quad \mathcal{G}(v) \le 0
$$

admits a unique minimizer $(a_*, v_*) \in \mathbb{R} \times \mathcal{H}$. We will make this point more precise in the next paragraph.

The key to such an equivalence between the nonlinear system (1.2) and infinite-dimensional optimization problem (3.1) is the existence of the Lagrange multiplier associated with the constraint $\mathcal{G}(v) \leq 0$. By [\[2,](#page-7-2) Proposition 27.31], an element $(a_*, v_*) \in \mathbb{R} \times \mathcal{H}$ solves the constrained optimization problem $\min_{(a,v): \mathcal{G}(v) \leq 0} \mathcal{L}(v)$ if and only if there exists a Lagrange multiplier $\mu_* \geq 0$ such that the KKT condition

$$
(3.2) \qquad -\mu_*\partial \mathcal{G}(v_*) \cap \partial_v \mathcal{L}(a_*, v_*) \neq \emptyset, \quad 0 \in \partial_a \mathcal{L}(a, v), \quad \mu_*\mathcal{G}(v_*) = 0, \quad \text{and} \quad \mathcal{G}(v_*) \leq 0
$$

is satisfied, where $\partial \mathcal{G}, \partial \mathcal{L}$ are subdifferentials of the convex functions \mathcal{G}, \mathcal{L} . Furthermore, we will argue that the Lagrange multiplier μ_* is strictly positive. Combined with $\mu_* \mathcal{G}(v_*) = 0$ in the KKT condition [\(3.2\)](#page-6-1), this means that $\mathcal{G}(v_*) = 0$, i.e., the constraint $\mathcal{G}(v) \leq 0$ is binding. With this positive Lagrange multiplier $\mu_* > 0$ and the binding condition $\mathcal{G}(v_*) = 0$, we claim the following equivalence between the minimizer of the optimization problem and the solution to the nonlinear system of equations [\(1.2\)](#page-1-0).

Theorem 3.1 (Equivalence).

• Suppose $(a_*, v_*) \in \mathbb{R} \times \mathcal{H}$ solves the constrained optimization problem with $||v_*|| > 0$. Let μ_* be the Lagrange multiplier satisfying the KKT condition. Let (γ_*, σ_*) be the positive scalar defined by

$$
\sigma_* = \|v_*\|/\sqrt{1-\delta^{-1}} > 0, \quad \gamma_* = \|v_*\|/\mu_* > 0.
$$

Then the pair $(a_*, \sigma_*, \gamma_*)$ solves the nonlinear system of equation.

• If $(a_*, \gamma_*, \sigma_*) \in \mathbb{R} \times \mathbb{R}_{>0} \times \mathbb{R}_{>0}$ satisfies the nonlinear system, letting

$$
v_* = \text{prox}[\gamma_* \ell_Y](a_* U + \sigma_* G) - a_* U
$$

 (a_*, v_*) solves the optimization problem (3.1) with $||v_*|| = \sigma_*$ √ $=\sigma_*\sqrt{1-\delta^{-1}}>0$ and the KKT condition is satisfied by the Lagrange multiplier $\mu_* = \sigma_* \sqrt{1 - \delta^{-1}}/\gamma_* > 0$.

Theorem [3.1](#page-6-2) implies that the nonlinear system of equations [\(1.2\)](#page-1-0) admits a unique solution $(a_*, \gamma_*, \sigma_*) \in \mathbb{R} \times \mathbb{R}_{>0} \times \mathbb{R}_{>0}$ if and only if the optimization problem $\min_{\mathcal{G}(v) < 0} \mathcal{L}(a, v)$ admits a unique solution $(a_*, v_*) \in \mathbb{R} \times \mathcal{H}$ with $v_* \neq 0$ and a unique Lagrange multiplier $\mu_* > 0$ satisfying the KKT condition [\(3.2\)](#page-6-1). The next lemma implies that the degenerate case $v_* = 0$ cannot happen.

Lemma 3.2. (Non-degeneracy) If (a_*, v_*) solves the optimization problem [\(3.1\)](#page-6-0), then $v_* \neq 0$.

In the proof of Lemma [3.2,](#page-6-3) the differentiability of the loss ℓ_y is crucial in preventing the degenerate case $v_* = 0$. When the loss is not differentiable, another different threshold, δ_{perfect} , emerges to determine whether $v_* = 0$ or $v_* \neq 0$ occurs (see [\[3\]](#page-7-1)).

Lemma 3.3. (Uniqueness) The minimizer of the optimization problem (3.1) is unique if it exists. Furthermore, the Lagrange multiplier satisfying the KKT condition [\(3.2\)](#page-6-1) is also unique.

Combining Theorem [3.1,](#page-6-2) Lemma [3.2,](#page-6-3) and Lemma [3.3,](#page-7-3) we conclude that the system of nonlinear equations has a unique solution if and only if the infinite-dimensional optimization problem admits a minimizer.

Finally, the following theorem establishes that the existence of a minimizer for the optimization problem is governed by the threshold δ_{∞} , thus completing the proof of Theorem [2.5.](#page-4-2)

Theorem 3.4. Let δ_{∞} be the threshold defined in [\(2.2\)](#page-4-1). Then we have the following:

- If $\delta \leq \delta_{\infty}$, the minimizer of [\(3.1\)](#page-6-0) does not exist.
- If $\delta > \delta_{\infty}$, the minimizer of [\(3.1\)](#page-6-0) exists.

4. Numerical simulation

We generate the covariates $(x_i)_{i=1}^n \stackrel{\text{iid}}{\sim} N(\mathbf{0}_p, I_p)$ and responses $y_i | x_i$ according to the Poisson model

(4.1)
$$
\forall k \in \mathbb{N}, \quad \mathbb{P}(y_i = k \mid \boldsymbol{x}_i) = \frac{\lambda_i^k}{k!} \exp(-\lambda_i) \text{ where } \lambda_i = \exp(-\kappa \boldsymbol{e}_1^\top \boldsymbol{x}_i).
$$

Here, e_1 is the first canonical basis vector, and $\kappa \geq 0$ is the signal strength. We fix $n = 1000$ and for varying values of $(p/n, \kappa)$, we generate 20 datasets of $(\bm{x}_i, y_i)_{i=1}^n$. For each dataset $(\bm{x}_i, y_i)_{i=1}^n$, we solve the optimization problem $\inf_{b \in \mathbb{R}^p} \sum_{i=1}^n \ell_{y_i}(\boldsymbol{x}_i^{\top}b)$ using the Poisson loss $\ell_{y_i}(t) = \exp(t)$ y_i and record whether a minimizer exists using linear programming. In Figure [2,](#page-8-5) we normalize the count of instances where a minimizer exists by dividing by 20, with the black points indicating higher rates of existence. Additionally, we plot the theoretical threshold $1/\delta_{\infty}$ defined in [\(2.2\)](#page-4-1) and compare it with the empirical result. The theoretical threshold effectively separates the two regions, delineating where the M-estimator exists and where it does not.

REFERENCES

- [1] Dennis Amelunxen, Martin Lotz, Michael B McCoy, and Joel A Tropp. Living on the edge: Phase transitions in convex programs with random data. Information and Inference: A Journal of the IMA, 3(3):224–294, 2014.
- [2] Heinz H Bauschke and Patrick L Combettes. Convex Analysis and Monotone Operator Theory in Hilbert Spaces. Springer, 2017.
- [3] Pierre C Bellec and Takuya Koriyama. Existence of solutions to the nonlinear equations characterizing the precise error of m-estimators. arXiv preprint arXiv:2312.13254, 2023.
- [4] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A* nonasymptotic theory of independence. Oxford university press, 2013.
- [5] Emmanuel J Cand`es and Pragya Sur. The phase transition for the existence of the maximum likelihood estimate in high-dimensional logistic regression. The Annals of Statistics, 48(1):27–42, 2020.
- [6] Qiyang Han and Huachen Ren. Gaussian random projections of convex cones: approximate kinematic formulae and applications. $arXiv$ preprint $arXiv:2212.05545$, 2022.
- [7] Bruno Loureiro, Cedric Gerbelot, Hugo Cui, Sebastian Goldt, Florent Krzakala, Marc Mezard, and Lenka Zdeborová. Learning curves of generic features maps for realistic datasets with a teacher-student model. Advances in Neural Information Processing Systems, 34:18137–18151, 2021.
- [8] Fariborz Salehi, Ehsan Abbasi, and Babak Hassibi. The impact of regularization on highdimensional logistic regression. Advances in Neural Information Processing Systems, 32, 2019.
- [9] Pragya Sur and Emmanuel J Cand`es. A modern maximum-likelihood theory for highdimensional logistic regression. *Proceedings of the National Academy of Sciences*, 116(29): 14516–14525, 2019.

FIGURE 2. Count of instances where minimizer exists for varying p/n and signal strength. Simulation parameter: $n = 1000$, 20 replicate, $\ell_y(u) = e^u - yu$ is the Poisson loss, $y_i | x_i$ satisfies the Poisson model [\(4.1\)](#page-7-4).

- 10 PHASE TRANSITION FOR UNREGULARIZED M-ESTIMATION IN SINGLE INDEX MODEL
- [10] Pragya Sur, Yuxin Chen, and Emmanuel J Cand`es. The likelihood ratio test in highdimensional logistic regression is asymptotically a rescaled chi-square. *Probability theory* and related fields, 175:487–558, 2019.
- [11] Wenpin Tang and Yuting Ye. The existence of maximum likelihood estimate in highdimensional binary response generalized linear models. Electronic Journal of Statistics, 2020.
- [12] Christos Thrampoulidis, Ehsan Abbasi, and Babak Hassibi. Precise error analysis of regularized m-estimators in high dimensions. IEEE Transactions on Information Theory, 64 (8):5592–5628, 2018.

Appendix A. Derivation of threshold from convex geometry

Define $\delta_{\infty} \in (0, +\infty]$ as

$$
1/\delta_{\infty}:=\inf_{t\in\mathbb{R}}\varphi(t)
$$

where $\varphi : \mathbb{R} \to \mathbb{R}$ is the convex function defined as

$$
\varphi(t) := \mathbb{E}\big[(G+Ut)^2I\{\Omega_V(Y)\}\big] + \mathbb{E}\big[(G+Ut)^2+I\{\Omega_V(Y)\}\big] + \mathbb{E}\big[(G+Ut)^2-I\{\Omega_V(Y)\}\big].
$$

Here $\Omega_{\vee}, \Omega_{\nearrow}, \Omega_{\searrow}$ are $\sigma(Y)$ -measurable events defined in [\(2.1\)](#page-2-3). We denote by H the Hilbert space which consists of measurable function of (G, Y, U) with finite second moments. Let $\mathcal{C} \subset \mathbb{R} \times \mathcal{H}$ be the cone defined as

(A.1)
$$
(t, p) \in \mathcal{C} \Leftrightarrow tU + p \begin{cases} \leq 0 & \text{under } \Omega_{\nearrow}(Y) \\ = 0 & \text{under } \Omega_{\vee}(Y) \\ \geq 0 & \text{under } \Omega_{\searrow}(Y) \end{cases}
$$

Lemma A.1. The threshold δ_{∞} can be represented as

$$
\delta_{\infty}^{-1} = \inf_{t \in \mathbb{R}} \varphi(t) = \inf_{t \in \mathbb{R}} \mathbb{E}[(G - p(t))^2] = \inf_{(t,p) \in \mathcal{C}} \mathbb{E}[(G - p)^2]
$$

where $(t, p(t)) \in C$ for all $t \in \mathbb{R}$ and $p(t)$ is given by

$$
p(t) := -tU + \begin{cases} 0 & under \ \Omega_{\vee}(Y) \\ (G + Ut)_{-} & under \ \Omega_{\nearrow}(Y) \\ (G + Ut)_{+} & under \ \Omega_{\searrow}(Y). \end{cases}
$$

Suppose that the law of (U, Y) satisfies

(A.2)
$$
\mathbb{E}\Big[U^2\big(I\{\Omega_{\vee}(Y)\} + I\{\Omega_{\searrow}(Y), U > 0\} + I\{\Omega_{\nearrow}(Y), U < 0\}\big)\Big] > 0,
$$

$$
\mathbb{E}\Big[U^2\big(I\{\Omega_{\vee}(Y)\} + I\{\Omega_{\searrow}(Y), U < 0\} + I\{\Omega_{\nearrow}(Y), U > 0\}\big)\Big] > 0.
$$

Then the map φ is coercive, i.e., $\lim_{|t|\to+\infty} \varphi(t) = +\infty$ and $\inf_{t\in\mathbb{R}} \varphi(t)$ admits a minimizer $t_* \in \mathbb{R}$. Furthermore, the optimal $p_* = p(t_*) \in \mathcal{H}$ satisfies

$$
\mathbb{E}[p_*^2] = \mathbb{E}[p_*G], \quad ||p_*|| = \sqrt{1 - \delta_{\infty}^{-1}}.
$$

Proof. Fix $t \in \mathbb{R}$. By the definition of the cone C and φ , it easily follows that

$$
\inf_{p \in \mathcal{H}: (t,p) \in \mathcal{C}} \mathbb{E}[(G-p)^2] = \mathbb{E}[(G-p(t))^2] = \varphi(t)
$$

This proves the representation $\inf_{t\in\mathbb{R}} \varphi(t) = \inf_{(t,p)\in\mathcal{C}} \mathbb{E}[(G-p)^2] = \inf_{t\in\mathbb{R}} \mathbb{E}[(G-p(t))^2]$.

Next, let us show that the map $\varphi : \mathbb{R} \to \mathbb{R}$

$$
\varphi(t) = \mathbb{E}\big[(G+Ut)^2I\{\Omega_\vee(Y)\}\big] + \mathbb{E}\big[(G+Ut)^2I\{\Omega_\nearrow(Y)\}\big] + \mathbb{E}\big[(G+Ut)^2I\{\Omega_\searrow(Y)\}\big]
$$

is coercive. Since $t \mapsto \varphi(t)$ is convex, it suffices to show the coercivity, i.e., $\lim_{|t|\to+\infty} \varphi(t) =$ $+\infty$. For the first term, expanding the square of $(G+Ut)^2$, it immediately follows that

$$
\mathbb{E}[(G+Ut)^2I{\Omega_V(Y)}]=t^2\mathbb{E}[U^2I{\Omega_V(Y)}]+O(t)
$$

For the second term $\mathbb{E}[(G+Ut)^2+I{\Omega}_{\check{c}}(Y)]$,

$$
\mathbb{E}[(G+Ut)^{2}_{+}I{\Omega}_{\nearrow}(Y)\}] = \mathbb{E}[(G+Ut)^{2}I{\Omega}_{\nearrow}(Y), G+Ut>0\}]
$$

= $t^{2}\mathbb{E}[U^{2}I{\Omega}_{\nearrow}(Y)\}I{\G+Ut>0}] + O(|t|)$

as $|t| \to +\infty$. By $\mathbb{E}[U^2] = 1 < +\infty$, the dominated convergence theorem implies

$$
\mathbb{E}[U^2I\{\Omega_{\nearrow}(Y)\}I\{G+Ut>0\}] = \mathbb{E}[U^2I\{\Omega_{\nearrow}(Y)\}I\{G+Ut>0,U>0\}] \n+ \mathbb{E}[U^2I\{\Omega_{\nearrow}(Y)\}I\{G+Ut>0,U<0\}] \n\rightarrow \begin{cases} \mathbb{E}[U^2I\{\Omega_{\nearrow}(Y),U>0\}] & t\rightarrow +\infty \\ \mathbb{E}[U^2I\{\Omega_{\nearrow}(Y),U<0\}] & t\rightarrow -\infty \end{cases}
$$

Thus,

$$
\frac{1}{t^2} \mathbb{E}[(G+Ut)_+^2 I\{\Omega_\nearrow(Y)\} \to \begin{cases} \mathbb{E}[U^2 I\{\Omega_\nearrow(Y), U > 0\}] & t \to +\infty \\ \mathbb{E}[U^2 I\{\Omega_\nearrow(Y), U < 0\}] & t \to -\infty \end{cases}
$$

By the same argument, we have

$$
\frac{1}{t^2} \mathbb{E}[(G+Ut)^2 \cdot I\{\Omega_\searrow(Y)\}] \to \begin{cases} \mathbb{E}[U^2 I\{\Omega_\searrow(Y), U < 0\}] & t \to +\infty \\ \mathbb{E}[U^2 I\{\Omega_\searrow(Y), U > 0\}] & t \to -\infty \end{cases}
$$

Putting them together, we obtain

$$
\frac{\varphi(t)}{t^2} \to \begin{cases} \mathbb{E}[U^2I\{\Omega_\vee(Y)\}] + \mathbb{E}[U^2I\{\Omega_\nearrow(Y), U > 0\}] + \mathbb{E}[U^2I\{\Omega_\searrow(Y), U < 0\}] & t \to +\infty \\ \mathbb{E}[U^2I\{\Omega_\vee(Y)\}] + \mathbb{E}[U^2I\{\Omega_\nearrow(Y), U < 0\}] + \mathbb{E}[U^2I\{\Omega_\searrow(Y), U > 0\}] & t \to -\infty \end{cases}.
$$

Since the limit is strictly positive by the condition $(A.2)$, the map $t \mapsto \varphi(t)$ is coercive, and hence $\inf_{t \in \mathbb{R}} \varphi(t)$ is attained at some $t = t_* \in \mathbb{R}$.

Finally, we prove $\mathbb{E}[p^2_*] = \mathbb{E}[p_*G]$ and $||p_*|| = \sqrt{1 - \delta_{\infty}^{-1}}$. The stationary condition of $\varphi'(t_*) =$ 0 gives

$$
0 = 2 \mathbb{E}[(G + Ut_*)UI\{\Omega_V(Y)\}] + 2 \mathbb{E}[(G + Ut_*) + UI\{\Omega_V(Y)\}] + 2 \mathbb{E}[(G + Ut_*) - UI\{\Omega_V(Y)\}]
$$

= 2 \mathbb{E}[(G - p_*)U]

where

$$
p_* := p(t_*) = -t_*U + (G + Ut_*) - I\{\Omega_\nearrow(Y)\} + (G + Ut_*)_+I\{\Omega_\searrow(Y)\}
$$

Here, $\mathbb{E}[GU] = 0$ since G and U are independent standard normal. Thus, the last equation gives $\mathbb{E}[p_*U] = 0$. Then, we have

$$
\mathbb{E}[p_*^2] = -t_*^2 + t_*^2 + \mathbb{E}[p_*^2] + 2t_* \mathbb{E}[p_* U] \qquad \mathbb{E}[p_* U] = 0
$$

\n
$$
= -t_*^2 + \mathbb{E}[(p_* + Ut_*)^2]
$$

\n
$$
= -t_*^2 + \mathbb{E}\Big[I\{\Omega_{\nearrow}(Y)\}(G + Ut_*)^2\Big] + \mathbb{E}\Big[I\{\Omega_{\searrow}(Y)\}(G + Ut_*)^2\Big],
$$

\n
$$
\mathbb{E}[p_*G] = \mathbb{E}[(p_* + Ut_*)G]
$$

\n
$$
= \mathbb{E}[(t_*U + G)_{-}GI\{\Omega_{\nearrow}(Y)\}] + \mathbb{E}[(t_*U + G)_{+}GI\{\Omega_{\searrow}(Y)\}] \qquad \mathbb{E}[UG] = 0
$$

\n
$$
(G + t \text{ } U)^2 = (G + t \text{ } U) \cdot (G + t \text{ } U) \text{ and the definition of } n \text{, we get}
$$

By
$$
(G + t_*U)^2_{\pm} = (G + t_*U)_{\pm}(G + t_*U)
$$
 and the definition of p_* , we get
\n
$$
\mathbb{E}[p_*^2] - \mathbb{E}[p_*G]
$$
\n
$$
= -t_*^2 + \mathbb{E}[(t_*U + G)_{-t*}UI{\Omega_{\nearrow}(Y)}] + \mathbb{E}[(t_*U + G)_{+t*}UI{\Omega_{\searrow}(Y)}]
$$
\n
$$
= -t_*^2 + t_* \mathbb{E}[U((G + Ut_*)_{-1}{\Omega_{\nearrow}(Y)} + (G + Ut_*)_{+1}{\Omega_{\searrow}(Y)}])]
$$
\n
$$
= -t_*^2 + t_* \mathbb{E}[U(t_*U + p_*)]
$$
\nby the definition of p_*
\n
$$
= -t_*^2 + t_*^2 + 0
$$
\n
$$
= 0.
$$

The equation $||p_*|| = \sqrt{1 - \delta_{\infty}^{-1}}$ follows from $||p_*||^2 = \mathbb{E}[p_*G]$ and $\delta_{\infty}^{-1} = \mathbb{E}[(G - p_*)^2]$ \Box

A.1. Proof of Theorem [2.4.](#page-4-0)

Lemma A.2. Suppose that ℓ_{y_i} is strictly convex, C^1 , and $\inf_u \ell_{y_i}(u)$ is finite. Then the Mestimator does not exist if and only if

(A.3)
$$
\exists \mathbf{b}_{*} \in \mathbb{R}^{p} \setminus \{0\} \text{ such that } \left(\forall i \in [n], \quad \mathbf{x}_{i}^{\top} \mathbf{b}_{*} \begin{cases} \geq 0 & \text{under } \Omega_{\searrow}(y_{i}) \\ = 0 & \text{under } \Omega_{\vee}(y_{i}) \\ \leq 0 & \text{under } \Omega_{\nearrow}(y_{i}) \end{cases} \right)
$$

Proof. [\(A.3\)](#page-11-0) holds \Rightarrow M-estimator does not exist:. Let $L(\boldsymbol{b}) := \sum_{i=1}^n \ell_{y_i}(\boldsymbol{x}_i^{\top} \boldsymbol{b})$ be the objective function. Suppose there exists $\mathbf{b}_* \in \mathbb{R}^p \setminus \{\mathbf{0}_p\}$ such that $(A.3)$ is satisfied. Then, the map $\mathbb{R}_{\geq 0} \ni t \mapsto L(t\mathbf{b}_*)$ is uniformly bounded by $\sum_{i=1}^n \ell_{y_i}(0)$. For all $\mathbf{b} \in \mathbb{R}^p$, let \mathbf{b}_{ν} be the convex combination

$$
\bm{b}_{\nu} = (1 - 1/\nu)\bm{b} + (1/\nu)(\nu \bm{b}_{*}), \quad \nu > 1.
$$

Note $b_{\nu} \to b + b_*$ as $\nu \to +\infty$. By the convexity of $L(b)$ and uniform bound $\sup_{t\geq 0} L(t\mathbf{b}_*) \leq$ $\sum_{i=1} \ell_{y_i}(0)$, we have

$$
L(\boldsymbol{b}_{\nu}) \leq (1 - 1/\nu)L(\boldsymbol{b}_{*}) + (1/\nu)L(\nu \boldsymbol{b}_{*}) \leq (1 - 1/\nu)L(\boldsymbol{b}_{*}) + (1/\nu)\sum_{i=1}^{n} \ell_{y_{i}}(0)
$$

Taking $\nu \to +\infty$, the RHS converges to $L(\boldsymbol{b}_*)$, while the LHS converges to $L(\boldsymbol{b} + \boldsymbol{b}_*)$ by the continuity of $L(\cdot)$ and $\mathbf{b}_{\nu} \to \mathbf{b} + \mathbf{b}_{*}$, so we are left with

$$
L(\boldsymbol{b}+\boldsymbol{b}_*)\leq L(\boldsymbol{b}),\quad \forall \boldsymbol{b}\in\mathbb{R}^p.
$$

Now we suppose that the M-estimator $\hat{\boldsymbol{b}} \in \operatorname{argmin}_{\boldsymbol{b} \in \mathbb{R}^p} L(\boldsymbol{b})$ exists. Setting $\boldsymbol{b} = \hat{\boldsymbol{b}}$ in the above display, since $b_* \neq 0_p$, we know that $\tilde{b} := b_* + \hat{b}$ is also a minimizer with $\tilde{b} \neq \hat{b}$. For all $t \in (0, 1)$, by the convexity of $L(\cdot)$, we have

$$
L(t\hat{\boldsymbol{b}} + (1-t)\tilde{\boldsymbol{b}}) \leq tL(\hat{\boldsymbol{b}}) + (1-t)L(\hat{\boldsymbol{b}})
$$

Since $\hat{\boldsymbol{b}}$ and $\tilde{\boldsymbol{b}}$ minimize L, this holds in equality. With $L(b) = \sum_i \ell_{y_i}(\boldsymbol{x}_i^\top b)$ and by the convexity of ℓ_{y_i} , the equality condition reads to

$$
\forall i \in [n], \quad \ell_{y_i}(t\boldsymbol{x}_i^\top \hat{\boldsymbol{b}} + (1-t)\boldsymbol{x}_i^\top \tilde{\boldsymbol{b}}) = t\ell_{y_i}(\boldsymbol{x}_i^\top \hat{\boldsymbol{b}}) + (1-t)\ell_{y_i}(\boldsymbol{x}_i^\top \tilde{\boldsymbol{b}})
$$

for all $t \in (0,1)$. By the strict convexity of ℓ_{y_i} , we must have $\mathbf{x}_i^{\top} \hat{\boldsymbol{b}} = \mathbf{x}_i^{\top} \tilde{\boldsymbol{b}}$ for all $i \in [n]$, i.e., $\hat{\boldsymbol{b}} - \tilde{\boldsymbol{b}} \in \text{Ker}(\boldsymbol{X})$. However, since $\boldsymbol{X} \in \mathbb{R}^{n \times p}$ has iid $N(0, 1)$ entry, \boldsymbol{X} is an $n \times p$ matrix with rank $(X) = p < n$. This implies $\text{Ker}(\mathbf{X}) = \{0_p\}$ and $\mathbf{b} - \mathbf{b} = 0_p$, which is a contradiction with $\mathbf{b} \neq \mathbf{b}$. Thus, if $(A.3)$ holds then the M-estimator does not exist.

M-estimator does not exist \Rightarrow [\(A.3\)](#page-11-0) holds: Suppose that the M-estimator does not exist. Then, there exists a sequence $(b_k)_{k=1}^{\infty}$ such that as $k \to +\infty$, we have $L(b_k) \to \inf_{b \in \mathbb{R}^p} L(b)$ and $||\mathbf{b}_k|| \to +\infty$. By the compactness of the unit sphere in \mathbb{R}^p , we can extract a subsequence $(b'_k)_{k=1}^{\infty}$ of $(b_k)_{k=1}^{\infty}$ such that $b'_k/||b'_k|| \to v$ for a unit sphere vector $v \in \mathbb{R}^p$. Therefore, we can assume without loss of generality that $\mathbf{b}_k/||\mathbf{b}_k||$ converges to a unit sphere vector \mathbf{v} .

We proceed by contradiction; suppose $(A.3)$ is not satisfied. Then we can find an index $i = i(v) \in [n]$ associated with the unit sphere vector v such that

$$
\boldsymbol{x}_i^{\top} \boldsymbol{v} \begin{cases} < 0 & \text{under } \Omega_{\searrow}(y_i) \\ \neq 0 & \text{under } \Omega_{\vee}(y_i) \\ > 0 & \text{under } \Omega_{\nearrow}(y_i) \end{cases}
$$

If $\Omega_{\searrow}(y_i)$, or $\Omega_{\nearrow}(y_i)$, then the derivative of $t \mapsto \ell_{y_i}(t\boldsymbol{x}_i^{\top}\boldsymbol{v})$ is positive for all $t \geq 1$. If Ω_{\vee} , it is not necessarily positive right away at $t = 1$, but eventually positive for t large enough, say some t_* : for any $t > t_*$, the derivative of $t \mapsto \ell_i(t\mathbf{x}_i^{\top}\mathbf{v})$ is positive. Call this derivative at t_* , say $A = \boldsymbol{x}_i^{\top} \boldsymbol{v} \ell'_{y_i} (t_* \boldsymbol{x}_i^{\top} \boldsymbol{v})$ and $A > 0$.

Let $v_k = b_k / ||b_k||$ so that $x_i^{\top} v_k \to x_i^{\top} v$. Since ℓ_{y_i} is C^1 , for k large enough we have that the derivative of $t \mapsto \ell_{y_i} (t x_i^{\top} v_k)$ at t_* is larger than $A/2$. Call this derivative $A_k = x_i^{\top} v_k \ell'_{y_i} (t_* x_i^{\top} v_k)$ so that $A_k > A/2 > 0$.

By the convexity of $t \mapsto \ell_{y_i}(t\boldsymbol{x}_i^{\top}\boldsymbol{v}_k)$, we have

$$
\ell_{y_i}(\boldsymbol{x}_i^{\top}\boldsymbol{b}_k) = \ell_{y_i}(\|\boldsymbol{b}_k\|_2\boldsymbol{x}_i^{\top}\boldsymbol{v}_k) \geq \ell_{y_i}(t_*\boldsymbol{x}_i^{\top}\boldsymbol{v}_k) + (\|\boldsymbol{b}\|_k - t_*)A_k \geq \ell_{y_i}(t_*\boldsymbol{x}_i^{\top}\boldsymbol{v}_k) + (\|\boldsymbol{b}_k\| - t_*)A/2.
$$

This gives

$$
L(\boldsymbol{b}_k) = \sum_{j=1}^n \ell_{y_j}(\boldsymbol{x}_j^\top \boldsymbol{b}_k) \ge \sum_{j \neq i} \inf_u \ell_{y_j}(u) + \ell_{y_i}(t_* \boldsymbol{x}_i^\top \boldsymbol{v}_k) + (\|\boldsymbol{b}_k\| - t_*)A,
$$

where the RHS goes to ∞ since $||\boldsymbol{b}_k|| \to +\infty$ and $A > 0$. This is a contradiction with $L(\mathbf{b}_k) \to \inf_{\mathbf{b}} L(\mathbf{b}) < +\infty$. Thus, if the M-estimator does not exist then $(A.3)$ holds.

Proof of Theorem [2.4.](#page-4-0) By the rotational invariance of $x_i \sim N(\mathbf{0}_p, I_p)$, we can assume without loss of generality that y_i depend on $\boldsymbol{x}_i = (x_{i1}, x_{i2}, \dots, x_{ip}) \in \mathbb{R}^p$ through its first coordinate x_{i1} , i.e.,

$$
(y_i,x_{i1}) \perp (x_{i2},\cdots x_{ip}).
$$

By Lemma [A.2](#page-11-1) the M-estimator does not exist if and only if

$$
\mathrm{Span}(\boldsymbol{x}_2,\ldots,\boldsymbol{x}_n)\cap\mathcal{C}(\boldsymbol{x}_1,\boldsymbol{y})\neq\{\boldsymbol{0}_p\}
$$

where $\mathcal{C}(\boldsymbol{x}_1, \boldsymbol{y}) \subset \mathbb{R}^n$ is the cone defined as

$$
\boldsymbol{p} \in \mathcal{C}(\boldsymbol{x}_1, \boldsymbol{y}) \Leftrightarrow \exists t \in \mathbb{R} \text{ such that } \left(\forall i \in [n], \ tx_{i1} + p_i = \begin{cases} \geq 0 & \text{under } \Omega_{\searrow}(y_i) \\ = 0 & \text{under } \Omega_{\vee}(y_i) \\ \leq 0 & \text{under } \Omega_{\nearrow}(y_i) \end{cases} \right)
$$

Here $Span(\mathbf{x}_2, \ldots, \mathbf{x}_n)$ is a rotationally invariant random subspace in \mathbb{R}^n with dimension $p-1$, and it is independent of $\mathcal{C}(x_1, y)$. Therefore, by [\[1,](#page-7-0) Theorem I] we have that for $\eta \in (0, 1/2)$, conditionally on x_1, y ,

$$
p-1+\text{stat.dim}(\mathcal{C}) \ge n+n^{1/2+\eta} \Rightarrow \mathbb{P}(\text{M-estimator exists} \mid \mathbf{x}_1, \mathbf{y}) \to 0,
$$

$$
p-1+\text{stat.dim}(\mathcal{C}) \le n-n^{1/2+\eta} \Rightarrow \mathbb{P}(\text{M-estimator exists} \mid \mathbf{x}_1, \mathbf{y}) \to 1
$$

almost everywhere. Here stat. $\dim(\mathcal{C}(\boldsymbol{u}, \boldsymbol{y}))$ is the statistical dimension of the cone given by

$$
\text{stat. dim}(\mathcal{C}(\boldsymbol{x}_1,\boldsymbol{y}))=n-\mathbb{E}[\text{dist}(\boldsymbol{g},\mathcal{C}(\boldsymbol{x}_1,\boldsymbol{y}))^2\mid \boldsymbol{x}_1,\boldsymbol{y}]\quad\text{with}\quad \boldsymbol{g}\sim N(\boldsymbol{0}_n,I_n),
$$

where g is independent of (x_1, y) . Substituting this to the previous display with η set to $1/4$, we get

$$
p-1 \geq \mathbb{E}[\text{dist}(\boldsymbol{g}, \mathcal{C}(\boldsymbol{x}_1, \boldsymbol{y}))^2 \mid \boldsymbol{x}_1, \boldsymbol{y}] + n^{3/4} \Rightarrow \mathbb{P}(\text{M-estimator exists} \mid \boldsymbol{x}_1, \boldsymbol{y}) \to 0
$$

$$
p-1 \leq \mathbb{E}[\text{dist}(\boldsymbol{g}, \mathcal{C}(\boldsymbol{x}_1, \boldsymbol{y}))^2 \mid \boldsymbol{x}_1, \boldsymbol{y}] - n^{3/4} \Rightarrow \mathbb{P}(\text{M-estimator exists} \mid \boldsymbol{x}_1, \boldsymbol{y}) \to 1
$$

With $p/n \to \delta^{-1}$, if we prove the convergence

$$
n^{-1}\,\mathbb{E}[\text{dist}(\bm{g},\mathcal{C}(\bm{x}_1,\bm{y}))^2\mid \bm{x}_1,\bm{y}] \rightarrow^p \delta_{\infty}^{-1}
$$

then we complete the proof. Below we prove this. We will denote x_1 by u so that $(u, y) =$ $(u_i, y_i)_{i=1}^n \stackrel{\text{iid}}{\sim} (U, Y)$. By the explicit gradient identities [\[1,](#page-7-0) (B.7)-(B.9)], the Euclidean norm of the gradient of $g \mapsto \text{dist}(g, C)^2$ is bounded by $2\|g\|_2$. Thus, conditionally on u, y , the Gaussian Poincaré inequality $[4,$ Theorem 3.20] yields

$$
\mathbb{E}[\text{dist}(\boldsymbol{g}, \mathcal{C}(\boldsymbol{u}, \boldsymbol{y}))^2 | \boldsymbol{u}, \boldsymbol{y}] = \text{dist}(\boldsymbol{g}, \mathcal{C}(\boldsymbol{u}, \boldsymbol{y}))^2 + O_P(\sqrt{n}).
$$

Here dist $(g, \mathcal{C}(\boldsymbol{u}, \boldsymbol{y}))^2 = \inf_{\boldsymbol{p} \in \mathcal{C}(\boldsymbol{u}, \boldsymbol{y})} ||\boldsymbol{g} - \boldsymbol{p}||^2$ is equal to the optimal value of

$$
\inf_{(t,\mathbf{p}) \in \mathbb{R} \times \mathbb{R}^n} \sum_{i=1}^n (g_i - p_i)^2 \quad \text{subject to} \quad \left(\forall i \in [n], \ t u_i + p_i = \begin{cases} \geq 0 & \text{under } \Omega_{\searrow}(y_i) \\ = 0 & \text{under } \Omega_{\vee}(y_i) \\ \leq 0 & \text{under } \Omega_{\nearrow}(y_i) \end{cases} \right)
$$

For each t, we can solve the minimization with respect to $p \in \mathbb{R}^n$. The optimal $p = p(t)$ is given by

$$
p_i(t) = -tu_i + \begin{cases} (g_i + u_i t)_+ & \text{under } \Omega_{\searrow}(y_i) \\ 0 & \text{under } \Omega_{\vee}(y_i) \\ (g_i + u_i t)_- & \text{under } \Omega_{\nearrow}(y_i). \end{cases}
$$

Therefore, $\frac{1}{n}$ dist $(g, \mathcal{C}(\boldsymbol{u}, \boldsymbol{y}))^2$ can be written as

$$
\frac{1}{n}\operatorname{dist}(\boldsymbol{g}, \mathcal{C}(\boldsymbol{u}, \boldsymbol{y}))^2 = \frac{1}{n} \inf_{t \in \mathbb{R}} \sum_{i=1}^n (tu_i + g_i)_-^2 I\{\Omega_{\searrow}(y_i)\} + (tu_i + g_i)_-^2 I\{\Omega_{\vee}(y_i)\} + (tu_i + g_i)_+^2 I\{\Omega_{\nearrow}(y_i)\}
$$
\n
$$
= \inf_{t \in \mathbb{R}} \varphi_n(t),
$$

where

$$
\varphi_n(t) := \frac{1}{n} \sum_{i=1}^n (tu_i + g_i)^2 I\{\Omega_{\searrow}(y_i)\} + (tu_i + g_i)^2 I\{\Omega_{\vee}(y_i)\} + (tu_i + g_i)^2 I\{\Omega_{\nearrow}(y_i)\}.
$$

Notice that $\varphi_n(t)$ is a random and convex function, and by the law of large number, $\varphi_n(t) \to^p \varphi(t) = \mathbb{E}[(tU+G)^2 \mathcal{I}\{\Omega_{\searrow}(Y)\}] + \mathbb{E}[(tU+G)^2 \mathcal{I}\{\Omega_{\vee}(Y)\}] + \mathbb{E}[(tU+G)^2 \mathcal{I}\{\Omega_{\nearrow}(Y)\}]$ for each $t \in \mathbb{R}$. By Lemma [A.1,](#page-9-2) we have

$$
\delta_\infty^{-1}=\inf_{t\in\mathbb{R}}\varphi(t),
$$

and $\varphi(t)$ is coercive, i.e., $\lim_{|t|\to+\infty} \varphi(t) = +\infty$. Then, $\inf_{t\in\mathbb{R}} \varphi(t)$ can be reduced to $\min_{t\in K} \varphi(t)$ for a compact set $K \subset \mathbb{R}$, and if a convex function converges pointwisely then it converges uniformly over any compact set. This provides $\inf_{\xi \in \mathbb{R}} \varphi_n(t) \to^P \inf_{t \in \mathbb{R}} \varphi(t) = \delta_{\infty}^{-1}$ and completes the proof. \Box

Appendix B. Set up for infinite-dimensional optimization problem

Lemma B.1. Suppose that $\ell_Y : \mathbb{R} \to \mathbb{R}$ is a proper lower semicontinuous convex function. Then the map

$$
\mathcal{L}: \mathbb{R} \times \mathcal{H} \to \mathbb{R} \cup \{+\infty\}, \quad (a, v) \mapsto \begin{cases} \mathbb{E}[\ell_Y(aU+v)] & \text{if } \mathbb{E}[|\ell_Y(aU+v)|] < +\infty \\ +\infty & \text{otherwise} \end{cases}
$$

is again a proper lower semicontinuous convex function. Furthermore, for all $(a, v) \in \text{dom } \mathcal{L}$, the subderivative at (a, v) is given by

$$
\partial_a \mathcal{L} = \{ \mathbb{E}[Uh] : h \in \partial \ell_Y(aU + v) \} \cap \mathbb{R},
$$

$$
\partial_v \mathcal{L} = \partial \ell_Y(aU + v) \cap \mathcal{H}.
$$

Proof. [\[2,](#page-7-2) Proposition 16.63]. \square

Lemma B.2 ([\[3,](#page-7-1) Lemma A.1]). Define $\mathcal{G}: \mathcal{H} \to \mathbb{R}$ as

$$
\mathcal{G}: \mathcal{H} \to \mathbb{R}, \quad v \mapsto ||v|| - \frac{\mathbb{E}[vG]}{\sqrt{1 - \delta^{-1}}}.
$$

Then G is convex, Lipschitz, and finite valued. Furthermore, G is Fréchet differentiable at $\mathcal{H} \setminus \{0\}$ in the sense that $\mathcal{G}(v + h) = \mathcal{G}(v) + \mathbb{E}[\nabla \mathcal{G}(v)h] + o(||h||)$ for all $||v|| > 0$, where the gradient $\nabla \mathcal{G}(v)$ is given by

$$
\nabla \mathcal{G}(v) = \frac{v}{\|v\|} - \frac{G}{\sqrt{1 - \delta^{-1}}}
$$

.

Lemma B.3 (Existence of Lagrange multiplier). Assume $\mathbb{E}[\ell_Y(G)] < +\infty$. Then $(a_*, v_*) \in$ $dom \mathcal{L}$ solves the constrained optimization problem:

$$
\min_{(a,v)\in\mathbb{R}\times\mathcal{H}} \mathcal{L}(a,v) \quad subject \ to \quad \mathcal{G}(v) \le 0
$$

if and only if there exists an associated Lagrange multiplier $\mu_* \geq 0$ such that the KKT condition below is satisfied:

(B.1)
$$
-\mu_*\partial \mathcal{G}(v_*) \cap \partial_v \mathcal{L}(a_*, v_*) \neq \emptyset, \quad 0 \in \partial_a \mathcal{L}(a_*, v_*), \quad \mu_*\mathcal{G}(v_*) = 0, \quad \mathcal{G}(v_*) \leq 0.
$$

Now we further assume that

- $\mathbb{E}[\left|\inf_u \ell_Y(u)\right|] < +\infty.$
- ℓ_Y is strictly convex.
- $\mathbb{P}(\Omega_{\diagdown}) + \mathbb{P}(\Omega_{\diagup}) > 0$ and $\ell_Y(\cdot)$ is not constant with probability 1.
- There exists a positive constant b and a positive random variable $D(Y)$ such that under Ω_{V} ,

$$
\ell_Y(u) \ge b^{-1}|u| - D(Y), \quad \forall u \in \mathbb{R}
$$

with $\mathbb{E}[D(Y)^2 I\{\Omega_{\vee}\}] < +\infty$.

Then, the Lagrange multiplier is always strictly positive $\mu_* > 0$ and the constraint is binding, *i.e.*, $G(v_*) = 0$.

Proof. First we verify Slater's condition:

$$
\operatorname{lev}_{\leq 0}\mathcal{G}\subseteq\operatorname{int}\operatorname{dom}\mathcal{G},\quad\operatorname{dom}\mathcal{L}\cap\operatorname{lev}_{\leq 0}\mathcal{G}\neq\emptyset.
$$

Since $\mathcal{G}(v) = ||v|| - \mathbb{E}[vG]/\sqrt{2}$ $1-\delta^{-1}$ is finite valued the first condition lev_{≤0} $\mathcal{G} \subseteq \text{int dom } \mathcal{G}$ immediately follows. As for the second condition, $(a, v) = (0, G)$ satisfies $\mathcal{G}(G) = 1 - (1 \delta^{-1}$ $^{-1/2}$ < 0 and $|\mathcal{L}(a, v)| = |\mathbb{E}[\ell_Y(G)]| \leq \mathbb{E}[\ell_Y(G)] < +\infty$ by the assumption. Therefore, the objective function and the constraint $(\mathcal{L}, \mathcal{G})$ satisfy Slater's condition. With Slater' condition, the "if and only if " part follows from [\[2,](#page-7-2) Proposition 27.21].

Let us show $\mu_* > 0$. Suppose $\mu_* = 0$. Then, (a_*, v_*) solves $\min_{(a,v)} \mathbb{E}[\ell_Y(aU + v)]$. Now, for any $n \in \mathbb{N}$, define $v_n \in \mathcal{H}$ as

$$
v_n = \begin{cases} u_{\min} I\{\ell_Y(u_{\min}) \leq n\} & \text{under } \Omega_{\vee} \\ -n & \text{under } \Omega_{\nearrow} \\ n & \text{under } \Omega_{\searrow} \end{cases} \text{ where } u_{\min} \in \operatorname{argmin}_{u \in \mathbb{R}} \ell_Y(u)
$$

Note that v_n is in H since the coercivity assumption implies that under the event Ω_{\vee} ,

$$
|u_{\min}|I\{\ell_Y(u_{\min})\leq n\}\leq b(n+D(Y))I\{\ell_Y(u_{\min})\leq n\}
$$

and the RHS has a finite second moment under Ω_{\vee} by the moment assumption on $D(Y)$. Evaluating the objective function $\mathcal L$ at $(a, v) = (0, v_n)$, by the optimality of (a_*, v_*) , we are left with

$$
\mathbb{E}[\ell_Y(a_*U + v_*)] \leq \mathbb{E}[\ell_Y(0 \cdot U + v_n)]
$$

\n
$$
= \mathbb{E}[\ell_Y(u_{\min})I\{\ell_Y(u_{\min}) \leq n\}I\{\Omega_{\vee}\}]
$$

\n
$$
+ \mathbb{E}[\ell_Y(0)I\{\ell_Y(u_{\min}) > n\}I\{\Omega_{\vee}\}]
$$

\n
$$
+ \mathbb{E}[\ell_Y(-n)I\{\Omega_{\nearrow}\}]
$$

\n
$$
+ \mathbb{E}[\ell_Y(n)I\{\Omega_{\searrow}\}].
$$

Note that each integrand on RHS is uniformly bounded by $|\inf_u \ell_Y(u)| + |\ell_Y(0)|$, where $|\inf_u \ell_Y(u)|$ and $|\ell_Y(0)|$ have finite moments by the assumption and Lemma [B.4.](#page-17-0) Thus, by the dominated

convergence theorem, taking the limit $n \to +\infty$, we obtain

$$
\mathbb{E}[\ell_Y(a_*U + v_*)] \leq \mathbb{E}[\min_u \ell_Y(u)I\{\Omega_\vee\}] + 0 + \mathbb{E}[\inf_u \ell_Y(u)I\{\Omega_\nearrow\}] + \mathbb{E}[\inf_u \ell_Y(u)I\{\Omega_\searrow\}]
$$

=
$$
\mathbb{E}[\inf_u \ell_Y(u)]
$$

and $\mathbb{E}[\ell_Y(a_*U + v_*) - \inf_u \ell_Y(u)] \leq 0$. Since the integrand $\ell_Y(a_*U + v_*) - \inf_u \ell_Y(u)$ is nonnegative, we get

$$
\ell_Y(a_*U + v_*) = \inf_u \ell_Y(u)
$$

with probability 1. Let us consider the event Ω_{γ} . By the strict convexity of $\ell_Y(\cdot)$, under the event Ω_{λ} , we have always $\ell_Y(x) > \lim_{t \to +\infty} \ell_Y(-t) = \inf_u \ell_Y(u)$ for all x. This means that $\ell_Y(a_*U + v_*) = \inf_u \ell_Y(u)$ cannot occur under Ω_{λ} , and hence $\mathbb{P}(\Omega_{\lambda}) = 0$. By the same argument, we get $\mathbb{P}(\Omega_{\diagdown}) = 0$. This is a contradiction with $\mathbb{P}(\Omega_{\diagup}) + \mathbb{P}(\Omega_{\diagdown}) > 0$, so we must have $\mu_* > 0$.

Lemma B.4. Suppose $\mathbb{E}[\ell_Y(G)_+] < +\infty$ and $\mathbb{E}[\inf_u \ell_Y(u)_-] > -\infty$ where $G \sim N(0, 1)$ independent of Y. Then, $\mathbb{E}[\lvert \ell_Y(0) \rvert] < +\infty$.

Proof. Note

$$
|\ell_Y(0)| \le \max(-\left(\inf_u \ell_Y(u)\right)_-,\ell_Y(0)_+) \le -\left(\inf_u \ell_Y(u)\right)_- + \ell_Y(0)_+
$$

and the RHS has a finite moment by the assumption and Jensen's inequality $\mathbb{E}|\ell_Y(0)_+|$ = $\mathbb{E}[\ell_Y(\mathbb{E}[G])_+] \leq \mathbb{E}[\ell_Y(G)_+] < +\infty$. Here we have used the fact that $u \mapsto (\ell_Y(u))_+$ is convex and $G \sim N(0, 1)$ is independent of Y.

Appendix C. Equivalence between nonlinear system and infinite-dimensional optimization problem: Theorem [3.1](#page-6-2)

Lemma C.1. Suppose $\mathbb{E}[\inf_{u} \ell_Y(u)] < +\infty$ and $\mathbb{E}[\ell_Y(G)] < +\infty$. Then for any $a, \sigma, \gamma \in$ $\mathbb{R} \times \mathbb{R}_{>0} \times \mathbb{R}_{>0}$, prox $[\gamma \ell_Y](aU + \sigma G) \in \mathcal{H}$.

Proof. Denote $\text{prox}[\gamma \ell_Y](aU + \sigma G)$ by p_* . Since p_* minimizes $u \mapsto \frac{1}{2\gamma}(aU + \sigma G - u)^2 + \ell_Y(u)$, we have the upper estimate

$$
\frac{1}{2\gamma}(aU + \sigma G - p_*)^2 + \ell_Y(p_*) \le \frac{1}{2\gamma}(aU + \sigma G)^2 + \ell_Y(0).
$$

With $\ell_Y(p) \geq \inf_u \ell_Y(u)$, this implies

$$
(aU + \sigma G - p_*)^2 \le (aU + \sigma G)^2 + 2\gamma(\ell_Y(0) - \inf_u \ell_Y(u)),
$$

where the RHS has a finite moment. This means $aU + \sigma G - p_* \in \mathcal{H}$ and $p_* \in \mathcal{H}$.

□

Lemma C.2. Suppose $(a_*, v_*) \in \mathbb{R} \times \mathcal{H}$ solves the optimization problem with $||v_*|| > 0$. Let us take a Lagrange multiplier $\mu_* > 0$ satisfying the KKT condition [\(B.1\)](#page-15-2). Define the positive scalar $(\gamma_*, \sigma_*) \in \mathbb{R}^2_{>0}$ by √

$$
\sigma_* = \frac{\|v_*\|}{\sqrt{1 - \delta^{-1}}}, \quad \gamma_* = \frac{\sigma_* \sqrt{1 - \delta^{-1}}}{\mu_*}.
$$

Then v_* takes the form of

$$
v_* = \text{prox}[\gamma_* \ell_Y](a_* U + \sigma_* G) - a_* U
$$

and $(a_*, \sigma_*, \gamma_*)$ solves the nonlinear system of equations:

(C.1)
$$
\gamma^{-2} \delta^{-1} \sigma^2 = \mathbb{E}[\ell'_Y (\text{prox}[\gamma \ell_Y](aU + \sigma G))^2]
$$

(C.2)
$$
0 = \mathbb{E}[U\ell'_Y(\text{prox}[\gamma \ell_Y](aU + \sigma G))] \tag{C.2}
$$

(C.3)
$$
\sigma(1 - \delta^{-1}) = \mathbb{E}[G \operatorname{prox}[\gamma \ell_Y](aU + \sigma G)]
$$

Proof. The map $v \mapsto \mathcal{G}(v)$ is Fréchet differentiable at $v = v_* \neq 0$ with $\nabla \mathcal{G}(v_*) = \frac{v_*}{\|v_*\|} - \frac{G}{\sqrt{1-\epsilon}}$ $\overline{1-\delta^{-1}}$ (Lemma [B.2\)](#page-15-1), while the constraint is binding $\mathcal{G}(v_*) = 0$ (Lemma [B.3\)](#page-15-3). Thus, we have

$$
-\mu_*\nabla \mathcal{G}(v_*) = -\mu_*\left(\frac{v_*}{\|v_*\|} - \frac{G}{\sqrt{1-\delta^{-1}}}\right) \in \partial_v \mathcal{L}(a_*, v_*), \quad 0 \in \partial_a \mathcal{L}(a_*, v_*), \quad \mathcal{G}(v_*) = 0,
$$

By the definition of (σ_*, γ_*) , the condition $-\mu_* \nabla \mathcal{G}(v_*) \in \partial_v \mathcal{L}(a_*, v_*)$ yields

$$
\partial_v \mathcal{L}(a_*, v_*) \ni -(v_*-\sigma_*G)/\gamma_*
$$

This means that v_* also minimizes the map

$$
\mathcal{H} \ni v \mapsto \mathcal{L}(v) + \mathbb{E}\Big[\frac{(\sigma_* G - v)^2}{2\gamma_*}\Big] = \mathbb{E}\Big[\ell_Y(a_* U + v) + \frac{(\sigma_* G - v)^2}{2\gamma_*}\Big].
$$

Since $\frac{\text{prox}[\gamma_* \ell_Y](a_* U + \sigma_* G) - a_* U \text{ minimizes the integrand and belongs to } \mathcal{H}$ by Lemma [C.1,](#page-17-1) we have

$$
v_* = \text{prox}[\gamma_* \ell_Y](a_* U + \sigma_* G) - a_* U \in \mathcal{H}.
$$

With $\sigma G - v_* = \gamma \ell'_Y(v_* + a_* U)$, this also gives $\ell'_Y(v_* + a_* U) \in \mathcal{H}$. This in particular means $\mathbb{E}[U\ell'_Y(a_*U+v_*)] \in \mathbb{R}$. With $\partial_a \mathcal{L}(a,v) = \{\mathbb{E}[U\ell'_Y(aU+v)]\} \cap \mathbb{R}$, the condition $0 \in \partial_a \mathcal{L}(a_*,v_*)$ provides

$$
0 = \mathbb{E}[U\ell'_Y(a_*U + v_*)].
$$

With $v_* = \frac{\text{prox}[\gamma_* \ell_Y]}{(a_* U + \sigma_* G)} - a_* U$, we get $(C.2)$. As for $(C.1)$ and $(C.3)$, rearranging $\sigma_* = ||v_*|| / \sqrt{1 - \delta^{-1}}$ and $\mathcal{G}(v_*) = ||v_*|| - \mathbb{E}[v_*G] / \sqrt{1 - \delta^{-1}} = 0$ yields

$$
\|\sigma_* G - v_*\|^2 = \sigma_*^2 - 2\sigma_* \mathbb{E}[v_* G] + \|v_*\|^2
$$

= $\sigma_*^2 - 2\sigma_*^2 (1 - \delta^{-1}) + \sigma_*^2 (1 - \delta^{-1})$
= $\delta^{-1} \sigma_*^2$,

$$
\mathbb{E}[G(v_* + a_* U)] = \sigma_* (1 - \delta^{-1}) + 0.
$$

Substituting $v_* = \frac{\text{prox}[\gamma_* \ell_Y](a_* U + \sigma_* G) - a_* U$ to these two equations, we obtain [\(C.1\)](#page-18-1) and [\(C.3\)](#page-18-2). This completes the proof. \Box

Lemma C.3. Suppose $(a_*, \sigma_*, \gamma_*) \in \mathbb{R} \times \mathbb{R}_{>0}^2$ solves the nonlinear system [\(C.1\)](#page-18-1)-[\(C.3\)](#page-18-2). Then, $(a_*, v_*) \in \mathbb{R} \times \mathcal{H}$ with

$$
v_* = \text{prox}[\gamma_* \ell_Y](a_* U + \sigma_* G) - a_* U \in \mathcal{H}
$$

solves the infinite dimensional optimization problem with $||v_*|| = \sigma_*$ √ $\sigma_*\sqrt{1-\delta^{-1}} > 0$, and the KKT condition [\(B.1\)](#page-15-2) is satisfied by the Lagrange multiplier $\mu_* = \sigma_* \sqrt{1-\delta^{-1}}/\gamma_* > 0$

Proof. We know from Lemma [C.1](#page-17-1) that $v_* \in \mathcal{H}$ and $\sigma G - v_* = \gamma \ell_Y'(v_* + a_* U) \in \mathcal{H}$. In this case the subderivaitve of $\mathcal L$ at (a_*, v_*) is

$$
\partial_a \mathcal{L}(a_*, v_*) = \{ \mathbb{E}[U \ell'_Y(a_* + v_*)] \}, \quad \partial_v \mathcal{L}(a_*, v_*) = \{ \ell'_Y(a_* + v_*) \}.
$$

Noting $\mathbb{E}[UG] = 0$, the nonlinear system can be written as

$$
\delta^{-1}\sigma_*^2 = \mathbb{E}[(\sigma G - v_*)^2]
$$

$$
0 = \mathbb{E}[U\ell'_Y(a_*U + v_*)]
$$

$$
\sigma_*(1 - \delta^{-1}) = \mathbb{E}[Gv_*].
$$

Here, the second equation gives

$$
0 \in \partial_a \mathcal{L}(a_*, v_*).
$$

Rearranging the first and the third equations, we have

$$
||v_*||^2 = (1 - \delta^{-1})\sigma_*^2
$$
, $\mathbb{E}[v_*G] = \sigma_*(1 - \delta^{-1})$.

This implies $||v_*|| > 0$ and $\mathcal{G}(v_*) = 0$. Since $||v_*|| > 0$, \mathcal{G} is differentiable at v_* . The derivative formula gives

$$
-\frac{\|v_*\|}{\gamma_*} \cdot \nabla \mathcal{G}(v_*) = -\frac{\|v_*\|}{\gamma_*} \left(\frac{v_*}{\|v_*\|} - \frac{G}{\sqrt{1-\delta^{-1}}} \right) = \frac{\sigma_* G - v_*}{\gamma_*} = \ell'_Y(a_* U + \sigma_* G) \in \partial_v \mathcal{L}(a_*, v_*)
$$

Therefore, (a_*, v_*) satisfies the KKT condition [\(B.1\)](#page-15-2) with the Lagrange multiplier $\mu_* = \frac{\|\nu_*\|}{\gamma}$ $\frac{\nu_{*} \parallel}{\gamma} > 0,$ and (a_*, v_*) solves the constrained optimization problem.

Appendix D. Non-degeneracy and uniqueness

D.1. **Proof of Lemma [3.2.](#page-6-3)** The argument in this proof is inspired by the proof of $[3, \text{Lemma}]$ $[3, \text{Lemma}]$ 2.6]. Suppose $v_* = 0$. Let $\mu_* > 0$ be the associated Lagrange multiplier satisfying the KKT condition [\(B.1\)](#page-15-2) so that $v_* = 0$ solves the unconstrained optimization problem $\min_{v \in \mathcal{H}} \mathcal{L}(a_*, v) +$ $\mu_*\mathcal{G}(v)$. With $\mathcal{G}(0) = 0$, this gives

$$
\mathbb{E}[\ell_Y(a_*U)] \le \mathbb{E}[\ell_Y(a_*U + v)] + \mu_*(|v|| - \mathbb{E}[vG]/\sqrt{1 - \delta^{-1}}).
$$

for all $v \in \mathcal{H}$. Multiplying the both sides by $\lambda := \sqrt{1 - \delta^{-1}}/\mu_* > 0$ and denoting $f(\cdot) =$ $\lambda(\ell_Y(a_*U + \cdot) - \ell_Y(a_*U))$, we have

(D.1)
$$
0 \leq \mathcal{A}(v) := \mathbb{E}[f(v)] + \mathbb{E}[v^2]^{1/2}\sqrt{1-\delta^{-1}} - \mathbb{E}[vG] \text{ for all } v \in \mathcal{H}.
$$

We parametrize $v \in \mathcal{H}$ as $v_t = \text{prox}[tf](tG) \in \mathcal{H}$ for all $t > 0$ and show $\mathcal{A}(v_t) < 0$ for sufficiently small $t > 0$. Note that $t^{-1}(tG - v_t) \in \partial f(v_t)$ implies

$$
-f(v_t) = f(0) - f(v_t) \ge t^{-1}(tG - v_t)(0 - v_t) = -Gv_t + t^{-1}v_t^2.
$$

Substituting this to $(D.1)$, noting that $\mathbb{E}[v_t G]$ is cancelled out, we have

$$
(D.2)
$$

$$
\mathcal{A}(v_t) \leq \mathbb{E}[Gv_t - t^{-1}v_t^2] + \mathbb{E}[v_t^2]^{1/2}\sqrt{1-\delta^{-1}} - \mathbb{E}[v_tG] \leq -t \cdot ||t^{-1}v_t|| (||t^{-1}v_t|| - \sqrt{1-\delta^{-1}}).
$$

Now we identify the limit of $||v_t/t||$ as $t \to 0+$. The Moreau envelope constructed function $t \mapsto \mathsf{env}_f(tG, t) = \frac{1}{2t} (tG - v_t)^2 + f(v_t)$ has the derivative

$$
-\frac{1}{2t^2}(tG - v_t)^2 + \frac{1}{t}G(tG - v_t) = \frac{1}{2}\Big[G^2 - \left(\frac{v_t}{t}\right)^2\Big],
$$

which is increasing in t because the Moreau envelope $\mathsf{env}_f(x, y)$ is jointly convex in $(x, y) \in$ $\mathbb{R} \times \mathbb{R}_{>0}$ (cf. [\[12,](#page-9-3) Lemma D.1]). This means that v_t^2/t^2 is non-increasing in t and has a non-negative limit as $t \to 0^+$. By the monotone convergence theorem, we get

(D.3)
$$
\lim_{t \to 0^+} ||v_t/t||^2 = \mathbb{E}[\lim_{t \to 0^+} (v_t/t)^2] \in [0, +\infty]
$$

Let us compute the limit v_t/t . First, we claim $v_t \to 0$. By the optimality of $v_t = \text{prox}[tf](tG)$ with $f(\cdot) = \lambda(\ell_Y(a_*U + \cdot) - \ell_Y(a_*U))$, we have

$$
\frac{1}{2t}(tG - v_t)^2 + tf(v_t) \le \frac{(tG)^2}{2t} + f(0) = tG^2 + 0.
$$

This gives

$$
\frac{1}{2}(tG - v_t)^2 \le t^2 G^2 - t \inf_x f(x) = t^2 G^2 - t (\inf_u \ell_Y(u) - \ell_Y(a_* U))
$$

Since $(G, \inf_u \ell_Y(u), \ell_Y(a_* U))$ are all bounded in ℓ_1 , they are all finite with probability 1. This provides $\lim_{t\to 0^+} v_t = 0$. Combined with $G - v_t/t = f'(v_t) = \lambda \ell_Y'(a_* U + v_t)$, since $\ell_Y(\cdot)$ is C^1 , we get $v_t/t \to G - \lambda \ell'_Y(a_*U)$. Substituting this limit to [\(D.3\)](#page-20-1), we obtan

$$
||v_t/t||^2 \to \mathbb{E}[(G - \lambda \ell'_Y(a_*U))^2] \in [0, +\infty].
$$

Recall that $(D.1)$ and $(D.2)$ imply

$$
0 \leq \mathcal{A}(v_t)/t \leq -\|v_t/t\|(\|v_t/t\| - \sqrt{1-\delta^{-1}}), \quad \forall t > 0.
$$

This excludes the case $\mathbb{E}[(G - \lambda \ell_Y(a_* U))^2] = +\infty$ otherwise the RHS converges to $-\infty$ as $t \to 0$. Thus, we must have $\mathbb{E}[(G - \lambda \ell_Y'(a_* U))^2] < +\infty$ and $\ell_Y(a_* U) \in \mathcal{H}$. Expanding the square, since G and $\ell'_Y(a_*U)$ is independent, we get

$$
\lim_{t \to +\infty} ||v_t/t||^2 = 1 + \lambda^2 \mathbb{E}[\ell'_Y(a_*U)^2].
$$

Substituting this to the upper bound of $\mathcal{A}(v_t)/t$,

$$
\lim_{t \to 0} -||v_t/t||(||v_t/t|| - \sqrt{1 - \delta^{-1}}) = -\sqrt{(1 + \lambda^2 \mathbb{E}[\ell_Y'(a_*U)^2])}(\sqrt{1 + \lambda^2 \mathbb{E}[\ell_Y'(a_*U)^2]} - \sqrt{1 - \delta^{-1}})
$$

$$
\leq -(1 - \sqrt{1 - \delta^{-1}}) < 0,
$$

which implies that there exists a sufficiently small t' such that $\mathcal{A}(v_{t})/t' < 0$. This is a contradiction with $\mathcal{A}(v_t)/t \geq 0$ for all $t > 0$. Therefore, we must have $v_* \neq 0$.

D.2. **Proof of Lemma [3.3.](#page-7-3)** Suppose that there exists two minimizer $(a, v), (a', v') \in \mathbb{R} \times \mathcal{H}$. Let $(a_t, v_t) = t(a, v) + (1 - t)(a', v')$ for all $t \in [0, 1]$. Then, by the convexity of objective function and constraint, (a_t, v_t) solves the constrained optimization problem. This implies $\mathcal{L}(a_t, v_t) = t\mathcal{L}(a, v) + (1-t)\mathcal{L}(a', v')$ for all $t \in [0, 1]$. With $\mathcal{L}(a, v) = \mathbb{E}[\ell_Y(aU + v)],$ we must have

$$
\mathbb{E}\Big[\ell_Y(t(aU+v) + (1-t)(a'U+v')) - t\ell_Y(aU+v) - (1-t)\ell_Y(a'U+v')\Big] = 0.
$$

By the convexity of ℓ_Y ,

$$
\ell_Y(t(aU + v) + (1 - t)(a'U + v')) = t\ell_Y(aU + v) + (1 - t)\ell_Y(a'U + v')
$$

for all $t \in [0,1]$. By the strict convexity of ℓ_Y , we must have $aU + v = a'U + v'$. If $a = a'$ holds then we have $v = v'$, completing the proof of uniqueness. We now show that $a = a'$. The condition $aU + v = a'U + v'$ gives

$$
v_t = tv + (1-t)v' = t\{v + U(a-a')\} + (1-t)v' = v' + tU(a-a'),
$$

so that $\mathbb{E}[v_t G] = \mathbb{E}[v'G]$ by independence of G and U. Meanwhile, since (a_t, v_t) solves the constrained optimization problem for all $t \in [0, 1]$, the constraint is satisfied in equality $||v_t|| \mathbb{E}[v_t G]/\sqrt{1-\delta^{-1}} = 0$ for all $t \in [0,1]$ (see Lemma [B.3\)](#page-15-3). Then, $t \mapsto ||v_t||$ must be constant as well. The polynomial $||v_t||^2$ is given

$$
||v_t||^2 = ||v' + tU(a - a')||^2 = ||v'||^2 + t \mathbb{E}[v'U(a - a')] + t^2(a - a')^2.
$$

Since it is constant in t, the quadratic term must be 0, hence $(a - a')^2 = 0$.

Finally, let us show the uniqueness of Lagrange multiplier. Suppose that there exists two distinct Lagrange multipliers $\mu_* \neq \mu_{**} \in \mathbb{R}_{>0}$ associated with the minimizer (a_*, v_*) . Since the subderivative $\partial_v \mathcal{L}(a_*, v_*)$ is the singleton $\{\ell'_Y(a_*U + v_*)\}$ at the minimizer (a_*, v_*) , the KKT conditions

$$
-\mu_*\nabla \mathcal{G}(v_*), -\mu_{**}\nabla \mathcal{G}(v_*) \in \partial_v \mathcal{L}(a_*, v_*)
$$

lead to

$$
-\mu_*\nabla \mathcal{G}(v_*) = -\mu_{**}\nabla \mathcal{G}(v_*) = \ell'_Y(a_*U + v_*).
$$

Combined with $\mu_* \neq \mu_{**}$, this gives $\nabla \mathcal{G}(v_*) = 0$ and $\ell'_Y(a_*U + v_*) = 0$. Here $0 = \nabla \mathcal{G}(v_*) = v_*/\|v_*\| - G/\sqrt{1-\delta^{-1}}$ implies $v_* = \frac{\|v_*\|}{\sqrt{1-\delta^{-1}}}G$. Letting $\sigma_* = \|v_*\|/\sqrt{1-\delta^{-1}} > 0$, substituting √ $1-\delta^{-1} > 0$, substituting this to $\ell'_Y(a_*U + v_*) = 0$, we get

$$
\ell'_Y(a_*U + \sigma_*G) = 0
$$

Since ℓ_Y is strictly convex, this means $\mathbb{P}(\Omega_V) = 1$, which is a contradiction with $\mathbb{P}(\Omega_V) < 1$. This completes the proof of uniqueness of Lagrange multiplier.

Appendix E. Proof of phase transition: Theorem [3.4](#page-7-5)

Lemma E.1. If $\delta \leq \delta_{\infty}$, the optimization problem [\(3.1\)](#page-6-0) does not admit any minimizer.

Proof. Let us take $(t_*, p_*) \in \mathcal{C} \subset \mathbb{R} \times \mathcal{H}$ as in Lemma [A.1](#page-9-2) so that

$$
||p_*|| = \sqrt{1 - \delta_{\infty}^{-1}}, \quad \mathbb{E}[p_*^2] = \mathbb{E}[p_*G].
$$

Substituting this to $\mathcal{G}(p_*) = ||p_*|| - \mathbb{E}[p_*G]/\sqrt{2}$ $1 - \delta^{-1}$, using the condition $\delta \leq \delta_{\infty}$, we have

$$
\mathcal{G}(p_*) = ||p_*|| \left(1 - \frac{\sqrt{1 - \delta_{\infty}^{-1}}}{\sqrt{1 - \delta^{-1}}} \right) \leq 0,
$$

i.e., p_* satisfies the constraint $\mathcal{G} \leq 0$ under the assumption $\delta \leq \delta_{\infty}$.

Let us fix $(a, v) \in \text{dom } \mathcal{L} \cap \text{lev}_{\leq 0} \mathcal{G}$ so that $\mathbb{E}[|\ell_Y(aU + v)|] < +\infty$ and $\mathcal{G}(v) \leq 0$. For any $\nu \geq 1$, consider the convex combination with coefficients $(1 - 1/\nu)$ and $1/\nu$ given by

$$
(a_{\nu}, v_{\nu}) := (a, v)(1 - 1/\nu) + (1/\nu)(\nu t_*, \nu p_*).
$$

Note that $(a_{\nu}, v_{\nu}) \rightarrow (a, v) + (t_*, p_*)$ almost surely as $\nu \rightarrow +\infty$. By the convexity of G, the convex combination (a_{ν}, v_{ν}) also satisfies the constraint $\mathcal{G}(v_{\nu}) \leq 0$. On the other hand, the convexity of ℓ_Y implies

$$
\ell_Y(a_{\nu}U + v_{\nu}) \le (1 - 1/\nu)\ell_Y(aU + v) + (1/\nu) \cdot \ell_Y(\nu t_* U + \nu p_*).
$$

Here, by the definition of $v_*,$

$$
\ell_Y(\nu t_* U + \nu p_*) = \begin{cases} \ell_Y(\nu(G+Ut_*)_-) & \text{under } \Omega_{\nearrow} \\ \ell_Y(\nu(G+Ut_*)_+) & \text{under } \Omega_{\searrow} \\ \ell_Y(0) & \text{under } \Omega_{\vee} \end{cases}
$$

and hence $\ell_Y(\nu t_* U + \nu t_*)$ is uniformly upper bounded by $\ell_Y(0)$ for all $\nu \geq 1$. We have

(E.1)
$$
\ell_Y(a_\nu U + v_\nu) \le (1 - 1/\nu)\ell_Y(aU + v) + (1/\nu)\ell_Y(0).
$$

Taking expectation, we get

$$
\mathcal{L}(a_{\nu},v_{\nu}) \leq (1 - 1/\nu)\mathcal{L}(a,v) + (1/\nu)\mathbb{E}[\ell_Y(0)].
$$

Now we consider the limit $\nu \to +\infty$. The RHS converges to $\mathcal{L}(a, v)$ as $\nu \to +\infty$ since we took $(a, \nu) \in \text{dom }\mathcal{L}$ and $\mathbb{E}[\lvert \ell_Y(0) \rvert] < +\infty$ by Lemma [B.4.](#page-17-0) As for the LHS $\mathcal{L}(a_\nu, v_\nu) =$ $\mathbb{E}[\ell_Y(a_\nu U + v_\nu)]$, from $(E.1)$, the integrand is uniformly bounded as

$$
|\ell_Y(a_{\nu}U + v_{\nu})| \le |\inf_{u} \ell_Y(u)| + |\ell_Y(aU + v)| + |\ell_Y(0)|
$$

where the RHS has a finite expectation. Therefore, by the dominated convergence theorem, we get

$$
\lim_{\nu \to +\infty} \mathcal{L}(a_{\nu}, v_{\nu}) = \lim_{\nu \to +\infty} \mathbb{E}[\ell_Y(a_{\nu}U + v_{\nu})]
$$
\n
$$
= \mathbb{E}[\lim_{\nu \to +\infty} \ell_Y(a_{\nu}U + v_{\nu})]
$$
\n
$$
= \mathbb{E}[\ell_Y((a + t_*)U + v_*)]) \quad \text{by } (a_{\nu}, v_{\nu}) \to (a + t_*, v + p_*) \text{ and the continuity of } \ell_Y(\cdot)
$$
\n
$$
= \mathcal{L}(a + t_*, v + p_*).
$$

We have proved that for any $(a, v) \in \text{dom }\mathcal{L}$, the inequality $\mathcal{L}(a + t_*, v + p_*) \leq \mathcal{L}(a, v)$ holds.

Now we suppose that a minimizer (a_{\min}, v_{\min}) exists. Then it must be unique by Lemma [3.3.](#page-7-3) Applying the inequality $\mathcal{L}(a+t_*,v+p_*) \leq \mathcal{L}(a,v)$ we have established with $(a,v) = (a_{\min}, v_{\min}),$ we must have $t_* = p_* = 0$. Substituting this to the definition of $p_*,$ we are left with

$$
(G)_{-}I\{\Omega_{\nearrow}(Y)\} + (G)_{+}I\{\Omega_{\searrow}(Y)\} = 0.
$$

Taking the expectation of this, since G and Y are independent and $\mathbb{E}[(G)_+] = \mathbb{E}[(G)_-] =$ $\sqrt{2/\pi} > 0$, we obtain $\mathbb{P}(\Omega_{\geq}(Y)) + \mathbb{P}(\Omega_{\leq}) = 0$, which contradicts the assumption $\mathbb{P}(\Omega_{\leq}) =$ $1 - \mathbb{P}(\Omega_{\nearrow}(Y)) - \mathbb{P}(\Omega_{\searrow}) < 1$. Therefore, the minimizer does not exist. \Box

Lemma E.2. Assume that there exists a positive constant b and positive random variable $D(Y)$ with $\mathbb{E}[D(Y)] < +\infty$ such that

$$
\ell_Y(u) \ge -D(Y) + \frac{1}{b} \times \begin{cases} u & under \ \Omega_{\nearrow} \\ -u & under \ \Omega_{\searrow} \\ |u| & under \ \Omega_{\vee} \end{cases}
$$

Suppose $\delta > \delta_{\infty}$. Then for any deterministic $\xi > 0$, if (a, v) satisfies $\mathcal{G}(v) \leq 0$ and $\mathcal{L}(a, v) \leq \xi$, then $a + ||v|| \leq C(\xi)$ for some constant depending on ξ. Consequently, the objective function of the minimization problem is coercive and admits a minimizer (a_*, v_*) .

Proof. By Lemma [A.1,](#page-9-2) δ_{∞} can be represented as $\delta_{\infty}^{-1} = \min_{(t,p)\in\mathcal{C}} \mathbb{E}[(G-p)^2]$, where $\mathcal{C} \subset \mathbb{R} \times \mathcal{H}$ is the cone defined as

$$
(a, v) \in \mathcal{C} \Leftrightarrow (v + aU) \begin{cases} \leq 0 & \text{under } \Omega_{\nearrow} \\ \geq 0 & \text{under } \Omega_{\searrow} \\ = 0 & \text{under } \Omega_{\vee} \end{cases}
$$

and the optimal $(t_*, p_*) \in \mathcal{C}$ satisfies

(E.2)
$$
||p_*|| = \sqrt{1 - \delta_{\infty}^{-1}} < \sqrt{1 - \delta^{-1}}, \quad \mathbb{E}[p_*^2] = \mathbb{E}[p_*G]
$$

where we have used $\delta > \delta_{\infty}$. Now for all $(a, v) \in \mathbb{R} \times \mathcal{H}$, define $\tilde{v} \in \mathcal{H}$ as

$$
\tilde{v} = -aU + \begin{cases} (aU + v)_{-} & \Omega_{\nearrow} \\ (aU + v)_{+} & \Omega_{\searrow} \\ 0 & \Omega_{\vee} \end{cases}
$$

so that $(a, \tilde{v}) \in \mathcal{C}$ for all $(a, v) \in \mathbb{R} \times \mathcal{H}$. Furthermore,

$$
v - \tilde{v} = \begin{cases} (v + aU)_+ & \Omega \nearrow \\ (v + aU)_- & \Omega \searrow \\ v + aU & \Omega \vee \end{cases} \quad \text{and} \quad \|v - \tilde{v}\| \le \|v\| + |a|
$$

By the condition $\mathcal{L}(a, v) = \mathbb{E}[\ell_Y(aU + v)] \leq \xi$ and the coercivity assumption of ℓ_Y , we know

$$
|v - \tilde{v}| \le D(Y) + b \begin{cases} \ell_Y((v + aU)_+) & \Omega \nearrow \\ \ell_Y((v + aU)_-) & \Omega \searrow \\ \ell_Y(v + aU) & \Omega_V \end{cases}
$$

= D(Y) + b\ell_Y(aU + v) + b \begin{cases} \ell_Y(0) - \ell_Y(v + aU) & \Omega \nearrow, aU + v \le 0 \\ \ell_Y(0) - \ell_Y(v + aU) & \Omega \searrow, aU + v \ge 0 \\ 0 & \text{otherwise} \end{cases}
\le D(Y) + b\ell_Y(aU + v) + b|\ell_Y(0) - \inf_{u} \ell_Y(u)|

so that

$$
\mathbb{E}[|v - \tilde{v}|] \le b\xi + \mathbb{E}[D(Y)] + b(\mathbb{E}[|\ell_Y(0)|] + \mathbb{E}[|\inf_u \ell_Y(u)|]) = C^{(1)}(\xi).
$$

By Assumption [1,](#page-3-0) we can take a sufficiently small $u_0 > 0$ such that

$$
\Omega_{+} := \left\{ \{U > u_{0}\} \cap \Omega_{\nearrow} \right\} \cup \left\{ \{U < -u_{0}\} \cap \Omega_{\searrow} \right\} \cup \left\{ \{|U| > u_{0}\} \cap \Omega_{\vee} \right\}
$$

$$
\Omega_{-} := \left\{ \{U < -u_{0}\} \cap \Omega_{\nearrow} \right\} \cup \left\{ \{U > u_{0}\} \cap \Omega_{\searrow} \right\} \cup \left\{ \{|U| > u_{0}\} \cap \Omega_{\vee} \right\}
$$

have positive probabilities. Let $p_0 = \min(\mathbb{P}(\Omega_-), \mathbb{P}(\Omega_+)) > 0$.

Suppose $a > 0$. Under $\{U > u_0\} \cap \Omega_{\geq}$, it holds that $aU + v \leq b\ell_Y(aU + v) + D(Y)$ and hence

$$
au_0 \le aU \le b\ell_Y(aU + v) + D(Y) - v.
$$

Under $\{U < -u_0\} \cap \Omega_{\searrow}$, we have $-(aU + v) \leq b\ell_Y(aU + v) + D(Y)$ and hence $au_0 \le a(-U) \le b\ell_Y(aU + v) + D(Y) + v$

Combining them all together,

$$
au_0 \mathbb{P}(\Omega_+) \leq \mathbb{E}[I\{\Omega_+\}au_0]
$$

\n
$$
\leq b \mathbb{E}[(\ell_Y(aU+v) + D(Y) + |v|)I\{\Omega_+\}]
$$

\n
$$
= b \mathbb{E}[(\ell_Y(aU+v) - \inf_u \ell_Y(u) + \inf_u \ell_Y(u) + D(Y) + |v|)I\{\Omega_+\}]
$$

\n
$$
\leq b \mathbb{E}[\ell_Y(aU+v) - \inf_u \ell_Y(u)] + \mathbb{E}[\inf_u \ell_Y(u)I\{\Omega_+\}] + \mathbb{E}[D(Y)] + \mathbb{E}[|v|]
$$

\n
$$
= b \mathbb{E}[\ell_Y(aU+v)] - \mathbb{E}[\inf_u \ell_Y(u)I\{\Omega_+^c\}] + \mathbb{E}[D(Y)] + \mathbb{E}[|v|]
$$

\n
$$
\leq b\xi + \mathbb{E}[\inf_u \ell_Y(u)] + \mathbb{E}[D(Y)] + ||v||_2.
$$

By the same argument, if $a < 0$, considering the event $\Omega_-,$ we have

$$
(-a)u_0 \mathbb{P}(\Omega_-) \le b\xi + \mathbb{E}[\left|\inf_u \ell_Y(u)\right|] + \mathbb{E}[D(Y)] + \|v\|_2
$$

Combined with $\min(\mathbb{P}(\Omega_+), \mathbb{P}(\Omega_-)) = p_0 > 0$, we have that

$$
|a| \le (u_0 p_0)^{-1} \Big(\xi + \mathbb{E}[\|\inf_u \ell_Y(u)\|] + \mathbb{E}[D] + \|v\|_2 \Big) \le C_1(\xi)(1 + \|v\|_2)
$$

With $||v - \tilde{v}|| \leq |a| + ||v||$, we obtain

(E.3)
$$
||v - \tilde{v}|| \leq C^{(2)}(\xi)(1 + ||v||_2).
$$

Now we claim

(E.4)
$$
\mathbb{E}[(G - p_*)^2] + \mathbb{E}[(p_* - \tilde{v})^2] \le \mathbb{E}[(G - \tilde{v})^2]
$$

Define for $s \in [0, 1]$ the convex combination $v_s = p_*(1 - s) + s\tilde{v}$ and the function

$$
\varphi(s) = \mathbb{E}[(G - v_s)^2] - \mathbb{E}[(G - p_*)^2] - \mathbb{E}[(v_s - p_*)^2].
$$

Here, we have $0 \leq \mathbb{E}[(G-v_s)^2] - \mathbb{E}[(G-p_*)^2]$ by the optimality of $(t_*, p_*) \in \operatorname{argmin}_{(t,p) \in \mathcal{C}} \mathbb{E}[(G-p_*)^2]$ p)², and the last term $-\mathbb{E}[(v_s - p_*)^2]$ is $O(s^2)$ by definition of v_s . Note that $\varphi(0) = 0$ and the quadratic term in $\varphi(s)$ cancel out. Since $\varphi(s)$ is linear in s and $\varphi(s) \le O(s^2)$ as $s \to 0$, the slope $\varphi'(0)$ must be non-negative and $\varphi(0) \leq \varphi(1)$ which is exactly inequality [\(E.4\)](#page-25-0).

In both sides of [\(E.4\)](#page-25-0), $\mathbb{E}[\tilde{v}^2]$ cancel out, $\mathbb{E}[G^2] = 1$ and $\mathbb{E}[(G - p_*)^2] = 1 - ||p_*||^2$, so that the previous display can be rewritten $-2 \mathbb{E}[p_*\tilde{v}] \leq -2 \mathbb{E}[\tilde{v}G]$. From this, for all v satisfying the constraint $\mathcal{G}(v) = ||v|| - \mathbb{E}[vG] \leq 0$,

$$
\mathbb{E}[\tilde{v}G] \leq \mathbb{E}[p_*\tilde{v}].
$$

\n
$$
\Rightarrow \mathbb{E}[vG] + \mathbb{E}[(\tilde{v} - v)G] \leq \mathbb{E}[p_*v] + \mathbb{E}[(\tilde{v} - v)p_*],
$$

\n
$$
\Rightarrow \left(\sqrt{1 - \delta^{-1}} - ||p_*||\right) ||v|| \leq \mathbb{E}[v(G - p_*)] \leq \mathbb{E}[(\tilde{v} - v)(p_* - G)].
$$

The parenthesis on the left is positive thanks to $(E.2)$. On the right-hand side, considering the event $I\{|p_* - G| > C\}$ for some constant $C > 0$ to be specified later, the Cauchy–Schwartz inequality gives

$$
\mathbb{E}[(\tilde{v} - v)(p_{*} - G)] \le C \mathbb{E}[|\tilde{v} - v|] + \mathbb{E}[I\{|p_{*} - G| > C\}|p_{*} - G||\tilde{v} - v|]
$$

$$
\le C \mathbb{E}[|\tilde{v} - v|] + \sqrt{\mathbb{E}[I\{p_{*} - G| \ge C\}|p_{*} - G|^{2}]} \cdot \|\tilde{v} - v\|.
$$

Since $p_* - G$ is bounded in L2, we can take a sufficiently large $C^{(3)}(\epsilon)$ for all $\epsilon > 0$ such that $\mathbb{E}[I\{|p_*-G| > C^{(3)}(\epsilon)\}|p_*-G|^2] \leq \epsilon^2$. Thus, we have that for all $\epsilon > 0$,

(E.6)
$$
\left(\sqrt{1-\delta^{-1}}-\|p_*\|\right)\|v\| \leq C^{(3)}(\epsilon)\,\mathbb{E}\,|\tilde{v}-v|+\epsilon\|v-\tilde{v}\|_2.
$$

Recall the bound $||v - \tilde{v}|| \leq C^{(2)}(\xi)(1 + ||v||)$ in [\(E.3\)](#page-25-1), so if we take a sufficiently small $\epsilon = \epsilon_0$ such that $\epsilon_0 C^{(2)}(\xi) \leq$ ($\left| \begin{array}{c} v \\ v \end{array} \right|$ $(1 - \delta^{-1} - ||p_*||)/2$, we obtain

(E.7)
$$
\frac{1}{2}(\sqrt{1-\delta^{-1}} - ||p_*||) ||v|| \leq C(\epsilon_0) \mathbb{E} |\tilde{v} - v| + \epsilon_0 C^{(2)}(\xi)
$$

Since $\mathbb{E} |\tilde{v} - v| \leq C^{(1)}(\xi)$ was established at the beginning, this completes the proof. \Box