# FIXED-TERM DECOMPOSITIONS USING EVEN-INDEXED FIBONACCI NUMBERS

HÙNG VIỆT CHU, ANEY MANISH KANJI, AND ZACHARY LOUIS VASSEUR

ABSTRACT. As a variant of Zeckendorf's theorem, Chung and Graham proved that every positive integer can be uniquely decomposed into a sum of even-indexed Fibonacci numbers, whose coefficients are either 0, 1, or 2 so that between two coefficients 2, there must be a coefficient 0. This paper characterizes all positive integers that do not have  $F_{2k}$  ( $k \ge 1$ ) in their decompositions. This continues the work of Kimberling, Carlitz et al., Dekking, and Griffiths, to name a few, who studied such a characterization for Zeckendorf decomposition.

#### 1. INTRODUCTION

We define the Fibonacci sequence  $(F_n)_{n=1}^{\infty}$  as  $F_1 = F_2 = 1$  and  $F_{n+1} = F_n + F_{n-1}$ for  $n \ge 2$ . Zeckendorf's theorem [28] states that every positive integer can be uniquely written as a sum of nonadjacent Fibonacci numbers from  $(F_n)_{n=2}^{\infty}$ . The sum is called the Zeckendorf decomposition of a positive integer. Note that we start from  $F_2$  since otherwise,  $F_1 = F_2 = 1$  ruins uniqueness. Zeckendorf-type decompositions have been extensively studied in the literature: to name a few, see [5, 6, 8, 10, 11, 14, 15, 20, 24, 25] for various generalizations to other sequences, [3, 12, 13, 16, 17, 21, 27] for digits in the decomposition, and [1, 2, 23, 26] for Zeckendorf games.

A beautiful Zeckendorf-type decomposition that uses even-indexed Fibonacci numbers only is due to Chung and Graham [9]:

**Theorem 1.1.** [9, Lemma 1] Every positive integer n can be uniquely represented as a sum  $n = \sum_{i\geq 1} c_i F_{2i}$ , where  $c_i$ 's are in  $\{0, 1, 2\}$  so that if  $c_i = c_j = 2$  with i < j, then for some k, i < k < j, we have  $c_k = 0$ .

We call the decomposition in Theorem 1.1 the *Chung-Graham decomposition* of an integer. This paper answers the following question.

**Question 1.2.** Given an even-indexed Fibonacci number  $F_{2N}$ ,  $N \ge 1$ , what are the positive integers whose Chung-Graham decomposition contains neither  $F_{2N}$  nor  $2F_{2N}$ ?

The analog of Question 1.2 is well-known for Zeckendorf decomposition. Kimberling [22] studied numbers without 1 in their decomposition. Another pioneering paper is due to Carlitz et al. [4] who described the set Z(N) of all positive integers having the summand  $F_N$  in their Zeckendorf decomposition, depending on the parity of N. Later,

<sup>2020</sup> Mathematics Subject Classification. 11B39.

Key words and phrases. Zeckendorf decomposition, even-indexed Fibonacci numbers, fixed terms.

The work was partially supported by the College of Arts & Sciences at Texas A&M University. A. M. Kanji and Z. L. Vasseur are undergraduates at Texas A&M University.

Griffiths [19] gave

$$Z(N) = \left\{ F_N \left[ \frac{n + \phi^2}{\phi} \right] + nF_{N+1} + j : 0 \le j \le F_{N-1} - 1, n \ge 0 \right\}.$$

Dekking [13] characterized all integers that share the same *initial* Zeckendorf decomposition, using the so-called compound Wythoff sequences and generalized Beatty sequences.

It is worth mentioning that Griffiths' analysis [19] can also be used to determine all positive integers having  $\{F_N : N \in A\}$  for some certain sets A in their Zeckendorf decomposition. The idea is to analyze consecutive rows of the table of all numbers having  $F_N$  as the minimum summand in their Zeckendorf decomposition and employed properties of the golden string, which we shall discuss in Section 2. Recently, Chu [7] generalized the golden string to study a generalized Zeckendorf decomposition.

In the present paper, we answer Question 1.2 using the same method as in [7, 19] while dealing with a considerably more involved table due to the appearance of the coefficient 2 in the Chung-Graham decomposition. In the process, we need to utilize more properties of the golden string (see Propositions 2.1 and 2.3, for example). We state our main result.

**Theorem 1.3.** For  $N \ge 1$ , the set of all positive integers that do not have  $F_{2N}$  nor  $2F_{2N}$  in their Chung-Graham decomposition is given by

$$B_{2N} := [1, F_{2N} - 1] \cup \bigcup_{k=N+1}^{\infty} \left\{ j + F_{2k}, j + (n+2)F_{2k} + \left\lfloor \frac{n+1}{\phi} \right\rfloor F_{2k-1} : 0 \le j \le F_{2N} - 1, n \ge 0 \right\},$$

where  $\phi = (\sqrt{5} + 1)/2$ .

For example, we list integers at most 30 that belong to the following sets

$$B_{2} = \{3, 6, 8, 11, 14, 16, 19, 21, 24, 27, 29, \ldots\},\$$
  

$$B_{4} = \{1, 2, 8, 9, 10, 16, 17, 18, 21, 22, 23, 29, 30\},\$$
  

$$B_{6} = \{1, 2, 3, 4, 5, 6, 7, 21, 22, 23, 24, 25, 26, 27, 28\}.$$

To facilitate our writing, we introduce some notation that distinguish the two coefficients 1 and 2. Given  $n \in \mathbb{N}$ , let  $C\mathcal{G}(n)$  denote the set of all Fibonacci numbers in the Chung-Graham decomposition of n. Let  $C\mathcal{G}_1(n)$  be the set of all numbers in  $C\mathcal{G}(n)$  that have coefficient 1 in the Chung-Graham decomposition of n, and let  $C\mathcal{G}_2(n) := C\mathcal{G}(n) \setminus C\mathcal{G}_1(n)$  be the set of all numbers in  $C\mathcal{G}(n)$  that have coefficient 2. For example,

$$\mathcal{CG}(2F_2 + F_4 + 2F_8 + F_{14}) = \{F_2, F_4, F_8, F_{14}\}, \\ \mathcal{CG}_1(2F_2 + F_4 + 2F_8 + F_{14}) = \{F_4, F_{14}\}, \text{ and} \\ \mathcal{CG}_2(2F_2 + F_4 + 2F_8 + F_{14}) = \{F_2, F_8\}.$$

For example,  $\max CG(n) = \max CG_1(n) = F_{14}$  means that the largest Fibonacci number that appears in the Chung-Graham decomposition of n is  $F_{14}$  whose coefficient is 1.

The paper is structured as follows: in Section 2, we define the golden string S and collect several properties that will be used in due course; Section 3 investigates the ordered list of integers having  $F_{2k}$  as the smallest Fibonacci number in their Chung-Graham decomposition; finally, Section 4 gathers some auxiliary results before proving Theorem 1.3.

#### 2. The Golden string

For two finite strings of symbols X and Y, we write X : Y to mean the concatenation of X and Y. The golden string, denoted by S, is an infinite string consisting of the letters A and B, built recursively as follows:  $S_1 = B$ ,  $S_2 = BA$ , and  $S_k = S_{k-1} : S_{k-2}$ for  $k \geq 3$ . For example,

$$S_3 = S_2 : S_1 = BAB,$$
  

$$S_4 = S_3 : S_2 = BABBA,$$
  

$$S_5 = S_4 : S_3 = BABBABAB$$

The first few letters of S are

We record several properties of S and  $(S_n)_{n=1}^{\infty}$ :

- a) The length of  $S_n$ , denoted by  $|S_n|$ , is equal to  $F_{n+1}$ .
- b) The substring  $S_n$  gives the first  $F_{n+1}$  letters of S.
- c) For each  $n \ge 2$ , the substring consisting of the first  $F_n$  letters of S is the same as the substring consisting of all the letters of S between the  $F_{n+1} + 1^{\text{th}}$  and the  $F_{n+2}$ <sup>th</sup> positions, inclusively. This claim follows immediately from the following:
  - by Property b), the first  $F_n$  letters of S are given by  $S_{n-1}$ ;
  - also by Property b), the first  $F_{n+2}$  letters of S are given by

$$S_{n+1} := S_n : S_{n-1};$$
 and

• by Property a),  $|S_n| = F_{n+1}$ .

- d) Griffiths [18] proved a neat relation between the Zeckendorf decomposition and the golden string: if  $n = F_{c_1} + F_{c_2} + \cdots + F_{c_\ell}$  is the Zeckendorf decomposition of n, then  $S_{c_k}: S_{c_{k-1}}: \cdots : S_{c_1}$  gives the first n letters of the golden string.
- e) Let  $N_B(n)$  denote the number of B's in the first n letters of S. We recall Griffiths' [18, Lemma 3.3], which states that

$$N_B(n) = \left\lfloor \frac{n+1}{\phi} \right\rfloor, \tag{2.1}$$

where  $\phi = (1 + \sqrt{5})/2$ , the golden ratio.

- f) The following claims are easily proved by induction: for  $m \ge 1$ ,
  - (f1) The  $(F_{2m+1}-1)^{\text{th}}$  letter of S is B.

  - (f1) The  $(2^{2m+1})$  is  $F_{2m+1}$  better of S is A. (f3) The  $(2F_{2m+1}-1)$ <sup>th</sup> letter of S is B.
  - (f4) The  $2F_{2m+1}^{\text{th}}$  letter of S is A.
  - Due to Proposition 2.1 below, (f3) and (f4) follow immediately from (f1) and (f2).

**Proposition 2.1.** For  $n \ge 4$ , the substring of S consisting of the first  $F_n$  letters of S is the same as the substring of S consisting of the next  $F_n$  letters.

*Proof.* Fix  $n \ge 4$ . Since for  $k < \ell$ ,  $S_k$  gives the initial letters of  $S_\ell$ , we can write  $S_{n-1} = S_{n-3} : L$  for some finite string L. We have

$$S_{n+1} = S_n : S_{n-1} = (S_{n-1} : S_{n-2}) : (S_{n-3} : L)$$
  
=  $S_{n-1} : (S_{n-2} : S_{n-3}) : L$   
=  $S_{n-1} : S_{n-1} : L.$ 

Since  $S_{n+1}$  gives the initial letters of S and  $|S_{n-1}| = F_n$ , we are done.

We shall use the following notation. For a string W, we write W - 2 to mean the string formed by deleting the last two letters of W.

**Lemma 2.2.** For  $n \ge 1$ , we have

$$S_n: S_{n+1} - 2 = S_{n+1}: S_n - 2.$$

*Proof.* We prove by induction. The equality is true for n = 1. Inductive hypothesis: suppose that it is true for  $n = \ell \ge 1$ . We show that

$$S_{\ell+1}: S_{\ell+2} - 2 = S_{\ell+2}: S_{\ell+1} - 2.$$

We have

$$S_{\ell+2}: S_{\ell+1} - 2 = (S_{\ell+1}: S_{\ell}): S_{\ell+1} - 2 = S_{\ell+1}: (S_{\ell}: S_{\ell+1} - 2)$$
  
=  $S_{\ell+1}: (S_{\ell+1}: S_{\ell} - 2) = S_{\ell+1}: S_{\ell+2} - 2.$ 

**Proposition 2.3.** For  $n \ge 5$ , the substring of S consisting of the first  $(F_{n-1} - 2)$  letters is the same as the substring of S consisting of all the letters between the  $(2F_n + 1)^{th}$  letter and the  $(F_{n+2} - 2)^{th}$  letter, inclusively.

*Proof.* Pick  $n \ge 5$ . We have

$$S_{n+1} = S_n : S_{n-1} = S_{n-1} : S_{n-2} : S_{n-2} : S_{n-3}$$
  
=  $S_{n-1} : S_{n-2} : S_{n-3} : S_{n-4} : S_{n-3}$   
=  $S_{n-1} : S_{n-1} : S_{n-4} : S_{n-3}.$ 

Observe that  $|S_{n-1}| = F_n$  and  $|S_{n-4} : S_{n-3}| = F_{n-3} + F_{n-2} = F_{n-1}$ . Hence, the substring of S consisting of all the letters between the  $(2F_n+1)^{\text{th}}$  letter and the  $(F_{n+2}-2)^{\text{th}}$  letter, inclusively is

$$W := S_{n-4} : S_{n-3} - 2.$$

By Lemma 2.2,

$$W = S_{n-3} : S_{n-4} - 2 = S_{n-2} - 2,$$

which gives the first  $F_{n-1} - 2$  letters of S.

## 3. The ordered list of positive integers n with $F_{2k} = \min CG(n)$

For  $k \ge 1$ , let  $A_{2k} = \{n : F_{2k} = \min CG(n)\}$ . We form a table whose rows are numbers in  $A_{2k}$  arranged in increasing order  $q(1) < q(2) < q(3) < \cdots$ . Let us look at the first few rows of the table.

q(1)	$F_{2k}$		
q(2)	$2F_{2k}$		
q(3)	$F_{2k}$	$F_{2k+2}$	
q(4)	$2F_{2k}$	$F_{2k+2}$	
q(5)	$F_{2k}$	$2F_{2k+2}$	
q(6)	$F_{2k}$		$F_{2k+4}$
q(7)	$2F_{2k}$		$F_{2k+4}$
q(8)	$F_{2k}$	$F_{2k+2}$	$F_{2k+4}$
q(9)	$2F_{2k}$	$F_{2k+2}$	$F_{2k+4}$
q(10)	$F_{2k}$	$2F_{2k+2}$	$F_{2k+4}$
q(11)	$F_{2k}$		$2F_{2k+4}$
q(12)	$2F_{2k}$		$2F_{2k+4}$
q(13)	$F_{2k}$	$F_{2k+2}$	$2F_{2k+4}$
:			

Table 1. The numbers in  $A_{2k}$  in increasing order

This section shows a way to construct new rows of Table 1 recursively. First, we prove that a number with a larger maximum Fibonacci number in its Chung-Graham decomposition belongs to a lower row. Consequently, the numbers n with the same  $\max CG(n)$  form consecutive rows in the table.

**Lemma 3.1.** For  $m \ge 0$ , the largest positive integer  $n \in A_{2k}$  with  $\max CG(n) = F_{2k+2m}$ , denoted by N(m), is

$$F_{2k} + F_{2k+2} + \dots + F_{2k+2m-2} + 2F_{2k+2m}$$

*Proof.* Write N(m) as  $\sum_{i=0}^{m} c_i F_{2k+2i}$ , where the  $c_i$ 's are in  $\{0, 1, 2\}$  and satisfy the Chung-Graham condition. Suppose, for a contradiction, that  $c_j = 0$  for some  $1 \leq j \leq m-1$ . If changing  $c_j$  to 1 does not violate the Chung-Graham condition, then  $N(m) + F_{2k+2j}$  is greater than N(m), while  $\max C\mathcal{G}(N(m) + F_{2k+2j}) = F_{2k+2m}$ . This contradicts the maximality of N(m). Hence, changing  $c_j$  to 1 violates the Chung-Graham condition; that is, there are  $1 \leq j' < j < j'' \leq m$  such that  $c_{j'} = c_{j''} = 2$ . Here we choose the largest j' and the smallest j'' that satisfy these conditions. We change both  $c_{j'}$  and  $c_j$  to 1. Then the new coefficients still satisfy the Chung-Graham condition, but since

$$F_{2k+2j'} + F_{2k+2j} > 2F_{2k+2j'} + 0F_{2k+2j},$$

the new number is greater than N(m). This again contradicts the maximality of N(m). Therefore,  $c_i \ge 1$  for all *i*, which clearly implies that

$$N(m) = F_{2k} + F_{2k+2} + \dots + F_{2k+2m-2} + 2F_{2k+2m}$$

as desired.

**Corollary 3.2.** Given  $n, m \in A_{2k}$ , if  $\max CG(n) < \max CG(m)$ , then n < m.

*Proof.* Suppose that  $\max CG(n) = F_{2k+2j_1}$  and  $\max CG(m) = F_{2k+2j_2}$  with  $j_1 < j_2$ . Then  $m \ge F_{2k} + F_{2k+2j_2}$ . By Lemma 3.1,

$$n \leq F_{2k} + F_{2k+2} + \dots + F_{2k+2j_{1}-2} + 2F_{2k+2j_{1}}$$
  
=  $(F_{2k-1} + F_{2k} + F_{2k+2} + \dots + F_{2k+2j_{1}-2}) + 2F_{2k+2j_{1}} - F_{2k-1}$   
=  $F_{2k+2j_{1}-1} + 2F_{2k+2j_{1}} - F_{2k-1} = F_{2k+2j_{1}+2} - F_{2k-1} < F_{2k+2j_{2}} < m.$ 

Thanks to Corollary 3.2, we know that the numbers having  $F_{2k+2\ell}$  ( $\ell \ge 1$ ) (either coefficient 1 or 2) as the maximum Fibonacci number in its decomposition form consecutive rows in Table 1. The next result tells us their location.

**Lemma 3.3.** For  $\ell \geq 1$ , the numbers in  $\{n \in A_{2k} : \max CG(n) = F_{2k+2\ell}\}$  lie between the  $(F_{2\ell+1}+1)^{th}$  and the  $F_{2\ell+3}^{th}$  rows, inclusively.

*Proof.* We prove by induction. Base case: it is easy to verify that the lemma holds for  $\ell = 1$ . Inductive hypothesis: suppose the lemma is true for  $\ell \leq m$  for some  $m \geq 1$ . We prove that the lemma holds for  $\ell = m + 1$ ; that is, the numbers in  $\{n \in A_{2k} : \max C\mathcal{G}(n) = F_{2k+2m+2}\}$  lie between the  $(F_{2m+3}+1)^{\text{th}}$  to the  $F_{2m+5}^{\text{th}}$  rows, inclusively. By the inductive hypothesis,

$$\begin{aligned} &|\{n \in A_{2k} : \max \mathcal{CG}(n) \le F_{2k+2m}\}| \\ &= \sum_{\ell=0}^{m} |\{n \in A_{2k} : \max \mathcal{CG}(n) = F_{2k+2\ell}\}| \\ &= 2 + \sum_{\ell=1}^{m} |\{n \in A_{2k} : \max \mathcal{CG}(n) = F_{2k+2\ell}\}| \\ &= F_3 + \sum_{\ell=1}^{m} (F_{2\ell+3} - F_{2\ell+1}) = F_3 + \sum_{\ell=1}^{m} F_{2\ell+2} = F_{2m+3}. \end{aligned}$$

Therefore, the numbers with the largest summand  $F_{2k+2m+2}$  start at the  $(F_{2m+3}+1)^{\text{th}}$  row.

It suffices to prove that

$$|\{n \in A_{2k} : \max \mathcal{CG}(n) = F_{2k+2m+2}\}| = F_{2m+4}$$

Note that

$$\{n \in A_{2k} : \max \mathcal{CG}(n) = F_{2k+2m+2}\}$$
$$= \{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_1(n) = F_{2k+2m+2}\}$$
$$\cup \{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_2(n) = F_{2k+2m+2}\}.$$

Since all numbers in  $\{n \in A_{2k} : \max CG(n) = \max CG_1(n) = F_{2k+2m+2}\}$  are created by adding  $F_{2k+2m+2}$  to the numbers in  $\{n \in A_{2k} : \max CG(n) \leq F_{2k+2m}\}$ , we know that

$$|\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_1(n) = F_{2k+2m+2}\}| = F_{2m+3}.$$

It remains to show

$$|\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_2(n) = F_{2k+2m+2}\}| = F_{2m+2}.$$

All numbers in  $\{n \in A_{2k} : \max C\mathcal{G}(n) = \max C\mathcal{G}_2(n) = F_{2k+2m+2}\}$  are formed by adding  $2F_{2k+2m+2}$  to a subset of  $\{n \in A_{2k} : \max C\mathcal{G}(n) \le F_{2k+2m}\}$ . Pick  $s \in \{n \in A_{2k} : \max C\mathcal{G}(n) \le F_{2k+2m}\}$  such that  $s + 2F_{2k+2m+2} \in \{n \in A_{2k} : \max C\mathcal{G}(n) = \max C\mathcal{G}_2(n) = F_{2k+2m+2}\}$ . Equivalently, if the Chung-Graham decomposition of s is  $\sum_{i=0}^m c_i F_{2k+2i}$ , then  $(c_0, c_1, \ldots, c_m)$  must have one of the following forms, based on the largest i (if any) with  $c_i = 0$ :

$$\begin{array}{rcl}
f_1 & (c_0, c_1, \dots, c_{m-3}, c_{m-2}, c_{m-1}, 0) \\
f_2 & (c_0, c_1, \dots, c_{m-3}, c_{m-2}, 0, 1) \\
f_3 & (c_0, c_1, \dots, c_{m-3}, 0, 1, 1) \\
\vdots \\
f_m & (c_0, 0, \dots, 1, 1, 1, 1) \\
f_{m+1} & (1, 1, \dots, 1, 1, 1, 1).
\end{array}$$

For  $1 \leq i \leq m-1$ , the number of s having the form  $f_i$  is equal to  $|\{n \in A_{2k} : \max CG(n) \leq F_{2k+2m-2i}\}|$ , which, by the inductive hypothesis, is

$$\sum_{j=0}^{m-i} |\{n \in A_{2k} : \max \mathcal{CG}(n) = F_{2k+2j}\}| = 2 + \sum_{j=1}^{m-i} F_{2j+2} = F_{2m-2i+3}.$$

The number of s having the form  $f_m$  and  $f_{m+1}$  is 2 and 1, respectively. Therefore,

$$|\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_2(n) = F_{2k+2m+2}\}| = \sum_{i=1}^{m-1} F_{2m-2i+3} + 2 + 1$$
$$= F_4 + F_5 + \dots + F_{2m+1}$$
$$= F_{2m+2}.$$

This completes our proof.

The proof of Lemma 3.3 also reveals the counts of numbers having  $F_{2k+2\ell}$  and  $2F_{2k+2\ell}$  as the maximum summand in their decomposition.

## **Corollary 3.4.** *For* $\ell \geq 1$ *, we have*

$$|\{n \in A_{2k} : \max C\mathcal{G}(n) = \max C\mathcal{G}_1(n) = F_{2k+2\ell}\}| = F_{2\ell+1}, \text{ and} \\ |\{n \in A_{2k} : \max C\mathcal{G}(n) = \max C\mathcal{G}_2(n) = F_{2k+2\ell}\}| = F_{2\ell}.$$

We now show that the numbers in  $\{n \in A_{2k} : \max CG(n) = \max CG_2(n) = F_{2k+2\ell}\}$  must lie below (are larger than) the numbers in  $\{n \in A_{2k} : \max CG(n) = \max CG(n) = \max CG_1(n) = F_{2k+2\ell}\}$ .

Lemma 3.5. For 
$$\ell \geq 0$$
,

$$\min\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_2(n) = F_{2k+2\ell}\}$$
  
> 
$$\max\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_1(n) = F_{2k+2\ell}\}.$$

*Proof.* When  $\ell = 0$ , we are comparing  $F_{2k}$  and  $2F_{2k}$ , and the lemma is obviously true. Suppose that  $\ell \ge 1$ . Since  $\{n \in A_{2k} : \max CG(n) = \max CG_1(n) = F_{2k+2\ell}\}$  is formed

by adding  $F_{2k+2\ell}$  to each number in  $\{n \in A_{2k} : \max CG(n) \leq F_{2k+2\ell-2}\}$ , Lemma 3.1 gives

$$\max\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_1(n) = F_{2k+2\ell}\} = F_{2k} + F_{2k+2\ell} + \dots + F_{2k+2\ell-4} + 2F_{2k+2\ell-2} + F_{2k+2\ell}.$$

On the other hand,

$$\min\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_2(n) = F_{2k+2\ell}\} = F_{2k} + 2F_{2k+2\ell}$$

We need only to verify that

$$F_{2k} + 2F_{2k+2\ell} > F_{2k} + F_{2k+2} + \dots + F_{2k+2\ell-4} + 2F_{2k+2\ell-2} + F_{2k+2\ell}.$$

Equivalently,

$$F_{2k+1} + F_{2k+2\ell} > F_{2k+1} + F_{2k+2} + \dots + F_{2k+2\ell-4} + 2F_{2k+2\ell-2}$$

which is true because the right side of the inequality is equal to  $F_{2k+2\ell}$ .

We have used in the proof of Lemmas 3.3 and 3.5 the fact that for  $\ell \ge 1$ ,

$$\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_1(n) = F_{2k+2\ell}\}$$
$$= \{n \in A_{2k} : \max \mathcal{CG}(n) \le F_{2k+2\ell-2}\} + F_{2k+2\ell}.$$

In other words, integers in  $\{n \in A_{2k} : \max CG(n) = \max CG_1(n) = F_{2k+2\ell}\}$  are formed by adding  $F_{2k+2\ell}$  to the all previous rows in Table 1. We now describe how to form integers in  $\{n \in A_{2k} : \max CG(n) = \max CG_2(n) = F_{2k+2\ell}\}$ .

**Lemma 3.6.** For  $\ell \geq 1$ , we have

$$\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_2(n) = F_{2k+2\ell}\}\$$
$$= \left\{n \in A_{2k} : n \le \sum_{i=0}^{\ell-1} F_{2k+2i}\right\} + 2F_{2k+2\ell}.$$

*Proof.* By Lemma 3.1, the largest integer  $n \in A_{2k}$  with  $\max CG(n) = \max CG_2(n) = F_{2k+2\ell}$  is

$$\sum_{i=0}^{\ell-1} F_{2k+2i} + 2F_{2k+2\ell};$$

hence,

$$\{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_2(n) = F_{2k+2\ell} \}$$
$$\subset \left\{ n \in A_{2k} : n \le \sum_{i=0}^{\ell-1} F_{2k+2i} \right\} + 2F_{2k+2\ell}.$$

We prove the reverse inclusion. Pick  $n \in A_{2k}$  such that

$$n \leq \sum_{i=0}^{\ell-1} F_{2k+2i}.$$
 (3.1)

Write the Chung-Graham decomposition of n as

$$n = c_0 F_{2k} + c_1 F_{2k+2} + \dots + c_{\ell-1} F_{2k+2\ell-2},$$

for  $c_i \in \{0, 1, 2\}$ . Suppose, for a contradiction, that

$$c_0 F_{2k} + c_1 F_{2k+2} + \dots + c_{\ell-1} F_{2k+2\ell-2} + 2F_{2k+2\ell}$$

is not a Chung-Graham decomposition. Then there is  $0 \le j \le \ell - 1$  such that  $c_j = 2$ and  $c_i = 1$  for all  $i \in [j + 1, \ell - 1]$ . It follows that

$$n \ge 2F_{2k+2j} + \sum_{i=j+1}^{\ell-1} F_{2k+2i}$$
(3.2)

From (3.1) and (3.2),

$$\sum_{i=0}^{\ell-1} F_{2k+2i} \geq 2F_{2k+2j} + \sum_{i=j+1}^{\ell-1} F_{2k+2i}.$$

Equivalently,

$$\sum_{i=0}^{j} F_{2k+2i} \geq 2F_{2k+2j},$$

which is a contradiction. Hence,

$$c_0 F_{2k} + c_1 F_{2k+2} + \dots + c_{\ell-1} F_{2k+2\ell-2} + 2F_{2k+2\ell}$$

is a Chung-Graham decomposition. Therefore,

$$n + 2F_{2k+2\ell} \in \{n \in A_{2k} : \max \mathcal{CG}(n) = \max \mathcal{CG}_2(n) = F_{2k+2\ell}\}.$$

This completes our proof.

# 4. PROOF OF THE MAIN THEOREM

Using what we know about Table 1 from Section 3, we now prove an identity that equates the difference between two earlier consecutive rows with the difference between two later consecutive rows in the table.

**Proposition 4.1.** Fix 
$$\ell \ge 1$$
. For  $1 + F_{2\ell+1} \le j \le F_{2\ell+3} - 1$ , we have  
 $q(j+1) - q(j) = q(j - F_{2\ell+1} + 1) - q(j - F_{2\ell+1}).$  (4.1)

*Proof.* Choose  $\ell \geq 1$ . By Lemma 3.3, the numbers in  $\{n : \max CG(n) \leq F_{2k+2\ell-2}\}$  lie from the 1<sup>st</sup> row to the  $F_{2\ell+1}$ <sup>th</sup> row, inclusively. Since

$$\{n : \max \mathcal{CG}(n) \le F_{2k+2\ell-2}\} + F_{2k+2\ell}$$
  
=  $\{n : \max \mathcal{CG}(n) = \max \mathcal{CG}_1(n) = F_{2k+2\ell}\},\$ 

we know that

$$q(i+1) - q(i) = q(i + F_{2\ell+1} + 1) - q(i + F_{2\ell+1})$$
(4.2)

whenever  $1 \le i \le F_{2\ell+1} - 1$ . Applying the change of variable  $j = i + F_{2\ell+1}$ , we obtain  $q(j+1) - q(j) = q(j - F_{2\ell+1} + 1) - q(j - F_{2\ell+1}), 1 + F_{2\ell+1} \le j \le 2F_{2\ell+1} - 1$ . (4.3) Furthermore, by Corollary 3.4 and Lemmas 3.5 and 3.6,

$$q(i+1) - q(i) = q(i+2F_{2\ell+1}+1) - q(i+2F_{2\ell+1}), 1 \le i \le F_{2\ell} - 1.$$
(4.4)

From (4.2) and (4.4),

$$\begin{split} q(i+F_{2\ell+1}+1)-q(i+F_{2\ell+1}) \ &= \ q(i+2F_{2\ell+1}+1)-q(i+2F_{2\ell+1}), 1 \le i \le F_{2\ell}-1. \\ \text{Using the change of variable } j = i+2F_{2\ell+1}, \text{ we have} \end{split}$$

$$q(j+1) - q(j) = q(j - F_{2\ell+1} + 1) - q(j - F_{2\ell+1}), 1 + 2F_{2\ell+1} \le j \le F_{2\ell+3} - 1.$$
(4.5)

Thanks to (4.3) and (4.5), it remains to verify (4.1) when  $j = 2F_{2\ell+1}$ ; that is,

$$q(2F_{2\ell+1}+1) - q(2F_{2\ell+1}) = q(F_{2\ell+1}+1) - q(F_{2\ell+1}).$$
(4.6)

By Lemmas 3.1, 3.3,3.5, and Corollary 3.4,

$$q(F_{2\ell+1}) = \sum_{i=0}^{\ell-2} F_{2k+2i} + 2F_{2k+2\ell-2},$$
  

$$q(F_{2\ell+1}+1) = F_{2k} + F_{2k+2\ell},$$
  

$$q(2F_{2\ell+1}) = F_{2k} + F_{2k+2} + \dots + F_{2k+2\ell-4} + 2F_{2k+2\ell-2} + F_{2k+2\ell},$$
  

$$q(2F_{2\ell+1}+1) = F_{2k} + 2F_{2k+2\ell}.$$

These confirm (4.6), and we are done.

We are now ready to prove the following key lemma to describe all integers in  $A_{2k}$ .

## **Lemma 4.2.** *For* $j \ge 2$ *,*

$$q(j+1) - q(j) = \begin{cases} F_{2k} \text{ if the } (j-1)^{th} \text{ letter of } \mathcal{S} \text{ is } A, \\ F_{2k+1} \text{ if the } (j-1)^{th} \text{ letter of } \mathcal{S} \text{ is } B. \end{cases}$$

$$(4.7)$$

*Proof.* It suffices to prove that (4.7) is true for all  $j \leq F_{2m+1} - 1$  for any arbitrary  $m \in \mathbb{N}$ . We do so by induction.

Base case: for m = 3, we can see from Table 1 that (4.7) is true for all  $j \le F_7 - 1$ . Inductive hypothesis: suppose that (4.7) is true for  $j \le F_{2m+1} - 1$  for some  $m \ge 3$ .

We need to show that it is true for all  $j \le F_{2m+3} - 1$ . We proceed by a case analysis. Case 1:  $j = F_{2m+1}$ . By Lemmas 3.1 and 3.3, we have

$$q(F_{2m+1}+1) - q(F_{2m+1}) = F_{2k} + F_{2k+2m} - (F_{2k} + \dots + F_{2k+2m-4} + 2F_{2k+2m-2})$$
  
=  $F_{2k+1}$ .

By Property f) in Section 2, (4.7) is true when  $j = F_{2m+1}$ .

Case 2:  $j = F_{2m+1} + 1$ . By Lemma 3.3,

$$q(F_{2m+1}+2) - q(F_{2m+1}+1) = (2F_{2k} + F_{2k+2m}) - (F_{2k} + F_{2k+2m}) = F_{2k}.$$

By Property f) in Section 2, (4.7) is true when  $j = F_{2m+1} + 1$ .

Case 3:  $F_{2m+1} + 2 \le j \le 2F_{2m+1} - 1$ . It follows from Proposition 4.1 that

$$q(j+1) - q(j) = q(j+1 - F_{2m+1}) - q(j - F_{2m+1})$$

Thanks to Proposition 2.1 and the fact that

$$j - F_{2m+1} \leq 2F_{2m+1} - 1 - F_{2m+1} = F_{2m+1} - 1,$$

the inductive hypothesis guarantees that (4.7) is true for  $F_{2m+1} + 2 \le j \le 2F_{2m+1} - 1$ .

Case 4:  $j = 2F_{2m+1}$ . It follows from Lemmas 3.1, 3.3, 3.5, and Corollary 3.4 that

$$q(2F_{2m+1}+1) - q(2F_{2m+1})$$
  
=  $(F_{2k} + 2F_{2k+2m}) - (F_{2k} + \dots + 2F_{2k+2m-2} + F_{2k+2m}) = F_{2k+1}$ 

Property f) in Section 2 confirms that (4.7) is true for  $j = 2F_{2m+1}$ .

Case 5:  $j = 2F_{2m+1} + 1$ . By Lemmas 3.3, 3.5, and Corollary 3.4,

$$q(2F_{2m+1}+2) - q(2F_{2m+1}+1) = (2F_{2k}+2F_{2k+2m}) - (F_{2k}+2F_{2k+2m}) = F_{2k}.$$

Property f) in Section 2 confirms that (4.7) is true for  $j = 2F_{2m+1} + 1$ .

Case 6:  $2F_{2m+1} + 2 \le j \le F_{2m+3} - 1$ . According to Proposition 4.1,

$$q(j+1) - q(j) = q(j+1-2F_{2m+1}) - q(j-2F_{2m+1}).$$

Since

$$j - 2F_{2m+1} \leq F_{2m+3} - 1 - 2F_{2m+1} = F_{2m} - 1,$$

the inductive hypothesis can be applied. Together with Proposition 2.3, we know that (4.7) holds for  $2F_{2m+1} + 2 \le j \le F_{2m+3} - 1$ .

This completes our proof.

**Proposition 4.3.** *For*  $k \ge 1$ *, we have* 

$$A_{2k} = \left\{ F_{2k}, (n+2)F_{2k} + \left\lfloor \frac{n+1}{\phi} \right\rfloor F_{2k-1} : n \ge 0 \right\}.$$

Proof. By Lemma 4.2, we have

$$A_{2k} = \{F_{2k}, 2F_{2k} + a(n)F_{2k} + b(n)F_{2k+1} : n \ge 0\},\$$

where a(n) and b(n) are the number of A's and B's, respectively, among the first n letters of S. Due to (2.1),

$$A_{2k} = \left\{ F_{2k}, 2F_{2k} + \left(n - \left\lfloor \frac{n+1}{\phi} \right\rfloor \right) F_{2k} + \left\lfloor \frac{n+1}{\phi} \right\rfloor F_{2k+1} : n \ge 0 \right\}$$
$$= \left\{ F_{2k}, (n+2)F_{2k} + \left\lfloor \frac{n+1}{\phi} \right\rfloor F_{2k-1} : n \ge 0 \right\}.$$

*Proof of Theorem 1.3.* For  $1 \le N < k$ , the set of integers whose Chung-Graham decomposition has  $F_{2k}$  but none of  $F_{2N}, \ldots, F_{2k-2}$  is

$$\left\{j + F_{2k}, j + (n+2)F_{2k} + \left\lfloor \frac{n+1}{\phi} \right\rfloor F_{2k-1} : 0 \le j \le F_{2N} - 1, n \ge 0 \right\}.$$

Indeed, all Chung-Graham decompositions of the form  $\sum_{i=1}^{N-1} c_i F_{2i}$ , when added to an integer in  $A_{2k}$ , gives an integer whose Chung-Graham decomposition has  $F_{2k}$  but none of  $F_{2N}, \ldots, F_{2k-2}$ . Meanwhile,

$$\left\{\sum_{i=1}^{N-1} c_i F_{2i} : c_i$$
's satisfy the Chung-Graham decomposition 
$$\right\} = [0, F_{2N} - 1].$$

Therefore,

$$B_{2N} := [1, F_{2N} - 1] \cup \bigcup_{k=N+1}^{\infty} \left\{ j + F_{2k}, j + (n+2)F_{2k} + \left\lfloor \frac{n+1}{\phi} \right\rfloor F_{2k-1} : 0 \le j \le F_{2N} - 1, n \ge 0 \right\}.$$

#### REFERENCES

- P. Baird-Smith, A. Epstein, K. Flint, and S. J. Miller, The generalized Zeckendorf game, *Fibonacci Quart.* 57 (2019), 1–14.
- [2] P. Baird-Smith, A. Epstein, K. Flint, and S. J. Miller, The Zeckendorf game, Springer Proc. Math. Stat. 297 (2020), 25–38.
- [3] C. Ballot, On Zeckendorf and base b digit sums, Fibonacci Quart. 51 (2013), 319-325.
- [4] L. Carlitz, V. E. Hoggatt, Jr., and R. Scoville, Fibonacci representations, *Fibonacci Quart.* 10 (1972), 1–28.
- [5] M. Catral, P. Ford, P. Harris, S. J. Miller, and D. Nelson, Generalizing Zeckendorf's Theorem: The Kentucky Sequence, *Fibonacci Quart.* 52 (2014), 68–90.
- [6] S. Chang, Average number of Zeckendorf integers, J. Number Theory 186 (2018), 452–472.
- [7] H. V. Chu, On generalized Zeckendorf decompositions and generalized golden strings, *Fibonacci Quart.* 59 (2021), 254–261.
- [8] H. V. Chu, D. C. Luo, and S. J. Miller, On Zeckendorf related partitions using the Lucas sequence, *Fibonacci Quart.* 60 (2022), 111-119.
- [9] F. R. K. Chung and R. L. Graham, On irregularities of distribution, *Colloq. Math. Soc. János Bolyai* 37 (1984), 181–222.
- [10] D. E. Daykin, Representation of natural numbers as sums of generalised Fibonacci numbers, J. London Math. Soc. 35 (1960), 143–160.
- [11] D. E. Daykin, Representation of natural numbers as sums of generalized Fibonacci numbers. II, *Fibonacci Quart.* 7 (1969), 494–510.
- [12] M. F. Dekking, The sum of digits functions of the Zeckendorf and the base phi expansions, *Theoret. Comput. Sci.* 859 (2021), 70–79.
- [13] M. F. Dekking, The structure of Zeckendorf expansions, Integers 21 (2021), 10 pp.
- [14] P. Demontigny, T. Do, A. Kulkarni, S. J. Miller, D. Moon, and U. Varma, Generalizing Zeckendorf's Theorem to *f*-decompositions, *J. Number Theory* 141 (2014), 136–158.
- [15] R. Dorward, P. L. Ford, E. Fourakis, P. E. Harris, S. J. Miller, E. Palsson, and H. Paugh, A generalization of Zeckendorf's theorem via circumscribed *m*-gons, *Involve* 10 (2017), 125–150.
- [16] M. Drmota and M. Skałba, The parity of the Zeckendorf sum-of-digits function, *Manuscripta Math.* 101 (2000), 361–383.
- [17] M. Griffiths, Digit proportions in Zeckendorf representations, *Fibonacci Quart.* 48 (2010), 168– 174.
- [18] M. Griffiths, The golden string, Zeckendorf representations and the sum of a series, Am. Math. Mon. 118 (2011), 497–507.
- [19] M. Griffiths, Fixed-term Zeckendorf representations, Fibonacci Quart. 52 (2014), 331–335.
- [20] N. Hamlin and W. A. Webb, Representing positive integers as a sum of linear recurrence sequences, *Fibonacci Quart.* 50 (2012), 99–105.
- [21] Y. Hosten, On the variation of the sum of digits in the Zeckendorf representation: an algorithm to compute the distribution and mixing properties, *J. Integer Seq.* **27** (2024), 48 pp.
- [22] C. Kimberling, One-Free Zeckendorf sums, Fibonacci Quart. 21 (1983), 53–57.
- [23] R. Li, X. Li, S. J. Miller, C. Mizgerd, C. Sun, D. Xia, and Z. Zhou, Deterministic Zeckendorf games, *Fibonacci Quart.* 58 (2020), 152–160.
- [24] T. Martinez, S. J. Miller, C. Mizgerd, and C. Sun, Generalizing Zeckendorf's theorem to homogeneous linear recurrences, I, *Fibonacci Quart.* 60 (2022), 222–230.

- [25] T. Martinez, S. J. Miller, C. Mizgerd, J. Murphy, and C. Sun, Generalizing Zeckendorf's theorem to homogeneous linear recurrences, II, *Fibonacci Quart.* 60 (2022), 231–254.
- [26] S. J. Miller, E. Sosis, and J. Ye, Winning strategies for generalized Zeckendorf games, *Fibonacci Quart.* 60 (2022), 270–292.
- [27] A. Shutov, On the sum of digits of the Zeckendorf representations of two consecutive numbers, *Fibonacci Quart.* **58** (2020), 203–207.
- [28] E. Zeckendorf, Representation des nombres naturels par une somme de nombres de Fibonacci ou de nombres de Lucas, *Bull. Soc. Roy. Sci. Liege* **41** (1972), 179–182.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA

*Email address*: hungchu1@tamu.edu

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA

Email address: aneykanji\_tamu@tamu.edu

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA

*Email address*: zachary.l.v@tamu.edu