

A comparison of three kinds of monotonic proof-theoretic semantics and the base-incompleteness of intuitionistic logic

Antonio Piccolomini d’Aragona

University of Tübingen, Tübingen, Germany
antonio.piccolomini-daragona@uni-tuebingen.de

Abstract

I deal with two approaches to proof-theoretic semantics: one based on argument structures and justifications, which I call *reducibility semantics*, and one based on consequence among (sets of) formulas over atomic bases, called *base semantics*. The latter splits in turn into a *standard reading*, and a variant of it put forward by Sandqvist. I prove some results which, when suitable conditions are met, permit one to shift from one approach to the other, and I draw some of the consequences of these results relative to the issue of completeness of (recursive) logical systems with respect to proof-theoretic notions of validity. This will lead me to focus on a notion of base-completeness, which I will discuss with reference to known completeness results for intuitionistic logic. The general interest of the proposed approach stems from the fact that reducibility semantics can be understood as a labelling of base semantics with proof-objects typed on (sets of) formulas for which a base semantics consequence relation holds, and which witness this very fact. Vice versa, base semantics can be understood as a type-abstraction of a reducibility semantics consequence relation obtained by removing the witness of the fact that this relation holds, and by just focusing on the input and output type of the relevant proof-object.

Keywords Proof-theoretic semantics, proof, completeness, base-completeness, intuitionism

1 Introduction

The name *proof-theoretic semantics* (PTS for short) indicates a family of constructive semantics whose core notion for explaining meaning and for defining the notions of (logical) validity is not that of truth, as happens in model theory, but that of proof; the meaning of the non-logical vocabulary is accordingly fixed, not via model-theoretic mappings from the language onto suitable (typically set-theoretic) structures, but through sets of (sets of) rules governing deduction at the atomic level—see [25] for an overview.

PTS stems from Prawitz’s work in proof theory, in particular from Prawitz’s normalisation theorems for Gentzen’s Natural Deduction [3, 17]. The first version of PTS is due to Prawitz himself; in this formulation, PTS is based on *argument structures* and *reductions* [18, 19], so I will call it *reducibility semantics*. Later on—maybe starting from the influential [24]—argument structures were left aside, and PTS became a theory of consequence for formulas over sets of (sets of) atomic rules; the constructivist burden was put entirely on such sets, and this is why—partly following [21]—I will indicate this approach as *standard base semantics*. A variant of base semantics was provided by Sandqvist [21], whence I shall call it *Sandqvist’s base semantics*. It differs from the standard reading—and from Prawitz’s original approach—in that it deals with disjunction in an elimination-like way, rather than in an introduction-based fashion.

Both reducibility semantics and base semantics (in its two variants) are expected to be semantics for constructive logics, and many completeness and incompleteness results have been obtained so far—see [14] for an overview, whereas more recent results can be found in [16, 27, 28]. Here, I prove some general results on the relation between reducibility semantics and base semantics, and use them to obtain further insights on completeness and incompleteness issues. In particular, I shall be concerned by the conditions or the extent under which one is allowed to go from a consequence relation (possibly relative to specific formulas and over a specific “model”) in one of the proof-theoretic semantics versions at issue here, to the same consequence in another version.

The interest of comparing the three approaches is twofold. As regards the relation between reducibility semantics and base semantics in the standard reading, both are ultimately based on the idea that the meaning of a logical constant

κ should be given by the conditions for *introducing* κ in formulas—thus coping with BHK semantics [29] and Gentzen’s semantic insight about introduction rules of Natural Deduction [3]. However, as already said above, Prawitz’s picture is at first sight richer, since there the constructivist spirit is expressed, not only through the kind of “models” which the notion of consequence is defined over, but additionally through “witnesses” for the consequence relation itself—i.e., A being a consequence of Γ is *defined* as existence of a *valid argument* from Γ to A . As also remarked above, standard base semantics instead drops valid arguments out, and one might at that point wonder whether the original constructivist spirit of Prawitz’s approach is respected. If the consequence relation is no longer “witnessed” by suitable proof-objects, we are left with “models” of a special kind only, so we may ask whether some “degrees of constructiveness” got lost in the pruning. I shall prove below that, to a *large* extent, nothing is lost, and hence that Prawitz’s semantics, when some constraints are met, is in fact equivalent to standard base semantics. It follows that all the (mostly in)completeness results which hold for the former apply to a certain understanding of the latter as well. This is admittedly not a major result, as it can be looked at as nothing but a “decoration” of the general theorems established in [16], and was stated by [10] in the context of Prawitz’s *theory of grounds* [20]—for the non-monotonic approach, but easily extendable to the monotonic one. However, this “decoration” involves steps which show that, in the specific framework of reducibility semantics, a detailed proof involves some not-so-trivial aspects.

The second point of interest touches upon the relation between Prawitz’s reducibility semantics and Sandqvist’s base semantics—hence, given the equivalence between the former and base semantics in the standard reading, also between standard base semantics and its Sandqvist variant. Since Sandqvist’s base semantics does without argument structures and reductions too, one may have also here issues about dropping out the “witnesses” of consequence that one had in Prawitz. But there is more than that. The elimination-based approach to disjunction employed by Sandqvist induces a structural difference with respect to Prawitz’s introduction-based approach, which triggers intuitionistic completeness in some most relevant cases, *contra* the provable intuitionistic incompleteness that we have for (base semantics in the standard reading, hence for) Prawitz’s reducibility semantics (see below for details). Given this (and given, incidentally, the acknowledged *harmony* between introduction and elimination rules in Natural Deduction), it seems thus to be worth exploring to what extent Prawitz’s reducibility semantics and base semantics in Sandqvist’s reading can be connected to each other. Below, I shall prove that the two approaches *can* be compared, although only on a “global” scale, and that this, together with the completeness-incompleteness mismatch, implies that they *are not* comparable at the level of “models”—meaning that the models verifying certain pairs (Γ, A) in one approach are not always models verifying the same pairs in the other, and vice versa. The “global” comparability also provides a sufficient condition for equivalence to hold between Prawitz’s and Sandqvist’s pictures at the level of logical validity. This may be of interest since, e.g., while intuitionistic logic (IL for short) is incomplete over standard base semantics (and hence, via the equivalence mentioned above, over reducibility semantics too) relative to “models” of *any* kind (more precisely, as we shall see below, relative to atomic bases of either limited or unlimited complexity), IL is instead complete over Sandqvist’s base semantics relative to “models” of a specific kind (in particular, it is not known what happens with atomic bases of level ≤ 1).

The result about the sufficient condition for equivalence between Prawitz’s and Sandqvist’s approaches at the level of logical validity requires introducing a notion of “point-wise” soundness and completeness of given logics Σ relative to proof-theoretic semantics. Here “point-wise” means, roughly, that validity over a model is implied (soundness) or implies (completeness) derivability in Σ plus the model (a notion which makes sense given, as we shall see, the “deductive” nature of models in proof-theoretic semantics). Besides illuminating the relation between Prawitz’s and Sandqvist’s frameworks, the notions of “point-wise” soundness and completeness will be shown to be of interest in themselves. In this connection, I shall prove a result of “point-wise” incompleteness (with respect to all the three proof-theoretic frameworks at issue here) for a class of super-intuitionistic logics, which includes IL.

The structure of the paper is as follows. By limiting myself to a propositional language, I start with an overview of atomic rules (Section 2). Then I provide an outline of base semantics and reducibility semantics (Section 3). In Section 4, I prove the general results mentioned above and draw some consequences from them, both relative to the general issue of completeness of given (recursive) systems, and relative to a notion of base-completeness (Section 5).

2 Language and atomic bases

Definition 1. *The language \mathcal{L} is given by the grammar*

$$X := p \mid \perp \mid X \wedge X \mid X \vee X \mid X \rightarrow X$$

where \perp is a constant atom for absurdity and $\neg X := X \rightarrow \perp$.

I will use the following notation:

- $\text{ATOM}_{\mathcal{L}}$ is the set of the atomic formulas of \mathcal{L} , i.e., $\{p_i \mid i \in \mathbb{N}\} \cup \{\perp\}$;
- $\text{FORM}_{\mathcal{L}}$ is the set of the formulas of \mathcal{L} ;
- A, B, C, \dots indicate arbitrary formulas;
- $\Gamma, \Delta, \Theta, \dots$ indicate arbitrary sets of formulas. It is important to remark that the latter will be *always* assumed to be *finite*. This is to avoid some issues, pointed out by [27], concerning properties of compactness, monotonicity, and compact monotonicity of the consequence relations in the comparison of reducibility semantics and base semantics. The limitation to finite sets of formulas will be on the other hand sufficient for raising my points—in particular, for transferring completeness or incompleteness results from one approach to the other.

I now define the notion of atomic base over \mathcal{L} . Atomic bases, however, require a preliminary definition of the notion of atomic rule over \mathcal{L} . The definition of atomic rules is by induction on what [15] calls the level of atomic rules.

Definition 2. Any atom is an atomic rule of level 0. An atomic rule of level 1 is

$$\frac{A_1 \quad \dots \quad A_n}{A}$$

with $A_1, \dots, A_n, A \in \text{ATOM}_{\mathcal{L}}$. Given sets of atomic rules $\mathfrak{R}_1, \dots, \mathfrak{R}_n$ whose maximal level is $\kappa - 1 \geq 0$, where each \mathfrak{R}_i may be empty ($i \leq n$), we say that

$$\frac{[\mathfrak{R}_1] \quad \dots \quad [\mathfrak{R}_n]}{A_1 \quad \dots \quad A_n} \frac{}{A}$$

with $A_1, \dots, A_n, A \in \text{ATOM}_{\mathcal{L}}$ is an atomic rule of level $\kappa + 1$.

Square brackets in Definition 2 indicate that the rule discharges lower level atomic rules in its premises. Thus, an atomic rule of level 3 has the form

$$\frac{[\mathfrak{R}_{1,1}] \quad \dots \quad [\mathfrak{R}_{1,m_1}] \quad \dots \quad [\mathfrak{R}_{n,1}] \quad \dots \quad [\mathfrak{R}_{n,m_n}]}{A_1 \quad \dots \quad A_n} \frac{}{A}$$

where, for every $i \leq n, j \leq m_i$, either $R_{j,m_i} = B \in \text{ATOM}_{\mathcal{L}}$, or

$$R_{j,m_i} = \frac{B_1 \quad \dots \quad B_s}{B}$$

with $B_1, \dots, B_s, B \in \text{ATOM}_{\mathcal{L}}$.

I will adopt the convention that atomic bases are—borrowing the terminology of [15]—of an *intuitionistic* kind, namely, that they contain a rule of *atomic explosion* for every atom in the language.

Convention 1. Every \mathfrak{B} contains the rules

$$\frac{\perp}{A} \text{AtExp}$$

for every $A \in \text{ATOM}_{\mathcal{L}}$.

Definition 3. An atomic base of level n is a set of atomic rules $\mathfrak{B} = \{\text{AtExp}, R_1, \dots, R_n\}$ with $\max\{\mathfrak{L}(R_1), \dots, \mathfrak{L}(R_n)\} = n$, where $\mathfrak{L}(R_i)$ indicates the level of R_i ($i \leq n$, see Convention 1 for *AtExp*).

I will use the following notation:

- \mathfrak{B}^n indicates that \mathfrak{B} has level n ;

- $\mathbb{B}^n := \{\mathfrak{B}^m \mid m \leq n\}$.

Whenever it is possible, I will omit the indication of the level of an atomic base or of a set of atomic bases. The empty atomic base, written \mathfrak{B}^0 , is assumed to be the atomic base which only contains AtExp . Observe that, as per Definition 3, the latter is disregarded when counting the level of the atomic base, so \mathfrak{B}^0 has level 0—i.e., $\max(\emptyset) = 0$.

Definition 4. The derivations-set $DER_{\mathfrak{B}}$ of \mathfrak{B} is defined inductively as follows:

- any $A \in \text{ATOM}_{\mathcal{L}}$ is a single-node derivation in $DER_{\mathfrak{B}}$. It is a derivation of A from \emptyset if the node applies an axiom $A \in \mathfrak{B}$, or a derivation of A from A if A is assumed as an atomic rule (whether or not $A \in \mathfrak{B}$);
- if the following are derivations in $DER_{\mathfrak{B}}$,

$$\begin{array}{c} \mathfrak{C}_i, \mathfrak{R}_i \\ \mathcal{D}_i \\ A_i \end{array}$$

($i \leq n$) where \mathfrak{C}_i and \mathfrak{R}_i are sets of atomic rules and $A_i \in \text{ATOM}_{\mathcal{L}}$ is the premise of an atomic rule R of the form

$$\frac{\begin{array}{ccc} [\mathfrak{R}_1] & & [\mathfrak{R}_n] \\ A_i & \dots & A_n \end{array}}{A} R$$

then

$$\frac{\begin{array}{ccc} \mathfrak{C}_1, [\mathfrak{R}_1] & & \mathfrak{C}_n, [\mathfrak{R}_n] \\ \mathcal{D}_1 & & \mathcal{D}_n \\ A_i & \dots & A_n \end{array}}{A} R$$

is a derivation of A from $\mathfrak{C}_1, \dots, \mathfrak{C}_n$ in $DER_{\mathfrak{B}}$ if $R \in \mathfrak{B}$, or a derivation of A from $\mathfrak{C}_1, \dots, \mathfrak{C}_n, R$ if $R \notin \mathfrak{B}$.

Definition 5. A is derivable from \mathfrak{K} in \mathfrak{B} —written $\mathfrak{K} \vdash_{\mathfrak{B}} A$ —iff there is $\mathcal{D} \in DER_{\mathfrak{B}}$ from \mathfrak{K} to A .

Definition 6. \mathfrak{C} is an extension of \mathfrak{B} iff $\mathfrak{B} \subseteq \mathfrak{C}$.

I use the following notation: $\mathfrak{C} \supseteq_n \mathfrak{B}$ indicates that $\mathfrak{C} \supseteq \mathfrak{B}$ and $\mathfrak{C} \in \mathbb{B}^n$.

Proposition 1. $\{\mathfrak{B} \mid \mathfrak{B} \supseteq_n \mathfrak{B}^0\} = \mathbb{B}^n$.

3 Base semantics and reducibility semantics

Since I am restricting myself to propositional logic, I will occasionally allow myself to indicate the universal and existential meta-quantifiers by the usual object-linguistic notations \forall and \exists .

3.1 Base semantics

Definition 7. That A is a consequence of Γ on \mathfrak{B}^m of level n in base semantics is indicated by $\Gamma \models_{\mathfrak{B}^m, n} A$. It holds iff $m \leq n$ and

1. $\Gamma = \emptyset \implies$

- $A \in \text{ATOM}_{\mathcal{L}} \implies \vdash_{\mathfrak{B}^m} A$;
- $A = B \wedge C \implies \vdash_{\mathfrak{B}^m, n} B$ and $\vdash_{\mathfrak{B}^m, n} C$;
- $A = B \vee C \implies \vdash_{\mathfrak{B}^m, n} B$ or $\vdash_{\mathfrak{B}^m, n} C$;
- $A = B \rightarrow C \implies B \vdash_{\mathfrak{B}^m, n} C$;

$$2. \Gamma \neq \emptyset \implies \forall \mathcal{C} \supseteq_n \mathfrak{B}^m (\models_{\mathcal{C}, n} \Gamma \implies \models_{\mathcal{C}, n} A)$$

where $\models_{\mathcal{C}, n} \Gamma$ means that $\forall B \in \Gamma (\models_{\mathcal{C}, n} B)$.

A variant of base semantics was introduced by Sandqvist [21]. In the present framework, it runs as follows.

Definition 8. *That A is a consequence of Γ on \mathfrak{B}^m of level n in base semantics in Sandqvist sense is indicated by $\Gamma \models_{\mathfrak{B}^m, n}^s A$. It holds iff $m \leq n$ and*

$$1. \Gamma = \emptyset \implies$$

$$(a) A \in \text{ATOM}_{\mathcal{L}} \implies \vdash_{\mathfrak{B}^m} A;$$

$$(b) A = B \wedge C \implies \models_{\mathfrak{B}^m, n}^s A \text{ and } \models_{\mathfrak{B}^m, n}^s B;$$

$$(c) A = B \vee C \implies \forall \mathcal{C} \supseteq_n \mathfrak{B}^m \forall D \in \text{ATOM}_{\mathcal{L}} (B \models_{\mathcal{C}, n}^s D \text{ and } C \models_{\mathcal{C}, n}^s D \implies \models_{\mathcal{C}, n}^s D);$$

$$(d) A = B \rightarrow C \implies B \models_{\mathfrak{B}^m, n}^s C;$$

$$2. \Gamma \neq \emptyset \implies \forall \mathcal{C} \supseteq_n \mathfrak{B} (\models_{\mathcal{C}, n} \Gamma \implies \models_{\mathcal{C}, n} A)$$

We remark that, in Sandqvist, \perp is understood as a *non-atomic* constant, so we should add a new sign to our language, say \perp^* , as distinguished from \perp . However, over intuitionistic bases, \perp^* and \perp are equivalent [15, but see also Proposition 2 below]. In what follows, \Vdash_n means \models_n or \models_n^s —possibly with an index for atomic bases.

Definition 9. *That A is a logical consequence of Γ of level n in base semantics (in Sandqvist sense) is indicated by $\Gamma \Vdash_n A$. It holds iff $\forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \Vdash_{\mathfrak{B}, n} A)$.*

Proposition 2. $\Vdash_{\mathfrak{B}, n} \perp \iff \forall A \in \text{ATOM}_{\mathcal{L}} (\Vdash_{\mathfrak{B}, n} A)$.

Proposition 3 (Monotonicity of \Vdash). $\Gamma \Vdash_{\mathfrak{B}, n} A \iff \forall \mathcal{C} \supseteq_n \mathfrak{B} (\Gamma \Vdash_{\mathcal{C}, n} A)$.

Proposition 4. $\Gamma \Vdash_n A \iff \Gamma \Vdash_{\mathfrak{B}^0, n} A$.

Corollary 1. $\Gamma \Vdash_n A \iff \forall \mathfrak{B} \in \mathbb{B}^n (\Vdash_{\mathfrak{B}, n} \Gamma \implies \Vdash_{\mathfrak{B}, n} A)$.

Proof. By Proposition 4, $\Gamma \Vdash_n A$ is tantamount to $\Gamma \Vdash_{\mathfrak{B}^0, n} A$ which in turn, by points 2 in Definition 7 or 8, is tantamount to $\forall \mathfrak{B} \supseteq_n \mathfrak{B}^0 (\Vdash_{\mathfrak{B}, n} \Gamma \implies \Vdash_{\mathfrak{B}, n} A)$. By Proposition 1, this is tantamount to $\forall \mathfrak{B} \in \mathbb{B}^n (\Vdash_{\mathfrak{B}, n} \Gamma \implies \Vdash_{\mathfrak{B}, n} A)$. \square

3.2 Reducibility semantics

As said, reducibility semantics is instead based on the notions of argument structure and reduction. The latter are then used for defining the notions of argumental validity over a base, and of argumental validity in general. The presentation below is mainly based on [17, 19], and partly on [23].

Definition 10. *An argument structure over \mathcal{L} is a pair $\langle T, \langle f, h, g \rangle \rangle$ such that*

- *T is a finite rooted tree (let $<$ denote the order relation over T) whose nodes are labelled by formulas of \mathcal{L} . Let us assume that the labels of the top-nodes of T are partitioned into two groups:*
 - *assumption-labels, which I indicate by T^{as} , and which can be formulas of any kind, and*
 - *atomic-axiom-labels, which I indicate by T^{ax} , and which can only be atoms;*
- *f is a function defined on some $\Gamma \subseteq T^{\text{as}}$ and is such that, $\forall \mu \in \Gamma$, it holds that $\mu < f(\mu)$;*
- *h is a function defined on some $H \subseteq T^{\text{ax}}$ and is such that, $\forall \mu \in H$, it holds that $\mu < h(\mu)$, $h(\mu)$ and all its children are labelled by atoms, and there is no $\nu \in \Gamma$ such that $f(\nu) = h(\mu)$;*
- *g is a function defined on some $J \subseteq \mathcal{P}(<)$ such that, for every $K \in J$,*
 - *K contains all and only the edges that link a given node μ to all its children, and*
 - *both μ and its children are labelled by atoms, and*
 - *there is no $\nu \in \Gamma$ such that $f(\nu) = \mu$, and*

- the function is such that $\forall v \in J$, it holds that $\mu < g(v)$, $g(v)$ and all its children are labelled by atoms, and there is no $\xi \in \Gamma$ such that $f(\xi) = g(v)$.

Definition 11. Given $\mathcal{D} = \langle T, \langle f, h, g \rangle \rangle$ where T has top-nodes $T^{\text{as}} = \Gamma$ and root A , the elements of Γ are the assumptions of \mathcal{D} and A is the conclusion of \mathcal{D} .

The intended meaning of T^{ax} in Definition 10 is that the atoms which label the given top-nodes in T are not assumptions, but axioms. The meaning of f, h, g in the same definition is as follows. In Natural Deduction terminology, they are discharge functions—see [17, 22, 24]. Top-nodes $\mu \in T^{\text{as}}$ in the domain of f are assumptions discharged throughout T , while top-nodes $\mu \in T^{\text{ax}}$ and sets of edges $\mu \in J$ in the domain of h, g respectively are assumed atomic rules discharged by an atomic rule. In all cases, the discharge takes place at the node $f(\mu)$.

Definition 12. \mathcal{D} is closed iff all its assumptions are discharged, and it is open otherwise.

Definition 13. Where Γ is the set of the undischarged assumptions of \mathcal{D} and A is the conclusion of \mathcal{D} , \mathcal{D} is an argument structure from Γ to A .

The notion of (immediate) sub-structure of \mathcal{D} can be defined as usual. The substitution of the sub-structure \mathcal{D}^* with the structure \mathcal{D}^{**} in \mathcal{D} —written $\mathcal{D}[\mathcal{D}^{**}/\mathcal{D}^*]$ —indicates the argument structure obtained from \mathcal{D} by replacing its sub-structure \mathcal{D}^* with the argument structure \mathcal{D}^{**} . Since $\mathcal{D}[\mathcal{D}^{**}/\mathcal{D}^*]$ might not be a sub-structure of \mathcal{D} , and since \mathcal{D} is defined as a tree plus discharge functions, when replacing \mathcal{D}^* with \mathcal{D}^{**} in \mathcal{D} one may need to re-define the discharge functions of \mathcal{D} , so that assumption formulas or assumed atomic rules discharged at some node μ in the tree of \mathcal{D} are discharged in $\mathcal{D}[\mathcal{D}^{**}/\mathcal{D}^*]$ —if they occur in it—at a node μ^* which μ is “mapped onto” in $\mathcal{D}[\mathcal{D}^{**}/\mathcal{D}^*]$, whereas some other discharges in \mathcal{D} might “disappear” in $\mathcal{D}[\mathcal{D}^{**}/\mathcal{D}^*]$. I assume that this rough description can be made suitably precise, but I shall abstract from this and only give one example. Take an argument structure

$$\mathcal{D} = \frac{\frac{\frac{[A]}{\mathcal{D}_1} \quad B}{B \vee C} \quad \frac{[B_1] \quad [B_2]}{\mathcal{D}_2} \quad D}{D} \quad \frac{[C]}{\mathcal{D}_3} \quad D}{\frac{D}{A \rightarrow D}}$$

Here, B_1 and B_2 are meant to indicate that the assumption B is used twice for deriving D through \mathcal{D}_2 . Then, the reduced form of \mathcal{D} is $\mathcal{D}[\vee_\rho(\mathcal{D}^*)/\mathcal{D}^*]$, where \mathcal{D}^* is the immediate sub-derivation of \mathcal{D} , and $\vee_\rho(\mathcal{D}^*)$ is the reduction of \mathcal{D}^* via the standard reduction for elimination of disjunction, i.e.,

$$\mathcal{D}[\vee_\rho(\mathcal{D}^*)/\mathcal{D}^*] = \frac{\frac{[A] \quad [A]}{\mathcal{D}_1} \quad B_1 \quad B_2}{\mathcal{D}_2} \quad D}{\frac{D}{A \rightarrow D}}$$

Thus, the red-to-brown discharge in \mathcal{D} is re-defined as the double red-to-brown discharge in $\mathcal{D}[\vee_\rho(\mathcal{D}^*)/\mathcal{D}^*]$, while the blue-to-purple discharges in \mathcal{D} “disappear” in the discharges of $\mathcal{D}[\vee_\rho(\mathcal{D}^*)/\mathcal{D}^*]$ —see [17] for more.

Definition 14. Given \mathcal{D} from $\Gamma = \{\mu_1, \dots, \mu_n\}$ to A and σ a function from and to argument structures such that $\sigma(\mu_i)$ is a (closed) argument structure with the same conclusion as μ_i ($i \leq n$), $\mathcal{D}^\sigma = \mathcal{D}[\sigma(\mu_1), \dots, \sigma(\mu_n)/\mu_1, \dots, \mu_n]$ is called the (closed) σ -instance of \mathcal{D} .

Definition 15. An inference is a triple $\langle \langle \mathcal{D}_1, \dots, \mathcal{D}_n \rangle, A, \delta \rangle$, where δ is an extension of the discharge functions associated to the \mathcal{D}_i -s ($i \leq n$). The argument structure associated to the inference, indicated by the figure

$$\frac{\mathcal{D}_1, \dots, \mathcal{D}_n}{A} \delta$$

is obtained by conjoining the trees of \mathcal{D}_i -s ($i \leq n$) through a root node A , and by adding δ to the discharges of assumptions of the \mathcal{D}_i -s ($i \leq n$). A rule is a set of inferences, whose elements are called instances of the rule

(this definition is inspired by [17]). I shall assume that rules can be described through meta-linguistic schemes, as in the case of standard introduction rules in Gentzen's Natural Deduction:

$$\frac{A \quad B}{A \wedge B} (\wedge_I) \quad \frac{A_i}{A_1 \vee A_2} (\vee_{I,i}), i = 1, 2 \quad \frac{[A] \quad B}{A \rightarrow B} (\rightarrow_I)$$

Definition 16. \mathcal{D} is canonical iff it is associated to an instance of an introduction rule. It is non-canonical otherwise.

Definition 17. Given a rule R , a reduction for R is a mapping ϕ from and to argument structures such that ϕ is defined on the set of the argument structures \mathbb{D} associated to instances of R and, $\forall \mathcal{D} \in \mathbb{D}$,

- (a) \mathcal{D} is from Γ to $A \implies \phi(\mathcal{D})$ is from $\Gamma^* \subseteq \Gamma$ to A ;
- (b) $\forall \sigma$, ϕ is defined on \mathcal{D}^σ and $\phi(\mathcal{D}^\sigma) = \phi(\mathcal{D})^\sigma$.

I will use the following notation: \mathfrak{J} indicates a set of reductions, and \mathfrak{J}^+ indicates an extension of \mathfrak{J} , i.e., $\mathfrak{J} \subseteq \mathfrak{J}^+$.

Definition 18. \mathcal{D} immediately reduces to \mathcal{D}^* relative to \mathfrak{J} iff, for some \mathcal{D}^{**} sub-structure of \mathcal{D} , there is $\phi \in \mathfrak{J}$ such that $\mathcal{D}^* = \mathcal{D}[\phi(\mathcal{D}^{**})/\mathcal{D}^{**}]$. The relation of \mathcal{D} reducing to \mathcal{D}^* relative to \mathfrak{J} is the reflexive-transitive closure of the relation of immediate reducibility of \mathcal{D} to \mathcal{D}^* .

I will use the following notation: $\mathcal{D} \leq_{\mathfrak{J}} \mathcal{D}^*$ means that \mathcal{D} reduces to \mathcal{D}^* relative to \mathfrak{J} .

Definition 19. An argument is a pair $\langle \mathcal{D}, \mathfrak{J} \rangle$.

Definition 20. $\langle \mathcal{D}, \mathfrak{J} \rangle$ is n -valid on \mathfrak{B}^m iff $m \leq n$ and

- \mathcal{D} is closed \implies
 - the conclusion of \mathcal{D} is atomic $\implies \mathcal{D} \leq_{\mathfrak{J}} \mathcal{D}^*$ with $\mathcal{D}^* = \langle T, \emptyset \rangle$ and $T \in \text{DER}_{\mathfrak{B}^m}$ closed;
 - the conclusion of \mathcal{D} is not atomic $\implies \mathcal{D} \leq_{\mathfrak{J}} \mathcal{D}^*$ for \mathcal{D}^* canonical with immediate sub-structures n -valid on \mathfrak{B}^m when paired with \mathfrak{J} ;
- \mathcal{D} is open $\implies \forall \sigma, \forall \mu \in \Gamma, \forall \mathfrak{J}^+ \supseteq \mathfrak{J}$ and $\forall \mathfrak{C} \supseteq_n \mathfrak{B}^m$, if $\langle \sigma(\mu), \mathfrak{J}^+ \rangle$ is n -valid on \mathfrak{C} , then $\langle \mathcal{D}^\sigma, \mathfrak{J}^+ \rangle$ is n -valid on \mathfrak{C} .

I will use the following notation: $\Gamma \models_{\mathfrak{B}, n}^\alpha A$ means that there is $\langle \mathcal{D}, \mathfrak{J} \rangle$ from Γ to A which is n -valid on \mathfrak{B} .

Proposition 5. For every n and every $\mathfrak{B} \in \mathbb{B}^n$:

- (a) $\forall A \in \text{ATOM}_{\mathcal{L}} (\models_{\mathfrak{B}, n}^\alpha A \iff \vdash_{\mathfrak{B}} A)$;
- (b) $\models_{\mathfrak{B}, n}^\alpha \perp \iff \forall A \in \text{ATOM}_{\mathcal{L}} (\vdash_{\mathfrak{B}} A)$;
- (c) $\models_{\mathfrak{B}, n}^\alpha A \wedge B \iff \models_{\mathfrak{B}, n}^\alpha A$ and $\models_{\mathfrak{B}, n}^\alpha B$;
- (d) $\models_{\mathfrak{B}, n}^\alpha A \vee B \iff \models_{\mathfrak{B}, n}^\alpha A$ or $\models_{\mathfrak{B}, n}^\alpha B$;
- (e) $\models_{\mathfrak{B}, n}^\alpha A \rightarrow B \iff A \models_{\mathfrak{B}, n}^\alpha B$;
- (f) $\Gamma \models_{\mathfrak{B}, n}^\alpha A \iff \forall \mathfrak{C} \supseteq_n \mathfrak{B} (\Gamma \models_{\mathfrak{C}, n}^\alpha A)$ —i.e. the notion is monotonic;
- (g) $\Gamma \models_{\mathfrak{B}, n}^\alpha A \iff \forall \mathfrak{C} \supseteq_n \mathfrak{B} (\models_{\mathfrak{C}, n}^\alpha \Gamma \implies \models_{\mathfrak{C}, n}^\alpha A)$.

Proof. The only non-trivial case is direction (\iff) of (g)¹. Suppose $\Gamma = \{A_1, \dots, A_m\}$, and suppose that $\forall \mathfrak{C} \supseteq_n \mathfrak{B} (\models_{\mathfrak{C}, n}^\alpha \Gamma \implies \models_{\mathfrak{C}, n}^\alpha A)$. This means that, for every $\mathfrak{C} \supseteq_n \mathfrak{B}$, if there is a closed $\langle \mathcal{D}_i, \mathfrak{J}_i \rangle$ for A_i which is n -valid on \mathfrak{C} ($i \leq m$), then there is a closed $\langle \mathcal{D}_A, \mathfrak{J}_A \rangle$ for A which is n -valid on \mathfrak{C} . Then, take any $\langle \mathcal{D}_i, \mathfrak{J}_i \rangle$ for A_i ($i \leq m$), and consider the rule $\{\langle \langle \mathcal{D}_1, \dots, \mathcal{D}_m \rangle, A \rangle\}$, to whose only element the argument structure

¹A similar proof is to be found in [27, Lemma 4.11]. Since Stafford's discussion has no restrictions on Γ being finite, in that context the result amounts to proving that Prawitz's reducibility semantics enjoys compact monotonicity. Another similar proof is found in [13, pp. 153-154], although there the proof is referred to non-monotonic reducibility semantics, and used for showing that, when suitable conditions are met, classical logic can be justified over Prawitz's reducibility semantics.

$$\mathcal{D}^* = \frac{\mathcal{D}_1 \quad \dots \quad \mathcal{D}_m}{A_1 \quad \dots \quad A_m} \quad A$$

is associated, and let ϕ^* be the mapping such that $\phi^*(\mathcal{D}^*) = \mathcal{D}_A$. Let \mathbb{F} be the set of all the reductions which satisfy Definition 17. Then, $\phi^* \in \mathbb{F}$: for, both \mathcal{D}^* and \mathcal{D}_A are from assumptions \emptyset to conclusion A , so condition (a) in Definition 17 is satisfied and, for every $\mathcal{D}^{*,\sigma}$ and \mathcal{D}_A^σ , it holds that $\mathcal{D}^{*,\sigma} = \mathcal{D}^*$ and $\mathcal{D}_A^\sigma = \mathcal{D}_A$ respectively, so condition (b) in Definition 17 is satisfied. Observe also that $\mathfrak{J}_i \subseteq \mathbb{F}$ ($i \leq m$) and $\mathfrak{J}_A \subseteq \mathbb{F}$. Consider now the argument structure

$$\mathcal{D} = \frac{A_1 \quad \dots \quad A_m}{A}$$

We have that $\langle \mathcal{D}, \mathbb{F} \rangle$ is n -valid on \mathfrak{B} . For, suppose we are given $\langle \mathcal{D}_i, \mathfrak{J}_i \rangle$ for A_i n -valid on any $\mathfrak{C} \supseteq_n \mathfrak{B}$ ($i \leq m$). We have to prove that $\langle \mathcal{D}[\mathcal{D}_1, \dots, \mathcal{D}_m/A_1, \dots, A_m], \mathbb{F} \cup \bigcup_{i \leq m} \mathfrak{J}_i \rangle$ is n -valid on \mathfrak{C} . But $\bigcup_{i \leq m} \mathfrak{J}_i \subseteq \mathbb{F}$, so $\mathbb{F} \cup \bigcup_{i \leq m} \mathfrak{J}_i = \mathbb{F}$, so we must prove that $\langle \mathcal{D}[\mathcal{D}_1, \dots, \mathcal{D}_m/A_1, \dots, A_m], \mathbb{F} \rangle$ is n -valid on \mathfrak{C} . By Definition 19, this obtains when $\mathcal{D}[\mathcal{D}_1, \dots, \mathcal{D}_m/A_1, \dots, A_m]$ reduces relative to \mathbb{F} to a closed argument structure for A which, when paired with \mathbb{F} , be n -valid on \mathfrak{C} . Since \mathbb{F} will contain a ϕ^* of the kind described above, we have $\phi^*(\mathcal{D}[\mathcal{D}_1, \dots, \mathcal{D}_m/A_1, \dots, A_m]) = \mathcal{D}_A$ for some \mathcal{D}_A such that, for some \mathfrak{J}_A , $\langle \mathcal{D}_A, \mathfrak{J}_A \rangle$ is n -valid on \mathfrak{C} . Since $\mathfrak{J}_A \subseteq \mathbb{F}$, $\langle \mathcal{D}_A, \mathbb{F} \rangle$ is also n -valid on \mathfrak{C} .² \square

Definition 21. $\langle \mathcal{D}, \mathfrak{J} \rangle$ is n -valid iff, $\forall \mathfrak{B} \in \mathbb{B}^n$, $\langle \mathcal{D}, \mathfrak{J} \rangle$ is n -valid on \mathfrak{B} .

I will use the following notation: $\Gamma \models_n^\alpha A$ means that there is an n -valid $\langle \mathcal{D}, \mathfrak{J} \rangle$ from Γ to A .

Proposition 6 (Schroeder-Heister, [24]). $\Gamma \models_n^\alpha A \iff \Gamma \models_{\mathfrak{B}^0}^\alpha A$.

Corollary 2. *The following facts hold:*

- (a) $\Gamma \models_n^\alpha A \iff \forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^\alpha A)$;
- (b) $\forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^\alpha A) \iff \forall \mathfrak{B} \in \mathbb{B}^n (\models_{\mathfrak{B}, n}^\alpha \Gamma \implies \models_{\mathfrak{B}, n}^\alpha A)$.

Proof. As for (a), (\implies) is trivial. For (\impliedby) , if $\forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^\alpha A)$ then, in particular, $\Gamma \models_{\mathfrak{B}^0}^\alpha A$ and, by Proposition 6, $\Gamma \models_n^\alpha A$. As for (b), (\implies) if $\forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^\alpha A)$, then $\Gamma \models_{\mathfrak{B}^0}^\alpha A$. By point (g) of Proposition 5 plus Proposition 1, this means $\forall \mathfrak{B} \in \mathbb{B}^n (\models_{\mathfrak{B}, n}^\alpha \Gamma \implies \models_{\mathfrak{B}, n}^\alpha A)$. (\impliedby) It is sufficient to observe (by using Definition 20) that the first implication in the proof of the inverse direction also holds from right to left, that is, $\Gamma \models_{\mathfrak{B}^0}^\alpha A \implies \forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^\alpha A)$. \square

4 Comparison of the three approaches

In what follows, I shall be comparing the three approaches to proof-theoretic semantics mentioned so far.

4.1 Reducibility semantics and standard base semantics

Definition 7 and Proposition 5 have the obvious effect of making standard base semantics and reducibility semantics equivalent, both at the level of consequence over a base, and at the level of logical consequence.

Theorem 1. $\Gamma \models_{\mathfrak{B}, n}^\alpha A \iff \Gamma \models_{\mathfrak{B}, n}^\alpha A$.

Theorem 2. $\Gamma \models_n^\alpha A \iff \Gamma \models_n^\alpha A$.

²Observe that, in principle, there might be a different ϕ^* for each different sequence $\mathcal{D}_1, \dots, \mathcal{D}_m$ such that $\langle \mathcal{D}_i, \mathfrak{J}_i \rangle$ is closed valid for A_i on \mathfrak{C} ($i \leq m$), since there might in principle be a different \mathcal{D}_A such that $\langle \mathcal{D}_A, \mathfrak{J}_A \rangle$ is closed valid for A on \mathfrak{C} associated to each such sequence. With classical logic in the meta-language, however, the situation is much smoother: either there is A_i such that there is no closed argument for A_i valid on \mathfrak{C} , in which case ϕ^* is the empty function, or for every A_i there is a closed argument for A_i valid on \mathfrak{C} , in which case there will surely be at least one closed argument for A valid on \mathfrak{C} , so ϕ^* can be the constant function. An additional remark is the following. One may complain that the reduction \mathbb{F} is not “constructive enough”. More constructive examples of reductions for the one-step argument structure from Γ to A can be given, in such a way as to prove again the direction (\impliedby) of (g) in Proposition 5—these are essentially adaptations from the incompleteness results proved in [2, 15, 16]. The discussion of this topic would have led me too far away, so I may want to deal with it in future works—for a partial treatment, see [13]. Concerning the issue of what a “good” reduction is, the reader may refer to [1]. Let me point out that, as done in [24], reductions could be also defined in terms of (sequences of) pairs of argument structures, i.e., sequence of pairs $\langle \mathcal{D}, \mathcal{D}^* \rangle$ where the first element of the $i + 1$ -th pair has at most the assumptions and the same conclusion as the second element in i -th pair. One final observation is that, besides few very quick remarks, I shall not deal here with “deviant” reducibility semantics where potential readings of what a reduction should be, stricter than those mentioned above, make the bi-implication (g) of Proposition 5 fail—leaving us with the left-to-right direction only.

So, we can transfer to reducibility semantics all the (in)completeness results which have been proved for base semantics in the standard reading. Below, we shall discuss some incompleteness results for intuitionistic logic IL. For the moment the reader may refer to [15, 16, 27], and rely on the following result.

Theorem 3. $\exists \Gamma \exists A (\Gamma \models_n^\alpha A \text{ and } \Gamma \not\vdash_{\text{IL}} A)$.

Despite their simplicity, Theorem 1 and Theorem 2 provide an interesting connection between Prawitz’s reducibility semantics and base semantics in the standard reading. For, the latter can be looked at as “extracted” from the former by dropping argument structures and reductions out, via a consequence relation defined over (sets of) formulas outright, rather than in a derivative way as existence of suitable valid arguments. Conversely, Prawitz’s reducibility semantics can be understood as obtained from base semantics in the standard reading, by decorating the formula-based consequence relation with argument structures and reductions which “witness” that such a relation holds. Hence, one may naturally wonder whether, whenever $\Gamma \models_{\mathfrak{B}, n} A$ or $\Gamma \models_n A$ hold, a “witness” of this can be found so that $\Gamma \models_{\mathfrak{B}, n}^\alpha A$ or $\Gamma \models_n^\alpha A$ hold too, and conversely whether, whenever $\Gamma \models_{\mathfrak{B}, n}^\alpha A$ or $\Gamma \models_n^\alpha A$ hold, the “witness” can be safely removed without loss of constructivity.

As I see it, the point raised here is best appreciated when formulated in a style similar to the one at play in another major constructivist approach, i.e., Martin-Löf’s (intuitionistic) type theory [8]. Here, following the formulas-as-types conception and the Curry-Howard correspondence inspiring it [7], we start with forms of *judgement*

$$a : A$$

where a is a proof-object and A is a type/proposition, so what the judgement expresses is that the proof-object a is of type A , i.e., it is a proof-object of proposition A and, conversely, that proposition A , understood as a type, i.e., as a class of proof-objects, is inhabited. Next to this, we have dependent proof-objects involved in conditional judgements

$$b(\mathbf{x}) : A (\mathbf{x} : \Gamma)$$

—with Γ set $\{A_1, \dots, A_n\}$ of types/propositions A_i , and \mathbf{x} sequence x_1, \dots, x_n such that $x_i : A_i$ ($i \leq n$)—meaning that $b(a)$ is a proof-object of type A if \mathbf{a} is a proof-object of type Γ —i.e., for every $i \leq n$, a_i is a proof-object of type A_i —and, conversely, that type A is inhabited provided all the types in Γ are. Based on this, one can introduce forms of judgement

$$A \text{ true}$$

where **true** is explained by the rule

$$\frac{a : A}{A \text{ true}} \text{ T}$$

and, hence, forms of judgement

$$A \text{ true } (\Gamma \text{ true})$$

—where $\Gamma \text{ true}$ means that, for every $i \leq n$, $A_i \text{ true}$ —which (with a type-theoretic abuse of notation, that is however harmless in the present context) might be explained by the rule

$$\frac{b(\mathbf{x}) : A (\mathbf{x} : \Gamma)}{A \text{ true } (\Gamma \text{ true})} \text{ Td}$$

Now, Prawitz’s reducibility semantics runs in very much the same way, in that $\models_{\mathfrak{B}}^\alpha$ and \models^α are always explained in terms of (existence of) proof-objects witnessing them, similarly to what happens in Martin-Löf’s intuitionistic type theory, when explaining **true** through the rules T and Td. In standard base semantics, instead, $\models_{\mathfrak{B}}$ and \models are defined at the sentence level directly so that, to put it in type-theoretic terms, it is as if we started with **true** outright, without requiring it to be explained by T and Td. To distinguish this picture from the previous one, we may write **true*** instead of **true**. Now, the question I raised above—whether from $\Gamma \models_{\mathfrak{B}}$ or $\Gamma \models A$ one can go to $\Gamma \models_{\mathfrak{B}}^\alpha A$ or $\Gamma \models^\alpha A$ and vice versa—would become the question whether at least one of the following rules is valid—leaving atomic bases aside:

$$\frac{A \text{ true } (\Gamma \text{ true})}{A \text{ true}^* (\Gamma \text{ true}^*)} \quad \frac{A \text{ true}^* (\Gamma \text{ true}^*)}{A \text{ true } (\Gamma \text{ true})}$$

—where Γ might be empty—namely, recalling the explanation of \mathbf{true} , whether one of the following rules is valid:

$$\frac{b(\mathbf{x}) : A(\mathbf{x} : \Gamma)}{A \mathbf{true}^*(\Gamma \mathbf{true}^*)} \quad \frac{A \mathbf{true}^*(\Gamma \mathbf{true}^*)}{b(\mathbf{x}) : A(\mathbf{x} : \Gamma)}$$

for suitable $b(\mathbf{x})$. Theorem 1 and Theorem 2 answer positively to these questions. At the same time, they imply that, under the given limitations (mostly concerning the restriction to finite Γ -s), Prawitz’s reducibility semantics and standard base semantics are “structurally” identical, meaning that the former enjoys *mutatis mutandis* the same properties as those employed in [15, 16] to prove incompleteness of intuitionistic propositional logic with respect to the latter—properties which I shall come back to below. Thus, as stated by Theorem 3, intuitionistic propositional logic is also incomplete over Prawitz’s reducibility semantics in the version at issue here—essentially the same line of thought is used in [10] to prove incompleteness of IL over Prawitz’s theory of grounds [20].

These full-equivalence results are mainly due to the fact that both Prawitz’s reducibility semantics and base semantics in the standard reading are “introduction-based”, i.e., they both ultimately rely on the idea that the meaning of a logical constant κ is given by the *conditions* for formulas having κ as main sign to hold. This is the reason why Theorem 1 can be proved by easy induction on the complexity of formulas in the closed case, and then by “closure” in the open case, using point (g) in Proposition 5—then, Theorem 2 follows by monotonicity, and Theorem 3 follows straightforwardly from Theorem 2 plus the incompleteness results proved in [15, 16].

4.2 Reducibility semantics and Sandqvist’s base semantics

However, this is also the reason why such a smooth inductive reasoning, hence full-equivalence, do not apply when comparing Prawitz’s reducibility semantics and base semantics *à la* Sandqvist. The reasoning breaks down since Sandqvist explains \vee in an “elimination-based” way. For, clause (c) in Definition 8 is in fact nothing but a (monotonic) semantic rendering of the standard elimination rule for \vee , restricted to atomic minor premises. But the question may arise also here whether, as in the previous case, whenever $\Gamma \models_{\mathfrak{B}, n}^s A$ or $\Gamma \models_n^s A$ hold, a “witness” of this can be found so that $\Gamma \models_{\mathfrak{B}, n}^\alpha A$ or $\Gamma \models_n^\alpha A$ hold too, and vice versa, whether from $\Gamma \models_{\mathfrak{B}, n}^\alpha A$ or $\Gamma \models_n^\alpha A$ one can conclude $\Gamma \models_{\mathfrak{B}, n}^s A$ or $\Gamma \models_n^s A$. Because smooth induction and full-equivalence are now ruled out, anyway, the most one can expect in this context is establishing *to what extent* the back-and-forth is possible. In turn, this can be understood as establishing the extent to which the smooth inductive reasoning used for proving Theorem 1 can be applied in the case of a comparison between Prawitz’s reducibility semantics and Sandqvist’s base semantics.

Going inductively from Prawitz’s reducibility semantics to base semantics in Sandqvist’s variant seems not beyond reach. Omitting details to be found in the proof of Theorem 4, suppose $A \vee B$ holds in Prawitz’s sense on some base. Then, for every extension of the base, either A or B hold on that extension. Assume that atom C is consequence of both A and B in Sandqvist’s sense on that extension. If we can grant inductively that the holding of A and B in Prawitz’s sense on the extension implies the holding of A and B in Sandqvist’s sense on the extension, we can conclude C holds in Sandqvist’s sense on the extension too which, by arbitrariness of the extension, means that $A \vee B$ holds in Sandqvist’s sense in the original base.

The inverse, however, seems not to hold: from the fact that, for every atom C which is consequence of A and B in Sandqvist’s sense on some extension of a given base, C holds in Sandqvist’s sense on the extension, we cannot conclude that A or B hold on the base in Sandqvist’s sense, so we cannot apply any potential induction for going from Sandqvist’s base semantics to Prawitz’s reducibility semantics—which is, incidentally, one of the selling points of Sandqvist’s approach as concerns completeness of intuitionistic logic, see [21].

It is therefore clear that, since for \vee we only have the Prawitz-to-Sandqvist direction, an inductive full-equivalence proof is broken also for the other propositional constants. What is missing is, more specifically, the possibility of going from Sandqvist to Prawitz. But this does not exclude that *assuming* the condition that we can go from Sandqvist to Prawitz everywhere, we can keep conditionally, for *all* propositional constants, the possibility of going the other way around everywhere as well.

The aforementioned condition is required mainly because of how the clause for \rightarrow and that for $\Gamma \neq \emptyset$ work. Again omitting details to be found in the proof of Theorem 4, and limiting ourselves to the implicational case, suppose that $A \rightarrow B$ holds in Prawitz’s sense on some base. Therefore, for any extension of the base, if A holds in Prawitz’s sense on the extension, also B holds in Prawitz’s sense on the extension. We must prove that the same holds on the extension also for Sandqvist’s variant, namely that, whenever A holds in Sandqvist’s sense on the extension, then also B holds in Sandqvist’s sense on the extension. This obtains in turn when we assume that, for every extension and every Γ and

A , we can go from A being consequence of Γ in Sandqvist's sense on the extension to A being consequence of Γ in Prawitz's sense on the extension.³

Thus, Prawitz's reducibility semantics and base semantics in Sandqvist's reading *are* connected, but not “point-wise”, i.e., assigning to *each* instance $\Gamma \vDash_{\mathfrak{B}, n}^s A$ of Sandqvist's consequence notion the corresponding instance $\Gamma \vDash_{\mathfrak{B}, n}^\alpha A$ of Prawitz's consequence notion, and vice versa. The connection is, so to say, “conditionally global”, in the sense described above. As said, this certainly depends on the structural difference given by Prawitz's “introduction-based”, and Sandqvist's “elimination-based” explanations of \vee . This “conditionally global” connection is however not without value, as it implies at least two seemingly interesting consequences. Before turning to that, however, let me prove the results which I have been illustrating informally so far. I will use the following notation: $A^* \leq A$ indicates that the logical complexity of A^* is smaller than or equal to that of A . Consider now the following statement:

$$\forall \Gamma \forall A \forall \mathfrak{C} \supseteq_n \mathfrak{B} (\Gamma \vDash_{\mathfrak{C}, n}^s A \implies \Gamma \vDash_{\mathfrak{C}, n}^\alpha A). \quad (1)$$

Theorem 4. $(1) \implies \forall \Gamma \forall A \forall \mathfrak{C} \supseteq_n \mathfrak{B} (\Gamma \vDash_{\mathfrak{C}, n}^\alpha A \implies \Gamma \vDash_{\mathfrak{C}, n}^s A)$.

Proof. Assume (1), and suppose $\Gamma = \emptyset$. We proceed by induction on complexity of A :

- take any arbitrary $\mathfrak{C} \supseteq_n \mathfrak{B}$ and suppose $A \in \text{ATOM}$. By point (a) of Proposition 5 and point (a) of Definition 8,

$$\vDash_{\mathfrak{C}, n}^\alpha A \text{ iff } \vdash_{\mathfrak{C}} A \text{ iff } \vDash_{\mathfrak{C}, n}^s A.$$

Hence, *a fortiori*,

$$\vDash_{\mathfrak{C}, n}^\alpha A \implies \vDash_{\mathfrak{C}, n}^s A.$$

Hence, by arbitrariness of \mathfrak{C} ,

$$\forall \mathfrak{C} \supseteq_n \mathfrak{B} (\vDash_{\mathfrak{C}, n}^\alpha A \implies \vDash_{\mathfrak{C}, n}^s A);$$

Observe that we do not have to worry about $A = \perp$ via Proposition 2.

- assume the induction hypothesis

$$\forall A^* < A \forall \mathfrak{C} \supseteq_n \mathfrak{B} (\vDash_{\mathfrak{C}, n}^\alpha A^* \implies \vDash_{\mathfrak{C}, n}^s A^*).$$

The case for \wedge is trivial. As for the others:

- (\vee) take any arbitrary $\mathfrak{C} \supseteq_n \mathfrak{B}$ and suppose $A = B \vee C$. So, by points (d) and (f) of Proposition 5

$$\vDash_{\mathfrak{C}, n}^\alpha B \vee C \text{ iff } \forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\vDash_{\mathfrak{C}^*, n}^\alpha B \vee C) \text{ iff } \forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\vDash_{\mathfrak{C}^*, n}^\alpha B \text{ or } \vDash_{\mathfrak{C}^*, n}^\alpha C).$$

For any arbitrary $\mathfrak{C}^* \supseteq_n \mathfrak{C}$ and any arbitrary $D \in \text{ATOM}_{\mathcal{L}}$, assume $B \vDash_{\mathfrak{C}^*, n}^s D$ and $C \vDash_{\mathfrak{C}^*, n}^s D$. Instantiate on \mathfrak{C}^* the previous

$$\forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\vDash_{\mathfrak{C}^*, n}^\alpha B \text{ or } \vDash_{\mathfrak{C}^*, n}^\alpha C)$$

and assume $\vDash_{\mathfrak{C}^*, n}^\alpha B$. Instantiate on B the induction hypothesis, so to obtain $\vDash_{\mathfrak{C}^*, n}^s B$ which, with $B \vDash_{\mathfrak{C}^*, n}^s D$, yields $\vDash_{\mathfrak{C}^*, n}^s D$. The same can be obtained by assuming $\vDash_{\mathfrak{C}^*, n}^\alpha C$. Discharging the assumptions $\vDash_{\mathfrak{C}^*, n}^\alpha B$ and $\vDash_{\mathfrak{C}^*, n}^\alpha C$, we obtain $\vDash_{\mathfrak{C}^*, n}^s D$ and, discharging the assumptions $B \vDash_{\mathfrak{C}^*, n}^s D$ and $C \vDash_{\mathfrak{C}^*, n}^s D$, we have

$$B \vDash_{\mathfrak{C}^*, n}^s D \text{ and } C \vDash_{\mathfrak{C}^*, n}^s D \implies \vDash_{\mathfrak{C}^*, n}^s D.$$

Since now \mathfrak{C}^* and D do not occur free in any undischarged assumption, we can universally quantify over them, and obtain

$$\forall \mathfrak{C}^* \supseteq_n \mathfrak{C} \forall D \in \text{ATOM}_{\mathcal{L}} (B \vDash_{\mathfrak{C}^*, n}^s D \text{ and } C \vDash_{\mathfrak{C}^*, n}^s D \implies \vDash_{\mathfrak{C}^*, n}^s D)$$

³This is a stronger claim than what is actually needed for proving Theorem 4. We could limit ourselves to assuming that the implication holds only for valid formulas, rather than for consequences (i.e., with $\Gamma = \emptyset$). However, the stronger claim clearly implies the weaker, and is needed in the proof of Theorem 13, which is why I favoured it.

which, by point (c) in Definition 8, means $\models_{\mathfrak{C}, n}^s B \vee C$. Let us now discharge $\models_{\mathfrak{C}, n}^\alpha B \vee C$, to obtain

$$\models_{\mathfrak{C}, n}^\alpha B \vee C \implies \models_{\mathfrak{C}, n}^s B \vee C.$$

Our only undischarged assumptions are (1) and the induction hypothesis, in none of which \mathfrak{C} occurs free. Thus, we can universally quantify over it, and obtain

$$\forall \mathfrak{C} \supseteq_n \mathfrak{B} (\models_{\mathfrak{C}, n}^\alpha B \vee C \implies \models_{\mathfrak{C}, n}^s B \vee C);$$

(\rightarrow) take any arbitrary $\mathfrak{C} \supseteq_n \mathfrak{B}$ and suppose $A = B \rightarrow C$. Then, by point (e) and (g) of Proposition 5,

$$\models_{\mathfrak{C}, n}^\alpha B \rightarrow C \text{ iff } B \models_{\mathfrak{C}, n}^\alpha C \text{ iff } \forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\models_{\mathfrak{C}^*, n}^\alpha B \implies \models_{\mathfrak{C}^*, n}^\alpha C).$$

Take now any arbitrary $\mathfrak{C}^* \supseteq_n \mathfrak{C}$ and assume $\models_{\mathfrak{C}^*, n}^s B$. Instantiating (1) on \emptyset, B and $\mathfrak{C}^* \supseteq_n \mathfrak{C} \supseteq_n \mathfrak{B}$, we get

$$\models_{\mathfrak{C}^*, n}^s B \implies \models_{\mathfrak{C}^*, n}^\alpha B$$

which, with the assumption $\models_{\mathfrak{C}^*, n}^s B$ above, yields $\models_{\mathfrak{C}^*, n}^\alpha B$. The latter, when

$$\forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\models_{\mathfrak{C}^*, n}^\alpha B \implies \models_{\mathfrak{C}^*, n}^\alpha C)$$

above is instantiated on \mathfrak{C}^* , yields $\models_{\mathfrak{C}^*, n}^\alpha C$. Now, the induction hypothesis instantiated on C yields

$$\forall \mathfrak{C} \supseteq_n \mathfrak{B} (\models_{\mathfrak{C}, n}^\alpha C \implies \models_{\mathfrak{C}, n}^s C)$$

which, when instantiated on $\mathfrak{C}^* \supseteq_n \mathfrak{C} \supseteq_n \mathfrak{B}$, yields $\models_{\mathfrak{C}^*, n}^s C$ with $\models_{\mathfrak{C}^*, n}^\alpha C$ above. We now discharge $\models_{\mathfrak{C}^*, n}^s B$, to obtain

$$\models_{\mathfrak{C}^*, n}^s B \implies \models_{\mathfrak{C}^*, n}^s C.$$

Since \mathfrak{C}^* is no longer free in any assumption, we conclude

$$\forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\models_{\mathfrak{C}^*, n}^s B \implies \models_{\mathfrak{C}^*, n}^s C)$$

which, by point 2 of Definition 8, is equivalent to $B \models_{\mathfrak{C}, n}^s C$ which in turn, by point (d) of Definition 8, is equivalent to $\models_{\mathfrak{C}, n}^s B \rightarrow C$. Discharging $\models_{\mathfrak{C}, n}^\alpha B \rightarrow C$, we obtain

$$\models_{\mathfrak{C}, n}^\alpha B \rightarrow C \implies \models_{\mathfrak{C}, n}^s B \rightarrow C.$$

By arbitrariness of \mathfrak{C} (which, again, is not free in any pending assumption), we obtain

$$\forall \mathfrak{C} \supseteq_n \mathfrak{B} (\models_{\mathfrak{C}, n}^\alpha B \rightarrow C \implies \models_{\mathfrak{C}, n}^s B \rightarrow C).$$

Suppose now $\Gamma \neq \emptyset$. Take any arbitrary $\mathfrak{C} \supseteq_n \mathfrak{B}$ and suppose $\Gamma \models_{\mathfrak{C}, n}^\alpha A$. By point (g) of Proposition 5, this means

$$\forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\models_{\mathfrak{C}^*, n}^\alpha \Gamma \implies \models_{\mathfrak{C}^*, n}^\alpha A).$$

Take any arbitrary $\mathfrak{C}^* \supseteq_n \mathfrak{C}$ and assume $\models_{\mathfrak{C}^*, n}^s \Gamma$. Let (1) be instantiated on \emptyset , the elements of Γ , and $\mathfrak{C}^* \supseteq_n \mathfrak{C} \supseteq_n \mathfrak{B}$, so

$$\models_{\mathfrak{C}^*, n}^s \Gamma \implies \models_{\mathfrak{C}^*, n}^\alpha \Gamma$$

which, with the assumption $\models_{\mathfrak{C}^*, n}^s \Gamma$ above, yields $\models_{\mathfrak{C}^*, n}^\alpha \Gamma$. The latter, when

$$\forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\models_{\mathfrak{C}^*, n}^\alpha \Gamma \implies \models_{\mathfrak{C}^*, n}^\alpha A)$$

is instantiated on \mathfrak{C}^* , yields $\models_{\mathfrak{C}^*, n}^\alpha A$. By what proved for the case with $\Gamma = \emptyset$, we obtain that

$$\forall \mathfrak{C} \supseteq_n \mathfrak{B} (\models_{\mathfrak{C}, n}^\alpha A \implies \models_{\mathfrak{C}, n}^s A).$$

Instantiate this on $\mathfrak{C}^* \supseteq_n \mathfrak{C} \supseteq_n \mathfrak{B}$, and get $\models_{\mathfrak{C}^*, n}^s A$ through $\models_{\mathfrak{C}^*, n}^\alpha A$. Hence, by discharging $\models_{\mathfrak{C}^*, n}^s \Gamma$, we have

$$\models_{\mathfrak{C}^*, n}^s \Gamma \implies \models_{\mathfrak{C}^*, n}^s A.$$

We can now bind \mathfrak{C}^* , and get

$$\forall \mathfrak{C}^* \supseteq_n \mathfrak{C} (\models_{\mathfrak{C}^*, n}^s \Gamma \implies \models_{\mathfrak{C}^*, n}^s A),$$

which means $\Gamma \models_{\mathfrak{C}, n}^s A$ by point 2 of Definition 8. Now we discharge the assumption $\Gamma \models_{\mathfrak{C}, n}^\alpha A$, and obtain

$$\Gamma \models_{\mathfrak{C}, n}^\alpha A \implies \Gamma \models_{\mathfrak{C}, n}^s A.$$

As usual, \mathbb{C} is not free in any open assumption, so we bind it and conclude

$$\forall \mathbb{C} \supseteq_n \mathfrak{B} (\Gamma \models_{\mathbb{C}, n}^\alpha A \implies \Gamma \models_{\mathbb{C}, n}^s A).$$

To obtain the theorem it is now just sufficient to discharge (1). \square

Corollary 3. $\forall \Gamma \forall A \forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^s A \implies \Gamma \models_{\mathfrak{B}, n}^\alpha A) \implies \forall \Gamma \forall A \forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^\alpha A \implies \Gamma \models_{\mathfrak{B}, n}^s A).$ ⁴

Let us now turn to the first of the two relevant consequences of Theorem 4 and derived Corollary 3 that I mentioned above, and let us observe to begin with that the different ways in which Prawitz’s reducibility semantics and Sandqvist’s base semantics deal with \forall do not by themselves exclude that the two approaches might be somehow “ordered” relative to consequence over an atomic base. Let us say that \models^α and \models^s are *base-comparable* if and only if either the antecedent or the consequent of Corollary 3 hold. Base-comparability can be also read as the property that \models^α and \models^s are “models-monomorphic”, i.e., either all models of \models^α are also models of \models^s , or vice versa, i.e., again, more precisely, if we set

$$\mathbb{M}_{\Gamma, A, n}^\alpha = \{\mathfrak{B} \mid \Gamma \models_{\mathfrak{B}, n}^\alpha A\} \text{ and } \mathbb{M}_{\Gamma, A, n}^s = \{\mathfrak{B} \mid \Gamma \models_{\mathfrak{B}, n}^s A\},$$

then for all n, Γ and A either $\mathbb{M}_{\Gamma, A, n}^\alpha \subseteq \mathbb{M}_{\Gamma, A, n}^s$ or $\mathbb{M}_{\Gamma, A, n}^s \subseteq \mathbb{M}_{\Gamma, A, n}^\alpha$.

However, given what we know today at the level of *logical validity*, we can also immediately rule out that, in some (very relevant) cases, Prawitz’s reducibility semantics is “models-monomorphic” over Sandqvist’s base semantics. For, IL is known to be complete over the latter when atomic bases have level ≥ 2 —see [21] and Theorem 9 below—while we have just proved that IL is *never* complete over the former—see Theorem 3 above. Therefore, since logical validity means validity over all atomic bases, the consequent of Theorem 4 and Corollary 3 fails when atomic bases have level ≥ 2 . And now Theorem 4 and Corollary 3 come to play an active role by themselves, for via them we can infer in turn, by contraposition, that under the given conditions their antecedent fails too, namely, that Sandqvist’s base semantics is not “models-monomorphic” over Prawitz’s reducibility semantics either with atomic bases of level ≥ 2 . So, under the given conditions, Prawitz’s reducibility semantics and Sandqvist’s base semantics are not base-comparable or, to put it in another way, their classes of models $\mathbb{M}_{\Gamma, A, n}^\alpha$ and $\mathbb{M}_{\Gamma, A, n}^s$ diverge when $n \geq 2$. In turn, this may speak in favour of Schroeder-Heister’s proposal to consider Sandqvist’s approach as belonging to a family of proof-theoretic semantics which prioritise elimination rules *in general* [26]—for an elimination-based approach, see also [5, 9].⁵

As we shall see, however, it is part of the second aforementioned consequence of Theorem 4 and Corollary 3 that the latter can be also used to get some *positive* information. Before turning to that, let me first make precise the informal remarks that I have been carrying out thus far. For doing this, I must refer to some concepts and results presented in [15, 16] and [21]—some of which will be used also in Section 5. Piecha, de Campos Sanz and Schroeder-Heister [15] observed that any disjunction-free formula A can be associated to a set of atomic rules. The association requires first of all to transform such an A into a suitable $\#(A)$ via a number of step-wise replacements—with respect to which $\#(A)$ is irreducible—as follows: any sub-formula of the form $B \rightarrow C \wedge D$ is replaced by $(B \rightarrow C) \wedge (B \rightarrow D)$, and any sub-formula of the form $B \rightarrow (C \rightarrow D)$ is replaced by $(B \wedge C) \rightarrow D$.

Definition 22 (Piecha, de Campos Sanz & Schroeder-Heister [15]). *To any disjunction-free A we associate a set of atomic rules via a function $^\circ$ defined as follows:*

- $\#(A) \in \text{ATOM}_{\mathcal{L}} \implies A^\circ = \{A\};$
- $\#(A) = B_1 \wedge \dots \wedge B_n \rightarrow C$ with $C \in \text{ATOM}_{\mathcal{L}} \implies A^\circ = \{R\}$, where R has the form

⁴On a disjunction-free propositional language, reducibility semantics and Sandqvist’s base semantics *are* equivalent.

Proposition 7. *On a disjunction-free propositional language, $\Gamma \models_{\mathfrak{B}, n}^\alpha A \iff \Gamma \models_{\mathfrak{B}, n}^s A$.*

Proposition 8. *On a disjunction-free propositional language, $\Gamma \models_n^\alpha A \iff \Gamma \models_n^s A$.*

Let me also remark that, if we want to stick to the “deviant” version of reducibility semantics of footnote 2—i.e., with only the left-to-right direction in (g) of Proposition 5—we can still prove, in very much the same way, results similar to Theorem 4 and Corollary 3.

Theorem 5. $\forall \Gamma \forall A \forall \mathbb{C} \supseteq_n \mathfrak{B} (\Gamma \models_{\mathbb{C}, n}^\alpha A \implies \Gamma \models_{\mathbb{C}, n}^\alpha A) \implies \forall \Gamma \forall A \forall \mathbb{C} \supseteq_n \mathfrak{B} (\Gamma \models_{\mathbb{C}, n}^\alpha A \implies \Gamma \models_{\mathbb{C}, n} A).$

Corollary 4. $\forall \Gamma \forall A \forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n} A \implies \Gamma \models_{\mathfrak{B}, n}^\alpha A) \implies \forall \Gamma \forall A \forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^\alpha A \implies \Gamma \models_{\mathfrak{B}, n} A).$

⁵This was pointed out to me by Peter Schroeder-Heister, whom I am therefore indebted to for what follows in this section. It should be remarked that Oliveira’s approach abstracts from the kind of atomic bases one uses, whereas the interest of the comparison at issue here lies in the fact that the compared approaches use the *same* atomic bases.

$$\frac{[\mathfrak{K}_1] \quad \dots \quad [\mathfrak{K}_n]}{D_1 \quad \dots \quad D_n} C$$

and

$$\mathfrak{K}_1 \quad \dots \quad \mathfrak{K}_n \\ D_1 \quad \dots \quad D_n$$

correspond to $B_1^\circ, \dots, B_n^\circ$ respectively;

- $\#(A) = B_1 \wedge \dots \wedge B_n \implies A^\circ = \{B_1^\circ, \dots, B_n^\circ\}$.

Definition 23. Given a disjunction-free Γ , we set $\Gamma^\circ = \bigcup_{A_i \in \Gamma} A_i^\circ$.

Let us now drop for a moment the constraint that the level of the atomic bases has an upper bound n . This notion can be easily obtained from Definitions 7, 8, 9, 20 and 21, by removing the requirement that the level of the atomic base is $m \leq n$, and that the extensions of the base have level at most n . I will indicate this notion by \models , while the class of atomic bases with no upper bound will be written \mathbb{B} . The extension-relation will be just \supseteq (this is the only point where I use atomic bases with unlimited complexity, although I come back to this in the concluding remarks).

Definition 24 (Piecha, de Campos Sanz & Schroeder-Heister [15]). \models enjoys the import principle iff, for every $\mathfrak{B} \in \mathbb{B}$, $\models_{\mathfrak{B} \cup \Gamma^\circ} A \iff \Gamma \models_{\mathfrak{B}} A$ for every disjunction-free Γ .

Let us now consider what Piecha and Schroeder-Heister, in [16], call the *generalised disjunction property* (shortly, GDP): for every $\mathfrak{B} \in \mathbb{B}$, if Γ is disjunction-free, then

$$\Gamma \models_{\mathfrak{B}} A \vee B \implies (\Gamma \models_{\mathfrak{B}} A \text{ or } \Gamma \models_{\mathfrak{B}} B).$$

I will now appeal to some notions and results concerning soundness and completeness of (recursive) systems—and of IL in particular—over proof-theoretic validity. Both these notions and results will be more thoroughly explained in the next section. We know that, for every n , IL is sound with respect to \models_n . This also holds in the case when the level of atomic bases has no upper bound. Using this fact, the following can be proved.

Theorem 6 (Piecha & Schroeder-Heister, [16]). *If GDP holds on every $\mathfrak{B} \in \mathbb{B}$, Harrop’s rule is valid in \models namely, since Harrop’s rule is not derivable in IL [6], the latter is incomplete over \models .*

Proof. Take any arbitrary $\mathfrak{B} \in \mathbb{B}$, and $\mathfrak{C} \supseteq \mathfrak{B}$. Suppose $\models_{\mathfrak{C}} \neg A \rightarrow B \vee C$. Then, $\neg A \models_{\mathfrak{C}} B \vee C$. It is well-known that there is a disjunction-free A^* such that $\neg A \vdash_{\text{IL}} A^*$ and $A^* \vdash_{\text{IL}} \neg A$. So, by soundness of IL, $A^* \models_{\mathfrak{C}} B \vee C$. By GDP, $A^* \models_{\mathfrak{C}} B$ or $A^* \models_{\mathfrak{C}} C$ and, again by soundness of IL, $\neg A \models_{\mathfrak{C}} B$ or $\neg A \models_{\mathfrak{C}} C$. Hence, $\models_{\mathfrak{C}} \neg A \rightarrow B$ or $\models_{\mathfrak{C}} \neg A \rightarrow C$, so $\models_{\mathfrak{C}} (\neg A \rightarrow B) \vee (\neg A \rightarrow C)$. Quantify over every $\mathfrak{C} \supseteq \mathfrak{B}$, and every $\mathfrak{B} \in \mathbb{B}$. \square

Theorem 7 (Piecha & Schroeder-Heister [16]). *If \models enjoys the import principle, then GDP holds on every $\mathfrak{B} \in \mathbb{B}$.*

Proof. Assume $\Gamma \models_{\mathfrak{B}} A \vee B$ for Γ disjunction-free. Then $\models_{\mathfrak{B} \cup \Gamma^\circ} A \vee B$, whence $\Gamma \models_{\mathfrak{B}} A$ or $\Gamma \models_{\mathfrak{B}} B$. \square

Corollary 5 (Piecha & Schroeder-Heister [16]). *If \models enjoys the import principle, then IL is incomplete over \models .*

Piecha, de Campos Sanz and Schroeder-Heister also proved that \models does enjoy the import principle and that, via Sanqvist’s coding [21], the usage of atomic rules of level ≥ 3 can be reduced to that of atomic rules of level 2 [15].

Theorem 8 (Piecha, de Campos Sanz & Schroeder-Heister [15]). *IL is incomplete over \models_2 .*

Theorem 8 implies incompleteness of IL over \models_2^α , a result I already mentioned (Theorem 3).

Theorem 9 (Sandqvist [21]). *IL is complete over \models_2^s .*⁶

⁶As said, Sandqvist’s completeness proof applies to a language where \perp is a nullary logical constant equipped with a special clause stating that \perp holds on a base iff every atom holds on the base. The proof can be adapted to a language where \perp is instead an atomic constant, and where bases come with atomic explosion. Via Proposition 2 above, what one obtains in this way is just Sandqvist’s special clause for \perp . Then, in the construction of the “tailored” base for any valid sequent, \perp is mapped onto itself. The “tailored” base will at that point contain, by default, a rule which infers any atomic image under such a mapping from derivations of (the image of) \perp (under the mapping). For further details see [21].

Theorem 10. *With $n = 2$, both the consequent and the antecedent of Corollary 3 fail.*

Proof. By Theorem 3, for some Γ and A , $\Gamma \models_2^\alpha A$ and $\Gamma \not\models_{\text{IL}} A$. Now $\Gamma \models_2^\alpha A$ implies that $\forall \mathfrak{B} \in \mathbb{B}^2 (\Gamma \models_{\mathfrak{B}, 2} A)$. If the consequent of Corollary 3 holds on $n = 2$, we have $\forall \mathfrak{B} \in \mathbb{B}^2 (\Gamma \models_{\mathfrak{B}, 2}^s A)$, i.e. $\Gamma \models_2^s A$. But Theorem 9 implies $\Gamma \not\models_2^s A$. So, the consequent of Corollary 3 fails on $n = 2$. Hence, the antecedent of Corollary 3 fails on $n = 2$ too. \square

If we allow for classical logic in the meta-language, we can now appeal to Theorem 10 to refine the informal description provided above, about the first relevant consequence of Theorem 4 and Corollary 3. I.e., we can infer from Theorem 10 (by classical meta-logic) that there are Γ, A and $\mathfrak{B} \in \mathbb{B}^2$ such that $\Gamma \models_{\mathfrak{B}, 2}^\alpha A$ and $\Gamma \not\models_{\mathfrak{B}, 2}^s A$, and Γ^*, A^* and $\mathfrak{B}^* \in \mathbb{B}^2$ such that $\Gamma^* \models_{\mathfrak{B}^*, 2}^s A^*$ and $\Gamma^* \not\models_{\mathfrak{B}^*, 2}^\alpha A^*$.

As a final observation, let me also stress that, via Theorems 1 and 2, the comparative results above hold also for a comparison between standard base semantics and Sanqvist’s reading.

Theorem 11. $\forall \Gamma \forall A \forall \mathfrak{C} \supseteq_n \mathfrak{B} (\Gamma \models_{\mathfrak{C}, n}^s A \implies \Gamma \models_{\mathfrak{C}, n} A) \implies \forall \Gamma \forall A \forall \mathfrak{C} \supseteq_n \mathfrak{B} (\Gamma \models_{\mathfrak{C}, n} A \implies \Gamma \models_{\mathfrak{C}, n}^s A)$.

Corollary 6. $\forall \Gamma \forall A \forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n}^s A \implies \Gamma \models_{\mathfrak{B}, n} A) \implies \forall \Gamma \forall A \forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \models_{\mathfrak{B}, n} A \implies \Gamma \models_{\mathfrak{B}, n}^s A)$.

Theorem 12. *With $n = 2$, both the antecedent and the consequent of Corollary 6 fail.⁷*

5 On base-completeness

Besides implying that Prawitz’s reducibility semantics and Sandqvist’s base semantics are not base-comparable, Theorem 4 and Corollary 3 have as said also a second, “positive” implication. They provide a sufficient condition for both Prawitz’s reducibility semantics and Sandqvist’s base semantics to be equivalent relative to *logical* consequence, and for a logic to be complete over Prawitz’s reducibility semantics—which is relevant in itself, but especially when we restrict to atomic bases of level ≤ 1 . To see this, however, one must introduce notions of “point-wise” soundness and completeness of given logics over proof-theoretic semantics. The interest of these notions, however, does not stem from interactions with Theorem 4 and Corollary 3 only, since the very same (in)completeness phenomena that motivated the negative consequences of these results, also imply limit-results as concerns “point-wise” completeness of given logics over proof-theoretic semantics in general—in particular, that IL is not “point-wise” complete with respect to the three kinds of proof-theoretic semantics at issue here for any complexity-bound on the atomic bases.

By a *system* Σ I shall understand a recursive set of super-intuitionistic rules (over \mathcal{L}). The derivability of A from Γ in Σ is indicated as usual with the notation $\Gamma \vdash_\Sigma A$. For example, Σ may be IL. To deal with derivability at the atomic level, Σ may be required to incorporate rules from an atomic base \mathfrak{B} . This yields the following general definition.

Definition 25. *The extended derivations-set $DER_{\Sigma \cup \mathfrak{B}}$ of $\Sigma \cup \mathfrak{B}$ is defined inductively as follows:*

- *the single node labelled by $A \in FORM_{\mathcal{L}}$, possibly used as an axiom when $A \in ATOM_{\mathcal{L}}$, is a derivation in $DER_{\Sigma \cup \mathfrak{B}}$;*
- *if the following is a derivation in $DER_{\Sigma \cup \mathfrak{B}}$,*

$$\begin{array}{c} \Gamma_i, \mathfrak{C}_i, \mathfrak{R}_i \\ \mathcal{D}_i \\ A_i \end{array}$$

where $\mathfrak{C}_i, \mathfrak{R}_i$ are sets of atomic rules used in \mathcal{D}_i and A_i is the premise of an atomic rule R of the form

$$\frac{\begin{array}{ccc} [\mathfrak{R}_1] & & [\mathfrak{R}_n] \\ A_1 & \dots & A_n \end{array}}{B} R$$

⁷The same applies to the results mentioned in footnote 4.

Proposition 9. *On a disjunction-free propositional language, $\Gamma \models_{\mathfrak{B}, n} A \iff \Gamma \models_{\mathfrak{B}, n}^s A$.*

Proposition 10. *On a disjunction-free propositional language, $\Gamma \models_n A \iff \Gamma \models_n^s A$.*

The proofs of Theorems 11 and 12, Corollaries 7, and Propositions 9 and 10 are the same as in the original case.

($i \leq n$)—where it is not necessarily required that $R \in \mathfrak{B}$ —then

$$\frac{\begin{array}{c} \Gamma_1, \mathfrak{C}_1, [\mathfrak{K}_1] \\ \mathcal{D}_1 \\ A_1 \end{array} \quad \dots \quad \begin{array}{c} \Gamma_n, \mathfrak{C}_n, [\mathfrak{K}_n] \\ \mathcal{D}_n \\ A_n \end{array}}{B} R$$

is a derivation in $DER_{\Sigma \cup \mathfrak{B}}$:

- the case of the logical rules of Σ runs in a standard inductive way.

So, for example, when Σ is IL, the last clause means that derivations in $IL \cup \mathfrak{B}$ are defined by smooth induction when standard introduction and elimination rules are at issue.

Definition 26. Let \mathfrak{K} be a set of atomic rules such that $\mathfrak{K} \cap \mathfrak{B} = \emptyset$. That A is derivable from Γ, \mathfrak{K} in $\Sigma \cup \mathfrak{B}$ is indicated by $\Gamma, \mathfrak{K} \vdash_{\Sigma \cup \mathfrak{B}} A$. It holds iff there is a derivation in $DER_{\Sigma \cup \mathfrak{B}}$ from Γ to A whose only additional rules besides those in \mathfrak{B} are those in \mathfrak{K} .

In the following, \Vdash_n can indifferently be either \models_n , or \models_n^α , or \models_n^s .

Definition 27. Σ is base-complete over \Vdash_n iff $\forall \mathfrak{B} \in \mathbb{B}^n, \Gamma \Vdash_{\mathfrak{B}, n} A \implies \Gamma \vdash_{\Sigma \cup \mathfrak{B}} A$.

Definition 28. Σ is base-sound over \Vdash_n iff $\forall \mathfrak{B} \in \mathbb{B}^n, \Gamma \vdash_{\Sigma \cup \mathfrak{B}} A \implies \Gamma \Vdash_{\mathfrak{B}, n} A$.

Definition 29. Σ is sound over \Vdash_n iff $\Gamma \vdash_{\Sigma} A \implies \Gamma \Vdash_n A$.

Definition 30. Σ is complete over \Vdash_n iff $\Gamma \Vdash_n A \implies \Gamma \vdash_{\Sigma} A$.

Proposition 11. Σ base-sound over $\Vdash_n \implies \Sigma$ sound over \Vdash_n .

Proof. Take $\Gamma \vdash_{\Sigma} A$. This means $\Gamma \vdash_{\Sigma \cup \mathfrak{B}^0} A$. By assumption of base-soundness of Σ , $\Gamma \Vdash_{\mathfrak{B}^0, n} A$, i.e. $\Gamma \Vdash_n A$. \square

Proposition 12. Σ base-complete over $\Vdash_n \implies \Sigma$ complete over \Vdash_n .

Proof. Take $\Gamma \Vdash_n A$. This means $\Gamma \Vdash_{\mathfrak{B}, n} A$ for every $\mathfrak{B} \in \mathbb{B}^n$ and, by assumption of base-completeness of Σ , for every $\mathfrak{B} \in \mathbb{B}^n, \Gamma \vdash_{\Sigma \cup \mathfrak{B}} A$. By instantiating this on \mathfrak{B}^0 , $\Gamma \vdash_{\Sigma \cup \mathfrak{B}^0} A$, i.e., $\Gamma \vdash_{\Sigma} A$. \square

From this, we can immediately infer base-incompleteness of IL over \Vdash_n , where \Vdash_n is either \models_n or \models_n^α .

Proposition 13. IL is not base-complete over \Vdash_n .

Proof. From Proposition 12 plus Theorem 3 or Theorem 8. \square

Also, we can provide a sufficient condition for reducibility semantics and Sandqvist's base semantics to be equivalent.

Theorem 13. $\exists \Sigma (\Sigma \text{ base-complete over } \models_n^s \text{ and base-sound over } \models_n^\alpha) \implies \forall \Gamma \forall A (\Gamma \models_n^s A \iff \Gamma \models_n^\alpha A)$.

Proof. Let Σ be as required, and take any arbitrary Γ and A . Let us prove the direction (\implies). Assume $\Gamma \models_n^s A$. By assumption of base-completeness, $\Gamma \vdash_{\Sigma} A$. Again, by assumption of base-soundness, $\Gamma \models_n^\alpha A$. Let us prove the direction (\impliedby). Take arbitrary Δ, B and $\mathfrak{B} \in \mathbb{B}^n$, and assume $\Delta \models_{\mathfrak{B}, n}^s B$. By assumption of base-completeness, we have $\Delta \vdash_{\Sigma \cup \mathfrak{B}} B$ and, by assumption of base-soundness, we have $\Delta \models_{\mathfrak{B}, n}^\alpha B$. Hence,

$$\Delta \models_{\mathfrak{B}, n}^s B \implies \Delta \models_{\mathfrak{B}, n}^\alpha B.$$

We can now introduce universal quantification over Δ, B and $\mathfrak{B} \in \mathbb{B}^n$, so to obtain

$$\forall \Delta \forall B \forall \mathfrak{B} \in \mathbb{B}^n (\Delta \models_{\mathfrak{B}, n}^s B \implies \Delta \models_{\mathfrak{B}, n}^\alpha B).$$

By Corollary 3, this implies

$$\forall \Delta \forall B \forall \mathfrak{B} \in \mathbb{B}^n (\Delta \models_{\mathfrak{B}, n}^\alpha B \implies \Delta \models_{\mathfrak{B}, n}^s B).$$

By instantiating on our previously chosen arbitrary Γ and A , we obtain

$$\forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \vDash_{\mathfrak{B}, n}^\alpha A \implies \Gamma \vDash_{\mathfrak{B}, n}^s A).$$

Assume now $\Gamma \vDash_n^\alpha A$. This means $\forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \vDash_{\mathfrak{B}, n}^\alpha A)$, whence $\forall \mathfrak{B} \in \mathbb{B}^n (\Gamma \vDash_{\mathfrak{B}, n}^s A)$, which in turn means $\Gamma \vDash_n^s A$. \square

Corollary 7. Σ base-complete over \vDash_n^s and base-sound over $\vDash_n^\alpha \implies \Sigma$ complete over \vDash_n^α .

Proof. Take any arbitrary Σ as required and suppose $\Gamma \vDash_n^\alpha A$. Theorem 13 yields $\Gamma \vDash_n^s A$ which, by base-completeness of Σ over \vDash_n^s , implies $\Gamma \vdash_\Sigma A$.⁸ \square

Besides their conceptual interest (and their link with the results proved below), Theorem 13 and Corollary 7 might have a number of interesting applications.

For example, suppose that some logic Σ is found to be base-complete (thus, complete) on Sandqvist’s base semantics when the complexity of atomic bases has upper bound 1, and suppose further that under the same circumstances Σ is base-sound (thus, sound) over Prawitz’s reducibility semantics. Then we can immediately infer that Prawitz’s and Sandqvist’s approaches are equivalent at the level of logical consequence when the complexity of atomic bases has upper bound 1, and additionally that Σ is complete over Prawitz’s reducibility semantics with bases of this kind—it makes no sense to investigate the same when atomic bases have level ≥ 2 , as we know that in that case IL is complete over Sandqvist (and incomplete over Prawitz), so there can be no $\Sigma \neq \text{IL}$ which is base-complete over Sandqvist when atomic bases have level ≥ 2 .⁹

The latter observation, however, holds not only for Σ -s such that $\Sigma \neq \text{IL}$. For, whatever the bound on the complexity of the atomic bases is, base-completeness fails for IL and a number of other logics too. Before seeing this more in detail, let me first comment a little bit more upon the notions of base-soundness and base-completeness.

Contrarily to model-theory, where the models of one’s language are given by mappings from the language onto (mostly set-theoretic) structures, the “models” of proof-theoretic semantics are deductive in nature, that is, they are just proof-systems. In a sense—above all if one looks at Prawitz’s first semantic works [18, 19]—proof-theoretic semantics might be even understood as a sort of “semantic generalisation” of normalisation theory for Natural Deduction, namely, as a semantics where certain normalisation properties *provable* relative to given Natural Deduction systems, are turned into semantic *requirements* which determine the meaning of the logical terminology, and which provide validity criteria for argument-structures. An example of this is what Schroeder-Heister called the *fundamental corollary* of normalisation theory [24], stating that closed normal derivations in constructive systems end in introduction form. In proof-theoretic semantics, this becomes the tenet that a closed argument structure is valid (on an atomic base) when it reduces (relative to some set of reductions) to a canonical form whose immediate substructures are also valid (on the same atomic base and relative to the same set of reductions)—for more on this, see [11, 12]. More in general, this complies with the idea that meaning and validity are essentially embedded into, or even stemming from, deduction and structural features of deduction.

From this point of view, it may be natural to expect that logics which are sound or complete over a given variant \vDash of proof-theoretic consequence, are also base-sound or base-complete over \vDash (although, as proved in Proposition 11 and 12, neither soundness nor completeness imply their base version). Given the intertwining between meaning and validity, on the one hand, and deduction and its structural properties, on the other hand, it may be in other words natural to expect that, if a logic Σ is sound over \vDash , derivations in Σ from Γ to A can be turned into argument-structures from Γ to A which are valid relative to \vDash_n (and which are extracted from $\Gamma \vDash_n A$, when \vDash is \vDash or \vDash^s). Since derivations

⁸Via Theorems 1 and 2, Theorem 13 and Corollary 7 can be also formulated for standard base semantics.

Theorem 14. $\exists \Sigma (\Sigma \text{ base-complete over } \vDash_n^s \text{ and base-sound over } \vDash_n) \implies \forall \Gamma \forall A (\Gamma \vDash_n^s A \iff \Gamma \vDash_n A)$.

Corollary 8. Σ base-complete over \vDash_n^s and base-sound over $\vDash_n \implies \Sigma$ complete over \vDash .

Let me also stress that in the “deviant” reading of reducibility semantics mentioned in footnote 2—i.e., with only the left-to-right direction in (g) of Proposition 5—something similar to Theorem 13 and Corollary 7 can be obtained for a relation between \vDash_n and \vDash_n^α .

Theorem 15. $\exists \Sigma (\Sigma \text{ base-complete over } \vDash_n \text{ and base-sound over } \vDash_n^\alpha) \implies \forall \Gamma \forall A (\Gamma \vDash_n A \iff \Gamma \vDash_n^\alpha A)$.

Corollary 9. Σ base-complete over \vDash_n and base-sound over $\vDash_n^\alpha \implies \Sigma$ complete over \vDash_n^α .

The proofs of Theorem 15 and Corollary 9 are similar to those of Theorem 13 and Corollary 7, using the results mentioned in footnote 4.

⁹Let me remark in passing that the completeness or incompleteness of IL—or of other logics—is as of yet unsettled in the case of the “deviant” reading of reducibility semantics mentioned in footnote 2. Since Theorem 4 and Corollary 3 can be also proved when comparing such a “deviant” reading with Sandqvist’s base semantics, one might reason as follows: if IL turned out to be, not only complete, but more strongly *base-complete* over base semantics in Sandqvist’s reading with atomic bases of level ≥ 2 , then the completeness of IL under the same condition would follow for “deviant” reducibility semantics as well. But this strategy is unattainable since, as I shall prove below, IL is *never* base-complete over the kinds of proof-theoretic semantics at issue in this paper.

are invariant under addition of atomic bases $\mathfrak{B} \in \mathbb{B}^n$ to Σ , one may expect the same to happen when the corresponding valid argument-structures are evaluated over $\Vdash_{\mathfrak{B}, n}$. Conversely, if Σ is complete over \Vdash , one may expect that argument-structures from Γ to A which are valid relative to \Vdash (extracted from $\Gamma \Vdash_n A$, when \Vdash is \models or \models^s), can be “represented” by derivations in Σ , and that this “representation” property is stable relative to validity over $\mathfrak{B} \in \mathbb{B}^n$, namely, that an argument-structure valid relative to $\Vdash_{\mathfrak{B}, n}$ has a “representative” derivation in $\Sigma \cup \mathfrak{B}$.

Thus, besides being required for formulating and establishing Theorem 13 and Corollary 7, base-soundness and base-completeness might have an independent interest, and be worth being investigated on their own. In what follows, however, I shall not concentrate on base-soundness and base-completeness in general, but just on some results one can draw relative to Sandqvist’s proof-theoretic semantics, based on principles and facts to be found in the proof-theoretic literature—mostly due to de Campos Sanz, Piecha and Schroeder-Heister. To begin with, let us state the following basic facts—where \Vdash means again as before \models_n , or \models_n^α , or \models_n^s .

Proposition 14. *IL is base-sound over \Vdash_n .*

Corollary 10. *IL is sound over \Vdash_n .*

Let me now introduce what Piecha, de Campos Sanz and Schroeder-Heister have called the *export principle* [15]—see also [4]. This requires the preliminary definition of translation-function from an atomic base \mathfrak{B} into a set of disjunction-free formulas \mathfrak{B}^* (and vice versa).

Definition 31 (Piecha, de Campos Sanz & Schroeder-Heister, [15]). *To any rule R of level n we associate a disjunction-free formula via a function $*$ defined by induction as follows:*

- $\mathcal{Q}(R) = 0 \implies R = A$ with $A \in \text{ATOM}_{\mathcal{L}}$. Then $R^* = A$;
- $\mathcal{Q}(R) = k + 1 \implies R$ has the form

$$\frac{\begin{array}{ccc} [\mathfrak{R}_1] & & [\mathfrak{R}_n] \\ A_1 & \dots & A_n \end{array}}{A}$$

where

$$\begin{array}{ccc} \mathfrak{R}_1 & & \mathfrak{R}_n \\ A_1 & \dots & A_n \end{array}$$

are rules R_1, \dots, R_n of level $\leq k$. Then $R^* = \bigwedge_{i \leq n} R_i^* \rightarrow A$.

Definition 32 (Piecha, de Campos Sanz & Schroeder-Heister, [15]). *Given $\mathfrak{B} = \{R_1, \dots, R_n\}$, we set $\mathfrak{B}^* = \{R_1^*, \dots, R_n^*\}$.*

Piecha, de Campos Sanz and Schroeder-Heister’s export principle was defined for standard consequence over atomic bases with unlimited complexity. However, their definition can be easily adapted, both to consequence over bases with upper-bounded level, and to consequence in Sandqvist’s sense. In what follows, \Vdash_n accordingly stands for \models_n or \models_n^s (or else \models_n^α via the equivalence between the latter and \models_n).

Definition 33 (Piecha, de Campos Sanz & Schroeder-Heister, [15]). *\Vdash_n enjoys the export principle iff, for every $\mathfrak{B} \in \mathbb{B}^n$, $\Gamma \Vdash_{\mathfrak{B}, n} A \iff \Gamma, \mathfrak{B}^* \Vdash_n A$.*

Likewise, the generalised disjunction property, mentioned in Section 4 as referred to bases with unlimited level, can be now restricted to a consequence over bases with upper bound (shortly, GDP^n): for every $\mathfrak{B} \in \mathbb{B}^n$, if Γ is disjunction-free, then

$$\Gamma \Vdash_{\mathfrak{B}, n} A \vee B \implies (\Gamma \Vdash_{\mathfrak{B}, n} A \text{ or } \Gamma \Vdash_{\mathfrak{B}, n} B).$$

A result similar to Theorem 6 can be proved for GDP^n . The proof runs in very much the same way as that of Theorem 6 itself, except that, when \Vdash_n is \models_n^s , one has to use the following fact.

Proposition 15 (Sandqvist [21]). *$(\models_{\mathfrak{B}, n}^s A \text{ or } \models_{\mathfrak{B}, n}^s B) \implies \models_{\mathfrak{B}, n}^s A \vee B$.*

Theorem 16. *If GDP^n holds on every $\mathfrak{B} \in \mathbb{B}^n$, Harrop’s rule is valid in \Vdash_n namely, since Harrop’s rule is not derivable in IL, the latter is incomplete over \Vdash_n .*

Of course, since IL is complete over \models_2^s , GDP^2 fails for \models_2^s .¹⁰ However, IL is *not* base complete over \models_2^s and, more in general, it is not base-complete over \models_n^s for any n . Let us see why.

Let us recall some further results by Piecha and Schroeder-Heister [16]. Once again, Piecha and Schroeder-Heister’s proofs live in a very general setting where, not only atomic bases have not an upper-bounded level but, additionally, no constraint is put on the set of the atomic bases which the notion of consequence is defined over. Their proofs can be however adapted (or, better, restricted) to the present framework, as well as to proof-theoretic consequence in Sandqvist’s sense.

Theorem 17 (Piecha & Schroeder-Heister, [16]). *Export principle plus completeness of IL over \Vdash_n imply that GDP^n holds for every $\mathfrak{B} \in \mathbb{B}^n$.*

Proof. Suppose \vee does not occur in Γ and, for arbitrary $\mathfrak{B} \in \mathbb{B}^n$, assume $\Gamma \Vdash_{\mathfrak{B}, n} A \vee B$. Export implies that $\Gamma, \mathfrak{B}^* \Vdash_n A \vee B$ and, by completeness of IL over \Vdash_n , we have $\Gamma, \mathfrak{B}^* \vdash_{\text{IL}} A \vee B$. Given that \vee does not occur in \mathfrak{B}^* either, we have $\Gamma, \mathfrak{B}^* \vdash_{\text{IL}} A$ or $\Gamma, \mathfrak{B}^* \vdash_{\text{IL}} B$. Hence, by Corollary 10, we have $\Gamma, \mathfrak{B}^* \Vdash_n A$ or $\Gamma, \mathfrak{B}^* \Vdash_n B$. Again by export, this yields $\Gamma \Vdash_{\mathfrak{B}, n} A$ or $\Gamma \Vdash_{\mathfrak{B}, n} B$. \square

Corollary 11 (Piecha & Schroeder-Heister, [16]). *If \Vdash_n enjoys the export principle, then IL is incomplete over \Vdash_n .*

Proof. By Theorem 17, export plus completeness of IL over \Vdash_n imply that GDP^n holds for every $\mathfrak{B} \in \mathbb{B}^n$. By Theorem 16, the latter implies that Harrop’s rule is valid over \Vdash_n , hence that IL is incomplete over \Vdash_n . Hence, export plus completeness of IL over \Vdash_n imply incompleteness of IL over \Vdash_n , which means that we can reject completeness of IL over \Vdash_n under the assumption that \Vdash_n enjoys the export principle. \square

Observe, in passing, that the latter implies what follows.

Proposition 16. \models_2^s does not enjoy the export principle.

The more general point, however, is that base-completeness of IL at a given level is tantamount to export principle plus completeness of IL at that level. Let us first of all establish an export principle for IL—see [23], and see [16, Remark 3.8] for a similar point. Starting with an example, let \mathfrak{B} consist of the rules—leaving atomic explosion aside—

$$\frac{}{p} \quad \frac{p}{v} \quad \frac{q}{z} \quad \frac{r}{z} \quad \frac{[s]}{u} \quad \frac{v}{q}$$

and let \mathfrak{R} consist of the rules

$$\frac{s}{t} R_1 \quad \frac{}{r} R_2$$

then the following

$$\frac{\frac{[q \vee (t \rightarrow u)]^1}{z} \quad \frac{[q]^2}{z} \quad \frac{}{r} R_2 \quad \frac{\frac{[s]^3}{t} R_1 \quad \frac{[t \rightarrow u]^4}{u} \quad \frac{p}{v}}{q} 3 \quad \frac{}{r} R_2}{z} 2, 4}{\frac{z}{(q \vee (t \rightarrow u)) \rightarrow z} 1} \quad \frac{w \rightarrow \perp}{((q \vee (t \rightarrow u)) \rightarrow z) \wedge (w \rightarrow \perp)}$$

¹⁰Consider to this end the base $\mathfrak{B} = \{R\}$ where R is the following (schematic) rule: given $A, B, C \in \text{ATOM}_{\mathcal{L}}$, for every $D \in \text{ATOM}_{\mathcal{L}}$,

$$\frac{A \quad \frac{[B]}{D} \quad \frac{[C]}{D}}{D}$$

We have $A \models_{\mathfrak{B}} B \vee C$, but neither $A \models_{\mathfrak{B}} B$, nor $A \models_{\mathfrak{B}} C$. Incidentally, this example shows that atomic bases need not be *finite* sets of atomic rules, but *recursive* sets of atomic rule. I abstracted from specifying this above. I thank Hermógenes Oliveira for useful discussions on this topic.

witnesses that $w \rightarrow \perp, \mathfrak{K} \vdash_{\text{IL} \cup \mathfrak{B}} ((q \vee (t \rightarrow u)) \rightarrow z) \wedge (w \rightarrow \perp)$. Observe that this can be turned into

$$p, p \rightarrow v, w \rightarrow \perp, r, q \wedge r \rightarrow z, s \rightarrow t, ((s \rightarrow u) \wedge v) \rightarrow q \vdash_{\text{IL}} ((q \vee (t \rightarrow u)) \rightarrow z) \wedge (w \rightarrow \perp)$$

where each assumption but $w \rightarrow \perp$ is ρ^* with $\rho \in \mathfrak{K} \cup \mathfrak{B}$. This is what the following proposition establishes in general.

Proposition 17. $\Gamma, \mathfrak{K} \vdash_{\text{IL} \cup \mathfrak{B}} A \iff \Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} A$.

Proof. (\implies) We proceed by induction on the length $\lambda(\mathcal{D})$ of the derivation \mathcal{D} in $\text{IL} \cup \mathfrak{B}$:

- $\lambda(\mathcal{D}) = 0 \implies$ there are two cases. \mathcal{D} is an application of the assumption rule, hence $A \in \Gamma$, hence $\Gamma \vdash_{\text{IL}} A$, hence $\Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} A$. Or \mathcal{D} is an instance of an atomic rule R of level 0 in \mathfrak{B} or in \mathfrak{K} , hence $A \in \text{ATOM}_{\mathcal{D}}$ and $A = R^* \in \mathfrak{B}^*$ or $A = R^* \in \mathfrak{K}$. But then clearly $\mathfrak{B}^* \vdash_{\text{IL}} A$, resp. $\mathfrak{K}^* \vdash_{\text{IL}} A$, hence $\Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} A$;

- $\lambda(\mathcal{D}) = k + 1 \implies \mathcal{D}$ ends by

– application of $(\vee_E) \implies \mathcal{D}$ has the form

$$\frac{\begin{array}{ccc} \Gamma, \mathfrak{K} & [B], \Gamma, \mathfrak{K} & [C], \Gamma, \mathfrak{K} \\ \mathcal{D}_1 & \mathcal{D}_2 & \mathcal{D}_3 \\ \hline B \vee C & A & A \end{array}}{A}$$

So, $\Gamma, \mathfrak{K} \vdash_{\text{IL} \cup \mathfrak{B}} B \vee C$ via \mathcal{D}_1 with $\lambda(\mathcal{D}_1) \leq k$, $B, \Gamma, \mathfrak{K} \vdash_{\text{IL} \cup \mathfrak{B}} A$ via \mathcal{D}_2 with $\lambda(\mathcal{D}_2) \leq k$, and $C, \Gamma, \mathfrak{K} \vdash_{\text{IL} \cup \mathfrak{B}} A$ via \mathcal{D}_3 with $\lambda(\mathcal{D}_3) \leq k$. By induction hypothesis, there are $\mathcal{D}_1^*, \mathcal{D}_2^*$ and \mathcal{D}_3^* in IL such that $\Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} B \vee C$ holds via \mathcal{D}_1^* , $B, \Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} A$ holds via \mathcal{D}_2^* , and $C, \Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} A$ holds via \mathcal{D}_3^* . But then, $\Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} A$ holds too, via

$$\frac{\begin{array}{ccc} \Gamma, \mathfrak{K}^*, \mathfrak{B}^* & [B], \Gamma, \mathfrak{K}^*, \mathfrak{B}^* & [C], \Gamma, \mathfrak{K}^*, \mathfrak{B}^* \\ \mathcal{D}_1^* & \mathcal{D}_2^* & \mathcal{D}_3^* \\ \hline B \vee C & A & A \end{array}}{A}$$

- application of $(\wedge_I), (\wedge_E), (\vee_I), (\rightarrow_I), (\rightarrow_E)$, or $(\perp) \implies$ similar to the case for (\vee_E) ;
- application of some rule R of level n in $\mathfrak{B} \implies A \in \text{ATOM}_{\mathcal{D}}$ and \mathcal{D} has the form

$$\frac{\begin{array}{ccc} [\mathfrak{C}_1], \Gamma, \mathfrak{K} & & [\mathfrak{C}_n], \Gamma, \mathfrak{K} \\ \mathcal{D}_1 & & \mathcal{D}_n \\ \hline B_1 & \dots & B_n \end{array} R}{A}$$

with $B_i \in \text{ATOM}_{\mathcal{D}}$ and \mathfrak{C}_i set of rules of level $\leq n - 2$ discharged by R ($i \leq n$). By induction hypothesis, there is \mathcal{D}_i^* such that $\mathfrak{C}_i^*, \Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} B_i$ holds via \mathcal{D}_i^* ($i \leq n$). Observe now that we have

$$R^* = \bigwedge_{i \leq n} (\bigwedge_{\rho \in \mathfrak{C}_i} \rho^* \rightarrow B_i) \rightarrow A \in \mathfrak{B}^*.$$

So, $\Gamma, \mathfrak{K}^*, \mathfrak{B}^* \vdash_{\text{IL}} A$ holds too via

$$\frac{\frac{\frac{[\bigwedge_{\rho \in \mathfrak{C}_1} \rho^*]}{\mathfrak{C}_1^*} \quad \Gamma, \mathfrak{K}^*, \mathfrak{B}^*}{\mathcal{D}_1^*} \quad B_1}{\bigwedge_{\rho \in \mathfrak{C}_1} \rho^* \rightarrow B_1} \quad \dots \quad \frac{\frac{[\bigwedge_{\rho \in \mathfrak{C}_n} \rho^*]}{\mathfrak{C}_n^*} \quad \Gamma, \mathfrak{K}^*, \mathfrak{B}^*}{\mathcal{D}_n^*} \quad B_n}{\bigwedge_{\rho \in \mathfrak{C}_n} \rho^* \rightarrow B_n}}{\bigwedge_{i \leq n} (\bigwedge_{\rho \in \mathfrak{C}_i} \rho^* \rightarrow B_i) \rightarrow A} A$$

- application of some rule R of level n in $\mathfrak{K} \implies$ similar to the previous case, except that now we have

$$R^* = \bigwedge_{i \leq n} (\bigwedge_{\rho \in \mathfrak{C}_i} \rho^* \rightarrow B_i) \rightarrow A \in \mathfrak{K}^*.$$

(\impliedby) We proceed by induction on the length $\lambda(\mathcal{D})$ of the derivation \mathcal{D} in IL :

- $\lambda(\mathcal{D}) = 0 \implies$ there are three cases. First, $A \in \Gamma$, hence $\Gamma \vdash_{\text{IL} \cup \mathfrak{B}} A$, hence $\Gamma, \mathfrak{X} \vdash_{\text{IL} \cup \mathfrak{B}} A$. The other two cases are $A \in \mathfrak{X}^*$ or $A \in \mathfrak{B}^*$. We proceed for both by induction on the level of the rule R_A such that $R_A^* = A$. Suppose first that $A \in \mathfrak{X}^*$. We show $R_A \vdash_{\text{IL} \cup \mathfrak{B}} A$. If $R_A \in \mathfrak{X}$ has level 0, the result holds trivially. Suppose the theorem proved for all rules of level $\leq n$, and suppose that R_A has level $n + 1$. Hence, R_A has the form

$$\frac{[\mathfrak{C}_1] \quad \dots \quad [\mathfrak{C}_m]}{B} R_A$$

where \mathfrak{C}_i is a set of rules of level $\leq n - 1$ discharged by R_A ($i \leq m$). So,

$$A = R_A^* = \bigwedge_{i \leq m} (\bigwedge_{\rho \in \mathfrak{C}_i} \rho^* \rightarrow B_i) \rightarrow B.$$

By induction hypothesis, for every $i \leq m$ and every $\rho \in \mathfrak{C}_i$, $\rho \vdash_{\text{IL} \cup \mathfrak{B}} \rho^*$. This means that we have a \mathcal{D}_i in $\text{IL} \cup \mathfrak{B}$ proving $\mathfrak{C}_i \vdash_{\text{IL} \cup \mathfrak{B}} \bigwedge_{\rho \in \mathfrak{C}_i} \rho^*$. But then we can build

$$\frac{\frac{[\mathfrak{C}_1] \quad \mathcal{D}_1}{\bigwedge_{\rho \in \mathfrak{C}_1} \rho^*} \quad \frac{[\bigwedge_{i \leq m} (\bigwedge_{\rho \in \mathfrak{C}_i} \rho^* \rightarrow B_i)]}{\bigwedge_{\rho \in \mathfrak{C}_1} \rho^* \rightarrow B_1}}{B_1} \quad \dots \quad \frac{[\mathfrak{C}_m] \quad \mathcal{D}_m}{\bigwedge_{\rho \in \mathfrak{C}_m} \rho^*} \quad \frac{[\bigwedge_{i \leq m} (\bigwedge_{\rho \in \mathfrak{C}_i} \rho^* \rightarrow B_i)]}{\bigwedge_{\rho \in \mathfrak{C}_m} \rho^* \rightarrow B_m}}{B_m} R_A}{\frac{B}{\bigwedge_{i \leq m} (\bigwedge_{\rho \in \mathfrak{C}_i} \rho^* \rightarrow B_i) \rightarrow B}}$$

whence $R_A \vdash_{\text{IL} \cup \mathfrak{B}} A$. Since $R_A \in \mathfrak{X}$, hence, $\Gamma, \mathfrak{X} \vdash_{\text{IL} \cup \mathfrak{B}} A$. As regards the case when $A \in \mathfrak{B}^*$, the reasoning is the same as in the previous case, except that rules have not to be assumed, but drawn from \mathfrak{B} , so we obtain $\vdash_{\text{IL} \cup \mathfrak{B}} A$, hence $\Gamma, \mathfrak{X} \vdash_{\text{IL} \cup \mathfrak{B}} A$.

When $\lambda(\mathcal{D}) = k + 1$, \mathcal{D} will end by applying some rule from IL , so induction can proceed smoothly. \square

Corollary 12. $\Gamma \vdash_{\text{IL} \cup \mathfrak{B}} A \iff \Gamma, \mathfrak{B}^* \vdash_{\text{IL}} A$

Proof. By Proposition 16 with $\mathfrak{X} = \emptyset$. \square

Theorem 18. IL is base-complete over $\Vdash_n \iff \Vdash_n$ enjoys the export principle and IL is complete over \Vdash_n .

Proof. (\implies) That base-completeness generally implies completeness has been already established in Proposition 12. Let us show that base-completeness of IL over \Vdash_n implies that the latter enjoys the export principle. Suppose $\Gamma \Vdash_{\mathfrak{B}, n} A$. By base-completeness, $\Gamma \vdash_{\text{IL} \cup \mathfrak{B}} A$ and, by Corollary 12, $\Gamma, \mathfrak{B}^* \vdash_{\text{IL}} A$. By Corollary 10, then, $\Gamma, \mathfrak{B}^* \Vdash_n A$. (\impliedby) Suppose $\Gamma \Vdash_{\mathfrak{B}, n} A$. Since we are assuming that \Vdash_n enjoys the export principle, we have $\Gamma, \mathfrak{B}^* \Vdash_n A$ and, since we are assuming completeness of IL over \Vdash_n , we have $\Gamma, \mathfrak{B}^* \vdash_{\text{IL}} A$. But then, by Corollary 12, we have $\Gamma \vdash_{\text{IL} \cup \mathfrak{B}} A$. \square

Corollary 13. IL is not base-complete over \Vdash_n^s .

Observe that, since our proofs above were independent of the fact that atomic bases are of an intuitionistic kind, they also hold in the case when they are not, i.e., they also apply to Sandqvist's framework properly understood—that is, where \perp is understood as a nullary operator defined by a special clause, and by taking atomic bases which are not necessarily required to contain the atomic explosion rules. Similarly, we have what follows.

Theorem 19. Σ is base-sound over \Vdash , enjoys the export principle and does not derive Harrop's rule \implies for no n is Σ base-complete over \Vdash_n .

Let me just remark, to conclude, that base-incompleteness of IL over \Vdash_2^s might have been proved more directly through the observation that GDP^2 fails over Sandqvist's consequence with atomic bases of level ≥ 2 , while Corollary 12 implies the following.

Proposition 18. With Γ disjunction free, $\Gamma \vdash_{\text{IL} \cup \mathfrak{B}} A \vee B \implies \Gamma \vdash_{\text{IL} \cup \mathfrak{B}} A$ or $\Gamma \vdash_{\text{IL} \cup \mathfrak{B}} B$.

Another quicker proof of the same obtains by combining Proposition 16 and Corollary 12. Be that as it may, it seems to me to have a broader interest the fact that, for every Σ enjoying the conditions of Theorem 19, base-completeness of Σ over \Vdash_n is tantamount to export principle of Σ plus completeness of Σ over \Vdash_n —additionally, this holds for every n , and not just for $n = 2$. This is why I decided to pursue this demonstrative strategy.

6 Concluding remarks

The crucial incompleteness results obtained by de Campos Sanz, Piecha and Schroeder-Heister [14, 15, 16] are referred to a broad framework where, either atomic bases are ordered by inclusion but without requiring them to have an upper-bounded level, or else no constraint is put on the structure of the underlying set of atomic bases. The same applies to more recent findings [27, 28, 26]. The results presented in this paper are on the contrary referred to sets of atomic bases where the extension-relation is given in terms of inclusion, and where the rules-level must not be greater than a fixed bound. The last constraint was forced by the fact that, among the aims of the paper, there was that of applying the results of Section 4 to (in)completeness issues, which in turn required in some cases—such as Sandqvist’s—to focus on atomic rules of level 2. The upper bound limitation can be however dropped out, while retaining most of the results presented in this paper.¹¹ It remains to be settled whether similar results would hold also under more liberal orders on atomic bases.

Acknowledgments I am grateful to Ansten Klev, Hermógenes Oliveira, Thomas Piecha, Dag Prawitz, Peter-Schroeder-Heister, Will Stafford, and the anonymous reviewers, for precious remarks which helped me improve previous versions of this paper.

Conflict of interests The author declares that there is no conflict of interests.

References

- [1] S. Ayhan. What are acceptable reductions? Perspectives from proof-theoretic semantics and type theory. *Australasian journal of logic*, 20(3):412–428, 2023. <https://doi.org/10.26686/ajl.v20i3.7692>.
- [2] W. de Campos Sanz and T. Piecha. A critical remark on the BHK interpretation of implication. *Philosophia Scientiae*, 18(3):13–22, 2014. <https://doi.org/10.4000/philosophiascientiae.965>.
- [3] G. Gentzen. Untersuchungen über das logische Schließen i, ii. *Mathematische Zeitschrift*, 39:176–210, 405–431, 1935. <https://doi.org/10.1007/BF01201353>.

¹¹As done in Section 4, I shall indicate these unlimited notions by \models , \models^α and \models^s respectively, and I shall use the notation \Vdash to refer to any of them. So, we have what follows—I will just limit myself to the main results, but the reader should understand these as given in a context where all the other notions come with the limited consequence relations replaced by their unlimited counter-parts.

Theorem 20. $\Gamma \models_{\mathfrak{B}} A \iff \Gamma \models_{\mathfrak{B}}^\alpha A$.

Theorem 21. $\Gamma \models A \iff \Gamma \models^\alpha A$

Theorem 22. $\exists \Gamma \exists A (\Gamma \models^\alpha A \text{ and } \Gamma \not\models_{\text{IL}} A)$.

Theorem 23. $\forall \Gamma \forall A \forall \mathfrak{C} \supseteq \mathfrak{B} (\Gamma \models_{\mathfrak{C}}^s A \implies \Gamma \models_{\mathfrak{C}}^\alpha A) \implies \forall \Gamma \forall A \forall \mathfrak{C} \supseteq \mathfrak{B} (\Gamma \models_{\mathfrak{C}}^\alpha A \implies \Gamma \models_{\mathfrak{C}}^s A)$

Corollary 14. $\forall \Gamma \forall A \forall \mathfrak{B} \in \mathbb{B} (\Gamma \models_{\mathfrak{B}}^s A \implies \Gamma \models_{\mathfrak{B}}^\alpha A) \implies \forall \Gamma \forall A \forall \mathfrak{B} \in \mathbb{B} (\Gamma \models_{\mathfrak{B}}^\alpha A \implies \Gamma \models_{\mathfrak{B}}^s A)$

Theorem 24. Both the antecedent and the consequent of Corollary 15 fail.

Theorem 25. $\exists \Sigma (\Sigma \text{ base-complete over } \models^s \text{ and base-sound over } \models^\alpha) \implies \forall \Gamma \forall A (\Gamma \models^s A \iff \Gamma \models^\alpha A)$.

Corollary 15. $\Sigma \text{ base-complete over } \models^s \text{ and base-sound over } \models^\alpha \implies \Sigma \text{ complete over } \models^\alpha$.

Proposition 19. $\forall \mathfrak{B} \in \mathbb{B} (\Gamma, \mathfrak{R} \vdash_{\text{IL} \cup \mathfrak{B}} A \iff \Gamma, \mathfrak{R}^*, \mathfrak{B}^* \vdash_{\text{IL}} A)$.

Corollary 16. $\forall \mathfrak{B} \in \mathbb{B} (\Gamma \vdash_{\text{IL} \cup \mathfrak{B}} A \iff \Gamma, \mathfrak{B}^* \vdash_{\text{IL}} A)$.

Theorem 26. $\text{IL is base-complete over } \Vdash \iff \Vdash \text{ enjoys the export principle and IL is complete over } \Vdash$.

Corollary 17. $\text{IL is not base-complete over } \Vdash$.

Theorem 27. *If Σ enjoys the export principle and does not derive Harrop’s rule, then Σ is not base-complete over \Vdash .*

The proofs of these results run in very much the same way as those for their limited counter-parts throughout Sections 4 and 5 in this paper.

- [4] A. Gheorghiu and D. Pym. Definite formulae, negation-as-failure, and the base-extension semantics of intuitionistic propositional logic. *Bulletin of the section of logic*, 52(2):239–266, 2023. <https://doi.org/10.18778/0138-0680.2023.16>.
- [5] A. Gheorghiu and D. Pym. From proof-theoretic validity to base-extension semantics for intuitionistic propositional logic. [arXiv:2210.05344](https://arxiv.org/abs/2210.05344), 2024.
- [6] R. Harrop. Concerning formulas of the types $A \rightarrow B \vee C$, $A \rightarrow (\exists x)B(x)$ in intuitionistic formal systems. *Journal of symbolic logic*, 25:27–32, 1960. doi : 10.2307/2964334.
- [7] W. Howard. The formulae-as-types notion of construction. In J. Seldin and R. Hindley, editors, *To H.B. Curry: Essays on combinatory logic, lambda calculus and formalism*, pages 479–490. Academy Press, 1980.
- [8] P. Martin-Löf. *Intuitionistic type theory*. Bibliopolis, 1984.
- [9] H. Oliveira. On Dummett’s pragmatist justification procedure. *Erkenntnis*, 86:429–455, 2021. <https://doi.org/10.1007/s10670-019-00112-7>.
- [10] A. Piccolomini d’Aragona. *Prawitz’s epistemic grounding. An investigation into the power of deduction*. Synthese Library, Springer, 2023.
- [11] A. Piccolomini d’Aragona. The proof-theoretic square. *Synthese*, 201(219), 2023. <https://doi.org/10.1007/s11229-023-04203-5>.
- [12] A. Piccolomini d’Aragona. Four constructivist attitudes in Prawitzian semantics. In I. Sedlar, editor, *The Logica Yearbook 2022*. College Publications, 2024.
- [13] A. Piccolomini d’Aragona. A note on schematic validity and completeness in Prawitz’s semantics. In F. Bianchini, V. Fano, and P. Graziani, editors, *Current topics in logic and the philosophy of science. Papers from SILFS 2022 postgraduate conference*. College Publications, 2024.
- [14] T. Piecha. Completeness in proof-theoretic semantics. In T. Piecha and P. Schroeder-Heister, editors, *Advances in proof-theoretic semantics*, pages 231–251. 2016. https://doi.org/10.1007/978-3-319-22686-6_15.
- [15] T. Piecha, W. de Campos Sanz, and P. Schroeder-Heister. Failure of completeness in proof-theoretic semantics. *Journal of philosophical logic*, 44:321–335, 2015. <https://doi.org/10.1007/s10992-014-9322-x>.
- [16] T. Piecha and P. Schroeder-Heister. Incompleteness of intuitionistic propositional logic with respect to proof-theoretic semantics. *Studia Logica*, 107(1):47–62, 2019. <https://doi.org/10.1007/s11225-018-9823-7>.
- [17] D. Prawitz. *Natural deduction. A proof-theoretical study*. Almqvist & Wiskell, 1965.
- [18] D. Prawitz. Ideas and results in proof theory. In J. E. Fenstad, editor, *Proceedings of the second Scandinavian logic symposium*, pages 235–307. Elsevier, 1971. [https://doi.org/10.1016/S0049-237X\(08\)70849-8](https://doi.org/10.1016/S0049-237X(08)70849-8).
- [19] D. Prawitz. Towards a foundation of a general proof-theory. In P. Suppes, L. Henkin, A. Joja, and G. C. Moisil, editors, *Proceedings of the Fourth International Congress for Logic, Methodology and Philosophy of Science, Bucharest, 1971*, pages 225–250. Elsevier, 1973. [https://doi.org/10.1016/S0049-237X\(09\)70361-1](https://doi.org/10.1016/S0049-237X(09)70361-1).
- [20] D. Prawitz. Explaining deductive inference. In H. Wansing, editor, *Dag Prawitz on proofs and meaning*, pages 65–100. Springer, 2015.
- [21] T. Sandqvist. Base-extension semantics for intuitionistic sentential logic. *Logic journal of the IGPL*, 23(5):719–731, 2015. <https://doi.org/10.1093/jigpal/jzv021>.
- [22] P. Schroeder-Heister. A natural extension for natural deduction. *Journal of symbolic logic*, 49(4):1284–1300, 1984.
- [23] P. Schroeder-Heister. A natural extension of natural deduction. *The journal of symbolic logic*, 49(4):1284–1300, 1984.

- [24] P. Schroeder-Heister. Validity concepts in proof-theoretic semantics. *Synthese*, 148:525–571, 2006. <https://doi.org/10.1007/s11229-004-6296-1>.
- [25] P. Schroeder-Heister. Proof-theoretic semantics. In E. N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. 2018.
- [26] P. Schroeder-Heister. Prawitz’s completeness conjecture: a reassessment. *Theoria*, 90(5):515–527, 2024. <https://doi.org/10.1111/theo.12541>.
- [27] W. Stafford. Proof-theoretic semantics and inquisitive logic. *Journal of philosophical logic*, 50:1199–1229, 2021. <https://doi.org/10.1007/s10992-021-09596-7>.
- [28] W. Stafford and V. Nascimento. Following all the rules: intuitionistic completeness for generalised proof-theoretic validity. *Analysis*, 2023. <https://doi.org/10.1093/analys/anac100>.
- [29] A. S. Troelstra and D. van Dalen. *Constructivism in mathematics. Vol. I*. North-Holland, 1988.