FIXED AND RANDOM COVARIANCE REGRESSION ANALYSES

BY TAO ZOU^{1,a}, WEI LAN^{2,b}, Runze Li^{3,c} and Chih-Ling Tsai^{4,d}

¹The Australian National University, ^atao.zou@anu.edu.au

²Southwestern University of Finance and Economics, ^blanwei@swufe.edu.cn

³The Pennsylvania State University, ^crzli@psu.edu

⁴University of California, Davis, ^dcltsai@ucdavis.edu

Covariance regression analysis is an approach to linking the covariance of responses to a set of explanatory variables X, where X can be a vector, matrix, or tensor. Most of the literature on this topic focuses on the "Fixed-X" setting and treats X as nonrandom. By contrast, treating explanatory variables X as random, namely the "Random-X" setting, is often more realistic in practice. This article aims to fill this gap in the literature on the estimation and model assessment theory for Random-X covariance regression models. Specifically, we construct a new theoretical framework for studying the covariance estimators under the Random-X setting, and we demonstrate that the quasi-maximum likelihood estimator and the weighted least squares estimator are both consistent and asymptotically normal. In addition, we develop pioneering work on the model assessment theory of covariance regression. In particular, we obtain the bias-variance decompositions for the expected test errors under both the Fixed-X and Random-X settings. We show that moving from a Fixed-X to a Random-X setting can increase both the bias and the variance in expected test errors. Subsequently, we propose estimators of the expected test errors under the Fixed-X and Random-X settings, which can be used to assess the performance of the competing covariance regression models. The proposed estimation and model assessment approaches are illustrated via extensive simulation experiments and an empirical study of stock returns in the US market.

1. Introduction. Regression is one of the most popular and well-known methods in statistics, and it has been widely used across various fields. Classical regression focuses on modeling the mean of a response variable Y as a function of a set of explanatory variables X. One common assumption is to treat X as fixed, i.e., nonrandom, which Breiman and Spector (1992) and Rosset and Tibshirani (2020) refer to as "Fixed-X". This assumption is reasonable if X is constructed, such as in experimental design. Otherwise, due to real-world data collection and predictive inferences, it may be more realistic to treat X as random, namely "Random-X". The seminal work of Rosset and Tibshirani (2020) provided a detailed review of the literature for both the Fixed-X and Random-X settings in classical mean regression models. In addition, they analyzed the difference between expected test errors under these two settings, and proposed the estimators of expected test errors for model assessment of mean regression models.

Instead of classical mean regression, this paper focuses on covariance regression models, which link the covariance of the *p*-dimensional response vector Y_i to the explanatory variables X_i for $i = 1, \dots, n$. Note that X_i can be a vector, matrix, or tensor, depending on the

MSC2020 subject classifications: Primary 62H12; secondary 62J02.

Keywords and phrases: Bias-variance decomposition, covariance regression, expected test error, random-*X* regression.

context of the study, although our main focus here is on the matrix form. In addition, we assume that Y_i and X_i are random variables defined on the probability space (Ω, \mathcal{F}, P) , where Ω is the sample space, \mathcal{F} is the σ -algebra of the subsets of Ω , and P is a probability measure on (Ω, \mathcal{F}) . To ensure the covariance regression model is well-defined, we next introduce the σ -algebra $\mathcal{F}_{\mathcal{X}} = \sigma \langle X_1, \cdots, X_n \rangle \subset \mathcal{F}$, which is generated by X_1, \cdots, X_n . Then we propose the covariance regression model given below:

(1.1)
$$\operatorname{Cov}[Y_i|\mathcal{F}_{\mathcal{X}}] = C_{\beta}(X_i), \text{ for } i = 1, \cdots, n,$$

where $C_{\beta}(\cdot)$ is a pre-specified parametric and measurable map characterized by a Kdimensional parameter vector $\beta = (\beta^{(1)}, \dots, \beta^{(K)})^{\top}$ such that $C_{\beta}(X_i) \in S_p^+ \subset \mathbb{R}^{p \times p}$ for all $i = 1, \dots, n$, and $S_p^+ = \{C \in \mathbb{R}^{p \times p} : C \text{ is positive definite}\}$. This covariance regression model encompasses various covariance matrix structures such as sphericity, compound symmetry, and banding; see examples given below.

1.1. Fixed-X and Random-X settings for covariance regression models with five examples. We introduce five examples of covariance regression models. The first is the linear covariance regression model $\text{Cov}[Y_i|\mathcal{F}_{\mathcal{X}}] = C_{\beta}(X_i) = \sum_{k=1}^{K} \beta^{(k)} X_i^{(k)}$, where $X_i = (X_i^{(1)}, \dots, X_i^{(K)})$, $\beta = (\beta^{(1)}, \dots, \beta^{(K)})^{\top}$, and the $X_i^{(k)}$ s are $p \times p$ network adjacency matrices or similarity matrices constructed from their associated features. The detailed structure of this model can be found in Zou et al. (2017, 2022).

The second example is the network autoregressive model, which can be used to describe the relationship between response variables of different network nodes. Let $X_i = (X_{i,jj'})_{p \times p}$ be a symmetric $p \times p$ row-normalized network adjacency matrix with $j, j' \in \{1, \dots, p\}$ representing the *j*-th and *j'*-th nodes of the underlying network, and $X_{i,jj} = 0$. Then, the network autoregressive model has the expression of $Y_{ij} = \beta^{(1)} \sum_{j' \neq j} X_{i,jj'} Y_{ij'} + \varepsilon_{ij}$ for $j = 1, \dots, p$, whose vector-matrix form is $Y_i = \beta^{(1)} X_i Y_i + \mathcal{E}_i$ $(i = 1, \dots, n)$, where the $Y_i =$ $(Y_{i1}, \dots, Y_{ip})^{\top}$ consists of *p* measurements associated with the *p* nodes in the network, $\beta^{(1)}$ is the network autoregressive coefficient, $\mathcal{E}_i = (\varepsilon_{i1}, \dots, \varepsilon_{ip})^{\top}$ is a *p*-dimensional random vector that is independent of $\{X_i : i = 1, \dots, n\}$ with mean zero and covariance matrix $\sigma^2 I_p$, and I_p is the *p*-dimensional identity matrix; see, e.g., Zhou et al. (2017), Huang et al. (2019) and Ma et al. (2019). Based on this model, we have $\operatorname{Cov}[Y_i | \mathcal{F}_{\mathcal{X}}] = C_{\beta}(X_i) = \sigma^2 (I_p - \beta^{(1)} X_i)^{-2}$, where $\beta = (\beta^{(1)}, \sigma^2)^{\top}$.

The third example is the covariance regression model $\operatorname{Cov}[Y_i|\mathcal{F}_{\mathcal{X}}] = C_{\beta}(X_i) = \Psi + BX_iX_i^{\top}B^{\top}$ proposed by Hoff and Niu (2012). For each $i = 1, \dots, n, X_i$ is a *d*-dimensional random covariate vector. In addition, Ψ is a $p \times p$ nonrandom positive definite matrix, B is a $p \times d$ nonrandom coefficient matrix, and β consists of all the parameters in Ψ and B. Based on Hoff and Niu (2012), this model has an interpretation as a type of random-effects model $Y_i = BX_i\gamma_i + \mathcal{E}_i$, where γ_i is a one-dimensional normally distributed random coefficient with mean zero and variance 1, \mathcal{E}_i is a p-dimensional normally distributed random error vector with mean zero and covariance matrix Ψ , and $\{\gamma_i : i = 1, \dots, n\}$, $\{\mathcal{E}_i : i = 1, \dots, n\}$ and $\{X_i : i = 1, \dots, n\}$ are mutually independent.

The fourth example is that $C_{\beta}(X_i)$ is an exponential function of the random matrix (see Chiu et al., 1996). Finally, the fifth example is that $C_{\beta}(X_i)$ is a function obtained from a Bayesian nonparametric covariance regression model (Fox and Dunson, 2015); see Section S.2.1.2 in the supplementary material for details.

The above five examples can be either Fixed-X or Random-X covariance models, depending on the settings of the X_i s. For example, the second example is a random model based on the X-setting in its cited literature. On the other hand, it can represent a spatial autoregressive model (Lee, 2004 and LeSage and Pace, 2009), where $X_i = x_i$ is a spatial-weights matrix constructed by non-random geographical coordinates. Specifically, $X_1 = x_1, \dots, X_n = x_n$ are fixed in this case, and $\mathcal{F}_{\mathcal{X}}$ and the covariance model reduce to the trivial σ -algebra $\{\emptyset, \Omega\}$ and $\operatorname{Cov}[Y_i|\mathcal{F}_{\mathcal{X}}] = \operatorname{Cov}(Y_i) = C_{\beta}(x_i)$, respectively. Similar examples include x_i having different structures of geographical coordinates in spatial statistics (see, e.g., Cressie, 1993 and Wikle et al., 2019), and x_i having a form of basis functions (see, e.g., Scott and Handcock, 2001).

A special case of the Fixed-X setting is where the x_i s are all equal to x for $i = 1, \dots, n$, which leads to $\text{Cov}(Y_i) = C_\beta(x)$. This comprises various structured covariance matrices as its special cases. For example, set $x = 1_p 1_p^\top - I_p$, where 1_p is the p-dimensional vector of ones. Then it yields the compound symmetry covariance matrix $\text{Cov}(Y_i) = C_\beta(x) = \beta^{(1)}I_p + \beta^{(2)}x$, where $\beta = (\beta^{(1)}, \beta^{(2)})^\top$. Analogously, the banded covariance matrix can be represented by the form $C_\beta(x)$ with a properly designed and fixed x (see Zou et al., 2017, p. 268). This special setting of x constitutes another scenario of the Fixed-X setting.

In contrast to the Fixed-X setting, it is common to consider the X_i s as random along with Y_i . Based on the literature in the above five examples, it is not surprising to find that these examples are Random-X covariance models. For the sake of simplicity, however, some of them assumed that, once observed, $X_i = x_i$ is fixed, and the ensuing theoretical results relied on that assumption. The main difference between the fixed $X_i = x_i$ and the random X_i settings is that, the fixed $X_i = x_i$ are implicitly assumed to be independent for $i = 1, \dots, n$. This is because any nonrandom constants x_i are mutually independent for $i = 1, \dots, n$, which largely simplifies the derivation of asymptotic results for parameter estimators of covariance regression models. On the other hand, X_i is commonly dependent in real applications of these examples. Accordingly, this motivates us to propose the Random-X covariance regression model, which can largely broaden the applications of these examples.

Based on our understanding, there is no rigorous study of estimation theory under the Random-X covariance setting. One possible reason is that the covariance regression model (1.1) under the Random-X setting induces an intrinsically complicated dependence structure among the p components of Y_i . This can be even more challenging when the X_i s are dependent for $i = 1, \dots, n$, which commonly appears in the literature; see, e.g., the above five examples. Consequently, it becomes a challenging task to develop the asymptotic properties of parameter estimators, in particular when $p \to \infty$.

As for the impact of Fixed-X and Random-X on the model assessment theory and expected test errors, it has only been studied in a seminal paper of Rosset and Tibshirani (2020), who considered a univariate response variable Y_i with p = 1 for mean regression models. In contrast, the covariance regression involves a multivariate response vector Y_i with dimension p > 1. Hence, it is intrinsically challenging to develop model assessment theory based on covariance regression models. The usefulness for establishing this theory can be illustrated via an empirical application in Section S.18 of the supplementary material. This application is based on the linear covariance regression model; see the first example above. By the assessment theory of covariance models, one can select better models and then make accurate inferences. In Section S.18, we demonstrate that the selected model based on the developed model assessment theory can yield an optimal portfolio, via the Markowitz optimization approach (Markowitz, 1952), which is superior to the market portfolio in terms of the Sharpe ratio. Some references on portfolio estimation or selection can be found in DeMiguel et al. (2019), Fan et al. (2012), and Cai et al. (2020).

1.2. Our contributions. The first aim of this paper is to construct a framework for studying the estimation theory under the Random-X covariance setting without imposing a normality assumption on the data while allowing the dimension K to go to infinity. Within this framework, we propose two types of parameter estimators for random covariance regression (1.1). One is the quasi-maximum likelihood estimator (QMLE) and the other is the weighted least squares (WLS) estimator. We then demonstrate the consistency and asymptotic normality of these two estimators under the following three scenarios: (i) p is fixed and $n \to \infty$; (ii) n is fixed and $p \to \infty$; and (iii) $n \to \infty$ and $p \to \infty$; where p is the dimension of the response vector Y_i , and $i = 1, \dots, n$. Subsequently, we show that the resulting covariance matrix estimators are consistent. We further compare the asymptotic variances of the WLS estimator and QMLE, and demonstrate that the WLS estimators, and it is asymptotically as efficient as the QMLE under the normality assumption.

In sum, this paper has made three novel contributions to the estimation theory of Random-X covariance regression. (I) In deriving estimation theory, we do not impose any distributional or independent and identically distributed (i.i.d.) conditions on (Y_i, X_i) , nor do we impose any independence condition on X_i , for $i = 1, \dots, n$. Instead, we only assume some moment conditions on $C_{\beta}(X_i)$ and its first three derivatives with respect to β , and impose the convergence in probability conditions on the averages of some functions of the random X_i s. The former conditions are used to demonstrate the important distinctions between the Fixed-X and Random-X settings for estimation theory. In addition, we allow a general error form for the data generating process, the model to be misspecified, and the conditional dependence of Y_i . (II) The parametric form of $C_{\beta}(X_i)$ in (1.1) can be linear or nonlinear, which contains various models in the literature as special cases, such as all the models introduced in Section 1.1. In addition, we allow the dimensions of $C_{\beta}(X_i)$ (i.e., p) and β (i.e., K), to go to infinity. The formulations of (I) and (II) are general under the Random-X setting, and hence result in technical challenges and complexities in establishing estimation theory. (III) Compared to classical large sample approaches (see, e.g., White, 1982 and van der Vaart, 1998) that commonly assume that the (Y_i, X_i) are i.i.d. with fixed p and $n \to \infty$, we need to develop a new central limit theorem (CLT) and a new law of large numbers (LLN) for general quadratic forms with random matrices to show our asymptotic results; hence, new technical ingredients are constructed to demonstrate the new CLT and LLN.

In addition to establishing the estimation theory for the Random-X covariance regression model, the second aim of this paper is to develop pioneering work on the model assessment theory. To this end, we obtain the bias-variance decompositions for the expected test errors under both Fixed-X and Random-X settings. This allows us to propose model assessment criteria. In sum, our contributions to the model assessment theory are given below. (I) We demonstrate that randomness in X can yield additional bias and variance in the expected test errors based on the WLS estimator. Furthermore, treating the random X as fixed can cause misleading estimators of expected test errors. (II) Developing model assessment theory of covariance regression with p > 1 is more complicated than that of mean regression with p = 1 in Rosset and Tibshirani (2020); taking into account the complex dependence structure from covariance regression results in technical challenges and complexities. Hence, we need to develop new approaches to demonstrate our model assessment theories.

In conclusion, the pioneering work of this paper opens a new avenue in studying random covariance regression models. Based on the estimation and model assessment theories developed under the Random-X covariance setting, one can not only enhance the accuracy of estimation and model assessment, but can also explore new applications in the related fields.

1.3. Organization of the paper. The remainder of this paper is organized as follows. Section 2 establishes the estimation structure for the random covariance regression model. Two parameter estimators, QMLE and WLS, are proposed and their corresponding theoretical properties are obtained. Section 3 builds up the framework of model assessment, and analyzes the difference in expected test errors between the Random-X and Fixed-X settings. Section

4 provides simulation studies and discussions. Section 5 gives concluding remarks. To save space, additional simulation results, an empirical study of stock returns in the US market, and additional tables and all figures obtained from simulation studies and the empirical example, are presented in Section S.17, Section S.18, and Section S.20 of the supplementary material, respectively. All theoretical proofs, and some lemmas, propositions, theorems, assumptions and remarks are also relegated to the supplementary material.

2. Estimation and Inference with Random Covariance Regression. In this section, we propose two types of estimation approaches when the random covariance regression is correctly specified by the parametric model (1.1). Denote the true covariance regression model by

(2.1)
$$\operatorname{Cov}[Y_i|\mathcal{F}_{\mathcal{X}}] = C_{\beta_0}(X_i), \text{ for } i = 1, \cdots, n,$$

where $\beta_0 = (\beta_0^{(1)}, \dots, \beta_0^{(K)})^{\top}$ is the *K*-dimensional true parameter vector of β . Since our main interest is covariance regression, the response Y_i has been standardized to have conditional mean zero. Before studying β 's estimation and inference in the following subsections, we introduce some notation below. Let $\|\cdot\|_{\zeta}$ be the vector ζ -norm or the matrix ζ -norm for $1 \leq \zeta \leq \infty$. Specifically, for any generic vector $\alpha = (\alpha_1, \dots, \alpha_q)^{\top} \in \mathbb{R}^q$, $\|\alpha\|_{\zeta} = (\sum_{l=1}^q |\alpha_l|^{\zeta})^{1/\zeta}$, and, for any generic matrix $Z \in \mathbb{R}^{m \times q}$,

(2.2)
$$||Z||_{\zeta} = \sup\left\{\frac{||Z\alpha||_{\zeta}}{||\alpha||_{\zeta}} : \alpha \in \mathbb{R}^{q} \text{ and } \alpha \neq 0\right\}.$$

In addition, we utilize the notation $Z = (z_{k_1k_2})_{m \times q}$ to denote an $m \times q$ matrix Z with the (k_1, k_2) -th element $z_{k_1k_2}$, for $k_1 = 1, \dots, m$ and $k_2 = 1, \dots, q$, and let $|Z| = (|z_{k_1k_2}|)_{m \times q}$. If $z_{k_1k_2}$ is a function of $\beta \in \mathfrak{B}$, namely $Z = Z(\beta) = (z_{k_1k_2}(\beta))_{m \times q}$, then we denote $\sup_{\beta \in \mathfrak{B}} |Z(\beta)| = (\sup_{\beta \in \mathfrak{B}} |z_{k_1k_2}(\beta)|)_{m \times q}$. If matrix $Z = (z_{k_1k_2})_{m \times q}$ is random, then, for $1 \leq \zeta \leq \infty$, denote $||z_{k_1k_2}||_{L^{\zeta}} = (\mathbb{E}|z_{k_1k_2}|^{\zeta})^{1/\zeta}$ as the L^{ζ} -norm of the random variable $z_{k_1k_2}$. Then define

(2.3)
$$||Z||_{L^{\zeta}} = (||z_{k_1k_2}||_{L^{\zeta}})_{m \times q}$$

which is an $m \times q$ matrix consisting of the L^{ζ} -norm of $z_{k_1k_2}$. Moreover, denote

(2.4)
$$|||Z|||_{L^{\zeta_1},\zeta_2} = ||||Z||_{L^{\zeta_1}}||_{\zeta_2}$$

for $\zeta_1, \zeta_2 \in [1, \infty]$, where ζ_1 and ζ_2 correspond to the L^{ζ_1} -norm (2.3) and the matrix ζ_2 norm (2.2), respectively. Lastly, let diag $\{g_1, \dots, g_p\}$ denote a generic diagonal matrix with diagonal elements g_1, \dots, g_p , and let \circ represent the Hadamard product of any two matrices with the same dimensions.

2.1. Quasi-maximum likelihood estimation under the Random-X setting. Under parametric model (1.1), the covariance coefficients β can be estimated via the quasi-maximum likelihood estimation (QMLE) approach (e.g., see Bollerslev and Wooldridge, 1992 and Tsay, 2014). We denote this estimator by $\hat{\beta}_{QMLE}$, and it maximizes the log-likelihood function

(2.5)
$$\ell(\beta) = -\frac{np}{2}\log(2\pi) - \frac{1}{2}\sum_{i=1}^{n}\log\left[\det\left\{C_{\beta}(X_{i})\right\}\right] - \frac{1}{2}\sum_{i=1}^{n}Y_{i}^{\top}C_{\beta}^{-1}(X_{i})Y_{i},$$

where det{·} is the matrix determinant. It is worth noting that even though this log-likelihood function is formulated via the normal distribution, the consistency and asymptotic normality of QMLE $\hat{\beta}_{\text{QMLE}}$ can be established without assuming that Y_i is Gaussian distributed.

In order to establish the asymptotic normality of $\hat{\beta}_{\text{QMLE}}$, we introduce the following assumption.

ASSUMPTION 1. Assume that the data generating process is $Y_i = C_{\beta_0}^{1/2}(X_i)\mathcal{E}_i$ for $i = 1, \dots, n$, where $C_{\beta_0}^{1/2}(X_i)$ is symmetric and the square root of covariance matrix $C_{\beta_0}(X_i)$, $\mathcal{E}_i = (\varepsilon_{i1}, \dots, \varepsilon_{ip})^\top$, ε_{ij} is conditionally independent relative to the σ -algebra $\mathcal{F}_{\mathcal{X}}$ in both i and j (see the definition of conditional independence in Chung, 2001, p. 322), $E[\varepsilon_{ij}|\mathcal{F}_{\mathcal{X}}] = 0$, and $E[\varepsilon_{ij}^2|\mathcal{F}_{\mathcal{X}}] = 1$ for all $i = 1, \dots, n$ and $j = 1, \dots, p$. In addition, assume that there exists a finite constant $\eta_1 > 0$ such that $\sup_{i,j} E|\varepsilon_{ij}|^{4+\eta_1} < \infty$.

Under Assumption 1, model (2.1) is the true model and not misspecified. In addition, if the \mathcal{E}_i s are independent and also independent of X_i , then $Y_i = C_{\beta_0}^{1/2}(X_i)\mathcal{E}_i$ is similar to the independent component model (ICM, see, e.g., Bai and Silverstein, 2004, Pan and Zhou, 2008, and Zheng, 2012) with random and exogenous X_i involved. Although the main results presented in this section are based on the multiplicative error form $Y_i = C_{\beta_0}^{1/2}(X_i)\mathcal{E}_i$, a more general error form for obtaining the estimation theory of model (2.1) is given in Section S.8 of the supplementary material. In addition, by Assumption 1, we have that $E[\mathcal{E}_i \mathcal{E}_i^\top | \mathcal{F}_{\mathcal{X}}] = I_p$. Accordingly,

(2.6)
$$\operatorname{E}[Y_i Y_i^{\top} | \mathcal{F}_{\mathcal{X}}] = C_{\beta_0}(X_i),$$

which determines the conditional dependence structure for the p components in the response vector Y_i . Furthermore, Section S.2.1.1 of the supplementary material demonstrates that $\operatorname{Cov}\left(\operatorname{vec}(Y_{i_1}Y_{i_1}^{\top}), \operatorname{vec}(Y_{i_2}Y_{i_2}^{\top})\right) = \operatorname{Cov}\left(\operatorname{vec}\left\{C_{\beta_0}(X_{i_1})\right\}, \operatorname{vec}\left\{C_{\beta_0}(X_{i_2})\right\}\right) \neq 0$ for $i_1, i_2 \in$ $\{1, \dots, n\}$ and $i_1 \neq i_2$, where $\operatorname{vec}(G)$ denotes the vectorization for any generic matrix G; this is because we do not impose any independent condition on X_i for $i = 1, \dots, n$. Accordingly, Y_i is dependent for $i = 1, \dots, n$. Although Y_i is conditionally independent by Assumption 1, the estimation theory of model (2.1) can be established by relaxing this conditional independence assumption; see Section S.9 of the supplementary material. More remarks on Assumption 1, such as interpretations, the examination of Assumption 1 with regard to the five examples in Section 1.1, and the significance and necessity of this assumption, are provided in Section S.2.1 of the supplementary material. We next introduce the assumption for $C_{\beta}(X_i)$ in model (2.1).

ASSUMPTION 2. Assume that, for $i = 1, \dots, n$, $C_{\beta}(X_i)$ denotes an indexed series of positive definite matrices, uniformly over β that lies in a compact parameter space $\mathfrak{B} \subset \mathbb{R}^K$, $K < \infty$, and β_0 is an interior point of \mathfrak{B} . In addition, assume that $C_{\beta}(x_i)$ is three times differentiable with respect to β for P_{X_i} -almost every x_i (see van der Vaart, 1998, p. 53), and $C_{\beta}(x_i)$ and its first three derivatives with respect to β are all measurable functions of x_i , where $P_{X_i} = \mathbb{P} \circ X_i^{-1}$ (see Chung, 2001, p. 36) is the induced measure from the probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Based on Assumption 2, the main results in this section focus on the dimension of β , K, being fixed. The estimation theory of model (2.1) allowing $K \to \infty$ is given in Section S.7 of the supplementary material. Under Assumptions 1 - 2, one can derive the quasi-score function evaluated at β_0 , which is

$$\frac{\partial \ell(\beta_0)}{\partial \beta^{(k)}} = \frac{1}{2} \sum_{i=1}^n Y_i^\top C_{\beta_0}^{-1}(X_i) \frac{\partial C_{\beta_0}(X_i)}{\partial \beta^{(k)}} C_{\beta_0}^{-1}(X_i) Y_i - \frac{1}{2} \sum_{i=1}^n \operatorname{tr} \left\{ C_{\beta_0}^{-1}(X_i) \frac{\partial C_{\beta_0}(X_i)}{\partial \beta^{(k)}} \right\}$$

(2.7)
$$= \frac{1}{2} \sum_{i=1}^n \mathcal{E}_i^\top C_{\beta_0}^{-1/2}(X_i) \frac{\partial C_{\beta_0}(X_i)}{\partial \beta^{(k)}} C_{\beta_0}^{-1/2}(X_i) \mathcal{E}_i - \frac{1}{2} \sum_{i=1}^n \operatorname{tr} \left\{ C_{\beta_0}^{-1}(X_i) \frac{\partial C_{\beta_0}(X_i)}{\partial \beta^{(k)}} \right\}$$

7

for $k = 1, \dots, K$, where, for any generic differentiable function $q(\beta)$ of β , we denote $\partial g(\beta)/\partial \beta^{(k)}$ evaluated at β_0 by $\partial g(\beta_0)/\partial \beta^{(k)}$ to save space. It is worth noting that Random-X raises challenges for obtaining the asymptotic properties of the parameter estimators under scenarios of "fixed n and $p \to \infty$ " and " $n \to \infty$ and $p \to \infty$ ". For instance, to derive the asymptotic normality of QMLE, the central limit theorem (CLT) needs to be developed for $(np)^{-1/2}\partial\ell(\beta_0)/\partial\beta^{(k)}$. Note that (2.7) includes a summation of the quadratic forms in \mathcal{E}_i with the random matrices $C_{\beta_0}^{-1/2}(X_i) \{\partial C_{\beta_0}(X_i)/\partial \beta^{(k)}\} C_{\beta_0}^{-1/2}(X_i)$, and $i = 1, \dots, n$. Accordingly, the standard asymptotic theories for quadratic forms with nonstochastic matrices (Kelejian and Prucha, 2001) cannot be directly applied due to the random X_i . In addition, to show the asymptotic normality of β_{QMLE} , the law of large numbers (LLN) is needed for the second derivative evaluated at β_0 , i.e., $(np)^{-1}\partial^2 \ell(\beta_0)/(\partial\beta^{(k_1)}\partial\beta^{(k_2)})$, which also contains the quadratic forms in \mathcal{E}_i with random matrices; see equation (S.50) in the supplementary material. To solve the problem, we propose a new CLT and a new LLN for the summation of quadratic forms in \mathcal{E}_i with general random matrices. Note that the new CLT and LLN can be applied under any one of the following three scenarios: (i) p is fixed and $n \to \infty$; (ii) n is fixed and $p \to \infty$; and (iii) $n \to \infty$ and $p \to \infty$; see Lemma S.10 and its extensive proof in Section S.11 of the supplementary material. Furthermore, the proof sketch for the CLT and the new technical ingredients required in the proof are presented in Sections S.13 of the supplementary material. By Lemma S.10, we demonstrate that β_{QMLE} is consistent and asymptotically normal under the two additional assumptions (i.e., Assumptions 3 - 4) on the random matrices $C_{\beta}(X_i)$ given below. To save space, the remarks of interpretations, examinations, significances, and necessities of Assumption 2 as well as Assumptions 3 - 4are given in Sections S.2.2 - S.2.4 of the supplementary material.

ASSUMPTION 3. Assume that there exists finite $\eta_2 > 8/\eta_1$ such that

$$(2.8) \max\left\{\sup_{p\geq 1}\sup_{i}\left\|\left\|C_{\beta_{0}}^{1/2}(X_{i})\right\|\right\|_{L^{\delta_{c}(2+\eta_{2})},1},\sup_{p\geq 1}\sup_{i}\left\|\left\|\sup_{\beta\in\mathfrak{B}}\left|C_{\beta}^{-1}(X_{i})\right|\right\|_{L^{\delta_{0}(2+\eta_{2})},1}\right.\right.\right.$$
$$\left.\sup_{p\geq 1}\sup_{i}\left\|\left\|\sup_{\beta\in\mathfrak{B}}\left|\frac{\partial C_{\beta}(X_{i})}{\partial\beta^{(k)}}\right|\right\|\right\|_{L^{\delta_{1}(2+\eta_{2})},1},\sup_{p\geq 1}\sup_{i}\left\|\left\|\sup_{\beta\in\mathfrak{B}}\left|\frac{\partial^{2} C_{\beta}(X_{i})}{\partial\beta^{(k_{1})}\partial\beta^{(k_{2})}}\right|\right\|_{L^{\delta_{2}(2+\eta_{2})},1}\right.\right\}<\infty,$$

for any $k, k_1, k_2, k_3 \in \{1, \dots, K\}$, and for some $\delta_c, \delta_0, \delta_1, \delta_2, \delta_3 \in (0, \infty)$ that satisfy

$$\max\left\{ \left(\frac{2}{\delta_c} + \frac{4}{\delta_0} + \frac{3}{\delta_1}\right), \left(\frac{2}{\delta_c} + \frac{3}{\delta_0} + \frac{1}{\delta_1} + \frac{1}{\delta_2}\right), \left(\frac{2}{\delta_c} + \frac{2}{\delta_0} + \frac{1}{\delta_3}\right) \right\} \le 1,$$

where η_1 is defined in Assumption 1.

Before introducing the next assumption, define

(2.9)
$$A_{i,Q}^{(k)} = \frac{1}{2} C_{\beta_0}^{-1/2}(X_i) \frac{\partial C_{\beta_0}(X_i)}{\partial \beta^{(k)}} C_{\beta_0}^{-1/2}(X_i)$$

and $\Upsilon_i^{(4)} = \operatorname{diag}\{\mu_{i1}^{(4)} - 3, \cdots, \mu_{ip}^{(4)} - 3\} \in \mathbb{R}^{p \times p}$, where $\mu_{ij}^{(4)} = \operatorname{E}[\varepsilon_{ij}^4 | \mathcal{F}_{\mathcal{X}}]$, and "Q" used in the subscript is an abbreviation of "QMLE" to save space. In addition, denote $V_{np,Q} = 2(np)^{-1} \sum_{i=1}^n \left(\operatorname{tr}(A_{i,Q}^{(k_1)} A_{i,Q}^{(k_2)}) \right)_{K \times K}$ and $\Omega_{np,Q} = (np)^{-1} \sum_{i=1}^n \left(\operatorname{tr}(A_{i,Q}^{(k_1)} \circ \Upsilon_i^{(4)} \circ \Lambda_i^{(4)}) \right)_{K \times K}$.

ASSUMPTION 4. As $\max\{n, p\} \to \infty$, we have that $V_{np,Q} \xrightarrow{P} V_Q$ and $\Omega_{np,Q} \xrightarrow{P} \Omega_Q$, where V_Q and Ω_Q are finite $K \times K$ matrices. In addition, assume both V_Q and $V_Q + \Omega_Q$ are positive definite.

It is worth noting that Assumptions 1 - 4 do not impose any distributional and i.i.d. conditions on (Y_i, X_i) , or an independent condition on X_i , for $i = 1, \dots, n$. Instead, we only assume some moment conditions on $C_{\beta}(X_i)$ and its first three derivatives with respect to β in Assumption 3, and impose the convergence in probability conditions on the averages of some functions of random X_i s in Assumption 4. The conditions in Assumption 3 can be used to demonstrate the important distinctions between the Fixed-X and Random-X settings for estimation theory; see Remark S.1 in Section S.2.3.2 of the supplementary material. In the two examples demonstrating the justification for Assumptions 3 - 4 presented in Section S.19 of the supplementary material, we find that, although Assumption 4 does not directly involve a distributional condition on X_i , the justification for Assumption 4 possibly requires some distributional and finite moment conditions on the functions of X_i . More remarks on Assumptions 3 and 4 are presented in Sections S.2.3 and S.2.4, respectively, of the supplementary material.

The above assumptions lead to the following conclusion.

THEOREM 1. Under Assumptions
$$1 - 4$$
, we have that, as $\max\{n, p\} \to \infty$,
 $\sqrt{np} \left(\hat{\beta}_{\text{QMLE}} - \beta_0\right) \xrightarrow{d} N(0, V_{\text{Q}}^{-1} + V_{\text{Q}}^{-1}\Omega_{\text{Q}}V_{\text{Q}}^{-1}).$

In Theorem 1 and the theorems developed below, we unify the following three scenarios by $\max\{n, p\} \to \infty$: (i) p is fixed and $n \to \infty$; (ii) n is fixed and $p \to \infty$; and (iii) $n \to \infty$ and $p \to \infty$. Note that the proofs of all the theorems and propositions in this paper are presented in Sections S.14 – S.15 of the supplementary material. In order to help better understand the key steps of showing the theorems and propositions, we have provided the flow charts and discussions in Section S.12 of the supplementary material. In addition to the QMLE, we next consider a least squares type estimator.

2.2. Weighted least squares estimation under the Random-X setting. Without the distribution assumption, it is natural to consider the weighted least squares (WLS) approach to obtain parameter estimators. To this end, let $\mathbf{Y}_i = \operatorname{vec}(Y_i Y_i^{\top})$ and $\mathbf{C}_{\beta}(X_i) = \operatorname{vec} C_{\beta}(X_i)$. For the given β , we define a p^2 -dimensional error vector $\mathbf{Y}_i - \mathbf{C}_{\beta}(X_i)$. In order to aggregate each component of $\mathbf{Y}_i - \mathbf{C}_{\beta}(X_i)$ and obtain a total error, we further introduce a $p^2 \times p^2$ positive semidefinite weight matrix $\mathbf{W}(X_i)$, which depends on X_i . Afterwards, we define the total error by

(2.10)
$$Tr(\beta) = \frac{1}{n} \sum_{i=1}^{n} \left\{ \mathbf{Y}_{i} - \mathbf{C}_{\beta}(X_{i}) \right\}^{\top} \mathbf{W}(X_{i}) \left\{ \mathbf{Y}_{i} - \mathbf{C}_{\beta}(X_{i}) \right\}.$$

Accordingly, the WLS estimator $\hat{\beta}_{WLS}$ can be obtained by minimizing $Tr(\beta)$. First let $\mathbf{W}(X_i) = I_{p^2}$. Then minimizing (2.10) leads to the ordinary least squares (OLS) estimator. To improve the efficiency of the OLS estimator, let $\mathbf{W}(X_i) = (\text{Cov}[\mathbf{Y}_i|X_i])^-$ be the generalized (Moore-Penrose) inverse of $\text{Cov}[\mathbf{Y}_i|X_i]$. We then obtain a generalized least squares (GLS) estimator $\hat{\beta}_{\text{GLS}}$ by minimizing (2.10). Theorem S.2 in Section S.5.2 of the supplementary material shows that $\hat{\beta}_{\text{GLS}}$ is asymptotically the most efficient among all $\hat{\beta}_{\text{WLS}}$. It is worth noting that $(\text{Cov}[\mathbf{Y}_i|X_i])^-$ is only positive semidefinite but not positive definite, which motivates us to consider a positive semidefinite matrix $\mathbf{W}(X_i)$ in obtaining the WLS estimator.

REMARK 1. The closed form of $\text{Cov}[\mathbf{Y}_i|X_i] = \mathbf{V}(X_i)$ in finding the GLS estimator is given in Theorem S.1 of Section S.5.1 of the supplementary material, and it depends on β_0 . Substituting β_0 with the OLS estimator yields an estimator $\hat{\mathbf{V}}(X_i)$ of $\mathbf{V}(X_i)$; the details of obtaining $\hat{\mathbf{V}}(X_i)$ are given in Section S.5.1 of the supplementary material. Following the GLS standard procedure in classical mean regression models (see, e.g., Wooldridge, 2012), we then obtain a feasible GLS (FGLS) estimator by minimizing (2.10) with $\mathbf{W}(X_i) = \hat{\mathbf{V}}^-(X_i)$. In fact, this procedure can be applied for any $\mathbf{W}(X_i)$ that depends on β_0 .

Analogous to the derivation of the asymptotic normality of QMLE, the asymptotic property of WLS relies on the first- and second-order derivatives of $Tr(\beta)$ evaluated at β_0 , i.e., $\partial Tr(\beta_0)/\partial\beta$ and $\partial^2 Tr(\beta_0)/(\partial\beta\partial\beta^{\top})$. Since both derivatives involve a summation of quadratic forms with random matrices, Lemma S.10 in Section S.11 of the supplementary material is used to establish the asymptotic normality of $\hat{\beta}_{WLS}$, which is given below. Note that the required Assumptions 3' and 4' for WLS are parallel to Assumptions 3 and 4 for QMLE. To save space, they are presented in Section S.3 of the supplementary material.

THEOREM 2. Under Assumptions 1, 2, 3' and 4', we have that, as $\max\{n, p\} \to \infty$,

$$\sqrt{np}\left(\hat{\beta}_{\mathrm{WLS}}-\beta_{0}\right) \xrightarrow{d} N\left(0, M_{\mathrm{W}}^{-1}(V_{\mathrm{W}}+\Omega_{\mathrm{W}})M_{\mathrm{W}}^{-1}\right),$$

where V_W , Ω_W and M_W are $K \times K$ matrices defined in Assumption 4', and "W" used in the subscript is an abbreviation of "WLS" to save space.

It is worth noting that Theorem 2 is based on a known weight matrix $W(X_i)$, and we can demonstrate that this theorem is also valid for the FGLS estimator mentioned in Remark 1.

REMARK 2. Theorems 1 and 2 can lead to the asymptotic variances of $\hat{\beta}_{QMLE}$, $\hat{\beta}_{WLS}$, and $\hat{\beta}_{GLS}$, and they are, correspondingly, $\operatorname{Avar}(\hat{\beta}_{QMLE}) = V_Q^{-1} + V_Q^{-1}\Omega_Q V_Q^{-1}$, $\operatorname{Avar}(\hat{\beta}_{WLS}) = M_W^{-1}(V_W + \Omega_W)M_W^{-1} |_{\mathbf{W}(X_i)=\mathbf{V}^-(X_i)}$. We then compare these asymptotic variances, and show that the GLS estimator is asymptotically the most efficient among all WLS estimators, and it is asymptotically as efficient as the QMLE under the normality assumption; see Theorems S.2 – S.3 in Section S.5.2 of the supplementary material. The connection between the above optimality of GLS and the standard GLS' optimality result in classical mean regression models (see, e.g., Wooldridge, 2012) is given in Remark S.2 of the supplementary material.

REMARK 3. Based on the above estimation approaches, the covariance matrix estimator can be obtained by substituting β in $C_{\beta}(X_i)$ with $\hat{\beta}$, where $\hat{\beta}$ can be either $\hat{\beta}_{QMLE}$ or $\hat{\beta}_{WLS}$. However, to assure the positive definiteness of the covariance matrix estimator, we need to impose the constraint $\beta \in B^+ = \{\beta : C_{\beta}(X_i) \succ 0, \text{ for all } i = 1, \dots, n\}$, where $C_{\beta}(X_i) \succ 0$ indicates that $C_{\beta}(X_i)$ is positive definite. Under this constraint, we obtain the constrained estimator $\hat{\beta}^c = \arg \min_{\beta \in B^+} M(\beta)$, where $M(\beta)$ can be $-\ell(\beta)$ and $n Tr(\beta)$ used for obtaining the constrained QMLE and WLS estimators, respectively. The algorithm for obtaining the constrained estimator is presented in Section S.16 of the supplementary material. In addition, employing Theorems 1 - 2, we can show that the unconstrained estimator is identical to the constrained estimator with probability tending to one, as $\max\{n, p\} \rightarrow \infty$; see the proof after proving Theorem 2 in Section S.15 of the supplementary material. Consequently, $\hat{\beta}$ and $\hat{\beta}^c$ are asymptotically equivalent. 10

REMARK 4. Theorems 1-2 are established under Assumption 1, which indicates that model (2.1) is the true model and not misspecified. When model (2.1) is misspecified, we have demonstrated that the proposed QMLE and WLS estimators are asymptotically normal; see Theorems S.4 – S.5 in Section S.6 of the supplementary material. The numerical studies of parameter estimation under the misspecified model are presented in Section S.17 of the supplementary material.

Recall that the first example in Section 1.1 introduces a linear covariance regression model,

(2.11)
$$C_{\beta}(X_i) = \sum_{k=1}^{K} \beta^{(k)} X_i^{(k)}.$$

Under this model, both of Zou et al.'s (2017) estimators (OLS and FGLS) are special cases of our proposed WLS estimator. Denote $\mathbf{X}_i = (\operatorname{vec}(X_i^{(1)}), \cdots, \operatorname{vec}(X_i^{(K)}))$. Then, the resulting WLS estimator has a closed form

(2.12)
$$\hat{\beta}_{\text{WLS}} = \left\{ \sum_{i=1}^{n} \mathbf{X}_{i}^{\top} \mathbf{W}(X_{i}) \mathbf{X}_{i} \right\}^{-1} \sum_{i=1}^{n} \mathbf{X}_{i}^{\top} \mathbf{W}(X_{i}) \mathbf{Y}_{i}$$

The WLS reduces to OLS by setting $W(X_i) = I_{p^2}$, and the resulting OLS estimator has a closed form

$$\hat{\beta}_{\text{OLS}} = \left(\sum_{i=1}^{n} \mathbf{X}_{i}^{\top} \mathbf{X}_{i}\right)^{-1} \sum_{i=1}^{n} \mathbf{X}_{i}^{\top} \mathbf{Y}_{i} = \left(\sum_{i=1}^{n} \operatorname{tr}(X_{i}^{(k_{1})} X_{i}^{(k_{2})})\right)^{-1}_{K \times K} \left(\sum_{i=1}^{n} Y_{i}^{\top} X_{i}^{(k)} Y_{i}\right)_{K \times 1}.$$

Based on the theoretical properties of $\hat{\beta}_{\text{QMLE}}$ and $\hat{\beta}_{\text{WLS}}$, we next show the stochastic convergence rate of the estimated covariance matrix.

2.3. Inference with Random-X covariance regression. Applying the results of Theorems 1 and 2, we can make inferences on the conditional covariance matrix $Cov[Y_i|\mathcal{F}_{\mathcal{X}}] = C_{\beta_0}(X_i)$. Let $||G||_F$ be the Frobenius norm for any generic matrix G. To study the asymptotic properties of the estimators of $C_{\beta_0}(X_i)$ and its inverse under the Frobenius norm, we introduce an assumption below.

ASSUMPTION 5. For any fixed *i*, as $p \to \infty$, assume that

$$p^{-1}\left(\operatorname{tr}\left\{C_{\beta_{0}}^{l}(X_{i})\frac{\partial C_{\beta_{0}}(X_{i})}{\partial\beta^{(k_{1})}}C_{\beta_{0}}^{l}(X_{i})\frac{\partial C_{\beta_{0}}(X_{i})}{\partial\beta^{(k_{2})}}\right\}\right)_{K\times K} \xrightarrow{\mathrm{P}} Q_{l,i},$$

for l = -2 and l = 0, where $C^0_{\beta_0}(X_i) \triangleq I_p$ and $i = 1, \dots, n$.

This assumption is a generalization of Condition (C3) from Zou et al. (2017) to random matrices $C_{\beta_0}(X_i)$ and $\partial C_{\beta_0}(X_i)/\partial \beta^{(k)}$, and it is analogous to the standard condition imposed on the asymptotic covariance matrix of the OLS estimator obtained from classical mean regression models under the Random-X setting.

We next present the theoretical properties of the estimators of $C_{\beta_0}(X_i)$ and the precision matrix given below, under Assumptions 1 – 3 and Assumption 5.

THEOREM 3. Under Assumptions 1 - 3 and Assumption 5, for any fixed *i*, as $p \to \infty$, and with *n* either fixed or going to infinity, we have that

$$n \left\| C_{\hat{\beta}}(X_i) - C_{\beta_0}(X_i) \right\|_F^2 \xrightarrow{d} Z^\top Q_{0,i}Z, \text{ and } n \left\| C_{\hat{\beta}}^{-1}(X_i) - C_{\beta_0}^{-1}(X_i) \right\|_F^2 \xrightarrow{d} Z^\top Q_{-2,i}Z,$$

given that $C_{\hat{\beta}}(X_i)$ is invertible a.s., where $\hat{\beta}$ is an estimator satisfying $\sqrt{np}(\hat{\beta} - \beta_0) \xrightarrow{d} Z$ with Z following a K-dimensional multivariate normal distribution.

Theorem 3 indicates that the orders of $p^{-1/2} \|C_{\hat{\beta}}(X_i) - C_{\beta_0}(X_i)\|_F$ and $p^{-1/2} \|C_{\hat{\beta}}^{-1}(X_i) - C_{\beta_0}(X_i)\|_F$ are of $(np)^{-1/2}$. Under the spectral norm $\|\cdot\|_2$, we can obtain the orders of $\|C_{\hat{\beta}}(X_i) - C_{\beta_0}(X_i)\|_2$ and $\|C_{\hat{\beta}}^{-1}(X_i) - C_{\beta_0}^{-1}(X_i)\|_2$ given below.

THEOREM 4. Under Assumptions 1 - 3, for any fixed *i*, we have that, as $\max\{n, p\} \rightarrow \infty$,

$$\left\| C_{\hat{\beta}}(X_i) - C_{\beta_0}(X_i) \right\|_2 = O_{\mathcal{P}}\left(\frac{1}{\sqrt{np}}\right), \text{ and } \left\| C_{\hat{\beta}}^{-1}(X_i) - C_{\beta_0}^{-1}(X_i) \right\|_2 = O_{\mathcal{P}}\left(\frac{1}{\sqrt{np}}\right),$$

given that $C_{\hat{\beta}}(X_i)$ is invertible a.s., where $\hat{\beta}$ is described in Theorem 3.

Based on Theorems 1 and 2, the estimators $\hat{\beta}_{QMLE}$ and $\hat{\beta}_{WLS}$ satisfy the asymptotic distribution assumptions in Theorems 3 and 4, respectively. Thus, the theoretical results in Theorems 3 and 4 are respectively applicable to these estimators. In addition, Theorems 1 – 4 are developed under the Random-X setting. Since Fixed-X is a special case of Random-X, the asymptotic properties in Theorems 1 – 4 still hold under the Fixed-X setting.

It is worth noting that the asymptotic variances of the QMLE and the WLS estimator in Theorems 1 and 2, respectively, are unknown in practice. Hence, we need to find their corresponding estimators. For the sake of convenience, let $\hat{\beta}$ denote the consistent estimator of β_0 . In both Theorems 1 and 2, we also let V, Ω and M denote the quantities that appeared in the asymptotic variance, $\operatorname{Avar}(\hat{\beta})$. To estimate V, Ω and M, we replace β_0 in (2.9) and (S.21) in Section S.3 of the supplementary material by $\hat{\beta}$, which yields the corresponding estimators $\widehat{A}_{i,Q}^{(k)}$ and $\widehat{A}_{i,W}^{(k)}$. Moreover, let $\widehat{\Upsilon}_i^{(4)} = \operatorname{diag}\{\widehat{\varepsilon}_{i1}^4 - 3, \cdots, \widehat{\varepsilon}_{ip}^4 - 3\} \in \mathbb{R}^{p \times p}$, where $(\widehat{\varepsilon}_{i1}, \cdots, \widehat{\varepsilon}_{ip})^{\top} \triangleq \widehat{\mathcal{E}}_i = C_{\hat{\beta}}^{-1/2}(X_i)Y_i$. Then, we obtain the estimators of V, Ω and M, which are given below:

(2.13)

$$\begin{split} \widehat{V}_{np,\mathbf{Q}} &= \frac{2}{np} \sum_{i=1}^{n} \left(\operatorname{tr}(\widehat{A}_{i,\mathbf{Q}}^{(k_{1})} \widehat{A}_{i,\mathbf{Q}}^{(k_{2})}) \right)_{K \times K}, \widehat{\Omega}_{np,\mathbf{Q}} = \frac{1}{np} \sum_{i=1}^{n} \left(\operatorname{tr}(\widehat{A}_{i,\mathbf{Q}}^{(k_{1})} \circ \widehat{\Upsilon}_{i}^{(4)} \circ \widehat{A}_{i,\mathbf{Q}}^{(k_{2})}) \right)_{K \times K}, \\ \widehat{V}_{np,\mathbf{W}} &= \frac{2}{np} \sum_{i=1}^{n} \left(\operatorname{tr}(\widehat{A}_{i,\mathbf{W}}^{(k_{1})} \widehat{A}_{i,\mathbf{W}}^{(k_{2})}) \right)_{K \times K}, \widehat{\Omega}_{np,\mathbf{W}} = \frac{1}{np} \sum_{i=1}^{n} \left(\operatorname{tr}(\widehat{A}_{i,\mathbf{W}}^{(k_{1})} \circ \widehat{\Upsilon}_{i}^{(4)} \circ \widehat{A}_{i,\mathbf{W}}^{(k_{2})}) \right)_{K \times K}, \\ \text{and } \widehat{M}_{np,\mathbf{W}} &= \frac{1}{np} \sum_{i=1}^{n} \left(\operatorname{vec}^{\top} \left\{ \frac{\partial C_{\hat{\beta}}(X_{i})}{\partial \beta^{(k_{1})}} \right\} \mathbf{W}(X_{i}) \operatorname{vec} \left\{ \frac{\partial C_{\hat{\beta}}(X_{i})}{\partial \beta^{(k_{2})}} \right\} \right)_{K \times K}. \end{split}$$

Note that Theorem 3 and Theorem 4 separately show that $C_{\hat{\beta}}(X_i)$ and $C_{\hat{\beta}}^{-1}(X_i)$ are consistent estimators of $C_{\beta_0}(X_i)$ and $C_{\beta_0}^{-1}(X_i)$. Using these results, we can verify that $\widehat{A}var(\hat{\beta}_{QMLE}) = \widehat{V}_{np,Q}^{-1} + \widehat{V}_{np,Q}^{-1}\widehat{\Omega}_{np,Q}\widehat{V}_{np,Q}^{-1}$ and $\widehat{A}var(\hat{\beta}_{WLS}) = \widehat{M}_{np,W}^{-1}\widehat{V}_{np,W}\widehat{M}_{np,W}^{-1} + \widehat{M}_{np,W}^{-1}\widehat{\Omega}_{np,W}\widehat{M}_{np,W}^{-1}$ are consistent estimators of $Avar(\hat{\beta}_{QMLE})$ and $Avar(\hat{\beta}_{WLS})$, respectively, which are the asymptotic variances of $\hat{\beta}_{QMLE}$ and $\hat{\beta}_{WLS}$. Using them, we can make inferences regarding β such as constructing confidence intervals and testing hypotheses.

Finally, the quantities $Q_{0,i}$ and $Q_{-2,i}$, which appear in the asymptotic distributions of Theorem 3, are unknown. In practice, they can be replaced by the consistent estimators. Specifically, replacing β_0 , which appears in Assumption 5, by $\hat{\beta}_{\text{QMLE}}$ (or $\hat{\beta}_{\text{WLS}}$), yields the consistent estimators of $Q_{0,i}$ and $Q_{-2,i}$. After studying parameter estimation and inference, we next focus on model assessment.

3. Model Assessment for Random-X Covariance Regression. To assess the performance of Random-X covariance regression models, we adopt the model assessment approach described in Chapter 7 of Hastie et al. (2009) to analyze the training error and test error. Based on the training data $(\mathbb{X}, \mathbb{Y}) = \{(X_i, Y_i) : i \in \{1, \dots, n\}\}$, we can estimate the unknown covariance regression parameter β by $\hat{\beta}$, where $\hat{\beta}$ can be either $\hat{\beta}_{\text{QMLE}}$ or $\hat{\beta}_{\text{WLS}}$ proposed in Section 2. Recall that in Section 2.2, we denoted $C_{\beta}(X_i) = \text{vec} C_{\beta}(X_i)$ and $\mathbf{Y}_i = \text{vec}(Y_i Y_i^{\top})$ for the training data $i \in \{1, \dots, n\}$. For the sake of simplicity, we let $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}}(\cdot)$, while $\hat{\mathbf{C}}(\cdot)$ is sometimes denoted by $\hat{\mathbf{C}}_{\mathbb{X},\mathbb{Y}}(\cdot)$ to avoid confusion. To assess the model fit, we adopt (2.10) and define the training error as follows:

(3.1)
$$\operatorname{Tr}(\mathbb{X},\mathbb{Y}) = \frac{1}{n} \sum_{i=1}^{n} L(\mathbf{Y}_{i}, \hat{\mathbf{C}}(X_{i})),$$

where the loss function $L(\mathbf{Y}_i, \hat{\mathbf{C}}(X_i)) = {\mathbf{Y}_i - \hat{\mathbf{C}}(X_i)}^\top \mathbf{W}(X_i) {\mathbf{Y}_i - \hat{\mathbf{C}}(X_i)}$ measures the error between \mathbf{Y}_i and $\hat{\mathbf{C}}(X_i)$. Unfortunately, the training error $\operatorname{Tr}(\mathbb{X}, \mathbb{Y})$ cannot be used to assess the performance of covariance regression models; this is because the training error does not properly account for model complexity (Hastie et al., 2009, p. 38). To this end, we propose the test error for covariance regression models, which is the average loss over the test data set $(\mathbb{X}_0, \mathbb{Y}_0) = \{(X_{0i}, Y_{0i}) : i \in \{1, \dots, m\}\}$, i.e.,

(3.2)
$$\operatorname{Te}(\mathbb{X}_0, \mathbb{Y}_0, \mathbb{X}, \mathbb{Y}) = \frac{1}{m} \sum_{i=1}^m L(\mathbf{Y}_{0i}, \hat{\mathbf{C}}(X_{0i})),$$

where $\mathbf{Y}_{0i} = \operatorname{vec}(Y_{0i}Y_{0i}^{\top})$ and $Y_{0i} \in \mathbb{Y}_0 = \{Y_{0i} : i \in \{1, \dots, m\}\}$. It is worth noting that the aim of this section only focuses on model assessment rather than utilizing $\hat{\mathbf{C}}(X_{0i})$ to predict \mathbf{Y}_{0i} . Hence, our focus is to assess the covariance regression models by analyzing the test error (3.2) and its expected value given below. In addition, the test error $\operatorname{Te}(\cdot)$ depends on both training data (\mathbb{X}, \mathbb{Y}) and test data $(\mathbb{X}_0, \mathbb{Y}_0)$, where $(\mathbb{X}_0, \mathbb{Y}_0)$ is not used to train the model $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}}(\cdot)$, and we employ the same $\hat{\mathbf{C}}(\cdot)$ for both training and test errors. This is similar to the definitions of training and test errors in classical mean regression models.

In practice, test observations (X_0, Y_0) may not be available. In this case, we can only employ the training data (X, Y) to estimate the expected test error

(3.3)
$$\operatorname{Err} = \operatorname{E} \left\{ \operatorname{Te}(\mathbb{X}_0, \mathbb{Y}_0, \mathbb{X}, \mathbb{Y}) \right\},$$

and then use the estimator of Err to assess covariance regression models. This is similar to treating Mallows' Cp (Mallows, 1973) and other model assessment criteria as the estimators of the expected test error in classical mean regression models (see, e.g., Hastie et al., 2009, p. 230). In these models, Rosset and Tibshirani (2020) found that the Random-X and Fixed-X settings can result in different expected test errors. This motivates us to analytically evaluate such differences for covariance regression models. To this end, we first study the expected test errors under both Random-X and Fixed-X settings. Then, we make comparisons between these two errors. We next adapt the definitions of Rosset and Tibshirani (2020) and consider the following three settings for both training data X and test data X_0 .

Fixed-X Setting. Assume that the explanatory variables in the training data set are fixed, i.e., $X_1 = x_1, \dots, X_n = x_n$. In addition, assume that the explanatory variables in the test data are the same as those in the training data, i.e., $X_{01} = x_1, \dots, X_{0n} = x_n$.

Random-X Setting. Assume that $(X_1, Y_1), \dots, (X_n, Y_n), (X_{01}, Y_{01}), \dots, (X_{0m}, Y_{0m})$ are i.i.d. according to some joint distribution of (X, Y).

Same-X Setting. Assume that $(X_1, Y_1), \dots, (X_n, Y_n)$ are i.i.d. according to some joint distribution of (X, Y). In addition, assume that $X_{0i} = X_i$ for $i = 1, \dots, m$ with m = n, and the corresponding responses Y_{0i} are independent distributed according to the conditional distribution of $Y_i|X_i$ for $i = 1, \dots, n$.

It is clear that both Random-X and Same-X settings have the random training data $\mathbb{X} = \{X_i : i \in \{1, \dots, n\}\}$ and the random test data $\mathbb{X}_0 = \{X_{0i} : i \in \{1, \dots, m\}\}$. Without causing confusion, we also use $\{(X_\alpha, Y_\alpha) : \alpha \in \{1, \dots, n\}\}$ and $\{(X_\alpha, Y_\alpha) : \alpha \in \{01, \dots, 0m\}\}$ to denote the training data and the test data, respectively.

We next evaluate the expected test error (3.3) under the aforementioned three settings. To make fair comparisons, we let m = n in the Random-X setting. However, the theoretical results that we have developed for expected test errors under the Random-X setting are valid for the test data with general size m. In this section, we also assume that the true covariance regression model is $Cov[Y_{\alpha}|\mathcal{F}_{\mathcal{X}}] = C(X_{\alpha})$ for both training and test data (i.e., $\alpha \in \{1, \dots, n\} \cup \{01, \dots, 0n\}$). Note that $C(\cdot)$ does not necessarily have the same parametric structure as $C_{\beta}(\cdot)$, and this allows us to study the impact of model misspecification on the resulting expected test errors. With a slight abuse of notation, we incorporate the training and test data together and denote $\mathcal{F}_{\mathcal{X}} = \sigma \langle X_1, \dots, X_n, X_{01}, \dots, X_{0n} \rangle$. In what follows, we introduce the main theoretical results with regard to the expected test errors. Additional theoretical results and remarks are provided in Section S.4 of the supplementary material.

3.1. *Bias-variance decomposition and optimism.* To study the expected test error Err in (3.3), we introduce a technical assumption below. This assumption is not only applied for both the training and test data, but also allows us to compare the bias-variance decompositions of Err across the Fixed-X, Random-X and Same-X settings.

ASSUMPTION 6. Assume that the data are generated by $Y_{\alpha} = C^{1/2}(X_{\alpha})\mathcal{E}_{\alpha}$ for all $\alpha = 1, \dots, n$ and $01, \dots, 0n$, $\mathcal{E}_{\alpha} = (\varepsilon_{\alpha 1}, \dots, \varepsilon_{\alpha p})^{\top}$ is independent of \mathcal{F}_X , where $\varepsilon_{\alpha j}s$ are i.i.d. in both α and j with $E(\varepsilon_{\alpha j}) = 0$, $E(\varepsilon_{\alpha j}^2) = 1$, $E(\varepsilon_{\alpha j}^4) = \mu^{(4)}$, $E|\varepsilon_{\alpha j}|^{4+\eta_1} < \infty$ for $j = 1, \dots, p$, and $\eta_1 > 0$ is defined in Assumption 1.

A condition similar to Assumption 6 is also given in Rosset and Tibshirani (2020) for establishing the model assessment theory of classical mean regression models. Assumption 6 is required for studying the bias-variance decomposition under the Fixed-X, Same-X, and Random-X settings. To this end, we first obtain the closed form of the conditional covariance $\mathbf{V}(X_{\alpha}) = \operatorname{Cov}[\mathbf{Y}_{\alpha}|X_{\alpha}]$ for $\alpha \in \{1, \dots, n\} \cup \{01, \dots, 0n\}$; see Proposition S.1 of Section S.4 in the supplementary material. In order to tackle the technical challenges and complexities of establishing the model assessment theory due to the complicated dependence structure $\operatorname{Cov}[\mathbf{Y}_{\alpha}|X_{\alpha}]$, the training error (3.1) and test error (3.2) defined earlier have incorporated the general weight matrix $\mathbf{W}(X_{\alpha})$. This is a new idea compared to the training error and test error considered by Rosset and Tibshirani (2020), who used the identity weight matrix. Then, let $\nu^2(X_{\alpha}) = \operatorname{tr}\{\mathbf{W}(X_{\alpha})\mathbf{V}(X_{\alpha})\}$, which is non-negative. In the bias-variance decomposition, we let \mathcal{B} and \mathcal{V} correspondingly represent the bias and variance measures of the estimator $\hat{\mathbf{C}}(X_{\alpha})$ of $\tilde{\mathbf{C}}(X_{\alpha})$, where $\tilde{\mathbf{C}}(X_{\alpha}) = \mathbf{W}^{1/2}(X_{\alpha})\mathbf{C}(X_{\alpha})$ with $\mathbf{C}(X_{\alpha}) = \operatorname{vec} C(X_{\alpha})$, and $\hat{\mathbf{C}}(X_{\alpha}) = \mathbf{W}^{1/2}(X_{\alpha})\hat{\mathbf{C}}(X_{\alpha})$. We further denote $\tilde{\mathbf{Y}}_{\alpha} = \mathbf{W}^{1/2}(X_{\alpha})\mathbf{Y}_{\alpha}$. Finally, we evaluate

the expected test errors Err in (3.3) under the Fixed-X, Random-X and Same-X settings, and obtain ErrF, ErrR and ErrS, respectively; see Proposition S.2 in Section S.4 of the supplementary material. Based on the results in Propositions S.1 and S.2, we next show the bias-variance decompositions for ErrF, ErrR and ErrS, respectively.

THEOREM 5. Suppose Assumption 6 is satisfied. (i) Under the Fixed-X setting, we have $\operatorname{ErrF} = n^{-1} \sum_{i=1}^{n} \nu^2(x_i) + \mathcal{B}_{\operatorname{ErrF}} + \mathcal{V}_{\operatorname{ErrF}}$, where

$$\mathcal{B}_{\mathrm{ErrF}} = \frac{1}{n} \sum_{i=1}^{n} \left\| \mathrm{E} \, \hat{\widetilde{\mathbf{C}}}(x_i) - \widetilde{\mathbf{C}}(x_i) \right\|_2^2 \text{ and } \mathcal{V}_{\mathrm{ErrF}} = \frac{1}{n} \sum_{i=1}^{n} \mathrm{tr} \left\{ \mathrm{Cov} \left(\hat{\widetilde{\mathbf{C}}}(x_i) \right) \right\};$$

(ii) Under the Random-X setting, we have $\operatorname{ErrR} = \operatorname{E} \left\{ \nu^2(X_1) \right\} + \mathcal{B}_{\operatorname{ErrR}} + \mathcal{V}_{\operatorname{ErrR}}$, where

$$\mathcal{B}_{\mathrm{ErrR}} = \mathrm{E} \left\| \mathrm{E} \left[\hat{\widetilde{\mathbf{C}}}(X_{01}) \middle| \mathbb{X}, X_{01} \right] - \widetilde{\mathbf{C}}(X_{01}) \right\|_{2}^{2} \text{ and } \mathcal{V}_{\mathrm{ErrR}} = \mathrm{E} \operatorname{tr} \left\{ \mathrm{Cov} \left[\hat{\widetilde{\mathbf{C}}}(X_{01}) \middle| \mathbb{X}, X_{01} \right] \right\};$$

(iii) Under the Same-X setting, we have $\operatorname{ErrS} = \operatorname{E} \left\{ \nu^2(X_1) \right\} + \mathcal{B}_{\operatorname{ErrS}} + \mathcal{V}_{\operatorname{ErrS}}$, where

$$\mathcal{B}_{\mathrm{ErrS}} = \mathrm{E} \left\| \mathrm{E} \left[\hat{\widetilde{\mathbf{C}}}(X_1) \middle| \mathbb{X} \right] - \widetilde{\mathbf{C}}(X_1) \middle\|_2^2 \text{ and } \mathcal{V}_{\mathrm{ErrS}} = \mathrm{E} \operatorname{tr} \left\{ \mathrm{Cov} \left[\hat{\widetilde{\mathbf{C}}}(X_1) \middle| \mathbb{X} \right] \right\}.$$

It is worth noting that \mathcal{B}_{ErrF} and \mathcal{V}_{ErrF} in (i) represent the mean squared bias and the average trace of $Cov(\hat{C}(x_i))$, respectively. This interpretation is also applicable to $(\mathcal{B}_{ErrR}, \mathcal{V}_{ErrR})$ in (ii) and $(\mathcal{B}_{ErrS}, \mathcal{V}_{ErrS})$ in (iii). For the sake of simplicity, we refer to \mathcal{B} and \mathcal{V} as "bias" and "variance", respectively.

To assess the difference between the Random-X and Same-X settings, we employ Theorem 5 and obtain that

(3.4)
$$\operatorname{ErrR} - \operatorname{ErrS} = \mathcal{B}^{(d)} + \mathcal{V}^{(d)},$$

where $\mathcal{B}^{(d)} = \mathcal{B}_{\text{ErrR}} - \mathcal{B}_{\text{ErrS}}$ and $\mathcal{V}^{(d)} = \mathcal{V}_{\text{ErrR}} - \mathcal{V}_{\text{ErrS}}$ represent the bias and variance differences, respectively. If $\mathcal{B}^{(d)} + \mathcal{V}^{(d)} \ge 0$, then the Random-X error, ErrR, is no smaller than the Same-X error, ErrS. The difference between ErrR and ErrS in (3.4) allows us to make comparisons between ErrR and ErrF, which is our main interest and is discussed below.

We first adapt the concept of optimism from classical mean regression (see, e.g., Hastie et al., 2009), and then define the optimism in covariance regression as follows:

$$(3.5) Opt = E \{ Te(\mathbb{X}_0, \mathbb{Y}_0, \mathbb{X}, \mathbb{Y}) - Tr(\mathbb{X}, \mathbb{Y}) \} = Err - E \{ Tr(\mathbb{X}, \mathbb{Y}) \},$$

which is the mean of the difference between the test error and training error. We next evaluate Opt under the Fixed-X, Same-X and Random-X settings and obtain OptF, OptS and OptR, respectively.

PROPOSITION 1. Suppose Assumption 6 is satisfied. (i) Under the Fixed-X setting, we have $\operatorname{OptF} = 2n^{-1} \sum_{i=1}^{n} \operatorname{tr} \operatorname{Cov}(\widetilde{\mathbf{Y}}_{i}, \hat{\widetilde{\mathbf{C}}}(x_{i}))$; (ii) under the Same-X setting, we obtain $\operatorname{OptS} = 2n^{-1} \sum_{i=1}^{n} \operatorname{tr} \operatorname{E} \operatorname{Cov}[\widetilde{\mathbf{Y}}_{i}, \hat{\widetilde{\mathbf{C}}}(X_{i}) | \mathbb{X}]$; and (iii) under the Random-X setting, we have $\operatorname{OptR} = \operatorname{OptS} + \mathcal{B}^{(d)} + \mathcal{V}^{(d)}$.

Using Proposition 1, we can further explore the relationships between the estimators of ErrF, ErrS and ErrR. Let $U(\mathbb{X}, \mathbb{Y})$ be an unbiased estimator of OptF under the Fixed-X setting, and a typical example of the unbiased estimator of OptF is presented in Section 3.2. Based on (3.5), ErrF can be naturally estimated by $\widehat{\text{ErrF}} = \text{Tr}(\mathbb{X}, \mathbb{Y}) + U(\mathbb{X}, \mathbb{Y})$, which is also the unbiased estimator of ErrS demonstrated below.

PROPOSITION 2. Suppose Assumption 6 is satisfied. (i) $U(\mathbb{X}, \mathbb{Y})$ is an unbiased estimator of OptS. In addition, the unbiased estimator of the expected test error in (3.3) is the same under the Fixed-X and Same-X settings. Specifically,

(3.6)
$$\widehat{\mathrm{Err}} \widehat{\mathrm{F}} = \widehat{\mathrm{Err}} \widehat{\mathrm{S}} = \mathrm{Tr}(\mathbb{X}, \mathbb{Y}) + U(\mathbb{X}, \mathbb{Y})$$

(ii) If OptF does not depend on x_1, \dots, x_n under the Fixed-X setting, then it is equal to OptS under the Same-X setting.

Let $\widehat{\mathcal{B}}^{(d)}$ and $\widehat{\mathcal{V}}^{(d)}$ be the estimators of $\mathcal{B}^{(d)}$ and $\mathcal{V}^{(d)}$, respectively. An example of the estimators of $\mathcal{B}^{(d)}$ and $\mathcal{V}^{(d)}$ is discussed in Section 3.2. By Propositions 1 – 2 and (3.5), we finally obtain that

$$\widehat{\operatorname{OptR}} = U(\mathbb{X}, \mathbb{Y}) + \widehat{\mathcal{B}}^{(d)} + \widehat{\mathcal{V}}^{(d)}, \text{ and}$$

(3.7) $\widehat{\operatorname{ErrR}} = \operatorname{Tr}(\mathbb{X}, \mathbb{Y}) + \widehat{\operatorname{OptR}} = \operatorname{Tr}(\mathbb{X}, \mathbb{Y}) + U(\mathbb{X}, \mathbb{Y}) + \widehat{\mathcal{B}}^{(d)} + \widehat{\mathcal{V}}^{(d)},$

In sum, equations (3.6) and (3.7) clearly indicate the components that are due to, first, the difference $\mathcal{B}^{(d)}$ between the biases in the estimators of ErrF and ErrR and, second, the difference $\mathcal{V}^{(d)}$ between the variances in the estimators of ErrF and ErrR.

In classical mean regression models, ErrF can refer to Mallows' Cp (Mallows, 1973), which is commonly used as a model assessment criterion. Analogously, ErrR can be used as a model assessment criterion for random covariance regressions, and that will be discussed in the next subsection.

3.2. *Mallows' Cp in linear covariance regression models.* In this section, we mainly focus on model assessment based on fitting linear covariance regression model (2.11) with the estimator $\hat{\beta}_{WLS}$ defined in (2.12). However, the true covariance regression model $Cov[Y_{\alpha}|\mathcal{F}_{\mathcal{X}}] = C(X_{\alpha})$ for both training and test data (i.e., $\alpha \in \{1, \dots, n\} \cup \{01, \dots, 0n\}$) is not necessarily correctly specified by (2.11).

Note that in model (2.11), the $X_i^{(k)}$ s are $p \times p$ matrices for $i = 1, \dots, n$ and $k = 1, \dots, K$. Denote $\mathbf{X}_i = (\operatorname{vec}(X_i^{(1)}), \dots, \operatorname{vec}(X_i^{(K)})) \in \mathbb{R}^{p^2 \times K}$. In addition, let $X_i = (X_i^{(1)}, \dots, X_i^{(K)})$, which was defined in the first example in Section 1.1. Under the Fixed-X setting, consider $X_i = x_i = (x_i^{(1)}, \dots, x_i^{(K)})$ and $\mathbf{x}_i = (\operatorname{vec}(x_i^{(1)}), \dots, \operatorname{vec}(x_i^{(K)})) \in \mathbb{R}^{p^2 \times K}$. Let $x = \{x_1, \dots, x_n\}$ and $\mathbb{W}(x) = \operatorname{diag}\{\mathbf{W}(x_1), \dots, \mathbf{W}(x_n)\}$ be a $np^2 \times np^2$ matrix whose diagonal blocks are $\mathbf{W}(x_1), \dots, \mathbf{W}(x_n)$ and the other elements are zeros. Similarly, let $\mathbb{V}(x) = \operatorname{diag}\{\mathbf{V}(x_1), \dots, \mathbf{V}(x_n)\} \in \mathbb{R}^{np^2 \times np^2}$, and the detailed expression of $\mathbf{V}(\cdot) \in \mathbb{R}^{p^2 \times p^2}$ is presented in Proposition S.1 of Section S.4 of the supplementary material. Finally, let $\widetilde{\mathbf{X}}_i = \mathbf{W}^{1/2}(X_i)\mathbf{X}_i \in \mathbb{R}^{p^2 \times K}$, $\widetilde{\mathbf{x}}_i = \mathbf{W}^{1/2}(x_i)\mathbf{x}_i \in \mathbb{R}^{p^2 \times K}$, $\widetilde{\mathbf{X}} = (\widetilde{\mathbf{X}}_1^\top, \dots, \widetilde{\mathbf{X}}_n^\top)^\top \in \mathbb{R}^{np^2 \times K}$, and $\widetilde{\mathbb{V}}(x) = \mathbb{W}^{1/2}(x)\mathbb{V}(x)\mathbb{W}^{1/2}(x) \in \mathbb{R}^{np^2 \times np^2}$. Analogously, for $\alpha \in \{01, \dots, 0n\}$, we denote $\mathbf{X}_\alpha = (\operatorname{vec}(X_\alpha^{(1)}), \dots, \operatorname{vec}(X_\alpha^{(K)})) \in \mathbb{R}^{p^2 \times K}$, $X_\alpha = (X_\alpha^{(1)}, \dots, X_\alpha^{(K)})$, and $\widetilde{\mathbf{X}}_\alpha = \mathbf{W}^{1/2}(X_\alpha)\mathbf{X}_\alpha \in \mathbb{R}^{p^2 \times K}$. The above notation will be used in the following proposition and theorem.

PROPOSITION 3. Suppose Assumption 6 is satisfied. In addition, consider fitting a linear covariance regression model (2.11) with $\hat{\beta} = \hat{\beta}_{WLS}$ defined in (2.12). Under the Fixed-X setting, we have that $OptF = 2n^{-1}tr\{(\widetilde{\mathbf{x}}^{\top}\widetilde{\mathbf{x}})^{-1}\widetilde{\mathbf{x}}^{\top}\widetilde{\mathbb{V}}(x)\widetilde{\mathbf{x}}\}$. In addition, under the Same-X setting, we have that $OptS = 2n^{-1}Etr\{(\widetilde{\mathbf{x}}^{\top}\widetilde{\mathbf{x}})^{-1}\widetilde{\mathbf{x}}^{\top}\widetilde{\mathbb{V}}(\mathbb{X})\widetilde{\mathbf{x}}\}$.

16

As discussed in Remark 1, set $W(X_{\alpha}) = V^{-}(X_{\alpha})$. Then Proposition 3 leads to

(3.8)
$$\operatorname{OptF} = \operatorname{OptS} = \frac{2K}{n}$$

which is a special case of the general result, OptF = OptS, as demonstrated in Proposition 2. Note that 2K/n is a constant and does not need to be estimated. In addition, consider the case that $V(\cdot)$ is known. This, together with Proposition 3, yields an unbiased estimator for both OptF and OptS as given below.

(3.9)
$$U(\mathbb{X},\mathbb{Y}) = \frac{2}{n} \operatorname{tr}\left\{ (\widetilde{\mathbf{X}}^{\top} \widetilde{\mathbf{X}})^{-1} \widetilde{\mathbf{X}}^{\top} \widetilde{\mathbb{V}}(\mathbb{X}) \widetilde{\mathbf{X}} \right\} \triangleq U(\mathbb{X}).$$

Note that this estimator does not depend on \mathbb{Y} . Using the above result and Proposition 2, we define Mallows' Cp for linear covariance regression models under the Fixed-X and Same-X settings as follows:

(3.10)
$$\operatorname{Cp} \triangleq \widehat{\operatorname{ErrF}} = \widehat{\operatorname{ErrS}} = \operatorname{Tr}(\mathbb{X}, \mathbb{Y}) + U(\mathbb{X}),$$

which is an unbiased estimator of ErrF or ErrS based on Proposition 2.

REMARK 5. It is worth noting that the estimator $U(\mathbb{X})$ in (3.9) depends on $\mathbf{V}(\cdot)$. In practice, $\mathbf{V}(\cdot)$ may be unknown and needs to be estimated. If $C(\cdot)$ is correctly specified by the linear covariance regression model (2.11), Section S.5.1 of the supplementary material provides an estimator $\hat{\mathbf{V}}(\cdot) \triangleq \hat{\mathbf{V}}_{\mathbb{X},\mathbb{Y}}(\cdot)$ of $\mathbf{V}(\cdot)$ based on the training data (\mathbb{X},\mathbb{Y}) . If we replace $\mathbf{V}(\cdot)$ by $\hat{\mathbf{V}}_{\mathbb{X},\mathbb{Y}}(\cdot)$ in the expression of $U(\mathbb{X})$, we then obtain the estimator $\hat{U}_{\mathbb{X},\mathbb{Y}}(\mathbb{X})$. This replacement yields Cp for the unknown $\mathbf{V}(\cdot)$, which is

(3.11)
$$\widehat{\operatorname{Cp}} = \operatorname{Tr}(\mathbb{X}, \mathbb{Y}) + \widehat{U}_{\mathbb{X}, \mathbb{Y}}(\mathbb{X}).$$

To evaluate the estimator of ErrR and compare it with Cp and \widehat{Cp} , we next employ Proposition 3 and Theorem 5 to demonstrate that the difference between the biases, which is $\mathcal{B}^{(d)}$, and the difference between the variances, which is $\mathcal{V}^{(d)}$, are both nonnegative.

THEOREM 6. Suppose Assumption 6 is satisfied. Consider fitting a linear covariance regression model (2.11) with $\hat{\beta} = \hat{\beta}_{WLS}$ defined in (2.12). We then have $\mathcal{V}_{ErrS} = n^{-1} \operatorname{Etr} \{ (\widetilde{\mathbf{X}}^{\top} \widetilde{\mathbf{X}})^{-1} \widetilde{\mathbf{X}}^{\top} \widetilde{\mathbb{V}}(\mathbb{X}) \widetilde{\mathbf{X}} \} = \operatorname{E} U(\mathbb{X})/2 = \operatorname{OptS}/2,$

$$\mathcal{V}_{\mathrm{ErrR}} = \frac{1}{n} \mathrm{tr} \left[\mathrm{E} \left\{ \left(\widetilde{\mathbf{X}}^{\top} \widetilde{\mathbf{X}} \right)^{-1} \widetilde{\mathbf{X}}^{\top} \widetilde{\mathbb{V}}(\mathbb{X}) \widetilde{\mathbf{X}} \left(\widetilde{\mathbf{X}}^{\top} \widetilde{\mathbf{X}} \right)^{-1} \right\} \mathrm{E} \left(\widetilde{\mathbf{X}}^{\top} \widetilde{\mathbf{X}} \right) \right]$$

$$(3.12) \qquad = \mathrm{E} \, \mathrm{tr} \left\{ \left(\widetilde{\mathbf{X}}^{\top} \widetilde{\mathbf{X}} \right)^{-1} \widetilde{\mathbf{X}}^{\top} \widetilde{\mathbb{V}}(\mathbb{X}) \widetilde{\mathbf{X}} \left(\widetilde{\mathbf{X}}^{\top} \widetilde{\mathbf{X}} \right)^{-1} \widetilde{\mathbf{X}}_{01}^{\top} \widetilde{\mathbf{X}}_{01} \right\}, and$$

 $\mathcal{B}^{(d)} = \mathcal{B}_{\text{ErrR}} - \mathcal{B}_{\text{ErrS}} \ge 0$. Furthermore, set $\mathbf{W}(X_{\alpha}) = \mathbf{V}^{-}(X_{\alpha})$. We obtain $\mathcal{V}^{(d)} = \mathcal{V}_{\text{ErrR}} - \mathcal{V}_{\text{ErrS}} \ge 0$, and hence $\text{ErrR} \ge \text{ErrS}$.

REMARK 6. Suppose that $C(\cdot)$ is correctly specified by the linear covariance regression model (2.11). Then we can verify that $\mathcal{B}_{ErrR} = \mathcal{B}_{ErrS} = 0$, which implies that the bias difference is $\mathcal{B}^{(d)} = 0$. In addition, consider the ideal case that $\hat{\beta} = \beta_0$. We then have $\tilde{\mathbf{C}}(X_{\alpha}) = \tilde{\mathbf{C}}(X_{\alpha})$. This, in conjunction with Proposition S.2 in Section S.4 of the supplementary material, leads to ErrR = ErrS. Moreover, Theorem 2 shows that $\hat{\beta}_{WLS}$ is the consistent estimator of β_0 as $\max\{n, p\} \to \infty$. Even though $\hat{\beta}_{WLS} \neq \beta_0$, $\hat{\beta}_{WLS}$ converges to β_0 with probability tending to 1. Using this result and Proposition S.2, one can expect that, as $\max\{n, p\} \to \infty$, the difference between ErrR and ErrS tends to be small under the correctly specified model.

Note that, under $\mathbf{W}(X_{\alpha}) = \mathbf{V}^{-}(X_{\alpha})$, (3.5) and (3.8) imply that $\operatorname{ErrS} = \mathrm{E}\{\operatorname{Tr}(\mathbb{X}, \mathbb{Y})\} + 2K/n$ and $\operatorname{ErrF} = \mathrm{E}\{\operatorname{Tr}(x, \mathbb{Y})\} + 2K/n$. Hence, ErrS and ErrF are different. In practice, however, ErrS and ErrF have the same estimator Cp in (3.10), and it is different from the ErrR estimator, $\widehat{\operatorname{ErrR}}$, in (3.7). From Theorem 6, one can expect that the estimators of the difference in biases and the difference in variances satisfy $\widehat{\mathcal{B}}^{(d)} \ge 0$ and $\widehat{\mathcal{V}}^{(d)} \ge 0$, respectively, which leads to $\widehat{\operatorname{ErrR}} \ge \operatorname{Cp}$. We next propose two estimation approaches to obtain $\widehat{\operatorname{ErrR}}$ explicitly.

3.2.1. Estimation of ErrR with $C(\cdot)$ correctly specified in model (2.11). Assume that $C(\cdot)$ is correctly specified by the linear covariance regression model (2.11). By Remark 6, we have $\mathcal{B}^{(d)} = \mathcal{B}_{\text{ErrR}} = \mathcal{B}_{\text{ErrS}} = 0$. Naturally, their corresponding estimators are $\widehat{\mathcal{B}}^{(d)} = \widehat{\mathcal{B}}_{\text{ErrR}} = \widehat{\mathcal{B}}_{\text{ErrS}} = 0$. Hence, based on (3.4) and (3.10), it suffices to estimate $\mathcal{V}^{(d)} = \mathcal{V}_{\text{ErrR}} - \mathcal{V}_{\text{ErrS}}$ and then obtain the estimator of ErrR. We next discuss the estimation of $\mathcal{V}_{\text{ErrS}}$ and $\mathcal{V}_{\text{ErrR}}$, separately.

Under the assumption that $\mathbf{V}(\cdot)$ is known, $U(\mathbb{X})$ in (3.9) is an unbiased estimator of OptS. In addition by Theorem 6, we obtain that $\widetilde{\mathcal{V}}_{ErrS} = U(\mathbb{X})/2$ is an unbiased estimator of \mathcal{V}_{ErrS} . As for the estimation of \mathcal{V}_{ErrR} in (3.12) of Theorem 6, we only have the training data \mathbb{X} and $\widetilde{\mathbf{X}}$, and the test data $\widetilde{\mathbf{X}}_{01}$ is not available. Hence, we consider using

(3.13) Etr
$$\left\{ \left(\widetilde{\mathbf{X}}^{(-i)\top} \widetilde{\mathbf{X}}^{(-i)} \right)^{-1} \widetilde{\mathbf{X}}^{(-i)\top} \widetilde{\mathbb{V}}(\mathbb{X}^{(-i)}) \widetilde{\mathbf{X}}^{(-i)} \left(\widetilde{\mathbf{X}}^{(-i)\top} \widetilde{\mathbf{X}}^{(-i)} \right)^{-1} \widetilde{\mathbf{X}}_{i}^{\top} \widetilde{\mathbf{X}}_{i} \right\}$$

to approximate (3.12), where $\widetilde{\mathbf{X}}^{(-i)} = (\widetilde{\mathbf{X}}_{1}^{\top}, \dots, \widetilde{\mathbf{X}}_{i-1}^{\top}, \widetilde{\mathbf{X}}_{i+1}^{\top}, \dots, \widetilde{\mathbf{X}}_{n}^{\top})^{\top}$ and $\mathbb{X}^{(-i)} = \{X_{1}, \dots, X_{i-1}, X_{i+1}, \dots, X_{n}\}$ for $i = 1, \dots, n$. Compared to the training data \mathbb{X} of size n in (3.12), $\mathbb{X}^{(-i)}$ in (3.13) is of size n-1. As a consequence, one expects that the approximation (3.13) is close to $\mathcal{V}_{\text{ErrR}}$ as n gets large. Thus, it is natural to consider a cross-validation estimator of $\mathcal{V}_{\text{ErrR}}$,

$$\widetilde{\mathcal{V}}_{\mathrm{ErrR}} = \frac{1}{n} \sum_{i=1}^{n} \mathrm{tr} \left\{ \left(\widetilde{\mathbf{X}}^{(-i)\top} \widetilde{\mathbf{X}}^{(-i)} \right)^{-1} \widetilde{\mathbf{X}}^{(-i)\top} \widetilde{\mathbb{V}}(\mathbb{X}^{(-i)}) \widetilde{\mathbf{X}}^{(-i)} \left(\widetilde{\mathbf{X}}^{(-i)\top} \widetilde{\mathbf{X}}^{(-i)} \right)^{-1} \widetilde{\mathbf{X}}_{i}^{\top} \widetilde{\mathbf{X}}_{i} \right\},\$$

which is an unbiased estimator of (3.13). Based on the estimators $\widetilde{\mathcal{V}}_{\text{ErrR}}$ and $\widetilde{\mathcal{V}}_{\text{ErrS}}$, we then obtain an estimator of $\mathcal{V}^{(d)}$, $\widetilde{\mathcal{V}}^{(d)} = \widetilde{\mathcal{V}}_{\text{ErrR}} - \widetilde{\mathcal{V}}_{\text{ErrS}}$.

The equation $\widetilde{\mathcal{V}}^{(d)} = \widetilde{\mathcal{V}}_{\text{ErrR}} - \widetilde{\mathcal{V}}_{\text{ErrS}}$, along with (3.4), (3.10) and the fact that $\widetilde{\mathcal{V}}_{\text{ErrS}} = U(\mathbb{X})/2$, yields an estimator of ErrR, which is

(3.15)
$$\operatorname{RCp} \triangleq \operatorname{Cp} + \widetilde{\mathcal{V}}^{(d)} = \operatorname{Tr}(\mathbb{X}, \mathbb{Y}) + \frac{1}{2}U(\mathbb{X}) + \widetilde{\mathcal{V}}_{\operatorname{ErrR}}.$$

This estimator is essentially Mallows' Cp adjusted by $\widetilde{\mathcal{V}}^{(d)}$, the estimator of the difference in variances, which is induced by the Random-X setting. Thus, we name it Random Cp (RCp) for covariance regression. Note that $\mathbf{V}(\cdot)$ may be unknown in practice. Analogously to Remark 5, we can replace $\mathbf{V}(\cdot)$ in (3.14) by its estimator $\hat{\mathbf{V}}(\cdot)$ defined in Section S.5.1 of the supplementary material. Then, we obtain the estimator of $\mathcal{V}_{\text{ErrR}}$ for the unknown $\mathbf{V}(\cdot)$, and denote it by $\hat{\mathcal{V}}_{\text{ErrR}}$. This, together with the estimator of $U(\mathbb{X})$ in Remark 5, namely $\hat{U}_{\mathbb{X},\mathbb{Y}}(\mathbb{X})$, yields the estimator of ErrR:

(3.16)
$$\widehat{\mathrm{RCp}} = \mathrm{Tr}(\mathbb{X}, \mathbb{Y}) + \frac{1}{2}\widehat{U}_{\mathbb{X}, \mathbb{Y}}(\mathbb{X}) + \widehat{\mathcal{V}}_{\mathrm{ErrR}}.$$

Based on Theorem 6, we expect that \widehat{RCp} is larger than \widehat{Cp} . It is worth noting that either \widehat{RCp} or \widehat{Cp} can be used to compare a set of covariance regression models in terms of their

expected test errors. Under the Random-X setting, however, $\widehat{\text{RCp}}$ should be a better choice since $\widehat{\text{Cp}}$ can underestimate expected test errors. Note that RCp and $\widehat{\text{RCp}}$ use the results from Proposition 3 and Theorem 6 obtained by fitting a linear covariance regression model (2.11). Hence, they are only applicable for the linear model. The following subsection discusses the second approach to estimating ErrR.

3.2.2. Estimation of ErrR with $C(\cdot)$ misspecified in model (2.11). Recall that before equation (3.1), we denoted $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}}(\cdot)$. If $C(\cdot)$ is misspecified by the linear covariance regression model (2.11), $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}}(\cdot)$ is a biased estimator of $\mathbf{C}(\cdot)$. Hence, $\mathcal{B}_{\text{ErrR}} \neq 0$ and $\mathcal{B}_{\text{ErrS}} \neq 0$, and $\widehat{\text{RCp}}$ is not an appropriate estimator of ErrR. Since the true covariance function $C(\cdot)$ is unknown, we cannot estimate $\mathcal{B}_{\text{ErrR}}$ and $\mathcal{B}_{\text{ErrS}}$ directly based on Theorem 5. Instead, we use the result $\text{ErrR} = \mathbb{E} \| \widetilde{\mathbf{Y}}_{01} - \widetilde{\mathbf{C}}(X_{01}) \|_2^2$ in Proposition S.2 of Section S.4 of the supplementary material, adopt a similar approach to obtaining the estimator (3.14), and then find a cross-validation estimator. Specifically, let $\hat{\mathbf{C}}^{(-i)}(\cdot)$ be the WLS estimator obtained from all training data (\mathbb{X}, \mathbb{Y}) except for (X_i, Y_i) . Then, calculate the ordinary crossvalidation (OCV) error,

(3.17)
$$OCV = \frac{1}{n} \sum_{i=1}^{n} \left\| \widetilde{\mathbf{Y}}_{i} - \widehat{\widetilde{\mathbf{C}}}^{(-i)}(X_{i}) \right\|_{2}^{2},$$

which is an approximately unbiased estimator of ErrR.

The advantage of using OCV is that it can estimate ErrR for a general and possibly nonlinear $C_{\beta}(\cdot)$. In addition, it does not require the true covariance function $C(\cdot)$ to be correctly specified. If $C(\cdot)$ is correctly specified by the linear covariance regression model (2.11), then either OCV or $\widehat{\text{RCp}}$ can be used to estimate ErrR. Under the random covariance regression models, we compare the performance of the statistics Cp, $\widehat{\text{Cp}}$, RCp, $\widehat{\text{RCp}}$ and OCV via simulation studies in Section 4 below.

4. Simulation Studies. In this section, we conduct simulation studies to investigate the finite sample performance of parameter estimations and expected test errors in covariance regression models. All the tables and figures in this section, except Tables 1 - 2, are provided in Section S.20 of the supplementary material to save space.

In these studies, for both training data $\alpha \in \{1, \dots, n\}$ and test data $\alpha \in \{01, \dots, 0n\}$, the response vectors Y_{α} are simulated by $Y_{\alpha} = C_{\beta_0}^{1/2}(X_{\alpha})\mathcal{E}_{\alpha}$, where the components of the random error \mathcal{E}_{α} are independent and identically generated from the standard normal distribution and the mixture of normal distributions 0.9N(0, 5/9) + 0.1N(0, 5), respectively. In addition, the following two different models are considered for $C_{\beta_0}(X_{\alpha})$ with $X_{\alpha} = (X_{\alpha}^{(2)}, \dots, X_{\alpha}^{(K)})$ and K = 5.

Model A. Let $C_{\beta_0}(X_{\alpha}) = \beta_0^{(1)} I_p + \sum_{k=2}^5 \beta_0^{(k)} X_{\alpha}^{(k)}$, where the $X_{\alpha}^{(k)}$'s are $p \times p$ matrices, the diagonal elements of $X_{\alpha}^{(k)}$ are zeros for $k = 2, \dots, 5$, the off-diagonal elements of $X_{\alpha}^{(2)}$ are independent and identically generated from the Bernoulli distribution with probability p^{-1} , and the off-diagonal elements of $X_{\alpha}^{(k)} = (X_{\alpha,j_1j_2}^{(k)})_{p \times p}$ for k = 3, 4, 5 are $X_{\alpha,j_1j_2}^{(k)} = \exp\{-(d_{\alpha,j_1j_2}^{(k)})^2\}$ with $d_{\alpha,j_1j_2}^{(k)}$ independent and identically generated from the uniform distribution $U(0, p^{1+(k-3)/6})$.

Model B. Let $C_{\beta_0}(X_{\alpha}) = \beta_0^{(1)} I_p + \sum_{k=2}^5 \beta_0^{(k)} X_{\alpha}^{(k)} X_{\alpha}^{(k)\top}$, where the $X_{\alpha}^{(k)}$ s are $p \times 1$ vectors, and the components of $X_{\alpha}^{(k)}$ are independent and identically generated from the normal distribution $N(0, p^{-1})$, for $k = 2, \dots, 5$.

Note that Models A and B, respectively, correspond to the first example and a variant of the third example of covariance regression in Section 1.1. In both models, we set $\beta_0^{(1)} = 5$ and $\beta_0^{(k)} = 1.4$ for $k = 2, \dots, 5$. In addition, we consider three sample sizes n = 25, 50, and 100, and two dimensions of the response vector p = 5 and 25. For each of the above models, a total of 1,000 simulation experiments are conducted. For the Fixed-X setting, we set X_{α} with $\alpha = 1, \dots, n$ and $01, \dots, 0n$ such that $X_{0i} = X_i$, for $i = 1, \dots, n$, and we only use one realization of $\{X_1 = x_1, \dots, X_n = x_n\}$ in all 1,000 simulation experiments. In contrast, for the Random-X setting, the X_{α} s are independent and identically generated for $\alpha = 1, \dots, n$ and $01, \dots, 0n$ across 1,000 realizations. Accordingly, these processes for generating X_{α} allow us to make a fair comparison between the Fixed-X and Random-X settings.

We first assess the performance of the constrained QMLE ($\hat{\beta}_{QMLE}^c$) and constrained WLS estimators based on the training data $i = 1, \dots, n$, where the constrained estimators are shown to be asymptotically equivalent to the unconstrained estimators in Remark 3. Among the constrained WLS estimators, we consider the constrained OLS estimator $\hat{\beta}_{OLS}^c$ by letting $\mathbf{W}(X_i) = I_{p^2}$, the constrained GLS estimator $\hat{\beta}_{GLS}^c$ by letting $\mathbf{W}(X_i) = \mathbf{V}^-(X_i)$, and the constrained FGLS estimator $\hat{\beta}_{FGLS}^c$ by letting $\mathbf{W}(X_i) = \hat{\mathbf{V}}^-(X_i)$, where the last two estimators can be obtained via Remark 1 and Remark 3. We also compare the constrained OLS estimate to the unconstrained OLS estimate. The numerical performance of the unconstrained estimate and some related discussion are given in Section S.17 of the supplementary material.

Under Model A with random errors generated from the mixture of normal distributions, Tables 1 – 2 present the average bias (BIAS), standard deviation (SD), and root mean squared error (RMSE) of the four constrained estimators via 1,000 realizations. In addition, Table S.1 reports two types averaged estimation errors in the covariance matrix estimate $C_{\hat{\beta}}(X_i)$. They are the S-Error $(\max_{1 \le i \le n} ||C_{\hat{\beta}}(X_i) - C_{\beta_0}(X_i)||_2$ with the spectral norm $|| \cdot ||_2$) and the Ferror $(p^{-1/2} \max_{1 \le i \le n} ||C_{\hat{\beta}}(X_i) - C_{\beta_0}(X_i)||_F$ with the Frobenius norm $|| \cdot ||_F$), where $\hat{\beta}$ can be any one of the four estimates, $\hat{\beta}_{\text{QMLE}}^c$, $\hat{\beta}_{\text{GLS}}^c$, and $\hat{\beta}_{\text{FGLS}}^c$. Moreover, Tables S.2 – S.4 present the results when random errors are generated from the standard normal distribution.

Tables 1, 2 and S.1 and Tables S.2 - S.4 reveal five important findings. (I) The BIAS, SD, RMSE, S-Error and F-Error generally decrease as n or p gets larger, which confirms the consistency of the parameter estimators demonstrated in Theorems 1-2, and the consistency of the covariance matrix estimators shown in Theorems 3 - 4. It is worth noting that the biases of the constrained QMLE, OLS and FGLS estimates generally decrease as n or p gets larger, but they cannot be zero. According to Remark 3, this finding is reasonable since the constrained estimators are asymptotically equivalent to the unconstrained estimators. Hence, the constrained estimators are not unbiased. Rather, they are asymptotically unbiased. In this example, however, the BIAS of the constrained GLS estimate is very close to zero across all settings of n and p. (II) The SD and RMSE of the GLS and FGLS estimates get closer to those of QMLE as either n or p increases, while these three estimates are generally superior to the OLS estimates. (III) The SD and RMSE of the GLS estimates are generally closer and comparable to those of QMLE under the normal random errors, while they are slightly inferior to those of QMLE under the mixture of normal random errors. (IV) The BIAS, SD, RMSE, S-Error and F-Error of the FGLS estimate gradually approach those of the GLS estimate as both n and p get larger, which is consistent with the comment below Theorem 2. It is of interest to note that the GLS estimates sometimes have higher RMSE than that of the OLS or FGLS estimates, especially when n and p are small. One possible explanation is due to the Monte Carlo approximation errors. Another possibility is that some errors may be due to small n and p. This is because the GLS is only asymptotically as efficient as the FGLS, and it is more efficient than the OLS as n or p goes to infinity. (V) There is not much difference between the Fixed-X and Random-X settings in the RMSE, S-Error and F-Error TABLE 1

The average bias (BIAS), standard deviation (SD), and root mean squared error (RMSE) of the constrained QMLE and OLS estimates obtained from Model A with the mixture of normal random errors under the Fixed-X and Random-X settings.

		p = 5					p = 25					
n		$\hat{\beta}_1$	$\hat{\beta}_2$	\hat{eta}_3	$\hat{\beta}_4$	$\hat{\beta}_5$	$\hat{\beta}_1$	$\hat{\beta}_2$	$\hat{\beta}_3$	$\hat{\beta}_4$	$\hat{\beta}_5$	
QMLE & Fixed-X Setting												
25	BIAS	0.34	0.01	-0.23	-0.19	-0.15	0.28	0.02	0.04	-0.04	-0.06	
	SD	0.96	0.72	0.81	0.84	0.90	0.41	0.20	0.28	0.32	0.47	
	RMSE	1.02	0.72	0.84	0.86	0.92	0.50	0.20	0.28	0.33	0.47	
50	BIAS	0.36	-0.05	0.06	-0.04	-0.18	0.23	0.00	0.01	0.05	0.05	
	SD	0.67	0.39	0.56	0.57	0.57	0.30	0.16	0.20	0.26	0.32	
	RMSE	0.76	0.39	0.56	0.57	0.60	0.38	0.16	0.20	0.27	0.32	
100	BIAS	0.32	0.08	0.08	-0.10	-0.05	0.23	0.04	0.04	-0.00	-0.00	
	SD	0.47	0.32	0.37	0.37	0.41	0.19	0.11	0.13	0.15	0.19	
	RMSE	0.57	0.33	0.38	0.38	0.41	0.30	0.12	0.13	0.15	0.19	
QMLE & Random-X Setting												
25	BIAS	0.11	-0.12	-0.11	-0.13	-0.11	0.15	-0.02	-0.00	-0.01	-0.01	
	SD	1.02	0.58	0.70	0.80	0.91	0.47	0.21	0.26	0.34	0.42	
	RMSE	1.03	0.60	0.71	0.81	0.92	0.49	0.22	0.26	0.34	0.42	
50	BIAS	0.17	-0.05	-0.03	-0.06	-0.07	0.18	0.01	0.01	0.03	0.04	
	SD	0.72	0.39	0.53	0.57	0.60	0.33	0.16	0.19	0.26	0.31	
	RMSE	0.74	0.39	0.53	0.57	0.60	0.38	0.16	0.19	0.26	0.31	
100	BIAS	0.17	-0.02	0.01	-0.02	-0.03	0.20	0.02	0.02	0.02	0.03	
	SD	0.48	0.28	0.36	0.38	0.40	0.20	0.11	0.13	0.17	0.23	
	RMSE	0.51	0.28	0.36	0.38	0.40	0.28	0.11	0.13	0.17	0.23	
OLS & Fixed-X Setting												
25	BIAS	-0.11	-0.33	-0.54	-0.43	-0.45	-0.10	-0.18	-0.14	-0.26	-0.24	
	SD	1.27	0.76	0.85	0.86	0.96	0.51	0.28	0.35	0.43	0.59	
	RMSE	1.27	0.83	1.01	0.96	1.06	0.52	0.33	0.38	0.50	0.64	
50	BIAS	-0.20	-0.37	-0.38	-0.31	-0.42	-0.07	-0.18	-0.16	-0.12	-0.16	
	SD	0.85	0.48	0.59	0.60	0.74	0.36	0.20	0.24	0.30	0.42	
	RMSE	0.88	0.60	0.70	0.67	0.85	0.37	0.27	0.29	0.32	0.45	
100	BIAS	-0.07	-0.16	-0.13	-0.31	-0.28	-0.04	-0.08	-0.08	-0.18	-0.07	
	SD	0.62	0.41	0.49	0.47	0.56	0.27	0.15	0.19	0.19	0.31	
	RMSE	0.62	0.44	0.51	0.56	0.62	0.27	0.17	0.21	0.26	0.32	
OLS & Random-X Setting												
25	BIAS	-0.17	-0.36	-0.37	-0.42	-0.53	-0.07	-0.17	-0.17	-0.20	-0.21	
	SD	1.16	0.67	0.83	0.94	0.97	0.53	0.28	0.35	0.44	0.56	
	RMSE	1.17	0.76	0.91	1.03	1.10	0.53	0.33	0.39	0.49	0.60	
50	BIAS	-0.15	-0.30	-0.29	-0.33	-0.37	-0.07	-0.13	-0.11	-0.16	-0.14	
	SD	0.87	0.50	0.65	0.67	0.73	0.39	0.20	0.27	0.33	0.41	
	RMSE	0.89	0.58	0.71	0.74	0.82	0.39	0.24	0.29	0.37	0.44	
100	BIAS	-0.10	-0.19	-0.24	-0.22	-0.27	-0.04	-0.10	-0.09	-0.09	-0.14	
	SD	0.59	0.37	0.46	0.54	0.55	0.26	0.15	0.19	0.24	0.30	
	RMSE	0.60	0.42	0.52	0.58	0.61	0.27	0.18	0.21	0.25	0.33	

measures across all four estimates. This finding is not surprising since Theorems 1 - 4 hold for both fixed and random settings.

After analyzing the estimation performance for Model A, we examine the performance of the parameter estimates under Model B. Tables S.5 - S.7 report the results when the random errors are generated from the mixture of normal distributions, while Tables S.8 - S.10 present the results when the random errors are simulated from the standard normal distribution. These tables show qualitatively similar findings to those in Tables 1, 2 and S.1 and Tables S.2 - S.4.

FIXED AND RANDOM COVARIANCE REGRESSION

TABLE 2

The average bias (BIAS), standard deviation (SD), and root mean squared error (RMSE) of the constrained GLS and FGLS estimates obtained from Model A with the mixture of normal random errors under the Fixed-X and Random-X settings.

				p = 5					p = 25		
n		$\hat{\beta}_1$	\hat{eta}_2	\hat{eta}_3	\hat{eta}_4	\hat{eta}_5	$\hat{\beta}_1$	\hat{eta}_2	\hat{eta}_3	\hat{eta}_4	\hat{eta}_5
GLS & Fixed-X Setting											
25	BIAS	0.06	-0.03	-0.00	0.00	-0.02	0.02	0.02	-0.01	-0.03	-0.04
	SD	1.30	0.62	0.64	0.64	0.73	0.55	0.21	0.27	0.29	0.44
	RMSE	1.30	0.62	0.64	0.64	0.73	0.55	0.21	0.27	0.29	0.44
50	BIAS	0.02	0.02	-0.00	-0.03	-0.03	-0.03	-0.02	-0.01	-0.01	-0.02
	SD	0.87	0.35	0.38	0.36	0.49	0.37	0.13	0.13	0.19	0.25
	RMSE	0.87	0.35	0.38	0.36	0.49	0.37	0.13	0.13	0.19	0.25
100	BIAS	0.02	0.02	0.02	-0.05	-0.02	-0.00	0.00	-0.00	-0.00	-0.00
	SD	0.61	0.28	0.34	0.30	0.37	0.27	0.09	0.09	0.08	0.14
	RMSE	0.61	0.29	0.34	0.30	0.37	0.27	0.09	0.09	0.08	0.14
GLS & Random-X Setting											
25	BIAS	-0.02	-0.03	-0.05	-0.03	-0.05	-0.02	-0.01	-0.02	-0.03	-0.02
	SD	1.28	0.56	0.64	0.68	0.75	0.53	0.21	0.25	0.29	0.39
	RMSE	1.28	0.56	0.64	0.68	0.75	0.53	0.21	0.25	0.29	0.39
50	BIAS	-0.04	-0.02	-0.03	-0.02	-0.06	-0.01	-0.02	0.01	-0.01	-0.02
	SD	0.86	0.37	0.46	0.46	0.50	0.38	0.14	0.17	0.21	0.26
	RMSE	0.86	0.37	0.46	0.46	0.50	0.38	0.14	0.17	0.21	0.26
100	BIAS	-0.01	-0.02	-0.01	-0.02	-0.01	-0.01	-0.00	-0.01	-0.00	-0.02
	SD	0.60	0.25	0.31	0.33	0.33	0.27	0.10	0.12	0.15	0.18
	RMSE	0.60	0.25	0.31	0.33	0.33	0.27	0.10	0.12	0.15	0.18
				FGI	LS & Fixe	ed-X Se	tting				
25	BIAS	-0.11	-0.29	-0.27	-0.34	-0.35	-0.07	-0.08	-0.09	-0.12	-0.17
	SD	1.26	0.84	1.04	0.92	1.05	0.54	0.25	0.32	0.39	0.63
	RMSE	1.26	0.89	1.08	0.98	1.10	0.55	0.26	0.33	0.41	0.65
50	BIAS	-0.13	-0.16	-0.18	-0.17	-0.32	-0.03	-0.04	-0.06	-0.05	-0.09
	SD	0.85	0.53	0.67	0.68	0.76	0.38	0.17	0.22	0.26	0.39
	RMSE	0.86	0.55	0.69	0.70	0.83	0.38	0.18	0.23	0.26	0.40
100	BIAS	-0.09	-0.09	-0.11	-0.14	-0.13	-0.04	-0.03	-0.04	-0.05	-0.06
	SD	0.61	0.37	0.51	0.50	0.55	0.27	0.13	0.18	0.24	0.22
	RMSE	0.61	0.38	0.52	0.52	0.56	0.28	0.13	0.18	0.24	0.23
	•	•		FGLS	& Rand	lom-X S	betting				
25	BIAS	-0.22	-0.29	-0.32	-0.34	-0.37	-0.06	-0.08	-0.10	-0.10	-0.12
	SD	1.24	0.74	0.87	1.04	1.24	0.52	0.25	0.32	0.40	0.55
	RMSE	1.26	0.79	0.93	1.09	1.30	0.53	0.27	0.33	0.41	0.57
50	BIAS	-0.11	-0.14	-0.18	-0.21	-0.23	-0.05	-0.05	-0.04	-0.06	-0.09
	SD	0.88	0.48	0.68	0.71	0.74	0.40	0.16	0.21	0.27	0.35
	RMSE	0.89	0.50	0.70	0.74	0.78	0.40	0.17	0.22	0.28	0.36
100	BIAS	-0.07	-0.10	-0.11	-0.13	-0.17	-0.04	-0.03	-0.04	-0.03	-0.05
	SD	0.63	0.36	0.56	0.58	0.84	0.27	0.13	0.19	0.22	0.30
	RMSE	0.63	0.37	0.57	0.60	0.85	0.27	0.13	0.19	0.22	0.30

In addition to parameter estimation, we next compare the expected test error of covariance regression models under the Fixed-X and Random-X settings. The expected test error is empirically calculated by averaging the test errors (3.2) from 1,000 realizations, which is the empirical version of Err defined in (3.3). Under the mixture of normal random errors, Figure S.1 depicts the corresponding expected test errors. Note that Figure S.1 is obtained under Model A, and the expected test error is calculated by setting $\mathbf{W}(X_{\alpha}) = \mathbf{V}^{-}(X_{\alpha})$ for $\alpha = 1, \dots, n$ and $01, \dots, 0n$, and $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}_{GLS}^{c}}(\cdot)$ in (3.2). To assess the impact of model misspecification under Model A, we consider the correctly-specified model $C_{\beta}(X_{\alpha}) = \beta^{(1)}I_{p} +$ $\sum_{k=2}^{5} \beta^{(k)} X_{\alpha}^{(k)} \text{ and a misspecified model } C_{\beta}(X_{\alpha}) = \beta^{(1)} I_p + \sum_{k=2}^{4} \beta^{(k)} X_{\alpha}^{(k)} + \beta^{(5)} \overline{X}_{\alpha}^{(5)},$ where $\overline{X}_{\alpha}^{(5)} = (\overline{X}_{\alpha,j_1j_2}^{(5)})_{p \times p}$ with $\overline{X}_{\alpha,j_1j_2}^{(5)} = (X_{\alpha,j_1j_2}^{(5)})^2$. In addition, Figure S.3 presents the results under the standard normal random errors.

For p = 5, Figure S.1 and Figure S.3 reveal the following important findings. (I) The Random-X expected test error, ErrR, is larger than the Fixed-X expected test error, ErrF, in both the mixture of normal and the normal random errors. Note that ErrF under our Fixed-X simulation setting converges to ErrS based on Proposition S.2 in Section S.4 of the supplementary material. Hence, this finding is consistent with the theoretical result in Theorem 6. In addition, the magnitude of the difference between ErrR and ErrF under the mixture of normal random errors is greater than that under normal random errors. (II) The difference between ErrR and ErrF becomes small as n increases when the model is correctly specified. This finding supports Remark 6. However, the difference increases when the model is misspecified; see the "Remark Regarding Theorem 6" in Section S.4 of the supplementary material. It is worth noting that Figure S.1 and Figure S.3 indicate that the aforementioned difference between ErrR and ErrF is not as strong for p = 25 as for p = 5. However, there still exist differences between ErrR and ErrF when the model is misspecified. These findings are consistent with the "Remark Regarding Theorem 6" in Section S.4.

Subsequently, we evaluate the performance of the estimates of expected test errors, Cp, \widehat{Cp} , RCp, \widehat{RCp} and OCV, when the data are generated under the Random-X setting. As mentioned in Section 3.2, Cp and \widehat{Cp} are obtained under Fixed-X, while RCp and \widehat{RCp} are obtained under Random-X. Accordingly, mis-using Cp and \widehat{Cp} under Random-X can underestimate the expected test error compared to that of RCp and \widehat{RCp} . In addition, we compute the average values of Cp, \widehat{Cp} , RCp, \widehat{RCp} , and OCV from 1,000 realizations, and then subtract the corresponding expected test errors. Accordingly, we obtain empirical biases that are a good measure for assessing performance. Figure S.2 reports the biases of Cp, \hat{C}_{p} , RC_{p} , $\hat{R}C_{p}$ and OCV in absolute value terms for the mixture of normal random errors. In addition, Figure S.4 presents the results for the standard normal random errors. These two figures reveal three important findings. (I) The absolute bias of RCp (RCp) is smaller than that of $C_{\rm D}$ ($\widehat{C_{\rm D}}$) across all settings. This finding is not surprising since $C_{\rm D}$ and $\widehat{C_{\rm D}}$ underestimate expected test errors under the Random-X setting; see the comment below equation (3.16). (II) RCp and \widehat{RCp} generally have a smaller absolute bias compared to OCV, though as n gets larger, the bias of OCV decreases. This finding is reasonable since OCV is designed for the general covariance regression model, while RCp and \widehat{RCp} are obtained for the linear covariance regression model, which is considered in this simulation study. (III) The absolute bias of $\widehat{\mathrm{RCp}}$ generally gets closer to RCp as n gets larger, which is sensible since the model is correctly specified; see the comment before equation (3.16) and in Remark 5. Based on these findings, we recommend using RCp in practice.

The above studies are based on Model A with the setting $\mathbf{W}(X_{\alpha}) = \mathbf{V}^{-}(X_{\alpha})$ and the estimate $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}_{GLS}^{c}}(\cdot)$. We next consider the setting $\mathbf{W}(X_{\alpha}) = I_{p}$ with the estimate $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}_{GLS}^{c}}(\cdot)$ under the same model. The comparisons between the expected test errors ErrR and ErrF as well as the examinations of the absolute biases of Cp, \widehat{Cp} , RCp, \widehat{RCp} and OCV, are presented in Figures S.5 – S.6 and Figures S.7 – S.8 for the mixture of normal random errors and the standard normal random errors, respectively. All of these figures are qualitatively similar to Figures S.1 – S.2 and Figures S.3 – S.4, although some of the differences between ErrR and ErrF and among Cp, \widehat{Cp} , RCp, \widehat{RCp} and OCV are not as strong as depicted in Figures S.1 – S.2 and Figures S.3 – S.4.

Finally, we compare expected test errors based on Model B under the following two different settings: (i) $\mathbf{W}(X_{\alpha}) = \mathbf{V}^{-}(X_{\alpha})$ with the estimate $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}_{cre}^{c}}(\cdot)$; and (ii)

 $\mathbf{W}(X_{\alpha}) = I_p$ with the estimate $\hat{\mathbf{C}}(\cdot) = \mathbf{C}_{\hat{\beta}_{OLS}^c}(\cdot)$. The comparisons between the expected test errors ErrR and ErrF as well as the examinations of the absolute biases are presented in Figures $\mathbf{S.9} - \mathbf{S.12}$ under setting (i), and in Figures $\mathbf{S.13} - \mathbf{S.16}$ under setting (ii). Compared to Figures $\mathbf{S.1} - \mathbf{S.4}$, the differences between ErrR and ErrF and among Cp, $\widehat{\mathbf{Cp}}$, RCp, $\widehat{\mathbf{RCp}}$ and OCV in Figures $\mathbf{S.9} - \mathbf{S.12}$ are less pronounced. On the other hand, the performances of Figures $\mathbf{S.13} - \mathbf{S.16}$ are qualitatively similar to those of Figures $\mathbf{S.5} - \mathbf{S.8}$. In addition to these results, additional simulation studies according to anonymous referees' suggestions are presented in Section $\mathbf{S.17}$ of the supplementary material.

5. Conclusion. In this article, we study two challenging tasks in covariance regression models. One is establishing asymptotic theories for parameter estimation in random covariance regression models; and the other is building up model assessment theories for assessing differences between the fixed and random covariance regression model settings. Specifically, under the Random-X setting, we demonstrate that both QMLE and WLS estimators are consistent and asymptotically normal as long as either n or p goes to infinity. As suggested by anonymous reviewers, we further obtain the asymptotic properties of the QMLE and WLS estimator by allowing several possibilities: a general error form for the data generating process, the dimension of β (i.e., K) to go to infinity, the model to be misspecified, and the conditional dependence of Y_i . In addition, the model assessment theory indicates that the difference between the expected test errors under the Fixed-X and Random-X settings is largely due to the bias and variance induced by the randomness in X. This finding allows us to propose two estimators, \widehat{RCp} and OCV, to estimate Random-X expected test errors. Simulation studies in Section 4 support both estimation and model assessment theories.

The pioneering work of this paper opens a new avenue in studying covariance regression models. For example, this framework for estimation and model assessment in the random covariance regression model (1.1) can be extended to analyze multivariate volatility models $\operatorname{Cov}[Y_i|\mathcal{G}_{i-1}] = C_{\beta}(X_{i-1})$ (see, e.g., Chapter 7 of Tsay, 2014), where \mathcal{G}_{i-1} is the σ -algebra generated from the observations up to time i-1, and X_i is a stochastic process. Specifically, we can estimate $Cov[Y_i|\mathcal{G}_{i-1}]$ and then construct time-varying portfolios for asset returns, which is an important subject in financial risk management; see, e.g., Tse and Tsui (2002), Laurent et al. (2012), and Poignard and Asai (2023). In addition, analogous estimation and model assessment approaches can be applied to mean-covariance regression models (see, e.g., Anderson, 1973 and Zou et al., 2022). For instance, we can adapt the mean-covariance regression model from Zou et al. (2022), and consider $Y_{\alpha} = BZ_{\alpha} + V_{\alpha}$, where $Z_{\alpha} \in \mathbb{R}^d$ is the *d*-dimensional explanatory vector, *B* is the $p \times d$ mean regression coefficient matrix, and V_{α} is standardized to have conditional mean zero and covariance matrix $\operatorname{Cov}[V_{\alpha}|\mathcal{F}_{\mathcal{X}}] = C_{\beta}(X_{\alpha})$. Furthermore, $\{(X_{\alpha}, Z_{\alpha}, Y_{\alpha}) : \alpha \in \{1, \cdots, n\}\}$ and $\{(X_{\alpha}, Z_{\alpha}, Y_{\alpha}) : \alpha \in \{01, \cdots, 0m\}\}$ are training data and test data, respectively, and $\mathcal{F}_{\mathcal{X}} =$ $\sigma((X_1, Z_1), \cdots, (X_n, Z_n), (X_{01}, Z_{01}), \cdots, (X_{0m}, Z_{0m}))$. Accordingly, $E[Y_{0i}|\mathcal{F}_{\mathcal{X}}] = BZ_{0i}$. Then, employing equation (3.2) of Zou et al. (2022, p. 321), we obtain the MLE of vec(B)below,

$$\operatorname{vec}(\hat{B}) = \left\{ \sum_{i=1}^{n} (Z_i \otimes I_p) C_{\hat{\beta}}^{-1}(X_i) (Z_i^{\top} \otimes I_p) \right\}^{-1} \sum_{i=1}^{n} (Z_i \otimes I_p) C_{\hat{\beta}}^{-1}(X_i) Y_i,$$

where $\hat{\beta}$ is the estimator of covariance regression parameter β . As a result, the prediction of Y_{0i} is $\hat{B}Z_{0i}$, for $i = 1, \dots, m$. It is clear that the prediction $\hat{B}Z_{0i}$ depends on the covariance regression structure $C_{\hat{\beta}}(\cdot)$. Hence, utilizing the estimation and model assessment approaches developed in this paper for covariance regression, one can select better models for $C_{\beta}(\cdot)$ and make more accurate inferences. In sum, a reliable $C_{\hat{\beta}}(\cdot)$ can play an important role in

It is also worth noting that in the context of nonparametric statistics, machine learning, and econometrics, theories are commonly established under the Random-X setting (see, e.g., Györfy et al., 2002, Vapnik, 1998 and Wooldridge, 2002). Accordingly, the proposed theoretical framework can facilitate extending covariance regression to nonparametric, machine learning, and econometric models for future research. Specifically, we can generalize the nonparametric variance function estimation (see, e.g., Wang et al., 2008) to nonparametric covariance function estimation, via nonparametric covariance regression models. Finally, Shen and Huang's (2020) two-level parametric bootstrap approach can be employed to estimate the bias difference $\mathcal{B}^{(d)}$ and variance difference $\mathcal{V}^{(d)}$ for covariance regression models. Compared to \widehat{RCp} , this type of approach does not require the model to be correctly specified, and the estimator of β can be different from the WLS estimator. We believe these extensions would strengthen the usefulness of the random covariance regression model.

References.

- Anderson, T. W. (1973). "Asymptotically efficient estimation of covariance matrices with linear structure," *The Annals of Statistics*, 1, 135-141.
- Bai, Z. D. and Silverstein, J. W. (2004). "Central limit theorem for linear spectral statistics of large dimensional sample covariance matrices," *The Annals of Probability*, 32, 553-605.
- Bollerslev, T. and Wooldridge, J. M. (1992). "Quasi-maximum likelihood estimation and inference in dynamic models with time-varying covariance," *Econometric Reviews*, 11, 143-173.
- Breiman, L. and Spector, P. (1992). "Submodel selection and evaluation in regression. The X-random case," *International statistical review/revue internationale de Statistique*, 60, 291-319.
- Cai, T. T., Hu, J., Li, Y. and Zheng, X. (2020). "High-dimensional minimum variance portfolio estimation based on high-frequency data," *Journal of Econometrics*, 214, 482-494.
- Chiu, T. Y., Leonard, T. and Tsui, K. W. (1996). "The matrix-logarithmic covariance model," *Journal of the American Statistical Association*, 91, 198-210.
- Chung, K. L. (2001). A Course in Probability Theory, Academic Press.
- Cressie, N. A. C. (1993). Statistics for Spatial Data, New York: Wiley.
- Demidenko, E. (2013). Mixed Models: Theory and Applications with R, John Wiley & Sons.
- DeMiguel, V., Garlappi, L. and Uppal, R. (2009). "Optimal versus naive diversification: How inefficient is the 1/N portfolio strategy?," *The Review of Financial Studies*, 22, 1915-1953.
- Fan, J., Li, Y. and Yu, K. (2012). "Vast volatility matrix estimation using high-frequency data for portfolio selection," *Journal of the American Statistical Association*, 107, 412-428.
- Fox, E. B. and Dunson, D. B. (2015). "Bayesian nonparametric covariance regression," *The Journal of Machine Learning Research*, 16, 2501-2542.
- Györfy, L., Kohler, M., Krzyzak, A. and Walk, H. (2002). A Distribution-Free Theory of Nonparametric Regression, New York: Springer.
- Hastie, T., Tibshirani, R. and Friedman, J. (2009). The Elements of Statistical Learning: Data Mining, Inference and Prediction, 2nd Edition, New York: Springer.
- Hoff, P. D. and Niu, X. (2012). "A covariance regression model," *Statistica Sinica*, 22, 729-753.
- Huang, D., Lan, W., Zhang, H. H. and Wang, H. (2019). "Least squares estimation of spatial autoregressive models for large-scale social networks," *Electronic Journal of Statistics*, 13, 1135-1165.
- Kelejian, H. and Prucha, I. (2001) "On the asymptotic distribution of the Moran I test statistic with applications," *Journal of Econometrics*, 104, 219-257.

- Laurent, S., Rombouts, J. V. and Violante, F. (2012). "On the forecasting accuracy of multivariate GARCH models," *Journal of Applied Econometrics*, 27, 934-955.
- Lee, L. F. (2004). "Asymptotic distributions of quasi-maximum likelihood estimators for spatial autoregressive models," *Econometrica*, 72, 1899-1925.
- LeSage, J. and Pace, R. K. (2009). *Introduction to Spatial Econometrics*, New York: Chapman & Hall.
- Ma, Y., Pan, R., Zou, T. and Wang, H. (2019). "A naive least squares method for spatial autoregression with covariates," *Statistica Sinica*, 30, 653-672.
- Mallows, C. L. (1973). "Some comments on C_p ," Technometrics, 15, 661-675.
- Markowitz, H. (1952). "Portfolio selection," The Journal of Finance, 7, 77-91.
- McCulloch, C. E., Searle, S.R. and Neuhaus, J. M. (2008). *Generalized, Linear, and Mixed Models*, 2nd Edition, John Wiley & Sons.
- Pan, G. and Zhou, W. (2008). "Central limit theorem for signal-to-interference ratio of reduced rank linear receiver," *The Annals of Applied Probability*, 18, 1232-1270.
- Poignard, B. and Asai, M. (2023). "High-dimensional sparse multivariate stochastic volatility models," *Journal of Time Series Analysis*, 44, 4-22.
- Rosset, S. and Tibshirani, R. J. (2020). "From Fixed-X to Random-X regression: Biasvariance decompositions, covariance penalties, and prediction error estimation," *Journal of the American Statistical Association*, 115, 138-151.
- Scott, M. A. and Handcock, M. S. (2001). "Covariance models for latent structure in longitudinal data," *Sociological Methodology*, 31, 265-303.
- Shen, X. and Huang, H. C. (2020). "Discussion of 'From Fixed-X to Random-X Regression: Bias-Variance Decompositions, Covariance Penalties, and Prediction Error Estimation'," *Journal of the American Statistical Association*, 115, 152-156.
- Tsay, R. S. (2014). Multivariate Time Series Analysis, Hoboken: John Wiley & Sons.
- Tse, Y. K. and Tsui, A. K. C. (2002). "A multivariate generalized autoregressive conditional heteroscedasticity model with time-varying correlations," *Journal of Business & Economic Statistics*, 20, 351-362.
- van der Vaart, A. W. (1998). Asymptotic Statistics, Cambridge University Press.
- Vapnik, N. V. (1998). Statistical Learning Theory, New York: Wiley.
- Wang, L., Brown, L. D., Cai, T. T. and Levine, M. (2008). "Effect of mean on variance function estimation in nonparametric regression," *The Annals of Statistics*, 36, 646-664.
- Wikle, C. K., Zammit-Mangion, A. and Cressie, N. (2019). *Spatio-Temporal Statistics with R*, Chapman and Hall.
- White, H. (1982). "Maximum likelihood estimation of misspecified models," *Econometrica*, 50, 1-25.
- Wooldridge, J. M. (2002). Econometric Analysis of Cross Section and Panel Data, MIT Press.
- Wooldridge, J. M. (2012). *Introductory Econometrics: A Modern Approach*, Mason, OH: Cengage Learning.
- Zheng, S. (2012). "Central limit theorems for linear spectral statistics of large dimensional F-matrices," *Annales de l'IHP Probabilités et statistiques*, 48, 444-476.
- Zhou, J., Tu, Y., Chen, Y. and Wang, H. (2017). "Estimating spatial autocorrelation with sampled network data," *Journal of Business & Economic Statistics*, 35, 130-138.
- Zou, T., Lan, W., Li, R. and Tsai, C.-L. (2022). "Inference on covariance-mean regression," Journal of Econometrics, 230, 318-338.
- Zou, T., Lan, W., Wang, H. and Tsai, C.-L. (2017). "Covariance regression analysis," *Journal* of the American Statistical Association, 112, 266-281.