GENERATORS OF TOP COHOMOLOGY

MANOJ KUMMINI AND MOHIT UPMANYU

ABSTRACT. Let *R* be a commutative noetherian ring and $f : X \longrightarrow$ Spec *R* a proper smooth morphism, of relative dimension *n*. From Hartshorne, *Residues and Duality*, Springer, 1966, one knows that the trace map $\text{Tr}_f : H^n(X, \omega_{X/R}) \longrightarrow R$ is an isomorphism when *f* has geometrically connected fibres. We construct an exact sequence that generates $\text{Ext}_X^n(\mathcal{O}_X, \omega_{X/R}) = H^n(X, \omega_{X/R})$ as an *R*-module in the following cases:

(1) when *R* is a DVR and *f* has a section;

(2) when $R = \mathbb{Z}$ and X is the Grassmannian $G_{2,m}$ for some $m \ge 4$.

This partially answers a question raised by Lipman.

1. INTRODUCTION

J. Lipman posed the following question. Let *R* be a (commutative) noetherian ring and $f : X \longrightarrow$ Spec *R* a proper smooth morphism, of relative dimension *n*. Write $\omega_{X/R} := \bigwedge^n \Omega_{X/R} = f!R[-n]$, the relative dualizing sheaf. By the duality theorem for proper morphisms [Har66, Chapter VII, Theorem 3.3] we get an isomorphism

 $\mathbf{R}f_*\mathbf{R}\mathcal{H}om_X(\omega_{X/R},\omega_{X/R}[n]) = \mathbf{R}\mathrm{Hom}_R(\mathbf{R}f_*\omega_{X/R},R).$

Since f is smooth, we also have a trace map [Har66, Chapter VII, Theorem 4.1]

 $\operatorname{Tr}_f: \operatorname{H}^n(X, \omega_{X/R}) \longrightarrow R$

which is an isomorphism when f has geometrically connected fibres [Har66, Chapter III, Theorem 11.2(g)].

Question 1.1 (Lipman). Can $\operatorname{Tr}_{f}^{-1}(1_R)$ be represented by a canonical exact sequence

$$0 \longrightarrow \omega_{X/R} \longrightarrow F_n \longrightarrow \cdots \longrightarrow F_1 \longrightarrow \mathscr{O}_X \longrightarrow 0 \in \operatorname{Ext}_X^n(\mathscr{O}_X, \omega_{X/R}) = \operatorname{H}^n(X, \omega_{X/R})?$$


For example, if $X = \mathbb{P}_{R}^{n}$, then the (exact) Koszul complex

$$0 \longrightarrow \omega_{X/R} \longrightarrow \mathscr{O}_X(-n)^{n+1} \longrightarrow \cdots \longrightarrow \mathscr{O}_X(-1)^{n+1} \longrightarrow \mathscr{O}_X \longrightarrow 0$$

generates $\operatorname{Ext}_X^n(\mathscr{O}_X, \omega_{X/R})$. One obtains this Koszul complex as follows. We have the Euler exact sequence (with $X = \mathbb{P}_R^n$)

$$0 \longrightarrow \Omega_{X/R} \longrightarrow \mathscr{O}_X(-1)^{n+1} \longrightarrow \mathscr{O}_X \longrightarrow 0$$

This corresponds to a global section *s* of $\mathcal{O}_X(+1)^{n+1}$, which defines a chain complex on $\bigwedge^* \mathcal{O}_X(-1)^{n+1}$. The exact Koszul complex given above is obtained when we take *s* is given by (x_0, \ldots, x_n) where the x_i are the homogeneous coordinates of \mathbb{P}_R^n . In general when X is a smooth connected projective scheme over R with a closed embedding $i : X \longrightarrow \mathbb{P}_{R}^{N}$, then we could consider the pull-back of the above Euler sequence on \mathbb{P}_{R}^{N} .



Here \mathcal{F} is the push-out and we denote the resulting map $\mathcal{F} \longrightarrow \mathcal{O}_X$ also by s. It is natural to consider whether the Koszul complex on X given by s is a generator of $\operatorname{Ext}_X^n(\mathcal{O}_X, \omega_{X/R})$. In Section 7, we show that this approach fails when $R = \mathbb{Z}$ and $X = G_{2,4}$. We show, more generally, that no element of $\operatorname{Ext}_X^1(\mathcal{O}_X, \Omega_{X/Z})$ will give a generator of $\operatorname{Ext}_X^n(\mathcal{O}_X, \omega_{X/Z})$.

However, in this paper, we directly describe a generator of $\operatorname{Ext}_X^n(\mathcal{O}_X, \omega_{X/R})$, in the following cases.

Theorem 1.2. *Question 1.1 has a positive answer if* R *is a DVR and* $f : X \longrightarrow$ Spec R *is a smooth projective morphism, with a section* Spec $R \longrightarrow X$.

Theorem 1.3. Question 1.1 has a positive answer when $R = \mathbb{Z}$ and X is the Grassmannian $G_{2,m}$, $m \ge 4$.

In the next section, we outline our strategy. Theorem 1.2 is proved as Theorem 5.1 in Section 5. Prior to that, in Section 4, we prove the analogous statement (Theorem 4.1) over fields, which is used in the proof of Theorem 5.1. Theorem 1.3 is proved in Section 6.

Acknowledgements. We thank Joseph Lipman for asking this question and providing us with some observations, Patrick Polo and Steven Sam for some clarification on the Grassmannian and Pramath Sastry for various discussions.

2. Strategy

Our starting point for finding a generator for Ext is the the following lemma. It might be well-known, but we include a proof at the end of this section for the sake of completeness.

Lemma 2.1. Let A be a Gorenstein local ring and $\mathbf{a} := a_1, \ldots, a_m$ a regular sequence in A. Then the augmented Koszul complex

 $0 \longrightarrow A \longrightarrow A^m \longrightarrow \cdots \longrightarrow A^m \longrightarrow A \longrightarrow A/\mathbf{a}A \longrightarrow 0$

is a generator of the A/**a**A-module $\operatorname{Ext}_{A}^{m}(A/\mathbf{a}A, A) \simeq A/\mathbf{a}A$.

To use the above lemma in our context, we look for a complete intersection inside X that is of relative dimension zero over R. For the strategy to work, this is not enough; we also need that $X \longrightarrow \operatorname{Spec} R$ has a section.

Notation 2.2. Let *R* be a Cohen-Macaulay ring with a dualizing module $\omega_R = R$ and *X* a smooth projective *R*-scheme, of relative dimension *n*. Assume that

- (1) the map $X \longrightarrow \operatorname{Spec} R$ has a section Z;
- (2) There exists a rank-*n* locally free sheaf \mathcal{F} on X and $\sigma \in H^0(X, \mathcal{F})$ such that the zero section X_n of σ is flat and of relative dimension 0 over R;
- (3) Z is a connected component of X_n .

Note that if *R* is not local, the hypothesis on *R* above might be stronger than requiring *R* to be Gorenstein, i.e., R_p is a Gorenstein local ring for all $p \in \text{Spec } R$.

Notation 2.3. We consider only relative dualizing sheaves in this paper. Therefore we abbreviate $\Omega_{X/R}$ as Ω_X and $\omega_{X/R}$ as ω_X .

Given such an \mathcal{F} and σ , we have the Koszul complex

$$0 \longrightarrow \det \mathcal{F}^* \longrightarrow \bigwedge^{n-1} \mathcal{F}^* \longrightarrow \cdots \longrightarrow \bigwedge^2 \mathcal{F}^* \longrightarrow \mathcal{F}^* \xrightarrow{\sigma} \mathcal{O}_X \longrightarrow 0,$$

which is a locally free resolution of \mathcal{O}_{X_n} , since X_n has codimension n inside X. Apply $\mathcal{H}om_{\mathcal{O}_X}(-, \omega_X)$ and note that $\omega_{X_n} = \mathcal{E}xt^n_{\mathcal{O}_X}(\mathcal{O}_{X_n}, \omega_X)$ to obtain an exact sequence

$$0 \longrightarrow \omega_X \longrightarrow \omega_X \otimes \mathcal{F} \longrightarrow \cdots \longrightarrow \omega_X \otimes \det \mathcal{F} \longrightarrow \omega_{X_n} \longrightarrow 0.$$

Since X_n is local complete intersection inside X and ω_X is a locally free \mathcal{O}_X -module of rank 1, we see that ω_{X_n} is a locally free \mathcal{O}_{X_n} -module of rank 1. We write \widetilde{K}_{\bullet} for this exact sequence. It is labelled in such a way that $\widetilde{K}_n = \omega_X \otimes \mathcal{F}$ and $\widetilde{K}_1 = \omega_X \otimes \det \mathcal{F}$.

Since Z is a connected component of X_n , we can write X_n as a disjoint union $Z \bigsqcup Z'$. Then we have a split exact sequence

$$(2.4) 0 \longrightarrow \mathscr{O}_Z \xrightarrow{i} \mathscr{O}_{X_n} \xrightarrow{\pi} \mathscr{O}_Z \longrightarrow 0$$

of \mathscr{O}_X -modules. Now consider the following commutative diagram. The lowest row is the exact sequence \widetilde{K}_{\bullet} . The middle row is the pull-back of the lowest row along the map *i*. The top row is the pull-back of the middle row along the natural surjective map $\mathscr{O}_X \longrightarrow \mathscr{O}_Z$.

$$(2.5) \qquad \begin{array}{c} 0 \longrightarrow \omega_X \longrightarrow \cdots \longrightarrow \widetilde{K}_2 \longrightarrow \mathcal{G}_1 \longrightarrow \mathcal{O}_X \longrightarrow 0 \\ & & & & & & \\ & & & & & & \\ 0 \longrightarrow \omega_X \longrightarrow \cdots \longrightarrow \widetilde{K}_2 \longrightarrow \mathcal{G}_0 \longrightarrow \mathcal{O}_Z \longrightarrow 0 \\ & & & & & & \\ 0 \longrightarrow \omega_X \longrightarrow \cdots \longrightarrow \widetilde{K}_2 \longrightarrow \widetilde{K}_1 \longrightarrow \omega_{X_n} \longrightarrow 0 \end{array}$$

Writing $I_{Z'}$ for the ideal sheaf of Z' in X, we see that $\mathcal{G}_0 = I_{Z'} \otimes \tilde{K}_1$.

The goal is to show that the top row of the above diagram generates $\operatorname{Ext}_X^n(\mathscr{O}_X, \omega_X)$ as an *R*-module. In the next proposition, we show that the middle row in the diagram (2.5) generates $\operatorname{Ext}_X^n(\mathscr{O}_Z, \omega_X)$ as an *R*-module.

Proposition 2.6. Let R and X be as above. Then

- (1) Let $x \in X_n$. Then the stalk of \widetilde{K}_{\bullet} at x generates the stalk of the sheaf $\mathcal{E}xt^n_{\mathscr{O}_X}(\mathscr{O}_{X_n},\omega_X)$ at x as an $\mathscr{O}_{X,x}$ -module.
- (2) The middle row in the diagram (2.5) generates $\operatorname{Ext}_X^n(\mathscr{O}_Z, \omega_X)$ as an R-module.

Proof. (1): Note that X and, therefore, X_n are Gorenstein. The proposition now follows from Lemma 2.1.

(2) From (2.4) we see that $i^*\pi^*$ is the identity map on $\mathcal{E}xt^n_{\mathcal{O}_X}(\mathcal{O}_Z,\omega_X)$. Hence, by (1), the middle row generates the stalk of $\mathcal{E}xt^n_{\mathcal{O}_X}(\mathcal{O}_Z,\omega_X)$ at every point on Z. Note that, since X is locally Cohen-Macaulay and $\operatorname{codim}_X(Z) = n$, $\mathcal{E}xt^i_{\mathcal{O}_X}(\mathcal{O}_Z,\omega_X) = 0$ for all i < n. Therefore $\operatorname{Ext}^n_X(\mathcal{O}_Z,\omega_X) = \Gamma(X, \mathcal{E}xt^n_{\mathcal{O}_X}(\mathcal{O}_Z,\omega_X))$. Since $Z = \operatorname{Spec} R$, we get the proposition.

We want to show that the top row in the diagram (2.5) generates $\operatorname{Ext}_X^n(\mathscr{O}_X, \omega_X)$ as an *R*-module (with suitable further hypothesis on *R*), by showing that the natural map $\operatorname{Ext}_X^n(\mathscr{O}_Z, \omega_X) \longrightarrow \operatorname{Ext}_X^n(\mathscr{O}_X, \omega_X)$ is an isomorphism. Write I_Z for the ideal sheaf of *Z* in *X*. Then we have the exact sequence

(2.7)
$$\operatorname{Ext}_{X}^{n}(\mathscr{O}_{Z},\omega_{X})\longrightarrow \operatorname{Ext}_{X}^{n}(\mathscr{O}_{X},\omega_{X})\longrightarrow \operatorname{Ext}_{X}^{n}(I_{Z},\omega_{X}).$$

By Proposition 2.6(2), $\operatorname{Ext}^{n}(\mathscr{O}_{Z}, \omega_{X})$ is a cyclic *R*-module. Hence it would suffice to show that

(2.8)
$$\operatorname{Ext}_X^n(\mathscr{O}_X,\omega_X) = R \text{ and } \operatorname{Ext}_X^n(I_Z,\omega_X) = 0.$$

We start with a few preliminary results.

Lemma 2.9. Let K be a field and Y a connected reduced projective K-scheme with a K-rational point. Then the natural map $K \longrightarrow H^0(Y, \mathcal{O}_Y)$ is an isomorphism.

Proof. Since Y is a connected reduced projective K-scheme, $H^0(Y, \mathcal{O}_Y)$ is a local reduced ring, finite over K, i.e, a finite extension field of K. Since Y has a K-rational point, there is a map $H^0(Y, \mathcal{O}_Y) \longrightarrow K$ of K-algebras. This proves the lemma.

Proposition 2.10. Assume that R is a normal domain. Then the natural maps $R \longrightarrow H^0(X, \mathcal{O}_X)$ and $H^0(X, \mathcal{O}_X) \longrightarrow H^0(Z, \mathcal{O}_Z)$ are isomorphisms.

Proof. Write $K = \operatorname{Frac}(R)$. We claim that $(R \longrightarrow \operatorname{H}^0(X, \mathscr{O}_X)) \otimes_R K$ is an isomorphism. Assume the claim. Note that $\operatorname{H}^0(X, \mathscr{O}_X)$ is an integral domain that is finite over R. By the claim, $\operatorname{Frac}(\operatorname{H}^0(X, \mathscr{O}_X)) = K$. Since R is normal, we get the first statement in proposition. The second statement follows immediately. To prove the claim, write X_K for the generic fibre $X \times_{\operatorname{Spec} R}$ Spec K. It is an integral scheme and has a K-rational point Z_K . By Lemma 2.9, the natural map $K \longrightarrow \operatorname{H}^0(X_K, \mathscr{O}_{X_K})$ is an isomorphism. By flat base-change, $\operatorname{H}^0(X_K, \mathscr{O}_{X_K}) = \operatorname{H}^0(X, \mathscr{O}_X) \otimes_R K$. This proves the claim.

We state the next proposition for one-dimensional PIDs, because the proofs of the ensuing lemmas assume that the ring is not a field. However the statement is true over fields also. In fact, over a field, we can prove the existence of the scheme X_n in Notation 2.2 with the desired properties. This is done in the next section.

Proposition 2.11. Assume that R is a PID that is not a field. Then the top row in the diagram (2.5) generates $\operatorname{Ext}_X^n(\mathcal{O}_X, \omega_X)$ as an R-module.

We prove this with a series of lemmas.

Lemma 2.12. $H^1(X, \mathcal{O}_X)$ is a free *R*-module.

Proof. Let *a* be an irreducible element of *R* and $\mathbb{k} = R/(a)$. Consider the exact sequence

$$0 \longrightarrow \mathrm{H}^{0}(X, \mathscr{O}_{X}) \xrightarrow{\cdot a} \mathrm{H}^{0}(X, \mathscr{O}_{X}) \longrightarrow \mathrm{H}^{0}(X_{\Bbbk}, \mathscr{O}_{X_{\Bbbk}}) \longrightarrow \mathrm{H}^{1}(X, \mathscr{O}_{X}) \xrightarrow{\cdot a} \mathrm{H}^{1}(X, \mathscr{O}_{X}) \longrightarrow$$

where $X_{\Bbbk} = X \times_{\text{Spec } R}$ Spec \Bbbk . Then $Z \times_{\text{Spec } R}$ Spec \Bbbk is a \Bbbk -rational point of the \Bbbk -scheme X_{\Bbbk} . Hence $H^0(X_{\Bbbk}, \mathscr{O}_{X_{\Bbbk}}) = \Bbbk$ by Lemma 2.9. By Proposition 2.10, $H^0(X, \mathscr{O}_X) = R$. Therefore the map $H^1(X, \mathscr{O}_X) \xrightarrow{\cdot a} H^1(X, \mathscr{O}_X)$ is injective. Since a is arbitrary, $H^1(X, \mathscr{O}_X)$ is torsion-free and, hence, free. \Box

Lemma 2.13. $H^0(X, I_Z) = 0$. $H^1(X, I_Z)$ is a free *R*-module.

Proof. Consider the exact sequence

 $0 \longrightarrow \mathrm{H}^{0}(X, I_{Z}) \longrightarrow \mathrm{H}^{0}(X, \mathscr{O}_{X}) \xrightarrow{p} \mathrm{H}^{0}(Z, \mathscr{O}_{Z}) \longrightarrow \mathrm{H}^{1}(X, I_{Z}) \longrightarrow \mathrm{H}^{1}(X, \mathscr{O}_{X}) \longrightarrow .$

The map p is an isomorphism, since $H^0(X, \mathcal{O}_X) \simeq R \simeq H^0(Z, \mathcal{O}_Z)$ and p(1) = 1. Hence, firstly, $H^0(X, I_Z) = 0$. Now note that $H^1(X, I_Z)$ is torsion-free, since it is a submodule of the free module $H^1(X, \mathcal{O}_X)$ (Lemma 2.12). Therefore $H^1(X, I_Z)$ is free.

Lemma 2.14. Let F be a quasi-coherent sheaf on X and q an integer. Then there is an exact sequence

$$0 \longrightarrow \operatorname{Ext}^{1}_{R}(\operatorname{H}^{q+1}(X,F),R) \longrightarrow \operatorname{Ext}^{n-q}(F,\omega_{X}) \longrightarrow \operatorname{Hom}_{R}(\operatorname{H}^{q}(X,F),R) \longrightarrow 0$$

of R-modules.

Proof. By \mathcal{D}_R^{\bullet} for the normalized dualizing complex of R. Note that \mathcal{D}_R^{\bullet} is isomorphic to the complex R[1] (i.e., the module R sitting in cohomological index -1) in the derived category of R. Write f for the map $X \longrightarrow \operatorname{Spec} R$. Then

$$f^{!}\mathcal{D}_{R}^{\bullet} := f^{*}\mathcal{D}_{R}^{\bullet} \otimes_{\mathscr{O}_{X}} \bigwedge^{n} \Omega_{X}[n] = \omega_{X}[n+1]$$

is a dualizing complex for X. By duality for proper morphisms, we have an isomorphism

$$\mathbf{R}f_*\mathbf{R}\mathcal{H}om_X(F, f^!\mathcal{D}_R^{\bullet}) = \mathbf{R}\mathcal{H}om_R(\mathbf{R}f_*F, \mathcal{D}_R^{\bullet})$$

in the derived category of *R*. Since $\mathbf{R}f_* = \mathbf{R}\Gamma$, we get $\mathbf{R}\operatorname{Hom}_X(F, f^!\mathcal{D}^{\bullet}_R) = \mathbf{R}\operatorname{Hom}_R(\mathbf{R}\Gamma F, \mathcal{D}^{\bullet}_R)$. Note that

$$\operatorname{Ext}^{n-q}(F,\omega_X) = \operatorname{H}^{n-q}(\operatorname{\mathbf{R}Hom}_X(F, f^!\mathcal{D}^{\bullet}_R[-n-1])) = \operatorname{H}^{-q-1}(\operatorname{\mathbf{R}Hom}_X(F, f^!\mathcal{D}^{\bullet}_R)).$$

We therefore compute $H^{-q-1}(\mathbb{R}\operatorname{Hom}_R(\mathbb{R}\Gamma F, \mathcal{D}_R^{\bullet}))$ using a spectral sequence. Fix a bounded-below complex G^{\bullet} of *R*-modules that represents $\mathbb{R}\Gamma F$. Without loss of generality, we can assume that \mathcal{D}_R^{\bullet} is an injective resolution of R[1]. Hence we have a double complex

$$E_0^{-i,-j} = \operatorname{Hom}_R(G^i, \mathcal{D}_R^{-j})$$

which is 0 whenever $-j \notin \{0, -1\}$. Taking the filtration by rows, we get a spectral sequence ${}_{h}E_{1}^{-i,-j} = \text{Hom}_{R}(\text{H}^{i}(X, F), \mathcal{D}_{R}^{-j})$ and, then

$${}_{h}E_{\infty}^{-i,-j} = {}_{h}E_{2}^{-i,-j} = \begin{cases} \operatorname{Ext}_{R}^{1}(\operatorname{H}^{i}(X,F),R), & -j = 0; \\ \operatorname{Hom}_{R}(\operatorname{H}^{i}(X,F),R), & -j = -1; \\ 0, & \text{otherwise} \end{cases}$$

The edge map is ${}_{h}E_{\infty}^{-i,0} \longrightarrow H^{-i}(\mathbb{R}Hom_{R}(\mathbb{R}\Gamma F, \mathcal{D}_{R}^{\bullet}))$; this is injective, and the cokernel is ${}_{h}E_{\infty}^{-i+1,-1}$. Thus we get an exact sequence

$$0 \longrightarrow \operatorname{Ext}^{1}_{R}(\operatorname{H}^{q+1}(X,F),R) \longrightarrow \operatorname{Ext}^{n-q}(F,\omega_{X}) \longrightarrow \operatorname{Hom}_{R}(\operatorname{H}^{q}(X,F),R) \longrightarrow 0$$

of *R*-modules.

Corollary 2.15. With notation as above,

(1) $\operatorname{Ext}^{n}(\mathcal{O}_{X}, \omega_{X}) \simeq R \text{ as } R \text{-modules.}$

(2) $\operatorname{Ext}^n(I_Z, \omega_X) = 0.$

Proof. (1): Apply Lemma 2.14 with q = 0 and $F = \mathcal{O}_X$. By Lemma 2.12, $\operatorname{Ext}^1_R(\operatorname{H}^1(X, \mathcal{O}_X), R) = 0$, so $\operatorname{Ext}^n(\mathcal{O}_X, \omega_X) \simeq \operatorname{Hom}_R(\operatorname{H}^0(X, \mathcal{O}_X), R) \simeq R$.

(2): Apply Lemma 2.14 with q = 0 and $F = I_Z$. Now use Lemma 2.13.

We can now prove Proposition 2.11.

Proof of Proposition 2.11. Corollary 2.15 establishes (2.8). Hence the proposition follows from (2.7). \Box

We conclude this section by giving a proof of Lemma 2.1.

Proof of Lemma 2.1. We prove this by induction on *m*. When m = 1, we have

$$(2.16) 0 \longrightarrow A \xrightarrow{a_1} A \longrightarrow A/a_1 A \longrightarrow 0$$

which, when applied $\text{Hom}_A(-, A)$, gives

$$0 \longrightarrow \operatorname{Hom}_{A}(A, A) \longrightarrow \operatorname{Hom}_{A}(A, A) \longrightarrow \operatorname{Ext}_{A}^{1}(A/a_{1}A, A) \longrightarrow 0$$

Then (2.16) corresponds to the image of id_A inside $Ext_A^1(A/a_1A, A)$ (see, e.g., [CE99, XIV, Theorem 1.1]) which is a generator of $Ext_A^1(A/a_1A, A)$ since $Hom_A(A, A)$ is a cyclic A-module generated by id_A .

Now assume that m > 1 and that the assertion holds for the sequence $\mathbf{b} := a_1, \ldots, a_{m-1}$. The Koszul complex $K_{\bullet}(\mathbf{a})$ (without the augmentation map $K_0(\mathbf{a}) \longrightarrow A/\mathbf{a}A$) is the mapping cone of the map

$$K_{\bullet}(\mathbf{b}) \xrightarrow{\cdot a_m} K_{\bullet}(\mathbf{b})$$

of complexes. Then we have an exact sequence

$$0 \longrightarrow K_{\bullet}(\mathbf{b}) \longrightarrow K_{\bullet}(\mathbf{a}) \longrightarrow K_{\bullet}(\mathbf{b})[-1] \longrightarrow 0$$

of complexes. An element $\xi \in \operatorname{Ext}_A^{m-1}(A/\mathbf{b}A, A)$ is given by a map $K_{\bullet}(\mathbf{b}) \longrightarrow A[-m+1]$ in the derived category of *A*-modules. (Here *A* is thought of as the complex $0 \longrightarrow A \longrightarrow 0$, with *A* in homological position 0.) The image of ξ under the connecting morphism $\operatorname{Ext}_A^{m-1}(A/\mathbf{b}A, A) \longrightarrow \operatorname{Ext}_A^m(A/\mathbf{a}A, A)$ is given by the composite map

$$K_{\bullet}(\mathbf{a}) \longrightarrow K_{\bullet}(\mathbf{b})[-1] \longrightarrow A[-m].$$

In particular, if we take the identity map $K_{m-1}(\mathbf{b}) \longrightarrow A$ (which gives a generator of $\operatorname{Ext}_A^{m-1}(A/\mathbf{b}A, A)$), we get the identity map $K_m(\mathbf{a}) \longrightarrow A$ (which gives a generator of $\operatorname{Ext}_A^m(A/\mathbf{a}A, A)$).

3. Example: the projective plane

In this section, we exhibit $\mathbb{P}^2_{\mathbb{Z}}$ as an example of the strategy described in the previous section. We can get the middle row of (2.5) directly (for a suitable choice of X_2 and section Z). We will then show that the top row of (2.5) equals the Koszul complex in $\text{Ext}^2(\mathcal{O}_X, \omega_X)$.

Write $X = \mathbb{P}^2_{\mathbb{Z}}$. Let x_0, x_1, x_2 be homogeneous coordinates on X and X_2 be the subscheme defined by (x_1, x_2) . Take $Z = X_2$. In other words, $\mathcal{F} = \mathcal{O}_X(1)^{\oplus 2} = \mathcal{O}_X \epsilon_1 \oplus \mathcal{O}_X \epsilon_2$ and $\sigma = (x_1, x_2) = x_1 \epsilon_1 + x_2 \epsilon_2$.

 $\omega_X = \mathcal{O}_X(-3) = \mathcal{O}_X \cdot (\epsilon_1^* \wedge \epsilon_2^* \wedge \epsilon_3^*)$. Since $X_2 = \text{Spec } \mathbb{Z}$, $\omega_{X_2} = \mathcal{O}_{X_2}$. Abbreviate $e_i \wedge e_j = e_{ij}$ etc. Hence the bottom and the middle rows of (2.5) are equal to

$$0 \to \mathcal{O}_X \epsilon_{123}^* \xrightarrow{\begin{bmatrix} x_1 \\ -x_2 \end{bmatrix}} \xrightarrow{\mathcal{O}_X \epsilon_{23}^*} \xrightarrow{\begin{bmatrix} x_2 & x_1 \end{bmatrix}} \mathcal{O}_X \epsilon_3^* \to \mathcal{O}_Z \epsilon_3^* \to 0$$

The Koszul complex on the section $\sum_{i=1}^{3} x_i \epsilon_i^*$ of $\bigoplus_{i=1}^{3} \mathcal{O}_X \epsilon_i^* = \mathcal{O}_X (-1)^{\oplus 3}$ is

$$0 \rightarrow \mathcal{O}_{X}\epsilon_{123}^{*} \xrightarrow{\begin{pmatrix} x_{1} \\ -x_{2} \\ x_{3} \end{pmatrix}} \stackrel{\mathcal{O}_{X}\epsilon_{23}^{*}}{\underset{\bigoplus}{\oplus}} \xrightarrow{\begin{pmatrix} x_{2} & x_{1} & 0 \\ -x_{3} & 0 & x_{1} \\ 0 & -x_{3} & -x_{2} \end{pmatrix}} \stackrel{\mathcal{O}_{X}\epsilon_{3}^{*}}{\underset{\bigoplus}{\oplus}} \xrightarrow{\begin{pmatrix} x_{2} & x_{1} \\ \oplus \\ \mathcal{O}_{X}\epsilon_{13}^{*} \\ \oplus \\ \mathcal{O}_{X}\epsilon_{12}^{*} \\ \end{pmatrix}} \stackrel{\mathcal{O}_{X}\epsilon_{2}^{*}}{\underset{\bigoplus}{\oplus}} \xrightarrow{\begin{pmatrix} x_{3} & x_{2} & x_{1} \\ \oplus \\ \mathcal{O}_{X}\epsilon_{12}^{*} \\ \end{pmatrix}} \stackrel{\mathcal{O}_{X} \leftarrow 0$$

Therefore we get a commutative diagram; for the sake of brevity we suppress the basis elements and the matrices. The right-most vertical map is the canonical surjection, while the other maps are the projection maps to the respective summands.

Now use [ML63, Proposition III.5.1] to see that the push-forward of the top row of (3.1) along the map $\mathrm{id}_{\mathscr{O}_X(-3)}$ is the same as the pull-back of the bottom row of (3.1) along the canonical map $\mathscr{O}_X \longrightarrow \mathscr{O}_Z$. Hence the top row of (3.1) is the top row of (2.5).

4. Over a field

In this section, we show that the strategy described in the previous section can be carried out over a field. This result will be used to prove the result for DVR.

Theorem 4.1. Let R be a field. Let X be a smooth projective R-subscheme of \mathbb{P}_R^N of relative dimension n. Assume that X has an R-rational point Z. Write $S = R[x_0, \ldots, x_N]$ for the homogeneous coordinate ring of \mathbb{P}_R^N . Then:

- (1) There exist homogeneous polynomials $f_1, \ldots, f_n \in S$ such that they define a complete intersection X_n inside X and Z is a point of X_n .
- (2) The top row of (2.5) generates $\operatorname{Ext}_X^n(\mathcal{O}_X, \omega_X)$ as an R-vector space. Moreover, the base-change of this exact sequence generates $\operatorname{Ext}_{X_S}^n(\mathcal{O}_{X_S}, \omega_{X_S})$ for all ring maps $R \longrightarrow S$ (where $X_S = X \times_{\operatorname{Spec} R} \operatorname{Spec} S$).

For a homogeneous ideal J of $R[x_0, \ldots, x_N]$, we write $J^{\text{sat}} = \bigcup_i (J : (x_0, \ldots, x_N)^i)$, i.e, the ideal of the projective subscheme of \mathbb{P}_R^N defined by J.

Proof of Theorem 4.1. Note that Z is defined by *n* linearly independent linear forms. After a suitable change of coordinates, we may assume that Z is defined by $\mathfrak{p} := (x_1, \ldots, x_n)$.

(1): We prove this by induction on k. Let k = 1. Note that I_X is a prime ideal of height N - n < N. Hence $\mathfrak{p} \not\subseteq (\mathfrak{p}^2 \cup I_X)$ and the assertion holds for k = 1.

Now assume that k > 1 and that we have found f_1, \ldots, f_{k-1} with the desired properties. Since X is non-singular, \bar{S}/I_X and, therefore, $\bar{S}/(I_X + (f_1, \ldots, f_{k-1}))^{\text{sat}}$ are Cohen-Macaulay except possibly at the irrelevant ideal. Since $\operatorname{ht}(I_X + (f_1, \ldots, f_{k-1})) = N - n + k - 1 < N$, we can find an $f_k \in \mathfrak{p} \setminus \mathfrak{p}^2$ that is a non-zero-divisor in $\bar{S}/(I_X + (f_1, \ldots, f_{k-1}))^{\text{sat}}$.

(2): Write I_Z for the ideal sheaf of Z in X. By Lemma 2.9, $H^0(X, \mathcal{O}_X) = R$, and arguing as in the proof of Lemma 2.13, $H^0(X, I_Z) = 0$. Hence, by Serre duality [Har77, Theorem III.7.6], (2.8) holds; therefore by (2.7), we get the first assertion. Since S is R-flat, we get the second assertion by flat base-change.

(The statement of Serre duality in [Har77, Theorem III.7.6] assumes that the base field is algebraically closed. However, this assumption is used only to see that local rings on projective spaces (over a field) are regular. This holds over arbitray fields, and not just over algebraically closed fields.)

5. Over a DVR

In this section, we show that the strategy described in the previous section can be carried out over a DVR. We assume that $X \longrightarrow \text{Spec } R$ has a section (with R a DVR) has a section Z, and show that there exists a complete intersection $X_n \subseteq X$ in which Z is a connected component.

Theorem 5.1. Let (R, \mathfrak{m}, \Bbbk) be a DVR. Let X be a projective R-subscheme of \mathbb{P}_R^N . Assume that the structure morphism $X \longrightarrow \operatorname{Spec} R$ is smooth and of relative dimension n. and that it has a section $\operatorname{Spec} R \longrightarrow Z \subseteq X$. Write $S = R[x_0, \ldots, x_N]$ for the homogeneous coordinate ring of \mathbb{P}_R^N . Then:

- (1) There exist homogeneous polynomials $\tilde{f}_1, \ldots, \tilde{f}_n \in S$ such that they define a complete intersection X_n inside X.
- (2) The natural map $X_n \longrightarrow \operatorname{Spec} R$ is flat and has zero-dimensional fibres.
- (3) Z is a connected component of X_n . Hence \mathcal{O}_Z is a direct summand of \mathcal{O}_{X_n} .
- (4) The top row of (2.5) generates $\operatorname{Ext}_X^n(\mathcal{O}_X, \omega_X)$ as a free R-module. Moreover, the base-change of this exact sequence generates $\operatorname{Ext}_{X_c}^n(\mathcal{O}_{X_S}, \omega_{X_S})$ for all ring maps $R \longrightarrow S$ (where $X_S = X \times_{\operatorname{Spec}} R$ Spec S).

Various parts of the above theorem will be proved as separate lemmas / propositions, in more generality than stated in the theorem: (1) and (2) as Proposition 5.4; (3) as Proposition 5.5; finally (4) is proved. We start with an observation.

Proposition 5.2. Without loss of generality, Z is defined by the ideal (x_1, \ldots, x_N) .

Proof. Without loss of generality, the homogeneous coordinate x_0 does not vanish at the closed point of Z. Hence it does not vanish at the generic point of Z also. Therefore we may assume that Z lies inside the open subset $\{x_0 \neq 0\} \simeq \mathbb{A}_R^N$. Write $y_i = \frac{x_i}{x_0}$.

Write $Z = \text{Spec}(R[y_1, \ldots, y_N]/I)$, or, equivalently, $R[y_1, \ldots, y_N]/I = R$. Let $a_i \in R$ be the image of $y_i, 1 \leq i \leq N$. Then $(y_1 - a_1, \ldots, y_N - a_N) \subseteq I$. Hence $I = (y_1 - a_1, \ldots, y_N - a_N)$. Therefore Z is defined inside \mathbb{P}_R^N by $(x_1 - a_1x_0, \ldots, x_N - a_nx_0)$. After a change of coordinates, we may assume that this is the ideal (x_1, \ldots, x_N) .

For subschemes *Y* of \mathbb{P}_{R}^{N} , we write *Y*^{cl} for its closed fiber over *R*.

Proposition 5.3. Write $\overline{S} := \mathbb{k}[x_0, \dots, x_N]$ and $\mathfrak{p} := (x_1, \dots, x_N)\overline{S}$. There exist homogeneous $f_1, \dots, f_n \in \mathfrak{p} \setminus \mathfrak{p}^2$ such that for all $1 \le k \le n$, f_k is a non-zero-divisor in $\overline{S}/(I_{X^{cl}} + (f_1, \dots, f_{k-1}))^{sat}$.

Proof. See the proof of Theorem 4.1(1).

Proposition 5.4. Assume that (R, \mathfrak{m}, \Bbbk) is a local Cohen-Macaulay ring. For $1 \le k \le n$, write

$$f_k = x_0^{\deg f_k - 1} l_k(x_1, \dots, x_N) + terms of lower degree in x_0$$

where l_k is linear. Let $\tilde{l}_k(x_1, \ldots, x_N)$ be a homogeneous linear lift of l_k to $R[x_1, \ldots, x_N]$. Consider any lift \tilde{f}_k of f_k to $R[x_0, \ldots, x_N]$ of the form

 $x_0^{\deg f_k-1}\tilde{l}_k(x_1,\ldots,x_N)$ + terms of lower degree in x_0 .

Let $X_k = X \cap D_{\tilde{f}_1} \cap \cdots \cap D_{\tilde{f}_k}$. (Here $D_{\tilde{f}_i}$ is the subscheme defined by \tilde{f}_1 and the intersection is the schemetheoretic intersection.) Then

(1) $\tilde{f}_k \in (x_1, ..., x_N) \setminus ((x_1, ..., x_N)^2 + I_{X_{k-1}}).$ (2) X_k is Cohen-Macaulay and flat over R, of relative dimension n - k.

Proof. (1): It is immediate from the definition that $\tilde{f}_k \in (x_1, ..., x_N)$. The other part follows from Proposition 5.3 since the reduction of the ideal $((x_1, ..., x_N)^2 + I_{X_{k-1}})$ modulo \mathfrak{m}_A lies inside $((x_1, ..., x_N)^2 + I_{X_{k-1}})$.

(2): *X* is Cohen-Macaulay and flat over *R*. Hence X^{cl} is Cohen-Macaulay [Mat89, Corollary to Theorem 23.3]. By Proposition 5.3, f_1, \ldots, f_k is a regular sequence on X^{cl} . Hence by [Mat89, Corollary to Theorem 22.5], $\tilde{f_1}, \ldots, \tilde{f_k}$ is a regular sequence on *X* and X_k is flat over *R*. Therefore X_k is additionally Cohen-Macaulay and of relative dimension n - k.

Proposition 5.5. Suppose that (R, \mathfrak{m}, \Bbbk) is a regular local ring. Then Z is a connected component of X_n .

Proof. X_n is finite and flat over R (Proposition 5.4). In order to prove the assertion, it suffices to show that Z is the only irreducible component containing the closed point of Z. Let (A, \mathfrak{n}) be the local ring of \mathcal{O}_{X_n} at the closed point of Z.

Since X_n is flat over R, A is flat over R, so every R-regular sequence in \mathfrak{m} is also A-regular. In particular A is Cohen-Macaulay. Since Z^{cl} is an isolated point of X_n^{cl} , it follows that $A/\mathfrak{m}A = \Bbbk$. Hence the multiplicity $e(\mathfrak{m}A, A)$ of the ideal $\mathfrak{m}A$ in A is 1, by [Mat89, Theorem 17.11]. Hence $e(\mathfrak{n}, A) = 1$, which implies that A is a regular local ring and, *a fortiori*, a domain which is what we needed to show.

By now, we have proved (1), (2) and (3) of Theorem 5.1.

Conclusion of the proof of Theorem 5.1. The first assertion of (4) follows from Proposition 2.11.

To complete the proof of Theorem 5.1, we need to show that the base-change of the top row in the diagram (2.5) gives a generator of $H^n(X_S, \omega_{X_S})$ for each ring map $R \longrightarrow S$. (Here $X_S := X \times_{\text{Spec } R} \text{Spec } S$.) We need to show two things:

(1) $(0 \longrightarrow \omega_X \longrightarrow \cdots \longrightarrow \widetilde{K}_2 \longrightarrow \mathcal{G}_1 \longrightarrow \mathcal{O}_X \longrightarrow 0) \otimes_R S$ is exact. For this, we need to show that the sheaves are flat over R. The terms other than \mathcal{G}_1 are locally free over X, so it remains to show that \mathcal{G}_1 is torsion-free over R. First note that $\mathcal{G}_0 = I_{Z'} \otimes \widetilde{K}_1$ is torsion-free. Let g be a (local) section of \mathcal{G}_1 that is torsion over R. The images of g in \mathcal{O}_X and in \mathcal{G}_0 are zero since both these sheaves are torsion-free over R. Since \mathcal{G}_1 is the pull-back, it follows that g = 0.

(2) $(0 \longrightarrow \omega_X \longrightarrow \cdots \longrightarrow \widetilde{K}_2 \longrightarrow \mathcal{G}_1 \longrightarrow \mathcal{O}_X \longrightarrow 0) \otimes_R S$ is non-zero in $\operatorname{Ext}_{X_S}^n(\mathcal{O}_{X_S}, \omega_{X_S})$. Suppose it is zero for some S. Then it would be zero for some field F with a ring map $S \longrightarrow F$, so we may

assume that *S* is a field. In this case, we immediately reduce (since maps between fields are faithfully flat) to the case that *S* is either the residue field of *R* or the fraction field of *R*. Since the fraction field of *R* is flat over *R* and $\operatorname{Ext}_X^n(\mathscr{O}_X, \omega_X) \simeq R$, the assertion follows in this case. Hence assume that $S = \Bbbk$.

We first argue that if we apply $-\otimes_R \Bbbk$, to the diagram (2.5) for R, we get diagram (2.5) for \Bbbk . Indeed, since the rows consist of sheaves that are torsion-free over R (Z = Spec R and X_n is flat over R), the rows remain exact after applying $-\otimes_R \Bbbk$. It is also immediate that $\mathcal{O}_X, \mathcal{O}_Z, \mathcal{O}_{X_n}$ are replaced by $\mathcal{O}_{X^{\text{cl}}}, \mathcal{O}_{Z^{\text{cl}}}, \mathcal{O}_{X_n^{\text{cl}}}$, respectively. Hence it remains to show that $\omega_X \otimes_R \Bbbk = \omega_{X^{\text{cl}}}$. This is indeed true, since X^{cl} is the pull-back of the divisor $\text{Spec } \Bbbk \subseteq \text{Spec } R$, and, therefore, the normal bundle of X^{cl} inside X is isomorphic to $\mathcal{O}_{X^{\text{cl}}}$. Therefore the base-change of the diagram (2.5) for R along the map $R \longrightarrow \Bbbk$ gives the diagram (2.5) for \Bbbk . By Theorem 4.1(2), the top row of the diagram (2.5) for \Bbbk generates $\text{Ext}_{X^{\text{cl}}}^n(\mathcal{O}_{X^{\text{cl}}}, \omega_{X^{\text{cl}}})$ as an \Bbbk -vector space.

This concludes the proof of Theorem 5.1.

6. Grassmannian

Let $m \ge 4$ and X the Grassmannian $G_{2,m}$ over \mathbb{Z} , i.e., the scheme whose R-points (for a ring R) classify the locally free (over R) quotient modules of rank m - 2 of $R^{\oplus m}$. It is a smooth \mathbb{Z} -scheme of relative dimension n := 2(m - 2). We show that the morphism $X \longrightarrow$ Spec \mathbb{Z} has a section Z and that there exists a complete intersection X_n inside X of relative dimension 0 over \mathbb{Z} satisfying the conditions of Notation 2.2. Since \mathbb{Z} is a PID, the strategy of Section 2 can be carried out to give a generator of $\text{Ext}_X^n(\mathscr{O}_X, \omega_X)$ as a free \mathbb{Z} -module.

Let $N = \binom{m}{2} - 1$. Then X can be embedded inside $\mathbb{P} := \mathbb{P}_{\mathbb{Z}}^{N}$ as a closed subscheme, defined by the Pluecker relations; see [EH00, Chapter III, §2.7]. Let $p_{i,j}$, $1 \le i < j \le m$ be homogeneous coordinates for \mathbb{P} . Let $S = \mathbb{Z}[p_{i,j}, 1 \le i < j \le m]$. The main result of this section is:

Theorem 6.1. For $3 \le k \le 2m - 1$, write $l_k = \sum_{i+j=k} p_{i,j}$. Write $V \subseteq \mathbb{P}$ for the linear subvariety defined by $l_3, l_4, \ldots, \widehat{l_{m+1}}, \ldots, l_{2m-1}$. Then $X \cap V$ has zero-dimensional fibres and there is a section Spec $\mathbb{Z} \longrightarrow X \cap V$ which defines an irreducible and reduced component of $X \cap V$.

To prove that $X \cap V$ has zero-dimensional fibres, we may do a base-change along $\mathbb{Z} \longrightarrow \Bbbk$ where \Bbbk is an algebraically closed field, and show that $X_{\Bbbk} \cap V_{\Bbbk}$ (the respective base-changes) is a zero-dimensional scheme. We do this now. We use [Ses07, Chapter 1] as the reference on Schubert varieties and standard monomial theory. We denote the Schubert varieties in X_{\Bbbk} by $Y_{i,j} 1 \le i < j \le n$. (This is the Schubert variety in X_{\Bbbk} corresponding to a permutation $\sigma \in S_n$ with $\{\sigma(1), \sigma(2)\} = \{1, 2\}$.)

Proposition 6.2. Let \Bbbk be an algebraically closed field. Then $X_{\Bbbk} \cap V_{\Bbbk}$ is a zero-dimensional scheme.

Proof. We think of X_{\Bbbk} as the quotient of the space of rank-two $m \times 2$ matrices under the natural action of $GL_2(\Bbbk)$ on the right. The homogeneous coordinate $p_{i,j}$ of \mathbb{P}_{\Bbbk} becomes, on X_{\Bbbk} , the determinant of the 2×2 minor consisting of the rows *i* and *j* and both the columns.

Note that V_{\Bbbk} is defined inside \mathbb{P}_{\Bbbk} by (the images in $S \otimes_{\mathbb{Z}} \Bbbk$) of the l_k . For each $m + 2 \le k \le 2m - 1$, let W_k be the linear subvariety defined by l_k, \ldots, l_{2m-1} . We claim that $X_{\Bbbk} \cap W_k$ is set-theoretically the union of all the Schubert subvarieties of X_{\Bbbk} of dimension k - 4. Note that $k - 4 = \dim X_{\Bbbk} - \operatorname{codim}_{\mathbb{P}_{\Bbbk}} W_k$. Therefore the irreducible components of $X_{\Bbbk} \cap W_{m+2}$ are precisely the Schubert varieties $Y_{k,m+1-k}$ for all $1 \le k < \frac{m}{2}$.

For each $3 \le k \le m$, let W'_k be the linear subvariety defined by l_3, \ldots, l_k . By symmetry, and assuming the above claim, $X_{\Bbbk} \cap W'_k$ is set-theoretically the union of all the opposite Schubert subvarieties of X_{\Bbbk}

of codimension k - 2. In particular, the irreducible components of $X_{\Bbbk} \cap W'_m$ are precisely the opposite Schubert varieties $Y^{k,m+1-k}$ for all $1 \le k < \frac{m}{2}$. Hence the irreducible components of $X_{\Bbbk} \cap V_{\Bbbk} = (X_{\Bbbk} \cap W_{m+2}) \cap (X_{\Bbbk} \cap W'_m)$ are the Richardson varieties $Y^{k,m+1-k}_{k,m+1-k}$ for all $1 \le k < \frac{m}{2}$, all of which are zerodimensional. (See, e.g., [Bri05, § 1.3].)

To prove the claim, we proceed by downward induction on k. When k = 2m - 1, $X_{\Bbbk} \cap W_k$ is the Schubert divisor $Y_{m-2,m}$, which is of dimension 2m-5. Assume that the claim has been proved up to k+1. Let Y be a Schubert variety of dimension k - 3. Hence Y is an irreducible component of $X_{\Bbbk} \cap W_{k+1}$. Say $Y = Y_{i,k-i}$. Then, in the terms of l_k , all but $l_{i,k-i}$ vanish along Y, since the ideal of $Y_{i,k-i}$ in X_{\Bbbk} is generated by $\{p_{j_1,j_2} \mid (j_1, j_2) \nleq (i, k - i)\}$. Moreover, $Y_{i,k-i} \cap \{p_{i,k-i} = 0\}$ is the union of the codimension-1 Schubert subvarieties of $Y_{i,k-i}$. Further, every (k - 4)-dimensional Schubert variety is a subvariety of a (k - 3)-dimensional Schubert variety. Hence $X_{\Bbbk} \cap W_k = X_{\Bbbk} \cap W_{k+1} \cap \{l_k = 0\}$ is set-theoretically the union of all the (k - 4)-dimensional Schubert varieties.

Proof of Theorem 6.1. As we remarked above, Proposition 6.2 implies that $X \cap V$ has zero-dimensional fibres. We now prove the existence of a section of the structure morphism $X \cap V \longrightarrow$ Spec \mathbb{Z} with the desired properties. Let $\mathbb{A} \subseteq \mathbb{P}$ be the affine space obtained by inverting $p_{1,n}$. More precisely, write $q_{i,j} = \frac{p_{i,j}}{p_{1,n}}$, $R = \mathbb{Z}[q_{i,j} : 1 \le i < j \le n, (i, j) \ne (1, n)]$ and \mathfrak{m} for the prime ideal of R generated by the $q_{i,j}$. Then $\mathbb{A} =$ Spec R. Note that X is defined by the the Pluecker relations, which are of the form

$$p_{i,j}p_{i',j'} \pm p_{i,i'}p_{j',j} \pm p_{i,j'}p_{i',j}$$
 where $i < i', j > j'$.

See, e.g., [EH00, Chapter III, §2.7]. (For the sake of convenience, we do not specify which terms on the right appear with positive sign and which with negative sign.) A Pluecker relation of the form

$$p_{1,n}p_{i',j'} \pm p_{1,k}p_{j',n} \pm p_{1,j'}p_{k,n}$$
, with $1 < i' < j' < n$

gives a generator

$$q_{i',j'} \pm q_{1,k}q_{j',n} \pm q_{1,j'}q_{k,n}$$

of the ideal of $X \cap \mathbb{A}$ inside \mathbb{A} . The other generators (i.e., with $i < i', j > j', (i, j) \neq (1, n)$) give the polynomials $q_{i,j}q_{i',j'} \pm q_{i,k}q_{j',j} \pm q_{i,j'}q_{k,j}$ inside the ideal of $X \cap \mathbb{A}$. For $3 \le k \le 2m - 1, k \ne n + 1$, the polynomial l_k gives the polynomial $q_{1,k-1} + q_{2,k-2} + \cdots \in R$.

We need to show that \mathfrak{m} is a minimal prime over the *R*-ideal \mathfrak{a} generated by

- (1) $q_{i',j'} \pm q_{1,i'}q_{j',n} \pm q_{1,j'}q_{i',n}, 1 < i' < j' < n;$
- (2) $q_{i,j}q_{i',j'} \pm q_{i,i'}q_{j',j} \pm q_{i,j'}q_{i',j}, i < i', j > j', (i, j) \neq (1, n);$
- (3) (a) $q_{1,k-1} + q_{2,k-2} + \cdots, 3 \le k \le n;$ (b) $\cdots + q_{k-n+1,n-1} + q_{k-n,n}, n+2 \le k \le 2m-1$
- and that $\mathfrak{a}R_{\mathfrak{m}} = \mathfrak{m}R_{\mathfrak{m}}$. Note that $\mathfrak{a} \subseteq \mathfrak{m}$. Therefore if we show that $\Omega_{(S/\mathfrak{a})/\mathbb{Z}}$ vanishes at \mathfrak{m} , it would follow that S/\mathfrak{a} is unramified over \mathbb{Z} at \mathfrak{m} , whence it would follow that $\mathfrak{a}R_{\mathfrak{m}} = \mathfrak{m}R_{\mathfrak{m}}$. Note that $\Omega_{(S/\mathfrak{a})/\mathbb{Z}}$ is

follow that S/\mathfrak{a} is unramified over \mathbb{Z} at \mathfrak{m} , whence it would follow that $\mathfrak{a}R_{\mathfrak{m}} = \mathfrak{m}R_{\mathfrak{m}}$. Note that $\Omega_{(S/\mathfrak{a})/\mathbb{Z}}$ is the cokernel of the jacobian matrix of the aforementioned generating set of \mathfrak{a} with respect to the variables $q_{i',j'}$. Further, for each (i', j'), there exists a unique generator in the above list in which $q_{i',j'}$ appears with coefficient 1, so the jacobian matrix has full rank in the residue field $\kappa(\mathfrak{m})$ at \mathfrak{m} . Therefore $\Omega_{(S/\mathfrak{a})/\mathbb{Z}}$ vanishes at \mathfrak{m} .

Proof of Theorem 1.3. Theorem 6.1 implies that the conditions on Notation 2.2 are met. Hence we can apply Proposition 2.11 to complete the proof of Theorem 1.3.

MANOJ KUMMINI AND MOHIT UPMANYU

7. An example

In this section we show that the approach of finding a generator of $\text{Ext}_X^1(\mathcal{O}_X, \Omega_{X/R})$ when $X = \mathbb{P}_R^n$ does work in the case of Grassmannians, as mentioned in the Introduction.

Let $R = \mathbb{Z}$ and $X = G_{2,4}$. Write $\mathbb{P}^5 = \mathbb{P}^5_{\mathbb{Z}}$. Let H_1, \ldots, H_4 be the hyperplanes in \mathbb{P}^5 defined by the linear polynomials l_3, l_4, l_6, l_7 (defined in the previous section) respectively. Let $D_i = X \cap H_i, 1 \le i \le 4$.

Proposition 7.1. $H^{i}(\mathbb{P}^{5}, \Omega_{\mathbb{P}^{5}}(m)) \simeq H^{i}(X, \Omega_{X}(m))$ for all *i* and for m = 0, 1.

Proof. Let *I* be the ideal defining *X* inside \mathbb{P}^5 . It is generated by a degree 2 polynomial. Hence $I/I^2 \simeq \mathcal{O}_X(-2)$. Take the conormal sequence twisted by $\mathcal{O}_X(m)$:

$$0 \longrightarrow \mathscr{O}_X(m-2) \longrightarrow \Omega_{\mathbb{P}^5}|_X(m) \longrightarrow \Omega_X(m) \longrightarrow 0.$$

We see that $H^*(X, \mathcal{O}_X(m-2)) = 0$ from the exact sequence

$$0 \longrightarrow \mathscr{O}_{\mathbb{P}^5}(m-4) \longrightarrow \mathscr{O}_{\mathbb{P}^5}(m-2) \longrightarrow \mathscr{O}_X(m-2) \longrightarrow 0.$$

Hence $\operatorname{H}^{*}(X, \Omega_{\mathbb{P}^{5}}|_{X}(m)) = \operatorname{H}^{*}(X, \Omega_{X}(m)).$

0

To calculate $H^*(X, \Omega_{\mathbb{P}^5}|_X(m))$, consider the exact sequence (obtained by applying $- \otimes \Omega_{\mathbb{P}^5}(2)$ to the above sequence)

$$\longrightarrow \Omega_{\mathbb{P}^5}(m-2) \longrightarrow \Omega_{\mathbb{P}^5}(m) \longrightarrow \Omega_{\mathbb{P}^5}|_X(m) \longrightarrow 0.$$

We want to show that $H^*(\mathbb{P}^5, \Omega_{\mathbb{P}^5}(m-2)) = 0$, which we get from the Euler exact sequence (after a twist)

$$0 \longrightarrow \Omega_{\mathbb{P}^5}(m-2) \longrightarrow \mathscr{O}_{\mathbb{P}^5}(m-3)^6 \longrightarrow \mathscr{O}_{\mathbb{P}^5}(m-2) \longrightarrow 0.$$

Corollary 7.2. $H^1(X, \Omega_X) = \mathbb{Z}$.

Proof. Follows from Proposition 7.1 and the Euler exact sequence.

Proposition 7.3. Let $1 \le i \le 4$. The natural map $\operatorname{Ext}^1(\mathcal{O}_{D_i}, \omega_X) \longrightarrow \operatorname{Ext}^1(\mathcal{O}_X, \omega_X) = \operatorname{H}^1(X, \mathcal{O}_X)$ is an isomorphism.

Proof. Write $D = D_i$. From the Euler exact sequence, we get that $H^0(\mathbb{P}^5, \Omega_{\mathbb{P}^5}(1)) = H^1(\mathbb{P}^5, \Omega_{\mathbb{P}^5}(1)) = 0$. Hence Proposition 7.1 gives that $H^0(X, \Omega_X(1)) = H^1(X, \Omega_X(1)) = 0$. From the exact sequence

$$0 \longrightarrow \mathscr{O}_X(-1) \longrightarrow \mathscr{O}_X \longrightarrow \mathscr{O}_D \longrightarrow 0$$

we get the exact sequence

$$\operatorname{Hom}_X(\mathscr{O}_X(-1),\Omega_X)\longrightarrow \operatorname{Ext}^1(\mathscr{O}_D,\omega_X)\longrightarrow \operatorname{Ext}^1(\mathscr{O}_X,\omega_X)\longrightarrow \operatorname{Ext}^1(\mathscr{O}_X(-1),\omega_X).$$

Since the first and the last terms are zero, we get the isomorphism.

Fix a generator

 $0\longrightarrow \Omega_X\longrightarrow \mathcal{F} \xrightarrow{s} \mathscr{O}_X\longrightarrow 0$

of $\operatorname{Ext}^1(\mathscr{O}_X, \omega_X)$. (Note that s is a global section of \mathcal{F}^* .) Let

$$(7.4) 0 \longrightarrow \Omega_X \longrightarrow \mathcal{F}_{D_i} \xrightarrow{s_i} \mathscr{O}_{D_i} \longrightarrow 0$$

be its inverse image in $\operatorname{Ext}^{1}(\mathscr{O}_{D_{i}}, \omega_{X})$.

In what follows, for a chain complex L_{\bullet} , $\mathbf{h}_{i}(L_{\bullet})$ denotes the *j*th homology of L_{\bullet} , i.e.,

$$\frac{\ker(L_j \longrightarrow L_{j-1})}{\operatorname{Im}(L_{j+1} \longrightarrow L_j)}.$$

Lemma 7.5. Let $1 \le i \le 4$. Let L_{\bullet} be a bounded complex such that $\operatorname{Tor}_{k}(L_{j}, \mathcal{O}_{D_{i}}) = 0$ for all j and k > 0. Write L'_{\bullet} for the total complex of $L_{\bullet} \otimes (\mathcal{F}_{D_{i}} \longrightarrow \mathcal{O}_{D_{i}})$. Then $\mathbf{h}_{j}(L'_{\bullet}) = \mathbf{h}_{j-1}(L_{\bullet}) \otimes \Omega_{X}$.

Proof. $L_{\bullet} \otimes (\mathcal{F}_{D_j} \longrightarrow \mathcal{O}_{D_j})$ gives a third-quadrant double complex

$$E^{-p,-q} \begin{cases} L_p \otimes \mathcal{O}_{D_j}, & q = 0\\ L_p \otimes \mathcal{F}_{D_j}, & q = 1\\ 0, & \text{otherwise.} \end{cases}$$

Since the vertical map $E^{-p,-q} \longrightarrow E^{-p,-q+1}$ are surjective and $\text{Tor}_1(L_p, \mathcal{O}_{D_j}) = 0$, the ${}_vE_1$ page of the spectral sequence is

$${}_{v}E_{1}^{-p,-q}\begin{cases} L_{p}\otimes\Omega_{X}, & q=1\\ 0, & \text{otherwise} \end{cases}$$

Hence L'_{\bullet} is quasi-isomorphic to $L_{\bullet} \otimes \Omega_X[-1]$.

Lemma 7.6. Let $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$ be coherent sheaves on X such that $\operatorname{Tor}_{\geq 1}^{\mathscr{O}_X}(\mathcal{F}_1, \mathcal{F}_2) = 0 = \operatorname{Tor}_{\geq 1}^{\mathscr{O}_X}(\mathcal{F}_2, \mathcal{F}_3)$. Then for all i, then $\operatorname{Tor}_i^{\mathscr{O}_X}(\mathcal{F}_1 \otimes \mathcal{F}_2, \mathcal{F}_3) = \operatorname{Tor}_i^{\mathscr{O}_X}(\mathcal{F}_1, \mathcal{F}_2 \otimes \mathcal{F}_3)$ for all i.

Proof. Let \mathcal{L}_1 and \mathcal{L}_3 be locally free resolutions of \mathcal{F}_1 and \mathcal{F}_3 respectively. Then we have an isomorphism $(\mathcal{L}_1 \otimes \mathcal{F}_2) \otimes \mathcal{L}_3 \longrightarrow \mathcal{L}_1 \otimes (\mathcal{F}_2 \otimes \mathcal{L}_3)$. By hypothesis, $\mathcal{L}_1 \otimes \mathcal{F}_2$ is quasi-isomorphic to $\mathcal{F}_1 \otimes \mathcal{F}_2$, so $\mathbf{h}_*((\mathcal{L}_1 \otimes \mathcal{F}_2) \otimes \mathcal{L}_3) = \operatorname{Tor}_*^{\mathscr{O}_X}(\mathcal{F}_1 \otimes \mathcal{F}_2, \mathcal{F}_3)$. Similarly, $\mathbf{h}_*(\mathcal{L}_1 \otimes (\mathcal{F}_2 \otimes \mathcal{L}_3)) = \operatorname{Tor}_*^{\mathscr{O}_X}(\mathcal{F}_1, \mathcal{F}_2 \otimes \mathcal{F}_3)$.

Lemma 7.7. *Let* $A \subseteq \{1, ..., 4\}$ *and* $b \in \{1, ..., 4\} \setminus A$ *. Then*

$$\operatorname{Tor}_{\geq 1}^{\mathscr{O}_X}\left(\bigotimes_{a\in A}\mathcal{G}_a,\mathcal{G}_b\right)=0.$$

for all $(\mathcal{G}_1,\ldots,\mathcal{G}_4) \in \prod_{i=1}^4 \{\mathscr{O}_{D_i},\mathcal{F}_{D_i}\}.$

Proof. We may assume that $\mathcal{G}_b = \mathcal{O}_{D_b}$, since for every quasi-coherent sheaf \mathcal{G} , $\operatorname{Tor}_{\geq 1}^{\mathcal{O}_X}(\mathcal{G}, \mathcal{F}_{D_b}) \subseteq \operatorname{Tor}_{\geq 1}^{\mathcal{O}_X}(\mathcal{G}, \mathcal{O}_{D_b})$ by (7.4). We proceed by induction on $(|A|, \delta_A)$, where $\delta_A = |\{a \in A \mid \mathcal{G}_a = \mathcal{F}_{D_a}\}|$. We order such pairs lexicographically.

Suppose that $\delta_A = 0$. Since the coordinate ring of X is Cohen-Macaulay, the homogeneous polynomials l_3 , l_4 , l_6 , l_7 form a regular sequence in any order. Therefore, irrespective of |A|, the assertion of the lemma follows. When (|A|, δ_A) = (1, 1), we use the fact that by (7.4),

$$\operatorname{Tor}_{\geq 1}^{\mathscr{O}_X}(\mathscr{F}_{D_a}, \mathscr{O}_{D_b}) \subseteq \operatorname{Tor}_{\geq 1}^{\mathscr{O}_X}(\mathscr{O}_{D_a}, \mathscr{O}_{D_b})$$

which is zero by the case $\delta_A = 0$.

Now assume that |A| > 1 and $\delta_A > 0$. Let $a_1 \in A$ be such that $\mathcal{G}_a = \mathcal{F}_{D_a}$ for some $a \in A' := A \setminus \{a_1\}$. By induction on |A|,

$$\operatorname{Tor}_{\geq 1}^{\mathscr{O}_{X}}\left(\bigotimes_{a\in A'}\mathcal{G}_{a},\mathcal{G}_{a_{1}}\right)=0 \quad \text{and} \quad \operatorname{Tor}_{\geq 1}^{\mathscr{O}_{X}}\left(\mathcal{G}_{a_{1}},\mathscr{O}_{D_{b}}\right)=0$$

Hence by Lemma 7.6,

$$\operatorname{Tor}_{i}^{\mathscr{O}_{X}}\left(\bigotimes_{a\in A}\mathcal{G}_{a},\mathscr{O}_{D_{b}}\right) = \operatorname{Tor}_{i}^{\mathscr{O}_{X}}\left(\bigotimes_{a\in A'}\mathcal{G}_{a},\mathcal{G}_{a_{1}}\otimes\mathscr{O}_{D_{b}}\right)$$

for all *i*. Repeating this, we find $\tilde{a} \in A$ such that $\mathcal{G}_{\tilde{a}} = \mathcal{F}_{D_{\tilde{a}}}$ and such that

$$\operatorname{Tor}_{i}^{\mathscr{O}_{X}}\left(\bigotimes_{a\in A}\mathcal{G}_{a},\mathscr{O}_{D_{b}}\right) = \operatorname{Tor}_{i}^{\mathscr{O}_{X}}\left(\mathcal{F}_{D_{\tilde{a}}},\bigotimes_{\substack{a\in A\\a\neq\tilde{a}}}\mathcal{G}_{a}\otimes\mathscr{O}_{D_{b}}\right) = 0$$

for all $i \ge 1$, since $\delta_{(A \setminus \{\tilde{a}\}) \cup \{b\}} = \delta_A - 1$.

Proposition 7.8. Tot $\left(\bigotimes_{k=1}^{4}(\mathcal{F}_{D_{k}} \xrightarrow{s_{i}} \mathscr{O}_{D_{k}})\right)$ gives a six-term exact sequence of the form

$$\alpha: \quad 0 \longrightarrow \Omega_X^{\otimes 4} \longrightarrow \cdots \longrightarrow \bigoplus_{i=0}^4 \mathcal{F}_i|_{D_1 \cap \cdots \widehat{D_i} \cdots \cap D_4} \longrightarrow \mathscr{O}_{D_1 \cap \cdots \cap D_4} \longrightarrow 0$$

Proof. We prove this by repeated application of Lemma 7.5. Let $L_{\bullet}^{(1)} = 0 \longrightarrow \mathcal{F}_{D_1} \longrightarrow \mathcal{O}_{D_1} \longrightarrow 0$ and, define, for $2 \leq i \leq 4$, Let $L_{\bullet}^{(i)} = \operatorname{Tot}(L_{\bullet}^{(i-1)} \otimes (\mathcal{F}_{D_i} \longrightarrow \mathcal{O}_{D_i}))$. Note that for each $1 \leq i \leq 3$, $L_{\bullet}^{(i)}$ satisfies the hypotheses of Lemma 7.5: Firstly, for each $j, L_j^{(i)}$ is a tensor product of a few \mathcal{F}_{D_k} and a few \mathcal{O}_{D_k} , so we can use Lemma 7.7. Secondly by induction on i, we see that $\mathbf{h}_j(L_{\bullet}^{(i)}) = 0$ unless j = i, and $\mathbf{h}_i(L_{\bullet}^{(i)}) \simeq \Omega_X^{\otimes i}$.


Proposition 7.9. Tot $\left(\bigotimes_{k=1}^{4} (\mathcal{F} \xrightarrow{s} \mathcal{O}_X)\right)$ gives a six-term exact sequence of the form

$$\beta: \quad 0 \longrightarrow \Omega_X^{\otimes 4} \longrightarrow \cdots \longrightarrow \bigoplus_{i=0}^4 \mathcal{F} \longrightarrow \mathcal{O}_X \longrightarrow 0.$$

Proof. Let L_{\bullet} be a bounded complex. Write $L'_{\bullet} = \operatorname{Tot}(L_{\bullet} \otimes (\mathcal{F} \xrightarrow{s} \mathcal{O}_X))$. Then $\mathbf{h}_j(L'_{\bullet}) = \mathbf{h}_{j-1}(L_{\bullet}) \otimes \Omega_X$. Now repeatedly apply this to $\operatorname{Tot}\left(\bigotimes_{k=1}^r (\mathcal{F} \xrightarrow{s} \mathcal{O}_X)\right), 2 \le r \le 4$.

We now obtain the following diagram of complexes, where the first row is α from Proposition 7.8 the second row is β from Proposition 7.9 and the last row γ is the *exact* Koszul complex for the map

14



 $(\pi \text{ and } \mu \text{ are the natural maps.})$ Note that $\alpha \in \operatorname{Ext}_X^4(\mathscr{O}_{D_1 \cap \dots \cap D_4}, \Omega_X^{\otimes 4}), \beta \in \operatorname{Ext}_X^4(\mathscr{O}_X, \Omega_X^{\otimes 4})$ and $\gamma \in \operatorname{Ext}_X^4(\mathscr{O}_X, \omega_X)$.

Lemma 7.10. $\pi^*(\alpha) = \beta$ and $\mu_*(\beta) = \gamma$.

Proof. Use [ML63, Proposition III.5.1] to see that $\pi^*(\alpha) = (\mathrm{id}_{\Omega_X^{\otimes 4}})_*(\beta) = \beta$ and that $\mu_*(\beta) = (\mathrm{id}_{\mathscr{O}_X})^*(\gamma) = \gamma$.

Lemma 7.11. $\pi^*(\mu_*(\alpha)) = \mu_*(\pi^*(\alpha)) = \gamma$.

Proof. The second equality is from Lemma 7.10. To prove the first equality, let \mathcal{A} be an abelian category with enough injectives. Write $Ch(\mathcal{A})$ for the abelian category of chain complexes over \mathcal{A} . Let A, B, C, D be objects in \mathcal{A} and $\pi : C \longrightarrow A$ and $\mu : B \longrightarrow D$ be morphisms. Let $n \in \mathbb{N}$. We want to show that $\pi^*\mu_* = \mu_*\pi^*$ as maps from $\operatorname{Ext}^n_{\mathcal{A}}(A, B)$ to $\operatorname{Ext}^n_{\mathcal{A}}(C, D)$. Let I_B and I_D be injective resolutions of B and D respectively. Then the following diagram commutes

Taking homology we see that $\pi^* \mu_* = \mu_* \pi^*$ as maps from $\operatorname{Ext}^n_{\mathcal{A}}(A, B)$ to $\operatorname{Ext}^n_{\mathcal{A}}(C, D)$.

Proposition 7.12. With notation as above:

- (1) $D_1 \cap \cdots \cap D_4$ is a reduced scheme $\{P_1, P_2\}$ where, for $i = 1, 2, P_i \simeq \text{Spec } \mathbb{Z}$.
- (2) The automorphism τ of X given by $p_{i,j} \mapsto p_{k,l}$ where $\{i, j, k, l\} = \{1, \dots, 4\}$ sends P_1 to P_2 and vice versa.

Proof. (1): In the Pluecker embedding, $X = G_{2,4}$ is defined inside $\mathbb{P}^5 := \operatorname{Proj} \mathbb{Z}[p_{1,2}, \dots, p_{3,4}]$ by the polynomial $p_{1,2}p_{3,4} - p_{1,3}p_{2,4} + p_{1,4}p_{2,3}$. Hence $D_1 \cap \dots \cap D_4$ is defined by $(p_{1,2}p_{3,4} - p_{1,3}p_{2,4} + p_{1,4}p_{2,3}, p_{1,2}, p_{1,3}, p_{2,4}, p_{3,4}, p_{1,4}p_{2,3})$.

(2): Note that

$$(p_{1,2}, p_{1,3}, p_{2,4}, p_{3,4}, p_{1,4}) \mapsto (p_{1,2}, p_{1,3}, p_{2,4}, p_{3,4}, p_{2,3})$$

and vice versa.

Proposition 7.13. γ is an even number when we identify $\text{Ext}_X^4(\mathcal{O}_X, \omega_X)$ with \mathbb{Z} . In particular it is not a generator of $\text{Ext}_X^4(\mathcal{O}_X, \omega_X)$.

Proof. Note that $\operatorname{Ext}_X^4(\mathscr{O}_{D_1\cap\cdots\cap D_4}, \omega_X) \simeq \operatorname{Ext}_X^4(\mathscr{O}_{P_1}, \omega_X) \oplus \operatorname{Ext}_X^4(\mathscr{O}_{P_2}, \omega_X) \simeq \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$. (The ε_i are basis elements.) The automorphism τ of X defined in Proposition 7.12 gives an isomorphism $\tau : \mathbb{Z}\varepsilon_1 \xrightarrow{\sim} \mathbb{Z}\varepsilon_2$, with $\varepsilon_1 \mapsto \varepsilon_2$ and $\varepsilon_2 \mapsto \varepsilon_1$. Write $\mu_*(\alpha) = m_1\varepsilon_1 + m_2\varepsilon_2$. Hence $\tau^*\mu_*(\alpha) = m_2\varepsilon_1 + m_1\varepsilon_2$. We see from Proposition 7.12 that $\mu_*(\tau^*\alpha) = m_2\varepsilon_1 + m_1\varepsilon_2$. On the other hand, note that τ permutes the divisors D_1, \ldots, D_4 . Hence, we see from the definition of α (Proposition 7.8) that $\tau^*\alpha = \alpha$. Therefore $m_1 = m_2$, so $\mu_*\alpha = m_1(\varepsilon_1 + \varepsilon_2)$. Thus we see that it is enough to show that $\pi^*(\varepsilon_1 + \varepsilon_2)$ is not a generator of $\operatorname{Ext}_X^4(\mathscr{O}_X, \omega_X)$.

Write π_i^* for the isomorphism $\operatorname{Ext}_X^4(\mathscr{O}_{P_i}, \omega_X) \longrightarrow \operatorname{Ext}_X^4(\mathscr{O}_X, \omega_X)$, i = 1, 2. Then $\pi^*(\varepsilon_1 + \varepsilon_2) = \sum_i \pi_i^*(\varepsilon_i)$. Since π_i is an isomorphism, $\pi_i^*(\varepsilon_i)$ is a generator of $\operatorname{Ext}_X^4(\mathscr{O}_X, \omega_X) \simeq \mathbb{Z}$. Therefore $\pi^*(\varepsilon_1 + \varepsilon_2)$ is an even multiple of a generator of $\operatorname{Ext}_X^4(\mathscr{O}_X, \omega_X)$.

References

- [Bri05] M. Brion. Lectures on the geometry of flag varieties. In *Topics in cohomological studies of algebraic varieties*, Trends Math., pages 33–85. Birkhäuser, Basel, 2005. arXiv:math/0410240 [math.AG]. 11
- [CE99] H. Cartan and S. Eilenberg. *Homological algebra*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1999. With an appendix by David A. Buchsbaum, Reprint of the 1956 original. 6
- [EH00] D. Eisenbud and J. Harris. The geometry of schemes, volume 197 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2000. 10, 11
- [Har66] R. Hartshorne. *Residues and duality*. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20. Springer-Verlag, Berlin, 1966. 1
- [Har77] R. Hartshorne. Algebraic geometry. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52. 8
- [Mat89] H. Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1989. Translated from the Japanese by M. Reid. 9
- [ML63] S. Mac Lane. Homology. Die Grundlehren der mathematischen Wissenschaften, Band 114. Springer-Verlag, Berlin-Göttingen-Heidelberg; Academic Press, Inc., Publishers, New York, 1963. 7, 15
- [Ses07] C. S. Seshadri. Introduction to the theory of standard monomials, volume 46 of Texts and Readings in Mathematics. Hindustan Book Agency, New Delhi, 2007. With notes by Peter Littelmann and Pradeep Shukla, Appendix A by V. Lakshmibai, Revised reprint of lectures published in the Brandeis Lecture Notes series. 10

CHENNAI MATHEMATICAL INSTITUTE, SIRUSERI, TAMILNADU 603103. INDIA *Email address:* mkummini@cmi.ac.in

CHENNAI MATHEMATICAL INSTITUTE, SIRUSERI, TAMILNADU 603103. INDIA *Email address*: mohit@cmi.ac.in