# The impact of memory-burdened primordial black holes on high-scale leptogenesis

Roberta Calabrese,  $^{1,\,2,\,*}$  Marco Chianese,  $^{3,\,2,\,\dagger}$  and Ninetta Saviano  $^{2,\,3,\,\ddagger}$ 

<sup>1</sup>Dipartimento di Fisica "Ettore Pancini", Università degli studi di Napoli

"Federico II", Complesso Univ. Monte S. Angelo, I-80126 Napoli, Italy

<sup>2</sup> INFN - Sezione di Napoli, Complesso Univ. Monte S. Angelo, I-80126 Napoli, Italy

<sup>3</sup>Scuola Superiore Meridionale, Università degli studi di Napoli "Federico II", Largo San Marcellino 10, 80138 Napoli, Italy

We explore the impact of the back-reaction of evaporation on the quantum state of Primordial Black Holes (PBHs), known as "memory burden", on the baryon asymmetry production in the Universe through high-scale leptogenesis. Focusing on PBH masses ranging from 1 to 1000 grams, we investigate the interplay between the non-thermal production of heavy sterile neutrinos and the entropy injection within this non-standard cosmological framework. By assuming appropriate values for the memory-burden parameters, q = 1/2 and k = 1, we derive mutual exclusion limits between PBHs and thermal leptogenesis in the mixed parameter space. Our analysis reveals that the primary contribution of PBHs to baryon asymmetry stems from entropy injection. Indeed, we find that, differently from earlier studies based on the semi-classical Hawking evaporation, the memory-burden effect suppresses the non-thermal source term in the PBH mass range explored. This has significant implications for understanding baryogenesis in such alternative cosmological scenarios.

# I. INTRODUCTION

The seminal idea that Primordial Black Holes (PBHs) have been formed in the very early universe from dense regions long before stars emerged [1-3], has been recently resumed, gaining new interest in relation to the context of dark matter [4–17], early Universe investigation [18–21], as well ad gravitational wave research [22–26]. The mass and spin distributions of PBHs can vary based on the specific cosmological scenario. PBHs with masses above  $10^{15}$  grams are stable over cosmological timescales, but lighter black holes may have already evaporated through Hawking radiation by the current era. This process of evaporation has been investigated in a variety of different contexts, playing important roles in the production of the baryon asymmetry of the Universe, gravitational wave emission, or dark matter generation, depending on their mass. Black hole evaporation, as described by the semiclassical framework, assumes that the black hole retains its classical nature throughout its entire lifetime [27]. However, it is becoming increasingly evident that this assumption may lead to an inconsistent model, pointing to the necessity for new physics, especially in relation to the information loss paradox [28]. Hawking's analysis fails to account for the back-reaction of emitted particles on the quantum state of the black hole. This omission becomes particularly important when the energy of the emitted quanta approaches the black hole's total energy. Recent research [29, 30] suggests that such back-reaction could give rise to a "memory burden", a phenomenon where the system's information resists being lost, driven by the response of quantum modes linked to its entropic degrees of freedom. As the black hole's mass decreases past a

certain point, the back-reaction becomes increasingly significant, potentially slowing the evaporation rate and extending the black hole's lifetime. Among the significant phenomenological consequences [31–47], the potential of the memory burden could also impact the baryogenesis via leptogenesis scenarios [48], where the lepton asymmetry, generated from lepton number violating interactions, is converted to a net excess of baryons via the non-perturbative (B+L)-violating sphaleron transitions [49–55]. Normally, depending on the mass and so on the temperature of the PBHs, the evaporation of primordial black holes in the early Universe could have impacted the process of leptogenesis through both the production of heavy right-handed neutrinos, and the injection of entropy in the plasma, with production of significant population of relativistic particles [56-60]. The presence of the memory-burden effect has the potential to alter this scenario, as it could extend the lifespan of PBHs. Consequently, considering certain values of their abundance, PBHs could come to dominate the Universe during a specific period when, under normal circumstances in the semiclassical regime, they would not have been the dominant component. In this study of thermal leptogenesis, we find that the primary influence of memory-burdened PBHs lies not in their role as a non-thermal source of heavy sterile neutrinos, which could otherwise enhance the baryon asymmetry production, but rather in their entropy injection, leading to a net reduction in the baryon asymmetry. We find that the memory-burden effect notably alters the parameter space, shifting the viable region toward lower PBH masses in comparison to previous studies.

The paper is organized as follows. In Sec. II, we describe the evaporation process of memory-burden PBHs and the corresponding impact on the cosmological history of the Universe. In Sec. III, we detail the particle model of the thermal high-scale leptogenesis. In Sec. IV, we discuss the Boltzmann equations dictating the evolution of the baryon asymmetry of the Universe. In Sec. V, we

<sup>\*</sup> rcalabrese@na.infn.it

 $<sup>^{\</sup>dagger}$  m.chianese@ssmeridionale.it

<sup>&</sup>lt;sup>‡</sup> nsaviano@na.infn.it

report the main results of the present analysis. Finally, in Sec. VI we draw our conclusions.

## II. COSMOLOGY WITH MEMORY-BURDENED PRIMORDIAL BLACK HOLES

In this Section, we describe how the existence of memory-burdened PBHs can influence the evolution of the Early Universe. Thanks to the memory-burden effect, PBHs live longer allowing their domination for smaller masses compared to the standard scenario. For the sake of concreteness, we consider neutral and non-rotating PBHs with a monochromatic mass distribution with initial mass  $M_{\rm PBH}^{\rm in}$ . Moreover, we assume they originate from the collapse of density perturbations when they reenter the horizon in a radiation-dominated Universe at the temperature

$$T_{\rm in} = \frac{1}{2} \left( \frac{5}{\pi^3 g_*} \right)^{1/4} \sqrt{\frac{3\gamma_{\rm PBH} M_{\rm Pl}^3}{M_{\rm PBH}^{\rm in}}} \,, \tag{1}$$

where  $\gamma_{\rm PBH} \sim 0.2$  is the gravitational collapse factor [18, 61], and  $M_{\rm Pl} = 1.22 \times 10^{19} \,\text{GeV}$  is the Planck mass. We refer to the initial PBHs abundance in terms of the quantity

$$\beta' = \gamma_{\rm PBH}^{1/2} \left( \frac{g_*(T_{\rm in})}{106.5} \right)^{-1/4} \left( \frac{h}{0.67} \right)^{-2} \left. \frac{\rho_{\rm PBH}}{\rho_{\rm cr.}} \right|_{\rm in} \,, \quad (2)$$

where h = 0.67 [62] is related to the Hubble constant, and  $\rho_{\text{PBH}}$  and  $\rho_{\text{cr.}}|_{\text{in}} = \frac{3\gamma_{\text{PBH}}^2}{32\pi} \frac{M_{\text{Pl}}^6}{(M_{\text{PBH}}^{\text{in}})^2}$  are the PBH and critical energy densities at  $T_{\text{in}}$ , respectively.

As soon as PBHs are formed, they radiate all the particles with a mass smaller their Hawking temperature  $\mathcal{T}_{\text{PBH}}$ , defined as [27, 63]

$$\mathcal{T}_{\rm PBH} \simeq 10^{10} \left( \frac{10^3 \text{ g}}{M_{\rm PBH}} \right) \text{GeV} \,.$$
 (3)

Owing to the memory-burden effect, the evaporation process occurs into two distinct phases. The first phase is the semi-classical (Hawking-like) one, where the PBH mass evolves according to

$$\frac{\mathrm{d}\ln M_{\mathrm{PBH}}}{\mathrm{d}\alpha} = -\frac{\kappa(M_{\mathrm{PBH}})}{HM_{\mathrm{PBH}}^3}\ln(10)\,,\tag{4}$$

where  $\alpha = \log_{10} a$  with *a* being the cosmic scale factor, and  $\kappa(M_{\rm PBH}) = 416.3 M_{\rm Pl}^4/(30720\pi)$  is constant for  $M_{\rm PBH} \lesssim 10^9$  g [64]. The second "memory-burdened phase", where the PBHs are stabilized by the memory burden, kicks in when  $M_{\rm PBH} = q M_{\rm PBH}^{\rm in}$ . In the second phase, quantum effects become relevant: the information stored on the event horizon back-reacts and slows down the evaporation process by a factor proportional to a negative power of the PBH entropy

$$S_{\rm PBH} = 4\pi G M_{\rm PBH}^2 \,. \tag{5}$$

Hence, the mass evaporates according to

$$\frac{\mathrm{d}\ln M_{\mathrm{PBH}}}{\mathrm{d}\alpha} = -\frac{1}{S_{\mathrm{PBH}}^k} \frac{\kappa(M_{\mathrm{PBH}})}{HM_{\mathrm{PBH}}^3} \ln(10), \qquad (6)$$

Taking into account both the two evaporation phases, the number of  $\chi$  particles emitted by a single PBH is

$$\frac{\mathrm{d}N_{\chi}}{\mathrm{d}E\mathrm{d}t} = \frac{g_{\chi}}{2\pi} \frac{\mathcal{F}_{\chi}(E, M_{\mathrm{PBH}})}{e^{E/\mathcal{T}_{\mathrm{PBH}}} + (-1)^{2s_{\chi}}} \\ \times \begin{cases} 1 & M_{\mathrm{PBH}} > qM_{\mathrm{PBH}}^{\mathrm{in}} \\ S_{\mathrm{PBH}}^{-k} & \mathrm{otherwise} \end{cases} .$$
(7)

where  $g_{\chi}$  and  $s_{\chi}$  are the degrees of freedom and the spin of  $\chi$  particles, respectively, and  $\mathcal{F}_{\chi}$  is the grey-body factor. By integrating this equation over the energy, we obtain the emission rate

$$\Gamma_{\rm PBH\to\chi} = \int_{m_{\chi}}^{\infty} \frac{\mathrm{d}N_{\chi}}{\mathrm{d}E\mathrm{d}t} \mathrm{d}E\,. \tag{8}$$

The parameter space of the memory-burden effect is therefore completely determined by the two additional parameters q and  $k \ge 0$ . In the present analysis, we fix q = 1/2 and k = 1 and leaves the study of different memory-burden parameters to future works. We note that such choice guarantees that PBHs with mass  $M_{\rm PBH}^{\rm in} \le 1000$  g have been completely evaporated before the onset of Big Bang Nucleosynthesis [41].

To follow the evolution of the main energy density components, we need to solve the following coupled Friedmann equations [56, 58, 65]

$$\frac{\mathrm{d}\varrho_{\mathrm{rad}}}{\mathrm{d}\alpha} = -f_{\mathrm{SM}}10^{\alpha}\frac{\mathrm{d}\ln M_{\mathrm{PBH}}}{\mathrm{d}\alpha}\varrho_{\mathrm{PBH}},$$

$$\frac{\mathrm{d}\varrho_{\mathrm{PBH}}}{\mathrm{d}\alpha} = \frac{\mathrm{d}\ln M_{\mathrm{PBH}}}{\mathrm{d}\alpha}\varrho_{\mathrm{PBH}},$$

$$H^{2} = \frac{8\pi}{3M_{\mathrm{Pl}}^{2}}\left(\frac{\varrho_{\mathrm{PBH}}}{a^{3}} + \frac{\varrho_{\mathrm{rad}}}{a^{4}}\right),$$
(9)

where *H* is the Hubble parameter,  $\rho_{\rm PBH} \equiv a^3 \rho_{\rm PBH}$  and  $\rho_{\rm rad} \equiv a^4 \rho_{\rm rad}$  are the comoving energy densities of PBHs and radiation, respectively, and  $f_{\rm SM}$  is the fraction of Hawking radiation composed of SM particles. In our framework,  $f_{\rm SM} \simeq 1$  since PBHs produce a negligible amount of right-handed neutrinos. Notice that we recover the standard evolution for  $\rho_{\rm PBH} = 0$ . The comoving entropy density of the Universe evolves as

$$\frac{\mathrm{d}\mathcal{S}}{\mathrm{d}\alpha} = -\frac{f_{\mathrm{SM}}}{T(\alpha)} \frac{\mathrm{d}\ln M_{\mathrm{PBH}}}{\mathrm{d}\alpha} \varrho_{\mathrm{PBH}} \,, \tag{10}$$

which is constant for  $\rho_{\text{PBH}} \rightarrow 0$ . Lastly, the Universe's temperature T evolves according to

$$\frac{\mathrm{d}T}{\mathrm{d}\alpha} \simeq -T \left[ \ln(10) - \frac{1}{4} f_{\mathrm{SM}} 10^{\alpha} \frac{\mathrm{d}\ln M_{\mathrm{PBH}}}{\mathrm{d}\alpha} \frac{\varrho_{\mathrm{PBH}}}{\varrho_{\mathrm{rad}}} \right] .$$
(11)

## III. HIGH-SCALE LEPTOGENESIS

We now describe in detail the high-scale thermal leptogenesis scenario that lays the foundation of our analysis. Thermal leptogenesis relies on a minimal extension of the Standard Model (SM) incorporating at least two singlet Majorana neutrinos. These neutrinos interact with the SM lepton and Higgs doublets, introducing dynamics that facilitate the generation of a matter-antimatter asymmetry. A particularly compelling feature of this minimal framework is its ability to simultaneously account for the experimentally observed neutrino masses and the baryon asymmetry of the Universe.

#### Type-I seesaw

The small masses of light, active neutrinos can be naturally explained via the type-I seesaw mechanism, characterized by the following Lagrangian terms

$$\mathcal{L} \supset -Y_{\alpha i} \overline{L}_{\alpha} \tilde{\phi} N_i - \frac{1}{2} \overline{N^c}_i \hat{M}_{ij} N_j + \text{h.c.}, \qquad (12)$$

where  $\tilde{\phi} = i\sigma_2\phi^*$ ,  $\phi$  is the Higgs doublet,  $L_{\alpha} = (\nu_{\alpha}, \ell_{\alpha})$ are the lepton doublets where  $\alpha = e, \mu, \tau$ , and  $N_i$ , i = 1, 2, 3 are the Majorana right-handed neutrinos.

The active neutrino mass matrix is then determined by the type-I seesaw relation [66, 67]

$$m_{\nu} \simeq -v_{\rm EW}^2 Y \cdot \frac{1}{\hat{M}} \cdot Y^T,$$
 (13)

where  $v_{\rm EW} = 174 \,{\rm GeV}$  is the Higgs vacuum expectation value,  $\hat{M}$  represents the mass matrix of the right-handed neutrinos, and Y is the Yukawa coupling matrix from Eq. (12). We consider the mass eigenbasis of the righthanded neutrinos so that  $\hat{M} = \text{diag}(M_1, M_2, M_3)$ . The  $3 \times 3 \,\text{matrix} \, m_{\nu}$  can be diagonalized by the lepton mixing matrix  $U_{\rm PMNS}$  [68–72], parametrized by three mixing angles, together with one Dirac and two Majorana phases.

We recast Eq. (13) in terms of the physical quantities [73]

$$Y = \frac{1}{v_{\rm EW}} \sqrt{\hat{M}} \cdot R \cdot \sqrt{\hat{m}_{\nu}} \cdot U_{\rm PMNS}^{\dagger} , \qquad (14)$$

where  $\hat{m}_{\nu} = \text{diag}(m_1, m_2, m_3)$  is the diagonalized lefthanded neutrino mass matrix, and R is the Casas-Ibarra orthogonal matrix parametrized by three complex angles  $z_{12}, z_{13}, z_{23}$ . While solar and atmospheric neutrino oscillation experiments have provided precise measurements of the PMNS mixing angles and the two mass-squared differences ( $\Delta m_{\text{sol}}^2$  and  $\Delta m_{\text{atm}}^2$ ), the absolute neutrino mass scale and the mass ordering remain unknown. The ordering may be either normal ( $m_1 < m_3$ ) or inverted ( $m_1 > m_3$ ) [74–76]. In the following discussion, we focus on the normal mass ordering [74], where the heaviest ( $m_h$ ) and the lightest ( $m_l$ ) active neutrino masses correspond to  $m_3$  and  $m_1$ , respectively. Additionally, we adopt a minimal model of thermal leptogenesis with a hierarchical mass spectrum for the right-handed neutrinos, such that  $M_1 \ll M_2$ ,  $M_3$ . As shown in Ref. [77], being  $m_1 \simeq m_2$  because of  $\Delta m_{\rm sol}^2 \ll \Delta m_{\rm atm}^2$ , the only relevant angle in the *R* matrix in Eq. (14) is  $z_{13} = x + iy$  whit x, y being real parameters. This reduces the *R* matrix to

$$R = \begin{pmatrix} \cos z_{13} & 0 & \sin z_{13} \\ 0 & 1 & 0 \\ -\sin z_{13} & 0 & \cos z_{13} \end{pmatrix}.$$
 (15)

## **CP** asymmetry

CP violation in the lepton sector is pivotal in generating the observed baryon asymmetry of the Universe through leptogenesis. Specifically, a non-zero CP asymmetry arises in the out-of-equilibrium decays of heavy right-handed Majorana neutrinos. In our scenario, the dominant contribution to the baryon asymmetry originates from the decay of  $N_1$  into SM leptons and Higgs  $(N_1 \rightarrow \ell_{\alpha} \phi)$ , provided that this decay process is CP asymmetric at the one-loop level [49]. The CP asymmetry for  $N_1$  decaying into lepton flavor  $\alpha$  is

$$\epsilon_{\alpha\alpha} = \frac{\Gamma(N_1 \to \ell_\alpha \phi) - \Gamma(N_1 \to \overline{\ell_\alpha} \overline{\phi})}{\Gamma(N_1 \to \ell_\alpha \phi) + \Gamma(N_1 \to \overline{\ell_\alpha} \overline{\phi})}, \qquad (16)$$

where  $\Gamma$  indicates the decay width. At temperatures  $T \gtrsim 10^{12} \text{ GeV}$ , the charged lepton Yukawa interactions are slower than the Hubble expansion rate, rendering leptogenesis insensitive to individual lepton flavors [49]. Since we mainly focus on  $M_1 \gtrsim 10^{11} \text{ GeV}$ , we can neglect flavor effects and sum over the lepton flavors  $\alpha = e, \mu, \tau$ . Hence, in the hierarchical limit  $(M_1 \ll M_{2,3})$ , the total CP asymmetry is [78]

$$\epsilon = \sum_{\alpha} \epsilon_{\alpha\alpha} = -\frac{3}{16\pi} \sum_{j=2,3} \frac{M_1}{M_j} \frac{\mathrm{Im}(YY^{\dagger})_{1j}}{(YY^{\dagger})_{11}} \,. \tag{17}$$

In this expression, the leptonic mixing matrix  $U_{\rm PMNS}$ drops out, indicating no direct link between the low-scale Dirac CP violation and high-scale CP asymmetry. Using the simplification in Eq. (15), the CP asymmetry parameter  $\epsilon$  can be expressed in terms of only four unknown parameters  $\{x, y, m_h, M_1\}$  [58]

$$|\epsilon| = \frac{3M_1}{16\pi v_{EW}^2} \frac{|\Delta m_{\rm atm}^2|}{m_h + m_l} \frac{|\sin(2x)\sinh(2y)|}{\cosh(2y) - f\cos(2x)}, \quad (18)$$

where  $f \equiv (m_h - m_l)/(m_h + m_l)$  with  $m_h = m_3$  and  $m_l = m_1 \approx m_2$ . The parameter  $\epsilon$  governs the CP violation in  $N_1$  decays, while thermal production of  $N_1$  violates CP with the same magnitude but opposite sign. In this study, we fix  $|x| = \pi/4$  and y = 044 which are a good approximation to maximize the baryon asymmetry

produced [58, 59]. This choice allows us to make a direct comparison between the scenario of memory-burden PBHs and the standard one explored in previous studies [58].

When the initial abundance of  $N_1$  is negligible, an initial "anti-asymmetry" forms in the lepton sector. As  $N_1$  reaches the thermal equilibrium, it decays when the Universe cools to  $T = M_1$ , provided that the decay rate exceeds the Hubble expansion rate. This condition is quantified as [49, 78, 79]

$$K \equiv \frac{\tilde{m}_1}{m^*} > 1, \tag{19}$$

where

$$m^* = \frac{8\pi v_{\rm EW}^2}{M_1^2} H \left(T = M_1\right) \,, \tag{20}$$

and

$$\tilde{m}_i = \sum_{\alpha} \frac{|Y_{\alpha i}|^2 v_{\rm EW}^2}{M_i} \,, \tag{21}$$

characterize the Hubble expansion rate at  $T = M_1$  and the decay rate, respectively.

If  $N_1$  decays occur out of equilibrium, the asymmetry produced by its decay would cancel the initial antiasymmetry unless suppressed by "washout" processes. These processes include inverse decays  $(\ell \phi^{\dagger} \rightarrow N_1 \text{ and }$  $\bar{\ell}\phi \to N_1$ ) and  $2 \to 2$  scatterings  $(\ell\phi^{\dagger} \to \bar{\ell}\phi)$  mediated by off-shell  $N_1$ . For K > 1, the washout processes efficiently suppress the anti-asymmetry, allowing a net asymmetry to survive, which defines the "strong-washout" regime. Conversely, in the "weak-washout" regime  $(K < 1), N_1$ decays occur later when washout processes are weak, causing the asymmetries to largely cancel unless an additional mechanism generates the  $N_1$  population [49]. Experimental constraints on neutrino masses favor the strong-washout regime, as the weak-washout regime requires  $m_h$  to be extremely close to  $\sqrt{\Delta m_{\rm atm}^2}$ . Therefore, in this analysis, we focus on the strong-washout regime. This also ensures that any contribution from  $N_2$  and  $N_3$ is negligible, as they are washed out at  $T \ll M_1$ .

#### IV. BOLTZMANN EQUATIONS

To determine the baryon asymmetry of the Universe, we solve the Boltzmann equations for the evolution of  $N_1$ and B-L incorporating the impact of memory-burdened PBHs. Focusing on the B-L asymmetry is particularly advantageous, as it is conserved during sphaleron processes, which subsequently convert a portion of the lepton asymmetry into baryon asymmetry. Our analysis accounts for the influence of PBHs, which contribute in three significant ways: (1) they can serve as a nonthermal source of  $N_1$ , (2) they modify the evolution of the Hubble parameter, and (3) they inject entropy at the end of their evaporation, diluting the final asymmetry. In our analysis, we assume an initial abundance of  $N_1$ and B - L asymmetry to be zero. The evolution is governed by the following processes:

- Decays  $(1 \rightarrow 2)$ : The decay of  $N_1$  into leptons and Higgs bosons  $(N_1 \rightarrow \bar{\ell}\phi)$  and their CP-conjugate processes  $(N_1 \rightarrow \bar{\ell}\phi)$ . The total decay rates are proportional to the number density of  $N_1$  and the thermally averaged decay width. These decays deplete the  $N_1$  population while generating a source term for the B - L asymmetry.
- Inverse Decays  $(2 \rightarrow 1)$ : Processes such as  $\ell \phi^{\dagger} \rightarrow N_1$  replenish the  $N_1$  population but act solely as a washout mechanism for the B-L asymmetry. The inverse decay rate is proportional to the equilibrium number densities of  $N_1$  and leptons and is related to the decay rate via  $\Gamma_{N_1}^{ID} = \Gamma_{N_1} n_{N_1}^{eq} / n_{\ell}^{eq}$ , where  $n_{N_1}^{eq}, n_{\ell}^{eq}$  are their equilibrium number densities.
- Scattering Processes  $(2 \leftrightarrow 2)$ : These include  $\Delta L = 2$  scatterings mediated by  $N_1$  exchange, such as  $\ell \phi^{\dagger} \rightarrow \bar{\ell} \phi$ . While these processes do not alter the number density of  $N_1$ , they contribute to the washout of the B L asymmetry. To avoid double counting with the s-channel contributions of inverse decays, we use a subtracted form of the scattering amplitude that includes only the off-shell  $N_1$  exchange. This ensures these processes remain effective at low temperatures and are not Boltzmann suppressed.

Although  $2 \leftrightarrow 2$  scatterings involving gauge bosons and top quarks also contribute, their impact is proportional to higher-order couplings  $\propto (Y^2 Y_t^2)$ , where  $Y_t$  is the top quark Yukawa coupling, and are therefore neglected in this analysis. Similarly, three-body decay processes are numerically insignificant, contributing only around 6%, and are also omitted [80].

Hence, we need to solve the Boltzmann equations for the thermal, non-thermal (from PBHs evaporation)  $N_1$ and *B-L* comoving number densities. These equations read as

$$\frac{\mathrm{d}\mathcal{N}_{N_1}}{\mathrm{d}\alpha} = \ln(10)\frac{\Gamma_{N_1}^{\mathrm{th.}}}{H}(\mathcal{N}_{N_1}^{\mathrm{eq.}} - \mathcal{N}_{N_1}), \qquad (22)$$

$$\frac{\mathrm{d}\mathcal{N}_{N_{i}}^{\mathrm{PBH}}}{\mathrm{d}\alpha} = \frac{\mathrm{ln}(10)}{H} \left( \Gamma_{\mathrm{PBH} \to N_{1}} \mathcal{N}_{\mathrm{PBH}} + -\Gamma_{N_{i}}^{\mathrm{PBH}} \mathcal{N}_{N_{i}}^{\mathrm{PBH}} \right) , \qquad (23)$$

$$\frac{\mathrm{d}\mathcal{N}_{\mathrm{B-L}}}{\mathrm{d}\alpha} = \frac{\mathrm{ln}(10)}{H} \left[ \epsilon (\mathcal{N}_{N_{1}} - \mathcal{N}_{N_{1}}^{\mathrm{eq.}}) \Gamma_{N_{1}}^{\mathrm{th.}} + \Gamma_{N_{1}}^{\mathrm{PBH}} \mathcal{N}_{N_{\ell}}^{\mathrm{PBH}} + \left( \frac{1}{2} \frac{\mathcal{N}_{N_{1}}^{\mathrm{eq.}}}{\mathcal{N}_{\ell}^{\mathrm{eq.}}} \Gamma_{N_{1}}^{\mathrm{th.}} + \gamma \frac{a^{3}}{\mathcal{N}_{\ell}^{\mathrm{eq.}}} \right) \mathcal{N}_{\mathrm{B-L}} \right],$$

$$(24)$$

where  $\mathcal{N}_{N_1} = n_{N_1}a^3$ ,  $\mathcal{N}_{\text{PBH}} = \rho_{\text{PBH}}/M_{\text{PBH}}$ , and  $\mathcal{N}_{\text{B-L}} = n_{\text{B-L}}a^3$ . The quantity  $\Gamma_{N_1}^{\text{th.}}$  is the thermally averaged,



FIG. 1. Evolution of the main quantities as a function of the auxiliary variable  $z = M_1/T$  for the benchmark case  $M_1 = 10^{14}$  GeV,  $m_h = 0.27 \text{ eV}$ ,  $M_{\text{PBH}}^{\text{in}} = 1.0 \text{ g}$  and  $\beta' = 10^{-4}$ , in case of the semi-classical Hawking evaporation scenario (left plot) and of the memory-burden effect one (right plot). The upper panels illustrate the evolution of the thermal and non-thermal yields of the right-handed neutrinos, shown by green dot-dashed and yellow dashed lines, respectively, alongside the absolute value of the B - L yield (solid orange lines). The horizontal dashed lines correspond to the observed baryon asymmetry. The middle panels display the evolution of the PBH mass, while the lower panels depict the radiation and PBH energy densities normalized by the total energy density. The vertical dotted lines correspond (from left to right) to the onset temperatures of the memory-burden phase and the sphaleron processes.

tree-level decay rate of  $N_1$ , given by

$$\Gamma_{N_1}^{\text{th.}} = \left\langle \frac{M_1}{E_{N_1}} \right\rangle \frac{(Y^{\dagger}Y)_{11}M_1}{8\pi} \,, \tag{25}$$

where the thermal averaging factor  $\langle M_1/E_{N_1} \rangle$  can be well approximated by the ratio  $K_1(z)/K_2(z)$ , with  $K_n$  being the *n*-th order modified Bessel function of the second kind. The quantity  $\Gamma_{N_1}^{\text{PBH}}$  in Eq. (23) is the decay width corrected by an average inverse dilatation factor

$$\Gamma_{N_1}^{\text{PBH}} \approx \frac{K_1(z_{\text{PBH}})}{K_2(z_{\text{PBH}})} \frac{(Y^{\dagger}Y)_{11}M_1}{8\pi},$$
(26)

where  $z_{\text{PBH}} = M_1/\mathcal{T}_{\text{PBH}}$ . The PBH production rate  $\Gamma_{\text{PBH}\to N_1}$  of right-handed neutrinos is reported in Eq. (8). The parameter  $\gamma$  in Eq. (25) quantifies the contribution to the washout from  $\Delta L = 2$  scattering processes. Ref. [58] shows that

$$\gamma = \frac{3T^6}{4\pi^5 v_{EW}^4} \text{Tr}[m_{\nu}^{\dagger} m_{\nu}].$$
 (27)

When the comoving number density of B - L freezes in when the sphaleron processes go out of equilibrium at  $T = T_{\rm sph}$  [81]. We quantify the final baryon asymmetry as

$$|Y_{\rm B}| = \eta_{\rm sph} \frac{|\mathcal{N}_{\rm B-L}|}{\mathcal{S}} \,. \tag{28}$$

where S is the comoving entropy density of the Universe obtained by solving Eq. (10) and  $\eta_{\rm sph} = 12/37$  is the sphaleron efficiency factor [53].

In Fig.s 1 and 2, we present two benchmark solutions for the Boltzmann and Friedmann equations under consideration, in case of the semi-classical Hawking evaporation with k = 0 (left plots) and the memory-burden effect with k = 0 and q = 1/2 (right plots). In both figures, we show the the evolution of thermal and non-thermal abundances of the right-handed neutrinos and the absolute value of the B-L yield (upper panels), the evolution of the PBH mass (middle panels), and the normalized radiation and PBH energy densities (lower panels), as a function of the auxiliary variable  $z = M_1/T$ . These plots highlight that the memory-burden effect extends the PBHs lifespan, implying a two-phase evaporation. As a result, for certain values of  $\beta'$ , where they would not have otherwise dominated the Universe, they instead become the dominant component (see e.g. Fig. 1).

As can be seen from these benchmark solutions, the



FIG. 2. Similar to Fig. 1, but assuming  $M_1 = 10^{11} \text{ GeV}$ ,  $m_h = 0.05 \text{ eV}$ ,  $M_{\text{PBH}}^{\text{in}} = 10^3 \text{ g}$ , and  $\beta' = 10^{-8}$ .



FIG. 3. Upper limits on  $\beta'$  as a function of the initial PBH mass for different values of the heaviest active neutrino mass  $m_h$  (shown with different colors). The left (right) plot corresponds to  $M_1 = 10^{13} \text{ GeV}$  ( $M_1 = 10^{14} \text{ GeV}$ ) of the lightest right-handed neutrinos.

primary effect of PBHs is not their role as a non-thermal source of right-handed neutrinos that would enhance the production of the baryon asymmetry as pointed out by Ref.s [56, 58]. Indeed, in the later stages of PBH evaporation, the production of right-handed neutrinos is significantly suppressed and contributes much less to the overall baryon asymmetry. Remarkably, the dominant effect of memory-burdened PBHs stems from the entropy injection, which results in a net reduction of the baryon asymmetry. It is interesting to note that PBHs, with masses in the range [1, 1000] g ,exhibit a behavior analogous to that obtained in Ref. [59] for larger masses, in the range  $[10^6, 10^9]$  g, where the memory-burden effect was not considered.

#### V. RESULTS

We here discuss the main results of our analysis obtained by performing a comprehensive scan of the parameter space for q = 1/2 and k = 1 to investigate how the memory-burden effect alters the mixed scenario in-

volving PBHs and high-scale leptogenesis. In Fig. 3, we present the upper limits on  $\beta'$  as a function of the initial PBH mass in case of  $M_1 = 10^{13} \,\text{GeV}$  (left plot) and  $M_1 = 10^{14} \,\text{GeV}$  (right plot). Higher values of  $\beta'$  are indeed ruled out due to a high entropy injection from memory-burdened evaporation of PBHs which dilutes the baryon asymmetry of the Universe produced by highscale leptogenesis to values smaller than the observed one [62]. Hence, the regions above the lines reported in the figure represent mutual exclusion limits of memoryburdened PBHs and high-scale leptogenesis. The limits are valid for initial PBH masses restricted to the range [1, 1000] g. Indeed, lighter PBHs are excluded by the constraints on the maximum allowed value of the inflationary energy scale according to CMB measurements, while heavier PBHs are ruled out by Big Bang Nucleosynthesis for k = 1 [41]. Moreover, we find that, as in the standard scenario [59] (see also Ref.s [56, 58]), the upper limit on  $\beta'$  remarkably depends on the heaviest active neutrino mass  $m_h$ .

In Fig. 4, we instead show the regions in the  $m_h$ - $M_1$  plane where the observed baryon asymmetry can be successfully reproduced for three benchmark values of the initial PBH mass ( $M_{\rm PBH} = 1, 10, 100 \text{ g}$ ). The colored lines indicate the parameters achieving the observed baryon asymmetry in case of different assumed values for the PBH abundance  $\beta'$ . These lines can be compared with the black dashed line corresponding to the standard leptogenesis scenario without memory-burdened PBHs. Higher values of the PBH abundance  $\beta'$  result in shrinking the viable region to the left due to a higher dilution from entropy injection.

From the results of the present analysis, we observe that a future determination of the absolute neutrino mass from direct measurements in laboratory experiments and from cosmological analyses will provide a strong constraints on memory-burdened PBHs in the framework of high-scale leptogensis.

## VI. CONCLUSIONS

We have investigated the impact of a non-standard cosmology driven by the presence and evaporation of light memory-burdened PBHs on the production of the baryon asymmetry of the Universe through high-scale leptogenesis. We have found that the primary contribution of PBHs is linked to entropy injection, whereas, in the absence of the memory-burden effect, the dominant contribution arises from the additional non-thermal source term of right-handed neutrinos. In this framework, we have explored the mixed parameter space assuming q = 1/2 and k = 1, deriving the mutual exclusion limits and projecting them in the PBHs and high-scale leptogenesis parameter spaces. We have demonstrated that the memory-burden effect significantly reshapes the parameter space, pushing the viable region to lower PBH masses compared to previous studies. This has important implications for understanding baryogenesis in nonstandard cosmologies.

### ACKNOWLEDGMENTS

We thank Stefano Morisi for useful comments and discussions. RC work was supported by the Italian grant 2022JCYC9E "Red+, a low-energy characterization for low-mass Dark Matter searches with Argon" (PRIN2022) funded by the Italian Ministero dell'Universit'a e della Ricerca (MUR). MC and NS acknowledge the support by the research project TAsP (Theoretical Astroparticle Physics) funded by the Istituto Nazionale di Fisica Nucleare (INFN). The work of NS is further supported by the research grant number 2022E2J4RK "PANTHEON: Perspectives in Astroparticle and Neutrino THEory with Old and New messengers" under the program PRIN 2022 funded by the Italian Ministero dell'Università e della Ricerca (MUR).

- Ya. B. Zel'dovich and I. D. Novikov, "The Hypothesis of Cores Retarded during Expansion and the Hot Cosmological Model," Soviet Astron. AJ (Engl. Transl.), 10, 602 (1967).
- [2] Stephen Hawking, "Gravitationally collapsed objects of very low mass," Mon. Not. Roy. Astron. Soc. 152, 75 (1971).
- [3] Bernard J. Carr and S. W. Hawking, "Black holes in the early Universe," Mon. Not. Roy. Astron. Soc. 168, 399– 415 (1974).
- [4] Nicolás Bernal and Óscar Zapata, "Self-interacting Dark Matter from Primordial Black Holes," JCAP 03, 007 (2021), arXiv:2010.09725 [hep-ph].
- [5] Paolo Gondolo, Pearl Sandick, and Barmak Shams Es Haghi, "Effects of primordial black holes on dark matter models," Phys. Rev. D 102, 095018 (2020), arXiv:2009.02424 [hep-ph].

- [6] Nicolás Bernal and Óscar Zapata, "Gravitational dark matter production: primordial black holes and UV freeze-in," Phys. Lett. B 815, 136129 (2021), arXiv:2011.02510 [hep-ph].
- [7] Nicolás Bernal and Óscar Zapata, "Dark Matter in the Time of Primordial Black Holes," JCAP 03, 015 (2021), arXiv:2011.12306 [astro-ph.CO].
- [8] Andrew Cheek, Lucien Heurtier, Yuber F. Perez-Gonzalez, and Jessica Turner, "Primordial black hole evaporation and dark matter production. I. Solely Hawking radiation," Phys. Rev. D 105, 015022 (2022), arXiv:2107.00013 [hep-ph].
- [9] Andrew Cheek, Lucien Heurtier, Yuber F. Perez-Gonzalez, and Jessica Turner, "Primordial black hole evaporation and dark matter production. II. Interplay with the freeze-in or freeze-out mechanism," Phys. Rev. D 105, 015023 (2022), arXiv:2107.00016 [hep-ph].



FIG. 4. Contour lines in the plane  $m_h-M_1$  for which the observed baryon asymmetry can be reproduced in case of different values for the PBH abundance  $\beta'$  (shown with different colors). The lines delimit to the right the regions where the baryon asymmetry is higher than the observed value. The panels correspond to different assumptions for the initial PBH mass.

- [10] Rome Samanta and Federico R. Urban, "Testing super heavy dark matter from primordial black holes with gravitational waves," JCAP 06, 017 (2022), arXiv:2112.04836 [hep-ph].
- [11] Nicolás Bernal, Fazlollah Hajkarim, and Yong Xu, "Axion Dark Matter in the Time of Primordial Black Holes," Phys. Rev. D 104, 075007 (2021), arXiv:2107.13575 [hepph].
- [12] Nicolás Bernal, Yuber F. Perez-Gonzalez, Yong Xu, and Óscar Zapata, "ALP dark matter in a primordial black hole dominated universe," Phys. Rev. D 104, 123536 (2021), arXiv:2110.04312 [hep-ph].
- [13] Pearl Sandick, Barmak Shams Es Haghi, and Kuver Sinha, "Asymmetric reheating by primordial black holes," Phys. Rev. D 104, 083523 (2021), arXiv:2108.08329 [astro-ph.CO].
- [14] Nicolás Bernal, Yuber F. Perez-Gonzalez, and Yong Xu, "Superradiant production of heavy dark matter from primordial black holes," Phys. Rev. D 106, 015020 (2022), arXiv:2205.11522 [hep-ph].
- [15] Andrew Cheek, Lucien Heurtier, Yuber F. Perez-Gonzalez, and Jessica Turner, "Evaporation of primordial black holes in the early Universe: Mass and spin distributions," Phys. Rev. D 108, 015005 (2023), arXiv:2212.03878 [hep-ph].

- [16] Thomas C. Gehrman, Barmak Shams Es Haghi, Kuver Sinha, and Tao Xu, "Recycled dark matter," JCAP 03, 044 (2024), arXiv:2310.08526 [hep-ph].
- [17] Enrico Bertuzzo, Yuber F. Perez-Gonzalez, Gabriel M. Salla, and Renata Zukanovich Funchal, "Gravitationally produced dark matter and primordial black holes," JCAP 09, 059 (2024), arXiv:2405.17611 [hep-ph].
- [18] Bernard Carr, Florian Kuhnel, and Marit Sandstad, "Primordial Black Holes as Dark Matter," Phys. Rev. D 94, 083504 (2016), arXiv:1607.06077 [astro-ph.CO].
- [19] Anne M. Green and Bradley J. Kavanagh, "Primordial Black Holes as a dark matter candidate," J. Phys. G 48, 043001 (2021), arXiv:2007.10722 [astro-ph.CO].
- [20] Bernard Carr, Kazunori Kohri, Yuuiti Sendouda, and Jun'ichi Yokoyama, "Constraints on primordial black holes," Rept. Prog. Phys. 84, 116902 (2021), arXiv:2002.12778 [astro-ph.CO].
- [21] Bernard Carr and Florian Kuhnel, "Primordial black holes as dark matter candidates," SciPost Phys. Lect. Notes 48, 1 (2022), arXiv:2110.02821 [astro-ph.CO].
- [22] Theodoros Papanikolaou, Vincent Vennin, and David Langlois, "Gravitational waves from a universe filled with primordial black holes," JCAP 03, 053 (2021), arXiv:2010.11573 [astro-ph.CO].
- [23] Guillem Domènech, Chunshan Lin, and Misao Sasaki, "Gravitational wave constraints on the primordial black

hole dominated early universe," JCAP **04**, 062 (2021), [Erratum: JCAP **11**, E01 (2021)], arXiv:2012.08151 [gr-qc].

- [24] Theodoros Papanikolaou, "Gravitational waves induced from primordial black hole fluctuations: the effect of an extended mass function," JCAP 10, 089 (2022), arXiv:2207.11041 [astro-ph.CO].
- [25] Aurora Ireland, Stefano Profumo, and Jordan Scharnhorst, "Primordial gravitational waves from black hole evaporation in standard and nonstandard cosmologies," Phys. Rev. D 107, 104021 (2023), arXiv:2302.10188 [grqc].
- [26] Guillem Domènech and Jan Tränkle, "From formation to evaporation: Induced gravitational wave probes of the primordial black hole reheating scenario," (2024), arXiv:2409.12125 [gr-qc].
- [27] S. W. Hawking, "Particle Creation by Black Holes," Commun. Math. Phys. 43, 199–220 (1975), [Erratum: Commun.Math.Phys. 46, 206 (1976)].
- [28] John Preskill, "Do black holes destroy information?" in International Symposium on Black holes, Membranes, Wormholes and Superstrings (1992) arXiv:hepth/9209058.
- [29] Gia Dvali, "A Microscopic Model of Holography: Survival by the Burden of Memory," (2018), arXiv:1810.02336 [hep-th].
- [30] Gia Dvali, Juan Sebastián Valbuena-Bermúdez, and Michael Zantedeschi, "Memory burden effect in black holes and solitons: Implications for PBH," Phys. Rev. D 110, 056029 (2024), arXiv:2405.13117 [hep-th].
- [31] Shyam Balaji, Guillem Domènech, Gabriele Franciolini, Alexander Ganz, and Jan Tränkle, "Probing modified Hawking evaporation with gravitational waves from the primordial black hole dominated universe," (2024), arXiv:2403.14309 [gr-qc].
- [32] Basabendu Barman, Md Riajul Haque, and Óscar Zapata, "Gravitational wave signatures of cogenesis from a burdened PBH," JCAP 09, 020 (2024), arXiv:2405.15858 [astro-ph.CO].
- [33] Nilanjandev Bhaumik, Md Riajul Haque, Rajeev Kumar Jain, and Marek Lewicki, "Memory burden effect mimics reheating signatures on SGWB from ultra-low mass PBH domination," (2024), arXiv:2409.04436 [astro-ph.CO].
- [34] Basabendu Barman, Kousik Loho, and Óscar Zapata, "Constraining burdened PBHs with gravitational waves," (2024), arXiv:2409.05953 [gr-qc].
- [35] Kazunori Kohri, Takahiro Terada, and Tsutomu T. Yanagida, "Induced Gravitational Waves probing Primordial Black Hole Dark Matter with Memory Burden," (2024), arXiv:2409.06365 [astro-ph.CO].
- [36] Yang Jiang, Chen Yuan, Chong-Zhi Li, and Qing-Guo Huang, "Constraints on the Primordial Black Hole Abundance through Scalar-Induced Gravitational Waves from Advanced LIGO and Virgo's First Three Observing Runs," (2024), arXiv:2409.07976 [astro-ph.CO].
- [37] Gia Dvali, Lukas Eisemann, Marco Michel, and Sebastian Zell, "Black hole metamorphosis and stabilization by memory burden," Phys. Rev. D 102, 103523 (2020), arXiv:2006.00011 [hep-th].
- [38] Gia Dvali, Florian Kühnel, and Michael Zantedeschi, "Primordial black holes from confinement," Phys. Rev. D 104, 123507 (2021), arXiv:2108.09471 [hep-ph].

- [39] Ana Alexandre, Gia Dvali, and Emmanouil Koutsangelas, "New mass window for primordial black holes as dark matter from the memory burden effect," Phys. Rev. D 110, 036004 (2024), arXiv:2402.14069 [hep-ph].
- [40] Valentin Thoss, Andreas Burkert, and Kazunori Kohri, "Breakdown of hawking evaporation opens new mass window for primordial black holes as dark matter candidate," Mon. Not. Roy. Astron. Soc. 532, 451–459 (2024), arXiv:2402.17823 [astro-ph.CO].
- [41] Md Riajul Haque, Suvashis Maity, Debaprasad Maity, and Yann Mambrini, "Quantum effects on the evaporation of PBHs: contributions to dark matter," JCAP 07, 002 (2024), arXiv:2404.16815 [hep-ph].
- [42] Marco Chianese, Andrea Boccia, Fabio Iocco, Gennaro Miele, and Ninetta Saviano, "The light burden of memory: constraining primordial black holes with high-energy neutrinos," (2024), arXiv:2410.07604 [astro-ph.HE].
- [43] Michael Zantedeschi and Luca Visinelli, "Memory-Burdened Primordial Black Holes as Astrophysical Particle Accelerators," (2024), arXiv:2410.07037 [astroph.HE].
- [44] Basabendu Barman, Kousik Loho, and Óscar Zapata, "Asymmetries from a charged memory-burdened PBH," (2024), arXiv:2412.13254 [hep-ph].
- [45] Debasish Borah and Nayan Das, "Successful cogenesis of baryon and dark matter from memory-burdened PBH," (2024), arXiv:2410.16403 [hep-ph].
- [46] Peter Athron, Marco Chianese, Satyabrata Datta, Rome Samanta, and Ninetta Saviano, "Impact of memoryburdened black holes on primordial gravitational waves in light of Pulsar Timing Array," (2024), arXiv:2411.19286 [astro-ph.CO].
- [47] Disha Bandyopadhyay, Debasish Borah, and Nayan Das, "Axion misalignment with memory-burdened PBH," (2025), arXiv:2501.04076 [hep-ph].
- [48] Masataka Fukugita and Tsutomu Yanagida, "Barygenesis without grand unification," Physics Letters B 174, 45–47 (1986).
- [49] Sacha Davidson, Enrico Nardi, and Yosef Nir, "Leptogenesis," Phys. Rept. 466, 105–177 (2008), arXiv:0802.2962 [hep-ph].
- [50] Chee Sheng Fong, Enrico Nardi, and Antonio Riotto, "Leptogenesis in the Universe," Adv. High Energy Phys. 2012, 158303 (2012), arXiv:1301.3062 [hep-ph].
- [51] W. Buchmuller, R. D. Peccei, and T. Yanagida, "Leptogenesis as the origin of matter," Ann. Rev. Nucl. Part. Sci. 55, 311–355 (2005), arXiv:hep-ph/0502169.
- [52] Pasquale Di Bari, "An introduction to leptogenesis and neutrino properties," Contemp. Phys. 53, 315–338 (2012), arXiv:1206.3168 [hep-ph].
- [53] Jeffrey A. Harvey and Michael S. Turner, "Cosmological baryon and lepton number in the presence of electroweak fermion number violation," Phys. Rev. D 42, 3344–3349 (1990).
- [54] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, "On the Anomalous Electroweak Baryon Number Nonconservation in the Early Universe," Phys. Lett. B 155, 36 (1985).
- [55] S. Yu. Khlebnikov and M. E. Shaposhnikov, "The Statistical Theory of Anomalous Fermion Number Nonconservation," Nucl. Phys. B 308, 885–912 (1988).
- [56] Yuber F. Perez-Gonzalez and Jessica Turner, "Assessing the tension between a black hole dominated early

- [57] Suruj Jyoti Das, Devabrat Mahanta, and Debasish Borah, "Low scale leptogenesis and dark matter in the presence of primordial black holes," JCAP 11, 019 (2021), arXiv:2104.14496 [hep-ph].
- [58] Nicolás Bernal, Chee Sheng Fong, Yuber F. Perez-Gonzalez, and Jessica Turner, "Rescuing high-scale leptogenesis using primordial black holes," Phys. Rev. D 106, 035019 (2022), arXiv:2203.08823 [hep-ph].
- [59] Roberta Calabrese, Marco Chianese, Jacob Gunn, Gennaro Miele, Stefano Morisi, and Ninetta Saviano, "Limits on light primordial black holes from high-scale leptogenesis," Phys. Rev. D 107, 123537 (2023), arXiv:2305.13369 [hep-ph].
- [60] Roberta Calabrese, Marco Chianese, Jacob Gunn, Gennaro Miele, Stefano Morisi, and Ninetta Saviano, "Impact of primordial black holes on heavy neutral leptons searches in the framework of resonant leptogenesis," Phys. Rev. D 109, 103001 (2024), arXiv:2311.13276 [hepph].
- [61] Bernard J. Carr, "The Primordial black hole mass spectrum," Astrophys. J. 201, 1–19 (1975).
- [62] Y. Akrami *et al.* (Planck), "Planck 2018 results. X. Constraints on inflation," Astron. Astrophys. **641**, A10 (2020), arXiv:1807.06211 [astro-ph.CO].
- [63] S. W. Hawking, "Black hole explosions," Nature 248, 30–31 (1974).
- [64] Satyabrata Datta, Ambar Ghosal, and Rome Samanta, "Baryogenesis from ultralight primordial black holes and strong gravitational waves from cosmic strings," JCAP 08, 021 (2021), arXiv:2012.14981 [hep-ph].
- [65] Cecilia Lunardini and Yuber F. Perez-Gonzalez, "Dirac and Majorana neutrino signatures of primordial black holes," JCAP 08, 014 (2020), arXiv:1910.07864 [hep-ph].
- [66] Tsutomu Yanagida, "Horizontal Symmetry and Masses of Neutrinos," Prog. Theor. Phys. 64, 1103 (1980).
- [67] Rabindra N. Mohapatra and Goran Senjanović, "Neutrino masses and mixings in gauge models with spontaneous parity violation," Phys. Rev. D 23, 165–180 (1981).
- [68] Bruno Pontecorvo, "Mesonium and antimesonium," Zhur. Eksptl'. i Teoret. Fiz. 33 (1957).
- [69] Bruno Pontecorvo, "Inverse beta processes and nonconservation of lepton charge," Zh. Eksp. Teor. Fiz 34, 247 (1957).

- [70] Ziro Maki, Masami Nakagawa, and Shoichi Sakata, "Remarks on the unified model of elementary particles," Prog. Theor. Phys. 28, 870–880 (1962).
- [71] B. Pontecorvo, "Neutrino Experiments and the Problem of Conservation of Leptonic Charge," Zh. Eksp. Teor. Fiz. 53, 1717–1725 (1967).
- [72] V. N. Gribov and B. Pontecorvo, "Neutrino astronomy and lepton charge," Phys. Lett. B 28, 493 (1969).
- [73] J. A. Casas and A. Ibarra, "Oscillating neutrinos and  $\mu \rightarrow e, \gamma$ ," Nucl. Phys. B **618**, 171–204 (2001), arXiv:hep-ph/0103065.
- [74] Francesco Capozzi, Eleonora Di Valentino, Eligio Lisi, Antonio Marrone, Alessandro Melchiorri, and Antonio Palazzo, "Unfinished fabric of the three neutrino paradigm," Phys. Rev. D 104, 083031 (2021), arXiv:2107.00532 [hep-ph].
- [75] Ivan Esteban, M. C. Gonzalez-Garcia, Michele Maltoni, Thomas Schwetz, and Albert Zhou, "The fate of hints: updated global analysis of three-flavor neutrino oscillations," JHEP 09, 178 (2020), arXiv:2007.14792 [hep-ph].
- [76] P. F. de Salas, D. V. Forero, S. Gariazzo, P. Martínez-Miravé, O. Mena, C. A. Ternes, M. Tórtola, and J. W. F. Valle, "2020 global reassessment of the neutrino oscillation picture," JHEP 02, 071 (2021), arXiv:2006.11237 [hep-ph].
- [77] Thomas Hambye, Yin Lin, Alessio Notari, Michele Papucci, and Alessandro Strumia, "Constraints on neutrino masses from leptogenesis models," Nucl. Phys. B 695, 169–191 (2004), arXiv:hep-ph/0312203.
- [78] Alessandro Strumia, "Baryogenesis via leptogenesis," in Les Houches Summer School on Theoretical Physics: Session 84: Particle Physics Beyond the Standard Model (2006) pp. 655–680, arXiv:hep-ph/0608347.
- [79] W. Buchmuller, P. Di Bari, and M. Plumacher, "Leptogenesis for pedestrians," Annals Phys. **315**, 305–351 (2005), arXiv:hep-ph/0401240.
- [80] Enrico Nardi, Juan Racker, and Esteban Roulet, "CP violation in scatterings, three body processes and the Boltzmann equations for leptogenesis," JHEP 09, 090 (2007), arXiv:0707.0378 [hep-ph].
- [81] Michela D'Onofrio, Kari Rummukainen, and Anders Tranberg, "Sphaleron Rate in the Minimal Standard Model," Phys. Rev. Lett. **113**, 141602 (2014), arXiv:1404.3565 [hep-ph].