Sequential Portfolio Selection under Latent Side Information-Dependence Structure: Optimality and Universal Learning Algorithms

Duy Khanh Lam* Scuola Normale Superiore

(Working paper)

January 22, 2025

Abstract

This paper investigates the investment problem of constructing an optimal no-short sequential portfolio strategy in a market with a latent dependence structure between asset prices and partly unobservable side information, which is often high-dimensional. The results demonstrate that a dynamic strategy, which forms a portfolio based on perfect knowledge of the dependence structure and full market information over time, may not grow at a higher rate infinitely often than a constant strategy, which remains invariant over time. Specifically, if the market is stationary, implying that the dependence structure is statistically stable, the growth rate of an optimal dynamic strategy, utilizing the maximum capacity of the entire market information, almost surely decays over time into an equilibrium state, asymptotically converging to the growth rate of a constant strategy.

Technically, this work reassesses the common belief that a constant strategy only attains the optimal limiting growth rate of dynamic strategies when the market process is identically and independently distributed. By analyzing the dynamic log-optimal portfolio strategy as the optimal benchmark in a stationary market with side information, we show that a random optimal constant strategy almost surely exists, even when a limiting growth rate for the dynamic strategy does not. Consequently, two approaches to learning algorithms for portfolio construction are discussed, demonstrating the safety of removing side information from the learning process while still guaranteeing an asymptotic growth rate comparable to that of the optimal dynamic strategy.

Keywords: Online Learning, Dynamic Strategy, Universality, Optimal Growth, Side Information, Latent Dependence Structure.

^{*}Some of the main results of this paper were presented in a poster titled "*Universal Investment Strategies by Online Learning under Latent Information-Dependence Structure*" at the workshop on "*Learning and Inference from Structured Data: Universality, Correlations, and Beyond*" held at the Abdus Salam International Centre for Theoretical Physics in Italy, from July 3 to 7, 2023. I would also like to express my gratitude to my advisor, Giulio Bottazzi, and to Daniele Giachini at the Scuola Superiore Sant'Anna for their valuable discussions.

1 Introduction

In sequential investment, the task of collecting and analyzing data on market events that correlate with future assets' prices, to make well-informed decisions at each investment period, is undoubtedly one of the most significant concerns. In practice, a market is often represented by latent dynamics, designed to capture the hidden dependence structures among assets' prices, their historical data, and other sources of side market information, such as energy prices, news, and other potentially correlated features. However, investors consistently face three fundamental challenges of uncertainty: the incompleteness of observable side market information, the computational difficulty associated with high-dimensional data, and the unknown nature of the market's dependence structure. The first challenge stems from the fact that valuable side market information is often costly and usually available only to a small subset of investors at any point in time, such as insiders, which makes most of the side information unobservable. The second challenge lies in the uncertainty and potentially large volume of market features correlated with future assets' prices, making estimation from collected data computationally expensive in terms of both time and processing power, which complicates high-frequency decision-making. Lastly, the third challenge arises from the hidden dependence structure of the market, where the statistical properties of its dynamics are unknown, leaving open the possibility that the process could either be identically and independently distributed (i.i.d.) or exhibit an undetermined memory length.

Motivation and overall result. In this paper, we explore the problem of constructing sequential no-short portfolios and address the three aforementioned challenges based on a general model of a market process with partially observable side market information. We demonstrate that the best dynamic strategy, defined as a sequence of portfolios determined using the complete history of market observations over time, including even the unobservable features, cannot infinitely often outperform an optimal constant strategy, represented as a unique and random portfolio, in terms of both growth rate and expected growth rate as time progresses to infinity. This finding implies that the constant strategy does not require knowledge of periodic side market information, yet it remains asymptotically optimal. Such a result contradicts conventional perspectives in the fields of information theory, finance, and potentially learning theory, which suggest that possessing full knowledge of market information and dependence structures should unequivocally outperform ignorance. Based on our findings, a class of learning algorithms that relies solely on past assets' prices, without any prior knowledge of latent dependence structures or side information, is shown to be optimal and comparable to another innovative learning algorithm that is more sophisticated and leverages complete market knowledge. This analysis underscores the safety of omitting side feature data in the learning process while still ensuring optimality in the asymptotic growth rate.

Paper organization and main novelties. We briefly describe the main novel results, which contradict popular conjectures and the status quo, along with the analysis techniques and literature references for comparison:

Section 2: The assets' returns and all side information, including unobservable features, are modeled as a pairwise process to capture various dependence structures. Then, the logoptimal strategy is introduced as the optimal benchmark among all strategies. Moreover, it is defined without requiring the integrability assumption for the variables, broadening the scope of market investigations. Note that integrability is a common assumption in the literature, ensuring that expectations are well-defined.

- *Section 3*: The optimality properties of the log-optimal strategy are established as a generalization of [Algoet and Cover \(1988](#page-32-0)). The analysis is performed under a measure transformation, which bypasses the required integrability by decomposing the portfolio's return into an integrable part. The properties demonstrate the advantages of utilizing full side market information and guarantee the optimal expected growth rate and asymptotic growth rate for the log-optimal strategy over other competitive strategies using less information.
- *Section* 4.1: This section reassesses the status quo in contemporary literature. Since the works of [Kelly \(1956](#page-32-1)); [Breiman](#page-32-2) [\(1960](#page-32-2), [1961](#page-32-3)), it has been widely accepted that a constant strategy of a single portfolio only achieves the optimal asymptotic growth rate under an i.i.d. process of assets' returns without side information, as reiterated in many later articles such as [Cover](#page-32-4) [\(1991](#page-32-4)); [Morvai](#page-32-5) [\(1991](#page-32-5)); [Algoet \(1992\)](#page-32-6); [Morvai](#page-32-7) [\(1992](#page-32-7)); [Cover \(1996](#page-32-8)); [Ordentlich and Cover](#page-32-9) [\(1996](#page-32-9)); [Cover and Ordentlich \(1996](#page-32-10)); [Gyorfi et al.](#page-32-11) [\(2006](#page-32-11)); [Cover and Thomas](#page-32-12) [\(2006](#page-32-12)); [Cesa-Bianchi and Lugosi \(2006](#page-32-13)); [Bhatt et al. \(2023\)](#page-32-14), etc. Additionally, after demonstrating the optimal limiting growth rate of the logoptimal strategy under stationary and ergodic processes without side information in [Algoet and Cover \(1988](#page-32-0)); the author conjectured that no constant strategy can match the optimal growth rate of a dynamic strategy under the same process in [Algoet](#page-32-6) $(1992)^{1}$ $(1992)^{1}$ $(1992)^{1}$ $(1992)^{1}$. However, we show that a random optimal constant strategy can almost surely asymptotically approach the optimal growth rate of the log-optimal strategy under a stationary market process with side information, becoming non-random if the process is ergodic.
- *Section* 4.2: This section establishes two equalities between the optimal limits of the growth rate and expected growth rate of the log-optimal strategy utilizing the infinitely large amount of market information, and those of the random optimal strategy, which is not conditioned on side information. It implies that under the stationarity condition, the advantages of utilizing the maximum capacity of market information will almost surely decay over time into an equilibrium state.
- *Section* 4.3: The general process is reduced to capture several models of market phenomena, including a market with a non-decaying impact of past information on future market, a market with finite-order memory, and one where future assets' returns depend only on side information but not on the past. Generally, if the latent dependence structure remains unchanged over time, a random optimal constant strategy could exist.
- *Section* 5: Under an unknown distribution, making the optimal strategies undetermined in advance, two approaches for universal learning algorithms to construct strategies are discussed. The first approach requires learning only past assets' returns to asymptotically attain the same rate as the random optimal constant strategy. This generalizes the optimality of the Universal portfolio by [Cover \(1991](#page-32-4)) beyond an i.i.d. process without side information. Alternatively, the second approach requires learning the entire past market information and dependence structure to asymptotically attain the same rate as the log-optimal strategy. This is based on approaches proposed by [Algoet \(1992\)](#page-32-6) and [Gyorfi et al. \(2006](#page-32-11)), but it can learn less information while still guaranteeing optimality.

¹See the last paragraph in Remark 4, pages 929-930, where the author also restates that a constant strategy attains the optimal asymptotic growth rate only under an i.i.d. process without side market information.

2 Model settings and formalizations

We consider modeling an investment scenario where an investor sequentially constructs no-short portfolios in a multi-asset market without frictions. Over time, the investor observes market information, including historical asset prices and side information such as oil prices, news, and events from other markets. While future asset prices remain inherently unpredictable, the revealed information can be utilized in various ways to develop investment strategies aimed at achieving favorable cumulative wealth across diverse market possibilities. However, not all side information correlated with asset prices is observable, and its relevance varies. Additionally, collecting extensive side information can be prohibitively expensive. Thus, an ideal model should be universal, capable of accommodating diverse dependence structures and incorporating unobservable side information, even when such information is infinite in scope. This universality naturally introduces the challenge of high dimensionality. Furthermore, the model should facilitate the formulation of an optimal strategy from the set of all accessible strategies, serving as a benchmark despite the uncertainties of future market behavior and incomplete information. Such a desired modeling framework, which addresses these challenges, is presented as follows.

Let us consider sequential portfolio construction for investment in a stock market with $m \geq 2$ risky assets, whose prices are causally affected by $k \geq 1$ features of observable and unobservable side information, over discrete time periods $n \in \mathbb{N}_+$. Assume that the assets' returns and side information are realized with respect to a pairwise random process $\{X_n, Y_n\}_{n=1}^{\infty}$ that is jointly defined on the canonical probability space $(\Omega, \mathbb{F}, \mathbb{P})$, i.e., $(X_n, Y_n)(\omega) = (X_n(\omega), Y_n(\omega))$ for all *n*, where $\omega \in \Omega$. Specifically, the positive real-valued vector $X_n := (X_{n,1},...,X_{n,m}) \in \mathbb{R}^m_{++}$ represents the positive returns of the assets at time *n*, which is defined as the ratio of the assets' prices at time *n* to those at the previous time *n* − 1; while the real-valued random vector $Y_n := (Y_{n,1},...,X_{n,k}) \in \mathbb{R}^k$ represents the side information related to the market's features at time *n*. We further denote $X_1^n :=$ $\{X_1,...,X_n\}$ and $Y_1^n := \{Y_1,...,Y_n\}$ as the finite sequences of asset returns and side information variables, respectively, from 1 to *n*. It should be remarked that the side information variables Y_n could take values in some Polish spaces rather than \mathbb{R}^k as the default setting.

At each time *n*, the sub- σ -field $\mathscr{F}_n := \sigma(X_1^n, Y_1^n) \subseteq \mathbb{F}$ embodies all past information up to the present of the random pairs (X_n, Y_n) , up to the limiting σ -field $\mathscr{F}_\infty := \sigma\left(\cup_{n=1}^\infty \mathscr{F}_n\right) \subseteq \mathbb{F}$, which encompasses the infinite past as $n \to \infty$ and is the smallest σ -field containing the largest possible information the investor can know. Consequently, it is possible to define a further sub- σ -field $\mathscr{F}_n^X = \sigma(X_1^{n-1}, Y_1^n) \subset \mathscr{F}_{n+1}$, which includes only the side information just before the realization of the assets' returns at each time *n*. Thus, the corresponding limiting information field generated by \mathscr{F}_{n}^{X} , as an infinite union, is denoted by the σ -field $\mathscr{F}_{\infty}^{X} := \sigma\left(\cup_{n=1}^{\infty} \mathscr{F}_{n}^{X}\right) \subseteq \mathbb{F}$. Noting that although we could model the side information events that occur between two consecutive X_n and X_{n+1} as Y_n instead of Y_{n+1} , which would make the market process simpler, it seems difficult to model a scenario where the present assets' returns depend solely on new market events but are independent of past ones. Hence, our modeling framework allows us to conveniently capture various market behaviors, as discussed in the section on investigating the stationary market.

For each *n*, given the σ -field \mathscr{F}_n^X , a portfolio $b_n : \mathbb{R}_{++}^{m \times (n-1)} \times \mathbb{R}^{k \times n} \to \mathscr{B}^m$, where the simplex $\mathscr{B}^m := \left\{ \beta := (\beta_1, ..., \beta_m) \in \mathbb{R}_{++}^m : \sum_{j=1}^m \beta_j = 1 \right\}$ denotes the domain of all no-short-constrained portfolios. This means each portfolio is causally selected based on the realizations of X_1^{n-1} and Y_1^n , but not X_1^n , in order to avoid the unrealistic and impossible case of the future assets' returns being always predicted perfectly. We subsequently denote the corresponding *strategy*, formed from the

portfolios b_n for all *n*, as the infinite sequence $(b_n) := \{b_n\}_{n=1}^{\infty}$ $\sum_{n=1}^{\infty}$. Accordingly, if a strategy uses only a fixed portfolio over time, such that $b_n := b$ for all *n*, it is termed a *constant strategy* and denoted as *b* without the time index. Additionally, the return of a portfolio *b* with respect to assets' returns X_n is denoted by $\langle b, X_n \rangle$, where $\langle \cdot, \cdot \rangle$ represents the scalar product of two vectors.

With the above settings, let the initial capital $S_0(b_0) =: S_0 = 1$ by convention for any strategy, and assume that the portfolios are constructed without commission fees for arbitrary fractions. The *cumulative wealth* and its corresponding exponential average *growth rate* after *n* periods of investment, yielded by a self-financed strategy (b_n) , are respectively defined as follows:

$$
S_n(b_n) := \prod_{i=1}^n \langle b_i, X_i \rangle \text{ and } W_n(b_n) := \frac{1}{n} \log S_n(b_n) = \frac{1}{n} \sum_{i=1}^n \log \langle b_i, X_i \rangle, \forall n,
$$

which shorthand for $S_n(\lbrace b_i \rbrace_{i=1}^n)$ $\binom{n}{i=1}$, X_1^n and W_n ($\{b_i\}_{i=1}^n$ $\sum_{i=1}^{n}$, *X*ⁿ), respectively. It is worth noting that the assumption of positive assets' returns is not only realistic but also serves to exclude the case where a strategy's cumulative wealth is depleted to zero, which would halt the investment.

2.1 The benchmark strategy and normalized assets' returns

In general, to investigate the intrinsic growth rate of a market, it is sufficient to analyze a single representative strategy that is guaranteed not to yield a lower growth rate than any other strategy as time evolves. Therefore, any potentially competitive strategy must be compared solely with this representative strategy for performance benchmarking. The so-called *log-optimal strategy*, as defined in Definition [1,](#page-4-0) provides such a strategy for the purpose of our investigation. However, to better understand the nature of the market under consideration, it should be noted that the benchmarking role of the log-optimal strategy may no longer hold when market frictions, such as commission fees, are introduced.

Definition 1. Given the filtration $\{\mathcal{F}_n^X\}_{n=1}^{\infty}$, consider the portfolios b_n^* over periods *n* that satisfy the following condition:

$$
\mathbb{E}\big(\log\langle b,X_n\rangle-\log\langle b_n^*,X_n\rangle|\mathscr{F}_n^X\big)\leq 0, \forall b\in\mathscr{B}^m, \forall n.
$$

Any sequence of such b_n^* defines a log-optimal strategy and is denoted as (b_n^*) henceforth.

We stress that Definition [1](#page-4-0) implies the log-optimal portfolios are ones maximizing the conditional expectation of the differences between logarithmic returns of two portfolios, given all past market assets' returns and side information up to the present period, regardless of their observability. This way of formalization does not necessitate the finite existence of individually conditional expected values of each portfolio's logarithmic returns but their well-defined conditional expected difference. Therefore, it allows us to broaden our investigation of a stochastic market to the general case of infinite or even ill-defined expectations by bypassing the required integrability of measurable variables, which will be discussed in the next sections.

Remark. In the literature, the integrability of X_n , i.e., the expected values $\mathbb{E}(\log \langle b, X_n \rangle)$, is often assumed to finitely exist in some form as a minimal requirement for analysis in several papers, such as [Morvai \(1991](#page-32-5)); [Cuchiero et al. \(2019\)](#page-32-15); [Gyorfi et al. \(2006](#page-32-11)), etc. In the paper [Algoet and Cover](#page-32-0) [\(1988](#page-32-0)), the definition of the log-optimal strategy is defined as in Definition [1;](#page-4-0) however, all subsequent analyses in that paper proceed under the assumption that integrability is satisfied, leading to its main theorems being stated for finite expectations. In fact, the technique we discuss next is also utilized in that paper, but only to investigate the continuity of the maximal expected logarithmic return of a portfolio and the existence of the maximizer. In contrast, we apply the same technique but use it to bypass the required integrability by bounding the variables, thus expanding the market scenarios in which expectations are not well-defined.

Variable transformation. In order to guarantee the well-definedness of the conditional expected difference between the logarithmic returns of portfolios, we show that any individual conditional expectation can be decomposed into the sum of a well-defined expected value and a common redundant part, which is then ignorable by subtraction. Specifically, by adapting the following approach in a manner similar to that originally used by [Algoet and Cover \(1988](#page-32-0)) to demonstrate the existence of log-optimal portfolios, any expected logarithmic portfolio return can be decomposed by performing a transformation of the probability measure through a change of variables.

Let us consider the equally allocated portfolio $\hat{b} := (1/m, ..., 1/m)$ as a reference portfolio and the variable of assets' returns *X*. We subsequently define the variable *U* of *normalized assets' returns* as the following scaling function of *X* with parameter \hat{b} :

$$
U := u(X) := X/\langle \hat{b}, X \rangle \in \mathcal{U}, \qquad (2.1)
$$

where $\mathscr U$ denotes the corresponding set of all normalized assets' returns as:

$$
\mathscr{U} := \{ u := (u_1, ..., u_m) \in \mathbb{R}_{++}^m : \langle \hat{b}, u \rangle = 1 \}.
$$
 (2.2)

Thus, for any distribution *P* of the random variable *X*, the corresponding distribution Q^P of the normalized random variable *U* is an image measure of *P*, defined as follows:

$$
Q^{P}(U \in A) = P(X : u(X) \in A), \forall A \subseteq \mathscr{U}.
$$

As a result, while the support of the original distribution P ranges over the whole space \mathbb{R}^m_+ , the support of the transformed distribution Q^P is constrained to a bounded region. This fact, along with other critical properties relevant to our analysis, which result directly from the transformation of the probability measure for decomposition, are summarized in Lemma [1.](#page-5-0)

Lemma 1. Consider the reference portfolio \hat{b} and the variable of normalized assets' returns *U*, defined in terms of the variable *X* according to [\(2.1\)](#page-5-1) and [\(2.2\)](#page-5-2). We have:

(a) Given a process of assets' returns $\{X_n\}_{n=1}^{\infty}$, the difference in growth rates between two generic strategies (\bar{b}_n) and (\tilde{b}_n) is always identical to the corresponding difference associ*ated with the normalized assets' returns* $U_n = u(X_n)$ *over time, i.e.,*

$$
W_n(\bar{b}_n)-W_n(\tilde{b}_n)=\frac{1}{n}\sum_{i=1}^n\log\langle\bar{b}_n,U_n\rangle-\frac{1}{n}\sum_{i=1}^n\log\langle\tilde{b}_n,U_n\rangle, \ \forall n.
$$

(b) *For any distribution, the corresponding expected difference between the logarithmic returns of two portfolios remains unchanged through the change of variable, as:*

$$
\mathbb{E} \big(\log \langle \bar{b}, X \rangle - \log \langle \tilde{b}, X \rangle \big) = \mathbb{E} \big(\log \langle \bar{b}, U \rangle - \log \langle \tilde{b}, U \rangle \big), \ \forall \bar{b}, \tilde{b} \in \mathscr{B}^m.
$$

Additionally, $0 \leq \max_{b \in \mathscr{B}^m} \mathbb{E} \big(\log \langle b, U \rangle \big) \leq \log(m).$

Proof. Since the scaling function is defined in [\(2.1\)](#page-5-1) as $u(X) = X/\langle \hat{b}, X \rangle$, given the variable *X*, the logarithmic returns of any portfolio can be decomposed into the following sum:

$$
\log \langle b, X \rangle = \log \langle b, u(X) \rangle + \log \langle \hat{b}, X \rangle, \ \forall b \in \mathcal{B}^m. \tag{2.3}
$$

Therefore, given any two generic strategies (\bar{b}_n) and (\tilde{b}_n) and the process $\{X_n\}_{n=1}^{\infty}$, we have:

$$
W_n(\bar{b}_n) - W_n(\tilde{b}_n) = \frac{1}{n} \sum_{i=1}^n \log \langle \bar{b}_n, X_n \rangle - \frac{1}{n} \sum_{i=1}^n \log \langle \tilde{b}_n, X_n \rangle
$$

=
$$
\frac{1}{n} \sum_{i=1}^n \log \langle \bar{b}_n, u(X_n) \rangle - \frac{1}{n} \sum_{i=1}^n \log \langle \tilde{b}_n, u(X_n) \rangle, \forall n,
$$

which establishes assertion (a), given that $U_n = u(X_n)$.

To prove assertion (b), we first note that the variable *U* is bounded as:

$$
0 < U_i = \frac{X_i}{\langle \hat{b}, X \rangle} = \frac{mX_i}{\sum_{i=1}^m X_i} \le m, \ \forall i \in \{1, ..., m\},
$$
 (2.4)

due to $\hat{b} = (1/m, ..., 1/m)$. As a result, for any portfolio, we have the following inequality:

$$
0 = \mathbb{E} \big(\log \langle \hat{b}, U \rangle \big) \leq \max_{b \in \mathscr{B}^m} \mathbb{E} \big(\log \langle b, U \rangle \big) \leq \mathbb{E} \big(\log \big(m \big) \big) = \log \big(m \big),
$$

which attests to the finite non-negativity of the maximal expected value $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U \rangle)$. Furthermore, we also obtain the decomposition deduced directly from [\(2.3\)](#page-5-3) as follows:

$$
\mathbb{E}\big(\log \big\langle b,X \big\rangle\big)=\mathbb{E}\big(\log \big\langle b,U \big\rangle\big)+\mathbb{E}\big(\log \big\langle \hat{b},X \big\rangle\big).
$$

Then, the needed equality of the assertion follows immediately as:

$$
\begin{aligned} \mathbb{E}\big(\log \big\langle \bar{b}, X\big\rangle - \log \big\langle \tilde{b}, X\big\rangle \big) &= \mathbb{E}\big(\log \big\langle \bar{b}, U\big\rangle \big) - \mathbb{E}\big(\log \big\langle \tilde{b}, U\big\rangle \big) \\ &= \mathbb{E}\big(\log \big\langle \bar{b}, U\big\rangle - \log \big\langle \tilde{b}, U\big\rangle \big), \ \forall \bar{b}, \tilde{b} \in \mathscr{B}^m, \end{aligned}
$$

which finalizes our proof.

The properties established in Lemma [1](#page-5-0) are helpful for our discussion in the sense that the log-optimal portfolios defined according to Definition [1,](#page-4-0) in which the random variables X_n and thus the associated logarithmic returns of portfolios $log \langle b_n^*, X_n \rangle$ might not be integrable, can be investigated through the equivalent conditional expected difference as:

 $\mathbb{E} \big(\log \langle b, X_n \rangle - \log \langle b_n^*, X_n \rangle | \mathscr{F}_n^X \big) = \mathbb{E} \big(\log \langle b, U_n \rangle - \log \langle b_n^*, U_n \rangle | \mathscr{F}_n^X \big) \leq 0, \forall b \in \mathscr{B}^m, \forall n, (2.5)$ in which the optimally conditional expected value $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U \rangle | \mathcal{F}_n^X)$ is always welldefined, i.e., the variables $log \langle b_n^*, U \rangle$ associated with the log-optimal portfolios b_n^* are integrable given any probability measure. Moreover, the approach of considering the difference between the logarithmic returns of two portfolios rather than their individual values allows us to remove the redundantly inherent term $\log \langle \hat{b}, X \rangle$, thus reducing the originally unbounded growth rate of the log-optimal strategy $W_n(b_n^*)$ to the upper-bounded growth rate $n^{-1} \sum_{i=1}^n \log \langle b_n^*, U_n \rangle$ over all periods. In this context, some important properties of the log-optimal strategy under the normalized assets' returns will be discussed in the next section.

3 Supporting lemmas on the utilization of market information

After introducing the log-optimal strategy and the normalized assets' returns variable, we investigate various properties of competitive strategies benchmarked against the log-optimal strategy in this section. Through these comparisons, the optimality of the benchmarked log-optimal strategy is demonstrated in terms of utilizing market information to asymptotically achieve superior growth rates and expected cumulative wealth. These properties also provide further insights into the maximum potential of a strategy when all side information is fully observed. The analysis is presented in the following three subsections, where some results are straightforward extensions of

 \Box

those by [Algoet and Cover \(1988\)](#page-32-0), adapted to the normalized assets' returns variable, while others are improvements, offering clearer proofs and novel deductions.

3.1 Optimality principles of the log-optimal strategy

In addition to being optimal in maximizing the conditional expected value of the logarithmic return of a portfolio, log-optimal portfolios are also optimal in guaranteeing the highest conditional expected ratio of return compared to all other portfolios, given any σ -field \mathscr{F}_n^X . Consequently, the growth rate of the log-optimal strategy will almost surely (a.s.) not be exceeded by that of any other competitive strategy infinitely often by an arbitrary magnitude as time evolves. These optimal properties of the log-optimal strategy are established in Lemma [2.](#page-7-0)

Lemma 2. Consider a log-optimal strategy (b_n^*) and any competitive strategy (b_n) . We have:

- $\int \mathbb{E}\left(\langle b_n, U_n \rangle / \langle b_n^*, U_n \rangle | \mathcal{F}_n^X\right) \leq 1, \forall n.$
- (b) $\limsup_{n \to \infty} (W_n(b_n) W_n(b_n^*)) \leq 0, \text{ a.s.}$

Noting that the growth rate of the log-optimal strategy $W_n(b_n^*)$ *does not necessarily have a limit.*

Proof. Assertion (a) is simply a direct result from applying the Kuhn-Tucker conditions for logoptimality in [Bell and Cover \(1980\)](#page-32-16) and [Algoet and Cover](#page-32-0) [\(1988](#page-32-0)). Specifically, noting that the log-optimal portfolios can be equivalently obtained as [\(2.5\)](#page-6-0) by Lemma [1,](#page-5-0) where the maximal conditional expected values $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U \rangle | \mathcal{F}_n^X)$ are finite, the log-optimal portfolios always exist, and the Kuhn-Tucker conditions yield the required inequality in the assertion.

To prove assertion (b), we modify the arguments originally presented by [Algoet and Cover](#page-32-0) [\(1988](#page-32-0)) to show that $(S_n(b_n)/S_n(b_n^*), \mathcal{F}_n^X)$ is a non-negative supermartingale. In detail, since $\langle \hat{b}, X_n \rangle > 0$ for any *n*, we obtain the following inequality using assertion (a):

$$
\mathbb{E}\Big(\frac{S_{n+1}(b_{n+1})}{S_{n+1}(b_{n+1}^*)}\Big|\mathscr{F}_{n+1}^X\Big)=\mathbb{E}\Big(\frac{S_n(b_n)\big\langle\hat{b},X_{n+1}\big\rangle\big\langle b_{n+1},U_{n+1}\big\rangle}{S_n(b_n^*)\big\langle\hat{b},X_{n+1}\big\rangle\big\langle b_{n+1}^*,U_{n+1}\big\rangle}\Big|\mathscr{F}_{n+1}^X\Big)\newline=\frac{S_n(b_n)}{S_n(b_n^*)}\mathbb{E}\Big(\frac{\big\langle b_{n+1},U_{n+1}\big\rangle}{\big\langle b_{n+1}^*,U_{n+1}\big\rangle}\Big|\mathscr{F}_{n+1}^X\Big)\leq \frac{S_n(b_n)}{S_n(b_n^*)},\forall n.
$$

Consequently, we obtain $\mathbb{E}(S_n(b_n)/S_n(b_n^*)) \leq \mathbb{E}(S_0/S_0) = 1$ and then deduce the required result of the assertion by invoking the Markov inequality as follows:

$$
\mathbb{P}(S_n(b_n) > n^2 S_n(b_n^*) < \frac{1}{n^2}
$$

\n
$$
\Rightarrow \qquad \sum_{n=1}^{\infty} \mathbb{P}\left(\frac{1}{n} \log \frac{S_n(b_n)}{S_n(b_n^*)} > \frac{2\log n}{n}\right) \le \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \text{ (the Basel problem)}
$$

\n
$$
\Rightarrow \mathbb{P}\left(\frac{1}{n} \log \frac{S_n(b_n)}{S_n(b_n^*)} > \varepsilon \text{ infinitely often}\right) = 0 \text{ (by the Borel-Cantelli lemma)},
$$

for any arbitrarily small $\varepsilon > 0$ as $n \to \infty$, i.e $\limsup_{n \to \infty} (W_n(b_n) - W_n(b_n^*)) \le 0$ almost surely. \Box

Besides the almost sure guarantee of optimality in asymptotic growth rate compared with other competitive strategies by Lemma [2,](#page-7-0) the log-optimal strategy is also the optimal one that maximizes the expected logarithmic cumulative wealth among all accessible strategies for any time horizon. This property is discussed in the next section, which evaluates the benefit of using a larger amount of information rather than a smaller amount when making decisions.

3.2 Positive impact of large information size

One might argue that the use of all information, including unobservable side information, is very costly and challenging due to the difficulty of accurately estimating the conditional distribution, particularly since the side information is often high-dimensional in reality. However, if there is a relationship between the assets' returns and all the side information, removing some of this information to reduce dimensionality only makes sense if its correlation with the assets' returns is insignificant - something that, in fact, remains unknown if not all information can be fully observed. In Lemma [3,](#page-8-0) we establish a comparison between two log-optimal portfolios that depend on different sizes of observed information. This comparison highlights the advantages of utilizing a larger set of information and justifies the optimality of the log-optimal strategies over any other kind of strategy among all possible ones.

Lemma 3. Assume an assets' returns variable X, defined on $(\Omega, \mathbb{F}, \mathbb{P})$, admits a regular condi*tional distribution given any sub-*σ*-field of* F*. Then, the normalized assets' returns variable U also admits the corresponding transformed conditional distribution, and the following hold:*

(a) *Given a sub-* σ *-field* $\mathscr{G} \subseteq \mathbb{F}$ *, there exists a* \mathscr{G} *-measurable optimal portfolio such that:*

$$
\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U\big\rangle|\mathscr{G}\big)=\max_{b\in\mathscr{G}}\mathbb{E}\big(\log\big\langle b,U\big\rangle|\mathscr{G}\big)
$$

,

 \Box

where the notation b [∈] ^G *denotes the* ^G *-measurability of a portfolio b. Consequently:*

$$
\mathbb{E}\big(\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U\big\rangle|\mathscr{G}\big)\big)=\max_{b\in\mathscr{G}}\mathbb{E}\big(\log\big\langle b,U\big\rangle\big).
$$

(b) *The following inequality holds for any two sub-* σ *-fields* $\mathscr{G} \subseteq \mathscr{H} \subseteq \mathbb{F}$:

$$
\mathbb{E}\big(\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U\big\rangle|\mathscr{G}\big)\big)\leq\mathbb{E}\big(\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U\big\rangle|\mathscr{H}\big)\big).
$$

Proof. Assertion (a) follows directly from Theorem 2 in [Algoet and Cover \(1988\)](#page-32-0), which guarantees the existence of a measurably selected optimal portfolio, depending on Q^P , that maximizes the expectation $\mathbb{E}(\log \langle b, U \rangle)$ among all portfolios in \mathcal{B}^m . Here, $U \sim Q^P$, and Q^P is the transformation of the distribution *P* of the variable *X*. Since the conditional distribution given the σ -field $\mathscr G$ is $\mathscr G$ -measurable, there exists an optimal portfolio, denoted as $b^*_{\mathscr G} \in \mathscr G$, as a maximizer for $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U \rangle | \mathcal{G})$. Then, taking the expectation on both sides of the equality $\max_{b \in \mathscr{B}^m} \mathbb{E} \left(\log \langle b, U \rangle | \mathscr{G} \right) = \max_{b \in \mathscr{G}} \mathbb{E} \left(\log \langle b, U \rangle | \mathscr{G} \right)$, we obtain the following:

$$
\mathbb{E}\big(\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U\big\rangle|\mathscr{G}\big)\big)=\mathbb{E}\big(\mathbb{E}\big(\log\big\langle b^*_\mathscr{G},U\big\rangle|\mathscr{G}\big)\big)\\=\max_{b\in\mathscr{G}}\mathbb{E}\big(\log\big\langle b,U\big\rangle\big),
$$

due to the law of total expectation applied to the random variable $\log \langle b_{\mathscr{G}}^*, U \rangle$.

Now, consider any two sub-σ-fields $\mathscr{G} \subseteq \mathscr{H} \subseteq \mathbb{F}$. There exist two optimal portfolios, denoted as $b^*_\mathscr{G} \in \mathscr{G}$ and $b^*_\mathscr{H} \in \mathscr{H}$, for $\max_{b \in \mathscr{B}^m} \mathbb{E} \big(\log \big\langle b, U \big\rangle | \mathscr{G} \big)$ and $\max_{b \in \mathscr{B}^m} \mathbb{E} \big(\log \big\langle b, U \big\rangle | \mathscr{H} \big),$ respectively. In addition, since $\mathscr{G} \subseteq \mathscr{H}$ implies $b^*_{\mathscr{G}} \in \mathscr{H}$, we have:

$$
\mathbb{E}\big(\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U\big\rangle|\mathscr{G}\big)\big)=\mathbb{E}\big(\log\big\langle b^*_\mathscr{G},U\big\rangle\big)\\ \leq\max_{b\in\mathscr{H}}\mathbb{E}\big(\log\big\langle b,U\big\rangle\big)=\mathbb{E}\big(\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U\big\rangle|\mathscr{H}\big)\big).
$$

This demonstrates assertion (b) and completes the proof.

It is worth stressing again here that we are considering all possible strategies (b_n) , where each causal portfolio *bⁿ* in the sequence is selected based solely on past market information up to time *n*. These strategies obviously include all the log-optimal strategies. Hence, for any time horizon *m*, a log-optimal strategy maximizes the expected logarithm of cumulative wealth among all strategies in terms of normalized asset returns, as shown by the following chain rule:

$$
\max_{(b_n)} \mathbb{E}\left(\sum_{i=1}^m \log \langle b_i, U_i \rangle\right) = \sum_{i=1}^m \max_{b_i} \mathbb{E}\left(\log \langle b_i, U_i \rangle\right)
$$

$$
= \sum_{i=1}^m \mathbb{E}\left(\max_{b \in \mathcal{B}^m} \mathbb{E}\left(\log \langle b, U_i \rangle | \mathcal{F}_i^X\right)\right), \forall m. \tag{3.1}
$$

It should be emphasized that the optimality property described above is general, in that it does not require specific functional characteristics or measurability for the log-optimal strategy or any other strategy. Consequently, any other strategy that causally uses only a smaller subset of past information as a sub- σ -field of \mathcal{F}_n^X at all times *n*, might be inferior in terms of the expected logarithm of cumulative wealth due to $\mathbb{E}(\max_{b \in \mathscr{B}^m} \mathbb{E}(\log \langle b, U_n \rangle | \mathscr{G})) \leq \mathbb{E}(\max_{b \in \mathscr{B}^m} \mathbb{E}(\log \langle b, U_n \rangle | \mathscr{F}_n^X))$ for any σ -field $\mathscr{G} \subseteq \mathscr{F}_n^X$, as shown in Lemma [3.](#page-8-0) Moreover, the corresponding expected growth rate of a strategy, in terms of normalized asset returns, is also maximized by the log-optimal strategies for any time horizon. This expected growth rate will almost surely converge to a finite limit as time progresses, provided the market is stationary, as demonstrated later in Corollary [1.](#page-16-0)

3.3 Limit behavior of the log-optimal strategy

Since the log-optimal portfolio is derived using the conditional distributions based on past observations over time, it is essential to investigate its limit behavior in terms of the distributions, as described in Lemma [4.](#page-9-0) Generally, the expected logarithmic returns of the portfolio, in terms of normalized asset returns, are continuous in the space of all distributions equipped with the weak topology. This implies that, given a sequence of converging distributions, any accumulation of the corresponding log-optimal portfolios must also be a log-optimal portfolio for the limiting distribution. Hence, as the regular conditional distribution, given an increasing information field, is a converging sequence, any choice of log-optimal strategy ensures that the portfolio's returns will almost surely converge to those of the limiting portfolio under the limiting distribution.

Moreover, another important consequence of the lemma is that the normalized return of the log-optimal portfolios $\langle b_n^*, U_n \rangle$ is uniquely defined for almost all outcomes of U_n under the associated conditional distributions, regardless of the choice of b_n^* . The log-optimal b_n^* is uniquely defined only if the associated regular conditional distribution $\mathbb{P}(U_n|\mathscr{F}_n^X)$ has full-dimensional support, which makes the corresponding expected values $\mathbb{E}(\log \langle b, U_n \rangle | \mathcal{F}_n^X)$ strictly concave in $b \in \mathcal{B}^m$. Consequently, any log-optimal strategy (b_n^*) must also converge to a limiting portfolio almost surely if the conditional distribution $\mathbb{P}\big(U_1|\mathscr{F}^X_\infty\big)$ almost surely has full-dimensional support.

Lemma 4. *With P-a.a X representing the extension notation for an event occurring for almost all outcomes of X according to a specific distribution P of X, we have:*

(a) *Consider two portfolios* \bar{b} *and* \tilde{b} *maximizing* $\mathbb{E}(\log \langle b, U \rangle)$ *, where* $U \sim Q$ *. Then:*

$$
\langle \bar{b}, U \rangle = \langle \tilde{b}, U \rangle, Q
$$
-a.a *U*.

(b) Let b^{*}(Q) denote a portfolio maximizing the expected portfolio's logarithmic return in terms of normalized assets' returns according to the distribution Q as $\mathbb{E}\big(\log \langle b^*(Q), U \rangle\big) =$ $\max_{b \in \mathscr{B}^m} \mathbb{E}(\log \langle b, U \rangle)$ where $U \sim Q$. Consider any sequence of distributions $\{Q_n\}_{n=1}^{\infty}$ *converging weakly to a limiting distribution* Q_{∞} *as* $n \to \infty$ *, then:*

$$
\begin{cases}\n\lim_{n\to\infty}\langle b^*(Q_n),U\rangle=\langle b^*(Q_\infty),U\rangle, Q_\infty\text{-}a.a U \\
\lim_{n\to\infty}\mathbb{E}\big(\log\langle b^*(Q_n),U\rangle\big)=\mathbb{E}\big(\log\langle b^*(Q_\infty),U\rangle\big)\n\end{cases}
$$

(c) Consider an assets' returns variable *X* defined on $(\Omega, \mathbb{F}, \mathbb{P})$ and admitting a regular condi*tional distribution given the* σ -fields $\mathscr{G}_n \subseteq \mathbb{F}$ *for all n and* $\mathscr{G}_\infty \coloneqq \sigma\left(\cup_{n=1}^\infty \mathscr{G}_n\right) \subseteq \mathbb{F}$ *, then the following equality holds for the normalized assets' returns U of X :*

> lim *n*→∞ $\mathbb{P}(X|\mathscr{G}_n) = \mathbb{P}(X|\mathscr{G}_\infty)$, weakly a.s, implying $\lim_{n \to \infty}$ *n*→∞ $\langle \bar{b}_n, U \rangle = \langle \bar{b}_\infty, U \rangle$, *a.s*,

where the portfolios \bar{b}_n are maximizers for $\max_{b\in\mathscr{B}^m} \mathbb{E}\big(\log \big\langle b, U\big\rangle|\mathscr{G}_n\big)$ with $n\in\mathbb{N}_+\cup\{\infty\}.$

Proof. To prove assertion (a), we utilize the Kuhn-Tucker condition for the log-optimal portfolio. Specifically, since $\mathbb{E}(\langle \bar{b}, U \rangle / \langle \tilde{b}, U \rangle) = 1$ where $U \sim Q$ by Lemma [2,](#page-7-0) we have:

$$
0 = \mathbb{E}\Big(\log \frac{\big<\bar{b}, U\big>}{\big<\tilde{b}, U\big>}\Big) \leq \log \mathbb{E}\Big(\frac{\big<\bar{b}, U\big>}{\big<\tilde{b}, U\big>}\Big) = 0,
$$

due to Jensen's inequality, where the equality holds if and only if $\langle \bar{b}, U \rangle / \langle \tilde{b}, U \rangle = 1$ for *Q*-almost all *U*, as required for the assertion.

Regarding assertion (b), we note that the bottom equality in the statement is directly justified by the continuity of the function $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U \rangle)$ in the distribution *Q* of *U*, as demonstrated Theorem 2 of [Algoet and Cover \(1988](#page-32-0)). Meanwhile, the upper equality in the statement is only a comment on Theorem 3 in this paper, but without a clear argument. Thus, we provide a clarified deduction. Specifically, suppose there exists a unique maximizer $b^*(Q_\infty)$ for the problem $\max_{b \in \mathcal{B}^m} \mathbb{E} (\log \langle b, U \rangle)$ where $U \sim Q_{\infty}$, then any sequence $\{b^*(Q_n)\}_{n=1}^{\infty}$ must converge to $b^*(Q_\infty)$, and so $\{\langle b^*(Q_n), U \rangle\}_{n=1}^\infty$ must converge to $\langle b^*(Q_\infty), U \rangle$ for almost all outcomes of *U*. Meanwhile, in the case where there are many maximizers for $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U \rangle)$, the sequence ${b^*(Q_n)}_{n=1}^{\infty}$ may not converge, but all its accumulation points must be among these maximizers. Furthermore, since $\langle b^*(Q_\infty), U \rangle$ must be unique for any choice of $b^*(Q_\infty)$, as guaranteed by assertion (a) for *Q*-almost all *U*, the sequence $\{\langle b^*(Q_n), U \rangle\}_{n=1}^{\infty}$ must converge to $\langle b^*(Q_\infty), U \rangle$ for *Q*-almost all *U*, as needed for the statement.

The proof for assertion (c) follows from Theorem 4 in the aforementioned paper, but with a clarified deduction. Specifically, since $\mathbb{P}(X|\mathscr{G}_n) \to \mathbb{P}(X|\mathscr{G}_\infty)$ weakly almost surely by Levy's martingale convergence theorem, we have $\langle \bar{b}_n, U \rangle \to \langle \bar{b}_\infty, U \rangle$ for $\mathbb{P}(U|\mathscr{G}_\infty)$ -almost all *U*, almost surely by assertion (b). Therefore, we conclude that $\langle \bar{b}_n, U \rangle \to \langle \bar{b}_\infty, U \rangle$ almost surely, completing the proof. \Box

4 Limit and equality theorems of growth rate in stationary market

Since information is gradually and continuously revealed, the amount of past data used for decisionmaking, including the unobservable events, increases to infinitely large amounts. This probably misleads us to think that a dynamic log-optimal strategy, which varies depending on increasing information fields and is optimal in both the sense of asymptotic growth rate and expected cumulative wealth and growth rate (as discussed in the previous sections), should also easily outperform any constant strategy, which is invariant under changing market information. In fact, as we mentioned in the introduction as our motivation, the status quo of the dominating opinion in the relevant contemporary literature reflects this evaluation, following the well-known theories established in [Algoet and Cover \(1988](#page-32-0)); [Algoet](#page-32-6) [\(1992](#page-32-6)); [Cover](#page-32-4) [\(1991](#page-32-4), [1996](#page-32-8)); [Cover and Ordentlich](#page-32-10) [\(1996](#page-32-10)); [Cover and Thomas \(2006](#page-32-12)). However, the results established in Theorem [1,](#page-12-0) Theorem [3](#page-22-0) and later Corollary [1](#page-16-0) demonstrate surprising findings that contradict this popular unsettled conjecture.

Before investigating the evolutionary behavior of the log-optimal strategy, let us remark on some properties of stationary processes and introduce a new notation. Given the operator *T* representing the metrically transitive and invertible measure-preserving transformation of the underlying probability space $(\Omega, \mathbb{F}, \mathbb{P})$, let $\mathcal{I} \subset \mathbb{F}$ denote the *invariant* sub- σ -field henceforth as:

$$
\mathscr{I} \coloneqq \left\{ F \in \mathbb{F}: \, T^{-1}F = F \right\}.
$$

A stationary pairwise process $\{X_n, Y_n\}_{n=1}^{\infty}$ is defined as $(X_n, Y_n)(\omega) = (X_1, Y_1)(T^{(n-1)}\omega)$ for all *n*. This process could be extended infinitely to the left as a two-sided process with the origin at $n = 1$ by Kolmogorov's extension theorem. Under this condition, the invariant σ -field $\mathscr I$ is contained in the completion of the σ -field \mathcal{F}_{∞} of infinite information (and thus also in the completion of the σ -field \mathscr{F}^X_∞), which implies that no finite amount of information, but only the infinite one, suffices to exactly determine the random ergodic mode of the process $\{X_n, Y_n\}_{n=1}^{\infty}$. Fortunately, the true mode among a mixture of stationary ergodic ones could be gradually identified by the log-optimal strategy, which depends on the continuously increasing information fields. For convenience, let us introduce the reduced case of a *discrete market* in Definition [2](#page-11-0) and consider it the implicit default market henceforth, unless otherwise specified.

Definition 2. A discrete market is defined as a pairwise process $\{X_n, Y_n\}_{n=1}^{\infty}$ in which the assets' returns variables X_n take values in a countable subset $\mathbb{X} \subset \mathbb{R}^m_{++}$, for all *n*. In other words, they follow discrete (purely atomic) distributions with support X.

Remark. It is worth noting that the introduction of the discrete market in Definition [2](#page-11-0) is merely to simplify the proof of Lemma [5,](#page-12-1) and it does not affect the subsequent theorems, which remain valid without this condition. However, since the assets' returns in the discrete market may take an infinite number of values, the loss of generality is minimal, particularly when considering realworld markets. Additionally, the normalized assets' returns variable *U* is bounded within a small set as in [\(2.4\)](#page-6-1), implying that a fine discretization can cover most, if not all, possible outcomes. In the proof of Proposition [1,](#page-18-0) which appears later, an approach to bypass Lemma [5](#page-12-1) is discussed, so the discrete market condition is not required, though this involves a few more auxiliary lemmas, making Theorem [1](#page-12-0) less straightforward.

4.1 Equilibrium state under infinitely increasing market information

Lemma [4](#page-9-0) provides critical tools to argue Theorem [1,](#page-12-0) which serves as the basis to affirm that the periodic portfolios' returns of the log-optimal strategy could not exceed those of all constant strategies infinitely often as time progresses. This implies that such a dynamically optimal strategy could outperform any constant strategy, by utilizing the entirety of the increasing past information, only over a finite time horizon; thereafter, this advantage of using past information gradually degrades into an equilibrium state where the difference between the portfolios' returns of the dynamically optimal strategy and some constant strategies becomes negligible over time. As the theorem states, the cause of this phenomenon is clearly the stationarity of the market process, which directs the evolution of the log-optimal portfolios toward some optimal ones with respect to the limiting distribution, given the infinite past events as the largest possible capacity of the information field. Roughly speaking, although the cumulative wealth of the log-optimal strategy could be significantly better, its growth rate will be asymptotically identical to that of some constant strategies once it enters the equilibrium state.

Lemma 5. Assume the market process $\{X_n, Y_n\}_{n=1}^{\infty}$ is stationary. Then, the following convergence *holds for any two generic strategies* (\bar{b}_n) *and* (\tilde{b}_n) *, such that* $(\log \langle \bar{b}_n, U_k \rangle - \log \langle \tilde{b}_n, U_k \rangle) \to 0$ *almost surely for all normalized assets' returns* U_k *with* $k \geq 1$ *:*

$$
\lim_{n\to\infty}\big(\log\big\langle\bar{b}_n,U_n\big\rangle-\log\big\langle\tilde{b}_n,U_n\big\rangle\big)=0,\ a.s.
$$

Proof. By stationarity, noting that any finite sample $\{X_1(\omega), X_2(\omega), ..., X_n(\omega)\}\)$ can be represented as $\{X_1(\omega), X_1(T\omega),..., X_1(T^{(n-1)}\omega)\}\$ for each $\omega \in \Omega$ and for all *n*, the ergodic mode of the market process is determined by the invariant σ -field $\mathscr I$. Hence, by applying Birkhoff's ergodic theorem, we obtain the following limit (with the discrete market as in Definition [2\)](#page-11-0):

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\mathbb{I}_{\{X_1=x\}}=\mathbb{P}\big(X_1=x|\mathscr{I}\big)>0,\text{ a.s.}
$$

This implies that for each realization of the process $\{X_n(\omega)\}_{n=1}^{\infty}$, there exists a corresponding subset $\mathbb{X}^{\omega} \subset \mathbb{X}$ such that all outcomes $x \in \mathbb{X}^{\omega}$ with $\mathbb{P}(X_1 = x | \mathscr{I})(\omega) > 0$ must occur infinitely often as $X_n(\omega) = x$ for $n \to \infty$, while other $x \notin \mathbb{X}^{\omega}$ occur only finitely many times. Therefore, the convergence $\big(\log\big\langle\bar b_n(\omega),u(x)\big\rangle-\log\big\langle\tilde b_n(\omega),u(x)\big\rangle\big)\to 0$ holds for $\mathbb{P}\big(X_1|\mathscr{I}\big)(\omega)$ -almost all $x \in \mathbb{X}^{\omega}$ almost surely, due to the assumption that $\big(\log\big\langle\bar{b}_n,U_k\big\rangle\big(\omega\big)-\log\big\langle\tilde{b}_n,U_k\big\rangle\big(\omega\big)\big)\to 0$ almost surely for all $k \ge 1$. This convergence implies that for any $\varepsilon > 0$ and each $x \in \mathbb{X}^{\omega}$, we have:

 $\exists N^{\omega,\varepsilon}(x) \text{ such that } |\log \langle \bar{b}_n(\omega), u(x) \rangle - \log \langle \tilde{b}_n(\omega), u(x) \rangle| < \varepsilon, \forall n \ge N^{\omega,\varepsilon}(x).$

Then, the required convergence $(\log \langle \bar{b}_n, U_n \rangle(\omega) - \log \langle \tilde{b}_n, U_n \rangle(\omega)) \to 0$ must hold. If not, then there exists $\mathcal{E}^{\omega} > 0$ such that $|\log \langle \bar{b}_t(\omega), u(x) \rangle - \log \langle \tilde{b}_t(\omega), u(x) \rangle| \geq \mathcal{E}^{\omega}$ for $t \geq K$ for some *x* as $K \to \infty$, i.e., some $x \in \mathbb{X}^{\omega}$ occurs only finitely many times because some $N^{\omega,\varepsilon^{\omega}}(x)$ are never visited, which contradicts the derived property above. Hence, we conclude that, with probability one, the convergence $\left(\log \langle \bar{b}_n, U_n \rangle - \log \langle \tilde{b}_n, U_n \rangle\right) \to 0$ holds, completing the proof. \Box

Theorem 1. If the market process $\{X_n, Y_n\}_{n=1}^{\infty}$ is stationary, there exists a random constant strategy, denoted as (b^{ω}) , such that the growth rate of any log-optimal strategy (b_n^*) does not exceed *that of the constant one infinitely often by any fixed magnitude, as:*

$$
\lim_{n\to\infty}\big(W_n(b^{\omega})-W_n(b_n^*)\big)=0,\ a.s.
$$

Moreover, such a random constant strategy (b^{ω}) can be chosen measurably.

Proof. Since the random process $\{X_n, Y_n\}_{n=1}^{\infty}$ is stationary, it can be extended to a two-sided process. Thus, any variable (X_n, Y_n) , for all *n*, admits a regular conditional distribution given any σ -field \mathscr{F}_n or the infinite σ -field \mathscr{F}_{∞} . This immediately results in $\mathbb{E}(\log \langle b, X_n \rangle | \mathscr{F}_{k}^X) =$ $\mathbb{E}(\log \langle b, X_1 \rangle | \mathcal{F}_k^X)$ for any $n > 1$ and $b \in \mathcal{B}^m$, given the same information field \mathcal{F}_k^X with $k \in$ $\mathbb{N}_+ \cup \{\infty\}$. Then, by Lemma [4,](#page-9-0) it follows that $\mathbb{P}(X_k|\mathscr{F}_n^X) \to \mathbb{P}(X_k|\mathscr{F}_\infty^X)$ weakly almost surely as $n \to \infty$ for all $k \ge 1$, and the limit below holds for any log-optimal strategy (b_n^*) :

$$
\lim_{n\to\infty}\log\langle b_n^*,U_k\rangle=\log\langle b_\infty^*,U_k\rangle, \forall k\geq 1, \text{ a.s.}
$$

with portfolio b^*_{∞} denoting an arbitrary maximizer for $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U_1 \rangle | \mathcal{F}^X_{\infty})$, and noting that b^*_{∞} is random. This convergence can be represented as $(\log \langle b^*_n, U_k \rangle - \log \langle b^*_\infty, U_k \rangle) \to 0$ for the log-optimal strategy (b_n^*) and the random constant strategy (b_{∞}^*) , for all $k \ge 1$. Hence, we have almost surely that $\left(\log \langle b_n^*, U_n \rangle - \log \langle b_\infty^*, U_n \rangle \right) \to 0$ by Lemma [5,](#page-12-1) and the following is obtained:

$$
\lim_{n \to \infty} (W_n(b^*_{\infty}) - W_n(b^*_{n})) = \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^n (\log \langle b^*_{\infty}, U_i \rangle - \log \langle b^*_{i}, U_i \rangle)
$$

=
$$
\lim_{n \to \infty} (\log \langle b^*_{\infty}, U_n \rangle - \log \langle b^*_{n}, U_n \rangle) = 0, \text{ a.s.}
$$

due to Cesaro's mean theorem and Lemma [1.](#page-5-0) As a consequence, by simply setting a portfolio $b^{\omega} = b_{\infty}^*$, which is random and can be measurably selected depending on the σ -field \mathscr{F}_{∞}^X by Lemma [3,](#page-8-0) we finish the proof of the existence of the optimal random constant strategy (b^{ω}) .

4.2 Optimal constant strategy unconditioned on side information

It is known from Theorem [1](#page-12-0) that there exist some random constant strategies, derived depending on infinitely many data of assets' returns and side information, that could asymptotically approach the growth rate and portfolio returns of the log-optimal strategy under a stationary market. However, one might argue a specific context in which there exists a non-random, i.e., independent from the market's possibilities, optimal constant strategy with such a property. In addition, an interesting question naturally arises from the existence of the random optimal constant strategy: whether the limiting optimal growth rate could be simultaneously represented for both the log-optimal strategy, which is dynamic, and the constant strategies. Theorem [2](#page-13-0) establishes such an equality for the limiting growth rate in terms of normalized assets' returns for the two regarded strategies under the stationary condition. Through this equality, the optimal strategy's growth rate when infinitely increasing market side information is used is equivalently measured by the maximal capacity of a constant strategy without depending on periodic market observation. Moreover, if the market is also ergodic, the existence of a non-random optimal constant strategy could be secured using the mentioned equality. Before the theorem statement, let us further introduce an optional condition for the market in Definition [3.](#page-13-1)

Definition 3. A market is said not to have *trash assets* if $\max_{1 \le i \le m} X_{n,i} / \min_{1 \le i \le m} X_{n,i} \ge c \gg 0$ for all *n* almost surely. This does not necessarily bound the returns of the assets, but instead excludes the trash assets from the considered portfolios, which are defined as those that might potentially collapse in value to nearly zero prices due to the bankruptcy of the firms.

Remark. It is important to highlight that the almost sure convergence of the log-optimal strategy in Theorem [2](#page-13-0) can be proven without the assumption of no-trash assets, by simply using Birkhoff's ergodic theorem, as in the original proof by [Algoet and Cover](#page-32-0) [\(1988](#page-32-0)). The key difference is that Theorem [2](#page-13-0) does not assume finite expected values, due to the transformation from assets' returns to the normalized ones. The additional no-trash assets condition is primarily used to ensure uniform integrability, which can be useful for certain later applications, such as more conveniently establishing Corollary [1](#page-16-0) and Theorem [4.](#page-27-0)

Theorem 2. If the process $\{X_n, Y_n\}_{n=1}^{\infty}$ is stationary, then there exists a random constant strategy (b^*) *that grows asymptotically as fast as the log-optimal strategy* (b^*_n) *as:*

$$
\lim_{n\to\infty}\big(W_n(b^*)-W_n(b_n^*)\big)=0,\ a.s.
$$

This follows from the limits holding in terms of normalized assets' returns:

$$
\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b^*, U_i \rangle = \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_i^*, U_i \rangle = \mathbb{E} \big(\log \langle b_{\infty}^*, U_1 \rangle | \mathcal{I} \big)
$$

$$
= \max_{b \in \mathcal{B}^m} \mathbb{E} \big(\log \langle b, U_1 \rangle | \mathcal{I} \big), a.s,
$$

where the portfolio b^*_{∞} is a measurable maximizer for $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U_1 \rangle | \mathcal{F}^X_{\infty})$. Moreover, *if the market does not have trash assets, then the convergences above are also in L*¹ *; and if the market is also ergodic, then the optimal constant strategy is non-random.*

Proof. By Lemma [2,](#page-7-0) an inequality $\limsup_{n\to\infty} (W_n(b^*) - W_n(b^*)) \le 0$ is established almost surely for any constant strategy (b^*) . Therefore, by using Lemma [1,](#page-5-0) it is sufficient to prove the following inequality for some constant strategy (b^*) :

$$
\liminf_{n\to\infty} \left(W_n(b^*) - W_n(b_n^*) \right) = \liminf_{n\to\infty} \frac{1}{n} \left(\sum_{i=1}^n \log \left\langle b^*, U_i \right\rangle - \sum_{i=1}^n \log \left\langle b_i^*, U_i \right\rangle \right) \ge 0, \text{ a.s.}
$$

Then, the remaining limits in the statement will follow.

First, we prove the condition of being L^1 -dominated for any sequence of logarithmic returns of a strategy (b_n) with the measurable portfolios b_n for all *n*, i.e., $\mathbb{E}(\sup_n |\log \langle b_n, U_n \rangle|)$ is finite. This is required to invoke Breiman's generalized ergodic theorem (noting that the original ergodic theorem by [Breiman \(1957](#page-32-17)) only modified the almost sure convergence from Birkhoff's ergodic theorem, which was later extended to convergence in L^1 by [Algoet](#page-32-18) [\(1994](#page-32-18)), with the same condition of integrability). Specifically, we have:

$$
|\log \langle b_n, U_n \rangle| \le \max \left\{ |\log (c)|, |\log (m)| \right\} < \infty, \ \forall n, \text{ a.s.}
$$
 (4.1)

This is due to the bounds $\log \langle b_n, U_n \rangle \leq \log(m)$ by Lemma [1,](#page-5-0) and $\log(c) \leq \log \langle b_n, U_n \rangle$ by the market condition without trash assets, as follows:

$$
U_{n,k} \geq \min_{1 \leq i \leq m} \frac{m X_{n,i}}{\sum_{j=1}^m X_{n,j}} \geq \min_{1 \leq i \leq m} \max_{1 \leq j \leq m} \frac{X_{n,i}}{X_{n,j}} \geq c, \ \forall n, \forall k \in \{1, ..., m\}, \ \text{a.s.}
$$

Next, we proceed to establish the limits, in terms of the normalized assets' returns, of the logoptimal strategy and a random constant strategy as follows.

(The log-optimal strategy). Consider a log-optimal strategy (b_n^*) with the \mathscr{F}_n^X -measurable portfolios b_n^* for all *n*, which can be selected by Lemma [3.](#page-8-0) By stationarity, for each *n*, the measurable portfolio b_n^* can be expressed as a shift from another portfolio:

$$
b_n^*(\omega) = b_n^{-*}(T^{(n-1)}\omega), \text{ where } \mathbb{E}(\log \langle b_n^{-*}, U_1 \rangle | T^{(1-n)}\mathscr{F}_n^X) = \max_{b \in \mathscr{B}^m} \mathbb{E}(\log \langle b, U_1 \rangle | T^{(1-n)}\mathscr{F}_n^X).
$$

The inverse filtration $\left\{T^{(1-n)}\mathscr{F}_{n}^X\right\}_{n=1}^\infty$ increases up to the following inverse limiting sub- σ -field: $\mathscr{F}_{-\infty}^X := T^{-\infty} \mathscr{F}_{\infty}^X = \sigma\left(\cup_{n=1}^{\infty} T^{(1-n)} \mathscr{F}_n^X\right) \subseteq \mathbb{F}.$

By Lemma [4,](#page-9-0) we then have $\langle b_n^{-*}, U_1 \rangle \to \langle b_\infty^{-*}, U_1 \rangle$ almost surely, where the portfolio b_∞^{-*} is a maximizer for $\max_{b \in \mathcal{B}^m} \mathbb{E} \left(\log \langle b, U_1 \rangle | \mathcal{F}_{-\infty}^X \right)$, which can also be $\mathcal{F}_{-\infty}^X$ -measurably selected by Lemma [3.](#page-8-0) Given the invariant σ -field \mathscr{I} , Breiman's generalized ergodic theorem yields:

$$
\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_i^*, U_i \rangle (\omega) = \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_i^{-*} (T^{(i-1)} \omega), U_1 (T^{(i-1)} \omega) \rangle
$$

\n
$$
= \mathbb{E} (\log \langle b_\infty^*, U_1 \rangle | \mathscr{I})
$$

\n
$$
= \mathbb{E} (\log \langle b_\infty^*, U_1 \rangle | \mathscr{I}), \text{ a.s and in } L^1,
$$
\n(4.2)

where b^*_{∞} is a \mathscr{F}^X_{∞} -measurable maximizer for $\max_{b \in \mathscr{B}^m} \mathbb{E} (\log \langle b, U_1 \rangle | \mathscr{F}^X_{\infty})$, and the *L*¹-domination condition is satisfied by the sequence $\left\{ \log \left\langle b_n^*, U_n \right\rangle \right\}_{n=1}^{\infty}$, as stated earlier.

(The random constant strategy). Consider a random optimal portfolio *b* ∗ as a maximizer for $\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U_1\big\rangle|\mathscr{I}\big),$ which forms the corresponding random constant strategy $\big(b^*\big).$ Given the well-defined $\mathbb{E} \big(\log \big\langle b^*, U_1 \big\rangle | \mathscr{I} \big)$ by Lemma [1,](#page-5-0) we have:

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b^*,U_i\right\rangle=\mathbb{E}\big(\log\left\langle b^*,U_1\right\rangle|\mathscr{I}\big)=\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\left\langle b,U_1\right\rangle|\mathscr{I}\big),\text{ a.s and in }L^1,\qquad(4.3)
$$

by invoking Birkhoff's ergodic theorem. Noting that the L^1 -convergence follows from the almost sure convergence by applying Lebesgue's dominated convergence theorem, given that the process $\left\{ n^{-1} \sum_{i=1}^{n} \log \left\langle b^*, U_i \right\rangle \right\}_{n=1}^{\infty}$ is uniformly integrable, as established by the bound in [\(4.1\)](#page-14-0).

(Equalities derivation). We now establish the equality between the obtained limits in [\(4.2\)](#page-14-1) and [\(4.3\)](#page-14-2). First, recall the following equality:

$$
\lim_{n\to\infty} \left(W_n(b^\omega) - W_n(b_n^*) \right) = \lim_{n\to\infty} \frac{1}{n} \sum_{i=1}^n \left(\log \left\langle b^\omega, U_i \right\rangle - \log \left\langle b_i^*, U_i \right\rangle \right) = 0, \text{ a.s.}
$$

for a random constant strategy (b^{ω}) by Theorem [1.](#page-12-0) As a result, we have:

$$
\liminf_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \left(\log \left\langle b^{\omega}, U_{i} \right\rangle - \log \left\langle b_{i}^{*}, U_{i} \right\rangle \right) \leq \liminf_{n \to \infty} \frac{1}{n} \left(\max_{b \in \mathcal{B}^{m}} \sum_{i=1}^{n} \log \left\langle b, U_{i} \right\rangle - \sum_{i=1}^{n} \log \left\langle b_{i}^{*}, U_{i} \right\rangle \right)
$$
\n
$$
= \max_{b \in \mathcal{B}^{m}} \mathbb{E} \left(\log \left\langle b, U_{1} \right\rangle | \mathcal{J} \right) - \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \left\langle b_{i}^{*}, U_{i} \right\rangle
$$
\n
$$
= \lim_{n \to \infty} \frac{1}{n} \left(\sum_{i=1}^{n} \log \left\langle b^{*}, U_{i} \right\rangle - \sum_{i=1}^{n} \log \left\langle b_{i}^{*}, U_{i} \right\rangle \right), \text{ a.s. (4.4)}
$$

The establishment of [\(4.4\)](#page-15-0) above follows from the limit in [\(4.3\)](#page-14-2) and the following:

$$
\lim_{n \to \infty} \max_{b \in \mathcal{B}^m} \frac{1}{n} \sum_{i=1}^n \log \langle b, U_i \rangle = \lim_{n \to \infty} \max_{b \in \mathcal{B}^m} \mathbb{E}^{P_n} \big(\log \langle b, U \rangle \big)
$$

=
$$
\max_{b \in \mathcal{B}^m} \mathbb{E} \big(\log \langle b, U_1 \rangle | \mathcal{J} \big), \text{ a.s.}
$$

which is due to Lemma [4](#page-9-0) regarding the sequence of distributions $\{P_n\}_{n=1}^{\infty}$ defined as:

$$
P_n(A) \coloneqq \frac{1}{n} \sum_{i=1}^n \mathbb{I}_{U_i}(A), \forall A \subseteq \mathscr{U}, \forall n,
$$

with the notation $\mathbb{E}^{P_n}(\cdot)$ denoting the expected value with respect to $U \sim P_n$. Importantly, noting that the sequence of empirical distributions $\{P_n\}_{n=1}^{\infty}$ converges weakly almost surely to the limiting distribution $\mathbb{P}(U_1|\mathscr{I})$ by the argument below:

$$
\lim_{n\to\infty}\mathbb{E}^{P_n}(f(U))=\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n f(U_i)=\mathbb{E}(f(U_1)|\mathscr{I}), \text{ a.s.}
$$

for any bounded continuous function $f(\cdot)$ by Birkhoff's ergodic theorem.

Therefore, we finally obtain the following needed almost sure equality:

$$
0 = \lim_{n \to \infty} (W_n(b^*) - W_n(b_n^*))
$$

=
$$
\max_{b \in \mathscr{B}^m} \mathbb{E} (\log \langle b, U_1 \rangle | \mathscr{I}) - \mathbb{E} (\log \langle b_{-\infty}^*, U_1 \rangle | \mathscr{I}), \text{ a.s.}
$$

which, along with the L^1 -convergence in [\(4.2\)](#page-14-1) and [\(4.3\)](#page-14-2), results in the following:

$$
0 = \lim_{n \to \infty} \mathbb{E}\Big(\Big|\frac{1}{n}\sum_{i=1}^{n} \log \big\langle b^*, U_i \big\rangle - \mathbb{E}\big(\log \big\langle b^*, U_1 \big\rangle | \mathscr{I}\big)\Big|\Big) + \lim_{n \to \infty} \mathbb{E}\Big(\Big|\frac{1}{n}\sum_{i=1}^{n} \log \big\langle b^*_i, U_i \big\rangle - \mathbb{E}\big(\log \big\langle b^*, U_1 \big\rangle | \mathscr{I}\big)\Big|\Big) \geq \lim_{n \to \infty} \mathbb{E}\big(\big|W_n(b^*) - W_n(b_n^*)\big|\big).
$$

Hence, the last L^1 -convergence is also obtained, and the proof is completely finished.

According to Theorem [2,](#page-13-0) although the best possible growth rate of all strategies in the market might not converge, or might even be infinite, there exists an unsophisticated constant strategy that is possibly not exceeded by any optimally sophisticated strategy, leveraged by knowledge of all market information, including unobservable ones, by a fixed magnitude infinitely often as time elapses. Specifically, recalling that by changing the variable to the normalized assets' returns, the

 \Box

growth rate of any strategy (b_n) is decomposed into two distinct terms as follows:

$$
W_n(b_n) = \frac{1}{n} \mathbb{E}\left(\sum_{i=1}^n \log \langle b_i, U_i \rangle\right) + \frac{1}{n} \mathbb{E}\left(\sum_{i=1}^n \log \langle \hat{b}, X_i \rangle\right),
$$

where the optimal rate of the first term converges almost surely to a finite limit as $n \to \infty$ (without additional assumptions), as asserted by Theorem [2.](#page-13-0) In contrast, the growth rate $W_n(\hat{b})$ associated with the reference constant strategy (\hat{b}) may diverge or oscillate infinitely, as it depends on the uncontrolled original variables X_n . Therefore, it is clear that the original growth rate $W_n(b_n^*)$ of the log-optimal strategy (b_n^*) will converge almost surely to a finite limit if the growth rate $W_n(\hat{b})$ has a finite limit. In particular, if the market is also ergodic, the following holds by Lemma [3:](#page-8-0)

$$
\mathbb{E}\left(\max_{b\in\mathscr{B}^m}\mathbb{E}\left(\log\left\langle b,U_1\right\rangle|\mathscr{F}_{\infty}^X\right)=\mathbb{E}\left(\log\left\langle b_{\infty}^*,U_1\right\rangle\right)\right)\\=\max_{b\in\mathscr{B}^m}\mathbb{E}\left(\log\left\langle b,U_1\right\rangle\right),\tag{4.5}
$$

so the optimal constant strategy (b^*) is a non-random maximizer for $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U_1 \rangle)$.

Furthermore, the expected difference in growth rates between the log-optimal strategy and the random optimal constant strategy, which becomes non-random if the ergodicity condition is introduced, is also depleted to zero, as established in Corollary [1.](#page-16-0) This surprising result highlights that such an optimal constant strategy is comparable to any other dynamic strategy among all the accessible ones that asymptotically maximize the expected growth rate of strategy, which might seem counterintuitive. Finally, in the case of a stationary and ergodic process $\{X_n, Y_n\}_{n=1}^{\infty}$, the limits of the growth rate and the expected growth rate of the regarded optimal strategies, as established in the corollary, are the same and equal to the terms in [\(4.5\)](#page-16-1).

Corollary 1. If the process $\{X_n, Y_n\}_{n=1}^{\infty}$ is stationary (the condition of a no-trash assets market *is not necessary), then the following limits hold for the log-optimal strategy* (b_n^*) and a random *constant strategy* (b^*) among all possible strategies (b_n) as:

$$
\lim_{k\to\infty}\max_{(b_n)}\mathbb{E}\big(W_k(b_k)-W_k(b^*)\big)=\lim_{k\to\infty}\mathbb{E}\big(W_k(b_k^*)-W_k(b^*)\big)=0,
$$

where the optimal expected growth rate in terms of normalized assets' returns has a limit:

$$
\lim_{k \to \infty} \max_{(b_n)} \frac{1}{k} \mathbb{E} \left(\sum_{i=1}^k \log \langle b_i, U_i \rangle \right) = \lim_{k \to \infty} \frac{1}{k} \mathbb{E} \left(\sum_{i=1}^k \log \langle b_i^*, U_i \rangle \right) = \lim_{k \to \infty} \frac{1}{k} \mathbb{E} \left(\sum_{i=1}^k \log \langle b^*, U_i \rangle \right)
$$
\n
$$
= \mathbb{E} \left(\max_{b \in \mathcal{B}^m} \mathbb{E} \left(\log \langle b, U_1 \rangle | \mathcal{F}_{\infty}^X \right) \right)
$$
\n
$$
= \mathbb{E} \left(\log \langle b^*, U_1 \rangle \right).
$$

Proof. If the market does not have trash assets, the proof is straightforward by applying Jensen's inequality for the absolute value function and invoking the L^1 -convergence in Theorem [2.](#page-13-0) We then have the following limit, given the random optimal constant strategy (b^*) as in [\(4.3\)](#page-14-2):

$$
\lim_{k\to\infty}|\max_{(b_n)}\mathbb{E}\big(W_k\big(b_k\big)-W_k\big(b^*\big)\big)|=\lim_{k\to\infty}|\mathbb{E}\big(W_k\big(b_k^*\big)-W_k\big(b^*\big)\big)|\\\leq\lim_{k\to\infty}\mathbb{E}\big(|W_k\big(b_k^*\big)-W_k\big(b^*\big)|\big)=0.
$$

By similar arguments, the limiting optimal expected growth rate of the best strategy among all possible ones, in terms of normalized assets' returns, is a direct consequence of the L¹-convergence in [\(4.2\)](#page-14-1) and [\(4.3\)](#page-14-2), which implies the convergence of expected values.

Now, consider the general case without the no-trash assets condition. We proceed to prove the needed convergence $\mathbb{E}(W_n(b_n^*) - W_n(b^*)) \to 0$ by establishing the equality between the individual

expected growth rates in terms of normalized assets' returns of the involved optimal strategies. Specifically, consider the random optimal constant strategy (b^*) as in [\(4.3\)](#page-14-2). We have:

$$
\lim_{n\to\infty}\frac{1}{n}\mathbb{E}\big(\sum_{i=1}^n\log\big\langle b^*,U_i\big\rangle\big)=\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\mathbb{E}\big(\log\big\langle b^*,U_i\big\rangle\big)=\mathbb{E}\big(\log\big\langle b^*,U_1\big\rangle\big),
$$

because the random portfolio $b^*(\omega)$ is constantly chosen depending on each $\omega \in \Omega$ over all random pairs $(X_n, Y_n)(\omega)$ for all *n*, of the stationary process $\{X_n, Y_n\}_{n=1}^{\infty}$.

Moreover, by taking the expectation of both sides of the almost sure equalities of the limiting growth rates of the two regarded optimal strategies in Theorem [2,](#page-13-0) and then applying Lemma [3](#page-8-0) on each of them (noting that b^* can be treated as a log-optimal portfolio that is measurably selected depending on the σ -field \mathscr{I}), we obtain the following equality:

$$
\mathbb{E} \big(\log \langle b^*, U_1 \rangle \big) = \mathbb{E} \big(\max_{b \in \mathscr{B}^m} \mathbb{E} \big(\log \langle b, U_1 \rangle | \mathscr{I} \big) \big) = \mathbb{E} \big(\max_{b \in \mathscr{B}^m} \mathbb{E} \big(\log \langle b, U_1 \rangle | \mathscr{F}_{\infty}^X \big) \big),
$$

where the expected value on the right-hand side is also the limiting expected growth rate, in terms of normalized assets' returns, of the log-optimal strategy, as shown in the following arguments.

Let us recall that $\mathbb{P}(U_1|\mathscr{F}_n^X)\to\mathbb{P}(U_1|\mathscr{F}_\infty^X)$ weakly almost surely, according to Lemma [4,](#page-9-0) so $\max_{b \in \mathcal{B}^m} \mathbb{E} \left(\log \langle b, U_n \rangle | \mathcal{F}_n^X \right) \to \max_{b \in \mathcal{B}^m} \mathbb{E} \left(\log \langle b, U_1 \rangle | \mathcal{F}_\infty^X \right)$ by the stationarity of the market process. Subsequently, by using Lebesgue's dominated convergence theorem (since the maximal conditional expected values $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U_n \rangle | \mathcal{F}_n^X)$ are non-negative and upper bounded for all *n* by Lemma [1\)](#page-5-0), we obtain the following convergence:

$$
\lim_{n\to\infty} \mathbb{E}\big(\max_{b\in\mathscr{B}^m} \mathbb{E}\big(\log\langle b,U_n\rangle\,|\,\mathscr{F}_n^X\big)\big)=\mathbb{E}\big(\lim_{n\to\infty}\max_{b\in\mathscr{B}^m} \mathbb{E}\big(\log\langle b,U_n\rangle\,|\,\mathscr{F}_n^X\big)\big)\newline=\mathbb{E}\big(\max_{b\in\mathscr{B}^m} \mathbb{E}\big(\log\langle b,U_1\rangle\,|\,\mathscr{F}_\infty^X\big)\big).
$$

Moreover, by further using the chain rule in [\(3.1\)](#page-9-1) and Cesaro's mean theorem, we have:

$$
\lim_{k \to \infty} \frac{1}{k} \mathbb{E} \left(\sum_{i=1}^{k} \log \left\langle b_i^*, U_i \right\rangle \right) = \lim_{k \to \infty} \max_{(b_n)} \frac{1}{k} \mathbb{E} \left(\sum_{i=1}^{k} \log \left\langle b_i, U_i \right\rangle \right)
$$
\n
$$
= \lim_{k \to \infty} \frac{1}{k} \sum_{i=1}^{k} \mathbb{E} \left(\max_{b \in \mathscr{B}^m} \mathbb{E} \left(\log \left\langle b, U_i \right\rangle | \mathscr{F}_i^X \right) \right)
$$
\n
$$
= \lim_{k \to \infty} \mathbb{E} \left(\max_{b \in \mathscr{B}^m} \mathbb{E} \left(\log \left\langle b, U_k \right\rangle | \mathscr{F}_k^X \right) \right)
$$
\n
$$
= \mathbb{E} \left(\max_{b \in \mathscr{B}^m} \mathbb{E} \left(\log \left\langle b, U_1 \right\rangle | \mathscr{F}_\infty^X \right) \right).
$$

Therefore, we finally obtain all the necessary equalities of the statement and complete the proof.

\Box

4.3 Market with latent dependence structures

In this section, we discuss market scenarios with latent dependence structures, which can be derived as consequences of Theorem [1](#page-12-0) and Theorem [2.](#page-13-0) This serves to demonstrate the generality of the modeling settings and the properties obtained in the previous sections for capturing various real-market behaviors. In a real market, such dependence is justified, as economic agents and institutions typically react to market events, which may be correlated within the same market or across different markets due to causal effects. The structure of dependence may vary due to factors such as technological advancements, regulations, and social features, which influence the spread, speed of reception, and perception of information across markets. It is important to note that the dependence structures in markets are often not fully known, as not all supplementary information

is publicly observable or may be prohibitively costly to collect and store. Additionally, some information may not be truly valuable due to weak correlations with asset returns. We now proceed to capture the dependence structure through the following three market phenomena.

Market with non-decaying impact of information. Consider a market process $\{X_n, Y_n\}_{n=1}^{\infty}$, where the current assets' returns X_n and other market features Y_n , including unobservable ones, depend on past information from the start to the previous period *n*−1. In other words, the market has unlimited memory of past events, such that their effects can persist infinitely, with non-decaying impacts on the future market over an unrestricted horizon. This phenomenon of unlimited memory can be modeled by a market process with a transition property, where the distribution of each pairwise random variable depends on all past events, or alternatively, it is adapted to the filtration $\left\{\mathscr{F}_n\right\}_{n=0}^\infty$ $\sum_{n=1}^{\infty}$ that encompasses increasing past market information. Proposition [1](#page-18-0) considers this type of market model with a mild assumption, enabling the result to be directly derived without the need for stationarity in the market process.

Remark. The introduction of the market model in Proposition [1](#page-18-0) is intended as a specific instance for modeling the non-decaying impact of information and is not central to the main discussion in this paper. However, it provides a weaker assumption than strict stationarity, under which the existence of a random optimal constant strategy remains valid, even though the growth rate in terms of normalized assets' returns of the log-optimal strategy $n^{-1} \sum_{i=1}^{n} \log \langle b_i^*, U_i \rangle$ might not be guaranteed. Another useful aspect is that it provides an alternative proof for Theorem [1,](#page-12-0) without the need for Lemma [5](#page-12-1) and the discrete market condition as specified in Definition [2.](#page-11-0)

Proposition 1. Consider the market process $\{X_n, Y_n\}_{n=1}^{\infty}$, where the distributions of the random *pairs* (X_n, Y_n) depend on past information, such that $(X_n, Y_n) \sim \mathbb{P}(X_1, Y_1 | \mathscr{F}_n)$ for all $n \geq 2$, where $\mathbb{P}(X_1,Y_1|\mathscr{F}_n)$ is the regular conditional distribution given the σ -field \mathscr{F}_n admitted by (X_1,Y_1) . *Then, there exists a random optimal constant strategy* (b^{ω}) such that $(W_n(b^{\omega}) - W_n(b_n^*)) \to 0$ *almost surely, which is also in L*¹ *under the additional no-trash assets condition.*

Proof. Based on Lemma [4,](#page-9-0) if the space of all distributions of normalized assets' returns U_1 is equipped with the weak topology, then for two distributions \hat{Q} and \tilde{Q} that are close to each other in the weak sense, i.e., their corresponding distance of expected values satisfies $\mathbb{E}^{\hat{Q}}(f(U_1))$ – $\mathbb{E}^{\tilde{Q}}(f(U_1))| < \varepsilon$ for arbitrarily small $\varepsilon > 0$ and any bounded continuous real function $f(\cdot)$, the portfolios $b^*(\hat{Q})$ maximizing $\mathbb{E}^{\hat{Q}}(\log \langle b, U_1 \rangle)$ and $b^*(\tilde{Q})$ maximizing $\mathbb{E}^{\tilde{Q}}(\log \langle b, U_1 \rangle)$ must satisfy $(\log \langle b^*(\hat{\mathcal{Q}}), U_1 \rangle - \log \langle b^*(\tilde{\mathcal{Q}}), U_1 \rangle) \to 0$ as $\epsilon \to 0$ for $\hat{\mathcal{Q}}$ -almost all and also $\tilde{\mathcal{Q}}$ -almost all *U*1. Therefore, by the assumption for the market process, since the regular conditional distribution $\mathbb{P}(U_1|\mathscr{F}_n^X)$ converges weakly almost surely to the limiting distribution $\mathbb{P}(U_1|\mathscr{F}_\infty^X)$ by Lemma [4,](#page-9-0) we have $\left(\log \langle b_n^*, U_n \rangle - \log \langle b_\infty^*, U_n \rangle\right) \to 0$ for $\mathbb{P}\left(U_1 | \mathscr{F}_n^X\right)$ -almost all U_n , almost surely. Thus, the desired almost sure convergence $(W_n(b^{\omega}) - W_n(b_n^*)) \to 0$ follows with the chosen $b^{\omega} := b_{\infty}^*$ by applying the Cesaro mean theorem as in the proof of Theorem [1.](#page-12-0) Additionally, the L^1 convergence can be easily derived using the Lebesgue's dominated convergence theorem.

Alternatively, the almost sure convergence $(W_n(b^{\omega}) - W_n(b_n^*)) \rightarrow 0$ can be demonstrated more clearly than previously described by adapting the arguments from [Lam \(2024\)](#page-32-19) (Lemma 1 and Lemma 2) to the portfolio's log-return. Specifically, we sketch the proof as follows: Define ${b^*(Q)}$ as the set of all maximizers for the corresponding $\mathbb{E}^Q(\log \langle b, U_1 \rangle)$. Since the regular conditional distributions $\mathbb{P}(U_n|\mathscr{F}_n^X) \to \mathbb{P}(U_1|\mathscr{F}_\infty^X)$, the Hausdorff distance between the involved sets, denoted by $d_H\big(b^*\big(\mathbb{P}\big(U_n|\mathscr{F}_n^X\big)\big), b^*\big(\mathbb{P}\big(U_1|\mathscr{F}_\infty^X\big)\big)\big),$ must converge to zero as $n\to\infty,$ where the involved sets are convex and compact. Therefore, for each maximizer b_{∞}^{*} , there always exists a \max_{n} $\min_{n} b_{n}^{*}$ such that $(b_{n}^{*}-b_{\infty}^{*}) \to 0$, so $\left(\log \left\langle b_{n}^{*}, U_{n}\right\rangle - \log \left\langle b_{\infty}^{*}, U_{n}\right\rangle\right) \to 0$ for $\mathbb{P}\left(U_{1} | \mathscr{F}_{n}^{X}\right)$ -almost all U_n (noting critically that all U_n are bounded, as in [\(2.4\)](#page-6-1)), which holds for any b_n^* . \Box

Market with decaying impact of information. Although a real market often reflects memories of the past, sometimes the impact of past events does not last very long on future market developments, which is probably caused by the limited memory of agents and institutions in the market. In this kind of market, future assets' returns do not immediately incorporate all available information but rather gradually absorb it over time until the past impact has no influence. A market process with a Markov property of finite order should be well-suited for modeling the persistence of past impacts on the future. Let us consider the market process $\{X_n, Y_n\}_{n=1}^{\infty}$ as a Markov chain of order $h \geq 1$, so that the variable (X_n, Y_n) depends on the previous *h* variables, and its conditional distribution given the σ -field \mathcal{F}_n is:

$$
\mathbb{P}\big(X_n,Y_n|\mathscr{F}_n\big)=\mathbb{P}\big(X_n,Y_n|\sigma\big(X_{n-h}^{n-1},Y_{n-h}^{n-1}\big)\big),\forall n>h,
$$
\n(4.6)

where the order *h* represents the length of the market memory of past information. The length of the market memory is higher when the impact of past information decays at a slower rate, i.e., it is more persistent. If the market is also stationary, then Theorem [2](#page-13-0) reduces to Corollary [2.](#page-19-0)

Memoryless or past-independent market. In the case of the market having no memory of past events, the side information available at the same period, revealed right before the realization of assets' returns, is the only factor affecting the assets' returns. This means the market is highly efficient in the sense that it absorbs all available side information immediately and leaves no capacity for past information to impact future market developments. The memoryless nature of the market could be modeled as a special case of the market with a Markov property of order $h = 0$ in [\(4.6\)](#page-19-1), which implies the (joint) distributions of the variables (X_n, Y_n) are independent of each other. Moreover, as in the case of a Markov process with finite order, Theorem [2](#page-13-0) also reduces to Corollary [2](#page-19-0) if the distributions of the variables are identical.

Corollary 2. Consider a process $\{X_n, Y_n\}_{n=1}^{\infty}$ which is either identically and independently dis*tributed or a stationary Markov chain of any finite order* $h \geq 1$ *according to [\(4.6\)](#page-19-1). Then, there* exists either a non-random or random optimal constant strategy (b^*) , respectively, such that $(W_n(b^*) - W_n(b_n^*)) \rightarrow 0$ *almost surely, which is also in* L^1 *under the no-trash assets condition.*

Proof. The result is simply a direct consequence of Theorem [2,](#page-13-0) taking into account that an i.i.d. process ${X_n, Y_n}_{n=1}^{\infty}$ is also stationary and ergodic, since the only invariant σ -field is the trivial one, i.e., $\mathscr{I} = {\Omega, \emptyset}$, so the constant optimal strategy (b^*) is non-random. 囗

According to Corollary [2,](#page-19-0) if the assets' returns depend solely on the side market information such that the random pairs are i.i.d., the non-random optimal constant strategy coincides with the log-optimal strategy, which is selected as a maximizer for $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U_1 \rangle)$. This result extends the well-known theorem for the i.i.d. process of sole assets' returns ${X_n}_{n=0}^{\infty}$ $\sum_{n=1}^{\infty}$ in the literature, which asserts the existence of an optimal constant strategy achieving the maximal limiting growth rate. Intuitively, the extended result means that the observation of i.i.d. side information, which affects the periodic assets' returns in a time-invariant dependency, almost surely does not improve decision-making over being unaware of the side information and dependence structure, in terms of growth rate over time; thus, even unobservable market features do not matter. However, if the side information is i.i.d., but the dependence structure between them and the assets' returns

changes over time, then the conditional distribution $\mathbb{P}(X_n | \sigma(Y_n))$ varies with *n*, and the results of Corollary [2](#page-19-0) are no longer valid due to the violated assumption.

Remark. In recent literature, [Bhatt et al. \(2023](#page-32-14)) propose a strategy construction using fully observable i.i.d. side market information, which can be considered from a different perspective for comparison. Additionally, [Cuchiero et al. \(2019](#page-32-15)) investigate a stationary and ergodic Markov chain (order $h = 1$) for the market process of only assets' returns, a sub-case of our model without side information. The authors considers the space of functions depending on assets' returns *X*, yielding a portfolio $b \in \mathcal{B}^m$, where there exists an optimal function generating a strategy with the maximal limiting growth rate, depending on the process $\{X_n\}_{n=1}^{\infty}$ $\sum_{n=1}^{\infty}$. They then propose a strategy by mixing all the functionally generated portfolios over time according to a fixed initial distribution on the function space, such that it approaches the maximal limiting growth rate of the strategy generated by the optimal function. However, this method of function mixing seems impractical. A closer inspection reveals that the log-optimal portfolios, defined as $\argmax_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U_1 \rangle | \sigma(X_0)),$ are optimal in the considered space of functions, as shown by Corollary [2.](#page-19-0) Moreover, the next section presents a learning algorithm that provides a practical way of mixing functions to create a strategy comparable to the log-optimal one for any finite-order $(h \ge 1)$ Markov or i.i.d. market process with side information.

5 Learning algorithms under unknown distributions

Although Theorem [2](#page-13-0) shows the existence of a random (or non-random, with additional ergodicity) optimal constant strategy, which does not rely on market events realized over time as the logoptimal strategy does, or on being conditioned on the infinitely past events, as the random optimal constant strategy in Theorem [1,](#page-12-0) it still depends on the knowledge of the conditional distribution based on the invariant events of the market process. As discussed in the previous section, the true ergodic mode of the stationary market process cannot be identified by observing any finite past information rather than the infinite past, thus an optimal constant strategy is also theoretically unidentified with any finite time horizon. Meanwhile, although the log-optimal strategy could detect the ergodic mode of the process, it requires knowing the infinite-dimensional distribution. However, the established existence of an optimal constant strategy clearly helps reduce the challenge of knowing such an infinite-dimensional distribution, which requires full observability of all market information, to a much easier and more feasible task. Therefore, in this section, two practical learning methods of algorithms are compared: one replicating the optimal constant strategy and the other replicating the log-optimal strategy. Additionally, the extra L^1 -convergence in the theorems is simply understood as a condition to also imply the optimality in terms of the expected growth rate of the proposed strategies, as in Corollary [1.](#page-16-0)

5.1 Market-knowledge-free learning algorithm

In a stationary market, since the random optimal constant strategy is invariant in time, it does not need to observe any side information or know the possibly latent dependence structures of the process. Although the optimal constant strategy cannot be identified beforehand, a learning algorithm-based dynamic strategy could gradually approach the growth rate of this constant strategy over time. Moreover, because this type of algorithm inherits the advantages of a constant strategy, it only requires learning from past assets' returns, which are always available and not costly, but not from any side information of past market events. It is worth noting that there are several algorithms in the literature for this problem, with different assumptions and complexities, which are not the focus of this paper, as surveyed in [van Erven](#page-33-0) et al. [\(2020](#page-33-0)). However, considering stationarity, the following two learning algorithms are proposed for strategy development.

Cover's Universal Strategy. The Universal strategy, originally proposed in [Cover](#page-32-4) [\(1991](#page-32-4), [1996\)](#page-32-8); [Cover and Thomas](#page-32-12) [\(2006](#page-32-12)); [Ordentlich and Cover](#page-32-9) [\(1996](#page-32-9)), and extended with side information in [Cover and Ordentlich](#page-32-10) [\(1996](#page-32-10)), aims to construct a strategy that grows asymptotically at the same rate as the best strategy within the restricted set of constant strategies, for each finite sequence of realizations of assets' returns, from the starting point to the last observations. This learning algorithm is model-free and does not impose any assumptions on the market process. Consequently, the growth rate of the Universal strategy will approach that of the log-optimal strategy as time progresses, provided that there exists an optimal constant strategy comparable to the log-optimal one in the market. Moreover, it is important to note that Cover and many subsequent works in the literature emphasize that this strategy will almost surely attain the optimal growth rate only in an i.i.d. market process of sole assets' returns, which is trivially a special case of our more general theorem. Hence, our paper extends the optimality of the Universal strategy to broader scenarios involving markets with side information, far beyond the simple i.i.d. process.

In detail, the construction of the Universal portfolio strategy is as follows:

$$
\bar{b}_1 := (1/m, ..., 1/m) \text{ and } \bar{b}_n := \frac{\int_{\mathcal{B}^m} bS_{n-1}(b)\mu(b) \, db}{\int_{\mathcal{B}^m} S_{n-1}(b)\mu(b) \, db}, \forall n \ge 2,
$$

where $\mu(\cdot)$ denotes the fixed uniform density. Essentially, the algorithm distributes the current cumulative wealth according to the previous weights of cumulative wealth of all constant strategies, thereby learning only from the past portfolio returns of these component strategies. Since the algorithm operates without additional assumptions on assets' returns, the Universal portfolio can be constructed directly using the variables X_n , rather than their normalizations. However, the main drawback of this algorithm is its computational difficulty in practice, particularly when the portfolio involves a large number of invested assets (for example, $m \geq 3$), which motivates the development of a simpler algorithm with the same optimality, as described next.

Empirical distribution-based log-optimal strategy. This algorithm is naturally developed from the argument in the proof of Theorem [2,](#page-13-0) but with an extension to include the market with non-decaying impact of information. In general, this learning algorithm is much simpler than the Universal strategy, as it constructs the strategy by sequentially solving for the log-optimal portfolios corresponding to the empirical distribution of realizations of assets' returns. Despite the algorithm's simplicity, such an empirical strategy, as proposed below, is comparable with the Universal strategy in general markets, as long as there exists an optimal constant strategy that is comparable to the log-optimal one, as demonstrated in Theorem [3.](#page-22-0)

Specifically, we introduce the dynamic strategy (\bar{b}_n) constructed as follows:

$$
\bar{b}_1 := (1/m, ..., 1/m) \text{ and } \mathbb{E}^{P_{n-1}}(\log \langle \bar{b}_n, U \rangle) := \max_{b \in \mathcal{B}^m} \mathbb{E}^{P_{n-1}}(\log \langle b, U \rangle), \forall n \ge 2,
$$
 (5.1)

where $\mathbb{E}^{P_n}(\cdot)$ denotes the expected value corresponding to $U \sim P_n$, which is defined as:

$$
P_n(A) := \frac{1}{n} \sum_{i=1}^n \mathbb{I}_{U_i}(A), \forall A \subseteq \mathcal{U}, \forall n,
$$
\n(5.2)

i.e., an empirical distribution of the sequence of distributions $\{P_n\}_{n=1}^{\infty}$. Noting that the empirical

distributions are computed simply using the normalized realizations of assets' returns.

Remark. In the literature, a similar learning algorithm with a comparable mechanism is proposed by [Morvai](#page-32-5) [\(1991](#page-32-5)). However, this paper only proposes a strategy that captures the growth rate of the log-optimal strategy in a stationary and ergodic market without side information, assuming integrability. Moreover, the strategy is stated to be optimal only if the market is i.i.d., as concluded by the author based on Cover's assessment. In comparison, Theorem [3](#page-22-0) demonstrates the optimality of the empirical distribution-based log-optimal strategy in a more general market, without requiring stationarity, and provides a simpler proof that does not rely on the integrability assumption.

Lemma 6. Consider a sequence of positive real values $\{a_n\}_{n=1}^{\infty}$. If the following holds:

$$
\lim_{n\to\infty}\big(\sum_{i=1}^n a_i - n\big(\prod_{i=1}^n a_i\big)^{1/n}\big) = 0,
$$

 \int *then* $\max_{1 \leq i, j \leq n} |a_i - a_j| \to 0$ *as* $n \to \infty$.

Proof. Let denote $\delta_n := (\sqrt{a_{i_n}} - \sqrt{a_{j_n}})^2 := \max_{1 \le i, j \le n} (\sqrt{a_i} - \sqrt{a_j})^2$ for all *n*, which are strictly positive if there exist two distinct terms in the finite sequence $\{a_i\}_{i=2}^n$. Then we have:

$$
\sum_{i=1}^{n} a_i - n \left(\prod_{i=1}^{n} a_i \right)^{1/n} = \delta_n + \sum_{i \notin \{i_n, j_n\}} a_i + 2\sqrt{a_{i_n} a_{j_n}} - n \left(\prod_{i=1}^{n} a_i \right)^{1/n}
$$

$$
\geq \delta_n + n \left(\prod_{i \notin \{i_n, j_n\}} a_i a_{i_n} a_{j_n} \right)^{1/n} - n \left(\prod_{i=1}^{n} a_i \right)^{1/n} = \delta_n,
$$

due to the AM-GM inequality. Moreover, since $(\sqrt{a_i} - \sqrt{a_j})^2 \ge |a_i - a_j|$ for all $i, j \in \{1, ..., n\}$, we consequently obtain the following limit:

$$
0 = \lim_{n \to \infty} \left(\sum_{i=1}^{n} a_i - n \left(\prod_{i=1}^{n} a_i \right)^{1/n} \right) \ge \lim_{n \to \infty} \max_{1 \le i, j \le n} \left(\sqrt{a_i} - \sqrt{a_j} \right)^2
$$

$$
\ge \lim_{n \to \infty} \max_{1 \le i, j \le n} |a_i - a_j| = 0,
$$

 \Box

which is the desired result and completes the proof.

Theorem 3. Assume that there exists an optimal constant strategy (b^{ω}) , which can be either *random or non-random, in the market process* $\{X_n, Y_n\}_{n=1}^{\infty}$ *(not necessarily stationary), such that* $(W_n(b^{\omega}) - W_n(b_n^*)) \rightarrow 0$ *almost surely. Then, the strategy* (b_n) *, constructed according to* [\(5.1\)](#page-21-0) *and [\(5.2\)](#page-21-1), is also optimal in terms of asymptotic growth rate, as the log-optimal strategy, satisfying:*

$$
\lim_{n\to\infty}\big(W_n(\bar{b}_n)-W_n(b_n^*)\big)=\lim_{n\to\infty}\big(W_n(\bar{b}_n)-W_n(b^\omega)\big)=0,\ a.s,
$$

*which are also in L*¹ *under the additional condition of no-trash assets. As a consequence, the* \bar{s} strategy (\bar{b}_n) is not only optimal in a stationary general market, but also in one with non-decaying *impact of past information on future market events, as described in Proposition [1.](#page-18-0)*

Proof. First, consider the case of a stationary market and provide a specific proof for the theorem statement. Recall the steps in [\(4.4\)](#page-15-0) of the proof of Theorem [2,](#page-13-0) we have:

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\left(\log\langle b^\omega,U_i\rangle-\log\langle b_i^*,U_i\rangle\right)=\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\left(\log\left\langle b^{*,n},U_i\right\rangle-\log\left\langle b_i^*,U_i\right\rangle\right)=0,
$$

where we define the constant strategy $(b^{*,n})$ as the portfolio $b^{*,n}$ that maximizes the following expectation with respect to the empirical distribution P_n of the normalized assets' returns U :

$$
\mathbb{E}^{P_n}(\log \langle b^{*,n},U \rangle) = \max_{b \in \mathscr{B}^m} \mathbb{E}^{P_n}(\log \langle b,U \rangle),
$$

with the empirical distribution P_n defined according to [\(5.2\)](#page-21-1) for all n . To avoid confusion, it should be noted that the strategy (\bar{b}_n) is dynamic, while the strategies $(b^{*,n})$ are constant and depend on P_n , which are denoted here for convenience in later deductions.

Next, we deduce the weak convergence of the empirical distribution sequence as $P_n \to \mathbb{P}(U_1|\mathscr{I})$ almost surely, due to Birkhoff's ergodic theorem. This results, almost surely, in that $\langle \bar{b}_n, U_1 \rangle \rightarrow$ $\langle b^*, U_1 \rangle$ for $\mathbb{P}(U_1|\mathscr{I})$ -almost all U_1 , with b^* maximizing $\mathbb{E}(\log \langle b, U_1 \rangle | \mathscr{I}) = \mathbb{E}(\log \langle b, U_k \rangle | \mathscr{I})$ for all $k > 1$, by Lemma [4.](#page-9-0) Thus, we also have $\left(\log \langle \bar{b}_n, U_k \rangle - \log \langle b^*, U_k \rangle\right) \to 0$ for all *k* almost surely, so $\left(\log \langle \bar{b}_n, U_n \rangle - \log \langle b^*, U_n \rangle\right) \to 0$ also holds almost surely, by Lemma [5.](#page-12-1) The needed equality is then obtained as follows, due to Cesaro's mean theorem and Theorem [2:](#page-13-0)

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\left(\log\left\langle\bar{b}_i,U_i\right\rangle-\log\left\langle b^*,U_i\right\rangle\right)=\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\left(\log\left\langle\bar{b}_i,U_i\right\rangle-\log\left\langle b^*_n,U_i\right\rangle\right)=0, \text{ a.s.}
$$

Hence, under the additional condition of no-trash assets, the almost sure convergence above is also in L^1 using Lebesgue's dominated convergence theorem.

(General case without the strict stationarity condition). If there exists a random optimal constant strategy but the market is not stationary, then the weak convergence of the empirical distribution $P_n \to \mathbb{P}(U_1|\mathscr{I})$ is not guaranteed, so the following arguments of the aforementioned proof fail. Instead, by utilizing the properties of the log-optimal portfolios, we can argue that the optimal constant strategy should also be almost optimal with respect to the empirical distribution P_n as $n \to \infty$. Indeed, this property can be established using Lemma [6](#page-22-1) through the following points.

Point 1. Since there exists a random optimal constant strategy (b^{ω}) by assumption, we have:

$$
0 = \lim_{n \to \infty} \left(\frac{1}{n} \sum_{i=1}^{n} \log \left\langle b^{\omega}, U_{i} \right\rangle - \max_{b \in \mathcal{B}^{m}} \frac{1}{n} \sum_{i=1}^{n} \log \left\langle b, U_{i} \right\rangle \right) = \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \left(\log \frac{\left\langle b^{\omega}, U_{i} \right\rangle}{\left\langle b^{*,n}, U_{i} \right\rangle} \right)
$$

$$
\leq \lim_{n \to \infty} \log \left(\frac{1}{n} \sum_{i=1}^{n} \frac{\left\langle b^{\omega}, U_{i} \right\rangle}{\left\langle b^{*,n}, U_{i} \right\rangle} \right)
$$

$$
= \lim_{n \to \infty} \log \mathbb{E}^{P_{n}} \left(\frac{\left\langle b^{\omega}, U \right\rangle}{\left\langle b^{*,n}, U \right\rangle} \right) \leq 0,
$$

due to Jensen's inequality and Lemma [2,](#page-7-0) taking into account that the portfolios $\bar{b}_{n+1} = b^{*,n}$ are the maximizers for $\max_{b \in \mathcal{B}^m} \mathbb{E}^{P_n}(\log \langle b, U \rangle)$ corresponding to the empirical distributions P^n . As a result, the obtained equality implies:

$$
\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \frac{\langle b^{\omega}, U_i \rangle}{\langle b^{*,n}, U_i \rangle} = \lim_{n \to \infty} \left(\prod_{i=1}^{n} \frac{\langle b^{\omega}, U_i \rangle}{\langle b^{*,n}, U_i \rangle} \right)^{1/n} = 1,
$$
\n(5.3)

which immediately leads to the following consequence by Lemma [6:](#page-22-1)

$$
\lim_{n \to \infty} \max_{1 \le i,j \le n} \left| \frac{\langle b^{\omega}, U_i \rangle}{\langle b^{*,n}, U_i \rangle} - \frac{\langle b^{\omega}, U_j \rangle}{\langle b^{*,n}, U_j \rangle} \right| = 0.
$$
\n(5.4)

This limit holds only if all terms of the sequence $\left\{ \frac{\langle b^{\omega}, U_i \rangle}{\langle b^{*,n}, U_i \rangle} \right\}_{i=1}^n$ converge uniformly to the same value 1 as $n \rightarrow \infty$. Indeed, since the equality [\(5.4\)](#page-23-0) also implies that:

$$
\lim_{n\to\infty}\Big(\max_{1\leq i\leq n}\Big\{\frac{\langle b^{\omega},U_i\rangle}{\langle b^{*,n},U_i\rangle}\Big\}-\min_{1\leq i\leq n}\Big\{\frac{\langle b^{\omega},U_i\rangle}{\langle b^{*,n},U_i\rangle}\Big\}\Big)=0,
$$

thus, by using the convergence in [\(5.3\)](#page-23-1), we obtain the needed result:

$$
1 = \lim_{n \to \infty} \min_{1 \le i \le n} \left\{ \frac{\langle b^{\omega}, U_i \rangle}{\langle b^{*,n}, U_i \rangle} \right\} \le \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^n \frac{\langle b^{\omega}, U_i \rangle}{\langle b^{*,n}, U_i \rangle} \le \lim_{n \to \infty} \max_{1 \le i \le n} \left\{ \frac{\langle b^{\omega}, U_i \rangle}{\langle b^{*,n}, U_i \rangle} \right\} = 1.
$$

Hence, $\left(\log \left\langle b^{\omega}, U_n \right\rangle - \log \left\langle b^{*,n}, U_n \right\rangle \right) \to 0$ as $n \to \infty$, due to $\left(\log \left\langle b^{\omega}, U_i \right\rangle - \log \left\langle b^{*,n}, U_i \right\rangle \right) \to 0$ uniformly for all $i \in \{1, ..., n\}$ as $n \to \infty$. In order to conclude the convergence of the strategy (\bar{b}_n) , we need to establish the following additional property.

Point 2. Noting that although the convergence of the sequence of distributions ${P_n}_{n=1}^{\infty}$ is not guaranteed, we still have the following convergence for all bounded continuous functions $f(\cdot)$:

$$
\lim_{n \to \infty} (\mathbb{E}^{P_{n+1}}(f(U)) - \mathbb{E}^{P_n}(f(U))) = \lim_{n \to \infty} \left(\frac{1}{n+1} \sum_{i=1}^{n+1} f(U_i) - \frac{1}{n} \sum_{i=1}^{n} f(U_i) \right)
$$

$$
= \lim_{n \to \infty} \left(\frac{1}{n+1} f(U_{n+1}) - \frac{1}{n(n+1)} \sum_{i=1}^{n} f(U_i) \right) = 0.
$$

Therefore, by Lemma [4,](#page-9-0) the function $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U \rangle)$ is continuous over the space of distributions equipped with the weak topology. This leads to the following:

$$
\lim_{n\to\infty}\big(\max_{b\in\mathscr{B}^m}\mathbb{E}^{P_{n+1}}\big(\log\big\langle b,U\big\rangle\big)-\max_{b\in\mathscr{B}^m}\mathbb{E}^{P_n}\big(\log\big\langle b,U\big\rangle\big)\big)=0.
$$

Moreover, Lemma [4](#page-9-0) also implies that if the log-optimal portfolios for $\max_{b \in \mathcal{B}^m} \mathbb{E}^{P_n}(\log \langle b, U \rangle)$ are unique for all *n*, then $(b^{*,n} - b^{*,n-1}) \to 0$ as $n \to \infty$. Additionally, recalling that this lemma further asserts that $\langle \hat{b}, U \rangle = \langle \bar{b}, U \rangle$ for any two maximizers \hat{b} and \bar{b} for max $_{b \in \mathcal{B}^m} \mathbb{E}^{P_n}(\log \langle b, U \rangle)$ for all P_n , the following convergence always holds:

$$
\lim_{n\to\infty}\left(\log\left\langle b^{*,n},U_n\right\rangle-\log\left\langle b^{*,n-1},U_n\right\rangle\right)=0,
$$

taking into account that all the normalized assets' returns $U_n \in \mathcal{U}$ are bounded for all *n*, as stated in [\(2.4\)](#page-6-1) in the proof of Lemma [1.](#page-5-0)

(Definitive statement of the optimality). Finally, we establish the convergence for the strategy (\bar{b}_n) , where $\bar{b}_n = b^{*,n-1}$ by construction of the strategy:

$$
\lim_{n\to\infty} \left(\log \left\langle b^{\omega}, U_n \right\rangle - \log \left\langle \bar{b}_n, U_n \right\rangle \right) = \lim_{n\to\infty} \left(\log \left\langle b^{\omega}, U_n \right\rangle - \log \left\langle b^{*,n}, U_n \right\rangle \right) \n+ \lim_{n\to\infty} \left(\log \left\langle b^{*,n}, U_n \right\rangle - \log \left\langle b^{*,n-1}, U_n \right\rangle \right) = 0.
$$

Then, invoking the Cesaro's mean theorem again, we obtain $(W_n(\bar{b}_n) - W_n(b_n^*)) \to 0$ as:

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\left(\log\left\langle b^\omega,U_i\right\rangle-\log\left\langle\bar{b}_n,U_i\right\rangle\right)=\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\left(\log\left\langle b_i^*,U_i\right\rangle-\log\left\langle\bar{b}_n,U_i\right\rangle\right)=0.
$$

This results in L^1 -convergence by the Lebesgues' dominated convergence theorem under the additional condition of no-trash assets. Thus, the proof is completed. \Box

5.2 Information-observation-based learning algorithm

After establishing the asymptotic optimality of the log-optimal strategy in [Algoet and Cover \(1988\)](#page-32-0), a problem arose concerning the construction of a strategy that asymptotically approaches the same finite growth rate as the log-optimal strategy, under a stationary and ergodic market with an unknown infinite-dimensional distribution. This problem was posed in Algoet's later work [Algoet](#page-32-6) [\(1992](#page-32-6)), where it was referred to as the desired universality of a so-called optimal strategy. In addition to formulating the problem, Algoet also sketched a learning algorithm to construct a strategy that could achieve the target limiting growth rate, based on the learning algorithms developed in the thesis of [Bailey \(1976](#page-32-20)) and later by [Ornstein \(1978\)](#page-32-21). However, this algorithm was known for its practical difficulties until the work by [Gyorfi et al.](#page-32-11) [\(2006](#page-32-11)), which made the original idea more practical. Since then, variants of Gyorfi's learning algorithm (a collective term for Gyorfi and his co-authors) have been regarded as the primary class of methods that fully implement Algoet's idea in contemporary literature. In this section, we extend the investigation of the properties of this method and enhance its practicality, especially when the market is stationary but not ergodic, with

side information and the guaranteed existence of an optimal constant strategy.

Anatomy of Algoet's and Gyorfi's learning algorithms. As mentioned in the previous section, since the true ergodic mode of a stationary process and the associated conditional distribution cannot be exactly identified from any finite observation of past events, a universal estimation scheme is formulated by learning the realizations of the process to approximate the true conditional distribution given the infinite past. In the paper [Algoet](#page-32-6) [\(1992](#page-32-6)), the author generalizes the universal estimation scheme for finite-valued processes developed by [Bailey](#page-32-20) [\(1976](#page-32-20)) and [Ornstein \(1978](#page-32-21)) to the Polish space, which guarantees the asymptotic optimal growth rate for a log-optimal strategy derived using the estimated conditional distribution. However, due to the complexity of its estimation, Algoet proposes another simpler algorithm that generates log-optimal portfolios based on conditional distributions estimated using finite-order Markov approximation functions, then combines them using the *bookkeeping technique*, commonly referred to as the *exponentially weighted average* (as in [Cesa-Bianchi and Lugosi](#page-32-13) [\(2006](#page-32-13))) in the present literature. The main practical issue with this algorithm is that the Markov approximations only capture similar events belonging to the same atoms in a σ -field, so the learning process often results in void sets. To address this problem, Gyorfi's algorithm uses the same mechanism, but the combination procedure is conducted with finite-order Markov approximations that capture various distances between two events, rather than requiring absolute equality.

Finite-order Markov approximation and the estimated log-optimal portfolio. Despite the improved practicality, it should be stressed that there are two inherent challenges in both the original Algoet's algorithm and the later Gyorfi's algorithm: first, they require complete observation of all available market side information, even though parts of it may be unobservable, which [Algoet](#page-32-6) [\(1992](#page-32-6)) explicitly mentions as a conjectured result for a possible extension of the algorithm to markets with side information 2 ; second, the Markov approximation must be taken to an infinite order, rendering the available data insufficient for the learning procedure to guarantee convergence of the estimation. Hence, although the optimality of the estimated log-optimal strategy is theoretically guaranteed, it seems unrealistic in practice. However, based on the results obtained in the previous section, we can expect some possible simplifications to the algorithm.

Without loss of generality, by assuming that all market side information is observable, consider the following estimated log-optimal strategy $(b_n^{(h,l)}(\theta))$ based on a Markov approximation of order *h* with respect to normalized assets' returns, and a parameter vector $\theta \in \mathbb{R}_{++}^{m \times (h-1)} \times \mathbb{R}^{k \times h}$:

$$
b_n^{(h,l)}(\boldsymbol{\theta}) \coloneqq \underset{b \in \mathcal{B}^m}{\operatorname{argmax}} \frac{1}{|\mathcal{D}_n^{(h,l)}(\boldsymbol{\theta})|} \sum_{X_i \in \mathcal{D}_n^{(h,l)}(\boldsymbol{\theta})} \log \langle b, U_i \rangle, \forall n \ge h+2,
$$
 (5.5)

which implies that $b_n^{(h,l)}(\theta)$ maximizes $\mathbb{E}^{P(\cdot|\mathscr{D}_n^{(h,l)}(\theta))}(\log \langle b, u(X) \rangle)$ for all $n > h$, where:

$$
X \sim P(\cdot | \mathscr{D}_n^{(h,l)}(\boldsymbol{\theta})), \text{ and } P(A | \mathscr{D}_n^{(h,l)}(\boldsymbol{\theta})) \coloneqq \frac{1}{|\mathscr{D}_n^{(h,l)}(\boldsymbol{\theta})|} \sum_{X_i \in \mathscr{D}_n^{(h,l)}(\boldsymbol{\theta})} \mathbb{I}_{X_i}(A), \forall A \subset \mathbb{R}_{++}^m,
$$

and $h, l \in \mathbb{N}_+$, while $c \in \mathbb{R}_{++}$ is a fixed small value. Also,

$$
\mathscr{D}_n^{(h,l)}(\theta) := \{ X_i : \left\| \left(X_{i-h}^{i-1}, Y_{i-h}^i \right) - \theta \right\| \le c/l, \, h+1 \le i \le n-1 \}, \, \forall n \ge h+2, \tag{5.6}
$$

 $\text{with } b_n := (1/m, ..., 1/m) \text{ and } P(\cdot | \mathscr{D}_n^{(h,l)}(\theta)) := \delta_{(1, ..., 1)}(\cdot)$ (Dirac measure) for an empty $\mathscr{D}_n^{(h,l)}(\theta)$ by convention.

²See Remark 2 in Section 4.4 for a discussion of the sufficiency of using side information for conditional distribution estimation, and Remark 2 in Section 5.2 for the conjectured result for a stationary and ergodic market.

Since the market process is assumed to be stationary, an investor could start investing after *h* periods, which is equivalent to moving the origin, and all estimated log-optimal portfolios will be formed from the first period onward. The sets $\mathcal{D}_n^{(h,l)}(\theta)$ represent collections of asset returns having finite sequences of past asset returns with similar patterns, measured by a norm (such as the Frobenius norm). Consider $\theta = (X_{n-h}^{n-1}, Y_{n-h}^n)$ as an instance. The norm of similarity between two events is computed using matrices as follows:

$$
\left\| (X_{i-h}^{i-1}, Y_{i-h}^i) - (X_{n-h}^{n-1}, Y_{n-h}^n) \right\| = \left\| \left[\begin{array}{cccc} Y_{i-h} & \cdots & Y_{i-1} & Y^i \\ X_{i-h} & \cdots & X_{i-1} & 0 \end{array} \right] - \left[\begin{array}{cccc} Y_{n-h} & \cdots & Y_{n-1} & Y^n \\ X_{n-h} & \cdots & X_{n-1} & 0 \end{array} \right] \right\|.
$$

Additionally, the value *c* is fixed, and $c/l \to 0$ as $l \to \infty$, which partitions the σ -field $\sigma(X_{n-h}^{n-1}, Y_{n-h}^n)$ into coarser approximation σ -fields for each level *c*/*l*. As *l* $\rightarrow \infty$, the approximation becomes finer, and all similar events must belong to the same atoms of the σ -field $\sigma(X_{n-h}^{n-1}, Y_{n-h}^n)$. Furthermore, it is worth noting that there are various methods to measure the similarity between events, as discussed in [Gyorfi et al. \(2006\)](#page-32-11). However, for representational purposes, we focus on the partition in [\(5.6\)](#page-25-1), which is commonly referred to as the *kernel-based criterion* in the literature, due to its empirically outstanding performance and implementability.

Strategy constructed by mixing kernels with fixed order. The mechanism of measuring similarity between events across several extents *c*/*l* allows the partitions to collect more elements during the scanning process of the past, thereby avoiding the empty sets $\mathscr{D}_n^{(h,l)}(\theta)$. This is especially important since the amount of available data is limited when the time horizon is small. Hence, similar to Gyorfi's argument, we construct the following combinatorial strategy by exponentially averaging the weighted strategies $(b_n^{(h,l)}(\theta_n))$ corresponding to all values of *l* ranging over \mathbb{N}_+ , where $\theta_n \coloneqq (X_{n-h}^{n-1}, Y_{n-h}^n)$, as:

$$
b_n^{(h)} := (1/m, ..., 1/m), \forall n < h+2, \text{ and } b_n^{(h)} := \frac{\sum_l b_n^{(h,l)}(\theta_n) S_{n-1}\left(b_{n-1}^{(h,l)}(\theta_{n-1})\right)}{\sum_l S_{n-1}\left(b_{n-1}^{(h,l)}(\theta_{n-1})\right)}, \forall n \ge h+2 \tag{5.7}
$$

Noting that the combination in [\(5.7\)](#page-26-0) fixes an *h*, which differs from the approach in Algoet's and Gyorfi's algorithms, which combine all orders of *h* over \mathbb{N}_+ . This distinction arises because the algorithm cannot learn from past data when $h > n$, and thus, the theoretically guaranteed optimality is not attainable in practice. This limitation leads to the restriction of choosing order *h* in the experiments in [Gyorfi et al.](#page-32-11) $(2006)^3$ $(2006)^3$ $(2006)^3$.

The asymptotic growth rate and optimality of the strategy $(b_n^{(h)})$ are established in Theorem [4](#page-27-0) with an arbitrarily fixed order *h*. The theorem asserts that the choice of order does not affect the optimality of the strategy $(b_n^{(h)})$, and therefore, there is no need to mix all orders *h* over \mathbb{N}_+ . Moreover, this result holds even if the algorithm is restricted to learning only from past asset returns. In this context, the limiting growth rate of the strategy $(b_n^{(h)})$ is given by $\mathbb{E}(\max_{b \in \mathscr{B}^m} \mathbb{E}(\log \langle b, U_1 \rangle | \sigma(X^0_{1-h})) | \mathscr{I})$ under a stationary but not ergodic market process. This statement is easily justified following the final derivation for the optimality of the strategy $(b_n^{(h)})$ in the proof of Theorem [4.](#page-27-0) Based on this property, it is safe to remove the side information from the learning procedure of the algorithm.

Notes on mixing kernels and markets with decaying impact of information. In the case

 3 In the remark on the "Validity of assumptions" and the second paragraph of Section 4.1 of the cited article, the authors note that certain component strategies with smaller order *h* perform better than others. Therefore, *h* can be restricted, as the market may be a low-order Markov process. Moreover, such low-order Markov component strategies effectively exploit the hidden dependence structure of the market, which is difficult to reveal. Clearly, this is the primary motivation of our paper: to address the challenge of unobservable latent market dependence.

where the stationary market is either an i.i.d. or a finite-order Markov process, with $h^* \geq 0$ denoting the true order of the market's memory, the strategy $(b_n^{(h)})$ must be constructed with a chosen $h \geq h^*$ to activate the Markov property during the learning procedure, thereby enabling Theorem [4](#page-27-0) with stationarity. In this market scenario, the combinatorial strategy $(b_n^{(h^*)})$ with true order h^* , learns the log-optimal strategy (b_n^*) , which is a sequence of functionally generated portfolios b_n^* defined by the function $\operatorname{argmax}_{b \in \mathcal{B}^m} \mathbb{E} \left(\log \langle b, U_1 \rangle | \sigma \left(X_{1-h}^0, Y_{1-h}^1 \right) \right)$ for all *n*. Let us recall that this result generalizes the work of [Cuchiero et al.](#page-32-15) [\(2019](#page-32-15)) for the same type of market, as mentioned in the last remark of the previous section. However, the strategy $(b_n^{(h^*)})$ is a mixing of estimated log-optimal strategies based on *h*^{*}-order Markov approximation functions with different kernel widths, rather than mixing all functions over their entire space, which seems impractical.

However, since the market's dependence structure is latent, the choice of order *h* for the strategy $(b_n^{(h^*)})$ should be large enough to cover all possibilities of the unknown true order h^* of the market's memory length. In practice, because the sets $\mathcal{D}_n^{(h,l)}(\theta)$ may become almost void if *h* is large, we should mix all orders within a threshold *H* to construct the strategy (\bar{b}_n) as follows:

$$
\bar{b}_n := (1/m, ..., 1/m), \forall n < h+2, \text{ and } \bar{b}_n := \frac{\sum_{h \leq H} b_n^{(h)} S_{n-1} \left(b_{n-1}^{(h)} \right)}{\sum_{h \leq H} S_{n-1} \left(b_{n-1}^{(h)} \right)}, \forall n \geq h+2.
$$

Then, due to the known property of the weighted exponential average (as demonstrated in [\(5.11\)](#page-29-0) of the proof of Theorem [4\)](#page-27-0), the growth rate of the combinatorial strategy (\bar{b}_n) in terms of normalized assets' returns will converge to that of the component strategy $(b_n^{(h^*)})$ with true order h^* . Finally, it should be noted that a critical weakness of the learning algorithm for the strategy $(b_n^{(h)})$ is that it requires $l \rightarrow \infty$, which makes the subroutine's computation very costly. Therefore, the marketknowledge-free learning algorithms described in the previous section, which capture the growth rate of the random optimal constant strategy, are generally still preferable.

Remark. The proof of Theorem [4](#page-27-0) follows a substantial portion of the arguments in the proof of [Gyorfi et al. \(2006](#page-32-11)), with a generalization to the case of a stationary but not ergodic market with side information, capturing a broader context where the optimal limiting growth rate of a strategy may be finitely random or not well-defined. Additionally, the theorem establishes the limiting growth rate for the strategy $(b_n^{(h)})$ with a fixed order *h*, in terms of normalized assets' returns, rather than mixing all orders *h*.

Theorem 4. If the market process $\{X_n, Y_n\}_{n=1}^{\infty}$ is stationary (not necessarily ergodic) and free of trash assets, then the strategy $(b_n^{(h)})$, defined according to [\(5.5\)](#page-25-2), [\(5.6\)](#page-25-1) and [\(5.7\)](#page-26-0), for any fixed $h > 0$ *, satisfies the following:*

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b_i^{(h)},U_i\right\rangle=\mathbb{E}\big(\log\left\langle b_{h+1}^*,U_1\right\rangle|\mathscr{I}\big),\ \ a.s\ and\ in\ L^1,
$$

where b_{h+1}^* is the \mathscr{F}_{h+1}^X -measurable log-optimal portfolio of the log-optimal strategy (b_n^*) . Therefore, the strategy $(b_n^{(h)})$ is also optimal in terms of asymptotic growth rate, as:

$$
\lim_{n\to\infty} (W_n(b_n^{(h)}) - W_n(b^*)) = \lim_{n\to\infty} (W_n(b_n^{(h)}) - W_n(b_n^*)) = 0, \text{ a.s and in } L^1.
$$

Proof. For the sake of readability, we divide the proof into smaller parts below.

(Letting time n increase to infinity). Firstly, recall that by stationarity, we have:

$$
\sigma(X_{n-h}^0, Y_{1-h}^1) = T^{-(n-1)} \sigma(X_{n-h}^{n-1}, Y_{n-h}^n), \forall n.
$$

Then, for any θ , we show that the sequence of distributions $\{P(\cdot | \mathcal{D}_n^{(h,l)}(\theta))\}_{n=1}^{\infty}$ converges

weakly almost surely to the following distributions as $n \to \infty$:

$$
\begin{cases} \mathbb{P}(X_1 | \|(X_{1-h}^0, Y_{1-h}^1) - \theta \| \le c/l, \mathcal{I}) & \text{if } \mathbb{P}(||(X_{1-h}^0, Y_{1-h}^1) - \theta \| \le c/l) > 0, \\ \delta_{(1, \dots, 1)} & \text{if } \mathbb{P}(||(X_{1-h}^0, Y_{1-h}^1) - \theta \| \le c/l) = 0. \end{cases}
$$
(5.8)

Indeed, consider any bounded continuous function $f(\cdot)$, we obtain:

$$
\lim_{n \to \infty} \mathbb{E}^{P(\cdot|\mathscr{D}_n^{(h,l)}(\theta))} f(X) = \lim_{n \to \infty} \frac{|h - n|}{|h - n| |\mathscr{D}_n^{(h,l)}(\theta)|} \sum_{X_i \in \mathscr{D}_n^{(h,l)}(\theta)} f(X_i)
$$

$$
= \frac{\mathbb{E}(f(X_1) \mathbb{I}_{||(X_{1-h}^0, Y_{1-h}^1) - \theta|| \le c/l} | \mathscr{I})}{\mathbb{E}(\mathbb{I}_{||(X_{1-h}^0, Y_{1-h}^1) - \theta|| \le c/l} | \mathscr{I})}
$$

$$
= \mathbb{E}(f(X_1) | || (X_{1-h}^0, Y_{1-h}^1) - \theta || \le c/l, \mathscr{I}), \text{ a.s.}
$$

 $\lim_{h \to 0} \frac{\ln \left(||(X_{1-h}^0, Y_{1-h}^1) - \theta|| \le c/l \right)}{h} > 0.$ Meanwhile, if $\mathbb{P}\left(||(X_{1-h}^0, Y_{1-h}^1) - s|| \le c/l \right) =$ 0, the sequence $\{P(\cdot | \mathcal{D}_n^{(h,l)}(\theta))\}_{n=1}^{\infty}$ is identical to the Dirac delta distribution $\delta_{(1,..,1)}$ almost surely, which implies $\mathbb{E}^{P(\cdot|\mathscr{D}_n^{(h,l)}(\theta))} f(X) = f(1,...,1)$ for all *n* almost surely.

(Letting extent l increase to infinity). As $l \rightarrow \infty$, the kernel width $c/l \rightarrow 0$, so we expect the approximated conditional distribution to converge to the true one as follows:

$$
\lim_{n \to \infty} \mathbb{P}(X_1 \mid \|(X_{1-h}^0, Y_{1-h}^1) - \theta \|\le c/l) = \mathbb{P}(X_1 \mid (X_{1-h}^0, Y_{1-h}^1) = \theta).
$$
\n(5.9)

This desired convergence is demonstrated as follows, for any $A \subset \mathbb{R}^m_{++}$ and θ :

$$
\lim_{l \to \infty} \mathbb{P}(X_l \in A \mid \left\| (X_{1-h}^0, Y_{1-h}^1) - \theta \right\| \le c/l) = \lim_{l \to \infty} \frac{\mathbb{P}(X_l \in A, \left\| (X_{1-h}^0, Y_{1-h}^1) - \theta \right\| \le c/l)}{\mathbb{P}(\left\| (X_{1-h}^0, Y_{1-h}^1) - \theta \right\| \le c/l)} \\
= \lim_{l \to \infty} \frac{\mathbb{E}(\mathbb{P}(X_l \in A | X_{1-h}^0, Y_{1-h}^1) - \theta \|\le c/l)}{\mathbb{E}(\mathbb{I}_{\left\| (X_{1-h}^0, Y_{1-h}^1) - \theta \right\| \le c/l)}} \\
= \mathbb{P}(X_l \in A | (X_{1-h}^0, Y_{1-h}^1) = \theta),
$$

for P-almost all $\theta \in \left\{ (X_{1-h}^0, Y_{1-h}^1)(\Omega) \right\}$ by Lebesgue's density theorem.

(Deriving the first needed inequality). Let the portfolio $b_{\infty}^{(h,l)}(\theta)$ denote a maximizer for:

$$
\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b(\theta),U_1\big\rangle|\,\big\|\big(X_{1-h}^0,Y_{1-h}^1\big)-\theta\big\|\leq c/l,\mathscr{I}\big),
$$

 $\inf_{(h, h) \in \mathcal{L}} \max_{t \in [h]} \mathbb{E} \left(||(X_{1-h}^0, Y_{1-h}^1) - \theta|| \le c/l \right) > 0.$ Otherwise, if $\mathbb{P} \left(||(X_{1-h}^0, Y_{1-h}^1) - \theta|| \le c/l \right) > 0$, let $b_{\infty}^{(h,l)}(\theta) \coloneqq (1,...,1)$.

Then, fixing *l*, due to the convergence of distributions in [\(5.8\)](#page-28-0) as $n \to \infty$, we obtain almost surely the convergence $\log \langle b_n^{(h,l)}(\theta), U_1 \rangle \to \log \langle b_\infty^{(h,l)}(\theta), U_1 \rangle$ using Lemma [\(4\)](#page-9-0). Hence, by applying Breiman's generalized ergodic theorem, we get:

$$
\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_i^{(h,l)}(\theta_i), U_i \rangle = \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_i^{(h,l)}(X_{i-h}^{i-1}, Y_{i-h}^i), U_i \rangle
$$

\n
$$
= \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_{\infty}^{(h,l)}(T^{(i-1)}(X_{1-h}^0, Y_{1-h}^1)), T^{(i-1)}U_1 \rangle
$$

\n
$$
= \mathbb{E} \big(\log \langle b_{\infty}^{(h,l)}(X_{1-h}^0, Y_{1-h}^1), U_1 \rangle | \mathcal{J} \big), \text{ a.s and in } L^1,
$$
 (5.10)

given that $\theta_n := (X_{n-h}^{n-1}, Y_{n-h}^n)$ as defined for the strategy $(b_n^{(h)})$ in [\(5.7\)](#page-26-0), and that the required L^1 domination is satisfied for the sequence $\left\{ \log \left\langle b_n^{(h,l)}, U_n \right\rangle \right\}_{n=1}^{\infty}$ due to [\(4.1\)](#page-14-0), under the assumption of no-trash assets.

Consequently, by letting $l \rightarrow \infty$, we obtain the convergence of distributions in [\(5.9\)](#page-28-1) for $\theta =$

 (X_{1-h}^0, Y_{1-h}^1) . Hence, by Lemma [4](#page-9-0) again, we have the following convergence:

$$
\lim_{n\to\infty}\log\left\langle b^{(h,l)}_{\infty}\left(X^0_{1-h},Y^1_{1-h}\right),U_1\right\rangle=\log\left\langle b^{(h,\infty)}_{\infty}\left(X^0_{1-h},Y^1_{1-h}\right),U_1\right\rangle, \text{ a.s.}
$$

where $b_{\infty}^{(h,\infty)}(X_{1-h}^0, Y_{1-h}^1)$ denotes a maximizer for $\max_{b \in \mathcal{B}^m} \mathbb{E}(\log \langle b, U_1 \rangle | \sigma(X_{1-h}^0, Y_{1-h}^1))$. Noting that this limit also holds for $\mathbb{P}\big(U_1|\mathscr{I}\big)$ -almost all U_1 , almost surely. Next, taking the limit as $l\to\infty$ of both sides of [\(5.10\)](#page-28-2), we obtain:

$$
\lim_{l \to \infty} \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_i^{(h,l)}(\theta_i), U_i \rangle = \lim_{l \to \infty} \mathbb{E} \big(\log \langle b_{\infty}^{(h,l)}(X_{1-h}^0, Y_{1-h}^1), U_1 \rangle | \mathcal{I} \big)
$$

= $\mathbb{E} \big(\log \langle b_{\infty}^{(h,\infty)}(X_{1-h}^0, Y_{1-h}^1), U_1 \rangle | \mathcal{I} \big)$
= $\mathbb{E} \big(\log \langle b_{h+1}^*, U_1 \rangle | \mathcal{I} \big), \text{ a.s.}$

where the last equality follows from the fact that $b_{\infty}^{(h,\infty)}(X_{1-h}^0,Y_{1-h}^1)(T^h\omega) = b_{h+1}^*(\omega)$, which is the measurable log-optimal portfolio corresponding to the σ -field $\mathscr{F}_{h+1}^X = T^h \sigma(X_{1-h}^0, Y_{1-h}^1)$ due to the stationarity of the market process. Then, by telescoping, we obtain the following:

$$
\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_n^{(h)}, U_i \rangle = \lim_{n \to \infty} \frac{1}{n} \log \left(\frac{1}{l} \sum_{i} \prod_{i=1}^{n} \langle b_i^{(h,l)}, U_i \rangle \right)
$$
\n
$$
\geq \lim_{n \to \infty} \sup_{l} \frac{1}{n} \log \left(\frac{1}{l} \prod_{i=1}^{n} \langle b_i^{(h,l)}, U_i \rangle \right)
$$
\n
$$
\geq \sup_{l} \lim_{n \to \infty} \frac{1}{n} \left(\log \frac{1}{l} + \sum_{i=1}^{n} \log \langle b_n^{(h)}, U_i \rangle \right)
$$
\n
$$
\geq \lim_{l \to \infty} \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \log \langle b_n^{(h)}, U_i \rangle
$$
\n
$$
= \mathbb{E} \left(\log \langle b_{h+1}^*, U_1 \rangle | \mathcal{J} \right), \text{ a.s.}
$$
\n(5.11)

 $\big)$

which is the first needed inequality for the statement of the limit.

(Deriving the remaining inequality). Let a strategy $(b_n^{*(h)})$ be defined such that at each time *n*, it looks back at the past $(X_{n-h}^{n-1}, Y_{n-h}^n)$ and selects the corresponding optimal portfolio, i.e.,

$$
\mathbb{E}\big(\log\big\langle b_n^{*(h)},U_n\big\rangle|\sigma\big(X_{n-h}^{n-1},Y_{n-h}^n\big)\big)=\max_{b\in\mathscr{B}^m}\mathbb{E}\big(\log\big\langle b,U_n\big\rangle|\sigma\big(X_{n-h}^{n-1},Y_{n-h}^n\big)\big),\ \forall n.
$$

This can be chosen from the origin $n = 1$ as the market is a two-sided infinite process. Clearly, all such optimal portfolios $b_n^{*(h)}$ could be represented by a single measurable log-optimal portfolio $b_{h+1}^* \in \mathscr{F}_{h+1}^X$. Hence, applying Birkhoff's ergodic theorem yields the following result:

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b_n^{*(h)},U_i\right\rangle=\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b_{h+1}^*,U_i\right\rangle=\mathbb{E}\big(\log\left\langle b_{h+1}^*,U_1\right\rangle|\mathscr{I}\big),\text{ a.s.}
$$

Moreover, since all the combinatorial portfolios $b_n^{(h)}$ also depend on past information $(X_{n-h}^{n-1}, Y_{n-h}^n)$, they are suboptimal to the portfolios $b_n^{*(h)}$. We can derive the following inequality using the Kuhn-Tucker condition for log-optimality, similar to the proof of Lemma [2:](#page-7-0)

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b_n^{(h)},U_i\right\rangle\leq \lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b_n^{*(h)},U_i\right\rangle=\mathbb{E}\big(\log\left\langle b_{h+1}^*,U_1\right\rangle|\mathscr{I}\big),\text{ a.s.}
$$

which finally constitutes the needed limit. Furthermore, the L^1 -convergence follows immediately from the application of Lebesgue's dominated convergence theorem.

(Definitive statement of the optimality). Using the same argument as before, and noting that any constant strategy based on a single portfolio is measurable with respect to any \mathscr{F}_{h+1}^X for all orders *h*, let us consider the optimal constant strategy (b^*) and the log-optimal strategy (b^*_n) in a

stationary market. We then obtain the following inequalities due to Lemma [2:](#page-7-0)

$$
\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b^*,U_i\right\rangle\leq \lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b_n^{(h)},U_i\right\rangle\leq \lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n\log\left\langle b_n^*,U_i\right\rangle, \text{ a.s.}
$$

Since the terms associated with the strategies (b^*) and (b^*_n) are equal by Theorem [2,](#page-13-0) we have:

$$
\lim_{n\to\infty}\big(W_n(b_n^{(h)})-W_n(b^*)\big)=\lim_{n\to\infty}\big(W_n(b_n^{(h)})-W_n(b_n^*)\big)=0, \text{ a.s.}
$$

which establishes the optimality (also in L^1) of the strategy $(b_n^{(h)})$ and completes the proof. \Box

6 Summary and concluding remarks

In this paper, we investigate the problem of sequential portfolio decision-making in a market with partially observable side information and a latent dependence structure. The established results demonstrate the existence of a random optimal constant strategy, which is time-invariant and does not require knowledge of the dependence structure or observation of additional market information. This strategy achieves an asymptotic growth rate as fast as the log-optimal strategy, which dynamically determines portfolios based on perfect market information over time. The reason for this phenomenon lies in the stationarity of the market process, which diminishes the advantages of fully utilizing market information for dynamic strategies over time, leading to an equilibrium state. Furthermore, if the market is also ergodic, a non-random optimal constant strategy exists, regardless of the market's evolutionary possibilities. These findings question conventional perspectives in the fields of information theory, learning theory, and finance, where the prevailing belief is that an optimal strategy should be dynamic and utilize perfect information, rather than being a simple, time-invariant one. The traditional view maintains that a constant strategy is optimal only when the process of sole assets' returns is i.i.d.

With the established equality theorems for the limiting growth rate between the log-optimal and random constant strategies, we also broaden and enhance the optimality guarantees of two approaches for learning algorithms. The first approach involves algorithms that replicate the asymptotic growth rate of the random optimal constant strategy, learning solely from past assets' returns. These algorithms have the advantage of simplicity, as they avoid the difficulties of requiring knowledge of side market information and the latent dependence structure, relying instead on publicly available data. In contrast, the second approach involves algorithms that replicate the asymptotic growth rate of the log-optimal strategy, which necessitate knowledge of the dependence structure and complete past side market information for estimation. These algorithms encounter challenges such as high-dimensional market feature data and the inaccessibility of all relevant information. Fortunately, the existence of an optimal constant strategy enables this type of algorithm to bypass the need for side information and reduce the memory length of past events, while still guaranteeing optimality. However, this approach is less favorable due to its computational complexity.

Connection between frameworks. Another significant contribution of this paper is the potential connection it establishes between existing frameworks in the literature. First, the validity of the "*online learning portfolio*" framework, as surveyed in [van Erven et al.](#page-33-0) [\(2020](#page-33-0)), is considerably strengthened. This framework investigates learning algorithms that asymptotically approach the growth rate of the best retrospective constant portfolios (i.e., constant strategies) using only past assets' returns. Its justification in finance, originally stated in the pioneering works of [Cover](#page-32-4) [\(1991](#page-32-4), [1996\)](#page-32-8); [Cover and Ordentlich](#page-32-10) [\(1996](#page-32-10)); [Cover and Thomas](#page-32-12) [\(2006](#page-32-12)); [Ordentlich and Cover \(1996](#page-32-9)), is that these algorithms can asymptotically achieve the optimal limiting growth rate among all dynamic strategies if the market of sole assets' returns is i.i.d. Now, we extend this framework by demonstrating that it guarantees the optimal limiting growth rate among all dynamic strategies in a stationary market with partially observable side information and latent dependence structure, while still relying solely on past assets' returns. Additionally, the problem of learning side information, as studied in [Cover and Ordentlich \(1996\)](#page-32-10); [Bhatt et al.](#page-32-14) [\(2023](#page-32-14)), which is potentially unobservable and high-dimensional, is reduced to an application of the online learning portfolio framework. Finally, a connection with the framework of universal learning and prediction posed by [Algoet \(1992](#page-32-6)), utilizing mixing functionally generated portfolios, is established, at least under the stationary condition of the market process. Besides, it should be remarked further that the work by [Cuchiero et al. \(2019\)](#page-32-15) also provides a connection between the stochastic portfolio theory and the log-optimal portfolio theory, restricted to the 1-order Markov process of sole assets' returns.

References

- Algoet, P. (1992), 'Universal schemes for prediction, gambling, and portfolio selection', *Annals of Probability* 20, 901–941.
- Algoet, P. (1994), 'The strong law of large numbers for sequential decisions under uncertainity', *IEEE Transactions on Information Theory* 40, 609–634.
- Algoet, P. and Cover, T. (1988), 'Asymptotic optimality asymptotic equipartition properties of log-optimum investments', *Annals of Probability* 16, 876–898.
- Bailey, D. H. (1976), 'Sequential schemes for classifying and predicting ergodic processes', *Thesis, Dept. Mathematics, Stanford Univ.* .
- Bell, R. and Cover, T. (1980), 'Competitive optimality of logarithmic investment', *Mathematics of Operations Research* 5(2), 161–166.
- Bhatt, A., Ryu, J. J. and Kim, Y.-H. (2023), 'On universal portfolios with continuous side information', *Proceedings of Machine Learning Research* .
- Breiman, L. (1957), 'The ndividual ergodic theorem of information theory', *The Annals of Mathematical Statistics* 28(3), 809–811.
- Breiman, L. (1960), 'Investment policies for expanding businesses optimal in a long run sense', *Naval Research Logistics Quarterly* .
- Breiman, L. (1961), 'Optimal gambling systems for favorable games', *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability* .
- Cesa-Bianchi, N. and Lugosi, G. (2006), 'Prediction, learning, and games', *Cambridge University Press* .
- Cover, T. M. (1991), 'Universal portfolios', *Mathematical Finance* 1, 1–29.
- Cover, T. M. (1996), 'Universal data compression and portfolio selection', *Proceedings of the 37th IEEE Symposium on Foundations of Computer Science* .
- Cover, T. M. and Ordentlich, E. (1996), 'Universal portfolios with side information', *IEEE Transactions on Information Theory* 42, 348–363.
- Cover, T. M. and Thomas, J. A. (2006), 'Elements of Information Theory, 2nd edition', *John Wiley and Sons, New York* .
- Cuchiero, C., Schachermayer, W. and Wong, T.-K. L. (2019), 'Cover's universal portfolio, stochastic portfolio theory, and the numeraire portfolio', *Mathematical Finance* 29.
- Gyorfi, L., Lugosi, G. and Udina, F. (2006), 'Nonparametric kernel-based sequential investment strategies', *Mathematical Finance* 16, 337–357.
- Kelly, J. L. (1956), 'A new interpretation of information rate', *Bell Systems Technical Journal* 35, 917–926.
- Lam, D. K. (2024), 'Mean-variance portfolio selection in long-term investments with unknown distribution: online estimation, risk aversion under ambiguity, and universality of algorithms', *ArXiv preprint* .
- Morvai, G. (1991), 'Empirical log-optimal portfolio selection', *Problems of Control and Information Theory* 20, 453–463.
- Morvai, G. (1992), 'Portfolio choice based on the empirical distribution', *Kybernetika* 28.
- Ordentlich, E. and Cover, T. M. (1996), 'Online portfolio selection', *Proceedings of the 9th Annual Conference on Computational Learning Theory* .
- Ornstein, D. (1978), 'Guessing the next output of a stationary process', *Israel Journal of Mathematics* .

van Erven, T., van der Hoeven, D., Kotlowski, W. and Koolen, W. M. (2020), 'Open problem: Fast and optimal online portfolio selection', *Proceedings of Thirty Third Conference on Learning Theory* .